

A NON-LINEAR DIFFERENTIAL EQUATION FOR THE PERIODS OF ELLIPTIC SURFACES

N.I. SHEPHERD-BARRON

Abstract

Suppose that $f : X \rightarrow C$ is a general Jacobian elliptic surface over \mathbb{C} . Then the primitive cohomology $H_{prim}^{1,1}(X)$ has, up to a sign, a natural orthonormal basis $(\eta_i)_{i \in [1, N]}$ given by certain meromorphic 2-forms η_i of the second kind, one for each ramification point of the classifying morphism ϕ from C to the stack $\overline{\mathcal{E}\ell\ell}$ of generalized elliptic curves. (Here N is both $h_{prim}^{1,1}(X)$ and the number of moduli of X .) A choice of local co-ordinate on $\overline{\mathcal{E}\ell\ell}$ provides, via the branch locus of ϕ , an étale local co-ordinate system $(t_i)_{i \in [1, N]}$ on the stack $\mathcal{J}\mathcal{E}$ of Jacobian elliptic surfaces.

The main result here is that truncation of the Gauss–Manin connexion yields the system

$$\{\partial_i H = (\partial_i \eta_i \wedge \eta_i) H\}_{i \in [1, N]}$$

of non-linear pde satisfied by $H = [\eta_1, \dots, \eta_N]$, where $\partial_i = \partial/\partial t_i$ and the skew tensor $\partial_i \eta_i \wedge \eta_i$ of rank 2 is the ecliptic of η_i (the plane in which the particle η_i is instantaneously moving with respect to t_i). Moreover, after rigidification of the integral cohomology, H can be interpreted as providing a period map for these surfaces with values in the complex orthogonal group $O_N(\mathbb{C})$, and we prove a generic infinitesimal Torelli theorem for this map. For rational elliptic surfaces this can be calculated explicitly.

AMS classification: 14C34, 32G20.

1 Review of results on elliptic surfaces

Suppose that $f : X \rightarrow C$ is an elliptic surface over \mathbb{C} that is Jacobian (that is, there is a given section C_0 of f). Assume that all its singular fibres are irreducible nodal curves; then f is determined by the corresponding classifying morphism $\phi : C \rightarrow \overline{\mathcal{E}\ell\ell}$. This morphism is dominant and is unramified over the locus $j = \infty$. Let $Z = \text{Ram}_\phi = \sum (e_P - 1)P$ denote the ramification divisor, where e_P is the ramification index of ϕ at $P \in C$. For any point $P \in C$, put $f^{-1}(P) = E_P$.

Put

$$h = p_g(X) = \dim H^0(X, \Omega_X^2) = h^{2,0}(X), \quad q = q(X) = \dim H^0(X, \Omega_X^1) = h^{1,0}(X).$$

So $K_X \sim f^*(K_C + \phi^*M)$, where M is the bundle of weight one modular forms on $\overline{\mathcal{E}\ell\ell}$, and $\deg \phi^*M = h - q + 1 = \chi(X, \mathcal{O}_X) \geq 1$, so that $h \geq q$. In particular

$|K_X| = f^*|K_C + \phi^*M|$ and, by Riemann–Roch on C , has no base points if $h > q$. If $h = q$ then $\text{Bs } |K_C + \phi^*M|$ is either empty or consists of a single point Q and $\phi^*M \sim Q$.

In [SB2] we showed that the stack $\mathcal{J}\mathcal{E}$ of Jacobian elliptic surfaces is smooth of the expected dimension $N = 10h + 8(1 - q)$, provided that $10h > 12(q - 1)$, where h is the geometric genus of X and q its irregularity, and the tangent bundle $T_{\mathcal{J}\mathcal{E}}$ is naturally a line bundle on the universal ramification divisor \mathcal{Z} . That is, the tangent space $T_{\mathcal{J}\mathcal{E}}(X)$ is given by $T_{\mathcal{J}\mathcal{E}}(X) = \oplus_{P \in Z} L_P$ where L_P is a skyscraper sheaf supported at P and its length is $e_P - 1$.

In fact the assumption that $10h > 12(q - 1)$ is unnecessary for the smoothness of $\mathcal{J}\mathcal{E}$, although it does imply that the morphism $\mathcal{J}\mathcal{E} \rightarrow \mathcal{M}_q$ is smooth.

Theorem 1.1 *$\mathcal{J}\mathcal{E}$ is smooth of dimension N .*

PROOF: The exact sequence

$$0 \rightarrow T_C \rightarrow \phi^*T_{\mathcal{E}\ell\ell} \rightarrow \mathcal{L}_Z \rightarrow 0,$$

where \mathcal{L}_Z is a line bundle on Z , shows that $T_{\mathcal{J}\mathcal{E}}(X) = H^0(Z, \mathcal{L}_Z)$, which is of dimension N . On the other hand, for each point $a \in Z$ of ramification index $e_a = e$ we constructed in [SB2] a variation of X over an $(e - 1)$ -dimensional polydisc Δ^{e-1} whose tangent space is the part of \mathcal{L} that is supported at a . The product of these polydiscs is then the base of an N -dimensional variation of X whose tangent space is $H^0(Z, \mathcal{L}_Z)$. \square

We also showed in [SB2] that, if $|K_X|$ has no base points¹ then the derivative of the period map defines a linear map $L_P : H^{2,0}(X) \rightarrow H_{\text{prim}}^{1,1}(X)$ whose image is an $(e_P - 1)$ -dimensional subspace V_P generated by certain explicit meromorphic 2-forms of the second kind that lie in $H^0(X, \Omega_X^2(e_P E_P))$. Moreover, $H_{\text{prim}}^{1,1}(X)$ is the orthogonal direct sum of the V_P . In particular, if also Z is reduced then there is an explicit meromorphic 2-form $\eta_P \in H^0(X, \Omega_X^2(2E_P))_{2^{\text{nd kind}}}$ that defines a class $[\eta_P]$ in the primitive cohomology $H_{\text{prim}}^{1,1}(X)$ and the classes $[\eta_P]$ form an orthogonal basis of $H_{\text{prim}}^{1,1}(X)$. This basis can then be rescaled to be orthonormal, up to a single choice of sign. The proof of orthogonality was of a variational nature and is only valid when $|K_X|$ has no base points; we shall show below that this orthogonality holds generally. We shall also give a description of the derivative of the period map even when there are base points.

2 Some formulae

The notation will be the same as above.

¹In *loc. cit.* we stated this under the weaker assumption that $h > 0$. The argument depended on the truth of the infinitesimal Torelli theorem, which holds if $\text{Bs } |K_X|$ is empty [S], [I], [Kl]. So the results of [SB2] hold if $h > q$. However, if there are base points then things are more subtle; we shall discuss this below.

Proposition 2.1 Suppose that $P \in Z$ and that z_P is a co-ordinate on C at P such that $z_P^{e_P}$ is a co-ordinate ζ_P on $\overline{\mathcal{E}\ell\ell}$. Assume also that v_P is a fibre co-ordinate on X along E_P that pulls back from a fibre co-ordinate on the universal elliptic curve and that vanishes along the zero section C_0 of $f : X \rightarrow C$. Suppose that $\eta \in H^0(X, \Omega_X^2(\sum_P e_P E_P))$ and consider the local expansion

$$\eta = \sum_{j > -e_P} a_{P,j} z_P^{j-1} dz_P \wedge dv_P$$

at P . Then η is of the second kind if and only if $a_{P,0} = 0$ for every $P \in Z$.

PROOF: Drop the subscript P , so that $v_P = v$, $z_P = z$ and $e_P = e$. Then there is a neighbourhood Δ of P in C such that, if $X_\Delta = f^{-1}(\Delta)$, the morphism $X_\Delta \rightarrow \Delta$ is defined by

$$y^2 = 4x^3 - g_4x - g_6$$

where g_4, g_6 are functions of z^e and $x = \wp(v)$, $y = \wp'(v)$. Therefore there is an action of the group μ_e of e 'th roots of unity on X_Δ defined in terms of the local co-ordinates (z, v) by $\chi^*(z, v) = (\chi z, v)$ for the generator $\chi = \exp(2\pi i/e)$ of μ_e .

Suppose that η is of the second kind and let η_Δ denote its restriction to X_Δ . Then $\gamma^* \eta_\Delta$ is of the second kind for all $\gamma \in \mu_e$. Take $\gamma = \chi$; then

$$\sum_{r=0}^{e-1} \gamma^{r*} \eta_\Delta = e \sum_{k \geq 0} a_{ke} z^{ke} \frac{dz}{z} \wedge dv,$$

so that $\sum_{r=0}^{e-1} \gamma^{r*} \eta_\Delta$ is of the second kind and has a simple pole. Therefore it is holomorphic, so that $a_0 = 0$.

Conversely, suppose that $\eta \in H^0(X, \Omega_X^2(eE))$ and that $\text{Res}_E \eta \neq 0$. Since μ_e acts trivially on E , and so on $H^1(E, \mathbb{C})$, it follows that

$$\text{Res}_E(\gamma^* \eta_\Delta) = \gamma^* \text{Res}_E \eta = \text{Res}_E \eta,$$

so that $\text{Res}_E(\sum_r \gamma^{r*} \eta_\Delta) = e \text{Res}_E \eta \neq 0$. But

$$\sum_r \gamma^{r*} \eta_\Delta = e \sum_{k \geq 0} a_{ke} z^{ke} \frac{dz}{z} \wedge dv$$

and so has a simple pole; since its residue along E is non-zero, it follows that $a_0 \neq 0$. \square

We let the subscript *2nd kind* stand for “of the second kind”.

Proposition 2.2 For every $P \in Z$ we have

$$\dim H^0(X, \Omega_X^2(e_P P))_{2nd \text{ kind}} / H^0(X, \Omega_X^2) = e_P - 1.$$

PROOF: Recall first that we can identify the linear system $|K_X|$ with $|K_C + \phi^* M|$ via f^* .

The cohomology of the short exact sequence

$$0 \rightarrow \mathcal{O}_C(K_C + \phi^*M) \rightarrow \mathcal{O}_C(K_C + \phi^*M + e_P P) \rightarrow \mathcal{O}_{e_P P} \rightarrow 0$$

of sheaves on C shows that

$$\dim H^0(C, \mathcal{O}_C(K_C + \phi^*M + e_P P)) / H^0(C, \mathcal{O}_C(K_C + \phi^*M)) = e_P.$$

By the previous proposition there is exactly one linear condition on a member of $H^0(X, \Omega_X^2(e_P P))$ to be of the second kind. \square

So we can write down a basis of $H^{1,1}(X)_{\text{prim}}$ that is represented by 2-forms of the second kind. After rescaling we can take this basis to be orthonormal.

Theorem 2.3 *If P_1, P_2 are distinct points in Z and $\eta_i \in H^0(X, \Omega_X^2(e_{P_i} P_i))_{2\text{nd kind}}$ for $i = 1, 2$ then $(\eta_1, \eta_2) = 0$.*

PROOF: After pulling back by some finite cover $D \rightarrow C$ that is unramified over Z we get a surface $Y \rightarrow D$ defined by $\phi_1 : D \rightarrow \overline{\mathcal{E}\ell\ell}$ such that $\deg \phi_1^* M > 1$. Then $|K_Y|$ has no base points so that, by the results of [SB2], the pull backs of the classes η_i are orthogonal on Y . Then the η_i are orthogonal on X . \square

3 The differential equation

Definition 3.1 *A surface is general if the ramification divisor Z is reduced and disjoint from the locus $j = \infty$. We denote the corresponding open substack of $\mathcal{J}\mathcal{E}$ by $\mathcal{J}\mathcal{E}^{\text{gen}}$.*

Any choice of local co-ordinate on $\overline{\mathcal{E}\ell\ell}$ gives an étale local co-ordinate system (t_1, \dots, t_N) on $\mathcal{J}\mathcal{E}^{\text{gen}}$ via the étale morphism $\mathcal{J}\mathcal{E}^{\text{gen}} \rightarrow \overline{\mathcal{E}\ell\ell}^{(N)}$, the N -fold symmetric product of $\overline{\mathcal{E}\ell\ell}$, that maps the surface $X \rightarrow C$ to the branch locus in $\overline{\mathcal{E}\ell\ell}$ of the classifying morphism $\phi : C \rightarrow \overline{\mathcal{E}\ell\ell}$.

Define $\widetilde{\mathcal{J}\mathcal{E}} \rightarrow \mathcal{J}\mathcal{E}^{\text{gen}}$ to be the \mathfrak{S}_N cover induced by $\overline{\mathcal{E}\ell\ell}^N \rightarrow \overline{\mathcal{E}\ell\ell}^{(N)}$. Then (t_1, \dots, t_N) is a local co-ordinate system on $\widetilde{\mathcal{J}\mathcal{E}}$. (We shall see below that if h is not too small compared to q then $\mathcal{J}\mathcal{E}$ and $\widetilde{\mathcal{J}\mathcal{E}}$ are irreducible.)

Moreover the rank N vector bundle $\mathcal{H}_{\text{prim}}^{1,1}$ on $\mathcal{J}\mathcal{E}^{\text{gen}}$ has a non-degenerate inner product and [SB2] the pull back of $\mathcal{H}_{\text{prim}}^{1,1}$ to $\widetilde{\mathcal{J}\mathcal{E}}$ is equipped with an orthonormal basis (η_i) indexed by the ramification divisor. The line $\mathbb{C}\eta_i$ is the image of $H^0(X, \Omega_X^2)$ under $\partial_i = \partial/\partial t_i$, provided that $|K_X|$ has no base points (which is always the case if $h > q$) and the kernel of ∂_i is $H^0(X, \Omega_X^2(-E_i))$. Write

$$H = [\eta_1, \dots, \eta_N].$$

We need some further notation: if x, y are elements of an inner product space then $x \wedge y$ acts on a third element z by

$$(x \wedge y).z = (y, z)x - (x, z)y,$$

the contraction of $x \wedge y$ against z . We extend this to an action of $x \wedge y$ on a matrix H column by column. We can truncate the implicit linear system of pde on H_{prim}^2 that is given by the Gauss–Manin connexion to get an explicit non-linear system of pde that constrains $H_{prim}^{1,1}$.

Theorem 3.2 *H satisfies the system*

$$\{\partial_i H = (\partial_i \eta_i \wedge \eta_i) H\}_{i \in [1, N]}$$

of non-linear pde, where there is no summation over the repeated index.

PROOF: We prove this column by column.

We showed in [SB2] that $\partial_i \eta_j$ is proportional to η_i when $i \neq j$. That is,

$$\partial_i \eta_j = \psi_{ij} \eta_i$$

for some function ψ_{ij} . Taking inner products shows that $(\partial_i \eta_j, \eta_i) = \psi_{ij}$, so that

$$\partial_i \eta_j = (\partial_i \eta_j, \eta_i) \eta_i.$$

On the other hand

$$(\partial_i \eta_i \wedge \eta_i) \cdot \eta_j = (\eta_i, \eta_j) \partial_i \eta_i - (\partial_i \eta_i, \eta_j) \eta_i = -(\partial_i \eta_i, \eta_j) \eta_i.$$

But $(\eta_i, \eta_j) = 0$, which differentiates to

$$(\partial_i \eta_i, \eta_j) + (\eta_i, \partial_i \eta_j) = 0,$$

and then

$$(\partial_i \eta_i \wedge \eta_i) \cdot \eta_j = (\eta_i, \partial_i \eta_j) \eta_i = \partial_i \eta_j.$$

So the equation holds when $i \neq j$.

When $i = j$ the situation might appear to be more complex, because $\partial_i \eta_i$ no longer lies in Fil^1 of the Hodge filtration. However, in this case the equation merely states that $\partial_i \eta_i = \partial_i \eta_i$, so that it holds in full. \square

We shall refer to this as the ecliptic pde.

Assume for the rest of this section that $4(1 + h - q) \geq 2q + 1$.

Lemma 3.3 *$\mathcal{J}\mathcal{E}$ dominates \mathcal{M}_q and is irreducible.*

PROOF: Suppose $C \in \mathcal{M}_q$ and $L \in \text{Pic}_C^{1+h-q}$. Then $4L$ and $6L$ are very ample, so that a general member of $H^0(C, 4L) \times H^0(C, 6L)$ defines a surface in $\mathcal{J}\mathcal{E}_{h,q}$. Since the universal Pic^{1+h-q} over \mathcal{M}_q is irreducible the lemma is proved. \square

Theorem 3.4 *H is generically immersive.*

PROOF: We need two lemmas.

Lemma 3.5 *Suppose that $N \geq 8$ and that G is a transitive subgroup of the symmetric group \mathfrak{S}_N which contains a transposition and a copy of the alternating group \mathfrak{A}_r for some $r > N/2$.*

Then $G = \mathfrak{S}_N$.

PROOF: Consideration of the group generated by the transpositions in G shows that there are integers n, k such that $n \geq 2$, $nk = N$ and G is a wreath product $\mathfrak{S}_n \wr K$ for some transitive subgroup K of \mathfrak{S}_k . Since $r > N/2 \geq k$ the resulting homomorphism $\mathfrak{A}_r \rightarrow K$ is trivial. Then there is an embedding $\mathfrak{A}_r \hookrightarrow \mathfrak{S}_n$, which is impossible unless $k = 1$ and $n = N$. \square

Lemma 3.6 *The Galois group G of the universal ramification divisor $\mathcal{Z} \rightarrow \mathcal{J}\mathcal{E}$ is \mathfrak{S}_N .*

PROOF: According to Cor. 10.3 of [SB2] \mathcal{Z} is irreducible, because it is dominated by the universal $|4L| \times |6L|$, so that G is a transitive subgroup of \mathfrak{S}_N .

To see that G contains a transposition, take a surface in $\mathcal{J}\mathcal{E}_{h,q}$ defined by $\phi : C \rightarrow \overline{\mathcal{E}}\ell\ell$ that has a point of triple ramification. The corresponding 2-dimensional variation of X that is constructed in [SB2] yields a point where $\mathcal{Z} \rightarrow \mathcal{J}\mathcal{E}$ is simply ramified, and so G contains a transposition.

Since $|6L|$ is very ample, the Galois group of $|6L|$ is \mathfrak{S}_r , where $r = 6(h+1-q) - q$. Then there is an injective homomorphism $\mathfrak{S}_r \rightarrow G$. Since $r > N/2$, from our assumption, we conclude by the previous lemma. \square

Corollary 3.7 *$\widetilde{\mathcal{J}\mathcal{E}}$ is irreducible.*

PROOF: $\widetilde{\mathcal{J}\mathcal{E}} \rightarrow \mathcal{J}\mathcal{E}^{gen}$ is the Galois closure of $\mathcal{Z} \rightarrow \mathcal{J}\mathcal{E}$. \square

The Galois group $G = \mathfrak{S}_N$ simultaneously permutes the co-ordinate frame $\partial_1, \dots, \partial_N$ on $\widetilde{\mathcal{J}\mathcal{E}}$ and the vectors η_1, \dots, η_N . Therefore the kernel of H_* at the generic point of $\widetilde{\mathcal{J}\mathcal{E}}$ is preserved by G , so that if H_* is not injective at the generic point then there is a function f on $\overline{\mathcal{E}}\ell\ell$ such that either $\ker H_*$ contains $\sum_1^N f_i \partial_i$ or it contains $f_i \partial_i - f_j \partial_j$ for every i, j , where f_i is the pull back of f to \mathcal{Z} under the i 'th projection $\mathcal{Z} \rightarrow \overline{\mathcal{E}}\ell\ell$.

Suppose first that $\sum f_i \partial_i \in \ker H_*$. There is an action of \mathbb{G}_m on $\overline{\mathcal{E}}\ell\ell$ (it is the integral of the vector field $\sum \partial_i$) and so on $\mathcal{J}\mathcal{E}$. If X is given by the equation

$$Y^2 = 4X^3 - g_4X - g_6$$

where $g_n \in H^0(C, \phi^* M^{\otimes n})$ then the closure of the \mathbb{G}_m -orbit through the point X of $\mathcal{J}\mathcal{E}$ is the *Gauss-Eisenstein* pencil

$$(Y^2 = 4X^3 - \lambda g_4X - \mu g_6)_{(\lambda, \mu) \in \mathbb{P}^1} ;$$

at $\lambda = 0$ and at $\mu = 0$ the surface has constant j -invariant and complex multiplication by $\mathbb{Q}(\zeta_6)$ and $\mathbb{Q}(\zeta_4)$, respectively. Moreover, since this \mathbb{G}_m -action is defined over \mathbb{Q} , the corresponding vector field on $\mathcal{J}\mathcal{E}$ is invariant under the monodromy group \mathfrak{S}_N and so is proportional to $\sum f_i \partial_i$. Then H is constant on each

Gauss–Eisenstein pencil while the ramification divisor Z is constant. Therefore, in each such pencil, there is no monodromy on the classes $[\eta_i]$. However, suppose that Z meets the locus ($j = \infty$) in the point P_i , so that the surface acquires a node. Let $\delta \in H^{1,1}(X)_{\text{prim}}$ be the corresponding vanishing cycle. Then η_i and δ are both orthogonal to η_j for every $j \neq i$, so that η_i is proportional to δ . So η_i is not invariant under monodromy.

Suppose next that every $f_i \partial_i - f_j \partial_j \in \ker H_*$. Say $Z = \{P_1, \dots, P_N\}$, $E_i = f^{-1}(P_i)$ and $\eta_i \in H^0(X, \Omega_X^2(2E_i))_{2^{\text{nd}} \text{ kind}}$. Suppose $j \geq 3$. Then the variational formulae proved above give

$$\partial_i(\eta_j) = \eta_j(P_i)\eta_i$$

for $i = 1, 2$. Since $f_1 \partial_1(\eta_j) = f_2 \partial_2(\eta_j)$ it follows that η_j vanishes at both P_1 and P_2 . Therefore η_j vanishes at P_k for every $k \neq j$. However, $H^0(X, \Omega_X^2(2E_j - \sum_{k \neq j} E_k)) = 0$, and the result is proved. \square

4 Period matrices

By regarding the η_i as periods and H as a period matrix we get a multi-valued holomorphic map $H : \mathcal{J}\mathcal{E} \rightarrow O_N(\mathbb{C})$, the complex orthogonal group.

The primitive cohomology $H^2(X, \mathbb{Z})_{\text{prim}}$ of a surface X in $\mathcal{J}\mathcal{E}$ is even, unimodular and of signature $(2h, N)$, and so is isometric to $\tilde{\Lambda} := U^h \perp E_8(-1)^{1+h-q}$, where U is the hyperbolic plane and E_8 is the usual root lattice.

Say that a sublattice L of $\tilde{\Lambda}$ is good for X if it is isotropic of rank h and there is no line $\ell = \mathbb{C}\omega \subseteq H^{2,0}(X)$ such that $L \subseteq \ell^\perp$.

Lemma 4.1 *There exists a sublattice L of $\tilde{\Lambda}$ that is good for X .*

PROOF: Suppose not. The $O_\Lambda(\mathbb{Z})$ -orbit of L is Zariski-dense in the Grassmannian Gr of isotropic h -planes in $\tilde{\Lambda}_\mathbb{C}$, and so for every $V \in Gr$ there is a line $\ell \subset H^{2,0}(X)$ such that $V \subseteq \ell^\perp$. An elementary argument using the incidence variety of such pairs (V, ℓ) gives a contradiction. \square

Write $\tilde{\Lambda} = U_1 \perp \dots \perp U_h \perp \Lambda$, where $\Lambda = U^h \perp E_8(-1)^{1+h-q}$, and pick a standard basis (a_i, b_i) of U_i for $i \in [1, h]$. Put $L = \oplus \mathbb{Z}a_i$, an isotropic sublattice of $\tilde{\Lambda}$ of rank h . Recall that any two such sublattices are equivalent under $O_{\tilde{\Lambda}}(\mathbb{Z})$. Let $\mathbb{J}\mathcal{E} \rightarrow \widetilde{\mathcal{J}\mathcal{E}}$ be the stack of pairs (X, ϕ) where X is a surface in $\mathcal{J}\mathcal{E}$ and $\phi : \tilde{\Lambda} \rightarrow H^2(X, \mathbb{Z})_{\text{prim}}$ is an isometry and let $\mathbb{J}\mathcal{E}^0$ be the substack of $\mathbb{J}\mathcal{E}$ defined by the condition that $\phi(L)$ is good for X ; the map $\mathbb{J}\mathcal{E}^0 \rightarrow \widetilde{\mathcal{J}\mathcal{E}}$ is surjective, by Lemma 4.1. If $(X, \phi) \in \mathbb{J}\mathcal{E}$ then there is a normalized basis $(\omega_i)_{i \in [1, h]}$ of $H^{2,0}(X)$ defined by $\int_{a_i} \omega_j = \delta_{ij}$ and then the orthonormal basis $(\eta_i)_{i \in [1, N]}$ of $\text{Fil}^1 / \text{Fil}^2$ can be further normalized by the requirement that $\int_{a_i} \eta_j = 0$ for all i, j . This gives a holomorphic splitting of the projection $\text{Fil}^1 \rightarrow \text{Fil}^1 / \text{Fil}^2$ of vector bundles on $\mathbb{J}\mathcal{E}^0$.

If we write $\eta_j = \sum \alpha_{ji} a_i + \sum \beta_{ij} b_i + \lambda_j$ with $\lambda_j \in \Lambda$ then each $\beta_{ij} = 0$ and then $(\eta_i, \eta_j) = (\lambda_i, \lambda_j)$. Moreover, λ_j is the column vector

$${}^t \left[\int_{c_1} \eta_j, \dots, \int_{c_N} \eta_j \right]$$

where (c_i) is a \mathbb{Z} -basis of Λ ; that is, λ_j is a vector of periods and we can regard the matrix $H = [\eta_1, \dots, \eta_N]$ as a period matrix, obtained by integrating the vectors η_j over cycles dual to a basis of Λ . Now H is an $N \times N$ matrix with orthonormal columns; that is, H is a holomorphic morphism $\mathbb{J}\mathbb{E}^0 \rightarrow O_N(\mathbb{C})$ that satisfies the ecliptic pde.

Remark:

- (1) The assumptions of Theorem 3.4 are enough to ensure that the forgetful morphism $\mathcal{J}\mathcal{E}_{h,q}^{gen} \rightarrow \mathcal{M}_q$ is smooth and surjective. However, \mathcal{M}_q does not appear to have an algebraic local co-ordinate system like that on $\mathcal{J}\mathcal{E}^{gen}$; instead, the algebraic curve on $\mathcal{J}\mathcal{E}^{gen}$ defined by setting all but one co-ordinate to be constant projects to an algebraic curve on \mathcal{M}_q at every point of which the derivative of the period map is constant (as is seen from the nature of the variation that we used), so that \mathcal{M}_q is covered by such curves. Neither does there seem to be an analogous differential equation on \mathcal{M}_q .
- (2) (Torelli) Is H injective? Or generically injective?
- (3) (Schottky) Can the ecliptic equation be integrated to describe the image of H ? That is, can the co-ordinates t_i be eliminated to give equations that define the image of H ?

5 Infinitesimal Torelli with base points

As already mentioned, Saito [S] (see also [Kl]) proved infinitesimal Torelli when $\text{Bs } |K_X|$, which we identify with $\text{Bs } |K_C + \phi^*M|$, is empty. Since $\deg \phi^*M > 0$, if $\text{Bs } |K_X|$ is not empty then it consists of a single point $Q \in C$, where $\phi^*M = \mathcal{O}_C(Q)$, and also $h = q$.

Theorem 5.1 *Assume that $\text{Bs } |K_X| = \{Q\}$. Then infinitesimal Torelli holds for X if and only if $Q \notin Z$. If $Q \in Z$ then the derivative of the period map is of corank 1.*

PROOF: Suppose that $Q \in Z$ and put $e_Q = e$. Then consider the 1-parameter variation of X constructed from the glueing

$$z^e = w^e + etw^{e-2};$$

this is a sub-variation of the $(e-1)$ -parameter variation $\mathcal{X} \rightarrow \Delta^{e-1}$ detailed in [SB2]. Observe that

$$w = z(1 - tz^{-2})$$

modulo t^2 . Consider differentials $\omega^{(j)}(t)$ on \mathcal{X}_t normalized by the requirement that $\int_{A_i} \omega^{(j)}(t) = \delta_i^j$ for appropriate 2-cycles A_i on X supported away from the fibre E_Q . Expand $\omega^{(j)}(t)$ as

$$\omega^{(j)}(t) = \sum b_{pq}^{(j)} w^p t^q dw \wedge dv$$

for some fibre co-ordinate v . Substituting $w = z(1 - tz^{-2})$ gives

$$\omega^{(j)}(t) = \left(\sum b_{pq}^{(j)} z^p t^q - \sum b_{pq}^{(j)} (p-1) z^{p-2} t^{q+1} \right) dz \wedge dv$$

modulo t^2 . Write $\omega^{(j)}(t) = \omega^{(j)} + t\eta^{(j)}$; then $\omega^{(j)} = \sum b_{p0}^{(j)} z^p dz \wedge dv$ and $\eta^{(j)} = b_{00}^{(j)} z^{-2} dz \wedge dv$ modulo $H^0(X, \Omega_X^2)$. Now $b_{00}^{(j)}$ is the constant coefficient of $\omega^{(j)}$ and so vanishes (Q is the base point of $|K_X|$); it follows that the class $[\eta^{(j)}]$ in $H_{prim}^{1,1}(X)$ also vanishes for every j , so that $\nabla_{\partial/\partial t} : H^{2,0}(X) \rightarrow H_{prim}^{1,1}(X)$ is the zero map and indeed infinitesimal Torelli fails.

For each $a \in Z$ there is [SB2] a 1-parameter variation of X , with parameter t_a , such that the kernel of $\nabla_{\partial/\partial t_a}$ equals $H^0(X, \Omega_X^2(-E_a))$. So if $a = Q$ then $\nabla_{\partial/\partial t_a} = 0$ and infinitesimal Torelli fails. If $Q \notin Z$ then the image of $\nabla_{\partial/\partial t_a}$ is the line $\mathbb{C} \cdot \eta_a$ and it follows that no linear combination of the $\nabla_{\partial/\partial t_a}$ can vanish.

□

6 Intersecting forms with curves

We give a formula for the cup product $[\eta] \cup [D]$ when η is a 2-form of the second kind and D is a curve in X that is unramified over C at the points of the ramification divisor Z .

Recall that X is provided with a zero section C_0 as part of its data.

Theorem 6.1 *Suppose that $\eta \in H^0(X, \Omega_X^2(\sum_{P \in Z} e_P E_P))_{2^{nd} \text{ kind}}$ and D is as above. Assume that we have chosen local co-ordinates as in Proposition 2.1. Assume further that D is defined locally at each point Q of $D \cap E_P$ by*

$$v_P = v_P(z_P) = \sum_{m \geq 0} c_{Q,m} z_P^m$$

and that the corresponding local expansion of η is

$$\eta = \sum_{0 \neq n > -e_P} a_{P,n} z_P^{n-1} dz_P \wedge dv_P$$

for every $P \in Z$. Then

$$[\eta] \cup_{dR} [D] = - \sum_{P \in Z} \sum_{m=1}^{e_P-1} a_{P,-m} \sum_{Q \in D \cap E_P} c_{P,m}.$$

PROOF: Without loss of generality we can assume that the polar divisor $(\eta)_\infty$ is supported on a single fibre E_P .

Choose small open discs $P \in \Delta' \subsetneq \Delta \subsetneq C$ and a small tubular neighbourhood X^0 of D in X . Set $R = \overline{\Delta} - \Delta'$, a closed annulus, and let $\gamma = \partial \overline{\Delta}$, $\gamma' = \partial \overline{\Delta}'$. Then there is a C^∞ $(1,0)$ -form Ψ on X^0 such that

$$(1) \quad \Psi = 0 \text{ on } f^{-1}(C - \Delta) \cap X^0 \text{ and}$$

$$(2) \quad \Psi = \sum_{n \neq 0} \frac{a_n}{n} z^n dv \text{ on } f^{-1}(\Delta') \cap X^0, \text{ so that } d\Psi = \eta \text{ there.}$$

Put $\tilde{\xi} = \eta - d\Psi$, a C^∞ 2-form on X^0 . Then $[\eta]|_{X^0} = [\tilde{\xi}]$, $\tilde{\xi}$ is zero on $f^{-1}(\Delta') \cap X^0$ and $\tilde{\xi}$ is holomorphic on $f^{-1}(C - \Delta) \cap X^0$. So

$$[\eta] \cup_{dR} [D] = [\eta]|_{X^0} \cup_{dR} [D] = [\tilde{\xi}] \cup_{dR} [D] = \frac{1}{2\pi i} \int_D \tilde{\xi}|_D = \frac{1}{2\pi i} \int_{D \cap f^{-1}(R)} \tilde{\xi}|_D,$$

where the last equality follows from the facts that $\tilde{\xi}|_D$ is zero on $D \cap f^{-1}(\Delta')$ and is a holomorphic 2-form on $D \cap f^{-1}(C - \Delta)$, so is zero there. So

$$\begin{aligned} 2\pi i [\eta] \cup_{dR} [D] &= \int_{D \cap f^{-1}(R)} (\eta - d\Psi)|_D \\ &= \int_{D \cap f^{-1}(R)} \eta|_D - \int_{D \cap f^{-1}(R)} d\Psi|_D \\ &= 0 - \int_{D \cap f^{-1}(R)} d\Psi|_D \\ &= \int_{D \cap f^{-1}(\gamma')} \Psi|_D - \int_{D \cap f^{-1}(\gamma)} \Psi|_D \text{ (Stokes)} \\ &= \int_{D \cap f^{-1}(\gamma')} \Psi|_D \text{ since } \Psi = 0 \text{ on } f^{-1}(C - \Delta) \\ &= 2\pi i \sum_{Q \in D \cap E_P} \text{Res}_Q \Psi|_D \text{ (Cauchy)}. \end{aligned}$$

Now

$$\Psi = \sum_{n \neq 0} \frac{a_n}{n} z^n dv = \sum_{n \neq 0, m \geq 0} \frac{m}{n} c_{Q,m} a_n z^{m+n-1} dz,$$

so that

$$[\eta] \cup_{dR} [D] = \sum_Q \text{Res}_Q \Psi = \sum_Q \sum_{m \geq 1} \frac{m}{-m} c_{Q,m} a_{-m} = - \sum_{m=1}^{e_P-1} a_{-m} \sum_Q c_{Q,m}.$$

□

Corollary 6.2 *If $f : X \rightarrow C$ is written in Weierstrass form as*

$$y^2 = 4x^3 - g_4x - g_6,$$

if $\phi : C \rightarrow \overline{\mathcal{E}\ell\ell}$ is simply ramified and if every v_P is chosen so that $dv_P = dx/y$ then

$$[\eta] \cup_{dR} [D] = - \sum_{P \in Z} a_{P,-1} \frac{dx}{dz_P}(P) \sum_{Q \in D \cap E_P} \frac{1}{y(Q)}.$$

PROOF: Put $z_P = z$ and $v_P = v$. Then, in terms of a period $\tau = \tau(z)$ of E_z we have

$$x = \wp(v, \tau), \quad y = \frac{\partial \wp(v, \tau)}{\partial v},$$

so that

$$\frac{dx}{dz} = y \frac{dv}{dz} + \frac{\partial \wp}{\partial \tau} \frac{d\tau}{dz}.$$

Now $\frac{d\tau}{dz}|_{z=0} = 0$, since ϕ is ramified at P , so that

$$c_{Q,1} = \frac{dv}{dz}(Q) = \frac{1}{y(Q)} \frac{dx}{dz}(P).$$

□

Remark: (1) If the ramification is not simple then we can calculate the coefficients $c_{P,n}$ in a similar way via Faà di Bruno's formula. We omit the details, however.

(2) In particular, $[\eta] \cup [C_0] = 0$, where C_0 is the zero section. Since also $[\eta] \cup [f] = 0$ for a fibre f , it follows that $[\eta]$ lies in $H_{prim}^2(X, \mathbb{C})$, and so in $\text{Fil}^1 H_{prim}^2(X, \mathbb{C})$.

7 Rational surfaces

Here we illustrate our results in the context of rational surfaces.

Suppose that $f : X \rightarrow C = \mathbb{P}^1$ is a generic Jacobian elliptic surface which is rational, that C_0 is the given zero section and that ϕ is a fibre. Then there are eight further sections C_1, \dots, C_8 , disjoint from each other and from C_0 , and the curves C_0, C_1, \dots, C_8 are the exceptional curves of a birational contraction $\pi : X \rightarrow \mathbb{P}^2$. The lattice $H_{prim}^2(X, \mathbb{Z})$ is isomorphic to the root lattice $E_8(-1)$ and if h is the class of a line in \mathbb{P}^2 and $\alpha_1, \dots, \alpha_8$ are defined by

$$\alpha_1 = C_1 - C_2, \quad \alpha_2 = 2h - C_1 - C_2 - C_3 \quad \text{while} \quad \alpha_j = C_{j-1} - C_j \quad \text{for } j \geq 3$$

then $(\alpha_1, \dots, \alpha_8)$ is a root basis of $H_{prim}^2(X, \mathbb{Z})$, which is isomorphic to the root lattice $E_8(-1)$. Note that $\phi = 3h - \sum_0^8 C_i$.

We can calculate each intersection number $[\eta_i] \cup_{dR} [C_j]$, and therefore each number $[\eta_i] \cup_{dR} [\alpha_j]$. Then the period matrix $H(X)$ is the 8×8 matrix

$$H(X) = ([\eta_i] \cup_{dR} [\alpha_j])$$

and we can calculate it as follows.

- (1) Take a field K of characteristic zero and take eight points $Q_1, \dots, Q_8 \in \mathbb{P}^2(K)$. Let h denote the class of a line in \mathbb{P}^2 .
- (2) Check that these points are in general position in the usual sense: they are distinct; no three are collinear; no six lie on a conic; there is no cubic through all of them that is singular at one of them.
- (3) Calculate a basis (U_1, U_2) of $H^0(\mathbb{P}_K^2, \mathcal{O}(3h - \sum Q_i))$.
- (4) Construct a basis $(U_1^2, U_1 U_2, U_2^2, V)$ of $H^0(\mathbb{P}_K^2, \mathcal{O}(6h - 2 \sum Q_i))$.
- (5) Construct a basis $(U_1^3, U_1^2 U_2, \dots, V U_2, W)$ of $H^0(\mathbb{P}_K^2, \mathcal{O}(9h - 3 \sum Q_i))$.
- (6) The forms (U_1, U_2, V, W) define a rational map

$$\rho : \mathbb{P}^2 - \rightarrow \mathbb{P}(1, 1, 2, 3) = \text{Proj } K[U_1, U_2, V, W]$$

whose image is a surface Σ . This is the anticanonical model of the smooth del Pezzo surface $\text{Bl}_{Q_1, \dots, Q_8} \mathbb{P}^2$ of degree 1 and is embedded as a sextic surface in $\mathbb{P}(1, 1, 2, 3)$.

- (7) After calculating the images $\rho(x)$ for sufficiently many K -points $x \in \mathbb{P}^2$ (or we could calculate $\rho(\eta)$ where η is the generic point) we can interpolate through those points to find the defining sextic equation $F = 0$ of Σ .
- (8) Make an explicit change of co-ordinates so that $F = W^2 - (4V^3 - g_4 V - g_6)$ where $g_n = g_n(U_1, U_2)$ is homogeneous of degree n .
- (9) Let Q_9 denote the ninth base point of the pencil $|3h - \sum_1^8 Q_i|$ and put $X = \text{Bl}_{Q_9} \Sigma$. Let C_0 denote the exceptional curve of the blow-up; then C_0 is a section of the elliptic fibration $f : X \rightarrow \mathbb{P}^1$ defined by $f(Q) = (U_1(Q), U_2(Q))$.
- (10) If $j \neq k$ then the line $D_{jk} = h - Q_j - Q_k$ is a section of f . The isomorphism $D_{ij} \rightarrow \mathbb{P}^1$ induced by f can be calculated.
- (11) The ramification locus Z in \mathbb{P}^1 is defined by the vanishing of the Jacobian covariant

$$J = \det \begin{bmatrix} \partial g_4 / \partial U_1 & \partial g_4 / \partial U_2 \\ \partial g_6 / \partial U_1 & \partial g_6 / \partial U_2 \end{bmatrix}.$$

- (12) Check the genericity of the surface $f : X \rightarrow \mathbb{P}^1$ by calculating the discriminant $D = g_4^3 - 27g_6^2$ and then checking that the discriminants of D and J and the resultant of D and J are all non-zero. (Of course, this could be carried out by reduction modulo p .)
- (13) Now $K(Z)$ is a separable octic extension of K and $e_P = 2$ for all geometric points $P_i \in Z$. Calculate each form $\eta_i = \eta_{P_i}$, up to a scalar μ_i , by writing down sections of the two-dimensional vector space $H^0(X, \Omega_X^2(2P_i))$ and using Proposition 2.1.
- (14) Calculate each intersection number $[\eta_i] \cup_{dR} [D_{jk}]$ via Corollary 6.2 and then calculate each $[\eta_i] \cup \alpha_l$ from the description of α_l as a \mathbb{Q} -linear combination of the D_{jk} .
- (15) Calculate each $[\eta_i]$ as a linear combination of the α_l from the 8×8 matrix $([\eta_i] \cup \alpha_l)$.
- (16) We know that the η_i are orthogonal; we can now make them orthonormal. That is, we can determine the scalars μ_i .
- (17) We have now calculated $H(X)$ up to a single sign if $K(Z)$ is a field.

References

- [EGW] P. Engel, F. Greer and A. Ward, *Periods of elliptic surfaces with $p_g = q = 1$* , arXiv:2308.04563
- [I] A. Ikeda, *Bielliptic curves of genus three and the Torelli problem for certain elliptic surfaces*, Adv. Math. **349** (2019), 125–161.
- [Kl] R. Kloosterman, *Infinitesimal Torelli for elliptic surfaces revisited*, J. Pure and Applied Algebra **226** (2022).
- [S] Masa-Hiko Saito, *On the infinitesimal Torelli problem of elliptic surfaces*, J. Math. Kyoto Univ. **23** (1983), 441–460.
- [SB1] N.I. Shepherd-Barron, *Asymptotic period relations for Jacobian elliptic surfaces*, arXiv:1904.13344, Proc. LMS., to appear.
- [SB2] ———, *Generic Torelli and local Schottky theorems for Jacobian elliptic surfaces*, arXiv:2009.03633, Compositio Math., to appear.

KING'S COLLEGE, STRAND, LONDON WC2R 2LS, U.K.
E-mail address: Nicholas.Shepherd-Barron@kcl.ac.uk