A SURVEY ON THE GENERALIZED FERMAT EQUATION OF VARIOUS SIGNATURES OVER TOTALLY REAL FIELDS

SATYABRAT SAHOO

ABSTRACT. Following the famous proof of Fermat's Last Theorem by Andrew Wiles using the modularity of elliptic curves over \mathbb{Q} , significant developments have been made in the study of Diophantine equations using the modularity method. This article presents a survey of numerous results on the solutions of the generalized Fermat equation of signatures (p, p, p), (p, p, 2), (p, p, 3), and (r, r, p) over totally real number fields using the modularity method.

1. Introduction

The study of solutions to the Diophantine equations is one of the most attractive areas in number theory. The most prominent case is the Fermat equation $x^p + y^p = z^p$. In [Wil95], Wiles proved Fermat's Last Theorem and established that the Fermat equation $x^n + y^n = z^n$ has no non-trivial primitive integer solutions for integers $n \geq 3$. Consider the generalized Fermat equation,

$$Ax^p + By^q + Cz^r = 0$$
, where $A, B, C, p, q, r \in \mathbb{Z} \setminus \{0\}$ (1.1)

with gcd(A, B, C) = 1 and primes $p, q, r \ge 2$ with $\frac{1}{p} + \frac{1}{q} + \frac{1}{r} < 1$. We say (p, q, r) as the **signature** of the generalized Fermat equation (1.1). The following conjecture is known for the solutions of the equation (1.1) (see [DG95]).

Conjecture 1.1. Fix $A, B, C \in \mathbb{Z}$ with gcd(A, B, C) = 1. Then, over all choices of prime exponents p, q, r with $\frac{1}{p} + \frac{1}{q} + \frac{1}{r} < 1$, the the generalized Fermat equation (1.1) has only finitely many non-trivial primitive integer solutions.

Here, a solution $(a, b, c) \in \mathbb{Z}^3$ to the equation (1.1) is called non-trivial if $abc \neq 0$, and primitive if gcd(a, b, c) = 1. In [DG95], Darmon and Granville proved Conjecture 1.1 for fixed p, q, r. More precisely:

Theorem 1.2. ([DG95, Theorem 2]) For fixed integers $A, B, C \in \mathbb{Z} \setminus \{0\}$ and fixed primes $p, q, r \geq 2$ with $\frac{1}{p} + \frac{1}{q} + \frac{1}{r} < 1$, the equation (1.1) has only finitely many non-trivial primitive integer solutions.

Throughout this article, we denote K as a totally real number field and \mathcal{O}_K as the ring of integers of K. This article presents a summary of various results on the solutions of the generalized Fermat equation (1.1) with signatures (p, p, p), (p, p, 2), (p, p, 3) and (r, r, p) over K using the modularity method. For an elliptic curve E defined over K and a rational prime p, let $\bar{\rho}_{E,p}: G_K := \operatorname{Gal}(\bar{K}/K) \to \operatorname{Aut}(E[p]) \simeq \operatorname{GL}_2(\mathbb{F}_p)$ be the mod-p Galois representation of G_K induced by the action of G_K on the p-torsion points E[p] of E. For any Hilbert modular newform f over K of

Date : December 5, 2025.

²⁰²⁰ Mathematics Subject Classification. 11D41, 11G05, 11R80.

Key words and phrases. Diophantine equations, Modularity, Totally real fields.

parallel weight k, level \mathfrak{n} with coefficient field \mathbb{Q}_f and for any non-zero prime ideal ω of $\mathcal{O}_{\mathbb{Q}_f}$, let $\bar{\rho}_{f,\omega}: G_K \to \mathrm{GL}_2(\mathbb{F}_{\omega})$ be the residual Galois representation attached to f, ω .

1.1. Structure of the article. In §2, §3, §4 and §5, we give the literature for the solutions of the generalized Fermat equation (1.1) with signatures (p, p, p), (p, p, 2), (p, p, 3) and (r, r, p), respectively over K. In §6, we give the strategy of proof in the modular method. In §7, we define Hilbert modular forms. Finally in §8, we present a summary of numerous results regarding the modularity of elliptic curves over K.

2. SIGNATURE
$$(p, p, p)$$
 CASE

In this section, we discuss various results on the solutions of the generalized Fermat equation (1.1) of signature (p, p, p), specifically $Ax^p + By^p + Cz^p = 0$ over totally real number fields K.

2.1. **Over** \mathbb{Q} . For A = B = C = 1, Wiles first proved Fermat's Last Theorem. More precisely:

Theorem 2.1. [Wil95, Theorem 0.5] For every positive integer $n \ge 3$, the Fermat equation $x^n + y^n + z^n = 0$ has no non-trivial primitive integer solution.

Let $L \in \{3, 5, 7, 11, 13, 17, 19, 23, 29, 53, 59\}$ and $r \geq 0$ be an integer. In [Ser87], Serre proved that the equation $x^p + y^p + L^r z^p = 0$ of exponent $p \geq 11$ with $p \neq L$ has no non-trivial primitive integer solutions. Furthermore, Serre proved that if $L \geq 3$ is a prime number which is neither a Fermat nor a Mersenne prime, then there exists a constant $C_L > 0$ (depending on L) such that for all primes $p > C_L$, the equation $x^p + y^p + L^r z^p = 0$ with integer r > 0 has no non-trivial primitive integer solutions. In [DM97], Darmon and Merel proved that the equation $x^n + y^n + 2z^n = 0$ with $n \geq 3$ has no non-trivial primitive integer solutions. In [Rib97], Ribet demonstrated that the equation $x^p + y^p + 2^r z^p = 0$ has no non-trivial primitive integer solutions for $2 \leq r < p$. Finally, in [Kra97], Kraus examined the integer solutions of the generalized Fermat equation $Ax^p + By^p + Cz^p = 0$ for various choices of non-zero integers A, B, C.

2.2. Over K. Let K be a totally real number field and \mathcal{O}_K be the ring of integers of K. Consider the Diophantine equation

$$Ax^p + By^p + Cz^p = 0, (2.1)$$

of prime exponent $p \geq 5$, where $A, B, C \in \mathcal{O}_K \setminus \{0\}$.

Definition 2.2 (Trivial solution). We say a solution $(a,b,c) \in \mathcal{O}_K^3$ to the equation (2.1) is trivial, if abc = 0, otherwise non-trivial. Further, we call it primitive if $a\mathcal{O}_K + b\mathcal{O}_K + c\mathcal{O}_K = \mathcal{O}_K$

Definition 2.3. We say the Diophantine equation $Ax^p + By^p + Cz^p = 0$ (resp. $Ax^p + By^p + Cz^r = 0$, $r \in \{2,3\}$) has no asymptotic solution in a subset $S \subseteq \mathcal{O}_K^3$, if there exists a constant $V := V_{K,A,B,C}$ (depending on K,A,B,C) such that for primes p > V, the equation $Ax^p + By^p + Cz^p = 0$ (resp. $Ax^p + By^p + Cz^r = 0$, $r \in \{2,3\}$) has no non-trivial primitive solutions in S.

Let P be the set of all non-zero prime ideals of \mathcal{O}_K . For any subset $S \subseteq P$, let $\mathcal{O}_S := \{\alpha \in K : v_{\mathfrak{P}}(\alpha) \geq 0 \text{ for all } \mathfrak{P} \in P \setminus S\}$ be the ring of S-integers in K and $\mathcal{O}_S^* := \{\alpha \in K : v_{\mathfrak{P}}(\alpha) = 0 \text{ for all } \mathfrak{P} \in P \setminus S\}$. We call the elements of \mathcal{O}_S^* as the S-units. Let $S_K := \{\mathfrak{P} \in P : \mathfrak{P}|2\}$, $T_K := \{\mathfrak{P} \in S_K : \mathbf{f}(\mathfrak{P},2) = 1\}$ and $U_K := \{\mathfrak{P} \in S_K : (3, v_{\mathfrak{P}}(2)) = 1\}$, where $\mathbf{f}(\mathfrak{P},2)$ denotes the residual degree of the prime ideal \mathfrak{P} lying above 2. We now state the Eichler-Shimura conjecture.

Conjecture 2.4 (Eichler-Shimura). Let f be a Hilbert modular newform defined over K of parallel weight 2, level \mathfrak{n} and with coefficient field $\mathbb{Q}_f = \mathbb{Q}$. Then, there exists an elliptic curve E_f/K of conductor \mathfrak{n} having the same L-function as f.

In [Dar04], Darmon proved that if either $[K:\mathbb{Q}]$ is odd or there exists some prime ideal $\mathfrak{q} \in P$ such that $v_{\mathfrak{q}}(\mathfrak{n}) = 1$, then Conjecture 2.4 holds over K. We write (ES) for "either $[K:\mathbb{Q}]$ is odd or $T_K \neq \varphi$ or Conjecture 2.4 holds for K."

In [FS15], Freitas and Siksek first studied the asymptotic solution of the Fermat equation $x^p + y^p + z^p = 0$ over K. More precisely:

Theorem 2.5. [FS15, Theorem 3] Let K be a totally real number field satisfying (ES). Suppose, for every solution (λ, μ) to the S_K -unit equation

$$\lambda + \mu = 1, \ \lambda, \mu \in \mathcal{O}_{S_K}^*, \tag{2.2}$$

either

- (1) there exists some $\mathfrak{P} \in T_K$ that satisfies $\max\{|v_{\mathfrak{P}}(\lambda)|, |v_{\mathfrak{P}}(\mu)|\} \leq 4v_{\mathfrak{P}}(2)$; or
- (2) there exists some $\mathfrak{P} \in U_K$ that satisfies both $\max\{|v_{\mathfrak{P}}(\lambda)|, |v_{\mathfrak{P}}(\mu)|\} \leq 4v_{\mathfrak{P}}(2)$ and $v_{\mathfrak{P}}(\lambda\mu) \equiv v_{\mathfrak{P}}(2) \pmod{3}$.

Then the equation $x^p + y^p + z^p = 0$ has no asymptotic solution in \mathcal{O}_K^3 .

Moreover, in [FS15], the authors also proved that the equation $x^p + y^p + z^p = 0$ has no asymptotic solution in \mathcal{O}_K^3 with $K = \mathbb{Q}(\sqrt{d})$ for a subset of $d \geq 2$ with density $\frac{5}{6}$ among the set of square-free integers $d \geq 2$. In [Dec16], Deconick extended the work of [FS15] to the equation $Ax^p + By^p + Cz^p = 0$, where $A, B, C \in \mathcal{O}_K$ with ABC is odd (in the sense that $\mathfrak{P} \nmid ABC$, for all prime $\mathfrak{P} \in S_K$). In [KS24a], Kumar and Sahoo studied the asymptotic solution of the equation $x^p + 2^r y^p + z^p = 0$ over K, for $r \in \mathbb{N}$. Finally, in [Sah24], the author extended the result of [KS24a] to the generalized Fermat equation $Ax^p + By^p + Cz^p = 0$, where $A, B, C \in \mathcal{O}_K \setminus \{0\}$ with ABC is even (in the sense that $\mathfrak{P}|ABC$, for some prime $\mathfrak{P} \in S_K$) and also calculated the density of all square free integers $d \geq 2$ such that the equation $Ax^p + By^p + Cz^p = 0$ has no asymptotic solution over $K = \mathbb{Q}(\sqrt{d})$. Similar to Theorem 2.5, all these results depend on some explicit bounds on the solution of the S-unit equation.

2.3. Construction of Frey elliptic curve. For any non-trivial solution $(a, b, c) \in \mathcal{O}_K^3$ to the equation $Ax^p + By^p + Cz^p = 0$, the Frey elliptic curve is given by

$$E/K: Y^{2} = X(X - Aa^{p})(X + Bb^{p}),$$
(2.3)

with $c_4 = 2^4 (A^2 a^{2p} - BCb^p c^p)$, $\Delta_E = 2^4 A^2 B^2 C^2 (abc)^{2p}$ and $j_E = 2^8 \frac{(A^2 a^{2p} - BCb^p c^p)^3}{A^2 B^2 C^2 (abc)^{2p}}$, where Δ_E denotes as the discriminant of E and j_E denotes as the j-invariant of E.

4

3. Signature (p, p, 2) case

In this section, we discuss various results on the solutions of the generalized Fermat equation (1.1) of signature (p, p, 2), i.e., $Ax^p + By^p + Cz^2 = 0$ over totally real number fields K.

- 3.1. Over Q. In [DM97], Darmon and Merel first proved that the equation $x^n +$ $y^n = z^2$ with $n \ge 3$ has no non-trivial primitive integer solutions. In [Ivo03], Ivorra studied the integer solutions of the equation $x^2 = y^p + 2^r z^p$ and $2x^2 = y^p + 2^r z^p$ of exponent $p \geq 5$ with $0 \leq r < p$. In [Sik03], Siksek established that the only nontrivial primitive integer solutions to the equation $x^2 = y^p + 2^r z^p$ of exponent p > 5with $r \ge 2$ are when r = 3, $x = \pm 3$, y = z = 1. In [BS04], Bennet and Skinner studied the integer solutions of the equation $Ax^2 = By^p + Cz^p$ for various choices of non-zero integers A, B, C. Recently in [Caz24], Cazorla Garcia studied the integer solutions of the equation $Ax^2 = q^k y^{2n} + z^n$ with 2|z, for any fixed positive integers A, q, k with A is square-free and $q \geq 3$ is a prime.
- 3.2. Over K. In [IKO20], Işik, Kara, and Ozman first studied the asymptotic solution of the equation $x^p + y^p = z^2$, and proved that $x^p + y^p = z^2$ has no asymptotic solution $(a,b,c) \in \mathcal{O}_K^3$ with 2|b when the narrow class number $h_K^+ = 1$ and $T_K :=$ $\{\mathfrak{P} \in S_K : \mathbf{f}(\mathfrak{P}, 2) = 1\} \neq \varphi$. In [Moc22], Mocanu generalized the result of [IKO20] by replacing the assumption $h_K^+=1$ in [IKO20] with $2 \nmid h_K^+$. In [KS24a], Kumar and Sahoo relaxed the assumptions in [IKO20] and proved that $x^p+y^p=z^2$ has no asymptotic solution $(a,b,c) \in \mathcal{O}_K^3$ with 2|ab without assuming the conditions $h_K^+ = 1$ and $T_K \neq \varphi$. Finally in [KS24b], Kumar and Sahoo extended the work of [Moc22] and studied the asymptotic solutions of the equation $x^2 = By^p + Cz^p$ over K, where B is an odd integer and C is either an odd integer or 2^r for some $r \in \mathbb{N}$. In [KS24b], the authors also studied the asymptotic solution of the equation $2x^2 = By^p + 2^r z^p$ over K, where B is an odd integer and $r \in \mathbb{N}$.

3.3. Construction of Frey elliptic curve.

• For any non-trivial solution $(a, b, c) \in \mathcal{O}_K^3$ to the equation $x^2 = By^p + Cz^p$, the Frey elliptic curve $E := E_{a,b,c}$ is given by

$$E/K: Y^2 = X(X^2 + 2aX + Bb^p), (3.1)$$

with $c_4 = 2^4 (Bb^p + 4Cc^p)$, $\Delta_E = 2^6 (B^2C)(b^2c)^p$ and $j_E = 2^6 \frac{(Bb^p + 4Cc^p)^3}{B^2C(b^2c)^p}$. • For any non-trivial solution $(a, b, c) \in \mathcal{O}_K^3$ to the equation $2x^2 = By^p + Cz^p$,

the Frey curve $E := E_{a,b,c}$ is given by

$$E/K: Y^{2} = X(X^{2} - 4aX + 2Bb^{p}), (3.2)$$

with $c_4 = 2^5 (Bb^p + 2^{r+2}c^p)$, $\Delta_E = 2^{9+r}B^2(b^2c)^p$ and $j_E = 2^{6-r} \frac{(Bb^p + 2^{r+2}c^p)^3}{B^2(b^2c)^p}$.

4. Signature
$$(p, p, 3)$$
 case

In this section, we discuss various results on the solutions of the generalized Fermat equation (1.1) of signature (p, p, 3), i.e., $Ax^p + By^p + Cz^3 = 0$ over totally real number fields.

- 4.1. Over \mathbb{Q} . In [DM97], Darmon and Merel proved that the equation $x^n + y^n = z^3$ with $n \geq 3$ has no non-trivial primitive integer solutions. In [BVY04], Bennett, Vatsal, and Yadzdani studied the integer solutions of the generalized Fermat equation of signature (n, n, 3), i.e., $Ax^n + By^n = Cz^3$, for various choices of non-zero integers A, B, C.
- 4.2. **Over** K. In [Moc22], Mocanu first studied the asymptotic solution $(a, b, c) \in \mathcal{O}_K^3$ to the equation $x^p + y^p = z^3$ with 3|b over K. Recently in [SK25], Kumar and Sahoo extended the work of [Moc22] to the generalized Fermat equation of signature (p, p, 3), i.e., $Ax^p + By^p + Cz^3 = 0$ over K, where $A, B, C \in \mathcal{O}_K \setminus \{0\}$, and studied the asymptotic solution of the equation $Ax^p + By^p + Cz^3 = 0$ over K.
- 4.3. Construction of Frey elliptic curve. For any non-trivial solution $(a, b, c) \in \mathcal{O}_K^3$ to the equation $Ax^p + By^p = Cz^3$, the Frey elliptic curve is given by

$$E/K: Y^{2} + 3CcXY + C^{2}Bb^{p}Y = X^{3}, (4.1)$$

where $c_4 = 3^2 C^3 c (9Aa^p + Bb^p)$, $\Delta_E = 3^3 AB^3 C^8 (ab^3)^p$ and $j_E = 3^3 \frac{Cc^3 (9Aa^p + Bb^p)^3}{AB^3 (ab^3)^p}$.

5. SIGNATURE
$$(r, r, p)$$
 CASE

In this section, we discuss various results on the solutions of the Fermat equation of signature (r, r, p), i.e.,

$$x^r + y^r = dz^p, (5.1)$$

over K, where $r, p \geq 5$ are rational primes and $d \in \mathcal{O}_K \setminus \{0\}$.

5.1. Over \mathbb{Q} . Throughout this subsection, we fix $d \in \mathbb{Z} \setminus \{0\}$. For r=5, the integer solutions of the equation (5.1) were initially studied by Billerey in [Bil07] and subsequently in [BD10], [DF14], [BCDF19]. In [Fre15], Freitas established a multi-Frey family of elliptic curves to study the integer solutions of the equation (5.1) and proved that the equation (5.1) (with r=7, d=3) has no non-trivial primitive integer solutions for primes $p > (1+3^{18})^2$. In [BCDF25b], Billerey et al. proved that the equation $x^7 + y^7 = 3z^n$ (resp. $x^7 + y^7 = z^n$) with integers $n \geq 2$ has non-trivial primitive integer solution (a,b,c) (resp. (a,b,c) with 2|a+b or 7|a+b). In [BCDF25a], Billerey et al. examined the integer solutions of the equation $x^{11} + y^{11} = z^n$ with integers $n \geq 2$ using Frey abelian varieties. The integer solutions of the equation (5.1) for r=13 were first studied in [DF13]. In [BCDF19] (resp. [BCDDF23]), the authors proved the equation (5.1) with r=13, d=3, and $p \neq 7$ (resp. for all p) has no non-trivial primitive integer solution.

In [FN24], Freitas and Najman proved that for a set of primes p with positive density and for fixed integers r, d with $r, p \nmid d$, the equation (5.1) has no non-trivial primitive integer solution (a, b, c) of the form 2|a + b or r|a + b. Recently in [KMO24], Kara, Mocanu and Özman studied the asymptotic integer solutions of the equation (5.1) by assuming Conjecture 2.4 and the weak Frey-Mazur conjecture (see [KMO24, Conjecture 3.11]).

5.2. **Over K.** For any prime $r \geq 5$, let ζ_r be a primitive rth root of unity in \mathbb{C} and $K^+ := K(\zeta_r + \zeta_r^{-1})$.

Definition 5.1 (Trivial solution). We say a solution $(a,b,c) \in \mathcal{O}_{K^+}^3$ to the equation (5.1) is trivial, if abc = 0, otherwise non-trivial. Further, we call it primitive if $a\mathcal{O}_{K^+} + b\mathcal{O}_{K^+} + c\mathcal{O}_{K^+} = \mathcal{O}_{K^+}$

6

Definition 5.2. We say the Diophantine equation $x^r + y^r = dz^p$ has no asymptotic solution in a subset $S \subseteq \mathcal{O}_{K^+}^3$ if there exists a constant $V := V_{K,r,d}$ (depending on K,r,d) such that for primes p > V, the equation $x^r + y^r = dz^p$ has no non-trivial primitive solutions in S.

In [Moc23], Mocanu first studied the asymptotic solution $(a, b, c) \in \mathcal{O}_{K^+}^3$ of the equation $x^r + y^r = z^p$ with 2|c. In [JS25], Jha and Sahoo extended the work of [Moc23] to the equation (5.1), and studied the asymptotic solution $(a, b, c) \in \mathcal{O}_{K^+}^3$ of the equation $x^r + y^r = dz^p$ with 2|c (resp. 2|a + b). In [JS25], the authors also studied the asymptotic solution $(a, b, c) \in \mathcal{O}_K^3$ of the equation $x^5 + y^5 = dz^p$ with $2 \nmid c$.

5.3. Construction of Frey elliptic curve for $x^r + y^r = dz^p$ over K^+ . In this subsection, we recall the Frey elliptic curve associated to any non-trivial primitive solution to the Diophantine equation $x^r + y^r = dz^p$ (see [Fre15] for more details). Let $L := K(\zeta_r)$. For any prime $r \geq 5$, choose

$$\phi_r(x,y) := \frac{x^r + y^r}{x + y} = \sum_{i=0}^{r-1} (-1)^i x^{r-1-i} y^i.$$

The factorization of the polynomial $\phi_r(x,y)$ over L is given by

$$\phi_r(x,y) = \prod_{i=1}^{r-1} (x + \zeta_r^i y).$$

For any k with $0 \le k \le \frac{r-1}{2}$, choose

$$f_k(x,y) := (x + \zeta_r^k y)(x + \zeta_r^{-k} y) = x^2 + (\zeta_r^k + \zeta_r^{-k})xy + y^2.$$

Then $f_k(x,y) \in K^+[x,y]$ for all k. The factorization of the polynomial $\phi_r(x,y)$ over K^+ is given by

$$\phi_r(x,y) = \prod_{i=1}^{\frac{r-1}{2}} f_k(x,y).$$

Since $r \ge 5$, we get $\frac{r-1}{2} \ge 2$. Fix three integers k_1, k_2, k_3 such that $0 \le k_1 < k_2 < k_3 \le \frac{r-1}{2}$. Let

$$\alpha := \zeta_r^{k_3} + \zeta_r^{-k_3} - \zeta_r^{k_2} - \zeta_r^{-k_2}, \beta := \zeta_r^{k_1} + \zeta_r^{-k_1} - \zeta_r^{k_3} - \zeta_r^{-k_3}, \gamma := \zeta_r^{k_2} + \zeta_r^{-k_2} - \zeta_r^{k_1} - \zeta_r^{-k_1}.$$

Then $\alpha, \beta, \gamma \in K^+$ and $\alpha f_{k_1} + \beta f_{k_2} + \gamma f_{k_3} = 0$. Let $A(x,y) := \alpha f_{k_1}(x,y), B(x,y) := \beta f_{k_2}(x,y)$, and $C(x,y) := \gamma f_{k_3}(x,y)$. Hence $A(x,y), B(x,y), C(x,y) \in K^+[x,y]$. Let $(a,b,c) \in \mathcal{O}_{K^+}^3$ be a non-trivial primitive solution of equation (5.1). Let A := A(a,b), B := B(a,b) and C := C(a,b). Then $A,B,C \in K^+$ and A+B+C=0. So the Frey elliptic curve is given by

$$E/K^{+}: Y^{2} = X(X - A)(X + B), \tag{5.2}$$

where $\Delta_E = 2^4 (ABC)^2$, $c_4 = 2^4 (AB + BC + CA)$ and $j_E = 2^8 \frac{(AB + BC + CA)^3}{(ABC)^2}$. The Frey curve in equation (5.2) depends on p since A, B, C depend on p.

5.4. Construction of Frey elliptic curve for $x^5+y^5=dz^p$ over K. For any non-trivial primitive solution $(a,b,c)\in\mathcal{O}_K^3$ to the Diophantine equation $x^5+y^5=dz^p$, the Frey elliptic curve is given by

$$E/K: y^2 = x^3 - 5(a^2 + b^2)x^2 + 5\phi_5(a, b)x,$$
(5.3)

where $\phi_5(a,b) := \frac{a^5 + b^5}{a + b}$, $c_4 = 2^4 5(5(a^2 + b^2)^2 - 3\phi_5(a,b))$, $\Delta_E = 2^4 5^3 (a + b)^2 (a^5 + b^5)^2$ and $j_E = 2^8 \frac{(5(a^2 + b^2)^2 - 3\phi_5(a,b))^3}{(a + b)^2 (a^5 + b^5)^2}$ (see [Bil07, §2] for the Frey curve).

6. Strategy of the proof

In this section, we provide a brief strategy of the proof in the modular method for solving the generalized Fermat equation over totally real fields K. The following are the key steps in the modular method:

- Step 1: For any non-trivial solution $(a, b, c) \in \mathcal{O}_K^3$ to the equation (1.1) of signature (p, p, p) or (p, p, 2) or (p, p, 3) or (r, r, p), associate a suitable **Frey elliptic** curve E/K.
- Step 2: Then prove the **modularity** of E/K for primes $p \gg 0$, E/K has **semi-stable reduction** (i.e., either good or multiplicative reduction) at all primes $\mathfrak{q}|p$, and the mod-p Galois representation $\bar{\rho}_{E,p}$ is **irreducible** for $p \gg 0$.
- Step 3: Using **level lowering results** of $\bar{\rho}_{E,p}$, we have $\bar{\rho}_{E,p} \sim \bar{\rho}_{f,p}$, for some Hilbert modular newform f of parallel weight 2 with rational eigenvalues of lower level \mathfrak{n}_p which strictly divides the conductor of E.
- Step 4: Finally to get a **contradiction**, prove that the finitely many Hilbert modular newforms of level \mathfrak{n}_p that occur in Step 3 do not correspond to $\bar{\rho}_{E,p}$ which contradicts $\bar{\rho}_{E,p} \sim \bar{\rho}_{f,p}$.

We now explain the proof of Theorem 2.5 briefly in terms of the above steps of the modular method.

• For any non-trivial solution $(a, b, c) \in \mathcal{O}_K^3$ to the equation $x^p + y^p + z^p = 0$, the Frey elliptic curve E/K is given by

$$E/K: Y^2 = X(X - a^p)(X + b^p).$$

- By [FS15, Corollary 2.1], the Frey curve E/K is modular for $p \gg 0$. By [FS15, Lemma 3.3], the Frey curve E/K is minimal and semistable at all prime ideals $\mathfrak{q}|p$. By [FS15, Theorem 6], we get $\bar{\rho}_{E,p}$ is irreducible for $p \gg 0$.
- Then using the level-loweing result, i.e., [FS15, Theorem 7], we get $\bar{\rho}_{E,p} \sim \bar{\rho}_{f,p}$, for some Hilbert modular newform f of parallel weight 2 with lower level \mathfrak{n}_p which strictly divides the conductor of E.
- We now explain the step 4 briefly. First, by using [FS15, Lemma 3.7], we conclude that the Frey curve E/K has potential multiplicative reduction at all primes $\mathfrak{P} \in T_K$, i.e., $v_{\mathfrak{P}}(j_E) < 0$ and $p | \# \bar{\rho}_{E,p}(I_{\mathfrak{P}})$ for $p > 4v_{\mathfrak{P}}(2)$. Next, by using [FS15, Theorem 7], there exists a constant $V_K > 0$ (depending on K) and an elliptic curve E' defined over K such that E'/K has good reduction away from S_K , E'/K has full 2-torsion points, $\bar{\rho}_{E,p} \sim \bar{\rho}_{E',p}$ for primes $p \geq V_K$.

Then, by using an image of inertia result, i.e., [FS15, Lemma 3.4], we conclude that E' has potential multiplicative reduction at all primes $\mathfrak{P} \in T_K$, i.e., $v_{\mathfrak{P}}(j_{E'}) < 0$ for all primes $\mathfrak{P} \in T_K$. Next, we express the j-invariant

 $j_{E'}$ of E' in terms of the solution (λ, μ) to the S_K -unit equation (2.2) by using the above properties of E'. Finally by using the explicit bound $\max\{|v_{\mathfrak{P}}(\lambda)|,|v_{\mathfrak{P}}(\mu)|\} \leq 4v_{\mathfrak{P}}(2)$, we get $v_{\mathfrak{P}}(j_{E'}) \geq 0$ for some $\mathfrak{P} \in T_K$, which is a contradiction (see [FS15, §5] for more details). A similar calculation also holds for primes $\mathfrak{P} \in U_K$.

7. Hilbert modular forms

In this section, we define Hilbert modular forms. Let F be a totally real field of degree n over \mathbb{Q} and let \mathcal{O}_F be the ring of integers of F. Let $k = (k_1, \ldots, k_n) \in \mathbb{Z}_{>0}^n$, and let $\mathbb{H} := \{z \in \mathbb{C} : \operatorname{im}(z) > 0\}$ be the upper half-plane. Let \mathfrak{D} be the absolute different of \mathcal{O}_F . For any non-zero prime ideal \mathfrak{p} of \mathcal{O}_F , let $F_{\mathfrak{p}}$ be the completion of F at \mathfrak{p} . For any ideals \mathfrak{a} and \mathfrak{b} of \mathcal{O}_F , we put

$$K_{\mathfrak{p}}(\mathfrak{a},\mathfrak{b}) := \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathrm{GL}_{2}(F_{\mathfrak{p}}) : \begin{matrix} a \in \mathcal{O}_{\mathfrak{p}}, & b \in \mathfrak{a}_{\mathfrak{p}}^{-1}\mathfrak{D}_{\mathfrak{p}}^{-1}, \\ c \in \mathfrak{b}_{\mathfrak{p}}\mathfrak{D}_{\mathfrak{p}}, & d \in \mathcal{O}_{\mathfrak{p}}, & |ad - bc|_{\mathfrak{p}} = 1 \end{matrix} \right\},$$

where $\mathfrak{m}_{\mathfrak{p}} = \mathfrak{m}\mathcal{O}_{\mathfrak{p}}$ for any ideal \mathfrak{m} of \mathcal{O}_F and $\mathfrak{D}_{\mathfrak{p}} = \mathfrak{D}\mathcal{O}_{\mathfrak{p}}$. Then $K_{\mathfrak{p}}(\mathfrak{a}, \mathfrak{b})$ is a subgroup of $\mathrm{GL}_2(F_{\mathfrak{p}})$. Furthermore, we put

$$K_0(\mathfrak{a},\mathfrak{b}) = \mathrm{SO}(2)^n \cdot \prod_{\mathfrak{b} < \infty} K_{\mathfrak{p}}(\mathfrak{a},\mathfrak{b}), \quad \mathrm{and} \quad W(\mathfrak{a},\mathfrak{b}) = \mathrm{GL}_2^+(\mathbb{R})^n K_0(\mathfrak{a},\mathfrak{b}).$$

In particular, if $\mathfrak{a} = \mathcal{O}_F$, we simply write $K_{\mathfrak{p}}(\mathfrak{b}) = K_{\mathfrak{p}}(\mathcal{O}_F, \mathfrak{b})$ and $W(\mathfrak{b}) = W(\mathcal{O}_F, \mathfrak{b})$. Let \mathbb{A}_F be the adèle ring of F. Let h be the narrow class number of F, and let $\{t_{\nu}\}_{\nu=1}^{h}$ taken to be a complete set of representatives of the narrow class group of F, where the archimedean part $t_{\nu,\infty}$ of t_{ν} is 1 for all ν . Then, we have the following disjoint decomposition of $GL_2(\mathbb{A}_F)$

$$GL_2(\mathbb{A}_F) = \bigcup_{\nu=1}^h GL_2(F) x_{\nu}^{-\iota} W(\mathfrak{b}), \tag{7.1}$$

where $x_{\nu}^{-\iota} = \begin{bmatrix} t_{\nu}^{-1} \\ 1 \end{bmatrix}$. For each ν , we define a congruence subgroup of $\mathrm{GL}_2(F)$ as follows:

$$\Gamma_{\nu}(\mathfrak{a},\mathfrak{b}) := \operatorname{GL}_{2}(F) \cap x_{\nu}W(\mathfrak{a},\mathfrak{b})x_{\nu}^{-1}$$

$$= \left\{ \begin{bmatrix} a & t_{\nu}^{-1}b \\ t_{\nu}c & d \end{bmatrix} \in \operatorname{GL}_{2}(F) : \begin{array}{l} a \in \mathcal{O}_{F}, & b \in \mathfrak{a}^{-1}\mathfrak{D}_{F}^{-1}, \\ c \in \mathfrak{b}\mathfrak{D}_{F}, & d \in \mathcal{O}_{F}, \end{array} \right. \quad ad - bc \in \mathcal{O}_{F}^{\times} \right\}.$$

In particular, if $\mathfrak{a} = \mathcal{O}_F$, we write $\Gamma_{\nu}(\mathfrak{b}) = \Gamma_{\nu}(\mathcal{O}_F, \mathfrak{b})$ as before.

Let $z = (z_1, \ldots, z_n) \in \mathbb{H}^n$. For a holomorphic function f on \mathbb{H}^n and an element $\gamma = (\gamma_1, \ldots, \gamma_n) \in \mathrm{GL}_2(\mathbb{R})^n$, define

$$f||_k \gamma(z) := \prod_{i=1}^n \det(\gamma_i)^{\frac{k_i}{2}} j(\gamma_i, z_i)^{-k_i} f(\gamma z),$$

where $\gamma z = (\gamma_1 z_1, \dots, \gamma_n z_n)$, $j(\gamma_i, z_i) = c_i z_i + d_i$ and $\gamma_i z_i = \frac{a_i z_i + b_i}{c_i z_i + d_i}$ for $\gamma_i = \begin{pmatrix} a_i & b_i \\ c_i & d_i \end{pmatrix}$.

Let $M_k(\Gamma_{\nu}(\mathfrak{b}))$ denote the space of all functions f_{ν} that are holomorphic on \mathbb{H}^n as well as at cusps and satisfying

$$|f_{\nu}||_{k}\gamma = f_{\nu}$$

for all γ in $\Gamma_{\nu}(\mathfrak{b})$. We note that such a function f_{ν} has a Fourier expansion

$$f_{\nu}(z) = \sum_{\xi \in F} a_{\nu}(\xi) \exp(2\pi i \xi z) \tag{7.2}$$

where ξ runs over all the totally positive elements in $t_{\nu}^{-1}\mathcal{O}_{F}$ and $\xi = 0$. We call $M_{k}(\Gamma_{\nu}(\mathfrak{b}))$ as the space of all Hilbert modular forms of weight k and level $\Gamma_{\nu}(\mathfrak{b})$. A Hilbert modular form $f_{\nu} \in M_{k}(\Gamma_{\nu}(\mathfrak{b}))$ is a cusp form if the constant term of $f_{\nu}|_{k}\gamma$ in its Fourier expansion is 0 for all $\gamma \in \mathrm{GL}_{2}^{+}(F)$, and the space of such cusp forms f_{ν} with respect to $\Gamma_{\nu}(\mathfrak{b})$ is denoted by $S_{k}(\Gamma_{\nu}(\mathfrak{b}))$.

Let $M_k(\mathfrak{b})$ denote the space of all functions $f := (f_1, \ldots, f_h)$ where $f_{\nu} \in M_k(\Gamma_{\nu}(\mathfrak{b}))$ for each $\nu \in \{1, \ldots, h\}$ and satisfying

$$f(g) = f(\gamma x_{\nu}^{-\iota} w) = (f_{\nu}||_{k} w_{\infty})(\mathbf{i})$$

where $g = \gamma x_{\nu}^{-\iota} w \in GL_2(F) x_{\nu}^{-\iota} W(\mathfrak{b})$ as in equation (7.1) and $\mathbf{i} = (i, \dots, i)$. We call $M_k(\mathfrak{b})$ the space of all Hilbert modular forms of weight k and level \mathfrak{b} . In particular, if $f_{\nu} \in S_k(\Gamma_{\nu}(\mathfrak{b}))$ for all ν , then the space of such f is denoted by $S_k(\mathfrak{b})$ and called the space of all Hilbert modular cusp forms of weight k and level \mathfrak{b} .

Let $\sigma_1, \ldots \sigma_n$ be the embeddings of F into \mathbb{R} . For any $m := (m_1, \ldots, m_n) \in \mathbb{Z}^n$ and $\alpha \in F$, we write $\alpha^m = \prod_{i=1}^n (\sigma_i(\alpha))^{m_i}$. Let \mathfrak{m} be an ideal of \mathcal{O}_F and write $\mathfrak{m} = \xi t_{\nu}^{-1} \mathcal{O}_F$ with a totally positive element ξ in F. We define the Fourier coefficients of f as follows:

$$C(\mathfrak{m}, f) := \begin{cases} N_{F/\mathbb{Q}}(\mathfrak{m})^{k_0/2} \xi^{-k/2} a_{\nu}(\xi) & \text{if } \mathfrak{m} = \xi t_{\nu}^{-1} \mathcal{O}_F \subset \mathcal{O}_F, \\ 0 & \text{if } \mathfrak{m} \text{ is not integral} \end{cases}$$
(7.3)

where $k_0 = \max\{k_1, \dots, k_n\}$ and $a_{\nu}(\xi)$ as in equation (7.2).

We denote $S_k^{\text{new}}(\mathfrak{n})$ for the space of all Hilbert modular newforms over F of weight k and level \mathfrak{n} . We say $f \in S_k^{\text{new}}(\mathfrak{n})$ is normalized if $C(\mathcal{O}_F, f) = 1$. A primitive Hilbert modular newform is a normalized Hecke eigenform in $S_k^{\text{new}}(\mathfrak{n})$ (see [Shi78] for more details).

8. Modularity of elliptic curves over K

In this section, we recall various results on the modularity of elliptic curves over totally real number fields. We first recall the definition of the modularity of an elliptic curve. To do this, we first review the definitions of the L-function attached to an elliptic curve and a Hilbert modular form.

Definition 8.1 (L-function attached to an elliptic curve). Let K be a number field and E/K be an elliptic curve. Then, the L-function attached to the elliptic curve E is defined by the Euler product

$$L(E,s) = \prod_{\mathfrak{p}} L_{\mathfrak{p}} \left(N_{K/\mathbb{Q}}(\mathfrak{p})^{-s} \right)$$

for $s \in \mathbb{C}$, where \mathfrak{p} runs over all prime ideals of \mathcal{O}_K ,

$$L_{\mathfrak{p}}(T)^{-1} = \begin{cases} 1 - a_{\mathfrak{p}}T + N_{K/\mathbb{Q}}(\mathfrak{p})T^2 & \text{if } E \text{ has good reduction at } \mathfrak{p}, \\ 1 - a_{\mathfrak{p}}T & \text{if } E \text{ has bad reduction at } \mathfrak{p}, \end{cases}$$

and

$$a_{\mathfrak{p}} = \begin{cases} N_{K/\mathbb{Q}}(\mathfrak{p}) + 1 - \#E(\mathbb{F}_{\mathfrak{p}}) & \text{if E has good reduction at \mathfrak{p},} \\ 0 & \text{if E has additive reduction at \mathfrak{p},} \\ 1 & \text{if E has split multiplicative reduction at \mathfrak{p},} \\ -1 & \text{if E has non-split multiplicative reduction at \mathfrak{p}.} \end{cases}$$

(see [Sil86, Appendix C, §16] for more details).

Definition 8.2 (*L*-function attached to a Hilbert modular form). Let F be a totally real number field. Let $f \in S_k^{\text{new}}(\mathfrak{n})$ be a primitive Hilbert modular form over F, of weight k and level \mathfrak{n} with Fourier coefficients $C(\mathfrak{m}, f)$. Then, the L-function attached to f is defined by

$$L(f,s) = \sum_{\mathfrak{m}} C(\mathfrak{m}, f) N_{K/\mathbb{Q}}(\mathfrak{m})^{-s}$$

for $s \in \mathbb{C}$, where \mathfrak{m} runs over all ideals of \mathcal{O}_F .

Then L(f, s) has an Euler product given by

$$L(f,s) = \prod_{\mathfrak{p}} \left(1 - C(\mathfrak{p}, f) N_{K/\mathbb{Q}}(\mathfrak{p})^{-s} + N_{K/\mathbb{Q}}(\mathfrak{p})^{k_0 - 1 - 2s} \right)^{-1},$$

where \mathfrak{p} runs over all prime ideals of \mathcal{O}_F (see [Tan12, §5.3] for more details).

Definition 8.3 (Modularity of elliptic curve). Let K be a totally real number field. We say an elliptic curve E/K is modular over K if there exists a primitive Hilbert modular newform f over K of parallel weight 2 and level $\mathfrak n$ with rational eigenvalues such that both E and f have same L-function, i.e., L(E,s) = L(f,s).

The Shimura-Taniyama conjecture states that every elliptic curve over \mathbb{Q} is modular. This conjecture was first proved by Wiles [Wil95], and Taylor and Wiles [TW95] for semi-stable elliptic curves over \mathbb{Q} . Later, in [BCDT01], Breuil, Conrad, Diamond, and Taylor proved this conjecture for all elliptic curves over \mathbb{Q} .

Theorem 8.4. Every elliptic curve E over \mathbb{Q} is modular.

The following results are known for the modularity of elliptic curves over K when $[K:\mathbb{Q}] \in \{2,3,4,5\}$. More precisely:

Theorem 8.5. ([FLHS15, Theorem 1]) Every elliptic curve E over a real quadratic field is modular.

Theorem 8.6. ([DNS20, Theorem 2]) Every elliptic curve E over a real cubic field is modular.

Theorem 8.7. [Box22, Theorem 1.1] Let K be a totally real quartic field not containing $\sqrt{5}$. Then, every elliptic curve E over K is modular.

Theorem 8.8. [IIY22, Theorem 1.2] For all but finitely many totally real fields K of degree 5 over \mathbb{Q} , every elliptic curve E over K is modular.

In general, for any totally real number field K, [FLHS15] proved that all but finitely many elliptic curves over K are modular. More precisely:

Theorem 8.9. [FLHS15, Theorem 5] Let K be a totally real number field. Then, all but finitely many \bar{K} -isomorphism classes of elliptic curves over K are modular.

ACKNOWLEDGMENTS

The author expresses his sincere gratitude to Prof. Narasimha Kumar and Prof. Nuno Freitas for the invaluable assistance in understanding the article [FS15]. The author sincerely thanks the reviewer for the insightful comments and valuable suggestions towards the improvement of this article.

REFERENCES

- [BS04] Bennett, Michael A.; Skinner, Chris M. Ternary Diophantine equations via Galois representations and modular forms. Canad. J. Math. 56 (2004), no. 1, 23–54.
- [BVY04] Bennett, Michael A.; Vatsal, Vinayak; Yazdani, Soroosh. Ternary Diophantine equations of signature (p, p, 3). Compos. Math. 140 (2004), no. 6, 1399–1416.
- [Bil07] Billerey, Nicolas. Équations de Fermat de type (5,5,p). (French) [[Fermat equations of type (5,5,p)] Bull. Austral. Math. Soc. 76 (2007), no. 2, 161–194.
- [BCDDF23] Billerey, Nicolas; Chen, Imin; Dembélé, Lassina; Dieulefait, Luis; Freitas, Nuno. Some extensions of the modular method and Fermat equations of signature (13, 13, n). Publ. Mat. 67 (2023), no. 2, 715–741.
- [BCDF19] Billerey, Nicolas; Chen, Imin; Dieulefait, Luis; Freitas, Nuno. A multi-Frey approach to Fermat equations of signature (r, r, p). Trans. Amer. Math. Soc. 371 (2019), no. 12, 8651–8677.
- [BCDF25a] Billerey, Nicolas; Chen, Imin; Dieulefait, Luis; Freitas, Nuno. On Darmon's program for the generalized Fermat equation, I. J. Reine Angew. Math. 822 (2025), 107–146.
- [BCDF25b] Billerey, Nicolas; Chen, Imin; Dieulefait, Luis; Freitas, Nuno. On Darmon's program for the generalized Fermat equation, II. Math. Comp. 94 (2025), no. 354, 1977–2003.
- [BD10] Billerey, Nicolas; Dieulefait, Luis V. Solving Fermat-type equations $x^5 + y^5 = dz^p$. Math. Comp. 79 (2010), no. 269, 535–544.
- [Box22] Box, Josha. Elliptic curves over totally real quartic fields not containing $\sqrt{5}$ are modular. Trans. Amer. Math. Soc. 375 (2022), no. 5, 3129–3172.
- [BCDT01] Breuil, Christophe; Conrad, Brian; Diamond, Fred; Taylor, Richard. On the modularity of elliptic curves over \mathbb{Q} : wild 3-adic exercises. J. Amer. Math. Soc. 14 (2001), no. 4, 843–939.
- [Caz24] Cazorla Garcia, Pedro-Josè. Asymptotic Fermat's last theorem for a family of equations of signature (2, 2n, n). Mathematika 70 (2024), no. 4, Paper No. e12279, 21 pp.
- [Dar04] Darmon, Henri. Rational points on modular elliptic curves. CBMS Regional Conference Series in Mathematics, 101. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 2004.
- [DG95] Darmon, Henri; Granville, Andrew. On the equations $z^m = F(x, y)$ and $Ax^p + By^q = Cz^r$. Bull. London Math. Soc. 27 (1995), no. 6, 513–543.
- [DM97] Darmon, Henri; Merel, Loïc. Winding quotients and some variants of Fermat's last theorem. J. Reine Angew. Math. 490 (1997), 81–100.
- [Dec16] Deconinck, Heline. On the generalized Fermat equation over totally real fields. Acta Arith. 173 (2016), no. 3, 225–237.
- [DF13] Dieulefait, Luis; Freitas, Nuno. Fermat-type equations of signature (13, 13, p) via Hilbert cuspforms. Math. Ann. 357 (2013), no. 3, 987–1004.
- [DF14] Dieulefait, Luis; Freitas, Nuno. The Fermat-type equations $x^5 + y^5 = 2z^p$ or $3z^p$ solved through \mathbb{Q} -curves. Math. Comp. 83 (2014), no. 286, 917–933.
- [Fre15] Freitas, Nuno. Recipes to Fermat-type equations of the form $x^r + y^r = Cz^p$. Math. Z. 279 (2015), no. 3-4, 605–639.
- [FN24] Freitas, Nuno; Najman, Filip. Two results on $x^r + y^r = dz^p$. Proc. Amer. Math. Soc. 152 (2024), no. 2, 591–598.
- [DNS20] Derickx, Maarten; Najman, Filip; Siksek, Samir Elliptic curves over totally real cubic fields are modular. Algebra & Number Theory 14 (2020), no. 7, 1791–1800.
- [FS15] Freitas, Nuno; Siksek, Samir. The asymptotic Fermat's last theorem for five-sixths of real quadratic fields. Compos. Math. 151 (2015), no. 8, 1395–1415.
- [FLHS15] Freitas, Nuno; Le Hung, Bao V.; Siksek, Samir. Elliptic curves over real quadratic fields are modular. Invent. Math. 201 (2015), no. 1, 159–206.
- [IIY22] Ishitsuka, Yasuhiro; Ito, Tetsushi; Yoshikawa, Sho. The modularity of elliptic curves over all but finitely many totally real fields of degree 5. Res. Number Theory 8 (2022), no. 4, Paper No. 82, 23 pp.
- [IKO20] Işik, Erman; Kara, Yasemin; Özman, Ekin. On ternary Diophantine equations of signature (p, p, 2) over number fields. Turkish J. Math. 44 (2020), no. 4, 1197–1211.

- [Ivo03] Ivorra, Wilfrid. Sur les équations $x^p + 2^\beta y^p = z^2$ et $x^p + 2^\beta y^p = 2z^2$. (French) [[On the equations $x^p + 2^\beta y^p = z^2$ and $x^p + 2^\beta y^p = 2z^2$] Acta Arith. 108 (2003), no. 4, 327–338.
- [JS25] Jha, Somnath; Sahoo, Satyabrat. Asymptotic Fermat equation of signature (r, r, p) over totally real fields. Ramanujan J. 68 (2025), no. 1, Paper No. 30, 24 pp.
- [KMO24] Kara Yasemin; Mocanu Diana; Özman Ekin. Non-trivial Integer Solutions of $x^r + y^r = Dz^p$. To appear in Indagationes Mathematicae.
- [Kra97] Kraus, Alain. Majorations effectives pour l'équation de Fermat généralisée. (French) [[Effective upper bounds for the generalized Fermat equation]] Canad. J. Math. 49 (1997), no. 6, 1139–1161.
- [KS24a] Kumar, Narasimha; Sahoo, Satyabrat. On the solutions of $x^p + y^p = 2^r z^p$, $x^p + y^p = z^2$ over totally real fields. Acta Arith. 212 (2024), no. 1, 31–47.
- [KS24b] Kumar, Narasimha; Sahoo, Satyabrat. On the solutions of $x^2 = By^p + Cz^p$ and $2x^2 = By^p + Cz^p$ over totally real fields. Ramanujan J. 65 (2024), no. 1, 27–43.
- [Moc22] Mocanu, Diana. Asymptotic Fermat for signatures (p, p, 2) and (p, p, 3) over totally real fields. Mathematika 68 (2022), no. 4, 1233–1257.
- [Moc23] Mocanu, Diana. Asymptotic Fermat for signatures (r, r, p) using the modular approach. Res. Number Theory 9 (2023), no. 4, Paper No. 71, 17 pp.
- [Rib97] Ribet, Kenneth A. On the equation $a^p + 2^{\alpha}b^p + c^p = 0$. Acta Arith. 79 (1997), no. 1, 7–16.
- [Sah24] Sahoo, Satyabrat. On the solutions of the generalized Fermat equation over totally real number fields. arXiv:2404.09171.
- [SK25] Sahoo, Satyabrat; Kumar, Narasimha. Asymptotic solutions of the generalized Fermattype equation of signature (p,p,3) over totally real number fields. J. Number Theory 274 (2025), 56–71.
- [Ser87] Serre, Jean-Pierre. Sur les représentations modulaires de degré 2 de $Gal(\overline{\mathbb{Q}}/\mathbb{Q})$. Duke Math. J. 54 (1987), no. 1, 179–230.
- [Shi78] Shimura, Goro. The special values of the zeta functions associated with Hilbert modular forms. Duke Math. J. 45 (1978), no. 3, 637–679.
- [Sik03] Siksek, Samir. On the Diophantine equation $x^2 = y^p + 2^k z^p$. J. Théor. Nombres Bordeaux 15 (2003), no. 3, 839–846.
- [Sil86] Silverman, Joseph H. The arithmetic of elliptic curves. Graduate Texts in Mathematics, 106. Springer-Verlag, New York, 1986. xii+400 pp.
- [Tan12] Tanabe, Naomi. Arithmetic properties of L-functions attached to Hilbert modular forms. Thesis (Ph.D.)—Oklahoma State University. ProQuest LLC, Ann Arbor, MI, 2012. 119 pp.
- [TW95] Taylor, Richard; Wiles, Andrew. Ring-theoretic properties of certain Hecke algebras. Ann. of Math. (2) 141 (1995), no. 3, 553–572.
- [Wil95] Wiles, Andrew. Modular elliptic curves and Fermat's last theorem. Ann. of Math. (2) 141 (1995), no. 3, 443–551.
- (S. Sahoo) Yau Mathematical Sciences Center, Tsinghua University, Beijing 100084, China.

Email address: satyabrat.sahoo.94@gmail.com