

ON THE OUTER AUTOMORPHISM GROUPS OF THE ABSOLUTE GALOIS GROUPS OF 2-ADIC LOCAL FIELDS

YU NISHIO

DECEMBER 2025

ABSTRACT. In the present paper, we study the outer automorphism groups of the absolute Galois groups of 2-adic local fields from the point of view of anabelian geometry. Let us recall that it is well-known that the natural homomorphism from the automorphism group of a mixed-characteristic local field to the outer automorphism group of the absolute Galois group of the given mixed-characteristic local field is injective. Moreover, Hoshi and the author of the present paper proved that, for absolutely abelian mixed-characteristic local fields with odd residue characteristic p and even extension degree over \mathbb{Q}_p , this subgroup arising from field automorphisms is not normal in the outer automorphism group and has infinitely many distinct conjugates. As a result in this direction, one of the main results of the present paper is the assertion that if a 2-adic local field satisfies certain conditions, then the set of conjugates of the subgroup arising from field automorphisms in the outer automorphism group is infinite, which thus implies that this subgroup is not normal in the outer automorphism group. On the other hand, for an odd prime number p , Hoshi proved the existence of an irreducible Hodge-Tate p -adic representation of dimension two of the absolute Galois group of a p -adic local field and an automorphism of the absolute Galois group such that the p -adic Galois representation obtained by pulling back the given p -adic Galois representation by the given automorphism is not Hodge-Tate. As a result in a direction that is different from the above direction, another main result of the present paper is the existence of a pair of a representation and an automorphism that is similar to the above pair in the case where $p = 2$.

2020 *Mathematics Subject Classification.* 11S20.

Key words and phrases. mixed-characteristic local field, absolute Galois group, anabelian geometry, mono-anabelian geometry, group of MLF-type, 2-adic local field, 2-adic Galois representation, Hodge-Tate, Aut-intrinsically Hodge-Tate.

INTRODUCTION

Let p be a prime number, k a finite extension of \mathbb{Q}_p , and \bar{k} an algebraic closure of k . Write $G_k \stackrel{\text{def}}{=} \text{Gal}(\bar{k}/k)$ for the absolute Galois group of k determined by the algebraic closure \bar{k} and $\text{Out}(G_k)$ for the group of outer automorphisms of the group G_k . In the present paper, we study the outer automorphism group $\text{Out}(G_k)$ from the point of view of anabelian geometry. Write $\text{Aut}(k)$ for the group of automorphisms of the field k . Thus, we have a natural homomorphism $\text{Aut}(k) \rightarrow \text{Out}(G_k)$ of groups. Let us first recall that it is well-known [cf., e.g., [5], Proposition 2.1] that this homomorphism is injective. In the present paper, let us regard $\text{Aut}(k)$ as a [necessarily finite] subgroup of $\text{Out}(G_k)$ by means of this injective homomorphism:

$$\text{Aut}(k) \subseteq \text{Out}(G_k).$$

Here, we note that it is well-known [cf., e.g., the discussion given at the final portion of [12], Chapter VII, §5; [7], Corollary 1.6, (iv)] that, if p is odd, then the equality $\text{Aut}(k) = \text{Out}(G_k)$ never holds. In particular, one may conclude that a finite extension of \mathbb{Q}_p should be considered to be “not anabelian” [cf. also [12], Chapter XII, §2, Closing remark]. Therefore, one main interest from the point of view of anabelian geometry is in the investigation of the “difference” between $\text{Aut}(k)$ and $\text{Out}(G_k)$. Some results concerning the “characterization” of the subgroup $\text{Aut}(k)$ of $\text{Out}(G_k)$ may be found in [11], §3, and [2], §3. Moreover, some results concerning this “difference” may be found in [3], [7], and [9] for odd prime numbers p . On the other hand, no similar results were known in the case where $p = 2$. Thus, it is a natural question to ask how far $\text{Aut}(k)$ differs from $\text{Out}(G_k)$ in the case where $p = 2$. The present paper gives a certain answer to this question. Write $(\mathbb{Q}_p)_+ \subseteq k_+$ for the underlying additive modules of the fields $\mathbb{Q}_p \subseteq k$, respectively. Next, let us recall that, by applying a functorial group-theoretic reconstruction algorithm established in the study of the mono-anabelian geometry of mixed-characteristic local fields [cf., e.g., [5], Definition 3.10, (vi), and [5], Proposition 3.11, (iv)], one obtains an action of the group $\text{Out}(G_k)$ on the module k_+ whose restriction to the subgroup $\text{Aut}(k) \subseteq \text{Out}(G_k)$ coincides with the natural action of $\text{Aut}(k)$ on k_+ .

One main technical result of the present paper is as follows [cf. Theorem 2.5]:

Theorem A. *Suppose that the following two conditions are satisfied:*

- (1) *The field k contains a primitive 4-th root of unity.*
- (2) *The finite extension k/\mathbb{Q}_2 is Galois, and, moreover, the Galois group $\text{Gal}(k/\mathbb{Q}_2)$ is abelian.*

Then there exists an outer automorphism α of G_k such that, for each nonzero integer n , if one writes α_+^n for the action of α^n on k_+ , then $\alpha_+^n((\mathbb{Q}_2)_+) \neq (\mathbb{Q}_2)_+$.

Moreover, in the present paper, we prove the following result [cf. Theorem 3.2]:

Theorem B. *Suppose that the two conditions in the statement of Theorem A are satisfied. Then the set of $\text{Out}(G_k)$ -conjugates of the subgroup $\text{Aut}(k) \subseteq \text{Out}(G_k)$ is infinite.*

A formal consequence of Theorem B is as follows [cf. Corollary 3.3]:

Theorem C. *Suppose that the two conditions in the statement of Theorem A are satisfied. Then the following hold:*

- (i) *The subgroup $\text{Aut}(k) \subseteq \text{Out}(G_k)$ is not normal.*
- (ii) *There exist infinitely many distinct subgroups of $\text{Out}(G_k)$ isomorphic to $\text{Aut}(k)$.*

The final main result of the present paper is as follows [cf. Corollary 4.5]:

Theorem D. *There exist a finite extension K of \mathbb{Q}_2 , an algebraic closure \bar{K} of K , a \mathbb{Q}_2 -vector space V of dimension 2, and a continuous representation $\rho : \text{Gal}(\bar{K}/K) \rightarrow \text{Aut}_{\mathbb{Q}_p K}(V)$ that is irreducible, abelian, crystalline [hence also Hodge-Tate], but not Aut-intrinsically Hodge-Tate [cf. [6], Definition 1.3].*

In [9], Kondo proved similar results to Theorem A, Theorem B, Theorem C, and Theorem D in the case where k is absolutely Galois [i.e., where the extension k/\mathbb{Q}_p is Galois], and p is odd. Moreover, in [10], Kumpitsch studied the outer automorphism groups of the absolute Galois groups of mixed-characteristic local fields from the point of view of the study of mapping class groups and constructed large subgroups of the outer automorphism groups.

0. NOTATIONAL CONVENTIONS

TOPOLOGICAL GROUPS. If G is a topological group, then we shall write G^{ab} for the *abelianization* of G [i.e., the quotient of G by the closure of the commutator subgroup of G], $G^{\text{ab-tor}}$ for the closure of the subgroup of torsion elements of G^{ab} , and $G^{\text{ab/tor}}$ for the quotient of G^{ab} by $G^{\text{ab-tor}}$. If H is a profinite group, and p is a prime number, then we shall write $H^{(p)}$ for the *maximal pro- p quotient* of H .

RINGS. If R is a ring, then we shall write R_+ for the underlying additive module of R and R^\times for the multiplicative group of units of R .

MODULES. Let M be a module. We shall write

$$M^\wedge \stackrel{\text{def}}{=} \varprojlim M/(n \cdot M)$$

— where the projective limit is taken over the positive integers n .

FIELDS. We shall refer to a field isomorphic to a finite extension of \mathbb{Q}_p , for some prime number p , as a *mixed-characteristic local field*.

If k is a mixed-characteristic local field, then we shall write

- $k^{(d=1)} \subseteq k$ for the [uniquely determined] minimal mixed-characteristic local field contained in k ,
- f_k for the absolute residue degree of k ,
- $d_k \stackrel{\text{def}}{=} [k : k^{(d=1)}]$ for the degree of the finite extension $k/k^{(d=1)}$,
- $\mathcal{O}_k \subseteq k$ for the ring of integers of k ,
- $\mathfrak{m}_k \subseteq \mathcal{O}_k$ for the maximal ideal of \mathcal{O}_k ,
- $\mathcal{O}_k^\times \stackrel{\text{def}}{=} 1 + \mathfrak{m}_k \subseteq \mathcal{O}_k^\times$,
- p_k for the residue characteristic of k , and
- a_k for the largest nonnegative integer such that k contains a primitive $p_k^{a_k}$ -th root of unity.

We shall refer to a group isomorphic to the absolute Galois group of a mixed-characteristic local field as a group of *MLF-type* [cf. [4], Definition 1.1]. In the present paper, let us always regard a group of MLF-type as a profinite group by means of the profinite topology discussed in [4], Proposition 1.2, (i) [cf. also [4], Proposition 1.2, (ii)].

1. EXISTENCE OF AN AUTOMORPHISM WITH A CERTAIN UNIPOTENCY CONDITION OF A GROUP OF MLF-TYPE CORRESPONDING TO A 2-ADIC LOCAL FIELD CONTAINING PRIMITIVE 4-TH ROOTS OF UNITY

In the present §1, we prove that a certain group of MLF-type admits an automorphism that satisfies a certain unipotency condition [cf. Theorem 1.6 below]. In the present §1, let G be a group of MLF-type. Thus, by applying the various functorial group-theoretic reconstruction algorithms of [5], §3 [cf. [5], Definition 3.5, (i), (ii), (iii); [5], Definition 3.10, (ii), (iv), (vi)], to the group G of MLF-type, we obtain

- a prime number $p(G)$,
- positive integers $d(G)$ and $f(G)$,
- a nonnegative integer $a(G) \stackrel{\text{def}}{=} \log_{p(G)} \left(\# \left((k^\times(G)_{\text{tor}})^{(p(G))} \right) \right)$
- a normal closed subgroup $P(G) \subseteq G$ of G , and
- topological modules $\mathcal{O}^\times(G) \subseteq k^\times(G)$ and $k_+(G)$

[cf. also [5], Summary 3.15]. Here, let us recall [cf. [5], Proposition 3.6; [5], Proposition 3.11, (i), (iv); [3], Proposition 2.5] that if k is a mixed-characteristic local field, \bar{k} is an algebraic closure of k , and $G_k \stackrel{\text{def}}{=} \text{Gal}(\bar{k}/k)$ is the absolute Galois group of k determined by the algebraic closure \bar{k} , then

- $p(G_k), d(G_k), a(G_k), f(G_k), P(G_k)$ coincide with p_k, d_k, a_k, f_k , the wild inertia subgroup of G_k , respectively, and, moreover,
- there exist functorial isomorphisms $\mathcal{O}_k^\prec \xrightarrow{\sim} \mathcal{O}^\prec(G_k), k^\times \xrightarrow{\sim} k^\times(G_k)$, and $k_+ \xrightarrow{\sim} k_+(G_k)$.

Furthermore, let us also recall [cf. also [5], Proposition 3.6; [5], Definition 3.10, (i), (ii), (vi); [5], Proposition 3.11, (i)] that

- the module $\mathcal{O}^\prec(G)$ is defined to be the image of $P(G) \subseteq G$ in G^{ab} , and
- the module $k_+(G)$ is defined to be $\mathcal{O}^\prec(G) \otimes_{\mathbb{Z}} \mathbb{Q}$, i.e., the perfection [cf., e.g., the discussion entitled “Monoids” of [3], §0] of the module $\mathcal{O}^\prec(G)$ [cf. also the commutative diagram

$$\begin{array}{ccc} \mathcal{O}_k^\prec & \xrightarrow{\log_k} & k_+ \\ \wr \downarrow & & \wr \downarrow \\ \mathcal{O}^\prec(G_k) & \longrightarrow & k_+(G_k) \end{array}$$

determined by the commutative diagram of [5], Proposition 3.11, (iv)].

In the present §1, suppose, moreover, that $p(G) = 2$, and that $a(G) \geq 2$.

Proposition 1.1 (Diekert). *There exist*

- *topological generators $\sigma, \tau, x_0, \dots, x_{d(G)}$ of G ,*
- *positive integers s, g such that $g \neq 1$, and*

that satisfy the following three conditions:

- (1) *The normal closed subgroup $P(G)$ of G is topologically normally generated by $x_0, \dots, x_{d(G)}$.*
- (2) *The equality $\sigma\tau\sigma^{-1} = \tau^{q(G)}$ holds, where we write $q(G) \stackrel{\text{def}}{=} 2^{f(G)}$.*
- (3) *The equality*

$$\sigma x_0 \sigma^{-1} = (x_0 \tau)^{\pi^g x_1^{2^s}} [x_1, x_2] \cdots [x_{d(G)-1}, x_{d(G)}]$$

holds, where we write π for the unique element of $\hat{\mathbb{Z}} = \prod_p \mathbb{Z}_p$ whose image in \mathbb{Z}_p is given by 1 if $p = 2$ (resp. by 0 if $p \neq 2$).

Proof. This assertion follows from [1], Theorem 3.1, and [5], Proposition 3.6, [3], Proposition 2.5, (i). \square

In the remainder of the present §1, let us fix topological generators σ , τ , $x_0, \dots, x_{d(G)}$ of G as in Proposition 1.1. Write $S \stackrel{\text{def}}{=} \{0, 1, \dots, d(G)\}$. Moreover, for each $i \in S$, write

- $y_i \in k_+(G)$ for the image of $x_i \in P(G)$ [cf. the condition (1) of Proposition 1.1] in $k_+(G)$ and
- $z_i \in \mathcal{O}^{\prec}(G)^{\text{ab/tor}}$ for the image of $x_i \in P(G)$ [cf. the condition (1) of Proposition 1.1] in $\mathcal{O}^{\prec}(G)^{\text{ab/tor}}$

[cf. the constructions of $\mathcal{O}^{\prec}(G)$, $k_+(G)$ explained in the discussion preceding Proposition 1.1].

Lemma 1.2. *The topological module $k_+(G)$ has a natural structure of \mathbb{Q}_2 -vector space of dimension $d(G)$.*

Proof. It follows immediately from the definition of $k_+(G)$ [cf. [5], Definition 3.10, (vi)] that $k_+(G)$ has a natural structure of \mathbb{Q}_2 -vector space. Moreover, it follows from [5], Proposition 3.6, and [5], Proposition 3.11, (iv), that this \mathbb{Q}_2 -vector space $k_+(G)$ is of dimension $d(G)$. \square

Lemma 1.3. *The $d(G)$ elements $y_1, \dots, y_{d(G)}$ form a basis of the \mathbb{Q}_2 -vector space $k_+(G)$ [cf. Lemma 1.2].*

Proof. Since $\{x_i\}_{i \in S}$ topologically normally generates $P(G)$ [cf. the condition (1) of Proposition 1.1], $\{z_i\}_{i \in S}$ topologically generates $\mathcal{O}^{\prec}(G)^{\text{ab/tor}} (\subseteq G^{\text{ab/tor}})$ [cf. [5], Definition 3.10, (i), (ii)]. Moreover, since $\mathcal{O}^{\prec}(G)^{\text{ab/tor}} \otimes_{\mathbb{Z}_2} (\mathbb{Z}_2/2^n\mathbb{Z}_2)$ is a finite 2-group for every positive integer n [cf. [5], Proposition 3.11, (i)], $\{z_i\}_{i \in S}$ is also a generator of $\mathcal{O}^{\prec}(G)^{\text{ab/tor}}$ even if we regard $\mathcal{O}^{\prec}(G)^{\text{ab/tor}}$ as a \mathbb{Z}_2 -module. Next, let us observe that it follows from the condition (2) of Proposition 1.1 that the image of τ in $G^{\text{ab/tor}}$ is trivial. Thus, it follows from the condition (3) of Proposition 1.1 that the relation $1 = z_0^H z_1^{2^s}$ in $\mathcal{O}^{\prec}(G)^{\text{ab/tor}}$ holds for some nonzero [cf. the condition that $g \neq 1$ of Proposition 1.1] integer H . Therefore, if we write $T \stackrel{\text{def}}{=} S \setminus \{0\}$, then $\{z_i \otimes 1\}_{i \in T}$ is a generator of \mathbb{Q}_2 -vector space $\mathcal{O}^{\prec}(G)^{\text{ab/tor}} \otimes_{\mathbb{Z}_2} \mathbb{Q}_2$. Here, let us observe that we have a natural topological isomorphism $k_+(G) \xrightarrow{\sim} \mathcal{O}^{\prec}(G)^{\text{ab/tor}} \otimes_{\mathbb{Z}_2} \mathbb{Q}_2$, by definition, that maps $y_i \in k_+(G)$ to $z_i \otimes 1 \in \mathcal{O}^{\prec}(G)^{\text{ab/tor}} \otimes_{\mathbb{Z}_2} \mathbb{Q}_2$ for each $i \in S$. This isomorphism is also an isomorphism of \mathbb{Q}_2 -vector spaces by construction. Moreover, since $k_+(G) \simeq \mathcal{O}^{\prec}(G)^{\text{ab/tor}} \otimes_{\mathbb{Z}_2} \mathbb{Q}_2$ is a \mathbb{Q}_2 -vector space of dimension $d(G)$ [cf. Lemma 1.2], $\{z_i \otimes 1\}_{i \in T}$ is a basis of the \mathbb{Q}_2 -vector space $\mathcal{O}^{\prec}(G)^{\text{ab/tor}} \otimes_{\mathbb{Z}_2} \mathbb{Q}_2$. Therefore, it follows from the condition imposed on the isomorphism $k_+(G) \xrightarrow{\sim} \mathcal{O}^{\prec}(G)^{\text{ab/tor}} \otimes_{\mathbb{Z}_2} \mathbb{Q}_2$ that $\{y_i\}_{i \in T}$ is a basis of the \mathbb{Q}_2 -vector space $k_+(G)$. \square

Lemma 1.4. *The element $y_{d(G)-1}$ is a nonzero element of $k_+(G)$.*

Proof. This assertion follows from Lemma 1.3 [cf. also our assumption that $a(G) > 1$, hence $d(G) > 1$]. \square

Definition 1.5. We shall write $^*\alpha$ for the automorphism of G defined by the equalities $^*\alpha(\sigma) = \sigma$, $^*\alpha(\tau) = \tau$, $^*\alpha(x_{d(G)}) = x_{d(G)}x_{d(G)-1}$, and $^*\alpha(x_i) = x_i$ for $i \in S \setminus \{d(G)\}$. Since $^*\alpha$ induces a bijection between the set of generators of G and preserves the defining relations of G [cf. Proposition 1.1], it indeed defines an automorphism of G .

Theorem 1.6. *Let G be a group of MLF-type such that $p(G) = 2$, and that $a(G) \geq 2$. Then, for each nonzero integer n , if one writes $^*\alpha_+^n$ for the automorphism of the \mathbb{Q}_2 -vector space $k_+(G)$ induced by $^*\alpha^n$ [cf. Definition 1.5], then $^*\alpha_+^n \neq \text{id}$, and, moreover, the equality $(^*\alpha_+^n - \text{id})^2 = 0$ in the ring of endomorphisms of $k_+(G)$ holds.*

Proof. First, we prove that $^*\alpha_+^n \neq \text{id}$ for each nonzero integer n . If $^*\alpha_+^n = \text{id}$, then $y_{d(G)} = ^*\alpha_+^n(y_{d(G)}) = y_{d(G)} + ny_{d(G)-1}$, which thus implies that $y_{d(G)-1} = 0$ in $k_+(G)$. However, this contradicts Lemma 1.4. Thus, we conclude that $^*\alpha_+^n \neq \text{id}$. Next, let us observe that, for each nonzero integer n , it follows from the easily verified equality $(^*\alpha_+^n - \text{id})^2(y_i) = 0$ for every $i \in S$ and Lemma 1.3 that the equality $(^*\alpha_+^n - \text{id})^2 = 0$ in $\text{End}(k_+(G))$ holds, as desired. This completes the proof of Theorem 1.6. \square

Corollary 1.7. *Let G be a group of MLF-type such that $p(G) = 2$, and $a(G) \geq 2$. Then the following hold:*

- (i) *The image of the natural homomorphism from the outer automorphism group of G to the automorphism group of $k_+(G)$ is infinite.*
- (ii) *The image of the natural homomorphism from the outer automorphism group of G to the automorphism group of G^{ab} is infinite.*
- (iii) *The image of the natural homomorphism from the outer automorphism group of G to the automorphism group of $k^\times(G)$ is infinite.*
- (iv) *The outer automorphism group of G is infinite.*

Proof. Assertion (i) follows from Theorem 1.6. Assertion (ii) follows from assertion (i), together with the definition of $k_+(G)$ [cf. [5], Definition 3.10, (vi)]. Assertion (iii) follows from assertion (ii) and the [easily verified] density of $k^\times(G)$ in G^{ab} [cf. [5], Definition 3.10, (iv)]. Assertion (iv) follows from assertion (i). \square

Remark 1.8. Let us recall that it follows immediately from [3], Corollary 5.5, that each of the three images discussed in Corollary 1.7, (i), (ii), (iii), in the case where $d(G)$ is equal to 1 is trivial.

2. EXISTENCE OF A SPECIAL AUTOMORPHISM OF THE ABSOLUTE GALOIS GROUP OF AN ABSOLUTELY ABELIAN 2-ADIC LOCAL FIELD CONTAINING PRIMITIVE 4-TH ROOTS OF UNITY

In the present §2, we prove that the absolute Galois group of a certain 2-adic local field admits an automorphism that has an interesting property [cf. Theorem 2.5 below]. In the present §2, let k be a mixed-characteristic local field and \bar{k} an algebraic closure of k . We shall write $G_k \stackrel{\text{def}}{=} \text{Gal}(\bar{k}/k)$ for the absolute Galois group of k determined by the algebraic closure \bar{k} . Write, moreover, $\text{Aut}(G_k)$, $\text{Aut}(k_+)$, and $\text{Aut}(k^\times)$ for the groups of automorphisms of the group G_k , the module k_+ , and the module k^\times , respectively. Thus, it follows from [5], Proposition 3.11, (i), (iv), that we have natural homomorphisms

$$\text{Aut}(G_k) \longrightarrow \text{Aut}(k_+), \quad \text{Aut}(G_k) \longrightarrow \text{Aut}(k^\times).$$

Definition 2.1.

- (i) We shall say that $\alpha \in \text{Aut}(k_+)$ is $(\mathbb{Q}_{p_k})_+$ -characteristic if $\alpha(k_+^{(d=1)}) = k_+^{(d=1)}$.
- (ii) We shall say that $\alpha \in \text{Aut}(k_+)$ is $(\mathbb{Q}_{p_k})_+$ -preserving if α is $(\mathbb{Q}_{p_k})_+$ -characteristic, and $\alpha|_{k_+^{(d=1)}}$ is the identity automorphism of $k_+^{(d=1)}$.
- (iii) We shall say that $\alpha \in \text{Aut}(k_+)$ is *group-theoretic* if α is contained in the image of the first homomorphism $\text{Aut}(G_k) \rightarrow \text{Aut}(k_+)$ of the above display.
- (iv) We shall say that $\alpha \in \text{Aut}(k^\times)$ is *group-theoretic* if α is contained in the image of the second homomorphism $\text{Aut}(G_k) \rightarrow \text{Aut}(k^\times)$ of the above display.
- (v) We shall say that $\alpha \in \text{Aut}(G_k)$ is $(\mathbb{Q}_{p_k})_+$ -characteristic if the group-theoretic automorphism of k_+ induced by α is $(\mathbb{Q}_{p_k})_+$ -characteristic.
- (vi) We shall say that $\alpha \in \text{Aut}(G_k)$ is $(\mathbb{Q}_{p_k})_+$ -preserving if the group-theoretic automorphism of k_+ induced by α is $(\mathbb{Q}_{p_k})_+$ -preserving.

Lemma 2.2. *Let α be an automorphism of G_k . Write $\alpha_+ \in \text{Aut}(k_+)$ and $\alpha^\times \in \text{Aut}(k^\times)$ for the respective group-theoretic automorphisms induced by α , $\text{Nm}_{k/k^{(d=1)}}$ for the norm map with respect to the finite extension $k/k^{(d=1)}$, and $\text{Tr}_{k/k^{(d=1)}}$ for the trace map with respect to the finite extension $k/k^{(d=1)}$. Then the following hold:*

(i) The automorphism α^\times fits into a commutative diagram of groups

$$\begin{array}{ccc} k^\times & \xrightarrow{\text{Nm}_{k/k^{(d=1)}}} & k^{(d=1)\times} \\ \alpha^\times \downarrow & & \parallel \\ k^\times & \xrightarrow{\text{Nm}_{k/k^{(d=1)}}} & k^{(d=1)\times}. \end{array}$$

(ii) The automorphism α_+ fits into a commutative diagram of modules

$$\begin{array}{ccc} k_+ & \xrightarrow{\text{Tr}_{k/k^{(d=1)}}} & k_+^{(d=1)} \\ \alpha_+ \downarrow & & \parallel \\ k_+ & \xrightarrow{\text{Tr}_{k/k^{(d=1)}}} & k_+^{(d=1)}. \end{array}$$

In particular, the automorphism α_+ restricts to an automorphism of $\text{Ker}(\text{Tr}_{k/k^{(d=1)}})$, i.e., the equality $\alpha_+(\text{Ker}(\text{Tr}_{k/k^{(d=1)}})) = \text{Ker}(\text{Tr}_{k/k^{(d=1)}})$ holds.

Proof. This lemma is none other than [7], Lemma 2.3. \square

Lemma 2.3. Suppose that $p_k = 2$, and that k is isomorphic to the finite Galois extension of $k^{(d=1)}$ obtained by adjoining a primitive 4-th root of unity. Write $^*\alpha$ for the automorphism of G_k defined in Definition 1.5, i.e., in the case where we take the “ G ” of Definition 1.5 to be G_k . Then, for every nonzero integer n , the automorphism $^*\alpha^n$ is not $(\mathbb{Q}_2)_+$ -characteristic.

Proof. It follows from Theorem 1.6; [5], Proposition 3.6; [3], Proposition 2.5 that, for every nonzero integer n , the group-theoretic automorphism $^*\alpha_+^n$ of k_+ is not the identity automorphism but satisfies the equality $(^*\alpha_+^n - \text{id})^2 = 0$ in $\text{End}(k_+)$. Here, let us observe that we can write $k = k^{(d=1)}(i)$ for some primitive 4-th root of unity i . Assume that $^*\alpha_+^n$ is $(\mathbb{Q}_2)_+$ -characteristic for some nonzero integer n . Thus, $^*\alpha_+^n(1) = b$ for some $b \in k^{(d=1)}$. Moreover, it follows from the final portion of Lemma 2.2, (ii), that $^*\alpha_+^n(i) = ci$ for some $c \in k^{(d=1)}$. Thus, since $^*\alpha_+$ is an automorphism of \mathbb{Q}_2 -vector space, it follows that, for arbitrary $x, y \in k^{(d=1)}$, the equalities

$$0 = (^*\alpha_+^n - \text{id})^2(x + yi) = x(b - 1)^2 + y(c - 1)^2i$$

hold. Thus, we have $(b, c) = (1, 1)$. In particular, $^*\alpha_+^n$ is the identity automorphism. However, this is a contradiction. \square

Lemma 2.4. *Suppose that $a_k \geq 2$, and that k is absolutely abelian, i.e., that k is Galois over $k^{(d=1)}$, and, moreover, the Galois group $\text{Gal}(k/k^{(d=1)})$ is abelian [cf. [3], Definition 4.2, (ii)]. Write k' for the quadratic extension of $k^{(d=1)}$ obtained by adjoining a primitive 4-th root of unity in k and $G_{k'} \stackrel{\text{def}}{=} \text{Gal}(\bar{k}/k')$. Then the following hold:*

- (i) G_k is a characteristic subgroup of $G_{k'}$. In particular, we have a natural homomorphism $\phi: \text{Aut}(G_{k'}) \rightarrow \text{Aut}(G_k)$.
- (ii) Let α' be an automorphism of $G_{k'}$ that is not $(\mathbb{Q}_p)_+$ -characteristic. Then $\phi(\alpha') \in \text{Aut}(G_k)$ [cf. (i)] is not $(\mathbb{Q}_p)_+$ -characteristic.

Proof. First, we verify assertion (i). Let β be an automorphism of $G_{k'}$. Write $*k \subseteq \bar{k}$ for the finite extension of k' that corresponds to the open subgroup $\beta(G_k) \subseteq G_{k'}$. Thus, it is immediate that $[k : k'] = [*k : k']$. Moreover, since k is absolutely abelian, it follows immediately from the main theorem of [8] [cf. also [3], Theorem E, which is a generalization of the main theorem of [8]] that k is contained in $*k$. Thus, we conclude from the above equality $[k : k'] = [*k : k']$ that $k = *k$, which thus implies that $\beta(G_k) = G_k$, as desired. This completes the proof of assertion (i).

Next, we verify assertion (ii). Let us first observe that it follows immediately from the various definitions involved that the diagram

$$\begin{array}{ccccc} k_+^{(d=1)} & \hookrightarrow & k'_+ & \hookrightarrow & k_+ \\ \phi(\alpha')_+|_{k_+^{(d=1)}} \downarrow & & \alpha'_+ \downarrow & & \downarrow \phi(\alpha')_+ \\ \alpha'_+(k_+^{(d=1)}) & \hookrightarrow & k'_+ & \hookrightarrow & k_+ \end{array}$$

commutes, where the horizontal arrows are the natural inclusions, and we write α'_+ (resp. $\phi(\alpha')_+$) for the group-theoretic automorphism induced by $\alpha' \in \text{Aut}(G_{k'})$ (resp. $\phi(\alpha') \in \text{Aut}(G_k)$). Since α'_+ is not $(\mathbb{Q}_p)_+$ -characteristic, $\alpha'_+(k_+^{(d=1)}) \neq k_+^{(d=1)}$. Thus, we conclude from the above diagram that $\phi(\alpha')_+$, hence also $\phi(\alpha')$, is not $(\mathbb{Q}_p)_+$ -characteristic. This completes the proof of assertion (ii). \square

Theorem 2.5. *Let k be an absolutely abelian mixed-characteristic local field such that $p_k = 2$, and $a_k \geq 2$. Then there exists an automorphism $\alpha \in \text{Aut}(G_k)$ such that, for each nonzero integer n , the automorphism α^n is not $(\mathbb{Q}_2)_+$ -characteristic.*

Proof. Write k' for the quadratic extension of $k^{(d=1)}$ obtained by adjoining a primitive 4-th root of unity in k and $G_{k'} \stackrel{\text{def}}{=} \text{Gal}(\bar{k}/k')$. Then it follows from Lemma 2.3 that there exists an automorphism β of $G_{k'}$ such that, for every nonzero integer n , the automorphism β^n is not $(\mathbb{Q}_2)_+$ -characteristic.

Thus, we conclude from Lemma 2.4, (ii), that the restriction of β^n to G_k is not $(\mathbb{Q}_2)_+$ -characteristic. This completes the proof of Theorem 2.5. \square

3. THE OUTER AUTOMORPHISM GROUP OF THE ABSOLUTE GALOIS GROUP OF AN ABSOLUTELY ABELIAN 2-ADIC LOCAL FIELD CONTAINING PRIMITIVE 4-TH ROOTS OF UNITY

In the present §3, we discuss the outer automorphism group of the absolute Galois group of a certain 2-adic local field. In the present §3, we maintain the notational conventions introduced at the beginning of the preceding §2. Write, moreover, $\text{Aut}(k)$ for the group of automorphisms of the field k and $\text{Out}(G_k)$ for the group of outer automorphisms of the group G_k . Thus, we have a natural injective [cf. [5], Proposition 2.1] homomorphism $\text{Aut}(k) \hookrightarrow \text{Out}(G_k)$ of groups. In the present §3, let us regard $\text{Aut}(k)$ as a subgroup of $\text{Out}(G_k)$:

$$\text{Aut}(k) \subseteq \text{Out}(G_k).$$

Lemma 3.1. *Let α be an automorphism of G_k . Suppose that k is absolutely Galois. If the image of α in $\text{Out}(G_k)$ is contained in $N_{\text{Out}(G_k)}(\text{Aut}(k))$, then α is $(\mathbb{Q}_{p_k})_+$ -preserving.*

Proof. This lemma is none other than [7], Lemma 3.3. \square

Theorem 3.2. *Let k be an absolutely abelian mixed-characteristic local field such that $p_k = 2$, and $a_k \geq 2$. Then the set of $\text{Out}(G_k)$ -conjugates of the subgroup $\text{Aut}(k) \subseteq \text{Out}(G_k)$ is infinite.*

Proof. It is immediate that, to verify Theorem 3.2, it suffices to show that $N_{\text{Out}(G_k)}(\text{Aut}(k))$ is of infinite index in $\text{Out}(G_k)$. On the other hand, if $\alpha \in \text{Aut}(G_k)$ is an automorphism as in Theorem 2.5, then it follows from Lemma 3.1 that, for each nonzero integer n , the image of α^n in $\text{Out}(G_k)$ is not contained in $N_{\text{Out}(G_k)}(\text{Aut}(k))$. In particular, we conclude that $N_{\text{Out}(G_k)}(\text{Aut}(k))$ is of infinite index in $\text{Out}(G_k)$, as desired. This completes the proof of Theorem 3.2. \square

Corollary 3.3. *Let k be an absolutely abelian mixed-characteristic local field such that $p_k = 2$, and $a_k \geq 2$. Then the following hold:*

- (i) *The subgroup $\text{Aut}(k)$ of $\text{Out}(G_k)$ is not normal.*
- (ii) *There exist infinitely many distinct subgroups of $\text{Out}(G_k)$ isomorphic to $\text{Aut}(k)$.*

Proof. These assertions follow immediately from Theorem 3.2. \square

Corollary 3.4. *Let k be the quadratic mixed-characteristic local field obtained by adjoining a primitive 4-th root of unity to $k^{(d=1)}$ such that $p_k = 2$.*

Then the group-theoretic automorphism of k_+ induced by an automorphism of G_k which lifts an element of the center of $\text{Out}(G_k)$ is the identity automorphism of k_+ .

Proof. Let γ be an element of the center of $\text{Out}(G_k)$. Write $\gamma_+ \in \text{Aut}(k_+)$ for the group-theoretic automorphism of k_+ induced by an automorphism of G_k which lifts γ . [Note that one verifies easily that γ_+ does not depend on the choice of such a lifting.] Then it follows from Lemma 3.1 that γ_+ is $(\mathbb{Q}_2)_+$ -preserving. Next, let $\alpha_+ \in \text{Aut}(k_+)$ be a group-theoretic automorphism of k_+ which is not $(\mathbb{Q}_2)_+$ -characteristic [cf. Theorem 2.5]. Then since γ is an element of the center of $\text{Out}(G_k)$, one verifies immediately that γ_+ commutes with α_+ . In particular, since γ_+ is $(\mathbb{Q}_2)_+$ -preserving, γ_+ restricts to the identity automorphism of $\alpha_+(k_+^{(d=1)}) \subseteq k_+$. Thus, since $d_k = 2$, and $k_+^{(d=1)} \neq \alpha_+(k_+^{(d=1)})$, we conclude that γ_+ is the identity automorphism of k_+ , as desired. This completes the proof of Corollary 3.4. \square

4. EXISTENCE OF A SPECIAL IRREDUCIBLE CRYSTALLINE REPRESENTATIONS OF DIMENSION TWO

In the present §4, let k be a mixed-characteristic local field and \bar{k} an algebraic closure of k . We shall write $G_k \stackrel{\text{def}}{=} \text{Gal}(\bar{k}/k)$ for the absolute Galois group of k determined by the algebraic closure \bar{k} and $\text{rec}_k: (k^\times)^\wedge \xrightarrow{\sim} G_k^{\text{ab}}$ for the isomorphism induced by the reciprocity homomorphism $k^\times \rightarrow G_k^{\text{ab}}$ in local class field theory. Let us recall that, for a given \mathbb{Q}_{p_k} -vector space V of finite dimension and a given continuous representation $\rho: G_k \rightarrow \text{Aut}_{\mathbb{Q}_{p_k}}(V)$ of G_k , we say that ρ is *Aut-intrinsically Hodge-Tate* if, for an arbitrary automorphism α of G_k , the composite $\rho \circ \alpha: G_k \xrightarrow{\sim} G_k \rightarrow \text{Aut}_{\mathbb{Q}_{p_k}}(V)$ is Hodge-Tate [cf. [6], Definition 1.3].

In the present §4, we prove the existence of an irreducible crystalline [hence also Hodge-Tate] 2-adic representation of dimension 2 that is not Aut-intrinsically Hodge-Tate [cf. Corollary 4.5 below].

First, let us recall the functorial assignment “ $\mathcal{O}^\times(-)$ ” of [5], Definition 3.10, (i). Observe that it follows from the functoriality of the assignment “ $\mathcal{O}^\times(-)$ ” that each automorphism of G_k naturally induces a automorphism of $\mathcal{O}^\times(G_k)$. In particular, by conjugating this automorphism of $\mathcal{O}^\times(G_k)$ by the isomorphism $\mathcal{O}_k^\times \xrightarrow{\sim} \mathcal{O}^\times(G_k)$ of [5], Proposition 3.11, (i), one concludes that each automorphism of G_k naturally induces a automorphism of \mathcal{O}_k^\times .

Definition 4.1. For an automorphism α of G_k , we shall write α_\times for the automorphism of \mathcal{O}_k^\times induced by α .

Lemma 4.2. *Suppose that k is the quadratic mixed-characteristic local field obtained by adjoining a primitive 4-th root of unity to $k^{(d=1)}$, and that $p_k = 2$. Write $\text{Nm}: k^\times \rightarrow (k^{(d=1)})^\times$ for the norm map with respect to the finite extension $k/k^{(d=1)}$. Then the following hold:*

- (i) *There exists an open submodule $U \subseteq \mathcal{O}_k^\times$ such that*
 - (1) *the topological module U has a natural structure of free \mathbb{Z}_2 -module of rank 2, and, moreover,*
 - (2) *the submodule $U \subseteq \mathcal{O}_k^\times$ is preserved by an arbitrary automorphism of \mathcal{O}_k^\times .*
- (ii) *Let $U \subseteq \mathcal{O}_k^\times$ be as in (i). Then the topological modules $U \cap \mathcal{O}_{k^{(d=1)}}^\times$, $U \cap \text{Ker}(\text{Nm})$ have natural structures of free \mathbb{Z}_2 -modules of rank 1, respectively.*
- (iii) *Let $U \subseteq \mathcal{O}_k^\times$ be as in (i). Then the equality $U \cap \mathcal{O}_{k^{(d=1)}}^\times \cap \text{Ker}(\text{Nm}) = \{1\}$ holds.*
- (iv) *Let $U \subseteq \mathcal{O}_k^\times$ be as in (i). Then the closed submodule of U topologically generated by the closed submodules $U \cap \mathcal{O}_{k^{(d=1)}}^\times$ and $U \cap \text{Ker}(\text{Nm})$ is open.*
- (v) *There exists an automorphism α of G_k such that, for an arbitrary nonzero integer n , the intersection $\alpha_\times^n(\mathcal{O}_{k^{(d=1)}}^\times) \cap \mathcal{O}_{k^{(d=1)}}^\times$ is not open in $\mathcal{O}_{k^{(d=1)}}^\times$. In particular, the automorphism α_\times^n does not preserve the submodule $\mathcal{O}_{k^{(d=1)}}^\times \subseteq \mathcal{O}_k^\times$.*

Proof. Assertions (i), (ii) follow from [5], Lemma 1.2, (i) [cf. also our assumption that $d_k = p_k = 2$.] Assertion (iii) is immediate [cf. the fact that U is torsion-free, which follows from the condition (1) of assertion (i)]. Assertion (iv) follows from assertions (ii), (iii), together with the condition (1) of assertion (i).

Finally, we verify assertion (v). Let α be an automorphism of G_k as in Definition 1.5. [Note that since $p(G_k) = p_k = 2$, $a(G_k) = a_k = 2 > 1$ — cf. [5], Proposition 3.6; [3], Proposition 2.5, (i) — the group G_k [of MLF-type] satisfies the assumption imposed on “ G ” in the discussion preceding Proposition 1.1.] Write β for the automorphism of the submodule $U \subseteq \mathcal{O}_k^\times$ obtained by forming the restriction of α_\times^n [cf. the condition (2) of assertion (i)]. Thus, since it follows from Theorem 1.6 [cf. also [5], Definition 3.10, (vi)] that

- (a[†]) the automorphism of $\mathcal{O}_k^\times \otimes_{\mathbb{Z}} \mathbb{Q}$ induced by α_\times^n is not the identity automorphism, but
- (b[†]) the image of the square of the endomorphism of $\mathcal{O}_k^\times \otimes_{\mathbb{Z}} \mathbb{Q}$ induced by the endomorphism of \mathcal{O}_k^\times given by “ $a \mapsto \alpha_\times^n(a) \cdot a^{-1}$ ” consists of the identity element of $\mathcal{O}_k^\times \otimes_{\mathbb{Z}} \mathbb{Q}$,

one concludes that

- (a) the automorphism β is not the identity automorphism of U , but
- (b) the image of the square of the endomorphism of U given by “ $a \mapsto \beta(a) \cdot a^{-1}$ ” consists of the identity element of U .

Moreover, it follows immediately from Lemma 2.2, (i), that

- (c) the automorphism β preserves the submodule $U \cap \text{Ker}(\text{Nm})$ of U .

Thus, it follows immediately from assertion (iv) [cf. also the condition (1) of assertion (i)], together with (b) and (c), that if the automorphism β preserves some open submodule of the submodule $U \cap \mathcal{O}_{k^{(d=1)}}^\times$, then β is the identity automorphism of U — in contradiction to (a). In particular, the automorphism β does not preserve any open submodule of the submodule $U \cap \mathcal{O}_{k^{(d=1)}}^\times$, which thus implies [cf. assertion (ii)] that

$$\beta(U \cap \mathcal{O}_{k^{(d=1)}}^\times) \cap U \cap \mathcal{O}_{k^{(d=1)}}^\times = \{1\}.$$

Thus, it follows immediately from [5], Lemma 1.2, (i), that

$$\alpha_\times^n(\mathcal{O}_{k^{(d=1)}}^\times) \cap \mathcal{O}_{k^{(d=1)}}^\times$$

is not open in $\mathcal{O}_{k^{(d=1)}}^\times$, as desired. This completes the proof of assertion (v), hence also of Lemma 4.2. \square

Proposition 4.3. *Let k be a mixed-characteristic local field containing a primitive 4-th root of unity such that $p_k = 2$. Suppose, moreover, that k is absolutely abelian, i.e., that k is absolutely Galois, and the Galois group $\text{Gal}(k/k^{(d=1)})$ is abelian [cf. [3], Definition 4.2, (ii)]. Then there exists an automorphism α of G_k such that, for an arbitrary nonzero integer n , if one writes α_\times^n for the automorphism of \mathcal{O}_k^\times induced by α^n , then the intersection*

$$\alpha_\times^n(\mathcal{O}_{k^{(d=1)}}^\times) \cap \mathcal{O}_{k^{(d=1)}}^\times$$

is not open in $\mathcal{O}_{k^{(d=1)}}^\times$.

Proof. Let us first observe that since k contains a primitive 4-th root of unity, it is immediate that k contains the quadratic mixed-characteristic local field obtained by adjoining a primitive 4-th root of unity to $k^{(d=1)}$. Moreover, since k is absolutely abelian, it follows immediately from the implication (1) \Rightarrow (2) of [3], Theorem F, (i), that G_k is a characteristic subgroup of the absolute Galois group of the quadratic extension of $k^{(d=1)}$ determined by the algebraic closure \bar{k} . Thus, one may conclude that we may assume without loss of generality — by applying a similar argument to the argument applied in the proof of Lemma 2.4, (ii), and replacing k by the quadratic extension of $k^{(d=1)}$ — that $d_k = 2$. On the other hand, if $d_k = 2$, then the desired conclusion follows from Lemma 4.2, (v). This completes the proof of Proposition 4.3. \square

Theorem 4.4. *Let k be a mixed-characteristic local field containing a primitive 4-th root of unity such that $p_k = 2$. Assume, moreover, that k is absolutely abelian. Then there exist a \mathbb{Q}_{p_k} -vector space V of dimension d_k and a continuous representation $\rho : \text{Gal}(\bar{k}/k) \rightarrow \text{Aut}_{\mathbb{Q}_{p_k}}(V)$ that is irreducible, abelian, crystalline [hence also Hodge-Tate], but not Aut-intrinsically Hodge-Tate [cf. [6], Definition 1.3].*

Proof. Let $\pi \in \mathcal{O}_k$ be a uniformizer of \mathcal{O}_k . Write ρ for the continuous representation of G_k [necessarily of dimension d_k] obtained by forming the composite

$$G_k \twoheadrightarrow G_k^{\text{ab}} \xrightarrow{\chi_{\pi, \text{id}_k}} \mathcal{O}_k^\times \hookrightarrow \text{Aut}_{\mathbb{Q}_{p_k}}(k_+)$$

— where the first arrow is the natural surjective homomorphism, the second arrow is the character defined in [6], Definition 1.7, and the third arrow is the natural inclusion. Then one verifies easily that this continuous representation ρ is irreducible and abelian. Moreover, it follows immediately from [13], Chapter III, §A.4, Proposition 5, that this continuous representation ρ is crystalline.

Next, to verify that the continuous representation ρ is not Aut-intrinsically Hodge-Tate, let us recall that it follows immediately from the various definitions involved that the composite

$$\mathcal{O}_k^\times \xrightarrow{\text{rec}_k} G_k^{\text{ab}} \xrightarrow{\chi_{\pi, \text{id}_k}} \mathcal{O}_k^\times$$

is an automorphism that restricts to an automorphism of the submodule $\mathcal{O}_{k(d=1)}^\times \subseteq \mathcal{O}_k^\times$. In particular, if α is an automorphism of G_k as in Proposition 4.3, then it follows immediately from [6], Lemma 1.9, together with the various definitions involved, that the composite $\rho \circ \alpha : G_k \xrightarrow{\sim} G_k \rightarrow \text{Aut}_{\mathbb{Q}_{p_k}}(k_+)$ is not Hodge-Tate, which thus implies that the continuous representation ρ is not Aut-intrinsically Hodge-Tate, as desired. This completes the proof of Theorem 4.4. \square

Corollary 4.5. *There exist a mixed-characteristic local field K such that $p_K = 2$, an algebraic closure \bar{K} of K , a \mathbb{Q}_{p_K} -vector space V of dimension 2, and a continuous representation $\rho : \text{Gal}(\bar{K}/K) \rightarrow \text{Aut}_{\mathbb{Q}_{p_K}}(V)$ that is irreducible, abelian, crystalline [hence also Hodge-Tate], but not Aut-intrinsically Hodge-Tate.*

Proof. This assertion is a formal consequence of Theorem 4.4. \square

ACKNOWLEDGMENTS

The author would like to thank Yuichiro Hoshi for a discussion related to this thesis, and would like to express deepest gratitude to my partner, K. Nishio for providing constant support and warm encouragement.

REFERENCES

- [1] V. Diekert. “Über die absolute Galoisgruppe dyadischer Zahlkörper”. In: *J. Reine Angew. Math.* 350 (1984), pp. 152–172.
- [2] Y. Hoshi. “A note on the geometricity of open homomorphisms between the absolute Galois groups of p -adic local fields”. In: *Kodai Math. J.* 36.2 (2013), pp. 284–298.
- [3] Y. Hoshi. “Topics in the anabelian geometry of mixed-characteristic local fields”. In: *Hiroshima Math. J.* 49.3 (2019), pp. 323–398.
- [4] Y. Hoshi. “Mono-anabelian reconstruction of number fields”. In: *On the examination and further development of inter-universal Teichmüller theory*. Vol. B76. RIMS Kôkyûroku Bessatsu. Kyoto: Res. Inst. Math. Sci. (RIMS), 2019, pp. 1–77.
- [5] Y. Hoshi. “Introduction to mono-anabelian geometry”. In: *Publications mathématiques de Besançon. Algèbre et théorie des nombres*. 2021. Publ. Math. Besançon Algèbre Théorie Nr. Besançon: Presses Univ. Franche-Comté, 2021, pp. 5–44. [2022].
- [6] Y. Hoshi. “On intrinsic Hodge-Tate-ness of Galois representations of dimension two”. In: *Kodai Math. J.* 47.1 (2024), pp. 99–111.
- [7] Y. Hoshi and Y. Nishio. “On the Outer Automorphism Groups of the Absolute Galois Groups of Mixed-characteristic Local Fields”. In: *Res. Number Theory* 8.3, 56 (Apr. 2022).
- [8] W. Jenkner. “Les corps p -adiques dont les groupes de Galois absolus sont isomorphes”. fr. In: *Journées arithmétiques de Genève - 9-13 septembre 1991*. Ed. by Coray D. F. and Pétermann Y.-F. S. Astérisque 209. Société mathématique de France, 1992, pp. 221–226. URL: https://www.numdam.org/item/AST_1992__209__221_0/.
- [9] K. Kondo. “On the non-normality of the field-theoretic subgroups of the outer automorphism groups of the absolute Galois groups of mixed-characteristic local fields”. MA thesis. Kyoto University, 2025.
- [10] T. Kumpitsch. “Outer automorphisms of the absolute Galois group of local fields of mixed characteristic”. PhD thesis. Johann Wolfgang Goethe-Universität Frankfurt am Main, 2022. URL: https://publikationen.ub.uni-frankfurt.de/opus4/frontdoor/deliver/index/docId/70616/file/PhDThesis_Kumpitsch.pdf.
- [11] S. Mochizuki. “Topics in absolute anabelian geometry I: generalities”. In: *J. Math. Sci. Univ. Tokyo* 19.2 (2012), pp. 139–242.
- [12] J. Neukirch, A. Schmidt, and K. Wingberg. *Cohomology of number fields*. Second. Vol. 323. Grundlehren der Mathematischen Wissenschaften. Berlin: Springer-Verlag, 2008.

- [13] J.-P. Serre. *Abelian l -adic representations and elliptic curves. With the collaboration of Willem Kuyk and John Labute. Revised reprint of the 1968 original.* Vol. 7. Research Notes in Mathematics. Wellesley, MA: A K Peters, Ltd., 1998.

(Yu Nishio)

Email address: nishio.math@gmail.com