

Jet Charge with Global Event Shapes: Probing Quark Flavor Dynamics

Yang-Ting Chien¹ and Sonny Mantry²

¹*Physics and Astronomy Department, Georgia State University, Atlanta, GA 30303*

²*Department of Physics and Astronomy, University of North Georgia, Dahlonega, GA 30597, USA*

We propose measuring the jet electric charge of jet regions, defined within the framework of global event shapes, as a probe of quark flavor dynamics within the nucleon and the hadronization process. In particular, we consider a measurement of the jet region charge while simultaneously keeping track of the energy flow throughout the event, as characterized by the global event shape. As a concrete example, we focus on the measurement of the *1-Jettiness jet charge* (\mathcal{Q}), the jet charge of the jet region (J) defined within the framework of the 1-Jettiness global event shape (τ_1) for the Deep Inelastic Scattering (DIS) process, $e^- + p \rightarrow e^- + J + X$, with unpolarized or longitudinally polarized protons. The 1-Jettiness distribution, binned according to jet charge, allows for enhanced quark flavor separation of the initial state unpolarized or polarized PDFs. On the other hand, the jet charge distribution binned by 1-Jettiness can serve as a probe of quark flavor dynamics in the final state hadronization process. We derive a factorization theorem for simultaneous measurements of τ_1 and \mathcal{Q} in the resummation region, $\tau_1 \ll P_{J_T}$, where P_{J_T} denotes the transverse momentum of the jet region. The correlation between the jet region charge and the charge of the struck quark is expected to be the strongest in the resummation region. The factorization theorem contains a new universal *charged jet function*, generalizing the standard jet function to include a jet charge measurement. Therefore these universal functions can be extracted from a global analysis of N-jettiness and thrust at e^+e^- colliders. We provide simulation studies to demonstrate the sensitivity of the 1-Jettiness jet charge observable to quark flavor dynamics in nucleon structure and explore the possibility of probing the final state hadronization process. This observable is well-suited for applications in existing HERA data and the future Electron-Ion Collider (EIC).

I. INTRODUCTION

Global Event shapes are infrared and collinear safe observables that characterize the geometric pattern of hadronic energy flows in the final state. They act as a continuous measure of how pencil-like, planar, or spherical the “shape” of the final state radiation is. Examples of global event shapes include thrust [1, 2], heavy jet mass [3], sphericity [4], C-parameter [5], Broadening [6], N-Jettiness [7], and 1-Jettiness [8–15] for Deep Inelastic Scattering (DIS). They have been studied in a variety of applications, including precision extractions of the strong coupling constant α_s [16–18] and the top quark mass [19–21]. Such global event shapes have also been computed to very high accuracy and compared to data [16, 22, 23], providing important tests of perturbative QCD and hadronization models.

However, due to the light quark flavor symmetry of QCD, global event shapes do not provide much information on light quark flavor dynamics in initial state nucleon structure or final state hadronization. Typically, a global QCD analysis over a wide range of observables is employed to extract universal initial state structure functions and final state fragmentation functions, separated by flavor. These include inclusive observables as well as more exclusive ones that involve tagging a final state hadron. For example, the neutral current and charged current fully inclusive deep inelastic scattering (DIS) processes, $e^- + p \rightarrow e^- + X$ and $e^- + p \rightarrow \nu_e + X$, respectively, are sensitive to different flavor combinations of initial state parton distribution functions (PDFs). Simi-

larly, using polarized protons in these processes allows for disentangling initial state helicity PDFs by flavor. More exclusive observables, such as semi-inclusive DIS (SIDIS), $e^- + p \rightarrow e^- + h + X$, where a final state hadron, h , is tagged, allows for additional flavor separation of initial state nucleon structure functions. Additionally, it acts as a probe of fragmentation functions by flavor. Measurements at the Relativistic Heavy Ion Collider (RHIC) and Large Hadron Collider (LHC) have shown that jets can also be a powerful probe of nucleon structure, including nucleon spin structure [24–26]. Such jet based probes have now also been developed for use at the EIC [27–35]. However, without tagging final state hadrons, flavor separation remain challenging in inclusive jet measurements.

The idea of jet charge was first explored [36] more than four decades ago. More recently, the jet electric charge has been shown [37, 38] to be able to provide discrimination between jets initiated by up-type and down-type quarks at the LHC. Jets are initiated by energetic partons emerging from a hard collision process, subsequently producing a parton shower through perturbative soft and collinear emissions which is followed by hadronization into a cluster of hadrons. A suitable definition of the measured jet charge will correlate with the electric charge of the jet-initiating parton, providing a jet tagging tool that discriminates between jets initiated by different parton flavors. Jet charge was used in DIS studies [39–44], for measurements of the charge asymmetry [45, 46], tagging b -quarks in jets [47–50] and hadronically decaying W bosons [51, 52], testing aspects of per-

turbative [37, 38, 53–55] and nonperturbative [56] Quantum Chromodynamics (QCD), probing nuclear-medium-induced jet quenching effects in heavy ion collisions [57–63], and probing the quark flavor structure of unpolarized and polarized [64, 65] transverse momentum dependent parton distribution functions (TMDPDFs). The jet charge has also been used as one of a variety of jet substructure tools [62, 66–68] for jet discrimination [69–71], with enhanced performances using machine learning techniques [52, 55].

The standard definition of the jet charge [37, 38] for a jet with transverse momentum, P_{J_T} , is given by:

$$\mathcal{Q}_\kappa = \sum_{h \in \text{jet}} z_h^\kappa Q_h, \quad (1)$$

where the sum is over all the hadrons in the jet, Q_h denotes the electric charge of the hadron, h , and $z_h = p_{T_h}/P_{J_T}$ is the fraction of the jet transverse momentum carried by the hadron, where p_{T_h} is the transverse momentum of the hadron. The parameter $\kappa > 0$ ensures that the contribution to the jet charge from soft hadrons is suppressed, ensuring infrared safety. It was shown that $\kappa \sim 0.3$ provides the best discrimination [72] between jets initiated by different quark flavors and will be the default value used in our analysis. A modified definition of the standard jet charge, the Dynamic Jet Charge [71], allows for the κ parameter to be a function of some property of the jet constituents. It can be chosen as a function of z_h , $\kappa = \kappa(z_h)$, to improve discrimination between quark and gluon jets and make jet discrimination more resilient against the contamination from underlying events. We will often drop the κ subscript in Eq. (1) and refer to the jet charge as simply, \mathcal{Q} , for brevity in notation. In our analysis, we consider both the standard and dynamic jet charge definitions, as they have different jet discrimination capabilities and sensitivity to soft radiation, serving as complementary probes.

Recently, it was shown that the jet charge property can be used for flavor separation of the unpolarized and polarized Transverse Momentum Dependent PDFs (TMDPDFs) [64]. The unpolarized TMDPDFs describe the 3D momentum space distribution of partons in the nucleon. The polarized TMDPDFs, in addition, encode quantum correlations between the parton momentum distributions and the proton spin. The electron-proton scattering process, $e^- + p \rightarrow e^- + J + X$, in the limit of small electron-jet transverse momentum imbalance, $q_T = |\vec{p}_T^e + \vec{P}_{J_T}| \ll \vec{p}_T^e \sim P_{J_T}$, is sensitive to TMDPDFs. The factorization formula for the q_T -distribution is sensitive to a particular flavor combination of TMDPDFs, a universal soft function, and a universal jet function. The jet function describes the dynamics of the leading jet that recoils against the electron, and in general depends on the jet radius, R , and the jet clustering algorithm. Flavor separation of the TMDPDFs is achieved through a jet charge measurement on the leading jet and binning the q_T -distribution by jet charge. The positive and negative jet charge bins provide enhanced sensitivity to

the u -quark and the d, s quark TMDPDFs, respectively, compared to the q_T -distribution integrated over all jet charge values.

It was also recently explored that, charge and flavor correlation among hadrons is sensitive to details of underlying hadronization processes [73] therefore it can be used to test hadronization models in Monte Carlo implementations. While the studies require tagging individual hadron flavors, i.e. particle identifications, jet charge or subjet charges could provide more inclusive information about quark flavors.

In this work, we build on these ideas and propose a new class of observables that incorporates the flavor discrimination power of jet charges within the framework of global event shapes as a probe of quark flavor dynamics in nuclear structure and hadronization. We propose measuring the jet charge (\mathcal{Q}) of jet regions as defined by the global event shape, independent of the use of jet algorithms. This allows for measuring the jet region charge while simultaneously keeping track of the energy flow throughout the event, as quantified by the global event shape. In particular, one can bin event shape distributions according to the jet region charge. Conversely, one can bin jet region charge distributions according to the global event shape value. Such observables can allow for disentangling flavor dynamics in the initial state nucleon structure functions and in the final state hadronization process.

We focus on the 1-Jettiness [8–14] global event shape for the neutral current DIS process, $e^- + p \rightarrow e^- + J + X$. The transverse momentum and rapidity of the jet region, J , are denoted by P_{J_T} and y_J , respectively. The usual 1-Jettiness cross section, differential in (τ_1, P_{J_T}, y_J) , is denoted by:

$$d\sigma[\tau_1, P_{J_T}, y_J] \equiv \frac{d^3\sigma(e^- + p \rightarrow J + X)}{dy_J dP_{J_T} d\tau_1}. \quad (2)$$

The resummation region of the τ_1 -distribution, $\tau_1 \ll P_{J_T}$, corresponds to events with a single jet topology such that energetic radiation, $E \sim P_{J_T} \gg \tau_1$, is collimated either along the proton beam or the leading jet direction, and only soft radiation, $E \sim \tau_1 \ll P_{J_T}$ at wide angles from the beam and jet directions is allowed. It requires resummation of large Sudakov logarithms of the form $\alpha_s^n \ln^{2m}(\tau_1/P_{J_T})$, for $m \leq n$, arising from the $\tau_1 \ll P_{J_T}$ restriction on final state radiation. This restriction effectively acts as a veto on additional jets at wide angles from the leading jet and beam directions. In this resummation region, the differential cross section has the schematic, factorized form

$$d\sigma_{\text{resum}}[\tau_1, P_{J_T}, y_J] \sim H \otimes B \otimes J \otimes S, \quad (3)$$

where H denotes the hard function describing the hard partonic scattering, J denotes the jet function describing the energetic radiation collinear with jet region, S denotes the soft function describing the soft radiation throughout the event, and B denotes the beam function [74] describing initial state collinear radiation along

the beam. The beam function can be further factorized as [7, 74]

$$B \sim \mathcal{I} \otimes f, \quad (4)$$

where \mathcal{I} describes the perturbatively calculable initial state collinear radiation and f denotes the PDF.

On the other hand, the fixed-order region, $\tau_1 \sim P_{J_T}$, corresponds to events with additional energetic radiation at wide angles from the beam and leading jet directions. In this region, calculations are done in the fixed-order perturbative QCD framework. The resummation and fixed-order regions can be matched to give a smooth 1-Jettiness spectrum over the entire range of τ_1 . Most recently, results for the 1-Jettiness spectrum was obtained at the $N^3\text{LL} + \mathcal{O}(\alpha_s^2)$ level of accuracy [14].

We extend this 1-Jettiness framework by, in addition, measuring the charge, Q , of the jet region, J . We refer to it as the “1-Jettiness Jet Charge.” The cross section is now differential in four variables, $(Q, \tau_1, P_{J_T}, y_J)$, and is denoted as:

$$d\sigma[Q, \tau_1, P_{J_T}, y_J] \equiv \frac{d^3\sigma(e^- + p \rightarrow J + X)}{dy_J dP_{J_T} d\tau_1 dQ}. \quad (5)$$

The correlation between the 1-Jettiness jet charge, Q , with the electric charge of the hard-scattered quark, allows for enhanced flavor separation of the initial state PDFs through the jet-charge-binned 1-Jettiness distributions. This correlation is expected to be the strongest in the resummation region, $\tau_1 \ll P_{J_T}$, where the struck quark evolves into the final state jet region through collinear and soft splittings without any hard emissions at wide angles. In this resummation region, using the Soft-Collinear Effective Theory (SCET) [75–80], the differential cross section for the 1-Jettiness Jet Charge distribution has the schematic form

$$d\sigma_{\text{resum}}[Q, \tau_1, P_{J_T}, y_J] \sim H \otimes B \otimes \mathcal{G} \otimes \mathcal{S}, \quad (6)$$

which has the same form as Eq. (3), except for the replacement $J \rightarrow \mathcal{G}$, corresponding to replacing the standard jet function with a *jet charge jet function*. This formula will receive power suppressed contributions from the contribution of soft radiation to the jet region charge, as discussed in appendix B. The jet charge jet function, \mathcal{G} , is obtained through the insertion of a jet charge measurement operator in the standard jet function. The new jet charge jet function \mathcal{G} is in general nonperturbative since the jet charge measurement is made on final state hadrons, making it sensitive to the hadronization process. However, it is universal therefore it can be extracted from other jet-based global event shapes such as the N-Jettiness [7] global event shape at e^+e^- colliders or the LHC. The differential cross section in Eq. (5) can be integrated over Q -bins to give the jet-charge-binned 1-Jettiness distribution. In the resummation region, the jet-charge-binned 1-Jettiness distribution has the schematic form

$$d\sigma_{\text{resum}}[Q^{\text{bin}}, \tau_1, P_{J_T}, y_J] \sim H \otimes B \otimes \mathcal{G}_{\text{bin}} \otimes \mathcal{S}, \quad (7)$$

where \mathcal{G}_{bin} denotes the jet charge function integrated over a jet charge bin. The positive jet charge bin ($Q_J > 0.25$) will give more weight to the contribution of the u -quark to the 1-Jettiness spectrum, corresponding to its positive electric charge. The negative jet charge bin ($Q_J < -0.25$) will give more weight to the contribution of the d, s -quarks to the 1-Jettiness spectrum, corresponding to their negative electric charge. Thus, binning the 1-Jettiness distribution according to the 1-Jettiness jet charge allows for disentangling initial state proton structure functions by flavor. For scattering of unpolarized protons, this corresponds to separating the collinear PDFs by flavor. The longitudinal single spin asymmetry constructed from scattering off polarized protons will allow for flavor separation of helicity PDFs.

Alternatively, with the multi-differential cross section in Eq. (5) one can integrate over τ_1 -bins to give the 1-Jettiness-binned jet charge distribution. This allows one to study the jet charge distribution as a function of the shape of energy flow in the final state as characterized by the value of τ_1 . Such studies can probe flavor dynamics in the hadronization process, providing a new tool for distinguishing between different hadronization models.

We also note that this observable can be adapted to the charged current DIS events, $e^- + p \rightarrow \nu_e + J + X$ since the undetected neutrino, tagged as missing energy, does not contribute to the τ_1, Q, P_{J_T} , or y_J observables. Such a charge current DIS observable will complement the analysis of neutral current DIS as it probes different combinations of initial state proton structure functions, allowing for further separation of structure functions by flavor. We leave the charge current DIS analysis for future work.

Finally, we note that the formalism for the 1-Jettiness charge observable developed here is ideally suited for the proposed Electron-Ion Collider (EIC) and can also be readily applied to the existing HERA data.

II. 1-JETTINESS GLOBAL EVENT SHAPE

For completeness, we give a brief review of the formalism and notation for the 1-Jettiness global event shape for DIS which has been extensively studied [8–14]. For more detailed explanations of the relevant notation used in this work, see Ref. [14]. We consider the neutral current (NC) DIS process:

$$e^-(k) + p(P) \rightarrow e^-(k') + J + X, \quad (8)$$

in the center of mass frame, where the initial state proton and electron momenta, P^μ and k^μ respectively, are given by

$$\begin{aligned} P^\mu &= \frac{\sqrt{s}}{2} n^\mu, & n^\mu &= (1, 0, 0, 1), \\ k^\mu &= \frac{\sqrt{s}}{2} \bar{n}^\mu, & \bar{n}^\mu &= (1, 0, 0, -1). \end{aligned} \quad (9)$$




FIG. 1. Pythia 8.312 simulation data (blue dots) for the normalized 1-Jettiness τ_1 -distribution compared to the theoretical prediction at the $N^3\text{LL}$ level of accuracy with a soft function model that describes hadronization effects. The scale variations (pink band), describing the theoretical uncertainty, are normalized to the central curve (red) over the displayed range. The relevant EIC kinematics chosen are: $\sqrt{s} = 90.0$ GeV, $P_{JT} = [20.0, 30.0]$ GeV, and $y_J = [-2.5, 2.5]$.

The 1-Jettiness global event shape, τ_1 , is defined as

$$\tau_1 = \sum_k \min \left\{ \frac{2q_B \cdot p_k}{Q_B}, \frac{2q_J \cdot p_k}{Q_J} \right\}, \quad (10)$$

where the sum is over all final state particles except the scattered electron. The q_B^μ and q_J^μ denote the beam and jet reference vectors, respectively, whose explicit expressions will be given shortly below. The Q_B and Q_J are energy scales at the order of the hard scale. Different choices for these quantities correspond to different definitions of 1-Jettiness. Each final state particle of momentum p_k is grouped with either the beam region or the jet region according to the minimization condition in Eq. (10), and accordingly contributes to τ_1 . The largest contributions to τ_1 come from final state particles with large energies and at wide angles from both the beam and jet reference vectors. On the other hand, energetic particles closely aligned with either the beam or jet reference vectors give relatively small contributions to the τ_1 value. Thus, events with small τ_1 values have energetic radiation closely aligned with the beam or jet reference vectors. Events with large τ_1 have additional energetic radiation at wide angles from both the beam and jet reference vectors. Therefore, τ_1 quantifies the pattern of final state energy flow.

The beam and jet reference vectors can be determined through a minimization procedure, corresponding to the choice that minimizes the value of τ_1 in Eq. (10). However, we instead work with the choice for the beam reference vector

$$q_B^\mu = xP^\mu, \quad (11)$$

aligned with the beam direction. The jet reference vector is chosen as

$$q_J = (K_{JT} \cosh y_K, \vec{K}_{JT}, K_{JT} \sinh y_K), \quad (12)$$

where K_J^μ is the four-momentum of the leading jet refined by the anti- k_t algorithm using jet radius $R = 1$. Here K_{JT} and y_K denote the transverse momentum and rapidity of the leading jet, and are used to construct the null jet reference vector, q_J^μ . We choose the constants Q_B and Q_J as

$$Q_B = x\sqrt{s}, \quad Q_J = 2K_{JT} \cosh y_K. \quad (13)$$

The factorization formula for the τ_1 distribution in the resummation region is given by [14]

$$\begin{aligned} d\sigma_{\text{resum}}[\tau_1, P_{JT}, y_J] &= \sigma_0 H(\xi, \mu; \mu_H) \\ &\times \int ds_J \int dt_a \mathcal{S} \left(\tau_1 - \frac{t_a}{Q_a} - \frac{s_J}{Q_J}, \mu; \mu_S \right) \\ &\times \left[\sum_{q_i} L_{q_i} J^{q_i}(s_J, \mu; \mu_J) B_{q_i}(x_*, t_a, \mu; \mu_B) \right. \\ &\left. + \sum_{\bar{q}_i} L_{\bar{q}_i} J^{\bar{q}_i}(s_J, \mu; \mu_J) B_{\bar{q}_i}(x_*, t_a, \mu; \mu_B) \right], \quad (14) \end{aligned}$$

where H , \mathcal{S} , J^{q_i, \bar{q}_i} , and B_{q_i, \bar{q}_i} denote the hard function, soft function, quark and antiquark jet functions, and quark and antiquark beam functions, respectively, and

$$x_* = \frac{e^{y_J} P_{JT}}{\sqrt{s} - e^{-y_J} P_{JT}}, \quad \xi = x_* \sqrt{s} P_{JT} e^{-y_J}. \quad (15)$$

Note that the hard, soft, jet, and beam functions are all evaluated at the common renormalization scale μ after renormalization group evolution from the scales $\mu_H \sim P_{JT}$, $\mu_S \sim \tau_1$, $\mu_J \sim \mu_B \sim \sqrt{P_{JT} \tau_1}$ which eliminates large logarithms in each of the respective functions. The renormalization group evolution sums large Sudakov logarithms arising from ratios of these scales. The quark and anti-quark beam functions ($B_{q, \bar{q}}$) are matched [74] onto the PDFs as

$$\begin{aligned} B_{(q_i, \bar{q}_i)}(x, t_a, \mu; \mu_B) &= \sum_i \int_x^1 \frac{dz}{z} \mathcal{I}_{(q_i, \bar{q}_i)j} \left(\frac{x}{z}, t_a, \mu; \mu_B \right) \\ &\times f_{j/A}(z, \mu_B), \quad (16) \end{aligned}$$

where the $\mathcal{I}_{(q_i, \bar{q}_i)j}$ functions are perturbatively calculable. The hard [81, 82], soft [83, 84], jet [81, 82, 85], and beam functions [86, 87] are known to $\mathcal{O}(\alpha_s^2)$. Expressions for σ_0 and L_{q_i, \bar{q}_i} can be found in Refs. [10, 14]. Results for this 1-Jettiness spectrum are known [14] at the $N^3\text{LL} + \mathcal{O}(\alpha_s^2)$ level of accuracy.

III. 1-JETTINESS JET CHARGE FACTORIZATION FRAMEWORK

The 1-Jettiness jet charge distribution corresponds to the measurement of the jet charge of the jet region as

defined by the 1-Jettiness global event shape, in addition to the measurement of the 1-Jettiness global event shape itself. In the resummation region, $\tau_1 \ll P_{JT}$, its factorization formula is given by

$$\begin{aligned} d\sigma_{\text{resum}}[\mathcal{Q}, \tau_1, P_{JT}, y_J] &= \sigma_0 H(x_* \sqrt{s} P_{JT} e^{-y_J}, \mu; \mu_H) \\ &\times \int ds_J \int dt_a \mathcal{S}\left(\tau_1 - \frac{t_a}{Q_a} - \frac{s_J}{Q_J}, \mu; \mu_S\right) \\ &\times \left[\sum_{q_i} L_{q_i} \mathcal{G}^{q_i}(\mathcal{Q}, s_J, \mu; \mu_J) B_{q_i}(x_*, t_a, \mu; \mu_B) \right. \\ &\left. + \sum_{\bar{q}_i} L_{\bar{q}_i} \mathcal{G}^{\bar{q}_i}(\mathcal{Q}, s_J, \mu; \mu_J) B_{\bar{q}_i}(x_*, t_a, \mu; \mu_B) \right]. \quad (17) \end{aligned}$$

Compared to Eq. (14), the jet function $J^q(s_J, \mu)$ is now replaced with the *charged jet function*, $\mathcal{G}^q(\mathcal{Q}, s_J, \mu)$, which corresponds to the insertion of a jet charge measurement operator in the standard jet function. Its field-theoretic definition is given in Eq. (A2). Note that the jet charge contribution from soft particles in the jet region is power suppressed compared to that of the energetic collinear particles. This can be understood from the definition of jet charge in Eq. (1) which weighs the charge of each particle by its jet energy fraction. The dynamic jet charge allows for the contribution of the soft particles to be further suppressed by choosing the κ parameter as a dynamic function of the hadron energy fraction, z_h . Therefore, at leading power, the jet charge measurement operator only acts on the energetic collinear particles in the jet region. Consequently, only the jet function, $J^q(s_J, \mu)$, is affected and replaced with the jet charge jet function $\mathcal{G}^q(\mathcal{Q}, s_J, \mu)$. We discuss power suppressed corrections arising from soft particles in appendix B.

The charged jet function, $\mathcal{G}^q(\mathcal{Q}, s_J, \mu)$, is in general a non-perturbative object since the jet charge measurement operator is not collinear safe and sensitive to the hadronization process. However, $\mathcal{G}^q(\mathcal{Q}, s_J, \mu)$ has the same renormalization scale dependence as $J^q(s_J, \mu)$. This follows from the renormalization scale independence of the cross section and the fact that 1-Jettiness Jet Charge cross section in Eq. (17) can be obtained simply by replacing $J^q(s_J, \mu)$ with $\mathcal{G}^q(\mathcal{Q}, s_J, \mu)$ in Eq. (14). Integrating Eq. (17) over all \mathcal{Q} should give back the standard 1-Jettiness cross section in Eq. (14). In particular, integrating $\mathcal{G}^q(\mathcal{Q}, s_J, \mu)$ over \mathcal{Q} gives back the standard jet function

$$\int d\mathcal{Q} \mathcal{G}^q(\mathcal{Q}, s_J, \mu) = J^q(s_J, \mu), \quad (18)$$

which also follows from Eqs. (A1) and (A2). In position space, the jet function, $J^q(y, \mu)$, and charged jet function, $\mathcal{G}^q(\mathcal{Q}, y, \mu)$, are defined through the Fourier transforms

$$\begin{aligned} J^q(s, \mu) &= \int \frac{dy}{2\pi} e^{iys} J^q(y, \mu), \\ \mathcal{G}^q(\mathcal{Q}, s, \mu) &= \int \frac{dy}{2\pi} e^{iys} \mathcal{G}^q(\mathcal{Q}, y, \mu). \quad (19) \end{aligned}$$

It is useful to define the ratio of the position space charged and standard jet functions

$$R^q(\mathcal{Q}, y) \equiv \frac{\mathcal{G}^q(\mathcal{Q}, y, \mu)}{J^q(y, \mu)}. \quad (20)$$

Therefore $\mathcal{G}^q(\mathcal{Q}, y, \mu)$ and $J^q(y, \mu)$ are related through the multiplicative function $R^q(\mathcal{Q}, y)$ as

$$\mathcal{G}^q(\mathcal{Q}, y, \mu) = R^q(\mathcal{Q}, y) J^q(y, \mu). \quad (21)$$

Note that $R^q(\mathcal{Q}, y)$ is renormalization scale independent due to the cancellation between the renormalization group running, which is multiplicative in position space, of $\mathcal{G}^q(\mathcal{Q}, y, \mu)$ and $J^q(y, \mu)$ in the ratio. The multiplicative relation in Eq. (21) becomes a convolution in momentum space. Using Eq. (21) in Eq. (19), we get

$$\mathcal{G}^q(\mathcal{Q}, s, \mu) = \int ds' R^q(\mathcal{Q}, s - s') J^q(s', \mu), \quad (22)$$

where $R^q(\mathcal{Q}, s)$ corresponds to the momentum space definition

$$R^q(\mathcal{Q}, s) = \int \frac{dy}{2\pi} e^{iys} R^q(\mathcal{Q}, y). \quad (23)$$

Combining Eqs. (18) and (22) leads to the normalization conditions

$$\int d\mathcal{Q} R^q(\mathcal{Q}, s) = \delta(s), \quad \int d\mathcal{Q} R^q(\mathcal{Q}, y) = 1. \quad (24)$$

From Eq. (22), we see that the function $R(\mathcal{Q}, s)$ acts as the integral kernel that transforms the standard jet function, $J^q(s, \mu)$, distribution in the mass squared variable, ‘ s ’, to the new distribution, $\mathcal{G}^q(\mathcal{Q}, s, \mu)$, where the final state is restricted to, in addition, have a jet charge value of \mathcal{Q} . It is a universal, non-perturbative function that can be extracted from data on other jet-based global event shapes, as explained in section V.

We see from Eqs. (21) and (22) that determining the non-perturbative and universal charged jet function, $\mathcal{G}^q(\mathcal{Q}, y, \mu)$ or $\mathcal{G}(\mathcal{Q}, s, \mu)$, amounts to determining the function, $R^q(\mathcal{Q}, y)$ or $R^q(\mathcal{Q}, s)$, respectively, while the jet function, $J(y, \mu)$ or $J(s, \mu)$, is perturbatively calculable. At lowest order in $\alpha_s(\mu)$ the jet function is given by $J^{(0)}(y, \mu) = 1$ and $J^{(0)}(s, \mu) = \delta(s)$, and correspondingly

$$\begin{aligned} \mathcal{G}^{q(0)}(\mathcal{Q}, y, \mu) &= R^q(\mathcal{Q}, y), \\ \mathcal{G}^{q(0)}(\mathcal{Q}, s, \mu) &= R^q(\mathcal{Q}, s). \quad (25) \end{aligned}$$

i.e. at this order, the charged jet function, $\mathcal{G}^{q(0)}(\mathcal{Q}, y, \mu)$ or $\mathcal{G}^{q(0)}(\mathcal{Q}, s, \mu)$, is identical to the $R^q(\mathcal{Q}, y)$ or $R^q(\mathcal{Q}, s)$, respectively.

Using Eq. (22) in Eq. (17), the resummed factorization

formula differential in jet charge takes the form:

$$\begin{aligned} d\sigma_{\text{resum}}[\mathcal{Q}, \tau_1, P_{J_T}, y_J] &= \sigma_0 H(\xi, \mu; \mu_H) \\ &\times \int ds_J \int dt_a \int ds'_J \mathcal{S}\left(\tau_1 - \frac{t_a}{Q_a} - \frac{s_J}{Q_J}, \mu; \mu_S\right) \\ &\times \left[\sum_{q_i} L_{q_i} R^{q_i}(\mathcal{Q}, s_J - s'_J) J^{q_i}(s'_J, \mu; \mu_J) B_{q_i}(x_*, t_a, \mu; \mu_B) \right. \\ &\left. + \sum_{\bar{q}_i} L_{\bar{q}_i} R^{\bar{q}_i}(\mathcal{Q}, s_J - s'_J) J^{\bar{q}_i}(s'_J, \mu; \mu_J) B_{\bar{q}_i}(x_*, t_a, \mu; \mu_B) \right] \end{aligned} \quad (26)$$

From the above factorization formula, we can also obtain the 1-Jettiness distribution integrated over a jet charge bin. Correspondingly, we can define the charged jet function integrated over a jet charge bin in momentum and position space as

$$\begin{aligned} \mathcal{G}_{\text{bin}}^q(s, \mu) &= \int ds' R_{\text{bin}}^q(s - s') J^q(s', \mu), \\ \mathcal{G}_{\text{bin}}^q(y, \mu) &= R_{\text{bin}}^q(y) J^q(y, \mu), \end{aligned} \quad (27)$$

where the binned $R_{\text{bin}}^q(s)$ and $R_{\text{bin}}^q(y)$ functions are given by

$$\begin{aligned} R_{\text{bin}}^q(s) &= \int_{\text{bin}} d\mathcal{Q} R^q(\mathcal{Q}, s), \\ R_{\text{bin}}^q(y) &= \int_{\text{bin}} d\mathcal{Q} R^q(\mathcal{Q}, y). \end{aligned} \quad (28)$$

The factorization formula for the jet-charge-binned 1-Jettiness distribution is now given by

$$\begin{aligned} d\sigma_{\text{resum}}[\mathcal{Q}^{\text{bin}}, \tau_1, P_{J_T}, y_J] &= \sigma_0 H(\xi, \mu; \mu_H) \\ &\times \int ds_J \int dt_a \int ds'_J \mathcal{S}\left(\tau_1 - \frac{t_a}{Q_a} - \frac{s_J}{Q_J}, \mu; \mu_S\right) \\ &\times \left[\sum_{q_i} L_{q_i} R_{\text{bin}}^{q_i}(s_J - s'_J) J^{q_i}(s'_J, \mu; \mu_J) B_{q_i}(x_*, t_a, \mu; \mu_B) \right. \\ &\left. + \sum_{\bar{q}_i} L_{\bar{q}_i} R_{\text{bin}}^{\bar{q}_i}(s_J - s'_J) J^{\bar{q}_i}(s'_J, \mu; \mu_J) B_{\bar{q}_i}(x_*, t_a, \mu; \mu_B) \right] \end{aligned} \quad (29)$$

IV. 1-JETTINESS JET CHARGE DISTRIBUTION WITH POLARIZED PROTONS

The 1-Jettiness Jet Charge formalism can be straightforwardly extended to DIS with longitudinally polarized protons. The 1-Jettiness distribution of the longitudinally polarized nucleon asymmetry serves as a probe of the helicity PDFs. The \mathcal{Q} -binned 1-Jettiness distribution then allows enhanced flavor separation of the helicity PDFs.

The resummation formula for the 1-Jettiness distribution, τ_1 , of the longitudinally polarized nucleon asymmetry is schematically given by [88]

$$d\Delta\sigma_{\text{resum}}[\tau_1, P_{J_T}, y_J] \sim \Delta H \otimes \Delta B \otimes J \otimes \mathcal{S}, \quad (30)$$

where Δ denotes the polarization dependence in hard and beam functions needed to obtain the longitudinal nucleon

polarization asymmetry. The polarized beam function [7, 74, 88], ΔB , can be factorized as

$$\Delta B \sim \Delta \mathcal{I} \otimes \Delta f, \quad (31)$$

where $\Delta \mathcal{I}$ is perturbatively calculable and Δf denotes the helicity PDF. The corresponding factorization and resummation formula for the 1-Jettiness jet charge distribution, in addition to the standard 1-Jettiness distribution, is given by replacing the standard jet function, J , with the charged jet function, \mathcal{G} :

$$d\Delta\sigma_{\text{resum}}[\mathcal{Q}, \tau_1, P_{J_T}, y_J] \sim \Delta H \otimes \Delta B \otimes \mathcal{G} \otimes \mathcal{S}. \quad (32)$$

Note that the charged jet function \mathcal{G} that appears here is identical to the one that appears in the 1-Jettiness Jet Charge formalism with unpolarized protons, corresponding to its universality property. Similarly, studying the jet-charge-binned 1-Jettiness distribution of the longitudinally nucleon spin asymmetry facilitates disentangling the flavor structure of the helicity PDFs. We leave detailed phenomenological studies of the flavor structure of the helicity PDFs using the 1-Jettiness jet charge formalism for future work. In this work, we focus on introducing the main ideas and formalism of 1-Jettiness jet charge in DIS of unpolarized protons only.

V. UNIVERSALITY OF THE CHARGED JET FUNCTION

Eqs. (26) and (29) are key results, providing factorization formulae for the 1-Jettiness Jet Charge distribution. They give the fully differential cross section $d\sigma[\mathcal{Q}_J, \tau_1, P_{J_T}, y_J]$ and the jet-charge-binned cross section $d\sigma[\mathcal{Q}_J^{\text{bin}}, \tau_1, P_{J_T}, y_J]$, respectively, in the resummation region $\tau_1 \ll P_{J_T}$. The predictive power of this factorization framework is enhanced due to the universality of the various functions appearing in the factorization formulae. The hard function $H(\xi, \mu)$ describes universal QCD corrections to the hard scattering quark current vertex due to single photon or Z-boson exchange in standard fully inclusive DIS. It is perturbatively calculable with results known [81, 82] to $\mathcal{O}(\alpha_s^2)$. The universal quark beam functions [74], $B_q(x, t, \mu)$, describe the physics of perturbative initial state radiation, collinear with the proton beam direction, and the initial state collinear PDF, as seen in Eq. (16). The quark beam functions are also known [86, 87] to $\mathcal{O}(\alpha_s^2)$. The jet function describes the physics of the energetic radiation in the jet region, and collinear with the leading jet direction, and is known [81, 82, 85] to $\mathcal{O}(\alpha_s^2)$. Note that for the jet functions initiated by light quarks, the light quark flavor symmetry of QCD in the limit of massless quarks, leads to the additional universality, $J^{q_i}(s, \mu) = J^q(s, \mu)$, so that the jet functions are independent of quark flavor. The soft function $\mathcal{S}(\tau_1, \mu)$ describes the soft radiation, $E \sim \tau_1$, throughout the event. It is perturbatively calculable when $\mu \sim \tau_1 \gg \Lambda_{\text{QCD}}$ with results known [83, 84]

to $\mathcal{O}(\alpha_s^2)$. In the region $\mu \sim \tau_1 \sim \Lambda_{\text{QCD}}$ the soft function becomes non-perturbative and requires the introduction of a non-perturbative model. However, this soft function, and correspondingly the non-perturbative model, is universal in that it is the same soft function that appears in Eq. (14) for the standard 1-Jettiness distribution where no jet charge measurement is made.

The new ingredients in the 1-Jettiness Jet Charge factorization formula are the $R^q(Q, s)$ functions appearing in Eq. (26). In general, they are non-perturbative objects since the jet charge measurement is not collinear safe and sensitive to the hadronization process. However, these non-perturbative functions are universal and renormalization scale independent. They can be extracted from other observables that also make jet charge measurements on the jet regions of global event shapes. N-Jettiness [7] is an ideal global event shape for such purposes. It characterizes how closely the pattern of final state radiation resembles a N-jet like event. Small values of N-Jettiness, τ_N , correspond to events that closely resemble an exclusive N-Jet event. Large values of τ_N correspond to events with additional jets or energetic radiation in different directions. The resummation region corresponds to small values of N-Jettiness, $\tau_N \ll Q$, where Q denotes the hard scale in the problem. For e^+e^- colliders, in the resummation region, the N-Jettiness differential distribution has the schematic form

$$\frac{d\sigma_{\text{resum}}}{d\tau_N} \sim H \otimes J \otimes \cdots \otimes J \otimes S, \quad (33)$$

where H describes the hard scattering process, J denotes the standard jet function, and there are N such factors each corresponding to one of the N jet directions, and S denotes the soft function that describes soft radiation throughout the event. Note that the case of $N=2$ can be related to the thrust global event in e^+e^- colliders in the limit $\tau_2 \ll Q$. For hadron colliders, the factorization formula takes a similar form

$$\frac{d\sigma_{\text{resum}}}{d\tau_N} \sim H \otimes B \otimes B \otimes \cdots \otimes J \otimes S, \quad (34)$$

with the extra factors of the beam functions for each of the initial state hadrons. These formulae can be compared with Eq. (3) for the 1-Jettiness distribution at an electron-hadron collider, which involves only one beam function and one jet function.

In order to access the universal $\mathcal{G}^q(Q, s, \mu)$ jet charge function, and the corresponding $R^q(Q, s)$ functions needed for the 1-Jettiness Jet Charge distribution, one can generalize the N-Jettiness observables at e^+e^- and hadron colliders to include a jet charge measurement on one or more of the jet regions. For example, if a jet charge measurement is made on the jet region with the highest energy E_J or transverse momentum P_{J_T} , the formulae in Eqs. (33) and (34) become

$$\frac{d\sigma_{\text{resum}}}{d\tau_N dQ} \sim H \otimes \mathcal{G} \otimes \cdots \otimes J \otimes S, \quad (35)$$

and

$$\frac{d\sigma_{\text{resum}}}{d\tau_N dQ} \sim H \otimes B \otimes B \otimes \mathcal{G} \otimes \cdots \otimes J \otimes S, \quad (36)$$

respectively. The factor of the jet charge jet function, \mathcal{G} , replaces the standard jet function, J , in the jet region where the jet charge measurement is made. These formulae at e^+e^- and hadron colliders can be compared to Eq. (6) for the 1-Jettiness Jet Charge distribution, and the same universal jet charge jet function, \mathcal{G} , appears in all these observables.

One can also consider the N-Jettiness distributions in a give jet charge bin, where we integrate over the jet charge of measured jet region. The corresponding jet-charge-binned N-Jettiness distributions take the form

$$\frac{d\sigma_{\text{resum}}^{\mathcal{Q}\text{-bin}}}{d\tau_N} \sim H \otimes \mathcal{G}_{\text{bin}} \otimes \cdots \otimes J \otimes S, \quad (37)$$

for e^+e^- colliders and

$$\frac{d\sigma_{\text{resum}}^{\mathcal{Q}\text{-bin}}}{d\tau_N} \sim H \otimes B \otimes B \otimes \mathcal{G}_{\text{bin}} \otimes \cdots \otimes J \otimes S, \quad (38)$$

for hadron colliders. Here \mathcal{G}_{bin} denotes the jet charge jet function integrated over a jet charge bin and is the same as the one that appears in Eq. (7) for the jet-charge-binned 1-Jettiness distribution in electron-proton collisions.

Furthermore, one can also consider different definitions of the N-Jettiness event shapes, associated with the choice of parameters in their definitions. For example, the dimensionless τ_{1a} event shape [10, 12–14] differs from the definition of τ_1 in Eq. (10) in the choice of the $Q_{J,B}$ values. Similarly, one can work with dimension one τ_N or dimensionless \mathcal{T}_N . For all of these different definitions, corresponding to different partitioning of final state radiation into beam and jet regions, the same universal jet functions or charged jet functions appear in the factorization formulae.

Thus, the jet charge jet functions \mathcal{G}^q and $\mathcal{G}_{\text{bin}}^q$, and correspondingly the functions $R^q(Q, s)$ and $R_{\text{bin}}^q(s)$ are universal and can be extracted from a comprehensive and systematic global analysis of the wide range of N-Jettiness global event shape observables at e^+e^- and hadron colliders.

VI. NON-PERTURBATIVE MODEL FOR CHARGED JET FUNCTION

In order to calculate the 1-Jettiness jet charge distribution, we need to quantify the nonperturbative component of the charged jet function $\mathcal{G}^q(Q, y, \mu)$, or equivalently, its Fourier transform, $\mathcal{G}^q(Q, s, \mu)$. From Eqs. (21) and (22), this corresponds to specifying the nonperturbative functions, $R^q(Q, y)$, and their Fourier transform functions, $R^q(Q, s)$. In the next three subsections, we present the general parameterization of a phenomenological model

for $R^q(\mathcal{Q}, y)$ and $R^q(\mathcal{Q}, s)$ and consider the simplest case of these function being independent of y or s , as well as a phenomenological parameterization of additional y or s dependence beyond the simplest model.

A. General Model Parameterization

The form of the nonperturbative functions, $R^q(\mathcal{Q}, y)$ and $R^q(\mathcal{Q}, s)$, is constrained by the normalization conditions in Eq. (24). Given any arbitrary function $\mathcal{K}^q(\mathcal{Q}, y)$, the normalization condition on $R^q(\mathcal{Q}, y)$ can be satisfied by constructing it from $\mathcal{K}^q(\mathcal{Q}, y)$ as follows,

$$R^q(\mathcal{Q}, y) = \frac{\mathcal{K}^q(\mathcal{Q}, y)}{\int d\mathcal{Q} \mathcal{K}^q(\mathcal{Q}, y)}, \quad (39)$$

as long as $\int d\mathcal{Q} \mathcal{K}^q(\mathcal{Q}, y) \neq 0$. The corresponding $R^q(\mathcal{Q}, s)$ function is obtained via the Fourier transform of Eq. (39) with respect to the y variable, as seen in Eq. (23)

$$R^q(\mathcal{Q}, s) = \int \frac{dy}{2\pi} e^{iys} \left[\frac{\mathcal{K}^q(\mathcal{Q}, y)}{\int d\mathcal{Q} \mathcal{K}^q(\mathcal{Q}, y)} \right], \quad (40)$$

from which it follows that $R^q(\mathcal{Q}, s)$ satisfies the required normalization condition in Eq. (24).

B. Simple Model Parameterization

In this section, we consider the simplest case where the charged jet functions $\mathcal{G}^q(\mathcal{Q}, y, \mu)$ or $\mathcal{G}^q(\mathcal{Q}, s, \mu)$ having the same y or s dependence as the standard jet functions, $J^q(y, \mu)$ or $J^q(s, \mu)$, respectively. This corresponds to the $R^q(\mathcal{Q}, y)$ being independent of y , as seen from Eqs. (20). Equivalently, $R^q(\mathcal{Q}, s)$ will be proportional to $\delta(s)$, which follows from Eq. (23). In this special case, the $R^q(\mathcal{Q}, y)$ and $R^q(\mathcal{Q}, s)$ functions can be written as

$$R^q(\mathcal{Q}, y) = r^q(\mathcal{Q}), \quad R^q(\mathcal{Q}, s) = r^q(\mathcal{Q})\delta(s), \quad (41)$$

where the function $r^q(\mathcal{Q})$ satisfies the normalization condition

$$\int d\mathcal{Q} r^q(\mathcal{Q}) = 1. \quad (42)$$

From Eqs. (21) and (22), the charged jet functions then reduce to the separable form

$$\begin{aligned} \mathcal{G}^q(\mathcal{Q}, y, \mu) &= r^q(\mathcal{Q}) J^q(y, \mu), \\ \mathcal{G}^q(\mathcal{Q}, s, \mu) &= r^q(\mathcal{Q}) J^q(s, \mu), \end{aligned} \quad (43)$$

where the charged jet function, \mathcal{G} , and the ordinary jet function, J , having the same dependence in the y or s variables. Using this model in Eqs. (41) and (43), the

	u	\bar{u}	d	\bar{d}	s	\bar{s}
r_+^i	0.62	0.18	0.13	0.50	0.19	0.52
r_-^i	0.08	0.50	0.48	0.13	0.43	0.13
r_0^i	0.30	0.32	0.39	0.37	0.38	0.35

TABLE I. Standard Jet Charge binned fractions for different quark flavors. The relevant EIC kinematics chosen are: $\sqrt{s} = 90.0$ GeV, $P_{JT} = [20.0, 30.0]$ GeV, and $y_J = [-2.5, 2.5]$.

factorization formula in Eq. (26) can be written as

$$\begin{aligned} d\sigma_{\text{resum}}[\mathcal{Q}, \tau_1, P_{JT}, y_J] &= \sigma_0 H(\xi, \mu; \mu_H) \\ &\times \int ds_J \int dt_a \mathcal{S}\left(\tau_1 - \frac{t_a}{Q_a} - \frac{s_J}{Q_J}, \mu; \mu_S\right) \\ &\times \left[\sum_{q_i} L_{q_i} r^{q_i}(\mathcal{Q}) J^{q_i}(s_J, \mu; \mu_J) B_{q_i}(x_*, t_a, \mu; \mu_B) \right. \\ &\left. + \sum_{\bar{q}_i} L_{\bar{q}_i} r^{\bar{q}_i}(\mathcal{Q}) J^{\bar{q}_i}(s_J, \mu; \mu_J) B_{\bar{q}_i}(x_*, t_a, \mu; \mu_B) \right], \end{aligned} \quad (44)$$

where the jet charge information is now contained entirely in the $r^q(\mathcal{Q})$ functions. The jet-charge-binned cross section in Eq. (29) then takes the following form

$$\begin{aligned} d\sigma_{\text{resum}}[\mathcal{Q}_J^{\text{bin}}, \tau_1, P_{JT}, y_J] &= \sigma_0 H(\xi, \mu; \mu_H) \\ &\times \int ds_J \int dt_a \mathcal{S}\left(\tau_1 - \frac{t_a}{Q_a} - \frac{s_J}{Q_J}, \mu; \mu_S\right) \\ &\times \left[\sum_{q_i} L_{q_i} r_{\text{bin}}^{q_i} J^{q_i}(s_J, \mu; \mu_J) B_{q_i}(x_*, t_a, \mu; \mu_B) \right. \\ &\left. + \sum_{\bar{q}_i} L_{\bar{q}_i} r_{\text{bin}}^{\bar{q}_i} J^{\bar{q}_i}(s_J, \mu; \mu_J) B_{\bar{q}_i}(x_*, t_a, \mu; \mu_B) \right], \end{aligned} \quad (45)$$

where the r_{bin}^q constants are given by the integration of the $r^q(\mathcal{Q})$ function over the jet charge bin

$$r_{\text{bin}}^q = \int_{\text{bin}} d\mathcal{Q} r^q(\mathcal{Q}), \quad (46)$$

and satisfy the constraint

$$\sum_{\text{bins}} r_{\text{bin}}^q = 1, \quad (47)$$

for each quark and anti-quark flavor, q , as required by Eq. (42). Thus, the r_{bin}^q values can be interpreted as the fraction of jets initiated by a parton of flavor q in the given jet charge bin. Based on the PYTHIA 8.312 simulations of five million events of the unpolarized DIS process, $e^- + p \rightarrow e^- + J + X$, for the typical EIC kinematics of $\sqrt{s} = 90.0$ GeV, $P_{JT} = [20.0, 30.0]$ GeV, and $y_J = [-2.5, 2.5]$, we show in Tables I and II the r_{bin}^q bin fractions for the Standard and Dynamic jet charge definitions, respectively. The $r_{+, -, 0}^q$ bin fractions correspond to the fraction of jets in the positive ($\mathcal{Q} > 0.25$), negative ($\mathcal{Q} < -0.25$), and central ($-0.25 < \mathcal{Q} < 0.25$) jet charge bins, respectively.

	u	\bar{u}	d	\bar{d}	s	\bar{s}
r_+^i	0.39	0.15	0.15	0.29	0.15	0.35
r_-^i	0.11	0.36	0.29	0.15	0.33	0.14
r_0^i	0.49	0.49	0.55	0.55	0.52	0.51

TABLE II. Dynamic Jet Charge binned fractions for different quark flavors. The relevant EIC kinematics chosen are: $\sqrt{s} = 90.0$ GeV, $P_{JT} = [20.0, 30.0]$ GeV, and $y_J = [-2.5, 2.5]$.

We can write the jet-charge-binned 1-Jettiness Jet Charge cross section in the resummation region as

$$\begin{aligned} d\sigma_{\text{resum}}[Q^{\text{bin}}, \tau_1, P_{JT}, y_J] \\ = \sum_q r_{\text{bin}}^q d\sigma_{\text{resum}}^q[\tau_1, P_{JT}, y_J], \end{aligned} \quad (48)$$

where the sum over q runs over all the quark and anti-quark flavors. The cross section $d\sigma_{\text{resum}}^q[\tau_1, P_{JT}, y_J]$ corresponds to the contribution from the q -parton flavor to the standard 1-Jettiness distribution:

$$\begin{aligned} d\sigma_{\text{resum}}^q[\tau_1, P_{JT}, y_J] &= \sigma_0 L_q H(\xi, \mu; \mu_H) \\ &\times \int ds_J \int dt_a J^q(s_J, \mu; \mu_J) B_q(x_*, t_a, \mu; \mu_B) \\ &\times \mathcal{S}\left(\tau_1 - \frac{t_a}{Q_a} - \frac{s_J}{Q_J}, \mu; \mu_S\right), \end{aligned} \quad (49)$$

while r_{bin}^q determines the weight given to this contribution based on the fraction of jets initiated by the corresponding parton flavor in the given jet charge bin.

We note that the form of Eq. (48) is similar to that in Ref. [64] which studied $e^- + p \rightarrow e^- + J + X$, in the limit of small electron-jet transverse momentum imbalance, $q_T = |\vec{p}_T^e + \vec{P}_{JT}| \ll \vec{p}_T^e \sim P_{JT}$, in different jet charge bins to probe the flavor structure of TMDPDFs. There the jet was defined with a jet algorithm and fixed jet radius. In our case, where the jet charge binning is applied to a jet region of a global event shape, the form in Eq. (48) only applies in the special limit of Eq. (43) where the charged jet function has the same y or s dependence as the standard jet function. The more general parameterization allows for a more non-trivial dependence of the charged jet function, $\mathcal{G}(s, \mu)$ and $\mathcal{G}(y, \mu)$ on the variables s or y that goes beyond the dependence in the standard jet function $J(s, \mu)$ and $J(y, \mu)$, respectively. In general, this leads to the non-trivial convolution structure in Eqs. (26) or (29).

C. A Useful Model Parameterization

The form of the jet-charge-binned 1-Jettiness cross section in Eq. (48) is the direct result of the special case of the nonperturbative model, as shown in Eq. (41). In particular, the r_{bin}^q parameters are constants, independent of




FIG. 2. The fraction of jets initiated by the u -quark (top panel) and d -quark (bottom panel) in the positive ($Q_J > 0.25$), negative ($Q_J < -0.25$), and central ($-0.25 < Q_J < 0.25$) Standard Jet Charge bins as a function of the squared jet mass, M_J^2 . The dashed lines indicate the corresponding average values from the fraction of jets over all squared jet mass bins. The relevant EIC kinematics chosen for generating the Pythia 8.312 simulation data are: $\sqrt{s} = 90.0$ GeV, $P_{JT} = [20.0, 30.0]$ GeV, and $y_J = [-2.5, 2.5]$.

s . We test how well this model works using PYTHIA simulations. Note that the variable s is the mass squared of all the energetic collinear particles in the jet region, arising from the charged jet function, $\mathcal{G}(s, \mu)$. However, the true measurable jet mass squared of the jet region also includes the contribution of soft particles in the jet region. The jet mass squared can be written as $M_J^2 = (p_c + p_s)^2$ where p_c^μ and p_s^μ denote the total sum of the momenta of the collinear and soft particles, respectively. We can write $M_J^2 = s + 2p_c \cdot p_s + p_s^2$, where $s = p_c^2$. We note that while the soft particles in the jet region make a significant contribution to jet mass, M_J , via the $2p_c \cdot p_s \sim s$ term, their contribution to the jet charge is suppressed since the contribution of each particle is weighed by its energy as seen in Eq. (1). Thus, the fraction, r_{bin}^q , of jets initiated by parton flavor, q , in a given jet charge bin is primarily determined by the energetic collinear particles,




FIG. 3. The fraction of jets initiated by the u -quark (top panel) and d -quark (bottom panel) in the positive ($Q_J > 0.25$), negative ($Q_J < -0.25$), and central ($-0.25 < Q_J < 0.25$) Dynamic Jet Charge bins as a function of the squared jet mass, M_J^2 . The dashed lines indicate the corresponding average values from the fraction of jets over all squared jet mass bins. The relevant EIC kinematics chosen for generating the Pythia 8.312 simulation data are: $\sqrt{s} = 90.0$ GeV, $P_{JT} = [20.0, 30.0]$ GeV, and $y_J = [-2.5, 2.5]$.

with the contributions from soft particles being power suppressed. Thus, the behavior of r_{bin}^q as a function of $M_J^2 = (p_c + p_s)^2$ will be similar to that as a function of $s = p_c^2$ such that we can write

$$r_{\text{bin}}^q(M_J^2) \simeq r_{\text{bin}}^q(s). \quad (50)$$

In Figs. 2 and 3, we show the jet mass dependence of the jet charge bin fractions based on PYTHIA 8.312 simulations for $r_{+,0}^u(M_J^2) \simeq r_{+,0}^u(s)$ and $r_{+,0}^d(M_J^2) \simeq r_{+,0}^d(s)$ for the Standard and Dynamic jet charge definitions, respectively. We again choose typical EIC kinematics with $\sqrt{s} = 90.0$ GeV, $P_{JT} = [20.0, 30.0]$ GeV, and $y_J = [-2.5, 2.5]$.

The results in Figs. 2 and 3 suggest that the r_{bin}^q do have a non-trivial dependence on M_J^2 , and correspondingly suggest a non-trivial dependence on s . This dependence is stronger for the Dynamic Jet Charge compared

to the Standard Jet Charge. These results show that the simplified charged jet function model in Eq. (43) which predicts constant r_{bin}^q values that are independent of s , is only an approximation and the more general model parameterization in Eqs. (39) and (40) is appropriate, where $\mathcal{K}(\mathcal{Q}, y)$ has a non-trivial y -dependence.

However, as seen in Figs. 2 and 3, while the r_{bin}^q values are not constant, the dependence on M_J^2 , or equivalently on s , is still mild. This suggests a model parameterization for $R^q(\mathcal{Q}, y)$ as a sum of a constant term and a y -dependent correction term as

$$R^q(\mathcal{Q}, y) = r^q(\mathcal{Q}) + f^q(\mathcal{Q}, y). \quad (51)$$

The y -dependent function $f^q(\mathcal{Q}, y)$ characterizes the deviation of $R^q(\mathcal{Q}, y)$ from being a constant, independent of y as characterized by the $r^q(\mathcal{Q})$ term. Equivalently, as seen from Eq. (21), $f^q(\mathcal{Q}, y)$ describes the additional y -dependence in the charged jet function, $\mathcal{G}^q(\mathcal{Q}, y, \mu)$, compared to that of the standard jet function, $J^q(y, \mu)$. In order to satisfy the normalization condition on $R^q(\mathcal{Q}, y)$ in Eq. (24), and given the normalization condition already satisfied by $r^q(\mathcal{Q})$ in Eq. (42), the function $f^q(\mathcal{Q}, y)$ must satisfy

$$\int d\mathcal{Q} f^q(\mathcal{Q}, y) = 0. \quad (52)$$

Since this condition on $f^q(\mathcal{Q}, y)$ must be satisfied for all values of y , it must either be a function that is separable in the variables y and \mathcal{Q} , or more generally, be built out of a sum of separable functions as

$$f^q(\mathcal{Q}, y) = \sum_n g_n^q(y) \chi_n^q(\mathcal{Q}). \quad (53)$$

The functions $\chi_n^q(\mathcal{Q})$ can be chosen to form a complete set of basis functions, satisfy the conditions

$$\begin{aligned} \int d\mathcal{Q} \chi_n^q(\mathcal{Q}) &= 0, \\ \int d\mathcal{Q} \chi_n^q(\mathcal{Q}) \chi_m^q(\mathcal{Q}) &= \delta_{m,n}. \end{aligned} \quad (54)$$

The first condition ensures the required constraint in Eq. (52). The second orthonormality condition allows the coefficient functions, $g^q(y)$, for a given $f^q(\mathcal{Q}, y)$, to be determined as

$$g_n^q(y) = \int d\mathcal{Q} f^q(\mathcal{Q}, y) \chi_n^q(\mathcal{Q}). \quad (55)$$

In momentum space, the $R^q(\mathcal{Q}, s)$ function is parameterized as

$$R^q(\mathcal{Q}, s) = r^q(\mathcal{Q}) \delta(s) + f^q(\mathcal{Q}, s), \quad (56)$$

where $f^q(\mathcal{Q}, s)$ are the Fourier transforms of $f^q(\mathcal{Q}, y)$

$$f^q(\mathcal{Q}, s) = \int \frac{dy}{2\pi} e^{iys} f^q(\mathcal{Q}, y), \quad (57)$$

and $f^q(Q, s)$ satisfy the condition

$$\int dQ f^q(Q, s) = 0, \quad (58)$$

which follows from Eq. (52), and is required by the underlying normalization condition in Eq. (24). The momentum space function $f^q(Q, s)$ can also be expressed as a sum over separable functions in terms of a complete set of basis functions as

$$f^q(Q, s) = \sum_n g_n^q(s) \chi_n^q(Q), \quad (59)$$

where the coefficient functions $g_n^q(s)$ are related to the $g_n^q(y)$ functions via the Fourier transform as

$$g_n^q(s) = \int \frac{dy}{2\pi} e^{iys} g_n^q(y). \quad (60)$$

Having identified some of the key structural features of the nonperturbative functions, $R(Q, y)$ and $R(Q, s)$, we leave a more detailed model building analysis for future work.

VII. NUMERICAL RESULTS

In this section, we demonstrate the sensitivity of the 1-Jettiness Jet Charge, Q , to probe quark flavor dynamics in nucleon structure and explore ways of probing the hadronization process. All the numerical results presented here are based on the PYTHIA 8.312 simulations of five million events of the unpolarized DIS process, $e^- + p \rightarrow e^- + J + X$, for the typical EIC kinematics of $\sqrt{s} = 90.0$ GeV, $P_{JT} = [20.0, 30.0]$ GeV, and $y_J = [-2.5, 2.5]$. In this analysis, we only present results for unpolarized DIS which probe the unpolarized PDFs of the initial state nucleon. These results can be straightforwardly generalized to studies with longitudinally polarized initial state nucleons to probe the polarized PDFs, using the formalism described in section IV.

We consider the standard jet charge definition with constant $\kappa = 0.3$ in Eq. (1). We will also present results based on the dynamic jet charge [71] definition, corresponding to $\kappa = \kappa(z_h, \xi_{\text{cut}}, k_<, k_>)$ where

$$\kappa(z_h) = \begin{cases} k_<, & z_h < \xi_{\text{cut}} \\ k_>, & z_h \geq \xi_{\text{cut}} \end{cases} \quad (61)$$

and ξ_{cut} , $k_<$, and $k_>$ are three constant parameters. We choose the default values of $\xi_{\text{cut}} = 0.3$, $k_< = 1.0$, $k_> = 0.3$. The standard and dynamic jet charge definitions have different jet discrimination capabilities and sensitivity to power corrections from soft radiation effects, thereby serving as complementary probes of quark flavor dynamics in nuclear structure and hadronization.




FIG. 4. The normalized 1-Jettiness τ_1 -distribution (black) using Pythia data along with the relative contributions of different quark flavors. The relevant EIC kinematics chosen are: $\sqrt{s} = 90.0$ GeV, $P_{JT} = [20.0, 30.0]$ GeV, and $y_J = [-2.5, 2.5]$.




FIG. 5. The normalized 1-Jettiness τ_1 -distribution (black) using Pythia data along with the relative contributions from the negative (blue), central (gray), and positive (red) standard jet charge bins corresponding to $Q < -0.25$, $-0.25 < Q < 0.25$, and $Q > 0.25$, respectively. The relevant EIC kinematics chosen are: $\sqrt{s} = 90.0$ GeV, $P_{JT} = [20.0, 30.0]$ GeV, and $y_J = [-2.5, 2.5]$.

A. Probing Nucleon Flavor Structure

In Fig. 4, we show the normalized 1-Jettiness distribution (upper black curve). We also show the contributions to the 1-Jettiness distribution from scattering off quarks of different flavors in the initial state nucleon. We see that the 1-Jettiness distribution is overwhelmingly dominated by the u -quark flavor due to the dominance of its unpolarized PDF in the proton in the valence region. As a result, this distribution is not very sensitive to the dynamics of the d -quarks and the sea quarks and anti-quarks.




FIG. 6. The normalized 1-Jettiness τ_1 -distribution (black) using Pythia data in the positive ($Q > 0.25$, top panel), central ($-0.25 < Q < 0.25$, middle panel), and negative ($Q < -0.25$, bottom panel) Standard Jet Charge bins, along with the relative contributions of different quark flavors. The relevant EIC kinematics chosen are: $\sqrt{s} = 90.0$ GeV, $P_{JT} = [20.0, 30.0]$ GeV, and $y_J = [-2.5, 2.5]$.

One can gain enhanced sensitivity to the u -antiquark and the d - and s -quark unpolarized PDFs by binning the 1-Jettiness distribution according to the value of the 1-Jettiness Standard Jet Charge, Q . In Fig. 5, we show the 1-Jettiness distribution binned according to Q . We see the relative contribution to the total 1-Jettiness distribution from each jet charge bin. The dominant contribution comes from the positive ($Q > 0.25$) jet charge




FIG. 7. The normalized 1-Jettiness τ_1 -distribution (black) using Pythia data in the positive ($Q > 0.25$, top panel), the central ($-0.25 < Q < 0.25$, middle panel) and negative ($Q < -0.25$, bottom panel) Dynamic Jet Charge bins, along with the relative contributions of different quark flavors. The relevant EIC kinematics chosen are: $\sqrt{s} = 90.0$ GeV, $P_{JT} = [20.0, 30.0]$ GeV, and $y_J = [-2.5, 2.5]$.

bin, corresponding to the fact that the 1-Jettiness distribution is dominated by the contribution from the u -quark which initiates jets with a standard jet charge distribution peaked in the positive jet charge bin. Note that u -quark initiated jets still have a non-zero probability of having a standard jet charge value in the negative ($Q < -0.25$) or central ($-0.25 < Q < 0.25$) jet charge bins, as indicated by the fact that the contribution of the

positive jet charge bin is not as dominant compared to the contribution from the u -quark in Fig. 4.

In Fig. 6, we show the relative contributions to the 1-Jettiness distribution from different quark flavors in each standard jet charge bin. We see that in the positive (top panel) and central (middle panel) jet charge bin, the 1-Jettiness distribution is overwhelmingly dominated by the u -quark contribution. However, we see that in the negative (bottom panel) jet charge bin, the contribution from the d -quark is comparable to that from the u -quark, indicating that one has enhanced sensitivity to the d -quark PDF when the 1-Jettiness distribution is restricted to the negative jet charge bin. In Fig. 7, we show the corresponding results using the dynamic jet charge definition. Once again, we see enhanced sensitivity to the d -quark PDF in the negative dynamic jet charge bin. Similarly, studying the 1-Jettiness distribution for DIS on longitudinally polarized protons, according to jet charge bins can help disentangle the flavor structure of the helicity PDFs. In this manner, the 1-Jettiness jet charge measurement can be used as a new probe of the quark flavor structure in the nucleon.

B. Probing Flavor Dynamics in Hadronization

One can also consider the 1-Jettiness jet charge distribution binned according to the 1-Jettiness value characterizing the shape of the energy flow in each event. This allows us to study the jet charge, Q , as a function of the shape of the overall energy flow as characterized by τ_1 . Such studies can provide constraints on hadronization models, allowing us to probe correlations between flavor dynamics and energy flow. We use the same PYTHIA8.312 simulation data for the EIC kinematics of $\sqrt{s} = 90.0$ GeV, $P_{JT} = [20.0, 30.0]$ GeV, and $y_J = [-2.5, 2.5]$ to explore observables that can probe hadronization models.

In Fig. 8, we show the total Q -distribution (black) and the Q -distribution binned (colored) in different τ_1 -bins for the standard (top panel) and dynamic (bottom panel) jet charge definitions. The total unbinned Q -distribution (black) is normalized to unity. The τ_1 -binned Q -distributions in turn are normalized to the unbinned distribution so that we can see the relative fraction of events in the different τ_1 -bins, characterizing the pattern of energy flow, as a function of the jet region charge.

Another potentially interesting observable is the average jet charge as a function of 1-Jettiness. Theoretically, it can be defined via the probability distribution in the 1-Jettiness jet charge, Q :

$$P(Q, \tau_1^{\text{bin}}, P_{JT}^{\text{bin}}, y_J^{\text{bin}}) = \frac{d\sigma[Q, \tau_1^{\text{bin}}, P_{JT}^{\text{bin}}, y_J^{\text{bin}}]}{d\sigma[\tau_1^{\text{bin}}, P_{JT}^{\text{bin}}, y_J^{\text{bin}}]}, \quad (62)$$

where the bin superscript indicates an integration over the corresponding bin. Note that $d\sigma[Q, \tau_1 P_{JT}, y_J]$ and




FIG. 8. 1-Jettiness Jet Charge distributions according to 1-Jettiness bins for the Standard Jet Charge (top panel) and the Dynamic Jet Charge (bottom panel). The binned curves are normalized to the total unbinned jet charge distribution curve which in turn is normalized to unity over the displayed range. The relevant EIC kinematics chosen are: $\sqrt{s} = 90.0$ GeV, $P_{JT} = [20.0, 30.0]$ GeV, and $y_J = [-2.5, 2.5]$.

$d\sigma[Q, \tau_1, P_{JT}, y_J]$ are defined in Eqs. (5) and (2), respectively. This probability distribution satisfies the required normalization condition

$$\int dQ P(Q, \tau_1^{\text{bin}}, P_{JT}^{\text{bin}}, y_J^{\text{bin}}) = 1. \quad (63)$$

This probability distribution can be used to define moments of the jet charge. The n -th moment is defined as

$$\langle Q^n \rangle = \int dQ Q^n P(Q, \tau_1^{\text{bin}}, P_{JT}^{\text{bin}}, y_J^{\text{bin}}), \quad (64)$$

with the first moment corresponding to the average jet charge

$$Q_{\text{avg}}(\tau_1^{\text{bin}}, P_{JT}^{\text{bin}}, y_J^{\text{bin}}) = \langle Q \rangle. \quad (65)$$

Similarly, the standard deviation of the jet charge in any bin is given by

$$Q_{\text{std.dev}}(\tau_1^{\text{bin}}, P_{JT}^{\text{bin}}, y_J^{\text{bin}}) = \sqrt{\langle Q^2 \rangle - \langle Q \rangle^2}. \quad (66)$$

VIII. CONCLUSIONS




FIG. 9. The average (top panel) and standard deviation (bottom panel) values of jet charge distributions binned according to the value of 1-Jettiness, τ_1 . The relevant EIC kinematics chosen are: $\sqrt{s} = 90.0$ GeV, $P_{J_T} = [20.0, 30.0]$ GeV, and $y_J = [-2.5, 2.5]$.

Using PYTHIA 8.312 simulation data, in Fig. 9 we show the jet region charge average (top panel) and standard deviation (bottom panel) as a function of τ_1 for the standard (blue) and dynamic (red) jet charge definitions. We see that the standard and dynamic jet charge definitions give rise to different behavior of the jet charge moments as a function of τ_1 . Such correlations between the pattern of energy flow, characterized by the τ_1 global event shape, and the jet region charge, Q , could probe hadronization models. In particular, the distribution of various moments of the 1-Jettiness jet charge with different jet charge definitions, and over a wide range of (τ_1, P_{J_T}, y_J) bins could provide non-trivial constraints on hadronization models. We leave a more detailed study exploring these distributions for future work.

We have proposed bringing together the flavor discrimination power of the jet charge observable with global event shapes that characterize the shape and pattern of energy flow throughout the event. This framework provides a new way to probe the quark flavor structure of the nucleon and the quark flavor dynamics in the hadronization process. It allows for a measurement of the jet region charge while simultaneously keeping track of the energy flow throughout the event, as characterized by the global event shape.

We focused on the measurement of the 1-Jettiness jet charge (Q), the jet charge of the jet region (J) defined within the framework of the 1-Jettiness global event shape (τ_1) for the Deep Inelastic Scattering (DIS) process, $e^- + p \rightarrow e^- + J + X$, with unpolarized or longitudinally polarized protons. The 1-Jettiness distribution, τ_1 , binned according to jet charge, Q , allows for enhanced quark flavor separation of the initial state unpolarized or polarized PDFs. Alternatively, the jet charge distribution binned by 1-Jettiness serves as a probe of quark flavor dynamics in the final state hadronization process. We derived a factorization theorem that simultaneously measures τ_1 and Q in the resummation region, $\tau_1 \ll P_{J_T}$, where the correlation between the jet charge and the charge of the struck quark is the strongest (P_{J_T} denotes the transverse momentum of the jet region). We also discussed the formulation of power corrections to this factorization framework. We considered both the standard and dynamic jet charge definitions in our analysis, serving as complementary probes with different jet discrimination capabilities and sensitivity to power corrections from soft radiation.

The factorization theorem gives rise to new universal and field theoretically well-defined *charged jet functions* that describe the evolution of the struck quark into a final state of given jet mass and charge values. The charged jet functions are non-perturbative but have the same renormalization group evolution as the ordinary jet function. We developed a general model parameterization for the non-perturbative component of these charged jet functions and studied its general structure and properties. We described how these universal functions can be extracted from a wide range of N-Jettiness global event shape observables at e^+e^- colliders, including the thrust global event shape. We provided numerical results through simulations studies using PYTHIA 8.312 to demonstrate the sensitivity of this observable to probe flavor dynamics in nucleon structure and explore the possibility of probing quark flavor dynamics in the final state hadronization process.

The 1-Jettiness jet charge framework can be applied to longitudinal nucleon spin asymmetries to probe the flavor structure of helicity PDFs. It can also be applied to charged current DIS, providing additional handles on constraining quark flavor dynamics in the nucleon and the hadronization process. This observable is ideally

suit for the proposed EIC and can be applied to existing HERA data. We leave such further theoretical and phenomenological studies of the 1-Jettiness jet charge observable for future work.

IX. ACKNOWLEDGEMENTS

The work of YTC is supported in part by the US Department of Energy (DOE) Contract No. DEAC05-06OR23177, under which Jefferson Science Associates, LLC operates Jefferson Lab, and by the Department of Energy Early Career Award grant DE-SC0023304. SM thanks Jefferson Lab for their hospitality and support through the Visiting Faculty Program (VFP) during which part of this work was carried out.

Appendix A: Charged Jet Function

The field-theoretic definition of the jet functions is given by

$$J^{q_i}(s_J = \omega_J r^+, \mu) = \frac{1}{4\pi N_c \omega_J} \text{Im} \left[i \int d^4 x e^{ir \cdot x} \right. \\ \left. \times \langle 0 | T \{ \bar{\chi}_n^{q_i}(0) \delta(\omega_J - \bar{P}^\dagger) \not{n} \chi_n^{q_i}(x) \} | 0 \rangle \right]. \quad (\text{A1})$$

The charged jet function is defined with an additional insertion of a jet charge measurement function in the above expression,

$$\mathcal{G}^q(\mathcal{Q}, s_J = \omega_J r^+, \mu) = \frac{1}{4\pi N_c \omega_J} \text{Im} \left[i \int d^4 x e^{ir \cdot x} \right. \\ \left. \times \langle 0 | T \{ \bar{\chi}_n^{q_i}(0) \delta(\omega_J - \bar{P}^\dagger) \delta(\mathcal{Q} - \hat{\mathcal{Q}}) \not{n} \chi_n^{q_i}(x) \} | 0 \rangle \right], \quad (\text{A2})$$

where we have used the Soft-Collinear Effective Theory (SCET) notation and $\hat{\mathcal{Q}}$ denotes the jet charge measurement operator

$$\mathcal{Q}|X\rangle = Q_X|X\rangle, \quad \langle X|\mathcal{Q}^\dagger = \langle X|Q_X, \quad (\text{A3})$$

where Q_X denotes the total jet charge of the state $|X\rangle$ and the jet charge operator is Hermitian, $\mathcal{Q}^\dagger = \mathcal{Q}$, with real eigenvalues.

Appendix B: Power Corrections to 1-Jettiness Jet Charge

The factorization formula in Eq. (17), in the resummation region $\tau_1 \ll P_{J_T}$, will receive power corrections from the contribution of soft particles to the jet charge. Since the jet charge definition includes a weight of the energy fractions of the particles in the jet region, as seen in Eq. (1), the contribution of soft particles ($z_h \sim \tau_1/P_{J_T}$) to the jet charge is formally suppressed compared to that of the collinear particles ($z_h \sim 1$) by $(\tau_1/P_{J_T})^\kappa$. Thus, for given choices of the τ_1 region and the parameter, κ ,

the effects of the power corrections can be suppressed by choosing a large enough value of P_{J_T} . We also note that the effects of power corrections are expected to be further suppressed in the dynamic jet charge definition where $\kappa = \kappa(z_h)$ is a dynamic function of z_h that allows one further suppress the contributions of soft particles, compared to the standard jet charge definition. In fact, it has been shown [71] that the dynamic jet charge is very insensitive to soft radiation by noting that the jet discrimination power of the dynamic jet charge is essentially the same in pp-collisions and PbPb-collisions even though the latter have significantly more soft contamination from MPI. By contrast, some degradation in the discrimination power was observed for the standard jet charge in PbPb-collisions compared to pp-collisions.

In this appendix, we articulate the general form of the power correction due to the contribution to the jet region charge from soft particles, in the resummation region, $\tau_1 \ll P_{J_T}$. The 1-Jettiness jet charge, \mathcal{Q} , measurement is incorporated by the insertion the delta function $\delta(\mathcal{Q} - \mathcal{Q}_c^J - \mathcal{Q}_s^J)$ in the 1-Jettiness factorization framework that requires that the sum of the jet charge contributions from the collinear and soft particles add up to the measured jet region charge, \mathcal{Q} . Here \mathcal{Q}_c^J and \mathcal{Q}_s^J denote the contribution from the collinear and soft particles in the jet region. Correspondingly, the delta functions $\delta(\mathcal{Q}_c^J - \hat{\mathcal{Q}}_c^J)$ and $\delta(\mathcal{Q}_s^J - \hat{\mathcal{Q}}_s^J)$ are inserted in the field-theoretic matrix element definitions of the jet function and soft function to obtain the charged jet function, $\mathcal{G}^q(\mathcal{Q}_c^J, s, \mu)$, and charged soft function $\mathcal{S}(\tau_1, \mathcal{Q}_s^J, \mu)$, respectively. Here $\hat{\mathcal{Q}}_c^J$ and $\hat{\mathcal{Q}}_s^J$ are the jet region charge operators that act on the collinear and soft particles in the jet region to obtain the charges \mathcal{Q}_c^J and \mathcal{Q}_s^J , respectively. For the charged jet function, this is seen in Eq. (A2).

The standard soft function is given in terms of the generalized hemisphere soft function [8–10] as

$$\mathcal{S}(\tau_1, \mu) = \int dk_a \int dk_J \delta(\tau_1 - k_a - k_J) \\ \times \mathcal{S}(k_a, k_J, \mu), \quad (\text{B1})$$

where hemisphere soft function is given by

$$\mathcal{S}(k_a, k_J, \mu) = \frac{1}{N_c} \sum_{X_s} \text{Tr} \langle 0 | \bar{T} [Y_{n_A}^\dagger Y_{n_J}] (0) \\ \times \delta(k_a - \frac{q_A \cdot K_{X_s}^{(a)}}{Q_a}) \delta(k_J - \frac{q_J \cdot K_{X_s}^{(J)}}{Q_J}) | X_s \rangle \\ \times \langle X_s | T [Y_{n_J}^\dagger Y_{n_A}] (0) | 0 \rangle. \quad (\text{B2})$$

Here we have defined the total soft momentum in the jet and beam regions as

$$K_{X_s}^{(J)} = \sum_{k \in X_s} p_k \theta(\frac{2q_A \cdot p_k}{Q_a} - \frac{2q_J \cdot p_k}{Q_J}), \\ K_{X_s}^{(a)} = \sum_{k \in X_s} p_k \theta(\frac{2q_J \cdot p_k}{Q_J} - \frac{2q_A \cdot p_k}{Q_a}), \quad (\text{B3})$$

respectively. The Y_{n_J} and Y_{n_A} denote soft Wilson lines that sum up soft eikonal emissions along the jet and beam directions, respectively. Correspondingly, the charged soft function can be obtained via

$$\mathcal{S}(\tau_1, \mathcal{Q}_s^J, \mu) = \int dk_a \int dk_J \delta(\tau_1 - k_a - k_J) \times \mathcal{S}(k_a, k_J, \mathcal{Q}_s^J, \mu), \quad (\text{B4})$$

where $\mathcal{S}(k_a, k_J, \mathcal{Q}_s^J, \mu)$ denotes the charged hemisphere function which involves the jet charge measurement operator $\delta(\mathcal{Q}_s^J - \hat{\mathcal{Q}}_s^J)$ acting on the soft particles in the jet region as

$$\begin{aligned} \mathcal{S}(k_a, k_J, \mu) &= \frac{1}{N_c} \sum_{X_s} \text{Tr} \langle 0 | \bar{T} [Y_{n_A}^\dagger Y_{n_J}] | 0 \rangle \\ &\times \delta(k_a - \frac{q_A \cdot K_{X_s}^{(a)}}{Q_a}) \delta(k_J - \frac{q_J \cdot K_{X_s}^{(J)}}{Q_J}) | X_s \rangle \\ &\times \langle X_s | \delta(\mathcal{Q}_s^J - \hat{\mathcal{Q}}_s^J) T[Y_{n_J}^\dagger Y_{n_A}] | 0 \rangle | 0 \rangle. \end{aligned} \quad (\text{B5})$$

We then also integrate over all possible values of \mathcal{Q}_c^J and \mathcal{Q}_s^J , subject to the constraint $\delta(\mathcal{Q} - \mathcal{Q}_c^J - \mathcal{Q}_s^J)$. This results in the general factorization formula

$$\begin{aligned} d\sigma_{\text{resum}}[\mathcal{Q}, \tau_1, P_{JT}, y_J] &= \sigma_0 H(x_* \sqrt{s} P_{JT} e^{-y_J}, \mu; \mu_H) \\ &\int d\mathcal{Q}_c^J \int d\mathcal{Q}_s^J \delta(\mathcal{Q} - \mathcal{Q}_c^J - \mathcal{Q}_s^J) \\ &\times \int ds_J \int dt_a \mathcal{S}\left(\tau_1 - \frac{t_a}{Q_a} - \frac{s_J}{Q_J}, \mathcal{Q}_s^J, \mu; \mu_S\right) \\ &\times \left[\sum_{q_i} L_{q_i} \mathcal{G}^{q_i}(\mathcal{Q}_c^J, s_J, \mu; \mu_J) B_{q_i}(x_*, t_a, \mu; \mu_B) \right. \\ &\left. + \sum_{\bar{q}_i} L_{\bar{q}_i} \mathcal{G}^{\bar{q}_i}(\mathcal{Q}_c^J, s_J, \mu; \mu_J) B_{\bar{q}_i}(x_*, t_a, \mu; \mu_B) \right]. \end{aligned} \quad (\text{B6})$$

Next, in the above factorization formula, we can expand

$\delta(\mathcal{Q} - \mathcal{Q}_c^J - \mathcal{Q}_s^J)$ in the limit $Q_s^J \ll Q_c^J$ as

$$\begin{aligned} &\delta(\mathcal{Q} - \mathcal{Q}_c^J - \mathcal{Q}_s^J) \\ &= \delta(\mathcal{Q} - \mathcal{Q}_c^J) + \delta'(\mathcal{Q} - \mathcal{Q}_c^J) \mathcal{Q}_s^J + \dots \end{aligned} \quad (\text{B7})$$

where the ellipses denote terms suppressed by higher powers in \mathcal{Q}_s^J . The first term in Eq. (B7), $\delta(\mathcal{Q} - \mathcal{Q}_c^J)$, along with the normalization condition on the charged soft function

$$\begin{aligned} &\int d\mathcal{Q}_s^J \mathcal{S}\left(\tau_1 - \frac{t_a}{Q_a} - \frac{s_J}{Q_J}, \mathcal{Q}_s^J, \mu; \mu_S\right) \\ &= \mathcal{S}\left(\tau_1 - \frac{t_a}{Q_a} - \frac{s_J}{Q_J}, \mu; \mu_S\right), \end{aligned} \quad (\text{B8})$$

reproduces the leading power factorization formula in Eq. (17). The first power correction comes from the second term in Eq. (B7) and is given by

$$\begin{aligned} d\sigma_{\text{resum}}^{\text{PC}}[\mathcal{Q}, \tau_1, P_{JT}, y_J] &= \sigma_0 H(x_* \sqrt{s} P_{JT} e^{-y_J}, \mu; \mu_H) \\ &\times \int ds_J \int dt_a \int d\mathcal{Q}_s^J \mathcal{Q}_s^J \mathcal{S}\left(\tau_1 - \frac{t_a}{Q_a} - \frac{s_J}{Q_J}, \mathcal{Q}_s^J, \mu; \mu_S\right) \\ &\times \left[\sum_{q_i} L_{q_i} \int d\mathcal{Q}_c^J \delta'(\mathcal{Q} - \mathcal{Q}_c^J) \mathcal{G}^{q_i}(\mathcal{Q}_c^J, s_J, \mu; \mu_J) \right. \\ &\quad \times B_{q_i}(x_*, t_a, \mu; \mu_B) \\ &\quad \left. + \sum_{\bar{q}_i} L_{\bar{q}_i} \int d\mathcal{Q}_c^J \delta'(\mathcal{Q} - \mathcal{Q}_c^J) \mathcal{G}^{\bar{q}_i}(\mathcal{Q}_c^J, s_J, \mu; \mu_J) \right. \\ &\quad \left. \times B_{\bar{q}_i}(x_*, t_a, \mu; \mu_B) \right]. \end{aligned} \quad (\text{B9})$$

In this manner, power corrections from the contribution of soft particles in the jet region to the jet charge can be systematically incorporated. We note however that this power correction will also require incorporating a non-perturbative model for the new charged soft function. We also note that this power correction only involves the first jet charge moment of the new charged soft function, as seen in the second line of Eq. (B9).

-
- [1] E. Farhi, Phys. Rev. Lett. **39**, 1587 (1977).
 - [2] M. Dasgupta and G. P. Salam, JHEP **08**, 032 (2002), hep-ph/0208073.
 - [3] L. Clavelli, Phys. Lett. B **85**, 111 (1979).
 - [4] J. D. Bjorken and S. J. Brodsky, Phys. Rev. D **1**, 1416 (1970).
 - [5] G. Parisi, Phys. Lett. B **74**, 65 (1978).
 - [6] H. Georgi and M. Machacek, Phys. Rev. Lett. **39**, 1237 (1977).
 - [7] I. W. Stewart, F. J. Tackmann, and W. J. Waalewijn, Phys. Rev. Lett. **105**, 092002 (2010), 1004.2489.
 - [8] Z.-B. Kang, S. Mantry, and J.-W. Qiu, Phys. Rev. D **86**, 114011 (2012), 1204.5469.
 - [9] Z.-B. Kang, X. Liu, S. Mantry, and J.-W. Qiu, Phys. Rev. D **88**, 074020 (2013), 1303.3063.
 - [10] D. Kang, C. Lee, and I. W. Stewart, Phys. Rev. D **88**, 054004 (2013), 1303.6952.
 - [11] Z.-B. Kang, X. Liu, and S. Mantry, Phys. Rev. D **90**, 014041 (2014), 1312.0301.
 - [12] D. Kang, C. Lee, and I. W. Stewart, JHEP **11**, 132 (2014), 1407.6706.
 - [13] Z. Chu, Y. Wang, J.-H. Ee, J. Chen, and D. Kang, JHEP **06**, 111 (2022), 2202.08040.
 - [14] H. Cao, Z.-B. Kang, X. Liu, and S. Mantry, Phys. Rev. D **110**, 014045 (2024), 2401.01941.
 - [15] J.-H. Ee, D. Kang, C. Lee, and I. W. Stewart, JHEP **07**, 240 (2025), 2504.05234.
 - [16] T. Becher and M. D. Schwartz, JHEP **07**, 034 (2008), 0803.0342.
 - [17] R. Abbate, M. Fickinger, A. H. Hoang, V. Mateu, and I. W. Stewart, Phys. Rev. D **83** (2011).
 - [18] R. Abbate, M. Fickinger, A. H. Hoang, V. Mateu,

- and I. W. Stewart, Phys. Rev. D **86**, 094002 (2012), 1204.5746.
- [19] S. Fleming, A. H. Hoang, S. Mantry, and I. W. Stewart, Phys. Rev. D **77**, 074010 (2008), hep-ph/0703207.
- [20] S. Fleming, A. H. Hoang, S. Mantry, and I. Stewart, Phys. Rev. D (2008).
- [21] A. H. Hoang, S. Mantry, A. Pathak, and I. W. Stewart, Phys. Rev. D **100**, 074021 (2019), 1708.02586.
- [22] Y.-T. Chien and M. D. Schwartz, JHEP **08**, 058 (2010), 1005.1644.
- [23] A. H. Hoang, V. Mateu, M. D. Schwartz, and I. W. Stewart, JHEP **09**, 092 (2025), 2506.09130.
- [24] E.-C. Aschenauer *et al.*, (2016), 1602.03922.
- [25] STAR, L. Adamczyk *et al.*, Phys. Rev. D **97**, 032004 (2018), 1708.07080.
- [26] D. Boer and C. Pisano, Phys. Rev. D **91**, 074024 (2015), 1412.5556.
- [27] X. Liu, F. Ringer, W. Vogelsang, and F. Yuan, Phys. Rev. Lett. **122**, 192003 (2019), 1812.08077.
- [28] M. Arratia, Z.-B. Kang, A. Prokudin, and F. Ringer, Phys. Rev. D **102**, 074015 (2020), 2007.07281.
- [29] X. Liu, F. Ringer, W. Vogelsang, and F. Yuan, (2020), 2007.12866.
- [30] E.-C. Aschenauer, K. Lee, B. Page, and F. Ringer, Phys. Rev. D **101**, 054028 (2020), 1910.11460.
- [31] M. Arratia, Y. Song, F. Ringer, and B. Jacak, Phys. Rev. C **101**, 065204 (2020), 1912.05931.
- [32] M. Arratia, Y. Makris, D. Neill, F. Ringer, and N. Sato, (2020), 2006.10751.
- [33] Z.-B. Kang, K. Lee, and F. Zhao, Phys. Lett. B **809**, 135756 (2020), 2005.02398.
- [34] E. Aschenauer *et al.*, Rept. Prog. Phys. **82**, 024301 (2019), 1708.01527.
- [35] L. Zheng, E. Aschenauer, J. Lee, B.-W. Xiao, and Z.-B. Yin, Phys. Rev. D **98**, 034011 (2018), 1805.05290.
- [36] R. Field and R. Feynman, Nucl. Phys. B **136**, 1 (1978).
- [37] D. Krohn, M. D. Schwartz, T. Lin, and W. J. Waalewijn, Phys. Rev. Lett. **110**, 212001 (2013), 1209.2421.
- [38] W. J. Waalewijn, Phys. Rev. D **86**, 094030 (2012), 1209.3019.
- [39] Fermilab-Serpukhov-Moscow-Michigan, J. Berge *et al.*, Phys. Lett. B **91**, 311 (1980).
- [40] J. Berge *et al.*, Nucl. Phys. B **184**, 13 (1981).
- [41] Aachen-Bonn-CERN-Munich-Oxford, P. Allen *et al.*, p. 307 (1982).
- [42] European Muon, J. Albanese *et al.*, Phys. Lett. B **144**, 302 (1984).
- [43] Amsterdam-Bologna-Padua-Pisa-Saclay-Turin, S. Barlag *et al.*, Z. Phys. C **11**, 283 (1982), [Erratum: Z.Phys.C 14, 281 (1982)].
- [44] R. Erickson *et al.*, Phys. Rev. Lett. **42**, 822 (1979), [Erratum: Phys.Rev.Lett. 42, 1246 (1979)].
- [45] AMY, D. Stuart *et al.*, Phys. Rev. Lett. **64**, 983 (1990).
- [46] ALEPH, D. Decamp *et al.*, Phys. Lett. B **259**, 377 (1991).
- [47] TASSO, W. Braunschweig *et al.*, Z. Phys. C **48**, 433 (1990).
- [48] D0, V. Abazov *et al.*, Phys. Rev. Lett. **98**, 041801 (2007), hep-ex/0608044.
- [49] CDF, T. Aaltonen *et al.*, Phys. Rev. D **88**, 032003 (2013), 1304.4141.
- [50] ATLAS, G. Aad *et al.*, JHEP **11**, 031 (2013), 1307.4568.
- [51] CMS, V. Khachatryan *et al.*, JHEP **12**, 017 (2014), 1410.4227.
- [52] Y.-C. J. Chen, C.-W. Chiang, G. Cottin, and D. Shih, Phys. Rev. D **101**, 053001 (2020), 1908.08256.
- [53] ATLAS, G. Aad *et al.*, Phys. Rev. D **93**, 052003 (2016), 1509.05190.
- [54] CMS, A. M. Sirunyan *et al.*, JHEP **10**, 131 (2017), 1706.05868.
- [55] K. Fraser and M. D. Schwartz, JHEP **10**, 093 (2018), 1803.08066.
- [56] Z.-B. Kang, A. J. Larkoski, and J. Yang, Phys. Rev. Lett. **130**, 151901 (2023), 2301.09649.
- [57] M. Gyulassy and X.-n. Wang, Nucl. Phys. B **420**, 583 (1994), nucl-th/9306003.
- [58] X.-N. Wang, M. Gyulassy, and M. Plumer, Phys. Rev. D **51**, 3436 (1995), hep-ph/9408344.
- [59] Y.-T. Chien and I. Vitev, Phys. Rev. Lett. **119**, 112301 (2017), 1608.07283.
- [60] M. Connors, C. Nattrass, R. Reed, and S. Salur, Rev. Mod. Phys. **90**, 025005 (2018), 1705.01974.
- [61] H. T. Li and I. Vitev, Phys. Rev. D **101**, 076020 (2020), 1908.06979.
- [62] CMS, A. M. Sirunyan *et al.*, JHEP **07**, 115 (2020), 2004.00602.
- [63] Y.-L. Du, D. Pablos, and K. Tywoniuk, (2020), 2012.07797.
- [64] Z.-B. Kang, X. Liu, S. Mantry, and D. Y. Shao, Phys. Rev. Lett. **125**, 242003 (2020), 2008.00655.
- [65] Z.-B. Kang, K. Lee, D. Y. Shao, and J. Terry, (2020), 2008.05470.
- [66] J. Gallicchio and M. D. Schwartz, Phys. Rev. Lett. **107**, 172001 (2011), 1106.3076.
- [67] A. J. Larkoski, I. Moult, and B. Nachman, Phys. Rept. **841**, 1 (2020), 1709.04464.
- [68] J. Brewer, J. Thaler, and A. P. Turner, (2020), 2008.08596.
- [69] Y.-T. Chien and R. Kunnawalkam Elayavalli, (2018), 1803.03589.
- [70] A. Buckley, G. Callea, A. J. Larkoski, and S. Marzani, (2020), 2006.10480.
- [71] Z.-B. Kang, X. Liu, S. Mantry, M. Spraker, and T. Wilson, Phys. Rev. D **103**, 074028 (2021).
- [72] CERN Report No. ATLAS-CONF-2013086, (unpublished) (2013).
- [73] Y.-T. Chien, A. Deshpande, M. M. Mondal, and G. Sterman, Phys. Rev. D **105**, L051502 (2022), 2109.15318.
- [74] I. W. Stewart, F. J. Tackmann, and W. J. Waalewijn, Phys.Rev. **D81**, 094035 (2010), 0910.0467.
- [75] C. W. Bauer, S. Fleming, and M. E. Luke, Phys.Rev. **D63**, 014006 (2000), hep-ph/0005275.
- [76] C. W. Bauer, S. Fleming, D. Pirjol, and I. W. Stewart, Phys.Rev. **D63**, 114020 (2001), hep-ph/0011336.
- [77] C. W. Bauer and I. W. Stewart, Phys.Lett. **B516**, 134 (2001), hep-ph/0107001.
- [78] C. W. Bauer, D. Pirjol, and I. W. Stewart, Phys.Rev. **D65**, 054022 (2002), hep-ph/0109045.
- [79] C. W. Bauer, S. Fleming, D. Pirjol, I. Z. Rothstein, and I. W. Stewart, Phys.Rev. **D66**, 014017 (2002), hep-ph/0202088.
- [80] M. Beneke, A. Chapovsky, M. Diehl, and T. Feldmann, Nucl.Phys. **B643**, 431 (2002), hep-ph/0206152.
- [81] A. Idilbi, X. Ji, and F. Yuan, Nuclear Physics B **753**, 42 (2006).
- [82] T. Becher, M. Neubert, and B. D. Pecjak, JHEP **01**, 076 (2007), hep-ph/0607228.
- [83] T. T. Jouttenus, I. W. Stewart, F. J. Tackmann, and

- W. J. Waalewijn, Phys.Rev. **D83**, 114030 (2011), 1102.4344.
- [84] R. Boughezal, X. Liu, and F. Petriello, Phys. Rev. D **91**, 094035 (2015), 1504.02540.
- [85] T. Becher and M. Neubert, Phys. Lett. B **637**, 251 (2006).
- [86] I. W. Stewart, F. J. Tackmann, and W. J. Waalewijn, JHEP **1009**, 005 (2010), 1002.2213.
- [87] J. Gaunt, M. Stahlhofen, and F. Tackmann, JHEP **04**, 113 (2014).
- [88] R. Boughezal, F. Petriello, U. Schubert, and H. Xing, Phys. Rev. D **96**, 034001 (2017), 1704.05457.