

# Spin Polarization from Circularly Polarized Light Induced Charge Transfer

Sindhana Pannir-Sivajothi<sup>1</sup> and David T. Limmer<sup>1, 2, 3, 4, a)</sup>

<sup>1)</sup>Department of Chemistry, University of California, Berkeley, California 94720, United States

<sup>2)</sup>Kavli Energy Nanoscience Institute at Berkeley, Berkeley, California 94720, United States

<sup>3)</sup>Chemical Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720, United States

<sup>4)</sup>Material Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720, United States

We show how a spin polarization can be generated through the photo-induced electron transfer of an achiral donor-acceptor complex following chiral light excitation. In particular, we illustrate the basic energetic and symmetry requirements for chirality induced spin selectivity where the chirality emerges from the electronic degrees of freedom following excitation with circularly polarized light. We study this effect in a simple model of a metalloporphyrin complex with an axial acceptor ligand using quantum mechanical rate theories and numerical simulations. We find that the spin polarization emerges due to the selective excitation of a ring current within the porphyrin, breaking the degeneracy of the two degenerate spin states. The resultant spin polarization increases with the spin orbit coupling between the metal in the porphyrin and the axial ligand, and is transient, with a lifetime dependent on the rate of dephasing from the Jahn-Teller distortion mode. This proposed effect should be observable in spin-resolved photoemission spectroscopy.

The chirality-induced spin selectivity (CISS) effect, where electron transmission through chiral molecules is spin-dependent, has been observed in both photoelectron transport through ground-state chiral molecules<sup>1,2</sup> and photoinduced electron transfer.<sup>3</sup> CISS is typically attributed to chirality defined by nuclear geometry, however achiral donor-acceptor systems excited with circularly polarized light (CPL) can admit transient chirality when both nuclear and electronic degrees of freedom are considered.<sup>4–6</sup> This suggests that a transient spin polarization may be generated following CPL in achiral complexes. In this Letter, we illustrate that a spin polarization in CPL-driven electron transfer in achiral donor-acceptor complexes is indeed possible transiently and discuss the underlying mechanism. This presents a novel means of controlling electron spin in molecular systems with potential application for molecular qubits.<sup>7–9</sup>

When a molecule absorbs CPL, the photon's angular momentum is transferred to it, resulting in excited states with nonzero projection of electronic orbital angular momentum along the light axis. For concreteness, we consider an achiral donor molecule with CPL-addressable degenerate excited states  $|^1D_\lambda\rangle$  where the label  $\lambda = \pm$  depends on its orbital angular momentum, as sketched in Fig. 1. Examples of such donors include  $\pi$ -conjugated molecules like metalloporphyrins<sup>10</sup> and metallophthalocyanines,<sup>11,12</sup> as well as monolayer transition metal dichalcogenides.<sup>13</sup> We define a molecular frame ( $\mathbf{x}_{\text{mol}}, \mathbf{y}_{\text{mol}}, \mathbf{z}_{\text{mol}}$ ), with  $\mathbf{z}_{\text{mol}}$  pointing from donor to acceptor and the molecule is oriented such that the light axis aligns with  $\mathbf{z}_{\text{mol}}$  as in Fig. 1. To understand the subsequent charge-transfer dynamics, we construct a minimal electronic Hamiltonian for this donor-acceptor system using only its frontier orbitals. Within this framework, two charge-transfer pathways emerge: a spin-conserving channel ( $|^1D_\lambda\rangle \rightarrow |^1CT_\lambda\rangle$ ), and a spin-orbit-coupled channel ( $|^1D_\lambda\rangle \rightarrow |^3CT_\lambda\rangle$ ) that depends




FIG. 1. Electron transfer in an achiral donor-acceptor complex driven by circularly polarized light. Left- and right-circularly polarized light excite the donor singlet states  $|^1D_\lambda\rangle \equiv |^1D_\lambda^* - A\rangle$  with  $\lambda = \pm$  from the ground state  $|\Psi_0\rangle \equiv |^1D - A\rangle$ . Singlet and triplet charge transfer states  $|^1CT_\lambda\rangle \equiv |^1D_\lambda^{*+} - A^{*-}\rangle$  and  $|^3CT_\lambda\rangle \equiv |^3D_\lambda^{*+} - A^{*-}\rangle$  are nearly degenerate and detuned from  $|^1D_\lambda\rangle$  by  $\Delta$ , while  $v_0$  and  $v_{\text{SOC}_\lambda}$  mediate spin-conserving and spin-orbit coupled electron transfer.

on the helicity of the exciting CPL. The model predicts a net spin polarization that reverses under  $\lambda \rightarrow -\lambda$  and grows with the spin-orbit coupling (SOC) strength  $v_{\text{SOC}_\lambda}$ . This spin polarization decays due to vibronic coupling in this case originating in Jahn-Teller distortions that lift the  $\lambda = \pm$  degeneracy,<sup>14,15</sup> which we model using Bloch-Redfield equations. The transient generation of a spin polarization therefore, occurs when the charge-transfer rate exceeds the rate of  $\lambda = \pm$  orbital angular momentum state mixing. Since such ultrafast coherent charge transfer has been observed in dyads with a metalloporphyrin donor and an axially bound acceptor,<sup>16–18</sup> we take this as a representative system for our study but expect the basic mechanism to hold generally.

To build a minimal Hamiltonian, we consider complex-valued frontier orbitals for the donor porphyrin, with creation operators  $\hat{d}_{H\pm\sigma}^\dagger$  for the HOMOs and  $\hat{d}_{L\pm\sigma}^\dagger$  for the

<sup>a)</sup>Electronic mail: [dlimmer@berkeley.edu](mailto:dlimmer@berkeley.edu)

LUMOs. For the acceptor, we have real-valued LUMOs with creation operators  $\hat{a}_{L\sigma}^\dagger$ , where  $\sigma \in \{\alpha, \beta\}$  denotes spin. The complex donor orbitals are constructed from Gouterman's four-orbital model, the canonical model for porphyrin optical spectra.<sup>19</sup> The Gouterman model includes two exactly degenerate LUMOs and two accidentally degenerate HOMOs all of which are real-valued and have transition dipole moments along  $\mathbf{x}_{\text{mol}}$  and  $\mathbf{y}_{\text{mol}}$ . We construct complex frontier orbitals,  $\hat{d}_{H\pm\sigma}^\dagger$  and  $\hat{d}_{L\pm\sigma}^\dagger$ , as linear combinations of the degenerate Gouterman orbitals and singly excited states built from these yield transition dipoles  $\mathbf{x}_{\text{mol}} \pm i\mathbf{y}_{\text{mol}}$ , which are selectively addressable with CPL (see Supplementary S1). When donor-acceptor dyads have orthogonal  $\pi$ -systems, they can exhibit substantial SOC-assisted charge transfer,<sup>20,21</sup> therefore, we choose to study a  $\pi$ -conjugated dyad with perpendicular donor and acceptor  $\pi$ -planes. Since the acceptor is bound to the donor axially via a metal center, the metal forms part of both donor and acceptor  $\pi$ -systems and charge transfer occurs via it.

Using the frontier orbitals and the electronic ground state  $|\Psi_0\rangle$  of the donor-acceptor dyad, we define the donor singlet excited states as

$$|^1D_\lambda\rangle = \frac{1}{\sqrt{2}}(\hat{d}_{L\alpha}^\dagger \hat{d}_{H\lambda\alpha} + \hat{d}_{L\beta}^\dagger \hat{d}_{H\lambda\beta})|\Psi_0\rangle, \quad (1)$$

where  $\lambda = \pm$  corresponds to states excited with left- and right-CPL (see Supplementary S1). These will be our initial states. The donor excited states,  $|^1D_+\rangle$  and  $|^1D_-\rangle$ , constitute an enantiomeric pair linked by spatial inversion. Although the donor lacks continuous rotational symmetry about  $\mathbf{z}_{\text{mol}}$ , its discrete symmetry (e.g.,  $D_{4h}$  in metalloporphyrins) supports quasi-angular-momentum states that are approximate eigenstates of the  $z$ -component of orbital angular momentum  $\hat{L}_{z\text{mol}} = \hat{\mathbf{L}} \cdot \mathbf{z}_{\text{mol}}$  with  $\hat{L}_{z\text{mol}}|^1D_\lambda\rangle \approx \lambda m\hbar|^1D_\lambda\rangle$ . These quasi-angular momentum states are selectively excitable with CPL and carry opposite ring currents for  $\lambda = \pm$ .<sup>10,22</sup> Under spatial inversion, the orbital angular momentum transforms as  $\mathcal{P}\hat{L}_{z\text{mol}}\mathcal{P}^{-1} = -\hat{L}_{z\text{mol}}$  because it is the scalar product of a polar vector  $\mathbf{z}_{\text{mol}}$ , that under inversion transforms as  $\mathbf{z}_{\text{mol}} \rightarrow -\mathbf{z}_{\text{mol}}$ , and an axial vector operator  $\hat{\mathbf{L}}$ , with  $\mathcal{P}\hat{\mathbf{L}}\mathcal{P}^{-1} = \hat{\mathbf{L}}$ . Consequently, the excited states are related to each other through spatial inversion  $\mathcal{P}|^1D_+\rangle \propto |^1D_-\rangle$ , similar to enantiomers of chiral molecules. We will refer to these chiral states as *transient enantiomers*. In a randomly oriented ensemble of achiral donor-acceptor complexes, circularly polarized light generates equal populations of the transient enantiomers, yielding zero net spin polarization. Thus, a finite net spin polarization requires oriented complexes as in Fig. 1.

The singlet and  $M_S = 0$  triplet charge-transfer (CT) states are

$$|^1CT_\lambda\rangle = \frac{1}{\sqrt{2}}(\hat{a}_{L\alpha}^\dagger \hat{d}_{H\lambda\alpha} + \hat{a}_{L\beta}^\dagger \hat{d}_{H\lambda\beta})|\Psi_0\rangle, \quad (2a)$$

$$|^3CT_\lambda\rangle = \frac{1}{\sqrt{2}}(\hat{a}_{L\alpha}^\dagger \hat{d}_{H\lambda\alpha} - \hat{a}_{L\beta}^\dagger \hat{d}_{H\lambda\beta})|\Psi_0\rangle. \quad (2b)$$

The CT state retains the label  $\lambda = \pm$  because the donor hole carries a ring current, resulting in finite expectation value of  $\hat{L}_{z\text{mol}}$ . The singlet CT state is detuned by  $\Delta$  from the singlet donor excited state. The singlet charge transfer state  $|^1CT_\lambda\rangle$  is coupled to the donor singlet state  $|^1D_\lambda\rangle$  through a diabatic coupling with magnitude  $v_0$  mediate by the metal center. Due to the orientation of the  $\pi$ -planes, for the donor we include only the  $p_z$  atomic orbitals of its atoms and for the acceptor, only the  $p_x$  orbitals. We include the metal's  $d_{xz}$  and  $d_{yz}$  orbitals, where  $d_{xz}$  contributes to both  $\pi$ -systems and thus can mediate charge transfer directly.

To account for the SOC-assisted charge transfer, we include the mean-field Breit-Pauli form of the spin-orbit coupling.<sup>23</sup> Within our minimal model basis, only the metal center contributes to the one-center SOC.<sup>24,25</sup> The one-center SOC terms are defined as matrix elements where the angular momentum operator and both atomic orbitals are centered around the same nucleus. Because the angular momentum operators are pure imaginary operators and have zero diagonal matrix elements in a real-valued atomic orbital basis and since we retain only one atomic orbital per atom except for the metal, it is evident that only the metal center will contribute to the one-center SOC. Therefore, the diabatic states are additionally coupled through a single-electron SOC term of the Hamiltonian with magnitude  $v_{\text{SOC}\lambda}$ , as derived in Supplementary S2 using the mean-field SOC and Hückel theory.

The resultant electronic Hamiltonian,  $\hat{H}_e$ , spanned by  $|^1D_\lambda\rangle$ ,  $|^1CT_\lambda\rangle$ , and  $|^3CT_\lambda\rangle$  is thus

$$\begin{aligned} \hat{H}_e = \sum_{\lambda=\pm,-} \Delta (|^1CT_\lambda\rangle \langle^1CT_\lambda| + |^3CT_\lambda\rangle \langle^3CT_\lambda|) \\ + v_0 (|^1D_\lambda\rangle \langle^1CT_\lambda| + |^1CT_\lambda\rangle \langle^1D_\lambda|) \\ + v_{\text{SOC}\lambda} (|^1D_\lambda\rangle \langle^3CT_\lambda| + |^3CT_\lambda\rangle \langle^1D_\lambda|) \end{aligned} \quad (3)$$

where all energy differences are with respect to the donor singlet. Note  $v_{\text{SOC}\lambda}$  depends on helicity with  $v_{\text{SOC}-} = -v_{\text{SOC}+}$ . One can interpret this difference in sign of the SOC matrix element as due to an effective transient magnetic field produced by the ring current upon excitation with CPL. The spin-conserving electron transfer occurs solely via the  $d_{xz}$  orbital while the SOC charge transfer involves both the  $d_{xz}$  and  $d_{yz}$  orbitals of the metal center. Unlike previous work where the spin-preserving electron transfer matrix element is real and the SOC matrix element is purely imaginary,<sup>26</sup> in our case both the direct and SOC-assisted electron transfer matrix elements are real-valued as we have complex frontier orbitals with finite angular momentum.

The triplet donor states with spin projection  $M_S = 0$  along the quantization axis  $\mathbf{z}_{\text{mol}}$  are

$$|^3D_\lambda\rangle = \frac{1}{\sqrt{2}}(\hat{d}_{L\alpha}^\dagger \hat{d}_{H\lambda\alpha} - \hat{d}_{L\beta}^\dagger \hat{d}_{H\lambda\beta})|\Psi_0\rangle. \quad (4)$$

In a mean-field Hamiltonian, the singlet and triplet states are degenerate. However, the full Hamiltonian would in-

clude additional exchange interaction terms of the form  $J \sum_{\sigma=\alpha,\beta} \sum_{\tau=\alpha,\beta} \hat{d}_{L\lambda\sigma}^\dagger \hat{d}_{H\lambda\tau}^\dagger \hat{d}_{L\lambda\tau} \hat{d}_{H\lambda\sigma}$  that lower the energy of the triplet with respect to the singlet. For donor excited states, the singlet-triplet energy gap  $2J \sim 400$  meV for metalloporphyrins.<sup>27</sup> In the CT configuration, the electron and hole are spatially separated, so the exchange interaction is small and the singlet and triplet CT states are nearly degenerate.<sup>28</sup> For our calculations we use  $\hat{H}_e$  as defined in Eq. 3, where we take the energy difference between the singlet and triplet CT states to be zero as it will be much smaller than other energy scales in the problem, and we exclude the state  $|^3D_\lambda\rangle$  from our analysis as it will be far-detuned from  $|^1D_\lambda\rangle$ ,  $|^1CT_\lambda\rangle$ , and  $|^3CT_\lambda\rangle$ . The electronic Hamiltonian with  $|^3D_\lambda\rangle$  included is in Supplementary S2.

While the donor singlets and  $M_S = 0$  triplets  $|^{1(3)}D_\lambda\rangle$  are eigenstates of both  $\hat{S}_{Dz_{mol}}$  and  $\hat{S}_{Az_{mol}}$  with eigenvalue 0, the charge transfer states are not. The total  $\mathbf{z}_{mol}$  projected spin  $\hat{S}_{z_{mol}} = \hat{S}_{Dz_{mol}} + \hat{S}_{Az_{mol}}$ , where the spin of electrons localized on the donor is

$$\hat{S}_{Dz_{mol}} = \hbar/2 \sum_{\lambda=+,-} (\hat{d}_{H\lambda\alpha}^\dagger \hat{d}_{H\lambda\alpha} - \hat{d}_{H\lambda\beta}^\dagger \hat{d}_{H\lambda\beta} + \hat{d}_{L\lambda\alpha}^\dagger \hat{d}_{L\lambda\alpha} - \hat{d}_{L\lambda\beta}^\dagger \hat{d}_{L\lambda\beta}) \quad (5)$$

and of those in the acceptor is

$$\hat{S}_{Az_{mol}} = \hbar/2 (\hat{a}_{L\alpha}^\dagger \hat{a}_{L\alpha} - \hat{a}_{L\beta}^\dagger \hat{a}_{L\beta}). \quad (6)$$

The spin polarization is defined as  $\hat{P}_z = \hat{S}_{Az_{mol}} - \hat{S}_{Dz_{mol}}$  and will be the primary observable of our interest. In the representation of our electronic states, the spin polarization manifests as the real part of the coherence between the charge transfer singlets and triplets,

$$\hat{P}_z = \sum_{\lambda=+,-} (\hbar |^3CT_\lambda\rangle \langle ^1CT_\lambda| + \text{h.c.}) \quad (7)$$

as has been noted in previous work.<sup>26</sup>

We want to model ultrafast coherent electron transfer and will not explicitly include a vibrational mode that is displaced between the donor excited state and the charge transfer state, so our analysis works best for rigid donor-acceptor complexes with minimal outer and inner-sphere reorganization, small detunings  $\Delta$ , and large electronic couplings that facilitate this ultrafast charge transfer process.<sup>29–31</sup> For our calculations, we use  $v_0 = 75$  meV which is reasonable for systems that satisfy the above criteria,<sup>29</sup> and we use  $v_{SOC\pm} = \pm 25$  meV based on SOC matrix elements in donor-acceptor complexes with heavy transition metal atoms.<sup>32,33</sup>

Given an initial state  $|^1D_+\rangle$  the subsequent unitary dynamics for the population and spin polarization can be deduced exactly. These are shown in Fig. 2(a-b). The population of the donor,  $p_{|^1D_+\rangle}(t) = |\langle ^1D_+(t) | ^1D_+(0) \rangle|^2$ , singlet charge transfer state,  $p_{|^1CT_+\rangle}(t) = |\langle ^1CT_+(t) | ^1D_+(0) \rangle|^2$  and triple charge




FIG. 2. Unitary dynamics. (a) The populations in the singlet donor  $p_{|^1D_+\rangle}$  (black), singlet charge transfer  $p_{|^1CT_+\rangle}$  (yellow) and the triplet charge transfer  $p_{|^3CT_+\rangle}$  (red) states and (b) the spin polarization  $\langle \hat{P}_z(t) \rangle$  following excitation into  $|^1D_+\rangle$ . (c) Mean spin polarization  $\bar{P}_z$  for different detunings  $\Delta$  and its dependence on initially excited state. Black star indicates condition in (a) and (b).

transfer state,  $p_{|^3CT_+\rangle}(t) = |\langle ^3CT_+(t) | ^1D_+(0) \rangle|^2$  each beat with a Rabi frequency,  $\Omega = \sqrt{(\Delta/2)^2 + v_0^2 + v_{SOC+}^2}$ . The donor state is exactly out of phase with the two charge transfer states, whose relative amplitudes depend on the ratio of  $v_0^2/v_{SOC+}^2$ . The spin polarization has a time dependence that is given by

$$\langle \hat{P}_z(t) \rangle = \frac{\hbar v_0 v_{SOC+}}{\Omega^2} [1 - \cos(2\Omega t/\hbar)], \quad (8)$$

which we plot in Fig. 2(b). The spin polarization beats with the same Rabi frequency as the populations, and with an amplitude proportional to the size of the spin-orbit coupling. Therefore, the mean spin polarization  $\bar{P}_z = \lim_{\tau \rightarrow \infty} \frac{1}{\tau} \int_0^\tau dt \langle \hat{P}_z(t) \rangle$  is non-zero and takes on opposite sign depending on whether the initial state is  $|^1D_+\rangle$  or  $|^1D_-\rangle$

$$\bar{P}_z = \hbar \frac{v_0 v_{SOC\lambda}}{\Omega^2}, \quad (9)$$

that is, the mean spin-polarization flips sign when excited with right- versus left-circularly polarized light as  $v_{\text{SOC}+} = -v_{\text{SOC}-}$ . We plot  $\bar{P}_z$  in Fig. 2(c), when the initial state is  $|^1D_+\rangle$  and  $|^1D_-\rangle$  as a function of  $\Delta$ , the energy difference between the donor and charge transfer states. We find  $\sim 0.3\hbar$  mean spin polarization at  $\Delta = 0$ .

Ideal systems to test our predictions in Fig. 2(c) are donor-acceptor dyads with identical donor and acceptor molecules ( $\Delta = 0$ ). By introducing substituents on the acceptor, its LUMO—and hence  $\Delta$ —can be tuned.<sup>34,35</sup> Across such a series, a CPL pump followed by spin-resolved photoemission spectroscopy would provide an experimental plot of mean spin-polarization like Fig. 2(c). However, in reality the initial excitation will dephase through vibronic coupling. Specifically, it is known that in metalloporphyrins and related systems, the degenerate  $|^1D_+\rangle$  and  $|^1D_-\rangle$  are coupled through Jahn-Teller distortions with the typical depolarization rate  $\sim 1.6$  ps.<sup>36</sup> Since the spin polarization depends on the nature of the excited donor state, this dephasing should cause the mean spin polarization to decay.

We quantify these spin relaxation rates using Bloch-Redfield equations, assuming that the electron-phonon coupling is small compared to  $v_0$  and  $v_{\text{SOC}\lambda}$ .<sup>37–39</sup> Typically, for degenerate electronic states  $|^1D_{\pm}\rangle$  which are quasi-eigenstates of the orbital angular momentum operator, the  $(E \times e)$  Jahn-Teller Hamiltonian is used where a pair of degenerate vibrational modes couple them.<sup>40</sup> However, for molecules like metalloporphyrin that have an approximate four-fold symmetry, the Jahn-Teller active vibrational modes are non-degenerate.<sup>14,15</sup> We model this by coupling the degenerate  $\lambda = \pm$  orbital angular momentum states to a bath through a bilinear system-bath coupling of the form  $\hat{V}_{e-b} = \hat{V}_e \otimes \hat{B}$  where

$$\hat{V}_e = f_D |^1D_+\rangle \langle ^1D_-| + f_{\text{CT}} |^1\text{CT}_+\rangle \langle ^1\text{CT}_-| + f_{\text{CT}} |^3\text{CT}_+\rangle \langle ^3\text{CT}_-| + \text{h.c.}, \quad (10)$$

with dimensionless coupling constants  $f_D$  and  $f_{\text{CT}}$ . The bath operator  $\hat{B} = \sum_k (g_k \hat{b}_k^\dagger + g_k^* \hat{b}_k)$  where  $\hat{b}_k$  is the annihilation operator of the bath mode  $k$  and the bath has a Drude-Lorentz spectral density

$$J(\omega) = \sum_k \frac{|g_k|^2}{\hbar} \delta(\omega - \omega_k) = \frac{2\Lambda\omega\omega_c}{\omega^2 + \omega_c^2} \quad (11)$$

with reorganization energy  $\Lambda = 0.2$  meV so the coupling strength is much smaller than other energy scales in  $\hat{H}_e$  for a range of  $f_D, f_{\text{CT}} \in [-0.4, 0.4]$ . We consider a fast bath with cutoff frequency  $\hbar\omega_c = 30$  meV ( $1/\omega_c \approx 22$  fs), corresponding to a bath-correlation decay time in the typical condensed-phase range of  $\sim 20 - 150$  fs.<sup>41,42</sup>

Along with the spin polarization  $\bar{P}_z$ , we also study the relaxation rate of the difference in population between the two transient-enantiomers  $\Delta\hat{N} = \hat{N}_+ - \hat{N}_-$  where

$$\hat{N}_\lambda = |^1D_\lambda\rangle \langle ^1D_\lambda| + |^1\text{CT}_\lambda\rangle \langle ^1\text{CT}_\lambda| + |^3\text{CT}_\lambda\rangle \langle ^3\text{CT}_\lambda| \quad (12)$$




FIG. 3. Relaxational dynamics. (a) Population difference  $\langle \Delta\hat{N}(t) \rangle = \langle \hat{N}_+(t) \rangle - \langle \hat{N}_-(t) \rangle$  (solid light orange) and its exponential fit  $\exp(-\Gamma_N t)$  (dashed orange) and (b) the spin polarization  $\langle \hat{P}_z(t) \rangle$  (solid light blue) and the exponential fit to its envelope  $\exp(-\Gamma_P t)$  (dashed blue) for  $\Delta = -30$  meV and  $f_D = f_{\text{CT}} = 0.4$ . (c) The decay rates  $\Gamma_P$  (blue circles) and  $\Gamma_N$  (orange squares) as a function of dimensionless coupling strengths  $f_D$  and  $f_{\text{CT}}$ . The solid line is the expected rate from our analytic expression in Eq. 13.

is the population of the  $\lambda$  enantiomer. Within Redfield theory when the initial state is  $|^1D_+\rangle$ , the approximate equation of motion of the population difference  $\langle \Delta\hat{N}(t) \rangle$  has a simple form

$$\frac{d\langle \Delta\hat{N}(t) \rangle}{dt} \approx -\frac{2\pi k_B T}{\hbar} \frac{\Lambda}{\hbar\omega_c} F^2 \langle \Delta\hat{N}(t) \rangle, \quad (13)$$

where  $k_B T$  is Boltzmann's constant times temperature. We derive this approximate form for the relaxation rates in Supplementary S4. The decay rate is proportional to the reorganization energy and the temperature, and inversely proportional to the cutoff frequency. The rate also scales quadratically with an effective coupling



strength,  $F$ , where

$$F = f_D + f_{CT} \frac{v_0^2 - v_{SOC+}^2}{v_0^2 + v_{SOC+}^2} \quad (14)$$

is the effective system-bath coupling strength. We compute the relaxation rates  $\Gamma_N$  and  $\Gamma_P$  of the population difference and spin polarization, respectively, numerically by fitting an exponential to the time series of  $\langle \Delta \hat{N}(t) \rangle$  and to the envelope of  $\langle \hat{P}_z(t) \rangle$  propagated with a Redfield equation of motion, and shown in Figs 3(a-b). Over a range of couplings, we find that both decay with a similar rate that is approximately given by Eq. 13, as shown in Fig. 3(c). Through engineering the bath properties, the spin polarization could be preserved for 10s of ps.

We find spin-dependent electron transfer, when an achiral donor-acceptor dyad is excited with a circularly-polarized pump. Theoretical and experimental works in the 1960s-90s have established that photoelectrons ejected by circularly polarized light have spin angular momentum correlated with the light’s angular momentum due to spin-orbit coupling.<sup>43,44</sup> Contrary to such direct photoemission with circularly-polarized light, our initial excitation results in a singlet with the component of total spin angular momentum along the light’s propagation axis  $\hat{S}_{z_{mol}} = \hat{S}_{Dz_{mol}} + \hat{S}_{Az_{mol}}$  ( $\mathbf{z}_{mol}$  in Fig. 1) being zero and remaining zero during the dynamics as it is a conserved quantity. However, the difference between spin angular momentum of electrons localized on the donor and the acceptor  $\hat{P}_z$  becomes nonzero during electron transfer and its value depends on the helicity of the exciting light. The CPL excites a transiently chiral electronic state that results in spin-polarization of electrons during charge transfer. This suggests that our work is better understood as a transient-electronic CISS effect and is conceptually different from direct conversion of electronic orbital angular momentum to spin angular momentum via spin-orbit coupling during photoemission with a circularly-polarized pump.

In recent experimental work, Liu *et al.* have demonstrated that upon exciting a transition metal dichalcogenide with CPL followed by electron transfer to an organic molecule, the generated spin polarization decays more slowly.<sup>45</sup> However, in that work the spin of the electron in the conduction band is fixed at the time of excitation as up or down depending on the helicity of the light and this electron is then transferred due to spin-valley locking. Our proposal takes this a step further, as the CPL excitation in itself only generates a singlet and during the dynamics of electron transfer, the system develops spin polarization. This idea can be further developed to include other notions of transient nuclear chirality.<sup>46</sup> The CISS effect has been demonstrated in donor-bridge-acceptor system when the bridge is chiral,<sup>3</sup> and it is known that IR excitation of the bridge vibrational modes can modify electron transfer rates,<sup>47</sup> so in future work, it would be interesting to study spin-polarization during photoinduced electron transfer in a

donor-bridge-acceptor system when an IR pulse excites vibrational modes making the bridge transiently chiral.

**Acknowledgments.** We acknowledge funding from William M. Keck Foundation (8959) and LDRD program at LBNL under the U.S. Department of Energy Office of Science, Office of Basic Energy Sciences, under Contract No. DEAC02-05CH11231.

- <sup>1</sup>K. Ray, S. P. Ananthavel, D. H. Waldeck, and R. Naaman, “Asymmetric scattering of polarized electrons by organized organic films of chiral molecules,” *Science* **283**, 814–816 (1999).
- <sup>2</sup>B. Göhler, V. Hamelbeck, T. Markus, M. Kettner, G. Hanne, Z. Vager, R. Naaman, and H. Zacharias, “Spin selectivity in electron transmission through self-assembled monolayers of double-stranded dna,” *Science* **331**, 894–897 (2011).
- <sup>3</sup>H. J. Eckvahl, N. A. Tcyrulnikov, A. Chiesa, J. M. Bradley, R. M. Young, S. Carretta, M. D. Krzyaniak, and M. R. Wasielewski, “Direct observation of chirality-induced spin selectivity in electron donor–acceptor molecules,” *Science* **382**, 197–201 (2023).
- <sup>4</sup>N. Mayer, S. Patchkovskii, F. Morales, M. Ivanov, and O. Smirnova, “Imprinting chirality on atoms using synthetic chiral light fields,” *Physical Review Letters* **129**, 243201 (2022).
- <sup>5</sup>Y. Chen, D. Haase, J. Manz, H. Wang, and Y. Yang, “From chiral laser pulses to femto-and attosecond electronic chirality flips in achiral molecules,” *Nature Communications* **15**, 565 (2024).
- <sup>6</sup>T. Moitra, L. Konecny, M. Kadek, O. Neufeld, A. Rubio, and M. Repisky, “Light-induced persistent electronic chirality in achiral molecules probed with time-resolved electronic circular dichroism spectroscopy,” *The Journal of Physical Chemistry Letters* **16**, 9210–9216 (2025).
- <sup>7</sup>J. Ferrando-Soria, E. Moreno Pineda, A. Chiesa, A. Fernandez, S. A. Magee, S. Carretta, P. Santini, I. J. Vitorica-Yrezabal, F. Tuna, G. A. Timco *et al.*, “A modular design of molecular qubits to implement universal quantum gates,” *Nature communications* **7**, 11377 (2016).
- <sup>8</sup>M. R. Wasielewski, M. D. Forbes, N. L. Frank, K. Kowalski, G. D. Scholes, J. Yuen-Zhou, M. A. Baldo, D. E. Freedman, R. H. Goldsmith, T. Goodson III *et al.*, “Exploiting chemistry and molecular systems for quantum information science,” *Nature Reviews Chemistry* **4**, 490–504 (2020).
- <sup>9</sup>F. Santanni and A. Privitera, “Metalloporphyrins as building blocks for quantum information science,” *Advanced Optical Materials* **12**, 2303036 (2024).
- <sup>10</sup>I. Barth, J. Manz, Y. Shigeta, and K. Yagi, “Unidirectional electronic ring current driven by a few cycle circularly polarized laser pulse: Quantum model simulations for Mg-porphyrin,” *Journal of the American Chemical Society* **128**, 7043–7049 (2006).
- <sup>11</sup>H. Isago, “Prototypical” Optical Absorption Spectra of Phthalocyanines and Their Theoretical Background Springer Japan, Tokyo, (2015), pp. 21–40.
- <sup>12</sup>C. G. Claessens, U. Hahn, and T. Torres, “Phthalocyanines: From outstanding electronic properties to emerging applications,” *The Chemical Record* **8**, 75–97 (2008).
- <sup>13</sup>T. Cao, G. Wang, W. Han, H. Ye, C. Zhu, J. Shi, Q. Niu, P. Tan, E. Wang, B. Liu *et al.*, “Valley-selective circular dichroism of monolayer molybdenum disulphide,” *Nature communications* **3**, 887 (2012).
- <sup>14</sup>J. T. Hougen, “Vibronic interactions in molecules with a fourfold symmetry axis,” *Journal of Molecular Spectroscopy* **13**, 149–167 (1964).
- <sup>15</sup>D. Panariti, A. Carella, S. Ciuti, A. Barbon, N. Holzer, P. Podutoori, Y. Kandrashkin, A. van der Est, and M. Di Valentin, “Long-lived light-induced electron spin polarization in porphyrin triplet states and the dynamic jahn–teller effect,” *The Journal of Chemical Physics* **162** (2025).
- <sup>16</sup>A. Benitz, M. B. Thomas, I. de Silva, V. N. Nesterov, G. F. Verbeck IV, and F. D’Souza, “Photoinduced electron transfer in axially coordinated supramolecular zinc tetrapyrrole bis (styryl)

- bodipy donor-acceptor conjugates,” *ChemPhotoChem* **5**, 260–269 (2021).
- <sup>17</sup>M. Guragain, D. Pinjari, R. Misra, and F. D’Souza, “Zinc tetrapyrrole coordinated to imidazole functionalized tetracyanobutadiene or cyclohexa-2, 5-diene-1, 4-diylidene-expanded-tetracyanobutadiene conjugates: Dark vs. light-induced electron transfer,” *Chemistry–A European Journal* **29**, e202302665 (2023).
  - <sup>18</sup>J. Chen, F. Chen, X. Wang, H. Zhuang, M. Guo, L. Wang, J. Xie, L. Zhang, H. Liu, Y. Shi et al., “Ultra-fast photoelectron transfer in bimetallic porphyrin optoelectrode for single neuron modulation,” *Nature Communications* **15**, 10241 (2024).
  - <sup>19</sup>M. Gouterman, “Spectra of porphyrins,” *Journal of Molecular Spectroscopy* **6**, 138–163 (1961).
  - <sup>20</sup>Z. E. Dance, S. M. Mickley, T. M. Wilson, A. B. Ricks, A. M. Scott, M. A. Ratner, and M. R. Wasielewski, “Intersystem crossing mediated by photoinduced intramolecular charge transfer: julolidine- anthracene molecules with perpendicular  $\pi$  systems,” *The Journal of Physical Chemistry A* **112**, 4194–4201 (2008).
  - <sup>21</sup>T. P. Fay and D. T. Limmer, “Unraveling the mechanisms of triplet state formation in a heavy-atom free photosensitizer,” *Chemical Science* **15**, 6726–6737 (2024).
  - <sup>22</sup>M. Malley, G. Feher, and D. Mauzerall, “The zeeman effect in porphyrins,” *Journal of Molecular Spectroscopy* **26**, 320–334 (1968).
  - <sup>23</sup>C. M. Marian, “Spin-orbit coupling and intersystem crossing in molecules,” *Wiley Interdisciplinary Reviews: Computational Molecular Science* **2**, 187–203 (2012).
  - <sup>24</sup>M. Gouterman, F. P. Schwarz, P. D. Smith, and D. Dolphin, “Porphyrins. xxvii. spin-orbit coupling and luminescence of group iv complexes,” *The Journal of Chemical Physics* **59**, 676–690 (1973).
  - <sup>25</sup>R. L. Ake and M. Gouterman, “Porphyrins xiv. theory for the luminescent state in vo, co, cu complexes,” *Theoretica chimica acta* **15**, 20–42 (1969).
  - <sup>26</sup>T. P. Fay and D. T. Limmer, “Origin of chirality induced spin selectivity in photoinduced electron transfer,” *Nano letters* **21**, 6696–6702 (2021).
  - <sup>27</sup>H. L. Kee, J. Bhaumik, J. R. Diers, P. Mroz, M. R. Hamblin, D. F. Bocian, J. S. Lindsey, and D. Holten, “Photophysical characterization of imidazolium-substituted pd (ii), in (iii), and zn (ii) porphyrins as photosensitizers for photodynamic therapy,” *Journal of Photochemistry and Photobiology A: Chemistry* **200**, 346–355 (2008).
  - <sup>28</sup>E. A. Weiss, M. A. Ratner, and M. R. Wasielewski, “Direct measurement of singlet- triplet splitting within rodlike photogenerated radical ion pairs using magnetic field effects: Estimation of the electronic coupling for charge recombination,” *The Journal of Physical Chemistry A* **107**, 3639–3647 (2003).
  - <sup>29</sup>M. Ortiz, S. Cho, J. Niklas, S. Kim, O. G. Poluektov, W. Zhang, G. Rumbles, and J. Park, “Through-space ultrafast photoinduced electron transfer dynamics of a c70-encapsulated bisporphyrin covalent organic polyhedron in a low-dielectric medium,” *Journal of the American Chemical Society* **139**, 4286–4289 (2017).
  - <sup>30</sup>G. E. Pieslinger, I. Ramírez-Wierzbicki, and A. Cadranell, “The excited-state creutz-taube ion,” *Angewandte Chemie International Edition* **61**, e202211747 (2022).
  - <sup>31</sup>Y. K. Kang, P. Zhang, I. V. Rubtsov, J. Zheng, G. Bullard, D. N. Beratan, and M. J. Therien, “Orientational dependence of cofacial porphyrin-quinone electronic interactions within the strong coupling regime,” *The Journal of Physical Chemistry B* **123**, 10456–10462 (2019).
  - <sup>32</sup>J. Li, P. I. Djurovich, B. D. Alleyne, M. Yousufuddin, N. N. Ho, J. C. Thomas, J. C. Peters, R. Bau, and M. E. Thompson, “Synthetic control of excited-state properties in cyclometalated ir (iii) complexes using ancillary ligands,” *Inorganic Chemistry* **44**, 1713–1727 (2005).
  - <sup>33</sup>D. Gómez Bustos, S. Sreenivasan, and B. Pinter, “A computational study on the effect of structural isomerism on the excited state lifetime and redox energetics of archetype iridium photoredox catalyst platforms [ir (ppy) 2 (bpy)]+ and ir (ppy) 3,” *The Journal of Chemical Physics* **162** (2025).
  - <sup>34</sup>R. Ma, P. Guo, H. Cui, X. Zhang, M. K. Nazeeruddin, and M. Grätzel, “Substituent effect on the meso-substituted porphyrins: theoretical screening of sensitizer candidates for dye-sensitized solar cells,” *The Journal of Physical Chemistry A* **113**, 10119–10124 (2009).
  - <sup>35</sup>J. Barbee and A. E. Kuznetsov, “Revealing substituent effects on the electronic structure and planarity of ni-porphyrins,” *Computational and Theoretical Chemistry* **981**, 73–85 (2012).
  - <sup>36</sup>C. Galli, K. Wynne, S. M. LeCours, M. Therien, and R. Hochstrasser, “Direct measurement of electronic dephasing using anisotropy,” *Chemical physics letters* **206**, 493–499 (1993).
  - <sup>37</sup>D. T. Limmer, *Statistical mechanics and stochastic thermodynamics: A textbook on modern approaches in and out of equilibrium* Oxford University Press, (2024).
  - <sup>38</sup>J. Jeske, J. David, M. B. Plenio, S. F. Huelga, and J. H. Cole, “Bloch-redfield equations for modeling light-harvesting complexes,” *The Journal of chemical physics* **142** (2015).
  - <sup>39</sup>F. Campaioli, J. H. Cole, and H. Hapuarachchi, “Quantum master equations: Tips and tricks for quantum optics, quantum computing, and beyond,” *PRX Quantum* **5**, 020202 (2024).
  - <sup>40</sup>K. R. Nandipati and O. Vendrell, “Dynamical jahn-teller effects on the generation of electronic ring currents by circularly polarized light,” *Physical Review Research* **3**, L042003 (2021).
  - <sup>41</sup>A. Ishizaki and G. R. Fleming, “Theoretical examination of quantum coherence in a photosynthetic system at physiological temperature,” *Proceedings of the National Academy of Sciences* **106**, 17255–17260 (2009).
  - <sup>42</sup>D. Hu, B. X. Chng, W. Ying, and P. Huo, “Trajectory-based non-adiabatic simulations of the polariton relaxation dynamics,” *The Journal of Chemical Physics* **162** (2025).
  - <sup>43</sup>U. Fano, “Spin orientation of photoelectrons ejected by circularly polarized light,” *Physical Review* **178**, 131 (1969).
  - <sup>44</sup>N. Cherepkov, “Spin polarization of atomic and molecular photoelectrons,” *Advances in atomic and molecular physics* **19**, 395–447 (1983).
  - <sup>45</sup>Y. Liu, T. Handa, N. Olsen, C. Nuckolls, and X. Zhu, “Spin-polarized charge separation at two-dimensional semiconductor/molecule interfaces,” *Journal of the American Chemical Society* **146**, 10052–10059 (2024).
  - <sup>46</sup>Z. Zeng, M. Först, M. Fechner, M. Buzzi, E. Amuah, C. Putzke, P. Moll, D. Prabhakaran, P. Radaelli, and A. Cavalleri, “Photo-induced chirality in a nonchiral crystal,” *Science* **387**, 431–436 (2025).
  - <sup>47</sup>M. Delor, P. A. Scattergood, I. V. Sazanovich, A. W. Parker, G. M. Greetham, A. J. Meijer, M. Towrie, and J. A. Weinstein, “Toward control of electron transfer in donor-acceptor molecules by bond-specific infrared excitation,” *Science* **346**, 1492–1495 (2014).

# Supplementary information: Spin Polarization from Circularly Polarized Light Induced Charge Transfer

Sindhana Pannir-Sivajothi<sup>1</sup> and David Limmer<sup>1, 2, 3, 4, \*</sup>

<sup>1</sup>*Department of Chemistry, University of California, Berkeley, California 94720, United States*

<sup>2</sup>*Kavli Energy Nanoscience Institute at Berkeley, Berkeley, California 94720, United States*

<sup>3</sup>*Chemical Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720, United States*

<sup>4</sup>*Material Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720, United States*

## S1. GOUTERMAN FOUR-ORBITAL MODEL

Metalloporphyrins approximately belong to the  $D_{4h}$  point group and the Gouterman orbitals include two exactly degenerate LUMOs that have  $e_g$  symmetry and two accidentally degenerate HOMOs that have  $a_{1u}$  and  $a_{2u}$  symmetries [1, 2]. Let the HOMOs with  $a_{2u}$  symmetry have creation operator  $\hat{d}_{H_1\sigma}^\dagger$  and with  $a_{1u}$  symmetry have creation operator  $\hat{d}_{H_2\sigma}^\dagger$  where  $\sigma = \alpha, \beta$  is the spin and we use restricted spin orbitals (the convention is the same as Gouterman's in [1]). Let the LUMOs with  $e_{gy}$  symmetry (transforms as  $yz$ ) and  $e_{gx}$  symmetry (transforms as  $xz$ ) have creation operators  $\hat{d}_{L_1\sigma}^\dagger$  and  $\hat{d}_{L_2\sigma}^\dagger$ , respectively. Then the transition dipole  $\langle \Psi_0 | \hat{\mu} \hat{d}_{L_2\sigma}^\dagger \hat{d}_{H_1\sigma} | \Psi_0 \rangle$  is along  $+x$ , while  $\langle \Psi_0 | \hat{\mu} \hat{d}_{L_1\sigma}^\dagger \hat{d}_{H_2\sigma} | \Psi_0 \rangle$  is along  $-x$ . On the other hand, both  $\langle \Psi_0 | \hat{\mu} \hat{d}_{L_1\sigma}^\dagger \hat{d}_{H_1\sigma} | \Psi_0 \rangle$  and  $\langle \Psi_0 | \hat{\mu} \hat{d}_{L_2\sigma}^\dagger \hat{d}_{H_2\sigma} | \Psi_0 \rangle$  are along  $+y$ . The singlet Q-band electronic states in terms of the real-valued frontier molecular orbitals with creation operators  $\hat{d}_{H_1\sigma}^\dagger$ ,  $\hat{d}_{H_2\sigma}^\dagger$ ,  $\hat{d}_{L_1\sigma}^\dagger$ , and  $\hat{d}_{L_2\sigma}^\dagger$ , are

$$|{}^1Q_x\rangle = \frac{1}{2} \left[ (\hat{d}_{L_2\alpha}^\dagger \hat{d}_{H_1\alpha} + \hat{d}_{L_2\beta}^\dagger \hat{d}_{H_1\beta}) + (\hat{d}_{L_1\alpha}^\dagger \hat{d}_{H_2\alpha} + \hat{d}_{L_1\beta}^\dagger \hat{d}_{H_2\beta}) \right] | \Psi_0 \rangle, \quad (S1a)$$

$$|{}^1Q_y\rangle = \frac{1}{2} \left[ (\hat{d}_{L_1\alpha}^\dagger \hat{d}_{H_1\alpha} + \hat{d}_{L_1\beta}^\dagger \hat{d}_{H_1\beta}) - (\hat{d}_{L_2\alpha}^\dagger \hat{d}_{H_2\alpha} + \hat{d}_{L_2\beta}^\dagger \hat{d}_{H_2\beta}) \right] | \Psi_0 \rangle. \quad (S1b)$$

The donor  $\lambda = +, -$  singlet and triplet excited states are defined as [3]

$$|{}^1D_{\pm}\rangle = \frac{1}{\sqrt{2}} \left( |{}^1Q_x\rangle \pm i |{}^1Q_y\rangle \right) = \frac{1}{\sqrt{2}} (\hat{d}_{L_{\pm\alpha}}^\dagger \hat{d}_{H_{\pm\alpha}} + \hat{d}_{L_{\pm\beta}}^\dagger \hat{d}_{H_{\pm\beta}}) | \Psi_0 \rangle, \quad (S2a)$$

$$|{}^3D_{\pm}\rangle = \frac{1}{\sqrt{2}} \left( |{}^3Q_x\rangle \pm i |{}^3Q_y\rangle \right) = \frac{1}{\sqrt{2}} (\hat{d}_{L_{\pm\alpha}}^\dagger \hat{d}_{H_{\pm\alpha}} - \hat{d}_{L_{\pm\beta}}^\dagger \hat{d}_{H_{\pm\beta}}) | \Psi_0 \rangle, \quad (S2b)$$

where

$$\hat{d}_{L_{\pm\sigma}}^\dagger = \frac{1}{\sqrt{2}} (\hat{d}_{L_2\sigma}^\dagger \pm i \hat{d}_{L_1\sigma}^\dagger), \quad (S3a)$$

$$\hat{d}_{H_{\pm\sigma}}^\dagger = \frac{1}{\sqrt{2}} (\hat{d}_{H_1\sigma}^\dagger \pm i \hat{d}_{H_2\sigma}^\dagger). \quad (S3b)$$

## S2. MOLECULAR ORBITALS AND HÜCKEL THEORY

We write down a Hückel Hamiltonian in the basis of the atomic orbitals involved in the  $\pi$ -conjugation within the donor and acceptor. Each donor atom contributes a  $p_z$  orbital, each acceptor atom a  $p_x$  orbital, and the metal center contributes both  $d_{xz}$  and  $d_{yz}$  orbitals. Let the donor-acceptor complex contain  $N_a$  atoms, with indices  $j \in \{1, 2, \dots, N_a - 1\}$  labeling all atoms except the metal. Define  $n_D$  as the set of donor atom indices and  $n_A$  as those of the acceptor excluding the metal. Since only the metal is shared between donor and acceptor,  $n_D \cap n_A = \emptyset$  and  $n_D \cup n_A = \{1, 2, \dots, N_a - 1\}$ . We denote the atomic orbitals (AOs) as  $|p_{z,j}\rangle$  for  $j \in n_D$ ,  $|p_{x,j}\rangle$  for  $j \in n_A$ , and  $|d_{xz}\rangle, |d_{yz}\rangle$  for the metal center. Let  $nn_j$  denote the neighbors of atom  $j$  (excluding the metal) and  $nn_M$  the

---

\* [dlimmer@berkeley.edu](mailto:dlimmer@berkeley.edu)

neighbors of the metal. The single-electron form of the mean-field Hamiltonian  $\hat{h} = \hat{h}_D + \hat{h}_A + \hat{v} + \hat{v}_{\text{SOC}}$  contains a Hückel term that is partitioned into donor  $\hat{h}_D$ , acceptor  $\hat{h}_A$ , and coupling  $\hat{v}$ , and a mean-field SOC term  $\hat{v}_{\text{SOC}}$ , where

$$\begin{aligned}
\hat{h}_D &= \sum_{j \in n_D} \alpha_j |p_{z,j}\rangle \langle p_{z,j}| + \alpha_x |d_{xz}\rangle \langle d_{xz}| + \alpha_y |d_{yz}\rangle \langle d_{yz}| \\
&\quad + \sum_{j' \in nn_M \cap n_D} \left( \beta_{xj'} |d_{xz}\rangle \langle p_{z,j'}| + \beta_{yj'} |d_{yz}\rangle \langle p_{z,j'}| + \text{h.c.} \right) + \sum_{j \in n_D} \sum_{j' \in nn_j} \beta_{jj'} |p_{z,j}\rangle \langle p_{z,j'}|, \\
\hat{h}_A &= \sum_{j \in n_A} \alpha_j |p_{x,j}\rangle \langle p_{x,j}| + \alpha_x |d_{xz}\rangle \langle d_{xz}| + \sum_{j \in n_A} \sum_{j' \in nn_j} \beta_{jj'} |p_{x,j}\rangle \langle p_{x,j'}| + \sum_{j' \in nn_M \cap n_A} \left( \beta_{xj'} |d_{xz}\rangle \langle p_{x,j'}| + \text{h.c.} \right), \\
\hat{v} &= -\alpha_x |d_{xz}\rangle \langle d_{xz}|, \\
\hat{v}_{\text{SOC}} &= \left[ \xi_M |d_{yz}\rangle \langle d_{xz}| i\hbar + \text{h.c.} \right] \cdot \hat{s}_z,
\end{aligned} \tag{S4}$$

with  $\beta_{jj'} = \beta_{jj'}^*$ ,  $\beta_{j'x} = \beta_{xj'}^*$ , and  $\beta_{j'y} = \beta_{yj'}^*$ , and the term  $\hat{v} = -\alpha_x |d_{xz}\rangle \langle d_{xz}|$  accounts for the double counting of the  $d_{xz}$  orbital energy in both  $\hat{h}_D$  and  $\hat{h}_A$ . The form  $\hat{v}_{\text{SOC}}$  of the SOC coupling can be derived from the Breit-Pauli form  $\hat{H}_{\text{SOC}} = \sum_j \sum_I \xi(r_{jI}) \hat{\mathbf{l}}_{jI} \cdot \hat{\mathbf{s}}_j$  where  $j$  labels the electrons and  $I$  the nuclei [4] keeping only single-center integrals. Within our AO basis, only the metal center contributes to the one-center SOC [5, 6],  $\hat{H}_{\text{SOC}} \approx \sum_j \xi(r_{jM}) \hat{\mathbf{l}}_{jM, z_{\text{mol}}} \hat{s}_{j, z_{\text{mol}}}$  where  $M$  labels the metal nucleus. The one-center SOC terms are defined as matrix elements where the angular momentum operator  $\hat{\mathbf{l}}_{jI}$  and both AOs are centered around the same nucleus. Because the angular momentum operators are pure imaginary operators and have zero diagonal matrix elements in a real-valued AO basis and since we retain only one AO per atom except for the metal, it is evident that only the metal center will contribute to the one-center SOC. Therefore, we can write the single-electron SOC term of the Hamiltonian as  $\hat{v}_{\text{SOC}} = [\xi_M |d_{yz}\rangle \langle d_{xz}| i\hbar + \text{h.c.}] \hat{s}_z$  where  $\xi_M = \langle d_{yz} | \xi(r_{jM}) | d_{xz} \rangle$  is real-valued.

By diagonalizing  $\hat{h}_D$ , we can compute the degenerate pair of HOMOs and LUMOs of the donor given by creation operators  $\hat{d}_{H_1\sigma}^\dagger$ ,  $\hat{d}_{H_2\sigma}^\dagger$ ,  $\hat{d}_{L_1\sigma}^\dagger$  and  $\hat{d}_{L_2\sigma}^\dagger$  as a LCAO, and similarly by diagonalizing  $\hat{h}_A$ , we can obtain the LUMO of the acceptor given by creation operator  $\hat{a}_{L\sigma}^\dagger$ . The MOs are  $\hat{d}_{L_\lambda\sigma}^\dagger |0\rangle = |\phi_{D,L_\lambda}\rangle |\sigma\rangle$ ,  $\hat{d}_{H_\lambda\sigma}^\dagger |0\rangle = |\phi_{D,H_\lambda}\rangle |\sigma\rangle$ ,  $\hat{a}_{L\sigma}^\dagger |0\rangle = |\phi_{A,L}\rangle |\sigma\rangle$ , and they satisfy  $\hat{h}_D |\phi_{D,L_\lambda}\rangle = \epsilon_{D,L} |\phi_{D,L_\lambda}\rangle$ ,  $\hat{h}_D |\phi_{D,H_\lambda}\rangle = \epsilon_{D,H} |\phi_{D,H_\lambda}\rangle$ , and  $\hat{h}_A |\phi_{A,L}\rangle = \epsilon_{A,L} |\phi_{A,L}\rangle$ . Both the spin-preserving and the spin-orbit coupled electron transfer happens via the  $d_\pi$  orbitals of the metal; therefore, the contribution of the  $d_\pi$  orbitals of the metal to the LUMOs of the donor and acceptor determine the charge transfer matrix elements. Due to symmetry, the  $d_{xz}$  orbital only contributes to  $\hat{d}_{L_2\sigma}^\dagger$  and  $\hat{a}_{L\sigma}^\dagger$  MOs, and  $d_{yz}$  orbitals only to  $\hat{d}_{L_1\sigma}^\dagger$ . Since  $\hat{d}_{L_1\sigma}^\dagger$ ,  $\hat{d}_{L_2\sigma}^\dagger$  and  $\hat{a}_{L\sigma}^\dagger$  are creation operators for purely real MOs,  $B_{xz} = \langle d_{xz} | \phi_{D,L_2} \rangle$ ,  $B_{yz} = \langle d_{yz} | \phi_{D,L_1} \rangle$  and  $A_{xz} = \langle d_{xz} | \phi_{A,L} \rangle$  are purely real. Then we can show using Eq. S3 that  $\langle d_{xz} | \phi_{D,L_\pm} \rangle = B_{xz}/\sqrt{2}$  and  $\langle d_{yz} | \phi_{D,L_\pm} \rangle = \pm i B_{yz}/\sqrt{2}$ . The matrix elements of the frontier MOs are

$$\begin{aligned}
\langle \phi_{D,H_\lambda} | (\hat{h}_D + \hat{h}_A + \hat{v}) | \phi_{D,H_\lambda} \rangle &= \epsilon_{D,H} \langle \phi_{D,H_\lambda} | \phi_{D,H_\lambda} \rangle + \langle \phi_{D,H_\lambda} | (\hat{h}_A + \hat{v}) | \phi_{D,H_\lambda} \rangle \\
&= \epsilon_{D,H} \\
\langle \phi_{D,L_\lambda} | (\hat{h}_D + \hat{h}_A + \hat{v}) | \phi_{D,L_\lambda} \rangle &= \epsilon_{D,L} \langle \phi_{D,L_\lambda} | \phi_{D,L_\lambda} \rangle + \langle \phi_{D,L_\lambda} | (\hat{h}_A + \hat{v}) | \phi_{D,L_\lambda} \rangle \\
&\approx \epsilon_{D,L} \\
\langle \phi_{A,L} | (\hat{h}_D + \hat{h}_A + \hat{v}) | \phi_{A,L} \rangle &= \epsilon_{A,L} \langle \phi_{A,L} | \phi_{A,L} \rangle + \langle \phi_{A,L} | (\hat{h}_D + \hat{v}) | \phi_{A,L} \rangle \\
&\approx \epsilon_{A,L} \\
\langle \phi_{D,L_\lambda} | (\hat{h}_D + \hat{h}_A + \hat{v}) | \phi_{A,L} \rangle &= \langle \phi_{D,L_\lambda} | \hat{h}_D | \phi_{A,L} \rangle + \langle \phi_{D,L_\lambda} | \hat{h}_A | \phi_{A,L} \rangle + \langle \phi_{D,L_\lambda} | \hat{v} | \phi_{A,L} \rangle \\
&= \epsilon_{D,L} \langle \phi_{D,L_\lambda} | \phi_{A,L} \rangle + \epsilon_{A,L} \langle \phi_{D,L_\lambda} | \phi_{A,L} \rangle - \alpha_x \langle \phi_{D,L_\lambda} | d_{xz} \rangle \langle d_{xz} | \phi_{A,L} \rangle \\
&\approx (\epsilon_{D,L} + \epsilon_{A,L} - \alpha_x) \langle \phi_{D,L_\lambda} | d_{xz} \rangle \langle d_{xz} | \phi_{A,L} \rangle \\
&= (\epsilon_{D,L} + \epsilon_{A,L} - \alpha_x) B_{xz} A_{xz} / \sqrt{2}
\end{aligned} \tag{S5}$$



and

$$\begin{aligned}
\langle \phi_{D,H_\lambda} | (\hat{h}_D + \hat{h}_A + \hat{v}) | \phi_{A,L} \rangle &= \langle \phi_{D,H_\lambda} | \hat{h}_D | \phi_{A,L} \rangle + \langle \phi_{D,H_\lambda} | \hat{h}_A | \phi_{A,L} \rangle + \langle \phi_{D,H_\lambda} | \hat{v} | \phi_{A,L} \rangle \\
&= \epsilon_{D,H} \langle \phi_{D,H_\lambda} | \phi_{A,L} \rangle + \epsilon_{A,L} \langle \phi_{D,H_\lambda} | \phi_{A,L} \rangle - \alpha_x \langle \phi_{D,H_\lambda} | d_{xz} \rangle \langle d_{xz} | \phi_{A,L} \rangle \\
&\approx (\epsilon_{D,H} + \epsilon_{A,L} - \alpha_x) \langle \phi_{D,H_\lambda} | d_{xz} \rangle \langle d_{xz} | \phi_{A,L} \rangle \\
&= 0 \\
\langle \phi_{D,H_{\lambda_1}} | (\hat{h}_D + \hat{h}_A + \hat{v}) | \phi_{D,L_{\lambda_2}} \rangle &= 0 \\
\langle \phi_{D,H_+} | (\hat{h}_D + \hat{h}_A + \hat{v}) | \phi_{D,H_-} \rangle &= 0 \\
\langle \phi_{D,L_+} | (\hat{h}_D + \hat{h}_A + \hat{v}) | \phi_{D,L_-} \rangle &= \langle \phi_{D,L_+} | (\hat{h}_A + \hat{v}) | \phi_{D,L_-} \rangle \\
&\approx 0 \\
\langle \phi_{D,L_\pm} | \langle \sigma_1 | \hat{v}_{\text{SOC}} | \sigma_2 \rangle | \phi_{D,L_\pm} \rangle &= \langle \phi_{D,L_\pm} | \left[ i\hbar\xi_M |d_{yz}\rangle \langle d_{xz}| - i\hbar\xi_M |d_{xz}\rangle \langle d_{yz}| \right] | \phi_{D,L_\pm} \rangle \langle \sigma_1 | \hat{s}_z | \sigma_2 \rangle \\
&= \left[ i\hbar\xi_M (\mp iB_{yz})B_{xz}/2 - i\hbar\xi_M B_{xz}(\pm iB_{yz}/2) \right] \langle \sigma_1 | \hat{s}_z | \sigma_2 \rangle \\
&= \pm \langle \sigma_1 | \hat{s}_z | \sigma_2 \rangle \hbar\xi_M B_{yz}B_{xz} \\
\langle \phi_{D,L_+} | \langle \sigma_1 | \hat{v}_{\text{SOC}} | \sigma_2 \rangle | \phi_{D,L_-} \rangle &= \langle \phi_{D,L_+} | \left[ i\hbar\xi_M |d_{yz}\rangle \langle d_{xz}| - i\hbar\xi_M |d_{xz}\rangle \langle d_{yz}| \right] | \phi_{D,L_-} \rangle \langle \sigma_1 | \hat{s}_z | \sigma_2 \rangle \\
&= \left[ i\hbar\xi_M (\mp iB_{yz})B_{xz}/2 - i\hbar\xi_M B_{xz}(\mp iB_{yz}/2) \right] \langle \sigma_1 | \hat{s}_z | \sigma_2 \rangle \\
&= 0 \\
\langle \phi_{D,H_{\lambda_1}} | \langle \sigma_1 | \hat{v}_{\text{SOC}} | \sigma_2 \rangle | \phi_{D,H_{\lambda_2}} \rangle &= 0 \\
\langle \phi_{D,L_\pm} | \langle \sigma_1 | \hat{v}_{\text{SOC}} | \sigma_2 \rangle | \phi_{A,L} \rangle &= \langle \phi_{L_\pm} | \left[ i\hbar\xi_M |d_{yz}\rangle \langle d_{xz}| - i\hbar\xi_M |d_{xz}\rangle \langle d_{yz}| \right] | \phi_{L,a} \rangle \langle \sigma_1 | \hat{s}_z | \sigma_2 \rangle \\
&= \pm \langle \sigma_1 | \hat{s}_z | \sigma_2 \rangle \hbar\xi_M B_{yz}A_{xz}/\sqrt{2} \\
\langle \phi_{A,L} | \langle \sigma_1 | \hat{v}_{\text{SOC}} | \sigma_2 \rangle | \phi_{A,L} \rangle &= 0
\end{aligned} \tag{S6}$$

where  $\lambda_1, \lambda_2 \in \{+, -\}$  and we have assumed that the overlap of atomic orbitals of non-neighboring atoms is zero, so  $\langle p_{z,j} | p_{x,j'} \rangle = 0$ . The atomic orbital basis is non-orthogonal; therefore, the orbital  $|d_{xz}\rangle$  has non-zero overlap with  $|p_{z,j}\rangle$  of adjacent atoms, so  $\langle \phi_{D,L_\pm} | \phi_{A,L} \rangle \neq \langle \phi_{D,L_\pm} | d_{xz} \rangle \langle d_{xz} | \phi_{A,L} \rangle$  but it will be almost equal to it.

The many-electron mean-field Hamiltonian is  $\hat{H}_{\text{MF}} = \sum_j [\hat{h}_D(j) + \hat{h}_A(j) + \hat{v}(j) + \hat{v}_{\text{SOC}}(j)]$  where  $j$  labels the electrons. After projecting this Hamiltonian into the subspace of frontier orbitals using the matrix elements in Eq. S5 and S6, it decomposes into  $\lambda = \pm$  sectors  $\hat{\Pi}_{\text{frontier}} \hat{H}_{\text{MF}} \hat{\Pi}_{\text{frontier}} = \sum_{\lambda=+,-} \hat{H}_\lambda$  with

$$\begin{aligned}
\hat{H}_\lambda &= \sum_{\sigma=\alpha,\beta} \left( \epsilon_{D,L} \hat{d}_{L\lambda\sigma}^\dagger \hat{d}_{L\lambda\sigma} + \epsilon_{D,H} \hat{d}_{H\lambda\sigma}^\dagger \hat{d}_{H\lambda\sigma} + \epsilon_{A,L} \hat{a}_{L\sigma}^\dagger \hat{a}_{L\sigma} + v_0 \hat{d}_{L\lambda\sigma}^\dagger \hat{a}_{L\sigma} + v_0 \hat{d}_{L\lambda\sigma} \hat{a}_{L\sigma}^\dagger \right) \\
&\quad + v_{\text{ISC}_\lambda} \left( \hat{d}_{L\lambda\alpha}^\dagger \hat{d}_{L\lambda\alpha} - \hat{d}_{L\lambda\beta}^\dagger \hat{d}_{L\lambda\beta} \right) + v_{\text{SOC}_\lambda} \left( \hat{d}_{L\lambda\alpha}^\dagger \hat{a}_{L\alpha} - \hat{d}_{L\lambda\beta}^\dagger \hat{a}_{L\beta} + \text{h.c.} \right),
\end{aligned} \tag{S7}$$

where  $v_0 \approx (\epsilon_{D,L} + \epsilon_{A,L} - \alpha_x) A_{xz} B_{xz} / \sqrt{2}$ ,  $v_{\text{ISC}_\pm} = \pm \xi_M B_{xz} B_{yz} \hbar^2 / 2$  and  $v_{\text{SOC}_\pm} = \pm \xi_M A_{xz} B_{yz} \hbar^2 / 2\sqrt{2}$  are all real-valued.

The mean-field Hamiltonian  $\hat{H}_{\text{MF}}$  only contains single-electron terms and does not include exchange coupling that results in the singlet-triplet energy gap. We can phenomenologically include the two-electron exchange interaction term  $\hat{V}_{\text{ex}} = J \sum_{\lambda_1} \sum_{\lambda_2} \sum_{\sigma=\alpha,\beta} \sum_{\tau=\alpha,\beta} \hat{d}_{L\lambda_2\sigma}^\dagger \hat{d}_{H\lambda_1\tau}^\dagger \hat{d}_{L\lambda_2\tau} \hat{d}_{H\lambda_1\sigma}$  that results in an energy gap between  $|^1D_\lambda\rangle$  and  $|^3D_\lambda\rangle$ . The Hamiltonian  $\hat{H}_e$  defined in Eq. 3 of the main text represents  $\hat{H}_{\text{MF}} + \hat{V}_{\text{ex}}$  in the subspace spanned by  $|^1D_\lambda\rangle$ ,  $|^1CT_\lambda\rangle$ , and  $|^3CT_\lambda\rangle$  as the triplet donor is far detuned from the other states, and therefore all calculations in this work are performed with this Hamiltonian. For completeness, here we include the mean-field Hamiltonian along

with the exchange coupling in the space spanned by  $|^1D_\lambda\rangle$ ,  $|^3D_\lambda\rangle$ ,  $|^1CT_\lambda\rangle$ , and  $|^3CT_\lambda\rangle$ ,

$$\begin{aligned} \hat{H}_{e-4} = \sum_{\lambda=+,-} & \left( \Delta |^1CT_\lambda\rangle \langle ^1CT_\lambda| + \Delta |^3CT_\lambda\rangle \langle ^3CT_\lambda| - 2J |^3D_\lambda\rangle \langle ^3D_\lambda| \right. \\ & + v_0 |^1D_\lambda\rangle \langle ^1CT_\lambda| + v_0 |^1CT_\lambda\rangle \langle ^1D_\lambda| + v_0 |^3D_\lambda\rangle \langle ^3CT_\lambda| + v_0 |^3CT_\lambda\rangle \langle ^3D_\lambda| \\ & + v_{\text{SOC}_\lambda} |^1D_\lambda\rangle \langle ^3CT_\lambda| + v_{\text{SOC}_\lambda} |^3CT_\lambda\rangle \langle ^1D_\lambda| + v_{\text{SOC}_\lambda} |^3D_\lambda\rangle \langle ^1CT_\lambda| + v_{\text{SOC}_\lambda} |^1CT_\lambda\rangle \langle ^3D_\lambda| \\ & \left. + v_{\text{ISC}_\lambda} |^1D_\lambda\rangle \langle ^3D_\lambda| + v_{\text{ISC}_\lambda} |^3D_\lambda\rangle \langle ^1D_\lambda| \right). \end{aligned} \quad (\text{S8})$$

We omit the triplet states with spin projection  $M_S = \pm 1$ , because the initial state  $|^1D_\pm\rangle$  lies in the  $M_S = 0$  manifold, and given  $[\sum_\lambda \hat{H}_\lambda, \hat{S}_z] = 0$ , the dynamics conserves  $M_S$ .

The term with coefficient  $v_{\text{ISC}_\lambda}$  in Eq. S8 comes from the difference in energy between a spin-up and a spin-down electron occupying the  $\hat{d}_{L\pm,\sigma}^\dagger$  orbitals due to the magnetic field generated by the ring currents in these MOs. The orbital angular momentum of these states interacts with the spin analogous to an effective magnetic field via spin-orbit coupling. By comparison, due to the magnetic field generated by the hole current in the charge transfer states (which is why they retain the label  $\lambda$ ), we would expect intersystem crossing between charge transfer states  $|^1CT_\lambda\rangle$  and  $|^3CT_\lambda\rangle$ . However, since the HOMOs of the donor do not have any metal  $d_{xz}$  or  $d_{yz}$  character, they do not contribute to the spin-orbit coupling and therefore any orbital angular momentum of the hole does not produce an effective magnetic field for the electron spin in the charge transfer states.

### S3. DYNAMICS WITHOUT VIBRONIC COUPLING

In this section, we solve for the dynamics of the system under unitary evolution with  $\hat{H}_e$ , in the absence of vibronic coupling. As the charge transfer states are degenerate, we change basis such that only one of them couples to the singlet donor  $|^1D_\lambda\rangle$ . These new charge transfer states include one that has mostly singlet character  $|^{1\gg 3}CT_\lambda\rangle$  and one with mostly triplet character  $|^{3\gg 1}CT_\lambda\rangle$ ,

$$|^{1\gg 3}CT_\lambda\rangle = \frac{v_0}{\sqrt{v_0^2 + v_{\text{SOC}_\lambda}^2}} |^1CT_\lambda\rangle + \frac{v_{\text{SOC}_\lambda}}{\sqrt{v_0^2 + v_{\text{SOC}_\lambda}^2}} |^3CT_\lambda\rangle \quad (\text{S9a})$$

$$|^{3\gg 1}CT_\lambda\rangle = -\frac{v_{\text{SOC}_\lambda}}{\sqrt{v_0^2 + v_{\text{SOC}_\lambda}^2}} |^1CT_\lambda\rangle + \frac{v_0}{\sqrt{v_0^2 + v_{\text{SOC}_\lambda}^2}} |^3CT_\lambda\rangle \quad (\text{S9b})$$

In this basis, the Hamiltonian

$$\begin{aligned} \hat{H}_e = \sum_{\lambda=+,-} & \left[ \Delta |^{1\gg 3}CT_\lambda\rangle \langle ^{1\gg 3}CT_\lambda| + \Delta |^{3\gg 1}CT_\lambda\rangle \langle ^{3\gg 1}CT_\lambda| \right. \\ & \left. + \sqrt{v_0^2 + v_{\text{SOC}_\lambda}^2} \left( |^{1\gg 3}CT_\lambda\rangle \langle ^1D_\lambda| + |^1D_\lambda\rangle \langle ^{1\gg 3}CT_\lambda| \right) \right]. \end{aligned} \quad (\text{S10})$$

The spin polarization in this basis is

$$\begin{aligned} \hat{P}_z = \sum_\lambda & \frac{\hbar}{v_0^2 + v_{\text{SOC}_\lambda}^2} \left[ 2v_0 v_{\text{SOC}_\lambda} |^{1\gg 3}CT_\lambda\rangle \langle ^{1\gg 3}CT_\lambda| - 2v_0 v_{\text{SOC}_\lambda} |^{3\gg 1}CT_\lambda\rangle \langle ^{3\gg 1}CT_\lambda| \right. \\ & \left. + (v_0^2 - v_{\text{SOC}_\lambda}^2) \left( |^{1\gg 3}CT_\lambda\rangle \langle ^{3\gg 1}CT_\lambda| + \text{h.c.} \right) \right]. \end{aligned} \quad (\text{S11})$$

The eigenstates of  $\hat{H}_e$  are

$$|^1+\lambda\rangle = \cos\theta |^1D_\lambda\rangle + \sin\theta |^{1\gg 3}CT_\lambda\rangle, \quad (\text{S12a})$$

$$|^1-\lambda\rangle = -\sin\theta |^1D_\lambda\rangle + \cos\theta |^{1\gg 3}CT_\lambda\rangle, \quad (\text{S12b})$$

$$|^{3\gg 1}CT_\lambda\rangle = |^{3\gg 1}CT_\lambda\rangle, \quad (\text{S12c})$$

with energies  $E_{1+} = \Delta/2 + \Omega$ ,  $E_{1-} = \Delta/2 - \Omega$  and  $E_3 = \Delta$ , where  $\Omega = \sqrt{(\Delta/2)^2 + v_0^2 + v_{\text{SOC}\lambda}^2}$ ,  $\cos \theta = \sqrt{(\Omega - \Delta/2)/2\Omega}$ , and  $\sin \theta = \sqrt{(\Omega + \Delta/2)/2\Omega}$ . When the initial state is  $|^1\text{D}_\lambda\rangle$ , the donor singlet population is

$$\begin{aligned} p_{|^1\text{D}_+\rangle}(t) &= \langle ^1\text{D}_\lambda | e^{i\hat{H}_e t/\hbar} |^1\text{D}_\lambda \rangle \langle ^1\text{D}_\lambda | e^{-i\hat{H}_e t/\hbar} |^1\text{D}_\lambda \rangle \\ &= \left( e^{iE_{1+}t/\hbar} \cos \theta \langle ^1+_\lambda | - e^{iE_{1-}t/\hbar} \sin \theta \langle ^1-_\lambda | \right) |^1\text{D}_\lambda \rangle \langle ^1\text{D}_\lambda | \\ &\quad \left( e^{-iE_{1+}t/\hbar} \cos \theta |^1+_\lambda \rangle - e^{-iE_{1-}t/\hbar} \sin \theta |^1-_\lambda \rangle \right) \\ &= 1 - \frac{v_0^2 + v_{\text{SOC}\lambda}^2}{2\Omega^2} [1 - \cos(2\Omega t/\hbar)]. \end{aligned} \quad (\text{S13})$$

Similarly, when the donor singlet is the initial state, the charge-transfer states have populations

$$p_{|^1\text{CT}_+\rangle}(t) = \frac{v_0^2}{2\Omega^2} [1 - \cos(2\Omega t/\hbar)], \quad (\text{S14a})$$

$$p_{|^3\text{CT}_+\rangle}(t) = \frac{v_{\text{SOC}+}^2}{2\Omega^2} [1 - \cos(2\Omega t/\hbar)]. \quad (\text{S14b})$$

When the initial state is  $|^1\text{D}_\lambda\rangle$ , the spin polarization

$$\begin{aligned} \langle \hat{P}_z(t) \rangle &= \langle ^1\text{D}_\lambda | e^{i\hat{H}_e t/\hbar} \delta \hat{P}_{z_{\text{mol}}} e^{-i\hat{H}_e t/\hbar} |^1\text{D}_\lambda \rangle \\ &= \left( e^{iE_{1+}t/\hbar} \cos \theta \langle ^1+_\lambda | - e^{iE_{1-}t/\hbar} \sin \theta \langle ^1-_\lambda | \right) \frac{2\hbar v_0 v_{\text{SOC}\lambda}}{v_0^2 + v_{\text{SOC}\lambda}^2} |^1\gg^3\text{CT}\rangle \langle ^1\gg^3\text{CT}| \\ &\quad \left( e^{-iE_{1+}t/\hbar} \cos \theta |^1+_\lambda \rangle - e^{-iE_{1-}t/\hbar} \sin \theta |^1-_\lambda \rangle \right) \\ &= \frac{\hbar v_0 v_{\text{SOC}\lambda}}{\Omega^2} [1 - \cos(2\Omega t/\hbar)]. \end{aligned} \quad (\text{S15})$$

#### S4. RELAXATION RATES: REDFIELD TENSOR

The Bloch-Redfield equations in the Markovian approximation are,

$$\frac{d\rho_{ab}(t)}{dt} = -i\omega_{ab}\rho_{ab}(t) + \sum_{c,d} R_{ab,cd}\rho_{cd}(t), \quad (\text{S16})$$

where  $\rho_{ij} = \langle i | \hat{\rho} | j \rangle$  are matrix elements of the system's density operator with  $|i\rangle$  and  $|j\rangle$  being the eigenstates of  $\hat{H}_e$  with Bohr frequencies  $\omega_{ij} = \omega_i - \omega_j$ , and  $R_{ab,cd}$  is the Redfield tensor. The Redfield tensor

$$R_{ab,cd} = - \left( \delta_{db} \sum_k \Gamma_{ak,kc}^+ - \Gamma_{db,ac}^+ - \Gamma_{db,ac}^- + \delta_{ac} \sum_k \Gamma_{dk,kb}^- \right) \quad (\text{S17})$$

where

$$\begin{aligned} \Gamma_{ab,cd}^+ &= \frac{1}{\hbar^2} \langle a | \hat{V}_e | b \rangle \langle c | \hat{V}_e | d \rangle \tilde{C}(\omega_{dc}) \\ \Gamma_{ab,cd}^- &= \frac{1}{\hbar^2} \langle a | \hat{V}_e | b \rangle \langle c | \hat{V}_e | d \rangle \tilde{C}(\omega_{ab}), \end{aligned} \quad (\text{S18})$$

and  $|a\rangle, |b\rangle, |c\rangle$ , and  $|d\rangle$  are eigenstates of  $\hat{H}_e$ . The spectral representation of the bath correlation function

$$\begin{aligned} \tilde{C}(\omega) &= \int_0^\infty d\tau e^{i\omega\tau} \text{Tr} [\hat{B}(\tau) \hat{B}(0) \hat{\rho}_b], \\ \text{Re}[\tilde{C}(\omega)] &= \pi \hbar J(|\omega|) \left[ n(|\omega|) + \Theta(\omega) \right], \\ \text{Im}[\tilde{C}(\omega)] &= \mathcal{P} \int_0^\infty d\omega' J(\omega') \left[ \frac{n(\omega') + 1}{\omega - \omega'} + \frac{n(\omega')}{\omega + \omega'} \right]. \end{aligned} \quad (\text{S19})$$




FIG. S1. The real part of the spectral function  $S_{JT}(\omega) = \text{Re}[\tilde{C}(\omega)] = \pi\hbar J(|\omega|) [n(|\omega|) + \Theta(\omega)]$  for a Drude spectral density  $J(\omega) = \frac{2\Lambda\omega\omega_c}{\omega^2 + \omega_c^2}$  with  $\Lambda = 0.2$  meV and  $\omega_c = 30$  meV.

The  $\text{Im}[\tilde{C}(\omega)]$  term in the Redfield tensor results in Lamb shifts in the bare Hamiltonian and does not directly affect relaxation, so we ignore it. We define  $S_{JT}(\omega) = \text{Re}[\tilde{C}(\omega)] = \pi\hbar J(|\omega|) [n(|\omega|) + \Theta(\omega)]$  and use a Drude-Lorentz spectral density  $J(\omega) = \frac{2\Lambda\omega\omega_c}{\omega^2 + \omega_c^2}$  with  $\Lambda = 0.2$  meV and  $\omega_c = 30$  meV;  $S_{JT}(\omega)$  for this set of parameters is plotted in Fig. S1. We have  $S_{JT}(0) = 2\pi\Lambda k_B T / \omega_c$ . For numerical results with the Bloch-Redfield equation plotted in Fig. 3 of the main text and Fig. S2, the equations of motion in Eq. S16 were propagated using the adaptive BDF integrator (using `solve_ivp` in SciPy) with relative and absolute tolerances of  $10^{-8}$  and  $10^{-10}$ , respectively. We do not make the secular approximation for the numerical calculations.

Representing  $\hat{V}_e$  in the eigenbasis  $|^1+_{+}\rangle$ ,  $|^{3\gg 1}CT_{+}\rangle$ ,  $|^1-_{+}\rangle$ ,  $|^1+_{-}\rangle$ ,  $|^{3\gg 1}CT_{-}\rangle$ , and  $|^1-_{-}\rangle$  of  $\hat{H}_e$ , where the states are labeled by the numbers 1 through 6, we have

$$\hat{V}_e = \begin{pmatrix} 0 & 0 & 0 & V_{14} & V_{15} & V_{16} \\ 0 & 0 & 0 & V_{24} & V_{25} & V_{26} \\ 0 & 0 & 0 & V_{34} & V_{35} & V_{36} \\ V_{41} & V_{42} & V_{43} & 0 & 0 & 0 \\ V_{51} & V_{52} & V_{53} & 0 & 0 & 0 \\ V_{61} & V_{62} & V_{63} & 0 & 0 & 0 \end{pmatrix} \quad (\text{S20})$$

with

$$\begin{aligned} V_{14} &= \frac{1}{2}F + \frac{\Delta/2}{2\Omega}G \\ V_{15} &= f_{CT} \frac{v_0 v_{\text{SOC}+}}{v_0^2 + v_{\text{SOC}+}^2} \sqrt{2 + \frac{\Delta}{\Omega}} \\ V_{16} &= - \frac{\sqrt{v_0^2 + v_{\text{SOC}+}^2}}{2\Omega} G \\ V_{24} &= -V_{15} \\ V_{25} &= f_{CT} \frac{v_0^2 - v_{\text{SOC}+}^2}{v_0^2 + v_{\text{SOC}+}^2} \\ V_{26} &= f_{CT} \frac{v_0 v_{\text{SOC}+}}{v_0^2 + v_{\text{SOC}+}^2} \sqrt{2 - \frac{\Delta}{\Omega}} \\ V_{34} &= V_{16} \\ V_{35} &= -V_{26} \\ V_{36} &= \frac{1}{2}F - \frac{\Delta/2}{2\Omega}G \end{aligned} \quad (\text{S21})$$






FIG. S2. Relaxation rates. The decay rate (a) of the population difference between transient enantiomers,  $\Gamma_N$ , and (b) the spin-polarization,  $\Gamma_P$ , for different values of system-bath coupling parameters  $f_D$  and  $f_{CT}$  when the initial state is the singlet donor  $|^1D_+\rangle$ .

where

$$F = f_D + f_{CT} \frac{v_0^2 - v_{SOC+}^2}{v_0^2 + v_{SOC+}^2}, \quad (S22a)$$

$$G = -f_D + f_{CT} \frac{v_0^2 - v_{SOC+}^2}{v_0^2 + v_{SOC+}^2}. \quad (S22b)$$

Our initial density matrix  $\hat{\rho}(0) = |^1D_+\rangle \langle ^1D_+|$  in the  $\hat{H}_e$  eigenbasis would only contain non-zero density matrix elements  $\rho_{11}$ ,  $\rho_{33}$ ,  $\rho_{13}$ , and  $\rho_{31}$ . When  $|v_0| \gg |v_{SOC+}|$ , the terms  $V_{15}$ ,  $V_{35}$ ,  $V_{24}$ , and  $V_{26}$  will be much smaller than others; as a result, starting with populations  $\rho_{11}$  and  $\rho_{33}$  will lead to populations  $\rho_{44}$  and  $\rho_{66}$ , but far lower population would flow into  $\rho_{55}$  and  $\rho_{22}$ . For our parameters,  $v_0 = 75\text{meV}$ ,  $v_{SOC\pm} = \pm 25\text{meV}$ , and  $\Delta = -30\text{meV}$ , we have  $\Omega \approx 80\text{meV}$ . Therefore, the energy differences between eigenstates of  $\hat{H}_e$  are  $0\text{meV}$ ,  $\sim 80\text{meV}$  and  $\sim 160\text{meV}$ . From Fig. S1, we see that  $S_{JT}(\omega)$  is about four times larger for  $\omega = 0\text{meV}$  than  $\omega = 80\text{meV}$  and  $\omega = 160\text{meV}$ , so we can focus our attention on terms of the Redfield tensor that involve  $V_{14}$ ,  $V_{25}$  and  $V_{36}$  as they involve degenerate states. When both  $f_D$  and  $f_{CT}$  are positive or negative, the  $F$  dominates both  $V_{14}$  and  $V_{36}$  and we expect the relaxation rate to be  $\propto F^2$  as can be clearly seen from the contour lines in Fig. S2 and from Fig. 3(c) in the main text. When  $f_D \approx -f_{CT} \frac{v_0^2 - v_{SOC+}^2}{v_0^2 + v_{SOC+}^2}$ , we expect the relaxation rate to be  $\propto G^2$  and slower as  $F \approx 0$  and  $G$  has an additional coefficient of  $|\Delta/2\Omega|$  in  $V_{14}$  and  $V_{36}$ . We can qualitatively observe this from the dark blue region in Fig. S2(a-b); however, we have not quantified this, as we need to run longer simulations to obtain reliable  $\Gamma_N$  and  $\Gamma_P$  in this region because the relaxation is slow.

The population difference  $\langle \Delta \hat{N} \rangle = \rho_{11} + \rho_{22} + \rho_{33} - \rho_{44} - \rho_{55} - \rho_{66}$  in the eigenbasis of  $\hat{H}_e$ . As  $S_{JT}(\omega)$  is larger when  $\omega = 0$  than at other Bohr frequencies of  $\hat{H}_e$  for our parameters, within the partial-secular approximation, we only need to consider population transfer between degenerate  $\lambda = \pm$  eigenstates with  $\omega_1 = \omega_4$ ,  $\omega_2 = \omega_5$ , and  $\omega_3 = \omega_6$ . This would give us an approximate equation of motion for  $\Delta \hat{N}$ ,

$$\begin{aligned} \frac{d\langle \Delta \hat{N} \rangle}{dt} &\approx -2R_{11,44}(\rho_{11} - \rho_{44}) - 2R_{22,55}(\rho_{22} - \rho_{55}) - 2R_{33,66}(\rho_{33} - \rho_{66}) \\ &\approx -\frac{S_{JT}(0)}{\hbar^2} F^2 \langle \Delta \hat{N} \rangle \\ &= -\frac{2\pi k_B T}{\hbar} \frac{\Lambda}{\hbar \omega_c} F^2 \langle \Delta \hat{N} \rangle, \end{aligned} \quad (S23)$$

where we have ignored  $R_{22,55}(\rho_{22} - \rho_{55})$  as initially there is no population in the states involved  $\rho_{22}(0) = \rho_{55}(0) = 0$  and we have also assumed  $|\Delta G/2\Omega| \ll |F|$ , so  $V_{14} \approx V_{36} \approx F/2$ .

## REFERENCES

- [1] M. Gouterman, "Spectra of porphyrins," *Journal of Molecular Spectroscopy* **6**, 138–163 (1961).
- [2] M. Gouterman, G. H. Wagnière, and L. C. Snyder, "Spectra of porphyrins: Part ii. four orbital model," *Journal of Molecular Spectroscopy* **11**, 108–127 (1963).
- [3] I. Barth, J. Manz, Y. Shigeta, and K. Yagi, "Unidirectional electronic ring current driven by a few cycle circularly polarized laser pulse: Quantum model simulations for Mg-porphyrin," *Journal of the American Chemical Society* **128**, 7043–7049 (2006).
- [4] C. M. Marian, "Spin–orbit coupling and intersystem crossing in molecules," *Wiley Interdisciplinary Reviews: Computational Molecular Science* **2**, 187–203 (2012).
- [5] M. Gouterman, F. P. Schwarz, P. D. Smith, and D. Dolphin, "Porphyrins. xxvii. spin-orbit coupling and luminescence of group iv complexes," *The Journal of Chemical Physics* **59**, 676–690 (1973).
- [6] R. L. Ake and M. Gouterman, "Porphyrins xiv. theory for the luminescent state in vo, co, cu complexes," *Theoretica chimica acta* **15**, 20–42 (1969).