

Tidal Love numbers for regular black holes

Rui Wang, Qi-Long Shi, Wei Xiong, and Peng-Cheng Li*

*School of Physics and Optoelectronics, South China University of Technology,
Guangzhou 510641, People's Republic of China*

(Dated: December 8, 2025)

Tidal Love numbers (TLNs) characterize the response of compact objects to external tidal fields and vanish for classical Schwarzschild and Kerr black holes in general relativity. Nonvanishing TLNs therefore provide a potential observational window into new physics. In this work, we present a unified and fully analytic study of the TLNs of three representative classes of regular black holes—the Bardeen black hole, the black hole with sub-Planckian curvature, and the black hole arising in asymptotically safe gravity—under scalar, vector, and axial gravitational perturbations. Employing a Green’s function method combined with systematic perturbative expansions, we show that TLNs of regular black holes are generically nonzero and exhibit strong model and mode dependence. In many cases, higher-order corrections develop logarithmic scale dependence, closely resembling renormalization-group running in quantum field theory and revealing a scale-dependent tidal response absent in classical black holes. Our analysis demonstrates that the internal structure of regular black holes, including de Sitter or Minkowski cores and quantum-gravity-inspired modifications, leaves distinct fingerprints in their tidal properties. These results establish TLNs as promising probes for testing regular black hole models with future gravitational-wave observations.

I. INTRODUCTION

General Relativity (GR) predicts the existence of both gravitational waves (GWs) and black holes (BHs). The direct detection of GWs from binary BH mergers has provided strong observational confirmation of these predictions [1]. During the early inspiral phase of a binary coalescence, the orbital separation gradually decreases, and tidal interactions—if present—become increasingly important. Such tidal interactions are characterized by the *tidal Love numbers* (TLNs), which quantify the response of a compact object to an external tidal field by relating the induced multipole moments to the applied tidal potential. The concept of Love numbers was first introduced by A. E. H. Love in 1909 [2] to describe the Earth’s deformation under lunar and solar tides. In relativistic astrophysics, TLNs have become essential observables linking the internal structure of compact objects to GW signals, and are now used to constrain neutron star equations of state [3, 4].

In GR, a remarkable property of four-dimensional, asymptotically flat BHs is the vanishing of their TLNs. This was first shown for Schwarzschild BHs [5–10], the same result was then shown to apply to Reissner-Nordström [11–13] (however, fermionic tidal perturbations may induce non-zero TLNs [14, 15]), Kerr [16–24] and Kerr-Newman [25] BHs. However, deviations from this prediction can arise in modified gravity theories [11, 26–29] or when BHs are surrounded by matter [30–33]. On the other hand, exotic compact objects, including BH mimickers, wormholes, and ultracompact stars, typically possess non-zero TLNs [11, 34–37]. The existence of nonvanishing TLNs in such cases could there-

fore serve as a smoking gun for new gravitational physics [11, 38–43]

Regular BHs (RBHs)—nonsingular solutions that avoid curvature divergences at the center—have been extensively explored as phenomenological models of quantum-corrected spacetimes [44, 45]. Examples include the Bardeen BH, which introduces a de Sitter-like core to remove singularities [46, 47]; the BH with sub-Planckian curvature, which maintains finite curvature for all masses [48, 49]; and BHs arising in asymptotically safe gravity (ASG) [50–53], which embody key aspects of quantum-gravity renormalization effects. Despite increasing interest, a systematic investigation of TLNs in these RBH scenarios remains incomplete [54–57]. Understanding how their internal geometries and quantum-inspired corrections influence tidal responses is therefore crucial for connecting theoretical models to future GW observations.

In this work, we perform a comprehensive analysis of the TLNs of three representative classes of RBHs: the Bardeen BH [46], the BH with sub-Planckian curvature [48], and the BH formed through gravitational dynamics during collapse within ASG [52], under scalar, vector and axial gravitational perturbations. To obtain these results analytically, we adopt a Green’s function technique developed in [28]. This method enables closed-form analytic solutions to the perturbed master equations and allows us to extract the full asymptotic structure of the tidal response. Our results reveal several notable features. First, TLNs of RBHs are generically nonzero, with their magnitude and sign depending sensitively on both the background model and the perturbation sector. Second, higher-order corrections frequently exhibit logarithmic scale dependence in the asymptotic region, leading to a “running” of the TLNs reminiscent of renormalization-group flow in quantum field theory. This scale dependence constitutes a qualitative dis-

* pchli2021@scut.edu.cn

inction from classical GR BHs and offers new insights into how quantum-gravity-motivated modifications of the near-horizon or interior geometry manifest in observable tidal properties. Third, different classes of RBHs display characteristic tidal signatures—such as the presence or absence of logarithmic running or the pattern of mode dependence—that may enable their discrimination in future GW observations.

The structure of this paper is as follows. Sec. II introduces the background metrics of the three representative RBH models. In Sec. III presents the formalism and Green’s function technique used to compute the TLNs.. Section IV discusses the obtained results for scalar, vector, and axial perturbations, highlighting their physical implications. Finally, Sec. V concludes with a summary of our findings and prospects for future GW tests of RBH models. We use geometric units $G = c = 1$ everywhere.

II. BACKGROUND OF REGULAR BLACK HOLES

In this section, we briefly review the three aforementioned models of RBHs, focusing on their metric representations and fundamental properties. The metrics for all three can be uniformly expressed in Schwarzschild coordinates as:

$$ds^2 = -f(r) dt^2 + \frac{dr^2}{f(r)} + r^2 d\Omega^2, \quad (1)$$

where $d\Omega^2 = d\theta^2 + \sin^2\theta d\phi^2$ is the metric on the unit two-sphere. The distinguishing feature of each specific RBH model is then given by its unique metric function $f(r)$.

A. Bardeen black holes

The Bardeen BH stands as the first RBH in GR, which was initially introduced as a phenomenological model [46]. Ayón-Beato and García provided a physical interpretation of the Bardeen BH by showing that it can be sourced by a magnetic monopole within a nonlinear electrodynamics framework [47]. In addition, quantum-gravity corrections derived from the generalized uncertainty principle have also been suggested as an alternative mechanism for generating the Bardeen BH [58].

The metric function of the Bardeen BH takes the form

$$f(r) = 1 - \frac{2Mr^2}{(r^2 + q^2)^{3/2}}, \quad (2)$$

where M is the mass of the BH and the parameter q acts as a regularization parameter. The solution continuously reduces to the Schwarzschild spacetime in the limit $q \rightarrow 0$. As shown in [47], the spacetime has horizons only when $q \leq \frac{4M}{3\sqrt{3}} \approx 0.77M$. For larger values of q , the spacetime has no horizons. A notable feature

of the Bardeen BH is that the central singularity is replaced by a regular core, which behaves effectively as a de Sitter region. This becomes clear from the small- r expansion, where $f(r) \rightarrow 1 - 2Mr^2/q^3$, as $r \rightarrow 0$. Dymnikova [59] showed that, within GR, any static spherically symmetric BH with a regular center and matter obeying the weak energy condition must approach a de Sitter core at the center. Well-known examples include the Hayward BH [60] and the Frolov BH [61], in addition to the Bardeen BH.

To study the gravitational perturbations of all three RBHs within a unified framework, we adopt the perspective that the regularity of the Bardeen BH arises from effective quantum corrections rather than from nonlinear electrodynamics, in analogy with the mechanisms underlying the other two RBH models.

B. Regular black holes with sub-Planckian curvature

Alternative to the Bardeen BH, Ref. [49] analyzed the general conditions required for obtaining a finite Kretschmann scalar and proposed a RBH solution featuring a Minkowski core. In this scenario, as $r \rightarrow 0$, the metric functions satisfy $-g_{tt} = g_{rr} \rightarrow 1$, so that the central singularity is replaced with a flat Minkowski spacetime instead of a de Sitter core. However, Ling and Wu [48] pointed out that in the RBH proposed in Ref. [49], the maximum value of the Kretschmann scalar depends on the BH mass. For a fixed short-distance regulator, the peak curvature scales as M^2 , meaning that it can grow arbitrarily large and exceed the Planck scale M_p^4 for sufficiently massive BHs. This behavior conflicts with the expectation from quantum gravity that space-time curvature should remain bounded.

Motivated by this issue, Ling and Wu [48] constructed a new class of RBHs with a Minkowski core, based on the generalized uncertainty principle in curved spacetime, in which the Kretschmann scalar remains sub-Planckian for any BH mass. At large radii, this metric can reproduce the asymptotic behavior of Bardeen, Hayward, or Frolov BHs, depending on the choice of the potential form. The metric function of the RBH with a Minkowski core and bounded curvature is

$$f(r) = 1 + 2\psi(r) = 1 - \frac{2M}{r} e^{-\alpha_0 M^x / r^c}, \quad (3)$$

where $\psi(r)$ denotes the modified Newton potential, M is the mass of the BH, and α_0 , x and c are dimensionless parameters. The parameters α_0 characterizes the deviation from the Newtonian potential and encodes quantum-gravity corrections. To ensure the existence of a horizon and keep the curvature below the Planck scale, the parameters must satisfy $c \geq x \geq c/3$ and $c \geq 2$. Typically, when $\alpha_0 = 0$, the metric reduces to the Schwarzschild solution. For $x = 2/3$ and $c = 2$, the metric matches the large- r asymptotics of the Bardeen BH, while choosing $x = 1$ and $c = 3$ yields the Hayward-like asymptotic

behavior. In this work, we focus on the case $x = 1$ and $c = 3$.

C. A regular black hole in asymptotically safe gravity

The Asymptotic Safety scenario provides a compelling candidate framework for quantum gravity, whose defining feature is the existence of a non-Gaussian ultraviolet fixed point of the renormalization group (RG) flow. In this approach, the scale dependence of Newton's constant is governed by RG trajectories that approach the UV fixed point at high energies, thereby preventing divergences in the gravitational coupling and ensuring the predictive power of the theory [62].

Recently, Bonanno et al. [52] extended the proposal of Markov and Mukhanov [63] and investigated the gravitational collapse of dust (a pressureless ideal fluid) within the framework of ASG. Guided by the Reuter fixed point [50], the authors introduced a multiplicative coupling between the matter sector and gravity, which effectively implements the weakening of the gravitational interaction at high energies. This modification leads to an effective Lagrangian that yields a regular interior geometry, which can be smoothly matched to a static and asymptotically flat exterior spacetime. The matching is uniquely fixed by requiring that both the induced metric and the extrinsic curvature remain continuous at the boundary of the collapsing matter distribution when approached from either side.

The resulting static, spherically symmetric exterior line element takes the form

$$f(r) = 1 - \frac{r^2}{3\xi} \log \left(1 + \frac{6M\xi}{r^3} \right), \quad (4)$$

where M denotes the BH mass and ξ is a scale parameter characterizing the deviation from the Schwarzschild solution. The precise value of ξ is currently unknown and should be constrained observationally. In the limit $\xi \rightarrow 0$, the Schwarzschild geometry is recovered. A numerical analysis of the horizon condition $f(r_h) = f'(r_h) = 0$, reveals a critical value $\xi_{\text{cr}} \simeq 0.4565 M^2$ with $r_h \simeq 1.2516 M$. For $0 < \xi < \xi_{\text{cr}}$, the metric exhibits two event horizons, an inner and an outer one. The two horizons merge into a single degenerate surface for $\xi = \xi_{\text{cr}}$, corresponding to an extremal BH. For $\xi > \xi_{\text{cr}}$, no horizon forms.

III. TIDAL LOVE NUMBERS USING THE GREEN'S FUNCTION TECHNIQUE

In this section, we briefly review the definition of TLNs in GR and revisit the procedure presented in Ref. [28], where the TLNs of static, spherically symmetric BHs were computed using the Green's function technique.

A. Definition of relativistic tidal Love numbers in GR

For a static, spherically symmetric metric of the form as Eq. (1), we can fix the BH horizon radius to $r_h = 1$. To achieve this, the original horizon radius r_h can be reparameterized as

$$r_h(\eta) = 2M + \sum_{n=1}^N \eta^n c_n, \quad (5)$$

where η denotes the parameter characterizing the deviation from the Schwarzschild BH and is expected to be very small. For instance, $\eta = q$ in the Bardeen BH case. Substituting this expansion into the metric function $f(\eta)$ and expanding it in powers of η ,

$$\sum_{k=0}^N \eta^k \frac{f^{(k)}(0)}{k!} = 0, \quad (6)$$

we can solve for the coefficients c_1, c_2, c_3, \dots order by order by comparing the coefficients of η at each expansion order. When the horizon is fixed to $r_h = 1$, the mass M cannot be an independent quantity and can be determined by $f(r_h) = 0$. In some modified metrics, M cannot be obtained in closed form, so we treat M itself as a power series in the deformation parameter η ,

$$M(\eta) = \sum_{n=0}^N M_n \eta^n. \quad (7)$$

Plugging this into the condition $r_h(\eta) = 1$ and expanding this equation in powers of η

$$\sum_{k=0}^N \eta^k \frac{r_h^{(k)}(0)}{k!} = 1, \quad (8)$$

we can sequentially solve for the mass expansion coefficients M_0, M_1, M_2, \dots . In the subsequent calculations, we only retain terms up to the third order, i.e., $N = 3$. This procedure ensures that the horizon is always normalized at $r_h = 1$ in different modified gravity backgrounds, thereby providing a consistent framework for subsequent perturbation equations and the computation of TLNs.

Assuming that the external tidal field is weak, the linear perturbation can be used to study the tidal response of a self-gravitating object. Due to the spherical symmetry of the Schwarzschild spacetime, massless perturbations of integer spin s admit complete separation of variables, leaving a single radial master function to be determined. In particular, the perturbations of scalar fields ($s = 0$), electromagnetic fields ($s = 1$), and the odd-parity (axial) gravitational perturbations ($s = 2$) can all be written in a Schrödinger-like Regge-Wheeler (RW)-type equation, each characterized by a different effective potential. In the gravitational case, the axial perturbations are governed by the original RW equation

[64]. In contrast, the even-parity (polar) gravitational perturbations of a Schwarzschild BH satisfy a distinct Schrödinger-like master equation, known as the Zerilli equation [65]. In the static limit, the RW equation reads

$$f_S(r) \frac{d}{dr} \left(f_S(r) \frac{d}{dr} \Psi(r) \right) - f_S(r) V_{\text{RW}}(r) \Psi(r) = 0, \quad (9)$$

where the effective potential is

$$V_{\text{RW}}(r) = \frac{l(l+1)}{r^2} + \frac{1-s^2}{r^3}, \quad (10)$$

the metric function for the Schwarzschild BH is $f_S(r) = 1 - 1/r$ and s denotes the spin of the perturbing fields. It is known that the general solution to Eq. (9) can be expressed in terms of hypergeometric functions, and it is straightforward to find the solution that is regular at the horizon. The asymptotic behavior of the solution at large r is generally given by [10]

$$\Psi(r) \sim r^{l+1} [1 + \mathcal{O}(r^{-1})] + K_l^s r^{-l} [1 + \mathcal{O}(r^{-1})], \quad (11)$$

where K_l^s is a constant that encodes the tidal response of the BH, which directly corresponds to the TLNs. This solution can be interpreted as consisting of two parts in the asymptotic region $r \rightarrow \infty$: the growing mode, which represents the external tidal field, and the decaying mode, which describes the induced response. It has been rigorously proven that the TLNs of the Schwarzschild BH is vanishing [5–10]. This finding has since been extended to Reissner-Nordström [11–13], Kerr [16–24] and Kerr-Newman [25] BHs.

The TLNs defined in Eq. (11) are derived from the asymptotic expansion of linear perturbations at large distances, where the expansion consists solely of power-law terms in r . These values are constant and hence independent of the measurement scale. However, in the presence of matter fields [30], or within alternative backgrounds or modified gravity theories [8, 11, 27], the perturbation fields at large distances do not necessarily exhibit a simple power-law series. Instead, the asymptotic behavior may involve logarithmic terms,

$$\begin{aligned} \Psi(r) \sim & r^{l+1} [1 + \mathcal{O}(r^{-1})] \\ & + K_l^s r^{-l} [\log r + \mathcal{O}(1)] [1 + \mathcal{O}(r^{-1})]. \end{aligned} \quad (12)$$

The coefficient of the logarithmic term can be interpreted as a beta function in the context of a classical renormalization flow [8, 10, 11].

B. The Green's function technique

In the following we give a brief review of the Green's function technique adopted in Ref.[28] to extract the TLNs of BHs in modified gravities. Assume that the static limit of the perturbation equations shares the form of the RW equation (9) but with distinct functions for the

effective potential $V(r)$ and the metric $f(r)$. Performing the following transformation

$$\Psi(r) = \frac{\Phi(r)}{\sqrt{Z(r)}}, \quad Z(r) = \frac{f(r)}{f_S(r)}, \quad (13)$$

the perturbation equation becomes

$$f_S(r) \frac{d}{dr} \left(f_S(r) \frac{d}{dr} \Phi(r) \right) - f_S(r) U(r) \Phi(r) = 0. \quad (14)$$

We can find that all the information of the modified spacetime is encoded only in a new effective potential

$$U(r) = \frac{V}{Z} - \frac{1}{4Z^2} \left[f_S \left(\frac{dZ}{dr} \right)^2 - 2Z \frac{d}{dr} \left(f_S \frac{dZ}{dr} \right) \right]. \quad (15)$$

In this work, we mainly consider the scalar, vector, and axial gravitational perturbations of RBHs. For the scalar and vector fields, the effective potentials are uniformly given by [66]

$$V(r) = \frac{l(l+1)}{r^2} + \frac{1-s^2}{r} f'(r). \quad (16)$$

The case of axial gravitational perturbations is more complicated than that of scalar and vector fields. This is primarily because the former involves perturbing the underlying equations of motion. However, since the metric (1) is introduced as a phenomenological model—rather than being derived from a known action or as an exact solution to specific field equations—the underlying fundamental theory remains unknown. To address this issue, we adopt the viewpoint that quantum or regularization corrections are effectively modeled as arising from the stress-energy tensor of an anisotropic fluid within the framework of classical Einstein gravity [67–69]. In this framework the axial perturbations are particularly simple, as they decouple from the matter perturbations. As a result, the perturbation equation is of the standard form given by (9), and the effective potential is expressed as [70]¹

$$V(r) = \frac{l(l+1)}{r^2} - \frac{f'(r)}{r} + f''(r). \quad (17)$$

Next, we consider the following expansions for the effective potential and the master function:

$$U(r) = \sum_{k \geq 0} \eta^k U^{(k)}(r), \quad \Phi(r) = \sum_{k \geq 0} \eta^k \Phi^{(k)}(r), \quad (18)$$

¹ It is important to note that the effective potential derived in [70] applies to an anisotropic fluid with an arbitrary non-zero radial pressure. In contrast, the effective potential in [71] was obtained specifically for the case of vanishing background radial pressure. In the limit of $p_r^{(0)} = 0$, the rr -component of the Einstein field equations yields the following relation for the metric gradient: $a'(r) = \frac{a(r)}{r} \frac{1-b(r)}{b(r)}$. Using this relation, Eq. (17) consistently reproduces the results presented in [71] when the limit $p_r^{(0)} = 0$ is taken.

where as before the parameter η represents the deviation from the Schwarzschild BH. Obviously, the zeroth-order potential is $U^{(0)}(r) = V_{\text{RW}}(r)$.

Our strategy is to solve equation (14) order by order in η by the Green's function method. At the leading order of η , we recover the standard RW equation.

$$\left[f_S(r) \frac{d}{dr} \left(f_S(r) \frac{d}{dr} \right) - f_S(r) U^{(0)}(r) \right] \Phi^{(0)}(r) = 0, \quad (19)$$

This equation has three regular singular points and can be easily transformed into the standard hypergeometric equation. The two linearly independent solutions to this equation are written in terms of the hypergeometric function as

$$\Phi_+^{(0)}(r) = r^{l+1} {}_2F_1 \left(-l-s, -l+s; -2l; \frac{1}{r} \right), \quad (20)$$

$$\Phi_-^{(0)}(r) = r^{-l} {}_2F_1 \left(l+1-s, l+1+s; 2l+2; \frac{1}{r} \right). \quad (21)$$

As analyzed in [28, 56], for $l \in \mathbb{C}$, both the solutions are divergent in the form $\propto \log(r-1)$ at the horizon. However, a unique linear combination of the two solutions exactly cancels the logarithmic terms at the horizon. As a result, the horizon-regular solution can be written as

$$\Phi^{(0)}(r) = \Phi_{\text{hor-reg}}^{(0)}(r) = \Phi_+^{(0)}(r) + K_l^{s(0)} \Phi_-^{(0)}(r), \quad (22)$$

Here, $K_l^{s(0)}$ is the constant encoding the tidal response at zero order, whose explicit expression is

$$K_l^{s(0)} = -\frac{\Gamma(-2l) \Gamma(l+1+s) \Gamma(l+1-s)}{\Gamma(-l-s) \Gamma(-l+s) \Gamma(2l+2)}. \quad (23)$$

This value is zero for a physical multipole $l \in \mathbb{Z}$, which corresponds to the fact that the TLNs are vanishing for four-dimensional BHs in GR.

Next, we consider the higher-order expansion of the perturbation equation (14). At k -th order, $\mathcal{O}(\eta^k)$, $k \geq 1$, the equation takes the form

$$\begin{aligned} f_S(r) \frac{d}{dr} \left(f_S(r) \frac{d}{dr} \Phi^{(k)}(r) \right) - f_S(r) U^{(0)}(r) \Phi^{(k)}(r) \\ = f_S(r) \mathcal{S}^{(k)}(r), \end{aligned} \quad (24)$$

where the source term $\mathcal{S}^{(k)}(r)$ is given by

$$\mathcal{S}^{(k)}(r) = \sum_{i=0}^{k-1} U^{(k-i)}(r) \Phi^{(i)}(r). \quad (25)$$

By observing that the derivative operator of the above inhomogeneous equation is the same as that of the homogeneous equation (19), the solution to the inhomogeneous equation can be obtained using the Green's function method. The Green's function $G(r, r')$ is defined by

$$\frac{d}{dr} \left(f_S(r) \frac{dG(r, r')}{dr} \right) - U^{(0)}(r) G(r, r') = \delta(r - r'), \quad (26)$$

with $\delta(r)$ being the delta function. Here, appropriate boundary conditions must be imposed. At the horizon $r = 1$, the field $\Phi(r)$ is required to be regular. As $r \rightarrow \infty$, we demand that $\Phi(r) = \mathcal{O}(r^l)$, which is justified by the fact that the full-order solution $\Phi(r)$ can always be renormalized such that the coefficient of the growing mode r^{l+1} is set to unity. Then the Green's function can be constructed with the two linearly independent solutions (20) and (21) of the corresponding homogeneous equation (19)

$$\begin{aligned} G(r, r') = -\frac{1}{2l+1} \left[\Phi_-^{(0)}(r) \Phi_{\text{hor-reg}}(r') \Theta(r - r') \right. \\ \left. + \Phi_{\text{hor-reg}}(r) \Phi_-^{(0)}(r') \Theta(r' - r) \right], \end{aligned} \quad (27)$$

where $\Theta(r)$ denotes the Heaviside step function, and the factor $2l+1$ arises from the Wronskian of the homogeneous solutions [28].

Using this Green's function, the k -th order solution of the inhomogeneous equation (24) can be written as

$$\begin{aligned} \Phi^{(k)}(r) &= \int_1^\infty G(r, r') \mathcal{S}^{(k)}(r') dr' \\ &= -\frac{1}{2l+1} \left[\Phi_-^{(0)}(r) \int_1^r \Phi_{\text{hor-reg}}(r') \mathcal{S}^{(k)}(r') dr' \right. \\ &\quad \left. + \Phi_{\text{hor-reg}}(r) \int_r^\infty \Phi_-^{(0)}(r') \mathcal{S}^{(k)}(r') dr' \right]. \end{aligned} \quad (28)$$

With this solution, the TLNs can be determined by the coefficient of the decaying mode r^{-l} at large r . As pointed out in [28], for $l \in \mathbb{Z}$, only the constant part of the integral contributes to this term in the $r \rightarrow \infty$ limit. Accordingly, the TLNs are identified from the constant term in the following expression:

$$I[\mathcal{S}^{(k)}(r)] = -\frac{1}{2l+1} \int_1^r \Phi_+^{(0)}(r') \mathcal{S}^{(k)}(r') dr'. \quad (29)$$

We emphasize that when logarithmic behavior appears in the asymptotic expansion, the logarithmic terms are also regarded as part of the TLNs.

IV. TIDAL LOVE NUMBERS FOR REGULAR BLACK HOLES

In this section, we investigate the linear static response of the three types of RBHs introduced in Sec. II under the influence of external tidal fields (scalar, vector, and axial gravitational types). By employing the Green's function method introduced in Sec. III, we can extract the TLNs.

A. Bardeen black hole

Here, we extract the response of the Bardeen BH under scalar, vector, and axial gravitational perturbations.

Since we set the horizon radius to $r_h = 1$, the mass parameter can be expanded as

$$M(q) = \frac{1}{2} + \frac{3}{4}q^2 + \frac{3}{16}q^4 - \frac{1}{32}q^6 + \mathcal{O}(q^8). \quad (30)$$

Correspondingly, the metric function $f(r)$ admits a series expansion in powers of q given by

$$f(r) = 1 - \frac{1}{r} - \frac{3q^2(r^2 - 1)}{2r^3} - 2q^4 \left(\frac{3}{16r} - \frac{9}{8r^3} + \frac{15}{16r^5} \right) - 2q^6 \left(\frac{1}{32r} + \frac{9}{32r^3} - \frac{45}{32r^5} + \frac{35}{32r^7} \right) + \mathcal{O}(q^8). \quad (31)$$

Plugging this into Eq.(16) and then Eq. (15), and performing the expansion in powers of q . The power-series expansion of the effective potential for the scalar ($s = 0$) and vector ($s = 1$) perturbations can be collectively expressed as

$$U_1(r) = \frac{-3 + 6l(l+1) - 6s^2}{4r^4} + \frac{6l(l+1) - 6s^2}{4r^3} + \frac{12 + 12s^2}{4r^5}, \quad (32)$$

$$U_2(r) = \frac{3l(l+1) - 3s^2}{8r^3} + \frac{-6 + 21l(l+1) - 21s^2}{8r^4} + \frac{57 + 42l(l+1) + 30s^2}{16r^5} + \frac{99 + 6l(l+1) + 66s^2}{16r^6} - \frac{33 + 12s^2}{4r^7}, \quad (33)$$

$$U_3(r) = -\frac{l(l+1) - s^2}{16r^3} + \frac{-4 + 17l(l+1) - 17s^2}{16r^4} + \frac{-43 + 160l(l+1) - 124s^2}{32r^5} + \frac{219 + 160l(l+1) + 92s^2}{32r^6} + \frac{-41 + 34l(l+1) + 74s^2}{32r^7} - \frac{391 + 2l(l+1) + 106s^2}{32r^8} + \frac{27 + 6s^2}{2r^9}. \quad (34)$$

For the axial gravitational perturbation ($s = 2$), plugging (31) into Eq. (17) and then Eq. (15), and performing the expansion in powers of q , one obtains the effective-potential expansion coefficients

$$U_1(r) = \frac{-9 + 3l(l+1)}{2r^3} + \frac{-21 + 6l(l+1)}{4r^4} + \frac{21}{r^5}, \quad (35)$$

$$U_2(r) = \frac{-9 + 3l(l+1)}{8r^3} + \frac{-69 + 21l(l+1)}{8r^4} + \frac{363 + 42l(l+1)}{16r^5} + \frac{513 + 6l(l+1)}{16r^6} - \frac{165}{4r^7}, \quad (36)$$

$$U_3(r) = -\frac{-3 + l + l^2}{16r^3} + \frac{-55 + 17l + 17l^2}{16r^4} + \frac{-307 + 160l(1+l)}{32r^5} + \frac{1251 + 160l(1+l)}{32r^6} + \frac{-215 + 34l(1+l)}{32r^7} - \frac{1753 + 2l(1+l)}{32r^8} + \frac{123}{2r^9}. \quad (37)$$

1. Scalar field response

In this section, we compute the TLN of Bardeen BH under scalar perturbation. The lowest multipole number starts from $l = 0$. For $l = 0$, the zeroth-order $\mathcal{O}(q^0)$ growing and decaying modes are given by Eqs. (20) and (21), and take the following form:

$$\Phi_+^{(0)}(r) = \Phi^{(0)}(r) = \Phi_{\text{hor-reg}}(r) = r, \quad (38)$$

$$\Phi_-^{(0)}(r) = -r \log\left(1 - \frac{1}{r}\right). \quad (39)$$

We should point out that the first equality in the first equation follows from Eq. (22), which relies on the fact that K_l^s vanishes for integer multipole number l in the case of the classical Schwarzschild BH. For $s = 0$ and $l = 0$, the source terms at first order $\mathcal{O}(q^2)$ can be obtained from Eq. (25), yielding

$$S^{(1)}(r) = U^{(1)}(r) \Phi^{(0)}(r) = \frac{3}{r^4} - \frac{3}{4r^3}. \quad (40)$$

Furthermore, the first-order solution and the second-order source term at $\mathcal{O}(q^4)$ can be written as follows:

$$\Phi^{(1)}(r) = \int_1^\infty G(r, r') S^{(1)}(r') dr' = -\frac{3}{4} - \frac{3}{4r}, \quad (41)$$

$$S^{(2)}(r) = U^{(2)}(r) \Phi^{(0)}(r) + U^{(1)}(r) \Phi^{(1)}(r) = -\frac{21}{2r^6} + \frac{9}{2r^5} + \frac{33}{8r^4} - \frac{3}{4r^3}. \quad (42)$$

Moreover, the second-order solution and the third-order source term at $\mathcal{O}(q^6)$, following Eqs. (28) and (25), can be computed as follows:

$$\Phi^{(2)}(r) = \int_1^\infty G(r, r') S^{(2)}(r') dr' = -\frac{3}{16} + \frac{21}{32r^3} + \frac{3}{8r^2} - \frac{15}{32r}, \quad (43)$$

$$S^{(3)}(r) = U^{(3)}(r) \Phi^{(0)}(r) + U^{(2)}(r) \Phi^{(1)}(r) + U^{(1)}(r) \Phi^{(2)}(r) = \frac{693}{32r^8} - \frac{1285}{128r^7} - \frac{329}{32r^6} + \frac{579}{128r^5} - \frac{41}{64r^4} - \frac{1}{4r^3}. \quad (44)$$

For $l = 0$, the TLN is obtained from Eq. (29), which expanded in q^2 gives

$$I[\mathcal{S}^{(1)}](r) = -\frac{3}{4} + \frac{3}{2r^2} - \frac{3}{4r}, \quad (45)$$

$$I[\mathcal{S}^{(2)}](r) = -\frac{3}{16} - \frac{21}{8r^4} + \frac{3}{2r^3} + \frac{33}{16r^2} - \frac{3}{4r}, \quad (46)$$

$$I[\mathcal{S}^{(3)}](r) = \frac{1}{32} + \frac{231}{64r^6} - \frac{257}{128r^5} - \frac{329}{128r^4} + \frac{193}{128r^3} - \frac{41}{128r^2} - \frac{1}{4r}. \quad (47)$$

We find that, in the case of the scalar perturbation with $l = 0$, there exist non-vanishing TLNs and no logarithmic terms appear. Moreover, we find that the first- and second-order terms provide negative contributions to the TLNs, while the third-order term gives a positive contribution. Up to $\mathcal{O}(q^8)$, the TLN is given by

$$K_{l=0}^0 = -\frac{3}{4}q^2 - \frac{3}{16}q^4 + \frac{1}{32}q^6 + \mathcal{O}(q^8). \quad (48)$$

For $l = 1$, The zeroth-order solutions (20) and (21) take the following form:

$$\Phi_+^{(0)}(r) = \Phi^{(0)}(r) = \Phi_{\text{hor-reg}}(r) = -\frac{r}{2} + r^2, \quad (49)$$

$$\Phi_-^{(0)}(r) = -12r + 6r \log\left(1 - \frac{1}{r}\right) - 12r^2 \log\left(1 - \frac{1}{r}\right). \quad (50)$$

the corresponding integrals Eq. (29) expanded in q^2 are given by

$$I[\mathcal{S}^{(1)}](r) = \frac{15}{16} + \frac{1}{8r^2} - \frac{13}{16r} + \frac{r}{4} - \frac{r^2}{2} - \frac{\log r}{2}, \quad (51)$$

$$\begin{aligned} I[\mathcal{S}^{(2)}](r) = & -\frac{255}{64} - \frac{15\pi^2}{16} - \frac{7}{32r^4} + \frac{3}{2r^3} - \frac{305}{64r^2} \\ & - \frac{\pi^2}{8r^2} + \frac{331}{32r} + \frac{13\pi^2}{16r} - \frac{11r}{4} - \frac{\pi^2 r}{4} - \frac{r^2}{8} \\ & + \frac{\pi^2 r^2}{2} - \frac{3 \log r}{4} + \pi^2 \log r - \frac{15 \log r}{8r^2} + \frac{63 \log r}{8r} \\ & - 3r \log r - \frac{45 \log^2 r}{16} - \frac{3 \log^2 r}{8r^2} + \frac{39 \log^2 r}{16r} \\ & - \frac{3}{4}r \log^2 r + \frac{3}{2}r^2 \log^2 r + \frac{\log^3 r}{2} - \frac{45}{8} \text{Li}_2(1-r) \\ & - \frac{3 \text{Li}_2(1-r)}{4r^2} + \frac{39 \text{Li}_2(1-r)}{8r} - \frac{3}{2}r \text{Li}_2(1-r) \\ & + 3r^2 \text{Li}_2(1-r) + 3 \log r \text{Li}_2(r) - 6 \text{Li}_3(r) \\ & + 6\zeta(3). \end{aligned} \quad (52)$$

We can see that for $l = 1$, the TLNs is non-zero and logarithmic terms appear. In particular, the $I[\mathcal{S}^{(1)}]$ part is simple enough to extract TLN explicitly: the corresponding coefficient reads $K_{l=1}^{0(1)} = 15/16 - 1/2 \log r$. However, the $I[\mathcal{S}^{(2)}]$ part is more complicated. This complexity

arises from the multiple integrations of logarithmic terms during the computation. Such integrations unavoidably produce the polylogarithm function. Nevertheless, we can extract the constant and $\log r$ terms from $I[\mathcal{S}^{(2)}](r)$ at large r . The TLN at the second order of q^2 is then obtained as $K_{l=1}^{0(2)} = -303/64 + 6\zeta(3) - 3/4 \log r$, which is positive and has a negative logarithmic running at the second order. Note that we compute the TLN only up to second order, since the integration (28) cannot be performed analytically further, and consequently we cannot construct the third-order source term. Up to $\mathcal{O}(q^6)$, the TLN is given by

$$K_{l=1}^0 = \left(\frac{15}{16} - \frac{1}{2} \log r\right) q^2 + \left(-\frac{303}{64} + 6\zeta(3) - \frac{3}{4} \log r\right) q^4 + \mathcal{O}(q^6). \quad (53)$$

2. Vector field response

Now we compute the TLNs of the Bardeen BH under vector perturbation. Similar to the steps present in the previous subsection, for $s = 1$ and $l = 1$, the TLNs can be obtained from the following expressions:

$$I[\mathcal{S}^{(1)}](r) = \frac{1}{2} - \frac{r}{4} - \frac{r^2}{4} - 2 \log r, \quad (54)$$

$$\begin{aligned} I[\mathcal{S}^{(2)}](r) = & -\frac{411}{16} - \frac{\pi^2}{2} + \frac{3}{8r^2} + \frac{217}{8r} - \frac{7r}{4} + \frac{\pi^2 r}{4} \\ & - \frac{r^2}{16} + \frac{\pi^2 r^2}{4} - \frac{21 \log r}{4} + 4\pi^2 \log r + \frac{3 \log r}{r^2} \\ & + \frac{51 \log r}{4r} - \frac{3}{2}r \log r - \frac{3 \log^2 r}{2} + \frac{3}{4}r \log^2 r \\ & + \frac{3}{4}r^2 \log^2 r + 2 \log^2 r - 3 \text{Li}_2(1-r) + \frac{3}{2}r \text{Li}_2(1-r) \\ & + \frac{3}{2}r^2 \text{Li}_2(1-r) + 12 \log r \text{Li}_2(r) - 24 \text{Li}_3(r) \\ & + 24\zeta(3). \end{aligned} \quad (55)$$

From $I[\mathcal{S}^{(1)}](r)$ we obtain the first-order vector TLN $K_1^{1(1)} = 1/2 - 2 \log r$, which is positive and has a negative logarithmic running. From the asymptotic behavior of $I[\mathcal{S}^{(1)}](r)$ at large r , the second-order vector TLN is obtained as $K_1^{1(2)} = -381/16 + 24\zeta(3) - 3 \log r$, which is negative plus a positive logarithmic running. Up to $\mathcal{O}(q^6)$, the vector TLN for $l = 1$ is given by

$$K_{l=1}^1 = \left(\frac{1}{2} - 2 \log r\right) q^2 + \left(-\frac{381}{16} + 24\zeta(3) - 3 \log r\right) q^4 + \mathcal{O}(q^6). \quad (56)$$

Finally, for $l = 2$, the integration (29) for vector per-

turbation at each order of q becomes

$$I[\mathcal{S}^{(1)}](r) = -\frac{153}{160} + \frac{333r}{320} - \frac{3r^2}{320} + \frac{3r^3}{10} - \frac{3r^4}{8} - \frac{27 \log r}{40}, \quad (57)$$

$$\begin{aligned} I[\mathcal{S}^{(2)}](r) = & -\frac{9597}{256} + \frac{1377\pi^2}{160} + \frac{1161}{640r^2} + \frac{25713}{640r} + \frac{2829r}{640} \\ & - \frac{2997\pi^2 r}{320} + \frac{2847r^2}{256} + \frac{27\pi^2 r^2}{320} - \frac{399r^3}{20} \\ & - \frac{3r^4}{32} - \frac{27\pi^2 r^3}{10} + \frac{27\pi^2 r^4}{8} + \frac{18117 \log r}{320} \\ & + \frac{243\pi^2 \log r}{20} + \frac{81 \log r}{80r^2} + \frac{3753 \log r}{320r} + \frac{27r \log r}{32} \\ & + \frac{243r^2 \log r}{40} - \frac{81r^3 \log r}{4} + \frac{4131 \log^2 r}{160} \\ & - \frac{8991r \log^2 r}{320} + \frac{81r^2 \log^2 r}{320} - \frac{81r^3 \log^2 r}{10} \\ & + \frac{81r^4 \log^2 r}{8} + \frac{243 \log^3 r}{40} + \frac{4131}{80} \text{Li}_2(1-r) \\ & + \frac{729\zeta(3)}{10} - \frac{8991r}{160} \text{Li}_2(1-r) + \frac{81r^2}{160} \text{Li}_2(1-r) \\ & - \frac{81r^3}{5} \text{Li}_2(1-r) + \frac{81r^4}{4} \text{Li}_2(1-r) \\ & + \frac{729}{20} \log r \text{Li}_2(r) - \frac{729}{10} \text{Li}_3(r). \end{aligned} \quad (58)$$

The first-order correction of the vector TLN directly reads $K_2^{(1)} = -153/160 - 27/40 \log r$. The second-order correction of the vector TLN is obtained from the asymptotic behavior of $I[\mathcal{S}^{(1)}](r)$ at large r , which is given by $K_2^{(2)} = -24087/256 + 729/10 \zeta(3) + 27/80 \log r$. Up to $\mathcal{O}(q^6)$, the vector TLN for $l = 2$ turns out to be

$$\begin{aligned} K_{l=2}^1 = & \left(-\frac{153}{160} - \frac{27}{40} \log r \right) q^2 \\ & + \left(-\frac{24087}{256} + \frac{729}{10} \zeta(3) + \frac{27}{80} \log r \right) q^4 + \mathcal{O}(q^6). \end{aligned} \quad (59)$$

3. Axial gravitational field response

Here, we compute the TLNs of the Bardeen BH under axial gravitational perturbations for $l = 2$ and $l = 3$. We carry the calculations up to second order in q^4 . For $l = 2$ the expansion of the integration (29) in q^4 reads

$$I[\mathcal{S}^{(1)}](r) = \frac{103}{40} - \frac{21r^2}{10} - \frac{r^3}{4} - \frac{9r^4}{40}, \quad (60)$$

$$\begin{aligned} I[\mathcal{S}^{(2)}](r) = & \frac{3541}{128} - \frac{2163}{320r^2} - \frac{13081}{640r} + \frac{837r}{320} - \frac{231r^2}{80} \\ & - \frac{11r^3}{80} - \frac{9r^4}{160} + \frac{6193}{320} \log r. \end{aligned} \quad (61)$$

From these results we see that the first-order TLN is non-vanishing and contains no logarithmic running, whereas

logarithmic running appears at second order through the $\log r$ term in $I[\mathcal{S}^{(2)}](r)$. In this case, all the correction terms contribute positively to the TLN. The TLN up to $\mathcal{O}(q^6)$ can be expressed as

$$K_{l=2}^2 = \frac{103}{40} q^2 + \left(\frac{3541}{128} + \frac{6193}{320} \log r \right) q^4 + \mathcal{O}(q^6). \quad (62)$$

For the case of $l = 3$, we obtain the following results:

$$\begin{aligned} I[\mathcal{S}^{(1)}](r) = & \frac{1667}{10080} - \frac{25r^2}{24} + \frac{1255r^3}{1008} - \frac{73r^4}{224} + \frac{39r^5}{140} \\ & - \frac{9r^6}{28}, \end{aligned} \quad (63)$$

$$\begin{aligned} I[\mathcal{S}^{(2)}](r) = & \frac{1934113}{161280} - \frac{1667}{3840r^2} - \frac{1918717}{806400r} - \frac{600521r}{44800} \\ & + \frac{2393r^2}{2688} + \frac{70667r^3}{20160} - \frac{2081r^4}{4480} + \frac{3r^5}{8} \\ & - \frac{9r^6}{112} + \frac{18889 \log r}{403200}. \end{aligned} \quad (64)$$

Similarly, the TLN for $l = 3$ up to $\mathcal{O}(q^6)$ turns out to be

$$K_{l=3}^2 = \frac{1667}{10080} q^2 + \left(\frac{1934113}{161280} + \frac{18889}{403200} \log r \right) q^4 + \mathcal{O}(q^6). \quad (65)$$

The behavior of TLN for $l = 3$ closely resembles that of the $l = 2$ case: the first-order term contributes a finite and non-zero TLNs without logarithmic running, while the second-order correction introduces a logarithmic dependence in r . Both cases contribute positively to the TLNs and the running.

B. Regular black holes with sub-Planckian curvature

In this subsection, we compute the TLNs of the RBH with sub-Planckian curvature [48]. Without loss of generality, we choose the parameters $x = 1$ and $c = 3$. Similar to the Bardeen BH case, we consider the response under scalar, vector, and axial gravitational perturbations. Setting the horizon radius $r_h = 1$, the mass can be expanded according to Eq. (7) as

$$M(\alpha_0) = \frac{1}{2} + \frac{\alpha_0}{4} + \frac{3\alpha_0^2}{16} + \frac{\alpha_0^3}{6} + \mathcal{O}(\alpha_0^4). \quad (66)$$

Up to the same order in the regular parameter α_0 , the metric function can be expanded as

$$\begin{aligned} f(r) = & 1 - \frac{1}{r} + \frac{(1-r^3)\alpha_0}{2r^4} + \frac{(-1+4r^3-3r^6)\alpha_0^2}{8r^7} \\ & + \frac{(1-9r^3+24r^6-16r^9)\alpha_0^3}{48r^{10}} + \mathcal{O}(\alpha_0^4). \end{aligned} \quad (67)$$

The effective potentials (16) for scalar ($s = 0$) and vector ($s = 1$) perturbations can be expanded in the

same way as follows:

$$U_1(r) = \frac{2l(l+1) - 2s^2}{4r^3} + \frac{-1 + 2l(l+1) - 2s^2}{4r^4} + \frac{-2 + 2l(l+1) - 2s^2}{4r^5} + \frac{9 + 6s^2}{4r^6}, \quad (68)$$

$$U_2(r) = \frac{6l(l+1) - 6s^2}{16r^3} + \frac{-4 + 10l(l+1) - 10s^2}{16r^4} + \frac{-11 + 14l(l+1) - 14s^2}{16r^5} + \frac{27 + 10l(l+1) + 14s^2}{16r^6} + \frac{14 + 6l(l+1) + 6s^2}{16r^7} + \frac{22 + 2l(l+1) + 10s^2}{16r^8} - \frac{9}{16r^9}, \quad (69)$$

$$U_3(r) = \frac{32l(l+1) - 32s^2}{96r^3} + \frac{-25 + 68l(l+1) - 68s^2}{96r^4} + \frac{-83 + 116l(l+1) - 116s^2}{96r^5} + \frac{51 + 64l(l+1) + 8s^2}{48r^6} + \frac{86 + 116l(l+1) + 10s^2}{96r^7} + \frac{217 + 80l(l+1) + 82s^2}{96r^8} + \frac{30 + 19l(l+1) + 26s^2}{96r^9} + \frac{83 + 14l(l+1) + 40s^2}{96r^{10}} + \frac{19 + 2l(l+1) + 16s^2}{96r^{11}}. \quad (70)$$

The expansion of the effective potential (16) for axial gravitational perturbations ($s = 2$) is given as follows:

$$U_1(r) = \frac{-3 + l(l+1)}{2r^3} + \frac{-7 + 2l(l+1)}{4r^4} + \frac{-4 + l(l+1)}{2r^5} + \frac{51}{4r^6}, \quad (71)$$

$$U_2(r) = \frac{-9 + 3l(l+1)}{8r^3} + \frac{-17 + 5l(l+1)}{8r^4} + \frac{-53 + 14l(l+1)}{16r^5} + \frac{165 + 10l(l+1)}{16r^6} + \frac{40 + 3l(l+1)}{8r^7} + \frac{50 + l(l+1)}{8r^8} - \frac{45}{16r^9}, \quad (72)$$

$$U_3(r) = \frac{-3 + l(l+1)}{3r^3} + \frac{-229 + 68l(l+1)}{96r^4} + \frac{-431 + 116l(l+1)}{96r^5} + \frac{363 + 64l(l+1)}{48r^6} + \frac{155 + 29l(l+1)}{24r^7} + \frac{1111 + 80l(l+1)}{96r^8} + \frac{126 + 19l(l+1)}{48r^9} + \frac{311 + 14l(l+1)}{96r^{10}} + \frac{31 + 2l(l+1)}{96r^{11}}. \quad (73)$$

1. Scalar field response

First of all, we compute the TLNs for the RBH with sub-Planckian curvature under scalar field perturbations. For the case $l = 0$, The zeroth-order ($\mathcal{O}(\alpha_0^0)$) growing and decaying modes, corresponding to Eqs. (20) and (21), are given explicitly by

$$\Phi_+^{(0)}(r) = \Phi^{(0)}(r) = \Phi_{\text{hor-reg}}(r) = r, \quad (74)$$

$$\Phi_-^{(0)}(r) = -r \log\left(1 - \frac{1}{r}\right). \quad (75)$$

In the case $s = 0$ and $l = 0$, the first-order source term at $\mathcal{O}(\alpha_0^1)$ is derived from Eq. (25) as

$$S^{(1)}(r) = U^{(1)}(r) \Phi^{(0)}(r) = \frac{9}{4r^5} - \frac{1}{2r^4} - \frac{1}{4r^3}. \quad (76)$$

Applying the Green's function formalism, the first-order solution and the second-order source term ($\mathcal{O}(\alpha_0^2)$) are expressed as

$$\Phi^{(1)}(r) = \int_1^\infty G(r, r') S^{(1)}(r') dr' = -\frac{1}{4} - \frac{1}{4r^2} - \frac{1}{4r}, \quad (77)$$

$$S^{(2)}(r) = U^{(2)}(r) \Phi^{(0)}(r) + U^{(1)}(r) \Phi^{(1)}(r) = -\frac{9}{8r^8} + \frac{15}{16r^7} + \frac{1}{2r^6} + \frac{15}{8r^5} - \frac{5}{8r^4} - \frac{1}{4r^3}. \quad (78)$$

Similarly, the second-order solution together with the third-order source term at $\mathcal{O}(\alpha_0^3)$, following Eqs. (28) and (25), are given by

$$\Phi^{(2)}(r) = \int_1^\infty G(r, r') S^{(2)}(r') dr' = -\frac{3}{16} + \frac{1}{32r^5} - \frac{1}{32r^3} - \frac{1}{4r^2} - \frac{7}{32r}, \quad (79)$$

$$S^{(3)}(r) = U^{(3)}(r) \Phi^{(0)}(r) + U^{(2)}(r) \Phi^{(1)}(r) + U^{(1)}(r) \Phi^{(2)}(r) = \frac{27}{128r^{11}} - \frac{1}{48r^{10}} + \frac{35}{96r^9} - \frac{29}{32r^8} + \frac{275}{192r^7} + \frac{11}{24r^6} + \frac{185}{128r^5} - \frac{145}{192r^4} - \frac{25}{96r^3}. \quad (80)$$

For $l = 0$, the integral (29) is expanded in α_0 as

$$I[S^{(1)}](r) = -\frac{1}{4} + \frac{3}{4r^3} - \frac{1}{4r^2} - \frac{1}{4r}, \quad (81)$$

$$I[S^{(2)}](r) = -\frac{3}{16} - \frac{3}{16r^6} + \frac{3}{16r^5} + \frac{1}{8r^4} + \frac{5}{8r^3} - \frac{5}{16r^2} - \frac{1}{4r}, \quad (82)$$

$$I[S^{(3)}](r) = -\frac{1}{6} + \frac{3}{128r^9} - \frac{1}{384r^8} + \frac{5}{96r^7} - \frac{29}{192r^6} + \frac{55}{192r^5} + \frac{11}{96r^4} + \frac{185}{384r^3} - \frac{145}{384r^2} - \frac{25}{96r}. \quad (83)$$

From the above results, it can be seen that, similar to the Bardeen BH, the case with $l = 0$ exhibits non-vanishing TLN, and no logarithmic term appears. The first-, second- and third-order corrections all contribute negatively to the TLNs. Since we have chosen the parameters $x = 1$ and $c = 3$, this type of BH shows the same asymptotic behavior as the Hayward BH at large r . In this case, the scalar TLNs for $l = 0$ can be written as

$$K_{l=0}^0 = -\frac{1}{4}\alpha_0 - \frac{3}{16}\alpha_0^2 - \frac{1}{6}\alpha_0^3 + \mathcal{O}(\alpha_0^4). \quad (84)$$

For $l = 1$, the TLNs are computed up to third order of α_0 . The zeroth-order solutions (20) and (21) take the following form:

$$\Phi_+^{(0)}(r) = \Phi^{(0)}(r) = \Phi_{\text{hor-reg}}(r) = -\frac{r}{2} + r^2, \quad (85)$$

$$\Phi_-^{(0)}(r) = -12r + 6r \log\left(1 - \frac{1}{r}\right) - 12r^2 \log\left(1 - \frac{1}{r}\right). \quad (86)$$

The TLNs can be obtained from the following expressions:

$$I[\mathcal{S}^{(1)}](r) = -\frac{13}{48} + \frac{1}{16r^3} - \frac{17}{48r^2} + \frac{31}{48r} + \frac{r}{12} - \frac{r^2}{6}, \quad (87)$$

$$I[\mathcal{S}^{(2)}](r) = -\frac{35}{192} - \frac{1}{64r^6} + \frac{7}{64r^5} - \frac{19}{96r^4} + \frac{11}{96r^3} - \frac{65}{192r^2} + \frac{53}{96r} + \frac{r}{12} - \frac{r^2}{8}, \quad (88)$$

$$I[\mathcal{S}^{(3)}](r) = -\frac{289}{2016} + \frac{1}{512r^9} - \frac{313}{23040r^8} + \frac{5459}{161280r^7} - \frac{1181}{23040r^6} + \frac{3373}{23040r^5} - \frac{6209}{23040r^4} + \frac{3583}{23040r^3} - \frac{1591}{4608r^2} + \frac{587}{1152r} + \frac{25r}{288} - \frac{r^2}{9}. \quad (89)$$

Here, we can see that for the scalar perturbations with $l = 1$, the TLNs are non-zero and no logarithmic term appears. The contributions to the TLNs exhibit negative behavior similar to the case of $l = 0$. This is in contrast to the Bardeen BH, whose TLN is positive and has a negative logarithmic running for $l = 1$. All contributions up to the third order can be accurately expressed as

$$K_{l=1}^0 = -\frac{13}{48}\alpha_0 - \frac{35}{192}\alpha_0^2 - \frac{289}{2016}\alpha_0^3 + \mathcal{O}(\alpha_0^4). \quad (90)$$

2. Vector field response

Next, we compute the TLNs for vector perturbations ($s = 1$). Similarly, for the case $l = 1$, the TLNs can be

obtained from the following integrations

$$I[\mathcal{S}^{(1)}](r) = -\frac{13}{12} + \frac{5}{4r} - \frac{r}{12} - \frac{r^2}{12}, \quad (91)$$

$$I[\mathcal{S}^{(2)}](r) = -\frac{35}{48} - \frac{7}{16r^4} + \frac{7}{48r^3} + \frac{1}{8r^2} + \frac{25}{24r} - \frac{r}{12} - \frac{r^2}{16}, \quad (92)$$

$$I[\mathcal{S}^{(3)}](r) = -\frac{4687}{8064} + \frac{65}{896r^7} - \frac{103}{1152r^6} - \frac{317}{5760r^5} - \frac{3359}{5760r^4} + \frac{1531}{5760r^3} + \frac{209}{1152r^2} + \frac{1073}{1152r} - \frac{25r}{288} - \frac{r^2}{18}. \quad (93)$$

Similar to scalar TLNs, we can clearly see that the TLNs are non-zero and that no logarithmic running appears and the contributions are also negative. The contributions to the TLN up to third order can be explicitly written as

$$K_{l=1}^1 = -\frac{13}{12}\alpha_0 - \frac{35}{48}\alpha_0^2 - \frac{4687}{8064}\alpha_0^3 + \mathcal{O}(\alpha_0^4). \quad (94)$$

For $l = 2$, we compute the TLN only up to second order of α_0 , as the integral (28) cannot be performed analytically, so we cannot construct the third-order source term. The explicit TLNs for the vector perturbations can be extracted from the following expressions:

$$\begin{aligned} I[\mathcal{S}^{(1)}](r) &= \frac{3}{320} + \frac{27}{64r} - \frac{129r}{320} - \frac{r^2}{320} + \frac{r^3}{10} - \frac{r^4}{8} \\ &\quad + \frac{9 \log r}{10}, \quad (95) \\ I[\mathcal{S}^{(2)}](r) &= -\frac{32641}{256} + \frac{9\pi^2}{80} - \frac{189}{1280r^4} - \frac{1161}{1280r^3} - \frac{1449}{128r^2} \\ &\quad + \frac{21889}{160r} + \frac{81\pi^2}{16r} - \frac{627r}{640} - \frac{387\pi^2 r}{80} - \frac{6341r^2}{1280} \\ &\quad - \frac{3\pi^2 r^2}{80} + \frac{91r^3}{10} + \frac{6\pi^2 r^3}{5} - \frac{3r^4}{32} - \frac{3\pi^2 r^4}{2} \\ &\quad + \frac{2427 \log r}{80} + \frac{108\pi^2 \log r}{5} - \frac{9 \log r}{16r^3} - \frac{351 \log r}{80r^2} \\ &\quad + \frac{6333 \log r}{80r} - \frac{3r \log r}{8} - \frac{27r^2 \log r}{10} + 9r^3 \log r \\ &\quad + \frac{27 \log^2 r}{80} + \frac{243 \log^2 r}{16r} - \frac{1161r \log^2 r}{80} \\ &\quad - \frac{9r^2 \log^2 r}{80} + \frac{18r^3 \log^2 r}{5} - \frac{9r^4 \log^2 r}{2} + \frac{54 \log^3 r}{5} \\ &\quad + \frac{27}{40} \text{Li}_2(1-r) + \frac{243 \text{Li}_2(1-r)}{8r} - \frac{1161r \text{Li}_2(1-r)}{40} \\ &\quad - \frac{9r^2 \text{Li}_2(1-r)}{40} + \frac{36r^3 \text{Li}_2(1-r)}{5} - 9r^4 \text{Li}_2(1-r) \\ &\quad + \frac{324 \log r \text{Li}_2(r)}{5} - \frac{648 \text{Li}_3(r)}{5} + \frac{648\zeta(3)}{5}. \quad (96) \end{aligned}$$

Here, the explicit form of the first-order TLN is $K_{l=1}^{1(1)} = 3/320 - 9/10 \log r$. The second-order TLN is obtained

from the asymptotic behavior of $I[\mathcal{S}^{(2)}](r)$ at large r , which is given by $K_{l=1}^{1(2)} = -40025/256 + 648/5 \zeta(3) + 27/20 \log r$. We observe that at first order $K_{l=1}^{1(1)}$ is positive while its running is negative. In contrast, at second order the coefficient $K_{l=1}^{1(2)}$ becomes negative, yet its running turns positive. Up to $\mathcal{O}(\alpha_0^3)$, the TLN is given by

$$K_{l=2}^1 = \left(\frac{3}{320} - \frac{9}{10} \log r \right) \alpha_0 + \left(-\frac{40025}{256} + \frac{648}{5} \zeta(3) + \frac{27}{20} \log r \right) \alpha_0^2 + \mathcal{O}(\alpha_0^3). \quad (97)$$

3. Axial gravitational field response

In this subsection, we compute the TLNs of the RBH with sub-Planckian curvature under axial gravitational perturbations for $l = 2$ and $l = 3$. For the case of $l = 2$, expanding the integral (29) in α_0 , the leading-order and next-to-leading-order terms are given by

$$I[\mathcal{S}^{(1)}](r) = \frac{337}{120} - \frac{51r}{20} - \frac{r^2}{10} - \frac{r^3}{12} - \frac{3r^4}{40}, \quad (98)$$

$$I[\mathcal{S}^{(2)}](r) = \frac{21491}{2880} - \frac{5729}{1920r^3} - \frac{189}{160r^2} - \frac{3301}{5760r} - \frac{803r}{320} - \frac{7r^2}{80} - \frac{17r^3}{240} - \frac{9r^4}{160} + \frac{2149}{2880} \log r. \quad (99)$$

Similar to the case of the Bardeen BH, we find that the TLNs exhibit no logarithmic running at the first order, while a logarithmic dependence appears at the second order. And the contributions are both positive. The TLN for $l = 2$ up to the second order in α_0 can be expressed as

$$K_{l=2}^2 = \frac{337}{120} \alpha_0 + \left(\frac{21491}{2880} + \frac{2149}{2880} \log r \right) \alpha_0^2 + \mathcal{O}(\alpha_0^3). \quad (100)$$

For the case of $l = 3$, expanding the integral (29) in α_0 , the leading-order and second-order terms are given by

$$I[\mathcal{S}^{(1)}](r) = \frac{1691}{4320} - \frac{425r}{336} + \frac{95r^2}{72} - \frac{1301r^3}{3024} - \frac{r^4}{672} + \frac{13r^5}{140} - \frac{3r^6}{28}, \quad (101)$$

$$I[\mathcal{S}^{(2)}](r) = -\frac{1427983}{3628800} - \frac{28747}{69120r^3} + \frac{51497}{50400r^2} + \frac{3562691}{7257600r} - \frac{867421r}{403200} + \frac{74633r^2}{40320} - \frac{25831r^3}{60480} + \frac{13r^4}{13440} + \frac{29r^5}{280} - \frac{9r^6}{112} + \frac{669067}{518400} \log r. \quad (102)$$

Similar to the $l = 2$ case, however, we note that the second-order constant term takes a negative value, indicating a weaker tidal response in comparison to the $l = 2$

mode. The expansion of the TLN for $l = 3$ up to the second order is given by

$$K_{l=3}^2 = \frac{1691}{4320} \alpha_0 + \left(-\frac{1427983}{3628800} + \frac{669067}{518400} \log r \right) \alpha_0^2 + \mathcal{O}(\alpha_0^3). \quad (103)$$

It is worth noting that in Ref. [28] the authors realized the Hayward metric as a background solution within the effective field theory framework of scalar-tensor theories with a timelike scalar profile [72], and evaluated the TLNs associated with axial gravitational perturbations. In contrast, in the present work the deviations from the classical singular Schwarzschild BH are modeled entirely within classical GR by introducing a nontrivial stress-energy tensor corresponding to an anisotropic fluid. The RBH possessing sub-Planckian curvature for the parameter choice $x = 1$ and $c = 3$ exhibits the same asymptotic behavior as the Hayward solution. A remarkable feature emerging from our results is that the logarithmic running of the TLNs already appears at the lowest multipole $l = 2$, whereas in the EFT-based realization it only shows up for $l \geq 4$. This highlights that the TLNs are sensitive probes of the internal structure of RBHs and are capable of distinguishing between different RBH realizations.

C. The regular black hole in asymptotically safe gravity

In this subsection, we calculate the TLNs of the RBH [52] in ASG under scalar, vector, and axial gravitational perturbations. Similarly, setting $r_h = 1$ we can express the BH mass M as a series expansion in terms of the deviation parameter ξ ,

$$M(\xi) = \frac{1}{2} + \frac{3\xi}{4} + \frac{3\xi^2}{4} + \frac{9\xi^3}{16} + \mathcal{O}(\xi^4). \quad (104)$$

Up to the same order in the parameter ξ , the metric function can be expanded as

$$f(r) = 1 - \frac{1}{r} - \frac{3(-1+r^3)\xi}{2r^4} - \frac{3(-2+r^3)(-1+r^3)\xi^2}{2r^7} - \frac{9(-6+12r^3-7r^6+r^9)\xi^3}{8r^{10}} + \mathcal{O}(\xi^4). \quad (105)$$

The effective potential for scalar field ($s = 0$) and vector field ($s = 1$) can be expanded in the same way as

$$U_1(r) = \frac{3l(l+1)-3s^2}{2r^3} + \frac{-3+6l(l+1)-6s^2}{4r^4} + \frac{-3+3l(l+1)-3s^2}{2r^5} + \frac{9(3+2s^2)}{4r^6}, \quad (106)$$

$$U_2(r) = \frac{3l(l+1)-3s^2}{2r^3} + \frac{-21+60l(l+1)-60s^2}{16r^4} + \frac{-69+96l(l+1)-96s^2}{16r^5} + \frac{72+15l(l+1)+39s^2}{4r^6}$$

$$+ \frac{93 + 12l(l+1) + 42s^2}{8r^7} + \frac{273 - 12l(l+1) + 120s^2}{16r^8} - \frac{9(69 + 20s^2)}{16r^9}, \quad (107)$$

$$U_3(r) = \frac{9(2l + 2l^2 - 2s^2)}{16r^3} + \frac{9(-3 + 10l + 10l^2 - 10s^2)}{16r^4} - \frac{9(15 - 24l - 24l^2 + 24s^2)}{16r^5} + \frac{9(42 + 30l + 30l^2 + 12s^2)}{16r^6} + \frac{9(66 + 24l + 24l^2 + 24s^2)}{16r^7} + \frac{9(141 + 6l + 6l^2 + 60s^2)}{16r^8} + \frac{9(-243 - 60s^2)}{16r^9} + \frac{9(-82 - 2l - 2l^2 - 16s^2)}{16r^{10}} + \frac{9(-152 + 2l + 2l^2 - 38s^2)}{16r^{11}} + \frac{9(288 + 54s^2)}{16r^{12}}. \quad (108)$$

In addition, the first-, second-, and third-order effective potential contributions for axial gravitational perturbations can be written explicitly as follows:

$$U_1(r) = \frac{-12 + 3l(l+1)}{2r^3} + \frac{-27 + 6l(l+1)}{4r^4} + \frac{-15 + 3l(l+1)}{2r^5} + \frac{171}{4r^6}, \quad (109)$$

$$U_2(r) = \frac{-12 + 3l(l+1)}{2r^3} + \frac{-261 + 60l(l+1)}{16r^4} + \frac{-453 + 96l(l+1)}{16r^5} + \frac{444 + 15l(l+1)}{4r^6} + \frac{477 + 12l(l+1)}{8r^7} + \frac{1185 - 12l(l+1)}{16r^8} - \frac{2925}{16r^9}, \quad (110)$$

$$U_3(r) = \frac{-36 + 9l(l+1)}{8r^3} + \frac{-387 + 90l(l+1)}{16r^4} + \frac{27(-37 + 8l(l+1))}{16r^5} + \frac{27(43 + 5l(l+1))}{8r^6} + \frac{27(59 + 4l(l+1))}{8r^7} + \frac{9(645 + 6l(l+1))}{16r^8} - \frac{10395}{16r^9} - \frac{9(181 + l(l+1))}{8r^{10}} + \frac{9(-296 + l(l+1))}{8r^{11}} + \frac{1377}{2r^{12}}. \quad (111)$$

Next, we compute the TLNs of the RBH in ASG for scalar, vector, and axial gravitational perturbations.

1. Scalar field response

We first compute the TLNs of the RBH in ASG under scalar perturbations for $l = 0$ and $l = 1$. For the case $l = 0$, the zeroth-order ($\mathcal{O}(\xi^0)$) modes, representing the growing and decaying solutions in Eqs. (20) and (21),

take the explicit forms

$$\Phi_+^{(0)}(r) = \Phi^{(0)}(r) = \Phi_{\text{hor-reg}}(r) = r, \quad (112)$$

$$\Phi_-^{(0)}(r) = -r \log\left(1 - \frac{1}{r}\right). \quad (113)$$

When $s = 0$ and $l = 0$, the first-order source term at $\mathcal{O}(\xi^1)$ can be evaluated from Eq. (25), yielding

$$S^{(1)}(r) = U^{(1)}(r) \Phi^{(0)}(r) = \frac{27}{4r^5} - \frac{3}{2r^4} - \frac{3}{4r^3}. \quad (114)$$

Using the Green's function approach, the first-order solution and the corresponding second-order source term $\mathcal{O}(\xi^2)$ are obtained as

$$\Phi^{(1)}(r) = \int_1^\infty G(r, r') S^{(1)}(r') dr' = -\frac{3}{4} - \frac{3}{4r^2} - \frac{3}{4r}, \quad (115)$$

$$S^{(2)}(r) = U^{(2)}(r) \Phi^{(0)}(r) + U^{(1)}(r) \Phi^{(1)}(r) = -\frac{351}{8r^8} + \frac{105}{8r^7} + \frac{33}{4r^6} + \frac{315}{16r^5} - \frac{15}{4r^4} - \frac{21}{16r^3}. \quad (116)$$

Proceeding to the next order, the second-order solution along with the third-order source term at $\mathcal{O}(\xi^3)$, as indicated by Eqs. (28) and (25), are given by

$$\Phi^{(2)}(r) = \int_1^\infty G(r, r') S^{(2)}(r') dr' = -\frac{3}{4} + \frac{39}{32r^5} + \frac{15}{16r^4} + \frac{21}{32r^3} - \frac{21}{16r^2} - \frac{33}{32r}, \quad (117)$$

$$S^{(3)}(r) = U^{(3)}(r) \Phi^{(0)}(r) + U^{(2)}(r) \Phi^{(1)}(r) + U^{(1)}(r) \Phi^{(2)}(r) = \frac{25515}{128r^{11}} - \frac{1035}{16r^{10}} - \frac{2331}{64r^9} - \frac{729}{4r^8} + \frac{1755}{32r^7} + \frac{405}{16r^6} + \frac{3807}{128r^5} - \frac{441}{64r^4} - \frac{27}{16r^3}. \quad (118)$$

For $l = 0$, we can extract the TLNs from the following functions:

$$I[\mathcal{S}^{(1)}](r) = -\frac{3}{4} + \frac{9}{4r^3} - \frac{3}{4r^2} - \frac{3}{4r}, \quad (119)$$

$$I[\mathcal{S}^{(2)}](r) = -\frac{3}{4} - \frac{117}{16r^6} + \frac{21}{8r^5} + \frac{33}{16r^4} + \frac{105}{16r^3} - \frac{15}{8r^2} - \frac{21}{16r}, \quad (120)$$

$$I[\mathcal{S}^{(3)}](r) = -\frac{9}{16} + \frac{2835}{128r^9} - \frac{1035}{128r^8} - \frac{333}{64r^7} - \frac{243}{8r^6} + \frac{351}{32r^5} + \frac{405}{64r^4} + \frac{1269}{128r^3} - \frac{441}{128r^2} - \frac{27}{16r}. \quad (121)$$

From the above, we can see that the TLNs are non-zero and no logarithmic running appears. Moreover, the con-

tribution at each order of ξ to TLN is all negative, showing similar behavior to the RBH with sub-Planckian curvature. Thus, up to the third order of ξ the scalar TLN for $l = 0$ is expressed as

$$K_{l=0}^0 = -\frac{3}{4}\xi - \frac{3}{4}\xi^2 - \frac{9}{16}\xi^3 + \mathcal{O}(\xi^4). \quad (122)$$

For $l = 1$, the scalar TLNs can be obtained from the following expressions:

$$I[\mathcal{S}^{(1)}](r) = -\frac{13}{16} + \frac{3}{16r^3} - \frac{17}{16r^2} + \frac{31}{16r} + \frac{r}{4} - \frac{r^2}{2}, \quad (123)$$

$$I[\mathcal{S}^{(2)}](r) = -\frac{35}{32} - \frac{39}{64r^6} + \frac{13}{4r^5} - \frac{299}{64r^4} + \frac{71}{64r^3} - \frac{95}{32r^2} + \frac{323}{64r} + \frac{7r}{16} - \frac{r^2}{2}, \quad (124)$$

$$I[\mathcal{S}^{(3)}](r) = -\frac{591}{448} + \frac{945}{512r^9} - \frac{24609}{2560r^8} + \frac{7611}{560r^7} - \frac{6399}{1280r^6} + \frac{17247}{1280r^5} - \frac{3237}{160r^4} + \frac{9099}{2560r^3} - \frac{2475}{512r^2} + \frac{267}{32r} + \frac{9r}{16} - \frac{3r^2}{8}. \quad (125)$$

Similar to the $l = 1$ case, we can see that the TLNs at each order of ξ are all negative and do not exhibit running terms. The explicit expansion of the scalar TLN for $l = 1$ can be written as

$$K_{l=1}^0 = -\frac{13}{16}\xi - \frac{35}{32}\xi^2 - \frac{591}{448}\xi^3 + \mathcal{O}(\xi^4). \quad (126)$$

2. Vector field response

Next, we compute the TLNs of the RBH in ASG under vector perturbations, for $l = 1$ and $l = 2$. For $l = 1$, the TLNs can be obtained from the following expressions:

$$I[\mathcal{S}^{(1)}](r) = -\frac{13}{4} + \frac{15}{4r} - \frac{r}{4} - \frac{r^2}{4}, \quad (127)$$

$$I[\mathcal{S}^{(2)}](r) = -\frac{35}{8} - \frac{123}{16r^4} + \frac{13}{8r^3} + \frac{23}{16r^2} + \frac{155}{16r} - \frac{7r}{16} - \frac{r^2}{4}, \quad (128)$$

$$I[\mathcal{S}^{(3)}](r) = -\frac{4917}{896} + \frac{18063}{896r^7} - \frac{801}{128r^6} - \frac{93}{20r^5} - \frac{10221}{320r^4} + \frac{2439}{320r^3} + \frac{45}{8r^2} + \frac{2007}{128r} - \frac{9r}{16} - \frac{3r^2}{16}. \quad (129)$$

Similar to scalar TLNs, the contributions to the TLN at each order are all negative and no logarithmic running appears. Up to the third order, the explicit expansion of the vector TLN for $l = 2$ can thus be written as

$$K_{l=1}^1 = -\frac{13}{4}\xi - \frac{35}{8}\xi^2 - \frac{4917}{896}\xi^3 + \mathcal{O}(\xi^4). \quad (130)$$

For $l = 2$, we mainly compute the first two orders of the TLNs. As mentioned earlier, the third-order integral (29) cannot be obtained analytically. The TLN can be obtained from the following expressions:

$$I[\mathcal{S}^{(1)}](r) = \frac{9}{320} + \frac{81}{64r} - \frac{387r}{320} - \frac{3r^2}{320} + \frac{3r^3}{10} - \frac{3r^4}{8} + \frac{27\log r}{10}, \quad (131)$$

$$I[\mathcal{S}^{(2)}](r) = -\frac{73383}{64} + \frac{81\pi^2}{80} - \frac{3321}{1280r^4} - \frac{2457}{640r^3} - \frac{27135}{256r^2} + \frac{1576023}{1280r} + \frac{729\pi^2}{16r} - \frac{11271r}{1280} - \frac{3483\pi^2 r}{80} - \frac{28527r^2}{640} - \frac{27\pi^2 r^2}{80} + \frac{3261r^3}{40} + \frac{54\pi^2 r^3}{5} - \frac{3r^4}{8} - \frac{27\pi^2 r^4}{2} + \frac{21843\log r}{80} + \frac{972\pi^2 \log r}{5} - \frac{81\log r}{16r^3} - \frac{3159\log r}{80r^2} + \frac{56997\log r}{80r} - \frac{27r\log r}{8} - \frac{243r^2\log r}{10} + 81r^3\log r + \frac{243\log^2 r}{80} + \frac{2187\log^2 r}{16r} - \frac{10449r\log^2 r}{80} - \frac{81r^2\log^2 r}{80} + \frac{162r^3\log^2 r}{5} - \frac{81r^4\log^2 r}{2} + \frac{486\log^3 r}{5} + \frac{243}{40}\text{Li}_2(1-r) + \frac{2187\text{Li}_2(1-r)}{8r} - \frac{10449r\text{Li}_2(1-r)}{40} - \frac{81r^2\text{Li}_2(1-r)}{40} + \frac{324r^3\text{Li}_2(1-r)}{5} - 81r^4\text{Li}_2(1-r) + \frac{2916\log r\text{Li}_2(r)}{5} - \frac{5832}{5}\text{Li}_3(r) + \frac{5832\zeta(3)}{5}. \quad (132)$$

We find that the first-order contribution to the TLN is positive, while the logarithmic running behaves inversely. The second-order correction of the vector TLN can be obtained from the asymptotic behavior of $I[\mathcal{S}^{(2)}](r)$ at large r . Up to $\mathcal{O}(\xi^3)$, the vector TLN for $l = 2$ is given by

$$K_{l=2}^1 = \left(\frac{9}{320} - \frac{27}{10}\log r\right)\xi + \left(-\frac{89997}{64} + \frac{5832}{5}\zeta(3) + \frac{243}{20}\log r\right)\xi^2 + \mathcal{O}(\xi^3), \quad (133)$$

from which one can see that the second-order contribution is negative while the running is positive.

3. Axial gravitational field response

Finally, we compute the TLNs of the RBH under axial gravitational perturbations within the framework of ASG

for $l = 2$ and $l = 3$. For the case of $l = 2$, the expansion of the integral (29) in ξ is given by

$$I[\mathcal{S}^{(1)}](r) = 9 - \frac{171r}{20} - \frac{3r^2}{20} - \frac{3r^3}{20} - \frac{3r^4}{20}, \quad (134)$$

$$I[\mathcal{S}^{(2)}](r) = \frac{159}{2} - \frac{513}{16r^3} - \frac{927}{40r^2} + \frac{177}{80r} - \frac{411r}{16} - \frac{3r^2}{8} - \frac{21r^3}{80} - \frac{3r^4}{20}, \quad (135)$$

$$I[\mathcal{S}^{(3)}](r) = \frac{269379}{11200} + \frac{45333}{448r^6} + \frac{1832517}{22400r^5} + \frac{1538937}{22400r^4} - \frac{1060911}{5600r^3} - \frac{63063}{800r^2} + \frac{62289}{1600r} - \frac{28971r}{640} - \frac{441r^2}{640} - \frac{27r^3}{80} - \frac{9r^4}{80}. \quad (136)$$

Consequently, the TLN for $l = 2$ up to the third order can directly read as

$$K_{l=2}^2 = 9\xi + \frac{159}{2}\xi^2 + \frac{269379}{11200}\xi^3 + \mathcal{O}(\xi^4). \quad (137)$$

It is evident that the TLN is nonzero in this case. However, we find that, even when computed up to the third order, no logarithmic running term appears for the $l = 2$ mode under axial gravitational perturbations in ASG, in contrast to the previous two models. Moreover, their contributions to the TLN are all positive, which is consistent with the previous two types of RBHs.

For the case of $l = 3$, the expansion of the integral (29) in ξ is given as follows

$$I[\mathcal{S}^{(1)}](r) = \frac{9}{7} - \frac{475r}{112} + \frac{1535r^2}{336} - \frac{529r^3}{336} - \frac{r^4}{336} + \frac{r^5}{4} - \frac{2r^6}{7}, \quad (138)$$

$$I[\mathcal{S}^{(2)}](r) = -\frac{208181}{6720} - \frac{513}{112r^3} + \frac{10763}{2240r^2} + \frac{124613}{3360r} - \frac{19157r}{960} + \frac{1529r^2}{84} - \frac{907r^3}{192} - \frac{5r^4}{672} + \frac{7r^5}{16} - \frac{2r^6}{7} + \frac{237 \log r}{7}, \quad (139)$$

$$I[\mathcal{S}^{(3)}](r) = -\frac{143685019}{376320} - \frac{120825}{12544r^6} - \frac{1309101}{31360r^5} - \frac{4498499}{125440r^4} + \frac{3584485}{18816r^3} - \frac{8035661}{125440r^2} + \frac{2444089}{8960r} + \frac{57889r}{1280} + \frac{114045r^2}{3584} - \frac{19847r^3}{2688} + \frac{57r^4}{112} - \frac{69r^5}{560} - \frac{3r^6}{14} + \frac{320991 \log r}{4480}. \quad (140)$$

Unlike the $l = 2$ case, here the logarithmic running terms appear at both the second and third orders. Only the first-order term contributes positively to the TLNs, while

the second and third-order terms contribute negatively. Accordingly, the TLNs up to the third order can be expressed as

$$K_{l=3}^2 = \frac{9}{7}\xi + \left(-\frac{208181}{6720} + \frac{237 \log r}{7}\right)\xi^2 + \left(-\frac{143685019}{376320} + \frac{320991 \log r}{4480}\right)\xi^3 + \mathcal{O}(\xi^4). \quad (141)$$

This indicates that for $l = 3$, the logarithmic contributions emerge beyond the leading order, marking a qualitative difference from the $l = 2$ case in the axial gravitational perturbations of the RBH in ASG.

D. Comparison of three regular black holes

In Table I, we present the leading-order tidal Love numbers (C_{const} , C_{log}) for RBHs under scalar, vector, and axial gravitational perturbations. All regular parameters are evaluated at $q^2 = 0.1$, $\alpha_0 = 0.1$, and $\xi = 0.1$ for the Bardeen BH, the BH with sub-Planckian curvature and the RBH in ASG, respectively. We can clearly observe both the connections and distinctions among the three models. The key finding is that all RBH models exhibit generally nonzero TLNs under scalar, vector, and gravitational perturbations, signaling a breakdown of the classical “no-hair” theorem in the quantum gravity regime and confirming a dynamical response of the internal structure to external tidal fields.

From Table I we can identify several qualitative and potentially observable discriminants among the three RBH models.

(i) Scalar sector. For the monopole mode $l = 0$, all models exhibit negative, scale-independent TLNs, so this mode is not very diagnostic. However, the dipole mode $l = 1$ shows a clear separation: the Bardeen BH yields a positive TLN with non-vanishing logarithmic running, whereas both the sub-Planckian and ASG black holes give negative TLNs without any running. Thus, the sign and scale dependence of the scalar dipole TLN already distinguish de Sitter-core geometries from Minkowski-core and ASG BHs.

(ii) Vector sector. In the dipole mode $l = 1$, the Bardeen BH again stands out by having a positive TLN accompanied by a sizeable negative logarithmic term, while the other two models have negative, non-running TLNs. For the quadrupole mode $l = 2$, the three models occupy distinct regions in the (C_{const} , C_{log}), Bardeen has ($C_{\text{const}} < 0$, $C_{\text{log}} < 0$), the sub-Planckian BH has a very small positive constant and a moderate positive running, and the ASG BH shows a slightly positive constant but a much larger positive running. This pattern provides a two-dimensional “fingerprint” for discriminating the three scenarios.

(iii) Axial gravitational sector. The axial quadrupole TLN $l = 2$, which is expected to be the most relevant for

GW observations, exhibits a pronounced enhancement in the ASG case: at the same deformation parameters, the ASG BH has $C_{\text{const}} \approx 0.9$, about a factor of three larger than the Bardeen and sub-Planckian values $\sim 0.26\text{--}0.28$, while remaining free of logarithmic running. The octupole mode $l = 3$ shows a similar hierarchy of magnitudes. These features indicate that the ASG BH leaves a much stronger imprint in the axial gravitational channel, offering a particularly clean observational handle on this quantum-gravity motivated model.

These results indicate that TLNs not only serve as a criterion for distinguishing classical BHs from their quantum-corrected counterparts, but also act as a precision diagnostic probe, offering a new theoretical window into decoding the internal structure of BHs and the underlying quantum gravitational mechanisms. Moreover, they highlight promising directions for future GW observatories to test these physical phenomena.

V. CONCLUSIONS

This paper presents a systematic theoretical study of the tidal response of three important classes of RBH models—Bardeen BH, the BH with sub-Planckian curvature, and the BH in the ASG framework—under static tidal fields, with a focus on calculating and analyzing their TLNs. In GR, the TLNs of four-dimensional, asymptotically flat BHs vanish, indicating that they do not develop induced multipole moments under external tidal fields, exhibiting a unique “rigidity.” However, beyond classical GR—e.g., when quantum-gravity effects or matter fields are considered in RBH models—this zero property can be violated. Therefore, TLNs serve as an extremely sensitive probe for investigating BH interior structures, testing GR, and exploring quantum gravity effects.

The core methodology of this work is an analytic Green’s function technique developed in [28]. For consistent comparisons across different RBH models, all horizons were normalized to $r_h = 1$. The mass M and metric function $f(r)$ of each BH were then expanded as series around their respective deviation or regular parameters (q for Bardeen, α_0 for sub-Planckian, and ξ for ASG BH). By introducing a field redefinition, all corrections to GR are encapsulated in a new effective potential $U(r)$. The perturbation equations are then solved order by order: the zeroth-order solution corresponds to the Schwarzschild background and is expressed in terms of hypergeometric functions, while higher-order solutions are obtained systematically via the Green’s function formalism to handle the inhomogeneous terms. The TLNs are extracted from the coefficients of the r^{-l} terms in the asymptotic expansion at spatial infinity. Notably, when logarithmic terms appear (e.g., $\log r$), their co-

efficients are also interpreted as part of the TLNs and can be understood in a classical RG framework as a “ β -function” [8, 10, 11].

The study yields rich and revealing results. First, for all three classes of RBHs, the TLNs are generically non-zero under scalar, vector, and axial gravitational perturbations, in stark contrast to classical GR BHs. This directly confirms that the nontrivial core structures of RBHs (e.g., the de Sitter core of Bardeen BH or the Minkowski core of sub-Planckian curvature BH) significantly affect their tidal response. Second, the TLNs strongly depend on the perturbation type and multipole l . For instance, in scalar perturbations of Bardeen BH, the $l = 0$ TLNs are negative and free of logarithms, whereas $l = 1$ exhibits logarithmic structure. Third, a key finding is the ubiquitous presence of logarithmic running. In many cases, first-order corrections do not contain logarithms, but second- or third-order corrections produce TLNs of the form $K \sim \text{constant} + \beta \log r$. This logarithmic term β renders the TLNs scale-dependent, analogous to running couplings in quantum field theory, suggesting an underlying renormalization-group structure even in classical gravity systems.

These findings have significant physical implications. Non-zero TLNs provide direct theoretical guidance for distinguishing RBHs from classical ones via GW observations. Future observatories, such as LISA [73], Taiji [74] or TianQin [75], may detect these tidal imprints during binary black hole mergers. The emergence of logarithmic terms further links TLNs to quantum gravity concepts like asymptotic safety, indicating that BHs’ tidal responses may encode high-energy quantum gravitational signatures.

Looking forward, this work opens multiple promising directions. Theoretically, it would be interesting to study more realistic rotating RBHs or to study responses under dynamical (non-static) tidal fields. For higher-order computations, combining numerical and analytic techniques becomes necessary due to the complexity of integrals. Observationally, incorporating these TLNs into waveform models of binary mergers will be crucial for assessing their detectability with future GW detectors. Finally, understanding the microscopic origin of logarithmic terms and their potential role in BH thermodynamics and the information paradox will likely provide a frontier connection between gravity, quantum theory, and differential geometry.

ACKNOWLEDGMENTS

The work is in part supported by NSFC Grant No. 12205104 and the startup funding of South China University of Technology.

TABLE I. Leading-order tidal Love numbers (C_{const} , C_{log}) for regular black holes under scalar, vector, and axial gravitational perturbations. All coefficients are evaluated at $q^2 = 0.1$, $\alpha_0 = 0.1$, and $\xi = 0.1$ for the Bardeen, Sub-Planckian, and ASG black holes, respectively.

Perturbation Type	Mode l	Bardeen BH ($q^2 = 0.1$)	Sub-Planckian BH ($\alpha_0 = 0.1$)	ASG BH ($\xi = 0.1$)
Scalar	$l = 0$	$C_{\text{const}} = -0.075$ $C_{\text{log}} = 0$	$C_{\text{const}} = -0.025$ $C_{\text{log}} = 0$	$C_{\text{const}} = -0.075$ $C_{\text{log}} = 0$
	$l = 1$	$C_{\text{const}} = 0.094$ $C_{\text{log}} = -0.05$	$C_{\text{const}} = -0.027$ $C_{\text{log}} = 0$	$C_{\text{const}} = -0.081$ $C_{\text{log}} = 0$
	$l = 2$	$C_{\text{const}} = 0.05$ $C_{\text{log}} = -0.2$	$C_{\text{const}} = -0.11$ $C_{\text{log}} = 0$	$C_{\text{const}} = -0.33$ $C_{\text{log}} = 0$
	$l = 3$	$C_{\text{const}} = -0.096$ $C_{\text{log}} = -0.068$	$C_{\text{const}} = 0.00094$ $C_{\text{log}} = 0.09$	$C_{\text{const}} = 0.0028$ $C_{\text{log}} = 0.27$
Vector	$l = 1$	$C_{\text{const}} = 0.26$ $C_{\text{log}} = 0$	$C_{\text{const}} = 0.28$ $C_{\text{log}} = 0$	$C_{\text{const}} = 0.9$ $C_{\text{log}} = 0$
	$l = 2$	$C_{\text{const}} = 0.017$ $C_{\text{log}} = 0$	$C_{\text{const}} = 0.039$ $C_{\text{log}} = 0$	$C_{\text{const}} = 0.13$ $C_{\text{log}} = 0$
	$l = 3$	$C_{\text{const}} = 0.017$ $C_{\text{log}} = 0$	$C_{\text{const}} = 0.039$ $C_{\text{log}} = 0$	$C_{\text{const}} = 0.13$ $C_{\text{log}} = 0$
	$l = 4$	$C_{\text{const}} = 0.017$ $C_{\text{log}} = 0$	$C_{\text{const}} = 0.039$ $C_{\text{log}} = 0$	$C_{\text{const}} = 0.13$ $C_{\text{log}} = 0$
Axial Grav.	$l = 2$	$C_{\text{const}} = 0.26$ $C_{\text{log}} = 0$	$C_{\text{const}} = 0.28$ $C_{\text{log}} = 0$	$C_{\text{const}} = 0.9$ $C_{\text{log}} = 0$
	$l = 3$	$C_{\text{const}} = 0.017$ $C_{\text{log}} = 0$	$C_{\text{const}} = 0.039$ $C_{\text{log}} = 0$	$C_{\text{const}} = 0.13$ $C_{\text{log}} = 0$
	$l = 4$	$C_{\text{const}} = 0.017$ $C_{\text{log}} = 0$	$C_{\text{const}} = 0.039$ $C_{\text{log}} = 0$	$C_{\text{const}} = 0.13$ $C_{\text{log}} = 0$
	$l = 5$	$C_{\text{const}} = 0.017$ $C_{\text{log}} = 0$	$C_{\text{const}} = 0.039$ $C_{\text{log}} = 0$	$C_{\text{const}} = 0.13$ $C_{\text{log}} = 0$

- [1] B. P. Abbott *et al.* (LIGO Scientific, Virgo), Phys. Rev. Lett. **116**, 061102 (2016), arXiv:1602.03837 [gr-qc].
- [2] A. E. H. Love, Proceedings of the Royal Society of London. Series A **82**, 273 (1909).
- [3] E. E. Flanagan and T. Hinderer, Phys. Rev. D **77**, 021502 (2008), arXiv:0709.1915 [astro-ph].
- [4] B. P. Abbott *et al.* (LIGO Scientific, Virgo), Phys. Rev. Lett. **121**, 161101 (2018), arXiv:1805.11581 [gr-qc].
- [5] H. Fang and G. Lovelace, Phys. Rev. D **72**, 124016 (2005), arXiv:gr-qc/0505156.
- [6] T. Binnington and E. Poisson, Phys. Rev. D **80**, 084018 (2009), arXiv:0906.1366 [gr-qc].
- [7] T. Damour and A. Nagar, Phys. Rev. D **80**, 084035 (2009), arXiv:0906.0096 [gr-qc].
- [8] B. Kol and M. Smolkin, JHEP **02**, 010, arXiv:1110.3764 [hep-th].
- [9] N. Gürlebeck, Phys. Rev. Lett. **114**, 151102 (2015), arXiv:1503.03240 [gr-qc].
- [10] L. Hui, A. Joyce, R. Penco, L. Santoni, and A. R. Solomon, JCAP **04**, 052, arXiv:2010.00593 [hep-th].
- [11] V. Cardoso, E. Franzin, A. Maselli, P. Pani, and G. Raposo, Phys. Rev. D **95**, 084014 (2017), [Addendum: Phys.Rev.D 95, 089901 (2017)], arXiv:1701.01116 [gr-qc].
- [12] D. Pereñíguez and V. Cardoso, Phys. Rev. D **105**, 044026 (2022), arXiv:2112.08400 [gr-qc].
- [13] M. Rai and L. Santoni, JHEP **07**, 098, arXiv:2404.06544 [gr-qc].
- [14] S. Chakraborty, P. Heidmann, and P. Pani, (2025), arXiv:2508.20155 [gr-qc].
- [15] X. Pang, Y. Tian, H. Zhang, and Q. Jiang, (2025), arXiv:2510.10036 [gr-qc].
- [16] P. Landry and E. Poisson, Phys. Rev. D **91**, 104018 (2015), arXiv:1503.07366 [gr-qc].
- [17] P. Pani, L. Gualtieri, and V. Ferrari, Phys. Rev. D **92**, 124003 (2015), arXiv:1509.02171 [gr-qc].
- [18] E. Poisson, Phys. Rev. D **91**, 044004 (2015), arXiv:1411.4711 [gr-qc].
- [19] A. Le Tiec, M. Casals, and E. Franzin, Phys. Rev. D **103**, 084021 (2021), arXiv:2010.15795 [gr-qc].
- [20] A. Le Tiec and M. Casals, Phys. Rev. Lett. **126**, 131102 (2021), arXiv:2007.00214 [gr-qc].
- [21] H. S. Chia, Phys. Rev. D **104**, 024013 (2021), arXiv:2010.07300 [gr-qc].
- [22] P. Charalambous, S. Dubovsky, and M. M. Ivanov, Phys. Rev. Lett. **127**, 101101 (2021), arXiv:2103.01234 [hep-th].
- [23] P. Charalambous, S. Dubovsky, and M. M. Ivanov, JHEP **05**, 038, arXiv:2102.08917 [hep-th].
- [24] M. M. Ivanov and Z. Zhou, Phys. Rev. Lett. **130**, 091403 (2023), arXiv:2209.14324 [hep-th].
- [25] L. Ma, Z.-H. Wu, Y. Pang, and H. Lu, Phys. Rev. D **111**, 044003 (2025), arXiv:2408.10352 [gr-qc].
- [26] V. Cardoso, M. Kimura, A. Maselli, and L. Senatore, Phys. Rev. Lett. **121**, 251105 (2018), [Erratum: Phys.Rev.Lett. 131, 109903 (2023)], arXiv:1808.08962 [gr-qc].
- [27] V. De Luca, J. Khoury, and S. S. C. Wong, Phys. Rev. D **108**, 044066 (2023), arXiv:2211.14325 [hep-th].
- [28] C. G. A. Barura, H. Kobayashi, S. Mukohyama, N. Oshita, K. Takahashi, and V. Yingcharoenrat, JCAP **09**, 001, arXiv:2405.10813 [gr-qc].
- [29] T. Katagiri, V. Cardoso, T. Ikeda, and K. Yagi, Phys. Rev. D **111**, 084081 (2025), arXiv:2410.02531 [gr-qc].
- [30] V. Cardoso and F. Duque, Phys. Rev. D **101**, 064028 (2020), arXiv:1912.07616 [gr-qc].
- [31] V. De Luca, A. Maselli, and P. Pani, Phys. Rev. D **107**, 044058 (2023), arXiv:2212.03343 [gr-qc].
- [32] E. Cannizzaro, V. De Luca, and P. Pani, Phys. Rev. D **110**, 123004 (2024), arXiv:2408.14208 [astro-ph.HE].
- [33] S. Chakraborty, G. Compère, and L. Machet, Phys. Rev. D **112**, 024015 (2025), arXiv:2412.14831 [gr-qc].
- [34] V. Cardoso and P. Pani, Living Rev. Rel. **22**, 4 (2019), arXiv:1904.05363 [gr-qc].
- [35] C. Chirenti, C. Posada, and V. Guedes, Class. Quant.

- Grav. **37**, 195017 (2020), arXiv:2005.10794 [gr-qc].
- [36] M. Collier, D. Croon, and R. K. Leane, Phys. Rev. D **106**, 123027 (2022), arXiv:2205.15337 [gr-qc].
- [37] S. Nair, S. Chakraborty, and S. Sarkar, Phys. Rev. D **107**, 124041 (2023), arXiv:2208.06235 [gr-qc].
- [38] A. Maselli, P. Pani, V. Cardoso, T. Abdelsalhin, L. Gualtieri, and V. Ferrari, Phys. Rev. Lett. **120**, 081101 (2018), arXiv:1703.10612 [gr-qc].
- [39] S. Datta, Class. Quant. Grav. **39**, 225016 (2022), arXiv:2107.07258 [gr-qc].
- [40] H. S. Chia, T. D. P. Edwards, D. Wadekar, A. Zimmerman, S. Olsen, J. Roulet, T. Venumadhav, B. Zackay, and M. Zaldarriaga, Phys. Rev. D **110**, 063007 (2024), arXiv:2306.00050 [gr-qc].
- [41] G. A. Piovano, A. Maselli, and P. Pani, Phys. Rev. D **107**, 024021 (2023), arXiv:2207.07452 [gr-qc].
- [42] T. Zi and P.-C. Li, Phys. Rev. D **108**, 024018 (2023), arXiv:2303.16610 [gr-qc].
- [43] M. Andrés-Carcasona and G. Caneva Santoro, (2025), arXiv:2512.01918 [gr-qc].
- [44] C. Lan, H. Yang, Y. Guo, and Y.-G. Miao, Int. J. Theor. Phys. **62**, 202 (2023), arXiv:2303.11696 [gr-qc].
- [45] C. Bambi, ed., *Regular Black Holes. Towards a New Paradigm of Gravitational Collapse*, Springer Series in Astrophysics and Cosmology (Springer, 2023) arXiv:2307.13249 [gr-qc].
- [46] J. M. Bardeen, (1968).
- [47] E. Ayon-Beato and A. Garcia, Phys. Lett. B **493**, 149 (2000), arXiv:gr-qc/0009077.
- [48] Y. Ling and M.-H. Wu, Class. Quant. Grav. **40**, 075009 (2023), arXiv:2109.05974 [gr-qc].
- [49] L. Xiang, Y. Ling, and Y. G. Shen, Int. J. Mod. Phys. D **22**, 1342016 (2013), arXiv:1305.3851 [gr-qc].
- [50] A. Bonanno and M. Reuter, Phys. Rev. D **62**, 043008 (2000), arXiv:hep-th/0002196.
- [51] A. Platania, Black Holes in Asymptotically Safe Gravity (2023) arXiv:2302.04272 [gr-qc].
- [52] A. Bonanno, D. Malafarina, and A. Panassiti, Phys. Rev. Lett. **132**, 031401 (2024), arXiv:2308.10890 [gr-qc].
- [53] A. Spina, Int. J. Grav. Theor. Phys. **1**, 8 (2025), arXiv:2510.14552 [gr-qc].
- [54] M. Motaharfar and P. Singh, Phys. Rev. D **111**, 106018 (2025), arXiv:2501.09151 [gr-qc].
- [55] C. Coviello, V. Vellucci, and L. Lehner, Phys. Rev. D **111**, 104073 (2025), arXiv:2503.04287 [gr-qc].
- [56] M. Motaharfar and P. Singh, Phys. Rev. D **112**, 066008 (2025), arXiv:2505.14784 [gr-qc].
- [57] Y. Liu and X. Zhang, (2025), arXiv:2509.12013 [gr-qc].
- [58] R. V. Maluf and J. C. S. Neves, Int. J. Mod. Phys. D **28**, 1950048 (2018), arXiv:1801.08872 [gr-qc].
- [59] I. Dymnikova, Gen. Rel. Grav. **24**, 235 (1992).
- [60] S. A. Hayward, Phys. Rev. Lett. **96**, 031103 (2006), arXiv:gr-qc/0506126.
- [61] V. P. Frolov, Phys. Rev. D **94**, 104056 (2016), arXiv:1609.01758 [gr-qc].
- [62] M. Niedermaier, Class. Quant. Grav. **24**, R171 (2007), arXiv:gr-qc/0610018.
- [63] M. A. Markov and V. F. Mukhanov, Nuovo Cim. B **86**, 97 (1985).
- [64] T. Regge and J. A. Wheeler, Phys. Rev. **108**, 1063 (1957).
- [65] F. J. Zerilli, Phys. Rev. Lett. **24**, 737 (1970).
- [66] R. A. Konoplya and O. S. Stashko, Phys. Rev. D **111**, 104055 (2025), arXiv:2408.02578 [gr-qc].
- [67] B. Toshmatov, C. Bambi, B. Ahmedov, A. Abdujabbarov, and Z. Stuchlík, Eur. Phys. J. C **77**, 542 (2017), arXiv:1702.06855 [gr-qc].
- [68] A. Ashtekar, J. Olmedo, and P. Singh, Phys. Rev. D **98**, 126003 (2018), arXiv:1806.02406 [gr-qc].
- [69] C.-Y. Chen and P. Chen, Phys. Rev. D **99**, 104003 (2019), arXiv:1902.01678 [gr-qc].
- [70] Q.-L. Shi, R. Wang, W. Xiong, and P.-C. Li, (2025), arXiv:2506.16217 [gr-qc].
- [71] V. Cardoso, K. Destounis, F. Duque, R. Panosso Macedo, and A. Maselli, Phys. Rev. Lett. **129**, 241103 (2022), arXiv:2210.01133 [gr-qc].
- [72] S. Mukohyama, K. Takahashi, K. Tomikawa, and V. Yingcharoenrat, JCAP **07**, 050, arXiv:2304.14304 [gr-qc].
- [73] P. Amaro-Seoane *et al.* (LISA), (2017), arXiv:1702.00786 [astro-ph.IM].
- [74] W.-H. Ruan, Z.-K. Guo, R.-G. Cai, and Y.-Z. Zhang, Int. J. Mod. Phys. A **35**, 2050075 (2020), arXiv:1807.09495 [gr-qc].
- [75] J. Mei *et al.* (TianQin), PTEP **2021**, 05A107 (2021), arXiv:2008.10332 [gr-qc].