

INVARIANT POLYNOMIALS, GAPS, AND SPARSENESS

JOHN P. D'ANGELO, DUSTY E. GRUNDMEIER, AND DANIEL A. LICHTBLAU

ABSTRACT. We consider each of the three classes of representations of cyclic groups that arise in the study of rational sphere maps. We study the possible number of terms for invariant polynomials with non-negative coefficients that are constant on the appropriate line or hyperplane. Our result provides crucial information about gaps in the possible target dimensions for certain invariant polynomial sphere maps. We interpret our results in terms of sparseness for solutions of certain linear systems.

AMS Classification Numbers: 32H35, 32M99, 32V99, 11B13, 11B05.

Key words: Rational sphere maps, Hermitian invariant groups, gaps, sparse linear systems, Frobenius number, degree estimates.

1. INTRODUCTION

Let Γ denote a fixed-point free finite subgroup of the unitary group $\mathbf{U}(n)$. Two of the authors proved (see [22] and [9]) that the existence of a non-constant rational sphere map invariant under Γ restricts Γ considerably. The group must be cyclic and represented in one of only three ways. Each of these three situations leads to questions about the possible minimal target dimensions for such maps. These questions lead us to study those invariant polynomials with non-negative coefficients that are constant on a line or plane and ask how many terms such polynomials can have. In our situation the set of possible values (of the number of terms) satisfies a Frobenius or postage stamp property; see Proposition 2.3. Not every value is possible. Thus there are **gaps** in the target dimensions for (so-called minimal) invariant maps. Gaps in the possible target dimensions for rational sphere maps also exist. See [19] and [20] for various results about such gaps. Because of invariance, the combinatorial results in this paper differ considerably. The invariance also has consequences for sparseness of linear systems.

The three possible groups are described as follows.

Example 1.1. Let η be a primitive m -th root of unity. Let $\Gamma(m, 1)$ be the cyclic group of order m generated by ηI , where I denotes the identity map on complex Euclidean space \mathbb{C}^n for some n .

Example 1.2. Let η be a primitive p -th root of unity where p is odd. Let $\Gamma(p, 2)$ be the cyclic group of order p generated by $\eta I_1 \oplus \eta^2 I_2$, where I_1 and I_2 denote identity matrices on complex Euclidean spaces of arbitrary dimension.

Example 1.3. Let η be a primitive 7-th root of unity. Let Γ be the cyclic group of order 7 generated by $\eta I_1 \oplus \eta^2 I_2 \oplus \eta^4 I_3$. Here each I_k denotes the identity matrix in an arbitrary dimension (not necessarily equal).

The most interesting case is perhaps Example 1.2, but Example 1.3 involves a new phenomenon we discuss in Section 5. In each case we will assume that the

source dimension is minimal. Hence in Example 1.1 the source dimension will equal one, and this situation is quite simple. In Example 1.2 the minimum source dimension is 2, and in Example 1.3 the minimum source dimension is 3. If we allow the identity matrices in the above examples to be in larger dimensions, the combinatorics becomes more difficult. The natural source dimensions in the three cases are 1, 2, 3 respectively.

Convention. We henceforth use the following notation: $\Gamma(m, 1)$ denotes the group in Example 1.1 where the source dimension equals 1 and $\Gamma(2r + 1, 2)$ denotes the group in Example 1.2 where the source dimension equals 2. Finally $\Gamma(7)$ denotes the group in Example 1.3 where the source dimension equals 3. We discuss the group from Example 1.1 in source dimension 2 in Section 4, where we explain and use the notation $\Gamma_2(m, 1)$. We discuss $\Gamma(7)$ in Section 5.

Remark 1.1. The groups $\Gamma(p, q)$, defined as in Example 1.2 in source dimension 2, but with the exponent 2 replaced by an exponent q relatively prime to p , fit to some extent in this story. They arise for sphere maps only when q can be chosen to equal 1 or 2. The condition (required for sphere mappings) that the coefficients be non-negative will thus be a crucial assumption.

For each of the above groups we are therefore interested in real polynomials satisfying the following three properties:

Definition 1.1. Let Γ denote any of the groups from Examples 1.1, 1.2, and 1.3. A real polynomial is Γ -**special** if

- f is Γ -invariant. In the first case $f(\eta x) = f(x)$ in source dimension 1 and $f(\eta x, \eta y) = f(x, y)$ in source dimension 2. In the second case $f(\eta x, \eta^2 y) = f(x, y)$. In the third case, $f(\eta x, \eta^2 y, \eta^4 z) = f(x, y, z)$. In each case η is a root of unity as in Examples 1.1, 1.2, and 1.3.
- In the first case, $f(1) = 1$. In the second case, $f(x, y) = 1$ on the line $x + y = 1$. In the third case, $f(x, y, z) = 1$ on the plane given by $x + y + z = 1$.
- The coefficients of f are non-negative, and its constant term is 0.

We say that f is a **good** polynomial if it satisfies the second and third items. Throughout this paper, the phrase “has non-negative coefficients” means “has **only** non-negative coefficients.”

When the group is understood we just say *special*. Our main result determines precise information on the gaps in the possible number of terms in special polynomials. See Theorem 1.2.

The following general uniqueness result holds for polynomials invariant under the group $\Gamma(p, q)$, but (unless $q = 1$ or $q = 2$) the coefficients must be of both signs. It is a special case of a result from 1992 appearing in [9].

Theorem 1.1. Put $\Gamma = \Gamma(p, q)$. There is a unique polynomial $\Phi(x, y) = \Phi_\Gamma(x, y)$ satisfying the following properties:

- Φ is Γ -invariant. Thus $\Phi(\eta x, \eta^q y) = \Phi(x, y)$.
- $\Phi(x, y) = 1$ on the line given by $x + y = 1$.
- $\Phi(0, 0) = 0$.
- Φ is of degree p .

Furthermore Φ is given by the following formula:

$$(1) \quad \Phi(x, y) = 1 - \prod_{j=1}^p (1 - \eta^j x - \eta^{qj} y).$$

Remark 1.2. The constant polynomial 1 satisfies the first two properties, but it is not important in our discussion. It is precluded by our assumption that $f(0, 0) = 0$. This assumption is required for uniqueness results.

Remark 1.3. By [9], a general uniqueness result holds for polynomials invariant under any fixed-point free subgroup Γ of $\mathbf{U}(n)$, and a formula analogous to (1) defines Φ_Γ . Except in the three cases noted, however, the coefficients of the corresponding real polynomials must be of both signs. Allowing coefficients of both signs arises when we map spheres to hyperquadrics, but will not be a major concern in this paper. For general q , combinatorial formulas for the integers arising in (1) appear in [23]. See also [15], [18], and [16] for additional information in this case.

Remark 1.4. For the groups $\Gamma(m, 1)$ and $\Gamma(2r+1, 2)$, the polynomial Φ_Γ includes all the non-constant invariant terms that generate the algebra of invariant polynomials. For the group $\Gamma(7)$, the polynomial Φ_Γ includes all of these terms and also several of their products. In general, for $\Gamma(p, q)$, not all the monomials generating the algebra appear in Φ_Γ . See [16].

We call Φ_Γ the **basic** polynomial for Γ . When $\Gamma = \Gamma(m, 1)$, the basic polynomial is given by x^m in source dimension 1 and by $(x + y)^m$ in source dimension 2. Applying Theorem 1.1 to $\Gamma(2r+1, 2)$ yields Corollary 1.1 below. Reference [4] includes an entire chapter on these polynomials, including proofs of the various statements made in the subsequent two paragraphs. See also [5] and [6] for further results. The basic polynomial for $\Gamma(7)$ appears in Proposition 1.1.

We recall some information about the group $\Gamma(2r+1, 2)$ in the following corollary of Theorem 1.1.

Corollary 1.1. *For each r there is a unique polynomial $f(x, y) = f_{2r+1}(x, y)$ satisfying the following properties:*

- f is $\Gamma(2r+1, 2)$ -invariant. Thus $f(\eta x, \eta^2 y) = f(x, y)$.
- $f(x, y) = 1$ on the line $x + y = 1$.
- The coefficients of f are non-negative and $f(0, 0) = 0$.
- f is of degree $2r+1$.

The polynomial $f_{2r+1}(x, y)$ has the following explicit formulas:

$$(2.1) \quad f_{2r+1}(x, y) = \left(\frac{x + \sqrt{x^2 + 4y}}{2} \right)^{2r+1} + \left(\frac{x - \sqrt{x^2 + 4y}}{2} \right)^{2r+1} + y^{2r+1}$$

$$(2.2) \quad f_{2r+1}(x, y) = x^{2r+1} + y^{2r+1} + \sum_{j=1}^r c(r, j) x^{2r+1-2j} y^j.$$

The coefficients $c(r, j)$ are positive integers that satisfy

$$(2.3) \quad c(r, j) = \frac{2r+1}{j} \binom{2r-j}{j-1}.$$

This family of polynomials f_{2r+1} satisfies many intriguing properties, of which we mention several here. The polynomial $f_{2r+1}(x, y)$ is congruent to $x^{2r+1} + y^{2r+1} \pmod{2r+1}$ if and only if $2r+1$ is either prime or 1. The family of polynomials satisfies a third-order recurrence relation and has an explicit rational generating function. See [4] for more information.

The number of terms in a polynomial will be of particular importance. Given a polynomial f (in any number of variables), $N(f)$ denotes the number of distinct monomials arising in f . By [8], the polynomials f_{2r+1} satisfy the following optimization property, or sharp **degree estimate**. Consider the collection of good polynomials f of degree $d(f)$. Thus $f(x, y) = 1$ on the line $x + y = 1$, the coefficients of f are non-negative, and $f(0, 0) = 0$. Then $d(f) \leq 2N(f) - 3$ and equality holds for the polynomial f_{2r+1} , where $N = r + 2$. The real polynomial f_{2r+1} is the squared Euclidean norm of a holomorphic polynomial $p(z, w) : \mathbb{C}^2 \rightarrow \mathbb{C}^{r+2}$ that maps the unit sphere in the source to the unit sphere in the target. The number of terms is the minimum embedding dimension of such a map, and thus the sharp degree estimate $d \leq 2N - 3$ holds for these polynomials. By [21], the sharp degree estimate $d \leq \frac{N-1}{n-1}$ holds in dimensions at least 3. These bounds therefore limit the number of unknown coefficients one must consider.

We mention the formula for Φ_Γ for the group $\Gamma(7)$, where the source dimension is 3. Write the variables as (x, y, z) . This formula appears in both [2] and [9].

Proposition 1.1. *The basic polynomial for the group $\Gamma(7)$ is given by*

$$(3) \quad \begin{aligned} \Phi(x, y, z) = & x^7 + y^7 + z^7 + 14(x^3y^2 + x^2z^3 + y^3z^2 + xyz) \\ & + 7(x^5y + xz^5 + y^5z + xy^3 + x^3z + yz^3) \\ & + 7(xy^2z^4 + x^2y^4z + x^4yz^2 + x^2y^2z^2) \end{aligned}$$

The main result in this paper is the following theorem on the gaps for invariant polynomials. The first part of the theorem is easy. The second part answers a question from [1]. The third part is completely new.

Theorem 1.2. *Assume f is a Γ -special polynomial; thus it satisfies the properties from Definition 1.1 for the group Γ . The following statements hold and are sharp.*

- For $\Gamma(m, 1)$, the number of terms $N(f)$ is an arbitrary non-negative integer.
- For $\Gamma(2r + 1, 2)$, the number of terms $N(f)$ can be either $r + 2$ or any positive integer at least $2r + 3$. No other values are possible.
- For $\Gamma(7)$, the situation is considerably more complicated. See Theorem 5.1.

In Section 6 we interpret these results in terms of sparse solutions to linear systems. Finding special polynomials amounts to solving a linear system for the coefficients, and finding the minimum number of possible non-zero coefficients seeks the sparsest solution. Gaps in the possible numbers of non-zero coefficients have a related interpretation.

In Section 7 we note several possible related directions of research.

The main new results in this paper are the proof of the second statement in Theorem 1.1 and almost all of Section 5, especially Theorem 5.1 and the methods used to obtain it. The authors acknowledge Franc Forstneric for useful discussions from years ago, Xiaojun Huang for his work on related problems, Jiri Lebl for his contributions to this area, and Ming Xiao for more recent conversations.

2. INVARIANCE AND $N(f)$

The following lemma about the number of terms in polynomials is obvious.

Lemma 2.1. *Suppose F_1 and F_2 are polynomials with no common terms. Then $N(F_1 + F_2) = N(F_1) + N(F_2)$. In particular, if all the monomials in F_2 are of higher degree than those of F_1 , then $N(F_1 + F_2) = N(F_1) + N(F_2)$.*

Let \mathbf{x} denote either (x, y) or (x, y, z) . We will be considering polynomials that equal 1 on the sets $x + y = 1$ or $x + y + z = 1$. In the first case such a polynomial must be divisible by $x + y - 1$, and in the second case by $x + y + z - 1$.

When the polynomial is invariant, it must be also divisible by the terms obtained by replacing the variable \mathbf{x} by $\gamma\mathbf{x}$, for each γ in the group. Hence we have the following result.

Proposition 2.1. *Assume G is special for Γ . Then there is an invariant polynomial H such that $G = \Phi_\Gamma - H + H \cdot \Phi_\Gamma$.*

Proof. Since $G - \Phi_\Gamma$ is invariant and vanishes where $\Phi_\Gamma = 1$, it is divisible by $\Phi_\Gamma - 1$. Since it is invariant, the quotient H also is and thus $G - \Phi_\Gamma = H \cdot (\Phi_\Gamma - 1)$. \square

The operation of replacing F with $F - H + FH$ is a special case of a partial tensor product operation introduced by the first author in his study of rational sphere maps. See [4]. We will use this operation throughout. The condition that our polynomials have non-negative coefficients requires care. We write $g \preceq h$ when all the coefficients of $h - g$ are non-negative. Suppose $0 \preceq F$. Put $G = F - H + HF$. If $0 \preceq H \preceq F$, then obviously $0 \preceq G$. It is possible, however, that $0 \preceq G$, but that $0 \not\preceq F - H$ is false.

Example 2.1. Put $F = \Phi_\Gamma$ for any of the three groups. For each k , the polynomial F^k is special. For $k \geq 3$, put $H = F + F^2 + \dots + F^{k-1}$. Then **none** of the coefficients of $F - H$ are positive, and yet

$$F - H + HF = F - (F + F^2 + \dots + F^{k-1}) + (F^2 + F^3 + \dots + F^k) = F^k.$$

We remind the reader that our primary purpose is to determine precisely the possible values of $N(G)$ for special G . We begin with two easy lemmas.

Lemma 2.2. *Suppose $F = 1$ on the line defined by $x + y = 1$. Let H be an arbitrary polynomial. Put $G = F - H + FH = F + H(F - 1)$. Then $G = 1$ on this line. If F and H are group invariant, then so is G . Analogous conclusions hold when $F = 1$ on the plane $x + y + u = 1$.*

Lemma 2.3. *Suppose F has non-negative coefficients. Let H consist of some terms of F , with corresponding coefficients that are positive but strictly smaller than those of F . Then $N(F - H) = N(F)$.*

Proposition 2.2. *Given a good polynomial f and a non-negative integer a , there are good polynomials g and h such that*

$$N(g) = aN(f)$$

$$N(h) = aN(f) - (a - 1).$$

When f is group-invariant we may choose g and h to be invariant.

Proof. Given f , find a monomial m in f of largest degree. For $0 < \lambda < 1$, we form

$$g = (f - \lambda m) + \lambda m f,$$

$$h = (f - m) + m f.$$

By the above lemmas, $N(g) = N(f) + N(f) = 2N(f)$ and $N(h) = 2N(f) - 1$. Iterating each of these operations (always on a monomial of largest degree), we obtain the conclusion. The invariance follows because f is invariant if and only if each monomial in f is invariant. \square

The proof yields the following useful result, which we call the Frobenius property or the postage stamp property.

Proposition 2.3. *Given special polynomials f and g , and non-negative integers a, b , there is a special polynomial H with $N(H) = aN(f) + bN(g)$. Furthermore, if h any special polynomial, then there is a special polynomial H with*

$$(4) \quad N(H) = N(h) + aN(f) + bN(g).$$

Proof. Choose a monomial m in h of highest degree, and (for $0 < t < 1$), define h^* by $h^* = h - tm + tmf$. By Lemmas 2.2 and 2.3, h^* is invariant, equals 1 on the line $x + y = 1$, and has non-negative coefficients. Furthermore, $N(h^*) = N(h) + N(f)$. Iterating this process, each time on a monomial of highest degree, after a steps we obtain a polynomial h^{**} with $N(h^{**}) = N(h) + aN(f)$. Now do the same using g to obtain a special polynomial H satisfying (4). The first statement follows from the second by using $h = f$ and replacing a by $a - 1$. \square

Proposition 2.4. *Let $p(t)$ be a polynomial of the form*

$$p(t) = 1 + \sum_{j=1}^K c_j t^j$$

where each $c_j \geq 0$. Suppose at least k of the c_j are positive. Put $F(t) = (1 + t)^m$. Then $N(Fp) \geq m + 1 + k$.

Proof. The product Fp contains the $m + 1$ monomials t^j for $0 \leq j \leq m$. Suppose a_1, \dots, a_k are distinct indices with $c_{a_i} > 0$. Then Fp also contains the monomials whose exponents are $m + a_1, \dots, m + a_k$. Hence there are at least $m + 1 + k$ terms. \square

The product can contain a larger number of monomials, but no better estimate is possible. If $p(t) = (1 + t)^k$, for example, then $N(Fp) = N((1 + t)^{m+k}) = m + k + 1$.

Example 2.2. Put $F(t) = (1 + t)^m$ for $m \geq 2$. Consider $p(t) = 1 + t + t^{m+1}$. Then $k = 2$ but $N(Fp) = 2m + 2 > m + 3$.

Remark 2.1. The proof of Theorem 1.2 is easy for $\Gamma(m, 1)$ in source dimension 1. We pause to make this observation now. Given an integer m at least 2, let η be a primitive m -th root of unity. For $\lambda_j > 0$ and $\sum_{j=1}^d \lambda_j = 1$, put

$$p(x) = \sum_{j=1}^d \lambda_j x^{mj}.$$

Then $p(\eta x) = p(x)$, its coefficients are non-negative, $p(1) = 1$, and p has no constant term. Thus p is **special** for $\Gamma(m, 1)$ in source dimension one, and $N(p) = d$. Since d is an arbitrary positive integer, there are no restrictions on $N(p)$ for p special.

We next describe the invariant monomials for each of our three classes of groups. In any source dimension, the algebra of non-constant monomials invariant under $\Gamma(m, 1)$ is generated by the constant 1 and the homogeneous polynomials of degree m . The degree of an invariant polynomial with $f(\mathbf{0}) = 0$ must be a multiple of m .

The algebra of polynomials invariant under $\Gamma(2r + 1, 2)$ is generated by 1, the monomials x^{2r+1} , y^{2r+1} , and for $1 \leq j \leq r$, the monomials $x^{2r+1-2j}y^j$.

For both of these classes of groups, the terms of the basic polynomial Φ_Γ are the basis elements of the algebra. That no longer holds for the group $\Gamma(7)$. Several products are required. For the third group, with variables x, y, z , the algebra of invariant polynomials is generated by the following:

$$x^7, y^7, z^7, x^5y, x^3y^2, xy^3, x^3z, y^5z, y^3z^2, yz^3, xyz.$$

But the basic polynomial from Proposition 1.1 has 17 terms, including for example $x^2y^2z^2$ and x^4yz . This situation, which was mentioned in Remark 1.4, changes the behavior for the gaps in possible values for $N(f)$.

For $\Gamma(2r + 1, 2)$ the next easy result follows by combining the lemmas above.

Proposition 2.5. *Fix r . Let $F = f_{2r+1}$ denote the basic polynomial defined by Theorem 1.1. Formula (2.3), for certain positive integers $c(r, j)$, writes F as:*

$$F(x, y) = x^{2r+1} + y^{2r+1} + \sum_{j=1}^r c(r, j) x^{2r+1-2j} y^j.$$

Let H be defined by $\sum_{j=1}^r \lambda_j x^{2r+1-2j} y^j$ where $0 \leq \lambda_j < c(r, j)$ for each j . Put $G = F - H + FH$. Then $N(G) = N(F) + N(FH)$.

Proof. Note that all terms in FH are of higher degree than those of F . Thus Lemma 2.1 implies $N(G) = N(F - H) + N(FH) = N(F) + N(FH)$. \square

We end this section with a warning. When the coefficients of polynomials g, h can be of both signs, there is no simple way to relate $N(gh)$ to $N(g)$ and $N(h)$.

Example 2.3. We give several examples.

- Put $g(x) = x^2 + 1 + \sqrt{2}x$ and $h(x) = x^2 + 1 - \sqrt{2}x$. Then $N(g) = N(h) = 3$, but $N(gh) = 2$.
- Put $g(x, y) = x - y$ and $h(x, y) = x^m + x^{m-1}y + \cdots + y^m$. Then their product $x^{m+1} - y^{m+1}$ has only two terms.
- There are polynomials p in one variable such that p^2 has fewer terms than p . See [3]. (Homogenizing then gives examples of homogeneous polynomials in two variables with this property.)

3. PROOF OF FIRST TWO PARTS OF THEOREM 1.2

Proof. In Remark 2.1 we have already observed the result for $\Gamma(m, 1)$ in source dimension 1. We next give the proof for $\Gamma(2r + 1, 2)$. Let F denote the basic map $\Phi_\Gamma = f_{2r+1}$. Recall that $N(F) = r + 2$ and that

$$F(x, y) = x^{2r+1} + y^{2r+1} + \sum_{j=1}^r c(r, j) x^{2r+1-2j} y^j.$$

We first show that there is a special polynomial G with $2r + 3$ terms. In fact there are $r + 2$ such examples.

We choose precisely one of the terms $c(r, j)x^{2r+1-2j}y^j$ and call it H . Thus $G = F - H + FH$, where

$$G(x, y) = F(x, y) - c(r, j)x^{2r+1-2j}y^j + (F(x, y))c(r, j)x^{2r+1-2j}y^j.$$

We have multiplied a term of degree $2r + 1 - j$ by F , and hence $d(G)$ satisfies

$$d(G) = 2r + 1 - j + (2r + 1) = 4r + 2 - j \leq 4r + 1.$$

Since $F(x, y) = 1$ on $x + y = 1$, so does G . Each monomial occurring in G is invariant. The non-negativity of the coefficients is clear; since the coefficient of $x^{2r+1-2j}y^j$ in F is precisely $c(r, j)$, we have eliminated one term and replaced it by $r + 2$ terms with positive coefficients. We claim the resulting terms are distinct. Except for the term Hy^{2r+1} , all the exponents of x in each of the new terms is even, and the exponents of x in the old terms are all odd. The term Hy^{2r+1} is of degree larger than $2r + 1$ and hence it is also new. Thus G has $(r + 1) + (r + 2) = 2r + 3$ terms (the old plus the new). This construction works for each j with $1 \leq j \leq r$ and yields a different G . Hence there are at least r such polynomials. We could also use the monomials x^{2r+1} and y^{2r+1} for H to obtain two more examples with $2r + 3$ terms.

Next we show how to get $N = 2r + 4$. The idea is similar, but this time we choose a number λ with $0 < \lambda < c(r, j)$ and form

$$G(x, y) = F(x, y) - \lambda x^{2r+1-2j}y^j + (F(x, y))\lambda x^{2r+1-2j}y^j.$$

For this polynomial G , we have $N(G) = (r + 2) + (r + 2) = 2r + 4$.

To obtain higher values, we use $k - j + 1$ consecutive terms to define H . Thus we choose λ_s such that $0 < \lambda_s < c(r, s)$. Put $G = F - H + FH$, where

$$H(x, y) = \lambda_j x^{2r+1-2j}y^j + \dots + \lambda_k x^{2r+1-2k}y^k.$$

Now $F - H$ has $r + 2$ terms, and HF has $(r + 2 + k - j)$ additional terms. Thus $N(G) = (r + 2) + (r + 2 + k - j)$. We get all possible values for $N(G)$ for G of degree up to $4r + 2$ in this way. To obtain higher values, we use these invariant polynomials in the same manner.

Suppose that $N(G) \leq 2r + 2$. We wish to show that $G = F$. By the degree estimate $d \leq 2N - 3$ from [8], if $N(g) \leq 2r + 2$, then

$$d(G) \leq 2(2r + 2) - 3 = 4r + 1.$$

If H contained a term of degree $2r + 1$, then HF and hence G would have a term of degree $4r + 2$. But $d(G) \leq 4r + 1$, and thus $d(H) < 2r + 1$. Since H is invariant, and H contains neither x^{2r+1} nor y^{2r+1} ,

$$(5) \quad H(x, y) = \sum_{j=1}^r \lambda_j x^{2r+1-2j}y^j.$$

To preserve non-negativity of the coefficients of G , we must have $0 \leq \lambda_j \leq c(r, j)$ for each j . Thus we have found the general special polynomial of at most this degree.

As before, write $G = F - H + FH$. If all the λ_j in (5) are 0, we obtain $G = F$ and $N(G) = r + 2$. Otherwise, find the largest j for which $\lambda_j > 0$. If $\lambda_j = c(r, j)$, then $N(G) \geq N(F) - 1 + N(F)$. If $\lambda_j < c(r, j)$, then $N(G) \geq N(F) + N(F)$. In either case, we obtain $N(G) \geq 2(r + 2) - 1 = 2r + 3$. Thus $N(G)$ cannot lie in the integer interval $[r + 3, 2r + 2]$. We have already shown that $N(G)$ must be at least

$r + 2$ and that this value is possible. We have shown that each value at least $2r + 3$ is possible. Thus we have proved the theorem for $\Gamma(2r + 1, 2)$. \square

Example 3.1. Put $r = 5$. Then $F = f_{2r+1}$ satisfies

$$(6) \quad F(x, y) = x^{11} + y^{11} + 11x^9y + 44x^7y^2 + 77x^5y^3 + 55x^3y^4 + 11xy^5.$$

Put $H = 44x^7y^2 + 77x^5y^3 + 55x^3y^4$ and put $G = F - \frac{H}{11} + \frac{FH}{11}$. Then G equals:

$$\begin{aligned} x^{11} + 11x^9y + 40x^7y^2 + 4x^{18}y^2 + 70x^5y^3 + 51x^{16}y^3 + 50x^3y^4 + 258x^{14}y^4 + 11xy^5 + 671x^{12}y^5 \\ + 979x^{10}y^6 + 814x^8y^7 + 352x^6y^8 + 55x^4y^9 + y^{11} + 4x^7y^{13} + 7x^5y^{14} + 5x^3y^{15}. \end{aligned}$$

The polynomial G has 18 terms; 7 come from F and 11 more arise from multiplying three consecutive terms by F , creating $7 + 4 = 11$ more terms.

4. THE GROUP $\Gamma(m, 1)$ IN SOURCE DIMENSION 2

We consider the group $\Gamma(m, 1)$ when the source dimension is 2.

Theorem 4.1. *Suppose G is special for the group $\Gamma(m, 1)$ and G is of degree Km . Then there are homogeneous polynomials H_k such that*

$$G = F - H_m + FH_m - H_{2m} + \cdots + FH_{(K-1)m} - H_{Km} + FH_{Km}.$$

If the H_{jm} have non-negative coefficients, then $N(G) \geq Km + 1$.

Proof. The result holds when $K = 1$ because the only special polynomial of degree m is the basic polynomial F , namely $(x + y)^m$, which has $m + 1$ terms. Otherwise, by Proposition 2.1, we write $G = (F - H) + HF$ for some invariant H . Note that $F(0, 0) = G(0, 0) = 0$ implies $H(0, 0) = 0$. Write

$$H = H_m + H_{2m} + \cdots + H_{(K-1)m}$$

in terms of its homogeneous expansion. The term H_{Km} must vanish, because otherwise G would be degree $(K + 1)m$. From $G = (F - H) + HF$, we obtain

$$G = (F - H_m) + (FH_m - H_{2m}) + \cdots + (FH_{(K-2)m} - H_{(K-1)m}) + FH_{(K-1)m}.$$

Each of the terms in parentheses is of different degree and hence $N(G)$ equals the following sum:

$$N(F - H_m) + N(FH_m - H_{2m}) + \cdots + N(FH_{(K-2)m} - H_{(K-1)m}) + N(FH_{(K-1)m}).$$

Also each term has non-negative coefficients. We group the terms in this sum in three parts: the first term, the last term, and the middle terms. Thus

$$N(G) = N(F - H_m) + \sum_{j=1}^{K-2} N(FH_{jm} - H_{(j+1)m}) + N(FH_{(K-1)m})$$

Let $a_j = N(H_{mj})$. Since $H_{(K-1)m} \neq 0$, Proposition 2.4 implies that

$$N(FH_{(K-1)m}) \geq m + a_{m-1}.$$

For the first term, we have

$$N(F - H_m) \geq N(F) - a_1 = m + 1 - a_1.$$

For the other terms, again by Proposition 2.4, we have

$$N(FH_{jm} - H_{(j+1)m}) \geq N(FH_{jm}) - a_{j+1} \geq m + a_j - a_{j+1}.$$

The sum telescopes. Adding all the inequalities together yields

$$N(G) \geq (m+1) + m(K-2) + m = Km + 1.$$

□

The restrictive assumption that the H_j have non-negative coefficients implies Theorem 1.2 for the group $\Gamma(m, 1)$ in source dimension 2. First we give an example showing that it can happen that some H_j have at least one negative coefficient and yet G is special.

Example 4.1. Put $H_2(x, y) = x^2 - xy + y^2$. Define $G(x, y)$ by

$$G(x, y) = (x+y)^2 - (x^2 - xy + y^2) + (x+y)^2(x^2 - xy + y^2) = x^4 + 3xy + x^3y + xy^3 + y^4.$$

Thus G has non-negative coefficients, $G = F - H_2 + FH_2$, and H_2 has a negative coefficient.

We prove our result for $\Gamma(m, 1)$ under the restrictive condition that the polynomials H_{jm} are non-negative.

Proof. Let G be special for $\Gamma(m, 1)$. Then G is of degree Km for some positive integer K . By Theorem 4.1, $N(G) = Km + 1$. When $K = 1$, the only map is F , and $N(F) = m + 1$. If $K \geq 2$, then $N(G) \geq Km + 1 \geq 2m + 1$; it follows that $N(G)$ is in neither the integer interval $[1, m]$ nor the interval $[m + 2, 2m]$.

It remains to show that all larger integers are possible, even allowing this restrictive assumption. We construct examples with N terms for $N \geq 2m + 1$. To obtain an example where $N = 2m + 1$, we put

$$G(x, y) = (x + y)^m - x^m + x^m(x + y)^m = F - H + HF.$$

Then $N(G) = (m) + (m + 1) = 2m + 1$. To obtain an example where $N = 2m + 2$, we put for $0 < \lambda < 1$,

$$G(x, y) = (x + y)^m - \lambda x^m + \lambda x^m(x + y)^m$$

and hence $N(G) = (m + 1) + (m + 1) = 2m + 2$. In general, let H be the first k terms in the expansion of $(x + y)^m$ and let $0 < \lambda < 1$. Again put $G = F - \lambda H + \lambda FH$. Then $N(G) = (m + 1) + (m + k)$. This procedure realizes all N up to $3m + 2$. For any G with N terms, we may choose a monomial h of highest degree in G , and replace G with $G - h + hG$, and achieve $N(G) + m + 1$ terms. If we replace G with $G - \lambda h + \lambda hG$, then we achieve $N(G) + m + 2$ terms. Thus we can achieve everything in the interval $2m + 1 \leq N \leq 4m + 3$. Continuing in this way, or by using the Frobenius property, we obtain examples with N terms for each N at least $2m + 1$. We conclude that $N(G)$ cannot satisfy $1 \leq N(G) \leq m$ nor $m + 2 \leq N(G) \leq 2m$, but is otherwise arbitrary.

□

Remark 4.1. When one allows H to have negative coefficients, interesting cancellations can occur. The second author has an example where $p = 4$ and the target dimension $N = 8$, so the bound in Theorem 4.1 does not hold in general.

5. THE GROUP $\Gamma(7)$.

The squared Euclidean norm of the minimal invariant sphere mapping for this group is a real polynomial F analogous to the polynomials $(x+y)^m$ and $f_{2r+1}(x,y)$ we have considered earlier. By Proposition 1.1 the basic polynomial is given by

$$(7) \quad \begin{aligned} F(x, y, z) = & x^7 + y^7 + z^7 + 14(x^3y^2 + x^2z^3 + y^3z^2 + xyz) \\ & + 7(x^5y + xz^5 + y^5z + xy^3 + x^3z + yz^3) \\ & + 7(xy^2z^4 + x^2y^4z + x^4yz^2 + x^2y^2z^2). \end{aligned}$$

The invariance is evident: $F(\eta x, \eta^2 y, \eta^4 z) = F(x, y, z)$. The connection to the group $\Gamma(7, 2)$ is also evident. More subtle is that monomials other than the generators of the algebra of invariant polynomials arise in F . For example, $x^2y^2z^2$ is the square of the basic invariant monomial xyz . This fact impacts the number of terms in a subtle way. Note also that the restriction to each of the two dimensional subspaces defined by setting a coordinate equal to 0 gives the map f_7 , but with a different choice of seventh root of unity. Thus we see where the coefficients of 7 and 14 arise. It is a bit surprising that these integers are also coefficients of the terms that are products of all three variables. We note that F has 6 terms of degree seven, 4 terms of degree six, 3 terms of degree five, 3 terms of degree four, and 1 term of degree three.

Let $G : \mathbb{R}^3 \rightarrow \mathbb{R}$ be a Γ -invariant polynomial with non-negative coefficients. Suppose $G(x, y, z) = 1$ on the plane defined by $x + y + z = 1$ and $G(0, 0, 0) = 0$. We are interested in the possible values for $N(G)$. Proposition 2.1 provides a method to construct such G . Begin with F and let m be any collection of terms of F such that all the coefficients of both m and $F - m$ are non-negative. Thus $0 \preceq m \preceq F$. Then the polynomial $F - m + mF$ satisfies all of our conditions. More generally, if G satisfies all of our conditions, then $G - p + pG$ also does as long as $0 \preceq p \preceq G$. All examples are constructed in this fashion. Things are subtle because a term might not satisfy $p \preceq F$, but does satisfy $p \preceq F - m + mF$.

The mapping F from (7) consists of 17 terms all of degree at most 7. There can be no smaller number of terms because each invariant polynomial that equals 1 on the plane must be divisible by $F - 1$. Thus $1 \leq N(G) \leq 16$ is impossible.

Put $H(x, y, z) = 14xyz$. Consider the polynomial $G = F - H + HF$. We have multiplied each term in F by the same monomial, and hence $N(HF) = 17$. But $F - H$ and HF overlap because there are four monomials in F of degree at most 4, namely xyz, x^3z, xy^3, yz^3 . When these get multiplied by xyz , we get no new monomials. The monomial xyz is missing from G . Therefore

$$N(G) = (17 - 1) + (17 - 4) = 29.$$

We continue by listing maps G for which $N(G)$ takes on all the known possible values up to 57. In each case, the map G has the form $G = F - H + HF$ where H is some invariant polynomial. Rather than writing all the lengthy formulas for G , we write the (much shorter) formulas for H . Given $k \geq 58$, the postage stamp principle implies that examples for which $N(G) = k$ exist.

- Put $H = 0$. Then $G = F$ and $N(G) = 17$.
- Put $H = 14xyz$. Then $N(G) = 29$.
- Put $H = \lambda 14xyz$ for $0 < \lambda < 1$. Then $N(G) = 30$.
- Put $H = 7x^3z$. Then $N(G) = 32$.

- Put $H = \lambda 7x^3z$ for $0 < \lambda < 1$. Then $N(G) = 33$.
- Put $H = \lambda z^7$ for $0 < \lambda < 1$. Then $N(G) = 34$.
- Put $H = 7xy^3 + 14xyz$. Then $N(G) = 37$.
- Put $H = 7xy^3 + \lambda 14xyz$ for $0 < \lambda < 1$. Then $N(G) = 38$.
- Put $H = \lambda(7xy^3 + 14xyz)$ for $0 < \lambda < 1$. Then $N(G) = 39$.
- Put $H = 14xyz + 14x^3y^2$. Then $N(G) = 40$.
- Put $H = 14xyz + 203x^2y^2z^2$. Then $N(G) = 41$.
- Put $H = 7xy^3 + 7yz^3$. Then $N(G) = 42$.
- Put $H = 7xy^3 + 14xyz + 7yz^3$. Then $N(G) = 43$.
- Put $H = 7xy^3 + \lambda 14xyz + 7yz^3$ for $0 < \lambda < 1$. Then $N(G) = 44$.
- Put $H = \lambda 7xy^3 + \lambda 14xyz + 7x^3z$ for $0 < \lambda < 1$. Then $N(G) = 45$.
- Put $H = \lambda(7xy^3 + 14xyz + 7yz^3)$ for $0 < \lambda < 1$. Then $N(G) = 46$.
- Put $H = 7xy^3 + 7x^3z + 14xyz + 7yz^3$. Then $N(G) = 47$.
- Put $H = 7xy^3 + 7x^3z + \lambda 14xyz + 7yz^3$ for $0 < \lambda < 1$. Then $N(G) = 48$.
- Put $H = \lambda 7xy^3 + 7x^3z + \lambda 14xyz + 7yz^3$ for $0 < \lambda < 1$. Then $N(G) = 49$.
- Put $H = \lambda 7xy^3 + 7x^3z + \lambda 14xyz + \lambda 7yz^3$ for $0 < \lambda < 1$. Then $N(G) = 50$.
- Put $H = \lambda(7xy^3 + 7x^3z + 14xyz + 7yz^3)$ for $0 < \lambda < 1$. Then $N(G) = 51$.
- Put $H = 14x^3y^2 + 7xy^3 + 7x^3z + 14xyz + \lambda 14x^2z^3$ for $0 < \lambda < 1$. Then $N(G) = 52$.
- Put $H = 14x^3y^2 + 7xy^3 + 7x^3z + \lambda 14xyz + \lambda 14x^2z^3$ for $0 < \lambda < 1$. Then $N(G) = 53$.
- Put $H = \lambda 14x^3y^2 + 7xy^3 + 7x^3z + \lambda 14xyz + \lambda 14x^2z^3$ for $0 < \lambda < 1$. Then $N(G) = 54$.
- Put $H = \lambda 14x^3y^2 + \lambda 7xy^3 + 7x^3z + \lambda 14xyz + \lambda 14x^2z^3$ for $0 < \lambda < 1$. Then $N(G) = 55$.
- Put $H = \lambda(14x^3y^2 + 7xy^3 + 7x^3z + 14xyz + 14x^2z^3)$ for $0 < \lambda < 1$. Then $N(G) = 56$.
- Put $H = \frac{x^7}{2} + 13xyz + 182x^2y^2z^2$. Then $N(G) = 57$.

Remark 5.1. The map for which $N(G) = 41$ exhibits an interesting phenomenon. In this example, $F - 203x^2y^2z^2 + (203x^2y^2z^2)F$ has the term $-196x^2y^2z^2$. But when we subtract the term $14xyz$ and add back in $14xyzF$, this negative coefficient goes away. The arithmetic comes from $7 - 203 = -196 = -(14)^2$. Hence all coefficients of G are non-negative. If we perform the operations in the opposite order, we never see a negative coefficient! The map for which $N(G) = 57$ also illustrates this phenomenon. It occurs because the basic polynomial F includes terms such as $x^2y^2z^2$ that are not basis elements of the algebra of invariant polynomials.

Example 5.1. To illustrate non-uniqueness, we give a different way to achieve $N(G) = 51$. Put $H = 14x^3y^2 + 7xy^3 + 7x^3z + 14xyz + 14x^2z^3$. Then $N(G) = 51$. The example above used fewer terms in the polynomial H .

Let G be special for $\Gamma(7)$. We will show, unless $N(G) = 17$, it must be true that $N(G) \geq 29$. We recall the degree estimate from [21]; when $n \geq 3$, we have $d \leq \frac{N-1}{n-1}$. Here $n = 3$ and hence $N \geq 2d + 1$. To show that $N(G) \geq 29$ for all special polynomials G , it suffices to prove the result for G of degree at most 13.

By Proposition 1.1, the basic polynomial F for the group $\Gamma(7)$ is given by (7). Therefore there are no special polynomials of degree less than 7. We will show that there are none of degree 8 or 9 and that there is a one-parameter family of degree 10. We then go on to study degrees 11, 12, 13.

Let $G : \mathbb{R}^3 \rightarrow \mathbb{R}$ be a $\Gamma(7)$ -invariant special polynomial. By Proposition 2.1, $G = F + H(F - 1) = F - H + HF$ where H is a $\Gamma(7)$ -invariant polynomial. We first show the simple statement that the degree of G must be at least 10 unless $G = F$.

Lemma 5.1. *Let G be a $\Gamma(7)$ -invariant special polynomial. Then either $G = F$ or $\deg(G) \geq 10$.*

Proof. Each non-constant invariant monomial invariant under $\Gamma(7)$ has degree at least 3. Hence, if $H \neq 0$, we must have

$$\deg(G) = \deg(FH) = 7 + \deg(H) \geq 10.$$

□

Corollary 5.1. *There is no invariant example G with $18 \leq N(G) \leq 20$.*

Proof. By the lemma, either $G = F$ and $N(G) = 17$, or $N(G) \geq 2(10) + 1 = 21$. □

Next we find all examples of degree 10. The only possibility is that $H = \lambda xyz$ for some $\lambda \neq 0$. Since all the coefficients of G are positive, in fact $0 < \lambda \leq 14$. The resulting maps have the form

$$G(x, y, z) = F(x, y, z) + \lambda xyz F(x, y, z) - \lambda xyz.$$

Then $N(G)$ is 29 or 30, depending on whether $\lambda = 14$.

Remark 5.2. It is useful to think of the collection \mathbf{S}_{10} of all examples of degree at most 10. We identify this set with the closed interval $[0, 14]$ on the real line. For larger degrees d , the set of examples \mathbf{S}_d of degree at most d will always be a compact convex subset of some \mathbf{R}^m . One must systematically study $N(G)$ for G in the boundary of \mathbf{S}_d . When the degree is 10, the dimension m equals 1 and the boundary consists of $\lambda = 0$ and $\lambda = 14$. When $\lambda = 0$, $N(G) = 17$. When $\lambda = 14$, $N(G) = 29$, and otherwise $N(G) = 30$.

The authors regard degree 11 as a particularly illustrative case; things are simple enough to be illuminating, but complicated enough to exhibit the key issues. We describe all examples of degree 11. From the HF term, we must have $\deg(H) \leq 4$ since F has degree 7. Thus, we can give the general form of H by taking all non-constant $\Gamma(7)$ -invariant monomials of degree at most 4; we put

$$H(x, y, z) = Uxyz + Byz^3 + Cx^3z + Dxy^3$$

The set \mathbf{S}_{11} is a compact subset of \mathbb{R}^4 . We list all the terms in G in terms of the coefficients of H :

z^7	x^7	y^7
$14y^3z^2$	$14x^2z^3$	$14x^3y^2$
$7y^5z$	$7xz^5$	$7x^5y$
$(7 - B)yz^3$	$(7 - C)x^3z$	$(7 - D)xy^3$
$7By^2z^6$	$7Cx^6z^2$	$7Dx^2y^6$
$14By^4z^5$	$14Cx^5z^4$	$14Dx^4y^5$
$7By^6z^4$	$7Cx^4z^6$	$7Dx^6y^4$
Byz^{10}	$Cx^{10}z$	Dxy^{10}

By^8z^3	Cx^3z^8	Dx^8y^3
$(7B + D)xy^3z^7$	$(B + 7C)x^7yz^3$	$(C + 7D)x^3y^7z$
$(7B + 7D)x^2y^5z^4$	$(7B + 7C)x^4y^2z^5$	$(7C + 7D)x^5y^4z^2$
$(7 + 7U + 14B)xy^2z^4$	$(7 + 7U + 14C)x^4yz^2$	$(7 + 7U + 14D)x^2y^4z$
$(U + 7B)xyz^8$	$(U + 7C)x^8yz$	$(U + 7D)xy^8z$
$(7U + 14B)x^2yz^6$	$(7U + 14C)x^6y^2z$	$(7U + 14D)xy^6z^2$
$(14U + 7B + 7D)xy^4z^3$	$(14U + 7B + 7C)x^3yz^4$	$(14U + 7C + 7D)x^4y^3z$
$(7U + 7B + 7D)x^2y^3z^5$	$(7U + 7B + 7C)x^5y^2z^3$	$(7U + 7C + 7D)x^3y^5z^2$
$(14 - U)xyz$	$(14U + 7)x^2y^2z^2$	$(7U + 14B + 14C + 14D)x^3y^3z^3$

We have listed the terms in three columns. The last row is exceptional. In each other row, the three columns are isomorphic. One sends x to y , y to z , z to x , and one sends B to C , C to D , and D to B . This symmetry is a consequence of the group invariance, and something analogous holds in all degrees.

Note also that the coefficient U plays a different role than B, C, D . The non-negativity of *all* the coefficients puts upper and lower bounds on U, B, C, D . All the coefficients of G must be non-negative, and hence the lower bounds for B, C, D are 0. But U can be negative.

We first discuss the case $U < 0$. Each of $U + 7B, U + 7C, U + 7D$ is a coefficient. If $U < 0$, then we must have $B, C, D > 0$. Of the coefficients, B occurs alone twice, $7B$ alone twice, and $14B$ alone once. That gives 5 positive coefficients. By the symmetry in B, C, D , we see that there are at least 15 positive coefficients of this type. The rows beginning with $7B + D$ and $7B + 7D$ create 6 more terms. The first three rows generate 9 more. Therefore the total number of positive coefficients is at least 30. The last row also has a positive coefficient. Thus $N(G) \geq 31$. In fact, at least three more terms must be positive. The coefficients $U + 7B$ and $7U + 14B$ cannot both vanish, since $U < 0$. By symmetry the same holds for C and D . We conclude finally that $N(G) \geq 34$.

Next we discuss the case $U > 0$: The five rows preceding the final row generate 15 terms. The coefficients in the second and third columns in the last row are also positive. Thus there are at least 17 terms in addition to the 9 terms in the first three rows. If $B = C = D = 0$, the degree is 10 but we are in degree 11. Hence at least one of these is non-zero; say B . The five terms in which B appears alone then guarantee at least $17 + 9 + 5 = 31$ terms. Thus $N(G) \geq 31$.

We're left with the case where $U = 0$. If also $B = C = D = 0$, then $H = 0$ and $N(G) = 17$. If none of B, C, D are 0, then there is a huge number of terms, as only four of the above terms have a minus sign anywhere. If one is 0, say $D = 0$, with $B, C \neq 0$, we see also a profusion of terms. Finally $D = C = 0$ and $B > 0$ yields at least 32 terms. To see this, set $U = C = D = 0$ in the chart of terms. The first three rows create 9 positive coefficients. The next rows create

$$2 + 1 + 1 + 1 + 1 + 1 + 2 + 2 + 3 + 1 + 1 + 2 + 2 + 3 = 23$$

additional terms, Thus $N(G) \geq 32$. In fact 32 is possible by putting $B = 7$. If $0 < B < 7$, then we get 33 terms.

We have proved the following.

Proposition 5.1. *Let G be a special polynomial for $\Gamma(7)$. If the degree of G is 7, then $N(G) = 17$. There is no such G of degree 8 or 9. If the degree of G is 10, then then either $N(G) = 29$ or $N(G) = 30$. If the degree of G is 11, then $N(G) \geq 31$.*

As the degree of H increases, more $\Gamma(7)$ -invariant monomials are allowed. The coefficients of G become more complicated and a more delicate analysis is required. From the structure of the coefficient expressions in G , the coefficient U of xyz in H is the only coefficient that could be negative until we include the coefficient V of $x^2y^2z^2$ in H . Thus, the case analysis for $\deg(H) \geq 6$ is more involved than above. We now analyze the case where H is of degree 5. We have 7 parameters.

$$H(x, y, z) = Uxyz + Byz^3 + Cx^3z + Dxy^3 + Ry^3z^2 + Sx^2z^3 + Tx^3y^2.$$

As before we list all the terms:

z^7	x^7	y^7
$7y^5z$	$7xz^5$	$7x^5y$
$(7 - B)y^3z^3$	$(7 - C)x^3z$	$(7 - D)xy^3$
$7By^2z^6$	$7Cx^6z^2$	$7Dx^2y^6$
Byz^{10}	$Cx^{10}z$	Dxy^{10}
$(14 - R)y^3z^2$	$(14 - S)x^2z^3$	$(14 - T)x^3y^2$
Ry^3z^9	Sx^9z^3	Tx^3y^9
$Ry^{10}z^2$	Sx^2z^{10}	$Tx^{10}y^2$
$7Rxy^5z^6$	$7Sx^6yz^5$	$7Tx^5y^6z$
$(B + 7R)y^8z^3$	$(C + 7S)x^3z^8$	$(D + 7T)x^8y^3$
$(14B + 7R)y^4z^5$	$(14C + 7S)x^5z^4$	$(14D + 7T)x^4y^5$
$(7B + 14R)y^6z^4$	$(7C + 14S)x^4z^6$	$(7D + 14T)x^6y^4$
$(7B + D + 7R)xy^3z^7$	$(B + 7C + 7S)x^7yz^3$	$(C + 7D + 7T)x^3y^7z$
$(7B + 7D + 7R + 7S)x^2y^5z^4$	$(7B + 7C + 7S + 7T)x^4y^2z^5$	$(7C + 7D + 7R + 7T)x^5y^4z^2$
$(7R + S)x^2y^7z^3$	$(7S + T)x^3y^2z^7$	$(7T + R)x^7y^3z^2$
$(7 + 7U + 14B)xy^2z^4$	$(7 + 7U + 14C)x^4yz^2$	$(7 + 7U + 14D)x^2y^4z$
$(14U + 7B + 7D + 14R)xy^4z^3$	$(14U + 7B + 7C + 14S)x^3yz^4$	$(14U + 7C + 7D + 14T)x^4y^3z$
$(7U + 14D + 7R)xy^6z^2$	$(7U + 14B + 7S)x^2yz^6$	$(7U + 14C + 7T)x^6y^2z$
$(U + 7B)xy^8z$	$(U + 7C)x^8yz$	$(U + 7D)xy^8z$
$(7U + 7B + 7D + 14R + 14S)x^2y^3z^5$	$(7U + 7B + 7C + 14S + 14T)x^5y^2z^3$	$(7U + 7C + 7D + 14R + 14T)x^3y^5z^2$
$(14 - U)xyz$		
$(14U + 7)x^2y^2z^2$		
$(7U + 14B + 14C + 14D + 7R + 7S + 7T)x^3y^3z^3$		
$(7R + 7S + 7T)x^4y^4z^4$		

We need to count the possible number of positive coefficients. As before we consider $U < 0$, $U > 0$, and $U = 0$ separately. Since the degree of H is five, at least one of R, S, T is positive.

Case $U < 0$: There are 6 terms with positive constant coefficients. Next, since the coefficients $U + 7B, U + 7C, U + 7D$ appear, we must have $B, C, D > 0$. There are 2 terms involving B alone; replacing B with C and D we get a total of 6 terms (with positive coefficients) of this type. The terms $B + 7R, 14B + 7R, 7B + 14R, 7B + D + 7R, 7B + 7D + 7R + 7S$ each generates 3 terms in this way. These give 15 additional positive coefficients. Since not all R, S, T are 0, we may suppose $R > 0$. There are 3 terms with R alone, generating 3 more positive coefficients. The row starting $7R + S$ generates 2 more. The coefficient of $x^4y^4z^4$ in the last row must be positive. The term $14 - U$ must be positive. We therefore have at least

$$6 + 6 + 15 + 3 + 2 + 1 = 33$$

terms with positive coefficients.

Case $U > 0$: Assume $R > 0$ but that $B, C, D = 0$. Let $\mu = 2$ if $R = 14$ and $\mu = 3$ if $0 < R < 14$. Let $\eta = 1$ if $U = 14$ and equal 0 if $0 < U < 14$. Going down the rows, we get at least

$$3+3+3+0+0+\mu+1+1+1+1+1+1+1+2+3+3+3+3+3+\eta+1+1+1 = 35+\mu+\eta \geq 37.$$

terms with positive coefficients.

Case $U = 0$: In order that the degree is 12, at least one of R, S, T is positive. Assume $R > 0$. If $B > 0$, but $C = D = 0$, we proceed down the rows, getting

$$3+3+2+1+1+2+1+1+1+1+1+1+2+3+2+3+2+1+3+1+1+1+1 = 38$$

terms with positive coefficients. We get even more such terms when C and/or D is positive. It remains to check the case where $U = B = C = D = 0$. Let $\mu = 2$ if $R = 14$ and $\mu = 3$ if $0 < R < 14$. Going down the rows gives

$$3+3+3+0+0+\mu+1+1+1+1+1+1+1+2+2+3+1+1+0+2+1+1+1+1 = 31+\mu.$$

We get at least 33 terms and thus have proved the following result.

Proposition 5.2. *Let G be a special polynomial for $\Gamma(7)$ of degree 12. Then $N(G) \geq 33$.*

Finally we must deal with the harder case when G has degree 13. Thus H has degree 6. There are 11 (non-constant) $\Gamma(7)$ -invariant monomials of degree at most 6. Thus we put

$$(8) \quad H = Uxyz + Byz^3 + Cxy^3 + Dx^3z + Ry^3z^2 + Sx^2z^3 + Tx^3y^2 \\ + Ky^5z + Lxz^5 + Mx^5y + Vx^2y^2z^2.$$

We note the symmetries in these parameters. The numbers U and V play special roles. The parameters B, C, D behave symmetrically, the parameters R, S, T behave symmetrically, and K, L, M also do. We illustrate this symmetry now.

The invariant monomials are listed in such a way that B is the coefficient of the monomial of degree 4 omitting x . Then permuting the variables by $x \rightarrow y \rightarrow z \rightarrow x$ requires permuting the parameters by $B \rightarrow C \rightarrow D \rightarrow B$. The same holds for R, S, T and K, L, M . This symmetry explains the table of terms; the second and third terms in each row are obtained from the first through these permutations. On the next page, we give the terms of the general example of degree at most 13. In the following list, it is convenient to write $\phi(a, b, c) = a + 7b + 14c$.

z^7	x^7	y^7
$(7 - B)yz^3$	$(7 - C)x^3z$	$(7 - D)xy^3$
$7By^2z^6$	$7Cx^6z^2$	$7Dx^2y^6$
Byz^{10}	$Cx^{10}z$	Dxy^{10}
$(14 - R)y^3z^2$	$(14 - S)x^2z^3$	$(14 - T)x^3y^2$
Ry^3z^9	Sx^9z^3	Tx^3y^9
$(14B + 7R)y^4z^5$	$(14C + 7S)x^5z^4$	$(14D + 7T)x^4y^5$
$(7K + L)xy^7z^5$	$(7L + M)x^5yz^7$	$(K + 7M)x^7y^5z$
$(7K + 7L + 7R)xy^5z^6$	$(7L + 7M + 7S)x^6yz^5$	$(7K + 7M + 7T)x^5y^6z$
$(7 - K)y^5z$	$(7 - L)xz^5$	$(7 - M)x^5y$
Ky^5z^8	Lx^8z^5	Mx^5y^8
$Ky^{12}z$	Lxz^{12}	$Mx^{12}y$
$(7B + 7K + 14R)y^6z^4$	$(7C + 7L + 14S)x^4z^6$	$(7D + 7M + 14T)x^6y^4$
$(7K + R)y^{10}z^2$	$(7L + S)x^2z^{10}$	$(7M + T)x^{10}y^2$
$(B + 14K + 7R)y^8z^3$	$(C + 14L + 7S)x^3z^8$	$(D + 14M + 7T)x^8y^3$
$(7B + D + 14L + 7R)xy^3z^7$	$(B + 7C + 14M + 7S)x^7yz^3$	$(C + 7D + 14K + 7T)x^3y^7z$
$(7 + 7U + 14B)xy^2z^4$	$(7 + 7U + 14C)x^4yz^2$	$(7 + 7U + 14D)x^2y^4z$
$(14U + 7B + 7D + 14R)xy^4z^3$	$(14U + 7B + 7C + 14S)x^3yz^4$	$(14U + 7C + 7D + 14T)x^4y^3z$
$(7U + 14D + 14K + 7R)xy^6z^2$	$(7U + 14B + 14L + 7S)x^2yz^6$	$(7U + 14C + 14M + 7T)x^6y^2z$
$(U + 7D + 7K)xy^8z$	$(U + 7B + 7L)xyz^8$	$(U + 7C + 7M)x^8yz$
$(7K + 7R + S + 7V)x^2y^7z^3$	$(7L + 7S + T + 7V)x^3y^2z^7$	$(7M + 7T + R + 7V)x^7y^3z^2$
$\phi(0, B + D + R + S, K + V)x^2y^5z^4$	$\phi(0, C + B + S + T, L + V)x^4y^2z^5$	$\phi(0, C + D + R + T, M + V)x^5y^4z^2$
$\phi(V, U + B + D + L, R + S)x^2y^3z^5$	$\phi(V, U + C + B + M, S + T)x^5y^2z^3$	$\phi(V, U + D + C + K, T + R)x^3y^5z^2$
$(7K + V)x^2y^9z^2$	$(7L + V)x^2y^2z^9$	$(7M + V)x^9y^2z^2$
$(7K + 7V)x^4y^6z^3$	$(7L + 7V)x^3y^4z^6$	$(7M + 7V)x^6y^3z^4$
$(14 - U)xyz$		
$(14U - V + 7)x^2y^2z^2$		
$(7U + 14B + 14C + 14D + 7R + 7S + 7T + 14V)x^3y^3z^3$		
$(7R + 7S + 7T + 7V)x^4y^4z^4$		

Theorem 5.1. *Let $G : \mathbb{R}^3 \rightarrow \mathbb{R}$ be a $\Gamma(7)$ -invariant special polynomial. Then the number $N(G)$ of terms satisfies the following:*

- *There is no G with $0 \leq N(G) < 17$.*
- *There is no G with $17 < N(G) < 29$.*
- *There are G with $N(G) = 29, 30, 32, 33, 34$ and $N(G) = k$ for each $k \geq 37$.*
- *The authors believe that the numbers 31, 35, 36 are impossible but have not proved this statement.*

Proof. Our list of examples include those of degree 17, 29, 30, 33, 34 as well as all degrees from 37 to 57. By the postage stamp principle from Proposition 2.3, there are examples for all integers exceeding 57. Since $G(0) = 0$ by definition of a special polynomial, degree 0 is impossible. The minimum possible value of $N(G)$ occurs when G is the basic polynomial F and therefore $0 \leq N(G) < 17$ is impossible.

Let G be a $\Gamma(7)$ -invariant special polynomial. We write $G = F - H + HF$, as usual, where H is some $\Gamma(7)$ -invariant polynomial. Since G is non-constant, we have $N(G) \neq 0$. When the degree of G is at most 12, the previous Propositions show that either $H = 0$ and $N(G) = 17$, or that $N(G) \geq 29$. The degree estimate $N \geq 2d+1$ implies that $N(G) \geq 29$ when the degree of G is at least 14. Therefore it remains only to check the case when $\deg(G) = 13$, or equivalently when $\deg(H) = 6$.

We proceed as before. Define H as in equation (8). Before the statement of the Theorem we have listed all the terms in the resulting $G = F - H + HF$. Here U and V are the only coefficients that can be negative since all other coefficients appear alone as coefficients of G .

The case $V < 0$: Three of the coefficients are $V + 7K, V + 7L, V + 7M$. Thus K, L, M must be positive. The first row gives 3 terms. Rows 8, 9, 10-15 give 24 more terms. From $(7R + 7S + 7T + 7V)$, at least one of R, S, T must be positive. Suppose $R > 0$, then the coefficients R and $(14B + 7R)$ must be positive. Thus $N(G) \geq 3 + 24 + 2 = 29$.

The case $V \geq 0$: The 3 possibilities are $U < 0$, $U = 0$, and $U > 0$. Furthermore, at least one of K, L, M must be non-zero for G to be degree 13. Without loss of generality, assume $K > 0$. When $U > 0$, we get $N(G) \geq 33$.

When $U = 0$, there are 17 terms with K that must appear in G . We also have the 3 constant terms. The $(14 - U)$ coefficient yields another term in G . The coefficients in each column of the second and third rows cannot simultaneously vanish, and we get another 3 terms in $N(G)$. Similarly, rows 5 and 6 give another 3 and rows 10 and 11 give 2 more terms. Thus, $N(G) \geq 3 + 17 + 1 + 3 + 3 + 2 = 29$.

When $U < 0$, we have the initial 3 constant terms and 15 terms with K and no U . Including also the $14 - U$ coefficient gives us 19 positive coefficients so far. We pair the corresponding coefficients in the second and third rows, fifth and sixth rows to get an additional 6 terms. Pairing the 10th and 11th rows gives 2 more. Thus we have at least 27 terms. Finally using $U < 0$ with the $U + 7B + 7L$ coefficient gives $B > 0$ or $L > 0$. Thus the $(7B + D + 14L + 7R)$ and $\phi(0, C + B + S + T, L + V)$ coefficients must be positive. Again we have $N(G) \geq 29$. \square

This theorem leaves undecided three possible values, namely 31, 35, 36. The authors believe these values are not possible. To find an example with 36 terms, degree estimates force G to be of degree at most 17, and hence H to be of degree at most 10. There are 39 (non-constant) monomials that are $\Gamma(7)$ -invariant and degree at most 10. The resulting G has 161 coefficients. The problem then is to determine how many of the coefficients of G can be made to simultaneously vanish. The authors hope to address this computational problem in a future paper.

6. INTERPRETATION IN TERMS OF SPARSENESS

In applied mathematics it is often important to know how many components of an n -tuple of numbers are non-zero. This number is often called its ℓ^0 norm; see [12] for its use in compressed sensing. See also [7] for the connection with CR

geometry. The ℓ^0 norm $\|\mathbf{w}\|_0$ of a vector \mathbf{w} is **not** a norm; it is the number of non-vanishing components of \mathbf{w} with respect to a fixed basis.

Consider an affine map $\mathbf{v} \rightarrow W(\mathbf{v})$ between vector spaces with bases chosen. It is natural to study the ℓ^0 norm of $W(\mathbf{v})$ as a function of \mathbf{v} , but this simple sounding question is annoyingly complicated. The results in this paper can be interpreted in this manner. The set of coefficients in the charts in Section 5 can be regarded as images of an affine map. We describe the easier situation for $\Gamma(2r+1, 2)$. The proof of Theorem 1.2 for this group includes finding all special polynomials of degree at most $4r+1$. We show that such polynomials depend on r non-negative parameters. Let us denote these r parameters by a vector \mathbf{v} with respect to the standard basis in \mathbb{R}^r . This vector \mathbf{v} lies in an explicit closed rectangle in \mathbb{R}^r . We use its components as coefficients in defining an invariant polynomial $h_{\mathbf{v}}$. All special polynomials of degree at most $4r+1$ then have the form

$$(9) \quad W(\mathbf{v}) = f_{2r+1}(x, y) + h_{\mathbf{v}}(x, y) (f_{2r+1}(x, y) - 1).$$

Note that W is an affine map. In (9), the coefficients of the polynomial $h_{\mathbf{v}}$ are the r parameters. Theorem 1.2 for $\Gamma(2r+1, 2)$ has the following interpretation:

- The minimum ℓ^0 norm satisfies $\|W(\mathbf{v})\|_0 = r + 2$.
- $\|W(\mathbf{v})\|_0$ cannot lie in the integer interval $[1, r+1]$ nor in $[r+3, 2r+2]$.

Furthermore, if we allow polynomials of larger degree, we can regard W as an affine map $h \mapsto W(h)$ from a vector space of invariant polynomials (of dimension larger than r) to itself. Then

$$W(h) = f_{2r+1} + (f_{2r+1} - 1)h.$$

The last conclusion of Theorem 1.2 for this group then states:

- For any $K \geq 2r+3$, there is an h with $\|W(h)\|_0 = K$.

Similar interpretations apply in the other cases; for $\Gamma(7)$ we have listed several explicit examples of this situation where the number of source variables and the dimension of the image of the affine map are large.

Remark 6.1. Given an affine map from \mathbb{R}^r to some \mathbb{R}^K , it seems to be quite difficult in general to describe the set of values of the ℓ^0 norm. The situation in this paper involves an additional difficulty which we illustrate in the following example.

Example 6.1. The affine map here arises when considering Example 1.1 in source dimension 2. Define $W : \mathbb{R}^3 \rightarrow \mathbb{R}^8$ by

$$W(A, B, C) = (1 - A, 2 - B, 1 - C, A, 2A + B, A + 2B + C, B + 2C, C).$$

The ℓ^0 norm of $W(A, B, C)$ is generically equal to 8. We note the following additional values and the corresponding invariant polynomial p . Each such polynomial equals 1 when $x + y = 1$.

- $\|W(0, 0, 0)\|_0 = 3$. Here $p(x, y) = (x + y)^2$.
- $\|W(1, -2, 1)\|_0 = 4$. Here $p(x, y) = 4xy + x^4 - 2x^2y^2 + y^4$.
- $\|W(1, 2, 1)\|_0 = 5$. Here $p(x, y) = (x + y)^4$.

The invariant polynomial in the second item has a negative coefficient, and hence it is *not special*. There is no other (A, B, C) for which $\|W(A, B, C)\|_0 = 4$. There are no (A, B, C) for which $\|W(A, B, C)\|_0$ equals either 1 or 2. Thus 1, 2 are gaps in the possible values in general, and 4 is also a gap if we insist that $W(A, B, C)$ lies in the orthant Q where all the target variables are non-negative.

We now summarize the situation from this perspective. Assume that the basic polynomial F is known. We consider a collection \mathbf{S} of polynomials H . (In this paper, H must be group-invariant.) We regard the coefficients of H as parameters and form $G = F - H + HF$. When the degree of H is bounded, the list of coefficients of G can be thought of as defining an affine map from some real Euclidean space to one of typically larger dimension. Let us write $W(H)$ for the image of this map. The condition that F has non-negative coefficients forces these coefficients to satisfy a system of linear inequalities. We want to describe the set of possible values of $N(G)$, equivalently $\|W(H)\|_0$, as H ranges over \mathbf{S} . For the group $\Gamma(2r+1, 2)$, for example, the number of terms $N(G)$ can be either $r+2$ or any positive integer at least $2r+3$, and no other values are possible. The result for $\Gamma(7)$ is considerably harder to state and appears in Theorem 5.1.

7. CONCLUDING REMARKS

The coefficients of special polynomials in our charts can be regarded as defining an affine map from $\mathbb{R}^N \rightarrow \mathbb{R}^K$, where N is the number of parameters in our polynomial H and K is the number of coefficients. Here typically $K \gg N$. These coefficients must be nonnegative and the structure of the equations leads to further constraints. The result is, for each degree k , a compact convex subset \mathbf{S}_k of \mathbb{R}^N in which the coefficients must lie. To analyze the possible values of $N(G)$ for $G \in \mathbf{S}_k$ one must study the boundary of \mathbf{S}_k . Doing so is essentially equivalent to understanding sparseness for linear systems. We saw one simple example where the set consisted of the interval $[0, 14]$. In this case $N(G_\lambda) = 17$ for $\lambda = 0$, $N(G_\lambda) = 30$ for $0 < \lambda < 1$, and $N(G_\lambda) = 29$ when $\lambda = 14$. It is possible in principle to regard all of our calculations from this point of view. Perhaps the most important observation is that invariance reduces the size of the linear algebra problem in two ways. First, the number of invariant monomials of up to a given degree is smaller than the number of all monomials up to that degree. Second, the invariance allows for a somewhat simplified manner of analyzing the boundary.

We next mention further directions one could take along the lines of this paper.

The paper [13] studies the Bergman kernel function for the quotient of the ball by the first two classes of groups we have mentioned. It seems interesting to consider the case of the group in Example 1.3.

Since the possibilities for group-invariant rational sphere maps are so limited, it is natural to replace invariance with equivariance. In [10] and [11], one assigns a certain group Γ_f to a rational sphere map f . This group consists of all automorphisms α of the source ball for which there exists an automorphism β of the target ball with $f \circ \alpha = \beta \circ f$. There are many possibilities for Γ_f . For example, given a finite group G , there always is a polynomial sphere map f for which Γ_f is isomorphic to G . Furthermore, it is known precisely when Γ_f is the full automorphism group, when it is the unitary group, when it contains the n -torus, and so on. See [14] and [17] for further results in this direction. There remain combinatorial questions about the relationship between the degree of a polynomial map and the number of terms for a map with given group Γ_f .

Similar questions apply for the groups $\Gamma(p, q)$, but one must be cognizant of the warning preceding Example 2.1. When coefficients can be of both signs, it is extremely difficult to count the number of terms in the product of two polynomials. It seems nonetheless worthwhile to study the gaps that arise in the number of possible terms for polynomials invariant under $\Gamma(p, q)$.

REFERENCES

- [1] J. Brooks, S. Curry, D. Grundmeier, P. Gupta, V. Kunz, A. Malcom, and K. Palencia, Constructing group-invariant CR mappings, *Complex Anal. Synerg.* 8 (2022), no. 4, Paper No. 20, 7 pp.
- [2] S. Chiappari, Proper holomorphic mappings of positive codimension in several complex variables. PhD thesis, University of Illinois at Urbana-Champaign, 1990.
- [3] D. Coppersmith and J. Davenport, Polynomials Whose Powers Are Sparse, *Acta Arith.* 58 (1991), 79-87.
- [4] J. D'Angelo, Rational Sphere Maps, *Progress in Mathematics*, Volume 341, Birkhäuser, (2021).
- [5] J. D'Angelo, Number-theoretic properties of certain CR mappings. *J. Geom. Anal.* 14(2): (2004), 215-229.
- [6] J. D'Angelo, The combinatorics of certain group invariant maps. *Complex variables and elliptic equations*, Vol. 58, Issue 5 (2013), 621-634.
- [7] J. D'Angelo, D. Grundmeier, and J. Lebl, Rational sphere maps, linear programming, and compressed sensing, *Complex Analysis and its Synergies* 6, 4 (2020). <https://doi.org/10.1007/s40627-020-0041-5>.
- [8] J. D'Angelo, S. Kos, and E. Riehl, A sharp bound for the degree of proper monomial mappings between balls. *J. Geom. Anal.* 13 (2003), no. 4, 581-593.
- [9] J. D'Angelo and D. Lichtblau, Spherical space forms, CR Maps, and proper maps between balls, *J. Geom. Anal.* 2 (1992), 391-415.
- [10] J. D'Angelo and M. Xiao, Symmetries in CR complexity theory, *Adv. Math.* 313 (2017), 590-627.
- [11] J. D'Angelo and M. Xiao, Symmetries and regularity for holomorphic maps between balls, *Math. Res. Lett.*, Vol. 25, no. 5 (2018), 1389-1404.
- [12] D. Donoho, Compressed sensing, *IEEE Trans. Inform. Theory* 52 (2006), no. 4, 1289-1306.
- [13] P. Ebenfelt, M. Xiao, and H. Xu, On the classification of normal Stein spaces and finite ball quotients with Bergman-Einstein metrics, *Int. Math. Res. Not. (IMRN)*, 2021.
- [14] E. Gevorgyan, H. Wang, A. Zimmer, A rigidity result for proper holomorphic maps between balls, *Proc. Amer. Math. Soc.* 152(2024), no.4, 1573-1585.
- [15] D. Grundmeier, Signature pairs for group-invariant Hermitian polynomials. *Internat. J. Math.* 22 (2011), no. 3, 311-343.
- [16] D. Grundmeier, Group-invariant CR mappings. PHD Thesis, University of Illinois at Urbana-Champaign (2011).
- [17] D. Grundmeier, J. Lebl, Rational Maps of Balls and their Associated Groups, to appear in *Sao Paulo J. Math.*
- [18] D. Grundmeier, K. Linsuain, B. Whitaker, Invariant CR mappings between hyperquadrics. *Illinois J. Math.* 62(2018), no.1-4, 321-340.
- [19] X. Huang, S. Ji, and D. Xu, Several results for holomorphic mappings from \mathbb{B}^n into \mathbb{B}^N , *Geometric analysis of PDE and several complex variables*, 267-292, *Contemp. Math.*, 368, Amer. Math. Soc., Providence, RI, 2005.
- [20] X. Huang, S. Ji, and W. Yin, On the third gap for proper holomorphic maps between balls, *Math. Ann.* 358 (2014), no. 1-2, 115-142.
- [21] J. Lebl and H. Peters, Polynomials constant on a hyperplane and CR maps of spheres, *Illinois J. Math.* 56 (2012), no. 1, 155-175 (2013).
- [22] D. Lichtblau, Invariant proper holomorphic maps between balls. *Indiana Univ. Math. J.* 41 (1992), no. 1, 213-231.
- [23] N. Loehr, G. Warrington, and H. Wilf, The combinatorics of a three-line circulant determinant. *Israel J. Math.* 143 (2004), 141-156.

DEPT. OF MATHEMATICS, UNIV. OF ILLINOIS, 1409 W. GREEN ST., URBANA IL 61801
Email address: `jpda@illinois.edu`

DEPT. OF MATHEMATICS, THE OHIO STATE UNIVERSITY, 231 WEST 18TH AVENUE, COLUMBUS,
OHIO 43210
Email address: `grundmeier.1@osu.edu`

WOLFRAM RESEARCH, CHAMPAIGN IL 61802
Email address: `danl@wolfram.com`