

# Inferring the breakdown scales of the chiral expansions for $g_A$ and $m_N$

Andreas Ekström,<sup>1,\*</sup> Daniel R. Phillips,<sup>2,†</sup> Lucas Platter,<sup>3,4,‡</sup> and Matthias R. Schindler<sup>5,§</sup>

<sup>1</sup>*Department of Physics, Chalmers University of Technology, SE-412 96, Göteborg, Sweden*

<sup>2</sup>*Institute of Nuclear and Particle Physics and Department of Physics and Astronomy,  
Ohio University, Athens, Ohio 45701, USA*

<sup>3</sup>*Department of Physics and Astronomy, University of Tennessee, Knoxville, Tennessee 37996, USA*

<sup>4</sup>*Physics Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, USA*

<sup>5</sup>*Department of Physics and Astronomy, University of South Carolina, Columbia, South Carolina 29208, USA*  
(Dated: December 9, 2025)

We apply Bayesian inference to the order-by-order chiral perturbation theory ( $\chi$ PT) expansions for the axial-vector coupling constant  $g_A$  and the nucleon mass  $m_N$ , and thereby infer the scales at which  $\chi$ PT breaks down for these two observables. Using a pointwise Bayesian analysis, we find that the inferred breakdown scales are notably different for the two observables. For the chiral expansion of  $g_A$ , we obtain  $251^{+20}_{-50}$  MeV and  $211^{+20}_{-30}$  MeV using two distinct sets of low-energy constants, while for the chiral expansion of  $m_N$  we infer a significantly larger breakdown scale of  $491^{+60}_{-90}$  MeV.

## I. INTRODUCTION

Chiral perturbation theory ( $\chi$ PT) [1–3] is among the most successful effective field theories in nuclear and particle physics.  $\chi$ PT is based on the fact that pions are Goldstone bosons, associated with the spontaneously broken  $SU(2)_L \times SU(2)_R$  chiral symmetry of quantum chromodynamics (QCD) that emerges in the limit of zero up and down quark masses—the chiral limit. The pions’ finite mass arises from explicit chiral symmetry breaking due to nonvanishing quark masses  $m_q$ . In  $\chi$ PT observables are expanded in powers of a ratio  $Q$  of light to heavy scales,  $Q = \Lambda_l/\Lambda_B$ , where the light scale  $\Lambda_l$  is identified with the pion mass  $M$  or a momentum  $p$  of similar size, and the heavy scale  $\Lambda_B$  denotes the breakdown scale of  $\chi$ PT. The pion mass itself has such an expansion, where  $M$  denotes the leading-order expression of the pion mass with  $M^2 \propto m_q$ . In the mesonic sector  $\Lambda_B$  is typically assumed to be either  $\sim 4\pi F_\pi$  (where  $F_\pi = 92.7$  MeV is the pion decay constant) [4] or the mass  $m_\rho = 0.770$  GeV of the  $\rho$  meson, the lightest QCD excitation not explicitly included in mesonic  $\chi$ PT.

This framework has also been extended to include nucleons. The non-vanishing of the nucleon mass in the chiral limit introduces additional complexities compared to the meson sector [5]. Different approaches to these issues have been proposed [6–10]; for reviews see, e.g., Refs. [7, 11, 12]. Here we focus on the formulation of nucleons as heavy baryons [6, 13]. At leading order in this approach, the nucleon is treated as a static field, which acts as a source of Goldstone bosons. Nucleon-pion interactions are then described by an effective Lagrangian which contains increasing powers of momenta and the pion mass, leading to expansions of nucleonic observables

in powers of  $M$  and  $p$ . Prominent examples include the nucleon mass, the nucleon axial-vector coupling, nucleon form factors, and nucleon polarizabilities [14].

Such expansions are of particular importance in lattice QCD, where simulations are frequently performed at pion masses larger than the physical value. Extrapolating these results to the physical point relies on the convergence of the  $\chi$ PT series. And, even in cases where lattice simulations are performed at the physical pion mass, it may still be advantageous to combine results from simulations carried out at different quark masses.

However, in the one-nucleon sector, the situation is complicated by the appearance of additional scales. In particular, the  $\Delta(1232)$  resonance is the lowest-lying (non-pionic) QCD excitation in that sector, and while baryon  $\chi$ PT without explicit  $\Delta$  degrees of freedom is commonly used, arguments for the inclusion of  $\Delta$  excitations as active degrees of freedom have been presented since the advent of baryon  $\chi$ PT [6, 15–17].

A prominent example for the need of including the  $\Delta$  is the chiral expansion of the axial-vector coupling constant  $g_A$ : the first nontrivial correction enters at relative order  $M^2$ , the next correction proportional to  $M^3$  is numerically very large. This slow convergence has long been attributed to the presence of  $\Delta$  excitations inside the loop corrections to  $g_A$  [18–23]. And indeed, such inclusion is mandatory in the large- $N_c$  limit of QCD, in order to preserve unitarity [20, 24]. This introduces the  $\Delta$ -nucleon mass difference as an additional scale of approximately 290 MeV—much lower than either  $4\pi F_\pi$  or  $m_\rho$ —into the expansion.

These considerations naturally lead us to ask: What information about the single-nucleon  $\chi$ PT breakdown scale or scales can be extracted from order-by-order expansions of observables in that theory? Earlier work on pion-nucleon scattering,  $m_N$ , and  $g_A$  [25–28] addressed this question by comparing individual higher-order contributions or the resummation of particular subsets of higher-order contributions to observables with lower-order terms. In general, it was found that baryon  $\chi$ PT

\* andreas.ekstrom@chalmers.se

† phillid1@ohio.edu

‡ lplatter@utk.edu

§ mschindl@mailbox.sc.edu

expansions cease to be reliable around pion masses of 300 MeV to 500 MeV, depending on the observable.

Recently, Bayesian statistical techniques have been developed to analyze the convergence pattern of effective field theories [29, 30], enabling quantitative estimates of their respective breakdown scales. In Refs. [31–36] these methods were applied to demonstrate that the breakdown scale of chiral EFT, applied to two-nucleon scattering, nuclei with mass number  $A = 3$  and 4, and neutron matter, has a most likely value of  $\Lambda_B = 600$  MeV. In Refs. [37, 38] these methods were applied to pionless EFT for two- and three-nucleon systems and a most likely breakdown scale of approximately the pion mass was inferred (but cf. Ref. [39] for a different perspective).

In this work, we apply these statistical techniques to single-nucleon  $\chi$ PT to infer the underlying breakdown scales of the  $\chi$ PT expansions of the nucleon mass  $m_N$  and the axial-vector coupling constant  $g_A$ . We achieve this as follows. First, we introduce the chiral expansions for  $m_N$  and  $g_A$ , see Sec. II. The Bayesian methodology employed to analyze their respective convergence patterns and the inference of the associated breakdown scales are described in Sec. III. We conclude with a discussion of our results and their implications in Sec. IV.

## II. CHIRAL EXPANSIONS OF $m_N$ AND $g_A$

We use established expressions for the  $\chi$ PT expansions of  $m_N$  and  $g_A$  as outlined below. The coefficients of these expansions generally depend on the pion-decay constant  $F$ , the nucleon mass  $\hat{m}$ , and the nucleon axial-vector coupling  $\hat{g}$  in their respective chiral limits, and a number of other low-energy constants (LECs):  $l_i$  from the fourth-order pionic Lagrangian and  $c_i, d_i, e_i$  from the pion-nucleon Lagrangian at second, third, and fourth order, respectively. Renormalized LECs are denoted by a superscript  $r$  and depend on the renormalization scale  $\mu$ . The renormalized fourth-order mesonic LECs  $l_i^r(\mu)$  are related to the scale-independent LECs  $\bar{l}_i$  by

$$l_i^r(\mu) = \frac{\beta_{l_i}}{32\pi^2} \left[ \bar{l}_i + 2 \ln \left( \frac{M_\pi^{(\text{phys})}}{\mu} \right) \right], \quad (1)$$

with  $\beta_{l_3} = -\frac{1}{2}$  and  $\beta_{l_4} = 2$  contributing in the following analyses. Analogously, the pion-nucleon LECs  $d_i^r(\mu)$  are connected to the scale-independent LECs  $\bar{d}_i$  by the expressions [40]

$$d_i^r(\mu) = \bar{d}_i + \frac{\beta_{d_i}}{(4\pi F)^2} \ln \left( \frac{M_\pi^{(\text{phys})}}{\mu} \right), \quad (2)$$

with the  $\beta_{d_i}$  given in Ref. [41]. For the physical pion mass we take  $M_\pi^{(\text{phys})} = 0.139$  GeV. For the pion-nucleon LECs we use the values listed in Table. I. They are obtained from a  $\chi$ PT analysis of the processes  $\pi N \rightarrow \pi N$  and  $\pi N \rightarrow \pi\pi N$  in Ref. [42]. The values referred to as “Set

Low-energy constants			
LEC	Set 1	Set 2	Set $m_N$
$c_1$ [GeV $^{-1}$ ]	-	-	-1.11
$c_2$ [GeV $^{-1}$ ]	3.51	4.89	3.13
$c_3$ [GeV $^{-1}$ ]	-6.63	-7.26	-5.61
$c_4$ [GeV $^{-1}$ ]	4.01	4.74	-
$(\bar{d}_1 + \bar{d}_2)$ [GeV $^{-2}$ ]	4.73	3.39	-
$\bar{d}_{10}$ [GeV $^{-2}$ ]	-0.8	10.9	-
$\bar{d}_{11}$ [GeV $^{-2}$ ]	-15.6	-30.9	-
$\bar{d}_{12}$ [GeV $^{-2}$ ]	5.9	-10.9	-
$\bar{d}_{13}$ [GeV $^{-2}$ ]	13.6	27.7	-
$\bar{d}_{14}$ [GeV $^{-2}$ ]	-7.43	-7.36	-
$\bar{d}_{16}$ [GeV $^{-2}$ ]	0.4	-3.0	-
$\bar{d}_{18}$ [GeV $^{-2}$ ]	-0.8	-0.8	-
$\bar{e}_1$ [GeV $^{-3}$ ]	-	-	12.6
$l_3^r(m)$	$1.4 \cdot 10^{-3}$	$1.4 \cdot 10^{-3}$	-
$l_4^r(m)$	$3.7 \cdot 10^{-3}$	$3.7 \cdot 10^{-3}$	-
$\hat{g}$	1.0	1.3	-
$\hat{m}$ [GeV]	0.87	0.87	0.8695
$F$ [GeV]	0.087	0.087	-
$g_A$	-	-	1.275
$F_\pi$ [GeV]	-	-	0.0927
$m_N$ [GeV]	-	-	0.939

TABLE I. Values of low-energy constants employed in the chiral expansions of the nucleon mass  $m_N$  (Set  $m_N$ ) and  $g_A$  (Sets 1 and 2).

1” and “Set 2” in Ref. [43] are the ones we employ in the chiral expansion of  $g_A$ . The LEC values used in the expansion for  $m_N$  are referred to as “Set  $m_N$ ” and are taken from Ref. [44], with the value of the fourth-order pion-nucleon LEC  $\bar{e}_1$  provided in Ref. [45].

The chiral expansion of the nucleon mass to order  $M^4$  as given in Ref. [46] is

$$m_N = \hat{m} - 4c_1 M^2 - \frac{3g_A^2 M^3}{32\pi F_\pi^2} + \left( k_1 \ln \frac{M}{m_N} + k_2 \right) M^4, \quad (3)$$

with the coefficients

$$k_1 = -\frac{3}{32\pi^2 F_\pi^2 m_N} (g_A^2 + m_N (-8c_1 + c_2 + 4c_3)) , \quad (4)$$

$$k_2 = \bar{e}_1 - \frac{3}{128\pi^2 F_\pi^2 m_N} (2g_A^2 - m_N c_2) . \quad (5)$$

The use of the physical values of the pion decay constant  $F_\pi$ , the nucleon mass  $m_N$  and the axial-vector coupling  $g_A$  in these expressions instead of the chiral limit values generates differences at order  $M^5$ , so beyond the order to which we are working. While  $m_N$  has been calculated to order  $M^5$  [26, 47, 48] and order  $M^6$  [27, 49, 50] in various renormalization schemes, using these expressions




FIG. 1. Pion mass dependence of the nucleon mass  $m_N$  using “Set  $m_N$ ” given in Table I.

would require different determinations of the LECs that are currently not available.

Figure 1 shows the pion mass dependence of  $m_N$  obtained in this expansion. This result will serve as our data for the Bayesian analysis of the order-by-order convergence of this expansion and enable us to infer the  $\chi$ PT breakdown scale for  $m_N$ . We will perform the same analysis for  $g_A$ , which we discuss next.

The chiral expansion of the axial-vector coupling constant to order  $M^4$  was recently derived in Ref. [43]<sup>1</sup>,

$$g_A = \dot{g} \left[ 1 + \left( \frac{\alpha_2}{(4\pi F)^2} \ln \left( \frac{M}{\mu} \right) + \beta_2 \right) M^2 + \alpha_3 M^3 + \frac{1}{\dot{g}} \left( \frac{\alpha_4}{(4\pi F)^4} \ln^2 \left( \frac{M}{\mu} \right) + \frac{\gamma_4}{(4\pi F)^4} \ln \left( \frac{M}{\mu} \right) + \beta_4 + \dot{g} C \right) M^4 \right]. \quad (6)$$

The parameters  $\alpha_2$  and  $\beta_2$  arising from the one-loop calculation were first given in Ref. [7],

$$\alpha_2 = -2 - 4\dot{g}^2 \quad (7)$$

$$\beta_2 = \frac{4}{\dot{g}} d_{16}^r(\mu) - \frac{\dot{g}^2}{(4\pi F)^2}. \quad (8)$$

The contribution cubic in the pion mass is known to give a large correction and was first calculated in Ref. [51],

$$\alpha_3 = \frac{1}{24\pi^2 F^2 \dot{m}} (3 + 3\dot{g}^2 - 4\dot{m}c_3 + 8\dot{m}c_4). \quad (9)$$

The expressions for the coefficients of the  $M^4$  contribution are given in Ref. [43]. The coefficient for the leading

non-analytic term is

$$\alpha_4 = -\frac{7}{3}\dot{g}(1 - \dot{g}^2) + 16\dot{g}^5. \quad (10)$$

The parameters  $\beta_4$  and  $\gamma_4$  can be arranged according to the power of  $\dot{g}$  they include,

$$\gamma_4 = \sum_{i=0}^5 \dot{g}^i \gamma_4^{(i)}, \quad (11)$$

$$\beta_4 = \sum_{i=0}^5 \dot{g}^i \beta_4^{(i)}, \quad (12)$$

with  $\gamma_4^{(4)} = \beta_4^{(4)} = 0$ . The remaining terms are given by [43]

$$\begin{aligned} (4\pi F)^4 \beta_4^{(0)} &= -4\pi^2 F^2 (2d_{10}^r + 4d_{11}^r + 3d_{12}^r + d_{13}^r), \\ (4\pi F)^4 \beta_4^{(1)} &= -8\pi^2 F^2 (d_{14}^r + 2(d_1^r + d_2^r)) - \frac{\pi^2}{3} \\ &\quad - 32\pi^2 l_3^r + \frac{3575}{864} + \frac{1}{2} \psi_{2/3}^{(1)} + \frac{16\pi^2 F^2}{\dot{m}} (c_2 + 4c_4 + \frac{1}{\dot{m}}), \\ (4\pi F)^4 \beta_4^{(2)} &= -64\pi^2 F^2 (3d_{16}^r - d_{18}^r), \\ (4\pi F)^4 \beta_4^{(3)} &= 32\pi^2 (-3l_3^r + l_4^r) - \frac{\pi^2}{27} (61 + 48 \log 3) \\ &\quad - \frac{335}{432} - \frac{1}{6} \psi_{2/3}^{(1)} + \frac{32\pi^2 F^2}{\dot{m}^2}, \\ (4\pi F)^4 \beta_4^{(5)} &= \frac{41}{36} + \frac{7}{3} \pi^2, \end{aligned} \quad (13)$$

where  $\psi_{2/3}^{(1)}$  denotes the first derivative of the digamma function evaluated at  $\frac{2}{3}$ , and

$$\begin{aligned} \gamma_4^{(0)} &= 16\pi^2 F^2 (-20d_{16}^3 + 8d_{18}^r + 14d_{10}^r \\ &\quad + 8d_{11}^r + 3d_{12}^r + d_{13}^r), \\ \gamma_4^{(1)} &= -32\pi^2 F^2 (d_{14}^r - 2(d_1^r + d_2^r)) - 64\pi^2 (l_3^r - l_4^r) \\ &\quad - \frac{389}{36} - \frac{64\pi^2 F^2}{\dot{m}} \left( c_2 + c_3 - c_4 - \frac{1}{2\dot{m}} \right), \\ \gamma_4^{(2)} &= 256\pi^2 F^2 (d_{18}^r - 3d_{16}^r), \\ \gamma_4^{(3)} &= -128\pi^2 (l_3^r - l_4^r) + \frac{1}{9} (13 - 16\pi^2) + \frac{48\pi^2 F^2}{\dot{m}^2}, \\ \gamma_4^{(5)} &= \frac{11}{3}. \end{aligned} \quad (14)$$

The constant  $C$  denotes a linear combination of LECs from the fifth-order pion-nucleon Lagrangian. Its value is currently not determined, with Ref. [43] exploring its impact by considering two non-zero values. We set  $C = 0$  in our analysis, but have checked that non-zero values such as those considered in Ref. [43] do not significantly impact our findings.

The pion mass dependence of  $g_A$  obtained in this expansion with LEC values from “Set 1” is shown in Fig. 2.

<sup>1</sup> We thank the authors of Ref. [43] for their help in identifying some typos in the original expressions for the expansion coefficients; these are corrected in the following equations.




FIG. 2. Pion mass dependence of the axial-vector coupling constant  $g_A$  for “Set 1” given in Ref. [43] of LECs and  $\hat{g} = 1$ . For the unknown LEC, we use  $C = 0$ .

### III. ANALYSIS AND RESULTS

We seek to infer the posterior probability densities  $p(\Lambda_B|\mathbf{y}, I)$  for the breakdown scale  $\Lambda_B$  of  $\chi$ PT conditioned on a set of order-by-order predictions for the single-nucleon observables  $m_N$  and  $g_A$ , respectively. The data vector  $\mathbf{y}$  contains the chiral expansion for either  $g_A$  or  $m_N$ , truncated at successive orders, as described in Sec. II. The inference is also conditioned on assumptions, denoted  $I$ , that are outlined in Sec. II and below. Intuitively, the posterior for  $\Lambda_B$  reflects how well the chiral expansion converges across successive orders and encodes the extent to which the observed convergence pattern is consistent with the natural scales of the  $\chi$ PT power counting, as reflected in the priors we place.

We use the point-wise method presented in Melendez et al. [30] and thus first use Bayes’ theorem to express the sought posterior as proportional to a likelihood times a prior

$$p(\Lambda_B|\mathbf{y}, I) \propto p(\mathbf{y}|\Lambda_B, I) \cdot p(\Lambda_B|I). \quad (15)$$

For the likelihood  $p(\mathbf{y}|\Lambda_B, I)$  we assume the  $n$ -th chiral order prediction for an observable  $X \in \{g_A, m_N\}$  to follow a formal expansion [29]

$$X^{(n)} = X_{\text{ref}} \sum_{i=0}^n c_i Q^i, \quad (16)$$

with dimensionless expansion parameter  $Q = M/\Lambda_B$  and coefficients  $c_i$ <sup>2</sup>. The breakdown scale for the observable  $X$  corresponds to the pion mass at which all terms in its  $\chi$ PT expansion give contributions of the same size. Therefore, once  $M$  actually reaches  $\Lambda_B$  we are well beyond the point at which  $\chi$ PT is a useful calculational

Order	$\Lambda_B(g_A; \text{Set 1})$	$\Lambda_B(g_A; \text{Set 2})$	$\Lambda_B(m_N; \text{Set } m_N)$
$M^2$	351 (201,461)	281 (171,361)	431 (271,551)
$M^3$	221 (181,251)	211 (171,241)	511 (391,601)
$M^4$	251 (201,271)	211 (181,231)	491 (401,551)

TABLE II. Median values and 68% degree of belief intervals (highest posterior density) for the breakdown scales, obtained at different orders in the chiral expansions for  $g_A$  and  $m_N$ . All values in MeV.

tool for  $X$ . We assume natural-sized coefficients by assigning a normal prior  $c_i \stackrel{\text{iid}}{\sim} \mathcal{N}(0, \bar{c}^2)$  with a conjugate inverse-chi-squared  $\chi^{-2}$  hyperprior for the variance  $\bar{c}^2$ , i.e., we have  $\bar{c}^2 \sim \chi^{-2}(\nu_0 = 2, \tau_0^2 = 1)$ . This choice places about 67% of the probability for  $\bar{c}^2$  within the natural range  $[1/3, 3]$ . For both observables, we set the reference scale  $X_{\text{ref}}$  equal to the leading-order contribution, so that  $c_0 = 1$ . We then quantify the expansion coefficients  $c_i$  in Eq. (16) from the order-by-order calculations of  $X$  at  $M = [9.85, 301.84]$  MeV. Here, we selected expansion data at two  $M$ -values to avoid over-confident posteriors by remaining consistent with the assumption of independent data<sup>3</sup>. For the prior  $p(\Lambda_B|I)$  we assume a log-uniform distribution with compact support on the interval  $[M_\pi^{(\text{phys})}/40, 40M_\pi^{(\text{phys})}]$ . From this we infer posterior probability distributions for the  $\chi$ PT breakdown scales for  $g_A$  and  $m_N$  as shown in Figs. 3-4, respectively. The order-by-order posteriors for  $\Lambda_B$  converge steadily, for both observables. The results for median values and 68% degree of belief intervals are summarized in Table II.

We checked that our Bayesian analysis is robust with respect to variations in the employed priors and to different choices of point-wise  $M$ -values used for extracting the order-by-order expansion data. As we increase the number of soft-scale evaluation points, the posterior modes of the breakdown scales remain unchanged, while the corresponding distributions become narrower. This is not a true reduction in posterior uncertainty since our analysis does not account for the finite correlation length of the EFT expansion coefficients  $c_i$ .

### IV. SUMMARY AND DISCUSSION

We have used Bayesian methods to infer a breakdown scale of approximately 230 MeV when analyzing the  $\chi$ PT expansion of the axial-vector coupling  $g_A$  and 490 MeV when considering the expansion of the nucleon mass,  $m_N$ . We note that these values for  $\Lambda_B$  were derived using point estimates of the LECs that appear in the  $\chi$ PT expansion of  $m_N$  and  $g_A$ . A complete Bayesian analysis of the

<sup>2</sup> Note that these  $c_i$  coefficients are distinct and not the same as the LECs entering the chiral expansions.

<sup>3</sup> One might be concerned that this interval includes  $M$  values that are potentially larger than  $\Lambda_B$ . However, this does not invalidate the use of Eq. (15) to calculate the  $\Lambda_B$  posterior.




FIG. 3. Posteriors for the breakdown scale  $\Lambda_B$  obtained from the analysis of the order-by-order expansion of the axial-vector coupling constant  $g_A$  and using the values from Set 1 (top panel) and Set 2 (bottom panel) in Table I for the LECs.

convergence of the EFT would estimate the LECs from data in the presence of truncation errors and then consistently determine  $\Lambda_B$  as part of that inference process. Refs. [32, 39] provide examples of such analyses for the case of  $\chi$ EFT applied to few-nucleon systems.

While it has been known for some time that the convergence of baryon  $\chi$ PT for  $g_A$  is worse than that for  $m_N$ , our Bayesian analysis quantifies the strength of this




FIG. 4. Posteriors for the breakdown scale  $\Lambda_B$  obtained from the analysis of the order-by-order expansion of nucleon mass  $m_N$  and using the values from “Set  $m_N$ ” in Table I for the LECs.

inference. The much lower number found for  $\Lambda_B$  in the case of the expansion of  $g_A$  appears to be related to the nucleon- $\Delta$  mass splitting. Meanwhile, the breakdown scale inferred in the case of  $m_N$  is closer to the mass scale of light mesonic degrees of freedom not explicitly included in  $\chi$ PT.

This emphasizes that the breakdown scale inferred from an effective field theory expansion of an observable will typically depend on the observable(s) used for inference. We note that the breakdown scale of nucleon-nucleon scattering in chiral effective theory has been inferred to be higher than both breakdown scales found here, at approximately 600 MeV [31–34, 36]. Even though there is evidence that the breakdown scale in nucleon-nucleon scattering is lower in a strictly renormalizable implementation of  $\chi$ EFT [52], nucleon-nucleon scattering does not appear to show strong sensitivity to the presence of the  $\Delta$ -excitation.

However, it remains an open question whether there are certain nuclear matrix elements that are influenced by  $\Delta$  excitation and so also exhibit a lower  $\Lambda_B$ . For example, the two-nucleon axial current involves the linear combination  $c_3 + 2c_4$  of pion-nucleon LECs. These are the same LECs responsible for the large  $M^3$  correction in the chiral expansion of  $g_A$ . However, in the case of the two-nucleon axial current they occur with the same sign, leading to cancellation instead of reinforcement. Thus, ultimately the contribution of this current to most nuclear  $\beta$ -decay matrix elements is of the size expected from the  $\chi$ EFT counting.

Our results also raise the important question whether the breakdown scale inferred for observables in powers of



the pion mass  $M$  is the same for observables that are also momentum-dependent and thus also expanded in powers of the momentum  $\mathbf{q}$ . This outcome would, for example, have important consequences for  $\chi$ PT calculations of form factors such as the axial form factor.

## ACKNOWLEDGMENTS

We are grateful to J. Gegelia and U.-G. Meißner for answering our questions regarding Ref. [43], and to V. Bernard for a detailed comparison of the expressions for the chiral expansion of  $g_A$ . MRS thanks

T. R. Richardson for helpful discussions. This work was supported by the National Science Foundation (Grant Nos. PHY-2111426 and PHY-2412612), the US Department of Energy (Contract No. DE-AC05-00OR22725), the U.S. Department of Energy, Office of Science, Office of Nuclear Physics (Award Nos. DE-SC0019647 and DE-FG02-93ER-40756) and by the Swedish Research Council (Grants No. 2020-05127 and No. 2024-04681). We thank the Institute for Nuclear Theory at the University of Washington for its kind hospitality and stimulating research environment. This research was supported in part by the INT's U.S. Department of Energy grant No. DE-FG02-00ER41132.

- 
- [1] S. Weinberg, *Physica A* **96**, 327 (1979).
  - [2] J. Gasser and H. Leutwyler, *Annals Phys.* **158**, 142 (1984).
  - [3] J. Gasser and H. Leutwyler, *Nucl. Phys. B* **250**, 465 (1985).
  - [4] A. Manohar and H. Georgi, *Nucl. Phys. B* **234**, 189 (1984).
  - [5] J. Gasser, M. E. Sainio, and A. Švarc, *Nucl. Phys. B* **307**, 779 (1988).
  - [6] E. E. Jenkins and A. V. Manohar, *Phys. Lett. B* **255**, 558 (1991).
  - [7] V. Bernard, N. Kaiser, J. Kambor, and U. G. Meissner, *Nucl. Phys. B* **388**, 315 (1992).
  - [8] T. Becher and H. Leutwyler, *Eur. Phys. J.* **9**, 643 (1999), [arXiv:hep-ph/9901384](#).
  - [9] J. Gegelia and G. Japaridze, *Phys. Rev. D* **60**, 114038 (1999), [arXiv:hep-ph/9908377](#).
  - [10] T. Fuchs, J. Gegelia, G. Japaridze, and S. Scherer, *Phys. Rev. D* **68**, 056005 (2003), [arXiv:hep-ph/0302117](#).
  - [11] V. Bernard, *Prog. Part. Nucl. Phys.* **60**, 82 (2008), [arXiv:0706.0312 \[hep-ph\]](#).
  - [12] S. Scherer and M. R. Schindler, *A Primer for Chiral Perturbation Theory*, Vol. 830 (2012).
  - [13] V. Bernard, N. Kaiser, and U. G. Meissner, *Phys. Rev. Lett.* **67**, 1515 (1991).
  - [14] V. Bernard, N. Kaiser, and U.-G. Meissner, *Int. J. Mod. Phys. E* **4**, 193 (1995), [arXiv:hep-ph/9501384](#).
  - [15] E. E. Jenkins and A. V. Manohar, in *Workshop on Effective Field Theories of the Standard Model* [34] (1991).
  - [16] T. R. Hemmert, B. R. Holstein, and J. Kambor, *Phys. Lett. B* **395**, 89 (1997), [arXiv:hep-ph/9606456](#).
  - [17] T. R. Hemmert, B. R. Holstein, and J. Kambor, *J. Phys. G* **24**, 1831 (1998), [arXiv:hep-ph/9712496](#).
  - [18] E. E. Jenkins and A. V. Manohar, *Phys. Lett. B* **259**, 353 (1991).
  - [19] M. A. Luty and M. J. White, *Phys. Lett. B* **319**, 261 (1993), [arXiv:hep-ph/9305203](#).
  - [20] R. Flores-Mendieta, C. P. Hofmann, E. E. Jenkins, and A. V. Manohar, *Phys. Rev. D* **62**, 034001 (2000), [arXiv:hep-ph/0001218](#).
  - [21] T. R. Hemmert, M. Procura, and W. Weise, *Phys. Rev. D* **68**, 075009 (2003), [arXiv:hep-lat/0303002](#).
  - [22] A. Calle Cordon and J. L. Goity, *Phys. Rev. D* **87**, 016019 (2013), [arXiv:1210.2364 \[nucl-th\]](#).
  - [23] Z. B. Hall et al., (2025), [arXiv:2503.09891 \[hep-lat\]](#).
  - [24] R. F. Dashen, E. E. Jenkins, and A. V. Manohar, *Phys. Rev. D* **49**, 4713 (1994), [Erratum: *Phys. Rev. D* **51**, 2489 (1995)], [arXiv:hep-ph/9310379](#).
  - [25] D. Djukanovic, J. Gegelia, and S. Scherer, *Eur. Phys. J. A* **29**, 337 (2006), [arXiv:hep-ph/0604164](#).
  - [26] J. A. McGovern and M. C. Birse, *Phys. Rev. D* **74**, 097501 (2006), [arXiv:hep-lat/0608002](#).
  - [27] M. R. Schindler, D. Djukanovic, J. Gegelia, and S. Scherer, *Phys. Lett. B* **649**, 390 (2007), [arXiv:hep-ph/0612164](#).
  - [28] V. Bernard and U.-G. Meissner, *Phys. Lett. B* **639**, 278 (2006), [arXiv:hep-lat/0605010](#).
  - [29] R. J. Furnstahl, N. Klco, D. R. Phillips, and S. Wesolowski, *Phys. Rev. C* **92**, 024005 (2015), [arXiv:1506.01343 \[nucl-th\]](#).
  - [30] J. A. Melendez, R. J. Furnstahl, D. R. Phillips, M. T. Pratala, and S. Wesolowski, *Phys. Rev. C* **100**, 044001 (2019), [arXiv:1904.10581 \[nucl-th\]](#).
  - [31] J. A. Melendez, S. Wesolowski, and R. J. Furnstahl, *Phys. Rev. C* **96**, 024003 (2017), [arXiv:1704.03308 \[nucl-th\]](#).
  - [32] S. Wesolowski, I. Svensson, A. Ekström, C. Forssén, R. J. Furnstahl, J. A. Melendez, and D. R. Phillips, *Phys. Rev. C* **104**, 064001 (2021), [arXiv:2104.04441 \[nucl-th\]](#).
  - [33] C. Drischler, R. J. Furnstahl, J. A. Melendez, and D. R. Phillips, *Phys. Rev. Lett.* **125**, 202702 (2020), [arXiv:2004.07232 \[nucl-th\]](#).
  - [34] P. J. Millican, R. J. Furnstahl, J. A. Melendez, D. R. Phillips, and M. T. Pratala, *Phys. Rev. C* **110**, 044002 (2024), [arXiv:2402.13165 \[nucl-th\]](#).
  - [35] P. J. Millican, R. J. Furnstahl, J. A. Melendez, and D. R. Phillips, (2025), [arXiv:2508.17558 \[nucl-th\]](#).
  - [36] B. McClung, C. Elster, and D. R. Phillips, *Phys. Rev. C* **111**, 064002 (2025), [arXiv:2501.09231 \[nucl-th\]](#).
  - [37] A. Ekström and L. Platter, *Phys. Lett. B* **860**, 139207 (2025), [arXiv:2409.08197 \[nucl-th\]](#).
  - [38] A. Ekström and L. Platter, *Phys. Rev. C* **112**, L031002 (2025), [arXiv:2507.08700 \[nucl-th\]](#).
  - [39] J. M. Bub, M. Piarulli, R. J. Furnstahl, S. Pastore, and D. R. Phillips, *Phys. Rev. C* **111**, 034005 (2025), [arXiv:2408.02480 \[nucl-th\]](#).
  - [40] J. Gasser, M. Ivanov, E. Lipartia, M. Mojzis, and A. Rusetsky, *The European Physical Journal C* **26**, 13–34

- (2002).
- [41] D. Siemens, V. Bernard, E. Epelbaum, A. Gasparyan, H. Krebs, and U.-G. Meißner, *Phys. Rev. C* **94**, 014620 (2016), [arXiv:1602.02640 \[nucl-th\]](#).
  - [42] D. Siemens, V. Bernard, E. Epelbaum, A. M. Gasparyan, H. Krebs, and U.-G. Meißner, *Phys. Rev. C* **96**, 055205 (2017), [arXiv:1704.08988 \[nucl-th\]](#).
  - [43] V. Bernard, J. Gegelia, S. Ghosh, and U.-G. Meißner, (2025), [arXiv:2505.05941 \[hep-ph\]](#).
  - [44] M. Hoferichter, J. Ruiz de Elvira, B. Kubis, and U.-G. Meißner, *Phys. Rept.* **625**, 1 (2016), [arXiv:1510.06039 \[hep-ph\]](#).
  - [45] U.-G. Meißner and A. Rusetsky, *Effective Field Theories* (Cambridge University Press, 2022).
  - [46] M. Hoferichter and J. R. de Elvira, “Nucleon mass: trace anomaly and  $\sigma$ -terms,” (2025) [arXiv:2506.23902 \[hep-ph\]](#).
  - [47] J. A. McGovern and M. C. Birse, *Phys. Lett. B* **446**, 300 (1999), [arXiv:hep-ph/9807384](#).
  - [48] Z.-R. Liang, H.-X. Chen, F.-K. Guo, Z.-H. Guo, and D.-L. Yao, (2025), [arXiv:2502.19168 \[hep-ph\]](#).
  - [49] M. R. Schindler, D. Djukanovic, J. Gegelia, and S. Scherer, *Nucl. Phys. A* **803**, 68 (2008), [Erratum: *Nucl.Phys.A* 1010, 122175 (2021)], [arXiv:0707.4296 \[hep-ph\]](#).
  - [50] N. D. Conrad, A. Gasparyan, and E. Epelbaum, *PoS CD2021*, 074 (2024).
  - [51] J. Kambor and M. Mojzis, *JHEP* **04**, 031 (1999), [arXiv:hep-ph/9901235](#).
  - [52] O. Thim, A. Ekström, and C. Forssén, *Phys. Rev. C* **109**, 064001 (2024), [arXiv:2402.15325 \[nucl-th\]](#).