

Photometric Analysis of TCP J20171288+1156589 - WZ Sge Type Dwarf Nova with Delayed Ordinary Superhumps Emergence

Alexander Tarasenkov^{1,2*}, Sergey Shugarov^{2,3}, Natalia Ikonnikova²,
Marina Burlak², Sergey Nazarov⁴, Sjoerd Dufoer⁵

¹Institute of Astronomy of the Russian Academy of Sciences, 48
Pyatnitskaya Str., Moscow, 119234, Russia.

²Sternberg Astronomical Institute, Lomonosov Moscow State University,
Universitetsky Pr., 13, Moscow, 119234, Russia.

³Astronomical Institute of the Slovak Academy of Sciences, Tatranská
Lomnica, 059 60, The Slovak Republic.

⁴Crimean Astrophysical Observatory, Nauchny, 298409, Crimea.

⁵Vereniging voor Sterrenkunde, Zeeweg 96, Brugge, 8200, Belgium.

*Corresponding author(s). E-mail(s): tarasenkov@inasan.ru;

Abstract

We present the results of photometric analysis of WZ Sge type dwarf nova TCP J20171288+1156589. This object exhibited an outburst with a large amplitude of > 7.9 magnitudes and was observed for over a month. The photometric evolution of the superoutburst was atypical for WZ Sge-type dwarf novae. Periodogram analysis reveals early superhumps with the most probable period of 0.0611 ± 0.0001 days during the initial decline. After a plateau phase of approximately 11 days, ordinary superhumps (likely stage B) emerged with a period of 0.0616 ± 0.0001 days, corresponding to a superhump excess of $\epsilon = 0.008$ correspondingly. This delay in the onset of ordinary superhumps is an unusual feature among WZ Sge stars. We evaluated the main parameters of the system: mass ratio $q = M_{RD}/M_{WD} = 0.06 \pm 0.005$, yielding component masses of $M_{WD} \sim 1.0 \pm 0.15 M_{\odot}$ for the white dwarf and $M_{RD} = 0.06 \pm 0.01 M_{\odot}$ for the donor. The estimated distance to the system is ~ 850 pc, and the binary separation is $a = 0.67 \pm 0.03 R_{\odot}$.

Keywords: cataclysmic variable stars, dwarf novae, WZ Sagittae stars, photometry, superhumps, TCP J20171288+1156589

1 Introduction

Dwarf novae are a class of cataclysmic variable stars, which demonstrate regular outbursts with an amplitude of several stellar magnitudes. They are close binary systems consisting of a white dwarf and a low-mass donor star that fills its Roche lobe. The material of the donor, flowing out through the inner Lagrange point, forms an accretion disk around the white dwarf (Warner 1995; Osaki 2005; Kato et al. 2009a). Depending on the morphology of the light curves during outbursts and the regularity of the outbursts, different types of dwarf novae are distinguished. In particular, SU UMa dwarf novae exhibit 2 types of outbursts: normal outbursts with amplitudes of 2 – 6 magnitudes and superoutbursts, which usually have amplitudes from several tenths to one magnitude larger (for example, see light curves in Sibgatullin et al. (2024); Pavlenko et al. (2025)), than normal outbursts, and last longer (several weeks) (Samus' et al. 2017). Superoutbursts are caused by a combination of thermal and tidal instabilities (Osaki 1989). Among the SU UMa dwarf novae, a subgroup of WZ Sge objects is distinguished, which exhibit a supercycle (the time between two successive superoutbursts) of great duration - years or tens of years, as well as a large superoutburst amplitude - up to 8-10 magnitudes (Kato 2015).

Some WZ Sge-type stars exhibit repeated outbursts, known as rebrightenings, shortly after the end of their initial outburst. These are short-term increases in the object's brightness, typically lasting 2-3 days and with an amplitude 1-3 magnitudes smaller than the primary outburst. Some stars have been observed to undergo as many as 11 such rebrightenings, but most often, only one rebrightening occurs. A more detailed description of rebrightenings can be found in the articles Uemura et al. (2008); Kato et al. (2009b); Namekata et al. (2017); Pavlenko et al. (2019); Shugarov and Malashevich (2021); Lipunov et al. (2024). A possible explanation for this phenomenon is given in paper Meyer and Meyer-Hofmeister (2015).

During superoutbursts, so-called superhumps in light curves of SU UMa-type dwarf novae are observed. Superhumps are periodic modulations in the light curve with an amplitude of several tenths of magnitude, superimposed on the main trend. The emergence of superhumps is explained by the development of tidally driven eccentric instability in the elliptical accretion disk around the white dwarf (Osaki 1996). The period of the superhumps corresponds to the period of beats between the orbital period of the system and the precession period of the accretion disk (Kato et al. 2009a). Superhumps are conventionally divided into "early", "ordinary", and "late". Early superhumps are observed only in dwarf novae of the WZ Sge subclass. They have small amplitudes (usually less than 0.1 magnitudes). Their shape is a double wave, and their period is almost equal to the orbital period. They then change to ordinary superhumps, with an amplitude of up to 0.3 magnitudes and a period several percent longer than the orbital period. The optical period usually changes slightly as the superoutburst develops. The evolution of superhumps is discussed in more detail in the following papers: Warner (1995); Osaki (2005); Kato et al. (2009a); Kato (2015); Kato et al. (2020).

However, there are stars in which the superhump evolution deviates from the standard. This paper presents the discovery and photometric study of

TCP J20171288+1156589 (hereafter referred to as TCP J2017), a recently discovered WZ Sge-type dwarf nova which showed atypical superhump development.

2 Discovery of TCP J2017 and Observational Campaign

TCP J2017 was discovered by Yuji Nakamura on 2025-07-19 using 0.10-m f/3 refractor equipped with a CMOS camera. The transient had an unfiltered magnitude of $15^m.4$, with a limiting magnitude of $16^m.5$. Discovery information is available at ¹. The object was also independently detected by the ASAS-SN (Shappee et al. 2014a; Kochanek et al. 2017a) survey system on 2025-07-23.35 and designated ASASSN-25de. Alert notice and primary information about TCP J2017 was acquired using Astro-COLIBRI platform (Reichherzer et al. 2021, 2023), after which monitoring photometric observations were started.

For photometric monitoring of TCP J2017 several telescopes were used. Photometric observations at the Crimean Astronomical Station were conducted using the 600-mm RC600 telescope. The telescope is equipped with a back-illuminated 2048x2048 camera with a $13.5 - \mu\text{m}$ pixel size, a field of view of approximately $22'$, and an image scale of $\approx 0.7''/\text{pixel}$. Observations of TCP J2017 were also carried out with the 500-mm f/8 Astrosib RC500 telescope of the RCO network (Ibrahimov et al. 2025). It is equipped with a FLI Kepler 4040 FSI camera with Johnson-Cousins-Bessel *UBVRI* filters and BACHES echelle spectrograph. The observer can switch between photometric and spectroscopic modes at any time. The photometric mode was used to study TCP J2017. Also, the 600-mm (f/12.5) Zeiss telescope of Astronomical Institute of the Slovak Academy of Sciences at Stará Lesná equipped with a FLI ML3041 CCD camera was used. The Newton telescope of the Crimean Astrophysical Observatory, with an aperture of 350 mm and a focal length of 1500 mm, equipped with a QHY 600M IMX455 CMOS camera, was also used for observations.

Table 1 Stellar magnitudes and colours of stars, used for photometry of TCP J2017

N	U	B	V	R	I	U-B	B-V	V-R	R-I
1	14.962	14.766	13.998	13.549	13.194	0.196	0.768	0.449	0.355
2	15.207	14.983	14.245	13.829	13.505	0.224	0.738	0.416	0.324
3	14.026	14.026	13.490	13.154	12.883	0.000	0.536	0.336	0.271
4	15.565	15.502	14.825	14.416	14.079	0.063	0.677	0.409	0.337
5	15.994	16.046	15.572	15.270	15.013	-0.052	0.474	0.302	0.257
6	15.843	15.849	15.146	14.727	14.376	-0.006	0.703	0.419	0.351
7	17.199	15.895	14.558	13.856	13.317	1.304	1.337	0.702	0.539
8	15.247	15.203	14.719	14.429	14.178	0.044	0.484	0.290	0.251
9	16.162	15.940	15.202	14.771	14.405	0.222	0.738	0.431	0.366
10	16.338	16.317	15.624	15.211	14.859	0.021	0.693	0.413	0.352
11	16.838	16.752	16.038	15.622	15.278	0.086	0.714	0.416	0.344
12	-	18.607	17.597	17.063	16.631	-	1.010	0.534	0.432

¹<http://www.cbat.eps.harvard.edu/unconf/followups/J20171288+1156589.html>

Table 2 Log of TCP J2017 observations. The designations of the telescopes are as follows: Z-600 - 600 mm Zeiss telescope (Stará Lesná), CRAO - 350 mm Newton (Crimea), RC600 - 600 mm Ritchey-Chrétien (Crimea), K500 - 500 mm Astrosib RC500 (RCO-Kislovodsk)

JD-2400000	U	NU	B	NB	V	NV	R	NR	I	NI	Telescope
60876.56	15.097	2	15.815	2	15.752	3	15.728	3	15.690	3	Z-600
60877.40	-	-	-	-	15.896 ¹	104	-	-	-	-	CR350
60877.40	15.265	3	15.847	3	15.906	303	15.815	3	15.773	2	RC600
60878.37	15.401	3	16.010	3	16.042	342	15.948	3	15.888	3	RC600
60878.42	-	-	-	-	16.043 ¹	56	-	-	-	-	CR350
60879.38	15.599	3	16.140	3	16.177	237	16.074	3	16.014	3	RC600
60879.41	-	-	-	-	16.184 ¹	96	-	-	-	-	CR350
60880.39	15.727	3	16.283	3	16.290	312	16.195	3	16.123	3	RC600
60882.40	-	-	-	-	16.490 ¹	100	-	-	-	-	CR350
60882.46	15.993	3	16.486	3	16.503	181	16.372	3	16.308	3	RC600
60883.35	-	-	-	-	16.598	105	-	-	-	-	K500
60883.38	-	-	-	-	16.602 ¹	79	-	-	-	-	CR350
60883.40	16.165	3	16.636	3	16.608	200	16.488	3	16.411	3	RC600
60884.33	-	-	-	-	16.686	135	-	-	-	-	K500
60884.39	16.248	3	16.723	3	16.695	166	16.553	3	16.468	3	RC600
60885.34	-	-	-	-	16.795	135	-	-	-	-	K500
60885.40	16.343	3	16.811	3	16.799	161	16.667	3	16.581	3	RC600
60886.37	-	-	-	-	16.915	105	-	-	-	-	K500
60887.38	-	-	-	-	16.977	105	-	-	-	-	K500
60888.37	-	-	-	-	17.087	120	-	-	-	-	K500
60888.39	16.679	3	17.150	3	17.092	140	16.951	3	16.828	3	RC600
60888.47	16.660	2	17.096	3	17.094	3	16.889	3	16.967	3	Z-600
60892.55	-	-	-	-	17.104	2	17.030	3	-	-	Z-600
60893.38	16.701	3	17.167	3	17.138	3	16.993	100	16.922	3	RC600
60893.41	16.730	7	17.231	9	17.120	9	17.053	9	16.894	9	Z-600
60894.36	16.791	3	17.266	3	17.204	3	17.108	92	17.047	3	RC600
60894.39	-	-	17.165	6	17.257	6	17.166	6	16.921	6	Z-600
60896.32	-	-	-	-	17.290	113	-	-	-	-	K500
60896.36	16.961	2	17.459	2	17.405	2	17.259	2	17.233	1	RC600
60898.29	-	-	18.214	5	18.388	5	18.064	5	-	-	RC600
60898.39	-	-	-	-	18.098	8	-	-	-	-	K500
60900.33	-	-	-	-	-	-	18.531	3	-	-	RC600
60900.34	-	-	-	-	18.370	13	-	-	-	-	K500
60902.40	-	-	-	-	-	-	19 ²	3	-	-	Z-600
60907.40	-	-	-	-	-	-	20 ²	3	-	-	Z-600

¹Unfiltered passband reduced to V.

²Limiting magnitude.

Also AAVSO observations were used for our analysis. These observations were obtained with the Fregenal de La Sierra (E-Eye) 40cm f/6.8 corrected Dall-Kirkham telescope in Spain, equipped with camera: QHY268M in unfiltered passband.

Photometry of the frames was carried out using the programs MaximDL and AstroimageJ (Collins et al. 2017). For a photometric study of the object, the *UBVRI* magnitudes of 12 nearby stars were determined using the Z-600 telescope (Slovakia) (see Figure 1). Magnitudes and colours of these stars are presented in Table 1. The

magnitudes were obtained by referencing the stars to a photometric standard near PU Vul (Henden and Munari 2006) over four nights. Note that the magnitudes of the stars around PU Vul were obtained by referencing the stars measured by Landolt (Landolt 1992). The same set of comparison stars was used for photometry of TCP J2017 on frames obtained by all telescopes. The data from all telescopes were converted to a single system, as close as possible to the standard Johnson-Cousins system, and corrected for systematic errors associated with differences in the instrumental systems of different instruments. However, we will designate this system as *UBVRI* further in the text. Log of photometric observations is shown in the Table 2.




Fig. 1 Finding charts for TCP J2017: left panel indicates TCP J2017 (marked as var) and comparison stars; right panel shows combined colour image, TCP J2017 is marked with an arrow and has a noticeable blue colour.

3 Historical Light Curve

TCP J2017 in the quiescent state is absent from the Gaia DR3 (Gaia Collaboration et al. 2023) catalogue, the ZTF (Bellm et al. 2019) observation archive (accessed via SNAD web tool (Malanchev et al. 2023)), and the Pan-STARRS (Chambers et al. 2016) catalogue. Given that Pan-STARRS has a limiting magnitude in the *g* filter of $23^m.3$, and TCP J2017 at the peak of the outburst reached $15^m.4$, we assume the lower limit of the outburst amplitude to be about $7^m.9$, which is a high value even for WZ Sge-type dwarf novae. ASAS-SN (Shappee et al. 2014b; Kochanek et al. 2017b) shows no evidence of previous outbursts over the past 13 years.

4 Observed Light Curve Features

TCP J2017 was observed for a month after its discovery, resulting in a multi-colour *UBVRI* light curve (See Figure 2). During about the first two weeks TCP J2017 demonstrated gradual fading at a rate of ~ 0.1 mag/day. Colour indices during this

phase of outburst are: $B - V = -0^m.03$, $V - R = 0^m.1$, $R - I = 0^m.06$, which are characteristic for dwarf nova outbursts. In early stage of outburst TCP J2017 has significant UV excess ($U - B = -0^m.58$), which decreases during outburst. To search for possible periodicities in the monotonic decay phase, the light curve combined from all telescopes was detrended by subtracting the polynomial trend. The search for periodicities was carried out using the WinEFK program using the Deeming (Deeming 1975) and Lafler-Kinman (Lafler and Kinman 1965) methods. Analysis reveals a low-amplitude periodic modulation with a period of 0.0576 ± 0.0001 or 0.0611 ± 0.0001 days. The periodogram with peaks corresponding to the identified periods is shown in Figure 3. A detailed justification for why we chose these periods for analysis, rather than those corresponding to other high peaks of the periodogram, is given in Section 6. This trend was present during the first 3-4 nights of monitoring and had an amplitude of about $0^m.02$ in V passband. In the following nights of the monotonic decline stage, the trend disappeared. The folded light curves for two intervals (JD 2460876–2460880 and 2460882–2460888) are shown in Figure 4 (left and right panels, respectively). For clarity, the averaged profile is overplotted on the observed points. We interpret this sinusoidal variations as early superhumps.




Fig. 2 Full lightcurve of TCP J2017. Data points are represented as circles and photometric limits - as triangles.

Figure 5 shows graphs of changes in colour indices from time. For better visualization on the lower part of Figure 5 the light curve in the V passband is shown, where only measurements in the filter, averaged over every night are plotted. One can see




Fig. 3 Periodogram of the light curve of TCP J2017 for the first 4 nights. Peaks corresponding to probable periods of early superhumps are marked.

a small (not more than 0.2 magnitudes) increase in all colour indices in the initial area of the fall of brightness (plateau). This is caused by gradual cooling, a decrease in the average temperature of the emitting areas of the accretion disk, a hot spot on it, and an accretion stream. These features make the main contribution to the total radiation. However, at the end of the plateau, the $B - V$ colour index began to decrease again. A similar effect was observed for V1006 Cyg (Pavlenko et al. 2024). We explain this change by the shrinking of the accretion disk, which exposed the hot white dwarf and allowed it to contribute to the system's total flux. Also, this effect can be explained by the re-appearance of accretion disk emission lines, which are suppressed at the initial stages of the outburst, in the outburst final stages. The Balmer lines contribute much more to the flux in the B passband compared to the V passband, leading to the observed "blueing" of the $B - V$ colour index. Differences in the spectra of dwarf novae in the outburst and quiescent state are shown, for example, in Han et al. (2020); Voikhanskaya (2018). Similar changes in colour indices occur in many SU UMa and WZ Sge-type stars (see Krushevska et al. (2024); Shugarov et al. (2021, 2015); Chochol et al. (2015)). We will describe the nature of these changes in more detail in Section 5.

After JD 2460888, the light curve reached a roughly constant brightness level ($17 - 17.2^m$) and ordinary superhumps began to develop. The most pronounced ordinary superhumps were observed on JD 2460893 and 2460894 (the 17th and 18th days after the start of observations). They had a characteristic sawtooth shape and an amplitude of $0^m.10 - 0^m.15$ in the R passband. The light curves for these two nights are shown in the Figure 6. Based on observations during these 2 nights, we determined the superhump period to be 0.0616 ± 0.0001 days. The superhumps profile folded with this period is shown in the Figure 7.




Fig. 4 Profiles of early superhumps of TCP J2017, folded with period 0.0576 ± 0.0001 days (upper panels) and 0.0611 ± 0.0001 (lower panels): first 4 nights (left panels) and other nights of primary decline phase (right panels). Data points are green circles, the average profile is shown by black connected dots.

After this stage, at JDs later than 2460896, the brightness of TCP J2017 began to decline rapidly at a rate of approximately 1 mag/day, and the object quickly faded below the detection limit of the telescopes used for the observations. No rebrightenings were detected after the main outburst. However, we do not exclude the possibility that TCP J2017 demonstrated the short-term decrease in brightness during plateau phase (so called "dip"), which is observed in some SU UMa-type dwarf novae (Kimura et al. 2016; Kato et al. 2023). This dip could be missed due to unfavourable weather conditions for observations.

A similar atypical development of superhumps was also observed in the star PNV J1842+4837 (V529 Dra) (Katysheva et al. 2013). This star also exhibited early superhumps in the first days after the outburst, after which the consistent periodic brightness fluctuations disappeared. Ordinary superhumps appeared only at the end of the outburst, during the rebrightening process. However, it can be assumed that this star did not undergo a rebrightening process, but rather a brief brightness drop of 5 magnitudes, after which the star's brightness returned to almost the same level and the monotonic brightness decline continued. No such brightness decline was observed in the star we studied. However, we do not rule out that it was missed due to bad weather conditions and its short duration.




Fig. 5 Colour index curves for TCP J2017 (upper panel). V passband light curve is shown in the lower panel.




Fig. 6 Ordinary superhumps of TCP J2017: 2025 August 5th (left panel) and 2025 August 6th (right panel).

5 Colour Variations

Analysis of the positions and tracks of cataclysmic variable stars in two-colour diagrams can provide important information about certain physical processes occurring in the system. The tracks were calculated theoretically in papers [Mayo et al. \(1980\)](#); [Echevarría and Jones \(1983\)](#); [Cannizzo and Kenyon \(1987\)](#) and others. The authors accounted for the variable emission from the accretion disk using the law found




Fig. 7 Ordinary superhumps of TCP J2017 folded with period 0.0616 ± 0.0001 days.

by (Shakura and Sunyaev 1973), as well as taking into account various emission sources in the system at different stages of the outburst and the influence of emission lines to calculate the integrated flux in the *UBVRI* passbands. Theoretical tracks and loops were plotted on various two-colour diagrams, which showed good agreement with the observational results (Bruch 1984; Echevarria and Jones 1984).

One of the goals of our work was to plot the tracks of TCP J2017 on two-colour diagrams and interpret them.

In Section 4, we already showed that the object’s colour indices change slightly during outbursts and brightness declines. To study these changes in more detail, we plotted the object’s positions and tracked its movement on two-colour diagrams (see Figure 8).

The $U-B/B-V$ diagram plots the positions of the field stars from Table 1 (see left panel of Figure 8). The interstellar reddening value, $E(B-V)$, calculated using various extinction calculators^{2 3 4}, ranges from 0.11 to 0.23 for distances between 0.6 and 1 kpc. Figure 8 (left panel) shows that the average reddening value is $E(B-V) = 0.18$.

Most of the stars we measured are most likely main-sequence (MS) stars of spectral class F-G. The distance to these stars can be estimated as no more than 1 kpc. The fact that nearer stars exhibit less reddening is consistent with the location of neighbouring stars on the two-colour diagram. However, if giants are present among

²<http://argonaut.skymaps.info/query>

³https://astro.acri-st.fr/gaia_dev/#extinction

⁴<https://ned.ipac.caltech.edu/forms/calculator.html>




Fig. 8 Two colour diagrams of TCP J2017: (a) $U - B - B - V$ (b) $B - V - V - R$ (c) $V - R - R - I$.

these stars, their distance and reddening would be greater. According to Kato (2015) (see Chapter 6), the absolute magnitude of dwarf novae during an outburst is approximately $+5^m$, which is comparable to that of F-G main-sequence stars. Therefore, considering the range of probable reddening values from the mentioned calculators and our own approximate estimate, we adopted a mean value of $E(B - V) = 0.18$.

The position of TCP J2017 and the track from the outburst to the onset of decay are shown by blue symbols and a dark green arrow. If we shift this position by the interstellar extinction value, the average colour temperature of the source will vary from $\sim 15,000K$ at the beginning of the outburst to $\sim 10,000K$ by the end of the plateau. We did not conduct later observations, during the dramatic fading and final decline in the U passband.

Middle and right panels of Figure 8 similarly show the positions of the comparison stars and TCP J2017. We conducted multicolour observations not only during the outburst and plateau, but also during the rapid and final brightness decline. During this decline, the $B - V$ colour index of TCP J2017 decreased by approximately 0.2 mag compared to the value at the beginning of the outburst. The $V - R$ and $R - I$ indices continued to increase.

After the outburst, the accretion disk's size decreases, its temperature and luminosity drop, and its colour indices increase. We assume that the white dwarf was obscured by the accretion disk during the outburst, preventing its radiation from reaching the observer. However, by the end of the outburst, we suppose that the disk's size has decreased, and the hot white dwarf has become visible. In this case, fluxes in the blue and UV spectral regions should increase due to the additional light from the white dwarf and also due to the manifestation of emission lines that were suppressed during the outburst. We indeed observe similar behaviour in the blue spectral region (B passband) (track to the right in the middle panel of Figure 8, but we have no observations in the U passband. In the VRI photometric passbands, the contribution from the hot white dwarf is small, with its maximum contribution occurring in the UV. Therefore, the cooling disk continues to increase the "redder" $V - R$ and $R - I$ colour indices, as seen in the right panel of Figure 8.

The average colour temperature we found during the outburst, 10,000-15,000K, is higher than the average for similar stars. Most SU and WZ dwarf novae have average temperatures between 8,000 and 12,000K. Some of these stars are ASASSN-19oc (Krushevska et al. 2024), AY Lac (Shugarov et al. 2021), MO Psc (Shugarov et al. 2015), OV Boo (Golysheva et al. 2017), RZ LMi (Shugarov et al. 2018), V1838 Aql (Golysheva and Shugarov 2014), etc. However, there are also hotter stars, with temperatures from 12,000 to 20,000K, these are CV Tri (Chochol et al. 2009), V627 Peg (Zemko et al. 2013), PR Her (Shugarov et al. 2015) and other objects.

We do not rule out the presence of a small systematic error in the determination of stellar magnitudes, although the calibration was carried out over several nights. This error may primarily affect the I passband. The long-wavelength cutoff of this band is determined by the sensitivity roll-off of the specific CCD chip, which varies among different CCD models and may potentially change over time, rather than being strictly constant. This is evident in the right panel of Figure 8.

It is impossible to correct the positions of the stars for interstellar reddening in such a way that they would lie precisely on the main sequence. However, this error does not exceed $0^m.05$ and does not alter our main conclusions or the estimates of the system parameters.

6 Main Results

For TCP J2017, we found two possible periods of the early superhumps - 0.0576 ± 0.0001 and 0.0611 ± 0.0001 days. This period is very close to the orbital period in similar systems (Kato et al. 2009a; Kato 2015). At the end of the outburst, we detected the period of ordinary superhumps (0.0616 ± 0.0001 days), probably in the "B" stage.

Since TCP J2017 is absent in the Gaia catalogues, the distance cannot be calculated using parallax, but it can be estimated using the standard formula

$$d = 10^{0.2 \cdot (V - M_v + 5 - A_v)}, \quad (1)$$

where A_v is the extinction in the V passband, V is the peak magnitude during the outburst in the V passband, M_v is the absolute magnitude of the object during the outburst. We put the value $V \approx 15.5$ based on the magnitude of the object at the time of detection and observations at early outburst stage (see Table 2). According to Kato (2015), the absolute magnitude at the maximum of a superoutburst of SU UMa-type stars is

$$M_v(max) = 5.640.259 \times P_{orb}(hr). \quad (2)$$

For P_{orb} of most stars of the WZ Sge-type systems and also TCP J2017 this formula gives $M_v(max) = 5.3$. So, $VM_v = 10.2$. The value of interstellar extinction $A_v = 3.1E(B - V) \sim 0.55$ mag. As a result, we get a distance to the system TCP J2017 $d \approx 850$ pc.

With known both the orbital period (P_{orb}) and the period of the ordinary superhumps (P_{ord}), it becomes possible to calculate the superhump excess value (ϵ):

$$\epsilon = (P_{ord}/P_{orb}) - 1. \quad (3)$$

In the case of TCP J2017 $\epsilon = (0.0616/0.0576) - 1 = 0.07$ or $\epsilon = (0.0616/0.0611) - 1 = 0.008$ for two possible early superhumps periods. These values, especially the second one, are typical for SU UMa and WZ Sge-type stars (see right panel of Figure 19 in Kato et al. (2009a) for the distribution of ϵ for SU UMa stars). The mass ratio of stellar components q ($q = M_{RD}/M_{WD}$, where M_{RD} is the mass of the donor star, and M_{WD} is the mass of the white dwarf) can be estimated from the value of ϵ . According to Kato (2022), we get $q = 0.29 \pm 0.03$ or $q = 0.06 \pm 0.005$ respectively.

Masses of red dwarf in TCP J2017 system can be approximately estimated from semi-empirical sequence as $0.11 \pm 0.03 M_{\odot}$ or $0.06 \pm 0.01 M_{\odot}$ respectively (see Tables 2, 6 and Figure 9 in Knigge et al. (2011)). Since we found the value $q = M_{RD}/M_{WD}$ for TCP J2017, the mass of the white dwarf in this system will be equal to $M_{WD} \sim 0.4 \pm 0.15 M_{\odot}$ or $M_{WD} \sim 1.0 \pm 0.15 M_{\odot}$ correspondingly.

From the component masses and orbital period, the distance between the components in the system can be obtained using Kepler's third law, modified by Newton. Thus, $a = 0.50 \pm 0.03 R_{\odot}$ in case of the first period and $a = 0.67 \pm 0.03 R_{\odot}$ in case of the second period.

Since WZ Sge-type systems are typically characterized by a mass ratio q of $0.07 - 0.08$ (Kato 2015), that's why we consider the period of early superhumps of 0.0611 ± 0.0001 days, corresponding to the value of $q = 0.06 \pm 0.005$, represents the actual behaviour of the system. The period 0.0576 ± 0.0001 days, which, although giving a slightly higher peak on the periodogram, is 1-day alias to the true one. Its obvious manifestation is due to the fact that all the instruments used are at similar longitudes, so the light curve has regular gaps that occur during daytime. Other alias periods were not considered because they yield values of the system parameters that are in principle impossible for dwarf novae.

7 Discussion

TCP J2017 is clearly a WZ Sge dwarf nova, as it exhibited an outburst with an amplitude of at least 7.9 magnitudes, during which early and ordinary superhumps were detected, and archival observations have not revealed previous outbursts for at least the last 13 years. However, its photometric behaviour during outburst is quite unusual. After the presence of early superhumps during the first 3-4 days, a phase of monotonous decline lasting more than 10 days before the emergence of ordinary superhumps sets in. Such a long waiting time between early and ordinary superhumps stages is not typical for outbursts of WZ Sge-type dwarf novae. Such photometric behaviour usually indicates an extremely energetic flare. In particular, the object MASTER OT J030227.28+191754.5, which demonstrates the most delayed occurrence of ordinary superhumps (30-32 days after outburst start) had an outburst amplitude of 10.2 magnitudes (Tampo et al. 2024). The authors suggest that such a large amplitude could be due to a massive, extremely low-viscosity disk, as well as a possible low inclination of the system. Another similar dwarf nova, GOTO065054+593624 (an ordinary superhumps event occurred 13.8 days after discovery), had a outburst amplitude of 8.5 magnitudes (Killestein et al. 2025). Considering the large amplitude of TCP J2017

(> 7.9 magnitudes) and low donor mass $M_{RD} = 0.06 \pm 0.01 M_{\odot}$, we assume a similar high-amplitude outburst mechanism of this object.

However, the TCP J2017 is distinguished by the absence of dips in the light curve, which often appear in the light curves of dwarf novae of the WZ Sge type. Although in our case the narrow dip could have been missed due to bad weather, its possible absence does not contradict the high amplitude of the outburst. Thus, MASTER OT J030227.28+191754.5 did not demonstrate a dip during the outburst. System TCP J18173469+1803499 (with an amplitude of 8 magnitudes) also did not exhibit a dip, although in this case, ordinary superhumps developed more quickly - on the 8th day after detection (Tarasenkova et al. 2024).

8 Conclusion

We present a comprehensive photometric study of the dwarf nova TCP J20171288+1156589 in outburst. Our analysis allows us to confidently classify this object as belonging to WZ Sge subgroup. The key parameters supporting this classification include a large outburst amplitude of more than 7.9 magnitudes and the presence of early superhumps with a period of 0.0611 ± 0.0001 days in the early stages of the outburst. The period of early superhumps in WZ Sge-type dwarf novae is very close to the orbital period of the system P_{orb} . Therefore, we consider them to be approximately equal and use them to estimate other fundamental parameters of the system.

The object exhibited an atypical superhump evolution. The early superhumps, observed during the first 3 – 4 nights of the outburst, ceased, and were followed by a plateau in the light curve. Ordinary superhumps, which we associate with stage B superhumps, appeared only after approximately 11 days, on JD 2460893 and JD 2460894, with a period of $P_{ord} = 0.0616 \pm 0.0001$ days. This delay in the onset of ordinary superhumps relative to the early superhump phase is an unusual feature among WZ Sge-type stars.

From the observed periods, we calculated the superhump excess to be $\epsilon = (P_{ord}/P_{orb}) - 1 = 0.008$. This value is typical for SU UMa and WZ Sge-type dwarf novae. This excess corresponds to system components mass ratio of $q = M_{RD}/M_{WD} = 0.06 \pm 0.005$. Using empirical relations, we estimated the component masses to be $M_{RD} = 0.06 \pm 0.01 M_{\odot}$ for donor, which appears to be a brown dwarf, as in some other WZ Sge-type systems (Kato 2015; Lipunov et al. 2024), and $M_{WD} \sim 1.0 \pm 0.15 M_{\odot}$ for the white dwarf primary.

The estimated distance to the system is $d \approx 850$ pc. Applying Kepler’s third law to the component masses and the orbital period, we derived the separation between the components in the binary system to be $a = 0.67 \pm 0.03 R_{\odot}$.

In summary, TCP J2017 is a WZ Sge-type dwarf nova that displays a non-typical superhump evolution, characterized by a significant delay between the early superhump phase and the development of late, ordinary superhumps. This makes it an interesting object for further studies aimed at understanding the accretion disk dynamics in extreme mass-ratio systems.

Data Availability Statement. All observational data are available upon reasonable request to the corresponding author.

Acknowledgements. The observations were performed at telescope Astrosib RC500 of shared research facility “Terskol observatory” of Institute of Astronomy of the Russian Academy of Sciences. The observations on RC600 telescopes of SAI MSU were carried out using the equipment purchased under the MSU Program of Development. AstroColibri was used to receive alert notice. The authors are grateful to Dr. V. P. Goranskij for providing the WinEfk program. The study was conducted under the state assignment of Lomonosov Moscow State University. A. Tarasenkov acknowledges the support of the Foundation for the Development of Theoretical Physics and Mathematics BASIS (project 25-2-1-39-1). The authors would like to thank the anonymous reviewer who provided useful and detailed comments which helped improve the paper.

Funding. Part of this work was supported by the Slovak Research and Development Agency under contract No. APVV-24-0160 and by Vedecká grantová agentúra (VEGA) No 2/0003/25 (S.Y.S.)

Conflict of interest. The authors declare no conflict of interest.

References

- Bellm EC, Kulkarni SR, Graham MJ, et al (2019) The Zwicky Transient Facility: System Overview, Performance, and First Results. *Publ. Astron. Soc. Pac.*131(995):018002. <https://doi.org/10.1088/1538-3873/aaecbe>, [arXiv:1902.01932](https://arxiv.org/abs/1902.01932) [astro-ph.IM]
- Bruch A (1984) A catalogue of UBV colours of cataclysmic variables. *Astron. Astrophys. Suppl.*56:441–448
- Cannizzo JK, Kenyon SJ (1987) The Spectral Evolution of Dwarf Nova Outbursts. *Astrophys. J.*320:319. <https://doi.org/10.1086/165545>
- Chambers KC, Magnier EA, Metcalfe N, et al (2016) The Pan-STARRS1 Surveys. *arXiv e-prints* [arXiv:1612.05560](https://arxiv.org/abs/1612.05560). <https://doi.org/10.48550/arXiv.1612.05560>, [arXiv:1612.05560](https://arxiv.org/abs/1612.05560) [astro-ph.IM]
- Chochol D, Katysheva NA, Shugarov SY, et al (2009) CCD photometry of the dwarf nova Trianguli 2008 - a new WZ Sge-type object. *Contributions of the Astronomical Observatory Skalnaté Pleso* 39(1):43–65
- Chochol D, Shugarov S, Katysheva N, et al (2015) Superoutburst of a New Sub-Period-Minimum Dwarf Nova CSS130418 in Hercules. *Acta Polytechnica CTU Proceedings* 2(1):165–169. <https://doi.org/10.14311/APP.2015.02.0165>
- Collins KA, Kielkopf JF, Stassun KG, et al (2017) AstroImageJ: Image Processing and Photometric Extraction for Ultra-precise Astronomical Light Curves. *Astron.*

- J.153(2):77. <https://doi.org/10.3847/1538-3881/153/2/77>, [arXiv:1701.04817](https://arxiv.org/abs/1701.04817) [astro-ph.IM]
- Deeming TJ (1975) Fourier Analysis with Unequally-Spaced Data. *Astrophys. Space Sci.*36(1):137–158. <https://doi.org/10.1007/BF00681947>
- Echevarría J, Jones D (1983) A photometric study of dwarf novae. *Rev. Mexicana Astron. Astrofis.*5:301–308
- Echevarria J, Jones DHP (1984) A period-colour relation for dwarf novae at minimum. *Mon. Not. R. Astron. Soc.*206:919–928. <https://doi.org/10.1093/mnras/206.4.919>
- Gaia Collaboration, Vallenari A, Brown AGA, et al (2023) Gaia Data Release 3. Summary of the content and survey properties. *Astron. Astrophys.*674:A1. <https://doi.org/10.1051/0004-6361/202243940>, [arXiv:2208.00211](https://arxiv.org/abs/2208.00211) [astro-ph.GA]
- Golysheva P, Shugarov S (2014) Multicolor photometric monitoring of a new WZ Sge-type star in Aquila. *Contributions of the Astronomical Observatory Skalnat Pleso* 43(3):312–318
- Golysheva P, Shugarov S, Goranskij V, et al (2017) Multicolor light curve analysis of eclipsing cataclysmic star OV Boo. *Central European Astrophysical Bulletin* 41:79–90
- Han Z, Boonruksar S, Qian S, et al (2020) Spectroscopic properties of the dwarf nova-type cataclysmic variables observed by LAMOST. *Publ. Astron. Soc. Jpn*72(5):76. <https://doi.org/10.1093/pasj/psaa065>, [arXiv:2007.14016](https://arxiv.org/abs/2007.14016) [astro-ph.SR]
- Henden A, Munari U (2006) UBV(RI)C photometric sequences for symbiotic stars. III. *Astron. Astrophys.*458(1):339–340. <https://doi.org/10.1051/0004-6361:20065662>
- Ibrahimov M, Sachkov M, Nalivkin M, et al (2025) The new 50-cm multi-purpose telescope of the Russian-Cuban observatory. *Contributions of the Astronomical Observatory Skalnat Pleso* 55(1):7–17. <https://doi.org/10.31577/caosp.2025.55.1.7>
- Kato T (2015) WZ Sge-type dwarf novae. *Publ. Astron. Soc. Jpn*67(6):108. <https://doi.org/10.1093/pasj/psv077>, [arXiv:1507.07659](https://arxiv.org/abs/1507.07659) [astro-ph.SR]
- Kato T (2022) Evolution of short-period cataclysmic variables: implications from eclipse modeling and stage A superhump method (with New Year’s gift). *arXiv e-prints* [arXiv:2201.02945](https://arxiv.org/abs/2201.02945). <https://doi.org/10.48550/arXiv.2201.02945>, [arXiv:2201.02945](https://arxiv.org/abs/2201.02945) [astro-ph.SR]
- Kato T, Imada A, Uemura M, et al (2009a) Survey of Period Variations of Superhumps in SU UMa-Type Dwarf Novae. *Publ. Astron. Soc. Jpn*61:S395–S616. <https://doi.org/10.1093/pasj/61.sp2.S395>, [arXiv:0905.1757](https://arxiv.org/abs/0905.1757) [astro-ph.SR]

- Kato T, Pavlenko EP, Maehara H, et al (2009b) SDSS J080434.20+510349.2: Eclipsing WZ Sge-Type Dwarf Nova with Multiple Rebrightenings. *Publ. Astron. Soc. Jpn*61:601. <https://doi.org/10.1093/pasj/61.3.601>, [arXiv:0903.1685](https://arxiv.org/abs/0903.1685) [astro-ph.SR]
- Kato T, Isogai K, Wakamatsu Y, et al (2020) Survey of period variations of superhumps in SU UMa-type dwarf novae. X. The tenth year (2017). *Publ. Astron. Soc. Jpn*72(1):14. <https://doi.org/10.1093/pasj/psz134>, [arXiv:1911.04645](https://arxiv.org/abs/1911.04645) [astro-ph.SR]
- Kato T, Itoh H, Vanmunster T, et al (2023) MASTER OT J055845.55+391533.4: SU UMa star with a dip and long rebrightening. *arXiv e-prints* [arXiv:2304.07695](https://arxiv.org/abs/2304.07695). <https://doi.org/10.48550/arXiv.2304.07695>, [arXiv:2304.07695](https://arxiv.org/abs/2304.07695) [astro-ph.SR]
- Katysheva N, Shugarov S, Chochol D, et al (2013) Multicolour Investigation of PNV J1842+4837 - a New WZ Sge-type Dwarf Nova in Draco. *Central European Astrophysical Bulletin* 37:335–344
- Killestein TL, Ramsay G, Kennedy M, et al (2025) GOTO065054+593624: An 8.5 mag amplitude dwarf nova identified in real time via Kilonova Seekers. *Astron. Astrophys.*699:A8. <https://doi.org/10.1051/0004-6361/202553823>, [arXiv:2501.11524](https://arxiv.org/abs/2501.11524) [astro-ph.SR]
- Kimura M, Isogai K, Kato T, et al (2016) ASASSN-15jd: WZ Sge-type star with intermediate superoutburst between single and double ones. *Publ. Astron. Soc. Jpn*68(4):55. <https://doi.org/10.1093/pasj/psw054>, [arXiv:1604.06344](https://arxiv.org/abs/1604.06344) [astro-ph.SR]
- Knigge C, Baraffe I, Patterson J (2011) The Evolution of Cataclysmic Variables as Revealed by Their Donor Stars. *Astrophys. J. Suppl. Ser.*194(2):28. <https://doi.org/10.1088/0067-0049/194/2/28>, [arXiv:1102.2440](https://arxiv.org/abs/1102.2440) [astro-ph.SR]
- Kochanek CS, Shappee BJ, Stanek KZ, et al (2017a) The All-Sky Automated Survey for Supernovae (ASAS-SN) Light Curve Server v1.0. *Publ. Astron. Soc. Pac.*129(980):104502. <https://doi.org/10.1088/1538-3873/aa80d9>, [arXiv:1706.07060](https://arxiv.org/abs/1706.07060) [astro-ph.SR]
- Kochanek CS, Shappee BJ, Stanek KZ, et al (2017b) The All-Sky Automated Survey for Supernovae (ASAS-SN) Light Curve Server v1.0. *Publ. Astron. Soc. Pac.*129(980):104502. <https://doi.org/10.1088/1538-3873/aa80d9>, [arXiv:1706.07060](https://arxiv.org/abs/1706.07060) [astro-ph.SR]
- Krushevska V, Shugarov S, Ochner P, et al (2024) Investigation of WZ Sge-type Dwarf Nova ASASSN-19oc: Optical Spectroscopy and Multicolor Light Curve Analysis. *Research in Astronomy and Astrophysics* 24(8):085002. <https://doi.org/10.1088/1674-4527/ad59ec>
- Lafler J, Kinman TD (1965) An RR Lyrae Star Survey with the Lick 20-INCH Astrograph II. The Calculation of RR Lyrae Periods by Electronic Computer. *Astrophys. J. Suppl. Ser.*11:216. <https://doi.org/10.1086/190116>

- Landolt AU (1992) UBVRI Photometric Standard Stars in the Magnitude Range 11.5 < V < 16.0 Around the Celestial Equator. *Astron. J.*104:340. <https://doi.org/10.1086/116242>
- Lipunov VM, Balanutsa PV, Pavlenko EP, et al (2024) Detection and Study of a Number of Transients on Telescopes of the MASTER Global Network and MASTER OT J044907.58+705812.7 as an Example. *Astronomy Reports* 68(12):1364–1375. <https://doi.org/10.1134/S1063772925701318>
- Malanchev K, Kornilov MV, Pruzhinskaya MV, et al (2023) The SNAD Viewer: Everything You Want to Know about Your Favorite ZTF Object. *Publ. Astron. Soc. Pac.*135(1044):024503. <https://doi.org/10.1088/1538-3873/acb292>, [arXiv:2211.07605](https://arxiv.org/abs/2211.07605) [astro-ph.IM]
- Mayo SK, Wickramasinghe DT, Whelan JAJ (1980) Theory and observations of the optical continuum and line spectra of accretion discs around white dwarfs. *Mon. Not. R. Astron. Soc.*193:793–824. <https://doi.org/10.1093/mnras/193.4.793>
- Meyer F, Meyer-Hofmeister E (2015) SU UMa stars: Rebrightenings after super-outburst. *Publ. Astron. Soc. Jpn*67(3):52. <https://doi.org/10.1093/pasj/psv023>, [arXiv:1504.00469](https://arxiv.org/abs/1504.00469) [astro-ph.SR]
- Namekata K, Isogai K, Kato T, et al (2017) Superoutburst of WZ Sge-type dwarf nova below the period minimum: ASASSN-15po. *Publ. Astron. Soc. Jpn*69(1):2. <https://doi.org/10.1093/pasj/psw107>, [arXiv:1610.04941](https://arxiv.org/abs/1610.04941) [astro-ph.SR]
- Osaki Y (1989) A Model for the Superoutburst Phenomenon of SU Ursae Majoris Stars. *Publ. Astron. Soc. Jpn*41(5):1005–1033. <https://doi.org/10.1093/pasj/41.5.1005>
- Osaki Y (1996) Dwarf-Nova Outbursts. *Publ. Astron. Soc. Pac.*108:39. <https://doi.org/10.1086/133689>
- Osaki Y (2005) The disk instability model for dwarf nova outbursts. *Proceedings of the Japan Academy, Series B* 81:291–305. <https://doi.org/10.2183/pjab.81.291>
- Pavlenko E, Nijima K, Mason P, et al (2019) ASASSN-18fk: A new WZ Sge-type dwarf nova with multiple rebrightenings and a new candidate for a superhumping intermediate polar. *Contributions of the Astronomical Observatory Skalnat Pleso* 49(2):204–216. <https://doi.org/10.48550/arXiv.1907.00623>, [arXiv:1907.00623](https://arxiv.org/abs/1907.00623) [astro-ph.SR]
- Pavlenko E, Shugarov S, Kato T, et al (2024) V1006 Cyg: SU UMa-type dwarf nova in the period gap that wobbles between subclasses. *Contributions of the Astronomical Observatory Skalnat Pleso* 54(2):117–127. <https://doi.org/10.31577/caosp.2024.54.2.117>

- Pavlenko EP, Sosnovskij AA, Antonyuk KA, et al (2025) SU UMa-type Dwarf Nova MASTER OT J212624.16+253827.2 in the Period Gap: Orbital Period and Negative Superhumps. *Astronomy Reports* 69(10):961–970. <https://doi.org/10.1134/S1063772925702208>
- Reichherzer P, Schüssler F, Lefranc V, et al (2021) Astro-COLIBRI-The COincidence LIBrary for Real-time Inquiry for Multimessenger Astrophysics. *Astrophys. J. Suppl. Ser.* 256(1):5. <https://doi.org/10.3847/1538-4365/ac1517>, [arXiv:2109.01672](https://arxiv.org/abs/2109.01672) [astro-ph.IM]
- Reichherzer P, Schüssler F, Lefranc V, et al (2023) Astro-COLIBRI 2—An Advanced Platform for Real-Time Multi-Messenger Discoveries. *Galaxies* 11(1):22. <https://doi.org/10.3390/galaxies11010022>, [arXiv:2212.00805](https://arxiv.org/abs/2212.00805) [astro-ph.IM]
- Samus’ NN, Kazarovets EV, Durlevich OV, et al (2017) General catalogue of variable stars: Version GCVS 5.1. *Astronomy Reports* 61(1):80–88. <https://doi.org/10.1134/S1063772917010085>
- Shakura NI, Sunyaev RA (1973) Black holes in binary systems. Observational appearance. *Astron. Astrophys.* 24:337–355
- Shappee BJ, Prieto JL, Grupe D, et al (2014a) The Man behind the Curtain: X-Rays Drive the UV through NIR Variability in the 2013 Active Galactic Nucleus Outburst in NGC 2617. *Astrophys. J.* 788(1):48. <https://doi.org/10.1088/0004-637X/788/1/48>, [arXiv:1310.2241](https://arxiv.org/abs/1310.2241) [astro-ph.HE]
- Shappee BJ, Prieto JL, Grupe D, et al (2014b) The Man behind the Curtain: X-Rays Drive the UV through NIR Variability in the 2013 Active Galactic Nucleus Outburst in NGC 2617. *Astrophys. J.* 788(1):48. <https://doi.org/10.1088/0004-637X/788/1/48>, [arXiv:1310.2241](https://arxiv.org/abs/1310.2241) [astro-ph.HE]
- Shugarov S, Malashevich S (2021) New Cataclysmic Variable in Pisces: A WZ Sge-subtype Star with Rebrightenings. *Peremennye Zvezdy* 41(6):6
- Shugarov S, Katysheva N, Gladilina N (2015) PR Her — a WZ Sge-type cataclysmic variable. *Contributions of the Astronomical Observatory Skalnat Pleso* 45(2):119–130
- Shugarov SY, Katysheva NA, Chochol D, et al (2018) Superhump and outburst activity of the cataclysmic variable RZ LMi in the U- and optical passbands. *Astrophys. Space Sci.* 363(5):100. <https://doi.org/10.1007/s10509-018-3299-9>
- Shugarov SY, Afonina MD, Zharova AV (2021) Multicolor Photometry of the WZ Sge-Type Cataclysmic Variable AY Lac. *Astrophysics* 64(4):458–472. <https://doi.org/10.1007/s10511-021-09707-x>
- Sibgatullin AB, Dodon VI, Galiullin II, et al (2024) An X-ray and Optical Study of the

- Dwarf Nova Candidate OGLE-BLG-DN-0064. *Astronomy Letters* 50(11):676–686. <https://doi.org/10.1134/S1063773725700033>, [arXiv:2504.05263](https://arxiv.org/abs/2504.05263) [astro-ph.HE]
- Tampo Y, Kato T, Isogai K, et al (2024) MASTER OT J030227.28+191754.5: An unprecedentedly energetic dwarf nova outburst. *Publ. Astron. Soc. Jpn*76(6):1228–1245. <https://doi.org/10.1093/pasj/psae082>, [arXiv:2408.13783](https://arxiv.org/abs/2408.13783) [astro-ph.SR]
- Tarasenkova AN, Sokolovsky KV, Korotkiy SA, et al (2024) Photometric study of dwarf nova TCP J18173469+1803499. *INASAN Science Reports* 9(4):160–164. <https://doi.org/10.51194/INASAN.2024.9.4.008>
- Uemura M, Arai A, Krajci T, et al (2008) Discovery of a WZ Sge-Type Dwarf Nova, SDSS J102146.44+234926.3: Unprecedented Infrared Activity during a Rebrightening Phase. *Publ. Astron. Soc. Jpn*60:227. <https://doi.org/10.1093/pasj/60.2.227>, [arXiv:0711.2831](https://arxiv.org/abs/0711.2831) [astro-ph]
- Voikhanskaya NF (2018) Outburst Cycle of the Dwarf Nova SS Cygni. *Astrophysical Bulletin* 73(1):84–97. <https://doi.org/10.1134/S1990341318010078>
- Warner B (1995) *Cataclysmic variable stars*, vol 28
- Zemko PO, Andreev MV, Chochol D, et al (2013) Photometric Investigation of the Dwarf Nova OT J213806.6+261957 in Pegasus. In: Di Stefano R, Orio M, Moe M (eds) *Binary Paths to Type Ia Supernovae Explosions*, pp 130–131, <https://doi.org/10.1017/S1743921312014858>