

# Odd viscosity and anomalous Hall effect in two dimensional electron systems with smooth disorder

D.S. Zohrabyan<sup>1,2</sup> and M.M. Glazov<sup>3</sup>

<sup>1</sup>*L. D. Landau Institute for Theoretical Physics, 142432 Chernogolovka, Russia*

<sup>2</sup>*Moscow Institute of Physics and Technology, 141701 Dolgoprudny, Russia*

<sup>3</sup>*Ioffe Institute, 194021 St. Petersburg, Russia*

A microscopic theory of odd viscosity in two-dimensional electron systems with smooth disorder and spin-orbit interaction is developed. It is shown that spin-orbit scattering gives rise to an off-diagonal component of the viscosity tensor. Hydrodynamic equations for spin and electric currents are derived for electrons interacting with smooth disorder. The contribution of odd viscosity to the anomalous Hall effect is calculated.

**1. Introduction.** In recent years, significant attention from researchers has been drawn to transport phenomena and effects in ultra-pure two-dimensional electronic systems, where scattering processes on static defects and phonons are suppressed compared to electron-electron scattering, scattering on rough channel boundaries, and smooth disorder [1–6]. In such systems, both ballistic and hydrodynamic regimes of electron propagation can be realized; in the latter, dissipation processes are controlled by electronic viscosity [7, 8]. Phenomena where the differences between ultra-pure systems and ordinary (diffusive) ones manifest most clearly are of particular interest. Among these are the normal and anomalous Hall effects and similar other spin-dependent phenomena [9–22].

Usually, the hydrodynamic regime of quasiparticle propagation is associated with a situation where interparticle collisions dominate over any other dissipative processes, allowing the quasiparticle distribution function to be represented as an equilibrium one (Fermi-Dirac function for electrons or Bose-Einstein for excitons), whose parameters (temperature, chemical potential, average – hydrodynamic – velocity, i.e., a shift of the distribution function in  $\mathbf{k}$ -space) are determined by external influences [23, 24]. However, a quasi-hydrodynamic transport regime can be realized if quasiparticles interact with smooth disorder, see, e.g., [15] and references therein. In this case, the relaxation times of the angular harmonics of the distribution function sharply decrease with increasing harmonic number, and the relaxation of the second angular harmonic, which determines viscosity, occurs four times faster than that of the first.

In connection with studies of hydrodynamic and quasi-hydrodynamic regimes of electronic transport, particular attention has been drawn to systems where there is an odd or off-diagonal component of the viscosity tensor,  $\eta_{xy} = -\eta_{yx}$  [25–30]. It is known that such a component of the viscosity tensor arises due to the action of the Lorentz force on electrons in an external magnetic field [6, 15, 31]. Contributions to  $\eta_{xy}$  due to electron spin polarization and spin-orbit interaction are also possible [16, 20]. Our goal is to develop an analytical theory of odd viscosity caused by spin-orbit interaction and scattering on smooth disorder. In this paper we derive macroscopic quasi-

hydrodynamic equations for a spin-polarized electron gas interacting with smooth disorder in the presence of spin-orbit coupling, obtaining a microscopic expression for the odd viscosity  $\eta_{xy}$ , and as an example, considering the anomalous Hall effect in the presence of odd viscosity.

**2. Model.** Consider a two-dimensional degenerate electron gas in the  $(xy)$  plane. An external electric field  $\mathbf{E}$  is applied in the plane of electron motion  $(xy)$ , and an external magnetic field is perpendicular to it  $\mathbf{B} \parallel z$ . Due to the Zeeman effect, the magnetic field creates a non-zero spin polarization of the electrons and also induces cyclotron motion of charge carriers under the action of the Lorentz force. In this work, we neglect quantum effects associated with the formation of Landau levels. The Hall effect arising in confined systems leads, as known, to the formation of a Hall field  $\mathbf{E}_H$ , perpendicular to the external  $\mathbf{E}$  and  $\mathbf{B}$  fields [32, 33]. As known, the Hall field  $\mathbf{E}_H = \mathbf{E}_H^n + \mathbf{E}_H^a$  has two contributions: a "normal" one ( $\mathbf{E}_H^n$ ), due to the action of the Lorentz force on electrons, and an anomalous one ( $\mathbf{E}_H^a$ ), which is proportional to the degree of electron spin polarization and related to spin-orbit interaction [34–42]. The latter will interest us here.

Let us introduce the  $2 \times 2$  spin density matrix  $\hat{\rho}_{\mathbf{k}} = f_{\mathbf{k}} \hat{I} + \mathbf{s}_{\mathbf{k}} \cdot \hat{\boldsymbol{\sigma}}$ , where  $\mathbf{k}$  is the electron quasi-wave vector,  $f_{\mathbf{k}} = (1/2) \text{Tr}\{\hat{\rho}_{\mathbf{k}}\}$  is the spin-averaged occupancy of state  $\mathbf{k}$ ,  $\mathbf{s}_{\mathbf{k}} = (1/2) \text{Tr}\{\hat{\rho}_{\mathbf{k}} \hat{\boldsymbol{\sigma}}\}$  is the distribution function of electrons by spin. Here  $\hat{I}$  is the identity matrix (hereafter omitted for brevity),  $\hat{\boldsymbol{\sigma}} = (\hat{\sigma}_x, \hat{\sigma}_y, \hat{\sigma}_z)$  is a pseudovector composed of the Pauli spin matrices. In this work, we neglect the wave-vector-odd splitting of conducting band states by spin, so the density matrix for  $\mathbf{B} \parallel z$  is diagonal and parametrized by two distribution functions  $f_{\mathbf{k}}^{\pm} = f_{\mathbf{k}} \pm s_{z,\mathbf{k}}$ . For what follows, this representation of the density matrix will be most convenient. The density matrix obeys the Boltzmann kinetic equation, from which one can obtain equations for the distribution functions  $f_{\mathbf{k}}^{\pm} \equiv f_{\mathbf{k}}^{\pm}(\mathbf{r}, t)$  in the following form [16]:

$$\frac{\partial f_{\mathbf{k}}^{\pm}}{\partial t} + \mathbf{v}_{\mathbf{k}} \cdot \frac{\partial f_{\mathbf{k}}^{\pm}}{\partial \mathbf{r}} + \mathbf{F} \cdot \frac{1}{\hbar} \frac{\partial f_{\mathbf{k}}^{\pm}}{\partial \mathbf{k}} = Q_{dis}^{\pm}\{f_{\mathbf{k}}^{\pm}\} \quad (1)$$

Here  $\mathbf{v}_{\mathbf{k}} \equiv \hbar^{-1} \partial \varepsilon_{\mathbf{k}} / \partial \mathbf{k} = \hbar \mathbf{k} / m$  is the "kinematic" electron velocity,  $\varepsilon_{\mathbf{k}} = \hbar^2 k^2 / 2m$  is its dispersion, and  $m$  is

the effective mass,

$$\mathbf{F} = e(\mathbf{E} + \mathbf{E}_H) + \frac{e}{c}[\mathbf{v}_\mathbf{k} \times \mathbf{B}], \quad (2)$$

is the force acting on the electron from the total electric field  $\mathbf{E} + \mathbf{E}_H$  arising in the channel due to charge accumulation at its edges and the magnetic field  $\mathbf{B}$ ,  $e < 0$  is the electron charge,  $c$  is the speed of light in vacuum. Here we are interested only in effects related to the emergence of odd viscosity due to scattering, so we do not take into account here contributions to the anomalous Hall effect due to Berry curvature of energy bands (anomalous electron velocity) and associated wave packet shift contributions (side jump) [16, 17, 43, 44]. On the right side of Eq. (1)  $Q_{dis}\{f_\mathbf{k}\}$  is the collision integral describing electron scattering on disorder.

We will consider only those scattering processes that leave electrons in the conduction band, and we will also neglect contributions from spin-flip processes. The scattering matrix element from a state with wave vector  $\mathbf{k}'$  to state  $\mathbf{k}$  in the first non-vanishing approximation in spin-orbit interaction can be written as

$$V_{\mathbf{k}'\mathbf{k}}^\pm = U_c(\mathbf{k} - \mathbf{k}') \pm i\xi U_v(\mathbf{k} - \mathbf{k}')[\mathbf{k}' \times \mathbf{k}]_z, \quad (3)$$

where the upper indices  $\pm$  number states with spin projection  $s_z = \pm 1/2$ ,  $U_{c,v}(\mathbf{q})$  are the Fourier transforms of the impurity potentials in the conduction and valence bands, respectively,  $\xi$  is a quantity characterizing the strength of spin-orbit interaction [43, 45]. The dimensionless small parameter characterizing the strength of spin-orbit interaction is  $\xi k_F^2 \ll 1$ , also  $\varepsilon_F \ll E_g$ , where  $\varepsilon_F$  and  $k_F$  are the Fermi energy and Fermi wave vector of the electrons.

The electron-impurity collision integral can be written in general form as [46]

$$Q_{dis}^\pm\{f_\mathbf{k}\} = \sum_{\mathbf{k}'} W_{\mathbf{k}'\mathbf{k}}^\pm (f_{\mathbf{k}'}^\pm - f_\mathbf{k}^\pm), \quad (4)$$

where

$$W_{\mathbf{k}'\mathbf{k}}^\pm = \frac{2\pi}{\hbar} \delta(\varepsilon_\mathbf{k} - \varepsilon_{\mathbf{k}'}) \{t_0(\theta) \pm t_1(\theta) \sin \theta\}, \quad \theta = \angle \mathbf{k}'\mathbf{k}. \quad (5)$$

Here the first term describes spin-independent scattering, and the second term describes asymmetric spin-dependent scattering (Mott effect) [36, 47–49],  $t_0(\theta)$ ,  $t_1(\theta)$  are even functions of angle  $\theta$ , with  $t_0(\theta) = n_{imp}|U_c(\theta)|^2$ , where  $n_{imp}$  is the impurity concentration,

$$t_1(\theta) = 2\pi D k_F^2 \xi n_{imp} \cdot U_v(\theta) \int_0^{2\pi} \frac{d\varphi}{2\pi} U_c(\varphi) U_c(\theta - \varphi), \quad (6)$$

where we introduced the notation  $U_{c,v}(\theta) \equiv U_{c,v}(2k \sin \theta/2)$ . Represent the nonequilibrium, i.e., induced by external electric and magnetic fields  $\mathbf{E}$  and  $\mathbf{B}$ , correction to the distribution function in the form

$$\delta f_\mathbf{k}^\pm = \sum_{n=1}^{\infty} \{f_{c,n}^\pm \cos n\varphi + f_{s,n}^\pm \sin n\varphi\}, \quad (7)$$

where  $f_{c,n}^\pm$ ,  $f_{s,n}^\pm$  are Fourier expansion coefficients depending on electron energy,  $\varphi_\mathbf{k} = \angle \mathbf{k}, Ox$ , and obtain

$$Q_{dis}^\pm\{f_\mathbf{k}^\pm\} = - \sum_{n=1}^{\infty} \frac{f_{c,n}^\pm \cos n\varphi + f_{s,n}^\pm \sin n\varphi}{\tau_n} \pm \sum_{n=1}^{\infty} \gamma_n (f_{s,n}^\pm \cos n\varphi - f_{c,n}^\pm \sin n\varphi). \quad (8)$$

The sum in the first line of Eq. (8) is the symmetric part of the collision integral, describing the relaxation of angular harmonics of the distribution function with relaxation times  $\tau_n$ , introduced in the standard way. The contribution in the second line of Eq. (8) corresponds to the asymmetric contribution to the collision integral, and the coefficients  $\gamma_n$  describe the conversion of even and odd harmonics due to spin-orbit interaction

$$\gamma_n = \frac{2\pi}{\hbar} D \int \frac{d\theta}{2\pi} t_1(\theta) \sin \theta \sin n\theta. \quad (9)$$

The quantities  $\tau_n$  and  $\gamma_n$  depend on electron energy. Note that in the case of a short-range potential, only the first harmonic of the distribution function is present in the asymmetric part of the collision integral [16, 17, 50], and the other contributions vanish:  $\gamma_n = \gamma_1 \delta_{1,n}$ . In the case of a long-range potential, all coefficients are non-zero  $\gamma_n \neq 0$ , in particular, the coefficient at the second angular harmonic  $\gamma_2$  determines the odd viscosity.

3. *Odd viscosity.* The components of the viscosity tensor for two-dimensional electrons can be derived following the standard approach outlined, e.g., in [51]. Considering a locally equilibrium distribution of electrons with hydrodynamic velocity in each spin branch  $\mathbf{u}^\pm \equiv \mathbf{u}^\pm(\mathbf{r}, t) = N_\pm^{-1} \sum_{\mathbf{k}} \mathbf{v}_\mathbf{k} f_\mathbf{k}^\pm$ , smoothly (on the scale of the mean free path) depending on coordinates, where  $N^\pm = \sum_{\mathbf{k}} f_\mathbf{k}^\pm$  is the concentration of electrons with a given spin projection, one can represent the components of the momentum flux tensor as  $\Pi_{\alpha\beta}^\pm = \sum_{\mathbf{k}} v_{\alpha,\mathbf{k}} v_{\beta,\mathbf{k}} f_\mathbf{k}^\pm$ . In the case of axial symmetry of interest to us, the term with the gradient of the momentum flux can be represented in the following convenient form

$$-\nabla \hat{\Pi}^\pm = \Delta(\eta_{xx}^\pm N^\pm \mathbf{u}^\pm) + \Delta[\eta_{xy}^\pm N^\pm \mathbf{u}^\pm \times \hat{\mathbf{z}}], \quad (10)$$

where two independent components of the (first) viscosity tensor are introduced:  $\eta \equiv \eta_{xx}^\pm$  (diagonal viscosity) and  $\eta_{xy}^\pm = -\eta_{yx}^\pm$  – the off-diagonal (odd) or Hall viscosity [25–29]. The latter is the subject of our interest. Note that in our problem formulation, the second viscosity does not manifest itself (general analysis is given, e.g., in [52]).<sup>1</sup>

<sup>1</sup> In Eq. (10), unlike the standard representation of the Navier-Stokes equations [53], the terms with the product of viscosity and concentration are under the Laplace operator. This is related to the specifics of scattering on static disorder. For an incompressible electron liquid, this difference is not essential.

Expanding the distribution function in angular harmonics, Eq. (7), and taking into account spatial gradients and magnetic field in the lowest (first) order, we obtain the diagonal viscosity

$$\eta_{xx}^{\pm} \equiv \eta^{\pm} = \frac{(v_F^{\pm})^2 \tau_2 (\varepsilon_F^{\pm})}{4}, \quad (11)$$

where  $v_F^{\pm} = \sqrt{2\varepsilon_F^{\pm}/m}$  is the electron velocity at the Fermi surface in the corresponding spin branch, and  $\tau_2$ , as indicated earlier, is the relaxation time of the second angular harmonic of the distribution function. In the relevant case of a small degree of electron polarization, the differences between  $\eta^+$  and  $\eta^-$  are negligible. The off-diagonal viscosity contains two contributions:

$$\eta_{xy}^{\pm} = \eta_{xy,H}^{\pm} + \eta_{xy,SO}^{\pm}, \quad (12)$$

where

$$\eta_{xy,H}^{\pm} = 2\omega_c \tau_2 \cdot \eta \quad (13)$$

is the Hall component of odd viscosity due to the action of the Lorentz force on electrons, where  $\tau_2$  and  $\eta$  are taken at the Fermi level [6, 15, 31];  $\eta_{xy,SO}^{\pm}$  is the contribution to odd viscosity due to spin-orbit interaction [16, 20, 54, 55]. The latter can be represented in a form similar to (13),

$$\eta_{xy,SO}^{\pm} = \pm \gamma_2 (\varepsilon_F^{\pm}) \tau_2 (\varepsilon_F^{\pm}) \eta^{\pm} \equiv \pm \eta_{xy,SO}^s (\varepsilon_F^{\pm}). \quad (14)$$

The odd viscosity of the electron gas due to spin-orbit interaction and averaged over spin branches takes the form

$$\eta_{xy,SO} = \frac{\eta_{xy,SO}^+ + \eta_{xy,SO}^-}{2} = P_s \varepsilon_F \frac{\partial \eta_{xy,SO}^s}{\partial \varepsilon_F}, \quad (15)$$

where  $P_s \ll 1$  is the degree of electron spin polarization

$$P_s \equiv \frac{N^+ - N^-}{N^+ + N^-} = -\frac{g\mu_B B}{2\varepsilon_F}, \quad (16)$$

$g$  is the electron g-factor,  $\mu_B$  is the Bohr magneton. The odd viscosity (15) depends on the type of impurity potential. In particular, as already noted above, for short-range scatterers  $\eta_{xy,SO} = 0$ . To analyze the dependence of  $\eta_{xy,SO}$  on the disorder correlation length, consider, as an example, Gaussian impurities

$$U_{c,v}(q) = U_0 \cdot e^{-q^2 r_0^2/4}, \quad (17)$$

where  $U_0 = \int U_{c,v}(\mathbf{r}) d^3r$  is the "strength" of the impurity potential,  $r_0$  is the correlation length. The dependence of odd viscosity on the dimensionless correlation length  $k_F r_0$  is shown in Fig. 1. In the limit  $k_F r_0 \gg 1$ , we have

$$\eta_{xy,SO}^{gauss} = \frac{4P_s}{\sqrt{3}} \cdot \xi \cdot k_F^2 r_0^2 \cdot \frac{\varepsilon_F^2}{U_0 \hbar n_{imp}}. \quad (18)$$




Figure 1. The graph shows the dependence of odd viscosity  $\eta_{xy,SO}$  (blue line) in the case of a Gaussian impurity potential (17) on the screening parameter  $k_F r_0$ . The odd viscosity is normalized to its asymptotic value  $\eta_{xy,SO}^{gauss}$  (18) at  $k_F r_0 \rightarrow \infty$ . The "effective" odd viscosity  $\eta_{xy,SO}^H$  (red line), which enters the expression for the Hall field (22), is also shown.

The expression for odd viscosity (15), due to the presence of smooth disorder and related to spin-orbit interaction, is one of the main results of this work.

4. *Spin-dependent hydrodynamic equations.* The kinetic equation (1), supplemented by appropriate boundary conditions at the system edges and the Poisson equation relating  $\mathbf{E}_H$  and the electron distribution function, allows, in principle, investigation of magnetotransport effects, including the anomalous Hall effect, for any type of disorder. In the case of smooth disorder,  $\tau_n$  rapidly decreases with increasing angular harmonic number  $n$ , so in the expansion (7) of the nonequilibrium distribution function in angular harmonics, it is sufficient to retain contributions with  $n = 1$  and 2. This significantly simplifies the problem and reduces the description of electronic transport to a hydrodynamic model. In particular, summing (1) over momenta, one can obtain continuity equations for  $N_{\pm}$  in standard form. Equations for fluxes are conveniently presented as a system of equations for electric current  $\mathbf{j} \equiv eN^+ \mathbf{u}^+ + eN^- \mathbf{u}^-$  and for spin current (current of the  $z$ -component of spin)  $\mathbf{i}^z \equiv eN^+ \mathbf{u}^+ - eN^- \mathbf{u}^-$ . The factor  $e$  in the expression for  $\mathbf{i}^z$  is introduced for convenience. Moreover, we represent each of the currents as a sum of a "normal" component, which does not contain spin-orbit contributions and can be calculated at  $\xi = 0$ , and an "anomalous" (spin-orbit) one, linear in  $\xi$ :  $\mathbf{j} = \mathbf{j}_n + \mathbf{j}_a$ ,  $\mathbf{i}^z = \mathbf{i}_n^z + \mathbf{i}_a^z$ . Correspondingly, the Hall field  $\mathbf{E}_H = \mathbf{E}_H^n + \mathbf{E}_H^a$ . Ultimately, we will be interested in  $\mathbf{j}_a$  – the spin-orbit contribution to the electric current and the corresponding component of the Hall field  $\mathbf{E}_H^a$ .

In particular, the "normal" component of electric cur-

rent obeys the standard equation,

$$\frac{\mathbf{j}_n}{\tau_1} - \omega_c[\mathbf{j}_n \times \hat{\mathbf{z}}] - \frac{e^2 N}{m}(\mathbf{E} + \mathbf{E}_H^n) = \eta \Delta \mathbf{j}_n + \eta_{xy,H}[\Delta \mathbf{j}_n \times \hat{\mathbf{z}}], \quad (19a)$$

containing the Lorentz force and odd viscosity related to the Lorentz force (13). All parameters  $\tau_1, \eta, \eta_{xy,H}$  are taken at energy equal to the Fermi energy; thus, the current depends on  $\varepsilon_F$  as a parameter. Similarly, the "normal" component of spin current, taking into account possible coordinate dependence of the degree of spin polarization  $P_s = P_s(\mathbf{r})$ , obeys the equation

$$e \frac{\varepsilon_F N}{m} \nabla P_s - P_s \frac{N}{m} e^2 \mathbf{E} - \zeta \cdot [e^2 \mathbf{E} \times [\nabla \times \mathbf{i}_n^z]] = \eta \Delta \mathbf{i}_n^z + \eta_s \Delta \mathbf{j}_n - \frac{\mathbf{i}_n^z}{\tau_1} - \frac{\mathbf{j}_n}{\tau_1^s}, \quad (19b)$$

where the following notations are introduced:  $\zeta = \eta/\varepsilon_F + \partial\eta/\partial\varepsilon_F$ ,  $\eta_s = P_s \varepsilon_F \cdot \partial\eta/\partial\varepsilon_F$  and  $(\tau_1^s)^{-1} = P_s \varepsilon_F \cdot \partial\tau_1^{-1}/\partial\varepsilon_F$ . Note that the flow of electric current is accompanied by the emergence of spin current both directly due to the presence of polarization in the system and due to the dependencies of momentum relaxation time and viscosity on Fermi energy, leading to their difference for electrons with different spin projections.

The equation for the "anomalous" component of electric current, taking into account the induced anomalous Hall field  $\mathbf{E}_H^a$ , has a somewhat more complex form:

$$\begin{aligned} & \frac{\mathbf{j}_a}{\tau_1} + \frac{\nabla p}{m} - \frac{N}{m} e^2 \mathbf{E}_H^a \\ &= \eta \Delta \mathbf{j}_a + \eta_s \Delta \mathbf{i}_a^z + \eta_{xy,SO}[\Delta \mathbf{j}_n \times \hat{\mathbf{z}}] + \eta_{xy,SO}^s[\Delta \mathbf{i}_n^z \times \hat{\mathbf{z}}] \\ & \quad + \Gamma_1[\mathbf{j}_n \times \hat{\mathbf{z}}] + \gamma_1[\mathbf{i}_n^z \times \hat{\mathbf{z}}] - \frac{\mathbf{i}_a^z}{\tau_1^s}, \end{aligned} \quad (19c)$$

where  $\Gamma_1 = P_s \varepsilon_F \cdot \partial\gamma_1/\partial\varepsilon_F$  and

$$\frac{\nabla p}{m} = P_s \left( \eta \Delta \mathbf{i}_a^z + \eta_{xy,SO}^s[\Delta \mathbf{j}_n \times \hat{\mathbf{z}}] + \gamma_1[\mathbf{j}_n \times \hat{\mathbf{z}}] - \frac{\mathbf{i}_a^z}{\tau_1} \right), \quad (19d)$$

is the "anomalous" pressure contribution from the zeroth harmonic of the distribution function, due to the fact that  $N_+ - N_-$  may depend on coordinates. This contribution is important but leads only to renormalizations of the corresponding coefficients on the right side of the equation for anomalous current (19c). Finally, the equation for the anomalous part of spin current has the form

$$\begin{aligned} & e \frac{\varepsilon_F N}{m} \nabla P_s - \omega_c[\mathbf{i}_a^z \times \hat{\mathbf{z}}] \\ &= \eta \Delta \mathbf{i}_a^z + \eta_{xy,H}[\Delta \mathbf{i}_a^z \times \hat{\mathbf{z}}] + \eta_{xy,SO}^s[\Delta \mathbf{j}_n \times \hat{\mathbf{z}}] \\ & \quad + \gamma_1[\mathbf{j}_n \times \hat{\mathbf{z}}] - \frac{\mathbf{i}_a^z}{\tau_1} - \frac{\mathbf{j}_a}{\tau_1^s}. \end{aligned} \quad (19e)$$




Figure 2. Schematic of an electron channel, infinite along the  $y$  axis and of width  $w$  along the  $x$  axis. The electric field is directed along the channel ( $\mathbf{E} \parallel y$ ), the magnetic field is perpendicular to the plane of particle motion ( $\mathbf{B} \parallel z$ ). The Hall effect leads to charge accumulation at the channel edges and  $\mathbf{E}_H \parallel x$ . The figure shows the parabolic current profile along the channel in the limit of hydrodynamic electron transport.

As above, all quantities depend on  $\varepsilon_F$  as a parameter. The system of equations for spin and electric currents (19) is one of the main results of this work.

In an unbounded homogeneous system, electric and spin currents do not depend on coordinates. For  $\mathbf{B} = 0$ , formula (19e) describes the relationship between spin current and electric current due to asymmetric scattering  $\mathbf{i}_a^z = \gamma_1[\mathbf{j}_n \times \hat{\mathbf{z}}]$  [43]. If  $\mathbf{B} \neq 0$ , then formula (19a) describes the relationship between field and current due to the normal Hall effect, and from (19c) one can derive the anomalous Hall conductivity for two-dimensional electrons in the form

$$\sigma_{xy}^a = -\sigma_{yx}^a = P_s \frac{e^2 N}{m} \frac{\partial}{\partial \varepsilon_F} (\gamma_1 \tau_1^2 \varepsilon_F), \quad (20)$$

where  $N = N^+ + N^-$ .

5. *Anomalous Hall effect in a conducting channel.* As an illustration of the developed theory, consider a conducting channel of width  $w$  along the  $x$  direction, along which an external electric field  $\mathbf{E} \parallel y$  is directed. The magnetic field  $\mathbf{B} \parallel z$  is perpendicular to the plane of electron motion. In this case, a Hall field  $\mathbf{E}_H \parallel x$  appears. In this geometry, there are no currents along the  $x$  axis. Let's find the  $y$ -component of electric current in the quasi-hydrodynamic limit, when  $l_1 l_2 \gg w$ , where  $l_{1,2} = v_F \tau_{1,2}$  are the relaxation lengths of momentum and the second angular harmonic of the distribution function. Solving (19a) with boundary conditions of zero current at the channel walls, we obtain: [16, 17, 50]

$$\mathbf{j}_n = \frac{e^2 N \mathbf{E}}{2m\eta} \left[ \left( \frac{w}{2} \right)^2 - x^2 \right], \quad (21)$$

which corresponds to Poiseuille flow with a parabolic profile. Substituting (21) into (19), taking into account boundary conditions and the continuity equation, we obtain the following expression for the anomalous contribution to the Hall field:

$$\mathbf{E}_H^a = [\hat{\mathbf{z}} \times \mathbf{E}] \left\{ \left( \frac{\Gamma_1}{\eta} - \gamma_1 \frac{\eta_s}{\eta^2} \right) \left[ \left( \frac{w}{2} \right)^2 - x^2 \right] - \frac{\eta_{xy,SO}^H}{\eta} \right\}, \quad (22)$$



where the first contribution corresponds to the contribution of asymmetric scattering [16, 50], and the second contribution is from odd viscosity, written in the form in which Hall viscosity usually enters the field expression [15]. With such notation, the "effective" odd viscosity will have the form  $\eta_{xy,SO}^H = \eta \cdot P_s \varepsilon_F \partial \gamma_2 \tau_2 / \partial \varepsilon_F$ , see the red curve in Fig. 1.

In the case of a short-range potential, the contribution of odd viscosity is absent. In the quasi-hydrodynamic regime, when  $l_2 \ll w \ll l_1$ , the contribution of odd viscosity due to impurity scattering can be significant.

**6. Conclusion.** The work develops a microscopic theory of odd (off-diagonal) viscosity of electrons in two-dimensional systems with smooth disorder and spin-orbit interaction. Analytical expressions for the components of the viscosity tensor are obtained, including the contribution of spin-orbit interaction  $\eta_{xy,SO}$ , which is absent in

the case of short-range scattering but significant for a long-range impurity potential. It is shown that this contribution is proportional to the spin polarization of carriers, the spin-orbit interaction constant, and depends on disorder parameters.

A closed system of equations for spin and electric currents is derived, taking into account both normal and anomalous contributions for the quasi-hydrodynamic transport regime, where the relaxation time of the first angular harmonic of the distribution function significantly exceeds the relaxation times of higher harmonics. The contribution of odd viscosity to the anomalous Hall field is obtained. This work is the first step towards constructing a microscopic theory of odd viscosity in interacting electronic systems.

*Acknowledgments.* The work was supported by the RSF Project No. 22-12-00211-Continuation.

- 
- [1] L. Levitov and G. Falkovich, Electron viscosity, current vortices and negative nonlocal resistance in graphene, *Nature Physics* **12**, 672 (2016).
  - [2] J. Crossno, J. K. Shi, K. Wang, X. Liu, A. Harzheim, A. Lucas, S. Sachdev, P. Kim, T. Taniguchi, K. Watanabe, T. A. Ohki, and K. C. Fong, Observation of the Dirac fluid and the breakdown of the Wiedemann-Franz law in graphene, *Science* **351**, 1058 (2016).
  - [3] D. A. Bandurin, I. Torre, R. K. Kumar, M. Ben Shalom, A. Tomadin, A. Principi, G. H. Auton, E. Khestanova, K. S. Novoselov, I. V. Grigorieva, L. A. Ponomarenko, A. K. Geim, and M. Polini, Negative local resistance caused by viscous electron backflow in graphene, *Science* **351**, 1055 (2016).
  - [4] P. J. W. Moll, P. Kushwaha, N. Nandi, B. Schmidt, and A. P. Mackenzie, Evidence for hydrodynamic electron flow in PdCoO<sub>2</sub>, *Science* **351**, 1061 (2016).
  - [5] R. Krishna Kumar, D. A. Bandurin, F. M. D. Pellegrino, Y. Cao, A. Principi, H. Guo, G. H. Auton, M. Ben Shalom, L. A. Ponomarenko, G. Falkovich, K. Watanabe, T. Taniguchi, I. V. Grigorieva, L. S. Levitov, M. Polini, and A. K. Geim, Superballistic flow of viscous electron fluid through graphene constrictions, *Nature Physics* **13**, 1182 (2017).
  - [6] P. S. Alekseev, Negative magnetoresistance in viscous flow of two-dimensional electrons, *Phys. Rev. Lett.* **117**, 166601 (2016).
  - [7] R. N. Gurzhi, Minimum of resistance in impurity-free conductors, *JETP* **17**, 521 (1963).
  - [8] R. N. Gurzhi, Hydrodynamic effects in solids at low temperatures, *Soviet Physics Uspekhi* **11**, 255 (1968).
  - [9] T. Scaffidi, N. Nandi, B. Schmidt, A. P. Mackenzie, and J. E. Moore, Hydrodynamic electron flow and Hall viscosity, *Phys. Rev. Lett.* **118**, 226601 (2017).
  - [10] F. M. D. Pellegrino, I. Torre, and M. Polini, Nonlocal transport and the hall viscosity of two-dimensional hydrodynamic electron liquids, *Phys. Rev. B* **96**, 195401 (2017).
  - [11] R. J. Doornenbal, M. Polini, and R. A. Duine, Spin-vorticity coupling in viscous electron fluids, *Journal of Physics: Materials* **2**, 015006 (2019).
  - [12] A. I. Berdyugin, S. G. Xu, F. M. D. Pellegrino, R. Krishna Kumar, A. Principi, I. Torre, M. Ben Shalom, T. Taniguchi, K. Watanabe, I. V. Grigorieva, M. Polini, A. K. Geim, and D. A. Bandurin, Measuring Hall viscosity of graphene's electron fluid, *Science* **364**, 162 (2019).
  - [13] P. S. Alekseev and M. A. Semina, Hall effect in a ballistic flow of two-dimensional interacting particles, *Phys. Rev. B* **100**, 125419 (2019).
  - [14] Y. O. Alekseev and A. P. Dmitriev, Giant Hall effect in the ballistic transport of two-dimensional electrons, *Phys. Rev. B* **104**, 085434 (2021).
  - [15] A. N. Afanasiev, P. S. Alekseev, A. A. Danilenko, A. P. Dmitriev, A. A. Greshnov, and M. A. Semina, Hall effect in Poiseuille flow of two-dimensional electron fluid, *Phys. Rev. B* **106**, 245415 (2022).
  - [16] K. K. Grigoryan, D. S. Zohrabyan, and M. M. Glazov, Anomalous hall effect in ultraclean electronic channels (2023).
  - [17] M. M. Glazov, Valley and spin accumulation in ballistic and hydrodynamic channels, *2D Materials* **9**, 015027 (2022).
  - [18] A. N. Afanasiev, P. S. Alekseev, A. A. Danilenko, A. A. Greshnov, and M. A. Semina, Rotational viscosity in spin resonance of hydrodynamic electrons, *Phys. Rev. B* **106**, L041407 (2022).
  - [19] K. S. Denisov, K. A. Baryshnikov, and P. S. Alekseev, Spin imaging of poiseuille flow of a viscous electronic fluid, *Phys. Rev. B* **106**, L081113 (2022).
  - [20] H. Funaki, R. Toshio, and G. Tatara, Vorticity-induced anomalous Hall effect in an electron fluid, *Phys. Rev. Res.* **3**, 033075 (2021).
  - [21] G. Tatara, Hydrodynamic theory of vorticity-induced spin transport, *Phys. Rev. B* **104**, 184414 (2021).
  - [22] E. H. Hasdeo, J. Ekström, E. G. Idrisov, and T. L. Schmidt, Electron hydrodynamics of two-dimensional anomalous Hall materials, *Phys. Rev. B* **103**, 125106 (2021).
  - [23] V. F. Gantmakher and Y. B. Levinson, *Carrier Scattering in Metals and Semiconductors* (North-Holland Publishing Company, 1987).
  - [24] V. N. Mantsevich and M. M. Glazov, Viscous hydro-

- dynamics of excitons in van der waals heterostructures, *Phys. Rev. B* **110**, 165305 (2024).
- [25] J. E. Avron, Odd viscosity, *Journal of Statistical Physics* **92**, 543 (1998).
- [26] D. Banerjee, A. Souslov, A. G. Abanov, and V. Vitelli, Odd viscosity in chiral active fluids, *Nature Communications* **8**, 1573 (2017).
- [27] S. Ganeshan and A. G. Abanov, Odd viscosity in two-dimensional incompressible fluids, *Phys. Rev. Fluids* **2**, 094101 (2017).
- [28] M. Fruchart, M. Han, C. Scheibner, and V. Vitelli, The odd ideal gas: Hall viscosity and thermal conductivity from non-hermitian kinetic theory (2022).
- [29] M. Fruchart, C. Scheibner, and V. Vitelli, Odd viscosity and odd elasticity, *Annual Review of Condensed Matter Physics* **14**, 471 (2023).
- [30] D. Zverevich, D. B. Gutman, and A. Levchenko, Hydrodynamic coulomb drag in odd electron liquids, *Phys. Rev. B* **112**, 045309 (2025).
- [31] S. Chapman and T. Cowling, *The Mathematical Theory of Non-uniform Gases: An Account of the Kinetic Theory of Viscosity, Thermal Conduction and Diffusion in Gases*, Cambridge Mathematical Library (Cambridge University Press, 1990).
- [32] E. H. Hall, On a new action of the magnet on electric currents, *American Journal of Mathematics* **2**, 287 (1879).
- [33] E. H. Hall, XXXVIII. On the new action of magnetism on a permanent electric current, *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science* **5**, 157 (1881).
- [34] E. N. Adams and E. I. Blount, Energy bands in the presence of an external force field—II: Anomalous velocities, *Journal of Physics and Chemistry of Solids* **10**, 286 (1959).
- [35] L. E. Gurevich and I. N. Yassievich, Theory of ferromagnetic Hall effect, *Sov. Phys. Solid. State* **4**, 2091 (1963).
- [36] V. Abakumov and I. Yassievich, Anomalous Hall effect for polarized electrons in semiconductors, *JETP* **34**, 1375 (1972).
- [37] Nozières, P. and Lewiner, C., A simple theory of the anomalous Hall effect in semiconductors, *J. Phys. France* **34**, 901 (1973).
- [38] N. A. Sinitsyn, Semiclassical theories of the anomalous Hall effect, *Journal of Physics: Condensed Matter* **20**, 023201 (2007).
- [39] V. Dugaev, M. Taillefumier, B. Canals, C. Lacroix, and P. Bruno, Strongly correlated systems, coherence and entanglement (World Scientific) Chap. Anomalous Hall effect, p. 145.
- [40] N. Nagaosa, J. Sinova, S. Onoda, A. H. MacDonald, and N. P. Ong, Anomalous Hall effect, *Rev. Mod. Phys.* **82**, 1539 (2010).
- [41] I. A. Ado, I. A. Dmitriev, P. M. Ostrovsky, and M. Titov, Anomalous Hall effect with massive Dirac fermions, *EPL* **111**, 37004 (2015).
- [42] V. K. Kozin, V. A. Shabashov, A. V. Kavokin, and I. A. Shelykh, Anomalous exciton hall effect, *Phys. Rev. Lett.* **126**, 036801 (2021).
- [43] M. M. Glazov and L. E. Golub, Valley Hall effect caused by the phonon and photon drag, *Phys. Rev. B* **102**, 155302 (2020).
- [44] M. M. Glazov and L. E. Golub, Skew Scattering and Side Jump Drive Exciton Valley Hall Effect in Two-Dimensional Crystals, *Physical Review Letters* **125**, 157403 (2020).
- [45] N. S. Averkiev, L. E. Golub, and M. Willander, Spin relaxation anisotropy in two-dimensional semiconductor systems, *J. Phys.: Condens. Matter* **14**, R271 (2002).
- [46] B. I. Sturman, Collision integral for elastic scattering of electrons and phonons, *Soviet Physics Uspekhi* **27**, 881 (1984).
- [47] M. I. Dyakonov and V. I. Perel', Current induced spin orientation of electrons in semiconductors, *Phys. Lett. A* **35A**, 459 (1971).
- [48] M. B. Lifshits and M. I. Dyakonov, Swapping spin currents: Interchanging spin and flow directions, *Phys. Rev. Lett.* **103**, 186601 (2009).
- [49] M. I. Dyakonov, ed., *Spin physics in semiconductors*, 2nd ed., Springer Series in Solid-State Sciences 157 (Springer International Publishing, 2017).
- [50] D. S. Zohrabyan and M. M. Glazov, Diffusive-hydrodynamic transition in the anomalous hall effect, *Physics of the Solid State* **65**, 2205 (2023).
- [51] L. Landau and E. Lifshitz, *Physical Kinetics* (Butterworth-Heinemann, Oxford, 1981).
- [52] V. A. Zakharov and I. S. Burmistrov, Residual bulk viscosity of a disordered two-dimensional electron gas, *Phys. Rev. B* **103**, 235305 (2021).
- [53] L. D. Landau and E. M. Lifshitz, *Fluid Mechanics* (Pergamon Press, 1987).
- [54] M. Matsuo, Y. Ohnuma, and S. Maekawa, Theory of spin hydrodynamic generation, *Phys. Rev. B* **96**, 020401 (2017).
- [55] R. Takahashi, H. Chudo, M. Matsuo, K. Harii, Y. Ohnuma, S. Maekawa, and E. Saitoh, Giant spin hydrodynamic generation in laminar flow, *Nature Communications* **11**, 3009 (2020).