

---

# The Hurwitz existence problem and the prime-degree conjecture: A computational perspective

---

Yiru Wang<sup>1</sup>, Bingqian Li<sup>2</sup>, Yi Zhou<sup>2</sup>, Zhiqiang Wei<sup>3</sup>, Yu Ye<sup>1</sup>, Yiqian Shi<sup>1</sup>, Bin Xu<sup>1</sup>

<sup>1</sup>School of Mathematical Sciences, University of Science and Technology of China(USTC)

<sup>2</sup>National Engineering Laboratory for Brain-inspired Intelligence Technology and Applications,  
School of Information Science and Technology, USTC

<sup>3</sup>School of Mathematics and Statistics, Henan University

## Abstract

We investigate the Hurwitz existence problem from a computational viewpoint. Leveraging the symmetric-group algorithm by Zheng and building upon implementations originally developed by Baroni, we achieve a complete and non-redundant enumeration of all non-realizable partition triples for positive integers up to 31. These results are further categorized into four types according to their underlying mathematical structure; it is observed that nearly nine-tenths of them can be explained by known theoretical results. As an application, we verify the prime-degree conjecture for all primes less than 32. In light of the exponential memory growth inherent in existing computational approaches—which limits their feasibility at higher degrees—we propose a novel software architecture designed to stabilize memory usage, thereby facilitating further detection of exceptional cases in the Hurwitz existence problem. The complete dataset of non-realizable partition triples, along with our implementation, will be made public on GitHub.

MSC 2010: 57M12 (primary), 20B35, 05A17.

## 1 Introduction

The classical Hurwitz existence problem concerns the realizability of prescribed branch data by branched coverings of the Riemann sphere  $\mathbb{CP}^1$  [1, 2]. Given a collection  $\Lambda$  of  $k \geq 3$  nontrivial partitions of a positive integer  $d$ , one asks whether there exist a compact connected Riemann surface  $S$  together with a branched covering  $f : S \rightarrow \mathbb{CP}^1$  whose branch data agree with  $\Lambda$ . When such a map  $f$  exists, we say that  $\Lambda$  is *realizable*. Any realizable collection must satisfy an algebraic necessary condition, given by the Riemann-Hurwitz formula [3, §7.2.1, p. 101]. We refer to this as the *Riemann-Hurwitz condition*. A collection satisfying this condition but not realizable will be called *exceptional*.

Hurwitz [1] reduced this problem to an equivalent one of constructing  $k$  suitable permutations in the symmetric group  $S_d$  associated with the collection  $\Lambda$ . See [3, §4.2] for a proof of this reduction. Along this line, Mednykh [4] gave a conceptual solution to the Hurwitz existence problem in the sense that he obtained a complicated formula for the number of equivalence classes of branched covers realizing  $\Lambda$ . However, it seems difficult in practice to determine the realizability of a given collection  $\Lambda$  from Mednykh’s formula. Therefore, beyond the Riemann-Hurwitz condition, the problem remains highly intricate and is known to encode deep interactions between complex analysis, algebraic geometry, and the representation theory of symmetric groups. The reader may consult [5–7] and the references therein for up-to-date theoretical results on the Hurwitz existence problem.

A particularly appealing case is the prime-degree conjecture proposed by Edmonds-Kulkarni-Stong [2], which asserts that whenever the degree  $d$  is prime, any collection  $\Lambda$  of  $k$  nontrivial partitions of  $d$  satisfying the Riemann-Hurwitz condition must be realizable. They also reduced the conjecture to the case  $k = 3$ . This conjecture was given strong support by the work of Zheng [8], who reformulated the realizability problem in terms of generating functions involving secondary partitions and characters of  $S_d$ , and confirmed the conjecture for primes less than 20. Building on Zheng’s algorithm, Baroni [7, 9] carried out further verifications, confirming the conjecture for all primes  $d \leq 29$ . However, Baroni’s computational approach suffers from a fundamental obstacle: it exhibits exponential memory growth, which limits the applicability for larger degrees.

In this work we revisit the Hurwitz existence problem from a computational perspective. Building on Zheng’s algorithm, we undertake a new computational architecture of the Hurwitz existence problem that overcomes this barrier. For simplicity, we restrict ourselves to the case where  $\Lambda$  consists of exactly three partitions.

Our contributions in this manuscript are threefold as follows.

First, by extending Baroni’s implementation and conducting computations on high-performance clusters, we obtain a complete and non-redundant enumeration of all non-realizable partition triples for every  $d \leq 31$ , both prime and composite. This improves the known range from  $d \leq 29$  and provides for the first time a full catalog of all exceptional data up to degree 31. As an immediate consequence, our enumeration verifies the prime-degree conjecture for all primes less than 32.

Second, the resulting data allow us to perform a systematic structural classification of the exceptional cases. We find that approximately 90% of all non-realizable triples fall into two families of the four—each corresponding to explicit theoretic constraints. Thus, the overwhelming majority of exceptional triples admit a clear conceptual explanation rather than arising from sporadic computational accidents.

Third, we introduce a new software architecture, which is inspired by Baroni’s work, designed to overcome the memory limitations inherent in previous approaches. Our design employs a two-layer architecture: a Python layer manages workflow and I/O, while computationally intensive kernels are JIT-compiled for performance. This separation, combined with hash-table data structure and batch processing, reduces peak memory consumption dramatically, enabling computations that were previously unattainable. This reduces the peak memory requirement for  $d = 32$  from multi-terabyte scales to under 400 GB in practice. Our design also emphasizes sustainability and resumability by providing detailed logging and checkpointing mechanisms suitable for long-running computations.

The complete dataset of exceptional partition triples for  $d \leq 31$  provides a refined picture of the exceptional landscape, and we hope that both the full dataset and its structural organization will facilitate future theoretical work on the Hurwitz existence problem.

The paper is organized as follows. Section 2 recalls Zheng’s character-theoretic algorithm and establishes notation. Section 3 describes the false-positive removal method. Section 4 classifies these triples into four families. Section 5 describes the new computational architecture. All exceptional data and code will be deposited in the repository.

## 2 Preliminaries

Throughout this section we recall the algorithm for realizability introduced by Zheng [8]. We fix a positive integer  $d$ , and let  $\mathcal{P}(d)$  denote the set of all partitions of  $d$ . For a partition  $\lambda = (\lambda_1, \dots, \lambda_r) \in \mathcal{P}(d)$ , we use the following notation:

- $S^\lambda$  denotes the irreducible representation of the symmetric group  $S_d$  corresponding to  $\lambda$ ;
- $\chi^\lambda$  denotes the irreducible character of  $S_d$  afforded by  $S^\lambda$ ;
- $C_\lambda \subseteq S_d$  denotes the conjugacy class consisting of permutations whose cycle-type is  $\lambda$ .

### 2.1 Secondary partitions

Following Zheng, a secondary partition of  $d$ :  $\omega = [\mu_1, \dots, \mu_n] \vdash d$ , where  $\mu_j \vdash d_j$ ,  $d_j \in \mathbb{Z}_{>0}$ ,  $\forall j$  such that  $d_1 + \dots + d_n = d$ . For such  $\omega$ , let  $n_1, \dots, n_m$  be the multiplicities of the distinct components among the set of  $\{\mu_{i_j}, j = 1, \dots, m\}$ , so that  $n_1 + \dots + n_m = n$ .

## 2.2 The coefficients $r(\omega)$

Define

$$r(\omega) = \frac{(-1)^{n-1}}{n} \frac{n!}{n_1! \dots n_m!} \prod_{j=1}^n \left( \frac{\dim S^{\mu_j}}{d_j!} \right)^2. \quad (1)$$

## 2.3 The polynomials $s(\omega; \mathbf{t})$

Let  $k \geq 1$  be the number of branch points. For  $D = (\mu_1, \dots, \mu_k)$ , denote

$$\mathbf{t}^D = \mathbf{t}_1^{\mu_1} \dots \mathbf{t}_k^{\mu_k}.$$

For  $\mu = [a_1, \dots, a_r]$ ,  $i \in \{1, \dots, k\}$ , denote

$$\mathbf{t}_i^\mu = t_{i,a_1} \dots t_{i,a_r}.$$

Given a secondary partition  $\omega = [\mu_1, \dots, \mu_n]$  with  $\mu_j \vdash d_j$ , define

$$s(\omega; t_{i1}, \dots, t_{id}) = \prod_{j=1}^n \left( \sum_{\nu \vdash d_j} \frac{\chi^{\mu_j}(C_\nu) |C_\nu|}{\dim S^{\mu_j}} \mathbf{t}_i^\nu \right). \quad (2)$$

## 2.4 Realizability polynomial

Define

$$F_{d,k}(\mathbf{t}) = \sum_{\omega \vdash \vdash d} r(\omega) \prod_{i=1}^k s(\omega; t_{i1}, \dots, t_{id}). \quad (3)$$

Zheng proved that  $F_{d,k}$  contains all information needed to determine realizability of branch data over  $k$  branch points.

**Theorem 2.1** (Zheng [8, Theorem 4]) Let  $\Lambda = (\lambda_1, \dots, \lambda_k)$  be a  $k$ -tuple of nontrivial partitions of  $d$ . Then  $\Lambda$  is realizable by a branched covering of compact Riemann surfaces if and only if the coefficient of the monomial

$$\mathbf{t}_1^{\lambda_1} \dots \mathbf{t}_k^{\lambda_k}$$

in  $F_{d,k}(\mathbf{t})$  is nonzero.

Thus the Hurwitz problem reduces to detecting which coefficients in the polynomial  $F_{d,k}$  vanish.

## 3 Eliminating False Positives

The detection of non-realizable branch data in Zheng's symmetric-group framework relies on evaluating the polynomial

$$F_{d,k}(\mathbf{t}) = \sum_{\omega \vdash \vdash d} r(\omega) \prod_{i=1}^k s(\omega; t_{i1}, \dots, t_{id})$$

where  $k = 3$  for three branch data case. That means,  $\forall \omega \vdash \vdash d$ , we should estimate

$$\begin{aligned} & \frac{(n-1)!}{n_1! \dots n_m!} \cdot \left( \prod_{j=1}^n \frac{(\dim S^{\mu_j})^2}{(d_j!)^2} \right) \cdot \prod_{i=1}^3 \prod_{j=1}^n \left( \sum_{\nu \vdash d_j} \frac{\chi^{\mu_j}(C_\nu) |C_\nu|}{\dim S^{\mu_j}} \right) \\ &= \frac{(n-1)!}{n_1! \dots n_m!} \cdot \prod_{j=1}^n \left( \frac{(\dim S^{\mu_j})^2}{(d_j!)^2} \cdot \frac{\left( \sum_{\nu \vdash d_j} \chi^{\mu_j}(C_\nu) |C_\nu| \right)^3}{(\dim S^{\mu_j})^3} \right) \\ &= \frac{\frac{n!}{n_1! \dots n_m!} \cdot \left( \sum_{\nu \vdash d_j} \chi^{\mu_j}(C_\nu) |C_\nu| \right)^3}{n \cdot \prod_{j=1}^n \dim S^{\mu_j} \cdot (d_j!)^2} \end{aligned} \quad (4)$$

$\Lambda$  is exceptional precisely when the coefficient of its corresponding monomial in  $F_{d,3}$  vanishes.

In actual computation, however, to both avoid the explosion of intermediate coefficients and guarantee exact arithmetic without floating-point errors, our implementation evaluates these coefficients modulo a fixed prime  $p$ . Thus a genuine zero coefficient certainly appears as zero mod  $p$ , but the converse need not hold: a nonzero integer coefficient may also vanish modulo  $p$ . Such spurious outputs are false positives, and removing them is essential for producing a complete and non-redundant list of exceptional triples.

To eliminate false positives rigorously, we employ the following procedure, justified by a bound on the numerator of coefficients.

### 3.1 Bounding the coefficients of $F_{d,3}$

For a secondary partition  $\omega = [\mu_1, \dots, \mu_n] \vdash d$ , the polynomial

$$s(\omega; \mathbf{t}) = \prod_{j=1}^n \left( \sum_{\nu \vdash d_j} \frac{\chi^{\mu_j}(C_\nu) |C_\nu|}{\dim S^{\mu_j}} \mathbf{t}_i^\nu \right).$$

is a product of  $n$  polynomials.

First, we shall estimate the numerator part of the above unexpanded form.

Using the first-orthogonality relation

$$\sum_{g \in S_{d_j}} |\chi^{\mu_j}(g)|^2 = d_j!, \quad \sum_{g \in S_{d_j}} 1 = d_j!,$$

and the Cauchy-Schwarz inequality, we have

$$\sum_{g \in S_{d_j}} |\chi^{\mu_j}(g)| \leq \left( \sum_{g \in S_{d_j}} |\chi^{\mu_j}(g)|^2 \right)^{1/2} \left( \sum_{g \in S_{d_j}} 1 \right)^{1/2} = d_j!.$$

Since  $|\chi^{\mu_j}(g)|$  is constant on conjugacy classes, we write  $\chi^{\mu_j}(C_\nu)$  for  $g \in C_\nu$  ( $\nu \vdash d_j$ ) and obtain an upper bound

$$\sum_{\nu \vdash d_j} |\chi^{\mu_j}(C_\nu)| |C_\nu| \leq d_j!.$$

Next, we derive an upper bound for the numerators of the rational coefficients after expressed with a common denominator.

Fix a secondary partition  $\omega = [\mu_1, \dots, \mu_n] \vdash d$ . When  $k = 3$ , the un-estimated part of  $|r(\omega) \prod_{i=1}^3 s(\omega; t^{(i)})|$  can be written as

$$\frac{\frac{n!}{n_1! \cdots n_m!}}{n \cdot \prod_{j=1}^n \dim S^{\mu_j} \cdot (d_j!)^2}. \quad (5)$$

Using  $n_1 + \dots + n_m = n$ , we have that

$$\frac{n!}{n_1! \cdots n_m!} \in \mathbb{Z}_{>0},$$

and

$$\frac{n!}{n_1! \cdots n_m!} \leq n! < d!.$$

Since

$$\dim S^{\mu_j} \leq d_j!,$$

together with

$$d_1 + \dots + d_n = d,$$

it follows that the numerator of (4) can be bounded by  $(d!)^4$ , while the denominator can be bounded by  $n \cdot (d!)^3$ .

Now, we consider the sum over all  $\omega$ . In order to apply the Chinese Remainder Theorem(CRT), we must first find a common denominator.

Because

$$\dim S^{\mu_j} \mid d_j!,$$

we may take  $(d!)^4$  as a common denominator. With this denominator, the numerator of each summand corresponding to a  $\omega$  is bounded by  $(d!)^8$ . Therefore, if there are  $p_2(d)$  secondary partitions  $\omega \vdash d$ , summing over the number of  $\omega$  we obtain

**Lemma 3.1 (Numerator bound)** *After using a common denominator as  $(d!)^4$ , every coefficient of the polynomial  $F_{d,3}$  is a rational number whose numerator (in absolute value) is at most*

$$p_2(d) \cdot (d!)^8.$$

### 3.2 Testing modulo sufficiently many primes

Let

$$M_d := p_2(d) \cdot (d!)^8.$$

If we choose primes  $p_1, \dots, p_N$  whose product exceeds  $M_d$ , then the following holds:

**Proposition 3.2 (Chinese remainder theorem for false positives)** *Suppose a set of three branch data  $\Lambda$  gives zero coefficient in  $F_{d,3}$  modulo each of the primes  $p_1, \dots, p_N$  ( $p_i \neq p_j, \forall i \neq j$ ). If*

$$p_1 p_2 \cdots p_N > M_d,$$

*then the true integer coefficient of  $\Lambda$  in  $F_{d,3}$  is zero. In particular,  $\Lambda$  is genuinely non-realizable.*

Thus by running the exceptional-checking program under a list of primes whose product exceeds  $M_d$ , any triple that survives all tests is confirmed to be a true exceptional triple.

### 3.3 Summary on False Positives for $d \leq 30$

We initially compute all the 'exceptional data' under the prime number 1,000,000,007. Then, using the aforementioned verification method, we identify all the false positives for  $d \leq 30$ , which are not genuine exceptional data, as shown in the table below.

$d$	$\lambda_1$	$\lambda_2$	$\lambda_3$
25	[ 8 8 2 2 2 1 1 1 ]	[ 7 4 4 2 2 2 2 1 1 ]	[ 11 5 3 2 2 2 ]
26	[ 10 5 3 2 1 1 1 1 1 1 ]	[ 7 6 2 2 2 2 2 1 1 1 ]	[ 10 9 4 3 ]
28	[ 4 3 3 3 3 2 2 2 2 1 ]	[ 15 5 4 2 1 1 ]	[ 11 7 4 3 3 ]
	[ 10 4 4 2 2 1 1 1 1 1 1 ]	[ 7 5 3 3 3 3 3 1 ]	[ 11 10 3 1 1 1 1 ]
	[ 7 6 3 2 2 2 1 1 1 1 1 1 ]	[ 4 3 3 3 3 2 2 2 2 2 2 ]	[ 19 4 3 1 1 ]
	[ 7 5 3 2 2 2 1 1 1 1 1 1 1 ]	[ 6 6 4 3 3 2 2 1 1 ]	[ 18 4 4 2 ]
30	[ 12 8 4 1 1 1 1 1 1 1 ]	[ 6 6 5 5 5 3 ]	[ 19 7 4 ]
	[ 11 6 6 1 1 1 1 1 1 1 ]	[ 11 5 4 4 2 2 2 ]	[ 7 6 4 4 4 3 2 ]
	[ 8 4 4 4 2 2 2 1 1 1 1 ]	[ 14 5 3 2 2 2 1 1 ]	[ 14 6 6 1 1 1 1 ]
	[ 4 4 4 4 3 3 2 2 1 1 1 1 ]	[ 5 4 4 4 3 3 2 2 2 1 ]	[ 14 10 1 1 1 1 1 1 ]
	[ 11 4 3 2 2 2 1 1 1 1 1 1 ]	[ 9 7 6 3 3 1 1 ]	[ 14 5 4 2 2 2 1 ]
	[ 6 6 3 3 2 2 1 1 1 1 1 1 1 ]	[ 16 5 4 2 1 1 1 ]	[ 14 8 4 3 1 ]

## 4 Classification of Exceptional Triples

Having obtained the complete and cleaned catalogue of exceptional partition triples for degrees  $d \leq 31$ , we organise them into four disjoint types.

**Type 0.**

The sum of the lengths of the three partitions is less than  $d + 2$ . Note that if a collection of three non-trivial partitions of  $d$  is realizable and the sum of the lengths of these three partitions is less than  $d + 2$ , then this collection is realizable by a branched cover from a compact Riemann surface with positive genus to  $\mathbb{CP}^1$ .

**Type I.**

The sum of the lengths equals  $d + 2$ , and exists one partition which contains a unique part greater than 1. Both G. Boccara [10] and Song-Xu [5] proved more general theorems, predicting that all the data of Type I are exceptional.

**Type II.**

If the triple is not of Type 0 or I, and there exists a common divisor  $c > 1$  of all the parts of two partitions, while the third partition cannot be combined into  $c$  partitions  $\mu_1, \dots, \mu_c$ , such that

$$\mu_i \vdash \frac{d}{c}, \forall i = 1, \dots, c,$$

then the triple is assigned to *Type II*. The main theorem in Wei-Wu-Xu [11] showed such data are exceptional.

**Type III.**

All exceptional triples that do not satisfy the previous conditions are collected in Type III.

We tag every exceptional triple with one of the four labels Types 0, I, II, III. Empirically, Types I and II account for roughly 90% of all exceptions. In fact, the conditions defining Types I and II are known to be sufficient to guarantee exceptionality, providing theoretical justification for these two classes. The authors have no theoretical understanding about the data of Type 0 or III.

## 5 Computational Architecture

Two aspects of formula (3) have a critical impact on performance:

1. The number of secondary partitions grows extremely rapidly with  $d$ , reaching several million already at  $d = 32$ .
2. Expanding each  $s(\omega; \mathbf{t})$  generates a large number of monomials. Each coefficient requires sums over the partitions, making the computation time-consuming, and the sheer number of coefficients makes storage memory-intensive.

Our computational architecture mitigates these bottlenecks through batched processing, a hash-table-based data structure, and JIT-compiled parallel kernels for compute-intensive loops.

### 5.1 Compatibility requirements

Let  $\mathcal{P}(d)$  denote the set of all partitions of  $d$ , ordered in reverse lexicographic order. For each  $\lambda \in \mathcal{P}(d)$ , we define its length

$$\ell(\lambda) = \text{the number of parts of } \lambda.$$

A triple  $(\lambda_1, \lambda_2, \lambda_3)$  with  $\lambda_i \vdash d$  that is realized as the branch data of a branched covering must satisfy the numerical constraints imposed by the Riemann-Hurwitz formula. For a branch point of cycle type  $\lambda$ , the ramification contribution equals  $d - \ell(\lambda)$ . Substituting these contributions into the Riemann-Hurwitz formula yields

$$2g(S) - 2 = -2d + \sum_{i=1}^3 (d - \ell(\lambda_i)). \quad (6)$$

Imposing the necessary condition  $g(S) \geq 0$  gives

$$\ell(\lambda_1) + \ell(\lambda_2) + \ell(\lambda_3) \leq d + 2. \quad (7)$$

Taking (6) modulo 2, we also obtain the parity condition

$$\ell(\lambda_1) + \ell(\lambda_2) + \ell(\lambda_3) \equiv d \pmod{2}. \quad (8)$$

(7) and (8) are the constraints compatible with the Riemann-Hurwitz formula.

The subsections below follow the execution flow of the algorithm. Each subsection corresponds to a major stage in the computational pipeline.

## 5.2 Precomputation

Before the coefficient computation begins, we construct several combinatorial and representation-theoretic tables associated with the symmetric groups  $S_1, \dots, S_d$ .

- **Partition table.**  
For each  $0 \leq n \leq d$ , we enumerate all partitions of  $n$  in reverse lexicographic order, together with their lengths, offsets, and cumulative counts. This provides constant-time access to every partition and enables binary search within each layer for  $n$ .
- **Conjugacy-class sizes.**  
For each partition  $\nu \vdash n$  we compute the size of the conjugacy class  $C_\nu$ .
- **Partition multiplication table.**  
We construct a table encoding the product of monomials. Concretely, given two monomials corresponding to partitions  $\lambda$  and  $\mu$ , their product corresponds to the sorted multiset of  $\lambda \cup \mu$ . The table stores the offset of this resulting partition, enabling constant-time lookup during polynomial multiplication.
- **Irreducible character table.**  
The values  $\chi^\lambda(C_\nu)$  for all  $\lambda, \nu \vdash n$ ,  $0 \leq n \leq d$ , are computed recursively using the Murnaghan-Nakayama rule via rim-hook removal, implemented with all arithmetic performed modulo a fixed prime  $p$ .
- **Factorials and inverse factorials modulo a prime  $p$ .**  
The values  $n! \pmod{p}$  and  $(n!)^{-1} \pmod{p}$  for  $0 \leq n \leq d$  are computed.

For  $d = 32$ , the full precomputation of these tables takes only 7 seconds in our experiment.

## 5.3 Batch Computation for $r(\omega)$ and $s(\omega)$

To compute  $r(\omega)$  and  $s(\omega; \mathbf{t})$  efficiently, we process the primary partitions in batches, where each batch contains  $B \in \mathbb{N}$  partitions (typically  $100 \leq B \leq 200$ ).

For each primary partition  $\lambda$ , all secondary partitions  $\omega$  are enumerated recursively by subdividing each primary partition; in practice, to control stack depth, we use stack to replace recursion. For every  $\omega$ , we compute  $r(\omega)$  and a compressed form of the  $s(\omega)$ , without storing  $\omega$ . To avoid memory blow-up, polynomials  $s(\omega)$  are never accumulated globally; instead, each batch is streamed directly to a disk-backed storage.

A batch size  $B = 150$  for  $d = 32$  occupies approximately 40GB on disk, allowing computations for  $d \geq 30$  without terabyte-scale memory.

## 5.4 Accumulation for Each Fixed $\lambda_i$

To test the realizability of triples with first component  $\lambda_i$ , we stream each batch from disk and update partial sums according to (3).

Instead of storing all  $(j, k)$  coefficients explicitly, we adopt a cache-friendly segmented hash table, with each segment roughly corresponding to a CPU cache line. The hash table uses the Cantor pairing

function to map each key  $(j, k)$  to a 64-bit integer. Collisions are resolved via quadratic probing, which searches for an available slot, backed by a small overflow buffer for robustness.

Each  $\lambda_i$  owns an independent accumulator, enabling embarrassingly parallel work distribution and high memory efficiency.

### 5.5 Zero-Detection and Exceptional Triples

Within the hash-table structure, a triple is classified as exceptional if:

1. it satisfies the necessary conditions mentioned in 5.1 , and
2. its key is either absent from the hash table or has value zero.

## Acknowledgements

B.X. would like to express his sincere gratitude to Yufei Bai, Xiaoyang Chen, Tielin Dai, Dun Liang, Sicheng Lu, Xin Nie, Zijin Peng, Qingyu Xu, Junyi Yang, Wenyan Yang, Xiaoya Zhai and Youliang Zhong for many valuable and stimulating conversations during the course of this project. The authors are also grateful to Chao Wu at the Supercomputing Center of USTC for kindly answering their questions on the supercomputing system there.

Both B.L. and Y.Z. are supported in part by Anhui Province Science and Technology Tackle Plan Project (No. 202423k09020008). B.X. is supported in part by the Project of Stable Support for Youth Team in Basic Research Field, CAS (Grant No. YSBR-001) and NSFC (Grant No. 12271495). The numerical calculations in this manuscript have been carried out on the supercomputing system in the Supercomputing Center of USTC.



## References

- [1] Adolf Hurwitz. Ueber Riemann'sche Flächen mit gegebenen Verzweigungspunkten. *Math. Ann.*, 39 (1):1–60, 1891. ISSN 0025-5831,1432-1807. doi: 10.1007/BF01199469. URL <https://doi.org/10.1007/BF01199469>. 1
- [2] Allan L. Edmonds, Ravi S. Kulkarni, and Robert E. Stong. Realizability of branched coverings of surfaces. *Trans. Amer. Math. Soc.*, 282(2):773–790, 1984. ISSN 0002-9947,1088-6850. doi: 10.2307/1999265. URL <https://doi.org/10.2307/1999265>. 1, 2
- [3] Simon Donaldson. *Riemann surfaces*, volume 22 of *Oxford Graduate Texts in Mathematics*. Oxford University Press, Oxford, 2011. ISBN 978-0-19-960674-0. doi: 10.1093/acprof:oso/9780198526391.001.0001. URL <https://doi.org/10.1093/acprof:oso/9780198526391.001.0001>. 1
- [4] Aleksandr D. Mednykh. Nonequivalent coverings of riemann surfaces with a prescribed ramification type. *Siberian Mathematical Journal*, 25(4):606–625, 1984. doi: 10.1007/BF00968900. URL <https://doi.org/10.1007/BF00968900>. 1
- [5] Jijian Song and Bin Xu. On rational functions with more than three branch points. *Algebra Colloq.*, 27(2):231–246, 2020. ISSN 1005-3867,0219-1733. doi: 10.1142/S100538672000019X. URL <https://doi.org/10.1142/S100538672000019X>. 1, 6
- [6] Carlo Petronio. The Hurwitz existence problem for surface branched covers. *Winter Braids Lect. Notes*, 7: Exp. No. 2, 43, 2020. ISSN 2426-0312.
- [7] Filippo G. Baroni and Carlo Petronio. Solution of the Hurwitz problem with a length-2 partition. *Illinois J. Math.*, 68(3):479–511, 2024. ISSN 0019-2082,1945-6581. doi: 10.1215/00192082-11417175. URL <https://doi.org/10.1215/00192082-11417175>. 1, 2
- [8] Hao Zheng. Realizability of branched coverings of  $S^2$ . *Topology Appl.*, 153(12):2124–2134, 2006. ISSN 0166-8641,1879-3207. doi: 10.1016/j.topol.2005.08.007. URL <https://doi.org/10.1016/j.topol.2005.08.007>. 2, 3
- [9] Filippo Baroni. Source code for the implementation of zheng's algorithm. <https://github.com/filippobaroni/hurwitz-existence-source-code>. Accessed: 2024-02-17. 2
- [10] Guy Boccara. Cycles comme produit de deux permutations de classes données. *Discrete Math.*, 38 (2-3):129–142, 1982. ISSN 0012-365X,1872-681X. doi: 10.1016/0012-365X(82)90282-5. URL [https://doi.org/10.1016/0012-365X\(82\)90282-5](https://doi.org/10.1016/0012-365X(82)90282-5). 6
- [11] Zhiqiang Wei, Yingyi Wu, and Bin Xu. A note on rational maps with three branching points on the riemann sphere, 2024. URL <https://arxiv.org/abs/2401.06956>. 6

YIYU WANG, YE YU, YIQIAN SHI, BIN XU  
SCHOOL OF MATHEMATICAL SCIENCES, USTC  
HEFEI 230026 CHINA  
wyr221072@mail.ustc.edu.cn  
yeyu@ustc.edu.cn  
yqshi@ustc.edu.cn  
bxu@ustc.edu.cn

BINGQIAN LI, YI ZHOU  
NATIONAL ENGINEERING LABORATORY FOR BRAIN-INSPIRED INTELLIGENCE  
TECHNOLOGY AND APPLICATIONS  
SCHOOL OF INFORMATION SCIENCE AND TECHNOLOGY, USTC  
HEFEI 230026 CHINA  
bqli315@gmail.com  
yi\_zhou@ustc.edu.cn

ZHIQIANG WEI  
SCHOOL OF MATHEMATICS AND STATISTICS

HENAN UNIVERSITY  
KAIFENG 475004 CHINA  
weizhiqiang15@mails.ucas.edu.cn