

ON WIENER-HOPF OPERATORS OVER LINEARLY ORDERED DISCRETE ABELIAN GROUPS

A. R. Mirotin

amirotin@yandex.ru

AMS 2010 Mathematics Subject Classification. Primary 43A15, 47B35.

Keywords: Toeplitz operator, Wiener-Hopf operator, Fredholm operator, Fredholm index, spectrum, essential spectrum, ordered Abelian group, compact Abelian group.

ABSTRACT

Let X denotes a discrete linearly ordered Abelian group, and let X_+ be the positive cone in X . In this note we compute the Fredholm index and study spectral properties of Wiener-Hopf operators $W_k g = 1_{X_+}(k * g)$, $k \in l_2(X_+)$ in the space $l_2(X_+)$ in terms of their symbols \check{k} where \check{k} stands for the inverse Fourier transform of k .

1. INTRODUCTION AND PRELIMINARIES

Everywhere below, X denotes a discrete linearly ordered Abelian group, X_+ is the positive cone in X . This means that X has a fixed subsemigroup X_+ containing the identity character 1, where $X_+ \cap X_+^{-1} = \{1\}$ and $X = X_+ \cup X_+^{-1}$. Then the semigroup X_+ induces a linear ordering in X compatible with the group structure, according to the rule $\xi \leq \chi := \chi \xi^{-1} \in X_+$. As is well known, a (discrete) Abelian group X can be linearly ordered if and only if it is torsion-free (see, for example, [7]), which, in turn, is equivalent to its character group G being compact and connected [6]. However, the linear ordering in X is, generally speaking, not unique. In what follows, we assume that this order is chosen and fixed. In applications, the role of X is often played by subgroups of the additive group \mathbb{R}^n , endowed with the discrete topology. In this situation, G is the Bohr compactification of X (see, for example, [2] and the references therein).

By $\hat{\varphi}$ (or $\mathcal{F}\varphi$) we denote the Fourier transform of a function φ from $L^1(G)$ or $L^2(G)$, i. e. (with the appropriate interpretation of the integral)

$$\hat{\varphi}(\xi) = \int_G \varphi(x) \overline{\xi(x)} dx, \quad \xi \in X,$$

where dx is the normalized Haar measure of G .

By \check{k} we denote the inverse Fourier transform $\mathcal{F}^{-1}k$ of a function k on X .

Consider the space

$$l_2(X_+) = \{f : X_+ \rightarrow \mathbb{C} : \sum_{\chi \in X_+} |f(\chi)|^2 < \infty\}$$

(here each function f has at most countable support) with the scalar product

$$\langle f, g \rangle = \sum_{\chi \in X_+} f(\chi) \overline{g(\chi)}.$$

The spaces $l_p(X_+)$ are defined similarly. Obviously, the system $\{1_{\{\chi\}}\}_{\chi \in X_+}$ (we denote by 1_A the indicator of the set A) of indicators of one-point subsets is an orthonormal basis of the space $l_2(X_+)$.

In the following we assume that $l_p(X_+)$ is embedded in $l_p(X)$ in a natural way. The notion of a Wiener-Hopf operator over X is due to Adukov [1].

Definition 1.1. Let $k \in l_2(X)$. A Wiener-Hopf operator W_k on $l_2(X_+)$ is defined by the equality

$$W_k g = 1_{X_+}(k * g)$$

where $*$ stands for the convolution of functions on X (as is known, $l_2(X) * l_2(X) \subset l_1(X) \subset l_2(X)$).

Thus,

$$W_k g(\chi) = \sum_{\xi \in X_+} k(\chi \xi^{-1}) g(\xi) \quad (\chi \in X_+).$$

Definition 1.2. We call the inverse Fourier transform of a function k , i. e., the function

$$\check{k}(x) = \sum_{\xi \in X} k(\xi) \xi(x),$$

the symbol of the operator W_k if the right-hand side makes sense.

The goal of this note is to study the Fredholm and spectral properties of Wiener-Hopf operators in the space $l_2(X_+)$ in terms of their symbol, if $k \in l_2(X_+)$, $\check{k} \in C(G)$. In the case of $k \in l_1(X_+)$, the Wiener-Hopf operators in the spaces $l_p(X_+)$, $p \geq 1$ and several others were considered by Adukov in the paper [1], where the issues of invertibility and Fredholm property of these operators were discussed in detail, but a formula for the index was not obtained and the spectrum of such operators was not studied.

Remark 1. Note that our conditions $k \in l_2(X_+)$, $\check{k} \in C(G)$ is less restrictive than $k \in l_1(X_+)$.

Main results of this note, namely, Theorems 2.4 and 3.12 presents the discrete analog of the Gohberg-Krein formula for the Fredholm index and, respectively, the description of the spectrum and essential spectra of Wiener-Hopf operators $W_k g = 1_{X_+}(k * g)$ with continuous symbols that are inverse Fourier transforms of functions $k \in l_2(X)$ in terms of the symbol.

2. FREDHOLM PROPERTIES OF WIENER-HOPF OPERATORS

Let us recall the concepts and facts of the theory of Fredholm operators needed for what follows. For a bounded operator A in a Banach space Y , the notation $A \in \Phi_+(Y)$ means that its image $\text{Im}A$ is closed and the kernel $\text{Ker}A$ is finite-dimensional, while the notation $A \in \Phi_-(Y)$ means that the quotient space $\text{Coker}A := Y/\text{Im}A$ is finite-dimensional; operators in $\Phi_-(Y) \cup \Phi_+(Y)$ are called *semi-Fredholm*, and operators in $\Phi(Y) := \Phi_-(Y) \cap \Phi_+(Y)$ are called *Fredholm* in Y .

The index of the operator $A \in \Phi(Y)$ is the number

$$\text{Ind}A := \dim \text{Ker}A - \dim \text{Coker}A.$$

We will need the following concepts introduced in [3].

Definition 2.3. *The rotation index (wading number) of a character $\chi \in X$ is defined as follows (the sign $\#$ denotes the number of elements of a finite set):*

- 1) $\text{ind}\chi = \#(X_+ \setminus \chi X_+)$, if $\chi \in X_+$ and the set $X_+ \setminus \chi X_+$ is finite;
- 2) $\text{ind}\chi = \text{ind}\chi_1 - \text{ind}\chi_2$, if $\chi = \chi_1 \chi_2^{-1}$, where $\chi_j \in X_+$, and both sets $X_+ \setminus \chi_j X_+$ are finite ($j = 1, 2$).

In other cases, the character is assumed to have an infinite index.

The set of characters with a finite index is denoted by X^i .

Recall that by the Bohr-van Kampen theorem each function φ from the group $C(G)^{-1}$ of invertible elements of the algebra $C(G)$ has the representation

$$\varphi = \chi e^g$$

where $\chi \in X$ and $g \in C(G)$. Moreover, the character χ in this representation is unique.

We will set

$$\Phi(G) := X^i e^{C(G)}, \text{ind}\varphi := \text{ind}\chi, \text{ if } \varphi \in \Phi(G), \varphi = \chi e^g.$$

The index formula for the Wiener-Hopf operator looks as follows.

Theorem 2.4. *Let $k \in l_2(X_+)$, $\check{k} \in C(G)$. The Wiener-Hopf operator W_k on $l_2(X_+)$ is Fredholm if and only if $\check{k} \in \Phi(G)$. In this case,*

$$\text{Ind}W_k = -\text{ind}\check{k}.$$

For the proof, it will be convenient to generalize the definition of a Toeplitz operator over a group G (see, e.g., [3]) to the case of a symbol $\varphi \in L^2(G)$.

Definition 2.5. Let $1 \leq p \leq \infty$. The Hardy space $H^p(G)$ over a group G is defined as follows:

$$H^p(G) = \{f \in L^p(G) : \widehat{f}(\chi) = 0 \ \forall \chi \notin X_+\}.$$

We endow $H^p(G)$ with L^p norm.

Definition 2.6. Let $\varphi \in L^2(G)$. We define the Toeplitz operator T_φ in $H^2(G)$ initially on the set of analytic polynomials $\text{Pol}_+(G)$ (the linear span of X_+) by the equality

$$T_\varphi q = P_+(\varphi q) \ (q \in \text{Pol}_+(G)),$$

where $P_+ : L^2(G) \rightarrow H^2(G)$ is the orthogonal projection.

Theorem 2.4 follows from [3, Theorem 4] and the following proposition.

Proposition 2.7. The operator T_φ in $H^2(G)$ with symbol $\varphi \in L^2(G)$ is unitary equivalent to the operator W_k in $l_2(X)$, where $k = \widehat{\varphi}$, and vice versa, for any $k \in l_2(X)$, the operator W_k is unitary equivalent to the operator T_φ with symbol $\varphi = \check{k}$.

Proof. The assertion of the proposition follows from the fact that for any analytic polynomial $q \in \text{Pol}_+(G)$ and any function $\varphi \in L^2(G)$, the equality

$$T_\varphi q = \mathcal{F}^{-1} W_{\widehat{\varphi}} \mathcal{F} q$$

is valid. For proof, note that for any function $g \in l_2(X)$, the equality

$$\varphi \mathcal{F}^{-1} g = \mathcal{F}^{-1}(\widehat{\varphi} * g)$$

holds. Furthermore, by Plancherel's theorem

$$\mathcal{F} P_+ \mathcal{F}^{-1} g = 1_{X_+} g \ (g \in l_2(X)).$$

Therefore, for $g \in l_2(X)$ we have

$$\mathcal{F} T_\varphi \mathcal{F}^{-1} g = \mathcal{F} P_+(\varphi \mathcal{F}^{-1} g) = \mathcal{F} P_+ \mathcal{F}^{-1}(\widehat{\varphi} * g) = W_{\widehat{\varphi}} g.$$

To complete the proof, it remains to set $q = \mathcal{F}^{-1} g$ in the last equality and note that q runs over $\text{Pol}_+(G)$ when g runs over the space $l_{00}(X_+)$ of finitely supported functions on X_+ . \square

Proposition 2.7 allows us to extend other properties of Toeplitz operators over groups to the case of Wiener-Hopf operators.

Corollary 2.8. The operator W_k is bounded if $k \in l_2(X)$, $\check{k} \in L^\infty(G)$. Moreover, $\|W_k\| = \|\check{k}\|_\infty$.

Proof. This follows immediately from the fact that the operator T_φ is bounded if $\varphi \in L^\infty(G)$, and also from the equality $\|T_\varphi\| = \|\varphi\|_\infty$. \square

Corollary 2.9. *For any function $\varphi \in L^2(G)$ satisfying the condition $\widehat{\varphi} \in l_1(X)$, the operator T_φ is bounded.*

Proof. Indeed, under the conditions of the corollary, the operator $W_{\widehat{\varphi}}$ is bounded by the inequality

$$\|\widehat{\varphi} * f\|_2 \leq \|\widehat{\varphi}\|_1 \|f\|_2.$$

\square

Corollary 2.10. *Let $\check{k} \in C(G)$. If the operator W_k is semi-Fredholm, then \check{k} is invertible in $L^\infty(G)$.*

This follows directly from Proposition 2.7 and [3, Theorem 3]. \square

3. SPECTRAL PROPERTIES OF WIENER-HOPF OPERATORS

Proposition 2.7 allows also to extend the results of Section 4 in [3] to Wiener-Hopf operators. First of all, we state the following corollary of this Proposition.

Corollary 3.11. *Let $\check{k} \in L^\infty(G)$ and $R(\check{k})$ denotes the essential range of \check{k} . Then*

$$R(\check{k}) \subseteq \sigma(W_k) \subseteq \overline{\text{conv}}(R(\check{k}))$$

where $\overline{\text{conv}}$ stands for the closed convex hole of a set in the complex plane.

Indeed, this follows from a corresponding result of Murphy for Toeplitz operators [5] and Proposition 2.7.

Hereinafter, by *holes* of a compact connected set $V \subset \mathbb{C}$ we mean the bounded components of its complement $\mathbb{C} \setminus V$. The following theorem describe the spectra and essential spectra of Wiener-Hopf operators.

Theorem 3.12. *Let $\check{k} \in C(G)$. Then*

1) *the essential Fredholm spectrum $\sigma_e(W_k)$ is obtained from the set $\check{k}(G)$ by adding those of its holes Λ (if they exist) for which $\check{k} - \lambda \notin \Phi(G)$ for all (for one) $\lambda \in \Lambda$;*

2) *the spectrum $\sigma(W_k)$ is obtained from the set $\sigma_e(W_k)$ by adding those holes Λ of the set $\check{k}(G)$ (if they exist) for such that $\check{k} - \lambda \in \Phi(G) \setminus \exp(C(G))$ for all (for one) $\lambda \in \Lambda$;*

3) *the essential Weyl spectrum $\sigma_w(W_k) = \sigma(W_k)$;*

4) *the spectra $\sigma_e(W_k)$ and $\sigma(W_k)$ are connected.*

Proof. Theorem 3.12 follows from Proposition 2.7, Theorem 2.4 and the corresponding results for Toeplitz operators on $H^p(G)$ [3, Theorem 5]. \square

Theorem 3.13. *Let $\check{k} \in H^\infty(G)$.*

(i) *Operator W_k is invertible in $l_2(X)$ if and only if \check{k} is invertible in $H^\infty(G)$.*

(ii) *The spectrum $\sigma(W_k)$ equals to the spectrum $\sigma_{H^\infty}(\check{k})$ of \check{k} in the algebra $H^\infty(G)$.*

Proof. (i) Operator W_k is invertible in $l_2(X_+)$ if and only if $T_{\check{k}}$ is invertible in $H^\infty(G)$ by Proposition 2.7.

If \check{k} is invertible in $H^\infty(G)$ then $\check{k}\psi = 1$ for some $\psi \in H^\infty(G)$. But

$$T_{\check{k}}T_\psi = T_{\check{k}\psi} = T_\psi T_{\check{k}}$$

by [3, Proposition 1] and $T_{\check{k}\psi} = I$ the unit operator in $H^2(G)$. Thus, W_k is invertible in $l_2(X_+)$.

Conversely, if W_k is invertible in $l_2(X_+)$, i. e. $T_{\check{k}}$ is invertible in $H^2(G)$, then $\check{k}H^2(G) = T_{\check{k}}H^2(G) = H^2(G)$, since $T_{\check{k}}f = \check{k}f$ for all $f \in H^2(G)$. It follows that $\check{k}\psi = 1$ for some $\psi \in H^\infty(G)$. Then $H^2(G) = \psi\check{k}H^2(G) = \psi H^2(G)$. Then $\psi \in H^\infty(G)$ by [3, Lemma 2], and hence \check{k} is invertible in $H^\infty(G)$.

(ii) This follows from (i) and the equality

$$\begin{aligned} (W_k - \lambda I)g &= 1_{X_+}(k * g) - \lambda g \\ &= 1_{X_+}((k - \lambda) * g) = W_{k-\lambda}g \end{aligned}$$

which is valid for all $g \in l_2(X_+)$. \square

Lemma 3.14. *Let $\check{k} \in L^\infty(G)$, $|\check{k}| \equiv 1$, and W_k is invertible in $l_2(X_+)$. If $\psi \in H^\infty(G)$ satisfies $\|\check{k} - \bar{\psi}\|_{L^\infty} < 1$ then ψ is invertible in $H^\infty(G)$.*

Proof. Theorem [3, Theorem 1] yields

$$\begin{aligned} \|I - T_{\check{k}\psi}\|_{H^2} &= \|T_{1-\check{k}\psi}\|_{H^2} \\ &\leq \|1 - \check{k}\psi\|_{L^\infty} = \|\bar{\check{k}} - \psi\|_{L^\infty} = \|\check{k} - \bar{\psi}\|_{L^\infty}. \end{aligned} \tag{3.1}$$

Since $\|\check{k} - \bar{\psi}\|_{L^\infty} < 1$, it follows that the operator $T_{\check{k}\psi} = T_{\check{k}}T_\psi$ is invertible. Since $T_{\check{k}}$ is also invertible, operator T_ψ is invertible, as well. Now Theorem 3.13 implies that ψ is invertible in $H^\infty(G)$. \square

Theorem 3.15. (cf. [1, Theorem 3.1]) *Let $\check{k} \in L^\infty(G)$, $|\check{k}| \equiv 1$.*

(i) *Operator W_k is left (right) invertible in $l_2(X_+)$ if and only if the inequality $\text{dist}_{L^\infty}(\check{k}, H^\infty(G)) < 1$ (respectively, $\text{dist}_{L^\infty}(\bar{\check{k}}, H^\infty(G)) < 1$) holds.*

(ii) *Operator W_k is invertible in $l_2(X_+)$ if and only if $\|\check{k} - \bar{\psi}\|_{L^\infty} < 1$ for some invertible ψ in $H^\infty(G)$.*

Proof. (i) Consider the Hankel operator $H_{\check{k}} : H^2(G) \rightarrow L^2(G) \ominus H^2(G)$ with symbol \check{k} (see [4]). Then $H_{\check{k}}f + T_{\check{k}}f = \check{k}f$, and therefore $\|H_{\check{k}}f\|_{L^2}^2 + \|T_{\check{k}}f\|_{L^2}^2 = \|f\|_{L^2}^2$. If W_k is left invertible then $T_{\check{k}}$ is left invertible too by Proposition 2.7. Then $\|T_{\check{k}}f\|_{L^2}^2 \geq C\|f\|_{L^2}^2$ for some constant $C > 0$. So, we conclude that $\|H_{\check{k}}f\|_{L^2} < 1$. It remains to note that $\|H_{\check{k}}f\|_{L^2} = \text{dist}_{L^\infty}(\check{k}, H^\infty(G))$ by [4, Theorem 1.5].

Now the statement for the right inverse follows from Proposition 2.7, and the following equality for the adjoint operator $T_{\check{k}}^* = T_{\check{k}}^-$.

(ii) If W_k is invertible in $l_2(X_+)$ then (i) yields that $\|\check{k} - \psi\|_{L^\infty} < 1$ for some $\psi \in H^\infty(G)$. Then ψ is invertible in $H^\infty(G)$ by Lemma 3.14.

Now let $\|\check{k} - \psi\|_{L^\infty} < 1$ for some invertible ψ in $H^\infty(G)$. Note that the inequality similar to (3.1) is valid if we replace \check{k} by \check{k} . Thus $\|I - T_{\check{k}\psi}^- \leq \|\check{k} - \psi\|_{L^\infty} < 1$. It follows that the operator $T_{\check{k}\psi}^* = T_{\check{k}}^*T_\psi$ is invertible. The operator T_ψ is invertible, too. Indeed, since $\psi\phi = 1$ for some $\phi \in H^\infty(G)$, one has $T_\phi T_\psi = T_\psi T_\phi = T_1 = I$. We conclude that the operator $T_{\check{k}}^*$ is invertible, and therefore $T_{\check{k}}$ and W_k are invertible, as well. \square

REFERENCES

- [1] Adukov, V. Wiener-Hopf operators on a subsemigroup of a discrete torsion free abelian group // Int. Eq. Oper. Th. – 1993. – 16. – P. 305 – 332.
- [2] T. Ehrhardt, C. van der Mee, L. Rodman, I. Spitkovski. Factorization in Weighted Wiener Matrix Algebras on Linearly Ordered Abelian Groups // Int. Eq. Oper. Th. – 2007 – 58, N 1. – P. 65 – 86.
- [3] A. R. Mirotn, Fredholm and spectral properties of Toeplitz operators on H^p spaces over ordered groups, Sb. Math.– 2011.– Vol. 202, N 5. – P. 721–737.
- [4] A. R. Mirotn, Compact Hankel operators on compact Abelian groups, St. Petersburg Math. J., Vol. 33 (2022), No. 3, P. 569–584, <https://doi.org/10.1090/spmj/1715>
- [5] Murphy, G.J. Spectral and index theory for Toeplitz operators, Proc. Roy. Irish Acad. Sect. A. – 1991. – Vol. 91, N 1. – P. 1–6.
- [6] Pontrjagin, L. Topological groups, Princeton, Princeton University Press, 1946.
- [7] Rudin, W. Fourier analysis on groups – New York and London: Interscience Publishers, 1962. – 285 p.