

THE 2-LOCAL HOMOTOPY TYPES OF G_2 -GAUGE GROUPS

MASAKI KAMEKO

ABSTRACT. We determine the 2-local homotopy types of G_2 -gauge groups over S^4 .

1. INTRODUCTION

Let G be a compact Lie group, and P a principal G -bundle over the 4-dimensional sphere S^4 . The G -gauge group over the base space S^4 is the topological group of G -bundle automorphisms of P . Suppose that G is simply-connected and simple. Its classifying space BG is 3-connected and $\pi_4(BG) \cong \mathbb{Z}$. Let us denote the homotopy class of a map f by the same symbol f . Then, the principal G -bundle P is classified by a map $k: S^4 \rightarrow BG$ in $\pi_4(BG) \cong \mathbb{Z}$ and there are infinitely many isomorphism classes of principal G -bundles over S^4 . In [4], Kono classified the homotopy types of $SU(2)$ -gauge groups over S^4 and showed that there are six homotopy types of these groups.

Since then, many classification results on the homotopy types of gauge groups of low-rank simple Lie groups have been obtained. Among compact Lie groups, $SU(2)$ is the rank 1 simply-connected simple Lie group. There are three simply-connected simple compact Lie groups of rank 2: $SU(3)$, $Sp(2)$, and G_2 . For $G = SU(3)$, Hamanaka and Kono [2] proved that $\mathcal{G}_k \simeq \mathcal{G}_{k'}$ if and only if $(k, 24) = (k', 24)$, where (m, n) is the greatest common divisor of m and n . For $G = Sp(2)$, Theriault [8] showed $\mathcal{G}_k \simeq_{(p)} \mathcal{G}_{k'}$ at any prime p if and only if $(k, 40) = (k', 40)$, where $\simeq_{(p)}$ means p -local homotopy equivalence. Then, for $G = G_2$, Kishimoto, Theriault and Tsutaya [3] showed that $\mathcal{G}_k \simeq_{(p)} \mathcal{G}_{k'}$ at any prime p if $(k, 168) = (k', 168)$ and if $\mathcal{G}_k \simeq_{(p)} \mathcal{G}_{k'}$ at any prime p , then $(k, 84) = (k', 84)$. The discrepancy between $(k, 168) = (k', 168)$ and $(k, 84) = (k', 84)$ has remained as an open problem.

In this paper, we complete the classification for rank 2 simple Lie groups up to p -local homotopy equivalence at any prime p by proving the following.

Theorem 1.1. *Let \mathcal{G}_k be the gauge group of a principal G_2 -bundle over S^4 whose classifying map is $k \in \mathbb{Z} \cong \pi_4(BG_2)$. The following holds.*

- (1) *If $\mathcal{G}_k \simeq \mathcal{G}_{k'}$, then $(k, 84) = (k', 84)$ and*
- (2) *$\mathcal{G}_k \simeq_{(p)} \mathcal{G}_{k'}$ at any prime p if $(k, 84) = (k', 84)$.*

Theorem 1.1 (2) is equivalent to the following local form.

Proposition 1.2. *Let \mathcal{G}_k be the gauge group of a principal G_2 -bundle over S^4 whose classifying map is $k \in \mathbb{Z} \cong \pi_4(BG_2)$. Then, we have*

- (1) *$\mathcal{G}_k \simeq_{(2)} \mathcal{G}_{k'}$ if $(k, 4) = (k', 4)$,*

2020 *Mathematics Subject Classification.* 55P15, 54C35.

Key words and phrases. homotopy type, gauge group, Lie group.

JSPS KAKENHI Grant Number 25K07013 supported this work.

- (2) $\mathcal{G}_k \simeq_{(3)} \mathcal{G}_{k'}$ if $(k, 3) = (k', 3)$,
- (3) $\mathcal{G}_k \simeq_{(7)} \mathcal{G}_{k'}$ if $(k, 7) = (k', 7)$ and
- (4) $\mathcal{G}_k \simeq_{(p)} \mathcal{G}_{k'} \simeq_{(p)} G_2 \times \Omega_0^4 G_2$ for $p \neq 2, 3, 7$.

Theorem 1.1 (1) was established in [3, Theorem 1.2 (a)] and Proposition 1.2 (2), (3), (4) were established in [3, Propositions 4.6 and 3.2]. In this paper, we prove Proposition 1.2 (1) and complete the proof of Theorem 1.1 assuming the results of [3].

First, we recall the basic method for determining the homotopy types of G -gauge groups over S^4 , which relates their homotopy types to Samelson products. We begin with the following homotopy fiber sequence of mapping spaces.

$$\Omega \text{Map}(S^4, BG)_k \rightarrow G \rightarrow \text{Map}_*(S^4, BG)_k \rightarrow \text{Map}(S^4, BG)_k \xrightarrow{\text{ev}} BG,$$

where $\text{Map}(S^4, BG)_k$ is the connected component of the space of continuous maps containing the map $k: S^4 \rightarrow BG$, $\text{Map}_*(S^4, BG)_k$ is its subspace consisting of base point preserving maps, and ev is the evaluation map. There is a homotopy equivalence $\Omega_0^3 G \simeq \text{Map}_*(S^4, BG)_0 \simeq \text{Map}_*(S^4, BG)_k$ and Gottlieb [1] showed that the classifying space of the gauge group $B\mathcal{G}_k$ is homotopy equivalent to the mapping space $\text{Map}(S^4, BG)_k$. Therefore, we have the following fiber sequence.

$$\mathcal{G}_k \rightarrow G \xrightarrow{\partial_k} \Omega_0^3 G \rightarrow B\mathcal{G}_k \rightarrow BG.$$

Thus, the G -gauge group \mathcal{G}_k is homotopy equivalent to the homotopy fiber of the map ∂_k . Let $i_3: S^3 \rightarrow G$ be the inclusion map of the bottom cell. Then, Lang [5] proved that the map ∂_k is the triple adjoint of the Samelson product $\langle k \cdot i_3, 1 \rangle$ where 1 is the identity map of G . By the linearity of the Samelson product, we have $\langle k \cdot i_3, 1 \rangle \simeq k \cdot \langle i_3, 1 \rangle$. Furthermore, in [8], Theriault showed that \mathcal{G}_k is p -locally homotopy equivalent to $\mathcal{G}_{k'}$ at all prime p if $(k, m) = (k', m)$ where m is the order of the Samelson product $\langle i_3, 1 \rangle$.

For an abelian group A , we denote by $A_{(p)}$ its localization at the prime p . If A is finite, then $A_{(p)}$ is the p -primary subgroup of A . By [3, Corollary 6.6], the order of $\langle i_3, 1 \rangle$ in $[\Sigma^3 G_2, G_2]_{(2)}$ is at least 4. Therefore, to show Proposition 1.2 (1), it suffices to prove the following proposition.

Proposition 1.3. *The order of the Samelson product $\langle i_3, 1 \rangle$ in $[\Sigma^3 G_2, G_2]_{(2)}$ is at most 4.*

As for the order of the Samelson product $\langle i_3, 1 \rangle$ in $[\Sigma^3 G_2, G_2]$, together with [3, Lemma 3.1, Proposition 4.5 and Corollary 6.6], by Proposition 1.3, we have the following theorem.

Theorem 1.4. *The order of the Samelson product $\langle i_3, 1 \rangle$ in $[\Sigma^3 G_2, G_2]$ is 84.*

Next, we recall some properties of a space $V := G_2/SU(2)$, which played an important role in [3] and is crucial in this paper. Let $q: G_2 \rightarrow V$ be the obvious projection map. By [3, Lemmas 7.1 and 7.2], the Samelson product $3 \cdot \langle i_3, 1 \rangle: \Sigma^3 G_2 \rightarrow G_2$ factors through $\Sigma^3 q: \Sigma^3 G_2 \rightarrow \Sigma^3 V$. Let us write this factorization as follows.

$$3 \cdot \langle i_3, 1 \rangle \simeq \psi \circ \Sigma^3 q.$$

Let $P^{n+1}(2)$ be the mapping cone of the degree 2 map $\times 2: S^n \rightarrow S^n$. The 6-skeleton of $V = S^5 \cup e^6 \cup e^{11}$ is $P^6(2)$ and, by [3] Lemma 7.4, there is a homotopy equivalence $\Sigma^3 V \simeq P^9(2) \vee S^{14}$. Let

$$p_{11}: V \rightarrow S^{11}$$

be the pinch map to the top cell, collapsing the 6-skeleton to the base point. We denote by

$$i_6: P^6(2) \rightarrow V$$

the inclusion map. Then, the above homotopy equivalence provides maps $s': \Sigma^3 V \rightarrow P^9(2)$ and $s'': S^{14} \rightarrow \Sigma^3 V$ such that the identity map of $\Sigma^3 V$ is homotopic to

$$\Sigma^3 i_6 \circ s' + s'' \circ \Sigma^3 p_{11}.$$

The following proposition is what we prove in the rest of this paper.

Proposition 1.5. *We have $4(\psi \circ s'') \simeq 0$ in $\pi_{14}(G_2)_{(2)}$.*

Proof of Proposition 1.3. We consider the following decomposition in $[\Sigma^3 G_2, G_2]$.

$$3 \cdot \langle i_3, 1 \rangle \simeq \psi \circ \Sigma^3 i_6 \circ s' \circ \Sigma^3 q + \psi \circ s'' \circ \Sigma^3 (p_{11} \circ q).$$

By [3, Lemma 5.2 (a) and (b)], the order of the identity map of $P^9(2)$ is 4. Hence, we have $4(\Sigma^3 i_6 \circ s') \simeq 0$. Therefore, by Proposition 1.5, the order of $\langle i_3, 1 \rangle$ in $[\Sigma^3 G_2, G_2]_{(2)}$ is at most 4. \square

In what follows, we use the symbol p to express the map $p: G_2 \rightarrow S^6 \simeq G_2/SU(3)$ only. When we need to indicate the generator of a cyclic group, we write $\mathbb{Z}/m\{a\}$ for the cyclic group of order m generated by a .

This paper is organized as follows. In Section 2, we collect some facts on homotopy groups of G_2 and S^6 and reduce the problem on the 2-primary homotopy group $\pi_{14}(G_2)_{(2)}$ to that of $\pi_{14}(S^6)_{(2)}$. In Section 3, we prove Proposition 1.5.

2. HOMOTOPY GROUPS

For $3 \leq i \leq 14$, homotopy groups $\pi_i(G_2)$ are computed by Mimura [6, Theorem 6.1].

$$\begin{aligned} \pi_3(G_2) &= \mathbb{Z}, & \pi_4(G_2) &= \{0\}, & \pi_5(G_2) &= \{0\}, & \pi_6(G_2) &= \mathbb{Z}/3, \\ \pi_7(G_2) &= \{0\}, & \pi_8(G_2) &= \mathbb{Z}/2, & \pi_9(G_2) &= \mathbb{Z}/6, & \pi_{10}(G_2) &= \{0\}, \\ \pi_{11}(G_2) &= \mathbb{Z} \oplus \mathbb{Z}/2, & \pi_{12}(G_2) &= \{0\}, & \pi_{13}(G_2) &= \{0\}, & \pi_{14}(G_2) &= \mathbb{Z}/168 \oplus \mathbb{Z}/2 \end{aligned}$$

For the same range, homotopy groups of S^6 are as follows.

$$\begin{aligned} \pi_3(S^6) &= \{0\}, & \pi_4(S^6) &= \{0\}, & \pi_5(S^6) &= \{0\}, & \pi_6(S^6) &= \mathbb{Z}, \\ \pi_7(S^6) &= \mathbb{Z}/2, & \pi_8(S^6) &= \mathbb{Z}/2, & \pi_9(S^6) &= \mathbb{Z}/24, & \pi_{10}(S^6) &= \{0\}, \\ \pi_{11}(S^6) &= \mathbb{Z}, & \pi_{12}(S^6) &= \mathbb{Z}/2, & \pi_{13}(S^6) &= \mathbb{Z}/60, & \pi_{14}(S^6) &= \mathbb{Z}/24 \oplus \mathbb{Z}/2 \end{aligned}$$

We collect here miscellaneous remarks on homotopy groups of spheres.

Proposition 2.1. *The following holds.*

- (1) $\pi_8(S^6) = \mathbb{Z}/2\{\eta_6^2\}$, $\pi_{11}(S^6) = \mathbb{Z}\{\Delta(\iota_{13})\}$, $\pi_{14}(S^6)_{(2)} = \mathbb{Z}/8\{\bar{\nu}_6\} \oplus \mathbb{Z}/2\{\varepsilon_6\}$,
- (2) $\pi_{14}(S^8) = \mathbb{Z}/2\{\nu_8^2\}$, $\pi_{14}(S^9) = \{0\}$, $\pi_{14}(S^{10}) = \{0\}$.
- (3) $\eta_n \nu_{n+1} \simeq 0$ for $n \geq 5$.

Proof. We refer the reader to the following part of Toda's book [9]. For (1), Proposition 5.3 on page 40, Proposition 5.9 on page 44, and Theorem 7.1 on page 61. For (2), Proposition 5.11 on page 46, Proposition 5.9 on page 44, and Proposition 5.8 on page 43. For (3), (5.9) on page 44. \square

Mimura computed the homotopy groups of Lie groups of low rank using the homotopy groups of spheres. For instance, fiber sequences $S^3 \rightarrow SU(3) \rightarrow S^5$, $SU(3) \xrightarrow{i} G_2 \xrightarrow{p} S^6$, where $S^3 = SU(2)$, $S^5 = SU(3)/SU(2)$, $S^6 = G_2/SU(3)$, give us exact sequences involving homotopy groups of spheres and homotopy groups of $SU(3)$ and G_2 . The following is a part of Theorem 6.1 in [6].

Proposition 2.2 (Mimura). *Let $i: SU(3) \rightarrow G_2$ be the inclusion map and $p: G_2 \rightarrow S^6 = G_2/SU(3)$ the projection map. By $\langle \alpha \rangle$, we denote an element in $\pi_i(G_2)$ such that $p_*(\langle \alpha \rangle) \simeq \alpha$. The following holds.*

- (1) $\pi_{11}(G_2) = \mathbb{Z}\{\langle 2\Delta(\iota_{13}) \rangle\} \oplus \mathbb{Z}/2\{i_*([\nu_5^2])\}$.
- (2) $\pi_{14}(G_2)_{(2)} = \mathbb{Z}/8\{\langle \bar{\nu}_6 + \varepsilon_6 \rangle\} \oplus \mathbb{Z}/2\{i_*([\nu_5^2]) \circ \nu_{11}\}$

We close this section by proving the following proposition. With this proposition, we may reduce the problem on $\pi_{14}(G_2)_{(2)}$ to that on $\pi_{14}(S^6)_{(2)}$.

Proposition 2.3. *$4(p \circ \psi \circ s'') \simeq 0$ in $\pi_{14}(S^6)_{(2)}$ if and only if $4(\psi \circ s'') \simeq 0$ in $\pi_{14}(G_2)_{(2)}$.*

Proof. The 2-primary component of the homotopy group $\pi_{14}(S^6)$ is $\mathbb{Z}/8\{\bar{\nu}_6\} \oplus \mathbb{Z}/2\{\varepsilon_6\}$. The 2-primary component of $\pi_{14}(G_2)$ is $\mathbb{Z}/8\{\langle \bar{\nu}_6 + \varepsilon_6 \rangle\} \oplus \mathbb{Z}/2\{i_*([\nu_5^2]) \circ \nu_{11}\}$, so that $p_*(\langle \bar{\nu}_6 + \varepsilon_6 \rangle) \simeq \bar{\nu}_6 + \varepsilon_6$ and $p_*(i_*([\nu_5^2]) \circ \nu_{11}) \simeq 0$. Suppose that $\psi \circ s'' \simeq a\langle \bar{\nu}_6 + \varepsilon_6 \rangle + bi_*([\nu_5^2]) \circ \nu_{11}$ in $\pi_{14}(G_2)_{(2)}$ where $a \in \mathbb{Z}/8$ and $b \in \mathbb{Z}/2$. Then, $4(\psi \circ s'') \simeq 4a\langle \bar{\nu}_6 + \varepsilon_6 \rangle$ and $4(p \circ \psi \circ s'') \simeq 4a\bar{\nu}_6$. Therefore, $4(\psi \circ s'') \simeq 0$ if and only if $4a = 0$ in $\mathbb{Z}/8$ and $4(p \circ \psi \circ s'') \simeq 0$ if and only if $4a = 0$ in $\mathbb{Z}/8$. It completes the proof. \square

3. PROOF OF PROPOSITION 1.5

In the following commutative diagram, $S^3 = SU(2)$, $S^5 = SU(3)/SU(2)$, $S^6 = G_2/SU(3)$ and all vertical and horizontal sequences are fiber sequences.

$$\begin{array}{ccccc}
 S^3 & \longrightarrow & SU(3) & \longrightarrow & S^5 \\
 \downarrow = & & \downarrow i & & \downarrow i_5 \\
 S^3 & \xrightarrow{i_3} & G_2 & \xrightarrow{q} & V \\
 \downarrow & & \downarrow p & & \downarrow p_6 \\
 * & \longrightarrow & S^6 & \xrightarrow{=} & S^6
 \end{array}$$

Let $C = S^6 \cup e^{11}$ be a space obtained by collapsing the bottom cell S^5 of $V = S^5 \cup e^6 \cup e^{11}$ to the base point and $j: V \rightarrow C$ the pinch map. We denote by \bar{i}_6 the inclusion map of the bottom cell of C . We consider the following homotopy commutative diagram whose horizontal and vertical sequences are all cofiber

sequences.

$$\begin{array}{ccccc}
P^6(2) & \xrightarrow{i_6} & V & \xrightarrow{p_{11}} & S^{11} \\
\downarrow & & \downarrow j & & \downarrow = \\
S^6 & \xrightarrow{\bar{i}_6} & C & \xrightarrow{\bar{p}_{11}} & S^{11} \\
\downarrow \times 2 & & \downarrow \bar{p}_6 & & \downarrow \\
S^6 & \xrightarrow{=} & S^6 & \longrightarrow & *
\end{array}$$

Since the attaching map $S^{10} \rightarrow S^6$ is null homotopic, $C = S^6 \cup e^{11} \simeq S^6 \vee S^{11}$ and there is a homotopy equivalence $i': C \rightarrow S^6 \vee S^{11}$, $\bar{s}: S^{11} \rightarrow C$ such that $i' \circ \bar{i}_6: S^6 \rightarrow S^6 \vee S^{11}$ is homotopic to the canonical inclusion of S^6 into $S^6 \vee S^{11}$ and \bar{p}_{11} is homotopic to the composition of i' followed by the canonical projection $S^6 \vee S^{11} \rightarrow S^{11}$. Let us choose a splitting $\bar{s}: S^{11} \rightarrow C$ such that $\bar{p}_{11} \circ \bar{s} \simeq \iota_{11}$ where ι_{11} is the identity map of S^{11} . Since i_5 is homotopic to the inclusion map of the bottom cell S^5 into V and since $p_6 \circ i_5$ is null homotopic, we have a map $\varphi: C \rightarrow S^6$ such that $\varphi \circ j \simeq p_6$. We recall the following theorem of Ōshima.

Theorem 3.1 (Ōshima [7, Theorem 2.1]). *There is a map $\gamma: S^{11} \rightarrow G_2$ such that*

- (1) $p \circ \gamma \simeq 2\Delta(\iota_{13})$,
- (2) $\langle i_3, \gamma \rangle \in \pi_{14}(G_2)$ has order $21 = 3 \cdot 7$.

Now, we deal with $p \circ \psi \circ \Sigma^3(q \circ \gamma) \in \pi_{14}(S^6)$ to prove Proposition 1.5. Since the identity map of $\Sigma^3 V$ is homotopic to $\Sigma^3 i_6 \circ s' + s'' \circ \Sigma^3 p_{11}$, we have

$$p \circ \psi \circ \Sigma^3(q \circ \gamma) \simeq p \circ \psi \circ \Sigma^3 i_6 \circ s' \circ \Sigma^3(q \circ \gamma) + p \circ \psi \circ s'' \circ \Sigma^3(p_{11} \circ q \circ \gamma).$$

To show that the first term $p \circ \psi \circ \Sigma^3 i_6 \circ s' \circ \Sigma^3(q \circ \gamma)$ is null homotopic, we prove the following proposition. It is an elementary proposition on the homotopy groups of spheres in low degrees, independent of the rest of Section 3, and we use only the results stated in Proposition 2.1.

Proposition 3.2. *Let $i_8: S^8 \rightarrow P^9(2)$ be the inclusion map of the bottom cell. Then, the following hold.*

- (1) $i_{8*}: \pi_{14}(S^8) \rightarrow \pi_{14}(P^9(2))$ is an isomorphism.
- (2) For maps $f: S^{14} \rightarrow S^8$, and $g: S^8 \rightarrow S^6$, $g \circ f \simeq 0$ in $\pi_{14}(S^6)$.
- (3) If a map $f: S^{14} \rightarrow S^6$ factors through $P^9(2)$, then $f \simeq 0$ in $\pi_{14}(S^6)$.

Proof. (1) Since S^8 and $P^9(2)$ are 7-connected, homotopy groups $\pi_{14}(S^8)$ and $\pi_{14}(P^9(2))$ are isomorphic to stable homotopy groups $\pi_{14}^S(S^8)$ and $\pi_{14}^S(P^9(2))$, respectively. Since $\pi_{14}^S(S^9) \simeq \{0\}$ and $\pi_{14}^S(S^8) \simeq \mathbb{Z}/2$, from the following long exact sequence of stable homotopy groups

$$\pi_{14}^S(S^8) \xrightarrow{\times 2} \pi_{14}^S(S^8) \xrightarrow{i_{8*}} \pi_{14}^S(P^9(2)) \rightarrow \pi_{14}^S(S^9)$$

associated with the cofiber sequence $S^8 \xrightarrow{i_8} P^9(2) \rightarrow S^9$, we obtain the desired result.

(2) $\pi_8(S^6) \simeq \mathbb{Z}/2\{\eta_6^2\}$ and $\pi_{14}(S^8) = \mathbb{Z}/2\{\nu_8^2\}$. Hence, $f \simeq a\eta_6^2 \simeq a\eta_6\eta_7$ and $g \simeq b\nu_8^2 \simeq b\nu_8\nu_{11}$ for some a, b . Therefore, we have $f \circ g \simeq ab(\eta_6\eta_7\nu_8\nu_{11}) \simeq 0$. since $\eta_7\nu_8 \simeq 0$.

(3) Suppose that $f \simeq g \circ h$ for some $g: P^9(2) \rightarrow S^6$ and $h: S^{14} \rightarrow P^9(2)$. By (1), there is a map $h': S^{14} \rightarrow S^8$ such that $h \simeq i_8 \circ h'$. Then, $f \simeq (g \circ i_8) \circ h'$. By (2), we have $f \simeq 0$. \square

To describe the second term $p \circ \psi \circ s'' \circ \Sigma^3(p_{11} \circ q \circ \gamma)$, we prepare the following proposition.

Proposition 3.3. *The following holds.*

- (1) $\pi_{11}(C) \simeq \mathbb{Z}\{\bar{i}_6 \circ \Delta(\iota_{13})\} \oplus \mathbb{Z}\{\bar{s}\}$,
- (2) $\text{Ker } \bar{p}_6_*: \pi_{11}(C) \rightarrow \pi_{11}(S^6)$ is generated by $w \simeq a\bar{i}_6 \circ \Delta(\iota_{13}) + b\bar{s}$ where $a \in \mathbb{Z}$ and $b = 1$ or a is odd and $b = 2$.
- (3) $j \circ q \circ \gamma \simeq cw$ where $c \in \{\pm 1, \pm 2\}$.
- (4) $p_{11} \circ q \circ \gamma \simeq bc(\iota_{11})$ where $bc \in \{\pm 1, \pm 2, \pm 4\}$.

Proof. (1) C is homotopy equivalent to $S^6 \vee S^{11}$. The pair $(S^6 \times S^{11}, S^6 \vee S^{11})$ is 16-connected. Let $i': S^6 \vee S^{11} \rightarrow S^6 \times S^{11}$ be the inclusion map and we may choose the homotopy equivalence so that $i' \circ \bar{i}_6$ and $i' \circ \bar{s}$ are homotopic to obvious inclusion maps of S^6 and S^{11} , respectively. Then, the induced homomorphism $i'_*: \pi_{11}(S^6 \vee S^{11}) \rightarrow \pi_{11}(S^6 \times S^{11})$ is an isomorphism. It is clear that $\pi_{11}(S^6 \times S^{11}) \simeq \pi_{11}(S^6) \oplus \pi_{11}(S^{11})$ and $i' \circ \bar{i}_6 \circ \Delta(\iota_{13})$ and $i' \circ \bar{s}$ generate $\pi_{11}(S^6 \times S^{11})$.

(2) Suppose that $\bar{p}_6 \circ \bar{s} \simeq a'\Delta(\iota_{13})$ for some integer a' . If a' is even, let $a = -a'/2$. Then, $a\bar{i}_6 \circ \Delta(\iota_{13}) + \bar{s}$ is a generator of the kernel. If a' is odd, let $a = -a'$. Then, $a(\bar{i}_6 \circ \Delta(\iota_{13})) + 2(\bar{s})$ is a generator of the kernel of $(\bar{p}_6)_*$.

(3) Since $\bar{p}_6 \circ j \simeq 0$, $j \circ q \circ \gamma \simeq cw$ for some $c \in \mathbb{Z}$. Suppose that $\varphi \circ w = d\Delta(\iota_{13})$ for some $d \in \mathbb{Z}$. Then, $\varphi \circ j \circ q \circ \gamma \simeq cd\Delta(\iota_{13})$. On the other hand, we have $\varphi \circ j \circ q \circ \gamma \simeq p_6 \circ q \circ \gamma \simeq p \circ \gamma \simeq 2\Delta(\iota_{13})$. Therefore, $cd = 2$. It implies that $c \in \{\pm 1, \pm 2\}$.

(4) We have $p_{11} \circ q \circ \gamma \simeq \bar{p}_{11} \circ j \circ q \circ \gamma \simeq c(\bar{p}_{11} \circ w) \simeq bc(\iota_{11})$. Since $b \in \{1, 2\}$ and $c \in \{\pm 1, \pm 2\}$, we have $bc \in \{\pm 1, \pm 2, \pm 4\}$. \square

Finally, with the above preparations, we are now ready to prove Proposition 1.5.

Proof of Proposition 1.5. By Proposition 2.3, to complete the proof, it suffices to show that $4(p \circ \psi \circ s'') \simeq 0$ in $\pi_{14}(S^6)_{(2)}$.

The identity map of $\Sigma^3 V$ is homotopic to $\Sigma^3 i_6 \circ s' + s'' \circ \Sigma^3 p_{11}$. Hence, we have

$$p \circ \psi \circ \Sigma^3(q \circ \gamma) \simeq p \circ \psi \circ \Sigma^3 i_6 \circ s' \circ \Sigma^3(q \circ \gamma) + p \circ \psi \circ s'' \circ \Sigma^3(p_{11} \circ q \circ \gamma).$$

By Proposition 3.2 (3), we have $p \circ \psi \circ \Sigma^3 i_6 \circ s' \circ \Sigma^3(q \circ \gamma) \simeq 0$ in $\pi_{14}(S^6)$. By Proposition 3.3 (4), we have $\Sigma^3(p_{11} \circ q \circ \gamma) \simeq bc(\iota_{14})$ in $\pi_{14}(G_2)$ and $bc \in \{\pm 1, \pm 2, \pm 4\}$. Therefore, in $\pi_{14}(S^6)$, we have

$$p \circ \psi \circ \Sigma^3(q \circ \gamma) \simeq bc(p \circ \psi \circ s'').$$

On the other hand, by Ōshima's theorem,

$$\psi \circ \Sigma^3(q \circ \gamma) \simeq \langle i_3, \gamma \rangle$$

has order 21, so that it is null homotopic in $\pi_{14}(G_2)_{(2)}$. Therefore, in $\pi_{14}(S^6)_{(2)}$, we have

$$p \circ \psi \circ \Sigma^3(q \circ \gamma) \simeq 0.$$

Hence, in $\pi_{14}(S^6)_{(2)}$, we have $bc(p \circ \psi \circ s'') \simeq 0$. If $bc \in \{\pm 1\}$, then $p \circ \psi \circ s'' \simeq 0$. If $bc \in \{\pm 2\}$, then $2(p \circ \psi \circ s'') \simeq 0$ and so $p \circ \psi \circ s'' \simeq 4e\bar{\nu}_6$ for some integer e . If $bc \in \{\pm 4\}$, $4(p \circ \psi \circ s'') \simeq 0$ and so $p \circ \psi \circ s'' \simeq \pm 2e\bar{\nu}_6$ for some integer

e. Therefore, in either case, we have $4(p \circ \psi \circ s'') \simeq 0$ in $\pi_{14}(S^6)_{(2)}$. Hence, by Proposition 2.3, we establish $4(\psi \circ s'') \simeq 0$ in $\pi_{14}(G_2)_{(2)}$, and it completes the proof of Proposition 1.5. \square

REFERENCES

- [1] D. H. Gottlieb, *Applications of bundle map theory*, Trans. Amer. Math. Soc. **171** (1972), 23–50, DOI 10.2307/1996373. MR0309111
- [2] H. Hamanaka and A. Kono, *Unstable K^1 -group and homotopy type of certain gauge groups*, Proc. Roy. Soc. Edinburgh Sect. A **136** (2006), no. 1, 149–155, DOI 10.1017/S0308210500004480. MR2217512
- [3] D. Kishimoto, S. Theriault, and M. Tsutaya, *The homotopy types of G_2 -gauge groups*, Topology Appl. **228** (2017), 92–107, DOI 10.1016/j.topol.2017.05.012. MR3679077
- [4] A. Kono, *A note on the homotopy type of certain gauge groups*, Proc. Roy. Soc. Edinburgh Sect. A **117** (1991), no. 3-4, 295–297, DOI 10.1017/S0308210500024732. MR1103296
- [5] G. E. Lang Jr., *The evaluation map and EHP sequences*, Pacific J. Math. **44** (1973), 201–210. MR0341484
- [6] M. Mimura, *The homotopy groups of Lie groups of low rank*, J. Math. Kyoto Univ. **6** (1967), 131–176, DOI 10.1215/kjm/1250524375. MR0206958
- [7] H. Ōshima, *Samelson products in the exceptional Lie group of rank 2*, J. Math. Kyoto Univ. **45** (2005), no. 2, 411–420, DOI 10.1215/kjm/1250281999. MR2161701
- [8] S. D. Theriault, *The homotopy types of $\mathrm{Sp}(2)$ -gauge groups*, Kyoto J. Math. **50** (2010), no. 3, 591–605, DOI 10.1215/0023608X-2010-005. MR2723863
- [9] H. Toda, *Composition methods in homotopy groups of spheres*, Annals of Mathematics Studies, vol. No. 49, Princeton University Press, Princeton, NJ, 1962. MR0143217

DEPARTMENT OF MATHEMATICAL SCIENCES, SHIBAURA INSTITUTE OF TECHNOLOGY, 307 MINUMAKU FUKASAKU, SAITAMA-CITY 337-8570, JAPAN
Email address: kameko@shibaura-it.ac.jp