

A note on the RAGE Theorem and phase-averaged dispersion for the Fibonacci Hamiltonian

Gaétan Leclerc

December 9, 2025

Abstract

We find a weaker condition on spectral measures, “eventual absolute continuity”, that ensure quantum delocalization as in the RAGE Theorem in the case of purely absolutely continuous spectrum. We then adapt these idea to strongly improve some phase-averaged delocalization bounds for the Fibonacci quasicrystal.

1 On the RAGE Theorem

In these notes, we discuss the spectral Theory of bounded selfadjoints operators, and in particular we discuss the Fibonacci Hamiltonian. A complete source on this topic is given by the books [DF22, DF24]. For an introduction, one can consult the survey [Da17].

Let \mathcal{H} be a Hilbert space. Let $H : \mathcal{H} \rightarrow \mathcal{H}$ be a Hamiltonian: to simplify, suppose that H is bounded and selfadjoint. For $\psi \in \mathcal{H}$, we denote by μ_ψ the spectral measure associated with H and ψ . This finite measure is supported on the spectrum $\sigma(H) \subset \mathbb{R}$ of H (we will denote $\mu_\psi \in \mathcal{P}(\sigma(H))$). We write $\psi \in \mathcal{H}_{ac}$ if μ_ψ is absolutely continuous with respect to the Lebesgue measure ($\mu_\psi \ll \lambda$). The set \mathcal{H}_{ac} is a closed linear subspace of \mathcal{H} .

The famous RAGE Theorem, in the (easy) absolutely continuous case, state that:

Theorem 1.1. *Let $\psi \in \mathcal{H}_{ac}$. Then for every $\varphi \in \mathcal{H}$,*

$$\langle e^{itH} \psi, \varphi \rangle \xrightarrow[t \rightarrow \infty]{} 0.$$

Proof. Let $\psi \in \mathcal{H}_{ac}$. Notice that for any Borel set E with zero lebesgue measure $|E| = 0$, we have vanishing of the spectral projector:

$$\|\mathbb{1}_E(H)\psi\|_{\mathcal{H}}^2 = \langle \mathbb{1}_E^2(H)\psi, \psi \rangle_{\mathcal{H}} = \int_{\sigma(H)} \mathbb{1}_E(t) d\mu_\psi(t) = \mu_\psi(E) = 0$$

by absolute continuity of μ_ψ . Now, consider the linear form:

$$f \in C^0(\sigma(H), \mathbb{C}) \mapsto \langle f(H)\psi, \varphi \rangle_{\mathcal{H}} \in \mathbb{C}.$$

The linear form is continuous for the topology of $L^\infty(\sigma(H), \mathbb{C})$, since $\|f(H)\|_{\mathcal{H} \rightarrow \mathcal{H}} \leq \|f\|_{L^\infty(\sigma(H), \mathbb{C})}$ by the usual properties of the fonctionnal calculus. By Riesz representation theorem, there exists a complex measure $\mu_{\psi, \varphi}$ such that $\langle f(H)\psi, \varphi \rangle_{\mathcal{H}} = \int f d\mu_{\psi, \varphi}$. But notice that for any borel set E with $|E| = 0$, we have

$$\mu_{\varphi, \psi}(E) = \langle \mathbb{1}_E(H)\psi, \varphi \rangle_{\mathcal{H}} = 0.$$

It follows that $\mu_{\varphi, \psi}$ is absolutely continuous with respect to the lebesgue measure. We then find

$$\langle e^{itH} \psi, \varphi \rangle = \hat{\mu}_{\psi, \varphi}(t) \rightarrow 0,$$

by the Riemann-Lebesgue Lemma. □

The condition that μ_ψ is absolutely continuous can actually be relaxed. For any $N \geq 1$, we define another Hamiltonian $H_k = I \otimes \cdots \otimes H \otimes \cdots \otimes I$ acting on $\mathcal{H}^{\otimes N}$, and we then define $\widehat{H} := \sum_k H_k$. One can then look at the spectral measure of $\psi^{\otimes N}$ associated to \widehat{H} , denoted $\mu_{\psi^{\otimes N}}$. Recall that we have the formula:

$$\mu_{\psi^{\otimes N}} = \mu_\psi^{*N},$$

where $*$ denotes a convolution. In fact, in general, we have $\mu_{\psi \otimes \varphi} = \mu_\psi * \mu_\varphi$. This is because:

$$\begin{aligned} \widehat{\mu_{\psi \otimes \varphi}}(t) &= \langle e^{it(I \otimes H + H \otimes I)} \psi \otimes \varphi, \psi \otimes \varphi \rangle_{\mathcal{H} \otimes \mathcal{H}} = \langle (e^{itH} \psi) \otimes (e^{itH} \varphi), \psi \otimes \varphi \rangle_{\mathcal{H} \otimes \mathcal{H}} \\ &= \langle e^{itH} \psi, \psi \rangle_{\mathcal{H}} \cdot \langle e^{itH} \varphi, \varphi \rangle_{\mathcal{H}} = \widehat{\mu_\psi}(t) \widehat{\mu_\varphi}(t) = \widehat{\mu_\psi * \mu_\varphi}(t). \end{aligned}$$

Definition 1.1. Let us define the eventually absolutely continuous subspace as:

$$\mathcal{H}_{eac} := \overline{\{\psi \in \mathcal{H} \mid \exists N \geq 1, \mu_{\psi^{\otimes N}} \ll \lambda\}}.$$

This is a closed linear subspace of \mathcal{H} .

Theorem 1.2 (RAGE). Suppose that $\psi \in \mathcal{H}_{eac}$. Then for every $\varphi \in \mathcal{H}$,

$$\langle e^{itH} \psi, \varphi \rangle \xrightarrow{t \rightarrow \infty} 0.$$

Proof. Let $\psi \in \mathcal{H}$ be such that $\mu_{\psi^{\otimes N}} = \mu_\psi^{*N}$ is absolutely continuous for some large enough N . Let $\varphi \in \mathcal{H}$. Then:

$$\langle e^{itH} \psi, \varphi \rangle_{\mathcal{H}}^N = \langle (e^{itH} \psi)^{\otimes N}, \varphi^{\otimes N} \rangle_{\mathcal{H}^{\otimes N}} = \langle e^{it\widehat{H}} \psi^{\otimes N}, \varphi^{\otimes N} \rangle_{\mathcal{H}^{\otimes N}} \xrightarrow{t \rightarrow \infty} 0$$

by the RAGE Theorem, since $\mu_{\psi^{\otimes N}}$ is absolutely continuous. We conclude for any $\psi \in \mathcal{H}_{eac}$ by density. \square

Question: What is the space $\mathcal{H}_{eac}^\perp \subset \mathcal{H}$?

2 On the Fibonacci Hamiltonian

Adapting the idea of the previous proof, we can establish strong phase-averaged delocalization for the Fibonacci Hamiltonian. Consider the Fibonacci Hamiltonian $H_\omega : \ell^2(\mathbb{Z}, \mathbb{C}) \rightarrow \ell^2(\mathbb{Z}, \mathbb{C})$, with phase $\omega \in \mathbb{T} = \mathbb{R}/\mathbb{Z}$, for some small enough fixed coupling constant $V > 0$. Recall that it is given by the formula:

$$H_\omega \psi(n) = \psi(n+1) + \psi(n-1) + V \mathbb{1}_{[1-\alpha, 1)}(\omega + n\alpha \bmod 1) \psi(n)$$

where $\alpha := (\sqrt{5} - 1)/2$ is the inverse of the golden ratio. These are (uniformly) bounded and selfadjoint linear operators. Notice that, denoting $S\psi(n) = \psi(n+1)$ the shift (unitary in $\ell^2(\mathbb{Z})$), we have $S^n H_\omega (S^*)^n = H_{\omega + \alpha n}$. In general, all the operators H_ω have the same spectrum $\Sigma_V \subset \mathbb{R}$, independantly on the choice of the phase. Define $\mu_\omega \in \mathcal{P}(\Sigma_V)$ to be the spectral measure associated with $\delta_0 \in \ell^2(\mathbb{Z}, \mathbb{C})$ and H_ω . Define also the density of state measure $\mu \in \mathcal{P}(\Sigma_V)$ by the formula

$$\mu = \int_{\omega \in \mathbb{T}} \mu_\omega d\omega.$$

It was proved in [Le25] that

$$|\widehat{\mu}(\xi)| \leq |\xi|^{-\varepsilon},$$

for some $\varepsilon > 0$ and for large enough $\xi > 0$. In other words, we have

$$\left| \int_{\omega \in \mathbb{T}} \langle e^{itH_\omega} \delta_0, \delta_0 \rangle_{\ell^2(\mathbb{Z}, \mathbb{C})} d\omega \right| \leq |t|^{-\varepsilon}$$

for large enough t . It follows that μ^{*N} is absolutely continuous for N large enough, since its Fourier transform will be e.g. in L^2 . Adapting the previous proof, we will show the following consequence:

Theorem 2.1 (Escape of mass). *For any ℓ^1 -localized state $\psi \in \ell^1(\mathbb{Z}, \mathbb{C})$, and for any $\varphi \in \ell^2(\mathbb{Z}, \mathbb{C})$, we have:*

$$\int_{\omega \in \mathbb{T}} \langle e^{itH_\omega} \psi, \varphi \rangle_{\ell^2(\mathbb{Z})} d\omega \xrightarrow{t \rightarrow \infty} 0.$$

Proof. Notice that, since

$$\mu = \int_{\omega} \mu_{\omega} d\omega,$$

we have

$$\mu^{*N} = \int_{\vec{\omega} \in \mathbb{T}^N} \mu_{\vec{\omega}} d\vec{\omega},$$

where $\mu_{\vec{\omega}} := \mu_{\omega_1} * \dots * \mu_{\omega_N}$. Define the naturally associated Hamiltonian $H_{\vec{\omega}} : \ell^2(\mathbb{Z}^N, \mathbb{C}) \rightarrow \ell^2(\mathbb{Z}^N, \mathbb{C})$ to be

$$H_{\vec{\omega}} = (H_{\omega_1} \otimes I \otimes \dots \otimes I) + \dots + (I \otimes \dots \otimes I \otimes H_{\omega_N}).$$

Notice then that, for any Borel set $E \subset \mathbb{R}$ with zero lenght $|E| = 0$, we have:

$$\int_{\vec{\omega} \in \mathbb{T}^N} \|\mathbb{1}_E(H_{\vec{\omega}}) \delta_0^{\otimes N}\|_{\ell^2(\mathbb{Z}^N)} d\vec{\omega} = 0.$$

Indeed, by Cauchy-Schwarz, by definition of the spectral measures, and by absolute continuity of μ^{*N} :

$$\begin{aligned} \left(\int_{\vec{\omega} \in \mathbb{T}^N} \|\mathbb{1}_E(H_{\vec{\omega}}) \delta_0^{\otimes N}\|_{\ell^2(\mathbb{Z}^N)} d\vec{\omega} \right)^2 &\leq \int_{\vec{\omega} \in \mathbb{T}^N} \|\mathbb{1}_E(H_{\vec{\omega}}) \delta_0^{\otimes N}\|_{\ell^2(\mathbb{Z}^N)}^2 d\vec{\omega} \\ &= \int_{\vec{\omega} \in \mathbb{T}^N} \langle \mathbb{1}_E^2(H_{\vec{\omega}}) \delta_0^{\otimes N}, \delta_0^{\otimes N} \rangle_{\ell^2(\mathbb{Z}^N)} d\vec{\omega} \\ &= \int_{\vec{\omega} \in \mathbb{T}^N} \int_{\Sigma_V} \mathbb{1}_E(t) d\mu_{\vec{\omega}}(t) d\vec{\omega} = \mu^{*N}(E) = 0. \end{aligned}$$

Be carefull that this doesn't imply that any of the measures μ_{ω} are absolutely continuous, and that they could very well all have atoms. Let us now fix any $\phi \in \ell^2(\mathbb{Z}^N, \mathbb{C})$, and consider the linear form:

$$f \in C^0(\Sigma_V, \mathbb{C}) \mapsto \int_{\vec{\omega} \in \mathbb{T}^N} \langle f(H_{\vec{\omega}}) \delta_0^{\otimes N}, \phi \rangle_{\ell^2(\mathbb{Z}^N)} d\vec{\omega} \in \mathbb{C}.$$

This linear form is continuous: by Riesz representation theorem, there exists a complex measure μ_{ϕ} supported on Σ_V such that

$$\int_{\vec{\omega} \in \mathbb{T}^N} \langle f(H_{\vec{\omega}}) \delta_0^{\otimes N}, \phi \rangle_{\ell^2(\mathbb{Z}^N)} d\vec{\omega} = \int_{\Sigma_V} f d\mu_{\phi}.$$

But notice that, for any Borel set $E \subset \mathbb{R}$ of zero lenght $|E| = 0$, we have

$$|\mu_{\phi}(E)| = \left| \int_{\vec{\omega} \in \mathbb{T}^N} \langle \mathbb{1}_E(H_{\vec{\omega}}) \delta_0^{\otimes N}, \phi \rangle_{\ell^2(\mathbb{Z}^N)} d\vec{\omega} \right| \leq \int_{\vec{\omega} \in \mathbb{T}^N} \|\mathbb{1}_E(H_{\vec{\omega}}) \delta_0^{\otimes N}\|_{\ell^2(\mathbb{Z}^N)} d\vec{\omega} \cdot \|\phi\|_{\ell^2(\mathbb{Z}^N)} = 0.$$

By the Radon-Nikodym theorem for complex measures, it follows that μ_{ϕ} is absolutely continuous. In particular, we have $\widehat{\mu_{\phi}}(t) \rightarrow 0$, which means that, for any $\phi \in \ell^2(\mathbb{Z}^d)$, we have:

$$\int_{\vec{\omega}} \langle e^{itH_{\vec{\omega}}} \delta_0^{\otimes N}, \phi \rangle_{\ell^2(\mathbb{Z}^N)} d\vec{\omega} \xrightarrow{t \rightarrow \infty} 0.$$

In particular, if ϕ is of the form $\phi = \varphi^{\otimes N}$, we find

$$\left(\int_{\omega \in \mathbb{T}} \langle e^{itH_{\omega}} \delta_0, \varphi \rangle_{\ell^2(\mathbb{Z})} d\omega \right)^N = \int_{\vec{\omega} \in \mathbb{T}^N} \langle e^{itH_{\vec{\omega}}} \delta_0^{\otimes N}, \varphi^{\otimes N} \rangle_{\ell^2(\mathbb{Z}^N)} d\vec{\omega} \xrightarrow{t \rightarrow \infty} 0.$$

Let us now conclude the proof. Take any $\psi \in \ell^1(\mathbb{Z}) \subset \ell^2(\mathbb{Z})$, and any $\varphi \in \ell^2(\mathbb{Z})$. We can write ψ as a sum (absolutely converging for $\|\cdot\|_{\ell^2(\mathbb{Z})}$):

$$\psi = \sum_{k \in \mathbb{Z}} \psi(k) S^{-k}(\delta_0),$$

where S is the shift. Hence, doing a change of variable in the integrals, using the relation $SH_\omega S^{-1} = H_{\omega+\alpha}$, and noticing that $\sum_{k \in \mathbb{Z}} \overline{\psi(k)} S^k \varphi \in \ell^2(\mathbb{Z})$:

$$\begin{aligned}
\int_{\omega \in \mathbb{T}} \langle e^{itH_\omega} \psi, \varphi \rangle d\omega &= \sum_{k \in \mathbb{Z}} \psi(k) \int_{\omega \in \mathbb{T}} \langle e^{itH_\omega} S^{-k} \delta_0, \varphi \rangle_{\ell^2(\mathbb{Z})} d\omega \\
&= \sum_{k \in \mathbb{Z}} \psi(k) \int_{\omega \in \mathbb{T}} \langle S^{-k} e^{itH_{\omega+k\alpha}} \delta_0, \varphi \rangle_{\ell^2(\mathbb{Z})} d\omega \\
&= \sum_{k \in \mathbb{Z}} \psi(k) \int_{\omega \in \mathbb{T}} \langle e^{itH_\omega} \delta_0, S^k \varphi \rangle_{\ell^2(\mathbb{Z})} d\omega \\
&= \int_{\omega \in \mathbb{T}} \left\langle e^{itH_\omega} \delta_0, \sum_{k \in \mathbb{Z}} \overline{\psi(k)} S^k \varphi \right\rangle_{\ell^2(\mathbb{Z})} d\omega \xrightarrow{t \rightarrow \infty} 0.
\end{aligned}$$

□

Acknowledgments

We thank Tuomas Sahlsten, Frederic Naud, Mostafa Sabri and Jake Fillman for discussions during the redaction of this note. It was mentioned to me that a “dual” work, in the sense that we restrict the absolutely continuous spectrum instead of extending it, can be found in the work of Avron and Simon [AS81].

References

- [AS81] J.E. Avron, B. Simon *Transient and recurrent spectrum* Journal of Functional Analysis Volume 43, Issue 1, August 1981, Pages 1-31 [doi:10.1016/0022-1236\(81\)90034-3](https://doi.org/10.1016/0022-1236(81)90034-3)
- [Le25] G. Leclerc, *Fourier decay of equilibrium states and the Fibonacci Hamiltonian*, Preprint [arXiv:2507.23731](https://arxiv.org/abs/2507.23731),
- [Da17] D. Damanik *Schrödinger operators with dynamically defined potentials*. Ergodic Theory and Dynamical Systems. 2017;37(6):1681-1764. [doi:10.1017/etds.2015.120](https://doi.org/10.1017/etds.2015.120). [arXiv:1410.2445](https://arxiv.org/abs/1410.2445)
- [DF22] D. Damanik, J. Fillman. *One-Dimensional Ergodic Schrödinger Operators: I. General Theory* Graduate Studies in Mathematics. Vol. 221, 2022. ISBN: 978-1-4704-5606-1 [doi:10.1090/gsm/221](https://doi.org/10.1090/gsm/221)
- [DF24] D. Damanik, J. Fillman. *One-Dimensional Ergodic Schrödinger Operators: II. Specific Classes* Graduate Studies in Mathematics. vol. 249, 2024. ISBN: 978-1-4704-6503-2