

HOLOMORPHIC ONE-FORMS WITHOUT ZEROS ON KÄHLER MANIFOLDS OF KODAIRA CODIMENSION ONE

SIMON PIETIG

ABSTRACT. We give a bimeromorphic classification of compact Kähler manifolds of Kodaira codimension one that admit a holomorphic one form without zeros.

CONTENTS

1. Introduction	1
2. Preliminaries	3
3. Deformations	9
4. Proof of the Main Result	10
References	13

1. INTRODUCTION

Let X be a smooth projective variety over \mathbb{C} with a nowhere vanishing holomorphic 1-form $\omega \in H^0(X, \Omega_X^1)$. In [PS17], Popa and Schnell showed that the Kodaira dimension of X is bounded by $\kappa(X) \leq \dim X - 1$. To prove this result, Popa-Schnell applied the theory of mixed Hodge modules. A fortiori, Hao gave the following geometric explanation of this behavior, for $\kappa(X) = \dim X - 1$.

Theorem 1.1. [Hao23, Thm. 1.3] *Let X be a smooth projective variety of Kodaira dimension $\kappa(X) = \dim X - 1$. If X has a nowhere vanishing holomorphic 1-form, then for any minimal model X^{\min} of X there exists a finite quasi-étale covering $X' \rightarrow X^{\min}$ such that any \mathbb{Q} -factorialization X'' of X' is a product $B \times E$, where B is a minimal model of general type and E is an elliptic curve.*

Recent work of Church [Chu24] and of Hao, Wang and Zhang [HWZ24] generalized Hao's result to a birational classification of smooth projective varieties with g everywhere linearly independent holomorphic 1-forms assuming the abundance conjecture without any restrictions of the Kodaira dimension of X .

The main result of our paper is the following generalization of Theorem 1.1 to the Kähler case. For a more precise version, see Theorem 4.1 down below.

Theorem 1.2. *Let X be a compact Kähler manifold of Kodaira dimension $\kappa(X) = \dim X - 1$. If X has a nowhere vanishing holomorphic 1-form, then there exist a finite étale cover $X' \rightarrow X$ such that X'*

LEIBNIZ UNIVERSITÄT HANNOVER, INSTITUT FÜR ALGEBRAISCHE GEOMETRIE, WELFENGARTEN 1, 30167 HANNOVER

PIETIG@MATH.UNI-HANNOVER.DE

ORCID ID

2020 *Mathematics Subject Classification.* 32Q15, 32J18, 32Q57, 32H04.

is bimeromorphic to an elliptic fiber bundle $Z \rightarrow B$ with trivial monodromy over a smooth projective base B of general type.

Remark 1.3. In contrast to the projective situation, one cannot expect that the finite étale covering space X' of X in Theorem 1.2 is bimeromorphic to a product $Z \cong B \times E$, see Theorem 2.7 below.

1.1. Outline of the argument. In the projective case, Hao argues as follows: Let X be a smooth projective variety over \mathbb{C} of Kodaira codimension 1 that admits a holomorphic 1-form $\omega \in H^0(X, \Omega_X^1)$ without zeros. Let X^{\min} be a good minimal model of X , and consider the Iitaka fibration $f: X^{\min} \rightarrow Y$. By [PS17, Thm. 2.1], the holomorphic 1-form ω cannot come from Y . In particular, the general fiber of f is not contracted by the Albanese morphism. Therefore, it is mapped to a translate of a fixed elliptic curve $E \subset \text{Alb}(X)$. As $\text{Alb}(X)$ is projective, we can dualize the inclusion $E \subset \text{Alb}(X)$ to get a morphism

$$\varphi: X^{\min} \longrightarrow \text{Alb}(X) \longrightarrow E.$$

The variety B in Theorem 1.1, as well as the splitting $X'' \cong B \times E$ of a \mathbb{Q} -factorialization $X'' \rightarrow X'$ of a finite quasi-étale cover $X' \rightarrow X^{\min}$ are then constructed from a general fiber of $\varphi: X^{\min} \rightarrow E$.

Now, let X be a compact connected Kähler manifold of Kodaira codimension 1 that admits a holomorphic 1-form $\omega \in H^0(X, \Omega_X^1)$ without zeros. Note that we cannot apply the result by Popa-Schnell [PS17, Thm. 2.1]. Moreover, $\text{Alb}(X)$ is not projective, and therefore, the morphism φ does not exist in general.

To overcome these obstacles, we invoke Lin's construction of an algebraic approximation via tautological models and families [Lin20] to obtain the following diagram

$$\begin{array}{ccccc} \mathcal{X} & \overset{\sim}{\dashrightarrow} & \mathcal{T}/G & & \\ \pi \searrow & & \swarrow \Pi/G & \searrow p/G & \\ & V & & Y/G. & \end{array}$$

Here, $\pi: \mathcal{X} \rightarrow V$ is a deformation of $X \cong \mathcal{X}_0 := \pi^{-1}(0)$ over a complex vector space V , and \mathcal{T} is a complex analytic variety on which a finite group G acts. The morphism

$$\Pi: \mathcal{T} \longrightarrow Y \times V \longrightarrow V$$

is an explicitly constructed G -equivariant family of elliptic fibrations $p_v: \Pi^{-1}(v) \rightarrow Y$ over a fixed smooth projective base Y , called the tautological family. The central elliptic fibration

$$p_0/G: \Pi^{-1}(0)/G \longrightarrow Y/G$$

is bimeromorphic to the Iitaka fibration of X . Moreover, the deformation $\pi: \mathcal{X} \rightarrow V$ has the property that the set of points $v \in V$ such that $\mathcal{X}_v := \pi^{-1}(v)$ is a projective manifold, is dense in the euclidean topology. The two families $\pi: \mathcal{X} \rightarrow V$ and $\Pi/G: \mathcal{T}/G \rightarrow V$ are bimeromorphic over V . The elliptic fibrations $p_v: \Pi^{-1}(v) \rightarrow Y$ are locally isomorphic over Y , i.e. there is an open cover $\{U_i\}_{i \in I}$ such that for every U_i the isomorphism class of the elliptic fibrations

$$p_v: \Pi^{-1}(v)|_{U_i} \longrightarrow U_i$$

does not depend on $v \in V$.

We then show that a small deformation of X also admits a holomorphic 1-form without zeros. In particular, there is a point $v_0 \in V$ such that \mathcal{X}_{v_0} is a projective manifold that admits a holomorphic 1-form without zeros. As the Iitaka fibration of \mathcal{X}_{v_0} is birational to $p_{v_0}/G: \Pi^{-1}(v_0)/G \rightarrow Y/G$, and the

elliptic fibrations $p_v: \Pi^{-1}(v) \rightarrow Y$ are locally isomorphic over Y , we conclude that the general fiber of the Iitaka fibration of any \mathcal{X}_v is isomorphic to a fixed elliptic curve. This overcomes the first obstacle.

To finish the argument, we show that the elliptic fibration $p_0: \Pi^{-1}(0) \rightarrow Y$ is locally trivial, i.e. locally isomorphic to a product. It remains to carefully describe the G -action on this fibration. Taking a carefully chosen subquotient, yields the desired elliptic fiber bundle $Z \rightarrow B$.

1.2. Conventions and notation. By a complex analytic variety, we mean a Hausdorff, second-countable, irreducible, and reduced complex space. A complex analytic variety is called Kähler if it admits a Kähler metric in the sense of [Var89, II. 1]. A compact complex analytic Kähler variety X is called minimal if it is \mathbb{Q} -factorial, has terminal singularities, and the canonical divisor K_X is nef. It is furthermore called a good minimal model, if a positive multiple of K_X is globally generated

Acknowledgments. I would like to thank Stefan Schreieder for suggesting this problem to me, for very helpful discussions, and for comments on this manuscript. The research was conducted in the framework of the DFG-funded research training group RTG 2965: From Geometry to Numbers.

2. PRELIMINARIES

2.1. Singular Kähler spaces. A complex analytic variety is called Kähler if it admits a Kähler metric in the sense of [Var89, II. 1]. We recall some standard results about compact Kähler varieties.

Proposition 2.1. [Var89, II. Prop. 1.3.1, Cor. 3.2.2], [GK20, Prop. 3.5] *Let $f: X \rightarrow Y$ be a morphism between compact complex analytic varieties. Then the following holds true.*

- (i) *If Y is Kähler and f is a closed embedding, then X is Kähler.*
- (ii) *If Y is Kähler and f is a projective morphism, then X is Kähler. In particular, by Hironaka's result on resolution of singularities, compact complex analytic Kähler varieties admit desingularizations by compact Kähler manifolds.*
- (iii) *If f is a finite surjective morphism, then X is Kähler if and only if Y is Kähler.*

2.2. Elliptic fibrations. If not stated otherwise, the following material can be found in [Nak02b, 1.1].

Definition 2.2. Let $f: X \rightarrow Y$ be a proper surjective morphism between normal complex analytic varieties.

- (i) The morphism f is called an elliptic fibration if it has connected fibers and the general fiber is a smooth curve of genus 1.
- (ii) The discriminant locus of an elliptic fibration is the not necessarily reduced closed complex subspace $\Delta \subset Y$ over which f is not smooth. The smooth locus is denoted by $Y^* := Y \setminus \Delta$.
- (iii) The elliptic fibration $f: X \rightarrow Y$ is called projective, if there exists an f -ample line bundle on X . Similarly, f is called locally projective, if there exists an open cover $Y = \bigcup U_i$ such that the elliptic fibrations $f^{-1}(U_i) \rightarrow U_i$ are projective for each U_i .

In the following, we will only consider elliptic fibrations $f: X \rightarrow Y$ over a connected complex manifold Y .

2.2.1. Jacobian fibrations.

Definition 2.3. Let $f: X \rightarrow Y$ be a smooth elliptic fibration.

- (i) The variation of Hodge structures H associated to f consists of the following data: the local system $H_{\mathbb{Z}} := R^1 f_* \mathbb{Z}$ on Y of free abelian groups of rank 2, the associated vector bundle $\mathcal{H} := H_{\mathbb{Z}} \otimes \mathcal{O}_Y$ on Y , the holomorphic subbundle $\mathcal{F}^1 := f_* \Omega_{X/Y}^1$, and the quotient $\mathcal{H}^{0,1} := \mathcal{H}/\mathcal{F}^1$.
- (ii) We say that f has trivial monodromy if the local system $H_{\mathbb{Z}}$ is isomorphic to \mathbb{Z}^2 .
- (iii) We define the Jacobian fibration associated to f as the complex manifold

$$J_H := \mathcal{H}^{0,1}/H_{\mathbb{Z}}$$

obtained by taking the quotient of the vector bundle $\mathcal{H}^{0,1} \rightarrow Y$ by the $H_{\mathbb{Z}}$ -action together with the natural morphism $\pi: J_H \rightarrow Y$.

- (iv) The zero section of $\mathcal{H}^{0,1} \rightarrow Y$ induces the global section $\sigma_0: Y \rightarrow J_H$. The sheaf of local sections of $\pi: J_H \rightarrow Y$ is a sheaf of abelian groups with addition of local sections and neutral element σ_0 . It is denoted by $\mathcal{J}_H = \mathcal{H}^{0,1}/H_{\mathbb{Z}}$, where we identify the vector bundle $\mathcal{H}^{0,1}$ with its sheaf of local sections.
- (v) Let $\eta \in H^1(Y, \mathcal{J}_H)$ be a cohomology class. The \mathcal{J}_H -torsor associated to η is denoted by

$$\pi^\eta: J_H^\eta \rightarrow Y.$$

- (vi) We denote by $\exp: H^1(Y, \mathcal{H}^{0,1}) \rightarrow H^1(Y, \mathcal{J}_H)$ and $c: H^1(Y, \mathcal{J}_H) \rightarrow H^2(Y, H_{\mathbb{Z}})$ the maps induced by the short exact sequence $0 \rightarrow H_{\mathbb{Z}} \rightarrow \mathcal{H}^{0,1} \rightarrow \mathcal{J}_H \rightarrow 0$, and call them the exponential, and Chern class map, respectively.

Corollary 2.4. *Let $f: X \rightarrow Y$ be a smooth elliptic fibration. Suppose that f is isotrivial with fiber $E \cong \mathbb{C}/\Lambda$ and that its monodromy is trivial. Then $J_H = Y \times E$ and $\mathcal{J}_H \cong \mathcal{O}_Y/\Lambda$.*

Definition 2.5. In the situation of Theorem 2.4, we denote the \mathcal{J}_H torsor associated to $\eta \in H^1(Y, \mathcal{J}_Y)$ by $\pi^\eta: (Y \times E)^\eta \rightarrow Y$.

Proposition 2.6. *[CHL19, Thm. 3.20, Prop. 3.23][Nak02b, Prop. 1.3.3] Let Y be a compact connected Kähler manifold, let $E = \mathbb{C}/\Lambda$ be an elliptic curve, and let $\eta \in H^1(Y, \mathcal{O}_Y/\Lambda)$.*

- (i) *The complex manifold $(Y \times E)^\eta$ is Kähler if and only if $c(\eta) \in H^2(Y, \Lambda)$ is torsion.*
- (ii) *Suppose that $(Y \times E)^\eta$ is Kähler. Let $\pi: (Y \times E)^\eta \rightarrow Y$ be the projection. Then for every $y \in Y$ the restriction map*

$$H^0((Y \times E)^\eta, \Omega_{(Y \times E)^\eta}^1) \longrightarrow H^0(\pi^{-1}(y), \Omega_{\pi^{-1}(y)}^1) \cong H^0(E, \Omega_E^1)$$

fits into a short exact sequence

$$0 \longrightarrow H^0(Y, \Omega_Y^1) \longrightarrow H^0((Y \times E)^\eta, \Omega_{(Y \times E)^\eta}^1) \longrightarrow H^0(E, \Omega_E^1) \longrightarrow 0.$$

In particular, any holomorphic 1-form $\omega \in H^0((Y \times E)^\eta, \Omega_{(Y \times E)^\eta}^1)$ that does not come from Y has no zeros.

- (iii) *The following is equivalent:*

- (a) *The smooth elliptic fibration $(Y \times E)^\eta \rightarrow Y$ admits a multisection.*
- (b) *There is a finite étale cover $Y' \rightarrow Y$ such that $(Y \times E)^\eta \times_Y Y' \cong Y' \times E$.*
- (c) *The cohomology class $\eta \in H^1(Y, \mathcal{O}_Y/\Lambda)$ is torsion.*

Remark 2.7. Theorem 2.6 allows us to construct a compact Kähler manifold X of Kodaira dimension $\kappa(X) = \dim X - 1$ that admits a holomorphic 1-form without zeros such that no finite étale cover of X splits into a product where one factor is an elliptic curve. This shows that Theorem 1.1 fails in the

Kähler case. Indeed, let C be a smooth projective curve of genus $g \geq 2$, and let $E = \mathbb{C}/\Lambda$ be an elliptic curve. Consider the exact sequence

$$H^1(C, \Lambda) \longrightarrow H^1(C, \mathcal{O}_C) \xrightarrow{\exp} H^1(C, \mathcal{O}_C/\Lambda) \xrightarrow{c} H^2(C, \Lambda),$$

and choose a class $\alpha \in H^1(C, \mathcal{O}_C)$ that is not in the image of

$$H^1(C, \Lambda) \otimes \mathbb{Q} \longrightarrow H^1(C, \mathcal{O}_C).$$

Since $H^1(C, \mathcal{O}_C)$ is a non-trivial complex vector space, and $H^1(C, \Lambda)$ is a finitely generated abelian group, such an α exists. Define $\eta := \exp(\alpha)$. Note that the choice of α implies that η is non-torsion. The exactness of the above sequence implies that the Chern class $c(\eta)$ vanishes. Thus, by Theorem 2.6 item (i) and item (ii), the space $X := (C \times E)^\eta$ is a compact Kähler manifold that admits a holomorphic 1-form without zeros. Since C is of genus $g \geq 2$, $\kappa(X) = \dim C = \dim X - 1$. We claim that no finite étale cover of X splits into a product where one factor is an elliptic curve. To show this, suppose that a finite étale cover $X' \rightarrow X$ splits into a product $X' \cong D \times F$, where F is an elliptic curve. Since $g(C) \geq 2$, the composition

$$\{x\} \times F \hookrightarrow D \times F \longrightarrow X \longrightarrow C$$

must be constant for every $x \in D$, and hence there exists a morphism $D \rightarrow C$ such that the composition $D \times F \rightarrow X \rightarrow C$ factors through $D \times F \rightarrow D \rightarrow C$. Since $D \times F \rightarrow X \rightarrow C$ is surjective, the morphism $D \rightarrow C$ cannot be constant. Therefore, the image $S \subset X$ of $D \times \{0\} \rightarrow X$ defines a multisection of $X \rightarrow C$. Thus, by Theorem 2.6 item (a), the cohomology class η is torsion, which is a contradiction.

2.2.2. Equivariant Weierstraß models.

Definition 2.8. [Nak87, Def. 1.1, 2], [CHL19, 3E] Let Y be a connected complex manifold and let \mathcal{L} be a line bundle on Y . Choose global sections $\alpha \in H^0(Y, \mathcal{L}^{-4})$ and $\beta \in H^0(Y, \mathcal{L}^{-6})$ such that $4\alpha^3 + 27\beta^2 \neq 0$ in $H^0(Y, \mathcal{L}^{-12})$.

- (i) The Weierstraß model $p: W(\mathcal{L}, \alpha, \beta) \rightarrow Y$ is then defined as

$$W(\mathcal{L}, \alpha, \beta) := \{y^2z - (x^3 + \alpha xz^2 + \beta z^3) = 0\} \subset \mathbb{P} := \mathbb{P}(\mathcal{O}_Y \oplus \mathcal{L}^2 \oplus \mathcal{L}^3) \longrightarrow Y,$$

where $x \in H^0(\mathbb{P}, \mathcal{O}_{\mathbb{P}}(1) \otimes p^*\mathcal{L}^{-2})$, $y \in H^0(\mathbb{P}, \mathcal{O}_{\mathbb{P}}(1) \otimes p^*\mathcal{L}^{-3})$, $z \in H^0(\mathbb{P}, \mathcal{O}_{\mathbb{P}}(1))$ are given by the direct sum inclusions $\mathcal{L}^2 \hookrightarrow \mathcal{O}_Y \oplus \mathcal{L}^2 \oplus \mathcal{L}^3$, $\mathcal{L}^3 \hookrightarrow \mathcal{O}_Y \oplus \mathcal{L}^2 \oplus \mathcal{L}^3$, $\mathcal{O}_Y \hookrightarrow \mathcal{O}_Y \oplus \mathcal{L}^2 \oplus \mathcal{L}^3$.

- (ii) The subspace $W(\mathcal{L}, \alpha, \beta)^\# \subset W(\mathcal{L}, \alpha, \beta)$ is defined as the locus where p is smooth.
- (iii) The canonical section of $W(\mathcal{L}, \alpha, \beta) \rightarrow Y$ is defined as $\Sigma := \{x = z = 0\}$. The sheaf of local meromorphic sections of $p: W(\mathcal{L}, \alpha, \beta) \rightarrow Y$ is denoted by $\mathcal{W}(\mathcal{L}, \alpha, \beta)^{mer}$ and the sheaf of local sections with image in $W(\mathcal{L}, \alpha, \beta)^\#$ is denoted by $\mathcal{W}(\mathcal{L}, \alpha, \beta)^\#$. Both sheaves are sheaves of abelian groups with addition of local (meromorphic) sections with respect to Σ .
- (iv) The Weierstraß model $p: W(\mathcal{L}, \alpha, \beta) \rightarrow Y$ is called minimal if there is no prime divisor Δ on Y such that $\text{div}(\alpha) - 4\Delta$ and $\text{div}(\beta) - 6\Delta$ are both effective divisors on Y .
- (v) Let G be a finite group. The Weierstraß model $p: W(\mathcal{L}, \alpha, \beta) \rightarrow Y$ is called G -equivariant if G acts on $W(\mathcal{L}, \alpha, \beta)$ and on Y such that p is G -equivariant and the canonical section Σ is fixed by the G -action.

In the case where \mathcal{L} , α , and β are evident from the context, we omit these symbols in the notation.

Remark 2.9. The morphism $p: W(\mathcal{L}, \alpha, \beta) \rightarrow Y$ is projective, flat, and surjective. Its discriminant locus is given by $\text{div}(4\alpha^3 + 27\beta^2)$, and the total space $W(\mathcal{L}, \alpha, \beta)$ is normal, see [Nak87, 1.2]. In the case that the discriminant locus is a normal crossing divisor, the Weierstraß model $p: W(\mathcal{L}, \alpha, \beta) \rightarrow Y$ is minimal

if and only if $W(\mathcal{L}, \alpha, \beta)$ has at worst rational singularities by [Nak87, Cor. 2.4]. Moreover, there is an isomorphism $R^1 p_* \mathcal{O}_{W(\mathcal{L}, \alpha, \beta)} \cong \mathcal{L}$ [Nak87, 1.2]. Suppose that a finite group G acts on $W(\mathcal{L}, \alpha, \beta)$ and on Y such that the Weierstraß model $p: W(\mathcal{L}, \alpha, \beta) \rightarrow Y$ is G -equivariant. Then we can apply the flat base change theorem [Har77, III. Prop. 9.3] to $R^1 p_* \mathcal{O}_{W(\mathcal{L}, \alpha, \beta)} \cong \mathcal{L}$ to get a G^{op} -action on \mathcal{L} .

Theorem 2.10. [Nak87, Thm. 2.5] *Let $f: X \rightarrow Y$ be an elliptic fibration over a connected complex manifold Y . Suppose that f admits a meromorphic section. Then there exists a unique minimal Weierstraß model $p: W(\mathcal{L}, \alpha, \beta) \rightarrow Y$ that is bimeromorphic to f .*

Definition 2.11. [CHL19, 3E] Let Y be a connected complex manifold and let G be a finite group acting on Y . Let $\Delta \subset Y$ be a G -equivariant normal crossing divisor and let H be a G -equivariant variation of Hodge structures on $j: Y^* := Y \setminus \Delta \hookrightarrow Y$ associated to some elliptic fiber bundle over Y^* . We define

$$\mathcal{E}_G(Y, \Delta, H)$$

to be the set of equivalence classes of G -equivariant elliptic fibrations $f: X \rightarrow Y$ that satisfy the following properties:

- (i) $f: X \rightarrow Y$ has local meromorphic sections over every point of Y ,
- (ii) the elliptic fibration $f^{-1}(Y^*) \rightarrow Y^*$ is bimeromorphic to an elliptic fiber bundle over Y^* , and
- (iii) this elliptic fibration induces H as its G -equivariant variation of Hodge structures.

Two such elliptic fibrations are equivalent if they are G -equivariantly bimeromorphic.

Lemma 2.12. [Nak87, Cor. 2.7] *Let Y be a connected complex manifold and let G be a finite group acting on Y . Let $\Delta \subset Y$ be a G -equivariant normal crossing divisor and let H be a G -equivariant variation of Hodge structures on $j: Y^* := Y \setminus \Delta \hookrightarrow Y$ associated to some elliptic fiber bundle over Y^* . Then there exists a unique minimal Weierstraß model $p: W(\mathcal{L}, \alpha, \beta) \rightarrow Y$ that is smooth over Y^* and induces H as its G -equivariant variation of Hodge structures.*

We can construct elements of $\mathcal{E}_G(Y, \Delta, H)$ as follows: Let G be a finite group and let $p: W(\mathcal{L}, \alpha, \beta) \rightarrow Y$ be a minimal G -equivariant Weierstraß model that is smooth over Y^* and has H as its associated variation of Hodge structures, which exists by Theorem 2.12. The subspace $W(\mathcal{L}, \alpha, \beta)^\# \subset W(\mathcal{L}, \alpha, \beta)$ defines a complex analytic group variety over Y with zero section $\Sigma \subset W(\mathcal{L}, \alpha, \beta)^\#$, where the group structure is given by the addition of local sections with respect to Σ . The action of $W(\mathcal{L}, \alpha, \beta)^\#$ on $W(\mathcal{L}, \alpha, \beta)^\#$ can be extended to an action of $W(\mathcal{L}, \alpha, \beta)^\#$ on $W(\mathcal{L}, \alpha, \beta)$ [Nak02a, 5.1.1]. By gluing local patches of $p: W(\mathcal{L}, \alpha, \beta) \rightarrow Y$ along a cocycle of local sections representing the cohomology class $\eta_G \in H_G^1(Y, \mathcal{W}(\mathcal{L}, \alpha, \beta)^\#)$, one can thus construct the twisted G -equivariant Weierstraß model $p^{\eta_G}: W(\mathcal{L}, \alpha, \beta)^{\eta_G} \rightarrow Y$ [CHL19, 3E]. This construction defines a map

$$(1) \quad H_G^1(Y, \mathcal{W}(\mathcal{L}, \alpha, \beta)^\#) \longrightarrow \mathcal{E}_G(Y, \Delta, H).$$

Proposition 2.13. [Nak87, Prop. 2.10][Nak02a, Prop. 5.5.1][CHL19, 3E] *Let $p: W(\mathcal{L}, \alpha, \beta) \rightarrow Y$ be a minimal G -equivariant Weierstraß model, such that the discriminant locus $\Delta := \text{div}(4\alpha^3 + 27\beta^2)$ is a normal crossing divisor. Denote by $j: Y^* \subset Y$ the inclusion and let $H := R^1 p_* \mathbb{Z}|_{Y^*}$ be the associated variation of Hodge structures.*

(i) *There is a G -equivariant short exact sequence*

$$(2) \quad 0 \longrightarrow j_* H_{\mathbb{Z}} \longrightarrow \mathcal{L} \longrightarrow \mathcal{W}^\# \longrightarrow 0.$$

(ii) There is an injective map

$$(3) \quad \mathcal{E}_G(Y, \Delta, H) \hookrightarrow H_G^1(Y, \mathcal{W}^{mer})$$

such that the composition

$$(4) \quad H_G^1(Y, \mathcal{W}(\mathcal{L}, \alpha, \beta)^\#) \longrightarrow \mathcal{E}_G(Y, \Delta, H) \hookrightarrow H_G^1(Y, \mathcal{W}^{mer})$$

equals the map in cohomology induced by the inclusion $\mathcal{W}^\# \subset \mathcal{W}^{mer}$.

Proposition 2.14. [CHL19, Thm. 3.20, Prop. 3.23] Let $p: W \rightarrow Y$ be a minimal Weierstraß model over a compact Kähler manifold such that the discriminant locus Δ is a normal crossing divisor. Denote by $j: Y^* := Y \setminus \Delta \hookrightarrow Y$ the inclusion and let $H := R^1 p_* \mathbb{Z}|_{Y^*}$ be the associated variation of Hodge structures. Let $\eta \in H^1(Y, \mathcal{W}^\#)$ be a cohomology class. Then the following are equivalent:

- (i) The total space $W(\mathcal{L}, \alpha, \beta)^\eta$ is bimeromorphic to a compact Kähler manifold.
- (ii) The boundary map associated to the short exact sequence $0 \rightarrow j_* H_{\mathbb{Z}} \rightarrow \mathcal{L} \rightarrow \mathcal{W}^\# \rightarrow 0$ sends η to a torsion class in $H^2(Y, j_* H_{\mathbb{Z}})$.

2.3. Tautological models and algebraic approximations. There are classes in $\mathcal{E}_G(Y, \Delta, H)$ that cannot be represented by a twisted Weierstraß model, the reason being that the map $H^1(Y, \mathcal{W}^\#) \rightarrow H^1(Y, \mathcal{W}^{mer})$ may not be surjective. Lin introduced the notion of a tautological model [Lin20, 3.4], which are finite covers of Weierstraß models and showed that every class in $\mathcal{E}_G(Y, \Delta, H)$ that comes from an elliptic fibration $f: X \rightarrow Y$, where X is bimeromorphic to a compact Kähler manifold, can be represented by a tautological model. Moreover, tautological models fit into a family.

Proposition 2.15. [CHL19, Lem. 3.19, Lem. 3.24][Lin20, 3.4] Let $f: X \rightarrow Y$ be a G -equivariant elliptic fibration from a compact complex analytic variety X bimeromorphic to a compact Kähler manifold to a connected complex manifold Y . Suppose that the discriminant divisor of f is a normal crossing divisor and that f admits local meromorphic sections over every point of Y . Denote by $H := R^1 f_* \mathbb{Z}|_{Y^*}$ the variation of Hodge structures induced by f and denote by $W(\mathcal{L}, \alpha, \beta) \rightarrow Y$ the unique G -equivariant minimal Weierstraß model associated to (Y, Δ, H) by Theorem 2.12.

- (i) Let $\eta_G \in H_G^1(Y, \mathcal{W}^{mer})$ be the image of the class of f under the map $\mathcal{E}_G(Y, \Delta, H) \rightarrow H_G^1(Y, \mathcal{W}^{mer})$ from eq. (3) above. Then there is a positive integer m such that $m\eta_G$ can be lifted to a class $\eta'_G \in H_G^1(Y, \mathcal{W}^\#)$.
- (ii) There is a G -equivariant commutative diagram

$$\begin{array}{ccc} T & \xrightarrow{m} & W(\mathcal{L}, \alpha, \beta)^{\eta'_G} \\ & \searrow g & \swarrow p \\ & Y & \end{array},$$

where $g: T \rightarrow Y$ is an elliptic fibration G -equivariantly bimeromorphic to $f: X \rightarrow Y$, and $m: T \rightarrow W(\mathcal{L}, \alpha, \beta)^{\eta'_G}$ is finite of degree m .

- (iii) The total space of T is normal and the discriminant locus of g equals the discriminant locus of p .
- (iv) Over the smooth locus of g and p , m is the multiplication by m -map.

Definition 2.16. In the situation and notation of Theorem 2.15, we call $g: T \rightarrow Y$ the tautological model associated to $f: X \rightarrow Y$, resp. to $\eta_G \in H_G^1(Y, \mathcal{W}^{mer})$.

Proposition 2.17. [Lin20, Prop.-Def. 3.17] *In the situation and notation of Theorem 2.15, there exists a family of elliptic fibrations*

$$\Pi: \mathcal{T} \xrightarrow{q} Y \times V \longrightarrow V := H^1(Y, \mathcal{L})$$

satisfying the following properties.

- (i) *For every $v \in V$ the fiber $\Pi^{-1}(v)$ is isomorphic to the tautological model that represents the class $\eta + \exp'(v) \in H^1(Y, \mathcal{W}^{mer})$, where η is the image of η_G via the natural map $H_G^1(Y, \mathcal{W}^{mer}) \rightarrow H^1(Y, \mathcal{W}^{mer})$, and \exp' denotes the composition $H^1(Y, \mathcal{L}) \rightarrow H^1(Y, \mathcal{W}^\#) \rightarrow H^1(Y, \mathcal{W}^{mer})$.*
- (ii) *Denote by $V^G \subset V$ the G -invariant subspace. Then there is a G -action on $\mathcal{T}^G := q^{-1}(Y \times V^G)$ such that the subfamily*

$$(5) \quad \Pi^G: \mathcal{T}^G \xrightarrow{q^G} Y \times V^G \longrightarrow V^G,$$

where $q^G := q|_{\mathcal{T}^G}$ is G -equivariantly locally trivial over Y , and the G -equivariant elliptic fibration parametrized by $v \in V$ is isomorphic to the tautological model that represents the class $\eta_G + \exp'_G(v) \in H_G^1(Y, \mathcal{W}^{mer})$, where \exp'_G denotes the composition $H_G^1(Y, \mathcal{L}) \rightarrow H_G^1(Y, \mathcal{W}^\#) \rightarrow H_G^1(Y, \mathcal{W}^{mer})$.

- (iii) *Let $\Pi': \mathcal{W} \rightarrow Y \times V \rightarrow V$ be the tautological family associated to the image $\eta' \in H^1(Y, \mathcal{W}^\#)$ of η'_G . Then there exists a map $m: \mathcal{T} \rightarrow \mathcal{W}$ over $Y \times V$ whose restriction to each fiber over V is the multiplication by m -map defined in Theorem 2.15. Over V^G , the map m is G -equivariant.*

$$\begin{array}{ccc} \mathcal{T} & \xrightarrow{m} & \mathcal{W} \\ & \searrow \Pi & \swarrow \Pi' \\ & Y \times V & \end{array}$$

Definition 2.18. In the situation and notation of Theorem 2.17, we call $\Pi^G: \mathcal{T}^G \rightarrow Y \times V^G \rightarrow V^G$ the G -equivariant tautological family associated to $f: X \rightarrow Y$, resp. to $\eta_G \in H_G^1(Y, \mathcal{W}^{mer})$.

Construction 2.19. Let X be a compact Kähler manifold that is bimeromorphic to the total space of an elliptic fibration $f_0: X_0 \rightarrow Y_0$ over a smooth projective variety Y_0 . Denote by $\Delta_0 \subset Y_0$ the discriminant divisor. Choose a log-resolution $(Y_1, \Delta_1) \rightarrow (Y_0, \Delta_0)$, and choose a resolution of singularities $X_1 \rightarrow X_0 \times_{Y_0} Y_1$ that is an isomorphism over the smooth locus of $X_0 \times_{Y_0} Y_1 \rightarrow Y_1$. Then the induced morphism

$$f_1: X_1 \longrightarrow X_0 \times_{Y_0} Y_1 \longrightarrow Y_1$$

is an elliptic fibration, which is smooth outside the normal crossing divisor $\Delta_1 \subset Y_1$. By construction, X_1 is Kähler. Thus, by [Nak02b, Thm. 3.3.3], f_1 is locally projective. Therefore, we can apply [CHL19, Prop. 3.11] to conclude that there exists a finite Galois cover $Y' \rightarrow Y_1$ by some projective manifold Y' with Galois group G such that the elliptic fibration

$$f': X' \longrightarrow Y'$$

from the normalization X' of $X_1 \times_{Y_1} Y'$ to Y' is G -equivariant, has local meromorphic sections over every point in Y' and is smooth outside a normal crossing divisor $\Delta' \subset Y'$. By Theorem 2.12 there exists a unique minimal Weierstraß model $p: W(\mathcal{L}, \alpha, \beta) \rightarrow Y'$ smooth over $Y' \setminus \Delta'$ that induces the same variation of Hodge structures as f' over $Y' \setminus \Delta'$. By Theorem 2.15 there exists a tautological model $g: T \rightarrow Y'$ of f' . Furthermore, by Theorem 2.17 the tautological model $g: T \rightarrow Y'$ fits into the tautological family associated to f'

$$\Pi: \mathcal{T} \longrightarrow Y' \times V \longrightarrow V$$

as the central fiber, i.e. $T \cong \mathcal{T}_0 := \Pi^{-1}(0)$, where $V := H^1(Y', \mathcal{L})$. The restriction of this family to the G -invariant subspace $V^G \subset V$ is G -equivariant by Theorem 2.17. We can therefore take the quotient to obtain the family

$$\Pi^G: \mathcal{T}^G \xrightarrow{q^G} Y' \times V^G \longrightarrow V^G.$$

Theorem 2.20. [Lin20, Proof of Thm. 1.2] *In the notation of Theorem 2.19, there exists a linear subspace $V' \subset V^G$ and a commutative diagram*

$$\begin{array}{ccc} \mathcal{X} & \xrightarrow{\sim} & \mathcal{T}^G/G \\ \pi \searrow & & \swarrow \Pi^G/G \\ & V' & \end{array}$$

such that

- (i) $\pi: \mathcal{X} \rightarrow V'$ is an algebraic approximation of X , i.e. π is a smooth and proper morphism such that $\pi^{-1}(0) \cong X$ and the subset

$$A := \{a \in V' \mid \pi^{-1}(a) \text{ is algebraic}\} \subset V'$$

is dense in the euclidean topology.

- (ii) The family \mathcal{X} is bimeromorphic over V' to \mathcal{T}/G .

2.4. Equivariant elliptic fiber bundles with trivial monodromy.

Proposition 2.21. [Pie25, Prop. 2.22] *Let $f: X \rightarrow Y$ be an elliptic fibration between compact connected complex manifolds, which has local meromorphic sections over every point of Y and is smooth outside a normal crossing divisor $\Delta \subset Y$. Suppose that a finite group G acts on X and Y such that f is G -equivariant. Suppose that the local system $\Lambda := R^1 f_* \mathbb{Z}|_{Y^*}$ of the variation of Hodge structures induced by f is G -equivariantly isomorphic to \mathbb{Z}^2 , where \mathbb{Z}^2 is endowed with the trivial G -action, and that the smooth fibers of f are isomorphic to \mathbb{C}/Λ . Then there is a cohomology class $\eta \in H^1(Y, \mathcal{O}_Y/\Lambda)^G$ and a group homomorphism $\sigma: G \rightarrow E$ such that $f: X \rightarrow Y$ is G -equivariantly bimeromorphic to $\pi^\eta: (Y \times E)^\eta \rightarrow Y$, where G acts diagonally: on Y by the given action and on E by translations with $\sigma(g)$. Moreover, if X and Y are Kähler, then $(Y \times E)^\eta$ is also Kähler.*

3. DEFORMATIONS

Let X be a compact Kähler manifold. A deformation of X is a smooth proper morphism $\pi: \mathcal{X} \rightarrow B$ of complex manifolds, where B has a distinguished base point $0 \in B$ such that the fiber $\mathcal{X}_0 := \pi^{-1}(0)$ is isomorphic to X .

Proposition 3.1. *Let X be a compact connected Kähler manifold that admits a holomorphic 1-form $\omega \in H^0(X, \Omega_X^1)$ without zeros, and let $\pi: \mathcal{X} \rightarrow B$ be a deformation of $\mathcal{X}_0 \cong X$. Then, after shrinking B , there exists a holomorphic 1-form $\tilde{\omega} \in H^0(\mathcal{X}, \Omega_{\mathcal{X}/B}^1)$ such that the isomorphism $\mathcal{X}_0 \cong X$ sends $\tilde{\omega}|_{\mathcal{X}_0}$ to ω . In particular, after shrinking B further if necessary, all fibers \mathcal{X}_b admit a holomorphic 1-form without zeros.*

Proof. We can assume without loss of generality that B is a polydisc centered around $0 \in \mathbb{C}^n$. Let I be the ideal sheaf of $0 \in B$. Tensoring the short exact sequence $0 \rightarrow I \rightarrow \mathcal{O}_B \rightarrow \mathbb{C}_0 \rightarrow 0$ with $\pi_* \Omega_{\mathcal{X}/B}^1$, yields the exact sequence

$$0 \longrightarrow \pi_* \Omega_{\mathcal{X}/B}^1 \otimes I \longrightarrow \pi_* \Omega_{\mathcal{X}/B}^1 \longrightarrow \pi_* \Omega_{\mathcal{X}/B}^1 \otimes \mathbb{C}_0 \longrightarrow 0.$$

As B is a polydisc, $H^1(B, \Omega_{\mathcal{X}/B}^1)$ vanishes. Thus, the induced map

$$H^0(B, \Omega_{\mathcal{X}/B}^1) \longrightarrow \pi_* \Omega_{\mathcal{X}/B}^1 \otimes \mathbb{C}_0$$

is surjective. By the semicontinuity theorem [Har77, Thm. 12.8], the functions

$$\begin{aligned} \varphi: B &\longrightarrow \mathbb{Z} \\ b &\longmapsto \dim H^0(\mathcal{X}_b, \Omega_{\mathcal{X}_b}) \end{aligned}$$

$$\begin{aligned} \psi: B &\longrightarrow \mathbb{Z} \\ b &\longmapsto \dim H^1(\mathcal{X}_b, \mathcal{O}_{\mathcal{X}_b}) \end{aligned}$$

are upper semi-continuous. By the Hodge decomposition, the sum $\varphi + \psi$ is equal to the function $b \mapsto \dim H^1(\mathcal{X}_b, \mathbb{C})$, which is constant. Therefore, also φ and ψ must be constant. Using that φ is constant, Grauert's theorem on proper base change [Har77, Cor. 12.9] implies that there natural map

$$\pi_* \Omega_{\mathcal{X}/B}^1 \otimes \mathbb{C}_0 \xrightarrow{\sim} H^0(\mathcal{X}_0, \Omega_{\mathcal{X}_0}^1)$$

is an isomorphism. Therefore, there is a holomorphic 1-form $\tilde{\omega} \in H^0(\mathcal{X}, \Omega_{\mathcal{X}/B}^1)$ such that the composition

$$H^0(B, \Omega_{\mathcal{X}/B}^1) \longrightarrow \pi_* \Omega_{\mathcal{X}/B}^1 \otimes \mathbb{C}_0 \cong H^0(\mathcal{X}_0, \Omega_{\mathcal{X}_0}^1)$$

sends $\tilde{\omega}$ to ω . Let $Z \subset \mathcal{X}$ be the zero locus of $\tilde{\omega}$. This is a closed subvariety, which does not contain \mathcal{X}_0 by assumption. As π is proper, $\pi(Z) \subset B$ is a closed subset that does not contain $0 \in B$. In particular, for all $b \in B \setminus \pi(Z)$, the fiber \mathcal{X}_b admits a holomorphic 1-form without zeros. Therefore, shrinking B to $B \setminus \pi(Z)$ yields the desired result. \square

4. PROOF OF THE MAIN RESULT

Theorem 4.1. *Let X be a compact Kähler manifold of Kodaira dimension $\kappa(X) = \dim X - 1$ that admits a holomorphic 1-form without zeros. Then there is a smooth projective variety Y of general type and of dimension $\dim Y = \dim X - 1$, an elliptic curve $E = \mathbb{C}/\Lambda$, a cohomology class $\eta \in H^1(Y, \mathcal{O}_Y/\Lambda)$ and a finite étale cover $X' \rightarrow X$ such that X' is bimeromorphic to $(Y \times E)^\eta$.*

Proof. First, we claim that X is bimeromorphic to the total space of an elliptic fibration over a projective base. Note that X has algebraic dimension $a(X)$ in the range $\dim X - 1 \leq a(X) \leq \dim X$. If $a(X) = \dim X - 1$, using algebraic reduction shows that X is bimeromorphic to the total space of an elliptic fibration $f_0: X_0 \rightarrow Y_0$ over a projective variety Y_0 , [Uen75, Thm. 2.4]. If $a(X) = \dim X$, X is projective by Moishezon's theorem. Thus, X admits a good minimal model X_0 by [Lai11, Thm. 4.4], i.e. X_0 is a projective variety, which is \mathbb{Q} -factorial, has terminal singularities, the canonical divisor K_{X_0} is nef and a positive multiple nK_{X_0} of K_{X_0} is globally generated. Consider the Iitaka fibration $f_0: X_0 \rightarrow Y_0$ defined as the Stein factorization of $|nK_{X_0}|: X_0 \rightarrow \mathbb{P}^N$, where $N := \dim H^0(X_0, nK_{X_0}) - 1$:

$$\begin{array}{ccc} X_0 & \xrightarrow{|nK_{X_0}|} & \mathbb{P}^N \\ & \searrow f_0 & \nearrow \\ & Y_0 & \end{array}$$

By [Iit71, Thm. 5], f_0 is an elliptic fibration.

Denote by $\Delta_0 \subset Y_0$ the discriminant divisor. Choose a log-resolution $(Y_1, \Delta_1) \rightarrow (Y_0, \Delta_0)$, and choose a resolution of singularities $X_1 \rightarrow X_0 \times_{Y_0} Y_1$ that is an isomorphism over the smooth locus of $X_0 \times_{Y_0} Y_1 \rightarrow Y_1$. Then the induced morphism

$$f_1: X_1 \longrightarrow X_0 \times_{Y_0} Y_1 \longrightarrow Y_1$$

is an elliptic fibration, which is smooth outside the normal crossing divisor $\Delta_1 \subset Y_1$. By construction, X_1 is Kähler. Thus, by [Nak02b, Thm. 3.3.3], f_1 is locally projective. Therefore, we can apply [CHL19, Prop. 3.11] to conclude that there exists a finite Galois cover $Y_2 \rightarrow Y_1$ by some projective manifold Y_2 with Galois group G such that the elliptic fibration

$$f_2: X_2 \longrightarrow Y_2$$

from the normalization X_2 of $X_1 \times_{Y_1} Y_2$ to Y_2 is G -equivariant, has local meromorphic sections over every point in Y_2 and is smooth outside a normal crossing divisor $\Delta_2 \subset Y_2$.

Denote by $H := R^1(f_2)_*\mathbb{Z}|_{Y_2^*}$ the variation of Hodge structures induced by f_2 . By Theorem 2.12 there exists a unique minimal Weierstraß model $p: W(\mathcal{L}, \alpha, \beta) \rightarrow Y_2$ smooth over $Y_2 \setminus \Delta_2$ that induces the same variation of Hodge structures as f_2 over $Y_2 \setminus \Delta_2$. Let $\eta_G \in H_G^1(Y_2, \mathcal{W}^{mer})$ be the image of the class of f_2 under the map $\mathcal{E}_G(Y_2, \Delta_2, H) \rightarrow H_G^1(Y_2, \mathcal{W}^{mer})$ from eq. (3) above. Then, by Theorem 2.15, there is a positive integer m such that $m\eta_G$ can be lifted to a class $\eta'_G \in H_G^1(Y_2, \mathcal{W}^\#)$, and there is a commutative diagram

$$\begin{array}{ccc} T & \xrightarrow{m} & W(\mathcal{L}, \alpha, \beta)^{\eta'_G} \\ & \searrow g & \swarrow p \\ & Y & \end{array},$$

where $g: T \rightarrow Y$ is the tautological model of f_2 , and $m: T \rightarrow W(\mathcal{L}, \alpha, \beta)^{\eta'_G}$ is finite of degree m , which restricts on smooth fibers to the multiplication by m -map.

Furthermore, by Theorem 2.17, the tautological model $g: T \rightarrow Y_2$ fits into the tautological family associated to f_2 :

$$\Pi: \mathcal{T} \longrightarrow Y_2 \times V \longrightarrow V$$

as the central fiber, i.e. $T \cong \mathcal{T}_0 := \Pi^{-1}(0)$, where $V := H^1(Y_2, \mathcal{L})$. The restriction of this family to the G -invariant subspace $V^G \subset V$ is G -equivariant by Theorem 2.17. We can therefore take the quotient to obtain the family

$$\Pi^G: \mathcal{T}^G \xrightarrow{q^G} Y_2 \times V^G \longrightarrow V^G.$$

By Theorem 2.20, there exists linear subspace $V' \subset V^G$ and a commutative diagram

$$\begin{array}{ccc} \mathcal{X} & \overset{\sim}{\dashrightarrow} & \mathcal{T}^G/G \\ & \searrow \pi & \swarrow \Pi^G/G \\ & V' & \end{array}$$

such that the family \mathcal{X} is bimeromorphic over V' to \mathcal{T}^G/G , the central fiber $\mathcal{X}_0 := \pi^{-1}(0)$ is isomorphic to X , and such that the subset

$$A := \{a \in V' | \pi^{-1}(a) \text{ is algebraic}\} \subset V'$$

is dense in the euclidean topology.

By Theorem 3.1, there is a small open neighborhood $U \subset V'$ of $0 \in V'$ such that the fibers \mathcal{X}_u for $u \in U$ admit a holomorphic 1-form without zeros. As $A \subset V'$ is dense, we can fix a point $a \in A \cap U$.

By construction, \mathcal{X}_a is birational to the elliptic fibration $\Pi^{-1}(a)/G \rightarrow Y_1$. Thus, \mathcal{X}_a has Kodaira codimension 1. Let X_a^{\min} be a good minimal model of \mathcal{X}_a , which exists by [Lai11, Thm. 4.4]. By [Hao23, Lem. 3.1], the general fiber of the Iitaka fibration $f_a: X_a^{\min} \rightarrow Z$ is isomorphic to a fixed elliptic curve. As the Iitaka fibration is unique up to birational equivalence [Iit71, Thm. 5], $f_a: X_a^{\min} \rightarrow Z$ is birational to $\Pi^{-1}(a)/G \rightarrow Y_1$. Therefore, the general fiber of $\Pi^{-1}(a)/G \rightarrow Y_1$ is isomorphic to a fixed elliptic curve. Thus, the same holds true for $\Pi^{-1}(a) \rightarrow Y_2$, and consequently also for $\Pi'^{-1}(a) \rightarrow Y_2$, where Π' is the map in the diagram from Theorem 2.17 item (iii):

$$\begin{array}{ccc} \mathcal{T} & \xrightarrow{m} & \mathcal{W} \\ & \searrow \Pi \quad \swarrow \Pi' & \\ & Y_2 \times V. & \end{array}$$

As all elliptic fibrations $\Pi'^{-1}(v) \rightarrow Y_2$ are locally over Y_2 isomorphic to $W(\mathcal{L}, \alpha, \beta) \rightarrow Y_2$, we conclude that the general fiber of $W(\mathcal{L}, \alpha, \beta) \rightarrow Y_2$ must be isomorphic to a fixed elliptic curve $E = \mathbb{C}/\Lambda$. Write $E = \{y^2z - (x^3 + axz^2 + bz^3) = 0\} \subset \mathbb{P}^2$. Then $W(\mathcal{O}_{Y_2}, a, b) \cong Y_2 \times E \rightarrow Y_2$ is a minimal Weierstraß model that induces the variation of Hodge structures H . As minimal Weierstraß models are unique, it follows that $W(\mathcal{L}, \alpha, \beta) \cong W(\mathcal{O}_{Y_2}, a, b)$. In particular, $\mathcal{L} \cong \mathcal{O}_{Y_2}$. By [Nak02b, Lem. 1.3.5], every meromorphic section of $Y_2 \times E \rightarrow Y_2$ is holomorphic. Thus, the sheaves $\mathcal{W}^\#$ and \mathcal{W}^{mer} agree, and, by Theorem 2.4, they are isomorphic to $\mathcal{O}_{Y_2}/\Lambda$. In particular, the composition in eq. (4) is of the form

$$\text{id}: H_G^1(Y_2, \mathcal{O}_{Y_2}/\Lambda) \longrightarrow \mathcal{E}_G(Y_2, \Delta_2, H) \hookrightarrow H_G^1(Y_2, \mathcal{O}_{Y_2}/\Lambda).$$

Therefore, the class η_G lies in $H_G^1(Y_2, \mathcal{O}_{Y_2}/\Lambda)$, and thus $f_2: X_2 \rightarrow Y_2$ is G -equivariantly bimeromorphic to $(Y_2 \times E)^{\eta_G} \rightarrow Y_2$.

By Remark Theorem 2.9, we obtain a G -action on \mathcal{O}_{Y_2} . This action is determined by a group homomorphism $\rho: G \rightarrow \mathbb{C}^*$. Let $\mathcal{M} \in \text{Pic}(Y_1)$ be the quotient of \mathcal{O}_{Y_2} by this action. Note that this is a torsion line bundle. Denote by n the order, and define

$$Y_3 := \text{Spec} \left(\bigoplus_{i=0}^{n-1} \mathcal{M}^i \right) \longrightarrow Y_1.$$

This is a finite étale cover of Y_1 . Moreover, if we denote by G_1 the kernel of ρ , then $Y_3 \cong Y_2/G_1$. By construction, G_1 acts trivially on H . Thus, we can apply Theorem 2.21 to conclude that G_1 acts diagonally on $(Y_2 \times E)^\eta$ via the given action on Y_2 and via translations with $\sigma(g) \in E$ on the fibers, where $\sigma: G_1 \rightarrow E$ is a group homomorphism. Let G_2 be the kernel of σ . Then

$$(Y_2 \times E)^\eta/G_2 \cong (Y_2/G_2 \times E)^\eta.$$

Moreover, the quotient group G_1/G_2 acts fixed point freely on $(Y_2/G_2 \times E)^\eta$. In particular, the quotient map

$$(Y_2/G_2 \times E)^\eta \longrightarrow (Y_2/G_2 \times E)^\eta/(G_1/G_2)$$

is finite étale.

Note that by construction, X is bimeromorphic to X_1 . As $Y_3 \rightarrow Y_1$ is finite étale, also $X_3 := X_1 \times_{Y_1} Y_3 \rightarrow X_1$ is finite étale. As X_1 is smooth, this implies that there exists a finite étale cover $X' \rightarrow X$ such that X' is bimeromorphic to X_3 . By construction, $X_3 \cong (Y_2/G_2 \times E)^\eta/(G_1/G_2)$. As $(Y_2/G_2 \times E)^\eta \longrightarrow (Y_2/G_2 \times E)^\eta/(G_1/G_2)$ is finite étale, there exists a further finite étale cover $X'' \rightarrow X'$ such that X'' is bimeromorphic to $(Y_2/G_2 \times E)^\eta$, which concludes the proof of Theorem 4.1. \square

REFERENCES

- [Chu24] Benjamin Church. *Nowhere vanishing 1-forms on varieties admitting a good minimal model*. 2024. arXiv: [2410.22753](https://arxiv.org/abs/2410.22753) [math.AG]. URL: <https://arxiv.org/abs/2410.22753>.
- [CHL19] Benoît Claudon, Andreas Horing, and Hsueh-Yung Lin. The fundamental group of compact Kahler threefolds. In: *Geom. Topol.* 23.7 (2019), pp. 3233–3271.
- [GK20] Patrick Graf and Tim Kirschner. Finite quotients of three-dimensional complex tori. In: *Ann. Inst. Fourier (Grenoble)* 70 (2020), pp. 881–914.
- [Hao23] Feng Hao. Nowhere Vanishing Holomorphic One-Forms on Varieties of Kodaira Codimension One. In: *Int. Math. Res. Not. IMRN* 2024.6 (2023), pp. 4501–4515.
- [HWZ24] Feng Hao, Zichang Wang, and Lei Zhang. *Good minimal models with nowhere vanishing holomorphic 1-forms*. 2024. arXiv: [2412.12582](https://arxiv.org/abs/2412.12582) [math.AG]. URL: <https://arxiv.org/abs/2412.12582>.
- [Har77] Robin Hartshorne. *Algebraic Geometry*. Grad. Texts in Math. 52. Springer, 1977.
- [Iit71] Shigeru Iitaka. On D -dimensions of algebraic varieties. In: *J. Math. Soc. Japan* 23 (1971).
- [Lai11] Ching-Jui Lai. Varieties fibered by good minimal models. In: *Math. Ann.* 350 (2011), pp. 533–547.
- [Lin20] Hsueh-Yung Lin. Algebraic approximations of compact Kahler manifolds of algebraic codimension 1. In: *Duke Math. J.* 169.14 (2020), pp. 2767–2826.
- [Nak02a] Noboru Nakayama. Global Structure of an Elliptic Fibration. In: *Publ. Res. Inst. Math. Sci.* 38.3 (2002), pp. 451–649.
- [Nak02b] Noboru Nakayama. Local Structure of an Elliptic Fibration. In: *Adv. Stud. Pure Math.* 35 (2002), pp. 185–295.
- [Nak87] Noboru Nakayama. On Weierstrass models. In: *Algebraic Geometry and Commutative Algebra in Honor of Masayoshi Nagata* 2 (1987), pp. 405–431.
- [Pie25] Simon Pietig. *Holomorphic 1-forms without zeros on Kahler threefolds*. 2025. arXiv: [2506.22067](https://arxiv.org/abs/2506.22067) [math.AG]. URL: <https://arxiv.org/abs/2506.22067>.
- [PS17] Mihnea Popa and Christian Schnell. Kodaira dimension and zeros of holomorphic one-forms. In: *Ann. of Math.* 179.3 (2017), pp. 1109–1120.
- [Uen75] Kenji Ueno. *Classification Theory of Algebraic Varieties and Compact Complex Spaces*. Lecture Notes in Math. 439. Springer, 1975.
- [Var89] Jean Varouchas. Kahler spaces and proper open morphisms. In: *Math. Ann.* 283 (1989), pp. 13–52.