

Emergence of dynamical tensor fields in composite models of gravity

Yadikaer Maitiniyazi ^a, Masatoshi Yamada ^b

^a*Center for Theoretical Physics and College of Physics, Jilin University, Changchun, 130012, Jilin, China*

^b*Department of Physics and Astronomy, Kwansei Gakuin University, Sanda, 669-1330, Hyogo, Japan*

Abstract

We study the composite models of gravity and investigate how dynamical tensor fields emerge by utilizing the functional Renormalization Group. In this paper, we consider two models: One is the fermionic theory. Another is the scalar theory. In both cases, we introduce an auxiliary tensor field corresponding to a composite field of the energy-momentum tensor by means of the Hubbard-Stratonovich transformation. We derive the flow equations for the field renormalization factors of the auxiliary tensor field and show the tensor field becomes dynamical in the infrared regime. The resulting kinetic terms are interpreted as gauge-fixed terms.

1. Introduction

The nature of gravity remains one of the biggest mysteries in fundamental physics. While general relativity succeeds remarkably well in matching experimental observations, a corresponding quantum theory of gravity remains elusive. The quantization of the metric field based on perturbation theory within the Einstein-Hilbert action leads to ultraviolet divergences which cannot be eliminated by renormalization. Such an incompatibility with quantum mechanics has been motivating searches for a microscopic description of gravity beyond the framework of quantum field theory.

Although it is clear that on macroscopic scales gravitational interactions are mediated by the metric field, there is a question of whether the metric field is a fundamental field at the microscopic level. Among the various approaches to quantum gravity, composite models offer an intriguing possibility that the graviton is not an elementary particle, but rather a bound state or emergent excitation arising from more fundamental degrees of freedom.

The idea that gravity may be emergent is inspired by parallels with other known composite phenomena in physics. For instance, the pions in quantum chromodynamics and the Nambu–Jona-Lasinio model [1, 2] are low-energy excitations of composite fermions. These analogies have motivated the exploration of models in which the graviton arises from strong dynamics or large- N gauge theories, or collective excitations in analog spacetime systems. Such approaches can be viewed as possible realizations of pregeometry [3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28]. Moreover, some models may also provide a natural explanation for the weakness of gravity compared to other interactions, as gravitational interactions could be suppressed by compositeness or large scales associated with binding.

In this paper we study the emergence of dynamical tensor fields in composite gravity models by utilizing the functional Renormalization group (fRG). We begin by outlining general principles and constraints on graviton compositeness, followed

by a detailed analysis of representative models. We introduce auxiliary tensor fields corresponding to the energy-momentum tensor via the Hubbard-Stratonovich transformation. As a first step toward a consistent composite theory of gravity, we study how interactions with fundamental fields generate kinetic terms for these tensor fields through their interactions with the underlying dynamical fundamental fields. We also compare the induced kinetic terms to those in general relativity.

This paper is organized as follows: In [Section 2](#), we introduce the composite models of gravity and present the setups in this work. In [Section 3](#), we derive the flow equations for the mass parameters and the field renormalization factors of an auxiliary tensor field corresponding to the composite field. We solve these flow equations and analyze the structure of the kinetic terms of the composite field in [Section 4](#). Finally, [Section 5](#) is devoted to summarizing the results.

2. Models

We study two models: a fermionic theory and a bosonic (scalar) theory. In this section, we present their initial action and introduce an auxiliary tensor field via the Hubbard-Stratonovich transformation.

2.1. Fermionic model

The first case of composite models is based on a $U(N_f)$ fermionic theory in Euclidean space in which the metric is $\delta_{\rho\sigma} = (1, 1, 1, 1)$.¹ The starting action is given by

$$S = \int d^4x \left[\frac{1}{2} \bar{\psi} \overleftrightarrow{\partial} \psi - f_2 T_{\mu\nu}^{(f)} T_{\mu\nu}^{(f)} - f_0 (T_{\rho\rho}^{(f)})^2 \right], \quad (1)$$

¹We adopt the definition of Euclidean space as in [29].

where $\psi = (\psi_0, \dots, \psi_{N_f})$ is the N_f -flavor spinor fields and $T_{\mu\nu}^{(f)}$ is the energy-momentum tensor,

$$T_{\mu\nu}^{(f)} = \frac{1}{4} \bar{\psi} \left(\gamma_\mu \overleftrightarrow{\partial}_\nu + \gamma_\nu \overleftrightarrow{\partial}_\mu \right) \psi, \quad (2a)$$

$$T_{\rho\rho}^{(f)} = \delta_{\rho\sigma} T_{\rho\sigma}^{(f)}. \quad (2b)$$

Here, $\bar{\psi} \gamma_\nu \overleftrightarrow{\partial}_\mu \psi = \bar{\psi} \gamma_\nu (\partial_\mu \psi) - (\partial_\mu \bar{\psi}) \gamma_\nu \psi$ and $\not{\partial} = \delta_{\mu\nu} \gamma_\mu \partial_\nu = \gamma_\mu \partial_\mu$ with γ_μ being the Dirac matrices. Note that f_2 and f_0 have mass dimension -4 .

Introducing an auxiliary tensor field $H_{\mu\nu}$ in the path integral formalism, the action (1) can be rewritten as

$$S = \int d^4x \left[\frac{1}{2} \bar{\psi} \not{\partial} \psi + T_{\mu\nu}^{(f)} H_{\mu\nu} - \lambda'_2 H_{\mu\nu} H_{\mu\nu} - \lambda'_0 H^2 \right], \quad (3)$$

where $H = \delta_{\rho\sigma} H_{\rho\sigma} = H_{\rho\rho}$ is the trace mode in the tensor field $H_{\mu\nu}$. Here, the tensor field $H_{\mu\nu}$ is dimensionless, and λ'_2 and λ'_0 have mass dimension $+4$. Note that using the equation of motion for $H_{\mu\nu}$

$$H_{\mu\nu} = \frac{1}{2\lambda'_2} \left(T_{\mu\nu}^{(f)} - \frac{\lambda'_0}{\lambda'_2 + 4\lambda'_0} \delta_{\mu\nu} T_{\rho\rho}^{(f)} \right), \quad (4)$$

the action (3) returns to the original one (1) and it turns out that the relations between couplings are given by $f_2 = -1/(4\lambda'_2)$ and $f_0 = -\lambda'_0/(4\lambda'_2(\lambda'_2 + 4\lambda'_0))$.

2.2. Scalar model

The second case is the massless $O(N)$ scalar field whose action in Euclidean space is given by

$$S = \int d^4x \left[\frac{1}{2} (\partial_\mu \vec{\phi})^2 - g_2 T_{\mu\nu}^{(s)} T_{\mu\nu}^{(s)} - g_0 (T_{\rho\rho}^{(s)})^2 \right], \quad (5)$$

where $\vec{\phi} = (\phi_1, \dots, \phi_N)$ is an N -component scalar field, $(\partial_\mu \vec{\phi})^2 = \partial_\mu \vec{\phi} \cdot \partial_\mu \vec{\phi}$ and the energy-momentum tensor $T_{\mu\nu}^{(s)}$ is given by

$$T_{\mu\nu}^{(s)} = \partial_\mu \vec{\phi} \cdot \partial_\nu \vec{\phi} - \frac{\delta_{\mu\nu}}{2} (\partial_\rho \vec{\phi})^2, \quad (6a)$$

$$T_{\rho\rho}^{(s)} = \delta_{\rho\sigma} T_{\rho\sigma}^{(s)}. \quad (6b)$$

We introduce an auxiliary tensor field $C_{\mu\nu}$ and rewrite the action as

$$S = \int d^4x \left[\frac{1}{2} \partial_\mu \vec{\phi} \cdot \partial_\mu \vec{\phi} + T_{\mu\nu}^{(s)} C_{\mu\nu} - \xi'_2 C_{\mu\nu} C_{\mu\nu} - \xi'_0 C^2 \right], \quad (7)$$

where $C = \delta_{\rho\sigma} C_{\rho\sigma} = C_{\rho\rho}$. In the same manner as in the fermionic case, the use of the equation of motion for $C_{\mu\nu}$,

$$C_{\mu\nu} = \frac{1}{2\xi'_2} \left(T_{\mu\nu}^{(s)} - \frac{\xi'_0}{\xi'_2 + 4\xi'_0} \delta_{\mu\nu} T_{\rho\rho}^{(s)} \right), \quad (8)$$

allows us to reproduce (5) with the relations between couplings as $g_2 = -1/(4\xi'_2)$ and $g_0 = -\xi'_0/(4\xi'_2(\xi'_2 + 4\xi'_0))$. Note here that g_2 and g_0 are mass dimension -4 . Accordingly, ξ'_2 and ξ'_0 are mass dimension $+4$, while $C_{\mu\nu}$ is a dimensionless field.

3. Functional flow equations

In the actions (3) and (7), the tensor fields, $H_{\mu\nu}$ and $C_{\mu\nu}$, are non-dynamical fields, i.e., there are no kinetic terms. Our aim in this work is to investigate the “dynamicalization” of the tensor fields $H_{\mu\nu}$ and $C_{\mu\nu}$ thanks to quantum fluctuations of the fermionic field and scalar field in the models (3) and (7). To this end, we utilize the Wetterich equation [30] which describes the deformation of the one-particle irreducible effective action Γ_k defined at an infrared (IR) cutoff scale k . See also [31, 32]. Its form is given by

$$\partial_t \Gamma_k = \frac{1}{2} \text{STr} \left[\frac{1}{\Gamma_k^{(2)} + \mathcal{R}_k} \partial_t \mathcal{R}_k \right]. \quad (9)$$

Here $t = \ln(k/\Lambda)$ is the dimensionless scale with a reference scale Λ , $\Gamma_k^{(2)}$ is the full two-point function obtained by the second-order functional derivative for Γ_k with respect to the superfield $\Phi = (\psi, \bar{\psi}, \vec{\phi}, H_{\mu\nu}, C_{\mu\nu}, \dots)$ and STr denotes the functional supertrace acting on all spaces (spinor, momentum, etc.) in which the superfield Φ is defined. In the formulation (9), the coarse-graining process is realized by the regulator function \mathcal{R}_k which is written as

$$\mathcal{R}_k(p) = \begin{cases} i p r_k^\psi(|p|/k) & \text{for fermion,} \\ p^2 r_k^\phi(|p|/k) & \text{for scalar.} \end{cases} \quad (10)$$

In this work, for $r_k(|p|/k)$, we employ the Litim type regulator [33] for the fermionic and scalar fields respectively as

$$r_k^\psi(|p|/k) = \left(\frac{k}{|p|} - 1 \right) \theta(1 - |p|^2/k^2), \quad (11)$$

$$r_k^\phi(|p|/k) = \left(\frac{k^2}{|p|^2} - 1 \right) \theta(1 - |p|^2/k^2), \quad (12)$$

where $\theta(x)$ is the step function

$$\theta(x) = \begin{cases} 1, & x \geq 0, \\ 0, & x < 0. \end{cases} \quad (13)$$

In what follows, we present an ansatz for the effective action Γ_k in order to study the generation of dynamics of the tensor fields within the composite models (3) and (7).

3.1. Fermionic model

For the fermionic model (3) we make the following ansatz for the effective action in Euclidean space:

$$\begin{aligned} \Gamma_k^\psi = \int d^4x & \left[\frac{1}{2} \bar{\psi} \not{\partial} \psi + \kappa_\psi T_{\mu\nu}^{(f)} H_{\mu\nu} - \lambda_2 H_{\mu\nu} H_{\mu\nu} - \lambda_0 H^2 \right. \\ & + \frac{Z_{H0}}{4} (\partial_\lambda H_{\mu\nu})^2 + \frac{Z_{H1}}{4} \partial_\lambda H_{\mu\nu} \partial_\nu H_{\mu\lambda} \\ & \left. + \frac{Z_{H2}}{4} \partial_\mu H_{\mu\nu} \partial_\nu H + \frac{Z_{H3}}{4} (\partial_\mu H)^2 \right]. \end{aligned} \quad (14)$$

Here we use the local potential approximation (LPA), fixing the field renormalization factor of the fermionic field to unity at all RG scales. We also neglect the RG running of κ_ψ , since these dimensionless couplings would only exhibit logarithmic running and their loop corrections are subleading, as they rely on the quantum-induced tensor propagator. The initial condition at a certain UV scale Λ for (14) reads $\kappa_\psi|_\Lambda = 1$, $\lambda_2|_\Lambda = \lambda'_2$, $\lambda_0|_\Lambda = \lambda'_0$, and $Z_{Hi}|_\Lambda = 0$ for all i .

Using the Wetterich equation (9), we obtain the flow equations for the couplings in the action (14). At the truncation level of (14), the loop corrections to the two-point function of $H_{\mu\nu}$ arise from the fermionic fluctuations. The flow equation for the two-point function of $H_{\mu\nu}$ is depicted diagrammatically as

$$\partial_t \left(\Gamma_k^{\psi, (2)}(p) \right)_{\mu\nu\rho\sigma} := \frac{\delta^2(\partial_t \Gamma_k^\psi)}{\delta H_{\mu\nu}(p) \delta H_{\rho\sigma}(-p)}$$

$$= - \text{diagram}, \quad (15)$$

where p is the external momentum, the single and double solid lines denote the fermionic fields ψ , $\bar{\psi}$ and tensor field $H_{\mu\nu}$, respectively, and the cross circle is the cutoff insertion $\partial_t \mathcal{R}_k$. The use of the Wetterich equation (9) yields

$$\partial_t \left(\Gamma_k^{\psi, (2)}(p) \right)_{\mu\nu\rho\sigma} = -2\kappa_\psi^2 \int \frac{d^4 q}{(2\pi)^4} \mathcal{T}_{\mu\nu\rho\sigma}^\psi(q, p) f_k^\psi(q, p) \partial_t r_k^\psi(|q|/k). \quad (16)$$

Here r_k^ψ is defined in (11), and f^ψ and $\mathcal{T}_{\mu\nu\rho\sigma}^\psi$ are given respectively by

$$f_k^\psi(q, p) = \frac{1}{q^4(1+r_k^\psi)^2(p+q)^2(1+\tilde{r}_k^\psi)}, \quad (17)$$

$$\mathcal{T}_{\mu\nu\rho\sigma}^\psi(q, p) = \frac{N_f}{16} \text{tr} \left[\not{q} \left((2q+p)_\mu \gamma_\nu + (2q+p)_\nu \gamma_\mu \right) \right. \\ \left. \times (\not{p} + \not{q}) \left((2q+p)_\rho \gamma_\sigma + (2q+p)_\sigma \gamma_\rho \right) \not{q} \right], \quad (18)$$

with $\tilde{r}_k^\psi = r_k^\psi(|q+p|/k)$. Performing the Taylor expansion in terms of p and projecting out the tensor structure in (16), we extract the flow equations for each coupling. The left-hand side of (16) at $p=0$ is $\partial_t(\Gamma_k^{\psi, (2)}(0))_{\mu\nu\rho\sigma} = -\partial_t \lambda_2 (\delta_{\mu\rho} \delta_{\nu\sigma} + \delta_{\mu\sigma} \delta_{\nu\rho}) - 2\partial_t \lambda_0 \delta_{\mu\nu} \delta_{\rho\sigma}$. The flow equations for the mass parameters are obtained from the $O(p^0)$ term and are given by

$$\partial_t \lambda_0 = k^4 \frac{1}{36} \frac{N_f \kappa_\psi^2}{(4\pi)^2}, \quad (19a)$$

$$\partial_t \lambda_2 = -k^4 \frac{1}{36} \frac{N_f \kappa_\psi^2}{(4\pi)^2}. \quad (19b)$$

Next, we show the flow equations for the field renormalization

factors Z_{Hi} . The left-hand side of (16) at order $O(p^2)$ is given by

$$\partial_t \left(\Gamma_k^{\psi, (2)}(p) \right)_{\mu\nu\rho\sigma} \stackrel{O(p^2)}{=} \frac{\partial_t Z_{H0}}{2} p^2 \delta_{\mu\rho} \delta_{\nu\sigma} \\ + \frac{\partial_t Z_{H1}}{4} (p_\sigma p_\nu \delta_{\mu\rho} + p_\rho p_\mu \delta_{\nu\sigma}) \\ + \frac{\partial_t Z_{H2}}{4} (p_\mu p_\nu \delta_{\rho\sigma} + p_\rho p_\sigma \delta_{\mu\nu}) + \frac{\partial_t Z_{H3}}{2} p^2 \delta_{\mu\nu} \delta_{\rho\sigma}. \quad (20)$$

After performing the momentum integral over q in the right-hand side of (16) and identifying the corresponding tensor structures in $\mathcal{T}_{\mu\nu\rho\sigma}^\psi(q, p)$, we obtain the flow equations for the kinetic terms, i.e., the coefficients of the $O(p^2)$ contributions. They take the following form

$$\partial_t Z_{Hi} = k^2 A_i \frac{N_f \kappa_\psi^2}{(4\pi)^2}. \quad (21)$$

In the current setup, we obtain the factors A_i as

$$A_0 = -\frac{7}{48}, \quad A_1 = \frac{1}{4}, \quad A_2 = -\frac{5}{24}, \quad A_3 = \frac{7}{96}. \quad (22)$$

3.2. Scalar model

We assume the effective action for the scalar model (7) to be

$$\Gamma_k^\phi = \int d^4 x \left[\frac{1}{2} \partial_\mu \vec{\phi} \cdot \partial_\mu \vec{\phi} + \kappa_\phi T_{\mu\nu}^{(s)} C_{\mu\nu} - \xi_2 C_{\mu\nu} C_{\mu\nu} - \xi_0 C^2 \right. \\ \left. + \frac{Z_{C0}}{4} (\partial_\lambda C_{\mu\nu})^2 + \frac{Z_{C1}}{4} \partial_\lambda C_{\mu\nu} \partial_\nu C_{\mu\lambda} \right. \\ \left. + \frac{Z_{C2}}{4} \partial_\mu C_{\mu\nu} \partial_\nu C + \frac{Z_{C3}}{4} (\partial_\mu C)^2 \right], \quad (23)$$

where we neglect the running effect of κ_ϕ and the field renormalization of the scalar field as the same reason as the fermionic case.

In the same manner as in the fermionic case, let us derive the flow equations for (23). The flow equation for the two-point function of $C_{\mu\nu}$ is represented diagrammatically as

$$\partial_t \left(\Gamma_k^{\phi, (2)}(p) \right)_{\mu\nu\rho\sigma} := \frac{\delta^2(\partial_t \Gamma_k^\phi)}{\delta C_{\mu\nu}(p) \delta C_{\rho\sigma}(-p)}$$

$$= \frac{1}{2} \text{diagram}, \quad (24)$$

where the dashed and double solid lines denote the scalar field $\vec{\phi}$ and the tensor field $C_{\mu\nu}$, respectively. The flow equation for the scalar two-point function reads

$$\partial_t \left(\Gamma_k^{\phi, (2)}(p) \right)_{\mu\nu\rho\sigma} = \kappa_\phi^2 \int \frac{d^4 q}{(2\pi)^4} \mathcal{T}_{\mu\nu\rho\sigma}^\phi(q, p) f_k^\phi(q, p) \partial_t r_k^\phi(|q|/k), \quad (25)$$

where r_k^ϕ is given in (12) and we have defined

$$f_k^\phi(q, p) = \frac{1}{q^4(1+r_k^\phi)^4(q+p)^2(1+\tilde{r}_k^\phi)^2}, \quad (26)$$

$$\mathcal{T}_{\mu\nu\rho\sigma}^\phi(q, p) = N[q_\mu(q+p)_\nu + q_\nu(q+p)_\mu - \delta_{\mu\nu}q \cdot (q+p)] \\ \times [q_\rho(q+p)_\sigma + q_\sigma(q+p)_\rho - \delta_{\rho\sigma}q \cdot (q+p)]q^2, \quad (27)$$

with $\tilde{r}_k^\phi = r_k^\phi(|p+q|/k)$. From the $O(p^0)$ term, the flow equation governing the two-point function of $C_{\mu\nu}$ yields the corresponding flow of the mass parameters as

$$\partial_t \xi_2 = -k^4 \frac{1}{6} \frac{N \kappa_\phi^2}{(4\pi)^2}, \quad (28a)$$

$$\partial_t \xi_0 = -k^4 \frac{1}{12} \frac{N \kappa_\phi^2}{(4\pi)^2}, \quad (28b)$$

while from the $O(p^2)$ term, we obtain

$$\partial_t Z_{Ci} = k^2 B_i \frac{N \kappa_\phi^2}{(4\pi)^2}. \quad (29)$$

Here the factors are found to be

$$B_0 = -\frac{1}{3}, \quad B_1 = \frac{4}{3}, \quad B_2 = -\frac{2}{3}, \quad B_3 = -\frac{1}{6}. \quad (30)$$

4. Analysis

Solving the flow equations for λ_2 , λ_0 and Z_{Hi} with the initial condition $\lambda_2|_{k=\Lambda} = \lambda'_2$, $\lambda_0|_{k=\Lambda} = \lambda'_0$ and $Z_{Hi}|_{k=\Lambda} = 0$ in the fermionic case, we obtain

$$\lambda_2(k) = \lambda'_2 + \frac{N_f}{4(4\pi)^2} \frac{4}{9} (k^4 - \Lambda^4), \quad (31)$$

$$\lambda_0(k) = \lambda'_0 + \frac{N_f}{4(4\pi)^2} \frac{-4}{9} (k^4 - \Lambda^4), \quad (32)$$

$$Z_{Hi}(k) = \frac{N_f A_i}{2(4\pi)^2} (k^2 - \Lambda^2). \quad (33)$$

In the same manner, in the scalar case we have

$$\xi_2(k) = \xi'_2 + \frac{N}{4(4\pi)^2} \frac{-2}{3} (k^4 - \Lambda^4), \quad (34)$$

$$\xi_0(k) = \xi'_0 + \frac{N}{4(4\pi)^2} \frac{-1}{3} (k^4 - \Lambda^4), \quad (35)$$

$$Z_{Ci}(k) = \frac{N B_i}{2(4\pi)^2} (k^2 - \Lambda^2), \quad (36)$$

where the initial condition was set so as to be $\xi_2|_{k=\Lambda} = \xi'_2$, $\xi_0|_{k=\Lambda} = \xi'_0$ and $Z_{Ci}|_{k=\Lambda} = 0$.

First, we impose the massless condition for the tensor fields in the IR regime ($k = 0$). Thus, the bare mass parameters are chosen such that the loop corrections cancel. In the fermionic case, we give

$$\lambda'_2 = \frac{N_f}{4(4\pi)^2} \frac{4}{9} \Lambda^4, \quad \lambda'_0 = -\frac{N_f}{4(4\pi)^2} \frac{4}{9} \Lambda^4, \quad (37)$$

while in the scalar case, we give

$$\xi'_2 = \frac{N}{4(4\pi)^2} \frac{-2}{3} \Lambda^4, \quad \xi'_0 = \frac{N}{4(4\pi)^2} \frac{-1}{3} \Lambda^4. \quad (38)$$

These subtractions correspond to the additive renormalization of the power divergences.

Next, we consider the kinetic terms of the tensor fields in the IR regime. To this end, we renormalize the fields as $\sqrt{Z_{H0}(0)} H_{\mu\nu} = h_{\mu\nu}$ and $\sqrt{Z_{C0}(0)} C_{\mu\nu} = c_{\mu\nu}$ in the fermionic and scalar cases, respectively. Then, the kinetic terms of $h_{\mu\nu}$ and $c_{\mu\nu}$ are given respectively by

$$\mathcal{L}_h = \frac{1}{4} (\partial_\lambda h_{\mu\nu})^2 + \frac{Z_{H1}(0)}{4Z_{H0}(0)} \partial_\lambda h_{\mu\nu} \partial_\nu h_{\mu\lambda} \\ + \frac{Z_{H2}(0)}{4Z_{H0}(0)} \partial_\mu h_{\mu\nu} \partial_\nu h + \frac{Z_{H3}(0)}{4Z_{H0}(0)} (\partial_\mu h)^2, \quad (39)$$

$$\mathcal{L}_c = \frac{1}{4} (\partial_\lambda c_{\mu\nu})^2 + \frac{Z_{C1}(0)}{4Z_{C0}(0)} \partial_\lambda c_{\mu\nu} \partial_\nu c_{\mu\lambda} \\ + \frac{Z_{C2}(0)}{4Z_{C0}(0)} \partial_\mu c_{\mu\nu} \partial_\nu c + \frac{Z_{C3}(0)}{4Z_{C0}(0)} (\partial_\mu c)^2. \quad (40)$$

Note that this field renormalization does not change the massless condition for the tensor fields, since the operation is multiplicative rather than additive. Within these kinetic terms, the explicit dependence on the cutoff scale Λ , as well as common factors such as the number of the degrees of freedom (N_f and N) and $(4\pi)^{-2}$, cancel out, leaving the coefficients dependent only on the ratios given in (22) and (30). More specifically, in the fermionic case, we have

$$\frac{Z_{H1}(0)}{Z_{H0}(0)} = \frac{A_1}{A_0} = -\frac{12}{7}, \quad (41a)$$

$$\frac{Z_{H2}(0)}{Z_{H0}(0)} = \frac{A_2}{A_0} = \frac{10}{7}, \quad (41b)$$

$$\frac{Z_{H3}(0)}{Z_{H0}(0)} = \frac{A_3}{A_0} = -\frac{1}{2}, \quad (41c)$$

while in the scalar case, we have

$$\frac{Z_{C1}(0)}{Z_{C0}(0)} = \frac{B_1}{B_0} = -4, \quad (42a)$$

$$\frac{Z_{C2}(0)}{Z_{C0}(0)} = \frac{B_2}{B_0} = 2, \quad (42b)$$

$$\frac{Z_{C3}(0)}{Z_{C0}(0)} = \frac{B_3}{B_0} = \frac{1}{2}. \quad (42c)$$

Here, the diffeomorphism invariant kinetic terms of a metric field $g_{\mu\nu}$ around a flat background $\delta_{\mu\nu}$ read

$$\mathcal{L}_{\text{diff}} = \frac{1}{4} (\partial_\lambda g_{\mu\nu})^2 - \frac{1}{2} \partial_\lambda g_{\mu\nu} \partial_\nu g_{\mu\lambda} \\ + \frac{1}{2} \partial^\mu g_{\mu\nu} \partial_\nu g - \frac{1}{4} (\partial_\mu g)^2, \quad (43)$$

where $g = g_{\mu\nu} \delta_{\mu\nu}$. Obviously, both eq. (39) with (41) and eq. (40) with (42) differ from the diffeomorphism invariant kinetic term




Figure 1: Single graviton exchange process in the fermionic (left) and scalar (right) theories.

(43). This fact indicates that eq. (41) and eq. (42) are the gauge-fixed kinetic terms. To see this, let us employ the covariant gauge fixing condition

$$\mathcal{L}_{\text{gf}} = \frac{1}{2\alpha} \left(\partial_\mu g_{\mu\nu} - \frac{1+\beta}{4} \partial_\nu g \right)^2, \quad (44)$$

where α and β are gauge-fixing parameters. Note that the choice $\beta = 1$ corresponds to the so-called de Donder (or harmonic) gauge. Together with the gauge fixing term (44), eq. (43) reads

$$\begin{aligned} \mathcal{L}_{\text{diff}} + \mathcal{L}_{\text{gf}} &= \frac{1}{4} (\partial_\lambda g_{\mu\nu})^2 - \frac{1}{2} \left(1 - \frac{1}{\alpha} \right) \partial_\lambda g_{\mu\nu} \partial_\nu g_{\mu\lambda} \\ &\quad + \frac{1}{2} \left(1 - \frac{1+\beta}{2\alpha} \right) \partial_\mu g_{\mu\nu} \partial_\nu g \\ &\quad - \frac{1}{4} \left(1 - \frac{(1+\beta)^2}{8\alpha} \right) (\partial_\mu g)^2. \end{aligned} \quad (45)$$

However, comparing (45) with (39) and (40) one finds no appropriate choice of α and β matching with the induced kinetic terms in both fermionic and scalar cases.

Another possible gauge-fixing term may be given by

$$\mathcal{L}_{\text{gf}} = \frac{1}{2\zeta_1} \partial_\lambda g_{\mu\nu} \partial_\nu g_{\mu\lambda} - \frac{1}{\zeta_2} \partial_\mu g_{\mu\nu} \partial_\nu g + \frac{1}{2\zeta_3} (\partial_\mu g)^2, \quad (46)$$

with three gauge fixing parameters ζ_1 , ζ_2 and ζ_3 although it is not covariant. For this gauge-fixing term, the kinetic terms (39) and (40) correspond to (43) plus (46) with $\zeta_1 = 7$, $\zeta_2 = 7$ and $\zeta_3 = 4$ for the fermionic case, while $\zeta_1 = -1$, $\zeta_2 = \infty$ and $\zeta_3 = 4/3$ for the scalar case.

We close this section with remarks on the gravitational interaction with a matter field. After renormalizing the tensor fields, the couplings between the metric fluctuations and the energy-momentum tensor in (14) and (23) take the form

$$\mathcal{L}^{Th} = \frac{\kappa_\psi}{\sqrt{Z_{H0}(0)}} T_{\mu\nu}^{(f)} h_{\mu\nu}, \quad (47)$$

$$\mathcal{L}^{Tc} = \frac{\kappa_\phi}{\sqrt{Z_{C0}(0)}} T_{\mu\nu}^{(s)} c_{\mu\nu}. \quad (48)$$

From these expressions, the gravitational coupling extracted from the single-graviton exchange process (see Figure 1) is identified as $G_N = \kappa_\psi^2 / (4\pi Z_{H0}(0))$ for the fermionic case and $G_N = \kappa_\phi^2 / (4\pi Z_{C0}(0))$ for the scalar case.

5. Summary

In this paper, we have taken a first step toward constructing a consistent gravitational theory in which the graviton emerges

as a composite field, using the fRG framework. We examined the fermionic and scalar cases separately and derived the flow equations for the kinetic terms of the auxiliary fields associated with the composite operators. Quantum fluctuations of the matter fields generate finite kinetic terms for these auxiliary fields, which are nondynamical at the tree level. The induced kinetic terms can be interpreted as diffeomorphism-invariant kinetic terms supplemented by gauge-fixing contributions.

To advance our understanding of gravitational dynamics within composite models, it is essential to evaluate higher-order interaction vertices, such as the three-graviton vertex and matter-graviton vertices. The fRG method provides a powerful framework for this task. In particular, dynamical bosonization within the fRG offers a promising approach for obtaining accurate evaluations of these vertices [34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45]. A detailed study of this program lies beyond the scope of the present work and will be reported in future publications.

References

- [1] Y. Nambu and G. Jona-Lasinio, *Dynamical Model of Elementary Particles Based on an Analogy with Superconductivity. I.*, *Phys. Rev.* **122** (1961) 345.
- [2] Y. Nambu and G. Jona-Lasinio, *Dynamical model of elementary particles based on an analogy with superconductivity. II.*, *Phys. Rev.* **124** (1961) 246.
- [3] H. Terazawa, Y. Chikashige, K. Akama and T. Matsuki, *Simple Relation Between the Fine Structure and Gravitational Constants*, *Phys. Rev. D* **15** (1977) 1181.
- [4] H. Terazawa, Y. Chikashige, K. Akama and T. Matsuki, *Model of the Nambu-Jona-Lasinio Type for Gravity*, *J. Phys. Soc. Jap.* **43** (1977) 5.
- [5] K. Akama, Y. Chikashige, T. Matsuki and H. Terazawa, *Gravity and Electromagnetism as Collective Phenomena: A Derivation of Einstein's General Relativity*, *Prog. Theor. Phys.* **60** (1978) 868.
- [6] K. Akama, Y. Chikashige and T. Matsuki, *Unified Model of the Nambu-Jona-Lasinio Type for the Gravitational and Electromagnetic Forces*, *Prog. Theor. Phys.* **59** (1978) 653.
- [7] K. Akama, *An Attempt at Pregeometry: Gravity With Composite Metric*, *Prog. Theor. Phys.* **60** (1978) 1900.
- [8] H. Terazawa and K. Akama, *Subquark Pregeometry With Spontaneously Broken Conformal Invariance*, *Phys. Lett. B* **97** (1980) 81.
- [9] K. Akama, *PREGEOMETRY INCLUDING FUNDAMENTAL GAUGE BOSONS*, *Phys. Rev. D* **24** (1981) 3073.
- [10] K. Akama, *Quantum Mechanical Equivalence of Two Lagrangian Formalisms in 'Scalar Pregeometry'*, *Prog. Theor. Phys.* **61** (1979) 687.

- [11] K. Akama and H. Terazawa, *Pregeometric Origin of the Big Bang*, *Gen. Rel. Grav.* **15** (1983) 201.
- [12] K. Akama and I. Oda, *Topological pregauge pregeometry*, *Phys. Lett. B* **259** (1991) 431.
- [13] H. Terazawa, *Various actions for pregeometry*, *Prog. Theor. Phys.* **86** (1991) 337.
- [14] K. Akama and I. Oda, *Chern-Simons pregeometry*, *Prog. Theor. Phys.* **89** (1993) 215.
- [15] K. Akama and I. Oda, *BRST quantization of pregeometry and topological pregeometry*, *Nucl. Phys. B* **397** (1993) 727.
- [16] K. Akama and I. Oda, *BRST quantization of pregeometry and topological pregeometry*, *Adv. Appl. Clifford Algebras* **4** (1994) 259.
- [17] C. Wetterich, *Gravity from spinors*, *Phys. Rev. D* **70** (2004) 105004 [[hep-th/0307145](#)].
- [18] A. Hebecker and C. Wetterich, *Spinor gravity*, *Phys. Lett. B* **574** (2003) 269 [[hep-th/0307109](#)].
- [19] C. Wetterich, *Spontaneous symmetry breaking origin for the difference between time and space*, *Phys. Rev. Lett.* **94** (2005) 011602 [[hep-th/0405223](#)].
- [20] C. Wetterich, *Pregeometry and euclidean quantum gravity*, *Nucl. Phys. B* **971** (2021) 115526 [[2101.07849](#)].
- [21] C. Wetterich, *Cosmology from pregeometry*, *Phys. Rev. D* **104** (2021) 104040 [[2104.14013](#)].
- [22] C. Wetterich, *Pregeometry and spontaneous time-space asymmetry*, *JHEP* **06** (2022) 069 [[2101.11519](#)].
- [23] R. Floreanini and R. Percacci, *Topological pregeometry*, *Mod. Phys. Lett. A* **5** (1990) 2247.
- [24] S. Matsuzaki, S. Miyawaki, K.-y. Oda and M. Yamada, *Dynamically emergent gravity from hidden local Lorentz symmetry*, *Phys. Lett. B* **813** (2021) 135975 [[2003.07126](#)].
- [25] G. E. Volovik, *Gravity from Symmetry Breaking Phase Transition*, *J. Low Temp. Phys.* **207** (2022) 127 [[2111.07817](#)].
- [26] Y. Maitiniyazi, S. Matsuzaki, K.-y. Oda and M. Yamada, *Irreversible vierbein postulate: Emergence of spacetime from quantum phase transition*, *Phys. Rev. D* **109** (2024) 106018 [[2309.16230](#)].
- [27] Y. Maitiniyazi, S. Matsuzaki, K.-y. Oda and M. Yamada, *Spacetime and Planck mass generation from scale-invariant degenerate gravity*, *Phys. Rev. D* **111** (2025) 046002 [[2411.17238](#)].
- [28] I. Oda, *Emergence of General Relativity from Cosmological Constant Via Ghost Condensation*, **2509.23648**.
- [29] L. Huang, M. Kawaguchi, Y. Maitiniyazi, S. Matsuzaki, A. Tomiya and M. Yamada, *Functional renormalization group study of a four-fermion model with CP violation*, *Phys. Rev. D* **111** (2025) 074015 [[2411.07027](#)].
- [30] C. Wetterich, *Exact evolution equation for the effective potential*, *Phys. Lett. B* **301** (1993) 90 [[1710.05815](#)].
- [31] U. Ellwanger, *FLow equations for N point functions and bound states*, *Z. Phys. C* **62** (1994) 503 [[hep-ph/9308260](#)].
- [32] T. R. Morris, *The Exact renormalization group and approximate solutions*, *Int. J. Mod. Phys. A* **9** (1994) 2411 [[hep-ph/9308265](#)].
- [33] D. F. Litim, *Optimized renormalization group flows*, *Phys. Rev. D* **64** (2001) 105007 [[hep-th/0103195](#)].
- [34] H. Gies and C. Wetterich, *Renormalization flow of bound states*, *Phys. Rev. D* **65** (2002) 065001 [[hep-th/0107221](#)].
- [35] H. Gies and C. Wetterich, *Universality of spontaneous chiral symmetry breaking in gauge theories*, *Phys. Rev. D* **69** (2004) 025001 [[hep-th/0209183](#)].
- [36] J. M. Pawłowski, *Aspects of the functional renormalisation group*, *Annals Phys.* **322** (2007) 2831 [[hep-th/0512261](#)].
- [37] S. Floerchinger and C. Wetterich, *Exact flow equation for composite operators*, *Phys. Lett. B* **680** (2009) 371 [[0905.0915](#)].
- [38] J. Braun, L. Fister, J. M. Pawłowski and F. Rennecke, *From Quarks and Gluons to Hadrons: Chiral Symmetry Breaking in Dynamical QCD*, *Phys. Rev. D* **94** (2016) 034016 [[1412.1045](#)].
- [39] M. Mitter, J. M. Pawłowski and N. Strodthoff, *Chiral symmetry breaking in continuum QCD*, *Phys. Rev. D* **91** (2015) 054035 [[1411.7978](#)].
- [40] A. K. Cyrol, M. Mitter, J. M. Pawłowski and N. Strodthoff, *Nonperturbative finite-temperature Yang-Mills theory*, *Phys. Rev. D* **97** (2018) 054015 [[1708.03482](#)].
- [41] A. K. Cyrol, M. Mitter, J. M. Pawłowski and N. Strodthoff, *Nonperturbative quark, gluon, and meson correlators of unquenched QCD*, *Phys. Rev. D* **97** (2018) 054006 [[1706.06326](#)].
- [42] R. Alkofer, A. Maas, W. A. Mian, M. Mitter, J. París-López, J. M. Pawłowski et al., *Bound state properties from the functional renormalization group*, *Phys. Rev. D* **99** (2019) 054029 [[1810.07955](#)].

- [43] T. Denz, M. Mitter, J. M. Pawłowski, C. Wetterich and M. Yamada, *Partial bosonization for the two-dimensional Hubbard model*, *Phys. Rev. B* **101** (2020) 155115 [[1910.08300](#)].
- [44] W.-j. Fu, J. M. Pawłowski and F. Rennecke, *QCD phase structure at finite temperature and density*, *Phys. Rev. D* **101** (2020) 054032 [[1909.02991](#)].
- [45] F. Goertz, Á. Pastor-Gutiérrez and J. M. Pawłowski, *Gauge-fermion cartography: From confinement and chiral symmetry breaking to conformality*, *Phys. Rev. D* **112** (2025) 034029 [[2412.12254](#)].