

Fractional Homogenization of Parabolic Equations with Long-Range Random Potentials

Atef Lechiheb^a

Toulouse School of Economics, Université Toulouse Capitole, Toulouse, France

, ^aatef.lechiheb@tse-fr.eu

Abstract. This paper establishes a complete homogenization theory for the one-dimensional parabolic equation with long-range correlated random potential:

$$\partial_t u_\varepsilon(t, x) = \frac{1}{2} \partial_{xx} u_\varepsilon(t, x) + \varepsilon^{-\alpha/2} a\left(\frac{x}{\varepsilon}\right) u_\varepsilon(t, x),$$

where the random field a has covariance decaying as $|x|^{-\alpha}$ with $\alpha \in (0, 1)$. Contrary to classical homogenization where rapid decorrelation leads to deterministic limits, the non-integrable covariance preserves macroscopic randomness.

We prove that under the critical scaling $\varepsilon^{-\alpha/2}$, the solution converges in distribution to a stochastic limit described by a fractional Gaussian field with Hurst index $H = 1 - \alpha/2 > 1/2$:

$$u(t, x) = \mathbb{E}^B \left[\varphi(x + B_t) \exp \left(\beta \int_{\mathbb{R}} L_t^x(y) dW^H(y) \right) \right],$$

where W^H is fractional Brownian motion and the integral is a Young integral. Our contributions include: (i) functional convergence of the integrated potential to fBm, (ii) quantitative convergence rates in Wasserstein distance $W_2(u_\varepsilon, u) \leq C\varepsilon^{\min(\alpha, 1-\alpha)/4}$, (iii) a central limit theorem for rescaled fluctuations with scaling $\varepsilon^{-\alpha/4}$, and (iv) superdiffusive transport $\mathbb{E}[X_t^2] \sim t^{2H}$.

The results reveal a new homogenization mechanism driven by long-range dependence, connecting stochastic homogenization, fractional calculus, and anomalous diffusion theory.

1 Introduction

Homogenization theory characterizes the effective large-scale behavior of partial differential equations with rapidly oscillating coefficients. In classical stochastic homogenization under mixing conditions, microscopic randomness averages out through spatial ergodicity, leading to deterministic homogenized equations [Papanicolaou and Varadhan \(1981\)](#); [Bensoussan, Lions and Papanicolaou \(1978\)](#). However, numerous physical systems exhibit *long-range dependence* (LRD), where spatial correlations decay slowly as power laws rather than exponentially. This phenomenon, observed in hydrology [Hurst \(1951\)](#), turbulence [Frisch \(1995\)](#), geophysics [Sahimi \(2003\)](#), and finance [Mandelbrot \(1997\)](#), fundamentally alters transport properties and challenges classical averaging principles.

For parabolic equations with random potentials, the choice of scaling exponent is crucial. Consider

$$\partial_t u_\varepsilon(t, x) = \frac{1}{2} \partial_{xx} u_\varepsilon(t, x) + \varepsilon^{-\gamma} a\left(\frac{x}{\varepsilon}\right) u_\varepsilon(t, x), \quad u_\varepsilon(0, x) = \varphi(x). \quad (1)$$

When a has integrable correlations ($\int |R_a| < \infty$), $\gamma = 1/2$ yields Gaussian white noise limits [Pardoux and Piatnitski \(2006\)](#); [Gu and Bal \(2014\)](#). In contrast, we address the regime

Keywords and phrases. Stochastic homogenization, long-range dependence, fractional Brownian motion, anomalous diffusion, Young integral, quantitative convergence, central limit theorem.

$R_a(x) \sim |x|^{-\alpha}$ with $\alpha \in (0, 1)$, where we show $\gamma = \alpha/2$ produces fractional Gaussian limits.

1.1 Mathematical framework and scaling heuristics

We consider equation (1) with $\gamma = \alpha/2$, where $a(x) = \Phi(g(x))$ is obtained from a stationary Gaussian field g with covariance $R_g(x) \sim \kappa_g |x|^{-\alpha}$ via a nonlinear transformation Φ of Hermite rank one. The Feynman-Kac representation

$$u_\varepsilon(t, x) = \mathbb{E}^B \left[\varphi(x + B_t) \exp \left(\varepsilon^{-\alpha/2} \int_0^t a \left(\frac{x + B_s}{\varepsilon} \right) ds \right) \right]$$

reveals that asymptotics depend on the functional

$$Y_t^{\varepsilon, x} = \varepsilon^{-\alpha/2} \int_0^t a \left(\frac{x + B_s}{\varepsilon} \right) ds.$$

Heuristically, its variance scales as

$$\text{Var}(Y_t^{\varepsilon, x}) \sim \varepsilon^{-\alpha} \int_0^t \int_0^t R_a \left(\frac{B_s - B_r}{\varepsilon} \right) ds dr \sim \int_0^t \int_0^t |B_s - B_r|^{-\alpha} ds dr \sim t^{2-\alpha} = t^{2H},$$

where $H = 1 - \alpha/2$, justifying the critical scaling $\varepsilon^{-\alpha/2}$.

1.2 Main contributions

This paper establishes a complete homogenization theory for equation (1) with LRD potentials:

- (i) **Functional convergence:** $\varepsilon^{-\alpha/2} \int_0^x a(y/\varepsilon) dy \Rightarrow \beta W^H$ with $H = 1 - \alpha/2$ (Theorem 3.1).
- (ii) **Homogenization limit:**

$$u_\varepsilon(t, x) \Rightarrow u(t, x) = \mathbb{E}^B \left[\varphi(x + B_t) \exp \left(\beta \int_{\mathbb{R}} L_t^x(y) dW^H(y) \right) \right]$$

(Theorem 3.2), where the integral is a Young integral well-defined for $H > 1/2$.

- (iii) **Quantitative convergence:** $W_2(u_\varepsilon(t, x), u(t, x)) \leq C \varepsilon^{\min(\alpha, 1-\alpha)/4}$ (Theorem 3.4).
- (iv) **Fluctuation theory:** $\varepsilon^{-\alpha/4}(u_\varepsilon - \mathbb{E}[u_\varepsilon]) \Rightarrow \mathcal{N}(0, \sigma^2(t, x))$ (Theorem 3.6).
- (v) **Anomalous transport:** $\mathbb{E}[X_t^2] \sim t^{2H}$ (Theorem 3.8), contrasting with normal diffusion for short-range correlations.

1.3 Relation to existing literature

Our work connects three active research areas: stochastic homogenization [Pardoux and Piatnitski \(2006\)](#); [Gu and Bal \(2014\)](#), limit theorems for long-range dependent processes [Taqqu \(1975\)](#); [Pipiras and Taqqu \(2017\)](#), and fractional stochastic calculus [Duncan, Hu and Pasik-Duncan \(2000\)](#). Closest is [Bal, Garnier, Motsch and Perrier \(2008\)](#) on Helmholtz equations with LRD coefficients; we extend this to the dynamic parabolic setting where test functions are Brownian paths.

The proofs combine: (i) functional convergence via Hermite expansions and the Breuer-Major theorem; (ii) Young integration theory for $H > 1/2$; (iii) Malliavin calculus and Stein's method for quantitative bounds; (iv) analysis of fluctuations via conditional central limit theorems.

1.4 Organization

Section 2 presents the mathematical framework. Section 3 states all main theorems. Section 4 provides essential tools on fractional calculus and Young integration. Sections 5–8 contain detailed proofs of the main results. Section 9 discusses transport properties. Appendices A–B provide additional technical proofs and numerical methods.

2 Preliminaries and Assumptions

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a complete probability space supporting all random objects.

2.1 Gaussian Fields with Long-Range Dependence

Assumption 2.1 (Gaussian field with long-range dependence). Let $\{g(x)\}_{x \in \mathbb{R}}$ be a centered stationary Gaussian field with $\mathbb{E}[g(0)^2] = 1$ whose covariance satisfies:

$$R_g(x) := \mathbb{E}[g(0)g(x)] = \kappa_g |x|^{-\alpha} L(|x|), \quad x \neq 0,$$

where $\alpha \in (0, 1)$, $\kappa_g > 0$, and L is a slowly varying function at infinity. We assume the spectral measure μ_g of g satisfies:

$$\int_{\mathbb{R}} |\xi|^{\alpha-1} \mu_g(d\xi) < \infty,$$

which ensures the existence of a continuous version of the field [Giraitis, Koul and Surgailis \(2012\)](#).

Remark 2.2. For simplicity in asymptotic statements, we often take $L \equiv 1$, so $R_g(x) \sim \kappa_g |x|^{-\alpha}$ as $|x| \rightarrow \infty$. The spectral condition guarantees the existence of a continuous modification via Kolmogorov's continuity theorem.

2.2 Nonlinear Transformation

Assumption 2.3 (Nonlinear transformation and Hermite rank). Let $\Phi : \mathbb{R} \rightarrow \mathbb{R}$ be a measurable function with $\mathbb{E}[\Phi(Z)^2] < \infty$ for $Z \sim \mathcal{N}(0, 1)$. Define

$$a(x) := \Phi(g(x)).$$

Assume Φ has Hermite rank $m_\Phi = 1$, i.e.,

$$V_1 = \mathbb{E}[\Phi(Z)H_1(Z)] \neq 0,$$

where $H_1(z) = z$ is the first Hermite polynomial. Additionally, we require $\Phi \in C^2(\mathbb{R})$ with bounded second derivative $|\Phi''(z)| \leq M$ for all $z \in \mathbb{R}$.

2.3 Initial Condition

Assumption 2.4 (Initial condition). The initial condition $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ satisfies $\varphi \in C_b(\mathbb{R})$ (bounded continuous) and is Lipschitz continuous. For Theorem 3.6, we additionally require $\varphi \in L^1(\mathbb{R}) \cap L^2(\mathbb{R})$.

2.4 Covariance Structure

Lemma 2.5 (Covariance structure of transformed field). *Under Assumptions 2.1 and 2.3, the covariance of a satisfies:*

$$R_a(x) := \mathbb{E}[a(0)a(x)] \sim \kappa|x|^{-\alpha} \quad \text{as } |x| \rightarrow \infty,$$

where $\kappa = V_1^2 \kappa_g$.

Proof. From the Hermite expansion $\Phi(u) = \sum_{q \geq 0} \frac{V_q}{q!} H_q(u)$ and orthogonality of Hermite polynomials with respect to Gaussian measure,

$$R_a(x) = \sum_{q=1}^{\infty} \frac{V_q^2}{q!} R_g(x)^q.$$

Since $R_g(x) \rightarrow 0$ as $|x| \rightarrow \infty$, the dominant term is $q = 1$:

$$R_a(x) = V_1^2 R_g(x) + O(|R_g(x)|^2) \sim V_1^2 \kappa_g |x|^{-\alpha}.$$

□

2.5 Fractional Brownian Motion

Definition 2.6 (Fractional Brownian motion). A fractional Brownian motion (fBm) $\{W^H(x)\}_{x \in \mathbb{R}}$ with Hurst index $H \in (0, 1)$ is a centered Gaussian process with covariance

$$\mathbb{E}[W^H(x)W^H(y)] = \frac{1}{2} (|x|^{2H} + |y|^{2H} - |x - y|^{2H}).$$

For $H > 1/2$, fBm has positively correlated increments and exhibits long-range dependence.

For our setting, the relevant Hurst index is

$$H := 1 - \frac{\alpha}{2} \in \left(\frac{1}{2}, 1\right). \quad (2)$$

2.6 Young Integration

Definition 2.7 (Young integral). Let $f \in C^\beta([a, b])$, $g \in C^\gamma([a, b])$ be Hölder continuous functions with exponents $\beta, \gamma \in (0, 1]$ satisfying $\beta + \gamma > 1$. The Young integral $\int_a^b f dg$ is defined as the limit of Riemann sums:

$$\int_a^b f dg := \lim_{|\Pi| \rightarrow 0} \sum_{i=0}^{n-1} f(t_i) [g(t_{i+1}) - g(t_i)],$$

where $\Pi = \{a = t_0 < t_1 < \dots < t_n = b\}$ is a partition of $[a, b]$. The integral satisfies the bound:

$$\left| \int_a^b f dg \right| \leq C_{\beta, \gamma} \|f\|_{C^\beta} \|g\|_{C^\gamma} (b - a)^{\beta + \gamma},$$

where $C_{\beta, \gamma}$ depends only on β and γ [Young \(1936\)](#); [Friz and Hairer \(2014\)](#).

Since Brownian local time $L_t^x(y)$ is γ -Hölder for any $\gamma < 1/2$ and W^H is $(H - \varepsilon)$ -Hölder with $H > 1/2$, we have $\gamma + (H - \varepsilon) > 1$ for sufficiently small ε , making $\int L_t^x dW^H$ well-defined as a Young integral.

2.7 Rescaled Processes

Define the rescaled integrated processes and normalization constant:

$$W_\varepsilon(x) := \varepsilon^{-\alpha/2} \int_0^x a\left(\frac{y}{\varepsilon}\right) dy, \quad (3)$$

$$\beta := \sqrt{\frac{\kappa}{H(2H-1)}} = \sqrt{\frac{V_1^2 \kappa_g}{(1-\alpha/2)\alpha}}. \quad (4)$$

3 Main Results

3.1 Functional convergence to fractional Brownian motion

Theorem 3.1 (Functional convergence of integrated potential). *Under Assumptions 2.1 and 2.3, as $\varepsilon \rightarrow 0$,*

$$W_\varepsilon \Rightarrow \beta W^H \quad \text{in } C(\mathbb{R}),$$

where W^H is fractional Brownian motion with Hurst index $H = 1 - \alpha/2$, and β is given by (4). The convergence holds in the space of continuous functions equipped with uniform convergence on compact sets.

3.2 Homogenization limit

Theorem 3.2 (Homogenization limit). *For each $t > 0$, $x \in \mathbb{R}$, under Assumptions 2.1, 2.3, and 2.4,*

$$u_\varepsilon(t, x) \Rightarrow u(t, x) := \mathbb{E}^B \left[\varphi(x + B_t) \exp \left(\beta \int_{\mathbb{R}} L_t^x(y) dW^H(y) \right) \right],$$

where the convergence holds jointly for finitely many (t, x) . The integral is a Young integral, well-defined since $H > 1/2$.

Remark 3.3 (Interpretation as fractional SPDE). Formally, $u(t, x)$ satisfies the stochastic partial differential equation

$$\partial_t u(t, x) = \frac{1}{2} \partial_{xx} u(t, x) + \beta u(t, x) \circ \partial_x W^H(x), \quad u(0, x) = \varphi(x),$$

where $\circ \partial_x W^H$ denotes Stratonovich-type multiplication by the (distributional) derivative of fBm. The equation is non-Markovian due to the memory induced by long-range correlations.

3.3 Quantitative convergence rates

Theorem 3.4 (Quantitative convergence in Wasserstein distance). *There exists a constant $C = C(t, \alpha, \kappa) > 0$ such that for all sufficiently small $\varepsilon > 0$,*

$$W_2(u_\varepsilon(t, x), u(t, x)) \leq C \varepsilon^{\min(\alpha, 1-\alpha)/4},$$

where W_2 denotes the Wasserstein-2 distance.

Remark 3.5 (Optimality discussion). The rate $O(\varepsilon^{\min(\alpha, 1-\alpha)/4})$ arises from two competing mechanisms: for small α , the error is dominated by the nonlinear remainder in the Hermite expansion; for α near 1, the Malliavin-Stein approximation error dominates. Numerical evidence suggests the optimal rate may be $\varepsilon^{\min(\alpha, 1-\alpha)/2}$ in Kolmogorov distance, with the square-root loss coming from the Wasserstein metric conversion.

3.4 Fluctuation theory

Theorem 3.6 (Central limit theorem for fluctuations). *Define the rescaled fluctuation process*

$$\mathcal{U}_\varepsilon(t, x) := \varepsilon^{-\alpha/4} (u_\varepsilon(t, x) - \mathbb{E}[u_\varepsilon(t, x)]).$$

Then, under Assumptions 2.1, 2.3, and 2.4,

$$\mathcal{U}_\varepsilon(t, x) \Rightarrow \mathcal{U}(t, x) \sim \mathcal{N}(0, \sigma^2(t, x)),$$

where the variance is given by

$$\sigma^2(t, x) = (\mathbb{E}^B[\varphi(x + B_t)])^2 \beta^2 \mathbb{E}^B \left[\int_0^t \int_0^t |B_s - B_r|^{2H-2} ds dr \right]. \quad (5)$$

Remark 3.7 (Scaling interpretation). The scaling $\varepsilon^{-\alpha/4}$ differs from the classical $\varepsilon^{-1/2}$ for short-range correlations, reflecting weaker fluctuations due to correlation persistence. The limiting variance involves Brownian self-interaction through the kernel $|B_s - B_r|^{2H-2}$.

3.5 Anomalous transport properties

Theorem 3.8 (Superdiffusive scaling). *Let $p(t, x) = u(t, x) / \int_{\mathbb{R}} u(t, y) dy$ be the effective particle density. The mean squared displacement exhibits superdiffusive behavior:*

$$\text{MSD}(t) := \int_{\mathbb{R}} x^2 p(t, x) dx \sim Ct^{2H} \quad \text{as } t \rightarrow \infty,$$

where $C > 0$ depends on α , κ , and φ , and $H = 1 - \alpha/2$.

Remark 3.9 (Transport regimes). The scaling t^{2H} with $H > 1/2$ contrasts with normal diffusion (t^1) for short-range correlations. As $\alpha \rightarrow 0$ ($H \rightarrow 1$), transport becomes ballistic (t^2); as $\alpha \rightarrow 1$ ($H \rightarrow 1/2$), we recover normal diffusion.

4 Technical Tools: Fractional Calculus and Young Integration

4.1 Continuity of Young Integral Map

Lemma 4.1 (Continuity of Young integral). *Let $\beta, \gamma \in (0, 1]$ with $\beta + \gamma > 1$. The Young integral operator*

$$\Psi : C^\beta([a, b]) \times C^\gamma([a, b]) \rightarrow \mathbb{R}, \quad \Psi(f, g) = \int_a^b f dg$$

is bilinear and continuous. Specifically, if $f_n \rightarrow f$ in C^β and $g_n \rightarrow g$ in C^γ , then

$$\lim_{n \rightarrow \infty} \int_a^b f_n dg_n = \int_a^b f dg,$$

and the convergence is uniform on compact sets.

Proof. By the Young-Loeve inequality [Friz and Hairer \(2014\)](#), we have:

$$\begin{aligned} \left| \int_a^b f_n dg_n - \int_a^b f dg \right| &\leq \left| \int_a^b (f_n - f) dg_n \right| + \left| \int_a^b f d(g_n - g) \right| \\ &\leq C_{\beta, \gamma} \|f_n - f\|_{C^\beta} \|g_n\|_{C^\gamma} (b - a)^{\beta + \gamma} \\ &\quad + C_{\beta, \gamma} \|f\|_{C^\beta} \|g_n - g\|_{C^\gamma} (b - a)^{\beta + \gamma}. \end{aligned}$$

Since $\|g_n\|_{C^\gamma}$ is bounded and $\|f_n - f\|_{C^\beta}, \|g_n - g\|_{C^\gamma} \rightarrow 0$, the result follows. \square

4.2 Fractional Itô-Tanaka Formula

Theorem 4.2 (Fractional Itô-Tanaka formula). *Let W^H be fractional Brownian motion with $H > 1/2$, and let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a C^2 function with bounded derivatives. Then for any $t \geq 0$ and $x \in \mathbb{R}$,*

$$f(x + W_t^H) = f(x) + \int_0^t f'(x + W_s^H) dW_s^H + \frac{1}{2} \int_{\mathbb{R}} L_t^x(y) f''(y) dy,$$

where the stochastic integral is interpreted in the Young sense, and $L_t^x(y)$ is the local time of W^H at level y .

Proof. We provide a detailed proof using mollification and Young integration estimates.

Step 1: Mollification. Let ρ_n be a sequence of mollifiers, and define $W_n^H = \rho_n * W^H$, which are C^∞ approximations. Since W^H is $(H - \varepsilon)$ -Hölder continuous, $W_n^H \rightarrow W^H$ in $C^{H-\varepsilon}$ uniformly on compacts.

Step 2: Classical Itô formula. For the smooth process W_n^H , the classical Itô formula applies:

$$f(x + W_n^H(t)) = f(x) + \int_0^t f'(x + W_n^H(s)) dW_n^H(s) + \frac{1}{2} \int_0^t f''(x + W_n^H(s)) d[W_n^H]_s,$$

where $[W_n^H]_s$ denotes the quadratic variation.

Step 3: Convergence. Since f' is Lipschitz and $W_n^H \rightarrow W^H$ in $C^{H-\varepsilon}$, we have:

$$\left| \int_0^t f'(x + W_n^H(s)) dW_n^H(s) - \int_0^t f'(x + W^H(s)) dW^H(s) \right| \rightarrow 0$$

by Lemma 4.1. For the quadratic variation term, we use the occupation density formula:

$$\int_0^t f''(x + W_n^H(s)) d[W_n^H]_s = \int_{\mathbb{R}} L_t^{n,x}(y) f''(y) dy,$$

where $L_t^{n,x}$ is the local time of W_n^H . Since $W_n^H \rightarrow W^H$ uniformly and local times converge [Revuz and Yor \(1999\)](#), we obtain the limit.

Step 4: Identification. The limit of the quadratic variation term is $\int_{\mathbb{R}} L_t^x(y) f''(y) dy$, giving the desired formula. \square

4.3 Local Time Regularity

Lemma 4.3 (Hölder regularity of Brownian local time). *For fixed $t > 0$, the process $y \mapsto L_t^x(y)$ admits a version that is γ -Hölder continuous for any $\gamma < 1/2$. Moreover, for any $p \geq 1$,*

$$\mathbb{E}[|L_t^x(y) - L_t^x(z)|^p] \leq C_p |y - z|^{p/2}.$$

Proof. This is a classical result in stochastic analysis. The Hölder continuity follows from the occupation density formula and the Tanaka formula. See [\(Revuz and Yor, 1999, Chapter VI\)](#) for a complete proof. \square

5 Proof of Functional Convergence

Let $\Phi(u) = \sum_{q=0}^{\infty} \frac{V_q}{q!} H_q(u)$ be the Hermite expansion. Then

$$W_\varepsilon(x) = \sum_{q=1}^{\infty} \frac{V_q}{q!} W_\varepsilon^{(q)}(x), \quad W_\varepsilon^{(q)}(x) := \varepsilon^{-\alpha/2} \int_0^x H_q\left(g\left(\frac{y}{\varepsilon}\right)\right) dy.$$

Lemma 5.1 (Negligibility of higher-order chaoses). *For any $q \geq 2$ and compact $K \subset \mathbb{R}$,*

$$\lim_{\varepsilon \rightarrow 0} \mathbb{E} \left[\sup_{x \in K} |W_\varepsilon^{(q)}(x)|^2 \right] = 0.$$

Consequently, $W_\varepsilon - V_1 \widetilde{W}_\varepsilon \Rightarrow 0$ in $C(\mathbb{R})$, where $\widetilde{W}_\varepsilon(x) := \varepsilon^{-\alpha/2} \int_0^x g(y/\varepsilon) dy$.

Proof. For fixed $q \geq 2$,

$$\begin{aligned} \mathbb{E} \left[|W_\varepsilon^{(q)}(x)|^2 \right] &= \varepsilon^{-\alpha} \int_0^x \int_0^x \mathbb{E} \left[H_q\left(g\left(\frac{u}{\varepsilon}\right)\right) H_q\left(g\left(\frac{v}{\varepsilon}\right)\right) \right] dudv \\ &= q! \varepsilon^{-\alpha} \int_0^x \int_0^x R_g\left(\frac{u-v}{\varepsilon}\right)^q dudv \\ &= q! \varepsilon^{2-\alpha} \int_0^{x/\varepsilon} \int_0^{x/\varepsilon} R_g(u-v)^q dudv. \end{aligned}$$

Since $|R_g(z)|^q \leq C|z|^{-q\alpha}$ and $q\alpha > \alpha$, we have

$$\varepsilon^{2-\alpha} \int_0^{x/\varepsilon} \int_0^{x/\varepsilon} |u-v|^{-q\alpha} dudv = O(\varepsilon^{1+\alpha(1-q)}) \rightarrow 0.$$

Tightness follows via Kolmogorov's criterion: for $p > 1/H$,

$$\mathbb{E} \left[|W_\varepsilon^{(q)}(x) - W_\varepsilon^{(q)}(y)|^p \right] \leq C_p |x - y|^{pH},$$

using hypercontractivity for Wiener chaoses. □

Lemma 5.2 (Covariance asymptotics). *For any $x, y \in \mathbb{R}$,*

$$\lim_{\varepsilon \rightarrow 0} \mathbb{E} \left[\widetilde{W}_\varepsilon(x) \widetilde{W}_\varepsilon(y) \right] = \frac{\kappa_g}{H(2H-1)} \cdot \frac{1}{2} (|x|^{2H} + |y|^{2H} - |x-y|^{2H}).$$

Proof. We compute

$$\begin{aligned} \mathbb{E} \left[\widetilde{W}_\varepsilon(x) \widetilde{W}_\varepsilon(y) \right] &= \varepsilon^{-\alpha} \int_0^x \int_0^y R_g\left(\frac{u-v}{\varepsilon}\right) dudv \\ &= \varepsilon^{2-\alpha} \int_0^{x/\varepsilon} \int_0^{y/\varepsilon} R_g(u-v) dudv. \end{aligned}$$

Let $I(X, Y) := \int_0^X \int_0^Y R_g(u-v) dudv$. From (Taqqu, 1975, Theorem 2.1),

$$I(X, Y) \sim \frac{\kappa_g}{H(2H-1)} \cdot \frac{1}{2} (X^{2H} + Y^{2H} - |X-Y|^{2H}) \quad \text{as } X, Y \rightarrow \infty.$$

Substituting $X = x/\varepsilon$, $Y = y/\varepsilon$ and using $\varepsilon^{2-\alpha} = \varepsilon^{2H}$ yields the result. □

Lemma 5.3 (Moment estimates and tightness). *For any $p \geq 2$, there exists $C_p > 0$ such that for all $\varepsilon > 0$ and $x, y \in \mathbb{R}$,*

$$\mathbb{E} \left[|\widetilde{W}_\varepsilon(x) - \widetilde{W}_\varepsilon(y)|^p \right] \leq C_p |x - y|^{pH}.$$

Proof. Since $\widetilde{W}_\varepsilon(x) - \widetilde{W}_\varepsilon(y)$ is Gaussian,

$$\mathbb{E} \left[|\widetilde{W}_\varepsilon(x) - \widetilde{W}_\varepsilon(y)|^p \right] = \mu_p \left(\mathbb{E} \left[|\widetilde{W}_\varepsilon(x) - \widetilde{W}_\varepsilon(y)|^2 \right] \right)^{p/2},$$

where $\mu_p = \mathbb{E}[|Z|^p]$ for $Z \sim \mathcal{N}(0, 1)$. From Lemma 5.2,

$$\mathbb{E} \left[|\widetilde{W}_\varepsilon(x) - \widetilde{W}_\varepsilon(y)|^2 \right] \sim \frac{\kappa_g}{H(2H-1)} |x - y|^{2H},$$

giving the bound. \square

Proof of Theorem 3.1. By Lemma 5.1, it suffices to prove $V_1 \widetilde{W}_\varepsilon \Rightarrow \beta W^H$. Lemma 5.2 gives convergence of finite-dimensional distributions to those of $\sqrt{\kappa_g/(H(2H-1))} W^H = \beta/V_1 \cdot W^H$. Lemma 5.3 with $p > 1/H$ yields tightness in $C(\mathbb{R})$ via Kolmogorov's criterion. Prokhorov's theorem completes the proof. \square

6 Proof of Homogenization Limit

6.1 Occupation Time Representation

The key functional admits two useful representations:

Lemma 6.1 (Occupation time representation). *For each $\varepsilon > 0$, $t \geq 0$, $x \in \mathbb{R}$,*

$$Y_t^{\varepsilon, x} := \varepsilon^{-\alpha/2} \int_0^t a\left(\frac{x + B_s}{\varepsilon}\right) ds = \int_{\mathbb{R}} L_t^x(y) dW_\varepsilon(y),$$

where $L_t^x(y)$ is the Brownian local time of $x + B_t$ at level y , and the integral is interpreted as a Young integral.

Proof. Let $W_\varepsilon^*(y) = \varepsilon^{-\alpha/2} \int_0^y a(z/\varepsilon) dz$. By the occupation density formula,

$$\int_0^t a\left(\frac{x + B_s}{\varepsilon}\right) ds = \int_{\mathbb{R}} a\left(\frac{y}{\varepsilon}\right) L_t^x(y) dy.$$

Now, using integration by parts for Young integrals (valid since L_t^x is $(1/2 - \delta)$ -Hölder and W_ε is $(H - \delta)$ -Hölder),

$$\begin{aligned} \int_{\mathbb{R}} L_t^x(y) dW_\varepsilon(y) &= \lim_{R \rightarrow \infty} \left[L_t^x(y) W_\varepsilon(y) \Big|_{-R}^R - \int_{-R}^R W_\varepsilon(y) dL_t^x(y) \right] \\ &= - \int_{\mathbb{R}} W_\varepsilon(y) dL_t^x(y) \quad (\text{since } L_t^x \text{ has compact support}). \end{aligned}$$

But $dL_t^x(y)$ is the occupation density measure, so

$$- \int_{\mathbb{R}} W_\varepsilon(y) dL_t^x(y) = - \int_{\mathbb{R}} \left(\varepsilon^{-\alpha/2} \int_0^y a(z/\varepsilon) dz \right) dL_t^x(y).$$

By Fubini's theorem (justified by the Hölder regularity),

$$= -\varepsilon^{-\alpha/2} \int_{\mathbb{R}} a(z/\varepsilon) \left(\int_z^\infty dL_t^x(y) \right) dz = \varepsilon^{-\alpha/2} \int_{\mathbb{R}} a(z/\varepsilon) L_t^x(z) dz,$$

which equals $Y_t^{\varepsilon, x}$. \square

6.2 Itô Representation

Lemma 6.2 (Itô representation). *Let $\Phi_\varepsilon(x) := \int_0^x W_\varepsilon(y) dy$. Then*

$$Y_t^{\varepsilon,x} = 2[\Phi_\varepsilon(x + B_t) - \Phi_\varepsilon(x)] - 2 \int_0^t W_\varepsilon(x + B_s) dB_s.$$

Proof. Apply Itô's formula to $F(y) = \Phi_\varepsilon(x + y)$ evaluated at $y = B_t$:

$$\Phi_\varepsilon(x + B_t) - \Phi_\varepsilon(x) = \int_0^t W_\varepsilon(x + B_s) dB_s + \frac{1}{2} \int_0^t W'_\varepsilon(x + B_s) ds.$$

Since $W'_\varepsilon(y) = \varepsilon^{-\alpha/2} a(y/\varepsilon)$, we have

$$\int_0^t W'_\varepsilon(x + B_s) ds = Y_t^{\varepsilon,x}.$$

Solving for $Y_t^{\varepsilon,x}$ yields the result. \square

6.3 Exponential Moment Bounds

Lemma 6.3 (Exponential moment bounds). *For any $p \in \mathbb{R}$, there exists $C_p(t) < \infty$ such that*

$$\sup_{\varepsilon > 0} \mathbb{E}[\exp(pY_t^{\varepsilon,x})] \leq C_p(t).$$

Proof. Condition on the Brownian path B . Then $Y_t^{\varepsilon,x}$ is Gaussian conditional on B with variance

$$\sigma_\varepsilon^2(B) := \text{Var}(Y_t^{\varepsilon,x} \mid B) = \varepsilon^{-\alpha} \int_0^t \int_0^t R_a\left(\frac{B_s - B_r}{\varepsilon}\right) ds dr.$$

Since $|R_a(z)| \leq C(1 + |z|)^{-\alpha}$, we have

$$\sigma_\varepsilon^2(B) \leq C\varepsilon^{-\alpha} \int_0^t \int_0^t (1 + |B_s - B_r|/\varepsilon)^{-\alpha} ds dr \leq C't^{2H},$$

uniformly in ε , where the last inequality follows from scaling: $(1 + |z|/\varepsilon)^{-\alpha} \leq \varepsilon^\alpha |z|^{-\alpha}$ for $|z| \geq \varepsilon$. Thus $\mathbb{E}[\exp(pY_t^{\varepsilon,x}) \mid B] = \exp(p^2 \sigma_\varepsilon^2(B)/2) \leq \exp(C_p t^{2H})$. \square

6.4 Convergence in Hölder Spaces

Lemma 6.4 (Convergence in Hölder spaces). *The convergence $W_\varepsilon \Rightarrow \beta W^H$ in $C(\mathbb{R})$ implies convergence in $C^\gamma(K)$ for any compact $K \subset \mathbb{R}$ and any $\gamma < H$.*

Proof. From Lemma 5.3, the family $\{W_\varepsilon\}$ satisfies Kolmogorov's continuity criterion: for p large enough,

$$\mathbb{E}[|W_\varepsilon(x) - W_\varepsilon(y)|^p] \leq C_p |x - y|^{pH}.$$

This implies tightness in $C^\gamma(K)$ for $\gamma < H - 1/p$ by the Arzelà-Ascoli theorem. Since convergence in finite-dimensional distributions plus tightness in $C^\gamma(K)$ implies convergence in $C^\gamma(K)$, the result follows. \square

Proof of Theorem 3.2. From Lemma 6.1, $Y_t^{\varepsilon,x} = \Psi(W_\varepsilon)$, where $\Psi(f) = \int_{\mathbb{R}} L_t^x(y) df(y)$.

By Theorem 3.1 and Lemma 6.4, $W_\varepsilon \Rightarrow \beta W^H$ in $C^\gamma(K)$ for any $\gamma < H$ and compact K . Since L_t^x is $(1/2 - \delta)$ -Hölder, we can choose $\gamma > 1/2$ (possible since $H > 1/2$) such that $\gamma + (1/2 - \delta) > 1$.

By Lemma 4.1, Ψ is continuous from $C^\gamma(K)$ to \mathbb{R} when restricted to functions with compact support. Since L_t^x has compact support, the continuous mapping theorem implies:

$$Y_t^{\varepsilon,x} \Rightarrow \beta \int_{\mathbb{R}} L_t^x(y) dW^H(y) =: Y_t^{0,x}.$$

From the Feynman-Kac representation $u_\varepsilon(t, x) = \mathbb{E}^B[\varphi(x + B_t)e^{Y_t^{\varepsilon,x}}]$, Lemma 6.3 provides uniform integrability. Thus,

$$u_\varepsilon(t, x) \Rightarrow \mathbb{E}^B[\varphi(x + B_t)e^{Y_t^{0,x}}] = u(t, x).$$

Joint convergence follows from joint convergence of $(Y_{t_1}^{\varepsilon,x_1}, \dots, Y_{t_k}^{\varepsilon,x_k})$ and the continuous mapping theorem. \square

7 Proof of Quantitative Convergence

Let $\mathfrak{H} = L^2(\mathbb{R})$ be the Hilbert space associated with the Gaussian field g . For $F \in \mathbb{D}^{1,2}$, denote by DF its Malliavin derivative and by L^{-1} the pseudo-inverse of the Ornstein-Uhlenbeck operator.

Theorem 7.1 (Malliavin-Stein bound [Nourdin and Peccati \(2012\)](#)). *Let $F \in \mathbb{D}^{1,2}$ with $\mathbb{E}[F] = 0$, $\mathbb{E}[F^2] = 1$. For $Z \sim \mathcal{N}(0, 1)$,*

$$d_W(F, Z) \leq \mathbb{E} [|1 - \langle DF, -DL^{-1}F \rangle_{\mathfrak{H}}|],$$

where d_W is the Wasserstein-1 distance.

Consider the normalized first chaos component:

$$F_\varepsilon := \frac{Y_t^{\varepsilon,x,(1)}}{\sigma_\varepsilon}, \quad \sigma_\varepsilon^2 := \mathbb{E} \left[\left(Y_t^{\varepsilon,x,(1)} \right)^2 \right],$$

where $Y_t^{\varepsilon,x,(1)} = \varepsilon^{-\alpha/2} \int_0^t g\left(\frac{x+B_s}{\varepsilon}\right) ds$.

Lemma 7.2 (Malliavin derivative). *For F_ε as above,*

$$D_u F_\varepsilon = \frac{1}{\sigma_\varepsilon} \varepsilon^{-\alpha/2} \int_0^t \mathbf{1}_{[0, x+B_s]}(u) ds, \quad u \in \mathbb{R}.$$

Proof. Since $D_u g(y) = \delta(y - u)$ (in the sense of distributions), we have

$$D_u Y_t^{\varepsilon,x,(1)} = \varepsilon^{-\alpha/2} \int_0^t \delta\left(\frac{x+B_s}{\varepsilon} - u\right) ds = \varepsilon^{1-\alpha/2} \int_0^t \delta(x+B_s - \varepsilon u) ds.$$

Approximating the delta function by $\frac{1}{2\eta} \mathbf{1}_{[-\eta, \eta]}$ and taking $\eta \rightarrow 0$ gives the indicator representation. \square

Lemma 7.3 (Variance of the Malliavin derivative norm). *There exists $C = C(\alpha, \kappa_g, t) > 0$ such that for all $\varepsilon \in (0, 1]$,*

$$\text{Var}(\|DF_\varepsilon\|_{\mathfrak{H}}^2) \leq C \varepsilon^{2\min(\alpha, 1-\alpha)}.$$

Proof. Let $h_\varepsilon(u) = \varepsilon^{-\alpha/2} \int_0^t \mathbf{1}_{[0, x+B_s]}(u) ds$. Then

$$\|DF_\varepsilon\|_{\mathfrak{H}}^2 = \frac{1}{\sigma_\varepsilon^2} \int_{\mathbb{R}^2} h_\varepsilon(u) h_\varepsilon(v) R_g(u-v) du dv.$$

This is a double Wiener-Itô integral. Using the product formula for multiple integrals,

$$\begin{aligned} \text{Var}(\|DF_\varepsilon\|_{\mathfrak{H}}^2) &= \frac{2}{\sigma_\varepsilon^4} \int_{\mathbb{R}^4} h_\varepsilon(u_1) h_\varepsilon(v_1) h_\varepsilon(u_2) h_\varepsilon(v_2) \\ &\quad \times R_g(u_1 - v_1) R_g(u_2 - v_2) R_g(u_1 - u_2) R_g(v_1 - v_2) du_1 dv_1 du_2 dv_2. \end{aligned}$$

Integrate over Brownian paths. Since $\mathbb{E}^B[|h_\varepsilon(u)|^p] \leq C_p t^p \varepsilon^{-p\alpha/2} \mathbb{P}(u \in [0, x + B_{[0,t]}])^{1/p}$ and the Brownian path has measure $O(t^{1/2})$,

$$\int_{\mathbb{R}} \mathbb{E}^B[|h_\varepsilon(u)|^p] du \leq C'_p t^{p+1/2} \varepsilon^{-p\alpha/2}.$$

Combining with $|R_g(z)| \leq \kappa_g |z|^{-\alpha}$ and separating integration domains into $|u_i - u_j| \leq \varepsilon$ and $|u_i - u_j| > \varepsilon$,

$$\text{Var}(\|DF_\varepsilon\|_{\mathfrak{H}}^2) \leq C \left(\varepsilon^{2(1-\alpha)} + \varepsilon^{2\alpha} \right) \leq 2C \varepsilon^{2\min(\alpha, 1-\alpha)}.$$

□

Proof of Theorem 3.4. Since $Y_t^{\varepsilon, x, (1)}$ belongs to the first Wiener chaos, $L^{-1}Y_t^{\varepsilon, x, (1)} = Y_t^{\varepsilon, x, (1)}$. Thus

$$\langle DF_\varepsilon, -DL^{-1}F_\varepsilon \rangle_{\mathfrak{H}} = \frac{1}{\sigma_\varepsilon^2} \|DY_t^{\varepsilon, x, (1)}\|_{\mathfrak{H}}^2.$$

By Theorem 7.1,

$$d_W(F_\varepsilon, Z) \leq \mathbb{E} \left[\left| 1 - \frac{1}{\sigma_\varepsilon^2} \|DY_t^{\varepsilon, x, (1)}\|_{\mathfrak{H}}^2 \right| \right] \leq \frac{\sqrt{\text{Var}(\|DF_\varepsilon\|_{\mathfrak{H}}^2)}}{\sigma_\varepsilon^2}.$$

From Lemma 7.3 and $\sigma_\varepsilon^2 \sim \beta^2 t^{2H}$,

$$d_W(F_\varepsilon, Z) \leq C \varepsilon^{\min(\alpha, 1-\alpha)/2}.$$

For the full functional $Y_t^{\varepsilon, x} = V_1 Y_t^{\varepsilon, x, (1)} + R_\varepsilon$, we have $W_2(R_\varepsilon, 0) \leq \sqrt{\mathbb{E}[R_\varepsilon^2]} = O(\varepsilon^{\alpha/2})$. By triangle inequality,

$$W_2(Y_t^{\varepsilon, x}, \beta Y_t^{0, (1)}) \leq C' \varepsilon^{\min(\alpha, 1-\alpha)/2}.$$

Finally, using the Lipschitz property of $y \mapsto \mathbb{E}^B[\varphi(x + B_t)e^y]$ (with Lipschitz constant $L_\varphi e^{Ct^{2H}}$ by Lemma 6.3) and $W_2 \leq \sqrt{2}d_W$ for centered random variables with bounded second moments,

$$W_2(u_\varepsilon(t, x), u(t, x)) \leq C'' \varepsilon^{\min(\alpha, 1-\alpha)/4}.$$

□

8 Proof of Fluctuation Theorem

Lemma 8.1 (Linearization error). *Let $f_\varepsilon(y) := \mathbb{E}^B[\varphi(x + B_t)e^y]$. Then*

$$u_\varepsilon(t, x) - \mathbb{E}[u_\varepsilon(t, x)] = f'_\varepsilon(0)Y_t^{\varepsilon, x} + \mathcal{R}_\varepsilon,$$

with $\mathbb{E}[\mathcal{R}_\varepsilon^2] = O(\varepsilon^\alpha)$.

Proof. Taylor expansion gives

$$e^{Y_t^{\varepsilon, x}} = 1 + Y_t^{\varepsilon, x} + \int_0^1 (1 - \theta) e^{\theta Y_t^{\varepsilon, x}} (Y_t^{\varepsilon, x})^2 d\theta.$$

Thus

$$\begin{aligned} u_\varepsilon(t, x) &= f_\varepsilon(0) + f'_\varepsilon(0)Y_t^{\varepsilon, x} + \mathbb{E}^B \left[\varphi(x + B_t) \int_0^1 (1 - \theta) e^{\theta Y_t^{\varepsilon, x}} (Y_t^{\varepsilon, x})^2 d\theta \right] \\ &= \mathbb{E}[u_\varepsilon(t, x)] + f'_\varepsilon(0)Y_t^{\varepsilon, x} + \mathcal{R}_\varepsilon. \end{aligned}$$

Since $\mathbb{E}[Y_t^{\varepsilon, x}] = 0$ and $\mathbb{E}[(Y_t^{\varepsilon, x})^4] = O(1)$ (by Lemma 6.3 with $p = 4$), the remainder satisfies

$$\mathbb{E}[\mathcal{R}_\varepsilon^2] \leq \|\varphi\|_\infty^2 \mathbb{E} \left[\left(\int_0^1 (1 - \theta) e^{\theta Y_t^{\varepsilon, x}} (Y_t^{\varepsilon, x})^2 d\theta \right)^2 \right] \leq C \mathbb{E}[(Y_t^{\varepsilon, x})^4] = O(\varepsilon^\alpha),$$

where the last equality uses $\text{Var}(Y_t^{\varepsilon, x}) \sim \varepsilon^\alpha t^{2H}$ for small ε . \square

Lemma 8.2 (Conditional CLT via Breuer-Major). *Conditional on the Brownian path B ,*

$$\varepsilon^{-\alpha/4} Y_t^{\varepsilon, x} \Rightarrow \mathcal{N}(0, \sigma^2(B)) \quad \text{as } \varepsilon \rightarrow 0,$$

where

$$\sigma^2(B) = \beta^2 \int_0^t \int_0^t |B_s - B_r|^{2H-2} ds dr.$$

Proof. Condition on B . Then $\varepsilon^{-\alpha/4} Y_t^{\varepsilon, x}$ is a normalized sum of the stationary sequence $a((x + B_s)/\varepsilon)$. Since a has Hermite rank 1, the classical Breuer-Major theorem [Breuer and Major \(1983\)](#) applies, giving Gaussian limit with variance:

$$\text{Var}(\varepsilon^{-\alpha/4} Y_t^{\varepsilon, x} \mid B) = \varepsilon^{-\alpha/2} \int_0^t \int_0^t R_a \left(\frac{B_s - B_r}{\varepsilon} \right) ds dr \rightarrow \beta^2 \int_0^t \int_0^t |B_s - B_r|^{2H-2} ds dr.$$

\square

Proof of Theorem 3.6. By Lemma 8.1,

$$\mathcal{U}_\varepsilon(t, x) = \varepsilon^{-\alpha/4} f'_\varepsilon(0) Y_t^{\varepsilon, x} + \varepsilon^{-\alpha/4} \mathcal{R}_\varepsilon.$$

Since $\varepsilon^{-\alpha/4} \mathcal{R}_\varepsilon = O_P(\varepsilon^{\alpha/4}) \rightarrow 0$, we have

$$\mathcal{U}_\varepsilon(t, x) = \varepsilon^{-\alpha/4} f'_\varepsilon(0) Y_t^{\varepsilon, x} + o_{\mathbb{P}}(1).$$

Now $f'_\varepsilon(0) = \mathbb{E}^B[\varphi(x + B_t)] + O(\varepsilon^{\alpha/2})$ because

$$f'_\varepsilon(y) = \mathbb{E}^B[\varphi(x + B_t)e^y], \quad f'_\varepsilon(0) = \mathbb{E}^B[\varphi(x + B_t)],$$

and the error from replacing $f'_\varepsilon(0)$ by $\mathbb{E}^B[\varphi(x + B_t)]$ is $O(\varepsilon^{\alpha/2})$ by smoothness of Φ .

Thus

$$\mathcal{U}_\varepsilon(t, x) = \mathbb{E}^B[\varphi(x + B_t)] \cdot \varepsilon^{-\alpha/4} Y_t^{\varepsilon, x} + o_{\mathbb{P}}(1).$$

From Lemma 8.2, conditional on B , $\varepsilon^{-\alpha/4} Y_t^{\varepsilon, x}$ converges to $\mathcal{N}(0, \sigma^2(B))$. Taking expectation over B yields the unconditional limit $\mathcal{N}(0, \sigma^2(t, x))$ with

$$\sigma^2(t, x) = (\mathbb{E}^B[\varphi(x + B_t)])^2 \mathbb{E}^B[\sigma^2(B)].$$

Computing $\mathbb{E}^B[\sigma^2(B)]$ gives (5). □

9 Anomalous Diffusion and Transport Properties

Proof of Theorem 3.8. We provide a rigorous proof of the superdiffusive scaling using self-similarity arguments and Laplace method.

Step 1: Scaling properties. Recall that fractional Brownian motion is self-similar: $W^H(ty) \stackrel{d}{=} t^H W^H(y)$. Brownian local time scales as: $L_{at}(x) \stackrel{d}{=} \sqrt{a} L_t(x/\sqrt{a})$.

For the exponential functional, define:

$$Z_t := \exp \left(\beta \int_{\mathbb{R}} L_t^0(y) dW^H(y) \right).$$

Under scaling $t \mapsto ct$, we have:

$$\begin{aligned} \int_{\mathbb{R}} L_{ct}^0(y) dW^H(y) &\stackrel{d}{=} \int_{\mathbb{R}} \sqrt{c} L_t(y/\sqrt{c}) dW^H(y) \\ &\stackrel{d}{=} c^{H+1/2} \int_{\mathbb{R}} L_t(z) dW^H(z), \end{aligned}$$

where we used the change of variable $z = y/\sqrt{c}$ and self-similarity of fBm.

Step 2: Moment generating function. Consider the moment generating function:

$$M(\lambda) := \mathbb{E} \left[\exp \left(\lambda \int_{\mathbb{R}} L_t^0(y) dW^H(y) \right) \right].$$

Conditional on B , the integral is Gaussian with variance:

$$\sigma_t^2 = \beta^2 \int_0^t \int_0^t |B_s - B_r|^{2H-2} ds dr.$$

Thus,

$$M(\lambda) = \mathbb{E}^B \left[\exp \left(\frac{\lambda^2 \beta^2}{2} \int_0^t \int_0^t |B_s - B_r|^{2H-2} ds dr \right) \right].$$

Step 3: Large time asymptotics. For large t , by Brownian scaling:

$$\int_0^t \int_0^t |B_s - B_r|^{2H-2} ds dr \stackrel{d}{=} t^{2H} \int_0^1 \int_0^1 |B_u - B_v|^{2H-2} du dv.$$

Therefore,

$$M(\lambda) \sim \exp(C \lambda^2 t^{2H}) \quad \text{as } t \rightarrow \infty,$$

where $C = \frac{\beta^2}{2} \mathbb{E} \left[\int_0^1 \int_0^1 |B_u - B_v|^{2H-2} du dv \right]$.

Step 4: Mean squared displacement. The effective density is:

$$p(t, x) = \frac{\mathbb{E}^B [\varphi(x + B_t) Z_t]}{\int_{\mathbb{R}} \mathbb{E}^B [\varphi(y + B_t) Z_t] dy}.$$

By change of variables $x = t^H \xi$, and using the scaling properties from Step 1:

$$\text{MSD}(t) = \frac{\int_{\mathbb{R}} x^2 \mathbb{E}^B [\varphi(x + B_t) Z_t] dx}{\int_{\mathbb{R}} \mathbb{E}^B [\varphi(x + B_t) Z_t] dx} \sim t^{2H} \frac{\int_{\mathbb{R}} \xi^2 \Psi(\xi) d\xi}{\int_{\mathbb{R}} \Psi(\xi) d\xi},$$

where

$$\Psi(\xi) = \lim_{t \rightarrow \infty} \mathbb{E}^B [\varphi(t^H \xi + B_t) Z_t] t^{-H}.$$

The limit exists by the scaling properties and the boundedness of φ .

Step 5: Constant computation. The constant C in the theorem statement is:

$$C = \frac{\int_{\mathbb{R}} \xi^2 \Psi(\xi) d\xi}{\int_{\mathbb{R}} \Psi(\xi) d\xi},$$

which is positive and finite under our assumptions on φ . □

10 Conclusion

We have established a complete homogenization theory for the parabolic equation

$$\partial_t u_\varepsilon = \frac{1}{2} \partial_{xx} u_\varepsilon + \varepsilon^{-\alpha/2} a(x/\varepsilon) u_\varepsilon$$

with long-range correlated potential a (covariance $\sim |x|^{-\alpha}$, $\alpha \in (0, 1)$).

The main results are:

- The solution converges to a stochastic limit given by a fractional Feynman-Kac formula involving fractional Brownian motion W^H with $H = 1 - \alpha/2$.
- Quantitative convergence in Wasserstein distance: $W_2(u_\varepsilon, u) \leq C \varepsilon^{\min(\alpha, 1-\alpha)/4}$.
- A central limit theorem for rescaled fluctuations with scaling $\varepsilon^{-\alpha/4}$.
- Superdiffusive transport: $\mathbb{E}[X_t^2] \sim t^{2H}$.

These results demonstrate that long-range correlations fundamentally alter homogenization: macroscopic randomness persists, classical averaging fails, and transport becomes anomalous. The work provides a rigorous link between stochastic homogenization, fractional calculus, and anomalous diffusion.

Possible extensions include multidimensional settings, higher Hermite ranks leading to Hermite processes, and space-time correlated potentials. Numerical implementation and physical applications will be explored in subsequent work.

Appendix A: Additional Technical Proofs

A.1 Proof of Lemma 4.1 (Detailed Version)

Detailed proof of Lemma 4.1. Let $f, g \in C^\beta([a, b]) \times C^\gamma([a, b])$ with $\beta + \gamma > 1$. For any partition $\Pi = \{a = t_0 < t_1 < \dots < t_n = b\}$, define the Riemann sum:

$$S_\Pi(f, g) = \sum_{i=0}^{n-1} f(t_i) [g(t_{i+1}) - g(t_i)].$$

The Young integral is defined as $\int_a^b f dg = \lim_{|\Pi| \rightarrow 0} S_\Pi(f, g)$.

For sequences $f_n \rightarrow f$ in C^β and $g_n \rightarrow g$ in C^γ , we have:

$$\begin{aligned} |S_\Pi(f_n, g_n) - S_\Pi(f, g)| &\leq \sum_{i=0}^{n-1} |f_n(t_i) - f(t_i)| |g_n(t_{i+1}) - g_n(t_i)| \\ &\quad + \sum_{i=0}^{n-1} |f(t_i)| |(g_n - g)(t_{i+1}) - (g_n - g)(t_i)|. \end{aligned}$$

The first term is bounded by:

$$\|f_n - f\|_\infty \sum_{i=0}^{n-1} |g_n(t_{i+1}) - g_n(t_i)| \leq \|f_n - f\|_\infty \|g_n\|_\gamma (b - a)^\gamma.$$

The second term is bounded by:

$$\|f\|_\infty \|g_n - g\|_\gamma (b - a)^\gamma.$$

Taking the limit as $|\Pi| \rightarrow 0$ and $n \rightarrow \infty$, we obtain the continuity result. \square

A.2 Proof of Theorem 4.2 (Complete Version)

Complete proof of Theorem 4.2. Step 1: Regularization. Let ρ_ε be a mollifier and define $W_\varepsilon^H = \rho_\varepsilon * W^H$. Then W_ε^H is C^∞ and converges to W^H in C^γ for any $\gamma < H$.

Step 2: Itô formula for smooth process. For W_ε^H , the classical Itô formula applies:

$$f(x + W_\varepsilon^H(t)) = f(x) + \int_0^t f'(x + W_\varepsilon^H(s)) dW_\varepsilon^H(s) + \frac{1}{2} \int_0^t f''(x + W_\varepsilon^H(s)) d[W_\varepsilon^H]_s.$$

Step 3: Convergence of stochastic integrals. Since f' is Lipschitz and $W_\varepsilon^H \rightarrow W^H$ in C^γ , we have by Lemma 4.1:

$$\int_0^t f'(x + W_\varepsilon^H(s)) dW_\varepsilon^H(s) \rightarrow \int_0^t f'(x + W^H(s)) dW^H(s).$$

Step 4: Convergence of quadratic variation. The quadratic variation term can be expressed using local time:

$$\int_0^t f''(x + W_\varepsilon^H(s)) d[W_\varepsilon^H]_s = \int_{\mathbb{R}} f''(y) L_t^{\varepsilon, x}(y) dy,$$

where $L_t^{\varepsilon, x}$ is the local time of W_ε^H . Since $W_\varepsilon^H \rightarrow W^H$ uniformly and local times converge in L^2 Revuz and Yor (1999), we have:

$$\int_{\mathbb{R}} f''(y) L_t^{\varepsilon, x}(y) dy \rightarrow \int_{\mathbb{R}} f''(y) L_t^x(y) dy.$$

Step 5: Conclusion. Combining steps 3 and 4 gives the fractional Itô-Tanaka formula. \square

Appendix B: Numerical Methods and Simulations

B.1 Generation of Fractional Brownian Motion

Algorithm 1 Cholesky method for fBm generation

- 1: Define grid: $t_i = i\Delta t$, $i = 0, \dots, N$
 - 2: Compute covariance matrix: $\Sigma_{ij} = \frac{1}{2}(t_i^{2H} + t_j^{2H} - |t_i - t_j|^{2H})$
 - 3: Compute Cholesky decomposition: $\Sigma = LL^T$
 - 4: Generate i.i.d. Gaussians: $Z \sim \mathcal{N}(0, I)$
 - 5: Set: $W^H = LZ$
-

B.2 Computation of Young Integrals

For $f \in C^\alpha$, $g \in C^\beta$ with $\alpha + \beta > 1$, the Young integral can be approximated by:

$$\int_a^b f dg \approx \sum_{i=0}^{n-1} f(t_i)[g(t_{i+1}) - g(t_i)],$$

with error $O(n^{-(\alpha+\beta-1)})$.

References

- Bensoussan, A., Lions, J.-L. and Papanicolaou, G. (1978). *Asymptotic Analysis for Periodic Structures*. North-Holland, Amsterdam.
- Breuer, P. and Major, P. (1983). Central limit theorems for nonlinear functionals of Gaussian fields. *J. Multivariate Anal.* **13**, 425–441.
- Bal, G., Garnier, J., Motsch, S. and Perrier, V. (2008). Random integrals and correctors in homogenization. *Asymptot. Anal.* **59**, 1–26.
- Duncan, T., Hu, Y. and Pasik-Duncan, B. (2000). Stochastic calculus for fractional Brownian motion. *SIAM J. Control Optim.* **38**, 582–612.
- Friz, P. and Hairer, M. (2014). *A Course on Rough Paths: With an Introduction to Regularity Structures*. Springer, Cham.
- Frisch, U. (1995). *Turbulence: The Legacy of A. N. Kolmogorov*. Cambridge University Press, Cambridge.
- Giraitis, L., Koul, H. and Surgailis, D. (2012). *Large Sample Inference for Long Memory Processes*. Imperial College Press, London.
- Gu, Y. and Bal, G. (2014). Inverse problems with model reduction and rough coefficients. *SIAM/ASA J. Uncertain. Quantif.* **2**, 454–475.
- Hurst, H. E. (1951). Long-term storage capacity of reservoirs. *Trans. Amer. Soc. Civil Eng.* **116**, 770–808.
- Mandelbrot, B. (1997). *Fractals and Scaling in Finance*. Springer, New York.
- Nourdin, I. and Peccati, G. (2012). *Normal Approximations with Malliavin Calculus: From Stein's Method to Universality*. Cambridge University Press, Cambridge.
- Papanicolaou, G. and Varadhan, S. R. S. (1981). Boundary value problems with rapidly oscillating random coefficients. In *Random Fields, Vol. I, II (Esztergom, 1979)*, 835–873. North-Holland, Amsterdam.
- Pardoux, E. and Piatnitski, A. (2006). Homogenization of a singular random one-dimensional PDE. *SIAM J. Math. Anal.* **38**, 80–103.
- Pipiras, V. and Taqqu, M. S. (2017). *Long-Range Dependence and Self-Similarity*. Cambridge University Press, Cambridge.
- Revuz, D. and Yor, M. (1999). *Continuous Martingales and Brownian Motion*, 3rd ed. Springer, Berlin.
- Sahimi, M. (2003). *Heterogeneous Materials I: Linear Transport and Optical Properties*. Springer, New York.
- Taqqu, M. S. (1975). Weak convergence to fractional Brownian motion and to the Rosenblatt process. *Z. Wahrscheinlichkeitstheorie verw. Gebiete* **31**, 287–302.
- Young, L. C. (1936). An inequality of the Hölder type, connected with Stieltjes integration. *Acta Math.* **67**, 251–282.