

Phonon-assisted tunneling in Jahn-Teller $E \times e$ impurity centers in crystals

V. Hizhnyakov

Institute of Physics, University of Tartu, W. Ostwald Street 1, 50411 Tartu, Estonia

Abstract

Tunnel transitions between differently distorted states of impurity centers in crystals with the $E \times e$ Jahn-Teller effect are considered, taking into account the change in phonon dynamics during the transitions. Both linear and quadratic vibronic interactions are taken into account. It was found that the phonon scattering attending tunnelling, leads not only to a broadening of the energy spectrum of the transitions, but also to a deterioration in resonance. This strongly affects the temperature dependence of tunneling. The results obtained are consistent with measurements of ultrasound attenuation of Ni^{2+} ions in Al_2O_3 crystal. A notable feature of $E \times e$ tunneling is the existence of a window of values for quadratic vibronic interaction at which its effect disappears, and tunnelling remains coherent at sufficiently high temperatures.

1. Introduction

As shown by Jahn and Teller [1], symmetrical configurations of molecules are unstable in degenerate electronic states due to the interaction of valence electrons with vibrations of the molecule (vibronic interaction). Due to this instability the potential energy surfaces (PES) of molecules acquire several energetically equivalent minima of low symmetry separated by potential barriers. The states of the molecule in these minima, although they may be long-lived, usually are not completely stable: tunnel transitions between these states are possible [2-5]. At zero temperature, these transitions occur as coherent tunnel transitions and they can be considered as pseudo-rotations [3]. This holds also for impurity centers in crystals in degenerate electronic states [6-12]. However, in crystals, due to the presence of phonon continuum, tunnel transitions are accompanied by vibrational relaxation. As a result, as temperature increases, coherent tunnel motion is replaced by incoherent phonon-assisted tunnel transitions between the PES minima. Experimentally, these transitions can be manifested themselves as the phase relaxation, leading to temperature broadening of the zero phonon lines in optical spectra [13], attenuation of photon echo, and ultrasonic attenuation [14].

In this paper, tunnel transitions in a trigonal impurity center in a double degenerate electronic state of the E representation interacting with doubly degenerate vibrations of the e -representation are considered. Jahn-Teller effect in this case is a well-known archetype of the Jahn-Teller problem, called as the $E \times e$ problem. If only linear vibronic interaction is considered, then the PES has axial symmetry with a conical intersection and an equilateral circle of minima (the shape of the PES resembles a Mexican hat). However, the quadratic vibronic interaction leads to the violation of axial symmetry and to the formation of three energy-equivalent minima in the PES. The zero-point states in these minima are inherently unstable with respect to the under-barrier tunnel transitions. As a result, the ground state is split into one doubly degenerate tunnel state of E representation and one totally-symmetric tunnel state [2]. Usually, a doubly degenerate state is a ground state; however, the reverse order of the tunnel state is also possible if the quadratic vibronic interaction dominates [15,16]. Splitting is highly dependent on the strength of the vibronic interaction, rapidly decreasing as it increases (see Figs. 2 - 5 in [16], which present the results of calculating the energies of lower levels of the $E \times e$ state for different linear and quadratic vibronic interactions). In crystals, the upper tunnel state slowly decays with the emission of a phonon [17,18]. At higher temperatures, two-phonon relaxation processes of Raman type become significant, which leads to the replacement of coherent tunneling with incoherent tunneling.

Here we focus on incoherent tunneling and pay attention to the case of the strong Jahn-Teller effect. In this case, the phonon dynamics are greatly altered during the tunnel transition, followed by strong phonon mixing. To obtain the tunneling rate, we apply the method developed in [19,20] to

calculate vibronic transitions, which is valid for arbitrary linear and quadratic vibronic interactions. Using this method, it was possible to calculate the rate of tunnel transitions, taking into account, within the framework of the non-perturbative theory, two-phonon processes of the Raman type. It has been established that these processes lead not only to a broadening of the spectrum of tunnel transitions, but also to a change (increase) in the average frequency of the spectrum of these transitions. At low temperatures $T \ll T_D$, the width and the average frequency of the spectrum of tunnel transitions depend on temperature as $\propto T^7/T_D^7$ and $\propto T^4/T_D^4$, respectively, where T_D is the Debye temperature. As a result, the Raman type processes cause not only a weakening, but also deterioration of the resonance tuning. The reason for the change in average frequency lies in the zero-point vibrations, which make the creation of a phonon during the Raman process more likely than the destruction of a phonon of close frequency. As a result of both mentioned effects, the temperature dependence of the rate of the tunnel transitions at low temperatures became equal to $\Omega \propto T^{-1}$. The temperature dependences of the width and the mean frequency of the spectrum of the of tunnel transition are similar to those for zero-phonon lines (ZPL) in the optical spectra of impurity centers in crystals [21,22]. However, the reasons for the temperature dependence of the average frequency of tunnel transitions and ZPL are different: the average frequency of ZPL is mainly dependent on the first order quadratic vibronic interaction [23-25], but in tunnel transitions it is determined by the Raman processes described by quadratic vibronic interaction in the second order (the effect of the first-order quadratic interaction on tunneling in Jahn-Teller systems is negated due to symmetry arguments).

A notable phenomenon we found is that in the $E \times e$ case there is a window for the values of the quadratic vibronic interaction near the critical value $b_0 = 1/9$ (for the Debye frequency ω_D as a unit), at which the effect of this interaction on the tunnel transitions disappears. As a result, for such quadratic vibronic interaction, tunnel transitions remain coherent at a sufficiently high temperature. The reason is that in the $E \times e$ case quadratic vibronic interaction leads to the formation of tunnel barriers. As the barriers increase, vibrations along the trough in the PES minimums harden, but vibrations across the trough soften. Due to this, the change in phonon dynamics during the tunnel transition first decreases, then disappears at the critical value of the quadratic vibronic interaction b_0 , and then begins to increase with the increase in quadratic vibronic interaction.

At higher temperatures, transitions between different PES minimums with creation and destruction of multiple phonons became important, which leads to the replacement of a decrease in the rate of transitions with an increase in it with an increase in temperature [10-14]. At high temperatures, the transitions turn into the classical barrier jumps, described by Arrhenius' law [12]. Thus, the change in the tunneling rate from decreasing to increasing with decreasing temperature indicates the emergence of quantum tunneling. It should be noted, that in experiments on attenuation of ultrasound by $E \times e$ impurity centers of Ni^{2+} ions in Al_2O_3 crystals, such a change in the temperature dependences of the transition rate between different PES minima was actually observed at 2 K [14].

2. General

The vibronic Hamiltonian of the impurity center in the crystal in the $E \times e$ case is [9-12, 16-18]

$$H = H_0 \cdot I + V_1 + V_2 \quad (1)$$

Here $H_0 = (1/2) \sum_{ij} (\omega_j^2 x_{ji}^2 - \partial^2 / \partial x_{ji}^2)$ is the Hamiltonian of phonons of the crystal with the impurity center in the totally symmetric electronic state, $i = 1, 2$ is the line number of the e representation, $j = 1, 2, \dots, N$ is the phonon number, ω_j is frequency and x_{ij} is the normal coordinate of a e -phonon, $N \sim N_A$, N_A is the Avogadro's number, I is the 2×2 unit matrix, V_1 and V_2 are the linear and quadratic vibronic interactions, $\hbar = 1$. Using the electronic functions $|\mathcal{G}\rangle$ and $|\mathcal{E}\rangle$ of E representation, transforming as $3z^2 - r^2$ and $x^2 - y^2$, respectively, V_1 and V_2 get the form [9-12]

$$V_1 = a(-Q_1\sigma_x + Q_2\sigma_z), \quad V_2 = b((Q_2^2 - Q_1^2)\sigma_z + 2Q_1Q_2\sigma_x), \quad (2)$$

where σ_x and σ_z are Pauli matrices, a and b are the parameters of linear and quadratic vibronic interaction, respectively,

$Q_1 = (2R_1^x - R_2^x - R_3^x)/\sqrt{12} + (R_2^y - R_3^y)/2$, $Q_2 = (2R_1^y - R_2^y - R_3^y)/\sqrt{12} - (R_2^x - R_3^x)/2$ are symmetrized confogurational coordinates of the e representation, R_i^α ($\alpha = x, y$) is the Cartesian displacement in $\alpha = x, y$ direction of the corner atom (ion) number i belonging to one of three identical nearest atom (ion) to the impurity atom (ion). In the trigonal molecule AB_3 , the symmetrized coordinates Q_i are the normal coordinates. In contrast, in the case of an impurity center in a crystal, Q_i are not normal cootdinat but linear combinations of a huge number $\sim N_A$ of normal coordinates of the entire crystal, which are of the form [16-18]

$$Q_i = \sum_j \varsigma_{ji} x_{ji}. \quad (3)$$

Here $\varsigma_{ji} = \sqrt{M} \sum_{\alpha l} C_{il}^\alpha e_{ji}^\alpha$, M is the mass of the corner atoms of the AB_3 cluster, e_{ij}^α are the unite polarization vectors of phonons. The reduced displacement $r_i^\alpha = R_i^\alpha \sqrt{M}$ is linked to the normale coordinates as follows: $r_i^\alpha = \sum_j e_{ji}^\alpha x_{ji}$, where $\sum_j e_{ij}^\alpha e_{lj}^{\alpha'} = \delta_{\alpha\alpha'} \delta_{ll'}$, $\alpha = x, y, z$. The density of statates (DOS) of phonons contributing to the configurational coordinates of e – representation Q_i equals $\rho(\omega) = \sum_j (\varsigma_{ij}^\epsilon)^2 \delta(\omega - \omega_j)$.

Let us introduce the variables $Q = \sqrt{Q_1^2 + Q_2^2}$ and $\varphi = \arcsin(Q_1/Q)$ and consider the electronic states $|\psi_{1\varphi}\rangle = \cos\varphi|\mathcal{G}\rangle + \sin\varphi|\mathcal{E}\rangle$, $|\psi_{2\varphi}\rangle = -\sin\varphi|\mathcal{G}\rangle + \cos\varphi|\mathcal{E}\rangle$. These states bring the matrix of vibronic interaction into a diagonal form $V = \tilde{V}\sigma_z$, where [10]

$$\tilde{V} = -aQ\sqrt{1 + (b/a)Q\cos(3\varphi) + (b/a)^2Q^2}. \quad (4)$$

In the case $a > 0$ considered here, the lower energy brunch of the potential energy corresponds to the state $|\psi_{1\varphi}\rangle$. There are three values of φ angle, which correspond to PES minima. In the case $a, b > 0$, they are $\varphi_l = 2\pi(l-1)/3$, where $l = 1, 2, 3$ is the number of one of the theree PES minima.

Let us consider small vibrations at minimums of PES, and confine ourselves to the case $3(bQ_0/2a)^2 \ll 1$, where $Q_0 \sim a$ is the value of Q at the minimuma of the PES. (For important values of $b \leq 0.2$, the neglected term is at least 40 times smaler than the counted one; for the highest possible value $b = 0.5$ for a PES with a single conical intersection, it is 16 times smaller.) Then, near the PES minimum number l , $V_l \approx -aQ - (bQ^2/2)\cos(3\varphi_l)$, where the ratios

$$Q \cong Q_{1l} + Q_{2l}^2/(2Q_0), \quad Q^2 \cos(3\varphi) \cong Q_{1l}^2 - 9Q_{2l}^2/2 \quad (5)$$

are valid. Here $Q_{1,2;l} = \sum_j \varsigma_j (\cos(\varphi_l)x_{j1,2} \pm \sin(\varphi_l)x_{j2,1})$ stand for small displacements along ($Q_{1;l}$) and across ($Q_{2;l}$) the direction φ_l . Simple calculation gives

$$Q_0 \cong a\kappa/(1 + \kappa b), \quad \kappa = \sum_j \varsigma_{ij}^2/\omega_j^2. \quad (6)$$

Thus, for moderate values of b , the Hamiltonian of small vibrations near the minimum number l of PES, can be represented as

$$H_l \cong H_0 - aQ_{1;l} + b_1Q_{1;l}^2/2 + b_2Q_{2;l}^2/2, \quad (7)$$

where $b_1 = -b$, $b_2 = 7b/2 - \kappa^{-1}$. Using the shift operator e^{∇_l} , we obtain $H_l \cong e^{\nabla_l} \tilde{H}_l e^{-\nabla_l} - E_{JT}$, where, $E_{JT} = \kappa \bar{a}^2/2$ is the Jahn-Teller stabilization energy,

$$\tilde{H}_l \cong H_0 + b_1Q_{1;l}^2/2 + b_2Q_{2;l}^2/2, \quad (8)$$

$\nabla_l = \sum_j x_{0jl} (\cos(\phi_l) \partial/\partial x_{j1} + \sin(\phi_l) \partial/\partial x_{j2})$, $x_{0jl} = \bar{a} \zeta_{jl} / \omega_j^2$ is the equilibrium position of the normal coordinate x_{ji} in the PES minimum number l .

The vibrations of the impurity center near PES minima along and across the direction to the origin given by the Hamiltonians (8) are different. They also differ for different PES minima. The difference is especially large in the case of small b , when b_2 is close to $-\kappa^{-1}$ and the PES has the shape of a slightly deformed Mexican hat with deep trough and three shallow minima. In this case, the spectrum of Q_1 vibrations taking place across the trough (along the direction to the origin) practically does not changed, but the Q_2 vibrations taking place along the trough (across the direction to the origin) became soft with the amplitude of their zero-point vibrations being comparable to Q_0 , and small frequencies [26] (with the average frequency of order $\omega_0 \sim \sqrt{b}$). With an increase in b , the difference $b_1 - b_2$ in the quadratic vibronic interaction for both vibrations decreases, and in the critical value $b = b_0 = 2/(9\kappa)$ it disappears giving $\tilde{H}_l = \tilde{H}_{l'}$ ($l, l' = 1, 2, 3$). For this critical value of b , the phonon Hamiltonians in different PES minima have different equilibrium positions, but the same frequencies. With larger b , Q_1 vibrations become softer than Q_2 vibrations.

It should be emphasized that the equation (8) for phonon Hamiltonians is valid for the dominant linear vibronic interaction when $Q_0 \gg 1$ and $b < 1/2$. In this case, the PES has three minima and one conical intersection; for large b values, this may not be the case: three additional conical intersections can come to the one of the center [15]. In addition, in this case, the amplitude of zero-point vibrations $x_0 \sim 1/\sqrt{2\omega_0}$ is small as compared to the distance between the minima. Corresponding condition reads $b \gg (\hbar\omega_D/12E_{JT})^2$. If we assume $E_{JT}/\hbar\omega_D \sim 5$, then $b > 3 \cdot 10^{-4}$.

3. Phonons in different potential energy minima

To quantitatively characterize the change in the dynamics of phonons caused by the Jahn-Teller effect, we use the phonon Green's functions method [27]. According to this method, a change of the local elastic spring by b' leads to a change in the phonon Green's function, described by the following equation: $G'(\omega) = G(\omega) (1 - b'G'(\omega))$. Here

$$G(\omega) = \int_0^\infty d\omega' \rho(\omega') / (\omega^2 - (\omega' - i0_+)^2) \quad (9)$$

is the Green function of unperturbed phonons contributing to the corresponding elastic spring. According to this equation, the Green functions of phonons in the PES minimum for vibrations along ($i = 1$) and across ($i = 2$) the direction to the origin are equal to $G_i(\omega) = G(\omega)/(1 - b_i G(\omega))$.

The quantum tunneling considered here occurs at low temperatures, when only low-frequency acoustic phonons are present in the crystal. To take these phonons into account, we use the Debye-Van Hove model. In this model, the DOS of phonons, contributing to the reduced displacements R_i^α , equals to $\rho_0(\omega) = 8\pi^{-1}\omega^2\sqrt{1-\omega^2}$ (the Debye frequency ω_D is taken as a unit). To find the DOS of acoustic phonons that contribute to tunnel transitions, keep in mind that only even plane waves

contribute to Q_i . These waves have zero amplitude in the central atom and $\propto \omega$ amplitude in the neighboring (corner) atoms. Therefore, in this model, the DOS involved in tunnel transitions has an additional multiplier ω^2 , which gives $\rho(\omega) = 32\pi^{-1}\omega^4\sqrt{1-\omega^2}$. (Note, that in [17,18] this DOS was used to describe the phonon-induced relaxation of $E \times e$ Jahn-Teller systems; in [20] it was used to find the characteristics of ZPL in the optical spectra of impurity centers in crystals.) The corresponding Green function of unperturbed phonons is as follows:

$$G(\omega) = -2 - 8\omega^2 - 16\omega^3 \left(\sqrt{\omega^2 - 1 - i0_+} - \omega \right). \quad (10)$$

In this model $\kappa = -G(0) = 2$ and $b_0 = 1/9$.

The case of almost purely linear vibronic interaction corresponds to a significant a , small $b_1 \cong -b$, and negative and close to $-1/\kappa = -1/2$ b_2 , when the shape of PES has a slightly deformed Mexican hat with three shallow minima. In this case, the frequencies of the vibrations along the PES trough are small. At the same time, the frequencies of vibrations in the direction of the origin practically do not change. This is clearly seen in numerical calculations of DOS of perturbed phonons $\Gamma_i(\omega) \equiv \text{Im}G_i(\omega)$ (see Fig. 1): in the case of very small b , low frequency pseudolocal vibration is manifested in the spectrum of Q_2 vibrations occurring along the trough (across the direction to the origin). The frequency of this mode equals $\omega_0 \cong 1.5\sqrt{b}$. Unlikely to that, the spectrum of phonons, involved in the Q_1 vibrations occurring across the trough (along the direction to the origin) is not much changed due to the Jahn-Teller effect. However, with increasing b , the difference in $|b_1 - b_2|$ and in phonon spectra across and along the direction to the origin decreases, and at $b_0 = 2/(9\kappa) = 1/9$ it disappears. With larger b , Q_2 vibrations become even harder than Q_1 vibrations. For $b > 1/6$ local mode for Q_2 vibrations with the frequency above the phonon spectrum appears in the phonon spectrum. For b close to κ^{-1} (but less than κ^{-1}), in addition to the hard local Q_2 mode, a low frequency pseudolocal Q_1 mode also appears in the phonon spectrum. A strong difference in DOS of vibrations at PES minima in different directions leads to a strong change in the phonon dynamics during a tunnel transition, followed by strong phonon mixing. However, this difference disappears if b approaches b_0 .




Fig. 1. DOS of phonons in the $E \times e$ case contributing to tunnel transitions: Γ_1 (brown line) is the DOS of vibrations along the direction to the origin, Γ_2 (blue line) is DOS of vibrations across the direction to the origin. Left figure corresponds to quadratic vibronic interaction $b = 0.02$, left middle to $b = 0.109$, right middle to $b = 1/6$ and right to $b = 0.35$. Note that the distinction between Γ_1 and Γ_2 disappears in the vicinity of $b_0 = 1/9$. At $b = 0.35$, there is a local mode for vibrations across the direction to the origin with a frequency of 1.2 (the Debye frequency is taken as a unit of frequency).

4. Rate of tunnel transitions. Using Stratonovich's identity

To find the rate of tunnel transitions between two minima of PES, we apply Fermi's golden rule. We get the following equation for this rate [10]: $\Omega = 2\pi \left\langle \sum_f |H_{ll',nf}|^2 \delta(E_{l'n} - E_{l'f}) \right\rangle$, where $H_{ll',nf} \approx (E_{JT}/2) \langle \Psi_n^{(l)} | \Psi_f^{(l')} \rangle$ is the non-diagonal matrix element of the Hamiltonian for the initial ($|\Psi_n^{(l)}\rangle$) and final ($|\Psi_f^{(l')}\rangle$) phonon state of the system in the minimum number l and the minimum number l' , respectively. Using the integral representation for the delta-function $\delta(x) = \int e^{ixt} (dt/2\pi)$ and the Schrödinger equation, the transition rate can be presented as [28]

$$\Omega \approx (E_{JT}/2)^2 \int dt F(t), \quad (11)$$

where $F(t) = \langle e^{iH_f t} e^{-iH_l t} \rangle$ is the Fourier transform of the energy spectrum of the tunnel transitions, $\langle \dots \rangle$ is the quantum-statistical averaging over the initial vibrational states (here and below integrals are performed from $-\infty$ to ∞ , unless otherwise noted). The equation (11) to the constant factor is analogous to the equation for the optical spectrum of an impurity center [28]; the difference is that in the case of tunneling, H_l and $H_{l'}$ describe energy equivalent states, while in the case of optical transition there is no such equivalence. In real crystals, due to presence of the inhomogeneous field resulting from crystal defects, the PES minima have a slightly different energy. The influence of this field can be taken into account by adding the factor $e^{-\gamma_0|t|}$ in the Fourier integral in equation Eq. (11), where γ_0 is the width of the distribution of the energy differences in the PES minima. Equation (11) holds if the tunnel transitions are incoherent. This is the case when the spectral width of these transitions is large as compared to the tunnel splitting.

Using the shift operator e^{∇_l} , $F(t)$ can be represented as $F(t) = \langle e^{\nabla_l} e^{i\tilde{H}_l t} e^{-i\tilde{H}_l t} e^{\nabla_{l'}(t)} e^{-\nabla_{l'}} \rangle$, where $\nabla_{l'} = \nabla_l - \nabla_{l'}$, $\nabla_{l'}(t) = e^{i\tilde{H}_l t} \nabla_{l'} e^{-i\tilde{H}_l t}$ is the time-dependent operator $\nabla_{l'}$. At low temperatures, the largest contribution to Ω comes from zero-phonon transitions, having small spectral width [22,23]. This contribution is given by $F(t)$ in the large t limit. In this limit the operators at strongly different times decouple [23]. This gives $F(t)_{t \rightarrow \infty} = \langle e^{i\tilde{H}_l t} e^{-i\tilde{H}_l t} \rangle \langle e^{\nabla_{l'}} \rangle \langle e^{-\nabla_{l'}} \rangle$. Using the Bloch-de Dominicis theorem, we obtain $\langle e^{\pm \nabla_{l'}} \rangle_l = e^{-Z/2}$, where $Z = -\langle \nabla_{l'}^2 \rangle$ is the Huang-Rhys (Ham) factor, which describes a decrease in the probability of zero-phonon transitions due to changes in the equilibrium positions of atoms during the transition. As a result, we get $F(t)_{t \rightarrow \infty} = e^{-Z} \tilde{F}(t)_{t \rightarrow \infty}$, where

$$\tilde{F}(t) = \langle e^{i\tilde{H}_l t} e^{-i\tilde{H}_l t} \rangle \quad (12)$$

is the Fourier transform in the case of purely quadratic vibronic interaction.

Usually, $\tilde{F}(t)$ is calculated using a cumulant decomposition with respect to the quadratic vibronic interaction and taking into account up to the second order terms [22,23]. This is only correct if the quadratic vibronic interaction is weak. The asymptotic value $\tilde{F}(t)_{t \rightarrow \infty}$ can also be found for the arbitrary strength of the quadratic vibronic interaction by summing the cumulants if only one configurational coordinate contributes to this interaction [29-31] (in Ref. [32], the ZPL width in a special case of two configurational coordinates was also considered). However, in the case of tunnel transitions under consideration, there are at least two pairs of configuration coordinates describing vibrations along and across the direction to the origin at each PES minimum that contribute to tunnel transitions.

To find $\tilde{F}(t)_{t \rightarrow \infty}$ in the case of multiple configurational coordinates involved in the quadratic vibronic interaction, we apply a method developed in [19,20] to compute vibronic transitions, which does not use cumulant decomposition. This method is based on the use of the Dyson relation $e^{i\hat{H}_I + V} \cong \hat{T} e^{i\sum_n V(t_n)} e^{i\hat{H}_I}$ and 34560/ Here $t_n = nt/N$, $n = 1, 2, \dots, N \gg 1$, \hat{T} arranges the operators $V(t_n)$ from left to right according to the increasing t_n . We take $l = 1$ and $l' = 2$. Then $V = \bar{b}(Q_1^2 - \tilde{Q}_2^2)/2$, where $\bar{b} = b_2 - b_1$, $Q_i \equiv Q_{i,1}$, $\tilde{Q}_2 = (\sqrt{3}Q_2 - Q_1)/2$ (Q_1 and \tilde{Q}_2 are not orthogonal). This gives

$$\tilde{F}(t) \cong \left\langle \hat{T} e^{i\bar{b}/2N \sum_n (Q_1^2(t_n) - \tilde{Q}_2^2(t_n))} \right\rangle. \quad (13)$$

Using the Stratonovich identity, we can present $\tilde{F}(t)$ in the form

$$\tilde{F}(t) \cong \left(\prod_n (2\pi)^{-1} \int \int du_{1n} du_{2n} e^{-(u_{1n}^2 + u_{2n}^2)/2} \right) \left\langle \hat{T} e^{\sqrt{i\bar{b}/N} \sum_n (Q_1(t_n)u_{1n} - \tilde{Q}_2(t_n)u_{2n})} \right\rangle \quad (14)$$

The quantum-statistical averaging can be performed using the Bloch–De Dominicis theorem. We get

$$\tilde{F}(t) = \prod_n \left(\frac{1}{2\pi} \int \int du_{1n} du_{2n} e^{-(u_{1n}^2 + u_{2n}^2)/2} \right) \exp \left[\frac{-\bar{b}t}{8N} \sum_{nn'} \left(D_{nn'}^{(1)} u_{1n} (4u_{1n} + iu_{2n'}) - (D_{nn'}^{(1)} + 3D_{nn'}^{(2)}) u_{2n} u_{2n'} \right) \right].$$

Here

$$D_{nn'}^{(i)} \equiv D_i(t_n - t_{n'}) = i \left\langle \hat{T} Q_i(t_n) Q_i(t_{n'}) \right\rangle \quad (15)$$

is the causal correlation function of coordinate Q_i .

To find $\tilde{F}(t)_{t \rightarrow \infty}$, we take into account that the correlation function $D^{(i)}(t_n - t_{n'})$ tends to zero if $|t_n - t_{n'}| \rightarrow \infty$ [23]. As a result, the matrix elements $D_{nn'}^{(i)}$ in the $|t_n - t_{n'}| \rightarrow \infty$ limit depend on the difference $n - n'$. Therefore the eigenvectors and eigenvalues of the matrix $\hat{D}^{(i)}$ in the $|t_n - t_{n'}| \rightarrow \infty$ limit are equal to [19] $S_{kn} = N^{-1/2} e^{i\pi(2k+1)n/N}$ and $D_k^{(i)} = \sum_m e^{-i\omega_k m/N} D_i(tm/N)$, respectively, where $\omega_k = \pi(2k+1)/t$, $k = -N_0, -N_0 + 1, \dots, N_0$, $N = 2N_0$. Replacing variables u_{in} by $x_{ik} = \sum_n S_{kn} u_{in}$, we get

$$\tilde{F}(t)_{t \rightarrow \infty} = \prod_k (2\pi)^{-1} \int \int dx_{1k} dx_{2k} \exp \left[-(x_{1k}^2 + x_{2k}^2)/2 + (\bar{b}t/2N) \sum_{ii'} D_{ii',k} x_{ik} x_{i'k} \right]. \quad (16)$$

Here $D_{11,k} = D_k^{(1)}$, $D_{22,k} = -(D_k^{(1)} + 3D_k^{(2)})/4$, $D_{12,k} = D_{21,k} = D_k^{(1)}/4$. Integration over variables x_{1k} and x_{2k} gives $\exp(-2^{-1} \ln |I + t\bar{b}D_k/N|)$, where $|\dots|$ is the determinant of the 2×2 matrix. Having calculated this dererminant, we get

$$\ln \tilde{F}(t)_{t \rightarrow \infty} = -\frac{1}{2} \sum_k \ln \left(\left(1 - \frac{\bar{b}t D_k^{(1)}}{N} \right) \left(1 + \frac{\bar{b}t (D_k^{(1)} + 3D_k^{(2)})}{4N} \right) + \left(\frac{\bar{b}t D_k^{(1)}}{4N} \right)^2 \right). \quad (17)$$

Let us replace the sums over k and m with the integrals over ω and t' : $\sum_k \rightarrow \int dk = (t/2\pi) \int d\omega$, $\sum_m \rightarrow \int dm = (N/t) \int dt'$. We get

$$F(t)_{t \rightarrow \infty} = e^{-(\gamma + i\delta)t - Z}, \quad (18)$$

where

$$\gamma - i\delta = \frac{1}{2\pi} \int_0^\infty d\omega \ln \left((1 - \bar{b}D_1(\omega)) \left(1 + \frac{\bar{b}}{4} (D_1(\omega) + 3D_2(\omega)) \right) + \frac{\bar{b}^2}{4} D_1^2(\omega) \right) \quad (19)$$

describes the spectral width (γ) and the average value of the frequency of the tunnel transitions (δ), $D_i(\omega) = \int dt e^{i\omega t} D_i(t)$ is the spectral representation of the correlation function $D^{(i)}(t)$. Note that in the case of single configurational coordinate ($D_1(\omega) \neq 0$, $D_2(\omega) = 0$) equation (19) coincides, up to notation, with Levenson's equation (28) in [29] and equation (12.80) in [31] obtained using the cumulant decomposition of the Fourier transform.

5. Temperature dependence of incoherent tunnel transitions

Substituting equation (18) into (12) and adding inhomogeneous width γ_0 to γ , we obtain the following equation for the rate of incoherent tunnel transitions:

$$\Omega \approx (E_T/2)^2 e^{-Z} (\gamma + \gamma_0) / (\delta^2 + (\gamma + \gamma_0)^2). \quad (20)$$

This equation holds true if the spectral width of the tunnel transitions $\gamma + \gamma_0$ is larger than the rate $\Gamma = \Delta/3$ of coherent tunnel transitions (here Δ is tunnel splitting). To calculate Ω , we need to know the spectral functions $D_i(\omega)$. After simple calculations, we get

$$D_i(\omega) = G_i(\omega) + in_{\omega} \Gamma_i |\langle \omega \rangle|, \quad (21)$$

where $n_{\omega} = 1/(e^{\omega/k_B T} - 1)$ is the Planck population factor. At zero temperature $D_i(\omega)_{T=0} = G_i(\omega)$. Taking into account the relations $(1 - \bar{b}G_1)(1 + \bar{b}G_2) = 1$ and $(1 - \bar{b}G_1)G_2 = G_1$, we get

$$\ln \left((1 - \bar{b}G_1) \left(1 + (\bar{b}/4) (G_1 + 3G_2) \right) + (\bar{b}^2/4) G_1^2 \right) = 0.$$

As a result, at zero temperature γ and δ get to zero, as it should be for physical reasons: the spectrum of zero-phonon transitions between the zero-point states of the energy-equivalent minimums of PES does not broaden or shift.

Let us consider now γ and δ for nonzero but low temperature $T \ll T_D$. Keeping in Eq. (19) the terms linear with respect to n_{ω} , we obtain

$$\gamma \cong (3\bar{b}^2/4\pi) \int d\omega n_{\omega} \Gamma_1(\omega) \Gamma_2(\omega), \quad (22)$$

$$\delta \cong 3(\bar{b}/\pi) \int d\omega n_{\omega} \left(\Gamma_1(\omega) (1 - \bar{b}\Delta_2/4) - \Gamma_2(\omega) (1 - \bar{b}\Delta_1/4) \right), \quad (23)$$

where $\Delta_i(\omega) = \text{Re}G_i(\omega)$. For low frequency of phonons, $\Gamma_1 \cong 16\omega^3/(1 - 2b)^2$, $\Delta_1 \cong -2/(1 - 2b)$, $\Gamma_2 \cong 16\omega^3/(49b^2)$, $\Delta_2 \cong -2/(7b)$. We get (in ω_D units)

$$\gamma = \frac{34560\zeta(7)(1 - 9b)^2}{49\pi^2(1 - 2b)^2b^2} \left(\frac{T}{T_D} \right)^7, \quad (24)$$

$$\delta = \frac{2(29b + 3)(1 - 9b)^2\pi^3}{245b^2(1 - 2b)^2} \left(\frac{T}{T_D} \right)^4, \quad (25)$$

where $\zeta(7) = 1.008$ ($\zeta(x)$ is zeta-function). Here γ describes the width of the spectrum of tunnel transitions caused by the two-phonon (Raman) processes. At low temperatures the width of the spectrum increases with a temperature according to the law $\gamma \propto T^7/T_D^7$. Another effect of two-phonon (Raman) processes is a change δ (increase) in the average frequency of the spectrum of tunnel transitions with temperature that at low temperatures depends on the temperature as $\delta \propto (T/T_D)^4$. These dependences of γ and δ on temperature are similar to the temperature dependences of the width and the average frequency ZPL in the optical spectrum of the impurity center at low temperature [21-23]. However, the reasons for the temperature dependence of the average frequency of tunnel transitions and ZPL are different: the ZPL average frequency is determined mainly by the quadratic vibronic interaction in the first order [22,23]; but in tunnel transitions it is given by the quadratic vibronic interaction in the second order. The reason for the latter change lies in the zero-point vibrations, which makes the creation of a phonon more likely than the destruction of a phonon of the same frequency. The effect of the first-order quadratic vibronic interaction on frequency of tunneling in Jahn-Teller systems is negated due to symmetry arguments.

As a result of above two effects, in a crystal with a weak inhomogeneous field, the rate of incoherent tunnel transitions at low temperatures has the form

$$\Omega \approx \frac{1.05b^2(1-2b)^2 e^{-Z} E_{JT}^2}{(b+3/29)^2(1-9b)^2 \hbar^2 \omega_D} \frac{T_D}{T}. \quad (26)$$

Here $Z = \bar{a}^2 \int d\omega \omega^{-2} \Gamma_1(\omega) (2n_\omega + 1)$ is the Hung-Rhys factor, which at low temperature is equal to

$$Z \approx \frac{16E_{JT} (1 + \pi^2(T/T_D)^2)}{3\pi \hbar \omega_D (1-2b)^2}. \quad (27)$$

According to (26), the rate of tunnel transitions Ω decreases with an increase in temperature. If temperature is so low that the width γ is less than the inhomogeneous width γ_0 , then Ω decreases with temperature in accordance with the law $\Omega \propto (T^8 + \text{Const})^{-1}$ (see Fig. 2, left). At higher temperatures, for which $\gamma > \gamma_0$ the decrease becomes inversely linear: $\Omega \propto T^{-1}$. At even higher temperatures, multiphonon processes, which were not taken into account here, become important, causing an increase in Ω with increasing temperature. Finally, these transitions turn into classical barrier hops described by Arrhenius' law [12]. Thus, the decrease in Ω with increasing T is a characteristic feature of quantum tunnelling, distinguishing it from the classical hop processes, which intensify with increasing temperature. Note that the change in the transition rate between different PES minima from decrease to increase with increasing temperature was observed in [14] at a temperature of 2 K in experiments on ultrasound attenuation by $E \times e$ impurity centers of Ni^{2+} ions in Al_2O_3 crystals.

The dependences of the width γ and the average frequency δ on the quadratic vibronic interaction b are presented in Fig. 2 (right). It follows from this figure that these quantities decrease greatly with decreasing $|b - b_0|$ and reach zero at $b = b_0$. The reason is that in the $E \times e$ case with linear vibronic interaction, the PES has the shape of a Mexican hat with axial symmetry and with an equilateral circle of minima. However, the quadratic vibronic interaction leads to the formation of three minima in PES. In the case of small b , the minima are shallow, and the vibrations across and along the Jahn-Teller distortion vary greatly: the first ones are much softer than the previous ones. As b increases, the barriers also increase, and the vibrations across the distortion harden, but the vibrations along the distortion soften. As a result, the difference in the vibrations along and across the Jahn-Teller distortions decreases with increasing b . Accordingly, the change in the dynamics of phonons during the tunnel transition

also decreases, which leads to a decrease in γ and δ . For $b = b_0 = 1/9$, each PES minimum became symmetric. At the same time, the spectra of vibrations in the minima have become equal and only the positions of the minima remain different. In this case, both γ and δ are zero. With larger b , the difference in the phonon spectra appears again and begins to increase; at the same time, γ and δ also increase.




Fig. 2. The rate (Ω), average frequency (δ) and spectral width (γ) of tunnel transitions in the Jahn-Teller $E \times e$ impurity center in the crystal. Left: the dependence of $\log_{10} \Omega$ on temperature for $b = 0.05$, $Z=7$ and for inhomogeneous widths $\gamma_0 = 10^{-5}$, $\gamma_1 = 10^{-4}$, $\gamma_2 = 10^{-3}$, $\gamma_3 = 10^{-2}$; right: the dependence of $\log_{10}(\gamma)$ (lower line) and $\log_{10}(\delta)$ (upper line) on the quadratic vibronic interaction b (in ω_D units), for $T = 0.05T_D$.

We can see that there is a window for the values of the quadratic vibration interaction near $b_0 = 1/9$, in which the effect of this interaction on the tunnel transitions is turned off. Indeed, in the range of $b \approx b_0$, the spectral width γ and average frequency of tunnel transitions δ become very small even at relatively high temperature, so that the tunnel splitting begins to exceed $\delta \propto (T/T_D)^4$ and $\gamma \propto (T/T_D)^7$ already at a sufficiently high temperature. As a result, for such quadratic vibronic interactions, tunnel transitions remain coherent even at a relatively high temperatures.

6. Summary

Incoherent tunnel transitions between differently distorted states of trigonal impurity centers in crystals with $E \times e$ Jahn-Teller effect are considered, proceeding from the assumption that the vibronic interaction with phonons is strong enough to destroy coherent tunnelling. It has been established that the Raman type processes accompanying tunnelling, caused by the quadratic vibronic interaction, lead not only to a broadening of the energy spectrum of transitions, but also to a deterioration in resonance. This greatly affects the temperature dependence of tunneling. At low temperatures $T \ll T_D$, the width and the mean frequency of the spectrum of tunnel transitions depend on temperature as $\propto T^7/T_D^7$ and $\propto T^4/T_D^4$, respectively, where T_D is the Debye temperature. The reason for the change in average frequency lies in the zero-point vibrations, which make the creation of a phonon during the Raman process more likely than the destruction of a phonon of close frequency.

At very low temperatures (when the width γ is less than the inhomogeneous width γ_0), the inverse rate Ω^{-1} increases with increasing temperature as $(T/T_D)^8 + \text{Const}$. At higher, but still low temperatures (for which $\gamma > \gamma_0$) this increase becomes linear: $\Omega^{-1} \propto T$. At even higher temperature, multiphonon processes become important, causing an increase in Ω with increasing

temperature. The decrease of the rate Ω with increasing temperature is the characteristic feature of the quantum tunnelling, distinguishing it from the classical hopping processes, which intensify with increasing temperature. It should be noted that in experiments on ultrasound attenuation by $E \times e$ impurity centers of Ni^{2+} ions in Al_2O_3 crystals, a change in the decrease to an increase in the rate of transition between different PES minima at a temperature 2 K was observed [14].

In the $E \times e$ case, a remarkable property of tunneling was discovered, which is that there is a window of values of the quadratic vibronic interaction near the critical value of $b_0 = 1/9$ (in units of the Debye frequency), at which its effect on tunnel transitions disappears, and tunneling remains coherent at sufficiently high temperatures. The reason of existence of this window is that the quadratic vibronic interaction results not only in the tunneling barriers, but also in the hardening of vibrations along the trough of the PES and in the softening of vibrations across the through of the PES. As a result, with a small quadratic vibronic interaction b , the change in phonon dynamics during the tunnel transition decreases with increasing b (see Fig. 1). This abnormal dependence takes place up to the critical value $b = b_0$, when the decrease is replaced by an increase. For the critical value of b , phonon Hamiltonians in different PES minima have different equilibrium positions but the same spectra. Therefore, the quadratic vibronic interaction at $b = b_0$ does not lead to a change in phonon dynamics during the tunnel transitions. As a result, Raman processes remain weak for $b \approx b_0$, and tunnel transitions remain coherent even at a relatively high temperatures.

The presented theory is limited to the case of strong linear vibronic interaction and moderate quadratic vibronic interaction, which corresponds to $a^2 \gg 1$ and $|b| < 1/2$ (in this case, the PES has three minima and one conical intersection). We also assume that the temperature is not very low, so that the width of the spectrum of tunnel transitions $\gamma_t \propto T^7$ (see Eq. (24) and Refs. [33,34]) is large compared to the tunnel splitting; otherwise, the tunnel transitions are coherent (however, at $b \approx b_0$, this temperature may be relatively high). Thus, the incoherent quantum tunnelling considered here can only occur at moderately low temperature.

Acknowledgements

This work was partially supported by the Estonian Research Council project PRG347 (2019 – 2023).

References

1. H.A. Jahn and E. Teller, Instability of polyatomic molecules in degenerate electronic states. I. Orbital degeneracy, Proc. R. Soc. London, Ser. A, **161** 220 (1937).
2. I.B. Bersuker, Sov. Phys. JETP **16** 933 (1963).
3. Mary C. O'Brien, Proc. R. Soc. A. **281** (1384) 323 (1964).
4. F.S. Ham, Phys. Rev. **138** A1727 (1965).
5. R. Englman, "The Jahn-Teller Effect in Molecules and Crystals", Wiley London (1972).
6. C.A. Bates "Jahn–Teller effects in paramagnetic crystals". Physics Reports **35** 187 (1978).
7. M.D. Sturge, The Jahn-Teller effect in Solids, Solid State Physics **20** 91 (1968).
8. P. Garcia-Fernandez, A. Trueba, M.T. Barrius, J.A. Aramburu, Miguel Mareno, "Dynamic and Static Jahn-Teller Effect in Impurities: Determination of Tunnel Splitting", in: Progress in Theoretical Chemistry and Physics (PTCP) **23** Springer Dordrecht (2011).
9. I. B. Bersuker and V. Z. Polinger, "Vibronic interactions in Molecules and Crystals", Springer-Verlag Berlin-Heidelberg (1989).
10. I.B. Bersuker, "The Jahn-Teller Effect", Cambridge University Press, Cambridge (UK) (2006).
11. Yu.E. Perlin, B.S. Tsukerblat, "Electron-Vibrational Interaction in Optical Spectra of Impurity Paramagnetic ions", Stinicca, Kishinev (1974) (in Russian).
12. Y. Toyozawa, "Optical processes in Solids", Cambridge University Press (2003).

13. V. Hizhnyakov, H. Kaasik, and I. Sildos, Phys. Stat. Sol. **234** 644 (2002)); V. Hizhnyakov, V. Boltrushko, H. Kaasik and I. Sildos, J. Chem. Phys. **119** 6290 (2003).
14. M.D. Sturge, J.T. Krause, E.M. Gyorgy, R.C. LeCraw, F.R. Merrit, Phys. Rev. **155** 218 (1967).
15. H. Koizumi, I. Bersuker, Phys. Rev. Lett. **83** 3009 (2000).
16. Kaja Pae, V. Hizhnyakov, J. Chem. Phys. **147** 084107 (2017).
17. Kaja Pae, V. Hizhnyakov, J. Chem. Phys. **141** 234113 (2014).
18. Kaja Pae, V. Hizhnyakov, J. Chem. Phys. **145** 064108 (2016).
19. V. Hizhnyakov, Imbi Tehver, Chem. Phys. Lett. **422** 299 (2006).
20. V. Hizhnyakov, Chem. Phys. Lett. **493** 19 (2010); **808** 140092 (2022).
21. D.E. McCumber and M.D. Surge, J. Appl. Phys. **34** 1682 (1963).
22. D.E. McCumber Math. Phys. **5** 508 (1964).
23. M.A. Krivoglaz, Zh. Eksp. Teor. Fiz. **46** 637 (1964).
24. K. Rebane, "Impurity Spectra of Solids", Plenum Press, New York, 1970.
25. O. Sild, K. Haller (Eds.), "Zero-Phonon Lines", Springer Berlin Heidelberg (1988).
26. G. I. Bersuker and V. Z. Polinger, Zh. Eksp. Teor. Fiz. **80** 1798 (1981).
27. A. A. Maradudin, E. W. Montroll and G. H. Weiss, Theory of Lattice Dynamics in the Harmonic Approximation, Academic, New York, 1963.
28. M. Lax, J. Chem. Phys. **20** 1752 (1952).
29. G.F. Levenson, phys. stat. Sol. **43** 739 (1971).
30. I.S. Osad'ko, Sov. Phys. Solid State **13** 974 (1971); **17** 2098 (1975).
31. I.S. Osad'ko, "Selective Spectroscopy of Single Molecules", Springer (2003).
32. D. Hsu, J.L. Skinner, J. Lumines. **37**, 331 (1987).
33. T. Holstein, S.K. Lyo, R. Orbach, in "Laser Spectroscopy of Solids", ed. By W.M. Yen and P.M. Selzer, Springer-Verlag Berlin Heidelberg (1986).
34. R.S. Meltzer, "Line Broadening Mechanisms and Their Measurements", in Optic Materials, G. Liu, B. Jacquier (Eds.) Berlin Heidelberg (2005).