

PLANAR $W^{1,1}$ -EXTENSION DOMAINS

PEKKA KOSKELA, TAPIO RAJALA, AND YI RU-YA ZHANG

ABSTRACT. We show that a bounded planar simply connected domain Ω is a $W^{1,1}$ -extension domain if and only if for every pair x, y of points in Ω^c there exists a curve $\gamma \subset \Omega^c$ connecting x and y with

$$\int_{\gamma} \frac{1}{\chi_{\mathbb{R}^2 \setminus \partial\Omega}(z)} ds(z) \leq C|x - y|.$$

Consequently, a planar Jordan domain Ω is a $W^{1,1}$ -extension domain if and only if it is a BV -extension domain, and if and only if its complementary domain $\tilde{\Omega}$ is a $W^{1,\infty}$ -extension domain.

CONTENTS

1. Introduction	2
2. Preliminaries	4
2.1. Hyperbolic geodesics	5
2.2. Whitney-type sets	6
2.3. John domains	7
2.4. Quasiconvex sets	9
3. Sufficiency for Jordan domains	12
3.1. Decomposition of the domain and the complementary domain	12
3.2. Definition of the extension	14
3.3. Absolute continuity	19
3.4. Proof of Theorem 3.1	20
4. Piecewise hyperbolic geodesics and cut-points	21
5. Sufficiency for simply connected domains	27
5.1. Decomposition of the complement	27
5.2. Definition of the extension	32
5.3. Continuity at one-sided points	36
5.4. Absolute continuity	36
5.5. Conclusion of the proof	40

Date: December 11, 2025.

2000 Mathematics Subject Classification. 46E35.

Key words and phrases. Sobolev space, extension, quasiconvexity.

The first has been supported by the Academy of Finland via the Centre of Excellence in Analysis and Dynamics Research (project No. 307333) and Finnish Centre of Excellence in Randomness and Structures (project No. 364210). The second author has been funded by the Academy of Finland grant No. 274372. The third author is funded by National Key R&D Program of China (Grant No. 2025YFA1018400 & No. 2021YFA1003100), NSFC grant No. 12288201 & No. 12571128 and the Chinese Academy of Science.

6. Necessity for simply connected domains	40
Appendix A. Capacity estimates	43
A.1. Conformal capacity	43
A.2. A variant of the Gehring-Hayman theorem	48
A.3. A lemma on quasiconvex sets	51
References	52

1. INTRODUCTION

Given a domain $\Omega \subset \mathbb{R}^2$, define for $1 \leq p \leq \infty$ the Sobolev space $W^{1,p}(\Omega)$ as

$$W^{1,p}(\Omega) = \{u \in L^p(\Omega) \mid \nabla u \in L^p(\Omega, \mathbb{R}^2)\},$$

where ∇u denotes the distributional gradient of u . One usually employs $W^{1,p}(\Omega)$ with the non-homogeneous norm $\|u\|_{L^p(\Omega)} + \|\nabla u\|_{L^p(\Omega, \mathbb{R}^2)}$ or with the homogeneous (semi-)norm $\|\nabla u\|_{L^p(\Omega, \mathbb{R}^2)}$.

We call $E: W^{1,p}(\Omega) \rightarrow W^{1,p}(\mathbb{R}^2)$ an extension operator for Ω if there exists a constant $C \geq 1$ so that for every $u \in W^{1,p}(\Omega)$ we have

$$\|Eu\|_{W^{1,p}(\mathbb{R}^2)} \leq C\|u\|_{W^{1,p}(\Omega)}$$

and $Eu|_{\Omega} = u$. A domain $\Omega \subset \mathbb{R}^2$ is called a $W^{1,p}$ -extension domain if there exists an extension operator $E: W^{1,p}(\Omega) \rightarrow W^{1,p}(\mathbb{R}^2)$. By [18], a bounded domain Ω admits an extension operator with respect to the above non-homogeneous norm if and only if there is a bounded extension operator $F: W^{1,p}(\Omega) \rightarrow W_{\text{loc}}^{1,p}(\mathbb{R}^2)$ with respect to the homogeneous semi-norm when $1 \leq p < \infty$.

Geometric properties of simply connected planar $W^{1,p}$ -extension domains are by now rather well understood, see [14], [19], [28] and [21]. In particular, a full geometric characterization for bounded simply connected planar $W^{1,p}$ -extension domains is available for $1 < p \leq \infty$. For the range $1 < p < 2$ bounded simply connected planar $W^{1,p}$ -extension domains were characterized in [21] as those bounded domains $\Omega \subset \mathbb{R}^2$ for which any two points $x, y \in \mathbb{R}^2 \setminus \Omega$ can be connected with a curve $\gamma \subset \mathbb{R}^2 \setminus \Omega$ satisfying

$$\int_{\gamma} \text{dist}(z, \partial\Omega)^{1-p} ds(z) \leq C|x - y|^{2-p}. \quad (1.1)$$

Regarding $p = 1$, in [20, Corollary 1.2] it was shown that the complement of a bounded simply connected $W^{1,1}$ -extension domain is quasiconvex. This was obtained as a corollary to a characterization of bounded simply connected BV -extension domains. Recall that a set $E \subset \mathbb{R}^2$ is called *C-quasiconvex* if there exists a constant $C \geq 1$ such that any pair of points $z_1, z_2 \in E$ can be connected by a rectifiable curve $\gamma \subset E$ whose length $\ell(\gamma)$ satisfies $\ell(\gamma) \leq C|z_1 - z_2|$.

Quasiconvexity of the complement does not imply $W^{1,1}$ -extendability in general. Indeed, a slit disk

$$\Omega = \mathbb{D} \setminus \{(x, 0) \mid x \geq 0\}$$

has a quasiconvex complement but Ω fails to be a $W^{1,1}$ -extension domain. In this paper we show that in order to obtain a characterization of $W^{1,1}$ -extendability in the spirit of the curve estimate (1.1), in addition to the quasiconvexity of the complement, we have to take into account the size of the set of self-intersections of the boundary $\partial\Omega$. Our main result is the following theorem.

Theorem 1.1. *Let $\Omega \subset \mathbb{R}^2$ be a bounded simply connected domain. Then Ω is a $W^{1,1}$ -extension domain if and only if there exists a constant $C < \infty$, quantitatively, such that for each pair $x, y \in \Omega^c$ of points there exists a curve $\gamma \subset \Omega^c$ connecting x and y with*

$$\int_{\gamma} \frac{1}{\chi_{\mathbb{R}^2 \setminus \partial\Omega}(z)} ds(z) \leq C|x - y|. \quad (1.2)$$

In other words, (1.2) requires that $\ell(\gamma) \leq C|x - y|$ and that $\mathcal{H}^1(\gamma \cap \partial\Omega) = 0$, where \mathcal{H}^1 denotes the 1-dimensional Hausdorff measure.

Both the necessity and sufficiency in Theorem 1.1 are new. Our construction of an extension operator actually gives a linear operator.

Corollary 1.2. *Let $\Omega \subset \mathbb{R}^2$ be a bounded simply connected domain. If Ω is a $W^{1,1}$ -extension domain, then Ω admits a bounded linear extension operator for $W^{1,1}$.*

Corollary 1.2 gives the first linearity result for Sobolev extension operators for $p = 1$ beyond concrete cases like Lipschitz domains. Up to now it has not been clear if linearity could be expected; recall that there is no bounded linear extension operator from the trace space $L^1(\mathbb{R})$ of $W^{1,1}(\mathbb{R}^2)$ to $W^{1,1}(\mathbb{R}^2)$. The existence of a bounded linear extension operator in the case of $p > 1$ was established in [16] for $W^{1,p}$ -extension domains via a particular method that cannot be applied for $p = 1$.

In the case of a Jordan domain we do not have boundary self-intersections and hence our characterization easily reduces to quasiconvexity of the complementary domain. From Theorem 1.1 and (the proof of) Lemma 2.12 we obtain the following corollary.

Corollary 1.3. *Let Ω be a planar Jordan domain. Then Ω is a $W^{1,1}$ -extension domain if and only if its complementary domain is quasiconvex.*

Corollary 1.3 together with earlier results (see e.g. [4], [32], [20]) yields the following somewhat surprising corollary.

Corollary 1.4. *Let Ω be a planar Jordan domain and $\tilde{\Omega}$ be its complementary domain. Then the following are equivalent:*

- (1) Ω is a $W^{1,1}$ -extension domain.
- (2) Ω is a BV-extension domain.
- (3) $\tilde{\Omega}$ is quasiconvex.
- (4) $\tilde{\Omega}$ is a $W^{1,\infty}$ -extension domain.

This corollary together with Lemma 2.1 below also extends the duality result from [21, Corollary 1.3] to the case of all $1 \leq p \leq \infty$.

Corollary 1.5. *Let $1 \leq p, q \leq \infty$ be Hölder dual exponents and let $\Omega \subset \mathbb{R}^2$ be a Jordan domain. Then Ω is a $W^{1,p}$ -extension domain if and only if $\mathbb{R}^2 \setminus \bar{\Omega}$ is a $W^{1,q}$ -extension domain.*

One further obtains a monotonicity property via Theorem 1.1 and [21]: if Ω is a bounded simply connected $W^{1,p}$ -extension domain with $1 < p < 2$, then it is also a $W^{1,q}$ -extension domain for all $1 \leq q \leq p$. However, a $W^{1,1}$ -extension domain need not be a $W^{1,p}$ -extension domain for any $1 < p < 2$. To see this consider the Jordan domain

$$\Omega = \mathbb{D} \setminus \{(x, y) \mid y \leq x^2\}.$$

The complement of Ω is clearly quasiconvex, but for any curve γ connecting the origin and the point $(t, 0)$ with $0 < t < 1$, we have

$$\int_{\gamma} \text{dist}(z, \partial\Omega)^{1-p} ds(z) \gtrsim \int_0^t x^{2-2p} dx \gtrsim t^{3-2p}.$$

Since there is no constant $C > 0$ such that $t^{3-2p} < Ct^{2-p}$ for all $0 < t < 1$ when $1 < p < 2$, (1.1) fails, and thus Ω is not an extension domain for $W^{1,p}(\Omega)$ when $1 < p < 2$.

Resulting from [21, Corollary 1.5] and the argument above, we conclude that the set of all $1 \leq s \leq \infty$ for which a bounded simply connected planar domain Ω is a $W^{1,s}$ -extension domain, can only be one of:

$$\emptyset, \{1\}, [1, q] \text{ with } q < 2, [1, \infty], (q, \infty] \text{ with } q > 2 \text{ or } \{\infty\}.$$

Let us close this introduction by briefly discussing the ideas of the proofs of Theorem 1.1 and Corollary 1.3. First of all, we cannot use the approach in [21] which gave a characterization for $1 < p < 2$. This is because there both the sufficiency and necessity of (1.1) were first proven for approximating Jordan domains and the general case was concluded via a limiting argument. Our condition (1.2) is not stable under (uniform) convergence of the complement and $W^{1,1}$ is not weakly compact. For sufficiency, an approximation procedure would only give BV -extension. Nevertheless, we first prove sufficiency in the Jordan case via a hyperbolic triangulation of the complementary domain instead of a usual Whitney decomposition. For the general case we construct a generalized hyperbolic triangulation for the complement of Ω and apply the Jordan result for suitable subdomains of Ω to show that the extension obtained via this triangulation is a Sobolev extension. Namely, when Ω is simply connected, it becomes necessary to manually construct the extension, drawing upon the approximation techniques employed in the Jordan case in Section 3. This process inevitably gives rise to a range of non-trivial technical challenges, and has been done in Sections 4 and 5. The necessity of (1.2) is proven directly in the simply connected case and the necessity in Corollary 1.3 is deduced from the general case.

2. PRELIMINARIES

In this section we recall some definitions and properties. The notation in our paper is quite standard. When we make estimates, we sometimes write the constants as positive real numbers $C(\cdot)$ with the parenthesis including all the parameters on which the constant depends. The constant $C(\cdot)$ may vary between appearances, even within a chain of inequalities. Especially, C refers to denote an absolute constant. By $a \lesssim b$ we mean $a \leq Cb$, and $a \sim b$ means that $b/C \leq a \leq Cb$ for some constant $C \geq 2$. The Euclidean distance between two sets $A, B \subset \mathbb{R}^2$ is denoted by $\text{dist}(A, B)$. By \mathbb{D} we always mean the open unit disk in \mathbb{R}^2 and by $\partial\mathbb{D}$ its boundary. We denote by $\ell(\gamma)$ the length of a curve γ . For a set $A \subset \mathbb{R}^2$, we

denote by A° its interior, ∂A its boundary, $A^c = \mathbb{R}^2 \setminus A$ its complement, and \overline{A} its closure respectively with respect to the Euclidean topology, unless another specific explanation is given. For an injective curve $\gamma: [0, 1] \rightarrow \mathbb{R}^2$ we use γ° to denote $\gamma((0, 1))$. The line segment connecting x and y is referred to by $[x, y]$, and for an injective curve γ with $x, y \in \gamma$, the subarc of γ joining x and y is denoted by $\gamma[x, y]$. We denote the 1-dimensional Hausdorff measure by \mathcal{H}^1 . For a disk $B \subset \mathbb{R}^2$ and a constant $c > 0$, we denote by cB the disk with the same center as B but a radius that is c times the radius of B .

We have the following swapping lemma.

Lemma 2.1. *Let $\Omega \subset \mathbb{R}^2$ be a bounded domain. Take $x \in \Omega$ and define an unbounded domain $\tilde{\Omega} = i_x(\Omega)$ using the inversion mapping*

$$i_x: \mathbb{R}^2 \setminus \{x\} \rightarrow \mathbb{R}^2 \setminus \{x\}: y \mapsto x + \frac{y - x}{|y - x|^2}.$$

Then

- (1) *The domain Ω is a $W^{1,1}$ -extension domain if and only if $\tilde{\Omega}$ is such a domain.*
- (2) *The domain Ω is quasiconvex if and only if $\tilde{\Omega}$ is such a domain.*

This lemma is obtained via the proof of [21, Lemma 2.1].

Proof. The first part of the lemma follows from the part (1) of [21, Lemma 2.1]. Thus we only need to show part (2).

Let $R = 2 \operatorname{diam}(\Omega)$ and $2r = \operatorname{dist}(x, \partial\Omega)$. Then $\partial\Omega \subset A(x, r, R) := B(x, R) \setminus \overline{B(x, r)}$.

Notice that i_x is a biLipschitz map when restricted to $A(x, r, R)$, with the biLipschitz constant only depending on r and R . Additionally, the fact that outside $A(x, r, R)$ and its image one can always connect using curves satisfying (1.2), imply claim (2). Indeed, if Ω has curves satisfying (1.2), then let x_1, x_2 be any two points in $\hat{\Omega}$, and let $z_1 = i_x^{-1}(x_1)$ and $z_2 = i_x^{-1}(x_2)$. Then $z_1, z_2 \in \Omega$. If the curve $\gamma \subset \Omega$ connecting them lies in $\overline{A(x, r, R)}$, then the bi-Lipschitz property of i_x directly gives the desired inequality for the curve $i_x \circ \gamma$ up to a multiplicative constant depending only on p, r and R .

Next if $z_1, z_2 \in A(x, r, R)$ but the corresponding curve is not contained in $\overline{A(x, r, R)}$, since $r = \operatorname{dist}(x, \partial\Omega)/2$, then we can replace the part of curve inside $B(x, r)$ by the shorter subarc of the circle $S^1(x, r)$ connecting the corresponding points on the circle $S^1(x, r)$. The new curve that we still denote by γ satisfies inequality (1.2) with a constant that only depends on the original constant and p . The desired inequality for the curve $i_x \circ \gamma$ follows by the argument in the previous case.

The case where $z_1, z_2 \in B(x, r)$ is trivial, since then x_1, x_2 are contained in the complement of a disk, and this complement is contained in $\hat{\Omega}$ and far away from $\partial\hat{\Omega}$. The case $z_1 \in B(x, r)$ while $z_2 \in A(x, r, R)$ follows easily from the combination of previous cases, and by symmetry we finish the proof of one direction. The other direction is similar. \square

2.1. Hyperbolic geodesics. Recall that the hyperbolic geodesics in \mathbb{D} and in $\mathbb{R}^2 \setminus \overline{\mathbb{D}}$ are defined as the arcs of (generalized) circles that intersect the unit circle orthogonally. Moreover, hyperbolic geodesics are preserved under conformal maps.

The following theorem was established by Gehring and Hayman [10]; see also e.g. [25, Theorem 4.20, Page 88].

Theorem 2.2. *Let Ω be a bounded simply connected domain, and $\varphi : \mathbb{D} \rightarrow \Omega$ be a conformal map. Given a pair of points $x, y \in \mathbb{D}$, denoting the corresponding hyperbolic geodesic in \mathbb{D} by $\Gamma_{x,y}$, and by $\gamma_{x,y}$ any arc connecting x and y in \mathbb{D} , we have*

$$\ell(\varphi(\Gamma_{x,y})) \leq C\ell(\varphi(\gamma_{x,y})).$$

One also has the following exterior version of the Gehring-Hayman theorem for Jordan domains. We record a proof in the Appendix.

Theorem 2.3. *Let $\tilde{\Omega}$ be the exterior of a Jordan domain. If $\tilde{\varphi} : \tilde{\Omega} \rightarrow \mathbb{R}^2 \setminus \overline{\mathbb{D}}$ is conformal and Γ is a hyperbolic geodesic in $\tilde{\Omega}$ connecting x and y satisfying $\tilde{\varphi}(\Gamma) \subset B(0, 100)$, and $\gamma \subset \tilde{\Omega}$ is any curve joining x and y , then we have*

$$\ell(\Gamma) \leq C\ell(\gamma)$$

for some absolute constant C .

According to Theorem 2.2, hyperbolic geodesics in a (bounded) simply connected domain are essentially the shortest possible curves joining given endpoints. For the exterior domain case, one needs to rule out the case of geodesics with large diameters; see Theorem 2.3.

We need the following lemma that follows from [25, Corollary 4.18]. Also see [5] and [2, Theorem 0.1], where it is called the “ball separation property”.

Lemma 2.4. *Let $\varphi : \mathbb{R}^2 \setminus \overline{\mathbb{D}} \rightarrow G$ be a conformal map, and $x, y \in G$ be a pair of points and Γ a hyperbolic geodesic joining them. If B is a disk with $B \cap \Gamma \neq \emptyset$ and of λ -Whitney type, then any curve γ connecting x, y in G has non-empty intersection with cB , where $c = c(\lambda)$.*

2.2. Whitney-type sets. Let us recall the following properties of Whitney-type sets.

Definition 2.5. *A connected set $A \subset \Omega \subset \mathbb{R}^2$ is called a λ -Whitney-type set in Ω with $\lambda \geq 1$ if the following holds.*

- (1) *There exists a disk with radius $\frac{1}{\lambda} \text{diam}(A)$ contained in A ;*
- (2) *$\frac{1}{\lambda} \text{diam}(A) \leq \text{dist}(A, \partial\Omega) \leq \lambda \text{diam}(A)$.*

Conformal mappings preserve Whitney-type sets in the following sense.

Lemma 2.6. *Suppose $\varphi : \Omega \rightarrow \Omega'$ is conformal, where $\Omega, \Omega' \subset \mathbb{R}^2$ are domains conformally equivalent to \mathbb{D} or $\mathbb{R}^2 \setminus \overline{\mathbb{D}}$, and $A \subset \Omega$ is a λ_1 -Whitney-type set. Then $\varphi(A) \subset \Omega'$ is a λ_2 -Whitney-type set with $\lambda_2 = \lambda_2(\lambda_1)$. Moreover, the restriction of the conformal map $\varphi|_A$ is a $L(\lambda_1)$ -biLipschitz map up to a dilation factor.*

Proof. The first claim follows, e.g., from [21, Lemma 2.12]. Thus we only need to verify the second claim.

To begin, for any two points $w_1, w_2 \in A$, by [21, Lemma 2.5] the hyperbolic distance between w_1 and w_2 is smaller than $C(\lambda_1)$. Denote by $\gamma[w_1, w_2]$ the hyperbolic geodesic joining w_1 and w_2 . Since the hyperbolic distance is realized by hyperbolic geodesics, we conclude that for any $z \in \gamma[w_1, w_2]$ the hyperbolic distance between z and w_1 is smaller than $C(\lambda_1)$. Fix $w_0 \in A$. Then [1, Theorem 2.10.8] implies that

$$|\varphi'(z)| \sim |\varphi'(w_1)| \sim |\varphi'(w_0)|,$$

where the constants depend only on λ_1 . Thus we have

$$|\varphi(w_1) - \varphi(w_2)| \leq \int_{\gamma[w_1, w_2]} |\varphi'(z)| dz \leq C(\lambda_1) |\varphi'(w_0)| \int_{\gamma[w_1, w_2]} 1 dz \leq C(\lambda_1) |\varphi'(w_0)| |w_1 - w_2|,$$

where the last inequality is a consequence of the observation that $\gamma[w_1, w_2]$ can be encompassed by no more than $C(\lambda_1)$ -many Whitney squares, and within each of these squares, the length of the hyperbolic geodesic is comparable to the length of the straight-line segment connecting its endpoints. Similarly, since φ^{-1} is also conformal and $\varphi(A)$ is of λ_2 -Whitney-type with $\lambda_2 = \lambda_2(\lambda_1)$, by the above estimate we get

$$|w_1 - w_2| \leq \int_{\gamma[\varphi(w_1), \varphi(w_2)]} |\varphi'(z)|^{-1} dz \leq C(\lambda_1) |(\varphi^{-1})'(\varphi(w_0))| |\varphi(w_1) - \varphi(w_2)|.$$

Since φ is conformal, we have $|\varphi'(w_0)| \neq 0$ and

$$|(\varphi^{-1})'(\varphi(w_0))| = |\varphi'(w_0)|^{-1}.$$

To conclude, we have that

$$C(\lambda_1)^{-1} |\varphi'(w_0)| |w_1 - w_2| \leq |\varphi(w_1) - \varphi(w_2)| \leq C(\lambda_1) |\varphi'(w_0)| |w_1 - w_2|,$$

as desired. \square

Sometimes we omit the constant λ when we are dealing with a λ -Whitney-type set whose constant is clear from the context.

2.3. John domains. Let us recall some known results about John domains. The relation to the curve condition (1.2) is given by part (1) of Lemma 2.8.

Definition 2.7 (John domain). *An open subset $\Omega \subset \mathbb{R}^2$ is called a John domain provided it satisfies the following condition: There exist a distinguished point $x_0 \in \Omega$ and a constant $J > 0$ such that, for every $x \in \Omega$, there is a curve $\gamma : [0, \ell(\gamma)] \rightarrow \Omega$ parameterized by the arc length, such that $\gamma(0) = x$, $\gamma(\ell(\gamma)) = x_0$ and*

$$\text{dist}(\gamma(t), \mathbb{R}^2 \setminus \Omega) \geq Jt.$$

The curve γ is called a John curve, and J is called a John constant.

We collect a number of results related to John domains in the following lemma.

Lemma 2.8. *The following statements hold:*

- (1) *Let $\Omega \subset \mathbb{R}^2$ be a bounded simply connected domain. If Ω^c is C -quasiconvex, then Ω is a J -John domain with $J = J(C)$.*
- (2) *Given a simply connected (bounded) J -John domain $\Omega \subset \mathbb{R}^2$ together with a conformal map $\varphi : \mathbb{D} \rightarrow \Omega$ with $\varphi(0) = x_0$, where x_0 is the distinguish point of Ω , we can extend φ continuously up to the boundary.*
- (3) *The hyperbolic geodesic connecting $x \in \Omega$ and x_0 in Ω is a J' -John curve, where $J' = J'(J)$.*
- (4) *The Lebesgue measure of the boundary of a simply connected (bounded) J -John domain $\Omega \subset \mathbb{R}^2$ is zero.*

Proof. (1) is from [23, Theorem 4.5]; observe that the C -quasiconvexity property is stronger than the corresponding C -bounded turning property employed in [23, Theorem 4.5], i.e. every pair of points $x, y \in \mathbb{R}^2 \setminus \Omega$ can be joined by a curve γ with $\text{diam}(\gamma) \leq C|x - y|$. (2) follows from [23, Theorem 2.18] and [24, Theorem 4.7, Page 441], and (3) comes from [9, Theorem 4.1].

To show (4), let x be an arbitrary point in $\partial\Omega$. Define $B_r = B(x, r)$ for some $0 < r < \frac{1}{2}|x - x_0|$, where x_0 is the John center of Ω . Choose a point $z \in B$ such that $|z - x| < r/4$, and denote by γ the hyperbolic geodesic from x_0 to z . By (4), γ is a $C(J)$ -John curve, and hence by choosing a point $y \in \gamma$ such that $|y - z| = r/2$, the definition of John domain, shows that there exists a constant $c = c(J)$ such that the disk $B(y, cr)$ is compactly contained in $B_r \cap \Omega$ for every $0 < r < \frac{1}{2}|x - x_0|$. Hence x cannot be a point of density of $\partial\Omega$. By applying the Lebesgue differentiation theorem to the characteristic function of $\partial\Omega$ we then conclude (5) by the arbitrariness of x . \square

In fact we have a stronger version of (2) of Lemma 2.8. Define the *inner distance* of Ω between $x, y \in \Omega$ by

$$\text{dist}_\Omega(x, y) = \inf_{\gamma \subset \Omega} \ell(\gamma),$$

where the infimum runs over all curves joining x and y in Ω . If $x \in \partial\Omega$ and $y \in \overline{\Omega}$ and $\gamma: [0, 1] \rightarrow \overline{\Omega}$ is continuous with $\gamma(0) = x$, $\gamma(1) = y$ and $\gamma((0, 1)) \subset \Omega$, we say that γ joins x and y in Ω . Furthermore, if $\ell(\gamma)$ is finite, then we say that x and y are rectifiably joinable in Ω and define $\text{dist}_\Omega(x, y)$ via $\inf_\gamma \ell(\gamma)$ over all such γ . The inner diameter $\text{diam}_\Omega(E)$ of a set $E \subset \overline{\Omega}$ is then defined in the usual way.

Lemma 2.9. *Let $\Omega \subset \mathbb{R}^2$ be a simply connected (bounded) J -John domain and let $\varphi: \mathbb{D} \rightarrow \Omega$ be conformal. Then $\varphi: (\mathbb{D}, |\cdot|) \rightarrow (\Omega, \text{dist}_\Omega)$ is uniformly continuous, where $|\cdot|$ denotes the Euclidean metric.*

Proof. By [17], $\varphi: (\mathbb{D}, |\cdot|) \rightarrow (\Omega, d_\Omega)$ is uniformly continuous, where for each pair of points $x, y \in \Omega$,

$$d_\Omega(x, y) := \inf_{\gamma \subset \Omega} \text{diam}(\gamma),$$

and the infimum runs over all curves joining x and y in Ω .

By [12, Theorem 5.14], for a simply connected (bounded) J -John domain Ω , we have

$$d_\Omega(x, y) \sim \text{dist}_\Omega(x, y)$$

for any pair of two points $x, y \in \Omega$, where the constant depending only on J . Thus the claim follows from the previous uniform continuity result. \square

Recall that the extended complex plane consists of the complex plane together with a point at infinity. As a complex manifold it can be described by two charts via the stereographic projections with $\frac{1}{z}$ as the transition map. Combining the lemma above with results in [7], we have the following conclusion.

Lemma 2.10. *Let $\Omega \subset \mathbb{R}^2$ be a bounded simply connected domain whose complement is quasiconvex. Then Ω^c is locally connected and each bounded component of its interior is a*

Jordan domain. (In fact, every component of the interior of Ω^c is Jordan on the Riemann sphere.)

Proof. By Lemma 2.8 we know that a Riemann map $\varphi: \mathbb{D} \rightarrow \Omega$ can be extended continuously to the boundary. Hence [7, Page 14, Corollary] shows that Ω^c is locally connected.

To show the second part of the lemma, we may assume that $0 \in \Omega$. Then by letting $g(z) = \frac{z}{|z|^2}$ we have that $g(\Omega^c)$ is compact, locally connected and its complement is connected (on the Riemann sphere). Therefore by applying [7, Page 17, Proposition 2.3] to $g(\Omega^c)$ we conclude the second part of the lemma; note that g is a homeomorphism on the Riemann sphere. \square

2.4. Quasiconvex sets. In this subsection we list some important properties of quasiconvex sets in the plane.

We begin with the following corollary of Theorem 2.3.

Corollary 2.11. *Let $\tilde{\Omega}$ be a complementary domain of some Jordan domain and C_1 -quasiconvex. If $\tilde{\phi}: \mathbb{R}^2 \setminus \overline{\mathbb{D}} \rightarrow \tilde{\Omega}$ is conformal and Γ is a hyperbolic geodesic in $\mathbb{R}^2 \setminus \overline{\mathbb{D}}$ connecting x and y with $\gamma \subset B(0, 100)$, then*

$$\ell(\tilde{\phi}(\Gamma)) \leq C(C_1)|\tilde{\phi}(x) - \tilde{\phi}(y)|.$$

Proof. According to the definition of quasiconvexity, there exists a curve $\gamma \subset \tilde{\Omega}$ joining x and y with $\ell(\gamma) \leq C_1|x - y|$. Then by Theorem 2.3, we have

$$\ell(\tilde{\phi}(\Gamma)) \leq C\ell(\gamma) \leq C(C_1)|\tilde{\phi}(x) - \tilde{\phi}(y)|$$

with the constant depending only on C_1 . \square

Lemma 2.12. *Let $\tilde{\Omega}$ be the complementary domain of a Jordan domain. Then if $\tilde{\Omega}$ is c_1 -quasiconvex, its closure is c_2 -quasiconvex with c_2 depending only on c_1 . Conversely, if the closure of $\tilde{\Omega}$ is c_2 -quasiconvex, then $\tilde{\Omega}$ is c_1 -quasiconvex with c_1 depending only on c_2 .*

Proof. We first show the sufficiency of c_2 -quasiconvexity of the closure. Let $x, y \in \tilde{\Omega}$. To begin, let us consider the special case where

$$|x - y| \leq \delta \text{diam}(\partial\tilde{\Omega}), \quad \text{dist}(x, \partial\tilde{\Omega}) \leq \delta \text{diam}(\partial\tilde{\Omega}) \quad \text{and} \quad \text{dist}(y, \partial\tilde{\Omega}) \leq \delta \text{diam}(\partial\tilde{\Omega})$$

with the constant δ in Lemma 2.14. Then by applying Corollary 2.11 and Lemma 2.14 to the hyperbolic geodesic connecting x and y we get the desired control on the length of the geodesic.

Let us now prove the general case. Let $x, y \in \tilde{\Omega}$ be two distinct points. Since $\overline{\tilde{\Omega}}$ is c_2 -quasiconvex, there exists a curve $\gamma \subset \overline{\tilde{\Omega}}$ connecting x to y with $\ell(\gamma) \leq c_2|x - y|$. Let $\{x_i\}_{i=0}^k \subset \gamma$ be such that $x_0 = x$, $x_k = y$,

$$0 < |x_i - x_{i-1}| \leq \frac{1}{6} \delta \text{diam}(\partial\tilde{\Omega}) \quad \text{for all } i \in \{1, \dots, k\}. \quad (2.1)$$

The existence of these points follows from the finiteness of the length of γ . By the definition of the length of a curve, we further have

$$\sum_{i=1}^k |x_i - x_{i-1}| \leq \ell(\gamma) \leq c_2 |x - y|. \quad (2.2)$$

For each $i \in \{1, \dots, k-1\}$ take a point

$$x'_i \in \tilde{\Omega} \cap B(x_i, \min\{|x_i - x_{i-1}|, |x_i - x_{i+1}|\})$$

and define $x'_0 = x$ and $x'_k = y$. By the choice of x'_i we have for every $i \in \{1, \dots, k\}$ the estimate

$$|x'_i - x'_{i-1}| \leq |x'_i - x_i| + |x_i - x_{i-1}| + |x_{i-1} - x'_{i-1}| \leq 3|x_i - x_{i-1}|. \quad (2.3)$$

Combining this with (2.1) and (2.2) gives

$$|x'_i - x'_{i-1}| \leq \frac{1}{2} \delta \operatorname{diam}(\partial \tilde{\Omega}) \quad \text{and} \quad \sum_{i=1}^k |x'_i - x'_{i-1}| \leq \sum_{i=1}^k 3|x_i - x_{i-1}| \leq 3c_2 |x - y|. \quad (2.4)$$

Now for each $i \in \{1, \dots, k\}$ we define a curve $\gamma_i \subset \tilde{\Omega}$ connecting x'_{i-1} and x'_i . There are two cases. If

$$|x'_i - x'_{i-1}| \leq \delta \operatorname{diam}(\partial \tilde{\Omega}), \quad \operatorname{dist}(x'_i, \partial \tilde{\Omega}) \leq \delta \operatorname{diam}(\partial \tilde{\Omega}) \quad \text{and} \quad \operatorname{dist}(x'_{i-1}, \partial \tilde{\Omega}) \leq \delta \operatorname{diam}(\partial \tilde{\Omega}), \quad (2.5)$$

then by the special case we find $\gamma_i \subset \tilde{\Omega}$ connecting x'_{i-1} and x'_i with $\ell(\gamma_i) \leq C(c_2)|x'_i - x'_{i-1}|$. If (2.5) fails, then by (2.4)

$$\operatorname{dist}(x'_i, \partial \tilde{\Omega}) > \delta \operatorname{diam}(\partial \tilde{\Omega}) \quad \text{or} \quad \operatorname{dist}(x'_{i-1}, \partial \tilde{\Omega}) > \delta \operatorname{diam}(\partial \tilde{\Omega}).$$

In this case we define $\gamma_i := [x'_i, x'_{i-1}] \subset \tilde{\Omega}$. Then $\ell(\gamma_i) = |x'_i - x'_{i-1}|$. By concatenating all the γ_i , via (2.4) we finally get a curve γ' in $\tilde{\Omega}$ connecting x to y with

$$\ell(\gamma') \leq \sum_{i=1}^k \ell(\gamma_i) \lesssim \sum_{i=1}^k |x'_i - x'_{i-1}| \lesssim |x - y|,$$

with the constant depending only on c_2 as required.

For the necessity, we choose sequences of points with $x_n \rightarrow x$ and $y_n \rightarrow y$ with $x_n, y_n \in \tilde{\Omega}$. Then for every large $n \in \mathbb{N}$, there exists a curve γ_n of length uniformly bounded from above by a multiple of $|x - y|$ which joins x_n and y_n , according to the quasiconvexity of $\tilde{\Omega}$. Therefore up to reparametrizing γ_n 's, Arzelà-Ascoli lemma tells us that (a subsequence of) $\{\gamma_n\}$ converges to a curve $\gamma \subset \tilde{\Omega}$ joining x and y , whose length is bounded from above by a multiple of $|x - y|$. \square

Lemma 2.13. *Let Ω be a bounded simply connected domain so that Ω^c is C_1 -quasiconvex and let $\varphi : \mathbb{D} \rightarrow \Omega$ be conformal. Set $B_n = B(0, 1 - 2^{-n})$ and $\Omega_n = \varphi(B_n)$ for $n \geq 1$. Then Ω_n^c is $C(C_1)$ -quasiconvex for each n .*

Proof. Notice that, by part (1) of Lemma 2.8 Ω is a J -John domain with $J = J(C_1)$. Fix a pair of points $x, y \in \Omega_n^c$. If $x, y \in \Omega^c$, then we are done by the quasiconvexity of Ω^c . Hence, by symmetry we may assume that $x \in \Omega \setminus \Omega_n$.

Let $x_0 \in \partial\Omega$ be a point such that the line segment $[\varphi^{-1}(x), \varphi^{-1}(x_0)]$ joining $\varphi^{-1}(x)$ and $\varphi^{-1}(x_0)$ lies in the line segment $[0, \varphi^{-1}(x_0)]$. Notice that $\gamma_1 = \varphi([\varphi^{-1}(x), \varphi^{-1}(x_0)])$ is a hyperbolic geodesic.

Suppose first that $y \in \Omega^c$. Then we can join x_0 and y by a curve γ_2 in Ω^c of length at most $C_1|x_0 - y|$ because Ω^c is quasiconvex. Note that part (3) of Lemma 2.8 implies

$$|x - x_0| \leq \ell(\gamma_1) \leq C(J) \operatorname{dist}(x, \partial\Omega). \quad (2.6)$$

Then by the fact that $\operatorname{dist}(x, \partial\Omega) \leq |x - y|$ as $x \in \Omega$ and $y \in \Omega^c$, via (2.6) we have

$$\ell(\gamma_1) + \ell(\gamma_2) \leq C(J)|x - y| + C(C_1)|x_0 - y| \leq C(C_1)(|x - y| + |x - x_0|) \leq C(C_1)|x - y|,$$

where in the second inequality we applied the triangle inequality to get

$$|x_0 - y| \leq |x - y| + |x - x_0|.$$

Hence $\gamma_1 \cup \gamma_2$ is a desired curve.

Next consider the case where $y \in \Omega \setminus \Omega_n$. Here we have two subcases depending whether (2.7) below holds.

Firstly suppose $\varphi^{-1}(y) \in B(\varphi^{-1}(x), (1 - |\varphi^{-1}(x)|)/2) =: B$. Then the existence of the desired curve follows directly from Lemma 2.6; note that we can join $\varphi^{-1}(x)$ and $\varphi^{-1}(y)$ inside $B \setminus \overline{B_n}$ by a curve of length uniformly bounded by a multiple of $|\varphi^{-1}(x) - \varphi^{-1}(y)|$ for some absolute constant, and then by applying a change of variable we obtain the desired estimate from Lemma 2.6. A similar conclusion also holds if the roles of x, y are reversed. By Lemma 2.6 we know that $\varphi(B(z, (1 - |z|)/2))$ is of Whitney-type for any $z \in \mathbb{D}$, and hence there exists an absolute constant C such that if

$$C|x - y| \leq \max\{\operatorname{dist}(x, \partial\Omega), \operatorname{dist}(y, \partial\Omega)\},$$

then these two points fall into one of the two cases above.

Hence we only need to consider the case where $x, y \in \Omega \setminus \Omega_n$ and

$$C|x - y| \geq \max\{\operatorname{dist}(x, \partial\Omega), \operatorname{dist}(y, \partial\Omega)\}. \quad (2.7)$$

Recall the definition of x_0 above, and similarly define $y_0 \in \partial\Omega$. We can connect x_0 to y_0 by a curve γ_2 in Ω^c with length controlled from above by $C_1|x_0 - y_0|$. Also join y_0 to y by the hyperbolic geodesic γ_3 , analogous to γ_1 that joins x_0 to x . The desired curve is obtained by concatenating γ_1 with γ_2 and γ_3 , respectively. Indeed by (2.7) and (2.6)

$$\begin{aligned} \ell(\gamma_1) + \ell(\gamma_2) + \ell(\gamma_3) &\leq C(J) \operatorname{dist}(x, \partial\Omega) + C(J) \operatorname{dist}(y, \partial\Omega) + C(C_1)|x_0 - y_0| \\ &\leq C(C_1)(|x - y| + |x_0 - y_0|) \leq C(C_1)|x - y|, \end{aligned}$$

where again in the last inequality we apply the triangle inequality with (2.7) to have

$$|x_0 - y_0| \leq |x - y| + \operatorname{dist}(x, x_0) + \operatorname{dist}(y, y_0) \leq |x - y| + C(J) \operatorname{dist}(x, \partial\Omega) + C(J) \operatorname{dist}(y, \partial\Omega).$$

□

The proof of the following technical lemma is given in Section A.3 of the Appendix.

Lemma 2.14. *Given $C_1 > 0$ and an unbounded C_1 -quasiconvex domain $\tilde{\Omega}$ whose boundary is Jordan, there is a constant $\delta_1 = \delta_1(C_1) > 0$ so that the following holds: If $\tilde{\varphi}: \tilde{\Omega} \rightarrow \mathbb{R}^2 \setminus \overline{\mathbb{D}}$ is conformal and $z_1, z_2 \in \tilde{\Omega}$ satisfy $|z_1 - z_2| < \delta_1 \text{diam}(\partial\tilde{\Omega})$ and $\text{dist}(z_i, \partial\tilde{\Omega}) \leq \delta_1 \text{diam}(\partial\tilde{\Omega})$ for $i = 1, 2$, then $|\tilde{\varphi}(z_1) - \tilde{\varphi}(z_2)| < \frac{1}{4}$ and $\text{dist}(\varphi(z_i), \partial\mathbb{D}) < \frac{1}{4}$ for $i = 1, 2$.*

3. SUFFICIENCY FOR JORDAN DOMAINS

Let Ω be a Jordan domain satisfying (1.2). By Lemma 2.12, the complementary domain $\tilde{\Omega}$ of Ω is C_1 -quasiconvex, where C_1 depends only on C in (1.2). We prove the following theorem in this section.

Theorem 3.1. *Let $\Omega \subset \mathbb{R}^2$ be a Jordan domain whose complementary domain is C_1 -quasiconvex. Then there exists an extension operator $E: W^{1,1}(\Omega) \rightarrow W^{1,1}(\mathbb{R}^2)$ with the norm depending only on C_1 .*

We begin by decomposing a part of $\tilde{\Omega}$ into hyperbolic triangles (defined below), and then define auxiliary functions on them. After this we construct an extension operator based on these functions.

Recall first that Ω is a J -John domain with $J = J(C_1)$ by part (1) of Lemma 2.8. Let $\varphi: \overline{\mathbb{D}} \rightarrow \overline{\Omega}$ be a homeomorphism which is conformal in \mathbb{D} and satisfies $\varphi(0) = x_0$, where x_0 is the distinguished point of Ω . Such a homeomorphism exists by the Riemann mapping theorem and the Carathéodory-Osgood theorem [24] since Ω is Jordan. Moreover we denote by $\tilde{\varphi}: \tilde{\Omega} \rightarrow \mathbb{R}^2 \setminus \mathbb{D}$ a homeomorphism which is conformal in $\tilde{\Omega}$. The map $\tilde{\varphi}$ is obtained also from an extension of a Riemann mapping.

3.1. Decomposition of the domain and the complementary domain. Fix $k_0 \in \mathbb{N}$ to be determined later. Let $A_1 = B(0, \frac{1}{2})$ and

$$A_k = B(0, 1 - 2^{-k}) \setminus B(0, 1 - 2^{-k+1})$$

for $k \geq 2$. For each $k \geq 1$, the radial rays passing through the points $x_k^{(j)} = e^{j2^{-k-k_0}\pi i}$ with $1 \leq j \leq 2^{k+k_0+1}$ divide A_k into 2^{k+k_0+1} subsets. We denote them by $D_k^{(j)}$ with $1 \leq j \leq 2^{k+k_0+1}$. Here the upper indices are labeled anticlockwise from the positive real axis. Define $Q_k^{(j)} = \varphi(D_k^{(j)})$, and $z_k^{(j)} = \varphi(x_k^{(j)})$ with $x_k^{(0)} = x_k^{(2^{k+k_0+1})}$. Since $D_k^{(j)}$ is a Whitney-type set with the constant depending only on k_0 , we obtain that $Q_k^{(j)}$ is also of $\lambda(k_0)$ -Whitney-type by Lemma 2.6.

Moreover, we show that

$$\text{dist}_{\Omega}(z_k^{(j-1)}, z_k^{(j)}) \leq C(J, k_0) \text{diam}(Q_k^{(j)}) \quad (3.1)$$

Towards this, let x_1 be a point in $\Gamma_1 \cap Q_k^{(j)}$, where Γ_1 is the hyperbolic geodesic joining $z_k^{(j-1)}$ and the origin. Recall that Γ_1 is a J -John curve by part (3) of Lemma 2.8. Thus we have

$$\text{dist}(x_1, \partial\Omega) \geq C(J) \ell(\Gamma_1[x_1, z_k^{(j-1)}]) \geq C(J) \text{dist}_{\Omega}(x_1, z_k^{(j)})$$




FIGURE 1. The hyperbolic triangulation of the complementary domain $\tilde{\Omega}$ of Ω via conformal maps $\varphi: \mathbb{D} \rightarrow \Omega$ and $\tilde{\varphi}: \tilde{\Omega} \rightarrow \mathbb{R}^2 \setminus \overline{\mathbb{D}}$.

where $\Gamma_1[x_1, z_k^{(j-1)}]$ denotes the subarc of Γ_1 joining x_1 and $z_k^{(j-1)}$. Likewise we define Γ_2 as the hyperbolic geodesic joining $z_k^{(j)}$ and the origin, x_2 to be a point in $\Gamma_2 \cap Q_k^{(j)}$, and have the estimate

$$\text{dist}(x_2, \partial\Omega) \geq C(J)\ell(\Gamma_2[x_2, z_k^{(j)}]) \geq C(J)\text{dist}_\Omega(x_2, z_k^{(j)}).$$

Since $Q_k^{(j)}$ is a connected set of $\lambda(k_0)$ -Whitney-type, we have

$$\text{dist}_\Omega(x_1, x_2) \leq C(k_0)\text{dist}(x_1, \partial\Omega) \sim_{k_0} \text{dist}(x_2, \partial\Omega).$$

Therefore by the triangle inequality we conclude that

$$\begin{aligned} \text{dist}_\Omega(z_k^{(j-1)}, z_k^{(j)}) &\leq \text{dist}_\Omega(z_k^{(j-1)}, x_1) + \text{dist}_\Omega(z_k^{(j)}, x_2) + \text{dist}_\Omega(x_1, x_2) \\ &\leq \text{dist}(x_1, \partial\Omega) \sim \text{diam}(Q_k^{(j)}) \end{aligned}$$

with the constant depending only on k_0 and J , which concludes (3.1).

Denote by $\gamma_k^{(j)}$ the hyperbolic geodesic of $\tilde{\Omega}$ connecting $z_k^{(j)}$ and $z_k^{(j-1)}$. Since Ω is J -John, by Lemma 2.8 φ is uniformly continuous; also see [25, Page 100, Corollary 5.3]. Therefore we

can choose $k_0 = k_0(C_1) \in \mathbb{N}$ large enough such that

$$|z_k^{(j-1)} - z_k^{(j)}| \leq \delta \operatorname{diam}(\Omega)$$

for any $k, j \in \mathbb{N}$, where δ is the constant in Lemma 2.14. This allows us to apply Corollary 2.11 with (3.1) and conclude that

$$\ell(\gamma_k^{(j)}) \leq C(C_1)|z_k^{(j-1)} - z_k^{(j)}| \leq C(C_1) \operatorname{diam}(Q_k^{(j)}). \quad (3.2)$$

A set is called a *hyperbolic triangle* if it is enclosed by three hyperbolic geodesics, meeting at the three vertices. For $k \in \mathbb{N}$ and $1 \leq j \leq 2^{k_0+k+1}$, we denote by $R_k^{(j)}$ the relatively closed set (with respect to the topology of $\tilde{\Omega}$) enclosed by $\gamma_k^{(j)}$, $\gamma_{k+1}^{(2j-1)}$ and $\gamma_{k+1}^{(2j)}$. It is a closed hyperbolic triangle (with respect to the topology of $\tilde{\Omega}$); see Figure 1.

Moreover, notice that for $1 \leq j \leq 2^{k_0+2}$ the image of each $\gamma_1^{(j)}$ under $\tilde{\varphi}$ is a circular arc; recall that $\tilde{\varphi}: \tilde{\Omega} \rightarrow \mathbb{R}^2 \setminus \mathbb{D}$ is a homeomorphism which is conformal in $\tilde{\Omega}$. Denote the preimage under $\tilde{\varphi}$ of the midpoint of each arc $\tilde{\varphi}(\gamma_1^{(j)})$ by $z_0^{(j)}$, and the corresponding hyperbolic geodesic connecting $z_0^{(j)}$ and $z_0^{(j+1)}$ by $\gamma_0^{(j)}$; here $z_0^{(2^{k_0+2}+1)} = z_0^{(1)}$. Then with the help of $\gamma_1^{(j)}$ we obtain 2^{k_0+1} closed hyperbolic triangles $R_0^{(j)}$ such that every $R_0^{(j)}$ is enclosed by $\gamma_0^{(j)}$, $\gamma_1^{(j)}$ and $\gamma_1^{(j+1)}$; here $\gamma_1^{(2^{k_0+1}+1)} = \gamma_1^{(1)}$. Moreover $\tilde{\varphi}(\gamma_0^{(j)}) \subset B(0, 10)$ for any $1 \leq j \leq 2^{k_0+2}$ since we defined the constant δ as in Lemma 2.14. Therefore it is legitimate to apply Corollary 2.11 with (3.2) to conclude

$$\begin{aligned} \ell(\gamma_0^{(j)}) &\leq C_1|z_0^{(j+1)} - z_0^{(j)}| \leq C_1 \operatorname{diam}(\gamma_0^{(j+1)} \cup \gamma_0^{(j)}) \leq C_1 \ell(\gamma_0^{(j+1)}) + C_1 \ell(\gamma_0^{(j)}) \\ &\leq C(C_1)|z_1^{(j-1)} - z_1^{(j)}| + |z_1^{(j+1)} - z_1^{(j)}| \leq C(C_1) \operatorname{diam}(Q_1^{(j)}), \end{aligned} \quad (3.3)$$

where in the last inequality we used the fact that

$$\operatorname{diam}(Q_1^{(j+1)}) \sim_{C_1} \operatorname{diam}(Q_1^{(j)})$$

coming from Lemma 2.6 with a change of variable.

3.2. Definition of the extension. We define an auxiliary function as follows. Our construction is linear in the parameters a_1, a_2, a_3 .

Lemma 3.2. *Let R be a closed hyperbolic triangle with vertices z_1, z_2, z_3 and enclosed by three Jordan arcs $\gamma_1, \gamma_2, \gamma_3$. Then for $a_1, a_2, a_3 \in \mathbb{R}$, there exists a function ϕ , locally Lipschitz in $R \setminus \{z_1, z_2, z_3\}$, with the following properties:*

$$\min\{a_1, a_2, a_3\} \leq \phi \leq \max\{a_1, a_2, a_3\}, \quad (3.4)$$

and for $i = 1, 2, 3$, we have $\phi(x) = a_i$ when $x \in \gamma_i$. Moreover,

$$\|\nabla \phi\|_{L^1(R)} \lesssim \sum_{i,j \in \{1,2,3\}} |a_i - a_j| \min\{\ell(\gamma_i), \ell(\gamma_j)\} \quad (3.5)$$

with the constant only depending on C_1 .

Proof. Without loss of generality we may assume that γ_3 has the longest length among these three curves. We first consider the case where $\max\{a_1, a_2\} \leq a_3$. Define

$$\phi(x) = \min \left\{ a_1 + \inf_{\theta_1} \int_{\theta_1} \frac{20(a_3 - a_1)}{\text{dist}(z, \partial R \setminus \gamma_1)} ds(z), a_2 + \inf_{\theta_2} \int_{\theta_2} \frac{20(a_3 - a_2)}{\text{dist}(z, \partial R \setminus \gamma_2)} ds(z), a_3 \right\},$$

where the infima are taken over all curves $\theta_i \subset R$ connecting x to γ_i . Observe that ϕ is bounded by its definition and locally Lipschitz in the (Euclidean) interior of R via the triangle inequality; indeed for any $x \in R$,

$$|\phi(x) - \phi(y)| \leq C|x - y| \max_{i=1,2} (a_3 - a_i) \text{dist}(z, \partial R \setminus \gamma_i)^{-1}$$

for $y \in B(x, 1/3 \text{dist}(x, \partial R))$ with an absolute constant C . Moreover ϕ takes the correct boundary value by its definition; note that both of the integrals in the definition of ϕ have logarithmic growth towards the boundary.

Define

$$A = \bigcup_{i=1}^2 \bigcup_{y \in \gamma_i} B\left(y, \frac{1}{10} \text{dist}(y, \partial R \setminus \gamma_i)\right).$$

Note that for $x \in R \setminus A$ we have

$$\phi(x) = a_3. \quad (3.6)$$

By the $5r$ -covering theorem, we know that there exists a countable collection of pairwise disjoint disks

$$\left\{ B\left(y_{ij}, \frac{1}{10} \text{dist}(y_{ij}, \partial R \setminus \gamma_i)\right) \right\}_{y_{ij} \in \gamma_i}$$

such that

$$A \subset \bigcup_{i=1}^2 \bigcup_{y_{ij} \in \gamma_i} B\left(y_{ij}, \frac{1}{2} \text{dist}(y_{ij}, \partial R \setminus \gamma_i)\right) =: \bigcup_{i=1}^2 \bigcup_{y_{ij} \in \gamma_i} B_{ij}.$$

Let us estimate the gradient inside B_{ij} . Define

$$A_i = \bigcup_{y_{ij} \in \gamma_i} B_{ij}$$

for $i = 1, 2$. Notice that $A_1 \cap A_2 = \emptyset$ by the triangle inequality; B_{ij} are open disks. Then for every $x \in A_i$, $i = 1, 2$, we have

$$\phi(x) = \min \left\{ a_i + \inf_{\theta_i} \int_{\theta_i} \frac{20(a_3 - a_i)}{\text{dist}(z, \partial R \setminus \gamma_i)} dz, a_3 \right\},$$

and hence for $x \in A_i$, $i = 1, 2$,

$$|\nabla \phi(x)| \lesssim \frac{a_3 - a_i}{\text{dist}(z, \partial R \setminus \gamma_i)}. \quad (3.7)$$

By the definition of A_i , (3.7) together with (3.6) gives us the local Lipschitz property of ϕ in $R \setminus \{z_1, z_2, z_3\}$.

Notice that, for any $z \in B_{ij}$, $i = 1, 2$ by the triangle inequality we have

$$2 \text{dist}(z, \partial \Omega \setminus \gamma_i) \geq \text{dist}(y_{ij}, \partial \Omega \setminus \gamma_i).$$

Then by (3.6) and (3.7), via the triangle inequality we further have

$$\begin{aligned}
\|\nabla\phi\|_{L^1(R)} &\lesssim \sum_{i=1,2} \sum_j \int_{B_{ij}} \frac{|a_3 - a_i|}{\text{dist}(z, \partial R \setminus \gamma_i)} ds(z) \\
&\lesssim \sum_{i=1,2} \sum_j \int_{B_{ij}} \frac{|a_3 - a_i|}{\text{dist}(y_{ij}, \partial R \setminus \gamma_i)} ds(z) \\
&\lesssim \sum_{i=1,2} \sum_j |a_3 - a_i| \text{dist}(y_{ij}, \partial R \setminus \gamma_i) \lesssim |a_3 - a_1| \ell(\gamma_1) + |a_3 - a_2| \ell(\gamma_2),
\end{aligned}$$

where the last inequality follows from the fact that $\frac{1}{5}B_{ij}$ are centered on γ_i and pairwise disjoint.

For the other cases, we apply suitable modifications to ϕ ; the idea is to increase or decrease the value of $\phi(x)$ towards a longest curve according to the values on the other two curves. Namely if $a_3 \leq \min\{a_1, a_2\}$, we define

$$\phi(x) = \max \left\{ a_2 - \inf_{\theta_2} \int_{\theta_2} \frac{20(a_2 - a_3)}{\text{dist}(z, \partial R \setminus \gamma_1)} ds(z), a_1 - \inf_{\theta_1} \int_{\theta_1} \frac{20(a_1 - a_3)}{\text{dist}(z, \partial R \setminus \gamma_2)} ds(z), a_3 \right\}.$$

For the remaining case, by symmetry we only consider the case where $a_1 \leq a_3 \leq a_2$, and define

$$\phi(x) = \max \left\{ a_2 - \inf_{\theta_2} \int_{\theta_2} \frac{20(a_2 - a_3)}{\text{dist}(z, \partial R \setminus \gamma_1)} ds(z), \min \left\{ a_1 + \inf_{\theta_1} \int_{\theta_1} \frac{20(a_3 - a_1)}{\text{dist}(z, \partial R \setminus \gamma_2)} ds(z), a_3 \right\} \right\}.$$

An argument similar to the above one gives the estimate for the L^1 -norm of the gradient of ϕ . \square

With the help of Lemma 3.2, we are ready to construct our extension operator.

Fix a function

$$u \in W^{1,1}(\Omega) \cap C^\infty(\mathbb{R}^2). \quad (3.8)$$

Recall that Ω is J -John by Lemma 2.8. For $k \geq 1$ and $1 \leq j \leq 2^{k_0+k+1}$ we associate to $\gamma_k^{(j)}$ a constant

$$a_k^{(j)} = \int_{Q_k^{(j)}} u(x) dx,$$

where $Q_k^{(j)} \in W$ is the Whitney-type set defined in Section 3.1 above.

Let

$$R = \bigcup R_k^{(j)},$$

where the indices run over all the $k, j \in \mathbb{N}$ such that $R_k^{(j)}$ is defined. First, by Lemma 3.2 we associate to each $R_k^{(j)}$ a function $\phi_k^{(j)} \in W^{1,1}(R_k^{(j)})$ for every $k \geq 1$ and $1 \leq j \leq 2^{k_0+k+1}$ so that

$$\phi_k^{(j)}(x) = \begin{cases} a_k^{(j)} & x \in \gamma_k^{(j)} \cap R_k^{(j)} \\ a_{k+1}^{(2j-1)} & x \in \gamma_{k+1}^{(2j-1)} \cap R_k^{(j)} \\ a_{k+1}^{(2j)} & x \in \gamma_{k+1}^{(2j)} \cap R_k^{(j)} \end{cases};$$




FIGURE 2. A hyperbolic triangle is shown with the three Whitney-type sets corresponding to its three edges, respectively. The colors of the edges and the Whitney-type sets show how the boundary values are taken by the function in the hyperbolic triangle.

see Figure 2. When $k = 0$, again by Lemma 3.2, for each $1 \leq j \leq 2^{k_0+1}$ we get a function $\phi_0^{(j)} \in W^{1,1}(R_0^{(j)})$ satisfying

$$\phi_0^{(j)}(x) = \begin{cases} a & x \in \gamma_0^{(j)} \setminus (\gamma_1^{(j)} \cup \gamma_1^{(j+1)}) \\ a_1^{(j)} & x \in \gamma_1^{(j)} \cap R_0^{(j)} \\ a_1^{(j+1)} & x \in \gamma_1^{(j+1)} \cap R_0^{(j)} \end{cases},$$

where $a_1^{(2^{k_0+1}+1)} = a_1^{(1)}$ and

$$a = \oint_{\Omega} u \, dx = \frac{1}{|\Omega|} \int_{\Omega} u \, dx.$$

Finally set

$$Eu(x) = \begin{cases} u(x), & \text{if } x \in \bar{\Omega} \\ \phi_k^{(j)}(x), & \text{if } x \in R_k^{(j)} \cap \tilde{\Omega} \\ a, & \text{otherwise} \end{cases}.$$

Now let us estimate the homogeneous Sobolev norm of Eu . It is clear from the construction that $Eu \in W_{\text{loc}}^{1,1}(\tilde{\Omega})$ since Eu is locally Lipschitz in $\tilde{\Omega}$ by Lemma 3.2. Since by Lemma 2.6 the conformal map φ is uniformly $C(\lambda)$ -bi-Lipschitz on λ -Whitney-type sets up to a dilation factor, we have that $Q_k^{(j)}$'s are $C(C_1)$ -John domains, and so are, for example, also $Q_k^{(j)} \cup Q_k^{(j+1)}$ and $Q_k^{(j)} \cup Q_{k+1}^{(2j+1)}$. Recall the following (1, 1)-Poincaré inequality for John domains: For any J -John domain $G \subset \mathbb{R}^2$ and $v \in W^{1,1}(G)$ we have

$$\int_G |v - v_G| dx \leq C(J) \text{diam}(G) \int_G |\nabla v| dx,$$

where v_G is the integral average of v in G ; see e.g. [6] and the references therein. Also note that by the definition of a John domain, the facts that every $Q_1^{(j)}$ is of $\lambda(C_1)$ -Whitney-type and that the John center of Ω is contained in every $Q_1^{(j)}$, we have

$$\text{diam}(Q_1^{(j)}) \sim \text{diam}(\Omega) \quad \text{and then} \quad |Q_1^{(j)}| \sim |\Omega| \sim \text{diam}(\Omega)^2 \quad (3.9)$$

for each $1 \leq j \leq 2^{k_0+1}$, where the constants depend only on C_1 .

Now we can estimate the homogeneous Sobolev norm of Eu over $\tilde{\Omega}$. By recalling (3.2) and (3.3), we apply Lemma 3.2 and the above Poincaré inequality to get

$$\begin{aligned} & \|\nabla Eu(x)\|_{L^1(\tilde{\Omega})} \\ & \lesssim \sum_{\substack{k \geq 1 \\ 1 \leq j \leq 2^{k_0+k+1}}} \int_{R_k^{(j)}} |\nabla \phi_k^{(j)}(x)| dx + \sum_{1 \leq j \leq 2^{k_0+2}} \int_{R_0^{(j)}} |\nabla \phi_k^{(j)}(x)| dx \\ & \lesssim \sum_{\substack{k \geq 1 \\ 1 \leq j \leq 2^{k_0+k+1}}} \left(|a_k^{(j)} - a_{k+1}^{(2j-1)}| + |a_k^{(j)} - a_{k+1}^{(2j)}| + |a_{k+1}^{(2j-1)} - a_{k+1}^{(2j)}| \right) \text{diam}(Q_k^{(j)}) \\ & \quad + \sum_{j=1}^{2^{k_0+1}} \left(|a_1^{(j)} - a_1^{(j+1)}| + |a_1^{(j)} - a| + |a_1^{(j+1)} - a| \right) \text{diam}(Q_1^{(j)}) \\ & \lesssim \sum_{\substack{k \in \mathbb{N} \\ 1 \leq j \leq 2^{k_0+k+1}}} \left(\int_{Q_k^{(j)} \cup Q_{k+1}^{(2j-1)}} |\nabla u| dx + \int_{Q_k^{(j)} \cup Q_{k+1}^{(2j)}} |\nabla u| dx + \int_{Q_{k+1}^{(2j-1)} \cup Q_{k+1}^{(2j)}} |\nabla u| dx \right) \\ & \quad + \int_{\Omega} |\nabla u| dx, \end{aligned}$$

where the constants depend only on C_1 .

Additionally, by our construction, we only have uniformly finite overlaps for the Whitney-type sets in the summation. Therefore we have

$$\|\nabla Eu\|_{L^1(\tilde{\Omega})} \leq C(C_1) \int_{\Omega} |\nabla u| dx. \quad (3.10)$$

Let B be a ball of radius $4 \operatorname{diam}(\Omega)$ such that Ω is contained in $\frac{1}{2}B$. Then

$$\begin{aligned} \|Eu\|_{L^1(B)} &\lesssim \sum_{\substack{k \geq 1 \\ 1 \leq j \leq 2^{k_0+k+1}}} \int_{R_k^{(j)}} |Eu(x)| dx + \int_{\Omega} |u(x)| dx \\ &\lesssim \sum_{\substack{k \geq 1 \\ 1 \leq j \leq 2^{k_0+k+1}}} \int_{Q_k^{(j)}} |u(x)| dx + \int_{\Omega} |u(x)| dx \lesssim \int_{\Omega} |u(x)| dx \end{aligned} \quad (3.11)$$

where the constant depending only on C_1 . Here we used (3.4) with the fact that

$$|R_k^{(j)}| \leq C(C_1) |Q_k^{(j)}| \sim_{C_1} |Q_{k+1}^{(2j-1)}| \sim_{C_1} |Q_{k+1}^{(2j)}|$$

coming from (3.2) and the fact that $Q_k^{(j)}$ is of $C(C_1)$ -Whitney-type.

3.3. Absolute continuity. We first check that Eu is absolutely continuous along almost every line segment parallel to the coordinate axes for $u \in C^\infty(\mathbb{R}^2) \cap W^{1,1}(\Omega)$. Notice that u is already smooth in $\bar{\Omega}$. Denoting the collection of the countable number of points $z_k^{(j)}$ by Z , we first claim that the restriction of Eu to $\mathbb{R}^2 \setminus Z$ is continuous.

Indeed since u is Lipschitz in $\bar{\Omega}$ and Eu is continuous in $\tilde{\Omega}$ by our construction, we just need to verify continuity of Eu at points of $\partial\Omega \setminus Z$. Given a sequence of points $\{y_i\}_{i=1}^\infty$ in $\tilde{\Omega}$ converging to $y \in \partial\Omega \setminus Z$, we claim that there is a sequence of hyperbolic triangles $\{R_i\}_{i=1}^\infty$ among $R_k^{(j)}$ such that,

$$y_i \in R_i \quad \text{and} \quad R_i \rightarrow y \text{ when } i \rightarrow \infty. \quad (3.12)$$

If so, then by (3.4) we know that $Eu(y_i)$ is bounded from above and below respectively by one of the integral averages of u over the corresponding Whitney-type sets Q_i or over two neighbors of Q_i . Let us show that now also Q_i converge to y (inside Ω). Indeed since R_i converge to y , especially the three vertices of R_i also converge to y as $i \rightarrow \infty$, by the continuity of φ and $\tilde{\varphi}$ on the boundary, we know that $\operatorname{diam}(Q_i) \rightarrow 0$. Therefore Q_i also converge to y . Since u is Lipschitz in $\bar{\Omega}$, we get the continuity of $Eu(x)$ in $\mathbb{R}^2 \setminus Z$.

Now let us show the existence of the asserted sequence $\{R_i\}_{i=1}^\infty$ as in (3.12). Fix $y \in \partial\Omega \setminus Z$, and for any $s \in \mathbb{N}$ define

$$\Gamma_s = \overline{\bigcup_{0 \leq k \leq s} \bigcup_j \gamma_k^{(j)}}, \quad (3.13)$$

where the second union in j is over all the indices j such that $\gamma_k^{(j)}$ is defined. Notice that Γ_s is compact, and $y \notin \Gamma_s$. Therefore there exists $\delta_s > 0$ such that $\operatorname{dist}(y, \Gamma_s) \geq \delta_s$. Thus if $\{y_i\}_{i=1}^\infty$ converges to y , then the lower indices of our hyperbolic triangles $R_{k(i)}^{(j)} = R_{k(i)}$ containing y_i tend to ∞ when $i \rightarrow \infty$. This together with the fact that $\tilde{\varphi}: \bar{\Omega} \rightarrow \mathbb{R}^2 \setminus \mathbb{D}$ is homeomorphic implies that $\tilde{\varphi}(R_{k(i)})$ converges to a point on the unit circle as $i \rightarrow \infty$, and thus $R_{k(i)} \rightarrow y$ as desired since $y_i \rightarrow y$ and $y_i \in R_{k(i)}$. Thus (3.12) follows.

Since Z is countable, almost every line parallel to the x_1 -axis contains no point of Z . Additionally, for almost every such line S ,

$$\int_{S \cap \tilde{\Omega}} |\nabla Eu| dx < \infty \quad (3.14)$$

by Fubini's theorem. Fix S with these properties. Since Eu is locally Lipschitz in $\tilde{\Omega}$, $Eu|_S$ is absolutely continuous for every closed segment $I \subset S$ with $I \subset \tilde{\Omega}$.

Now for any line segment $I \subset S$, we first observe that $Eu|_I$ is certainly continuous as $Eu|_{\mathbb{R}^2 \setminus Z}$ is continuous. Next we show that $Eu|_I$ is absolutely continuous. Let $z, w \in I$.

If $z, w \in \overline{\Omega}$, then the L -Lipschitz continuity of u in $\overline{\Omega}$ gives

$$|Eu(z) - Eu(w)| \leq L|z - w|. \quad (3.15)$$

If $[z, w] \cap \overline{\Omega} = \emptyset$, then since Eu is absolutely continuous on line segments in S , we obtain

$$|Eu(z) - Eu(w)| \leq \int_{[z, w]} |\nabla Eu| dx. \quad (3.16)$$

For what is left, by symmetry we may assume that $z \in \overline{\Omega}$ while $w \in \tilde{\Omega}$. Then let z_0 be the closest point of $\overline{\Omega} \cap I$ to w between z and w . Then

$$|Eu(z) - Eu(w)| \leq |Eu(z) - Eu(z_0)| + |Eu(z_0) - Eu(w)| \leq L|z - z_0| + \int_{[z_0, w]} |\nabla Eu| dx. \quad (3.17)$$

For the last inequality, we used the facts that Eu is continuous on S and absolutely continuous on each closed subinterval $I \subset S$ when $I \subset \tilde{\Omega}$, together with (3.14).

Suppose that $\epsilon > 0$ and we are given $z_1, \dots, z_{2n} \in I$ so that the one-dimensional open intervals (z_{2i-1}, z_{2i}) , $1 \leq i \leq n$, are pairwise disjoint. By applying the relevant one of (3.15), (3.16) or (3.17) to each pair z_{2i-1}, z_{2i} , our assumption (3.14) together with the absolute continuity of integrals gives the existence of $\delta > 0$ so that $\sum_{i=1}^n |Eu(z_{2i-1}) - Eu(z_{2i})| \leq \epsilon$ provided $\sum_{i=1}^n |z_{2i-1} - z_{2i}| < \delta$. This implies the desired absolute continuity property. The case of lines parallel to the x_2 -axis is handled analogously.

Thus Eu is absolutely continuous along almost every line segment parallel to the coordinate axes for $u \in C^\infty(\mathbb{R}^2) \cap W^{1,1}(\Omega)$, and with the norm estimates (3.10) and (3.11) we conclude that $E: C^\infty(\mathbb{R}^2) \cap W^{1,1}(\Omega) \rightarrow W^{1,1}(B)$ is a (linear) extension operator.

3.4. Proof of Theorem 3.1. By subsection 3.3, the norm inequalities (3.10) and (3.11), and the fact that the Lebesgue measure of $\partial\Omega$ is zero (see part (4) of Lemma 2.8),

$$E: C^\infty(\mathbb{R}^2) \cap W^{1,1}(\Omega) \rightarrow W^{1,1}(B), \quad \text{where } \Omega \subset \frac{1}{2}B \subset B.$$

is both linear and bounded with respect to the non-homogeneous $W^{1,1}$ -norm. Since $C^\infty(\mathbb{R}^2) \cap W^{1,1}(\Omega)$ is dense in $W^{1,1}(\Omega)$ provided that Ω is a Jordan domain [22], linearity and boundedness of E allow us to extend the domain of E to the entire $W^{1,1}(\Omega)$. This concludes Theorem 3.1 because B is a $W^{1,1}$ -extension domain.

4. PIECEWISE HYPERBOLIC GEODESICS AND CUT-POINTS

Recall that we relied on a hyperbolic triangulation of the complementary domain in the construction of our extension operator in the Jordan case. In order to obtain a suitable variant of this for the (strongly quasiconvex) exterior of our simply connected domain, we need a counterpart of exterior hyperbolic geodesics.

Let $\varphi: \mathbb{D} \rightarrow \Omega$ be a conformal map. Our assumption that Ω^c is quasiconvex, say with constant C_1 , together with Lemma 2.8 allows us to extend φ continuously to entire $\overline{\mathbb{D}}$. As usual, the extension is also denoted by φ . We set $\tilde{\Omega} := \mathbb{R}^2 \setminus \overline{\Omega}$ and let $\{\tilde{\Omega}_i\}_{i=1}^N$ be an enumeration of the connected components of $\tilde{\Omega}$ with $N \in \mathbb{N} \cup \{+\infty\}$, so that $\tilde{\Omega}_1$ is the (unique) unbounded component. Then each $\tilde{\Omega}_i$ is Jordan for $i \geq 2$ by Lemma 2.10.

Because the boundary of Ω may have self-intersections, we eventually base our construction on labeling via $\partial\mathbb{D}$. Towards this end, let $x', y' \in \partial\mathbb{D}$ be distinct points such that $x := \varphi(x')$, $y := \varphi(y')$ satisfy $|x - y| < \delta$, where $\delta > 0$ will be fixed later. When $\delta < \pi/4$, this guarantees that the hyperbolic geodesic between x', y' in the exterior of the disk is contained in $B(0, 3)$. A *piecewise hyperbolic geodesic* (in Ω^c) associated to x, y via φ (between $x, y \in \partial\Omega$) in the complement of Ω is any curve obtained via the following construction.

Define $x'_n = (1 - 2^{-n})x'$, $y'_n = (1 - 2^{-n})y'$. Then $x_n := \varphi(x'_n)$, $y_n := \varphi(y'_n) \in \partial D_n$, where

$$D_n = \varphi(B(0, 1 - 2^{-n})),$$

for $n \geq 2$, is a Jordan domain. Denote by \tilde{D}_n the complementary domain of D_n . Since φ is continuous up to the boundary, the points $x_n, y_n \in \partial\tilde{D}_n$ converge to x, y , respectively. Also observe that $\Omega^c \subset \tilde{D}_{n+1} \subset \tilde{D}_n$, and $\Omega^c = \bigcap_n \tilde{D}_n$. Pick a hyperbolic geodesic $\gamma_n \subset \tilde{D}_n$ of \tilde{D}_n connecting x_n and y_n . Since Ω^c is C_1 -quasiconvex, Lemma 2.12 and Lemma 2.13 show that \tilde{D}_n is c_1 -quasiconvex with $c_1 = c_1(C_1)$. We now fix $\delta_1 = \delta_1(c_1) < \frac{1}{4c_1}$ as in Lemma 2.14. Then Lemma 2.14 guarantees that $|\tilde{\varphi}_n(z) - \tilde{\varphi}_n(w)| \leq \frac{1}{4}$ for any conformal map $\tilde{\varphi}_n: \tilde{D}_n \rightarrow \mathbb{R}^2 \setminus \overline{\mathbb{D}}$ and $z, w \in \partial D_n$ with $|z - w| < \delta_1 \text{diam}(\Omega)$; recall here that such a $\tilde{\varphi}_n$ extends continuously to the boundary and hence we may apply Lemma 2.14 even for points on the boundary. By the uniform continuity of φ on \mathbb{D} [25, Page 100, Corollary 5.3], we may choose $\delta = \delta(\delta_1, C_1)$ so that

$$|x_n - y_n| < \delta_1 \text{diam}(\Omega)$$

whenever $|x' - y'| < \delta$; notice that $|x'_n - y'_n| < |x' - y'|$. This determines the value of δ . Under this choice, $|\tilde{\varphi}_n(x_n) - \tilde{\varphi}_n(y_n)| < \frac{1}{4}$ and hence the c_1 -quasiconvexity of \tilde{D}_n together with Corollary 2.11 guarantees that

$$\ell(\gamma_n) \leq c_2 |x_n - y_n|, \tag{4.1}$$

where $c_2 = c_2(c_1)$.

Recalling that $x_n \rightarrow x$ and $y_n \rightarrow y$ when n tends to infinity, we conclude from (4.1) that $\gamma_n \subset B(\varphi(x), M)$ and $\ell(\gamma_n) \leq M$ for some M independently of n . Parametrize each γ_n by arc length. Then, by the Arzelà-Ascoli lemma, we obtain via uniform convergence a curve γ joining x and y in Ω^c . We call it a *piecewise hyperbolic geodesic joining x and y* . Also notice that by (4.1), if $\varphi(x') = \varphi(y')$, then the piecewise hyperbolic geodesic joining x and y is a constant curve.

Remark 4.1. A natural question arises: If the approximating sequences of $\varphi(x)$ and $\varphi(y)$ were chosen differently in the previous steps, especially if the preimages of x, y were chosen differently (but still distinct), could this result in a different curve γ ? This question is answered affirmatively by Lemma 4.5 below, modulo reparametrization.

Moreover we could also define piecewise hyperbolic geodesics joining sufficiently close points near $\partial\Omega$, for example, the points on the piecewise hyperbolic geodesics above; this is the reason why we chose $\delta_1 < \frac{1}{4c_1}$ when we applied Lemma 2.14 above. More precisely, the argument above applies whenever $x, y \in \Omega^c$ satisfy

$$|x - y| \leq \delta \operatorname{diam}(\Omega), \operatorname{dist}(x, \partial\Omega) \leq \delta \operatorname{diam}(\Omega), \text{ and } \operatorname{dist}(y, \partial\Omega) \leq \delta \operatorname{diam}(\Omega)$$

with our choice of δ . Indeed, regarding Lemma 2.14 with our approach above, it suffices to show that when n is large enough we have

$$|x - y| \leq \delta \operatorname{diam}(D_n), \operatorname{dist}(x, \partial D_n) \leq \delta \operatorname{diam}(D_n), \text{ and } \operatorname{dist}(y, \partial D_n) \leq \delta \operatorname{diam}(D_n);$$

notice that $x, y \in \tilde{D}_n$ for any $n \in \mathbb{N}$. This follows from the uniform continuity of φ .

Some of the lemmas below also hold for these piecewise hyperbolic geodesics. However, in order to simplify the statements of lemmas, and since those additional curves are not needed in what follows, we only prove the lemmas in the case where the end points of piecewise hyperbolic geodesics are on the boundary of Ω .

The following lemma justifies our terminology.

Lemma 4.2. *Let Ω be a bounded simply connected domain whose complement is C_1 -quasiconvex. Then there exist constants $C(C_1)$ and $\delta = \delta(C_1) > 0$ such that, for every pair of (distinct) $x, y \in \partial\Omega$ with*

$$|x - y| \leq \delta \operatorname{diam}(\Omega), \tag{4.2}$$

any piecewise hyperbolic geodesic γ joining x to y in Ω^c satisfies

$$\ell(\gamma[z, w]) \leq C(C_1)|z - w|,$$

whenever $z, w \in \gamma$ and $\gamma[z, w]$ is any subcurve of γ joining z and w . In particular, γ is an injective curve (when parametrized by arc length).

Additionally, for any $1 \leq i \leq N$ with $\gamma \cap \tilde{\Omega}_i \neq \emptyset$, we have that $\gamma \cap \tilde{\Omega}_i$ is a hyperbolic geodesic in $\tilde{\Omega}_i$.

Proof. Let $\gamma[z, w]$ be a subcurve of γ joining z and w . Then by our construction of the piecewise hyperbolic geodesics, there are $t_n < s_n$ such that the restrictions of γ_n to $[t_n, s_n]$ converge to $\gamma[z, w]$. We may assume that $z_n = \gamma_n(t_n)$ converges to z and that $w_n = \gamma_n(s_n)$ converges to w . Hence by the lower semi-continuity of length (as a functional on curves), Corollary 2.11 and Lemma 2.13 one has

$$\ell(\gamma[z, w]) \lesssim \liminf_{n \rightarrow \infty} \ell(\gamma_n[z_n, w_n]) \lesssim \liminf_{n \rightarrow \infty} |z_n - w_n| \lesssim |z - w|.$$

Hence our first claim follows.

For the final claim, let $\tilde{\varphi}_n: \mathbb{R}^2 \setminus \overline{\mathbb{D}} \rightarrow \tilde{D}_n$ to be conformal for each $n \in \mathbb{N}$, and extend it to $\overline{\mathbb{R}^2} \setminus \overline{\mathbb{D}}$ by setting $\tilde{\varphi}_n(\infty) = \infty$. Also when $i \geq 2$ and $\gamma \cap \tilde{\Omega}_i \neq \emptyset$, let $x_0 \in \tilde{\Omega}_i$ be a point of maximal distance to $\partial\tilde{\Omega}_i$; for $\tilde{\Omega}_1$ we set $x_0 = \infty$.

Denote by $\phi: \overline{\mathbb{R}^2} \setminus \overline{\mathbb{D}} \rightarrow \overline{\mathbb{R}^2} \setminus \overline{\mathbb{D}}$ a Möbius transformation such that $\phi(\infty) = \tilde{\varphi}_n^{-1}(x_0)$. Then the compositions $\tilde{\varphi}_n \circ \phi: \overline{\mathbb{R}^2} \setminus \overline{\mathbb{D}} \rightarrow \tilde{D}_n \cup \{\infty\}$ form a normal family by [30, Theorem 19.2]. Hence there is a subsequence that converges locally uniformly to a conformal map $\tilde{\varphi}: \overline{\mathbb{R}^2} \setminus \overline{\mathbb{D}} \rightarrow \tilde{\Omega}_i$; see [30, Theorem 21.1]. Notice that hyperbolic geodesics are invariant under conformal maps. Therefore γ_n can be regarded as a hyperbolic geodesic induced by $\tilde{\varphi}_n \circ \phi$ on the Riemann sphere by the uniqueness of the limit, and hence also γ as induced by $\tilde{\varphi} \circ \phi$. Thus the part of γ in $\tilde{\Omega}_i$ is a hyperbolic geodesic; it easily follows from the above argument that $\gamma \cap \tilde{\Omega}_i$ is connected. \square

We need the following further properties of piecewise hyperbolic geodesics.

Lemma 4.3. *Let Ω and δ be as in Lemma 4.2, and $x, y \in \partial\Omega$. Given any piecewise hyperbolic geodesic $\Gamma \subset \Omega^c$ associated to the pair x, y one has*

$$\Gamma \cap \partial\Omega \subset \gamma \cap \partial\Omega,$$

for any curve $\gamma \subset \Omega^c$ connecting x and y . Moreover, if the complement of Ω satisfies the curve condition (1.2), then

$$\int_{\Gamma} \frac{1}{\chi_{\mathbb{R}^2 \setminus \partial\Omega}(z)} ds(z) \leq C|x - y|,$$

where the constant C depends only on the constant in (1.2).

Proof. Let $\gamma \subset \Omega^c$ be an arbitrary curve joining $x, y \in \partial\Omega$. Denote the corresponding hyperbolic geodesics approximating Γ by $\Gamma_n \subset \tilde{D}_n$ with endpoints x_n, y_n , according to the definition of Γ .

Suppose $z \in \Gamma \cap \partial\Omega$. We claim that $z \in \gamma \cap \partial\Omega$. We may assume that $z \neq x, z \neq y$. Let $\{z_n\}, z_n \in \Gamma_n$ be a sequence of points converging to z , and define for every $n \geq 2$,

$$B_n = B\left(z_n, \frac{1}{2} \text{dist}(z_n, \partial D_n)\right).$$

Since each B_n is a 2-Whitney-type set in \tilde{D}_n , then Lemma 2.4 gives a constant c independent of n such that $\gamma' \cap cB_n \neq \emptyset$ for any $\gamma' \subset \tilde{D}_n$ joining x_n and y_n . Since $x_n \rightarrow x$ and $y_n \rightarrow y$, then when n is large enough so that

$$|x_n - z_n| \geq C(C_1)|x - x_n|, \quad |y_n - z_n| \geq C(C_1)|y - y_n|,$$

the $C(C_1)$ -quasiconvexity of \tilde{D}_n given by Lemma 2.13 implies that there exist curves in \tilde{D}_n joining x to x_n and y to y_n , respectively, such that these two curves do not intersect cB_n . By concatenating them with γ , we apply Lemma 2.4 to conclude that

$$\gamma \cap cB_n \neq \emptyset \tag{4.3}$$

when n is large enough; recall that $\gamma \subset \Omega^c \subset \tilde{D}_n$.

Observe that by $z \in \Gamma \cap \partial\Omega$, we have that

$$\lim_{n \rightarrow \infty} \text{dist}(z_n, \partial D_n) = 0.$$

Then by the assumption $z_n \rightarrow z$ when $n \rightarrow \infty$, we conclude that $cB_n \rightarrow z$, and hence $z \in \gamma$ by (4.3). Thus $z \in \gamma \cap \partial\Omega$. This proves the claim, and the first part of the lemma follows.

The second part follows from Lemma 4.2 together with the first part applied to a curve γ satisfying (1.2); note that Γ is an injective curve. \square

For further reference we record the following consequences of the above lemmas.

Lemma 4.4. *Given Ω as in Lemma 4.2, and a piecewise hyperbolic geodesic Γ , a subcurve $\Gamma[x, y]$ of Γ and a curve γ joining x to y in Ω^c , we have*

$$\Gamma[x, y] \cap \partial\Omega \subset \gamma \cap \partial\Omega.$$

Moreover $\Gamma \cap \partial\tilde{\Omega}_i$ consists of at most two points for each $1 \leq i \leq N$, and it is a doubleton if and only if $\Gamma \cap \tilde{\Omega}_i$ is a hyperbolic geodesic joining boundary points.

Proof. Let x_0, y_0 be the end points of Γ on $\partial\Omega$. Then the concatenation of three curves, the subcurve $\Gamma[x, x_0]$ of Γ , γ , and the subcurve $\Gamma[y, y_0]$ of Γ , is a curve β joining x_0 to y_0 . Then the first conclusion follows from Lemma 4.3 applied to Γ and β .

If $\Gamma \cap \partial\tilde{\Omega}_i$ has more than two points, then consider the first and the last points of Γ intersecting $\partial\tilde{\Omega}_i$ according to its parametrization, and join them by a hyperbolic geodesic α inside $\tilde{\Omega}_i$; recall that $\tilde{\Omega}_i$ is Jordan by Lemma 2.10. Then we obtain a new curve γ by rerouting the subarc of Γ via α . This contradicts Lemma 4.3 for γ since we have assumed that $\Gamma \cap \partial\tilde{\Omega}_i$ has more than two points. Thus $\Gamma \cap \partial\tilde{\Omega}_i$ has at most two points.

Let us show the last part of the lemma. We may assume that $\tilde{\Omega}_i$ is bounded; otherwise we just apply suitable Möbius transformations on the Riemann sphere. If $\Gamma \cap \tilde{\Omega}_i$ is a hyperbolic geodesic joining boundary points, then $\Gamma \cap \partial\tilde{\Omega}_i$ consists of two points since $\tilde{\Omega}_i$ is Jordan. For the other direction, by Lemma 4.2 it suffices to show that $\Gamma \cap \tilde{\Omega}_i$ is non-empty. On the contrary let us assume that $\Gamma \cap \tilde{\Omega}_i = \emptyset$. Then by joining the given two points $x, y \in \Gamma \cap \partial\tilde{\Omega}_i$ with the hyperbolic geodesic $\alpha \subset \tilde{\Omega}_i$, we obtain a Jordan curve γ' by concatenating $\Gamma[x, y]$ with α ; see Lemma 4.2 and note that the open subarc $\Gamma(x, y)$ is contained in the exterior of $\tilde{\Omega}$ by our assumption and the conclusion of the previous paragraph. However by our construction, $\partial\Omega$ intersects both the interior and the exterior of the Jordan domain given by γ' . This implies that there are points belonging to Ω on both sides of γ' . This contradicts the Jordan curve theorem since Ω is connected while $\gamma' \subset \Omega^c$. Therefore we conclude that $\Gamma \cap \tilde{\Omega}_i \neq \emptyset$. \square

The following corollary is a by-product of the lemmas above.

Lemma 4.5. *Given a domain Ω together with a pair of points $x, y \in \partial\Omega$ as in Lemma 4.2, there exists a unique piecewise hyperbolic geodesic joining them, up to a reparametrization.*

Proof. The existence follows directly from Lemma 4.2, and we only need to show the uniqueness.

Suppose that there are two piecewise hyperbolic geodesics γ_1, γ_2 connecting x, y . Then by Lemma 4.3 we have

$$\gamma_1 \cap \partial\Omega = \gamma_2 \cap \partial\Omega. \quad (4.4)$$

Since $\Omega^c = \partial(\Omega^c) \cup \tilde{\Omega} = \partial\Omega \cup \tilde{\Omega}$, we then only need to show that γ_1 coincides with γ_2 componentwise. This follows from (4.4), Lemma 4.2 and Lemma 4.4 according to the uniqueness of hyperbolic geodesics in Jordan domains. \square

In order to deal with self-intersections of the boundary, it is convenient to classify points on the boundary in terms of their preimages.

Definition 4.6. Let $\varphi: \mathbb{D} \rightarrow \Omega$ be conformal and assume that it extends as a continuous map (still denoted by φ) to entire $\overline{\mathbb{D}}$. A point $x \in \partial\Omega$ is called a *cut-point* if $\varphi^{-1}(x)$ is not a singleton. A point x that is not a cut-point is called *one-sided*.

We warn the reader that our terminology above is not standard. See e.g. [27, Chapter 14] for a discussion on the relation with the usual topological definition of cut-points. As stated, the definition appears to depend on the choice of the continuous conformal map. This is not the case since any other (continuous) conformal map differs from the chosen one only by a precomposition via a Möbius self-map of the (closed) disk.

Lemma 4.7. Let $\Omega \subset \mathbb{R}^2$ be a bounded simply connected John domain. Then the set T of cut-points can be characterized as

$$T = \bigcup_{\gamma} \gamma^o \cap \partial\Omega, \quad (4.5)$$

where the union is over all piecewise hyperbolic geodesics γ associated to pairs of (suitable) points in $\partial\mathbb{D}$ and γ^o denotes the curve γ without its endpoints. Moreover, the union can equivalently be taken over a countable set of curves.

Proof. First of all, by part (2) of Lemma 2.8 we have a continuous map $\varphi: \overline{\mathbb{D}} \rightarrow \overline{\Omega}$ which is conformal in \mathbb{D} .

Let $\{x_i\}_{i=1}^{\infty} \subset \partial\mathbb{D}$ be dense. We first claim that

$$T \subset \bigcup \gamma_{i,j}^o \cap \partial\Omega. \quad (4.6)$$

where the union is taken over all the piecewise hyperbolic geodesics $\gamma_{i,j}$ joining those $\varphi(x_i)$ to $\varphi(x_j)$ with

$$|\varphi(x_i) - \varphi(x_j)| \leq \delta \operatorname{diam}(\Omega).$$

Recall that $\gamma_{i,j}$ is obtained via a subsequence of the domains \tilde{D}_n . Since we only have countably many curves, by a usual diagonal argument, we may assume that each of them is obtained via the same subsequence. Observe that by Lemma 4.2, these curves are all injective.

Fix $z \in T$. By the definition of cut-points there exist $z_1, z_2 \in \partial\mathbb{D}$, $z_1 \neq z_2$, such that $\varphi(z_1) = \varphi(z_2) = z$. Recall here that the image of any nontrivial arc of $\partial\mathbb{D}$ under our conformal map that is continuous up to the boundary is connected and not a singleton. By the density of $\{x_i\}_{i=1}^{\infty}$ there exist $i, j \in \mathbb{N}$ such that

$$\varphi(x_i) \neq z \neq \varphi(x_j), \quad |\varphi(x_i) - \varphi(x_j)| \leq \delta \operatorname{diam}(\Omega)$$

with $\varphi(x_i) \neq \varphi(x_j)$ and x_i, x_j divide $\partial\mathbb{D}$ into two connected components; one contains z_1 and the other contains z_2 . The curve $\varphi([0, z_1] \cup [0, z_2])$ is Jordan since $z_1 \neq z_2$ while $\varphi(z_1) = \varphi(z_2)$, and φ is injective in \mathbb{D} , where $[0, z_i]$ is the (radial) line segment connecting the origin and z_i for $i = 1, 2$.

We show that the points $\varphi(x_i)$ and $\varphi(x_j)$ belong to different connected components of $\mathbb{R}^2 \setminus \varphi([z_1, 0] \cup [0, z_2])$. To begin with, since φ is continuous up to the boundary, then

$$\varphi([z_1, 0] \cup [0, z_2])$$

divides Ω into two components, coming from the two components of $\mathbb{D} \setminus ([z_1, 0] \cup [0, z_2])$. Recall that x_i and x_j are in different connected components of $\partial\mathbb{D} \setminus \{z_1, z_2\}$. Then by taking two sequences of points inside the components of $\mathbb{D} \setminus ([z_1, 0] \cup [0, z_2])$ approaching x_i and x_j , respectively, we conclude that $\varphi(x_i)$ and $\varphi(x_j)$ are not in the same component of $\mathbb{R}^2 \setminus \varphi([z_1, 0] \cup [0, z_2])$ since φ is continuous up to the boundary and neither $\varphi(x_i)$ nor $\varphi(x_j)$ belong to $\varphi([z_1, 0] \cup [0, z_2])$. Consequently, $\gamma_{i,j}$ has to intersect the Jordan curve $\varphi([z_1, 0] \cup [0, z_2])$. Since $\varphi([z_1, 0] \cup [0, z_2]) \cap \Omega^c = \{z\}$ and $\gamma_{i,j} \subset \Omega^c$, we know that

$$\gamma_{i,j} \cap \varphi([z_1, 0] \cup [0, z_2]) = \{z\}$$

and thus the claim is proved since $\varphi(x_i)$, z , and $\varphi(x_j)$ are distinct. Therefore we obtain (4.6).

Let us then show the other inclusion. Let γ be a piecewise hyperbolic geodesic joining two different points in $\partial\Omega^c$ and let $z \in \gamma \cap \partial\Omega$ while $\varphi(y_1) \neq z \neq \varphi(y_2)$. Let $y_1, y_2 \in \partial\mathbb{D}$ be such that $\varphi(y_1)$ and $\varphi(y_2)$ are the endpoints of γ . Now the curve

$$\gamma' := \varphi([y_1, 0] \cup [0, y_2]) \cup \gamma$$

is Jordan by Lemma 4.2 and it divides Ω into two connected components Ω_1 (the exterior component) and Ω_2 (the interior component). Let Ω' be the Jordan domain given by γ' .

Let us assume contrary to the claim that $z \notin T$. Then there exists $r > 0$ such that $B(z, r)$ intersects only one of the components, either Ω_1 or Ω_2 ; otherwise we have two sequences of points converging to z in different components, i.e. there exist two sequences in different component of $\mathbb{D} \setminus ([y_1, 0] \cup [0, y_2])$ approximating $\varphi^{-1}(z)$, which by the continuity of φ contradicts the assumption that $z \notin T$ but $z \in \varphi([y_1, 0] \cup [0, y_2])$.

Assume that $B(z, r) \cap \Omega_1 = \emptyset$. Since $z \in \gamma \cap \partial\Omega$, by the first part of Lemma 4.3 we know that there is no path connecting $\varphi(y_1)$ and $\varphi(y_2)$ in $\Omega^c \setminus \{z\}$. Towards a contradiction, we apply the fact that $\gamma \subset \partial\Omega'$ and the Jordan-Schoenflies theorem to Ω' .

Indeed, by the Jordan-Schoenflies theorem (see e.g. [29]) there is a homeomorphism $\phi: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that $\phi(\Omega') = \mathbb{D}$, and then $\phi(\gamma) \subset \partial\mathbb{D}$. Recall that $\varphi(y_1) \neq z \neq \varphi(y_2)$. Choose $0 < \epsilon < 1$ such that

$$\min\{|\varphi(y_1) - z|, |\varphi(y_2) - z|\} > \epsilon r.$$

Then $\varphi(y_1), \varphi(y_2) \notin B(z, \epsilon r)$, and $\phi(B(z, \epsilon r))$ is also a Jordan domain. Certainly $\phi(z)$ is an interior point of $\phi(B(z, \epsilon r))$, and hence there exists $\delta > 0$ such that $B(\phi(z), \delta) \subset \phi(B(z, \epsilon r))$. Denote by σ the subarc of $\partial(\mathbb{D} \cup B(\phi(z), \delta))$ which joins $\phi(\varphi(y_1))$ and $\phi(\varphi(y_2))$ and reroutes $\phi(\gamma)$, and let $\Gamma = \phi^{-1}(\sigma)$.

Observe first that $\Gamma \subset \Omega^c$ by our construction and the assumptions that $\Omega_2 \subset \Omega'$ and $B(z, \epsilon r) \cap \Omega_1 = \emptyset$. Also Γ joins $\varphi(y_1)$ and $\varphi(y_2)$ by not passing through z . Therefore we obtain the desired curve, which leads to a contradiction. A similar argument can be applied to the case where $B(z, r) \cap \Omega_2 = \emptyset$.

The existence of countably many curves follows directly from the proof above. \square

An immediate consequence of Lemma 4.3 and Lemma 4.7 is the following corollary.

Corollary 4.8. *Let $\Omega \subset \mathbb{R}^2$ be a simply connected bounded John domain whose complement satisfies the curve condition (1.2) in Theorem 1.1. Then the set of cut-points of $\partial\Omega$ has \mathcal{H}^1 -measure zero.*

Let us record the following consequence of the proof of Lemma 4.7.

Lemma 4.9. *Let $\Omega \subset \mathbb{R}^2$ be a simply connected bounded John domain whose complement is quasiconvex and $\varphi: \mathbb{D} \rightarrow \Omega$ be a conformal map. Also let γ be a piecewise hyperbolic geodesic joining two distinct points $\varphi(y_1), \varphi(y_2) \in \partial\Omega$, and let $z \in \gamma \cap \partial\Omega$ with $\varphi(y_1) \neq z \neq \varphi(y_2)$. Then z is a cut-point, and every open disk centered at z has non-trivial intersection both with Ω_1 and with Ω_2 , where Ω_1 and Ω_2 are the two components of $\Omega \setminus (\varphi([y_1, 0] \cup [0, y_2]) \cup \gamma) = \Omega \setminus (\varphi([y_1, 0] \cup [0, y_2]))$.*

5. SUFFICIENCY FOR SIMPLY CONNECTED DOMAINS

We construct the desired extension via a modification to our procedure for the Jordan case. The first obstacle is that we cannot anymore use hyperbolic triangles in the complementary domain since there need not be a complementary domain to work with. To overcome this, we use piecewise hyperbolic geodesics to obtain a desired decomposition in each of the components of the interior of the complement of Ω . This allows us to mimic the Jordan case, but we have to work harder to verify the regularity of our extension. Observe that by (1.2), Ω^c is C_1 -quasiconvex with the constant C_1 from (1.2).

5.1. Decomposition of the complement. Let $\varphi: \overline{\mathbb{D}} \rightarrow \overline{\Omega}$ be a conformal map which is continuous up to the boundary, given by (1) and (2) of Lemma 2.8. The decomposition of Ω is the same as in Section 3.1 for the Jordan case. The Whitney-type sets in Ω are denoted by $Q_k^{(j)}$ and $z_k^{(j)} = \varphi(x_k^{(j)})$ stand for the images of the endpoints on $\partial\mathbb{D}$ of the radial rays used in the decomposition of \mathbb{D} . Again our decomposition begins with a suitable k_0 according to the constant δ in the previous section such that

$$|z_{k_0}^{(j)} - z_{k_0}^{(j-1)}| \leq \delta \operatorname{diam}(\Omega).$$

This time the complement Ω^c will be simultaneously triangulated inside each connected component of $\tilde{\Omega} = \mathbb{R}^2 \setminus \overline{\Omega}$ using hyperbolic geodesics. For this purpose we consider all pairs $(z_k^{(j-1)}, z_k^{(j)})$ and pick for each of them a piecewise hyperbolic geodesic. It may happen that $z_k^{(j-1)} = z_k^{(j)}$ for some $j, k \in \mathbb{N}$, but recall that every piecewise hyperbolic geodesic is defined via labeling on $\partial\mathbb{D}$, namely $x_k^{(j-1)}$ and $x_k^{(j)}$ for this pair of points, and in this case the piecewise hyperbolic geodesic is a constant curve. Since we deal with a countable collection of curves and our piecewise hyperbolic geodesics are obtained via the Arzelà-Ascoli theorem, we may assume that each of them is obtained via the same subsequence of the conformal maps $\tilde{\varphi}_n$. In other words, all the components of $\tilde{\Omega}$ are triangulated at the same time.

Recall from Lemma 4.2 that a piecewise hyperbolic geodesic is a shortest curve, up to a multiplicative constant, such that its restriction to any of the connected components of $\tilde{\Omega}$ is a hyperbolic geodesic (or empty). Moreover by Lemma 4.2 and (3.2) we again have

$$\ell(\gamma_k^{(j)}) \lesssim |z_k^{j-1} - z_k^j| \lesssim \operatorname{diam}(Q_k^{(j)}), \quad (5.1)$$

with the constant depending only on C_1 .

Next we study the possible cases of degenerated hyperbolic triangles; see Figure 3.

Lemma 5.1. *For $k = 1, 2, 3$, let γ_k be a piecewise hyperbolic geodesics with two different endpoints $z_k, z_{k+1} \in \partial\Omega$ respectively; suppose that $z_4 = z_1$. Then there are two possibilities:*

- 1) each of the curves is contained in the union of the other two, and the common set of these three curves is a single point.
- 2) For some $1 \leq j_0 \leq N$, the intersection of each γ_k with $\tilde{\Omega}_{j_0}$ is a hyperbolic geodesic and these three geodesics form a hyperbolic triangle of $\tilde{\Omega}_{j_0}$. For any other $1 \leq j \leq N, j \neq j_0$, if two of the curves γ_k intersect $\tilde{\Omega}_j$, then the respective intersections overlap with each other.

Proof. First of all if $y \in \gamma_1 \cap \partial\Omega$, then γ_2 or γ_3 must contain y by Lemma 4.3. Namely

$$\gamma_1 \cap \partial\Omega \subset (\gamma_2 \cup \gamma_3) \cap \partial\Omega. \quad (5.2)$$

The analogous statements also hold for γ_2 and γ_3 .

Let y_0 be the last point along γ_1 from z_2 towards z_1 such that $y_0 \in \gamma_1 \cap \gamma_2$. We claim that $y_0 \in \partial\Omega$.

Suppose on the contrary that $y_0 \notin \partial\Omega$. Then $y_0 \in \tilde{\Omega}_j$ for some $1 \leq j \leq N$, and $\tilde{\Omega}_j$ is Jordan on the Riemann sphere by Lemma 2.10. Let y_1 be the first point on $\gamma_1 \cap \partial\tilde{\Omega}_j$ when we trace (along γ_1) towards z_2 from y_0 . Consider the subcurves $\gamma_{1,0}$ and $\gamma_{2,0}$ of γ_1, γ_2 , respectively, between y_0 and z_2 . By Lemma 4.4 we have that $y_1 \in \gamma_{2,0}$. It follows from Lemma 4.2 that the intersections of both γ_1 and γ_2 with $\tilde{\Omega}_j$ are hyperbolic geodesics that contain both y_1 and y_0 . By mapping $\tilde{\Omega}_j$ conformally to \mathbb{D} (or to the exterior of the closed unit disk if $\tilde{\Omega}_j$ is unbounded), the uniqueness of hyperbolic geodesics yields that the intersection of γ_1 with $\tilde{\Omega}_j$ equals to the intersection of γ_2 with $\tilde{\Omega}_j$. This contradicts the definition of y_0 .

Consider again the subcurves $\gamma_{1,0}$ and $\gamma_{2,0}$ from the previous paragraph. We conclude by Lemma 4.4 that

$$\gamma_{1,0} \cap \partial\Omega = \gamma_{2,0} \cap \partial\Omega. \quad (5.3)$$

Moreover, if $\gamma_{1,0}$ intersects $\tilde{\Omega}_j$, then this intersection β is by Lemma 4.2 a hyperbolic geodesic, say with endpoints w_1, w_2 . Then $w_1, w_2 \in \partial\Omega$ and it follows that $w_1, w_2 \in \gamma_{2,0}$. Consider the corresponding subcurve α of $\gamma_{2,0}$. Since $\beta \subset \tilde{\Omega}_j$, it follows from Lemma 4.4 that $\alpha \subset \tilde{\Omega}_i$ is a hyperbolic geodesic. As in the previous paragraph, we conclude that $\alpha = \beta$. Hence $\gamma_{1,0} \subset \gamma_{2,0}$. By reversing the roles of $\gamma_{1,0}$ and $\gamma_{2,0}$ we deduce that

$$\gamma_{1,0} = \gamma_{2,0}. \quad (5.4)$$

Suppose that $y_0 \in \gamma_3$. In this case, we repeat the argument from the previous paragraph first for the subcurve $\gamma_{1,2}$ of γ_1 between z_1 and y_0 and the subcurve $\gamma_{3,4}$ of γ_3 between $z_1 = z_4$ and y_0 and after that for the subcurve $\gamma_{3,3}$ of γ_3 between z_3 and y_0 and the subcurve $\gamma_{2,2}$ of γ_2 between z_3 and y_0 . This gives us $\gamma_{1,2} = \gamma_{3,4}$ and $\gamma_{3,3} = \gamma_{2,2}$ and we conclude that $\gamma_3 \subset \gamma_1 \cup \gamma_2$.

We are left with the case $y_0 \notin \gamma_3$. We claim that $y_0 \in \tilde{\Omega}_{j_0}$ for some j_0 such that $\gamma_1 \cap \tilde{\Omega}_{j_0}$ and $\gamma_2 \cap \tilde{\Omega}_{j_0}$ do not intersect. Indeed, by the definition of y_0 and (5.2) we have that

$$(\gamma_1 \setminus \gamma_{1,0}) \cap \gamma_2 = \emptyset$$

and

$$(\gamma_1 \setminus \gamma_{1,0}) \cap \partial\Omega \subset \gamma_3 \cap \partial\Omega.$$

Note that $\gamma_3 \cap \partial\Omega$ is a closed set, and $y_0 \in \partial\Omega$ while $y_0 \notin \gamma_3$. This implies that there is no sequence of points in $(\gamma_1 \setminus \gamma_{1,0}) \cap \partial\Omega$ converging to y_0 . Let y'_0 be the last point in




FIGURE 3. Here are the two possible cases of degenerated hyperbolic triangles.

$(\gamma_1 \setminus \gamma_{1,0}) \cap \partial\Omega$ towards y_0 . Then the open subcurve $\gamma_1(y_0, y'_0)$ is contained in the interior of Ω^c , and by the connectedness it is contained in some component $\tilde{\Omega}_{j_0}$ of $\tilde{\Omega}$. By applying Lemma 4.2 to γ_1 and the Jordan domain $\tilde{\Omega}_j$ we conclude the claim.

Now by Lemma 4.2, (5.2) and the definition of y_0 , the endpoint w_1 on $\partial\tilde{\Omega}_{j_0}$ of the part of γ_1 in $\tilde{\Omega}_{j_0}$ has to be contained in γ_3 . Arguing as above, we conclude that the remaining part of γ_1 coincides with a subcurve of γ_3 . Similarly, the endpoint $w_2 \in \partial\tilde{\Omega}$ of the part of γ_2 in $\tilde{\Omega}_{j_0}$ must be contained in γ_3 , and we conclude that the remaining part of γ_2 coincides with a subcurve of γ_3 . Finally, let $\gamma_{3,1,2}$ be the subcurve of γ_3 joining w_1, w_2 . Since the union of the (closures) of the subcurves of γ_1, γ_2 in $\tilde{\Omega}_{j_0}$ also joins w_1 to w_2 and only intersects $\partial\Omega$ in the set $\{y_0, w_1, w_2\}$ and $y_0 \notin \gamma_3$, Lemma 4.4 implies that $\gamma_{3,1,2}$ joins w_1 to w_2 in $\tilde{\Omega}_{j_0}$. By Lemma 4.2 all of $\gamma_{3,1,2}$ and the subcurves of γ_1, γ_2 in $\tilde{\Omega}_{j_0}$ are hyperbolic geodesics and it follows that they form a hyperbolic triangle. \square

By Lemma 5.1, we can define the (degenerated) hyperbolic triangles $R_k^{(j)}$ similarly as in the Jordan case, namely the bounded (relatively) closed set enclosed by the union of $\gamma_k^{(j)}$, $\gamma_{k+1}^{(2j-1)}$ and $\gamma_{k+1}^{(2j)}$, and there is at most one connected component $\tilde{\Omega}_i$ of $\tilde{\Omega}$ such that $R_k^{(j)} \cap \tilde{\Omega}_i$ is a (non-degenerated) hyperbolic triangle, denoted by $R_{k,i}^{(j)}$.

For every $k \geq 1$, $1 \leq j \leq 2^{k_0+k+1}$ and $1 \leq i \leq N$, set

$$\gamma_{k,i}^{(j)} = \gamma_k^{(j)} \cap \tilde{\Omega}_i.$$

Notice that $\gamma_{k,i}^{(j)}$ may well be empty for a choice of k, j, i . Nevertheless

$$\bigcup_{i=1}^N \gamma_{k,i}^{(j)} = \gamma_k^{(j)} \cap \tilde{\Omega};$$

see Lemma 4.7. We call $\gamma_{k,i}^{(j)}$ the i -th component of $\gamma_k^{(j)}$, call $\gamma_{k,i}^{(j)}$ the *mother* of $\gamma_{k+1,i}^{(2j-1)}$ and $\gamma_{k+1,i}^{(2j)}$, and refer to $\gamma_{k+1,i}^{(2j-1)}$ and $\gamma_{k+1,i}^{(2j)}$ (possibly empty) as *children* of $\gamma_{k,i}^{(j)}$. By construction each mother has two children, and every child has a mother; see Lemma 5.1. Moreover, $\gamma_{k,i}^{(j_1)}$ is called a *sibling* of $\gamma_{k,i}^{(j_2)}$ if they have the same mother. We say that $\gamma_{k,i}^{(j)}$ is in the *first generation* if $k = 1$, that two curves $\gamma_{k_1,i_1}^{(j_1)}$, $\gamma_{k_2,i_2}^{(j_2)}$ are in the *same generation* if $k_1 = k_2$, and that the generation of $\gamma_{k_1,i_1}^{(j_1)}$ is higher than that of $\gamma_{k_2,i_2}^{(j_2)}$ if $k_1 < k_2$.

Notice that $\gamma_{k,i}^{(j)}$ may potentially coincide with the i -th component of some other piecewise hyperbolic geodesic. The following lemma describes the potential overlaps.

Lemma 5.2. *If $k_1 < k_2$, then $\gamma_{k_1,i}^{(j_1)}$ and $\gamma_{k_2,i}^{(j_2)}$ are not siblings. Furthermore, if we have that $\gamma_{k_1,i}^{(j_1)}$ coincides with $\gamma_{k_2,i}^{(j_2)}$, $k_1 < k_2$ for some i , j_1 and j_2 , then the mother of $\gamma_{k_2,i}^{(j_2)}$ also coincides with $\gamma_{k_1,i}^{(j_1)}$.*

Similarly if $\gamma_{k,i}^{(j_1)}$ and $\gamma_{k,i}^{(j_2)}$ coincide with each other but they are not siblings, then the mother of $\gamma_{k,i}^{(j_2)}$ also coincides with $\gamma_{k,i}^{(j_1)}$.

Proof. It is clear that $\gamma_{k_1,i}^{(j_1)}$ and $\gamma_{k_2,i}^{(j_2)}$ are not in the same generation since $k_1 < k_2$. Thus they are not siblings.

Let us show the second claim where $\gamma_{k_1,i}^{(j_1)}$ coincides with $\gamma_{k_2,i}^{(j_2)}$, $k_1 < k_2$ for some i , j_1 and j_2 . Recall that piecewise hyperbolic geodesics are obtained by taking limits of hyperbolic geodesics in Jordan domains \tilde{D}_n . Suppose that the end points of $\gamma_{k_1,i}^{(j_1)}$ are x and y . Then for each $l = 1, 2$ there exists an approaching sequence of $\gamma_{n,l} \subset \tilde{D}_n$ for $\gamma_{k_l}^{(j_l)}$ such that $x_{n,l}, y_{n,l} \in \gamma_{n,l}$ approximate x and y , respectively. Let $\alpha_n \subset \tilde{D}_n$ be the hyperbolic geodesic joining $x_{n,1}$ and $x_{n,2}$. Likewise $\beta_n \subset \tilde{D}_n$ is the hyperbolic geodesic connecting $y_{n,1}$ and $y_{n,2}$.

Recall that our choice of δ_1 in Section 4 ensures the images under $\tilde{\varphi}_n: \tilde{D}_n \rightarrow \mathbb{R}^2 \setminus \overline{\mathbb{D}}$ of the hyperbolic geodesics α_n and β_n to be contained in $B(0, 10)$. Then according to Corollary 2.11

$$\ell(\alpha_n) \lesssim |x_{n,1} - x_{n,2}|, \quad \ell(\beta_n) \lesssim |y_{n,1} - y_{n,2}| \quad (5.5)$$

for sufficiently large n , where the constants depend only on C_1 . This allows us to take the limit of α_n and β_n by Arzelà-Ascoli lemma, up to relabeling the sequences. Since $x_{n,l} \rightarrow x$ and $y_{n,l} \rightarrow y$ as $n \rightarrow \infty$ for $l = 1, 2$, we conclude from (5.5) that $\alpha_n \rightarrow x$ and $\beta_n \rightarrow y$.

On the other hand, by the geometry of hyperbolic geodesics in the exterior of the unit disk, via conformal mappings, we conclude that α_n and β_n have non-trivial intersections with $\gamma'_{n,2}$, where the sequence $\{\gamma'_{n,2}\}$ approaches the mother of $\gamma_{k_2,i}^{(j_2)}$. Thus the mother of $\gamma_{k_2,i}^{(j_2)}$ also goes through x and y , and our conclusion follows from Lemma 4.4.

The second part of the lemma follows from a similar argument with notational changes. \square

Let $[\gamma_{k,i}^{(j)}]$ be the collection of all the i -th components of our piecewise hyperbolic geodesics which coincide with a non-empty curve $\gamma_{k,i}^{(j)}$ in $\tilde{\Omega}_i$ (see Lemma 4.2) and order the elements in it as follows: $\gamma_{k_1,i}^{(j_1)}$ is older than $\gamma_{k_2,i}^{(j_2)}$ if and only if either $k_1 < k_2$ or $k_1 = k_2$ but $j_1 < j_2$.

We denote the oldest one in $[\gamma_{k,i}^{(j)}]$ by $\mathcal{A}[\gamma_{k,i}^{(j)}]$, and call it *the ancestor of $\gamma_{k,i}^{(j)}$* . So as to define an extension operator conveniently later, we force $\gamma_{1,i}^{(1)} \in [\gamma_{k,i}^{(j)}]$ if $\gamma_{1,i}^{(j)} \in [\gamma_{k,i}^{(j)}]$ for some $2 \leq j \leq 2^{k_0+k+1}$; it is just an artificial curve which is only used in the next subsection, and we suppress the abuse of notation here.

To clarify the definition of an ancestor, consider the curves $\gamma_{k,i}^{(j)}$, $\gamma_{k+1,i}^{(2j-1)}$ and $\gamma_{k+1,i}^{(2j)}$ enclosing a non-degenerated hyperbolic triangle $R_{k,i}^{(j)}$. Then Lemma 5.2 shows that

$$\mathcal{A}[\gamma_{k+1,i}^{(2j-1)}] = \gamma_{k+1,i}^{(2j-1)} \quad \text{and} \quad \mathcal{A}[\gamma_{k+1,i}^{(2j)}] = \gamma_{k+1,i}^{(2j)}. \quad (5.6)$$

Lemma 5.3. *Each generation in $[\gamma_{k,i}^{(j)}]$ contains at most two distinct elements (except the first generation which has at most three elements). Moreover if the highest (but not the first) generation of $[\gamma_{k,i}^{(j)}]$ contains two distinct elements, then they are siblings.*

Proof. We prove the first claim by contradiction. Suppose that $[\gamma_{k,i}^{(j)}]$ contains three distinct elements γ_l , $l = 1, 2, 3$ from the same generation, and note that they coincide in the component $\tilde{\Omega}_i$ by definition. By joining each of the two end points of γ_l via a hyperbolic geodesic in Ω starting from the image of the origin under the conformal map $\varphi: \mathbb{D} \rightarrow \Omega$, we obtain a Jordan domain Ω_l via concatenating the two hyperbolic geodesics together with γ_l , for every $l = 1, 2, 3$ correspondingly; recall here that γ_l is injective by Lemma 4.2 for each $l = 1, 2, 3$.

We claim that these three domains are pairwise disjoint. Indeed recall that

$$D_n = \varphi(B(0, 1 - 2^{-n}))$$

with $\varphi: \mathbb{D} \rightarrow \Omega$ conformal, and that each $\tilde{\gamma}_l$ is the limit of a uniformly converging sequence, consisting of hyperbolic geodesics $\gamma_{l,n} \subset \tilde{D}_n$, where \tilde{D}_n is the exterior of D_n . Then the end points $x_{l,n}$, $y_{l,n}$ of $\gamma_{l,n}$ converge to the end points x_l , y_l of γ_l , respectively. By the uniform continuity of φ the hyperbolic geodesics $\varphi([0, \varphi^{-1}(x_{l,n})])$ joining $\varphi(0)$ to $x_{l,n}$ also converge uniformly to the hyperbolic geodesics joining $\varphi(0)$ to x_l (in D_n); an analogous statement holds for $\varphi([0, \varphi^{-1}(y_{l,n})])$. Therefore

$$\alpha_{l,n} := \varphi([0, \varphi^{-1}(x_{l,n})]) \cup \varphi([0, \varphi^{-1}(y_{l,n})]) \cup \gamma_{l,n}, \quad n \in \mathbb{N}$$

is a sequence of Jordan curves converging to a Jordan curve

$$\alpha_l = \varphi([0, \varphi^{-1}(x_l)]) \cup \varphi([0, \varphi^{-1}(y_l)]) \cup \gamma_l$$

uniformly as $n \rightarrow \infty$ for $l = 1, 2, 3$.

Note that for each $n \in \mathbb{N}$, the Jordan domains enclosed by $\alpha_{l,n}$'s are pairwise disjoint since γ_l are in the same generation. Then the claim follows from the uniform convergence of $\alpha_{l,n}$. Indeed if two of Ω_1 , Ω_2 , Ω_3 were to intersect, then there would exist a disk $B(z, 3r)$ contained in the intersection, say, of Ω_1 and Ω_2 . However, for each $l = 1, 2$, when n is large enough, every point in $\alpha_{l,n}$ has at most distance r to α_l . Since α_l and $\alpha_{l,n}$ are Jordan, we conclude by the triangle inequality that $B(z, r)$ is in the intersection of the Jordan domains given by $\alpha_{1,n}$ and $\alpha_{2,n}$. This is a contradiction.

Next we show that, since α_l 's coincide in $\tilde{\Omega}_i$ for each $l = 1, 2, 3$, at least two of the corresponding Jordan domains have non-trivial intersection; this gives the contradiction. In

fact, by the Jordan-Schoenflies theorem (see e.g. [29]) there is a homeomorphism $\phi: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that the curve $\gamma_l \cap \tilde{\Omega}_i$ is mapped to the interval $[-1, 1]$ on the real line; since γ_l , $l = 1, 2, 3$ coincide, our map ϕ does not depend on l . Notice that Ω_l are also mapped to Jordan domains. Then there exists a constant $0 < r < 1$ such that $B(0, r) \cap \phi(\Omega_l)$ is a half disk for each $l = 1, 2, 3$; indeed choose

$$r = \min_{l=1,2,3} \text{dist} \left(0, \phi(\partial\Omega_l \setminus (\gamma_l \cap \tilde{\Omega}_i)) \right).$$

This implies that at least two of the half disks have a non-trivial intersection and that the same holds for Ω_l 's since ϕ is a homeomorphism. All in all we have shown the first part of the lemma.

The second part of the lemma follows from the second part of Lemma 5.2. \square

Fix a curve $\gamma_{k,i}^{(j)}$. Recall that $[\gamma_{k,i}^{(j)}]$ is the (equivalence) class of (subarcs of) piecewise hyperbolic geodesics coinciding with $\gamma_{k,i}^{(j)}$. For each curve $\gamma \in [\gamma_{k,i}^{(j)}]$, we define the *family chain* $\mathcal{F}(\gamma)$ of γ as the finite ordered set $\mathcal{F}(\gamma) = \{\gamma_1, \gamma_2, \dots, \gamma_l\}$, consisting of certain elements in $[\gamma_{k,i}^{(j)}]$ and with the property that γ_n is the mother or sibling of γ_{n+1} for $1 \leq n \leq l-1$, obtained via the following procedure. Define $\hat{\gamma}_1 = \gamma$. Set $\hat{\gamma}_2$ to be the mother of $\hat{\gamma}_1$ and continue inductively; see Lemma 5.2. This procedure stops after a finite number of steps, when we reach $\hat{\gamma}_{l-1} = \gamma_{k_0,i}^{j_0}$ such that the mother of $\gamma_{k_0,i}^{j_0}$ is not in $[\gamma_{k,i}^{(j)}]$, or $k_0 = 1$; namely $\gamma_{k_0,i}^{j_0}$ is in the highest generation of $[\gamma_{k,i}^{(j)}]$ (by Lemma 5.2). Then define $\hat{\gamma}_l = \mathcal{A}[\gamma_{k,i}^{(j)}]$; note that it is possible that $\hat{\gamma}_l = \hat{\gamma}_{l-1}$. We now set $\gamma_n = \hat{\gamma}_{l-n+1}$, $1 \leq n \leq l$. Observe that for $n \geq 3$, γ_n is a child of γ_{n-1} by our construction.

We say that a family chain $\mathcal{F}(\gamma)$ is *maximal* if it is maximal (with respect to set inclusion) amongst all family chains of elements in the curve family $[\gamma]$; recall that $\mathcal{F}(\gamma') \subset [\gamma]$ as a set for every $\gamma' \in [\gamma]$. Lemma 5.3 together with Lemma 5.1 and Lemma 5.2 tells that, from each $[\gamma_{k,i}^{(j)}]$, one can extract at most two different maximal family chains. Now we are ready to construct our extension operator. See Figure 4.

5.2. Definition of the extension. Let $u \in W^{1,\infty}(\Omega)$; then $u \in W^{1,1}(\Omega)$. As in the Jordan case, we define

$$a_k^{(j)} = \int_{Q_k^{(j)}} u(x) dx.$$

We still associate this value to $\gamma_k^{(j)}$ and write it as $a_{\gamma_k^{(j)}}$; we associate this value also to all the subarcs $\gamma_{k,i}^{(j)}$ of $\gamma_k^{(j)}$. However, one cannot define our extension directly via them and Lemma 3.2 because of the degeneracy.

To overcome this, we use the terminology defined in the previous subsection. We define $Eu(x) = a_{\mathcal{A}[\gamma_{k,i}^{(j)}]}$ on $\gamma_{k,i}^{(j)}$, where $a_{\mathcal{A}[\gamma_{k,i}^{(j)}]}$ is the value associated to $\mathcal{A}[\gamma_{k,i}^{(j)}]$. Then Eu is well defined on these hyperbolic geodesics by the definition of $[\gamma_{k,i}^{(j)}]$.

Next we associate to each non-degenerated $R_{k,i}^{(j)}$ in $\tilde{\Omega}_i$ a function $\phi_k^{(j)} \in W^{1,1}(R_k^{(j)})$ so that (recall again that there is at most one possible i such that $R_{k,i}^{(j)}$ is non-degenerated by




FIGURE 4. A collection of hyperbolic geodesics (in the approximation to the piecewise hyperbolic geodesics in the picture (b)) related to the elements in $[\gamma_{k,i}^{(j)}]$ is illustrated in the figure (a). Here the collection $[\gamma_{k,i}^{(j)}]$ gives rise to two different maximal family chains: One corresponds to (the limit of) $\{\alpha_3, \alpha_3, \alpha_2, \alpha_1\}$ and the other corresponds to (the limit of) $\{\alpha_3, \alpha_4\}$.

Lemma 5.1.)

$$\phi_k^{(j)}(x) = \begin{cases} a_{\mathcal{A}[\gamma_k^{(j)}]}, & x \in \gamma_k^{(j)} \cap R_k^{(j)} \\ a_{\mathcal{A}[\gamma_{k+1}^{(2j-1)}]}, & x \in \gamma_{k+1}^{(2j-1)} \cap R_k^{(j)} \\ a_{\mathcal{A}[\gamma_{k+1}^{(2j)}]}, & x \in \gamma_{k+1}^{(2j)} \cap R_k^{(j)} \end{cases}.$$

We choose $\phi_k^{(j)}$ to be the function ϕ from Lemma 3.2 with the boundary values given above. Then we have the gradient estimate (3.5) that will be employed later. Observe that since $R_{k,i}^{(j)}$ is non-degenerated, via (5.6) we actually have

$$\mathcal{A}[\gamma_{k+1,i}^{(2j-1)}] = \gamma_{k+1,i}^{(2j-1)} \quad \text{and} \quad \mathcal{A}[\gamma_{k+1,i}^{(2j)}] = \gamma_{k+1,i}^{(2j)}.$$

On the rest of $\tilde{\Omega}$ we define $Eu = a_{\gamma_1^{(1)}}$. This is consistent because we forced $\gamma_{1,i}^{(1)} \in [\gamma_{k,i}^{(j)}]$ if $\gamma_{1,i}^{(j)} \in [\gamma_{k,i}^{(j)}]$ for some $2 \leq j \leq 2^{k_0+k+1}$. To conclude, the operator E is defined by

$$Eu(x) = \begin{cases} u(x), & x \in \Omega \\ a_{\mathcal{A}[\gamma_k^{(j)}]}, & x \in \gamma_k^{(j)} \\ \phi_k^{(j)}(x), & x \text{ is in the interior of a non-degenerated } R_{k,i}^j \\ a_{\gamma_1^{(1)}}, & \text{in the rest of } \tilde{\Omega} \end{cases}.$$

Observe that $Eu \in W_{loc}^{1,1}(\tilde{\Omega})$. Indeed by Lemma 3.2 and Lemma 5.1, Eu is locally Lipschitz continuous in $\tilde{\Omega}$; the consistency of boundary values between (non-degenerated) hyperbolic triangles is guaranteed by the definition of $\phi_k^{(j)}(x)$.

This time we estimate the Sobolev-norm of the extension over a non-degenerated hyperbolic triangle $R_{k,i}^{(j)} \subset \tilde{\Omega}_i$ by inserting the intermediate values given to the corresponding curves in the family chain. Namely by Lemma 3.2, (5.1) and (5.6),

$$\begin{aligned} \int_{R_{k,i}^{(j)}} |\nabla \phi_k^{(j)}(x)| dx &\lesssim \left(|a_{\mathcal{A}[\gamma_k^{(j)}]} - a_{k+1}^{(2j-1)}| \ell(\gamma_{k,i}^{(j)}) + |a_{\mathcal{A}[\gamma_k^{(j)}]} - a_{k+1}^{(2j)}| \ell(\gamma_{k,i}^{(j)}) \right. \\ &\quad \left. + |a_{k+1}^{(2j-1)} - a_{k+1}^{(2j)}| \min\{\ell(\gamma_{k+1,i}^{(2j-1)}), \ell(\gamma_{k+1,i}^{(2j)})\} \right) \\ &\lesssim \left(|a_k^{(j)} - a_{k+1}^{(2j-1)}| + |a_k^{(j)} - a_{k+1}^{(2j)}| + |a_{k+1}^{(2j-1)} - a_{k+1}^{(2j)}| \right) \text{diam}(Q_k^{(j)}) \\ &\quad + \sum_{\gamma_l \in \mathcal{F}(\gamma_{k,i}^{(j)})} |a_{\gamma_l} - a_{\gamma_{l+1}}| \ell(\mathcal{A}[\gamma_{k,i}^{(j)}]) \\ &= S_k^{(j)}(1) + S_k^{(j)}(2), \end{aligned}$$

where we used the fact $\ell(\mathcal{A}[\gamma_{k,i}^{(j)}]) = \ell(\gamma_{k,i}^{(j)})$. Here the constants depend only on C_1 in (1.2).

The first term $S_k^{(j)}(1)$ is the same as in the Jordan case, and so we know that, after summing over all the hyperbolic triangles, it can be controlled by $\|\nabla u\|_{L^1(\Omega)}$ up to a multiplicative

constant. Notice that here each $Q_k^{(j)}$ corresponds to at most one non-degenerated $R_{k,i}^{(j)}$ by Lemma 5.1.

Observe that

$$\sum_{i=1}^N \ell(\mathcal{A}[\gamma_{k,i}^{(j)}]) = \sum_{i=1}^N \ell(\gamma_{k,i}^{(j)}) \leq C(C_1) \text{diam}(Q_k^{(j)}), \quad (5.7)$$

by (5.1).

Fix i and two distinct non-degenerated triangles $R_{k,i}^{(j)}$, $R_{k',i}^{(j')}$. Then $\mathcal{F}(\gamma_{k,i}^{(j)})$ is disjoint from $\mathcal{F}(\gamma_{k',i}^{(j')})$ unless $[\gamma_{k,i}^{(j)}] = [\gamma_{k',i}^{(j')}]$, or equivalently $R_{k,i}^{(j)}$ and $R_{k',i}^{(j')}$ share the same boundary curve. Recall that in each $[\gamma_{k,i}^{(j)}]$ there are at most two different family chains starting from the highest generation to the lowest one, and hence the ancestor in each $[\gamma_{k,i}^{(j)}]$ is counted at most twice if $k \neq 1$; if $k = 1$ the multiplicity of $\gamma_{1,i}^{(1)}$ is at most 2^{k_0+3} . Hence, as a result of changing the order of summation together with (5.7) and (3.9) the following estimate holds:

$$\begin{aligned} \sum_{R_{k,i}^{(j)}} S_k^{(j)}(2) &= \sum_{R_{k,i}^{(j)}} \sum_{\gamma_l \in \mathcal{F}(\gamma_{k,i}^{(j)})} |a_{\gamma_l} - a_{\gamma_{l+1}}| \ell(\mathcal{A}[\gamma_{k,i}^{(j)}]) \\ &\lesssim \sum_{k,j} (|a_k^{(j)} - a_{k+1}^{(2j-1)}| + |a_k^{(j)} - a_{k+1}^{(2j-1)}| + |a_k^{(j)} - a_k^{(j-1)}| + |a_k^{(j)} - a_k^{(j+1)}|) \sum_{i=1}^N \ell(\mathcal{A}[\gamma_{k,i}^{(j)}]) \\ &\lesssim \sum_{k,j} (|a_k^{(j)} - a_{k+1}^{(2j-1)}| + |a_k^{(j)} - a_{k+1}^{(2j-1)}| + |a_k^{(j)} - a_k^{(j-1)}| + |a_k^{(j)} - a_k^{(j+1)}|) \text{diam}(Q_k^{(j)}) \\ &\lesssim \sum_{k,j} S_k^{(j)}(1), \end{aligned}$$

where the constants depend only on C_1 . Therefore the estimate for $\sum_{k,j} S_k^{(j)}(1)$ gives

$$\|\nabla Eu\|_{L^1(\tilde{\Omega})} \leq C(C_1) \|\nabla u\|_{L^1(\Omega)}.$$

Finally let B be a disk of radius $4 \text{diam}(\Omega)$ such that Ω is contained in $\frac{1}{2}B$. Similarly as in the Jordan case, by part (4) of Lemma 2.8, we have that

$$\begin{aligned} \|Eu\|_{L^1(B)} &\lesssim \sum_{R_{k,i}^{(j)}} \int_{R_{k,i}^{(j)}} |Eu(x)| dx + \int_{\Omega} |u(x)| dx \\ &\lesssim \sum_{\substack{k \geq 1 \\ 1 \leq j \leq 2^{k_0+k+1}}} \int_{Q_k^{(j)}} |u(x)| dx + \int_{\Omega} |u(x)| dx \lesssim \int_{\Omega} |u(x)| dx, \end{aligned} \quad (5.8)$$

where we used (3.4), (3.9) and the fact that

$$|R_k^{(j)}| \lesssim |Q_k^{(j)}|$$

coming from (3.2) and the fact that $Q_k^{(j)}$ is of $C(C_1)$ -Whitney-type. Here all the constants depend only on C_1 .

5.3. Continuity at one-sided points. By the curve condition (1.2) and Corollary 4.8, the set $T \subset \partial\Omega$ of cut-points has \mathcal{H}^1 -measure zero. Therefore if we can extend u continuously to the one-sided points, then we have defined Eu everywhere except for a set of \mathcal{H}^1 -measure zero.

Let us first show that we may assume u to be continuous at the one-sided points in the following sense:

Lemma 5.4. *Let Ω be a bounded simply connected J -John domain, and suppose that $u \in W^{1,\infty}(\Omega) \cap C(\Omega)$. Then we can extend u (uniquely) to \tilde{u} so that $\tilde{u} \in C(\bar{\Omega} \setminus T)$, where T is the collection of cut-points of $\partial\Omega$.*

Proof. Let $z \in \partial\Omega \setminus T$. Since $u \in C(\Omega)$, it suffices to show that $\lim_{i \rightarrow \infty} u(z_i)$ exists whenever $z_i \in \Omega$ satisfies $|z_i - z| \rightarrow 0$ as $i \rightarrow \infty$. Let $\varphi: \mathbb{D} \rightarrow \bar{\Omega}$ be continuous and conformal inside \mathbb{D} ; see part (2) of Lemma 2.8. By Lemma 2.9, we also have that $\varphi: (\mathbb{D}, |\cdot|) \rightarrow (\Omega, \text{dist}_\Omega)$ is uniformly continuous.

Since $z \notin T$, we have that $\varphi^{-1}(z)$ is a singleton; say $\varphi^{-1}(z) = w$. Our claim follows if $|\varphi^{-1}(z_i) - w| \rightarrow 0$ when $i \rightarrow \infty$; indeed since φ is uniformly continuous with respect to the dist_Ω -metric of Ω , then by the assumption that $u \in W^{1,\infty}(\Omega) \cap C(\Omega)$ we conclude that $\lim_{i \rightarrow \infty} u(z_i)$ exists. This is necessarily the case; otherwise a subsequence converges to some $w' \neq w$ and then $\varphi(w') = z$ by the continuity of φ at w' , contradicting the fact that $\varphi^{-1}(z)$ is a singleton. \square

Recall that by the definition of the ancestors and that of $\phi_k^{(j)}(x)$, Eu is (locally Lipschitz) continuous in $\tilde{\Omega}$. Then the continuity of Eu (under $u \in W^{1,\infty}(\Omega) \cap C(\Omega)$) restricted to $\mathbb{R}^2 \setminus H$ with $H := T \cup Z$ follows exactly as in the Jordan case, where Z is also defined like the previous one: the collection of the countably many points $z_k^{(j)}$ on the boundary. This time the fact that $y \notin \Gamma_s$ (defined in (3.13)) comes from Lemma 4.7, where we observed that, except for the endpoints, piecewise hyperbolic geodesics can meet the boundary only at cut-points. Therefore, by further defining $Eu(x) = \tilde{u}(x)$ for $x \in \partial\Omega$ for $u \in W^{1,\infty}(\Omega) \cap C(\Omega)$, we have that $Eu(x)$ is continuous in $\mathbb{R}^2 \setminus H$.

5.4. Absolute continuity. We show the absolute continuity of Eu along almost every horizontal or vertical line under the assumption that $\tilde{u} \in C(\Omega) \cap W^{1,\infty}(\Omega)$. By Lemma 5.4 we may assume that

$$u \in W^{1,\infty}(\Omega) \cap C(\bar{\Omega} \setminus T).$$

We begin by applying the Jordan case in the following way.

Since $\mathcal{H}^1(T) = 0$ by Corollary 4.8, for any $\epsilon > 0$, we find

$$U_\epsilon := \Omega \cap \bigcup_{z \in T} B(z, r_z)$$

with the property that the \mathcal{H}^1 -measure of its vertical and horizontal projections are less than ϵ ; the existence of such disks follows from the definition of \mathcal{H}^1 -measure and the fact that projections are 1-Lipschitz.

Since φ is a homeomorphism in \mathbb{D} , the set $\varphi^{-1}(U_\epsilon)$ is open. For each $z \in T$ and every $w \in \varphi^{-1}(\{z\})$, by the continuity of φ we may pick a small enough $\delta_w \in (0, \frac{1}{2})$ such that the




FIGURE 5. Here we show in red the disk $B(z, r_z)$ and its preimage under φ . The set V_z , whose boundaries are line segments inside the unit disk, is shown with blue color.

set

$$V_w := \{v \in \mathbb{D} : \langle w, v \rangle > (1 - \delta_w)\}$$

satisfies

$$\text{dist}(\varphi(V_w), z) \leq \frac{1}{2}r_z. \quad (5.9)$$

where $\langle w, v \rangle$ means the inner product of w and v . See Figure 5.

Finally we define

$$\Omega_\epsilon := \varphi \left(\mathbb{D} \setminus \overline{\bigcup_{w \in \varphi^{-1}(T)} V_w} \right).$$

We claim that $\Omega \setminus \Omega_\epsilon \subset U_\epsilon$. If this were to hold, then the \mathcal{H}^1 -measure of the projection of $\Omega \setminus \Omega_\epsilon$ on each coordinate axis would be less than ϵ .

Let us verify the claim. Since φ is a homeomorphism in \mathbb{D} and continuous up to the boundary, we have

$$\Omega \setminus \Omega_\epsilon = \varphi \left(\overline{\bigcup_{w \in \varphi^{-1}(T)} V_w} \right) \cap \Omega = \overline{\bigcup_{w \in \varphi^{-1}(T)} \varphi(V_w) \cap \Omega}.$$

Fix $y \in \Omega \setminus \Omega_\epsilon$ and let $y_n \in \bigcup_{w \in \varphi^{-1}(T)} \varphi(V_w)$ satisfy $y_n \rightarrow y$. Let $\delta := \text{dist}(y, \partial\Omega)$. Then $\delta > 0$. For n large enough, we have $|y_n - y| \leq \delta/4$, and hence

$$\text{dist}(y_n, \partial\Omega) \geq \text{dist}(y, \partial\Omega) - |y_n - y| \geq \frac{3}{4}\delta.$$

Moreover by (5.9) there exists $z_n \in T$ such that $|z_n - y_n| \leq r_{z_n}/2$, and then we obtain

$$\delta \leq \frac{2}{3}r_{z_n}.$$

Therefore

$$|y - z_n| \leq |y_n - y| + |y_n - z_n| \leq \frac{1}{4}\delta + \frac{1}{2}r_{z_n} \leq \frac{2}{3}r_{z_n},$$

and then $y \in B(z_n, r_{z_n}) \cap \Omega \subset U_\epsilon$. Thus the claim follows.

Since φ is a conformal map and continuous up to the boundary, then $\varphi|_{\partial(\varphi^{-1}(\Omega_\epsilon))}$ is continuous and injective as we took out neighborhoods of all the cut-points. Then by the compactness of $\partial(\varphi^{-1}(\Omega_\epsilon))$ we conclude that $\varphi|_{\partial(\varphi^{-1}(\Omega_\epsilon))}$ is an embedding to \mathbb{R}^2 and hence Ω_ϵ is Jordan. We claim that $\mathbb{R}^2 \setminus \overline{\Omega_\epsilon}$ is quasiconvex. If this claim were to hold, by the result on the $W^{1,1}$ -extension in the Jordan case (in Section 3) we could conclude that there exists an extension $u_\epsilon \in W^{1,1}(\mathbb{R}^2)$ of $u|_{\Omega_\epsilon}$.

Indeed, the proof of the claim is similar to the one of Lemma 2.13. If $x, y \in \Omega^c$, then (1.2) gives us the desired curve. If $x \in \Omega^c$ while $y \in \Omega \setminus \Omega_\epsilon$, then $|x - y| \geq \text{dist}(y, \partial\Omega)$. Hence by connecting y to $y_0 \in \partial\Omega$ such that the line segment $[\varphi^{-1}(y), \varphi^{-1}(y_0)]$ joining $\varphi^{-1}(y)$ and $\varphi^{-1}(y_0)$ lies in the line segment $[0, \varphi^{-1}(y_0)]$, (1.2) and part (1) and (3) of Lemma 2.8 give us the desired curve; note that $[\varphi^{-1}(y), \varphi^{-1}(y_0)]$ is contained in the union of V_z 's by definition. Thus by symmetry we may assume that $x, y \in \Omega \setminus \Omega_\epsilon$.

We claim that, there exists an absolute constant C such that, whenever

$$|x - y| \leq C \min\{\text{dist}(x, \partial\Omega), \text{dist}(y, \partial\Omega)\}, \quad (5.10)$$

we have

$$|\varphi^{-1}(x) - \varphi^{-1}(y)| \leq \frac{1}{16} \min\{\text{dist}(\varphi^{-1}(x), \partial\mathbb{D}), \text{dist}(\varphi^{-1}(y), \partial\mathbb{D})\}. \quad (5.11)$$

To show the existence of C , we apply Lemma 2.6 to the Whitney-type disk

$$B_1 = B(\varphi^{-1}(x), 1/16 \text{dist}(\varphi^{-1}(x), \partial\mathbb{D})),$$

which implies that $\varphi(B_1)$ is of Whitney-type. Thus there exists an absolute constant c such that

$$B(x, c \text{dist}(x, \partial\Omega)) \subset \varphi(B_1).$$

By choosing $C = \frac{1}{2}c$, we obtain the claim.

Suppose that x, y satisfy

$$|x - y| \geq C \min\{\text{dist}(x, \partial\Omega), \text{dist}(y, \partial\Omega)\}$$

where C is the one in (5.10). Then we can also define $x_0 \in \partial\Omega$ in a similar way, connect x_0 with x via a hyperbolic geodesic in Ω and join x_0 and y_0 in Ω^c by a curve with length controlled by $C(C_1)|x_0 - y_0|$. This gives us a desired curve by (1.2) and Lemma 2.8 again. Thus we only need to consider the case where

$$|x - y| \leq C \min\{\text{dist}(x, \partial\Omega), \text{dist}(y, \partial\Omega)\}.$$

Let $B = B(\varphi^{-1}(x), 4|\varphi^{-1}(x) - \varphi^{-1}(y)|)$. Then it is a disk contained in a 4-Whitney-type set because of (5.11). We claim that there exists a curve $\gamma \subset B \setminus \varphi^{-1}(\Omega_\epsilon)$ joining $\varphi^{-1}(x), \varphi^{-1}(y)$ in \mathbb{D} such that its length is controlled by $|\varphi^{-1}(x) - \varphi^{-1}(y)|$ up to a multiplicative constant. If this holds, by Lemma 2.6 the curve $\varphi(\gamma)$ joining x, y in $\varphi(B) \cap (\Omega \setminus \Omega_\epsilon)$ has length smaller than $|x - y|$ up to a multiplicative constant. Consequently $\mathbb{R}^2 \setminus \Omega_\epsilon$ is quasiconvex with the constant independent of ϵ .

Now let us show the existence of γ under the assumption (5.11). Observe first that by definition

$$\varphi^{-1}(\Omega_\epsilon) = \mathbb{D} \setminus \overline{\bigcup_{z \in T} V_z} = \left(\bigcap_{z \in T} (\mathbb{D} \setminus V_z) \right)^o$$

is a convex domain since it is the interior of the intersection of convex sets and the connectedness follows directly from the definition. Since we always require that $\delta_w \in (0, \frac{1}{2})$, then $B(0, \frac{1}{2}) \subset \varphi^{-1}(\Omega_\epsilon)$. This combined with convexity yields that for every $z \in \varphi^{-1}(\overline{\Omega}_\epsilon)$, locally there is a Euclidean (open) sector contained in $\varphi^{-1}(\Omega_\epsilon)$ with angle at least $\frac{\pi}{3}$, centered at z , towards the origin. Thus if we cannot join $\varphi^{-1}(x)$, $\varphi^{-1}(y)$ in $\frac{3}{4}B \setminus \varphi^{-1}(\Omega_\epsilon)$, then $\frac{3}{4}B \setminus \varphi^{-1}(\Omega_\epsilon)$ has at least two components. However this is impossible since $\varphi^{-1}(x)$, $\varphi^{-1}(y) \in \frac{1}{4}B$ but both $\varphi^{-1}(x)$ and $\varphi^{-1}(y)$ are not in the convex set $\varphi^{-1}(\Omega_\epsilon)$; planar geometry tells that such a sector with angle at least $\frac{\pi}{3}$ cannot exist. Therefore we can join $\varphi^{-1}(x)$, $\varphi^{-1}(y)$ in $\frac{3}{4}B \setminus \varphi^{-1}(\Omega_\epsilon)$ by a curve γ' with $\ell(\gamma') \leq (8 + \frac{3}{2}\pi)|\varphi^{-1}(x) - \varphi^{-1}(y)|$. Thus by letting $\gamma = \varphi(\gamma')$, we conclude from Lemma 2.6 and a change of variable that $\ell(\gamma) \lesssim |x - y|$ with an absolute constant, as desired.

Notice that Eu is absolutely continuous along almost every closed line segment parallel to the coordinate axes and contained in $\tilde{\Omega}$ since Eu is locally Lipschitz in $\tilde{\Omega}$ by our construction. Recall that by Lemma 5.4 we may assume that $u \in C(\overline{\Omega} \setminus T) \cap W^{1,\infty}(\Omega)$. Also take a representative of $u_{1/n} \in W^{1,1}(\mathbb{R}^2)$, the extension of the restriction of u to the Jordan domain $\Omega_{1/n}$ given by Theorem 3.1, so that it is absolutely continuous (and hence continuous) along almost every line segment parallel to the coordinate axes.

By Fubini's theorem,

$$\int_L |\nabla Eu| dx < \infty, \quad \int_L |\nabla u_{1/n}| dx < \infty \quad (5.12)$$

for \mathcal{H}^1 -almost every line segment L parallel to the coordinate axes. It follows that for almost every horizontal (or vertical) line S , (5.12) holds, each $u_{1/n}$ is absolutely continuous on every closed line segment of S , and so is also Eu on each of these closed line segments of S that is additionally contained in $\tilde{\Omega}$.

Define

$$Z_0 = \cap_{n \geq 1} (\Omega \setminus \Omega_{1/n}).$$

Since the vertical and horizontal projections of $\Omega \setminus \Omega_{1/n}$ have \mathcal{H}^1 -measure no more than $1/n$, we conclude that almost every line S parallel to the coordinate axes is disjoint from Z_0 . Equivalently, $S \cap \Omega \subset \Omega_{1/n}$ for some $n \in \mathbb{N}$. Furthermore, recalling that $H = T \cup Z$ has \mathcal{H}^1 -measure zero, we have that almost every S parallel to the coordinate axes is disjoint from H . By Lemma 5.4, we may assume that u is continuous on $S \cap \overline{\Omega}$ for each such S .

Fix a horizontal line segment S that satisfies the conclusions of the preceding two paragraphs. Let us verify that $Eu|_I$ is absolutely continuous, where $I \subset S$ is a closed line segment. By the definition of S , there exists $M \in \mathbb{N}$ such that $Eu|_{\overline{\Omega} \cap S} = u_{1/M}|_{\overline{\Omega} \cap S}$, and $Eu_{1/M}$ is continuous on S ; notice that by Lemma 5.4 and our assumption, $Eu(x) = u_{1/M}(x)$ for $x \in \partial\Omega \cap S$. Let $z_1, \dots, z_{2n} \in I$ be points whose first components are strictly increasing.

If $z_1, z_2 \in \overline{\Omega}$, then by the assumption that $u \in C(\overline{\Omega} \setminus T) \cap W^{1,\infty}(\Omega)$, we get

$$|Eu(z_1) - Eu(z_2)| = |u_{1/M}(z_1) - u_{1/M}(z_2)| \leq \int_{[z_1, z_2]} |\nabla u_{1/M}| dx.$$

If $[z_1, z_2] \subset \tilde{\Omega}$, then since Eu is absolutely continuous on closed line segments in S , we obtain

$$|Eu(z_1) - Eu(z_2)| \leq \int_{[z_1, z_2]} |\nabla Eu| dx.$$

For the remaining case, by symmetry we may assume that $z_1 \in \overline{\Omega}$ while $z_2 \in \tilde{\Omega}$. Then let z be the nearest point to z_2 on $\overline{\Omega} \cap I$ between z_1 and z_2 , and observe by the same absolute continuity properties as above that

$$\begin{aligned} |Eu(z_1) - Eu(z_2)| &\leq |Eu(z_1) - Eu(z)| + |Eu(z) - Eu(z_2)| \\ &= |u_{1/M}(z_1) - u_{1/M}(z)| + |Eu(z) - Eu(z_2)| \\ &\leq \int_{[z_1, z]} |\nabla u_{1/M}| dx + \int_{[z, z_2]} |\nabla Eu| dx, \end{aligned}$$

where in the last inequality we applied the continuity of Eu on S .

Using a similar argument for the other pairs, we get

$$\begin{aligned} \sum_{i=1}^n |Eu(z_{2i-1}) - Eu(z_{2i})| &\leq \sum_{i=1}^n 2 \int_{[z_{2i-1}, z_{2i}]} (|\nabla u_{1/M}| + |\nabla Eu|) dx \\ &= 2 \int_{\cup_i [z_{2i-1}, z_{2i}]} (|\nabla u_{1/M}| + |\nabla Eu|) dx \end{aligned}$$

from which we get the desired absolute continuity of Eu on I by the absolute continuity of the integral. The case of vertical lines is dealt with analogously, and we conclude the absolute continuity of Eu along almost every vertical and horizontal line.

5.5. Conclusion of the proof. Combining the discussions in all the subsections above, we have shown that there exists a linear extension operator

$$E: W^{1,1}(\Omega) \cap C(\Omega) \cap W^{1,\infty}(\Omega) \rightarrow W^{1,1}(B)$$

such that

$$\|\nabla Eu\|_{L^1(B)} \lesssim \|\nabla u\|_{L^1(\Omega)}$$

and

$$\|Eu\|_{L^1(B)} \lesssim \|u\|_{L^1(\Omega)},$$

where the constants depend only on the constant in (1.2); recall that the boundary of Ω as Lebesgue measure zero by part (4) of Lemma 2.8. Here B is a disk of radius $4 \operatorname{diam}(\Omega)$ such that Ω is contained in $\frac{1}{2}B$. By the fact that B is an $W^{1,1}$ -extension domain, and the fact that $C(\Omega) \cap W^{1,\infty}(\Omega)$ is dense in $W^{1,1}(\Omega)$ [22], we conclude the sufficiency of the curve condition (1.2).

6. NECESSITY FOR SIMPLY CONNECTED DOMAINS

In this section we prove the necessity of the curve condition (1.2) for bounded simply connected $W^{1,1}$ -extension domains. We use the following result stated in [20, Corollary 1.2], which was shown via results in [4].

Theorem 6.1 ([20, Corollary 1.2]). *Let $\Omega \subset \mathbb{R}^2$ be a bounded simply connected $W^{1,1}$ -extension domain. Then Ω^c is C_1 -quasiconvex with C_1 depending only on the norm of the extension operator (with respect to the homogeneous norm).*

Let $\varphi: \mathbb{D} \rightarrow \Omega$ be a conformal map. By Theorem 6.1 the complement of Ω is quasiconvex for some constant C_1 . Therefore Ω is $J(C_1)$ -John by part (1) of Lemma 2.8 and also resulting from part (2) of Lemma 2.8 the map φ extends as a continuous map (still denoted φ) to the boundary $\partial\mathbb{D}$. As before, let us denote $\tilde{\Omega} := \mathbb{R}^2 \setminus \overline{\Omega}$ and let $\{\tilde{\Omega}_i\}_{i=1}^N$ be an enumeration of the connected components of $\tilde{\Omega}$ with $N \in \mathbb{N} \cup \{+\infty\}$.

We show that (1.2) is necessary. Fix $x, y \in \Omega^c$. If

$$|x - y| \leq \frac{1}{2} \max\{\text{dist}(x, \partial\Omega), \text{dist}(y, \partial\Omega)\},$$

then we may take γ to be the line segment joining x and y , and then (1.2) holds for this γ . Thus we may assume that

$$|x - y| \geq \frac{1}{2} \max\{\text{dist}(x, \partial\Omega), \text{dist}(y, \partial\Omega)\}.$$

By Theorem 6.1 there exists a curve $\gamma' \subset \Omega^c$ joining x, y with length controlled by $C_1|x - y|$. If γ' does not touch the boundary, then we are done. Otherwise starting from $z_0 = x$, we choose z_1 such that

$$\ell(\gamma'[z_0, z_1]) = \delta \text{diam}(\Omega)$$

for the subcurve $\gamma'[z_0, z_1]$ of γ' joining z_0 to z_1 , with δ in Lemma 4.2; if there is no such a point, we let $z_1 = y$ and define γ_1 to be the piecewise hyperbolic geodesic joining z_0 and z_1 . If $\gamma'[z_0, z_1]$ touches the boundary at most once, then we define $\gamma_1 = \gamma'[z_0, z_1]$. If not, choose the first and last points x_1, y_1 of $\gamma'[z_0, z_1] \cap \partial\Omega$ according to the parametrization of γ' , and reroute $\gamma'[z_0, z_1]$ via the piecewise hyperbolic geodesic joining x_1 and y_1 to obtain a new curve γ_1 ; the existence of the piecewise hyperbolic geodesic follows from Lemma 4.2. Also according to Lemma 4.2,

$$\ell(\gamma_1) \lesssim \ell(\gamma'[z_0, z_1])$$

with the constant depending only on C_1 . Continue this procedure via replacing z_0 by z_1 to find z_2 and to define γ_2 , and iterate until we finally get the point y . Observe that this process stops after finitely many steps since γ' has finite length, and it produces a finite sequence of curves γ_i , $1 \leq i \leq N$ such that

$$\sum_{i=1}^N \ell(\gamma_i) \lesssim \sum_{i=1}^N \ell(\gamma'[z_{i-1}, z_i]) \lesssim \ell(\gamma') \lesssim |x - y|.$$

We claim that $\gamma = \cup_{i=1}^N \gamma_i$ satisfies (1.2) (with the constant depending only on C_1). By our construction and the injectivity of piecewise hyperbolic geodesics from Lemma 4.2, it suffices to show that for $x, y \in \partial\Omega$ with $|x - y| \leq \delta \text{diam}(\Omega)$ we have

$$\mathcal{H}^1(I) = 0$$

where I is the intersection between $\partial\Omega$ and the piecewise hyperbolic geodesic Γ joining x and y .

Let Γ be a piecewise hyperbolic geodesic joining $x, y \in \partial\Omega$ in Ω^c with

$$|x - y| \leq \delta \operatorname{diam}(\Omega).$$

By Lemma 4.9 any $z \in I \setminus \{x, y\}$ is a cut-point, and every neighborhood of z intersects both Ω_0 and Ω_1 , where Ω_1 and Ω_2 are the two components of $\Omega \setminus (\varphi([v_1, 0] \cup [0, v_2]) \cup \Gamma)$ with $x = \varphi(v_1)$ and $y = \varphi(v_2)$. Write $\alpha = [v_1, 0] \cup [0, v_2]$.

Suppose that $\mathcal{H}^1(I) > 0$. Let us next construct a test function that will give us a contradiction. Pick $z_0 \in I \setminus \{x, y\}$ with

$$\mathcal{H}^1(I \cap B(z_0, r)) > 0 \quad \text{for all } r > 0;$$

this is possible by the assumption that $\mathcal{H}^1(I) > 0$; see [8, Theorem 2, Page 72]. Since $\varphi(\alpha)$ is a compact set,

$$r_0 := \frac{1}{8} \operatorname{dist}(z, \varphi(\alpha)) > 0.$$

We define ψ by setting

$$\psi(z) := \max \left\{ 0, 1 - \frac{1}{4r_0} \operatorname{dist}(z, B(z_0, r_0)) \right\} \chi_{\Omega_1}(z).$$

By our definition of r_0 , we have $\psi \in W^{1,1}(\Omega)$; notice that $\partial\Omega_1 \cap \Omega \subset \varphi(\alpha)$, where χ_{Ω_1} is the characteristic function of Ω_1 .

Suppose that there exists an extension $E\psi \in W^{1,1}(\mathbb{R}^2)$ of ψ . By Lemma 4.7 and the John property of Ω , for every $z \in K := I \cap B(z_0, r_0)$ and every $0 < r_z < \delta < r_0$ with any $\delta > 0$, for each $i = 0, 1$ there is a point $z^{(i)} \in B(z, r_z)$ such that

$$B(z^{(i)}, cr_z) \subset B(z, r_z) \subset B(z_0, 2r_0) \cap \Omega_i \subset \Omega_i,$$

where the constant c depends only on the John constant J .

Therefore, by the Poincaré inequality,

$$1 \lesssim |E\psi_{B(z^{(0)}, cr_z)} - E\psi_{B(z^{(1)}, cr_z)}| \lesssim r_z \int_{B(z, r_z)} |\nabla E\psi(w)| dw \quad (6.1)$$

for every $z \in K$, where $E\psi_B$ denotes the integral average of $E\psi$ on B .

Notice that $K \subset \bigcup_{z \in K} B(z, r_z)$. By the $5r$ -covering theorem, there exists a collection of countably many pairwise disjoint disks

$$\{B(z_i, r_i)\} \subset \{B(z, r) : z \in K, 0 < r < \delta\}$$

such that

$$K \subset \bigcup_i B(z_i, 5r_i).$$

Moreover since $\mathcal{H}^2(K) = 0$, for any $\epsilon > 0$, there exists $\delta_0 > 0$ such that

$$\sum_i r_i^2 < \epsilon \quad (6.2)$$

whenever $0 < \delta < \delta_0$.

Now by (6.1) we have

$$\int_{\bigcup_i B(z_i, r_i)} |\nabla E\psi(w)| dw \gtrsim \sum_i r_i \gtrsim \mathcal{H}_\delta^1(K) \gtrsim \mathcal{H}^1(K).$$

for $\delta > 0$ small enough. However, since $|\nabla E\psi| \in L^1(\mathbb{R}^2)$, the absolute continuity of the integral and (6.2), give that

$$\int_{\bigcup_i B(z_i, r_i)} |\nabla E\psi(w)| dw$$

can be made arbitrarily small by choosing a sufficiently small ϵ . Thus we have obtained a contradiction and hence the necessity of (6.1) for $W^{1,1}$ -extension domains follows.

APPENDIX A. CAPACITY ESTIMATES

In this appendix we provide the proofs of Theorem 2.3 and Lemma 2.14. The proofs given here use capacity estimates.

A.1. Conformal capacity. Let Ω be a domain. For a given pair of continua $E, F \subset \overline{\Omega} \subset \mathbb{R}^2$, define the *conformal capacity between E and F in Ω* as

$$\text{Cap}(E, F, \Omega) = \inf\{\|\nabla u\|_{L^2(\Omega)}^2 : u \in \Delta(E, F, \Omega)\},$$

where $\Delta(E, F, \Omega)$ denotes the class of all $u \in W_{\text{loc}}^{1,2}(\Omega)$ that are continuous in $\Omega \cup E \cup F$ and satisfy $u = 1$ on E , and $u = 0$ on F . By definition, we see that the conformal capacity is increasing with respect to Ω .

In what follows, we introduce the properties of conformal capacity which are used in this paper; see e.g. [30, Chapter 1] for additional properties. We remark that, even though [30] (with some other references below) states the results for “modulus”, “modulus” is equivalent with conformal capacity in our setting below (see e.g. [26, Proposition 10.2, Page 54]).

Lemma A.1. *The conformal capacity is conformally invariant, that is, for domains Ω and Ω' in \mathbb{R}^2 , a conformal (onto) map $\varphi: \Omega \rightarrow \Omega'$ and continua E and F in Ω , we have*

$$\text{Cap}(\varphi(E), \varphi(F), \Omega') = \text{Cap}(E, F, \Omega). \quad (\text{A.1})$$

Moreover, if φ has a homeomorphic extension, still denoted by φ , $\varphi: \overline{\Omega} \rightarrow \overline{\Omega}'$, then (A.1) also holds for continua in $\overline{\Omega}$. Especially this is the case if both Ω and Ω' are Jordan.

Proof. Let $0 < \epsilon < \frac{1}{3}$ and fix a function $u \in W^{1,2}(\Omega)$ that is continuous in $\Omega \cup E \cup F$ and satisfies $u = 1$ on E and $u = 0$ on F . Let now u_ϵ be the function

$$u'_\epsilon = \frac{u - \epsilon}{1 - 2\epsilon}$$

in Ω . Then the sets $U_\epsilon = \{u_\epsilon > 1\}$ and $V_\epsilon = \{u_\epsilon < 0\}$ are relatively open in $\Omega \cup E \cup F$, and contain E and F , respectively. Define

$$u_\epsilon = \max\{\min\{u'_\epsilon, 1\}, 0\}.$$

Observe that $u_\epsilon \rightarrow u$ in $W^{1,2}(\Omega)$ as $\epsilon \rightarrow 0$.

For each $\epsilon > 0$, $u_\epsilon \circ \varphi^{-1}$ is an admissible function in $\Delta(\varphi(E), \varphi(F), \Omega')$. Thus, by the chain rule, conformality of φ^{-1} and a change of variables, we have the estimate

$$\begin{aligned} \text{Cap}(\varphi(E), \varphi(F), \Omega') &\leq \int_{\Omega'} |\nabla u(\varphi^{-1}(x))|^2 |D\varphi^{-1}(x)|^2 dx \\ &\leq \int_{\Omega} |\nabla u(x)|^2 |D\varphi^{-1}(\varphi(x))|^2 J_\varphi(x) dx \leq \int_{\Omega} |\nabla u|^2 dx. \end{aligned}$$

The claim now follows by taking an infimum over functions u above and symmetry. \square

In what follows, whenever we mention the conformal invariance of conformal capacity, we always refer to this lemma above.

In the unit disk \mathbb{D} , and the exterior unit disk $\mathbb{R}^2 \setminus \overline{\mathbb{D}}$, we have the following *Loewner estimate* for the conformal capacity. Let E and F be continua in $\overline{\mathbb{D}}$. Then

$$\text{Cap}(E, F, \mathbb{D}) \geq c \log \left(1 + \frac{\min\{\text{diam}(E), \text{diam}(F)\}}{\text{dist}(E, F)} \right) \quad (\text{A.2})$$

where $c > 0$ is a universal constant, and the analogous inequality holds for $E, F \subset \mathbb{R}^2 \setminus \mathbb{D}$; see e.g. [31, Lemma 7.38] together with [11, Remark 2.12].

We call a domain $A \subset \mathbb{R}^2$ a *ring domain* if its complement has exactly two components. If the components of A are U_0 and U_1 , then we denote $A = R(U_0, U_1)$. It follows from topology that ∂A also has two components, $V_0 = U_0 \cap \overline{A}$ and $V_1 = U_1 \cap \overline{A}$. If U_0, V_0 and V_1 are compact, we have

$$\text{Cap}(V_0, V_1, A) = \text{Cap}(U_0, V_1, A \cup U_0); \quad (\text{A.3})$$

indeed, “ \leq ” directly follows from the definition and “ \geq ” follows by extending each $u \in \Delta(V_0, V_1, A)$ as constant 1 to $U_0 \setminus V_0$, see also [30, Theorem 11.3] (and its proof). Furthermore, we have the following estimate for the capacity of boundary components in a ring domain.

Lemma A.2. *Let $A = R(U_0, U_1) \subset \mathbb{R}^2$ be a ring domain with U_1 unbounded. Assume that $V_0 = U_0 \cap \overline{A}$ and $V_1 = U_1 \cap \overline{A}$ are compact. There exist two universal increasing functions $\phi_i: (0, \infty) \rightarrow (0, \infty)$, $i = 1, 2$, so that $\lim_{t \rightarrow 0+} \phi_i(t) = 0$ and $\lim_{t \rightarrow \infty} \phi_i(t) = \infty$, and so that*

$$\phi_1 \left(\frac{\text{diam}(U_0)}{\text{dist}(U_0, U_1)} \right) \leq \text{Cap}(V_0, V_1, A) \leq \phi_2 \left(\frac{\text{diam}(U_0)}{\text{dist}(U_0, U_1)} \right). \quad (\text{A.4})$$

Proof. Define

$$\sigma = \frac{\text{diam}(U_0)}{\text{dist}(U_0, U_1)}.$$

The lower bound of (A.4) follows from [30, Theorem 11.7, Theorem 11.9]. For the upper bound, if $0 < \sigma < \frac{1}{2}$ we have $U_0 \subset B$, $2B \subset A \cup U_0$ for a suitable disk B , and then by the monotonicity of capacity and [30, Example 7.5] one obtains the upper bound $C(\log(1/\sigma))^{-1}$ for some absolute constant $C > 0$. When $\frac{1}{2} \leq \sigma < \infty$ one just applies the test function

$$u(x) = \min \left\{ 1, \max \left\{ 0, 1 - \frac{\text{dist}(x, U_0)}{\text{dist}(U_0, U_1)} \right\} \right\}.$$

Then u is Lipschitz, and

$$|\nabla u| \leq \frac{1}{\text{dist}(U_0, U_1)}.$$

Hence by letting $x_0 \in U_0$

$$\begin{aligned} \|\nabla u\|_{L^2(\Omega)}^2 &\leq \text{dist}(U_0, U_1)^{-2} |B(x_0, \text{diam}(U_0) + \text{dist}(U_0, U_1))| \\ &\lesssim \left(\frac{\text{diam}(U_0) + \text{dist}(U_0, U_1)}{\text{dist}(U_0, U_1)} \right)^2 \sim (1 + \sigma)^2. \end{aligned}$$

□

We record the following estimate, which states a kind of converse to (A.2); see e.g. [22, Lemma 2.2] for a weaker version of the claim.

Lemma A.3. *Let Ω be a domain and $E, F \subset \Omega$ be a pair of disjoint continua. Then if $\text{Cap}(E, F, \Omega) \geq c_0$, we have*

$$\min\{\text{diam}_\Omega(E), \text{diam}_\Omega(F)\} \gtrsim \text{dist}_\Omega(E, F), \quad (\text{A.5})$$

where the constant only depends on c_0 . Especially

$$\min\{\text{diam}_\Omega(E), \text{diam}_\Omega(F)\} \gtrsim \text{dist}(E, F),$$

and if $\Omega = \mathbb{R}^2$

$$\min\{\text{diam}(E), \text{diam}(F)\} \gtrsim \text{dist}(E, F). \quad (\text{A.6})$$

If we further assume that Ω is Jordan, then (A.5) also holds if $E \subset \partial\Omega$ and $F \subset \Omega$ (or $E \subset \partial\Omega$ and $F \subset \partial\Omega$) are continua with $\text{Cap}(E, F, \Omega) \geq c_0$.

Proof. Step 1: We begin with the case where $E, F \subset \Omega$. We may clearly assume that $\text{diam}_\Omega(E) \leq \text{diam}_\Omega(F)$ and also that $2 \text{diam}_\Omega(E) \leq \text{dist}_\Omega(E, F)$; otherwise the claim holds trivially. Choose $z \in E$, and write $\frac{\text{dist}_\Omega(E, F)}{\text{diam}_\Omega(E)} = \sigma$. We define

$$u(x) = \begin{cases} 1, & \text{if } \text{dist}_\Omega(x, z) \leq \text{diam}_\Omega(E) \\ 0, & \text{if } \text{dist}_\Omega(x, z) \geq \text{dist}_\Omega(E, F) \\ \frac{\log(\text{dist}_\Omega(E, F)) - \log(\text{dist}_\Omega(x, z))}{\log(\text{dist}_\Omega(E, F)) - \log(\text{diam}_\Omega(E))}, & \text{otherwise} \end{cases}$$

Then u is locally Lipschitz and

$$|\nabla u(x)| \leq (\log \sigma)^{-1} \text{dist}_\Omega(x, z)^{-1}.$$

Write

$$R = B_\Omega(z, \text{dist}_\Omega(E, F)) \setminus B_\Omega(z, \text{diam}_\Omega(E)),$$

and for $i \geq 1$

$$A_i = B_\Omega(z, 2^i \text{diam}(E)) \setminus B_\Omega(z, 2^{i-1} \text{diam}(E)),$$

where $B_\Omega(z, r)$ is the disk centered at z with radius r with respect to the inner distance. A direct calculation via a dyadic annular decomposition with respect to the inner distance gives

$$\begin{aligned}
c_0 &\leq \int_\Omega |\nabla u|^2 dx \lesssim (\log \sigma)^{-2} \int_R \text{dist}_\Omega(x, z)^{-2} dx \\
&\lesssim (\log \sigma)^{-2} \sum_{i=1}^{\infty} \int_{R \cap A_i} 4^{-i} \text{diam}(E)^{-2} dx \\
&\lesssim (\log \sigma)^{-2} \sum_{i=1}^{[\log \sigma] + 1} 1 \\
&\lesssim (\log \sigma)^{-2} \log \sigma \lesssim (\log \sigma)^{-1},
\end{aligned}$$

where $[\log \sigma]$ denotes the integer part of $[\log \sigma]$, and in the fourth inequality we used the fact that $B_\Omega(z, r) \subset B(z, r)$. Hence $\sigma \leq C(c_0)$, which means that $\text{dist}_\Omega(E, F) \lesssim \text{diam}_\Omega(E)$.

Step 2: We continue with the case $E, F \subset \partial\Omega$. To begin with, let us first show that

$$\text{dist}_\Omega(E, F) < \infty.$$

Let $\varphi: \overline{\mathbb{D}} \rightarrow \overline{\Omega}$ be a homeomorphism given by the Caratheodory-Osgood theorem. Since

$$\text{Cap}(E, F, \mathbb{R}^2) \geq \text{Cap}(E, F, \Omega) \geq c_0,$$

we conclude by (A.6) that neither E nor F is a singleton. Then $\varphi^{-1}(E)$ is an closed arc contained in the unit circle and of positive 1-Hausdorff measure. Since $\varphi \in W^{1,2}(\mathbb{D})$, then

$$\text{dist}_\Omega(E, \varphi(0)) < \infty;$$

otherwise by noticing that φ is a smooth map on $\overline{B(0, 1/2)}$ and applying Fubini's theorem in the annulus $\mathbb{D} \setminus \overline{B(0, 1/2)}$ with respect to the polar coordinates, we obtain a contradiction to the fact that $\varphi \in W^{1,2}(\mathbb{D})$. An analogous argument also shows that

$$\text{dist}_\Omega(F, \varphi(0)) < \infty,$$

and hence by the triangle inequality $\text{dist}_\Omega(E, F) < \infty$.

Let $\phi: \mathbb{H} \rightarrow \Omega$ be a conformal map (homeomorphically extended to \mathbb{R}), where \mathbb{H} denotes the upper half plane so that $\phi^{-1}(E)$ and $\phi^{-1}(F)$ are two continua in the real line. We define a sequence of continua $E_j \subset \phi^{-1}(E)$ (for large j) by setting

$$E_j = \{z \in \mathbb{R}: \text{dist}(z, \mathbb{R} \setminus \phi^{-1}(E)) \geq 2^{-j+1}\}.$$

The sets $F_j \subset \phi^{-1}(F)$ are defined similarly. Furthermore, define

$$\hat{E}_j = E_j + i2^{-j}, \quad \hat{F}_j = F_j + i2^{-j}.$$

Note that for every two points $z_1, z_2 \in E$, the hyperbolic geodesic Γ' joining them satisfies (see Lemma 2.2)

$$\ell(\Gamma') \lesssim \text{dist}_\Omega(z_1, z_2),$$

where the constant is absolute. Given $w_1, w_2 \in \phi(\hat{E}_j)$, denote the hyperbolic geodesic connecting them by Γ . Extend Γ to a full hyperbolic geodesic Γ_1 joining points $z_1, z_2 \in \partial\Omega$.

Since $w_1, w_2 \in \phi(\hat{E}_j)$, from the definition of \hat{E}_j and planar geometry we conclude via ϕ that $z_1, z_2 \in E$. Therefore

$$\text{dist}_\Omega(w_1, w_2) \leq \ell(\Gamma) \leq \ell(\Gamma_1) \lesssim \text{dist}_\Omega(z_1, z_2) \leq \text{diam}_\Omega(E).$$

By the arbitrariness of w_1, w_2 , we conclude that

$$\text{diam}_\Omega(\phi(\hat{E}_j)) \lesssim \text{diam}_\Omega(E) \quad (\text{A.7})$$

with an absolute constant. Similarly we get

$$\text{diam}_\Omega(\phi(\hat{F}_j)) \lesssim \text{diam}_\Omega(F). \quad (\text{A.8})$$

By the monotonicity of capacity, we have

$$\text{Cap}(\hat{E}_j, \hat{F}_j, \mathbb{H}) \geq \text{Cap}(\hat{E}_j, \hat{F}_j, \mathbb{H} + i2^{-j}).$$

Via the translation map $z \mapsto z - i2^{-j}$ and the conformal invariance of capacity, we further have

$$\text{Cap}(\hat{E}_j, \hat{F}_j, \mathbb{H} + i2^{-j}) = \text{Cap}(E_j, F_j, \mathbb{H}).$$

If we can show that

$$\text{Cap}(E_j, F_j, \mathbb{H}) \geq \frac{1}{16} \text{Cap}(\phi^{-1}(E), \phi^{-1}(F), \mathbb{H}) \quad (\text{A.9})$$

for j large enough, then by the conformal invariance of capacity together with the chain of inequalities above

$$\text{Cap}(\phi(\hat{E}_j), \phi(\hat{F}_j), \Omega) \geq \frac{1}{16} \text{Cap}(\phi^{-1}(E), \phi^{-1}(F), \mathbb{H}) = \frac{1}{16} \text{Cap}(E, F, \Omega) \geq \frac{c_0}{16}.$$

Thus Lemma A.3 implies that

$$\min\{\text{diam}_\Omega(\phi(\hat{E}_j)), \text{diam}_\Omega(\phi(\hat{F}_j))\} \gtrsim \text{dist}_\Omega(\phi(\hat{E}_j), \phi(\hat{F}_j)),$$

where the constant only depends on c_0 . This together with (A.7) and (A.8) gives that there is a rectifiable curve γ_j connecting $\phi(\hat{E}_j)$ and $\phi(\hat{F}_j)$ such that

$$\ell(\gamma_j) \lesssim \min\{\text{diam}_\Omega(\phi(\hat{E}_j)), \text{diam}_\Omega(\phi(\hat{F}_j))\} \lesssim \min\{\text{diam}_\Omega(E), \text{diam}_\Omega(F)\},$$

with the constant independent of j . We may assume that these curves are hyperbolic geodesics by Lemma 2.2. Then by parameterizing γ_j with arc length and applying the Arzelà-Ascoli lemma, we get a curve $\gamma \subset \overline{\Omega}$ joining E, F with the length bounded by $\min\{\text{diam}_\Omega(E), \text{diam}_\Omega(F)\}$ up to a multiplicative constant. Observe that, since φ is a homeomorphism, the uniform convergence of $\gamma_j \rightarrow \gamma$ implies that of $\varphi^{-1}(\gamma_j) \rightarrow \varphi^{-1}(\gamma)$. Then the definition of hyperbolic geodesics in the unit disk gives that $\varphi^{-1}(\gamma)$ is also a hyperbolic geodesic, and so is γ . The desired estimate (A.5) then follows. Thus it suffices to show (A.9).

Set

$$a = \min\{(\text{diam}(\phi^{-1}(E)), \text{diam}(\phi^{-1}(F)), \text{dist}(\phi^{-1}(E), \phi^{-1}(F)))\}.$$

Let the middle point of $\phi^{-1}(E)$ be z_1 and that of $\phi^{-1}(F)$ be z_2 . Write

$$b_E = \frac{1}{2} \text{diam}(\phi^{-1}(E)) + \frac{a}{8}, \quad b_F = \frac{1}{2} \text{diam}(\phi^{-1}(F)) + \frac{a}{8}. \quad (\text{A.10})$$

Recall that

$$\text{diam}(\phi^{-1}(E)) - \text{diam}(E_j) = 2^{-j+2}, \quad \text{diam}(\phi^{-1}(F)) - \text{diam}(F_j) = 2^{-j+2}.$$

For $j \in \mathbb{N}$, $2^{-j+5} \leq a$, we define a map f by setting

$$f(x) = \begin{cases} z_1 + \frac{\text{diam}(\phi^{-1}(E))}{\text{diam}(E_j)}(x - z_1), & |x - z_1| \leq \frac{1}{2} \text{diam}(E_j) \\ z_1 + \left(\frac{1}{2} \text{diam}(\phi^{-1}(E)) + \frac{a(|x - z_1| - \frac{1}{2} \text{diam}(E_j))}{2^{-j+4} + a} \right) \frac{(x - z_1)}{|x - z_1|}, & \frac{1}{2} \text{diam}(E_j) \leq |x - z_1| \leq b_E \\ z_2 + \frac{\text{diam}(\phi^{-1}(F))}{\text{diam}(F_j)}(x - z_2), & |x - z_2| \leq \frac{1}{2} \text{diam}(F_j) \\ z_2 + \left(\frac{1}{2} \text{diam}(\phi^{-1}(F)) + \frac{a(|x - z_2| - \frac{1}{2} \text{diam}(F_j))}{2^{-j+4} + a} \right) \frac{(x - z_2)}{|x - z_2|}, & \frac{1}{2} \text{diam}(F_j) \leq |x - z_2| \leq b_F \\ x, & \text{otherwise.} \end{cases}$$

Then f is a well-defined homeomorphism; recall (A.10) and notice that

$$b_E - \frac{1}{2} \text{diam}(E_j) = 2^{-j+1} + \frac{a}{8}, \quad b_F - \frac{1}{2} \text{diam}(F_j) = 2^{-j+1} + \frac{a}{8}.$$

By our choice of j , f maps \mathbb{H} to \mathbb{H} , E_j to $\phi^{-1}(E)$ and F_j to $\phi^{-1}(F)$. Moreover since $2^{-j+5} \leq a$,

$$\frac{1}{2} \leq \frac{a}{2^{-j+4} + a} \leq 1, \quad 1 \leq \frac{\text{diam}(\phi^{-1}(E))}{\text{diam}(E_j)} \leq 2, \quad 1 \leq \frac{\text{diam}(\phi^{-1}(F))}{\text{diam}(F_j)} \leq 2.$$

Thus f is 2-biLipschitz. For any $u \in \Delta(E_j, F_j, \mathbb{H})$ we have $u \circ f^{-1} \in \Delta(\phi^{-1}(E), \phi^{-1}(F), \mathbb{H})$ since f is a homeomorphism. Thus by the chain rule and a change of variables we get

$$\begin{aligned} \text{Cap}(\phi^{-1}(E), \phi^{-1}(F), \mathbb{H}) &\leq \int_{\mathbb{H}} |\nabla u(f^{-1}(x))|^2 |Df^{-1}(x)|^2 dx \\ &\leq 4 \int_{\mathbb{H}} |\nabla u(x)|^2 J_f(x) dx \leq 16 \int_{\mathbb{H}} |\nabla u|^2 dx, \end{aligned}$$

which implies (A.9) as desired, and then (A.5) follows.

Step 3: We are left with the case where $E \subset \partial\Omega$ and $F \subset \Omega$. In this case we only perform the above approximation for E (and consider j large enough so that $E_j \cap \phi^{-1}(F) = \emptyset$). The argument in the previous case applies with this modification and gives the remaining claim. \square

A.2. A variant of the Gehring-Hayman theorem. We employ following lemma on conformal annuli to prove Theorem 2.3.

Lemma A.4. *Let $\Omega \subset \mathbb{R}^2$ be a Jordan domain, and let a homeomorphism $\tilde{\varphi}: \mathbb{R}^2 \setminus \Omega \rightarrow \mathbb{R}^2 \setminus \mathbb{D}$ be conformal in $\mathbb{R}^2 \setminus \overline{\Omega}$. For $z_1 \in \partial\Omega$, define*

$$A(z_1, k) := \{x \in \mathbb{R}^2 \setminus \mathbb{D} \mid 2^{k-1} < |x - \tilde{\varphi}(z_1)| \leq 2^k\},$$

for $k \in \mathbb{Z}$. Furthermore, let $\Gamma \subset \mathbb{R}^2 \setminus \overline{\Omega}$ be a hyperbolic geodesic joining z_1 and $z_2 \in \partial\Omega$ such that $\tilde{\varphi}(\Gamma) \subset B(0, 100)$. Also let $\gamma \subset \mathbb{R}^2 \setminus \Omega$ be a curve connecting z_1 and z_2 . Set

$$\Gamma_k := \tilde{\varphi}^{-1}(A(z_1, k)) \cap \Gamma$$

and let γ_k be any subcurve of γ in $\tilde{\varphi}^{-1}(A(z_1, k))$ joining the inner and outer boundaries of $\tilde{\varphi}^{-1}(A(z_1, k))$, if such a subcurve exists. (Here the inner and outer boundaries of $\tilde{\varphi}^{-1}(A(z_1, k))$ are the preimages under $\tilde{\varphi}$ of the inner and outer boundaries of $A(z_1, k)$.) Then

$$\ell(\Gamma_k) \sim \text{dist}(\Gamma_k, \partial\Omega) \quad (\text{A.11})$$

and

$$\ell(\gamma_k) \gtrsim \ell(\Gamma_k) \sim \text{diam}(\Gamma_k).$$

Here all the constants are independent of Ω and the choice of $\tilde{\varphi}, z_1, \gamma, z_2, k$.

Proof. We may assume that Γ_k is non-empty; otherwise the claim of the lemma is trivial.

The fact that $\ell(\Gamma_k) \sim \text{dist}(\Gamma_k, \partial\Omega) \sim \text{diam}(\Gamma_k)$ follows immediately from Lemma 2.6 and a change of variable, since by definition $\tilde{\varphi}(\Gamma_k)$ is contained in a Whitney-type set in $\mathbb{R}^2 \setminus \overline{\mathbb{D}}$.

Hence we only need to prove that $\ell(\gamma_k) \gtrsim \ell(\Gamma_k)$. Observe that, since γ_k by definition joins the inner and outer boundaries of $\tilde{\varphi}^{-1}(A(z_1, k))$, then $\tilde{\varphi}(\gamma_k)$ joins the inner and outer boundaries of $A(z_1, k)$, and hence

$$\ell(\tilde{\varphi}(\gamma_k)) \gtrsim \text{diam}(\tilde{\varphi}(\Gamma_k)) \sim \text{dist}(\tilde{\varphi}(\Gamma_k), \partial\mathbb{D}). \quad (\text{A.12})$$

We next argue by case study.

Case 1: $\text{dist}(\tilde{\varphi}(\gamma_k), \tilde{\varphi}(\Gamma_k)) \geq \frac{1}{3} \text{dist}(\tilde{\varphi}(\Gamma_k), \partial\mathbb{D})$. By Lemma 2.6, the assumption and the fact that $\tilde{\varphi}(\Gamma_k)$ is contained in a Whitney-type set, we know that for any curve γ' joining γ_k and Γ_k , its length satisfies

$$\ell(\gamma') \gtrsim \text{diam}(\Gamma_k),$$

and hence

$$\text{dist}_\Omega(\gamma_k, \Gamma_k) \gtrsim \text{diam}(\Gamma_k). \quad (\text{A.13})$$

Moreover by (A.2) for the exterior of the unit disk, (A.12) and the monotonicity of the capacity we obtain

$$1 \lesssim \text{Cap}(\tilde{\varphi}(\overline{\gamma}_k), \tilde{\varphi}(\overline{\Gamma}_k), \mathbb{R}^2 \setminus \overline{\mathbb{D}}) = \text{Cap}(\overline{\gamma}_k, \overline{\Gamma}_k, \mathbb{R}^2 \setminus \overline{\Omega}) \leq \text{Cap}(\overline{\gamma}_k, \overline{\Gamma}_k, \mathbb{R}^2).$$

Hence by (A.13) and Lemma A.3 we know that

$$\ell(\gamma_k) \geq \text{diam}(\gamma_k) \gtrsim \text{diam}(\Gamma_k) \sim \ell(\Gamma_k).$$

Case 2: $\text{dist}(\tilde{\varphi}(\gamma_k), \partial\mathbb{D}) \geq \frac{1}{3} \text{dist}(\tilde{\varphi}(\Gamma_k), \partial\mathbb{D})$. This assumption implies that $\tilde{\varphi}(\gamma_k) \cup \tilde{\varphi}(\Gamma_k)$ is contained in a Whitney-type set. Then $\gamma_k \cup \Gamma_k$ is also contained in a Whitney-type set by Lemma 2.6, and then the desired estimate follows directly from Lemma 2.6 and (A.12) via a change of variable.

Case 3:

$$\text{dist}(\tilde{\varphi}(\gamma_k), \tilde{\varphi}(\Gamma_k)) < \frac{1}{3} \text{dist}(\tilde{\varphi}(\Gamma_k), \partial\mathbb{D})$$

and

$$\text{dist}(\tilde{\varphi}(\gamma_k), \partial\mathbb{D}) < \frac{1}{3} \text{dist}(\tilde{\varphi}(\Gamma_k), \partial\mathbb{D}).$$

In this case, there is a subcurve $\tilde{\gamma}_k \subset \gamma_k$ such that $\ell(\tilde{\varphi}(\tilde{\gamma}_k)) \gtrsim \ell(\tilde{\varphi}(\Gamma_k))$ and $\text{dist}(\tilde{\varphi}(\tilde{\gamma}_k), \partial\mathbb{D}) \gtrsim \text{dist}(\tilde{\varphi}(\Gamma_k), \partial\mathbb{D})$, as γ_k is a (connected) curve. Then we are reduced to a case similar to the second one, and it follows that

$$\ell(\gamma_k) \gtrsim \ell(\tilde{\gamma}_k) \geq \text{diam}(\tilde{\gamma}_k) \gtrsim \text{diam}(\Gamma_k) \sim \ell(\Gamma_k).$$

Consequently we obtain the desired estimate. \square

We call the set $\tilde{\varphi}^{-1}(A(z, k))$ defined above a *conformal annulus* centered at z . Now let us prove Theorem 2.3.

Proof of Theorem 2.3. We first deal with the case where $\tilde{\varphi}(x), \tilde{\varphi}(y)$ are not on the same radial ray. Extend the hyperbolic geodesic joining x and y to the boundary of $\tilde{\Omega}$ at z_1 and z_2 , and construct the conformal annuli as in Lemma A.4

We first consider the case where x, y are in the same conformal annulus (centered at z_1 or z_2). Then since $\tilde{\varphi}(x), \tilde{\varphi}(y)$ are contained in some Whitney-type set of the exterior of the unit disk by the planar geometry, via from Lemma 2.6 we know that x, y are in some Whitney-type set of $\tilde{\Omega}$, and the claim follows directly.

Then we show the claim in the case where x and y are in different annuli. We may assume that they lie on the boundary of different conformal annuli, since the general case follows from the triangle inequality with the conclusion of the first case.

We employ the notation in Lemma A.4. If $|\tilde{\varphi}(x) - \tilde{\varphi}(y)| \leq 1$, then by applying the Jordan curve theorem to the union of the subarc of $\partial\mathbb{D}$ inside $B(\tilde{\varphi}(z), 2^{k-1})$ with the inner boundary of the conformal annulus $A(z, k)$, with the help of the conformal map we conclude that, any curve joining x and y intersects the inner boundary of $\varphi(A(z, k))$ whenever Γ_k is non-empty. Similarly via the Jordan curve theorem we obtain that any curve joining x and y intersects the outer boundary of $\varphi(A(z, k))$ whenever Γ_k is non-empty. In other words, any curve joining x and y should go across the conformal annulus $\varphi(A(z, k))$ whenever Γ_k is non-empty. Then by Lemma A.4 we have

$$\ell(\Gamma) \leq \sum_k \ell(\Gamma_k) \lesssim \sum_k \ell(\gamma_k) \lesssim \ell(\gamma)$$

for some absolute constants. Therefore we obtain the desired conclusion.

Suppose $|\tilde{\varphi}(x) - \tilde{\varphi}(y)| > 1$. If $\Gamma_k \neq \emptyset$ but $\gamma_k = \emptyset$, then the Jordan curve theorem (with the argument above) implies that $k \geq 1$. By $|\tilde{\varphi}(x) - \tilde{\varphi}(y)| > 1$ and the fact that $\tilde{\varphi}(\Gamma) \subset B(0, 100)$, according to (A.2) we have

$$\text{Cap}(\Gamma_k, \gamma, \tilde{\Omega}) = \text{Cap}(\tilde{\varphi}(\Gamma_k), \tilde{\varphi}(\gamma), \mathbb{R}^2 \setminus \overline{\mathbb{D}}) \geq C.$$

Thus by Lemma A.3 and Lemma A.4 we have

$$\text{diam}(\Gamma_k) \leq C \text{diam}(\gamma) \leq C \ell(\gamma)$$

for some absolute constants. Therefore by Lemma A.4 again we conclude that

$$\ell(\Gamma) \leq \sum_{k < 0} \ell(\Gamma_k) + \sum_{0 \leq k \leq 7} \ell(\Gamma_k) \lesssim \sum_{k < 0} \ell(\gamma_k) + \sum_{0 \leq k \leq 7} \ell(\Gamma_k) \lesssim \ell(\gamma).$$

Hence we conclude the theorem in the case where $\tilde{\varphi}(x), \tilde{\varphi}(y)$ are not on the same radial ray.

If $\tilde{\varphi}(x), \tilde{\varphi}(y)$ are on the same radial ray, we extend the hyperbolic geodesic joining x and y to the boundary of $\tilde{\Omega}$ at z_1 , and only consider the conformal annuli centered at z_1 . Then the above argument also applies similarly to this case, and we conclude the theorem. \square

A.3. A lemma on quasiconvex sets.

Proof of Lemma 2.14. We may assume that $0 \notin \tilde{\Omega}$. Let $S = \partial B(0, 10 \operatorname{diam}(\partial \tilde{\Omega}))$, and Q be a $4\sqrt{2}$ -Whitney square of $\tilde{\Omega}$ intersecting S . Then $S \subset \tilde{\Omega}$ and $\operatorname{diam}(Q)/\operatorname{dist}(Q, \partial \tilde{\Omega})$ is bounded from below by an absolute constant. We claim that

$$\operatorname{Cap}(Q, \partial \tilde{\Omega}, \tilde{\Omega}) \gtrsim 1. \quad (\text{A.14})$$

To begin with, by (A.3) we have

$$\operatorname{Cap}(Q, \partial \Omega, \tilde{\Omega}) = \operatorname{Cap}(\partial Q, \partial \Omega, \tilde{\Omega} \setminus Q).$$

Denote the center of Q by x_0 . Since the Möbius transformation $\phi: z \mapsto \frac{\operatorname{diam}(Q)}{(z-x_0)}$ is uniformly biLipschitz in $\{z \in \mathbb{R}^2: 0 < \operatorname{dist}(z, Q) \leq 2 \operatorname{dist}(Q, \partial \Omega)\}$ as Q is a Whitney square, we have

$$\frac{\operatorname{diam}(\partial Q)}{\operatorname{dist}(Q, \partial \Omega)} \sim \frac{\phi(\operatorname{diam}(\partial Q))}{\operatorname{dist}(\phi(Q), \phi(\partial \Omega))}$$

and thus by (A.4) together with our assumption

$$\operatorname{Cap}(\phi(\partial Q), \phi(\partial \Omega), \phi(\tilde{\Omega} \setminus Q)) \geq C(\sigma).$$

Then (A.14) follows from the conformal invariance of capacity as ϕ is conformal in the ring domain $G \setminus Q$.

By the conformal invariance of capacity, we conclude from (A.14) that

$$\operatorname{Cap}(\tilde{\varphi}(Q), \partial \mathbb{D}, \mathbb{R}^2 \setminus \overline{\mathbb{D}}) \gtrsim 1.$$

Notice that by Lemma 2.6, $\tilde{\varphi}(Q)$ is of λ -Whitney-type for some absolute constant λ . Hence

$$\operatorname{diam}(\tilde{\varphi}(Q)) \sim_{\lambda} \operatorname{dist}(\partial \mathbb{D}, \tilde{\varphi}(Q)) \lesssim \operatorname{diam}(\mathbb{D}) = 2$$

according to Lemma A.3. By the triangle inequality and the arbitrariness of Q we have

$$\operatorname{dist}(\tilde{\varphi}(S), \partial \mathbb{D}) \leq C_2^{-1} \quad \text{and} \quad 1 \leq \operatorname{diam}(\tilde{\varphi}(S)) \leq C_2^{-1}. \quad (\text{A.15})$$

where $C_2 < 1$ is an absolute constant.

Since $\tilde{\Omega}$ is C_1 -quasiconvex, then for the points z_1, z_2 in the statement of the lemma, there exists a curve $\gamma \subset \tilde{\Omega}$ joining z_1 and z_2 such that

$$\ell(\gamma) \leq \delta_1 C_1 \operatorname{diam}(\partial \tilde{\Omega}).$$

By letting $\delta_1 < C_1^{-1} \leq 1$, with the definition of z_1, z_2 , we have $\gamma \subset B(0, 4 \operatorname{diam}(\partial \tilde{\Omega}))$ by the triangle inequality. Then by (A.4) again we obtain

$$\operatorname{Cap}(\gamma, S, \tilde{\Omega}) \leq \phi_2(\delta_1 C_1/6). \quad (\text{A.16})$$

By conformal invariance we have

$$\operatorname{Cap}(\tilde{\varphi}(\gamma), \tilde{\varphi}(S), \mathbb{R}^2 \setminus \overline{\mathbb{D}}) \leq \phi_2(\delta_1 C_1/6). \quad (\text{A.17})$$

Notice that since $\gamma \subset B(0, 4 \operatorname{diam}(\partial \tilde{\Omega}))$, then $\tilde{\varphi}(\gamma)$ is contained in the Jordan domain enclosed by $\tilde{\varphi}(S)$. Hence if

$$|\tilde{\varphi}(z_1) - \tilde{\varphi}(z_2)| \geq \frac{1}{4},$$

by (A.15), (A.4) and (A.17) we further have

$$\phi_1(C_3/4) \leq \text{Cap}(\tilde{\varphi}(\gamma), \tilde{\varphi}(S), \mathbb{R}^2 \setminus \overline{\mathbb{D}}) \leq \phi_2(\delta_1 C_1/6). \quad (\text{A.18})$$

Thus by choosing δ_1 small enough such that $\delta_1 < C_1^{-1}$ but (A.18) fails, we obtain the first part of the lemma.

Towards the second part, by symmetry we only need to prove the inequality for z_1 . If $z_1 \in \partial\tilde{\Omega}$, then by the Caratheodory-Osgood theorem we know that $\tilde{\varphi}(z_1) \in \partial\mathbb{D}$, and the desired claim follows. If z_1 is in the interior of $\tilde{\Omega}$, let Γ be the hyperbolic geodesic joining z_1 to ∞ . Then since $\text{dist}(z_i, \partial\tilde{\Omega}) \leq \delta_1 \text{diam}(\partial\tilde{\Omega})$, by (A.4) we have

$$\text{Cap}(\partial\tilde{\Omega}, \Gamma, \tilde{\Omega}) \geq \phi_1(\delta_1^{-1}) \quad (\text{A.19})$$

for some absolute constant. By the conformal invariance of capacity, we conclude from (A.19) that

$$\text{Cap}(\partial\mathbb{D}, \tilde{\varphi}(\Gamma), \mathbb{R}^2 \setminus \overline{\mathbb{D}}) \geq \phi_1(\delta_1^{-1}).$$

Then the desired estimate follows from Lemma A.3 by choosing δ_1 suitably. \square

REFERENCES

- [1] Kari Astala, Tadeusz Iwaniec, Gaven Martin, *Elliptic partial differential equations and quasiconformal mappings in the plane*. Princeton Mathematical Series, 48. Princeton University Press, Princeton, NJ, 2009.
- [2] Z. M. Balogh, S. M. Buckley *Geometric characterizations of Gromov hyperbolicity*. Invent. Math. **153** (2003), no. 2, 261–301.
- [3] D. Burago, Y. Burago, S. Ivanov, *A course in metric geometry*. Graduate Studies in Mathematics, 33. American Mathematical Society, Providence, RI, 2001.
- [4] Y. D. Burago, V. G. Maz'ya, *Certain questions of potential theory and function theory for regions with irregular boundaries (Russian)*. Zap. Nauchn. Semin. LOMI **3**, 152 p. (1967).
- [5] M. Bonk, J. Heinonen, P. Koskela, *Uniformizing Gromov hyperbolic spaces*. Astérisque, No. 270 (2001).
- [6] S. Buckley, P. Koskela, *Sobolev-Poincaré implies John*. Math. Res. Lett. **2** (1995), no. 5, 577–593.
- [7] Adrien Douady, John H. Hubbard, *Exploring the Mandelbrot set. The Orsay Notes*. [http://3dmandelbrot.com/files/Exploring the Mandelbrot set Douady and Hubbard.pdf](http://3dmandelbrot.com/files/Exploring%20the%20Mandelbrot%20set%20Douady%20and%20Hubbard.pdf)
- [8] L. C. Evans, R. F. Gariepy, *Measure theory and fine properties of functions*. Studies in Advanced Mathematics. CRC Press, Boca Raton, FL, 1992.
- [9] F. W. Gehring, K. Hag, O. Martio, *Quasihyperbolic geodesics in John domains*. Math. Scand. **65** (1989), no. 1, 75–92.
- [10] F. W. Gehring, W. K. Hayman, *An inequality in the theory of conformal mapping*. J. Math. Pures Appl. (9) **41** (1962) 353–361.
- [11] F. W. Gehring, O. Martio, *Quasixremal distance domains and extension of quasiconformal mappings*. J. Analyse Math. **45** (1985), 181–206.
- [12] M. Ghamisari, R. Näkki and J. Väisälä, *John disks and extension of maps*. Mh. Math. **117** (1994), 63–94.
- [13] V. M. Gol'dshtein and Yu. G. Reshetnyak, *Quasiconformal mappings and Sobolev spaces*, Mathematics and its Applications (Soviet Series), **54** (1990), Kluwer Academic Publishers Group, Dordrecht.
- [14] V. M. Gol'dshtein, T. G. Latfullin and S. K. Vodop'yanov, *A criterion for the extension of functions of the class L_2^1 from unbounded plain domains (Russian)*, Sibirsk. Mat. Zh. **20** (1979), 416–419.
- [15] H. Hakobyan and D. A. Herron, *Euclidean quasiconvexity*, Ann. Acad. Sci. Fenn. Math. **33** (2008), 205–230.
- [16] P. Hajlasz, P. Koskela and H. Tuominen, *Sobolev embeddings, extensions and measure density condition*, J. Funct. Anal. **254** (2008), 1217–1234.
- [17] D.A. Herron, *Riemann maps and diameter distance*. Amer. Math. Monthly **119** (2012), 140–147.

- [18] D.A. Herron and P. Koskela, *Uniform, Sobolev extension and quasiconformal circle domains*, J. Anal. Math. **57** (1991), 172–202.
- [19] P. W. Jones, *Quasiconformal mappings and extendability of Sobolev functions*, Acta Math. **47** (1981), 71–88.
- [20] P. Koskela, M. Miranda Jr. and N. Shanmugalingam, *Geometric properties of planar BV-extension domains*, International Mathematical Series (N.Y.) **11** (2010), no. 1, 255–272.
- [21] Pekka Koskela, Tapio Rajala, Yi Zhang, *A geometric characterization of planar Sobolev extension domains*, Sci. China Math. **68** (2025), 2347–2416.
- [22] Pekka Koskela, Yi Zhang, *A density problem on Sobolev spaces*, Arch. Ration. Mech. Anal. **222** (2016), no. 1, 1–14.
- [23] R. Näkki and J. Väisälä, *John disks*, Exposition. Math. **9** (1991), 3–43.
- [24] B. Palka, *An introduction to complex function theory*, Undergraduate Texts in Mathematics, New York: Springer-Verlag, 1991.
- [25] Ch. Pommerenke, *Boundary behaviour of conformal maps*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 299. Springer-Verlag, Berlin, 1992.
- [26] S. Rickman, *Quasiregular mappings*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], 26. Springer-Verlag, Berlin, 1993.
- [27] W. Rudin, *Real and complex analysis*. Second edition. McGraw-Hill Series in Higher Mathematics. McGraw-Hill Book Co., New York-Düsseldorf-Johannesburg, 1974.
- [28] P. Shvartsman, *On Sobolev extension domains in \mathbb{R}^n* , J. Funct. Anal. **258** (2010), no. 7, 2205–2245.
- [29] C. Thomassen, *The Jordan-Schönflies theorem and the classification of surfaces*. Amer. Math. Monthly **99** (1992), no. 2, 116–130.
- [30] Jussi Väisälä, *Lectures on n -dimensional quasiconformal mappings*. Lecture Notes in Mathematics, Vol. 229. Springer-Verlag, Berlin-New York, 1971.
- [31] M. Vuorinen, *Conformal geometry and quasiregular mappings*. Lecture Notes in Mathematics, 1319. Springer-Verlag, Berlin, 1988.
- [32] N. Zobin, *Extension of smooth functions from finitely connected planar domains*. J. Geom. Anal. **9** (1999), no. 3, 491–511.

UNIVERSITY OF JYVASKYLA, DEPARTMENT OF MATHEMATICS AND STATISTICS, P.O. Box 35 (MAD),
FI-40014 UNIVERSITY OF JYVASKYLA, FINLAND

Email address: pekka.j.koskela@jyu.fi

Email address: tapio.m.rajala@jyu.fi

ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, THE CHINESE ACADEMY OF SCIENCES, BEIJING
100190, CHINA

Email address: yzhang@amss.ac.cn