

The Space-Based Time-Domain Revolution in Astrophysics

Daniel Huber,^{1,2}

¹Institute for Astronomy, University of Hawai'i, 2680 Woodlawn Drive, Honolulu, HI 96822, USA; email: huberd@hawaii.edu

²Sydney Institute for Astronomy (SIfA), School of Physics, University of Sydney, NSW 2006, Australia

Xxxx. Xxx. Xxx. Yyyy. AA:1–50

[https://doi.org/10.1146/\(\(please add article doi\)\)](https://doi.org/10.1146/((please add article doi)))

Copyright © YYYY by the author(s). All rights reserved

Keywords

space telescopes, time domain astronomy, asteroseismology, stellar rotation, stellar granulation, stellar flares, young stellar objects, milky way galaxy, black holes, supernovae, active galactic nuclei, asteroids, solar system planets, citizen science

Abstract

Space-based time-domain telescopes such as CoRoT, Kepler/K2 and TESS have profoundly impacted astrophysics over the past two decades. Continuous light curves with high cadence and high photometric precision are now available for millions of sources within our galaxy and beyond. In addition to revolutionizing exoplanet science, the data have enabled breakthroughs ranging from the solar system to stellar interiors, the transient universe, and active galaxies.

- Stellar astrophysics has been transformed by the ability to probe the internal structures of stars, test the physics of stellar convection, connect stellar rotation and magnetic activity, and reveal complex variability in young stars.
- Ages of stellar populations probe the formation history of our Milky Way, and binary star variability enables the detection of “dark” galactic populations such as solar-mass black holes and neutron stars.
- Early-time observations of explosive transients provide new insights into the progenitors of supernovae, while the quasi-periodic variability of galaxies probes the physics of accretion processes onto supermassive black holes and the tidal disruption of stars.
- Observations of solar system objects reveal asteroid compositions through their rotation periods and amplitudes, constrain the cloud structure of ice giants, and allow the discovery of new objects in the outer solar system.
- Open data policies and software have contributed to remarkable scientific productivity and enabled discoveries by citizen scientists, including new exoplanets and exotic variability in mature Sun-like stars.

Contents

1. Introduction	2
1.1. Scope of this Review	2
1.2. A Brief History of Space-Based Time-Domain Photometry	3
1.3. Fundamental Parameters of Space-Based Time-Domain Missions	5
1.4. From Exoplanets to Astrophysics: the Evolution of Target Selection	9
2. Stellar Astrophysics	10
2.1. Asteroseismology	10
2.2. Convection and Granulation	13
2.3. Rotation and Stellar Activity	15
2.4. Flares and Outbursts	16
2.5. Dippers, Bursters, and Exotic Variables	18
2.6. Binary Stars	19
3. Galactic Astronomy	21
3.1. Galactic Archaeology	21
3.2. Neutron Stars and Black Holes	24
4. Extragalactic Science	25
4.1. Explosive Transients	25
4.2. Active Galactic Nuclei	26
4.3. Tidal Disruption Events	27
5. Solar System Science	28
5.1. Small Bodies	28
5.2. Dwarf Planets and Planet Nine	29
5.3. Ice Giant Planets	29
6. Citizen Science	30
7. The Future of Space Based Time-Domain Photometry	31

1. Introduction

1.1. Scope of this Review

Time-domain photometry — measuring the brightness of objects over multiple epochs — has a long history in astronomy. Examples include the first recorded supernova explosions by Chinese astronomers over 2000 years ago (“guest stars”, [Clark & Stephenson 1977](#)), the naked-eye detection of variability in Mira by David Fabricius in the 16th century ([Olbers 1850](#)), the discovery of eclipsing binary stars ([Goodricke 1783](#)) and the pioneering observations of Cepheid pulsations that led to the discovery of the expansion of the Universe ([Leavitt 1908](#)). Over the past 20 years, time-domain observations from dedicated space-based telescopes have led to a revolution in astrophysics. Uninterrupted, high-precision light curves for millions of sources have affected fields ranging from solar system science, stellar astrophysics, exoplanet science, galactic stellar populations, supernovae, and galaxy formation. Astronomy has now entered an era in which a substantial number of stars in our galaxy have continuous high-precision space-based light curves, allowing large-scale variability classification efforts akin to stellar spectral typing in the early 20th century.

This review will describe breakthroughs in astrophysics enabled by space-based time-domain telescopes. Particular focus will be given to results from NASA’s Kepler/K2 ([Borucki et al. 2010](#), [Howell et al. 2014](#)) and TESS missions ([Ricker et al. 2015](#)), as op-

posed to time-domain observations from general purpose observatories such as the Hubble space telescope or Gaia. The review is intentionally broad in scope, covering high-level results across a range of topics with the goal to provide sufficient information on the current state of each field, but not replacing in-depth reviews. The review will only briefly cover exoplanet science and asteroseismology, which are the traditional scientific drivers for dedicated space-based time-domain telescopes, and have been the subject of other reviews (e.g. [Aerts 2021](#), [Kurtz 2022](#), [Lissauer et al. 2023](#)).

1.2. A Brief History of Space-Based Time-Domain Photometry

Time-domain observations from space have long been recognized as a major opportunity to advance our understanding of exoplanets and stellar astrophysics. The first concepts for a dedicated time-domain monitoring of stars from space were put forward in Europe in the 1990's, including PRISMA ([Appourchaux et al. 1993](#)), STARS ([Fridlund et al. 1995](#)), MONS ([Kjeldsen et al. 2000](#)) and Eddington ([Favata 2004](#)). Eddington was an ESA-led mission with a planned 1.2 m aperture to detect exoplanets using the transit method and perform asteroseismology of Sun-like stars. While it was selected, none of the above missions were ultimately funded for launch.

The first successfully launched mission dedicated to space-based time-domain photometry was the Canadian space telescope MOST (Microvariability and Oscillations in Stars, [Walker et al. 2003](#), [Matthews 2007](#)). MOST was a 15 cm telescope in a sun-synchronous low-Earth orbit, targeting individual stars for up to 90 continuous days with cadences as short as one minute with the aim to probe the structure of stars using asteroseismology. MOST successfully discovered pulsations in a range of intermediate-mass and massive stars, as well as oscillations in red giants ([Kallinger et al. 2008](#)). MOST also measured rotational modulation in young stars ([Croll et al. 2006](#)) and detected some of the first exoplanet transits observed from space ([Rowe et al. 2006](#), [Winn et al. 2011](#)).

The start of the space-photometry revolution is generally associated with the launch of the French-led CoRoT (Convection Rotation and Planetary Transits) satellite in 2006 ([Baglin et al. 2006](#)). CoRoT was also launched in a sun-synchronous orbit and carried out the first space-based “long-stare” campaigns lasting for nearly five months in two dedicated fields. CoRoT furthermore made several important advancements in exoplanet science, including the detection of a transiting rocky exoplanet ([Léger et al. 2009](#)). CoRoT also detected oscillations in several main-sequence stars (see, e.g., [Michel et al. 2008](#)) and in several thousands of red-giant stars (see, e.g., [De Ridder et al. 2006](#), [Hekker et al. 2009](#)).

In parallel, other space telescopes were used for time-domain astronomy. For example, the Hubble space telescope was used for both stellar variability studies ([Edmonds & Gilliland 1996](#), [Gilliland 2008](#)) and the characterization of transiting exoplanets ([Nutzman & Charbonneau 2008](#)). Furthermore, innovative approaches used ancillary data for stellar astrophysics, including the star-tracker of the failed WIRE (Wide-Field Infrared Explorer) satellite ([Schou & Buzasi 2001](#), [Retter et al. 2003](#), [Bruntt et al. 2005](#), [Stello et al. 2008](#)), as well as the SMEI (Solar Mass Ejection Imager) experiment ([Tarrant et al. 2007](#)).

The launch of the Kepler space telescope in 2009 continued the revolution of space-based photometry. With the advantage of a large aperture, an Earth-trailing orbit, and continuous observations of a single field, Kepler provided light curves with unprecedented precision and duration for over 150,000 pre-selected stars ([Borucki et al. 2010](#), [Batalha et al. 2010](#)). Following the failure of two reaction wheels, Kepler ceased observations of




Figure 1

The number of stars with precise, continuous space-based light curves has increased by four orders of magnitude in the past 20 years. Approximate number of stars observed with precisions of <1 mmag per hour as a function of year of launch of space-based missions. Symbol areas scale with the typical time baseline of continuous observations. Dashed circles for TESS and PLATO denote the typical longest continuous observations. Color coding shows the wavelength of the central bandpass. BRITE Constellation is the only mission which observed with multiple bandpasses (CoRoT used a prism to obtain spectral information for exoplanet targets). See Table 1 for details on yield estimates.

its original field after nearly 4 years of operations. Thanks to engineering ingenuity, its successor mission K2 performed 70 day campaigns along the ecliptic plane, observing a total of nearly 400,000 pre-selected stars over the next 3 years (Howell et al. 2014). The end of the Kepler/K2 era came in 2018, when spacecraft operations were discontinued due to lack of fuel, which prevented it from maintaining precise pointing control.

BRITE constellation, a collaboration between Austria, Canada, and Poland, launched in 2013 to perform the first dedicated multi-color time-domain photometry from space (Weiss et al. 2014). The constellation consists of six nanosatellites with 3 cm apertures, each equipped either a red or blue filter, to perform precise two-colour photometry of naked-eye ($V < 6$) stars. Key results so far include the study of mass loss in Be stars (Baade et al. 2016), the discovery of the most massive star in a heartbeat binary system to date (Pablo et al. 2017), and the detection of oscillations in nearby red giants (Kallinger et al. 2019).

The next phase of the space-photometry revolution was enabled by TESS, which launched in a lunar resonance orbit in 2018 (Ricker et al. 2015, Winn 2024). TESS is largely conducting an all-sky survey strategy, obtaining on average 27-day light curves with continuous coverage for up to one year near the ecliptic poles. The proximity to Earth allows TESS the collection of full-frame images, thus removing the need for target selection except for stars which require the fastest cadences. At the time of writing, TESS is mon-

itoring 2000 stars per observing sector in 20-second cadence, tens of thousands of stars in 2-min cadence, and obtains full frame images at 200-second cadence¹.

Figure 1 summarizes the above “brief history” of space-based photometry by comparing the number of stars with <1mmag/hr photometric precision as a function of launch date of the mission². Over the past 20 years, the number of stars with such data has increased by 4 orders of magnitude. The next chapters of the space-photometry revolution will be written by ESA’s PLATO (Rauer et al. 2025), NASA’s Roman (Spergel et al. 2013), and the Chinese Earth 2.0 (Ge et al. 2022) missions, which are slated for launch in the second half of this decade. Chapter 7 provides a brief outlook on these new time-domain missions.

1.3. Fundamental Parameters of Space-Based Time-Domain Missions

Space-based time-domain missions can be characterized by a set of fundamental parameters, which set the framework for achievable science with a given mission. This section gives a pedagogical introduction into the most common parameters of time-domain missions.

- The *photometric precision* is the relative uncertainty of brightness measurements. The detected signal from a star is given by:

$$S = F_* \times t \times \pi(D/2)^2 \times (h\nu)^{-1} \times Q. \quad 1.$$

Here, S is the total number of electrons or counts received from the star, F_* is the flux emitted per unit time and unit area, t is the integration time, D is the telescope diameter, $h\nu$ is the photon energy and Q includes instrument characteristics such as detector quantum efficiency and system throughput. In the idealized limit of photon noise, the measurement error or noise is $N = \sqrt{S}$. The photometric precision, which is the relative error on a flux measurement, is thus given by $\sigma = N/S = S^{-1/2}$. For a fixed bandpass and detector we therefore have:

$$\sigma \propto F_*^{-1/2} \times t^{-1/2} \times D^{-1}. \quad 2.$$

As expected, better photometric precision (lower σ) is achieved for larger telescope apertures, longer integration times and brighter stars. Equation 2 allows simple scalings from one mission to another. For example, Kepler’s aperture is approximately ten times larger than TESS’s. Therefore, for a fixed brightness of a star, the photometric precision per unit time (fixed cadence) is 10 times better for Kepler compared to TESS. Equivalently, a star needs to be 100 times (5 magnitudes) brighter for TESS to reach the same photometric precision per unit time as Kepler. Equation 2 is a good approximation for space-based photometry, which does not suffer from scintillation noise, but breaks down for bright stars due to saturation and at the faint stars due to sky noise and detector read noise.

Space-based time-domain missions typically collect data in sub-exposures that are co-added on board to a given cadence, which is then downloaded to the ground. For example, for Kepler the exposure time for individual frames was 6 seconds, while for TESS individual exposure times are 2 seconds. This explains why the light curve

Photometric Precision: Fractional uncertainty of brightness measurements; in the limit of photon noise, the precision increases with telescope aperture, integration time, and stellar brightness.

¹Full frame images were obtained at 30-minute cadence during the first 2 years of the TESS Mission, and at 10-minute cadence during years 3-4.

²The figure excludes CHEOPS (Benz et al. 2021), which is a targeted follow-up mission.




Figure 2

For a given space-based time-domain telescope, the light curve scatter per unit time is constant. Each panel shows a 12 hour segment of the same star using a different TESS data product: 30-min sampling (top), 10-minute sampling (second from top), 2-min sampling (second from bottom) and 20-second sampling (bottom). The x-axis and y-axis scales are identical in all panels.

scatter for a given star increases as the time sampling of the data product increases (Figure 2). For example, comparing the light curve of the same star observed by TESS with 30-minute cadence and 20-second time sampling shows that the light curve scatter of the former is smaller by a factor $\sqrt{30 \times 60/20} \approx 10$, since each data point is based on a longer effective exposure time. In other words, the light curve scatter per unit time remains constant for a given instrument.

- The *time baseline* is the typical length of a light curve. It sets a limit on the lowest frequency (longest period) that can be unambiguously measured from a light curve. For evenly spaced data, the frequency resolution in the Fourier domain is given by:

$$\Delta f = 1/T. \quad 3.$$

In frequency analysis, Fourier spectra are commonly oversampled by a factor proportional to Δf (typically 5 – 10).

The maximum time baseline is determined by the spacecraft orbit. Continuous observations of a single field in a sun-synchronous orbit around Earth (such as for MOST, CoRoT, BRITE) are typically limited to ≈ 5 months due to Earth's orbital motion around the Sun. Kepler, located in an Earth-trailing orbit, allowed uninterrupted observations for fields off the ecliptic plane, with quarterly rolls to keep the spacecraft oriented towards the Sun. The innovative lunar resonance orbit of TESS allows

Time Baseline:

Length of uninterrupted observations; sets lowest measurable frequency (longest detectable period).

uninterrupted observations near the ecliptic poles, while engineering requirements related to illumination of solar panels and thermal stability restrict typical observing lengths to ≈ 27 days. Planned missions such as Roman, PLATO and Earth 2.0 will be deployed in an orbit around the second Lagrange point (L2), allowing long uninterrupted observations over large parts of the sky. Disadvantages of this solution are higher mission costs, driven by the requirement for larger antennas to download data back to Earth, and overall lower data rates.

The time baseline can refer to the length of continuous observations or the total length of timeseries. The difference is the duty cycle, which is the fraction of time observations have been taken between the first and the last datapoint. For example, Kepler has collected the (so-far) longest continuous timeseries to date (4 years with a duty cycle close to 90%), while TESS is currently obtaining the longest total timeseries (> 8 years with a maximum duty cycle of $\approx 50\%$).

- The *observing cadence* or *time sampling* is the number of observations per unit time that are downloaded from the spacecraft³. The cadence sets a fundamental limit on the highest frequency (or shortest period) of signal that can be recovered from a light curve. The upper limit for a time sampling Δt is given by the Nyquist frequency:

$$f_{\text{Nyq}} = 1/2\Delta t. \quad 4.$$

The Nyquist frequency is only strictly defined for equally sampled timeseries, and signals above the Nyquist frequency can be recovered from typical Kepler/K2 and TESS light curves (Murphy et al. 2013, Chaplin et al. 2014b, Liagre et al. 2025).

The time sampling also causes fractional amplitude attenuation caused by time-averaging of a signal with frequency f , which is given by:

$$A = \text{sinc}(\pi f t_{\text{exp}}), \quad 5.$$

where $\text{sinc}(x) = (\sin x)/x$ and t_{exp} is the exposure time. For observations with no dead time between exposures (such as those obtained by Kepler/K2 and TESS), the exposure time is equal to the sampling time and thus the fractional attenuation in power can be written as:

$$P = \text{sinc}^2\left(\frac{\pi}{2} \frac{f}{f_{\text{nyq}}}\right), \quad 6.$$

In the time-domain sampling can cause smearing effects, for example for the ingress and egress of transit or eclipse observations.

- The *number of targets* specifies the number of light curves produced per observing interval for a given cadence. This is predominantly set by the onboard storage capacity of the spacecraft, as well as the capability to download data to the ground and thus the orbit and communications antenna. Missions typically download smaller number of targets with faster sampling, and a larger number of targets at slower sampling.
- The *field of view and pixel size* set the spatial scales over which stars can be observed and resolved. The plate scale of an instrument is given by:

³“Cadence” can sometimes be used in a confusing manner. For example, light curves are frequently referred to as having “30-minute cadence” or “1-minute cadence”, but the latter has a higher cadence than the former. Alternative wording such as “sampling” or “faster/slower cadence” can avoid this ambiguity.

Cadence or

Sampling: Number of observations per unit time; determines highest detectable frequency (Nyquist limit).

Nyquist Frequency:

Highest frequency reliably detectable given the sampling cadence; signals above are aliased.

Field of View (FOV):

Angular sky area observed by a telescope; larger FOV enables more targets at lower spatial resolution.

Pixel Scale:

Angular size covered by one detector pixel; balances field of view with spatial resolution.

Bandpass:

Wavelength range of observations; affects photometric precision and stellar types most effectively studied.

Blending (Dilution):

Flux contamination from nearby stars within an aperture, reducing observed amplitudes.

$$s''/mm = \frac{206.265}{F[mm]} = \frac{206.265}{fD[mm]}, \quad 7.$$

where F is the focal length, D is the telescope diameter, $f = F/D$ is the f-ratio. For a given physical size and pixel dimensions of a detector, Equation 7 allows the calculation of the pixel size and field of view. For example, a single TESS detector has 4096×4096 pixels with a physical size of 62×62 mm. The focal length of $F = 146$ mm for a single TESS camera yields a pixel size of $62/4096 \times 206.265/146 \approx 21''$ and field of view of $62/3600 \times 206.265/146 \approx 24$ degrees⁴.

Blending of stars within a given aperture causes the dilution of a signal as follows:

$$A_d = A \times (1 - 10^{-0.4\Delta m}), \quad 8.$$

where A is the original signal amplitude, A_d is the diluted amplitude and Δm is the magnitude difference between the two stars in a given observing bandpass.

- The *observing bandpass* determines wavelength range over which photons are collected. The majority of space-based time-domain missions used broadband optical bandpasses in order to collect as many photons as possible, resulting in higher photometric precision (see Equation 2). TESS used a redder bandpass than Kepler/K2 to improve sensitivity for cool main-sequence stars. Multiple photometric bands can aid in a number of astrophysical contexts such as asteroseismic mode identification and temperature measurements of spots. The BRITE nanosatellite constellation is so far the only space-based time-domain mission to have used dedicated multiple filters. Some mission concepts that will make use of multiband photometry are currently under consideration (see Section 7).

In practice, many of the above parameters are linked. For example, there is a fundamental “deep and narrow” versus “shallow and wide” tradeoff between aperture size (photometric precision) and field of view, and therefore number of observable targets. Kepler’s larger aperture resulted in a smaller FOV allowed with a smaller pixel size ($4''$), whereas the large field of view from TESS comes at the cost of $21''$ pixels and a 10 times smaller aperture. On the other hand, the proximity of TESS orbit to Earth allows data to be downloaded at a higher rate for a given antenna size, while the number of stars observed by Kepler was limited to data downlink rate from an Earth-trailing orbit. Table 1 summarizes the parameters for missions shown in Figure 1.

Systematic errors are another important characteristic of space-based missions. For time-domain missions, important examples include:

- Gaps due to periodic downlinks of data to Earth, which can introduce aliasing and affect the recovery of periodicities in the data.
- Spacecraft motion, which causes periodic changes in the point-spread function, aperture flux losses and intrapixel sensitivity changes. This was a challenge for K2, which only operated with two reaction wheels and thus experienced periodic drifts.
- Pointing inaccuracies, which cause changes in the position of stars on the detector after each observing segment. This can lead to offsets in absolute flux, requiring

⁴The calculation does not take into account optical design considerations required to ensure aberrations remain small over a large FOV.

Table 1 Observational parameters of past, current and approved space-based time-domain missions.

Mission	Launch	T_{\max}	D	FOV	Res	Δt	T_{cont}	λ	n
—	—	yrs	cm	(deg ²)	"/pix	sec	days	nm	—
WIRE	1999	6	5	64	60	15	20	620	~ 200
MOST	2003	15.8	15	5	3	30	60	550	~ 3000
CoRoT	2006	7.4	27	7.6	2.3	32, 512	150	650	$\sim 4 \times 10^4$
Kepler	2009	4.0	95	105	4	60, 1800	1460	600	$\sim 2 \times 10^5$
BRITE	2013	$>12.6^\dagger$	3	24	28	10	180	420, 620	~ 700
K2	2014	4.4	95	105	4	60, 1800	70	600	$\sim 5 \times 10^5$
TESS	2018	$>7.5^\dagger$	4×10	2300	21	20, 120, 600	27	800	$\sim 10^6$
Roman	2026	5^\ddagger	240	0.28	0.11	720	70	1290	$\sim 10^7$
PLATO	2026	4^\ddagger	12×26	2250	15	2.5, 25, 50, 600	1460	750	$\sim 10^6$
Earth2.0	2028	4^\ddagger	6×30	500	4	30, 600, 1800	1460	675	$\sim 10^6$

Launch dates for Roman, PLATO and Earth 2.0 are estimated. K2 was a continuation of the Kepler mission, and thus the year refers to the start of operations rather than a launch date. T_{\max} denotes the mission duration, and thus the upper limit on the longest possible time baseline of observations († current missions that are still active, ‡ nominal durations for planned missions excluding possible extensions). FOV is the total field of view. Δt and T_{cont} are the typical cadence and duration of continuous light curves, n is the number of observed stars with $<1\text{mag/hr}$ photometry, λ is the central wavelength of the observing bandpass. For Roman, FOV and T_{cont} are taken from the Galactic Bulge Time Domain Survey. For missions up to and including K2 n is assumed to be approximately the total number of observed stars (since the magnitude corresponding to a $<1\text{mag/hr}$ photon noise limit is fainter than most targeted stars). For TESS, n was estimated as the total number of stars in the TESS Input Catalog down to $I < 12\text{ mag}$. For Roman, n was estimated as the total number of star counts with $H < 18\text{ mag}$ based on the photometric precision model from Penny et al. (2019) and the synthetic population from Gould et al. (2015). For PLATO and Earth 2.0, n was taken as the estimated number of stars with downloaded light curves from Rauer et al. (2025) and Ge et al. (2022).

“stitching” of data segments, and causing difficulty in measuring small amplitude signals with periods longer than an observing quarter/sector.

- Background variations due to zodiacal dust and straylight reflected off the Earth or the Moon. This causes periodic flux changes which introduce systematic errors.

Some systematic errors are influenced by mission parameters. For example, spacecraft orbits close to Earth increase the effects of background variations due to straylight entering the telescope aperture. Many of the systematics mentioned above can be successfully mitigated with sophisticated data reduction techniques (e.g. Reegen et al. 2006).

1.4. From Exoplanets to Astrophysics: the Evolution of Target Selection

A major component for space-based time-domain mission is target selection, since onboard storage capacities are frequently insufficient to download all the data that is collected. Figure 3 illustrates how target selection differs between missions. Missions such as WIRE, MOST and BRITE were predominantly magnitude limited, resulting in a preference for luminous, massive main-sequence or giant stars. Kepler targeted main-sequence solar-like stars to detect Earth-like planets (Batalha et al. 2010, Wolniewicz et al. 2021), but also observed many red-giant stars because of the difficulty of discerning dwarf from giant stars in the absence of parallaxes in the pre-Gaia era (Mann et al. 2012). While these stars




Figure 3

The evolution of target selection for space-based photometry missions. Gaia color versus absolute magnitude for stars observed by space-based time-domain mission. Color-coding shows the logarithmic number density in each bin. WIRE, MOST and BRITE distributions are approximated by stars with $G < 6$, and TESS distribution with $G < 12$. The Kepler and K2 panels use actual target lists. Values are not corrected for reddening.

yielded a wealth of stellar astrophysics results, they also contended with exoplanet science: many giant stars were dropped from the target list based on their variability characteristics in favor of smaller stars that are more suitable for the detection of exoplanets.

The rich astrophysics science yield from Kepler shaped the target selection for later missions. The K2 Mission targeted a broader distribution of spectral types (Huber et al. 2016), including compact stars and massive stars, driven by the exoplanet-science focus shifting to cooler dwarfs and more diverse general astrophysics investigations. TESS full-frame images have mostly removed the need for target selection. An exception are stars requiring very fast cadence such as white dwarfs, flare stars and oscillating Sun-like stars.

2. Stellar Astrophysics

2.1. Asteroseismology

Asteroseismology is the study of pulsations to probe the interior of stars. Stellar pulsations are observed across the Hertzsprung-Russell (HR) diagram and can be broadly divided into driving mechanisms due to opacity changes in the stellar interior (so-called classical pulsators such as δ Scuti stars and Cepheids) and stars which oscillate due to turbulent

surface convection (so-called solar-like oscillators). Stellar pulsations span a wide range of timescales (minutes to years) and amplitudes (few parts-per-million to several percent), and as such are exquisitely matched to the continuous, high-precision light curves produced from space-based telescopes. A number of reviews have been dedicated to the scientific advances in asteroseismology over the past decade, including solar-like oscillators (Chaplin & Miglio 2013, García & Ballot 2019, Jackiewicz 2021) and classical pulsators (Guzik 2021, Kurtz 2022). This section provides a brief historical overview on the impact of the space-based photometry revolution on asteroseismology, with a focus on solar-like oscillators.

Prior to space-based photometry, the detection of solar-like oscillations relied on ground-based radial velocities (see Bedding & Kjeldsen 2003, for a review). The first confirmed detection of oscillations in a star other than the Sun was made in Procyon by Brown et al. (1991), followed by the first detection of regularly spaced frequencies in η Boo by Kjeldsen et al. (1995). The greatly improved radial velocity precision for detecting exoplanets enabled the detection of oscillations in several nearby main sequence and subgiant stars such as β Hyi (Bedding et al. 2001, Carrier et al. 2001), α Cen A (Bouchy & Carrier 2001, Butler et al. 2004) and B (Carrier & Bourban 2003, Kjeldsen et al. 2005), as well as red giant stars such as ξ Hya (Frandsen et al. 2002) and ϵ Oph (De Ridder et al. 2006).

However, the short timeseries and gaps in ground-based observing campaigns limited their potential. Space-based photometry from MOST initially yielded a non-detection in Procyon (Matthews et al. 2004), with later datasets showing a detection consistent with radial velocity observations (Guenther et al. 2008, Huber et al. 2011). MOST also detected oscillations in red giants (Barban et al. 2007), including the first observational evidence for non-radial modes (Kallinger et al. 2008). CoRoT achieved a major breakthrough by detecting oscillations in a number of main sequence stars (e.g. Appourchaux et al. 2008, Michel et al. 2008) and several thousands of red giant stars (e.g. Hekker et al. 2009). Importantly, CoRoT unambiguously demonstrated that red giants oscillate in non-radial modes (De Ridder et al. 2009), which opened the door for detailed studies of the interior structure of red giants (see Hekker & Christensen-Dalsgaard 2017, for a recent review).

Kepler detected oscillations in over 600 main-sequence and subgiant stars (Chaplin et al. 2014a, Mathur et al. 2022, Sayeed et al. 2025) and over twenty thousand red giants (Hekker et al. 2011, Stello et al. 2013, Mathur et al. 2016, Yu et al. 2018), enabling the study of oscillations across the low-mass H-R diagram. The larger number of red giants with detected oscillations is due to a combination of two effects: First, oscillation amplitudes increase with luminosity (Bedding 1995), making a detection easier at a given apparent magnitude. Second, the majority of targets were observed with 30-minute sampling, setting an upper limit of $\log g \sim 3.5$ since less evolved stars oscillate above the Nyquist frequency.

TESS has continued the solar-like oscillator revolution in two important ways. First, TESS observes main-sequence and subgiant stars that are much brighter and nearby than those detected by *Kepler* (Figure 4). This allows the combination of asteroseismology with classical observations such as long-baseline interferometry, which can constrain important physics such as the convective mixing length parameter (e.g. Silva Aguirre et al. 2017) and initial Helium abundances (e.g. Verma et al. 2019). TESS's 20-second cadence mode, introduced in the first extended mission, has led to an increase in photometric precision of bright stars (Huber et al. 2022), enabling the detection of oscillations in hundreds of unevolved stars (Hatt et al. 2023, Lund et al. 2025). Second, TESS has enabled the near all-sky detection of oscillating red giants, which are critical probes for galactic archeology (Section 3.1). As of writing, TESS has detected oscillations in 160,000 red giants (Hon

Asteroseismology:
Study of stellar oscillations to probe internal structure, ages, and fundamental parameters.




Figure 4

Stars with detected solar-like oscillations as a function of distance. Following ground-based RV campaigns (≈ 20 stars), CoRoT yielded the first high-quality detections from space (≈ 10 stars). Kepler enabled ≈ 700 detections, but mostly in faint and distant stars. The K2 (≈ 200 detections) and TESS (≈ 900 detections) missions have bridged the gap to nearby stars, for which asteroseismology can be combined with complementary methods. Catalog detections are taken from [Sayeed et al. \(2025\)](#) (Kepler), [Lund et al. \(2024\)](#) (K2), [Hatt et al. \(2023\)](#) and [Lund et al. \(2025\)](#) (TESS).

[et al. 2021](#)), an order of magnitude increase over Kepler, based on the analysis of only single sectors of data over most of the sky.

Space-based time-domain missions also revolutionized asteroseismology of classical pulsators such as δ Scuti stars, γ Doradus stars, pulsating OB stars and compact pulsators. Examples include the first detection of regular gravity-mode period spacings in a massive star by CoRoT ([Degroote et al. 2010](#)) and breakthroughs in the Kepler/K2 era such as the discovery of hybrid δ Scuti stars - γ Doradus stars (e.g. [Grigahcène et al. 2010](#)), core rotation period measurements for hundreds of intermediate-mass stars ([Van Reeth et al. 2016](#), [Li et al. 2020](#)), and measurements of dozens of rotation periods for white dwarfs (e.g. [Hermes et al. 2017](#)). The TESS Mission expanded the revolution to a larger sample of classical pulsators, which were sparse in Kepler/K2, leading to the discovery of stochastic low-frequency variability in massive OB stars ([Bowman et al. 2019](#)), constraints on interior mixing mechanisms ([Pedersen et al. 2021](#)) and the discovery of regular spacings in young δ Scuti stars ([Bedding et al. 2020](#)). For more extensive reviews of classical pulsators the reader is referred to [Kurtz \(2022\)](#) and [Bowman & Bugnet \(2024\)](#) for general reviews, [Aerts & Tkachenko \(2024\)](#) for fast rotators and [Córscico et al. \(2019\)](#) for pulsating white dwarfs.




Figure 5

Space-based telescopes revealed that the time-domain variability due to stellar granulation and oscillations are closely linked. Correlation between surface gravities measured through asteroseismology and the autocorrelation of light curves for stars observed by Kepler. Figure reproduced from [Kallinger et al. \(2016\)](#).

2.2. Convection and Granulation

Convection is one of two major energy transport mechanisms in stars. In stellar layers that are unstable to convection, an equilibrium perturbation will result in gas parcels continuously rising and thus transporting energy from the interior to the surface. The conditions for unstable convection depend on the temperature and composition gradients within the star ([Schwarzschild 1958](#), [Ledoux 1947](#)). Broadly speaking, the conditions for convective energy transport are found in the cores of stars with masses $\gtrsim 1.2M_{\odot}$ and in the envelopes of stars with masses $\lesssim 1.2M_{\odot}$.

Stellar granulation is a manifestation of convection on the surfaces of stars. As convective cells rise and cool, the integrated brightness of stars as observed in light curves change over time. Granulation causes brightness variations over a broad range of timescales. [Harvey \(1985\)](#) postulated that the time evolution of granulation can be approximated by a series of Lorentzian functions with typical decay timescales, which correspond to granules with typical sizes. These timescales are commonly referred to as granulation (smallest scales), mesogranulation (larger scales), and supergranulation (largest scales).

Informed by numerical simulations ([Freytag et al. 1996](#)), [Kjeldsen & Bedding \(2011\)](#) proposed scaling relations for the timescales and amplitudes of stellar granulation. The assumption is that both stellar oscillations and granulation are driven by convection, and thus the typical timescales and amplitudes scale linearly with the pressure scale height H , which for an isothermal atmosphere is given by $H \propto g^{-1}$, with g denoting the surface gravity of the star. This is the same dependence as the typical timescale of stellar oscillations (the

Granulation: Surface brightness fluctuations caused by convection; linked to stellar surface gravity and oscillations.

frequency of maximum power, ν_{\max}).

A major breakthrough of the space-based photometry revolution was the observational confirmation that granulation indeed shows a tight scaling with the properties of stellar oscillations, and thus fundamental parameters of stars. Mathur et al. (2011) first demonstrated that both the timescales and amplitudes of granulation in Kepler red giants show a tight correlation to the frequency of maximum power and thus stellar surface gravity. For main-sequence and subgiant stars, Bastien et al. (2013) derived an empirical relation between the brightness variations at a fixed 8-hour timescale (so-called “flicker”). Further investigation demonstrated that granulation tracks surface gravity as precisely as oscillations (Kallinger et al. 2014), allowing the determination of surface gravities to a precision of ≈ 0.01 dex over an astounding 5 orders of magnitude in surface gravity (Figure 5).

The tight connection between granulation and oscillations opened up a flurry of new methods to precisely measure surface gravities from light curves alone. This is significant because granulation extends over a wide range of timescales, and thus does not require the fast sampling that is required to measure oscillations. Furthermore, the combination of surface gravities with precise radii of stars derived from Gaia parallaxes in principle allows the measurement of stellar masses without the use of stellar models (Stassun et al. 2018). Bastien et al. (2016) expanded their initial framework to use an iterative process to derive typical granulation timescales. Pande et al. (2018) and Bugnet et al. (2018) expanded the approach by Bastien et al. (2016) to work in the frequency domain using Kepler stars. The large number of available space-based light curves also inspired data-driven and machine-learning approaches such as local-linear regression (Sayeed et al. 2021) and transformer-based models (Pan et al. 2024).

In addition to stars with convective envelopes, variability with observational signatures akin to granulation has now been observed in more massive stars. First evidence for this was presented by Kallinger & Matthews (2010) in δ Scuti stars observed by MOST, and recent theoretical work suggests that stars without convective envelopes may indeed show signatures of granulation (Tschernitz & Bourdin 2025). For more massive stars, TESS has revealed ubiquitous stochastic low-frequency variability (Bowman et al. 2019), which for some stars has been linked to subsurface convection (Lecoanet et al. 2019) and/or internal gravity waves (Edelmann et al. 2019, Horst et al. 2020, Serebriakova et al. 2024). This new class of variability allows insights into mixing processes in massive stars (Varghese et al. 2025), which are the progenitors of neutron stars and black holes (see Section 3.2).

The observational breakthroughs in the study of convection from space-based photometry has significant implications for our understanding of stars. The remarkably simple scaling of granulation (and oscillation) with stellar properties stands in contrast to the complex nature of convection as a physical process, which requires three dimensional models to be described accurately. Modeling granulation and oscillation amplitudes from first principles is poorly understood owing to large uncertainties when modeling the convection that stochastically drives and damps these processes (Houdek & Dupret 2015), and attempts to connect models to observations often require ad-hoc correction factors (Cranmer et al. 2014). However, significant theoretical progress is being made (e.g. Trampedach et al. 2014, Zhou et al. 2024, Eitner et al. 2024, Witzke et al. 2024, Perdomo García et al. 2025) and space-based missions will continue to provide benchmark to improve our understanding of convection.

2.3. Rotation and Stellar Activity

Rotation is among the most fundamental properties of stars. After inheriting angular momentum from the collapse of a protostellar cloud, stars change their rotation periods due to astrophysical processes such as angular momentum loss due to stellar winds and tidal interactions with other stars and planets. The former leads to the powerful possibility of using rotation periods as a proxy for stellar age (gyrochronology, [Skumanich 1972](#), [Barnes 2007](#)). Stellar activity is closely related to rotation, being driven by magnetic fields that are generated through internal dynamo processes ([Charbonneau 2010](#), [Brun & Browning 2017](#)). Observational signatures include photospheric dark spots and bright faculae, flares and chromospheric emission. A review of stellar rotation and activity for main-sequence stars is given by [Santos et al. \(2024\)](#). Probes of interior rotation through asteroseismology, which are not covered here, have been reviewed by [Aerts & Tkachenko \(2024\)](#).

Space-based light curves are exquisitely suited to study stellar rotation and activity. The small spot sizes in typical Sun-like stars are inaccessible to ground-based observations. Space-based photometry from Kepler, K2 and TESS resulted in catalogs of rotation periods for tens of thousands of cool stars across the HR diagram ([McQuillan et al. 2014](#), [Ceillier et al. 2017](#), [Santos et al. 2021](#), [Reinhold & Hekker 2020](#), [Claytor et al. 2024](#)), including a large number of young stars ([Rebull et al. 2018](#), [Douglas et al. 2019](#)). These observations have yielded a range of surprises in our understanding of angular momentum transport, including weakened magnetic braking in Sun-like stars ([van Saders et al. 2016](#), [Hall et al. 2021](#)), stalled spin-down in K-dwarfs (e.g. [Curtis et al. 2019](#)), and a “gap” in the rotation period distribution ([McQuillan et al. 2014](#)). Space-based light curves have also yielded detections of apparent rotational modulations in unexpected parameter spaces such as A stars ([Balona 2017](#)), which may be caused by chemical spots.

Space-based data also showed that stellar activity is remarkably well traced by the amplitude of rotational modulation. Kepler data demonstrated that more active stars tend to oscillate with lower amplitudes ([Garcia et al. 2010](#), [Chaplin et al. 2011a](#)), which was later quantified through the suppression of oscillations in red giants in tight eclipsing binaries ([Gaulme et al. 2013](#)). Activity measurements in thousands of Kepler main-sequence stars confirmed that stellar activity closely traces stellar rotation up to a certain threshold, after which the activity level saturates, and revealed interesting differences to spectroscopic indicators such as the absence of stars with intermediate activity levels ([Santos et al. 2024](#)).

A fascinating debate enabled by space-based photometry concerns the activity level of the Sun. Early Kepler results hinted that other Sun-like stars appear to be noisier than the Sun ([Gilliland et al. 2011, 2015](#)), which provided the argument for an extended Kepler mission (since more data was required to detect Earth-like transiting planets). Other studies concluded that the Sun does not appear to be more active than other Sun-like stars ([Basri et al. 2010, 2013](#)). The precise stellar parameters from Gaia renewed interest in this topic, again concluding that stars that are nearly identical to the Sun are more variable ([Reinhold et al. 2020b](#), [Zhang et al. 2020](#)). This, however, may also be attributed to selecting stars with cooler temperatures and higher metallicities ([Metcalf & van Saders 2020](#), [Reinhold et al. 2020a](#)). Updated rotation period and activity measurements conclude that the Sun appears to be consistent with other Sun-like stars ([Mathur et al. 2023](#), [Santos et al. 2023](#)). Possible explanations consistent with current literature include that the Sun is transitioning into a lower activity state ([Metcalf et al. 2016](#)), that the Sun and other Sun-like stars occasionally experience epochs of high activity ([Reinhold et al. 2020b](#)), or that the Kepler sample does not include a sufficient number of stars that are truly Sun-like ([Herbst et al. 2025](#)).

Gyrochronology:

Method to determine stellar ages using the spin-down of stars with convective envelopes over time.

Stellar Flares:

Sudden brightenings resulting from magnetic reconnection on stellar surfaces; probe magnetic activity and habitability environments.

2.4. Flares and Outbursts

Stellar flares are sudden releases of energy on the surfaces of stars that occur on timescales of minutes, typically characterized by a steep flux increase followed by exponential decay. Understanding the frequency, duration, and energy distributions of flares for stars across the H-R diagram has wide-ranging implications in astrophysics, including the magnetic activity evolution of stars, the habitability of exoplanets, and high-energy space weather events on our own Sun that can affect the Earth.

The power of space-based time-domain data for the study of flares was realized as soon as the first quarter of 1-minute cadence Kepler data, which yielded clear detections in hundreds of cool M and K stars (Walkowicz et al. 2011). Subsequent studies initially focused on M dwarfs, finding that more active stars flare more often, but also demonstrating a large range of flare frequency and energy that blurred the distinction between active and inactive stars through classical spectroscopic activity indicators (Hawley et al. 2014). Larger scale catalogs enabled detections of hundreds of thousands of flares in thousands of stars, demonstrating that flare activity decreases for low-mass field stars as their rotation rates decrease, and revealing a strong correlation with Rossby number, in line with expectations that stellar activity is coupled to flare rates (Davenport 2016). Surprisingly, these studies also showed that while a single power law is unable to explain flare energies, the power law index is constant across spectral types, indicating a common physical mechanism (Davenport 2016, Davenport et al. 2019, Van Doorselaere et al. 2017, Yang & Liu 2019). Kepler also enabled the first detection of quasi-periodic pulsations in white light flares (Pugh et al. 2016).

Kepler light curves also enabled several surprising discoveries on flare properties of stars. An example is superflares on G-type stars (Maehara et al. 2012), which release flare energies an order of magnitude higher than those typically observed in the Sun (Figure 6, left). Follow-up studies demonstrated that superflares are correlated with large starspots, and surprisingly can occur for stars with rotation periods indicating a similar age to the Sun (Shibayama et al. 2013, Notsu et al. 2019). Another surprising discovery was that flares appear to be more wide-spread across the H-R diagram than previously thought, including A-type stars (Balona 2015) and giant stars (Yang & Liu 2019, Ong et al. 2024). The observed flare properties of A stars appear to indicate a different mechanism than for stars with convective envelopes (Yang & Liu 2019).

The K2 Mission extended the success from Kepler by observing stellar flares across a wider range of spectral types. In particular, K2 provided first insights into the flare rates of ultracool dwarfs (Gizis et al. 2017). K2 also opened up the study of flare rates in clusters with a wider range of ages than those available with Kepler (Ilin et al. 2019, 2021), largely confirming that flare rates show a strong dependence on age, and that flare rates decline faster with age for higher mass stars. The community-driven target selection by K2 also enabled simultaneous multiwavelength campaigns, combining space-based optical observations with ground-based spectroscopy and space-based near-UV observations with Swift (Paudel et al. 2024). These studies have laid important ground-work to understand the high-energy environment of low-mass stars, in particular ultracool dwarfs, which have become prime targets for detecting exoplanets in the habitable zones of stars.

The most recent milestone in flare studies has been enabled by the TESS Mission through two innovations: the unprecedented number of all-sky 2-minute cadence observations, and the introduction of 20-second cadence observations. The former enabled large-scale, all-sky catalogs of stellar flares, allowing an order of magnitude increase in stars with detected flares over Kepler and K2 (Günther et al. 2020, Feinstein et al. 2020). These stud-




Figure 6

Space-based time-domain photometry has revolutionized our understanding of stellar flares. Left, reproduced from [Maehara et al. \(2015\)](#): Kepler light curve of a main-sequence G-type star showing a superflare. The bottom panel shows a zoom-in on the flare. Right, reproduced from [Feinstein et al. \(2022\)](#): Flare rates across the H-R diagram derived from TESS light curves.

ies solidified that the flare frequency distributions in stars appear to be universal across a large range of stellar masses, and that photometric variations with flare-like morphologies in observations appear to be ubiquitous across the H-R diagram (Figure 6, right). At least part of these detections are likely unrelated to classical flares and may instead be related to pulsational phenomena such as stochastic outbursts, which were first revealed through Kepler/K2 observations of white dwarfs ([Bell et al. 2015](#), [Hermes et al. 2015](#)).

TESS 20-second cadence data provided spectacular new insights into the detailed morphology of flares, which was smeared out in most previous observations, and demonstrating that quasi-periodic pulsations are common ([Gilbert et al. 2022](#), [Howard 2022](#)). The detailed morphology of flares in M dwarfs revealed by TESS has led to connections with observations in the Sun, linking the observed multimodal variability to plasma condensation in the corona ([Yang et al. 2023](#)). Fast cadence data from TESS has also been used to study the effect of star-planet interactions on stellar flares ([Ilin & Poppenhaeger 2022](#)).

The long baselines of space-based time-domain have enabled the exploration of using flare variability as a tracer for stellar activity cycles ([Scoggins et al. 2019](#)). Evidence for such variability has been detected in TESS data spanning over five years ([Feinstein et al. 2024](#)) and when combining observations of the same star from Kepler and TESS ([Wainer et al. 2024](#)). As the TESS time baselines extend even further, flares could provide a powerful probe to measure long-term activity cycles of stars across the H-R diagram.




Figure 7

The remarkable variability of young stars revealed through space-based time-domain photometry. From [Cody et al. \(2022\)](#): each panels shows a 70-day light curve from K2 of disk-bearing young stars in the Taurus star-forming region. Abbreviations in square brackets denote the morphological classification scheme from [Cody et al. \(2014\)](#).

Young Stars: Stars in early formation stages (ages $\lesssim 10\text{-}100$ Myr), showing variability from accretion, circumstellar obscuration, or disks.

2.5. Dippers, Bursters, and Exotic Variables

A common theme in the space-based time-domain revolution is that astrophysical processes which previously appeared “simple” were revealed to have extraordinary complexity. One of the best examples is the variability in young stars (broadly defined as stars with ages $\lesssim 100$ Myr, with material from the forming disk typically disappearing within $\lesssim 10$ Myr) as well as the discovery of exotic variability in mature stars.

Time variability observations of young stellar objects have a long history ([Joy 1945](#)). Early space-based results from CoRoT and Spitzer of the open cluster NGC 2264 revealed a rich complexity of variability, broadly attributed to spots, obscuration events from circumstellar material that is left over from the formation, and accretion fluctuations ([Cody et al. 2014](#)). The rich data quality motivated a classification scheme, separating symmetric variability, predominant decreases in brightness (the “dippers”), and predominant increases in brightness (the “bursters”). Combined with the periodic nature of the signal, these observations motivated a morphological classification scheme using periodicity, stochasticity, and symmetry to separate the light curves.

Kepler provided few insights into the variability of young stars due to its target selection focusing on mature solar-type stars. A recent exception was the Gaia-based discovery of a

young moving group that was serendipitously observed by Kepler (Bouma et al. 2022). What Kepler lacked was recovered by K2, which, thanks to its pointing strategy in the ecliptic plane, covered numerous young associations with high-precision space-based photometry for the first time. K2 revealed spectacular variability in young star-forming regions, including upper Scorpius and Ophiuchus (Ansdell et al. 2016, Cody & Hillenbrand 2018, Hedges et al. 2018), the Lagoon Nebula Cluster (Venuti et al. 2021), and Taurus (Cody et al. 2022, Roggero et al. 2021). These observations revealed that younger populations are more dominated by phenomena attributed to accretion (e.g. bursts), while older populations show a large fraction of dippers consistent with more evolved discs, implying that accretion timescales are shorter than disc dispersal timescales (Cody et al. 2022).

Light curves also probe geometric effects such as disc inclinations. Surprisingly, some dipper stars were found to have face-on orientations, and samples with isotropically orientated outer discs (resolved with sub-mm observations by ALMA) show that the dipper phenomenon is unrelated to the outer disc geometry (Ansdell et al. 2020). This implies that either dipper events may not be predominantly caused by dust in the inner disc, or that inner and outer disc misalignments may be common. Cody et al. (2022) found that inclination correlations exist only for periodic and quasiperiodic sources, while aperiodic sources appear largely isotropic, suggesting that different subclasses of dipper stars may be attributed to astrophysical phenomena. Overall, while young star variability is shaped by age and geometry, a consistent physical picture is still elusive and likely related to the complex interplay of accretion, magnetic fields, and disc evolution. New observations by TESS have already started to add valuable data to this topic (Tajiri et al. 2020), including the surprising discovery of dipper stars substantially older than the canonical ≈ 10 Myr for which disks should have dispersed (Gaidos et al. 2022).

Space-based time-domain data has also revealed new classes of exotic variable stars. An example is the “complex periodic variables”, which show nearly periodic brightness decreases that are superimposed on rotational variability due to spots (Stauffer et al. 2017). The phenomenon occurs mostly on young pre-main sequence M dwarfs without evidence of disks, and at times can undergo “state-changes” during a cycle. Combined data from K2 and TESS are now favoring a picture in which this variability is caused by gas or dust that is co-rotating with the star (Bouma et al. 2024).

One of the most remarkable aspects of the space-based time-domain photometry revolution is the discovery of exotic variability in mature, solar-type stars. The most famous example is Boyajian’s star, a seemingly normal F-type main-sequence star demonstrating large, irregular brightness variations (Boyajian et al. 2016). Other examples include the “random transitter” HD 139139, a star exhibiting seemingly random transits without periodicities during the K2 observing window (Rappaport et al. 2019) that may have disappeared at later epochs based on observations with CHEOPS (Alonso et al. 2023). Consensus explanations for the exotic variability remain elusive, and the continuation of space-photometry revolution is set to uncover more of these exotic results.

2.6. Binary Stars

Binary stars are benchmark systems in astrophysics for determining fundamental properties of stars with no or little model dependence (e.g. Torres et al. 2010). The traditional prototype are eclipsing binary stars, which for the special case of double-lined spectroscopic (SB2) binaries allow dynamical measurements of radii and masses of the binary components.




Figure 8

Space-based time-domain missions have uncovered tens of thousands of binaries with orbital periods spanning over five orders of magnitude. Histogram of orbital periods of binaries discovered by Kepler and TESS, separated into contact/eclipsing binaries and binaries detected using pulsation timing. Data taken from [Kirk et al. \(2016\)](#), [Murphy et al. \(2018\)](#), [Green et al. \(2023\)](#), [Prša et al. \(2022\)](#), and [Kostov et al. \(2025\)](#).

Contact binaries are stars with close orbital periods for which the binary components are tidally deformed, yielding near sinusoidal light curve shapes. For binaries on wider orbits, astrometric and interferometric monitoring of binary orbits yields similar constraints. Binary stars with pulsating components provide unique laboratories for testing stellar physics (see Southworth & Bowman, this volume).

Eclipsing and contact binaries are accessible through ground-based photometry, and time-domain surveys such as OGLE, ZTF, ASAS-SN, and ATLAS have yielded large catalogs of such binary star systems ([Soszyński et al. 2017](#), [Heinze et al. 2018](#), [Chen et al. 2020](#), [Rowan et al. 2023](#)). The high precision and continuous coverage of space-based time-domain missions revealed physical effects that were previously not detected. Prominent examples include the first detection of Doppler beaming, a periodic modulation of the photon emission rate as the stars move towards the observer ([Bloemen et al. 2011](#)), the discovery of “heartbeat” stars, binaries on eccentric orbits which cause dynamically induced tidal distortion and pulsations ([Thompson et al. 2012](#)), and the discovery of tidally-tilted pulsators ([Handler et al. 2020](#)). The last two studies are examples of the new field of “tidal asteroseismology” that was sparked by the precision of space-based time-domain photometry ([Burkart et al. 2012](#), [Guo 2020](#)).

Another breakthrough was the discovery of high-order eclipsing systems. The first

Heartbeat Stars:

Eccentric binaries showing tidally induced brightness variations and pulsations near periastron.

examples from Kepler included hierarchical triply eclipsing triple systems with a main-sequence primary star (Carter et al. 2011) and an evolved red giant primary star (Derekas et al. 2011), the latter also showing tidally forced variability from the orbit of the lower mass binary. Eclipse timing variations using Kepler data revealed evidence for hundreds of triple systems, providing statistical constraints on the distributions of orbital parameters in higher order binaries (Borkovits et al. 2016). The large number of stars monitored by TESS has enabled the discovery of even higher order systems, including over a hundred eclipsing quadruple systems (Kostov et al. 2024), the first known sextuple system with three eclipsing binaries (Powell et al. 2021), and a triply eclipsing sextuple star system around a previously known eclipsing binary star (Zasche et al. 2023). Space-based missions have even yielded discoveries of new eclipsing binaries around naked-eye stars that have previously been missed by ground-based observations (Bedding et al. 2019, Hey et al. 2022).

A legacy product of the space-based time-domain missions is large catalogs of binaries with measured orbital parameters. The nominal Kepler mission discovered over 3000 eclipsing binaries (Prša et al. 2011, Kirk et al. 2016), followed by several hundred systems so far discovered in the K2 dataset (Armstrong et al. 2015, Maxted & Hutcheon 2018). A novel method to detect binaries discovered with Kepler was the timing of coherent pulsations, allowing measurements equivalent to radial velocities through frequency (Kurtz et al. 2015) and phase (Murphy et al. 2014) modulations. The signal strength of pulsation timing increases with orbital period (Hey et al. 2020), thus providing a complementary method to eclipsing binaries. Finally, the TESS mission has so far delivered over 10,000 contact binary candidates on close orbits (Green et al. 2023) and thousands of detached eclipsing binaries (Prša et al. 2022, IJspeert et al. 2024, Kostov et al. 2025). Overall, the space-based time-domain missions have so far discovered tens of thousands of new binaries with orbital periods spanning nearly five orders of magnitude (Figure 8).

Space-based telescopes also advanced our understanding of cataclysmic variables (CVs). CVs consist of a white dwarf primary which is accreting material from a donor star, leading to both irregular and periodic brightness variations. Kepler data allowed the discovery of new CVs (Howell et al. 2013), including examples showing eclipses (Scaringi et al. 2013). The continuous coverage also enabled the confirmation that periods that are slightly different than the orbital periods (so-called “superhumps”) can be attributed to hot spots in a tilted, precessing disk surrounding the white dwarf (Gies et al. 2013) and led to novel insights in the related phenomenon of superoutbursts (Osaki & Kato 2013). K2 and TESS increased the sample of CVs with space-based light curves to a few dozen, revealing hitherto unknown variability (Bruch 2022). CVs are an excellent example of stellar astrophysics where the precision and continuity of space-based light curves led to new insights even for variable stars with large amplitudes that have been studied from the ground for decades.

3. Galactic Astronomy

3.1. Galactic Archaeology

The study of the formation and evolution of the Milky Way has emerged as a key topic in astrophysics over the past decades (see Helmi 2020, for a review). Historically, the study of stellar populations has been driven by large-scale photometric and astrometric datasets such as the Geneva-Copenhagen survey (Nordström et al. 2004) and Hipparcos (van Leeuwen 2007). Over the past decades, multiplexed spectroscopic surveys have added high-precision chemical abundances, which are associated with the birth environment (“chemical tagging”,

see [Freeman & Bland-Hawthorn 2002](#)), while the Gaia mission provided exquisite astrometry for billions of stars ([Gaia Collaboration et al. 2023](#)).

The major contribution of space-based time-domain missions was to measure ages of stars on sufficiently large scales to provide a chronological order of galactic stellar populations and events - a subfield now commonly referred to as “galactic archaeology”. Since more luminous stars probe larger distances, the primary tool to provide this information is asteroseismology of red giants. The mass, metallicity, and internal physics of a red giant star directly define its age, and thus the combination of space-based light curves and spectroscopic ground-based data thus provides the possibility to measure ages for a significant number of stars on galactic scales.

The potential of asteroseismology for galactic archaeology was first realized by [Miglio et al. \(2009\)](#), who observed qualitative agreement between population synthesis models and asteroseismic detections in 800 stars observed by CoRoT. Later work analyzing multiple CoRoT pointings revealed significant differences in mass distributions, most likely due to different vertical scale heights being probed ([Miglio et al. 2013](#)). This was also observed in a sample of Kepler dwarfs and subgiant stars ([Chaplin et al. 2011b](#)). The large Kepler red giant sample provided strong potential and challenges for advancing galactic archaeology, the latter being predominantly due to the poorly understood selection function of red giants ([Sharma et al. 2016](#)). Building upon this lesson, the K2 galactic archaeology program ([Stello et al. 2017](#)) was the first large-scale coordinated effort to use space-based light curves as probes of galactic structure ([Zinn et al. 2020](#)). Finally, the full-frame images from TESS eliminated the need for target selection, yielding a near all-sky view for nearby oscillating red giants free of selection biases ([Mackereth et al. 2021](#), [Hon et al. 2021](#)).

In parallel, spectroscopic surveys specifically targeted stars with space-based detections of oscillations to provide complementary chemical abundances, which are critical to infer ages from asteroseismic detections. This includes APOGEE observations of the CoRoT ([Anders et al. 2017](#)), Kepler ([Pinsonneault et al. 2014](#), [Serenelli et al. 2017](#)), and K2 ([Schonhut-Stasik et al. 2024](#)) fields, AAT/HERMES observations of K2 and TESS stars ([Sharma et al. 2018, 2019](#)). LAMOST observations of predominantly the Kepler/K2 fields ([Zong et al. 2020](#)), and observations of the recently commissioned 4MOST telescope ([Bensby et al. 2019](#)).

Ages provided by space-based photometry yielded expected results and surprising discoveries. [Silva Aguirre et al. \(2018\)](#) used Kepler data to show that stars enriched in α -process elements relative to iron are on average ≈ 8 Gyr older than stars that are α poor, consistent with galactic chemical evolution models and distinct “thin” and “thick” disk populations in the Milky Way. [Chiappini et al. \(2015\)](#) and [Martig et al. \(2015\)](#) identified an unexpected population of young α -rich stars, which may be the result binary interactions which strip off mass from the primary star, thus yielding a lower mass and (seemingly) older age ([Grisoni et al. 2024](#)). Combining ages with radial metallicity gradients and kinematics support a picture in which stars in the galaxy undergo radial migration ([Anders et al. 2017](#), [Sharma et al. 2021](#)) and that the galaxy evolved “inside-out” and may have experienced a late burst of star formation ([Warfield et al. 2024](#)), confirming earlier results from photometric and spectroscopic surveys ([Casagrande et al. 2011](#), [Bensby et al. 2014](#), [Bergemann et al. 2014](#)).

Space-based photometry also provided first age estimates for non-disk components of our galaxy. Specifically, determining the age of a star that formed outside our galaxy puts a lower limit on the time that the star joined our Milky Way (e.g. a merger event). Kepler




Figure 9

Space-based photometry has provided a new tool to study old stellar populations in our galaxy. Top down view of an HST image of the Milky Way analog UGC 12158. Colored symbols show stars for which space-based light curves from different missions have yielded asteroseismic detections and for which spectroscopic follow-up data exists. Image credit: Friedrich Anders, based on work from [Anders et al. \(2023\)](#).

and TESS data showed that Gaia-Enceladus, a major merger of the Milky Way that is now commonly associated with the “birth” of the kinematic thick disk, occurred at least ≈ 8 years ago ([Chaplin et al. 2020](#), [Montalbán et al. 2021](#), [Grunblatt et al. 2021](#)). Furthermore, the dwarf galaxy merger that led to the Helmi stream occurred at least ≈ 12 Gyr ago ([Lindsay et al. 2025](#)).

TESS observations also detected oscillations in “halo” stars ([Borre et al. 2022](#)), the oldest component of our galaxy whose origin is still uncertain. A challenge is that reliable ages for metal-poor stars require modeling of individual frequencies, which has only been done for a small number of very-metal poor stars ([Huber et al. 2024](#), [Larsen et al. 2025](#)) and becomes challenging for more luminous giants. The oldest ages of stars in the Milky Way may be used to constrain cosmological models and tensions between local and distant measures of the Hubble constant ([Moresco 2024](#)). Improvements to stellar model physics

will be required to assess the accuracy of such estimates. Achieving this goal is a major motivation for the planned HAYDN Mission (Miglio et al. 2021, see Section 7).

A side product of the space-based revolution of galactic archaeology has been constraints on interstellar dust. This is enabled by the combination of high-precision spectroscopic parameters and asteroseismic surface gravities, which pin down the spectral energy distribution and allow extinction to be fitted as a free parameter when modeling broadband photometry. Application of this method to Kepler data (Rodrigues et al. 2014) yielded strong systematic differences to traditional 3D dust maps. Future extensions of spectroscopic surveys with surface gravities calibrated by asteroseismology (e.g. Godoy-Rivera et al. 2021) will provide the potential to constrain large-scale empirical dust maps.

3.2. Neutron Stars and Black Holes

The Milky Way is predicted to be home to millions of solar-mass black holes, several thousand of which are expected to reside in non-interacting binary systems (e.g. Shao & Li 2019). Determining the demographics of such compact “dark” objects in binaries is critical for our understanding of massive star evolution and the progenitors of gravitational wave events. While several hundred black holes have been detected as X-ray sources in interacting binaries, the search for detached systems has significantly increased thanks to large scale surveys. Currently only a handful of such black holes are known in the field, all of which have been identified and confirmed through the combination of Gaia astrometry and radial velocities (Chakrabarti et al. 2023, El-Badry et al. 2023a,b).

Neutron stars and black holes can be detected photometrically either through ellipsoidal variations (for short orbital periods), gravitational lensing as the compact object moves in front of the luminous companion (self-lensing binaries), or pulsation timing. The first method has yielded several candidate black hole detections from ground-based photometric surveys such as ASAS-SN ((Thompson et al. 2019, Jayasinghe et al. 2021, 2022). However, many of these could not be independently confirmed and are often compatible with non-single star evolution and binary mass transfer scenarios (e.g. El-Badry et al. 2022).

Space-based photometry holds strong promise to increase the known population of detached neutron stars and black holes. Kepler enabled the first discovery of a self-lensing binary star including a white dwarf and a main-sequence star (Kruse & Agol 2014), with additional systems discovered later on (Kawahara et al. 2018, Masuda et al. 2019). First systematic searches using all-sky data from TESS have started to constrain the occurrence rate of white dwarfs around Sun-like stars to $\approx 1\%$ by looking for self-lensing events (Yamaguchi et al. 2024), and constrained the occurrence of short-period black holes around Sun-like stars to less than 10^{-5} through a systematic search for ellipsoidal variability (Green et al. 2025). Future searches, using extended TESS light curves, may yield the first photometric detection of nearby neutron stars and black holes in detached binaries.

Space-based data can also provide important complementary data to characterize neutron stars and black holes, for example by vetting candidates identified from spectroscopy (Jayasinghe et al. 2023) and providing additional data to model ellipsoidal variations for candidates (Jayasinghe et al. 2022, 2021). TESS light curves have also enabled asteroseismology of red-giant stars with black holes detected using Gaia astrometry, providing precise characterization of the primary stars which “host” the dark companions (Hey et al. 2025).

4. Extragalactic Science

4.1. Explosive Transients

Transient phenomena produce variability over limited time spans and, in general, do not show repeatable behavior. The study of explosive transients such as supernovae has wide-ranging implications in astrophysics, from constraining the expansion history of our universe by using Type Ia supernovae as standard candles to understanding the chemical enrichment of galaxies through core-collapse (Type II) supernovae. Recent reviews of explosive transients can be found in [Burrows & Vartanyan \(2021\)](#) and [Ruiter & Seitzzahl \(2025\)](#).

The detection of supernovae has motivated ground-based imaging surveys such as the Palomar Transient Factory (PTF, [Law et al. 2009](#)), the Zwicky Transient Facility (ZTF, [Bellm et al. 2019](#)), the All-Sky Automated Survey for Supernovae (ASAS-SN, [Shappee et al. 2014](#)), and the recently commissioned Rubin Observatory Legacy Survey of Space and Time ([Ivezic et al. 2019](#)). Time-domain surveys to detect near-Earth asteroids and other rapid phenomena such as PanSTARRS ([Chambers et al. 2016](#)) and ATLAS ([Tonry et al. 2018](#)) also detect supernovae on a regular basis. All these surveys provide photometric coverage of large areas (and in the case of ASAS-SN and ATLAS the entire sky) every few nights on average. Space-based time-domain missions such as Kepler/K2 and TESS provide complementary data to these surveys with faster cadence and better photometric precision, at the cost of a more limited survey area.

The first detection of Type Ia supernovae with Kepler was enabled through a Guest Observer program targeting hundreds of galaxies in the Kepler field ([Olling et al. 2015](#)). The resulting light curves showed exquisite precision compared to available ground-based data. The light curves revealed no evidence for the interaction of supernova ejecta with a stellar companion, solidifying the now established theory that type Ia supernova progenitors rarely contain non-compact objects and predominantly result from white dwarf binaries. In contrast, Kepler results on two core-collapse supernovae resulted in evidence for shock waves that interact with pre-existing material from a red supergiant progenitor ([Garnavich et al. 2016](#)). Both findings were made possible by the early detection of the events thanks to continuous time-domain monitoring from space.

The success by Kepler led to systematic programs to search for explosive transients using the K2 Mission, which provided more extended sky coverage along the ecliptic. [Shappee et al. \(2019\)](#) and [Dimitriadis et al. \(2019\)](#) used the early-time data from K2 to discover two components in the rise morphology of a type Ia supernova (Figure 10), indicating that two distinct physical processes are at play during the explosion. Other Type Ia supernovae observed with similar precision were found to be consistent with a single power law rise ([Wang et al. 2021](#)), highlighting the complexity and diversity in “normal” supernovae. K2 data also yielded the discovery of fast-evolving, luminous transients that may be the result of an explosion hitting material left behind from a pre-explosion mass-loss episode ([Rest et al. 2018](#)). Finally, K2 allowed constraints on the progenitor of Type IIb supernova through high-cadence observations of the shock cooling light curve ([Armstrong et al. 2021](#)).

The large field of view of the TESS Mission (covering $\approx 5\%$ of the entire sky at a given pointing) further strengthened the synergy between supernova science and space-based time-domain missions. TESS data was used to characterize unusual Type Ia supernovae with emission features that may originate from circumstellar material ([Vallely et al. 2019](#)) and confirm the flux excess in early-rise light curves of Type Ia explosions, with data defying current models ([Wang et al. 2024](#)). TESS has allowed the first population study of Type

Explosive Transients:

Rapidly evolving events such as supernovae or kilonovae, probing progenitors and explosion physics.




Figure 10

Space-based time-domain missions have provided exquisite new insights into the optical behaviour of extragalactic sources. Left, reproduced from [Shappee et al. \(2019\)](#): K2 light curve of a Type Ia supernova with a double-peaked rise component. Right, reproduced from [Burke et al. \(2020\)](#): TESS light curve showing the stochastic variability caused by accretion processes onto a supermassive black hole in the center of a dwarf galaxy. Colors show a damped random walk (DRW) and Gaussian Process (CARMA) model.

Ia supernova characteristics using early-time light curves, hinting at two distinct types of Type Ia supernova populations based on the broad distribution of power law slopes ([Fausnaugh et al. 2021, 2023](#)). TESS also obtained early light curves of several core-collapse supernovae ([Vallely et al. 2019](#)), enabling more constraints on progenitors through shock-cooling ([Wang et al. 2023](#)) and the detections of optical flashes and afterglows of gamma-ray bursts associated with core-collapse supernovae ([Jayaraman et al. 2024, Roxburgh et al. 2024](#)). Finally, TESS is predicted to be able to provide optical counterpart measurements of kilonovae, transients caused by mergers of neutron stars and/or black holes that are sources of gravitational wave events ([Mo et al. 2023](#)). TESS is now the most prolific optical space-based time-domain telescope for studying explosive transients, leading to it sometimes being referred to as the “Transient Explorer Survey Satellite”.

Active Galactic Nuclei (AGN):

Galaxies with accreting supermassive black holes producing emission across a broad range of the electromagnetic spectrum.

4.2. Active Galactic Nuclei

The centers of most galaxies are home to supermassive black holes. Accretion of material onto these black holes produces variability across the electromagnetic spectrum, ranging from radio emission from electrons accelerated along magnetic field lines, X-rays due to hot gas close to the black hole and optical/IR radiation in the accretion disk. Galaxies with central black holes masses $\gtrsim 10^5 M_\odot$ show luminosities that dominate the galaxy spectrum and are generally referred to as “active galactic nuclei” (AGN, see [Netzer 2015](#), for a review).

Time-domain variability of AGN can link the emission sources to physical processes in the accretion disk. The first space-based optical studies of AGN using Kepler data showed a wide variety of variability previously unseen in ground-based data ([Mushotzky et al. 2011, Edelson et al. 2014, Revalski et al. 2014, Smith et al. 2018](#)). The power spectrum slopes of an AGN sample showed correlations with bolometric luminosity and yielded the first detection of optical quasi-periodic oscillations, which are commonly detected in X-ray observations of stellar-mass black hole binaries, supporting the idea that accretion behaves similarly across a large range of masses ([Smith et al. 2018](#)). The characteristic frequency for the optical variability was ≈ 40 days and provided hints at a relationship between the variability timescale and black hole mass. Differences to the power slopes found in X-

rays indicated that the optical variability is not just due to reprocessing of X-ray photons. Kepler also detected optical variability of a blazar, a subgroup of AGN with powerful radio emission due to jets that oriented towards the observer (Edelson et al. 2013).

K2 observations expanded the study of optical variability to dozens of blazars. The results showed variability that is consistent with turbulence in the relativistic jet and accretion disk (Wehrle et al. 2019, Carini et al. 2020), and hinted at differences between two subclasses of blazars based on their radio spectrum variability (Wehrle et al. 2023). K2 also provided the first opportunity for repeated observations of blazars using space-based photometry, showing significant differences in power spectrum slopes over time (Wehrle et al. 2023).

TESS enabled the combination of optical space-based photometry with simultaneous observations at other wavelengths. Weaver et al. (2020) studied multi-wavelength variability of prototype blazar BL Lac, finding remarkable similarity in variability across optical, UV, and X-ray wavelengths. The data revealed frequency-dependent time lags relative to TESS, which can be explained by a model involving energization of the radiating electrons at a shock front and provided first constraints on the strength of the involved magnetic field. Unlike earlier K2 results, TESS data showed no difference in optical variability properties of blazar subclasses, despite expectations that one class (Flat Spectrum Radio Quasars, FSRQs) has a substantial optical contribution from the accretion disk. Characteristic optical timescales were found to range from ≈ 0.8 -8 days, consistent with optical variability originating in the jet (Dingler & Smith 2024).

TESS has also enabled the study of AGN in lower-mass galaxies. The characteristic variability timescale of a dwarf AGN was found to be significantly shorter than the corresponding timescales for higher-mass objects, providing strong evidence that the variability timescales of accretion disks correlate with black hole mass (Figure 10, Burke et al. 2020). A consequence of this result is that TESS can also be used to identify AGN from light curve variability alone: Treiber et al. (2023) identified dozens of new AGN candidates based on TESS data and demonstrated that traditional selection criteria based on spectroscopy would fail for $\approx 40\%$ of the identified cases.

4.3. Tidal Disruption Events

When stars orbit too close to a supermassive black hole, tidal forces can exceed self-gravity, leading to a tidal disruption of the star. Remnants of the star can form an accretion disk around the black hole, leading to a luminous transient event. Such tidal disruption events (TDEs) provide insights into both black hole physics and stellar evolution, as well as identify dormant black holes that are otherwise undetectable (for a recent review see Gezari 2021).

Space-based time-domain photometry has provided exquisite new insights into the early time variability of TDEs that are discovered with ground-based surveys. Holoien et al. (2019) presented the first TESS light curve of a TDE, allowing constraints on the time of first light and peak brightness, as well as providing constraints on the expansion velocities. TESS provided important constraints on the first periodic TDE, which is believed to be the result of a partial disruption event (Payne et al. 2021). The exquisite precision of TESS revealed subtle changes from one flare event to the other, which may be linked due to systematic differences in outbursts in each orbit (Payne et al. 2023). Finally, TESS has also been used to shine light on the nature of ambiguous nuclear transients by providing early-time (pre-transient) light curves that can distinguish between TDEs and AGN by searching for stochastic variability that is more consistent with AGN (Hinkle et al. 2023).

Tidal Disruption

Events: Stellar disruption by a supermassive black hole, producing (at times periodic) flares and variability.

Small bodies:

Objects orbiting the Sun that are not planets, dwarf planets, or natural satellites (moons). These include comets, asteroids, and interstellar objects.

5. Solar System Science

5.1. Small Bodies

The study of small solar-system bodies such as asteroids and comets informs our understanding of the formation of our solar system, such as the delivery of water to inner terrestrial planets and the motion of giant planets (e.g. [Morbidelli et al. 2000](#)). Space-based time-domain photometry contributes to our understanding of small body populations by detecting light variations from rotationally modulated reflected sunlight. Rotation period and amplitude distributions give insight into the compositions of small bodies because for a given composition, asteroids reach a critical maximum rotation period beyond which they disintegrate. The composition in turn yields clues on their origin in the solar system.

The necessity for the K2 Mission to point along the ecliptic plane was fortuitous for solar system science. Early studies used asteroid data serendipitously collected within large superstamps that were downloaded to study clusters, gravitational microlensing events, and solar system planets ([Szabó et al. 2016](#), [Podlowska-Gaca et al. 2021](#)). K2 measured rotation rates of hundreds of main-belt asteroids, finding significantly longer rotation periods than ground-based surveys. This was interpreted as evidence that period distributions from such surveys are biased to short periods ([Molnár et al. 2018](#)), since ground-based observations are hampered by day/night cycles and weather.

K2 also obtained dedicated observations of Jovian Trojan asteroids. Early results constrained the binary fraction of Jovian Trojans to $\approx 20\%$, and a rotation period distribution with a spin barrier that implies a composition consistent with comets, favoring an outer solar system origin ([Szabó et al. 2017](#), [Ryan et al. 2017](#)). Later studies increased the sample size to over 100 Trojans ([Kalup et al. 2021](#)), revealing a bimodality in their rotation distribution. Similar results of bimodal rotation periods were also found for Centaurs, a population of small bodies with orbits between Jupiter and Neptune ([Marton et al. 2020](#)).

Reaching into the outer solar system, K2’s sensitivity allowed studies of some distant trans-Neptunian objects (TNOs). The analysis of 2007 OR₁, the second most distant known TNO, yielded a significantly larger size and slower rotation rate than previously thought, implying a larger surface gravity and thus more likely retention of volatiles ([Pál et al. 2016](#)). [Kecskeméthy et al. \(2023\)](#) analyzed a sample of 30 TNOs with K2 and found slower rotation and higher amplitudes for large TNOs compared to main belt asteroids. A possible explanation is that the compositions of TNOs allow the retention of irregular shapes at larger sizes than their inner solar system counterparts.

TESS has continued the success story by measuring rotation periods and amplitudes for thousands of main-belt asteroids, showing that long rotation periods were underestimated by previous ground-based surveys by up to an order of magnitude ([Pál et al. 2020](#), Figure 11). TESS also made discoveries that some small bodies rotate faster than expected: this includes fast-spinning medium-sized Hilda asteroids ([Takács et al. 2025](#)) and fast-rotating Jovian trojans ([Kiss et al. 2025](#)). These results imply either higher density compositions than previously assumed, collisions, or that the small body populations are more highly integrated with each other than previously thought.

A new class of solar system small bodies are interstellar objects, which are characterized by hyperbolic orbits. Three interstellar objects are currently known, discovered either by the ground-based surveys PanSTARRS ([Chambers et al. 2016](#), [Meech et al. 2017](#)) and ATLAS ([Tonry et al. 2018](#), [Seligman et al. 2025](#)) or amateur astronomers. TESS data was used to obtain pre-discovery imaging of the third interstellar object, I/ATLAS, providing




Figure 11

K2 and TESS data have provided new insights into the origin of small bodies and giant planets in the solar system. Left, reproduced from Pál et al. (2020): Distribution of rotation periods of main-belt asteroids from previous surveys (red) and TESS (black), demonstrating that many long-period asteroids have previously been missed. Right, reproduced from Simon et al. (2016): K2 observations of Neptune, showing both the full continuous light curve (top) and 5 day segments with rapidly evolving variability due to clouds (bottom).

constraints on its composition (Feinstein et al. 2025, Martinez-Palomera et al. 2025).

5.2. Dwarf Planets and Planet Nine

K2 Campaign 7 included observations of the Pluto and Charon system, providing a continuous high-precision light curve of the dwarf planet in our solar system (Benecchi et al. 2018). Pluto and Charon were unresolved within a K2 pixel, and a dedicated mask was used to obtain observations along the K2 detectors. The light curve showed clear variability due to the rotation of Pluto and Charon, covering 13 rotation cycles during the campaign. The data showed a smaller amplitude than predicted, consistent with a model in which Pluto continues to move material on its surface through evaporation/re-deposition processes. The variability was also found to be stable over a 3 month baseline, consistent with the small heliocentric distance change during the timespan of the observations.

Thanks to its all-sky coverage, TESS data has been suggested to be an effective tool for discovering distant solar system objects, including the hypothesized Planet Nine. By shifting and stacking individual FFI exposures, studies using a few sectors of TESS data have recovered TNOs down to $V \approx 22$ and discovered several new candidates (Rice & Laughlin 2020). This implies sensitivity to an object like the predicted Planet Nine to a distance of about 900 au (Holman et al. 2019, Payne et al. 2019).

5.3. Ice Giant Planets

The ecliptic pointing of the K2 Mission campaigns allowed uninterrupted observations of Neptune and Uranus. A key motivation was to detect intrinsic oscillations, which would provide insights into the interior structure of giant planets (Vorontsov et al. 1976), similar to the constraints on Saturn's interior from ring seismology (Mankovich & Fuller 2021). K2

obtained fast-cadence (1-min) observations of Neptune over 49 continuous days (Rowe et al. 2017). No oscillations were detected, but the data provided an upper limit of ≈ 5 ppm. The data did show remarkable variability on a range of timescales due to clouds in Neptune’s atmosphere (Simon et al. 2016, Figure 11), including consistent variability due to a single discrete storm and smaller or fainter clouds that contribute to short-term brightness variability. The observations provide context for spatially unresolved observations of substellar objects with cloud variability, such as directly imaged free-floating brown dwarfs.

A remarkable discovery from K2’s Neptune observations was the detection of solar oscillations in reflected light (Gaulme et al. 2016). This detection has important implications since solar reference values for global asteroseismic parameters are frequently used to infer fundamental properties of stars. The frequency of maximum power is expected to depend on wavelength (Howe et al. 2020), but until that point had not been measured in Kepler’s bandpass. The analysis of K2 data showed a 5% higher value than typical adopted values from other solar observations, potentially indicating a bias which would propagate to mass and radius estimates for stars other than the Sun. The difference may also be consistent with the stochastic nature of the excitation of solar-like oscillations.

6. Citizen Science

Space-based time-domain missions have helped usher in an era in which non-professional astronomers – citizen scientists – have made significant contributions to our understanding of astrophysics. This is predominantly enabled by two developments: the availability of well-documented, open-source software tools which lower the barrier of interacting with data products, and dedicated efforts by the research community to facilitate the analysis of data by citizen scientists. An example of the former is the `lightkurve` software tool (Lightkurve Collaboration et al. 2018), initially developed by the Kepler/K2 science office and now maintained by a growing number of community members. `lightkurve` has been cited over 1000 times in less than 10 years, illustrating its high impact within the community. Many other open-source software tools have been generated to help facilitate interactions with space-based time-domain data, ranging from specific applications (e.g. transit fitting, asteroseismology) to generating light curves from TESS full-frame images. A non-exhaustive list can be found on the TESS Science Support Center webpage (<https://heasarc.gsfc.nasa.gov/docs/tess/data-analysis-tools.html>).

A successful example to engage citizen scientists in the analysis of space-based time-domain data is the Planet Hunters project, launched in 2010 to analyze Kepler data. Planet Hunters used the Zooniverse platform to allow citizen scientists to visually inspect light curves via a web interface. Due to the large amount of light curves produced by Kepler, traditional automated approaches inevitably miss some interesting signal. Citizen scientists discovered new planet candidates (Fischer et al. 2012), including planets in interesting parameter spaces such as a transiting circumbinary planet in a quadruple star system (Schwamb et al. 2013). The most well-known example of a citizen science discovery from Kepler is Boyajian’s star (Boyajian et al. 2016, see Section 4), a non-standard signal that has since sparked new software tools to automatically look for similar signal of arbitrary shape (Wheeler & Kipping 2019). Citizen scientists also contributed several discoveries with data from the K2 Mission, including K2-138, a system of five sub-Neptune-sized planets in a near-resonant chain (Christiansen et al. 2018) and K2-288Bb, a super-Earth-sized planet possibly orbiting the secondary star in a binary system (Feinstein et al. 2019).

Citizen Science:

Participation of non-professionals in research, enabled by open data policies, open-source software, and community platforms.

The success of citizen science continued with the TESS Mission, including Planet Hunters TESS (Eisner et al. 2020) and Planet Patrol (Kostov et al. 2022). Both projects resulted in large numbers of classifications and used algorithms to combine or refine individual classifications (Tardugno Poleo et al. 2024). Notable TESS discoveries include a planet around an evolved subgiant star in a long-period orbit and a multi-planet system orbiting a bright G-type star (Eisner et al. 2021). The examples described above have sparked the community-organized Visual Survey Group, which aims to discover exoplanets using Kepler/K2 and TESS (Kristiansen et al. 2022). The success of online platforms to engage citizen scientists has also expanded beyond exoplanet science: the Zooniverse project “Cosmic Cataclysms” aims to vet transient events which are identified using automated machine-learning methods in TESS full-frame image data (Roxburgh et al. 2025).

The discovery of transiting planets around bright stars by TESS has also enabled data contributions from amateur astronomers for validating planet candidates. Ground-based, seeing-limited timeseries photometry of stars near TESS planet candidates are now routinely used to rule out eclipsing binaries that may be blended with the target star given TESS’s large (21”) pixels. These observations, which are coordinated through the TESS Follow-up Program (Collins 2019), are largely provided by the amateur astronomer community and have been invaluable to prioritize radial velocity follow-up resources. The TESS Follow-Up Program (TFOP) also provided a unique collaboration platform for professional astronomers, including data sharing via the Exoplanet Follow-up Observing Program (ExoFOP) website (Christiansen et al. 2025). TFOP has enabled mass measurements of hundreds of transiting exoplanets discovered by TESS, many of which are now targets for atmospheric studies with the James Webb Space Telescope.

7. The Future of Space Based Time-Domain Photometry

Several space-based time-domain telescopes are funded to launch before the end of this decade. ESA’s PLATO Mission, with a planned launch in late 2026, will obtain fast-cadence, high-precision photometry for approximately one million stars to detect transiting exoplanets and measure asteroseismic ages of the host stars (Rauer et al. 2025). PLATO’s field of view will be larger than Kepler’s, targeting brighter stars that are more amenable to asteroseismology and radial velocity follow-up. The planned L2 orbit will allow much longer continuous observing campaigns than TESS, thus opening up the discovery space for long-period planets. The current strategy for PLATO is to observe a field in the southern hemisphere for 2 years (Nascimbeni et al. 2025), with a possible extension of 4 years, or longer with an extended mission. PLATO’s observing strategy is thus intermediate to Kepler (narrow and deep) and TESS (shallow and wide).

The Roman Space Telescope, scheduled for launch in late 2026, is NASA’s next flagship mission (Spergel et al. 2015). While formally a multi-purpose observatory, Roman’s large field of view (one hundred times larger than Hubble) will provide powerful time-domain opportunities. The Galactic Bulge Time-Domain Survey, one of three core-community surveys, will obtain 420 days of high-cadence (12 min) photometry in the H-band over a 2 square degree field of view over 5 years along the galactic bulge and in the galactic center (Observations Time Allocation Committee & Community Survey Definition Committees 2025). The primary science goal is the detection of exoplanets using gravitational microlensing (Penny et al. 2019), but the data will enable a wide range of ancillary science, including asteroseismology of hundreds of thousands of red giants (Huber et al. 2023, Weiss

et al. 2025), the detection of tens of thousands of transiting exoplanet candidates (Wilson et al. 2023), the detection of stellar-mass black holes (Lam et al. 2023), and probing the variability of stellar populations in the galactic center (Terry et al. 2023). Additional time-domain surveys may be conducted as guest observer programs. Roman will provide the first high-precision space-based light curves of a large number of stars at infrared wavelengths.

The Earth 2.0 (ET) Mission, with a launch targeted for 2028, aims to detect Earth-analog planets using transits and to probe the demographics of planets on wide orbits using microlensing (Ge et al. 2022). The field of view covered by the transit telescopes will include the Kepler field. ET’s nominal mission duration is 4 years, thus providing a total of 8 years of continuous baseline for Kepler stars, and over a 20-year total baseline. The microlensing telescope will provide light curves in the galactic bulge, and use simultaneous ground-based observations to obtain masses of planets by measuring microlensing parallaxes.

Space telescopes that are not dedicated to time-domain photometry also provide powerful datasets. The ESA Gaia mission, in addition to revolutionizing our understanding of the kinematics of stars in our galaxy, has detected over ten million variable stars based on sparsely sampled ≈ 50 photometric epochs through DR3 (Eyer et al. 2023, Hey & Aerts 2024). The release of Gaia DR4, which includes light curves, is planned for the end of 2026. The James Webb Space Telescope (JWST) currently obtains light curves with the highest photometric precision per unit time ever obtained in the optical/IR as part of its goal to probe exoplanet atmospheres, with datasets showing evidence for stellar variability due to granulation on Sun-like stars (Splinter et al. 2025, Coulombe et al. 2025) or flares on low-mass stars (Howard et al. 2023). Finally, missions dedicated to the targeted follow-up study of exoplanet transits and atmospheres also enable to probe general stellar variability. This includes CHEOPS (Benz et al. 2021), which launched in 2019, as well as the future planned missions Pandora (Quintana et al. 2024), Twinkle (Stotesbury et al. 2022) and Ariel (Tinetti et al. 2018).

Important frontiers in time-domain astrophysics remain to be explored with future missions that are not yet funded. The HAYDN mission concept (Miglio et al. 2021) aims to perform dedicated optical high-precision continuous photometry of stars in crowded fields, such as galactic globular clusters and the galactic bulge, which were inaccessible to previous missions such as Kepler/K2 and TESS due to their large pixel size. Unlike Roman, HAYDN would provide dedicated long-term monitoring in crowded fields, enabling the most precise characterization of stars using asteroseismology and addressing open problems in stellar models that will affect nearly all fields in astrophysics. The Early Evolution Explorer mission (EVE, MacGregor et al. 2025) aims to fill the gap of obtaining multicolor space-based photometry to study the exoplanets, activity, and accretion in stars in young star-forming regions. Multicolor photometry will also provide strong benefits for a variety of other science cases, including mode identification in asteroseismology.

Dedicated space-based time-domain telescopes have had an astounding scientific impact in astronomy over the last two decades. Kepler/K2 and TESS data are currently contributing to publications at a rate comparable to NASA’s active Great Observatories such as Hubble and Chandra, and exceed the productivity of all major flagship missions when normalized by cost to launch (Figure 12). This remarkable impact is likely due to the breadth of science enabled by time-domain observations and the open data policy adopted by Kepler/K2 and TESS, which allows scientists from around the world to make scientific discoveries without proprietary data periods. This success highlights that open time-domain missions should remain a cornerstone for the future of space-based astronomy.




Figure 12

Space-based time-domain telescopes with open data policies show remarkable scientific productivity. Left: Number of papers per year using data from flagship missions (HST, Chandra, Gaia, JWST) and dedicated time-domain missions (Kepler/K2 and TESS) over the last 25 years. Numbers were estimated through a NASA/ADS query for mission names appearing in abstracts. An offset of -200 papers/year, estimated from pre-2000 publications, was applied for Kepler to account for abstracts unrelated to the spacecraft. Kepler/K2 is shown in lighter shading after 2018, when the mission stopped taking data. Right: Same as left but normalized by the cost to launch in millions of dollars. The following costs were assumed: HST=1.5B, JWST=10B, Chandra=2B, Gaia=1B, Kepler=600M, TESS=300M.

SUMMARY POINTS

1. Space-based time-domain photometry has transformed astrophysics by providing uninterrupted, high-precision light curves of millions of objects, with impacts across nearly all areas of astronomy.
2. Major advances in stellar astrophysics include the detection of solar-like oscillations in tens of thousands of stars with convective envelopes, the discovery of unexpected behavior in the spin-down of stars, a tight correlation between granulation and oscillations, the realization that the Sun may be an outlier in its activity level compared to other stars, the detection of a wide variety of flares and outbursts across different types of stars, and the discovery of exotic high-order binary systems.
3. Ages measured for stellar populations in the Milky Way have confirmed the basic picture of galactic chemical evolution and radial migration, and provided a chronology for merger events with satellite galaxies. Space-based light curves are also starting to constrain the demographics of “dark” galactic populations such as solar-mass black holes and neutron stars.
4. Extragalactic observations have yielded evidence for new physical processes in supernova explosions from early-time light curves, revealed that the correlation and typical variability timescale for active galaxies extends to low masses, and uncovered variations in the properties of repeating tidal disruption events.
5. Observations in the solar system demonstrated that rotation period distributions (and thus compositions) of asteroid populations are more diverse than previously thought, revealed complex cloud variability in ice giant planets, and yielded new candidates for trans-Neptunian objects in the far reaches of our solar system.

6. Open data policies and open-source software have enabled citizen scientists and amateur astronomers to play key roles in discoveries, ranging from new exoplanets to unusual stellar variability. They also contributed to the remarkable scientific impact of space-based time-domain missions, especially when normalized by mission cost to launch.

DISCLOSURE STATEMENT

The author is not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

ACKNOWLEDGMENTS

This review is dedicated to the Principal Investigators, Architects and Engineers of space-based time-domain missions, who paved the way for numerous discoveries and careers in astrophysics. I am grateful to Conny Aerts for the encouragement to write this review, and for comments on the manuscript. I am also grateful to Tim Bedding, Maria Berge-mann, Susan Benecchi, Ann Marie Cody, Christina Hedges, Adina Feinstein, Veselin Kostov, Savita Mathur, Armin Rest, Krista Lynne Smith and Josh Winn for helpful comments and suggestions. Funding support is acknowledged from the Alfred P. Sloan Foundation, the National Aeronautics and Space Administration (80NSSC19K0597, 80NSSC21K0652, 80NSSC24K0621, 80NSSC22K0781) and the Australian Research Council (FT200100871). The Kepler and TESS data used in this paper were obtained from the Mikulski Archive for Space Telescopes (MAST) at the Space Telescope Science Institute (STScI) and are freely available via the MAST archive: <https://archive.stsci.edu/>. MAST is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555. Support to MAST for these data is provided by the NASA Office of Space Science via grants NAG5-7584 and NNX09AF08G, and by other grants and contracts. Funding for the Kepler mission was provided by NASA’s Science Mission Directorate, and provided by the NASA Explorer Program for the TESS mission.

LITERATURE CITED

- Aerts C. 2021. *Reviews of Modern Physics* 93:015001ADS Bibcode: 2021RvMP...93a5001A
- Aerts C, Tkachenko A. 2024. *Astronomy and Astrophysics* 692:R1ADS Bibcode: 2024A&A...692R...1A
- Alonso R, Hoyer S, Deleuil M, Simon AE, Beck M, et al. 2023. *Astronomy and Astrophysics* 680:A78ADS Bibcode: 2023A&A...680A..78A
- Anders F, Chiappini C, Rodrigues TS, Miglio A, Montalbán J, et al. 2017. *Astronomy and Astrophysics* 597:A30ADS Bibcode: 2017A&A...597A..30A
- Anders F, Gispert P, Ratcliffe B, Chiappini C, Minchev I, et al. 2023. *Astronomy and Astrophysics* 678:A158ADS Bibcode: 2023A&A...678A.158A
- Andsell M, Gaidos E, Hedges C, Tazzari M, Kraus AL, et al. 2020. *Monthly Notices of the Royal Astronomical Society* 492:572–588Publisher: OUP ADS Bibcode: 2020MNRAS.492..572A
- Andsell M, Gaidos E, Williams JP, Kennedy G, Wyatt MC, et al. 2016. *Monthly Notices of the Royal Astronomical Society* 462:L101–L105Publisher: OUP ADS Bibcode: 2016MNRAS.462L.101A

- Appourchaux T, Gough DO, Catala C, Frandsen S, Frohlich C, et al. 1993. *Prisma - the First Space Mission to See Inside the Stars*. vol. 42. ADS Bibcode: 1993ASPC...42..411A
- Appourchaux T, Michel E, Auvergne M, Baglin A, Toutain T, et al. 2008. *Astronomy & Astrophysics* 488(2):705–714
- Armstrong DJ, Kirk J, Lam KWF, McCormac J, Walker SR, et al. 2015. *Astronomy and Astrophysics* 579:A19ADS Bibcode: 2015A&A...579A..19A
- Armstrong P, Tucker BE, Rest A, Ridden-Harper R, Zenati Y, et al. 2021. *Monthly Notices of the Royal Astronomical Society* 507:3125–3138ADS Bibcode: 2021MNRAS.507.3125A
- Baade D, Rivinius T, Pigulski A, Carciofi AC, Martayan C, et al. 2016. *Astronomy and Astrophysics* 588:A56Publisher: EDP ADS Bibcode: 2016A&A...588A..56B
- Baglin A, Auvergne M, Boisdard L, Lam-Trong T, Barge P, et al. 2006. *CoRoT: a high precision photometer for stellar evolution and exoplanet finding*. In *36th COSPAR Scientific Assembly*, vol. 36
- Balona LA. 2015. *Monthly Notices of the Royal Astronomical Society* 447:2714–2725Publisher: OUP ADS Bibcode: 2015MNRAS.447.2714B
- Balona LA. 2017. *Monthly Notices of the Royal Astronomical Society* 467:1830–1837Publisher: OUP ADS Bibcode: 2017MNRAS.467.1830B
- Barban C, Matthews JM, De Ridder J, Baudin F, Kuschnig R, et al. 2007. *Astronomy & Astrophysics* 468(3):1033–1038
- Barnes SA. 2007. *The Astrophysical Journal* 669:1167–1189Publisher: IOP ADS Bibcode: 2007ApJ...669.1167B
- Basri G, Walkowicz LM, Batalha N, Gilliland RL, Jenkins J, et al. 2010. *The Astrophysical Journal* 713(2):L155–L159
- Basri G, Walkowicz LM, Reiners A. 2013. *The Astrophysical Journal* 769:37Publisher: IOP ADS Bibcode: 2013ApJ...769...37B
- Bastien FA, Stassun KG, Basri G, Pepper J. 2013. *Nature* 500:427–430ADS Bibcode: 2013Natur.500..427B
- Bastien FA, Stassun KG, Basri G, Pepper J. 2016. *The Astrophysical Journal* 818(1):43
- Batalha NM, Borucki WJ, Koch DG, Bryson ST, Haas M, et al. 2010. *The Astrophysical Journal* 713(2):L109–L114
- Bedding K. 1995. Amplitude of stellar oscillations: the implications for Asteroseismology
- Bedding TR, Butler RP, Kjeldsen H, Baldry IK, O’Toole SJ, et al. 2001. *The Astrophysical Journal* 549:L105–L108Publisher: IOP ADS Bibcode: 2001ApJ...549L.105B
- Bedding TR, Hey DR, Murphy SJ. 2019. *Research Notes of the American Astronomical Society* 3:163Publisher: IOP ADS Bibcode: 2019RNAAS...3..163B
- Bedding TR, Kjeldsen H. 2003. *Publications of the Astronomical Society of Australia* 20:203–212ADS Bibcode: 2003PASA...20..203B
- Bedding TR, Murphy SJ, Hey DR, Huber D, Li T, et al. 2020. *Nature* 581(7807):147–151
- Bell KJ, Hermes JJ, Bischoff-Kim A, Moorhead S, Montgomery MH, et al. 2015. *The Astrophysical Journal* 809:14ADS Bibcode: 2015ApJ...809...14B
- Bellm EC, Kulkarni SR, Graham MJ, Dekany R, Smith RM, et al. 2019. *Publications of the Astronomical Society of the Pacific* 131:018002Publisher: IOP ADS Bibcode: 2019PASP..131a8002B
- Benecchi SD, Lisse CM, Ryan EL, Binzel RP, Schwamb ME, et al. 2018. *Icarus* 314:265–273ADS Bibcode: 2018Icar..314..265B
- Bensby T, Bergemann M, Rybizki J, Lemasle B, Howes L, et al. 2019. *The Messenger* 175:35–38ADS Bibcode: 2019Msngr.175...35B
- Bensby T, Feltzing S, Oey MS. 2014. *Astronomy and Astrophysics* 562:A71Publisher: EDP ADS Bibcode: 2014A&A...562A..71B
- Benz W, Broeg C, Fortier A, Rando N, Beck T, et al. 2021. *Experimental Astronomy* 51:109–151ADS Bibcode: 2021ExA....51..109B
- Bergemann M, Ruchti GR, Serenelli A, Feltzing S, Alves-Brito A, et al. 2014. *Astronomy and*

- Astrophysics* 565:A89Publisher: EDP ADS Bibcode: 2014A&A...565A..89B
- Bloemen S, Marsh TR, Østensen RH, Charpinet S, Fontaine G, et al. 2011. *Monthly Notices of the Royal Astronomical Society* 410:1787–1796Publisher: OUP ADS Bibcode: 2011MNRAS.410.1787B
- Borkovits T, Hajdu T, Sztakovics J, Rappaport S, Levine A, et al. 2016. *Monthly Notices of the Royal Astronomical Society* 455:4136–4165Publisher: OUP ADS Bibcode: 2016MNRAS.455.4136B
- Borre CC, Aguirre Børsen-Koch V, Helmi A, Koppelman HH, Nielsen MB, et al. 2022. *Monthly Notices of the Royal Astronomical Society* 514:2527–2544ADS Bibcode: 2022MNRAS.514.2527B
- Borucki WJ, Koch D, Basri G, Batalha N, Brown T, et al. 2010. *Science* 327:977–
- Bouchy F, Carrier F. 2001. *Astronomy and Astrophysics* 374:L5–L8Publisher: EDP ADS Bibcode: 2001A&A...374L...5B
- Bouma LG, Jayaraman R, Rappaport S, Rebull LM, Hillenbrand LA, et al. 2024. *The Astronomical Journal* 167:38Publisher: IOP ADS Bibcode: 2024AJ....167...38B
- Bouma LG, Kerr R, Curtis JL, Isaacson H, Hillenbrand LA, et al. 2022. *The Astronomical Journal* 164:215Publisher: IOP ADS Bibcode: 2022AJ....164..215B
- Bowman DM, Bugnet L. 2024. Asteroseismology. Publication Title: arXiv e-prints ADS Bibcode: 2024arXiv241001715B
- Bowman DM, Burssens S, Pedersen MG, Johnston C, Aerts C, et al. 2019. *Nature Astronomy* 3(8):760–765ArXiv: 1905.02120
- Boyajian TS, LaCourse DM, Rappaport SA, Fabrycky D, Fischer DA, et al. 2016. *Monthly Notices of the Royal Astronomical Society* 457:3988–4004Publisher: OUP ADS Bibcode: 2016MNRAS.457.3988B
- Brown TM, Gilliland RL, Noyes RW, Ramsey LW. 1991. *The Astrophysical Journal* 368:599Publisher: IOP ADS Bibcode: 1991ApJ...368..599B
- Bruch A. 2022. *Monthly Notices of the Royal Astronomical Society* 514:4718–4735Publisher: OUP ADS Bibcode: 2022MNRAS.514.4718B
- Brun AS, Browning MK. 2017. *Living Reviews in Solar Physics* 14:4ADS Bibcode: 2017LRSP...14....4B
- Bruntt H, Kjeldsen H, Buzasi DL, Bedding TR. 2005. *The Astrophysical Journal* 633(1):440–446
- Bugnet L, García RA, Davies GR, Mathur S, Corsaro E, et al. 2018. *Astronomy and Astrophysics* 620:A38ADS Bibcode: 2018A&A...620A..38B
- Burkart J, Quataert E, Arras P, Weinberg NN. 2012. *Monthly Notices of the Royal Astronomical Society* 421:983–1006Publisher: OUP ADS Bibcode: 2012MNRAS.421..983B
- Burke CJ, Shen Y, Chen YC, Scaringi S, Faucher-Giguere CA, et al. 2020. *The Astrophysical Journal* 899:136Publisher: IOP ADS Bibcode: 2020ApJ...899..136B
- Burrows A, Vartanyan D. 2021. *Nature* 589:29–39ADS Bibcode: 2021Natur.589...29B
- Butler RP, Bedding TR, Kjeldsen H, McCarthy C, O'Toole SJ, et al. 2004. *The Astrophysical Journal* 600:L75–L78Publisher: IOP ADS Bibcode: 2004ApJ...600L..75B
- Carini M, Wehrle AE, Wiita PJ, Ward Z, Pendleton K. 2020. *The Astrophysical Journal* 903:134Publisher: IOP ADS Bibcode: 2020ApJ...903..134C
- Carrier F, Bouchy F, Kienle F, Bedding TR, Kjeldsen H, et al. 2001. *Astronomy and Astrophysics* 378:142–145Publisher: EDP ADS Bibcode: 2001A&A...378..142C
- Carrier F, Bourban G. 2003. *Astronomy and Astrophysics* 406:L23–L26Publisher: EDP ADS Bibcode: 2003A&A...406L..23C
- Carter JA, Fabrycky DC, Ragozzine D, Holman MJ, Quinn SN, et al. 2011. *Science* 331(6017):562–565
- Casagrande L, Schönrich R, Asplund M, Cassisi S, Ramírez I, et al. 2011. *Astronomy and Astrophysics* 530:A138Publisher: EDP ADS Bibcode: 2011A&A...530A.138C
- Ceillier T, Tayar J, Mathur S, Salabert D, García RA, et al. 2017. *Astronomy and Astrophysics* 605:A111Publisher: EDP ADS Bibcode: 2017A&A...605A.111C

- Chakrabarti S, Simon JD, Craig PA, Reggiani H, Brandt TD, et al. 2023. *The Astronomical Journal* 166:6ADS Bibcode: 2023AJ....166....6C
- Chambers KC, Magnier EA, Metcalfe N, Flewelling HA, Huber ME, et al. 2016. The Pan-STARRS1 Surveys. ADS Bibcode: 2016arXiv161205560C
- Chaplin WJ, Basu S, Huber D, Serenelli A, Casagrande L, et al. 2014a. *The Astrophysical Journal Supplement Series* 210:1Publisher: IOP ADS Bibcode: 2014ApJS..210....1C
- Chaplin WJ, Bedding TR, Bonanno A, Broomhall AM, García RA, et al. 2011a. *The Astrophysical Journal* 732(1):L5
- Chaplin WJ, Elsworth Y, Davies GR, Campante TL, Handberg R, et al. 2014b. *Monthly Notices of the Royal Astronomical Society* 445:946–954Publisher: OUP ADS Bibcode: 2014MNRAS.445..946C
- Chaplin WJ, Kjeldsen H, Christensen-Dalsgaard J, Basu S, Miglio A, et al. 2011b. *Science* 332(6026):213–216
- Chaplin WJ, Miglio A. 2013. *Annual Review of Astronomy and Astrophysics* 51(1):353–392ArXiv: 1303.1957
- Chaplin WJ, Serenelli AM, Miglio A, Morel T, Mackereth JT, et al. 2020. *Nature Astronomy* 4(4):382–389ArXiv:2001.04653 [astro-ph]
- Charbonneau P. 2010. *Living Reviews in Solar Physics* 7:3ADS Bibcode: 2010LRSP....7....3C
- Chen X, Wang S, Deng L, de Grijs R, Yang M, Tian H. 2020. *The Astrophysical Journal Supplement Series* 249:18ADS Bibcode: 2020ApJS..249...18C
- Chiappini C, Anders F, Rodrigues TS, Miglio A, Montalbán J, et al. 2015. *Astronomy and Astrophysics* 576:L12ADS Bibcode: 2015A&A...576L..12C
- Christiansen JL, Crossfield IJM, Barentsen G, Lintott CJ, Barclay T, et al. 2018. *The Astronomical Journal* 155:57ADS Bibcode: 2018AJ....155...57C
- Christiansen JL, McElroy DL, Harbut M, Ciardi DR, Crane M, et al. 2025. *The Planetary Science Journal* 6:186ADS Bibcode: 2025PSJ.....6..186C
- Clark DH, Stephenson FR. 1977. *The historical supernovae*. Publication Title: Oxford [Eng.] ; New York : Pergamon Press ADS Bibcode: 1977hisu.book....C
- Claytor ZR, van Saders JL, Cao L, Pinsonneault MH, Teske J, Beaton RL. 2024. *The Astrophysical Journal* 962:47ADS Bibcode: 2024ApJ...962...47C
- Cody AM, Hillenbrand LA. 2018. *The Astronomical Journal* 156:71Publisher: IOP ADS Bibcode: 2018AJ....156...71C
- Cody AM, Hillenbrand LA, Rebull LM. 2022. *The Astronomical Journal* 163:212Publisher: IOP ADS Bibcode: 2022AJ....163..212C
- Cody AM, Stauffer J, Baglin A, Micela G, Rebull LM, et al. 2014. *The Astronomical Journal* 147:82Publisher: IOP ADS Bibcode: 2014AJ....147...82C
- Collins K. 2019. *TESS Follow-up Observing Program Working Group (TFOP WG) Sub Group 1 (SG1): Ground-based Time-series Photometry*. vol. 233. ADS Bibcode: 2019AAS...23314005C
- Coulombe LP, Radica M, Benneke B, D'Aoust E, Dang L, et al. 2025. *Nature Astronomy* 9:512–525ADS Bibcode: 2025NatAs...9..512C
- Cranmer SR, Bastien FA, Stassun KG, Saar SH. 2014. *The Astrophysical Journal* 781:124ADS Bibcode: 2014ApJ...781..124C
- Croll B, Walker GAH, Kuschnig R, Matthews JM, Rowe JF, et al. 2006. *The Astrophysical Journal* 648:607–613Publisher: IOP ADS Bibcode: 2006ApJ...648..607C
- Curtis JL, Agüeros MA, Douglas ST, Meibom S. 2019. *The Astrophysical Journal* 879:49ADS Bibcode: 2019ApJ...879...49C
- Córsico AH, Althaus LG, Miller Bertolami MM, Kepler SO. 2019. *Astronomy and Astrophysics Review* 27:7ADS Bibcode: 2019A&ARv..27....7C
- Davenport JRA. 2016. *The Astrophysical Journal* 829:23Publisher: IOP ADS Bibcode: 2016ApJ...829...23D
- Davenport JRA, Covey KR, Clarke RW, Boeck AC, Cornet J, Hawley SL. 2019. *The Astrophysical*

Journal 871:241Publisher: IOP ADS Bibcode: 2019ApJ...871..241D

De Ridder J, Barban C, Baudin F, Carrier F, Hatzes AP, et al. 2009. *Nature* 459(7245):398–400

De Ridder J, Barban C, Carrier F, Mazumdar A, Eggenberger P, et al. 2006. *Astronomy and Astrophysics* 448:689–695Publisher: EDP ADS Bibcode: 2006A&A...448..689D

Degroote P, Aerts C, Baglin A, Miglio A, Briquet M, et al. 2010. *Nature* 464:259–261ADS Bibcode: 2010Natur.464..259D

Derekas A, Kiss LL, Borkovits T, Huber D, Lehmann H, et al. 2011. *Science* 332:216ADS Bibcode: 2011Sci...332..216D

Dimitriadis G, Foley RJ, Rest A, Kasen D, Piro AL, et al. 2019. *The Astrophysical Journal* 870:L1Publisher: IOP ADS Bibcode: 2019ApJ...870L...1D

Dingler R, Smith KL. 2024. *The Astrophysical Journal* 973:10Publisher: IOP ADS Bibcode: 2024ApJ...973...10D

Douglas ST, Curtis JL, Agüeros MA, Cargile PA, Brewer JM, et al. 2019. *The Astrophysical Journal* 879:100ADS Bibcode: 2019ApJ...879..100D

Edelmann PVF, Ratnasingam RP, Pedersen MG, Bowman DM, Prat V, Rogers TM. 2019. *The Astrophysical Journal* 876:4Publisher: IOP ADS Bibcode: 2019ApJ...876...4E

Edelson R, Mushotzky R, Vaughan S, Scargle J, Gandhi P, et al. 2013. *The Astrophysical Journal* 766:16ADS Bibcode: 2013ApJ...766...16E

Edelson R, Vaughan S, Malkan M, Kelly BC, Smith KL, et al. 2014. *The Astrophysical Journal* 795:2ADS Bibcode: 2014ApJ...795....2E

Edmonds PD, Gilliland RL. 1996. *The Astrophysical Journal* 464:L157Publisher: IOP ADS Bibcode: 1996ApJ...464L.157E

Eisner NL, Barragán O, Aigrain S, Lintott C, Miller G, et al. 2020. *Monthly Notices of the Royal Astronomical Society* 494:750–763ADS Bibcode: 2020MNRAS.494..750E

Eisner NL, Nicholson BA, Barragán O, Aigrain S, Lintott C, et al. 2021. *Monthly Notices of the Royal Astronomical Society* 505:1827–1840ADS Bibcode: 2021MNRAS.505.1827E

Eitner P, Bergemann M, Hoppe R, Nordlund A, Plez B, Klevas J. 2024. *Astronomy and Astrophysics* 688:A52Publisher: EDP ADS Bibcode: 2024A&A...688A..52E

El-Badry K, Rix HW, Cendes Y, Rodriguez AC, Conroy C, et al. 2023a. *Monthly Notices of the Royal Astronomical Society* 521:4323–4348Publisher: OUP ADS Bibcode: 2023MNRAS.521.4323E

El-Badry K, Rix HW, Quataert E, Howard AW, Isaacson H, et al. 2023b. *Monthly Notices of the Royal Astronomical Society* 518:1057–1085Publisher: OUP ADS Bibcode: 2023MNRAS.518.1057E

El-Badry K, Seeburger R, Jayasinghe T, Rix HW, Almada S, et al. 2022. *Monthly Notices of the Royal Astronomical Society* 512:5620–5641ADS Bibcode: 2022MNRAS.512.5620E

Eyer L, Audard M, Holl B, Rimoldini L, Carnerero MI, et al. 2023. *Astronomy and Astrophysics* 674:A13ADS Bibcode: 2023A&A...674A..13E

Fausnaugh MM, Valley PJ, Kochanek CS, Shappee BJ, Stanek KZ, et al. 2021. *The Astrophysical Journal* 908:51Publisher: IOP ADS Bibcode: 2021ApJ...908...51F

Fausnaugh MM, Valley PJ, Tucker MA, Kochanek CS, Shappee BJ, et al. 2023. *The Astrophysical Journal* 956:108Publisher: IOP ADS Bibcode: 2023ApJ...956..108F

Favata F. 2004 538:3–11Conference Name: Stellar Structure and Habitable Planet Finding ADS Bibcode: 2004ESASP.538....3F

Feinstein AD, Montet BT, Ansdell M, Nord B, Bean JL, et al. 2020. *The Astronomical Journal* 160:219ADS Bibcode: 2020AJ....160..219F

Feinstein AD, Noonan JW, Seligman DZ. 2025. Precure Observations of 3I/ATLAS from TESS Suggests Possible Distant Activity. ADS Bibcode: 2025arXiv250721967F

Feinstein AD, Schlieder JE, Livingston JH, Ciardi DR, Howard AW, et al. 2019. *The Astronomical Journal* 157:40ADS Bibcode: 2019AJ....157...40F

Feinstein AD, Seligman DZ, France K, Gagné J, Kowalski A. 2024. *The Astronomical Journal*

- 168:60ADS Bibcode: 2024AJ....168...60F
- Feinstein AD, Seligman DZ, Günther MN, Adams FC. 2022. *The Astrophysical Journal* 925:L9Publisher: IOP ADS Bibcode: 2022ApJ...925L...9F
- Fischer DA, Schwamb ME, Schawinski K, Lintott C, Brewer J, et al. 2012. *Monthly Notices of the Royal Astronomical Society* 419:2900–2911ADS Bibcode: 2012MNRAS.419.2900F
- Frandsen S, Carrier F, Aerts C, Stello D, Maas T, et al. 2002. *Astronomy and Astrophysics* 394:L5–L8Publisher: EDP ADS Bibcode: 2002A&A...394L...5F
- Freeman K, Bland-Hawthorn J. 2002. *Annual Review of Astronomy and Astrophysics* 40:487–537ADS Bibcode: 2002ARA&A...40..487F
- Freytag B, Ludwig HG, Steffen M. 1996. *Astronomy and Astrophysics* 313:497–516ADS Bibcode: 1996A&A...313..497F
- Fridlund M, Gough DO, Jones A, Appourchaux T, Badiali M, et al. 1995. *STARS: A Proposal for a Dedicated Space Mission to Study Stellar Structure and Evolution*. vol. 76. ADS Bibcode: 1995ASPC...76..416F
- Gaia Collaboration, Vallenari A, Brown AGA, Prusti T, de Bruijne JHJ, et al. 2023. *Astronomy and Astrophysics* 674:A1Publisher: EDP ADS Bibcode: 2023A&A...674A...1G
- Gaidos E, Mann AW, Rojas-Ayala B, Feiden GA, Wood ML, et al. 2022. *Monthly Notices of the Royal Astronomical Society* 514:1386–1402Publisher: OUP ADS Bibcode: 2022MNRAS.514.1386G
- Garcia RA, Mathur S, Salabert D, Ballot J, Regulo C, et al. 2010. *Science* 329(5995):1032–1032
- García RA, Ballot J. 2019. *Living Reviews in Solar Physics* 16(1):4_eprint: 1906.12262
- Garnavich PM, Tucker BE, Rest A, Shaya EJ, Olling RP, et al. 2016. *The Astrophysical Journal* 820:23Publisher: IOP ADS Bibcode: 2016ApJ...820...23G
- Gaulme P, McKeever J, Rawls ML, Jackiewicz J, Mosser B, Guzik JA. 2013. *The Astrophysical Journal* 767:82Publisher: IOP ADS Bibcode: 2013ApJ...767...82G
- Gaulme P, Rowe JF, Bedding TR, Benomar O, Corsaro E, et al. 2016. *The Astrophysical Journal* 833:L13ADS Bibcode: 2016ApJ...833L..13G
- Ge J, Zhang H, Zang W, Deng H, Mao S, et al. 2022. ET White Paper: To Find the First Earth 2.0. ArXiv:2206.06693 [astro-ph]
- Gezari S. 2021. *Annual Review of Astronomy and Astrophysics* 59:21–58ADS Bibcode: 2021ARA&A...59...21G
- Gies DR, Guo Z, Howell SB, Still MD, Boyajian TS, et al. 2013. *The Astrophysical Journal* 775:64Publisher: IOP ADS Bibcode: 2013ApJ...775...64G
- Gilbert EA, Barclay T, Quintana EV, Walkowicz LM, Vega LD, et al. 2022. *The Astronomical Journal* 163:147Publisher: IOP ADS Bibcode: 2022AJ....163..147G
- Gilliland RL. 2008. *The Astronomical Journal* 136(2):566–579
- Gilliland RL, Chaplin WJ, Dunham EW, Argabright VS, Borucki WJ, et al. 2011. *The Astrophysical Journal Supplement Series* 197:6Publisher: IOP ADS Bibcode: 2011ApJS..197....6G
- Gilliland RL, Chaplin WJ, Jenkins JM, Ramsey LW, Smith JC. 2015. *The Astronomical Journal* 150:133Publisher: IOP ADS Bibcode: 2015AJ....150..133G
- Gizis JE, Paudel RR, Schmidt SJ, Williams PKG, Burgasser AJ. 2017. *The Astrophysical Journal* 838:22Publisher: IOP ADS Bibcode: 2017ApJ...838...22G
- Godoy-Rivera D, Tayar J, Pinsonneault MH, Rodríguez Martínez R, Stassun KG, et al. 2021. *The Astrophysical Journal* 915:19Publisher: IOP ADS Bibcode: 2021ApJ...915...19G
- Goodricke J. 1783. *Philosophical Transactions of the Royal Society of London Series I* 73:474–482ADS Bibcode: 1783RSPT...73..474G
- Gould A, Huber D, Penny M, Stello D. 2015. *Journal of Korean Astronomical Society* 48:93–104_eprint: 1410.7395
- Green MJ, Maoz D, Mazeh T, Faigler S, Shahaf S, et al. 2023. *Monthly Notices of the Royal Astronomical Society* 522:29–55Publisher: OUP ADS Bibcode: 2023MNRAS.522...29G
- Green MJ, Ziv Y, Rix HW, Maoz D, Hamoudy I, et al. 2025. *Astronomy and Astrophysics*

- 695:A210Publisher: EDP ADS Bibcode: 2025A&A...695A.210G
- Grigahcène A, Antoci V, Balona L, Catanzaro G, Daszyńska-Daszkiewicz J, et al. 2010. *The Astrophysical Journal* 713(2):L192–L197
- Grisoni V, Chiappini C, Miglio A, Brogaard K, Casali G, et al. 2024. *Astronomy and Astrophysics* 683:A111Publisher: EDP ADS Bibcode: 2024A&A...683A.111G
- Grunblatt SK, Zinn JC, Price-Whelan AM, Angus R, Saunders N, et al. 2021. *The Astrophysical Journal* 916(2):88
- Guenther DB, Kallinger T, Gruberbauer M, Huber D, Weiss WW, et al. 2008. *The Astrophysical Journal* 687(2):1448–1459
- Guo Z. 2020. Listening to the Heartbeat: Tidal Asteroseismology in Action. Conference Name: Stars and their Variability Observed from Space ADS Bibcode: 2020svos.conf..203G
- Guzik JA. 2021. *Frontiers in Astronomy and Space Sciences* 8:55ADS Bibcode: 2021FrASS...8...55G
- Günther MN, Zhan Z, Seager S, Rimmer PB, Ranjan S, et al. 2020. *The Astronomical Journal* 159:60Publisher: IOP ADS Bibcode: 2020AJ....159...60G
- Hall OJ, Davies GR, van Saders J, Nielsen MB, Lund MN, et al. 2021. *Nature Astronomy* 5:707–714ADS Bibcode: 2021NatAs...5..707H
- Handler G, Kurtz DW, Rappaport SA, Saio H, Fuller J, et al. 2020. *Nature Astronomy* 4:684–689ADS Bibcode: 2020NatAs...4..684H
- Harvey J. 1985. *High-resolution helioseismology*. In *Future Missions in Solar, Heliospheric & Space Plasma Physics*, ed. E. Rolfe & B. Battrock, vol. 235 of *ESA Special Publication*
- Hatt E, Nielsen MB, Chaplin WJ, Ball WH, Davies GR, et al. 2023. *Astronomy and Astrophysics* 669:A67ADS Bibcode: 2023A&A...669A..67H
- Hawley SL, Davenport JRA, Kowalski AF, Wisniewski JP, Hebb L, et al. 2014. *The Astrophysical Journal* 797:121Publisher: IOP ADS Bibcode: 2014ApJ...797..121H
- Hedges C, Hodgkin S, Kennedy G. 2018. *Monthly Notices of the Royal Astronomical Society* 476:2968–2998Publisher: OUP ADS Bibcode: 2018MNRAS.476.2968H
- Heinze AN, Tonry JL, Denneau L, Flewelling H, Stalder B, et al. 2018. *The Astronomical Journal* 156:241ADS Bibcode: 2018AJ....156..241H
- Hekker S, Christensen-Dalsgaard J. 2017. *Astronomy and Astrophysics Review* 25:1Publisher: Springer ADS Bibcode: 2017A&ARv..25....1H
- Hekker S, Gilliland RL, Elsworth Y, Chaplin WJ, De Ridder J, et al. 2011. *Monthly Notices of the Royal Astronomical Society* 414:2594–2601Publisher: OUP ADS Bibcode: 2011MNRAS.414.2594H
- Hekker S, Kallinger T, Baudin F, De Ridder J, Barban C, et al. 2009. *Astronomy and Astrophysics* 506:465–469Publisher: EDP ADS Bibcode: 2009A&A...506..465H
- Helmi A. 2020. *Annual Review of Astronomy and Astrophysics* 58(1):205–256ArXiv:2002.04340 [astro-ph]
- Herbst K, Amazo-Gómez EM, Papaioannou A. 2025. *Astronomy and Astrophysics* 697:A80Publisher: EDP ADS Bibcode: 2025A&A...697A..80H
- Hermes JJ, Gänsicke BT, Kawaler SD, Greiss S, Tremblay PE, et al. 2017. *The Astrophysical Journal Supplement Series* 232:23ADS Bibcode: 2017ApJS..232...23H
- Hermes JJ, Montgomery MH, Bell KJ, Chote P, Gänsicke BT, et al. 2015. *The Astrophysical Journal* 810:L5ADS Bibcode: 2015ApJ...810L...5H
- Hey D, Aerts C. 2024. *Astronomy and Astrophysics* 688:A93ADS Bibcode: 2024A&A...688A..93H
- Hey D, Li Y, Ong J. 2025. Asteroseismology of the red giant companions to Gaia BH2 and BH3. ADS Bibcode: 2025arXiv250309690H
- Hey DR, Kochoska A, Monier R, Kochukhov O, Johnston C, et al. 2022. *Monthly Notices of the Royal Astronomical Society* 511:2648–2658Publisher: OUP ADS Bibcode: 2022MNRAS.511.2648H
- Hey DR, Murphy SJ, Foreman-Mackey D, Bedding TR, Pope BJS, Hogg DW. 2020. *The Astro-*

- nomical Journal* 159:202Publisher: IOP ADS Bibcode: 2020AJ....159..202H
- Hinkle JT, Kochanek CS, Shappee BJ, Valley PJ, Auchettl K, et al. 2023. *Monthly Notices of the Royal Astronomical Society* 521:3517–3526Publisher: OUP ADS Bibcode: 2023MNRAS.521.3517H
- Holman MJ, Payne MJ, Pál A. 2019. *Research Notes of the American Astronomical Society* 3:160ADS Bibcode: 2019RNAAS...3..160H
- Holoien TWS, Valley PJ, Auchettl K, Stanek KZ, Kochanek CS, et al. 2019. *The Astrophysical Journal* 883:111Publisher: IOP ADS Bibcode: 2019ApJ...883..111H
- Hon M, Huber D, Kuzlewicz JS, Stello D, Sharma S, et al. 2021. *The Astrophysical Journal* 919(2):131
- Horst L, Edelmann PVF, Andr  ssy R, R  pke FK, Bowman DM, et al. 2020. *Astronomy and Astrophysics* 641:A18Publisher: EDP ADS Bibcode: 2020A&A...641A..18H
- Houdek G, Dupret MA. 2015. *Living Reviews in Solar Physics* 12:8ADS Bibcode: 2015LRSP...12...8H
- Howard WS. 2022. *Monthly Notices of the Royal Astronomical Society* 512:L60–L65Publisher: OUP ADS Bibcode: 2022MNRAS.512L..60H
- Howard WS, Kowalski AF, Flagg L, MacGregor MA, Lim O, et al. 2023. *The Astrophysical Journal* 959:64ADS Bibcode: 2023ApJ...959...64H
- Howe R, Chaplin WJ, Basu S, Ball WH, Davies GR, et al. 2020. *Monthly Notices of the Royal Astronomical Society* 493:L49–L53Publisher: OUP ADS Bibcode: 2020MNRAS.493L..49H
- Howell SB, Everett ME, Seabode SA, Szkody P, Still M, et al. 2013. *The Astronomical Journal* 145:109Publisher: IOP ADS Bibcode: 2013AJ....145..109H
- Howell SB, Sobeck C, Haas M, Still M, Barclay T, et al. 2014. *Publications of the Astronomical Society of the Pacific* 126:398Publisher: IOP ADS Bibcode: 2014PASP..126..398H
- Huber D, Bedding TR, Arentoft T, Gruberbauer M, Guenther DB, et al. 2011. *The Astrophysical Journal* 731(2):94
- Huber D, Bryson ST, Haas MR, Barclay T, Barentsen G, et al. 2016. *The Astrophysical Journal Supplement Series* 224(1):2
- Huber D, Pinsonneault M, Beck P, Bedding TR, Bland-Hawthorn J, et al. 2023. Asteroseismology with the Roman Galactic Bulge Time-Domain Survey. Publication Title: arXiv e-prints ADS Bibcode: 2023arXiv230703237H
- Huber D, Slumstrup D, Hon M, Li Y, B  rsen-Koch VA, et al. 2024. *The Astrophysical Journal* 975:19Publisher: IOP ADS Bibcode: 2024ApJ...975...19H
- Huber D, White TR, Metcalfe TS, Chontos A, Fausnaugh MM, et al. 2022. *The Astronomical Journal* 163(2):79
- IJspeert LW, Tkachenko A, Johnston C, Aerts C. 2024. *Astronomy and Astrophysics* 691:A242Publisher: EDP ADS Bibcode: 2024A&A...691A.242I
- Ilin E, Poppenhaeger K. 2022. *Monthly Notices of the Royal Astronomical Society* 513:4579–4586ADS Bibcode: 2022MNRAS.513.4579I
- Ilin E, Schmidt SJ, Davenport JRA, Strassmeier KG. 2019. *Astronomy and Astrophysics* 622:A133ADS Bibcode: 2019A&A...622A.133I
- Ilin E, Schmidt SJ, Poppenh  ger K, Davenport JRA, Kristiansen MH, Omohundro M. 2021. *Astronomy and Astrophysics* 645:A42ADS Bibcode: 2021A&A...645A..42I
- Ivezic Z, Kahn SM, Tyson JA, Abel B, Acosta E, et al. 2019. *The Astrophysical Journal* 873:111ADS Bibcode: 2019ApJ...873..111I
- Jackiewicz J. 2021. *Frontiers in Astronomy and Space Sciences* 7:102ADS Bibcode: 2021FrASS...7..102J
- Jayaraman R, Fausnaugh M, Ricker GR, Vanderspek R, Mo G. 2024. *The Astrophysical Journal* 972:162Publisher: IOP ADS Bibcode: 2024ApJ...972..162J
- Jayasinghe T, Rowan DM, Thompson TA, Kochanek CS, Stanek KZ. 2023. *Monthly Notices of the Royal Astronomical Society* 521:5927–5939Publisher: OUP ADS Bibcode: 2023MNR-

- RAS.521.5927J
- Jayasinghe T, Stanek KZ, Thompson TA, Kochanek CS, Rowan DM, et al. 2021. *Monthly Notices of the Royal Astronomical Society* 504:2577–2602Publisher: OUP ADS Bibcode: 2021MNRAS.504.2577J
- Jayasinghe T, Thompson TA, Kochanek CS, Stanek KZ, Rowan DM, et al. 2022. *Monthly Notices of the Royal Astronomical Society* 516:5945–5963Publisher: OUP ADS Bibcode: 2022MNRAS.516.5945J
- Joy AH. 1945. *The Astrophysical Journal* 102:168Publisher: IOP ADS Bibcode: 1945ApJ...102..168J
- Kallinger T, Beck PG, Hekker S, Huber D, Kuschnig R, et al. 2019. *Astronomy and Astrophysics* 624:A35Publisher: EDP ADS Bibcode: 2019A&A...624A..35K
- Kallinger T, De Ridder J, Hekker S, Mathur S, Mosser B, et al. 2014. *Astronomy & Astrophysics* 570:A41ArXiv:1408.0817 [astro-ph]
- Kallinger T, Guenther DB, Matthews JM, Weiss WW, Huber D, et al. 2008. *Astronomy and Astrophysics* 478:497–505Publisher: EDP ADS Bibcode: 2008A&A...478..497K
- Kallinger T, Hekker S, Garcia RA, Huber D, Matthews JM. 2016. *Science Advances* 2:1500654ADS Bibcode: 2016SciA....2E0654K
- Kallinger T, Matthews JM. 2010. *The Astrophysical Journal* 711:L35–L39Publisher: IOP ADS Bibcode: 2010ApJ...711L..35K
- Kalup CE, Molnár L, Kiss C, Szabó GM, Pál A, et al. 2021. *The Astrophysical Journal Supplement Series* 254:7Publisher: IOP ADS Bibcode: 2021ApJS..254....7K
- Kawahara H, Masuda K, MacLeod M, Latham DW, Bieryla A, Benomar O. 2018. *The Astronomical Journal* 155:144ADS Bibcode: 2018AJ....155..144K
- Kecskeméthy V, Kiss C, Szakáts R, Pál A, Szabó GM, et al. 2023. *The Astrophysical Journal Supplement Series* 264:18Publisher: IOP ADS Bibcode: 2023ApJS..264...18K
- Kirk B, Conroy K, Prša A, Abdul-Masih M, Kochoska A, et al. 2016. *The Astronomical Journal* 151:68Publisher: IOP ADS Bibcode: 2016AJ....151...68K
- Kiss C, Takács N, Kalup CE, Szakáts R, Molnár L, et al. 2025. *Astronomy and Astrophysics* 694:L17ADS Bibcode: 2025A&A...694L..17K
- Kjeldsen H, Bedding TR. 2011. *Astronomy and Astrophysics* 529:L8ADS Bibcode: 2011A&A...529L...8K
- Kjeldsen H, Bedding TR, Butler RP, Christensen-Dalsgaard J, Kiss LL, et al. 2005. *The Astrophysical Journal* 635:1281–1290Publisher: IOP ADS Bibcode: 2005ApJ...635.1281K
- Kjeldsen H, Bedding TR, Christensen-Dalsgaard J. 2000 203:73Conference Name: IAU Colloq. 176: The Impact of Large-Scale Surveys on Pulsating Star Research ADS Bibcode: 2000ASPC..203...73K
- Kjeldsen H, Bedding TR, Viskum M, Frandsen S. 1995. *The Astronomical Journal* 109:1313Publisher: IOP ADS Bibcode: 1995AJ....109.1313K
- Kostov VB, Kuchner MJ, Caciapuoti L, Acharya S, Ahlers JP, et al. 2022. *Publications of the Astronomical Society of the Pacific* 134:044401ADS Bibcode: 2022PASP..134d4401K
- Kostov VB, Powell BP, Fornear AU, Di Fraia MZ, Gagliano R, et al. 2025. *The Astrophysical Journal Supplement Series* 279:50ADS Bibcode: 2025ApJS..279...50K
- Kostov VB, Powell BP, Rappaport SA, Borkovits T, Gagliano R, et al. 2024. *Monthly Notices of the Royal Astronomical Society* 527:3995–4017Publisher: OUP ADS Bibcode: 2024MNRAS.527.3995K
- Kristiansen MHK, Rappaport SA, Vanderburg AM, Jacobs TL, Martin Schwengeler H, et al. 2022. *Publications of the Astronomical Society of the Pacific* 134:074401ADS Bibcode: 2022PASP..134g4401K
- Kruse E, Agol E. 2014. *Science* 344:275–277ADS Bibcode: 2014Sci...344..275K
- Kurtz DW. 2022. *Annual Review of Astronomy and Astrophysics* 60:31–71ADS Bibcode: 2022ARA&A..60...31K

- Kurtz DW, Hambleton KM, Shibahashi H, Murphy SJ, Prša A. 2015. *Monthly Notices of the Royal Astronomical Society* 446:1223–1233 Publisher: OUP ADS Bibcode: 2015MNRAS.446.1223K
- Lam CY, Abrams N, Andrews J, Bachelet E, Bahramian A, et al. 2023. Roman CCS White Paper: Characterizing the Galactic population of isolated black holes. ADS Bibcode: 2023arXiv230612514L
- Larsen JR, Rørsted JL, Aguirre Børsen-Koch V, Lundkvist MS, Christensen-Dalsgaard J, et al. 2025. *Astronomy and Astrophysics* 697:A153 ADS Bibcode: 2025A&A...697A.153L
- Law NM, Kulkarni SR, Dekany RG, Ofek EO, Quimby RM, et al. 2009. *Publications of the Astronomical Society of the Pacific* 121:1395 Publisher: IOP ADS Bibcode: 2009PASP..121.1395L
- Leavitt HS. 1908. *Annals of Harvard College Observatory* 60:87–108
- Lecoanet D, Cantiello M, Quataert E, Couston LA, Burns KJ, et al. 2019. *The Astrophysical Journal* 886:L15 ADS Bibcode: 2019ApJ...886L..15L
- Ledoux P. 1947. *The Astrophysical Journal* 105:305 ADS Bibcode: 1947ApJ...105..305L
- Li G, Van Reeth T, Bedding TR, Murphy SJ, Antoci V, et al. 2020. *Monthly Notices of the Royal Astronomical Society* 491:3586–3605 Publisher: OUP ADS Bibcode: 2020MNRAS.491.3586L
- Liagre B, García RA, Mathur S, Pinsonneault MH, Serenelli A, et al. 2025. Beyond the Nyquist frequency: Asteroseismic catalog of undersampled Kepler late subgiants and early red giants. ADS Bibcode: 2025arXiv250815654L
- Lightkurve Collaboration, Cardoso JVD, Hedges C, Gully-Santiago M, Saunders N, et al. 2018. Lightkurve: Kepler and TESS time series analysis in Python. Pages: ascl:1812.013 eprint: 1812.013
- Lindsay CJ, Hon M, Ong JMJ, García RA, Palakkatharappil DB, et al. 2025. Precise Asteroseismic Ages for the Helmi Streams. ADS Bibcode: 2025arXiv250701091L
- Lissauer JJ, Batalha NM, Borucki WJ. 2023. Exoplanet Science from Kepler. Conference Name: Protostars and Planets VII Volume: 534 ADS Bibcode: 2023ASPC..534..839L
- Lund MN, Basu S, Bieryla A, Casagrande L, Huber D, et al. 2024. *Astronomy and Astrophysics* 688:A13 ADS Bibcode: 2024A&A...688A..13L
- Lund MN, Chontos A, Grundahl F, Mathur S, García RA, et al. 2025. Luminaries in the Sky: The TESS Legacy Sample of Bright Stars. I. Asteroseismic detections in naked-eye main-sequence and sub-giant solar-like oscillators. ADS Bibcode: 2025arXiv250808699L
- Léger A, Rouan D, Schneider J, Barge P, Fridlund M, et al. 2009. *ap* 506:287–302 eprint: 0908.0241
- MacGregor M, Cody AM, Turner N, Scott V, Didion A, et al. 2025. *EVE - the Early eVolution Explorer*. vol. 245. ADS Bibcode: 2025AAS...24523409M
- Mackereth JT, Miglio A, Elsworth Y, Mosser B, Mathur S, et al. 2021. *Monthly Notices of the Royal Astronomical Society* 502:1947–1966 Publisher: OUP ADS Bibcode: 2021MNRAS.502.1947M
- Maehara H, Shibayama T, Notsu S, Notsu Y, Nagao T, et al. 2012. *Nature* 485(7399):478–481
- Maehara H, Shibayama T, Notsu Y, Notsu S, Honda S, et al. 2015. *Earth, Planets and Space* 67:59 Publisher: Springer ADS Bibcode: 2015EP&S...67...59M
- Mankovich CR, Fuller J. 2021. *Nature Astronomy* 5:1103–1109 ADS Bibcode: 2021NatAs...5.1103M
- Mann AW, Gaidos E, Lépine S, Hilton EJ. 2012. *The Astrophysical Journal* 753:90 Publisher: IOP ADS Bibcode: 2012ApJ...753...90M
- Martig M, Rix HW, Silva Aguirre V, Hekker S, Mosser B, et al. 2015. *Monthly Notices of the Royal Astronomical Society* 451:2230–2243 Publisher: OUP ADS Bibcode: 2015MNRAS.451.2230M
- Martinez-Palomera J, Tuson A, Hedges C, Dotson J, Barclay T, Powell B. 2025. Pre-discovery TESS Observations of Interstellar Object 3I/ATLAS. ADS Bibcode: 2025arXiv250802499M
- Marton G, Kiss C, Molnár L, Pál A, Farkas-Takács A, et al. 2020. *Icarus* 345:113721 Publisher: Elsevier ADS Bibcode: 2020Icar..34513721M
- Masuda K, Kawahara H, Latham DW, Bieryla A, Kunitomo M, et al. 2019. *The Astrophysical Journal* 881:L3 ADS Bibcode: 2019ApJ...881L...3M
- Mathur S, Claytor ZR, Santos ARG, García RA, Amard L, et al. 2023. *The Astrophysical Journal* 952:131 Publisher: IOP ADS Bibcode: 2023ApJ...952..131M

- Mathur S, García RA, Breton S, Santos ARG, Mosser B, et al. 2022. *Astronomy and Astrophysics* 657:A31Publisher: EDP ADS Bibcode: 2022A&A...657A..31M
- Mathur S, García RA, Huber D, Regulo C, Stello D, et al. 2016. *The Astrophysical Journal* 827:50ADS Bibcode: 2016ApJ...827...50M
- Mathur S, Hekker S, Trampedach R, Ballot J, Kallinger T, et al. 2011. *The Astrophysical Journal* 741:119Publisher: IOP ADS Bibcode: 2011ApJ...741..119M
- Matthews JM. 2007. *Communications in Asteroseismology* 150:333
- Matthews JM, Kusching R, Guenther DB, Walker GAH, Moffat AF, et al. 2004. *Nature* 430(6995):51–53
- Maxted PFL, Hutcheon RJ. 2018. *Astronomy and Astrophysics* 616:A38ADS Bibcode: 2018A&A...616A..38M
- McQuillan A, Mazeh T, Aigrain S. 2014. *The Astrophysical Journal Supplement Series* 211:24Publisher: IOP ADS Bibcode: 2014ApJS...211...24M
- Meech KJ, Weryk R, Micheli M, Kleyna JT, Hainaut OR, et al. 2017. *Nature* 552:378–381ADS Bibcode: 2017Natur.552..378M
- Metcalf TS, Egeland R, van Saders J. 2016. *The Astrophysical Journal* 826:L2Publisher: IOP ADS Bibcode: 2016ApJ...826L...2M
- Metcalf TS, van Saders J. 2020. Comment on "The Sun is less active than other solar-like stars". ADS Bibcode: 2020arXiv200704416M
- Michel E, Baglin A, Auvergne M, Catala C, Samadi R, et al. 2008. *Science* 322(5901):558–560
- Miglio A, Chiappini C, Morel T, Barbieri M, Chaplin WJ, et al. 2013. *EPJ Web of Conferences* 43:03004ArXiv: 1301.1515
- Miglio A, Girardi L, Grundahl F, Mosser B, Bastian N, et al. 2021. *Experimental Astronomy* 51:963–1001Publisher: Springer ADS Bibcode: 2021ExA....51..963M
- Miglio A, Montalbán J, Baudin F, Eggenberger P, Noels A, et al. 2009. *Astronomy & Astrophysics* 503(3):L21–L24
- Mo G, Jayaraman R, Fausnaugh M, Katsavounidis E, Ricker GR, Vanderspek R. 2023. *The Astrophysical Journal* 948:L3Publisher: IOP ADS Bibcode: 2023ApJ...948L...3M
- Molnár L, Pál A, Sárneczky K, Szabó R, Vinkó J, et al. 2018. *The Astrophysical Journal Supplement Series* 234:37Publisher: IOP ADS Bibcode: 2018ApJS...234...37M
- Montalbán J, Mackereth JT, Miglio A, Vincenzo F, Chiappini C, et al. 2021. *Nature Astronomy* 5(7):640–647ArXiv:2006.01783 [astro-ph]
- Morbidelli A, Chambers J, Lunine JI, Petit JM, Robert F, et al. 2000. *Meteoritics and Planetary Science* 35:1309–1320ADS Bibcode: 2000M&PS...35.1309M
- Moresco M. 2024. Measuring the expansion history of the Universe with cosmic chronometers. ADS Bibcode: 2024arXiv241201994M
- Murphy SJ, Bedding TR, Shibahashi H, Kurtz DW, Kjeldsen H. 2014. *Monthly Notices of the Royal Astronomical Society* 441:2515–2527Publisher: OUP ADS Bibcode: 2014MNRAS.441.2515M
- Murphy SJ, Moe M, Kurtz DW, Bedding TR, Shibahashi H, Boffin HMJ. 2018. *Monthly Notices of the Royal Astronomical Society* 474:4322–4346Publisher: OUP ADS Bibcode: 2018MNRAS.474.4322M
- Murphy SJ, Shibahashi H, Kurtz DW. 2013. *Monthly Notices of the Royal Astronomical Society* 430:2986–2998Publisher: OUP ADS Bibcode: 2013MNRAS.430.2986M
- Mushotzky RF, Edelson R, Baumgartner W, Gandhi P. 2011. *The Astrophysical Journal* 743:L12ADS Bibcode: 2011ApJ...743L..12M
- Nascimbeni V, Piotto G, Cabrera J, Montalto M, Marinoni S, et al. 2025. *Astronomy and Astrophysics* 694:A313ADS Bibcode: 2025A&A...694A..313N
- Netzer H. 2015. *Annual Review of Astronomy and Astrophysics* 53:365–408ADS Bibcode: 2015ARA&A..53..365N
- Nordström B, Mayor M, Andersen J, Holmberg J, Pont F, et al. 2004. *Astronomy & Astrophysics* 418(3):989–1019

- Notsu Y, Maehara H, Honda S, Hawley SL, Davenport JRA, et al. 2019. *The Astrophysical Journal* 876:58Publisher: IOP ADS Bibcode: 2019ApJ...876...58N
- Nutzman P, Charbonneau D. 2008. *Publications of the Astronomical Society of the Pacific* 120(865):317–327
- Observations Time Allocation Committee R, Community Survey Definition Committees C. 2025. Roman Observations Time Allocation Committee: Final Report and Recommendations. ADS Bibcode: 2025arXiv250510574O
- Olbers W. 1850. *Astronomische Nachrichten* 31:129ADS Bibcode: 1850AN.....31..129O
- Olling RP, Mushotzky R, Shaya EJ, Rest A, Garnavich PM, et al. 2015. *Nature* 521(7552):332–335
- Ong JMJ, Hon MTY, Soares-Furtado M, Stephan AP, van Saders J, et al. 2024. *The Astrophysical Journal* 966:42Publisher: IOP ADS Bibcode: 2024ApJ...966...42O
- Osaki Y, Kato T. 2013. *Publications of the Astronomical Society of Japan* 65:50Publisher: OUP ADS Bibcode: 2013PASJ...65...50O
- Pablo H, Richardson ND, Fuller J, Rowe J, Moffat AFJ, et al. 2017. *Monthly Notices of the Royal Astronomical Society* 467:2494–2503Publisher: OUP ADS Bibcode: 2017MNRAS.467.2494P
- Pan JS, Ting YS, Yu J. 2024. *Monthly Notices of the Royal Astronomical Society* 528:5890–5903Publisher: OUP ADS Bibcode: 2024MNRAS.528.5890P
- Pande D, Bedding TR, Huber D, Kjeldsen H. 2018. *Monthly Notices of the Royal Astronomical Society* 480:467–472Publisher: OUP ADS Bibcode: 2018MNRAS.480..467P
- Paudel RR, Barclay T, Youngblood A, Quintana EV, Schlieder JE, et al. 2024. *The Astrophysical Journal* 971:24Publisher: IOP ADS Bibcode: 2024ApJ...971...24P
- Payne AV, Auchtettl K, Shappee BJ, Kochanek CS, Boyd PT, et al. 2023. *The Astrophysical Journal* 951:134Publisher: IOP ADS Bibcode: 2023ApJ...951..134P
- Payne AV, Shappee BJ, Hinkle JT, Valley PJ, Kochanek CS, et al. 2021. *The Astrophysical Journal* 910:125ADS Bibcode: 2021ApJ...910..125P
- Payne MJ, Holman MJ, Pál A. 2019. *Research Notes of the American Astronomical Society* 3:172ADS Bibcode: 2019RNAAS...3..172P
- Pedersen MG, Aerts C, Pápics PI, Michielsen M, Gebruers S, et al. 2021. *Nature Astronomy* 5:715–722ADS Bibcode: 2021NatAs...5..715P
- Penny MT, Gaudi BS, Kerins E, Rattenbury NJ, Mao S, et al. 2019. *The Astrophysical Journal Supplement Series* 241:3ADS Bibcode: 2019ApJS..241....3P
- Perdomo García A, Vitas N, Khomenko E, Collados M. 2025. *Astronomy and Astrophysics* 700:A191Publisher: EDP ADS Bibcode: 2025A&A...700A.191P
- Pinsonneault MH, Elsworth Y, Epstein C, Hekker S, Mészáros S, et al. 2014. *The Astrophysical Journal Supplement Series* 215:19Publisher: IOP ADS Bibcode: 2014ApJS..215...19P
- Podlowska-Gaca E, Poleski R, Bartczak P, McDonald I, Pál A. 2021. *The Astrophysical Journal Supplement Series* 255:4Publisher: IOP ADS Bibcode: 2021ApJS..255....4P
- Powell BP, Kostov VB, Rappaport SA, Borkovits T, Zasche P, et al. 2021. *The Astronomical Journal* 161:162Publisher: IOP ADS Bibcode: 2021AJ....161..162P
- Prša A, Batalha N, Slawson RW, Doyle LR, Welsh WF, et al. 2011. *The Astronomical Journal* 141(3):83
- Prša A, Kochoska A, Conroy KE, Eisner N, Hey DR, et al. 2022. *The Astrophysical Journal Supplement Series* 258:16Publisher: IOP ADS Bibcode: 2022ApJS..258...16P
- Pugh CE, Armstrong DJ, Nakariakov VM, Broomhall AM. 2016. *Monthly Notices of the Royal Astronomical Society* 459:3659–3676Publisher: OUP ADS Bibcode: 2016MNRAS.459.3659P
- Pál A, Kiss C, Müller TG, Molnár L, Szabó R, et al. 2016. *The Astronomical Journal* 151:117Publisher: IOP ADS Bibcode: 2016AJ....151..117P
- Pál A, Szakáts R, Kiss C, Bódi A, Bognár Z, et al. 2020. *The Astrophysical Journal Supplement Series* 247:26Publisher: IOP ADS Bibcode: 2020ApJS..247...26P
- Quintana EV, Dotson JL, Colón KD, Barclay T, Supinski P, et al. 2024. *The Pandora SmallSat: multiwavelength characterization of exoplanets and their host stars*. vol. 13092. ADS Bibcode:

- 2024SPIE13092E..14Q
- Rappaport S, Vanderburg A, Kristiansen MH, Omohundro MR, Schwengeler HM, et al. 2019. *Monthly Notices of the Royal Astronomical Society* 488:2455–2465Publisher: OUP ADS Bibcode: 2019MNRAS.488.2455R
- Rauer H, Aerts C, Cabrera J, Deleuil M, Erikson A, et al. 2025. *Experimental Astronomy* 59:26Publisher: Springer ADS Bibcode: 2025ExA....59...26R
- Rebull LM, Stauffer JR, Cody AM, Hillenbrand LA, David TJ, Pinsonneault M. 2018. *The Astrophysical Journal* 155:196Publisher: IOP ADS Bibcode: 2018AJ....155..196R
- Reegen P, Kallinger T, Frast D, Gruberbauer M, Huber D, et al. 2006. *Monthly Notices of the Royal Astronomical Society* 367(4):1417–1431
- Reinhold T, Hekker S. 2020. *Astronomy and Astrophysics* 635:A43Publisher: EDP ADS Bibcode: 2020A&A...635A..43R
- Reinhold T, Shapiro AI, Solanki SK, Montet BT, Krivova NA, et al. 2020a. Reply to the comment of T. Metcalfe and J. van Saders on the Science report "The Sun is less active than other solar-like stars". ADS Bibcode: 2020arXiv200704817R
- Reinhold T, Shapiro AI, Solanki SK, Montet BT, Krivova NA, et al. 2020b. *Science* 368(6490):518–521ArXiv:2005.01401 [astro-ph]
- Rest A, Garnavich PM, Khatami D, Kasen D, Tucker BE, et al. 2018. *Nature Astronomy* 2:307–311ADS Bibcode: 2018NatAs...2..307R
- Retter A, Bedding TR, Buzasi DL, Kjeldsen H, Kiss LL. 2003. *The Astrophysical Journal* 591(2):L151–L154
- Revalski M, Nowak D, Wiita PJ, Wehrle AE, Unwin SC. 2014. *The Astrophysical Journal* 785:60ADS Bibcode: 2014ApJ...785...60R
- Rice M, Laughlin G. 2020. *The Planetary Science Journal* 1:81ADS Bibcode: 2020PSJ.....1...81R
- Ricker GR, Winn JN, Vanderspek R, Latham DW, Bakos GA, et al. 2015. *Journal of Astronomical Telescopes, Instruments, and Systems* 1:014003ADS Bibcode: 2015JATIS...1a4003R
- Rodrigues TS, Girardi L, Miglio A, Bossini D, Bovy J, et al. 2014. *Monthly Notices of the Royal Astronomical Society* 445:2758–2776Publisher: OUP ADS Bibcode: 2014MNRAS.445.2758R
- Roggero N, Bouvier J, Rebull LM, Cody AM. 2021. *Astronomy and Astrophysics* 651:A44ADS Bibcode: 2021A&A...651A..44R
- Rowan DM, Jayasinghe T, Stanek KZ, Kochanek CS, Thompson TA, et al. 2023. *Monthly Notices of the Royal Astronomical Society* 523:2641–2650ADS Bibcode: 2023MNRAS.523.2641R
- Rowe JF, Gaulme P, Lissauer JJ, Marley MS, Simon AA, et al. 2017. *The Astronomical Journal* 153:149Publisher: IOP ADS Bibcode: 2017AJ....153..149R
- Rowe JF, Matthews JM, Kuschnig R, Guenther DB, Mo AFJ, et al. 2006 :5
- Roxburgh H, Ridden-Harper R, Lane ZG, Rest A, Hubley L, et al. 2024. *The Astrophysical Journal* 963:89ADS Bibcode: 2024ApJ...963...89R
- Roxburgh H, Ridden-Harper R, Moore A, Montilla C, Leicester B, et al. 2025. TESSELLATE: Piecing Together the Variable Sky With TESS. ADS Bibcode: 2025arXiv250216905R
- Ruiter AJ, Seitzzahl IR. 2025. *Astronomy and Astrophysics Review* 33:1ADS Bibcode: 2025A&ARv..33....1R
- Ryan EL, Sharkey BNL, Woodward CE. 2017. *The Astronomical Journal* 153:116Publisher: IOP ADS Bibcode: 2017AJ....153..116R
- Santos ARG, Breton SN, Mathur S, García RA. 2021. *The Astrophysical Journal Supplement Series* 255:17Publisher: IOP ADS Bibcode: 2021ApJS..255...17S
- Santos ARG, Godoy-Rivera D, Finley AJ, Mathur S, García RA, et al. 2024. *Frontiers in Astronomy and Space Sciences* 11:1356379ADS Bibcode: 2024FrASS..1156379S
- Santos ARG, Mathur S, García RA, Broomhall AM, Egeland R, et al. 2023. *Astronomy and Astrophysics* 672:A56Publisher: EDP ADS Bibcode: 2023A&A...672A..56S
- Sayeed M, Huber D, Chontos A, Li Y. 2025. A Homogeneous Catalog of Oscillating Solar-Type Stars Observed by the Kepler Mission and a New Amplitude Scaling Relation Including Chromospheric

- Activity. ADS Bibcode: 2025arXiv250315599S
- Sayeed M, Huber D, Wheeler A, Ness MK. 2021. *The Astronomical Journal* 161(4):170
- Scaringi S, Groot PJ, Still M. 2013. *Monthly Notices of the Royal Astronomical Society* 435:L68–L72Publisher: OUP ADS Bibcode: 2013MNRAS.435L..68S
- Schonhut-Stasik J, Zinn JC, Stassun KG, Pinsonneault M, Johnson JA, et al. 2024. *The Astronomical Journal* 167:50Publisher: IOP ADS Bibcode: 2024AJ....167...50S
- Schou J, Buzasi DL. 2001. *Observations of p-modes in alpha Cen*. In *SOHO 10/GONG 2000 Workshop: Helio- and Asteroseismology at the Dawn of the Millennium*, ed. A. Wilson & P. L. Pallé, vol. 464 of *ESA Special Publication*
- Schwamb ME, Orosz JA, Carter JA, Welsh WF, Fischer DA, et al. 2013. *The Astrophysical Journal* 768:127ADS Bibcode: 2013ApJ...768..127S
- Schwarzschild M. 1958. *Structure and evolution of the stars*. Publication Title: Structure and Evolution of Stars. Martin Schwarzschild. ISBN: 9780691080444. Princeton University Press ADS Bibcode: 1958ses..book.....S
- Scoggins MT, Davenport JRA, Covey KR. 2019. *Research Notes of the American Astronomical Society* 3:137ADS Bibcode: 2019RNAAS...3..137S
- Seligman DZ, Micheli M, Farnocchia D, Denneau L, Noonan JW, et al. 2025. *The Astrophysical Journal* 989:L36Publisher: IOP ADS Bibcode: 2025ApJ...989L..36S
- Serebriakova N, Tkachenko A, Aerts C. 2024. *Astronomy and Astrophysics* 692:A245Publisher: EDP ADS Bibcode: 2024A&A...692A.245S
- Serenelli A, Johnson J, Huber D, Pinsonneault M, Ball WH, et al. 2017. *The Astrophysical Journal Supplement Series* 233(2):23
- Shao Y, Li XD. 2019. *The Astrophysical Journal* 885:151Publisher: IOP ADS Bibcode: 2019ApJ...885..151S
- Shappee BJ, Holoien TWS, Drout MR, Auchettl K, Stritzinger MD, et al. 2019. *The Astrophysical Journal* 870:13Publisher: IOP ADS Bibcode: 2019ApJ...870...13S
- Shappee BJ, Prieto JL, Grupe D, Kochanek CS, Stanek KZ, et al. 2014. *The Astrophysical Journal* 788:48Publisher: IOP ADS Bibcode: 2014ApJ...788...48S
- Sharma S, Hayden MR, Bland-Hawthorn J, Stello D, Buder S, et al. 2021. *Monthly Notices of the Royal Astronomical Society* 506:1761–1776ADS Bibcode: 2021MNRAS.506.1761S
- Sharma S, Stello D, Bland-Hawthorn J, Hayden MR, Zinn JC, et al. 2019. *Monthly Notices of the Royal Astronomical Society* 490:5335–5352Publisher: OUP ADS Bibcode: 2019MNRAS.490.5335S
- Sharma S, Stello D, Bland-Hawthorn J, Huber D, Bedding TR. 2016. *The Astrophysical Journal* 822:15ADS Bibcode: 2016ApJ...822...15S
- Sharma S, Stello D, Buder S, Kos J, Bland-Hawthorn J, et al. 2018. *Monthly Notices of the Royal Astronomical Society* 473:2004–2019ADS Bibcode: 2018MNRAS.473.2004S
- Shibayama T, Maehara H, Notsu S, Notsu Y, Nagao T, et al. 2013. *The Astrophysical Journal Supplement Series* 209:5Publisher: IOP ADS Bibcode: 2013ApJS..209....5S
- Silva Aguirre V, Bojsen-Hansen M, Slumstrup D, Casagrande L, Kawata D, et al. 2018. *Monthly Notices of the Royal Astronomical Society* 475:5487–5500ADS Bibcode: 2018MNRAS.475.5487S
- Silva Aguirre V, Lund MN, Antia HM, Ball WH, Basu S, et al. 2017. *The Astrophysical Journal* 835:173ADS Bibcode: 2017ApJ...835..173S
- Simon AA, Rowe JF, Gaulme P, Hammel HB, Casewell SL, et al. 2016. *The Astrophysical Journal* 817:162ADS Bibcode: 2016ApJ...817..162S
- Skumanich A. 1972. *The Astrophysical Journal* 171:565Publisher: IOP ADS Bibcode: 1972ApJ...171..565S
- Smith KL, Mushotzky RF, Boyd PT, Wagoner RV. 2018. *The Astrophysical Journal* 860:L10Publisher: IOP ADS Bibcode: 2018ApJ...860L..10S
- Soszyński I, Udalski A, Szymański MK, Wyrzykowski L, Ulaczyk K, et al. 2017. *Acta Astronomica* 67:297–316ADS Bibcode: 2017AcA....67..297S

- Spergel D, Gehrels N, Baltay C, Bennett D, Breckinridge J, et al. 2015. Wide-Field Infrared Survey Telescope-Astrophysics Focused Telescope Assets WFIRST-AFTA 2015 Report. ADS Bibcode: 2015arXiv150303757S
- Spergel D, Gehrels N, Breckinridge J, Donahue M, Dressler A, et al. 2013. *ArXiv e-prints* .eprint: 1305.5425
- Splinter J, Coulombe LP, Frazier RC, Cowan NB, Rauscher E, et al. 2025. *The Astronomical Journal* 170:323Publisher: IOP ADS Bibcode: 2025AJ....170..323S
- Stassun KG, Corsaro E, Pepper JA, Gaudi BS. 2018. *The Astronomical Journal* 155:22Publisher: IOP ADS Bibcode: 2018AJ....155...22S
- Stauffer J, Collier Cameron A, Jardine M, David TJ, Rebull L, et al. 2017. *The Astronomical Journal* 153:152Publisher: IOP ADS Bibcode: 2017AJ....153..152S
- Stello D, Bruntt H, Preston H, Buzasi D. 2008. *The Astrophysical Journal* 674(1):L53–L56
- Stello D, Huber D, Bedding TR, Benomar O, Bildsten L, et al. 2013. *The Astrophysical Journal* 765(2):L41
- Stello D, Zinn J, Elsworth Y, Garcia RA, Kallinger T, et al. 2017. *The Astrophysical Journal* 835:83Publisher: IOP ADS Bibcode: 2017ApJ...835...83S
- Stotesbury I, Edwards B, Lavigne JF, Pesquita V, Veilleux J, et al. 2022. *Twinkle: a small satellite spectroscopy mission for the next phase of exoplanet science*. vol. 12180. eprint: arXiv:2209.03337. ADS Bibcode: 2022SPIE12180E..33S
- Szabó GM, Pál A, Kiss C, Kiss LL, Molnár L, et al. 2017. *Astronomy and Astrophysics* 599:A44Publisher: EDP ADS Bibcode: 2017A&A...599A..44S
- Szabó R, Pál A, Sárneczky K, Szabó GM, Molnár L, et al. 2016. *Astronomy and Astrophysics* 596:A40Publisher: EDP ADS Bibcode: 2016A&A...596A..40S
- Tajiri T, Kawahara H, Aizawa M, Fujii MS, Hattori K, et al. 2020. *The Astrophysical Journal Supplement Series* 251:18Publisher: IOP ADS Bibcode: 2020ApJS..251...18T
- Takács N, Kiss C, Szakáts R, Plachy E, Kalup CE, et al. 2025. *The Astrophysical Journal* 986:L33ADS Bibcode: 2025ApJ...986L..33T
- Tardugno Poleo V, Eisner N, Hogg DW. 2024. *The Astronomical Journal* 168:100ADS Bibcode: 2024AJ....168..100T
- Tarrant NJ, Chaplin WJ, Elsworth Y, Sreckley SA, Stevens IR. 2007. *Monthly Notices of the Royal Astronomical Society: Letters* 382(1):L48–L52
- Terry SK, Hosek Jr. MW, Lu JR, Lam C, Abrams N, et al. 2023. The Galactic Center with Roman. ADS Bibcode: 2023arXiv230612485T
- Thompson SE, Everett M, Mullally F, Barclay T, Howell SB, et al. 2012. *The Astrophysical Journal* 753:86Publisher: IOP ADS Bibcode: 2012ApJ...753...86T
- Thompson TA, Kochanek CS, Stanek KZ, Badenes C, Post RS, et al. 2019. *Science* 366:637–640ADS Bibcode: 2019Sci...366..637T
- Tinetti G, Drossart P, Eccleston P, Hartogh P, Heske A, et al. 2018. *Experimental Astronomy* 46:135–209Publisher: Springer ADS Bibcode: 2018ExA....46..135T
- Tonry JL, Denneau L, Heinze AN, Stalder B, Smith KW, et al. 2018. *Publications of the Astronomical Society of the Pacific* 130:064505ADS Bibcode: 2018PASP..130f4505T
- Torres G, Andersen J, Gimenez A. 2010. *The Astronomy and Astrophysics Review* 18(1-2):67–126ArXiv: 0908.2624
- Trampedach R, Stein RF, Christensen-Dalsgaard J, Nordlund A, Asplund M. 2014. *Monthly Notices of the Royal Astronomical Society* 445:4366–4384ADS Bibcode: 2014MNRAS.445.4366T
- Treiber HP, Hinkle JT, Fausnaugh MM, Shappee BJ, Kochanek CS, et al. 2023. *Monthly Notices of the Royal Astronomical Society* 525:5795–5812Publisher: OUP ADS Bibcode: 2023MNRAS.525.5795T
- Tschernitz J, Bourdin PA. 2025. *The Astrophysical Journal* 979:L39Publisher: IOP ADS Bibcode: 2025ApJ...979L..39T
- Valley PJ, Fausnaugh M, Jha SW, Tucker MA, Eweis Y, et al. 2019. *Monthly Notices of the Royal*

- Astronomical Society* 487:2372–2384Publisher: OUP ADS Bibcode: 2019MNRAS.487.2372V
- Van Doorselaere T, Shariati H, Debosscher J. 2017. *The Astrophysical Journal Supplement Series* 232:26Publisher: IOP ADS Bibcode: 2017ApJS..232...26V
- van Leeuwen F. 2007. *Astronomy and Astrophysics* 474:653–664Publisher: EDP ADS Bibcode: 2007A&A...474..653V
- Van Reeth T, Tkachenko A, Aerts C. 2016. *Astronomy and Astrophysics* 593:A120ADS Bibcode: 2016A&A...593A.120V
- van Saders JL, Ceillier T, Metcalfe TS, Aguirre VS, Pinsonneault MH, et al. 2016. *Nature* 529(7585):181–184
- Varghese A, Ratnasingam RP, Ramírez-Galeano L, Mathis S, Rogers TM. 2025. *The Astrophysical Journal* 992:L5Publisher: IOP ADS Bibcode: 2025ApJ...992L...5V
- Venuti L, Cody AM, Rebull LM, Beccari G, Irwin MJ, et al. 2021. *The Astronomical Journal* 162:101Publisher: IOP ADS Bibcode: 2021AJ....162..101V
- Verma K, Raodeo K, Basu S, Silva Aguirre V, Mazumdar A, et al. 2019. *Monthly Notices of the Royal Astronomical Society* 483:4678–4694Publisher: OUP ADS Bibcode: 2019MNRAS.483.4678V
- Vorontsov SV, Zharkov VN, Lubimov VM. 1976. *Icarus* 27:109–118ADS Bibcode: 1976Icar...27..109V
- Wainer TM, Davenport JRA, Tovar Mendoza G, Feinstein AD, Wagg T. 2024. *The Astronomical Journal* 168:232ADS Bibcode: 2024AJ....168..232W
- Walker G, Matthews J, Kuschnig R, Johnson R, Rucinski S, et al. 2003. *Publications of the Astronomical Society of the Pacific* 115(811):1023–1035
- Walkowicz LM, Basri G, Batalha N, Gilliland RL, Jenkins J, et al. 2011. *The Astronomical Journal* 141:50Publisher: IOP ADS Bibcode: 2011AJ....141...50W
- Wang Q, Armstrong P, Zenati Y, Ridden-Harper R, Rest A, et al. 2023. *The Astrophysical Journal* 943:L15ADS Bibcode: 2023ApJ...943L..15W
- Wang Q, Rest A, Dimitriadis G, Ridden-Harper R, Siebert MR, et al. 2024. *The Astrophysical Journal* 962:17ADS Bibcode: 2024ApJ...962...17W
- Wang Q, Rest A, Zenati Y, Ridden-Harper R, Dimitriadis G, et al. 2021. *The Astrophysical Journal* 923:167Publisher: IOP ADS Bibcode: 2021ApJ...923..167W
- Warfield JT, Zinn JC, Schonhut-Stasik J, Johnson JW, Pinsonneault MH, et al. 2024. *The Astronomical Journal* 167:208Publisher: IOP ADS Bibcode: 2024AJ....167..208W
- Weaver ZR, Williamson KE, Jorstad SG, Marscher AP, Larionov VM, et al. 2020. *The Astrophysical Journal* 900:137Publisher: IOP ADS Bibcode: 2020ApJ...900..137W
- Wehrle AE, Carini M, Wiita PJ. 2019. *The Astrophysical Journal* 877:151Publisher: IOP ADS Bibcode: 2019ApJ...877..151W
- Wehrle AE, Carini M, Wiita PJ, Pepper J, Gaudi BS, et al. 2023. *The Astrophysical Journal* 951:58Publisher: IOP ADS Bibcode: 2023ApJ...951...58W
- Weiss TJ, Downing NJ, Pinsonneault MH, Zinn JC, Stello D, et al. 2025. Modeling Asteroseismic Yields for the Roman Galactic Bulge Time-Domain Survey. ADS Bibcode: 2025arXiv250304999W
- Weiss WW, Rucinski SM, Moffat AFJ, Schwarzenberg-Czerny A, Koudelka OF, et al. 2014. *Publications of the Astronomical Society of the Pacific* 126:573Publisher: IOP ADS Bibcode: 2014PASP..126..573W
- Wheeler A, Kipping D. 2019. *Monthly Notices of the Royal Astronomical Society* 485:5498–5510ADS Bibcode: 2019MNRAS.485.5498W
- Wilson RF, Barclay T, Powell BP, Schlieder J, Hedges C, et al. 2023. *The Astrophysical Journal Supplement Series* 269:5ADS Bibcode: 2023ApJS..269....5W
- Winn JN. 2024. The Transiting Exoplanet Survey Satellite. ADS Bibcode: 2024arXiv241012905W
- Winn JN, Matthews JM, Dawson RI, Fabrycky D, Holman MJ, et al. 2011. *The Astrophysical Journal* 737:L18Publisher: IOP ADS Bibcode: 2011ApJ...737L..18W
- Witzke V, Shapiro AI, Kostogryz NM, Mauviard L, Bhatia TS, et al. 2024. *Astronomy and Astrophysics* 681:A81Publisher: EDP ADS Bibcode: 2024A&A...681A..81W

- Wolniewicz LM, Berger TA, Huber D. 2021. *The Astronomical Journal* 161(5):231
- Yamaguchi N, El-Badry K, Sorabella NM. 2024. *Publications of the Astronomical Society of the Pacific* 136:124202Publisher: IOP ADS Bibcode: 2024PASP..136l4202Y
- Yang H, Liu J. 2019. *The Astrophysical Journal Supplement Series* 241:29Publisher: IOP ADS Bibcode: 2019ApJS..241...29Y
- Yang KE, Sun X, Kerr GS, Hudson HS. 2023. *The Astrophysical Journal* 959:54Publisher: IOP ADS Bibcode: 2023ApJ...959...54Y
- Yu J, Huber D, Bedding TR, Stello D, Hon M, et al. 2018. *The Astrophysical Journal Supplement Series* 236(2):42
- Zasche P, Henzl Z, Mašek M, Uhlař R, Kára J, et al. 2023. *Astronomy and Astrophysics* 675:A113Publisher: EDP ADS Bibcode: 2023A&A...675A.113Z
- Zhang J, Shapiro AI, Bi S, Xiang M, Reinhold T, et al. 2020. *The Astrophysical Journal* 894:L11Publisher: IOP ADS Bibcode: 2020ApJ...894L..11Z
- Zhou Y, Christensen-Dalsgaard J, Asplund M, Li Y, Trampedach R, et al. 2024. *The Astrophysical Journal* 962:118Publisher: IOP ADS Bibcode: 2024ApJ...962..118Z
- Zinn JC, Stello D, Elsworth Y, García RA, Kallinger T, et al. 2020. *The Astrophysical Journal Supplement Series* 251:23Publisher: IOP ADS Bibcode: 2020ApJS..251...23Z
- Zong W, Fu JN, De Cat P, Wang J, Shi J, et al. 2020. *The Astrophysical Journal Supplement Series* 251:15ADS Bibcode: 2020ApJS..251...15Z