

GAUSS–HEILBRONN SUMS AND COVERINGS OF DELIGNE–LUSZTIG TYPE CURVES

TETSUSHI ITO, DAICHI TAKEUCHI, AND TAKAHIRO TSUSHIMA

ABSTRACT. We study exponential sums on Witt vectors, known as Gauss–Heilbronn sums, and the curves whose Frobenius traces realize these sums via a Deligne–Lusztig type construction. For 3-typical Witt vectors of length two, we analyze Gauss–Heilbronn sums, from which we fully determine the Frobenius slopes of the associated curves.

1. INTRODUCTION

Exponential sums over finite fields have been studied extensively for their intrinsic arithmetic properties and their connections to the geometry of algebraic curves. Classical examples include cubic and Kloosterman-type sums; see, for instance, [3, 5, 6, 7, 9, 15, 16, 32]. Even in these basic yet nontrivial cases, such sums already exhibit rich arithmetic phenomena.

A natural generalization arises from the theory of Witt vectors, whose higher-level additive characters give rise to a richer family of exponential sums. Let q be a power of a prime p , let l be a prime, and let $W_{n,q}$ denote the ring of l -typical Witt vectors of length n over \mathbb{F}_q . Associated to characters of $W_{n,q}$, one can define exponential sums that carry significant arithmetic content and reveal intricate connections between number-theoretic and geometric structures; see, for example, [8, 18, 19, 20, 21, 25, 26, 27, 30, 31]. For an integer $m \geq 1$, which plays the role of a tame level when $p = l$, we introduce a Deligne–Lusztig type affine curve $C_{m,n,l,q}/\mathbb{F}_q$ whose Frobenius traces realize the exponential sums

$$\mathcal{S} = \left\{ - \sum_{x \in \mathbb{F}_q} \psi \left(\sum_{i=0}^{n-1} a_i x^{l^i m} \right) \mid (a_0, \dots, a_{n-1}) \in \mathbb{F}_q^n \setminus \{0\} \right\}, \quad \text{in the case } p \neq l,$$

and

$$\mathcal{T} = \left\{ - \sum_{x \in \mathbb{F}_q} \xi(a \cdot (x^m, 0, \dots, 0)) \mid a \in W_{n,q} \setminus \{0\} \right\}, \quad \text{in the case } p = l,$$

where ψ is a nontrivial additive character of \mathbb{F}_q and ξ is an additive character of $W_{n,q}$ of order p^n . If $m = 1$, the sums \mathcal{T} are referred to as Gauss–Heilbronn sums in [30]. In the following, we simply write $C_{m,n}$ for $C_{m,n,l,q}$.

In the case $n = 2$, we give an explicit formula for the L -polynomials of $\overline{C}_{1,2}$. Next, assuming $p = l$, we study the finite étale covering

$$C_{P,n} \longrightarrow C_{1,n},$$

which is associated with a separable \mathbb{F}_p -linearized polynomial $P(x) \in \mathbb{F}_q[x]$. For a smooth affine curve X , let \overline{X} denote its smooth compactification. If $n = 2$ and $p = 2$,

2020 *Mathematics Subject Classification.* Primary 11M38; Secondary 11L03, 11T24, 14F20, 11G20.

Key words and phrases. Gauss–Heilbronn sum, Witt vectors, Deligne–Lusztig type curve, L -polynomial, Frobenius slopes.

the curves $\{\overline{C}_{P,2}\}_P$ can be identified with the van der Geer–van der Vlugt curves [14]. In particular, we give an explicit formula for the L -polynomials of $\overline{C}_{P,2}$ in the case $n = 2$ and $p = 3$. Consequently, the Frobenius slopes of $\overline{C}_{P,2}$ are $\{1/3, 2/3\}$. By varying the choice of P , we enlarge the class of curves whose Frobenius slopes can be completely determined. We also determine the Frobenius slopes of the smooth compactification $\overline{C}_{2,2}$.

Let $W = W_{2,q} = \mathbb{F}_q^2$ with abelian group law

$$(a, b) + (a', b') = (a + a', b + b' - aa'(a + a')).$$

Fix a prime $\ell \neq p$. Let W_{prim}^\vee denote the set of $\overline{\mathbb{Q}}_\ell$ -valued characters whose restriction to $\{0\} \times \mathbb{F}_q$ is nontrivial. For a smooth projective curve X/\mathbb{F}_q , its L -polynomial is

$$L_{X/\mathbb{F}_q}(T) := \det(1 - \text{Fr}_q^* T; H^1(X_{\mathbb{F}}, \overline{\mathbb{Q}}_\ell)) \in \mathbb{Q}[T],$$

where \mathbb{F} is an algebraic closure of \mathbb{F}_q . For $\xi \in W_{\text{prim}}^\vee$, let

$$S_\xi := \sum_{x \in \mathbb{F}_q} \xi(x, 0).$$

As one of the principal results, we obtain an explicit factorization of the L -polynomial of $\overline{C}_{1,2}$. The statement is as follows.

Theorem A (Explicit L -polynomial for $n = 2$). *Assume $n = 2$ and $l = 3$. Then we have*

$$L_{\overline{C}_{1,2}/\mathbb{F}_q}(T) = \prod_{\xi \in W_{\text{prim}}^\vee} (1 + S_\xi T + qT^2).$$

This may be viewed as a generalization of Carlitz’s formula [5] for cubic exponential sums.

We now focus on the case $p = 3$ and study the 3-adic valuation of S_ξ .

Theorem B (Valuation of Gauss–Heilbronn sums). *Assume $n = 2$ and $p = l = 3$. Let $\zeta_9 := e^{2\pi i/9}$ and let \mathfrak{p} denote a prime of $\mathbb{Q}(\zeta_9)$ lying over (3) . Let $v_{\mathfrak{p}}(-)$ denote the valuation of $\mathbb{Q}(\zeta_9)$ at \mathfrak{p} . We write $q = 3^f$. Then we have*

$$v_{\mathfrak{p}}(S_\xi) = \frac{f}{3}.$$

This theorem is proved by comparing the exponential sums with the Frobenius traces of certain supersingular curves.

In [30, Theorem 1.3], Liu computes the Frobenius slopes of $\overline{C}_{1,n}$. While Liu states the result for general q , the proof crucially relies on the assumption $q = p$ (see Remark 5.8 for details). Our argument applies uniformly for all $q = 3^f$, including the case $f > 1$, where Liu’s method seems not to apply. Thus Theorem A provides a genuinely new valuation result for Gauss–Heilbronn sums.

As an application of Theorem B, we deduce the following result for $m = 2$.

Theorem C (Frobenius slopes). *Assume $m = 2$, $n = 2$ and $p = l = 3$. The Frobenius slopes of $\overline{C}_{2,2}$ with respect to Fr_q^* are*

$$\left\{ \frac{1}{2}, \frac{1}{3}, \frac{2}{3}, \frac{1}{6}, \frac{5}{6} \right\}.$$

Using Theorem B, we obtain

$$v_{\mathfrak{p}} \left(\sum_{x \in \mathbb{F}_q} \nu_q(x) \xi(x, 0) \right) = \frac{f}{6},$$

where $\nu_q(x) := x^{\frac{q-1}{2}}$ is the quadratic character and $\xi \in W_{\text{prim}}^{\vee}$. Combining this valuation with the product formula in [29] yields Theorem C.

For a finite extension $\mathbb{F}_{p^a}/\mathbb{F}_{p^b}$, let $\text{Tr}_{p^a/p^b}: W_{n,p^a} \rightarrow W_{n,p^b}$ denote the trace map.

We define the curve $C_{P,n}$ associated with a separable \mathbb{F}_p -linearized polynomial $P(x) \in \mathbb{F}_q[x]$. The curve $C_{P,n}$ is geometrically connected. We will study the L -polynomial and Frobenius slopes of the projective curve $\overline{C}_{P,n}$ in the case $n = 2$ and $p = l = 3$.

Using Theorems A and B, we show the following.

Theorem D (Explicit L -polynomial for $n = 2$, $p = l = 3$). *Assume $n = 2$ and $p = l = 3$. We take a nontrivial additive character ψ of \mathbb{F}_q . For a suitable extension \mathbb{F}_{q^s} one has*

$$(1.1) \quad L_{\overline{C}_{P,2}/\mathbb{F}_{q^s}}(T) = \prod_{\xi \in W_{\text{prim}}^{\vee}} \prod_{\alpha \in K_{P,\psi}} \left(1 + \sum_{x \in \mathbb{F}_{q^s}} \psi(\text{Tr}_{q^s/q}(\alpha x)) \xi(\text{Tr}_{q^s/q}(x, 0)) T + q^s T^2 \right),$$

where $K_{P,\psi} \subset \mathbb{F}_{q^s}$ is a finite explicitly determined subset depending on P and ψ . As a consequence, the Frobenius slopes of $\overline{C}_{P,2}$ over \mathbb{F}_q are precisely $\{1/3, 2/3\}$.

The proof proceeds by reducing the problem to Theorem A and then applying Theorem B to evaluate the resulting exponential sums.

Organization of the paper. Section 2 introduces the Deligne-Lusztig type curve $C_{m,n}$ and develops its basic properties for general levels. Section 3 proves Theorem A and clarifies the supersingularity phenomena for $n = 2$ and $l = 3$, showing that $\overline{C}_{m,n}$ is supersingular when $p = 2$ or 5 . In contrast, when $p = l$ (with l an arbitrary prime) or $p \equiv 1 \pmod{l}$, we show that $\overline{C}_{m,n}$ is not supersingular. The proofs of these cases rely on Katz's monodromy theory and Stickelberger's theorem, respectively, and are independent of our main valuation results. Section 4 treats the case $p = 3$, and proves Theorems B and C. Section 5 studies the coverings $C_{P,n}$ of $C_{1,n}$ and establishes Theorem D.

2. DELIGNE-LUSZTIG TYPE CONSTRUCTION

In this section, we introduce the curve $C_{m,n,l,q}$, which is the main object of study in this paper.

Let l be a fixed prime number and $n \geq 1$. The l -typical Witt vector scheme of length n , denoted by \mathbb{W}_n , is defined over \mathbb{Z} as the affine scheme

$$\mathbb{W}_n := \text{Spec } \mathbb{Z}[X_0, X_1, \dots, X_{n-1}],$$

where the coordinates X_i correspond to the components of a Witt vector.

To define the addition on \mathbb{W}_n , we use the ghost map

$$\text{gh}: \mathbb{W}_n \longrightarrow \mathbb{G}_{a,\mathbb{Z}}^n, \quad X \mapsto (w_0(X), \dots, w_{n-1}(X)),$$

where the ghost coordinates w_i are integer-coefficient polynomials in X_0, \dots, X_{n-1} given by

$$\begin{aligned} w_0(X) &= X_0, \\ w_1(X) &= X_0^l + lX_1, \\ w_2(X) &= X_0^{l^2} + lX_1^l + l^2X_2, \\ &\vdots \end{aligned}$$

The addition and multiplication on \mathbb{W}_n are defined uniquely by the conditions

$$\text{gh}(X + Y) = \text{gh}(X) + \text{gh}(Y), \quad \text{gh}(X \cdot Y) = \text{gh}(X) \cdot \text{gh}(Y),$$

so that both operations become coordinate-wise in the ghost components. In other words, the Witt vector sum and product

$$\begin{aligned} (2.1) \quad X + Y &:= (S_0(X, Y), \dots, S_{n-1}(X, Y)), \\ X \cdot Y &:= (P_0(X, Y), \dots, P_{n-1}(X, Y)) \end{aligned}$$

are determined by integer polynomials $S_i(X, Y), P_i(X, Y)$ satisfying the above identities (cf. [34, II§6]).

Let q be a power of a prime number p . We let

$$(2.2) \quad \mathbb{W}_{n,q} := \mathbb{W}_n \otimes_{\mathbb{Z}} \mathbb{F}_q, \quad W_{n,q} := \mathbb{W}_{n,q}(\mathbb{F}_q).$$

Let

$$L_q: \mathbb{W}_{n,q} \rightarrow \mathbb{W}_{n,q}, \quad (x_0, \dots, x_{n-1}) \mapsto (x_0^q, \dots, x_{n-1}^q) - (x_0, \dots, x_{n-1})$$

be the Lang torsor.

Definition 2.1. Let m be a positive integer prime to p . Let

$$i: \mathbb{A}_{\mathbb{F}_q}^1 \hookrightarrow \mathbb{W}_{n,q}, \quad t \mapsto (t^m, 0, \dots, 0)$$

be the closed immersion. Let $C_{m,n,l,q}$ denote the affine curve over \mathbb{F}_q defined by the cartesian diagram

$$\begin{array}{ccc} C_{m,n,l,q} & \longrightarrow & \mathbb{W}_{n,q} \\ \downarrow \pi & & \downarrow L_q \\ \mathbb{A}_{\mathbb{F}_q}^1 & \xrightarrow{i} & \mathbb{W}_{n,q}. \end{array}$$

Then $C_{m,n,l,q}$ admits a natural action of $W_{n,q} := \mathbb{W}_{n,q}(\mathbb{F}_q)$. The morphism

$$(2.3) \quad \pi: C_{m,n,l,q} \rightarrow \mathbb{A}_{\mathbb{F}_q}^1, \quad (t, x_0, \dots, x_{n-1}) \mapsto t,$$

is a finite étale Galois covering whose Galois group is $W_{n,q}$. If $m = 1$, the parameter t is expressed as $x_0^q - x_0$.

For $\xi \in W_{n,q}^\vee$, let $\mathcal{L}_\xi(x_0, \dots, x_{n-1})$ denote the sheaf of rank one on $\mathbb{W}_{n,q}$ defined by the $W_{n,q}$ -covering $L_q: \mathbb{W}_{n,q} \rightarrow \mathbb{W}_{n,q}$ and ξ . For a morphism of \mathbb{F}_q -schemes $(f_0, \dots, f_{n-1}): X \rightarrow \mathbb{W}_{n,q}$, $x \mapsto (f_0(x), \dots, f_{n-1}(x))$, let $\mathcal{L}_\xi(f_0, \dots, f_{n-1})$ denote the pull-back

$$(f_0, \dots, f_{n-1})^* \mathcal{L}_\xi(x_0, \dots, x_{n-1}).$$

We simply write $\mathcal{L}_\xi(t^m, 0)$ for $\mathcal{L}_\xi(t^m, 0, \dots, 0)$.

Let \mathbb{F} be an algebraic closure of \mathbb{F}_q . We write \mathbb{A}^1 for the affine line over \mathbb{F} . We take a prime number $\ell \neq p$. For an algebraic variety X over a field F and a field extension E/F , let X_E denote the base change of X to E . For an algebraic variety X over \mathbb{F}_q ,

we denote by $H_c^i(X)$ and $H^i(X)$ for $H_c^i(X_{\mathbb{F}}, \overline{\mathbb{Q}}_{\ell})$ and $H^i(X_{\mathbb{F}}, \overline{\mathbb{Q}}_{\ell})$, respectively. For a finite abelian group A , let A^{\vee} denote the character group $\text{Hom}_{\mathbb{Z}}(A, \overline{\mathbb{Q}}_{\ell}^{\times})$.

Lemma 2.2. *We have an isomorphism*

$$H_c^1(C_{m,n,l,q}) \simeq \bigoplus_{\xi \in W_{n,q}^{\vee}} H_c^1(\mathbb{A}^1, \mathcal{L}_{\xi}(t^m, 0's)).$$

Proof. Let π be as in (2.3). We have $\pi_* \overline{\mathbb{Q}}_{\ell} \simeq \bigoplus_{\xi \in W_{n,q}^{\vee}} \mathcal{L}_{\xi}(t^m, 0's)$. Taking $H_c^1(\mathbb{A}^1, -)$ gives the claim. \square

Lemma 2.3. *Assume $p \neq l$. The ghost morphism*

$$\text{gh}: \mathbb{W}_{n,q} \rightarrow \mathbb{G}_{a, \mathbb{F}_q}^n, \quad x = (x_0, \dots, x_{n-1}) \mapsto (w_0(x), \dots, w_{n-1}(x))$$

is an isomorphism of ring schemes.

Proof. This is well-known (cf. [34, II§6, Théorème 6]). \square

Lemma 2.4. *Assume $p \neq l$. The curve $C_{m,n,l,q}$ is defined by*

$$\begin{aligned} w_0^q - w_0 &= t^m, \\ w_1^q - w_1 &= t^{lm}, \\ &\vdots \\ w_{n-1}^q - w_{n-1} &= t^{l^{n-1}m}. \end{aligned}$$

Proof. We consider the defining equation of $C_{m,n,l,q}$:

$$(2.4) \quad (x_0^q, \dots, x_{n-1}^q) = (x_0, \dots, x_{n-1}) + (t^m, 0 \dots, 0).$$

We note $w_i(x_0^q, \dots, x_{n-1}^q) = w_i(x_0, \dots, x_{n-1})^q$. We have

$$(2.5) \quad w_i(t^m, 0's) := w_i(t^m, 0, \dots, 0) = t^{l^i m} \quad \text{for } 0 \leq i \leq n-1.$$

Let $x = (x_0, \dots, x_{n-1})$. By the definition of the sum of $\mathbb{W}_{n,q}$ and Lemma 2.3, the equality (2.4) is equivalent to

$$\begin{aligned} w_0(x)^q &= w_0(x) + t^m, \\ w_1(x)^q &= w_1(x) + t^{lm}, \\ &\vdots \\ w_{n-1}(x)^q &= w_{n-1}(x) + t^{l^{n-1}m}. \end{aligned}$$

\square

Lemma 2.5. *Assume $p \neq l$. We identify $W_{n,q} = \mathbb{W}_{n,q}(\mathbb{F}_q)$ with \mathbb{F}_q^n via gh . For $a = (a_0, \dots, a_{n-1}) \in \mathbb{F}_q^n$, let $f_a(t) := \sum_{i=0}^{n-1} a_i t^{l^i m}$. Let $\psi \in \mathbb{F}_q^{\vee} \setminus \{1\}$. Let $\xi \in W_{n,q}^{\vee}$, which corresponds to the character of \mathbb{F}_q^n defined by $x := (x_0, \dots, x_{n-1}) \mapsto \psi(\sum_{i=0}^{n-1} a_i x_i)$ with $a = (a_0, \dots, a_{n-1}) \in \mathbb{F}_q^n$. Then*

$$\mathcal{L}_{\xi}(t^m, 0's) \simeq \mathcal{L}_{\psi}(f_a(t)).$$

We also have

$$\sum_{t \in \mathbb{F}_q} \xi(t^m, 0's) = \sum_{t \in \mathbb{F}_q} \psi(f_a(t)).$$

Proof. By Lemma 2.3 and (2.5), we have

$$(2.6) \quad \mathcal{L}_\xi(t^m, 0's) \simeq \mathcal{L}_\psi \left(\sum_{i=0}^{n-1} a_i w_i(t^m, 0's) \right) \simeq \mathcal{L}_\psi(f_a(t)).$$

Finally, the map $gh: W_{n,q} \xrightarrow{\sim} \mathbb{F}_q^n$ is an isomorphism, and applying the same identity $w_i(t^m, 0's) = t^{l^i m}$ gives

$$\sum_{t \in \mathbb{F}_q} \xi(t^m, 0's) = \sum_{t \in \mathbb{F}_q} \psi(f_a(t)).$$

□

Corollary 2.6. *Assume $p \neq l$. We take a nontrivial character $\psi \in \mathbb{F}_q^\vee \setminus \{1\}$. For $a = (a_0, \dots, a_{n-1}) \in \mathbb{F}_q^n$, let $f_a(t) := \sum_{i=0}^{n-1} a_i t^{l^i m}$. Then we have an isomorphism*

$$H_c^1(C_{m,n,l,q}) \simeq \bigoplus_{a \in \mathbb{F}_q^n} H_c^1(\mathbb{A}^1, \mathcal{L}_\psi(f_a(t))).$$

Furthermore, we have

$$\dim H_c^1(\mathbb{A}^1, \mathcal{L}_\psi(f_a(t))) = \deg(f_a(t) - 1) \quad \text{for } a \neq 0,$$

$$\dim H_c^1(C_{m,n,l,q}) = \sum_{a \in \mathbb{F}_q^n \setminus \{0\}} \deg(f_a(t) - 1) = (q-1) \sum_{i=0}^{n-1} q^i (ml^i - 1).$$

Proof. The first claim follows from Lemmas 2.2 and 2.5.

The latter claim follows from $H_c^i(\mathbb{A}^1, \mathcal{L}_\psi(f_a(t))) = 0$ for $i \neq 1$ and the Grothendieck–Ogg–Shafarevich formula, since the Swan conductor exponent of $\mathcal{L}_\psi(f_a(t))$ is $\deg f_a(t)$ by $p \nmid lm$. The last equality is directly checked. □

Now, we treat the case $p = l$. We now recall a result of [4] and [18, Lemma 3.1]. Let $W_{n,p} := \mathbb{W}_n(\mathbb{F}_p)$ and let $\text{Tr}_{q/p}: W_{n,q} \rightarrow W_{n,p}$ denote the trace map. Let $W_{n,q,\text{prim}}$ denote the set of characters ξ of $W_{n,q}$ such that $\xi|_{\{0's\} \times \mathbb{F}_q} \neq 1$.

Lemma 2.7. *Let ${}^0\xi \in W_{n,p}^\vee$ be a faithful character. For $a \in W_{n,q}$, define $\xi_a \in W_{n,q}^\vee$ by*

$$\xi_a(x) = {}^0\xi \circ \text{Tr}_{q/p}(ax).$$

Then the map

$$W_{n,q} \longrightarrow W_{n,q}^\vee, \quad a \longmapsto \xi_a$$

is a bijection. For $a = (a_0, \dots, a_{n-1}) \in W_{n,q}$, we have

$$\xi_a \in W_{n,q,\text{prim}}^\vee \iff a_0 \neq 0.$$

Proof. The former claim is well-known.

The latter claim follows from $\xi_a(0's, x) = {}^0\xi(\text{Tr}_{q/p}(0's, a_0^{p^{n-1}} x))$ for $x \in \mathbb{F}_q$. □

Lemma 2.8 ([18, Lemma 3.1]). *If $a = (a_0, \dots, a_{n-1}) \in W_{n,q}$ is nonzero, then ξ_a has order p^{n-d} , where d is the largest integer such that $a_i = 0$ for all $i \leq d-1$. For such an a , the Swan conductor exponent of $\mathcal{L}_{\xi_a}(t^m, 0's)$ at ∞ is given by*

$$\text{Swan}_\infty(\mathcal{L}_{\xi_a}(t^m, 0's)) = mp^{n-1-d}.$$

We have $\dim H_c^1(\mathbb{A}^1, \mathcal{L}_{\xi_a}(t^m, 0's)) = mp^{n-1-d} - 1$. In particular, if $\xi_a \in W_{n,q,\text{prim}}^\vee$, we have $\dim H_c^1(\mathbb{A}^1, \mathcal{L}_{\xi_a}(t^m, 0's)) = mp^{n-1} - 1$.

Proof. The first claim is just [18, Lemma 3.1]. The second claim follows from the Grothendieck–Ogg–Shafarevich formula and $H_c^i(\mathbb{A}^1, \mathcal{L}_{\xi_a}(t^m, 0's)) = 0$ for $i \neq 1$. The last claim follows from Lemma 2.7. \square

Lemma 2.9. *Assume $p = l$ and write $q = p^f$. We have an isomorphism $W_{n,q} \simeq (\mathbb{Z}/p^n\mathbb{Z})^f$ as groups.*

Proof. We take a basis $\{v_1, \dots, v_f\}$ of \mathbb{F}_q over \mathbb{F}_p . Recall

$$\begin{aligned} F: \mathbb{W}_{n,q} &\rightarrow \mathbb{W}_{n,q}, \quad (x_0, \dots, x_{n-1}) \mapsto (x_0^p, \dots, x_{n-1}^p), \\ V: \mathbb{W}_{n,q} &\rightarrow \mathbb{W}_{n,q}, \quad (x_0, \dots, x_{n-1}) \mapsto (0, x_0, \dots, x_{n-2}). \end{aligned}$$

By $FV = VF = p$,

$$p^j(v_i, 0's) = (0, \dots, 0, v_i^{p^j}, 0, \dots, 0),$$

where $v_i^{p^j}$ appears in the $(j+1)$ -th component. Thus $(v_i, 0's)$ has order p^n . Let

$$\varphi: (\mathbb{Z}/p^n\mathbb{Z})^f \rightarrow W_{n,q}, \quad (a_1, \dots, a_f) \mapsto \sum_{i=1}^f a_i(v_i, 0's).$$

Assume $\varphi(a_1, \dots, a_f) = 0$. Since the Witt addition coincides with coordinatewise addition on the first component, the first coordinate of $\varphi(a_1, \dots, a_f)$ is

$$\sum_{i=1}^f (a_i \bmod p) v_i \in \mathbb{F}_q.$$

Since $\{v_1, \dots, v_f\}$ is an \mathbb{F}_p -basis, we obtain $a_i \equiv 0 \pmod{p}$ for all i . By applying the same argument to the higher p -power components, we obtain $a_i \equiv 0 \pmod{p^n}$ for every i . Thus φ is injective. Since the both sides of φ have the same cardinality, we obtain the claim. \square

For a smooth projective curve X , let $g(X)$ denote its genus.

Corollary 2.10. *We have*

$$g(\overline{C}_{m,n,l,q}) = \frac{q-1}{2} \sum_{i=0}^{n-1} q^i (ml^i - 1).$$

Proof. We show that the canonical map

$$H_c^1(C_{m,n,l,q}) \rightarrow H^1(C_{m,n,l,q})$$

is an isomorphism.

Assume $p \neq l$. Then the forgetful map $H_c^1(\mathbb{A}^1, \mathcal{L}_\psi(f_a(t))) \rightarrow H^1(\mathbb{A}^1, \mathcal{L}_\psi(f_a(t)))$ is an isomorphism if $a \neq 0$. Hence the claim follows from Lemma 2.6.

Assume $p = l$. For $a \in W_{n,q} \setminus \{0\}$, the canonical map

$$H_c^1(\mathbb{A}^1, \mathcal{L}_{\xi_a}(t^m, 0's)) \rightarrow H^1(\mathbb{A}^1, \mathcal{L}_{\xi_a}(t^m, 0's))$$

is an isomorphism since $\mathcal{L}_{\xi_a}(t^m, 0's)$ is ramified at ∞ . Hence the claim follows from Lemma 2.8.

By the claim, the natural map $H_c^1(C_{m,n,l,q}) \rightarrow H^1(\overline{C}_{m,n,l,q})$ is an isomorphism. The dimension of $H_c^1(C_{m,n,l,q})$ is computed using Lemmas 2.6 and 2.8. \square

Lemma 2.11. (1) *We have*

$$\mathrm{Tr}(\mathrm{Fr}_q^*; H^1_c(\mathbb{A}^1, \mathcal{L}_\xi(t^m, 0's))) = - \sum_{t \in \mathbb{F}_q} \xi(t^m, 0's).$$

(2) *We take an isomorphism $\overline{\mathbb{Q}}_\ell \simeq \mathbb{C}$. Then we have*

$$\left| \sum_{t \in \mathbb{F}_q} \xi(t^m, 0's) \right| \leq \dim H^1_c(\mathbb{A}^1, \mathcal{L}_\xi(t^m, 0's)) \sqrt{q}.$$

Proof. The claim (1) follows from the Grothendieck trace formula.

The claim (2) follows from the Weil conjectures for curves. \square

Lemma 2.12. *Let $1 \leq n' \leq n$. We write $q = p^f$. There exist finite morphisms*

$$\begin{aligned} \phi_1: C_{m,n,l,q} &\rightarrow C_{1,n,l,q}, \quad (t, x_0, \dots, x_{n-1}) \mapsto (x_0, \dots, x_{n-1}), \\ \phi_2: C_{1,n,l,q} &\rightarrow C_{1,n,l,p}, \quad (x_0, \dots, x_{n-1}) \mapsto \sum_{i=0}^{f-1} (x_0^{p^i}, \dots, x_{n-1}^{p^i}), \\ \phi_3: C_{1,n,l,p} &\rightarrow C_{1,n',l,p}, \quad (x_0, \dots, x_{n-1}) \mapsto (x_0, \dots, x_{n'-1}), \end{aligned}$$

where the sum in ϕ_2 is taken in the commutative algebraic group $\mathbb{W}_{n,q}$.

Proof. The second claim follows from, for $(x_0, \dots, x_{n-1}) \in \mathbb{W}_n$,

$$L_p(\phi_2(x_0, \dots, x_{n-1})) = (x_0^q, \dots, x_{n-1}^q) - (x_0, \dots, x_{n-1}) = L_q(x_0, \dots, x_{n-1}).$$

The first and third claims follow from definition. \square

3. WITT SUMS OF LENGTH TWO

3.1. Length two. In this section, we specialize the results of the previous section to the case $m = 1$ and $n = 2$, and discuss them in more detail for use in the next section.

We define

$$g_l(x, y) := \frac{x^l + y^l - (x + y)^l}{l} \in \mathbb{Z}[x, y],$$

which we regard as an element of $\mathbb{F}_q[x, y]$. This is nothing but $S_1(x, y)$ in (2.1).

The addition law of the algebraic group $\mathbb{W}_{2,q}$ is given by

$$(x, y) + (x', y') = (x + x', y + y' + g_l(x, x')).$$

If $l \neq 2$, we have $g_l(x, -x) = 0$. Thus

$$\begin{aligned} L_q(x, y) = (x^q, y^q) - (x, y) &= \begin{cases} (x^q, y^q) + (-x, -y + x^2) & \text{if } l = 2, \\ (x^q, y^q) + (-x, -y) & \text{if } l \neq 2 \end{cases} \\ &= \begin{cases} (x^q - x, y^q - y - x^{q+1} + x^2) & \text{if } l = 2, \\ (x^q - x, y^q - y + g_l(x^q, -x)) & \text{if } l \neq 2. \end{cases} \end{aligned}$$

Recall that $C_{1,2,l,q}$ is defined by the cartesian diagram

$$\begin{array}{ccc} C_{1,2,l,q} & \longrightarrow & \mathbb{W}_{2,q} \\ \downarrow \pi & & \downarrow L_q \\ \mathbb{A}_{\mathbb{F}_q}^1 & \xrightarrow{i} & \mathbb{W}_{2,q}, \end{array}$$

where $i: \mathbb{A}_{\mathbb{F}_q}^1 \hookrightarrow \mathbb{W}_{2,q}$ is the closed immersion $t \mapsto (t, 0)$. Then $C_{1,2,l,q}$ is a smooth affine curve defined by

$$\begin{cases} y^q - y = -x(x^q - x) & \text{if } l = 2, \\ y^q - y = -g_l(x^q, -x) & \text{if } l \neq 2. \end{cases}$$

We simply write C for $C_{1,2,l,q}$. The curve C admits a natural action of $W := \mathbb{W}_{2,q}(\mathbb{F}_q)$ as follows:

$$(3.1) \quad (x, y) \cdot (a, b) = (x + a, y + b + g_l(x, a)) \quad \text{for } (x, y) \in C, (a, b) \in W.$$

We recall that the morphism

$$\pi: C \rightarrow \mathbb{A}_{\mathbb{F}_q}^1, \quad (x, y) \mapsto x^q - x,$$

is a finite étale Galois covering whose Galois group is W (cf. (2.3)).

Lemma 3.1. *The curve $\overline{C}_{1,2,2,q}$ is supersingular.*

Proof. The defining equation of $C_{1,2,2,q}$ is $y^q - y = xR(x)$, where $R(x) := -x^q + x$ is \mathbb{F}_q -linearized. Hence the claim follows from [14, 35]. \square

Lemma 3.2. *Assume $p \neq l$. Then C is defined by $y^q - y = (x^q - x)^l$.*

Proof. By Lemma 2.4 for $m = 1$ and $n = 2$, the defining equation of C is $w_0^q - w_0 = t$ and $w_1^q - w_1 = t^l$. Setting $x = w_0$ and $y = w_1$, we obtain $y^q - y = (x^q - x)^l$, as claimed. \square

Let $\xi \in W_{\text{prim}}^\vee$. For simplicity, we set

$$\mathcal{L}_\xi := \mathcal{L}_\xi(x, 0).$$

Lemma 3.3. *We have $H_c^1(\mathbb{A}^1, \mathcal{L}_\xi) = l - 1$.*

Proof. The claim follows from Lemmas 2.5, 2.6 and 2.8. \square

Recall that W_{prim}^\vee is the set of characters of W whose restriction to the subgroup $\{0\} \times \mathbb{F}_q$ is nontrivial.

Lemma 3.4. *We have an isomorphism*

$$H_c^1(C) \simeq \bigoplus_{\xi \in W_{\text{prim}}^\vee} H_c^1(\mathbb{A}^1, \mathcal{L}_\xi).$$

Further, we have $\dim H_c^1(C) = (l - 1)q(q - 1)$.

Proof. By Lemma 2.2,

$$H_c^1(C) \simeq \bigoplus_{\xi \in W^\vee} H_c^1(\mathbb{A}^1, \mathcal{L}_\xi).$$

The $(\{0\} \times \mathbb{F}_q)$ -fixed part of $H_c^1(C)$ is zero because $C/(\{0\} \times \mathbb{F}_q)$ is isomorphic to $\mathbb{A}_{\mathbb{F}_q}^1$. Hence the former claim follows.

The latter claim follows from $\#W_{\text{prim}}^\vee = q(q - 1)$ and Lemma 3.3 (cf. Corollary 2.10). \square

We define

$$S_\xi := \sum_{x \in \mathbb{F}_q} \xi(x, 0).$$

By Lemmas 2.11(2) and 3.3,

$$(3.2) \quad |S_\xi| \leq (l-1)\sqrt{q}.$$

If $p \neq l$, by Lemma 2.3,

$$(3.3) \quad W \xrightarrow{\sim} \mathbb{F}_q^2, \quad (x_0, x_1) \mapsto (x_0, x_0^l + lx_1).$$

For $a \in \mathbb{F}_q$ and $\psi \in \mathbb{F}_q^\vee$, let $\psi_a \in \mathbb{F}_q^\vee$ denote the character $x \mapsto \psi(ax)$.

Lemma 3.5. *Assume $p \neq l$. Let $\xi \in W_{\text{prim}}^\vee$. Assume that via (3.3), the character ξ corresponds to $\psi_a \boxtimes \psi$ with $\psi \in \mathbb{F}_q^\vee \setminus \{1\}$ and $a \in \mathbb{F}_q$. We have*

$$S_\xi = \sum_{x \in \mathbb{F}_q} \psi(x^l + ax).$$

We write $q = p^f$. We define an equivalence relation on $\mathbb{F}_q^\times \times \mathbb{F}_q$ by

$$(a, b) \sim (a', b') \iff \text{there exists } c \in \mathbb{F}_p^\times \text{ such that } a' = ca \text{ and } b' = cb.$$

Lemma 3.6. *For $(a, b) \in \mathbb{F}_q^\times \times \mathbb{F}_q$, let $C_{a,b}$ denote the affine curve over \mathbb{F}_q defined by $y^p - y = ax^l + bx$. Assume $p \neq l$. Then we have*

$$H^1(\overline{C}) \simeq \bigoplus_{[(a,b)] \in (\mathbb{F}_q^\times \times \mathbb{F}_q)/\sim} H^1(\overline{C}_{a,b}).$$

Proof. We take ${}^0\psi \in \mathbb{F}_p^\vee \setminus \{1\}$ and set $\psi = {}^0\psi \circ \text{Tr}_{q/p}$. From $p \neq l$ and Lemma 2.6, it follows that

$$\begin{aligned} H^1(\overline{C}) &\simeq H_c^1(C) \simeq \bigoplus_{a_0 \in \mathbb{F}_q^\times} \bigoplus_{a_1 \in \mathbb{F}_q} H_c^1(\mathbb{A}^1, \mathcal{L}_{\psi}(a_0 x^l + a_1 x)), \\ H^1(\overline{C}_{a,b}) &\simeq H_c^1(C_{a,b}) \simeq \bigoplus_{c \in \mathbb{F}_p^\times} H_c^1(\mathbb{A}^1, \mathcal{L}_{\psi}(cax^l + cbx)). \end{aligned}$$

Thus the claim follows. \square

3.2. Stickelberger's theorem and Witt sum valuation. By computing the valuation of S_ξ when $q = p = l$, we deduce that $\overline{C}_{m,n,p,q}$ is not supersingular for $n \geq 2$ and arbitrary q .

Let $p \geq 3$ be a prime number. Let

$$\xi: W_2(\mathbb{F}_p) \rightarrow \mathbb{C}^\times$$

be a faithful character. Let $\psi(x) := \xi(0, x)$ for $x \in \mathbb{F}_p$.

For an integer $n \geq 1$, let $\zeta_n := e^{2\pi i/n}$. Fix a prime ideal \mathfrak{p}_0 of $\mathbb{Q}(\zeta_{p-1})$ lying above p . Then the mod \mathfrak{p}_0 reduction induces an isomorphism

$$\mu_{p-1}(\mathbb{C}) \rightarrow \mathbb{F}_p^\times.$$

Let ν denote the character

$$\mathbb{F}_p^\times \xrightarrow{\cong} \mu_{p-1}(\mathbb{C}) \hookrightarrow \mathbb{C}^\times.$$

We set $\nu(0) = 0$. Set

$$S_\xi := \sum_{x \in \mathbb{F}_p} \xi(x, 0),$$

$$U_\xi := \sum_{x \in \mathbb{F}_p} \nu(x)^{-1} \xi^{p-1}(x, 0).$$

Let \mathfrak{p}' be the unique prime ideal of $\mathbb{Q}(\zeta_{p-1}, \zeta_{p^2})$ lying above \mathfrak{p}_0 .

Proposition 3.7. *Let v denote the valuation of $\mathbb{Q}(\zeta_{p-1}, \zeta_{p^2})$ associated to \mathfrak{p}' , normalized by $v(p) = 1$. The following hold:*

- (1) $v(S_\xi U_\xi) = \frac{1}{p-1}$.
- (2) $v(S_\xi) = \frac{1}{p}$.
- (3) $v(U_\xi) = \frac{1}{p(p-1)}$.

Proof. (1): We simply write $g(x, y)$ for $g_p(x, y)$. We compute

$$\begin{aligned} S_\xi U_\xi &= \sum_{x, y \in \mathbb{F}_p} \xi(x, 0) \nu(y)^{-1} \xi((p-1)(y, 0)) \\ &= \sum_{x, y \in \mathbb{F}_p} \nu(y)^{-1} \xi((x, 0) + (-y, y^p)) \\ &= \sum_{x, y \in \mathbb{F}_p} \nu(y)^{-1} \xi(x - y, y^p + g(x, -y)) \\ &= \sum_{y, z \in \mathbb{F}_p} \nu(y)^{-1} \xi(z, y^p - g(y, z)) \quad (z := x - y) \\ &= \sum_z \xi(z, 0) \sum_y \nu(y)^{-1} \psi(y^p - g(y, z)). \end{aligned}$$

We claim that $v(\sum_y \nu(y)^{-1} \psi(y^p - g(y, z))) \geq \frac{1}{p-1}$ for any z . Indeed, we have

$$\sum_y \nu(y)^{-1} \psi(y^p - g(y, z)) \equiv \sum_y \nu(y)^{-1} = 0 \pmod{\mathfrak{p}'^{\frac{1}{p-1}}}.$$

Since the sum is contained in $\mathbb{Q}(\zeta_{p-1}, \zeta_p)$, we must have $v(\sum_y \nu(y)^{-1} \psi(y^p - g(y, z))) \geq \frac{1}{p-1}$.

For $x, y \in \mathbb{Q}(\zeta_{p-1}, \zeta_{p^2})$ and a rational number r , we write $x \equiv y \pmod{r+}$ if $v(x - y) > r$. We can proceed

$$\begin{aligned} S_\xi U_\xi &= \sum_z \xi(z, 0) \sum_y \nu(y)^{-1} \psi(y^p - g(y, z)) \\ &\equiv \sum_{z, y} \nu(y)^{-1} \psi(y^p - g(y, z)) \pmod{\frac{1}{p-1}+} \\ &= \sum_{z, y} \nu(y)^{-1} \psi(y^p(1 - g(1, z))) \quad (\text{replace } z \text{ by } yz) \end{aligned}$$

if $1 = g(1, z)$, then the sum \sum_y is zero. If $1 \neq g(1, z)$, we have

$$\begin{aligned} \sum_y \nu(y)^{-1} \psi(y^p(1 - g(1, z))) &= \sum_y \nu(y)^{-1} \psi(y(1 - g(1, z))) \quad (q = p) \\ &= \nu(1 - g(1, z)) \sum_y \nu(y)^{-1} \psi(y). \end{aligned}$$

Set $\alpha := \sum_y \nu(y)^{-1} \psi(y)$. By Stickelberger's formula for p -adic valuation of Gauss sum, we have $v(\alpha) = \frac{1}{p-1}$.

We obtain

$$S_\xi U_\xi \equiv \sum_z \nu(1 - g(1, z)) \alpha \pmod{\frac{1}{p-1}}.$$

To conclude, it remains to show that $\sum_z \nu(1 - g(1, z))$ is a \mathfrak{p}' -adic unit. This follows from the computation

$$\sum_z \nu(1 - g(1, z)) \equiv \sum_z (1 - g(1, z)) \pmod{\mathfrak{p}_0}$$

and the observations that $\sum_z z^i = 0$ for $i = 0, 1, \dots, p-2$ and that $g(1, z)$ is of the form $-z^{p-1} + (\text{lower term})$. Hence

$$\sum_z (1 - g(1, z)) \equiv - \sum_z z^{p-1} \equiv 1 \pmod{\mathfrak{p}_0}.$$

Thus the claim follows.

(2), (3): We show

$$v(S_\xi) \geq \frac{1}{p}, \quad v(U_\xi) \geq \frac{1}{p(p-1)}.$$

Then the claims follow from (1). For the estimate of S_ξ ,

$$S_\xi = \sum_x \xi(x, 0) \equiv 0 \pmod{\mathfrak{p}'}.$$

Let K be the unique subextension of $\mathbb{Q}(\zeta_{p^2})/\mathbb{Q}$ such that $[K : \mathbb{Q}] = p$. Note that the subgroup $\text{Gal}(\mathbb{Q}(\zeta_{p^2})/K) \subset \text{Gal}(\mathbb{Q}(\zeta_{p^2})/\mathbb{Q}) \cong W_2(\mathbb{F}_p)^\times$ consists of elements of the form $(a, 0)$. The action of $(a, 0)$ is given by $\xi(1, 0) \mapsto \xi(a, 0)$. Then $\xi(x, 0)$ is sent to $\xi(ax, 0)$ by the action of $(a, 0)$ for $x \in \mathbb{F}_p$. Hence $S_\xi \in K$. This proves $v(S_\xi) \geq \frac{1}{p}$ since the ramification index of K/\mathbb{Q} at p is p .

Similarly,

$$U_\xi = \sum_{x \in \mathbb{F}_p} \nu(x)^{-1} \xi^{p-1}(x, 0) \equiv \sum_{x \in \mathbb{F}_p} \nu(x)^{-1} = 0 \pmod{\mathfrak{p}'}.$$

Thus $v(U_\xi) \geq (p(p-1))^{-1}$. □

Corollary 3.8. *Assume that p is odd and $n \geq 2$. Then the curve $\overline{C}_{m,n,p,q}$ is not supersingular.*

Proof. By Lemma 2.12, we have finite morphisms

$$C_{m,n,p,q} \xrightarrow{\phi_1} C_{1,n,p,q} \xrightarrow{\phi_2} C_{1,n,p,p} \xrightarrow{\phi_3} C_{1,2,p,p}.$$

This induces a finite morphism $\overline{C}_{m,n,p,q} \rightarrow \overline{C}_{1,2,p,p}$. Hence it suffices to show the claim for $m = 1$, $n = 2$ and $q = p$. If $\overline{C}_{1,2,p,p}$ is supersingular, the valuation of the Frobenius trace $-S_\xi$ is greater than or equal to $1/2$. This does not occur by Proposition 3.7. \square

Remark 3.9. *The above corollary can also be deduced from using [30, Theorem 1.3]. Here we give a short proof for completeness.*

Lemma 3.10. *Write $q = p^f$ with a prime p , l an odd prime, and assume $p \equiv 1 \pmod{l}$.*

- (1) *The projective curve defined by $y^p - y = x^l$ is not supersingular.*
- (2) *The curve $\overline{C}_{m,n,l,q}$ is not supersingular.*

Proof. (1) For $\chi \in \mu_l(\mathbb{F}_p)^\vee \setminus \{1\}$ and $\psi \in \mathbb{F}_p^\vee \setminus \{1\}$, the Gauss sum $G(\chi, \psi) = \sum_{x \in \mathbb{F}_p^\times} \chi(x^{(p-1)/l}) \psi(x)$ is not of the form \sqrt{p} times a root of unity by Stickelberger's theorem (cf. [2, Remark 2.2]). Let D be the projective curve with affine equation $y^p - y = x^l$. Then $G(\chi, \psi)$ appears as a Frobenius eigenvalue on $H^1(D)$. Hence the claim follows.

(2) By Lemma 2.12, we have a finite morphism $C_{m,n,l,q} \rightarrow C_{1,2,l,q}$ and the curve $C_{1,2,l,q}$ is defined by $y^q - y = (x^q - x)^l$ by Lemma 3.2. Further, we have a finite morphism $\overline{C}_{1,2,l,q} \rightarrow D$ given by $(x, y) \mapsto (x^q - x, \sum_{i=0}^{f-1} y^{p^i})$. Hence the claim follows from (1). \square

3.3. Cubic case. In this section, we always assume that $m = 1$, $n = 2$ and $l = 3$. Then $g_3(x, x') = -xx'(x + x')$. We simply write C for $C_{1,2,3,q}$. Hence the curve C is defined by

$$y^q - y = -g_3(x^q, -x) = -x^{q+1}(x^q - x).$$

For a finite extension of finite fields $\mathbb{F}_{p^t}/\mathbb{F}_{p^s}$, let $\text{Tr}_{p^t/p^s} : \mathbb{F}_{p^t} \rightarrow \mathbb{F}_{p^s}$ denote the trace map.

3.4. Characteristic two. First, we treat the case $p = 2$.

Lemma 3.11. *Assume $p = 2$. For $(a, b) \in \mathbb{F}_q^\times \times \mathbb{F}_q$, let $E_{a,b}$ denote the elliptic curve over \mathbb{F}_q defined by $y^2 + y = ax^3 + bx$. Then*

$$\begin{aligned} L_{\overline{C}/\mathbb{F}_q}(T) &= \prod_{(a,b) \in \mathbb{F}_q^\times \times \mathbb{F}_q} L_{E_{a,b}/\mathbb{F}_q}(T) \\ &= \prod_{(a,b) \in \mathbb{F}_q^\times \times \mathbb{F}_q} \left(1 + \sum_{x \in \mathbb{F}_q} (-1)^{\text{Tr}_{q/2}(ax^3 + bx)} T + qT^2 \right) \\ &= \prod_{\xi \in W_{\text{prim}}^\vee} (1 + S_\xi T + qT^2). \end{aligned}$$

Hence \overline{C} is supersingular.

Proof. The first equality follows from Lemma 3.6. The second equality is well-known. The third equality follows from Lemma 3.5.

Since each $E_{a,b}$ is supersingular, the curve \overline{C} , whose L -polynomial is the product of $L_{E_{a,b}/\mathbb{F}_q}(T)$, is also supersingular. \square

Remark 3.12. *The sum $\sum_{x \in \mathbb{F}_q} (-1)^{\text{Tr}_{q/2}(ax^3 + bx)}$ is explicitly calculated in [6].*

3.5. Odd characteristic. Through this subsection, we assume $p \neq 2$. We count the number of fixed points.

Lemma 3.13. *Let $(a, b) \in W$. Then we have*

$$\#C_{\mathbb{F}}^{\text{Fr}_{q^2} \circ (a, b)} = \begin{cases} 2q^2, & \text{if } a \neq 0 \text{ and } b = -\frac{a^3}{4} + \zeta a \text{ with } \zeta \in \mathbb{F}_q^\times \setminus (\mathbb{F}_q^\times)^2, \\ q^2, & \text{if } a \neq 0 \text{ and } b = -\frac{a^3}{4}, \\ q^3, & \text{if } a = b = 0, \\ 0, & \text{otherwise.} \end{cases}$$

Proof. By (3.1), it suffices to count the number of pairs $(x, y) \in \mathbb{F}^2$ satisfying

$$(3.4) \quad \begin{aligned} x^{q^2} + a &= x, \\ y^{q^2} - ax^{q^2}(x^{q^2} + a) + b &= y, \\ y^q - y &= -x^{q+1}(x^q - x). \end{aligned}$$

Substituting the first equation into the second gives

$$y^{q^2} - y = ax^{q^2}(x^{q^2} + a) - b = ax^2 - a^2x - b.$$

Using the first and third equations of (3.4), we obtain

$$\begin{aligned} y^{q^2} - y &= (y^q - y)^q + (y^q - y) \\ &= -x^{q^2+q}(x^{q^2} - x^q) - x^{q+1}(x^q - x) \\ &= -x^q(x - a)(x - x^q - a) + x^{q+1}(x^q - x) \\ &= -ax^{2q} + 2ax^{q+1} - a^2x^q. \end{aligned}$$

Hence,

$$ax^{2q} - 2ax^{q+1} + a^2x^q + ax^2 - a^2x = b.$$

Set $z := x^q - x$. Then this implies

$$az^2 + a^2z = b.$$

From $x^{q^2} - x = -a$, we also have

$$z^q + z = x^{q^2} - x = -a.$$

Therefore (3.4) is equivalent to

$$(3.5) \quad \begin{aligned} z &= x^q - x, \\ z^q + z &= -a, \\ az^2 + a^2z &= b, \\ y^q - y &= -x^{q+1}(x^q - x). \end{aligned}$$

Case 1: $a \neq 0$. Let $z_1 = z + (a/2)$. Substituting this into $az^2 + a^2z = b$ gives

$$(3.6) \quad z_1^2 = \frac{a^2}{4} + \frac{b}{a}.$$

The equality $z^q + z = -a$ is equivalent to $z_1^q = -z_1$.

(i) Assume $b \neq -a^3/4$. Then $z_1 \neq 0$, and hence $z_1^{q-1} = -1$. Let $\nu_q(x) = x^{(q-1)/2}$ for $x \in \mathbb{F}_q^\times$. From (3.6), we have

$$\nu_q\left(\frac{a^2}{4} + \frac{b}{a}\right) = z_1^{q-1} = -1.$$

Thus, if a solution of (3.5) exists, we must have $\frac{a^2}{4} + \frac{b}{a} \in \mathbb{F}_q^\times \setminus (\mathbb{F}_q^\times)^2$. If this condition holds, there exist $2q^2$ solutions of (3.5).

(ii) Assume $b = -a^3/4$. Then $z_1 = 0$ by (3.6), and clearly $z_1^q = z_1$. Hence there are q^2 solutions of (3.5).

Case 2: $a = 0$. If $b \neq 0$, there is no solution of (3.5). Assume $b = 0$. Then we consider

$$x^{q^2} = x, \quad y^{q^2} = y, \quad y^q - y = -x^{q+1}(x^q - x).$$

If $x = 0$, there are q solutions. Assume $x \neq 0$. Since $x^{q+1} \in \mathbb{F}_q^\times$, we have

$$\left(\frac{y}{x^{q+1}}\right)^q - \frac{y}{x^{q+1}} = -x^q + x.$$

Thus $(y/x^{q+1}) + x \in \mathbb{F}_q$, and hence $y \in \mathbb{F}_{q^2}$. Namely, there are $q(q^2 - 1)$ such solutions. Therefore we obtain q^3 solutions of (3.5). \square

In the following, let $\xi \in W_{\text{prim}}^\vee$.

Corollary 3.14. *Let $\nu_q(x) := x^{(q-1)/2}$ for $x \in \mathbb{F}_q^\times$, and define*

$$U_\xi := \sum_{x \in \mathbb{F}_q^\times} \nu_q(x) \xi^2(x, 0).$$

Let $\psi(x) := \xi(0, x)$ for $x \in \mathbb{F}_q$, and put

$$G(\psi) := \sum_{x \in \mathbb{F}_q} \psi(x^2).$$

Then

$$\text{Tr}(\text{Fr}_{q^2}^*; H_c^1(\mathbb{A}^1, \mathcal{L}_\xi)) = -q + \nu_q(2) G(\psi) U_\xi.$$

Proof. Using Lemma 3.13 and the fixed point formula (cf. [11, Section 3]), we have

$$\begin{aligned} & \text{Tr}(\text{Fr}_{q^2}^*; H_c^1(\mathbb{A}^1, \mathcal{L}_\xi)) \\ &= \frac{1}{q^2} \sum_{(a,b) \in W} \xi^{-1}(a, b) \text{Tr}((\text{Fr}_{q^2} \circ (a, b))^*; H_c^1(C)) \\ &= \frac{1}{q^2} \sum_{(a,b) \in W} \xi^{-1}(a, b) (q^2 - \#C_{\mathbb{F}}^{\text{Fr}_{q^2} \circ (a, b)}) \\ (3.7) \quad &= - \sum_{x \in \mathbb{F}_q^\times} \xi^{-1}\left(x, -\frac{x^3}{4}\right) - 2 \sum_{x \in \mathbb{F}_q^\times, \zeta \in \mathbb{F}_q^\times \setminus (\mathbb{F}_q^\times)^2} \xi^{-1}\left(x, -\frac{x^3}{4} + \zeta x\right) - q \\ &= - \sum_{x \in \mathbb{F}_q^\times} \xi^{-2}\left(\frac{x}{2}, 0\right) - 2 \sum_{x \in \mathbb{F}_q^\times} \xi^{-2}\left(\frac{x}{2}, 0\right) \sum_{\zeta \in \mathbb{F}_q^\times \setminus (\mathbb{F}_q^\times)^2} \xi(0, -\zeta x) - q, \end{aligned}$$

where we used $2(x/2, 0) = (x, -x^3/4)$ in the last equality.

For $x \in \mathbb{F}_q^\times$, we have

$$\sum_{\zeta \in \mathbb{F}_q^\times \setminus (\mathbb{F}_q^\times)^2} \xi(0, -\zeta x) + \frac{1}{2} \sum_{\zeta \in \mathbb{F}_q^\times} \xi(0, -\zeta^2 x) = \sum_{\zeta \in \mathbb{F}_q^\times} \xi(0, -\zeta x) = -1,$$

and therefore

$$\sum_{\zeta \in \mathbb{F}_q^\times \setminus (\mathbb{F}_q^\times)^2} \xi(0, -\zeta x) = -\frac{1}{2} \left(\sum_{\zeta \in \mathbb{F}_q} \xi(0, -\zeta^2 x) + 1 \right) = -\frac{\nu_q(-x)G(\psi) + 1}{2}.$$

Using this, we obtain

$$\begin{aligned} & 2 \sum_{x \in \mathbb{F}_q^\times} \xi^{-2} \left(\frac{x}{2}, 0 \right) \sum_{\zeta \in \mathbb{F}_q^\times \setminus (\mathbb{F}_q^\times)^2} \xi(0, -\zeta x) \\ &= - \sum_{x \in \mathbb{F}_q^\times} \xi^{-2} \left(\frac{x}{2}, 0 \right) (\nu_q(-x)G(\psi) + 1) \\ &= -G(\psi) \sum_{x \in \mathbb{F}_q^\times} \nu_q(-x) \xi^{-2} \left(\frac{x}{2}, 0 \right) - \sum_{x \in \mathbb{F}_q^\times} \xi^{-2} \left(\frac{x}{2}, 0 \right) \\ &= -\nu_q(2) G(\psi) U_\xi - \sum_{x \in \mathbb{F}_q^\times} \xi^{-2} \left(\frac{x}{2}, 0 \right). \end{aligned}$$

Substituting this into (3.7) yields the claimed formula. \square

Lemma 3.15. *We have $S_\xi^2 = q + \nu_q(2) G(\psi) U_\xi$.*

Proof. Note that $2(x/2, 0) = (x, -x^3/4)$. We compute

$$\begin{aligned} S_\xi^2 &= \left(\sum_{x \in \mathbb{F}_q} \xi(x, 0) \right)^2 = \sum_{x, y \in \mathbb{F}_q} \xi(x + y, -xy(x + y)) \\ &= \sum_{x, z \in \mathbb{F}_q} \xi(z, -x(z - x)z) \quad (z := x + y) \\ &= \sum_{x, z \in \mathbb{F}_q} \xi \left(z, z \left(x - \frac{z}{2} \right)^2 - \frac{z^3}{4} \right) \\ &= \sum_{x_1, z \in \mathbb{F}_q} \xi \left(z, zx_1^2 - \frac{z^3}{4} \right) \quad (x_1 := x - \frac{z}{2}) \\ &= \sum_{z \in \mathbb{F}_q} \xi^2 \left(\frac{z}{2}, 0 \right) \sum_{x_1 \in \mathbb{F}_q} \psi(zx_1^2). \end{aligned}$$

For $z = 0$ the inner sum equals q , and for $z \neq 0$ we apply the Gauss-sum identity $\sum_{t \in \mathbb{F}_q} \psi(\alpha t^2) = \nu_q(\alpha)G(\psi)$ for $\alpha \in \mathbb{F}_q^\times$ to get

$$S_\xi^2 = q + \sum_{z \in \mathbb{F}_q^\times} \xi^2 \left(\frac{z}{2}, 0 \right) \nu_q(z) G(\psi).$$

Putting $w := z/2$, we obtain

$$S_\xi^2 = q + \nu_q(2) G(\psi) \sum_{w \in \mathbb{F}_q^\times} \nu_q(w) \xi^2(w, 0) = q + \nu_q(2) G(\psi) U_\xi.$$

This completes the proof. \square

The following formula is a generalization of [5, (1.3)].

Proposition 3.16. *We let*

$$T_\xi := \sum_{x \in \mathbb{F}_q} \xi(x^2, 0).$$

Then

$$S_\xi^2 = q + \nu_q(2) G(\psi) (T_{\xi^2} - S_{\xi^2}).$$

Proof. We have

$$T_\xi = 1 + \sum_{x \in \mathbb{F}_q^\times} \xi(x^2, 0) = 1 + \sum_{x \in \mathbb{F}_q^\times} (1 + \nu_q(x)) \xi(x, 0) = S_\xi + \sum_{x \in \mathbb{F}_q^\times} \nu_q(x) \xi(x, 0).$$

Replacing ξ by ξ^2 gives

$$U_\xi = \sum_{x \in \mathbb{F}_q^\times} \nu_q(x) \xi^2(x, 0) = T_{\xi^2} - S_{\xi^2}.$$

Thus Lemma 3.15 implies

$$S_\xi^2 = q + \nu_q(2) G(\psi) U_\xi = q + \nu_q(2) G(\psi) (T_{\xi^2} - S_{\xi^2}).$$

Hence we obtain the claim. \square

We recall $\dim H_c^1(\mathbb{A}^1, \mathcal{L}_\xi) = 2$ by Lemma 3.3.

Corollary 3.17. *Let α_ξ, β_ξ be the eigenvalues of Fr_q^* on $H_c^1(\mathbb{A}^1, \mathcal{L}_\xi)$. Then we have $\alpha_\xi + \beta_\xi = -S_\xi$ and $\alpha_\xi \beta_\xi = q$.*

Proof. By lemma 2.11(1) and Corollary 3.14, we have

$$\begin{aligned} \alpha_\xi + \beta_\xi &= - \sum_{x \in \mathbb{F}_q} \xi(x, 0) = -S_\xi, \\ \alpha_\xi^2 + \beta_\xi^2 &= -q + \nu_q(2) G(\psi) U_\xi. \end{aligned}$$

Hence

$$2\alpha_\xi \beta_\xi = (\alpha_\xi + \beta_\xi)^2 - (\alpha_\xi^2 + \beta_\xi^2) = S_\xi^2 + q - \nu_q(2) G(\psi) U_\xi = 2q,$$

where the last equality follows from Lemma 3.15. \square

We give an explicit formula for the L -polynomial $L_{\overline{C}/\mathbb{F}_q}(T)$.

Theorem 3.18. *For $\xi \in W_{\text{prim}}^\vee$, let $S_\xi = \sum_{x \in \mathbb{F}_q} \xi(x, 0)$. Then we have*

$$L_{\overline{C}/\mathbb{F}_q}(T) = \prod_{\xi \in W_{\text{prim}}^\vee} (1 + S_\xi T + qT^2).$$

Proof. By the proof of Corollary 2.10 and Lemma 3.4, we have

$$H^1(\overline{C}) \xleftarrow{\sim} H_c^1(C) \simeq \bigoplus_{\xi \in W_{\text{prim}}^\vee} H_c^1(\mathbb{A}^1, \mathcal{L}_\xi).$$

Therefore the determinant factors as the product of the determinants on the summands:

$$L_{\overline{C}/\mathbb{F}_q}(T) = \prod_{\xi \in W_{\text{prim}}^\vee} \det(1 - \text{Fr}_q^* T; H_c^1(\mathbb{A}^1, \mathcal{L}_\xi)).$$

For each $\xi \in W_{\text{prim}}^\vee$ let α_ξ, β_ξ be the Frobenius eigenvalues on $H_c^1(\mathbb{A}^1, \mathcal{L}_\xi)$. By Corollary 3.17 they satisfy

$$\alpha_\xi + \beta_\xi = -S_\xi, \quad \alpha_\xi \beta_\xi = q.$$

Hence the local factor for ξ is

$$\det(1 - \text{Fr}_q^* T; H_c^1(\mathbb{A}^1, \mathcal{L}_\xi)) = (1 - \alpha_\xi T)(1 - \beta_\xi T) = 1 + S_\xi T + qT^2.$$

Combining these factors for all $\xi \in W_{\text{prim}}^\vee$ yields the claimed formula. \square

Corollary 3.19. *For $\xi \in W_{\text{prim}}^\vee$, let $0 \leq \theta_\xi \leq \pi$ be the element satisfying*

$$\cos \theta_\xi = -\frac{S_\xi}{2\sqrt{q}}$$

(cf. (3.2)). Then we have

$$\#\overline{C}(\mathbb{F}_{q^n}) = q^n + 1 - q^{n/2} \sum_{\xi \in W_{\text{prim}}^\vee} 2 \cos n\theta_\xi.$$

Proof. Let the notation be as in the proof of Theorem 3.18. We have $\{\alpha_\xi, \beta_\xi\} = \{\sqrt{q}e^{\pm i\theta_\xi}\}$. Hence the claim follows from

$$\#\overline{C}(\mathbb{F}_{q^n}) = q^n + 1 - \sum_{\xi \in W_{\text{prim}}^\vee} (\alpha_\xi^n + \beta_\xi^n) = q^n + 1 - q^{n/2} \sum_{\xi \in W_{\text{prim}}^\vee} 2 \cos n\theta_\xi.$$

\square

Corollary 3.20. *Assume that $q \equiv 2 \pmod{3}$ and that p is odd. Let $\psi \in \mathbb{F}_q^\vee \setminus \{1\}$, and define*

$$S_{\psi,a} := \sum_{x \in \mathbb{F}_q} \psi(x^3 + ax).$$

Then

$$L_{\overline{C}/\mathbb{F}_q}(T) = \prod_{a \in \mathbb{F}_q} (1 + S_{\psi,a} T + qT^2)^{q-1}.$$

Proof. Let $\xi \in W_{\text{prim}}^\vee$. By Lemma 3.5, there exists $(a, b) \in \mathbb{F}_q^\times \times \mathbb{F}_q$ such that

$$S_\xi = \sum_{x \in \mathbb{F}_q} \psi(ax^3 + bx).$$

Since $q \equiv 2 \pmod{3}$, we can take $c \in \mathbb{F}_q^\times$ such that $c^3 = a$. Letting $d := a^{-1}b$, we then have $S_\xi = S_{\psi,d}$. The assertion follows from Theorem 3.18. \square

3.5.1. Katz's monodromy theory and the supersingularity of \overline{C} . Let \mathcal{F} be a smooth $\overline{\mathbb{Q}}_\ell$ -sheaf on $\mathbb{A}_{\mathbb{F}_q}^1$. Let $\rho_{\mathcal{F}}$ denote the ℓ -adic representation of $\pi_1(\mathbb{A}_{\mathbb{F}_q}^1)$ associated to \mathcal{F} . The Zariski closure of the image of $\pi_1(\mathbb{A}_{\mathbb{F}_q}^1)$ (resp. the normal subgroup $\pi_1(\mathbb{A}^1)$) by $\rho_{\mathcal{F}}$ is called the arithmetic (resp. geometric) monodromy group of \mathcal{F} , we denote by $G_{\mathcal{F},\text{arith}}$ (resp. $G_{\mathcal{F},\text{geom}}$). If $G_{\mathcal{F},\text{arith}}$ is finite, \mathcal{F} is said to have finite monodromy.

Choose and fix a square root $\sqrt{q} \in \overline{\mathbb{Q}}_\ell^\times$. Let $\text{Gal}(\mathbb{F}/\mathbb{F}_q)$ be the Galois group. Define the one-dimensional Galois representation $\overline{\mathbb{Q}}_\ell(\tfrac{1}{2})$ by requiring that a geometric Frobenius $\text{Frob}_q \in \text{Gal}(\mathbb{F}/\mathbb{F}_q)$ acts on it as \sqrt{q}^{-1} .

Let l be a prime and let $\psi \in \mathbb{F}_q^\vee \setminus \{1\}$. Let $\pi: \mathbb{A}_{\mathbb{F}_q}^2 \rightarrow \mathbb{A}_{\mathbb{F}_q}^1$, $(x, t) \mapsto t$. We define a local system on $\mathbb{A}_{\mathbb{F}_q}^1$:

$$\mathcal{F}_{\psi, l, q} := R^1\pi_! \mathcal{L}_\psi(x^l + tx)\left(\frac{1}{2}\right)$$

(cf. [22] or [29]).

Proposition 3.21. *Assume $p \neq l$. The curve $\overline{C}_{1,2,l,q}$ is supersingular for all q if and only if $\mathcal{F}_{0\psi \circ \text{Tr}_{q/p}, l, q}$ has finite monodromy for ${}^0\psi \in \mathbb{F}_p^\vee \setminus \{1\}$.*

Proof. We write $C_{l,q}$ for $C_{1,2,l,q}$. Let $t \in \mathbb{F}_q$. By the proper base change theorem,

$$(\mathcal{F}_{\psi, l, q})_{\bar{t}} \simeq H_c^1(\mathbb{A}^1, \mathcal{L}_\psi(x^l + tx))\left(\frac{1}{2}\right).$$

Hence, by the proof of Lemma 3.6,

$$H^1(\overline{C}_{l,q})\left(\frac{1}{2}\right) \simeq \bigoplus_{\psi \in \mathbb{F}_q^\vee \setminus \{1\}} \bigoplus_{t \in \mathbb{F}_q} (\mathcal{F}_{\psi, l, q})_{\bar{t}}.$$

The geometric monodromy group of $\mathcal{F}_{\psi, l, q}$ equals the one of $\mathcal{F}_{0\psi \circ \text{Tr}_{q/p}, l, q}$. Hence the claim follows from [17, Theorem 8.14.4] (cf. [33, Lemma 1 and Corollary 1]). \square

The *period* of a supersingular curve X over \mathbb{F}_q is defined as the smallest positive integer n such that the Frobenius map $\text{Fr}_{q^n}^*$ acts on $H^1(X)$ as multiplication by $q^{n/2}$.

Lemma 3.22. *Assume $q = 5^f$. Then the curve \overline{C} is supersingular with period 60.*

Proof. Let ${}^0\psi \in \mathbb{F}_5^\vee \setminus \{1\}$. By [23, Theorem 17.1(1)], both the arithmetic and geometric monodromy groups of $\mathcal{F}_{0\psi \circ \text{Tr}_{q/5}, 3, q}$ are $\text{SL}_2(\mathbb{F}_5)$. The least common multiple of the orders of elements of $\text{SL}_2(\mathbb{F}_5)$ is 60. Hence the required assertion follows. \square

4. CHARACTERISTIC THREE

We assume $p = 3$ and let $q = 3^f$. Let $\text{Tr}_{q/3}: W_{2,q} \rightarrow W_{2,3}$ denote the trace map.

Lemma 4.1. *Let ${}^0\xi \in W_{2,3}^\vee$ be a faithful character. Let $(a, b) \in \mathbb{F}_q^\times \times \mathbb{F}_q$, and put $c := a^{-1}b^{1/3}$. Then*

$$S_{\xi_{a,b}} = \sum_{x \in \mathbb{F}_q} {}^0\xi(\text{Tr}_{q/3}(x, cx)).$$

Proof. We have

$$S_{\xi_{a,b}} = \sum_{x \in \mathbb{F}_q} \xi_{a,b}(x, 0) = \sum_{x \in \mathbb{F}_q} {}^0\xi(\text{Tr}_{q/3}(ax, bx^3)) = \sum_{x \in \mathbb{F}_q} {}^0\xi(\text{Tr}_{q/3}(ax, b^{1/3}x)).$$

Setting $t = ax$, we obtain the claim. \square

Corollary 4.2. *Let ${}^0\xi \in W_{2,3}^\vee$ be a faithful character. For $c \in \mathbb{F}_q$, set*

$$S_{q,c} := \sum_{x \in \mathbb{F}_q} {}^0\xi(\text{Tr}_{q/3}(x, cx)).$$

Then

$$L_{\overline{C}/\mathbb{F}_q}(T) = \prod_{c \in \mathbb{F}_q} (1 + S_{q,c}T + qT^2)^{q-1}.$$

Proof. By Theorem 3.18 and Lemmas 2.7 and 4.1, we obtain

$$\begin{aligned} L_{\overline{C}/\mathbb{F}_q}(T) &= \prod_{\xi \in W_{\text{prim}}^\vee} (1 + S_\xi T + qT^2) \\ &= \prod_{(a,b) \in \mathbb{F}_q^\times \times \mathbb{F}_q} (1 + S_{\xi_{a,b}} T + qT^2) = \prod_{c \in \mathbb{F}_q} (1 + S_{q,c} T + qT^2)^{q-1}. \end{aligned}$$

□

Lemma 4.3. *Let $c \in \mathbb{F}_q^\times$. Let ${}^0\xi \in W_{2,3}^\vee$ be a faithful character and put $\xi := {}^0\xi \circ \text{Tr}_{q/3}$. Let $\psi(x) := \xi(0, x)$ for $x \in \mathbb{F}_q$, and define $G(\psi) := \sum_{t \in \mathbb{F}_q} \psi(t^2)$. Then*

$$S_{q,0} S_{q,c} = G(\psi) \sum_{x \in \mathbb{F}_q^\times} \xi(x, -(c+1)x - c^2 x^{-1}) \nu_q(x).$$

Proof. We compute

$$\begin{aligned} S_{q,0} S_{q,c} &= \sum_{x \in \mathbb{F}_q} \xi(x, 0) \sum_{y \in \mathbb{F}_q} \xi(y, cy) = \sum_{x,y} \xi(x+y, cy - xy(x+y)) \\ &= \sum_{x,z} \xi(z, c(z-x) - x(z-x)z) \quad (z := x+y) \\ &= \sum_x \xi(0, -cx) + \sum_x \sum_{z \neq 0} \xi(z, -zx^2 - (z^2 + c)x + cz), \end{aligned}$$

where we separated the summand with $z = 0$. The first term vanishes because $\psi = \xi(0, \cdot)$ is a nontrivial additive character and $c \neq 0$: $\sum_x \xi(0, -cx) = \sum_x \psi(-cx) = 0$.

For $z \neq 0$ complete the square in x :

$$zx^2 - (z^2 - c)x + cz = z \left(x - \frac{z^2 + c}{2z} \right)^2 - \frac{(z^2 - c)^2}{4z}.$$

Putting $x_1 := x - (2z)^{-1}(z^2 - c)$ (which is a bijection of \mathbb{F}_q), we get

$$\sum_{x \in \mathbb{F}_q} \xi(z, zx^2 - (z^2 + c)x + cz) = \xi \left(z, -\frac{(z^2 - c)^2}{4z} \right) \sum_{x_1 \in \mathbb{F}_q} \psi(zx_1^2).$$

Using the Gauss-sum identity $\sum_{t \in \mathbb{F}_q} \psi(\alpha t^2) = \nu_q(\alpha) G(\psi)$ for $\alpha \in \mathbb{F}_q^\times$, we obtain for $z \neq 0$:

$$\sum_{x \in \mathbb{F}_q} \xi(z, zx^2 - (z^2 + c)x + cz) = \xi \left(z, -\frac{(z^2 - c)^2}{4z} \right) \nu_q(z) G(\psi).$$

Hence

$$S_{q,0} S_{q,c} = G(\psi) \sum_{z \neq 0} \xi \left(z, -\frac{(z^2 - c)^2}{4z} \right) \nu_q(z).$$

Finally note

$$\text{Tr}_{q/3} \left(\frac{(z^2 - c)^2}{z} \right) = \text{Tr}_{q/3} \left(z^3 + cz + \frac{c^2}{z} \right) = \text{Tr}_{q/3} \left((c+1)z + \frac{c^2}{z} \right),$$

so renaming $z \mapsto x$ yields the claimed formula. □

4.1. Valuation of the Gauss-Heilbronn sum. Let $\zeta_9 = e^{2\pi i/9}$. Let \mathfrak{p} denote the prime of $\mathbb{Q}(\zeta_9)$ lying over (3). Let $v_{\mathfrak{p}}(-)$ denote the valuation of the completion $\mathbb{Q}(\zeta_9)$ at \mathfrak{p} normalized such that $v_{\mathfrak{p}}(3) = 1$. For $\xi \in W_{\text{prim}}^{\vee}$, we study $v_{\mathfrak{p}}(S_{\xi})$ using supersingular curves.

Lemma 4.4. *Let $(a, b) \in \mathbb{F}_q^{\times} \times \mathbb{F}_q$ and let ${}^0\psi \in \mathbb{F}_3^{\vee} \setminus \{1\}$. Put $\psi := {}^0\psi \circ \text{Tr}_{q/3}$ and*

$$S(a, b, \psi) := \sum_{x \in \mathbb{F}_q^{\times}} \nu_q(x) \psi(ax^3 + ax(x+b)b).$$

Then $v_{\mathfrak{p}}(S(a, b, \psi)) \geq f/2$.

Proof. First assume $b = 0$. By $p = 3$ and $\psi := {}^0\psi \circ \text{Tr}_{q/3}$, we have $\psi(ax^3) = \psi(a^{1/3}x)$. Hence

$$S(a, 0, \psi) = \sum_{x \in \mathbb{F}_q^{\times}} \nu_q(x) \psi(ax^3) = \sum_{x \in \mathbb{F}_q^{\times}} \nu_q(x) \psi(a^{1/3}x) = G(\psi_{a^{1/3}}) = \nu_q(a^{1/3}) G(\psi),$$

whose \mathfrak{p} -adic valuation is $f/2$.

Now assume $b \neq 0$. Writing

$$c := ab \in \mathbb{F}_q^{\times}, \quad d := ab^2 + a^{1/3},$$

we have

$$S(a, b, \psi) = \sum_{x \in \mathbb{F}_q^{\times}} \nu_q(x) \psi(cx^2 + dx).$$

Let $D_{c,d}$ be the affine curve defined by

$$y^3 - y = cx^4 + dx^2$$

over \mathbb{F}_q . The group $\{\pm 1\} \times \mathbb{F}_3$ acts on $D_{c,d}$ via

$$(\iota, \zeta) : (x, y) \mapsto (\iota x, y + \zeta) \quad \text{for } (\iota, \zeta) \in \{\pm 1\} \times \mathbb{F}_3.$$

Let $D_{c,d}^0 \subset D_{c,d}$ be the open subscheme defined by $x \neq 0$. The morphism

$$D_{c,d}^0 \rightarrow \mathbb{G}_{\text{m}, \mathbb{F}_q}, \quad (x, y) \mapsto x^2,$$

is a finite étale Galois covering with Galois group $\{\pm 1\} \times \mathbb{F}_3$. Let ν_0 be the nontrivial character of $\{\pm 1\}$, and put $\mathbb{G}_{\text{m}} = \text{Spec } \mathbb{F}[t^{\pm 1}]$. Then

$$H_c^1(D_{c,d})[\nu_0 \boxtimes {}^0\psi] \xleftarrow{\sim} H_c^1(D_{c,d}^0)[\nu_0 \boxtimes {}^0\psi] \simeq H_c^1(\mathbb{G}_{\text{m}}, \mathcal{K}_{\nu_0}(t) \otimes \mathcal{L}_{0\psi}(ct^2 + dt)),$$

where $\mathcal{K}_{\nu_0}(t)$ is the Kummer sheaf defined by the Kummer covering $y^2 = x$ and ν_0 . The sheaf $\mathcal{K}_{\nu_0}(t) \otimes \mathcal{L}_{0\psi}(ct^2 + dt)$ on $\mathbb{G}_{\text{m}, \mathbb{F}_q}$ is tamely ramified at 0 and wildly ramified at ∞ with Swan conductor 2. By the Grothendieck-Ogg-Shafarevich formula,

$$\dim H_c^1(\mathbb{G}_{\text{m}}, \mathcal{K}_{\nu_0}(t) \otimes \mathcal{L}_{0\psi}(ct^2 + dt)) = 2.$$

Since the defining equation of $D_{c,d}$ is $y^3 - y = x(cx^3 + dx)$ and the polynomial $cx^3 + dx$ is \mathbb{F}_3 -linearized, the smooth compactification $\overline{D}_{c,d}$ is supersingular by [13]. Hence all eigenvalues of Fr_q^* on $H^1(\overline{D}_{c,d})$ are of the form $\kappa \sqrt{q}$ with κ a root of unity.

Let α_1, α_2 be the eigenvalues of Fr_q^* on $H_c^1(D_{c,d})[\nu_0 \boxtimes {}^0\psi]$. Then $\alpha_i = \kappa_i \sqrt{q}$ with κ_i a root of unity, and therefore $v_{\mathfrak{p}}(\alpha_i) = f/2$. Since

$$S(a, b, \psi) = -\alpha_1 - \alpha_2,$$

we obtain

$$v_{\mathfrak{p}}(S(a, b, \psi)) = \min\{v_{\mathfrak{p}}(\alpha_1), v_{\mathfrak{p}}(\alpha_2)\} \geq f/2.$$

Hence we have proved the claim. \square

Proposition 4.5. *Let $\xi \in W_{\text{prim}}^\vee$. Then we have $v_{\mathfrak{p}}(S_\xi U_\xi) = f/2$.*

Proof. We write ν for ν_q . By $p = 3$, we have $\xi^2(x, 0) = \xi(-x, x^3)$. Thus

$$S_\xi U_\xi = \sum_{\substack{x \neq 0 \\ y \in \mathbb{F}_q}} \nu(x) \xi(-x, x^3) \xi(y, 0) = \sum_{\substack{x \neq 0 \\ y \\ y}} \nu(x) \xi(-x + y, x^3 + xy(-x + y)).$$

Put $z = -x + y$. Then $y = x + z$, and we obtain

$$S_\xi U_\xi = \sum_{x \neq 0} \sum_z \nu(x) \xi(z, x^3 + x(x + z)z).$$

We use the notation $\psi = {}^0\psi \circ \text{Tr}_{q/3}$ with ${}^0\psi \in \mathbb{F}_3^\vee \setminus \{1\}$, so that $\xi(x_0, x_1) = \xi(x_0, 0)\psi(ax_1)$ with $a \in \mathbb{F}_q^\times$. We get

$$S_\xi U_\xi = \sum_z \xi(z, 0) \sum_{x \neq 0} \nu(x) \psi(ax^3 + ax(x + z)z).$$

In the notation of Lemma 4.4,

$$S_\xi U_\xi = \sum_z \xi(z, 0) S(a, z, \psi).$$

For $x, y \in \mathbb{Q}_3(\zeta_9)$ and $r \in \mathbb{Q}$, we write $x \equiv y \pmod{r+}$ if $v_{\mathfrak{p}}(x - y) > r$. By Lemma 4.4 and $v_{\mathfrak{p}}(\xi(z, 0) - 1) > 0$ for any $z \in \mathbb{F}_q$, we have

$$(4.1) \quad S_\xi U_\xi = \sum_z \xi(z, 0) S(a, z, \psi) \equiv \sum_z S(a, z, \psi) \pmod{f/2+}.$$

Completing the square in z and using $p = 3$, we have

$$ax^3 + ax(x + z)z = ax(z - x)^2.$$

Hence

$$\sum_z S(a, z, \psi) = \sum_{x \neq 0} \nu(x) \sum_z \psi(ax(z - x)^2).$$

For fixed $x \neq 0$, the inner sum is a quadratic Gauss sum:

$$\sum_z \psi(ax(z - x)^2) = \nu(ax) G(\psi) \quad \text{for } a \in \mathbb{F}_q^\times.$$

Therefore,

$$\sum_z S(a, z, \psi) = \sum_{x \neq 0} \nu(x) \nu(ax) G(\psi) = \nu(a) \sum_{x \neq 0} G(\psi) = \nu(a)(q - 1)G(\psi).$$

Finally, since $v_{\mathfrak{p}}(G(\psi)) = f/2$ and (4.1), we obtain

$$v_{\mathfrak{p}}(S_\xi U_\xi) = v_{\mathfrak{p}}(\nu(a)(q - 1)G(\psi)) = f/2.$$

We obtain the claim. \square

We show one of our main theorems.

Corollary 4.6. *Let $\xi \in W_{\text{prim}}^\vee$. Then we have $v_{\mathfrak{p}}(S_\xi) = f/3$ and $v_{\mathfrak{p}}(U_\xi) = f/6$.*

Proof. By Lemma 3.15,

$$(4.2) \quad S_\xi^2 = q - \nu_q(2) G(\psi) U_\xi.$$

If $v_p(\nu_q(2) G(\psi) U_\xi) \geq f$, we have $v_p(S_\xi) \geq f/2$ and $v_p(U_\xi) \geq f/2$ by $v_p(G(\psi)) = f/2$. By Proposition 4.5, we have $v_p(U_\xi) = (f/2) - v_p(S_\xi) \leq 0$, which contradicts $v_p(U_\xi) \geq f/2$. Hence $v_p(\nu_q(2) G(\psi) U_\xi) < f$ and again by (4.2),

$$v_p(S_\xi^2) = v_p(\nu_q(2) G(\psi) U_\xi) = (f/2) + v_p(U_\xi) = f - v_p(S_\xi),$$

where we use Proposition 4.5 at the last equality. Thus the claim follows. \square

Corollary 4.7. *Let $\xi \in W_{\text{prim}}^\vee$ and let*

$$T_\xi = \sum_{x \in \mathbb{F}_q} \xi(x^2, 0), \quad V_{\xi, c} := \sum_{x \in \mathbb{F}_q^\times} \xi(x, (2c-1)x - c^2 x^{-1}) \nu_q(x) \quad \text{for } c \in \mathbb{F}_q^\times.$$

Then we have $v_p(T_\xi) = v_p(V_{\xi, c}) = f/6$ for any $c \in \mathbb{F}_q^\times$.

Proof. The assertion for T_ξ follows from Proposition 3.16 and Corollary 4.6. The assertion for $V_{\xi, c}$ follows from Lemma 4.3 and Corollary 4.6. \square

For $\alpha, \beta \in \overline{\mathbb{Q}_\ell}^\times$, if α/β is a root of unity, we write $\alpha \equiv \beta \pmod{\mu_\infty(\overline{\mathbb{Q}_\ell})}$. The following lemma is shown in a more general setting by a different method in Corollary A.2.

Lemma 4.8. *Let $\xi \in W_{\text{prim}}^\vee$. Let ν_0 be a nontrivial character of $\{\pm 1\}$ and let $\mathcal{K}_{\nu_0}(x)$ be the Kummer sheaf on $\mathbb{G}_{\text{m}, \mathbb{F}_q}$ defined by $y^2 = x$ and ν_0 . Then we have*

$$\det(-\text{Fr}_q^*; H_c^1(\mathbb{G}_{\text{m}}, \mathcal{K}_{\nu_0}(x) \otimes \mathcal{L}_\xi(x, 0))) \equiv q G(\psi) \pmod{\mu_\infty(\overline{\mathbb{Q}_\ell})}.$$

Proof. We simply write $\mathcal{L} := \mathcal{K}_{\nu_0}(x) \otimes \mathcal{L}_\xi(x, 0)$. Let D be the projective line over \mathbb{F}_q and we regard $\mathbb{G}_{\text{m}, \mathbb{F}_q} = \text{Spec } \mathbb{F}_q[x^{\pm 1}]$ as the open subscheme of D . Let $j: \mathbb{G}_{\text{m}, \mathbb{F}_q} \rightarrow D$ be the open immersion. We freely use the notation in [29]. We recall $H_c^i(\mathbb{G}_{\text{m}}, \mathcal{L}) = 0$ for $i \neq 1$. By the product formula [29, Théorème (3.2.1.1)] for $X = D$, $K = j_! \mathcal{L}$, we have

$$\det(-\text{Fr}_q^*; H_c^1(\mathbb{G}_{\text{m}}, \mathcal{L})) = q \epsilon(D_{(0)}, \mathcal{L}|_{\eta_0}, d\pi_0) \epsilon\left(D_{(\infty)}, \mathcal{L}|_{\eta_\infty}, -\frac{d\pi_\infty}{\pi_\infty^2}\right),$$

where $\pi_0 = x$ and $\pi_\infty = x^{-1}$ (cf. [29, (3.1.1.5)]). By [29, (3.1.5.6) and (3.5.3.1)], we have

$$\epsilon(D_{(0)}, \mathcal{L}|_{\eta_0}, d\pi_0) \equiv \epsilon(D_{(0)}, \mathcal{K}_{\nu_0}(x)|_{\eta_0}, d\pi_0) \equiv G(\psi) \pmod{\mu_\infty(\overline{\mathbb{Q}_\ell})}.$$

We recall that the Swan conductor exponent of \mathcal{L} is 3 by Lemma 2.8. We also have

$$\begin{aligned} \epsilon\left(D_{(\infty)}, \mathcal{L}|_{\eta_\infty}, -\frac{d\pi_\infty}{\pi_\infty^2}\right) &\equiv q^{-2} \epsilon(D_{(\infty)}, \mathcal{L}|_{\eta_\infty}, d\pi_\infty) \pmod{\mu_\infty(\overline{\mathbb{Q}_\ell})} \\ &= q^{-2} \epsilon(\chi_{\mathcal{L}}, \Psi_{d\pi_\infty}) \\ &= q^{-2} \int_{\pi_\infty^{-4} \mathcal{O}_\infty^\times} \chi_{\mathcal{L}}^{-1}(z) \Psi_{d\pi_\infty}(z) dz, \end{aligned}$$

where we use [29, (3.1.5.5)], [29, Théorème (3.1.5.4)(v)] and [29, (3.1.3.2)], respectively. Here $\chi_{\mathcal{L}}$ is a character of $\mathbb{F}_q((\pi_\infty))^\times$ induced by $\mathcal{L}|_{\eta_\infty}$ via the local class field theory, $\mathcal{O}_\infty := \mathbb{F}_q[[\pi_\infty]]$, and $\int_{\mathcal{O}_\infty} dz = 1$. Applying [1, Proposition 8.7(ii)] to the local field $K = \mathbb{F}_q((\pi_\infty))$ with $\chi = \chi_{\mathcal{L}}$ and $\psi = \Psi_{d\pi_\infty}$, we note that by Lemma 2.8 the character

$\chi_{\mathcal{L}}$ has Swan conductor exponent 3. In the notation of loc. cit. we therefore take $\text{ord}(\psi) = 0$, $\text{ord}(\beta) = 1$ and $\text{ord}(c) = -4$, so that the relevant integral is over $\pi_{\infty}^{-4}\mathcal{O}_{\infty}^{\times}$. With the normalization $\int_{\mathcal{O}_{\infty}} dz = 1$, Proposition 8.7(ii) gives

$$\epsilon(\chi_{\mathcal{L}}, \Psi_{d\pi_{\infty}}) \equiv q^2 \pmod{\mu_{\infty}(\overline{\mathbb{Q}}_{\ell})}.$$

□

Now we prove one of our main theorems.

Theorem 4.9. *We simply write D for $C_{2,2,3,q}$ and assume $q = 3^f$.*

(1) *Let ν_0 be a nontrivial character of $\{\pm 1\}$. Then*

$$H^1(\overline{D}) \simeq \bigoplus_{\xi \in W^{\vee}} H_{\text{c}}^1(\mathbb{A}^1, \mathcal{L}_{\xi}(x, 0)) \oplus H_{\text{c}}^1(\mathbb{G}_{\text{m}}, \mathcal{K}_{\nu_0}(x) \otimes \mathcal{L}_{\psi}(x, 0)).$$

Moreover, $g(\overline{D}) = (q-1)(5q+1)/2$.

(2) *The multiset of slopes of $H_{\text{c}}^1(\mathbb{G}_{\text{m}}, \mathcal{K}_{\nu_0}(x) \otimes \mathcal{L}_{\psi}(x, 0))$ with respect to Fr_q^* is*

$$\begin{cases} \left\{ \frac{1}{2} \right\} & \text{if } \xi \notin W_{\text{prim}}^{\vee} \text{ and } \xi \neq 1, \\ \left\{ \frac{1}{2}, \frac{1}{6}, \frac{5}{6} \right\} & \text{if } \xi \in W_{\text{prim}}^{\vee}. \end{cases}$$

(3) *The multiset of slopes of \overline{D} with respect to Fr_q^* is*

$$\left\{ \frac{1}{2}, \frac{1}{3}, \frac{2}{3}, \frac{1}{6}, \frac{5}{6} \right\}.$$

Proof. We prove (1). By the definition of D and the proof of Corollary 2.10, we have

$$H^1(\overline{D}) \simeq H_{\text{c}}^1(D) \simeq \bigoplus_{\xi \in W^{\vee}} H_{\text{c}}^1(\mathbb{A}^1, \mathcal{L}_{\xi}(x^2, 0)).$$

For $\mathcal{F} \in \{\mathcal{L}_{\xi}(x, 0), \mathcal{L}_{\xi}(x^2, 0)\}$, we have a short exact sequence

$$0 \rightarrow \overline{\mathbb{Q}}_{\ell} \rightarrow H_{\text{c}}^1(\mathbb{G}_{\text{m}}, \mathcal{F}) \rightarrow H_{\text{c}}^1(\mathbb{A}^1, \mathcal{F}) \rightarrow 0.$$

We simply write $\mathcal{L}_{\nu_0, \xi}$ for $\mathcal{K}_{\nu_0}(x) \otimes \mathcal{L}_{\xi}(x, 0)$. By the projection formula,

$$H_{\text{c}}^1(\mathbb{G}_{\text{m}}, \mathcal{L}_{\xi}(x^2, 0)) \simeq H_{\text{c}}^1(\mathbb{G}_{\text{m}}, \mathcal{L}_{\xi}(x, 0)) \oplus H_{\text{c}}^1(\mathbb{G}_{\text{m}}, \mathcal{L}_{\nu_0, \xi}).$$

The above short exact sequence then implies

$$H_{\text{c}}^1(\mathbb{A}^1, \mathcal{L}_{\xi}(x^2, 0)) \simeq H_{\text{c}}^1(\mathbb{A}^1, \mathcal{L}_{\xi}(x, 0)) \oplus H_{\text{c}}^1(\mathbb{G}_{\text{m}}, \mathcal{L}_{\nu_0, \xi}).$$

Thus the first claim follows. The genus formula follows from Lemma 2.10 for $l = p = 3$ and $n = 2$.

We now prove (2). Assume $\xi \notin W_{\text{prim}}^{\vee}$. If $\xi = 1$, then

$$H_{\text{c}}^1(\mathbb{G}_{\text{m}}, \mathcal{L}_{\nu_0, \xi}) \simeq H_{\text{c}}^1(\mathbb{G}_{\text{m}}, \mathcal{K}_{\nu_0}(x)) = 0.$$

Thus assume $\xi \neq 1$. It is well known that the Frobenius slope of $H_{\text{c}}^1(\mathbb{G}_{\text{m}}, \mathcal{L}_{\nu_0, \xi})$ with respect to Fr_q^* is $1/2$, since $\mathcal{L}_{\xi}(x, 0) \simeq \mathcal{L}_{\psi}(x)$ for some $\psi \in \mathbb{F}_q^{\vee} \setminus \{1\}$.

Assume now that $\xi \in W_{\text{prim}}^{\vee}$. Since $\text{Swan}_{\infty}(\mathcal{L}_{\nu_0, \xi}) = 3$ by Lemma 2.8,

$$\dim H_{\text{c}}^1(\mathbb{G}_{\text{m}}, \mathcal{L}_{\nu_0, \xi}) = 3.$$

Lemma 4.8 yields

$$(4.3) \quad v_{\mathfrak{p}} \left(\det(-\text{Fr}_q^*; H_c^1(\mathbb{G}_m, \mathcal{L}_{\nu_0, \xi})) \right) = \frac{3f}{2}.$$

Let α, β, γ be the Frobenius eigenvalues of Fr_q^* on $H_c^1(\mathbb{G}_m, \mathcal{L}_{\nu_0, \xi})$, ordered so that $a := v_{\mathfrak{p}}(\alpha) \leq b := v_{\mathfrak{p}}(\beta) \leq c := v_{\mathfrak{p}}(\gamma)$. Since $v_{\mathfrak{p}}(U_{\xi}) = f/6$ by Corollary 4.6, we have $a \leq v_{\mathfrak{p}}(\alpha + \beta + \gamma) = f/6$.

By Poincaré duality and $\mathcal{L}_{\nu_0, \xi}^{\vee} \simeq \mathcal{L}_{\nu_0, \xi^{-1}}$,

$$(4.4) \quad H_c^1(\mathbb{G}_m, \mathcal{L}_{\nu_0, \xi})^{\vee}(-1) \simeq H^1(\mathbb{G}_m, \mathcal{L}_{\nu_0, \xi^{-1}}) \xleftarrow{\sim} H_c^1(\mathbb{G}_m, \mathcal{L}_{\nu_0, \xi^{-1}}).$$

Thus, since $v_{\mathfrak{p}}(U_{\xi^{-1}}) = f/6$,

$$(4.5) \quad v_{\mathfrak{p}}(q\alpha^{-1} + q\beta^{-1} + q\gamma^{-1}) = \frac{f}{6}.$$

From (4.3), we have $a + b + c = 3f/2$.

Assume $a < f/6$. Then $a = b$, because $v_{\mathfrak{p}}(\alpha + \beta + \gamma) = f/6$. Hence both $v_{\mathfrak{p}}(q\alpha^{-1})$ and $v_{\mathfrak{p}}(q\beta^{-1})$ exceed $5f/6$. Then $c = (3f/2) - 2a > 5f/6$, giving $v_{\mathfrak{p}}(q\gamma^{-1}) = f - c < f/6$, which contradicts (4.5). Thus $a = f/6$.

Replacing ξ by ξ^{-1} and using (4.4), we obtain $\min\{v_{\mathfrak{p}}(q\alpha^{-1}), v_{\mathfrak{p}}(q\beta^{-1}), v_{\mathfrak{p}}(q\gamma^{-1})\} = v_{\mathfrak{p}}(q\gamma^{-1}) = f/6$, hence $c = 5f/6$. Since $a + b + c = 3f/2$, we get $b = f/2$.

This proves the claim.

Finally, (3) follows from (1) and (2). \square

5. COVERING OF THE CURVE

We simply write C_n for $C_{1,n,p,q}$. Let $P(x) \in \mathbb{F}_q[x]$ be an \mathbb{F}_p -linearized separable polynomial. We define $C_{P,n}$ by the cartesian diagram

$$\begin{array}{ccc} C_{P,n} & \xrightarrow{\varphi_n} & C_n \\ \downarrow \text{pr}'_1 & & \downarrow \text{pr}_1 \\ \mathbb{A}_{\mathbb{F}_q}^1 & \xrightarrow{x \mapsto P(x)} & \mathbb{A}_{\mathbb{F}_q}^1, \end{array}$$

where pr_1 is given by $(x_0, \dots, x_{n-1}) \mapsto x_0$. Our main aim in this section is to determine the L -polynomial of $\overline{C}_{P,n}$ in the case $n = 2$ and $p = 3$.

We note that $C_{P,1}$ is isomorphic to $\mathbb{A}_{\mathbb{F}_q}^1$.

We define an algebraic group $\mathbb{W}_{n,q}^P$ by the cartesian diagram

$$\begin{array}{ccc} \mathbb{W}_{n,q}^P & \xrightarrow{\phi_n} & \mathbb{W}_{n,q} \\ \downarrow & & \downarrow \text{pr}_1 \\ \mathbb{G}_{a, \mathbb{F}_q} & \xrightarrow{x \mapsto P(x)} & \mathbb{G}_{a, \mathbb{F}_q}, \end{array}$$

where pr_1 is given by $(x_0, \dots, x_{n-1}) \mapsto x_0$. We consider

$$\phi_n: \mathbb{W}_{n,q}^P(\mathbb{F}) \rightarrow \mathbb{W}_{n,q}(\mathbb{F})$$

and we define

$$W_{P,n} := \phi_n^{-1}(W_{n,q}).$$

The morphisms $C_{P,n} \xrightarrow{\varphi_n} C_n \subset \mathbb{W}_{n,q}$ and $\text{pr}'_1: C_{P,n} \rightarrow \mathbb{G}_{a,\mathbb{F}_q}$ induce a morphism $C_{P,n} \hookrightarrow \mathbb{W}_{n,q}^P$. By $\mathbb{W}_{n,q}^P = \mathbb{A}_{\mathbb{F}_q}^1 \times_{P, \mathbb{A}_{\mathbb{F}_q}^1} \mathbb{W}_{n,q}$, we have

$$\mathbb{W}_{n,q}^P \times_{\phi_n, \mathbb{W}_{n,q}} C_n \simeq (\mathbb{A}_{\mathbb{F}_q}^1 \times_{P, \mathbb{A}_{\mathbb{F}_q}^1} \mathbb{W}_{n,q}) \times_{\mathbb{W}_{n,q}} C_n \simeq \mathbb{A}_{\mathbb{F}_q}^1 \times_{P, \mathbb{A}_{\mathbb{F}_q}^1} C_n = C_{P,n}.$$

Hence, for $x \in \mathbb{W}_{n,q,\mathbb{F}}$,

$$x \in C_{P,n,\mathbb{F}} \iff \phi_n(x) \in C_{P,\mathbb{F}}.$$

For $x \in C_{P,n,\mathbb{F}}$ and $b \in W_{P,n}$, we have $x + b \in C_{P,n,\mathbb{F}}$ since $\phi_n(x + b) = \phi_n(x) + \phi_n(b) \in C_{P,\mathbb{F}}$ by $\phi_n(b) \in W_{n,q}$.

Lemma 5.1. *Let $K_P := \{x \in \mathbb{F} \mid P(x) = 0\}$. The group $W_{P,n}$ is isomorphic to a product group $K_P \times W_{n,q}$.*

Proof. We have a short exact sequence

$$0 \rightarrow K_P \rightarrow W_{P,n} \xrightarrow{\phi_n} W_{n,q} \rightarrow 0.$$

Let $x = (x_0, \dots, x_{n-1}) \in W_{P,n} \subset \mathbb{F}^n$. Then $\phi_n(x) = (P(x_0), x_1, \dots, x_{n-1}) \in W_{n,q}$. Let $x \in W_{P,n}$. We show $p^n x = 0$. We have $\phi_n(p^n x) = p^n \phi_n(x) = 0$ by Lemma 2.9. Hence $p^n x \in \text{Ker } \phi_n$. Then we can write $(y_0, 0's) = p^n x$. By $p^n x = (0, *, \dots, *)$, we know $y_0 = 0$. Hence $p^n x = 0$. Hence the short exact sequence is split. \square

We have a finite étale Galois covering

$$\pi_n: C_n \rightarrow \mathbb{A}_{\mathbb{F}_q}^1, \quad (x_0, \dots, x_{n-1}) \mapsto x_0^q - x_0,$$

whose Galois group is $W_{n,q} = \mathbb{W}_n(\mathbb{F}_q)$.

We take $\psi \in \mathbb{F}_q^\vee \setminus \{1\}$. Let r be the minimal positive integer such that $K_P \subset \mathbb{F}_{q^r}$. Let $Q(x) \in \mathbb{F}_q[x]$ be an \mathbb{F}_p -linearized polynomial such that $Q(P(x)) = P(Q(x)) = x^{q^r} - x$. We have a surjection

$$\mathbb{F}_{q^r} \twoheadrightarrow K_P, \quad x \mapsto Q(x).$$

Taking the dual, we have

$$K_P^\vee \hookrightarrow \mathbb{F}_{q^r}^\vee.$$

We have the bijection

$$\Psi: \mathbb{F}_{q^r} \rightarrow \mathbb{F}_{q^r}^\vee, \quad a \mapsto (x \mapsto \psi \circ \text{Tr}_{q^r/q}(ax)).$$

We define $K_{P,\psi}^0 \subset \mathbb{F}_{q^r}$ to be $\Psi^{-1}(K_P^\vee)$. We take a minimal extension $\mathbb{F}_{q^s}/\mathbb{F}_{q^r}$ such that, for any $a \in K_{P,\psi}^0$, there exist solutions $x \in \mathbb{F}_{q^s}$ such that $x^q - x = a$. We consider a map $g: \mathbb{F}_{q^s} \rightarrow \mathbb{F}_{q^s}$, $x \mapsto x^{1/q} - x$ and choose a subset $K_{P,\psi} \subset \mathbb{F}_{q^s}$ such that

$$g|_{K_{P,\psi}}: K_{P,\psi} \xrightarrow{\sim} K_{P,\psi}^0.$$

The morphism

$$\pi_n \circ \varphi_n: C_{P,n,\mathbb{F}_{q^s}} \rightarrow \mathbb{A}_{\mathbb{F}_{q^s}}^1, \quad (x_0, \dots, x_{n-1}) \mapsto P(x_0)^q - P(x_0)$$

is a finite étale Galois covering whose Galois group is $W_{P,n}$.

Proposition 5.2. *We have*

$$\pi_{n*} \varphi_{n*} \overline{\mathbb{Q}}_\ell \simeq \bigoplus_{\xi \in W_{n,q}^\vee, \alpha \in K_{P,\psi}} \mathcal{L}_\psi(\alpha x) \otimes \mathcal{L}_\xi(x, 0's).$$

Proof. We consider the commutative diagram

$$\begin{array}{ccccc}
 C_{P,n,\mathbb{F}} & \xrightarrow{\varphi_n} & C_{n,\mathbb{F}} & \xrightarrow{\pi_n} & \mathbb{A}^1 \\
 \downarrow \text{pr}'_1 & & \downarrow \text{pr}_1 & \nearrow f_0 & \\
 \mathbb{A}^1 & \xrightarrow{x \mapsto P(x)} & \mathbb{A}^1 & &
 \end{array}$$

where f_0 is given by $x \mapsto x^q - x$. Here, the left commutative diagram is cartesian. The morphism $\mathbb{A}^1 \rightarrow \mathbb{A}^1$, $x \mapsto x^{q^r} - x$ factors through

$$\mathbb{A}^1 \xrightarrow{x \mapsto Q(x)} \mathbb{A}^1 \xrightarrow{x \mapsto P(x)} \mathbb{A}^1.$$

The group K_P is regarded as a quotient of \mathbb{F}_{q^r} by $x \mapsto Q(x)$. For any $\psi' \in K_P^\vee \subset \mathbb{F}_{q^r}^\vee$, we have $P^* \mathcal{L}_{\psi'}(x) \simeq \overline{\mathbb{Q}}_\ell$. Thus we obtain a nonzero morphism $\mathcal{L}_{\psi'}(x) \rightarrow P_* \overline{\mathbb{Q}}_\ell$. Thus we obtain an injection $\bigoplus_{\psi' \in K_P^\vee} \mathcal{L}_{\psi'}(x) \rightarrow P_* \overline{\mathbb{Q}}_\ell$. By comparing the ranks, we obtain an isomorphism $\bigoplus_{\psi' \in K_P^\vee} \mathcal{L}_{\psi'}(x) \simeq P_* \overline{\mathbb{Q}}_\ell$. By the proper base change theorem,

$$\varphi_{n*} \overline{\mathbb{Q}}_\ell \simeq \text{pr}_1^* P_* \overline{\mathbb{Q}}_\ell \simeq \bigoplus_{\psi' \in K_P^\vee} \text{pr}_1^* \mathcal{L}_{\psi'}(x) \simeq \bigoplus_{a \in K_{P,\psi}^0} \text{pr}_1^* \mathcal{L}_\psi(ax).$$

Let $H \subset W_{n,q}$ denote the subgroup consisting of the elements $(0, a_1, \dots, a_{n-1})$. Let $\mathcal{L}_{H,\psi'}$ denote the sheaf defined by the H -covering φ_n and $\psi' \in H^\vee$. Applying π_{n*} and using the projection formula yield

$$\begin{aligned}
 \pi_{n*} \varphi_{n*} \overline{\mathbb{Q}}_\ell &\simeq \bigoplus_{a \in K_{P,\psi}^0} f_{0*} \text{pr}_{1*} \text{pr}_1^* \mathcal{L}_\psi(ax) \\
 &\simeq \bigoplus_{a \in K_{P,\psi}^0} f_{0*} (\mathcal{L}_\psi(ax) \otimes \text{pr}_{1*} \overline{\mathbb{Q}}_\ell) \\
 &\simeq \bigoplus_{a \in K_{P,\psi}^0} \bigoplus_{\psi' \in H^\vee} f_{0*} (\mathcal{L}_\psi(ax) \otimes \mathcal{L}_{H,\psi'}).
 \end{aligned}$$

We take $\alpha \in K_{P,\psi}$ such that $\alpha^{1/q} - \alpha = a$. Then $\mathcal{L}_\psi(ax) \simeq \mathcal{L}_\psi(\alpha(x^q - x)) \simeq f_0^* \mathcal{L}_\psi(\alpha x)$. Thus

$$\begin{aligned}
 f_{0*} (\mathcal{L}_\psi(ax) \otimes \mathcal{L}_{H,\psi'}) &\simeq f_{0*} (f_0^* \mathcal{L}_\psi(\alpha x) \otimes \mathcal{L}_{H,\psi'}) \\
 &\simeq \mathcal{L}_\psi(\alpha x) \otimes f_{0*} \mathcal{L}_{H,\psi'} \\
 &\simeq \bigoplus_{\xi \in W_{\psi'}^\vee} \mathcal{L}_\psi(\alpha x) \otimes \mathcal{L}_\xi(x, 0's),
 \end{aligned}$$

where $W_{\psi'}^\vee := \{\xi \in W_{n,q}^\vee \mid \xi|_H = \psi'\}$.

Hence we obtain

$$\pi_{n*} \varphi_{n*} \overline{\mathbb{Q}}_\ell \simeq \bigoplus_{\alpha \in K_{P,\psi}} \bigoplus_{\xi \in W_{n,q}^\vee} \mathcal{L}_\psi(\alpha x) \otimes \mathcal{L}_\xi(x, 0's).$$

□

The morphism $C_n \rightarrow C_{n-1}$, $(x_0, \dots, x_{n-1}) \mapsto (x_0, \dots, x_{n-2})$ induces a natural projection

$$r_n: C_{P,n} \rightarrow C_{P,n-1}.$$

Clearly we have $\pi_n \circ \varphi_n = \pi_{n-1} \circ \varphi_{n-1} \circ r_n$. The morphism r_n induces

$$r_n^*: H_c^1(C_{P,n-1}) \hookrightarrow H_c^1(C_{P,n}).$$

We have a surjective homomorphism $W_{n,q} \rightarrow W_{n-1,q}$, $(x_0, \dots, x_{n-1}) \mapsto (x_0, \dots, x_{n-2})$, which induces an injection $W_{n-1,q}^\vee \rightarrow W_{n,q}^\vee$.

Lemma 5.3. *Under the identifications in Proposition 5.2, the subspace*

$$r_n^*: H_c^1(C_{P,n-1}) \hookrightarrow H_c^1(C_{P,n})$$

corresponds to

$$\bigoplus_{\xi \in W_{n-1,q}^\vee, \alpha \in K_{P,\psi}} H_c^1(\mathbb{A}^1, \mathcal{L}_\psi(\alpha x) \otimes \mathcal{L}_\xi(x, 0's)),$$

which is exactly the morphism induced by the injection $W_{n-1,q}^\vee \hookrightarrow W_{n,q}^\vee$.

Proof. Let $\mathbb{F}_q \hookrightarrow W_{n,q}$, $t \mapsto (0's, t)$. We have an isomorphism $C_{P,n}/\mathbb{F}_q \simeq C_{P,n-1}$. Thus

$$H_c^1(C_{P,n-1}) = H_c^1(C_{P,n})^{\mathbb{F}_q},$$

where $(-)^{\mathbb{F}_q}$ denotes the \mathbb{F}_q -fixed part. We have $\xi|_{\mathbb{F}_q} = 1 \iff \xi \in W_{n-1,q}^\vee$. Hence the claim follows. \square

Corollary 5.4. *We define*

$$V_n := H_c^1(C_{P,n})/H_c^1(C_{P,n-1}).$$

Then we have an isomorphism

$$V_n \simeq \bigoplus_{\xi \in W_{n,q,\text{prim}}^\vee} \bigoplus_{\alpha \in K_{P,\psi}} H_c^1(\mathbb{A}^1, \mathcal{L}_\psi(\alpha x) \otimes \mathcal{L}_\xi(x, 0's)).$$

Proof. The claim follows from $W_{n,q,\text{prim}}^\vee = W_{n,q}^\vee \setminus W_{n-1,q}^\vee$ and Lemma 5.3. \square

We recall that $W_{n,q,\text{prim}}^\vee$ denotes the set of characters ξ of $W_{n,q}$ such that $\xi|_{\{0's\} \times \mathbb{F}_q} \neq 1$.

Lemma 5.5. *Let ${}^0\xi \in W_{n,p}^\vee$ be a faithful character and let $\xi \in W_{n,q,\text{prim}}^\vee$. Choose $a = (a_0, \dots, a_{n-1}) \in W_{n,q}$ such that $\xi = \xi_a$ in the notation of Lemma 2.7. Let ${}^0\psi := {}^0\xi|_{\{0's\} \times \mathbb{F}_p}$ and $\psi := {}^0\psi \circ \text{Tr}_{q/p}$. Let $a' := a + (0's, \alpha^{p^{n-1}})$. Then $\xi_{a'} \in W_{n,q,\text{prim}}^\vee$ and*

$$\mathcal{L}_\psi(\alpha x) \otimes \mathcal{L}_\xi(x, 0's) \simeq \mathcal{L}_{\xi_a}(a' \cdot (x, 0's)).$$

Proof. We recall

$$a \cdot (x, 0's) = (a_0 x, a_1 x^p, \dots, a_{n-1} x^{p^{n-1}}).$$

We have

$$\begin{aligned} \mathcal{L}_\psi(\alpha x) \otimes \mathcal{L}_\xi(x, 0's) &\simeq \mathcal{L}_{\xi_a}(0's, (\alpha x)^{p^{n-1}}) \otimes \mathcal{L}_{\xi_a}(a \cdot (x, 0's)) \\ &\simeq \mathcal{L}_{\xi_a}((0's, \alpha^{p^{n-1}}) \cdot (x, 0's)) \otimes \mathcal{L}_{\xi_a}(a \cdot (x, 0's)) \\ &\simeq \mathcal{L}_{\xi_a}(a' \cdot (x, 0's)). \end{aligned}$$

We write $a' = (a'_0, \dots, a'_{n-1})$. By $a'_0 = a_0 \neq 0$, we have $\xi_{a'} \in W_{n,q,\text{prim}}^\vee$ by Lemma 2.7. \square

Corollary 5.6. *Let V_n be as in Corollary 5.4.*

- (1) *We have $\dim V_n = q^{n-1}(p^{n-1} - 1)(q - 1) \deg P$.*
- (2) *The curve $\overline{C}_{n,P}$ is geometrically connected.*

(3) *The natural map $H_c^1(C_{P,n}) \rightarrow H^1(\overline{C}_{P,n})$ is an isomorphism. We have*

$$g(\overline{C}_{P,n}) = \deg P \cdot \frac{q-1}{2} \sum_{i=1}^n q^{i-1}(p^{i-1} - 1) = \deg P \cdot g(\overline{C}_n).$$

Proof. Let $\xi \in W_{n,q,\text{prim}}^\vee$ and $\alpha \in K_{P,\psi}$. We simply write $\mathcal{L}_{\xi,\alpha}$ for $\mathcal{L}_\psi(\alpha x) \otimes \mathcal{L}_\xi(x, 0)$'s. We show (1). The claim follows from Corollary 5.4 and

$$\dim H_c^1(\mathbb{A}^1, \mathcal{L}_{\xi,\alpha}) = p^{n-1} - 1, \quad \#W_{n,q,\text{prim}}^\vee = q^{n-1}(q-1), \quad \#K_{P,\psi} = \deg P,$$

where the first equality follows from Lemmas 2.8 and 5.5.

We show (2). By Poincaré duality, it suffices to show that $\dim H_c^2(C_{P,n}) = 1$. We show this by induction on n . If $\xi \in W_{n,q,\text{prim}}^\vee$, then $\mathcal{L}_{\xi,\alpha}$ is ramified, and $H_c^2(\mathbb{A}^1, \mathcal{L}_{\xi,\alpha}) = 0$. Assume $n = 2$. By Proposition 5.2,

$$\begin{aligned} H_c^2(C_{P,2}) &\simeq \bigoplus_{\xi \in W_{2,q}^\vee, \alpha \in K_{P,\psi}} H_c^2(\mathbb{A}^1, \mathcal{L}_{\xi,\alpha}) \\ &\simeq \bigoplus_{\alpha \in K_{P,\psi}, b \in \mathbb{F}_q} H_c^2(\mathbb{A}^1, \mathcal{L}_\psi((\alpha + b)x)) \simeq H_c^2(\mathbb{A}^1, \overline{\mathbb{Q}}_\ell) \simeq \overline{\mathbb{Q}}_\ell(-1). \end{aligned}$$

Assume $n \geq 3$. By (1) and $H_c^2(\mathbb{A}^1, \mathcal{L}_{\xi,\alpha}) = 0$ for $\xi \in W_{n,q,\text{prim}}^\vee$, we have an isomorphism $H_c^2(C_{P,n-1}) \xrightarrow{\sim} H_c^2(C_{P,n})$. Thus the claim follows from the induction hypothesis.

We show (3). By induction on n , we show that the forgetful map

$$H_c^1(C_{P,n}) \rightarrow H^1(C_{P,n})$$

is an isomorphism.

The forgetful map

$$H_c^1(\mathbb{A}^1, \mathcal{L}_{\xi,\alpha}) \rightarrow H^1(\mathbb{A}^1, \mathcal{L}_{\xi,\alpha})$$

is an isomorphism for $a \in K_{P,\psi}$ and $\xi \in W_{n,q,\text{prim}}^\vee$. The claim for $n = 2$ follows from this. By the induction hypothesis, the map

$$H_c^1(C_{P,n}) \rightarrow H^1(C_{P,n})$$

is an isomorphism.

Hence we obtain an isomorphism $H_c^1(C_{P,n}) \xrightarrow{\sim} H^1(\overline{C}_{P,n})$. The latter claim follows from (1) and

$$2g(\overline{C}_{P,n}) = \dim H_c^1(C_{P,n}) = \sum_{i=1}^n \dim V_i.$$

□

Corollary 5.7. (1) *We define*

$$L_{V_n/\mathbb{F}_{q^s}}(T) := \det(1 - \text{Fr}_{q^s}^* T; V_n).$$

Then we have

$$L_{\overline{C}_{P,n}/\mathbb{F}_{q^s}}(T) = \prod_{i=2}^n L_{V_i/\mathbb{F}_{q^s}}(T)$$

and

$$L_{V_n/\mathbb{F}_{q^s}}(T) = \prod_{\xi \in W_{n,q,\text{prim}}^\vee} \prod_{\alpha \in K_{P,\psi}} \det(1 - \text{Fr}_{q^s}^* T; H_c^1(\mathbb{A}^1, \mathcal{L}_\psi(\alpha x) \otimes \mathcal{L}_\xi(x, 0))).$$

(2) Assume that the set of the Frobenius slopes of V_n with respect to Fr_q^* are

$$\left\{ \frac{1}{p^{n-1}}, \dots, \frac{p^{n-1} - 1}{p^{n-1}} \right\}$$

for any n . Then the set of the Frobenius slopes of $\overline{C}_{P,n}$ with respect to Fr_q^* are

$$\left\{ \frac{j}{p^i} \mid 1 \leq i \leq n-1, 1 \leq j \leq p^i - 1 \right\}.$$

Proof. The claim (1) is a consequence of Corollary 5.4.

The claim (2) follows from the assumption and Corollary 5.6. \square

Remark 5.8. We consider Lemma 5.5. We have

$$(5.1) \quad \mathcal{L}_0(\xi)(a' \cdot (x, 0')) = \mathcal{L}_0(\xi)(a'_0 x, \dots, a'^{p^{n-1}}_{n-1} x) = \mathcal{L}_0(\xi) \left(\sum_{i=0}^{n-1} V^i ([a'^{p^{-i}}_i x]) \right),$$

where $[-]: \mathbb{A}_{\mathbb{F}_q}^1 \rightarrow \mathbb{W}_{n,q}$ denotes the Teichmüller lift and $V: \mathbb{W}_{n,q} \rightarrow \mathbb{W}_{n,q}$ is the Verschiebung (cf. [18, the proof of Lemma 3.1]). By $a'_0 \neq 0$ and (5.1), the Newton polygon of the polynomial

$$\det(1 - \text{Fr}_{q^s}^* T; H^1_c(\mathbb{A}^1, \mathcal{L}_\psi(ax) \otimes \mathcal{L}_\xi(x, 0)))$$

is determined if [30, Theorem 1.3] is correct. If [30, Theorem 1.3] holds for general q , the assumption in (2) in Corollary 5.7 is satisfied and the Frobenius slopes of \overline{C}_P are determined.

Although [30, Theorem 1.3] is stated for arbitrary q , the arguments in the proof of [30, Proposition 5.1] seem to be valid only in the case $q = p$. In particular, the precise Newton polygon information required in [30, Theorem 1.3] cannot, in general, be deduced from [30, Proposition 5.2]. Indeed, when $q > p$, the valuations of the coefficients of the reciprocal characteristic polynomial of the matrix appearing in the last paragraph of [30, p. 281] cannot, in general, be controlled using [30, Proposition 5.2] alone.

Corollary 5.9. Assume that p is odd. The curve $\overline{C}_{P,n}$ is not supersingular. If $q = p$ and a smooth projective curve X is a finite quotient of $\overline{C}_{P,n}$, the curve X is not supersingular.

Proof. The curve $\overline{C}_{P,n}$ admits a finite quotient \overline{C}_n . Since \overline{C}_n is not supersingular by Corollary 3.8, the curve $\overline{C}_{P,n}$ cannot be supersingular.

If X is supersingular, the Frobenius slope of X is $1/2$. Thus the latter claim follows from Corollary 5.7(3). \square

Corollary 5.10. Assume that $q = p$ and p is odd. Let $\xi \in W_{n,q,\text{prim}}^\vee$ and $\psi \in \mathbb{F}_p^\vee \setminus \{1\}$. Let $\pi: \mathbb{A}_{\mathbb{F}_q}^2 \rightarrow \mathbb{A}_{\mathbb{F}_q}^1$, $(x, t) \mapsto x$ and $\pi': \mathbb{A}_{\mathbb{F}_q}^2 \rightarrow \mathbb{A}_{\mathbb{F}_q}^1$, $(x, t) \mapsto t$. Let

$$\mathcal{F}_{\xi, \psi} := R^1 \pi'_! (\pi^* \mathcal{L}_\xi(x, 0') \otimes \mathcal{L}_\psi(tx)) \left(\frac{1}{2} \right).$$

Then $\mathcal{F}_{\xi, \psi}$ does not have finite monodromy.

Proof. Let $t \in \mathbb{F}_q$. By the proper base change theorem and Lemma 5.5, we have

$$(\mathcal{F}_{\xi, \psi})_{\bar{t}} \simeq H^1_c(\mathbb{A}^1, \mathcal{L}_\xi(x, 0') \otimes \mathcal{L}_\psi(tx)) \left(\frac{1}{2} \right) \simeq H^1_c(\mathbb{A}^1, \mathcal{L}_{\xi'}(x, 0')) \left(\frac{1}{2} \right)$$

with some $\xi' \in W_{n,q,\text{prim}}^\vee$. By [30, Theorem 1.3], the Frobenius eigenvalues of Frob_q^* on $(\mathcal{F}_{\xi, \psi})_{\bar{t}}$ are not a root of unity. The sheaf $\mathcal{L}_\xi(x, 0')$ is geometrically irreducible, its

determinant is arithmetically of finite order and pure of weight zero. Hence the claim follows (cf. [17, Theorem 8.14.4]). \square

Assume that $n = 2$ and $p = l = 3$. We simply write W for $W_{2,q}$ as before. For $\xi \in W_{\text{prim}}^\vee$ and $\alpha \in K_{P,\psi}$, let

$$S_{\xi,\alpha} := \sum_{x \in \mathbb{F}_{q^s}} \psi(\text{Tr}_{q^s/q}(\alpha x)) \xi(\text{Tr}_{q^s/q}(x, 0)).$$

Finally, we state our main theorem of this paper.

Corollary 5.11. *Assume $n = 2$ and $p = l = 3$.*

(1) *We have*

$$L_{\overline{C}_{P,2}/\mathbb{F}_{q^s}}(T) = \prod_{\xi \in W_{\text{prim}}^\vee} \prod_{a \in K_{P,\psi}} (1 + S_{\xi,a} T + q^s T^2).$$

(2) *The Frobenius slopes of $\overline{C}_{P,2}$ with respect to Fr_q^* are $\{1/3, 2/3\}$.*

Proof. We show (1). By $n = 2$, we have

$$L_{\overline{C}_{P,2}/\mathbb{F}_{q^s}}(T) = L_{V_2/\mathbb{F}_{q^s}}(T).$$

By Corollary 5.5, we have

$$\mathcal{L}_\psi(\alpha x) \otimes \mathcal{L}_\xi(x, 0) \simeq \mathcal{L}_{\xi_{a'}}(x, 0),$$

Thus, by Corollary 3.17,

$$\begin{aligned} \det(1 - \text{Fr}_{q^s}^* T; H_c^1(\mathbb{A}^1, \mathcal{L}_\psi(\alpha x) \otimes \mathcal{L}_\xi(x, 0))) &= \det(1 - \text{Fr}_{q^s}^* T; H_c^1(\mathbb{A}^1, \mathcal{L}_{\xi_{a'}}(x, 0))) \\ &= 1 + S_{\xi_{a'}} T + q^s T^2. \end{aligned}$$

Hence the claim follows from Corollary 5.7 and $S_{\xi,\alpha} = S_{\xi_{a'}}$. The claim (2) follows from Corollary 4.6. \square

Acknowledgement. T. I. is supported by JSPS KAKENHI Grant Numbers 23K20786, 24K21512 and 25K00905. D. T. is supported by JSPS KAKENHI Grant Number 25KJ0122. T. T. is supported by JSPS KAKENHI Grant Numbers 25K06959 and 23K20786.

APPENDIX A. DETERMINANT

In this appendix, we show that the Frobenius determinant on the first étale cohomology group with suitable coefficients is equal to a Gauss sum multiplied by a root of unity. This provides a generalization of Lemma 4.8.

Let q be a power of an odd prime p . The following is shown similarly to [18, the proof of Theorem 7.1].

Lemma A.1. *Let \mathcal{L} be a smooth $\overline{\mathbb{Q}}_\ell$ -sheaf pure of weight 0 on a smooth curve X^0 over \mathbb{F}_q . Let $X := X_{\mathbb{F}}^0$. Assume that*

$$\det(-\text{Fr}_q^*; H_c^1(X, \mathcal{L})) \in \mathbb{Z}[\zeta_{p^n}] \quad \text{with an integer } n \geq 0.$$

Furthermore, we suppose that the natural map

$$H_c^1(X, \mathcal{L}) \rightarrow H^1(X, \mathcal{L})$$

is an isomorphism. We take $\psi \in \mathbb{F}_q^\vee \setminus \{1\}$ and denote by

$$G(\psi) = \sum_{x \in \mathbb{F}_q} \psi(x^2)$$

the quadratic Gauss sum. If we set $s := \dim H_c^1(X, \mathcal{L})$, we have

$$\frac{\det(-\text{Fr}_q^*; H_c^1(X, \mathcal{L}))}{G(\psi)^s}$$

is a root of unity.

Proof. Put

$$\alpha := \det(-\text{Fr}_q^*; H_c^1(X, \mathcal{L})) \in \mathbb{Z}[\zeta_{p^n}].$$

Since \mathcal{L} is pure of weight 0, Weil II [10] implies that $H_c^1(X, \mathcal{L})$ and $H_c^1(X, \mathcal{L}^\vee)$ are mixed of weights ≤ 1 . By Poincaré duality and the assumption, the natural map

$$H_c^1(X, \mathcal{L}^\vee) \rightarrow H^1(X, \mathcal{L}^\vee)$$

is an isomorphism and

$$H_c^1(X, \mathcal{L}) \simeq H^1(X, \mathcal{L}^\vee)^\vee(-1) \simeq H_c^1(X, \mathcal{L}^\vee)^\vee(-1).$$

Hence $H_c^1(X, \mathcal{L})$ is mixed of weights ≥ 1 and therefore pure of weight 1. Thus for any embedding $\iota: \overline{\mathbb{Q}}_\ell \hookrightarrow \mathbb{C}$, the complex absolute value satisfies

$$|\iota(\alpha)| = q^{s/2}.$$

Similarly, for the Gauss sum, one has $G(\psi) \in \mathbb{Z}[\zeta_{p^n}]$ and $|\iota(G(\psi))| = q^{1/2}$ for any ι . Hence

$$(A.1) \quad \left| \iota \left(\frac{\alpha}{G(\psi)^s} \right) \right| = 1.$$

Let \mathfrak{p}' be a finite place of $\mathbb{Q}(\zeta_{p^n})$ with $\mathfrak{p}' \nmid p$. Then both α and $G(\psi)$ in $\mathbb{Z}[\zeta_{p^n}]$ are \mathfrak{p}' -units.

Since all archimedean absolute values of $\alpha/G(\psi)^s$ are 1 by (A.1), the product formula implies that the p -adic valuation is zero. Therefore

$$\frac{\alpha}{G(\psi)^s} \in \mathbb{Z}[\zeta_{p^n}],$$

i.e. it is an algebraic integer, and by (A.1) all of its complex absolute values are equal to 1.

By Kronecker's theorem, an algebraic integer whose Galois conjugates all have absolute value 1 is a root of unity. Thus $\alpha/G(\psi)^s$ is a root of unity. \square

Corollary A.2. *Let $\mathcal{K}_\chi(x)$ be the Kummer sheaf on $\mathbb{G}_{\text{m}, \mathbb{F}_q}$ defined by $y^2 = x$ and a character $\chi \in \mathbb{F}_2^\vee$. Let n be a positive integer and let $W_{n,q} := \mathbb{W}_n(\mathbb{F}_q)$, where \mathbb{W}_n denotes the p -typical Witt vector scheme of length n . For $\chi \in \mathbb{F}_2^\vee$ and a nontrivial character $\xi \in W_{n,q}^\vee \setminus \{1\}$, we define*

$$V_{\chi, \xi} := \begin{cases} H_c^1(\mathbb{A}^1, \mathcal{L}_\xi(x, 0's)) & \text{if } \chi = 1, \\ H_c^1(\mathbb{G}_{\text{m}}, \mathcal{K}_\chi(x) \otimes \mathcal{L}_\xi(x, 0's)) & \text{if } \chi \neq 1. \end{cases}$$

Then

$$\frac{\det(-\text{Fr}_q^*; V_{\chi, \xi})}{G(\psi)^{\dim V_{\chi, \xi}}}$$

is a root of unity.

Proof. We simply write \mathcal{L} for $\mathcal{K}_\chi(x) \otimes \mathcal{L}_\xi(x, 0's)$. We define X^0 to be $\mathbb{A}_{\mathbb{F}_q}^1$ if $\chi = 1$ and $\mathbb{G}_{m,\mathbb{F}_q}$ if $\chi \neq 1$. We apply Lemma A.1. By the trace formula,

$$S_m := -\text{Tr}(\text{Fr}_{q^m}^*; H_c^1(X, \mathcal{L})) \in \mathbb{Z}[\zeta_{p^n}] \quad \text{for an integer } m \geq 1.$$

We define the L -function by

$$L_{\mathcal{L}}(T) := \exp \left(\sum_{m=1}^{\infty} \frac{S_m}{m} T^m \right) \in \mathbb{Q}(\zeta_{p^n})[[T]].$$

We have $H_c^i(X, \mathcal{L}) = 0$ for $i \neq 1$. Hence the function $L_{\mathcal{L}}(T)$ equals $\det(1 - \text{Fr}_q^* T; H_c^1(X, \mathcal{L}))$ and the leading coefficient is

$$\alpha := \det(-\text{Fr}_q^*; H_c^1(X, \mathcal{L})) \in \mathbb{Q}(\zeta_{p^n}).$$

Since $V_{\chi, \xi} \subset H^1(Y)$ for a smooth projective curve Y , the eigenvalues of Fr_q^* on $V_{\chi, \xi}$ are algebraic integers by [10]. Hence α is an algebraic integer and $\alpha \in \mathbb{Z}[\zeta_{p^n}]$.

Since \mathcal{L} is ramified at every point of $\overline{X} \setminus X$ and has rank one, the natural map

$$H_c^1(X, \mathcal{L}) \rightarrow H^1(X, \mathcal{L})$$

is an isomorphism. The latter assertion follows from Lemma A.1. \square

REFERENCES

- [1] A. Abbes, T. Saito, Local Fourier transform and epsilon factors, Compos. Math. 146, No. 6, 1507–1551 (2010).
- [2] N. Aoki, On pure Gauss sums, Comment. Math. Univ. St. Pauli 61, No. 2, 133–165 (2012).
- [3] B. J. Birch, How the number of points of an elliptic curve over a fixed prime field varies, J. Lond. Math. Soc. (2) 43, 57–60 (1968).
- [4] J. L. Brylinski, Théorie du corps de classes de Kato et revêtements abéliens de surfaces, Ann. Inst. Fourier 33, No. 3, 23–38 (1983).
- [5] L. Carlitz, A note on exponential sums, Math. Scand. 42, 39–48 (1978).
- [6] L. Carlitz, Explicit evaluation of certain exponential sums, Math. Scand. 44, 5–16 (1979).
- [7] H. Davenport, On certain exponential sums, J. Reine Angew. Math. 169, 158–176 (1933).
- [8] C. Davis, D. Wan, L. Xiao, Newton slopes for Artin–Schreier–Witt towers, Math. Ann. 364, no. 3–4, 1451–1468 (2016).
- [9] P. Deligne, Cohomologie étale, Séminaire de Géométrie Algébrique du Bois-Marie SGA 4 1/2, Lecture Notes in Mathematics 569, Springer-Verlag, Berlin, 1977.
- [10] P. Deligne, La conjecture de Weil. II, Publ. Math., Inst. Hautes Étud. Sci. 52, 137–252 (1980).
- [11] P. Deligne, G. Lusztig, Representations of reductive groups over finite fields, Ann. Math. (2) 103, 103–161 (1976).
- [12] F. García-Cortés, A. Rojas-León, Finite monodromy of some two-parameter families of exponential sums, 2406.10385 [math.NT], preprint.
- [13] G. van der Geer and M. van der Vlugt, Reed–Muller codes and supersingular curves. I, Compos. Math. 84, no. 3, 333–367 (1992).
- [14] T. Ito, D. Takeuchi, T. Tsushima, The L -polynomials of van der Geer–van der Vlugt curves in characteristic 2, 2505.22036 [math.NT], preprint.
- [15] N. M. Katz, On the monodromy groups attached to certain families of exponential sums, Duke Math. J. 54, 41–56 (1987).
- [16] N. M. Katz, Gauss sums, Kloosterman sums, and monodromy groups, Annals of Mathematics Studies, 116.
- [17] N. M. Katz, Exponential sums and differential equations, Annals of Mathematics Studies, 124.
- [18] N. M. Katz, Witt vectors and a question of Keating and Rudnick, Int. Math. Res. Not. 2013, No. 16, 3613–3638 (2013).

- [19] N. M. Katz, Witt vectors and a question of Entin, Keating, and Rudnick, *Int. Math. Res. Not.* 2015, No. 14, 5959–5975 (2015).
- [20] N. M. Katz, Witt vectors and a question of Rudnick and Waxman, *Int. Math. Res. Not.* 2017, No. 11, 3377–3412 (2017).
- [21] N. M. Katz, G_2 and some exceptional Witt vector identities, *Finite Fields Appl.* 47, 125–144 (2017).
- [22] N. M. Katz, G. Laumon, Transformation de Fourier et majoration de sommes exponentielles, *Publ. Math., Inst. Hautes Étud. Sci.* 62, 145–202 (1985).
- [23] N. M. Katz, P. H. Tiep, Rigid local systems on \mathbb{A}^1 with finite monodromy, *Mathematika* 64, 785–846 (2018).
- [24] N. M. Katz, P. H. Tiep, Exponential Sums, Hypergeometric Sheaves and Monodromy Groups, 2024, <https://web.math.princeton.edu/~nmk/kt18final.pdf>
- [25] J. Kramer-Miller, p -adic estimates of abelian Artin L -functions on curves, *Forum Math. Sigma* 10, Paper No. e2, 30 p. (2022).
- [26] J. Kramer-Miller, J. Upton, Newton polygons of sums on curves II: Variation in p -adic families, *Int. Math. Res. Not.* 2023, No. 21, 18500–18536 (2023).
- [27] J. Kramer-Miller, J. Upton, Newton polygons of sums on curves I: Local-to-global theorems, *Math. Ann.* 391, No. 1, 1347–1394 (2025).
- [28] P. V. Kumar, T. Helleseth, A. R. Calderbank, An upper bound for some exponential sums over Galois rings with applications, *IEEE Trans. Inform. Theory* 41, no. 2, 456–468 (1995).
- [29] G. Laumon, Transformation de Fourier, constantes d'équations fonctionnelles et conjecture de Weil, *Publ. Math., Inst. Hautes Étud. Sci.* 65, 131–210 (1987).
- [30] C. Liu, The L -functions of twisted Witt extensions, *J. Number Theory* 125, No. 2, 267–284 (2007).
- [31] C. Liu, D. Wei, The L -functions of Witt coverings, *Math. Z.* 255, no. 1, 95–115 (2007).
- [32] R. Livné, The average distribution of cubic exponential sums, *J. Reine Angew. Math.* 375/376, 362–379 (1987).
- [33] A. Rojas-León, Finite monodromy of some families of exponential sums, *J. Number Theory* 197, 37–48 (2019).
- [34] J. P. Serre, *Corps Locaux*, 2nd ed., Hermann, Paris, 1968.
- [35] D. Takeuchi, T. Tsushima, Gauss sums and van der Geer–van der Vlugt curves, *Bull. Lond. Math. Soc.* 56, No. 2, 602–623 (2024).

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, KYOTO UNIVERSITY KYOTO, 606–8502, JAPAN

Email address: `tetsushi@math.kyoto-u.ac.jp`

DEPARTMENT OF MATHEMATICS, INSTITUTE OF SCIENCE TOKYO, 2-12-1 OOKAYAMA, MEGURO-KU, TOKYO, 152-8551, JAPAN

Email address: `daichi.takeuchi4@gmail.com`

KEIO UNIVERSITY SCHOOL OF MEDICINE, 4-1-1 HIYOSHI, KOHOKU-KU, YOKOHAMA, 223-8521, JAPAN

Email address: `tsushima@keio.jp`