

Contact lifts and Hölder lifts to central extension of Carnot groups

Yihan Cui

December 16, 2025

Abstract

We consider the existence problem of lift F of a map f between Carnot group with different smoothness, where we use central extension to define lifting. Our main result is the existence of the contact lifts of Lipschitz and Sobolev maps and the rigidity result for the contact lift of quasiconformal maps: a quasiconformal map admits a contact lift then it is bi-Lipschitz. We also show a necessary criterion for the extension of γ -Hölder lift when $\gamma > \frac{n}{n+1}$ for step- n Carnot group.

1 Introduction

The lifting problem is a useful tool to study and construct a contact map between higher step Carnot group. Let's illustrate this problem.

Let $V_1 \xrightarrow{\iota} G_1 \xrightarrow{\pi_1} H_1$ and $V_2 \xrightarrow{\iota} G_2 \xrightarrow{\pi_2} H_2$ be central extensions of Carnot group. A typical example is the Heisenberg group $R \xrightarrow{\iota} \mathbb{H}_n \xrightarrow{\pi} \mathbb{R}^{2n}$. The purpose is to find a contact lift $F : G_1 \rightarrow G_2$ of contact map $f : H_1 \rightarrow H_2$ such that $\pi_2 \circ F = f \circ \pi_1$ i.e. the following diagram commutes

$$\begin{array}{ccc} G_1 & \xrightarrow{\pi_1} & H_1 \\ \downarrow F & & \downarrow f \\ G_2 & \xrightarrow{\pi_2} & H_2 \end{array}$$

In the Heisenberg group case, we want to find a contact lift F of f such that the following diagram commutes

$$\begin{array}{ccc} \mathbb{H}_n & \xrightarrow{\pi} & \mathbb{R}^{2n} \\ \downarrow F & & \downarrow f \\ \mathbb{H}_n & \xrightarrow{\pi} & \mathbb{R}^{2n} \end{array}$$

When the domain is one dimensional, by control theory ([1], Section 8), a curve lays in the Euclidean space can be uniquely lifted into Carnot group. When the domain is higher dimensional, we are supposed to consider the integrability of submanifold.

A well studied topic is the Heisenberg group, a classical technique used by Allcock is the lift of a map $\mathbb{D} \rightarrow \mathbb{R}^{2n}$ in [2]. He construct a map $\mathbb{D} \rightarrow \mathbb{H}_n$ satisfying a Lagrangian condition.

For map $\mathbb{R}^k \rightarrow \mathbb{R}^k$, a type of problems is the existence of embedding from a domain in Euclidean space into Heisenberg group. Because of the Lie algebra graded structure, the dimension of the domain must be smaller or equal to n . For

more detail, readers can go to [5]. A useful way to avoid the restriction of Lie algebra homomorphism is to study maps with poor smoothness condition like Hölder maps. In [6], Piotr Hajłasz, Jacob Mirra and Armin Schikorra study the Gromov non-embedding theorem with differential form which is a crucial tool in our paper for studying the existence of Hölder lifts and their properties.

The early work is from Balogh–Hoefer–Isenegger–Tyson [3] considering the existence of contact lifts of Lipschitz maps $\mathbb{R}^2 \rightarrow \mathbb{R}^2$ into first Heisenberg group. Their methods provides a good example when study the existence problem of the contact lifts and even the Hölder lifts. As to the lifts of $\mathbb{R}^{2n} \rightarrow \mathbb{R}^{2n}$, this problem is originally motivated by the Mostow rigidity. A classical result is that the Heisenberg group \mathbb{H}_n is the ideal boundary of a Siegel domain in \mathbb{C}^{n+1} under hyperbolic metric induced by Bergman metric([10],Section 4). Quasi-isometries between such spaces correspond naturally to quasisymmetries between their ideal boundaries [11]. Capogna–Tang [4] study lifts of symplectomorphisms $\mathbb{R}^{2n} \rightarrow \mathbb{R}^{2n}$ to full-rank contact maps between Heisenberg groups \mathbb{H}_n .

If we just want to find a map F such that the diagram commutes is easy, the key is to find a lift F with the same smoothness of f . An important and enlightening result is given by Eero Hakavuori, Susanna Heikkila, and Toni Ikonen [7]. They study the existence problem of lifting a smooth contact map between Carnot group.

Follow their work, We consider the existence problem of lift F of a map f between Carnot group with different smoothness. Let $V_1 \xrightarrow{\iota} G_1 \xrightarrow{\pi_1} H_1$ and $V_2 \xrightarrow{\iota} G_2 \xrightarrow{\pi_2} H_2$ be fixed central extensions of Carnot groups given by 2-cocycles ρ_1 and ρ_2 , see Section 2 for details.

Since the completed result in [7] about the criterion for the existence of smooth contact lifts is proved with the help of Rumin complex, we can also use this for the existence of Lipschitz or Sobolev contact lifts as long as we can get a similar result like Lemma 5.3 in [7].

Therefore, for section 3 and 4, our purpose is to generalize the result for Lipschitz or Sobolev condition.

Lemma 1.1. *Let $U_1 \subset \mathbb{H}_1$ and $\tilde{U}_1 \subset \pi_1^{-1}(U_1) \subset \mathbb{G}_1$ be domains. A Lipschitz contact map $f : U_1 \rightarrow \mathbb{H}_2$ admits a Lipschitz contact lift $F : \tilde{U}_1 \rightarrow \mathbb{G}_2$ if and only if*

$$(f \circ \pi_1) \left(\Gamma_{LIP}(g, \tilde{U}_1) \right) \subset \Gamma_{LIP}^\rho(f(\pi_1(g)), \mathbb{H}_2) \quad (1.1)$$

for some $g \in \tilde{U}_1$.

and

Lemma 1.2. *Let $U_1 \subset \mathbb{H}_1$ and $\tilde{U}_1 \subset \pi_1^{-1}(U_1) \subset \mathbb{G}_1$ be domains. A Sobolev contact map $f : U_1 \rightarrow \mathbb{H}_2$ admits a Sobolev contact lift $F : \tilde{U}_1 \rightarrow \mathbb{G}_2$ if and only if*

$$(f \circ \pi_1) \left(\Gamma_{LIP}(g, \tilde{U}_1) \right) \subset \Gamma_{W^{1,p}}^\rho(f(\pi_1(g)), \mathbb{H}_2) \text{ for } \mathcal{H}^{Q_1-1} - \text{a.e.} \quad (1.2)$$

for some $g \in \tilde{U}_1$, where $f, F \in W^{1,p}$ and $p > Q_1$, Q_1 is homogeneous dimension of H_1 .

In these two lemma, we use $F(\tilde{\gamma}(1)) = f(\gamma(1)) \cdot \exp(\int_{\gamma} (f \circ \pi_1)^* \alpha_2)$ to define the lifting F of f , since (1.1) and (1.2) leads to F is well defined in this way. For the smoothness of lifting F , we find the upper gradient of F with the help of this definition.

In section 5, we study the rigidity of the contact lift of quasiconformal maps getting the following result.

Theorem 1.3. *Suppose that f is a K -quasiconformal contact map and it admits a contact lift F then f is bi-Lipschitz.*

Because of the restriction of the structure of lifting, the choice of maps are greatly limited. With the help lift invariant differential form, we can use the concept of weight to distinguish the different actions of different layers in the Lie algebra. To be specific, the i -th layer are expending at the rate of r^i when the horizontal layer are are expending at the rate of r .

A interesting case is the quaternionic Heisenberg group $\mathbb{H}_{\mathbb{H}}^n$, because the lifting is from a map form \mathbb{R}^{4n} to \mathbb{R}^{4n} , the second layer provides a very strong restriction for the algebra structure on Lie algebra.

Theorem 1.4. *Suppose f is a Sobolev continuous map on \mathbb{R}^{4n} which admits a contact lift F on $\mathbb{H}_{\mathbb{H}}^n$ and $D_F F$ is full-rank for every point, then f is an affine transformation.*

In section 6, we show a classical example Heisenberg group \mathbb{H}^n . Thanks to the standard symplectic form on \mathbb{R}^{2n} , we can find a family of smooth lifting to approximate the sobolev lifting.

Theorem 1.5. *For any sobolev map f on \mathbb{R}^{2n} and its contact lift F on \mathbb{H}^n , there are a series of smooth g_ε which admit smooth contact lift G_ε , $g_\varepsilon \rightarrow f$ and $G_\varepsilon \rightarrow F$ as $\varepsilon \rightarrow 0$.*

In section 7, we study the Hölder lift with the following theorem.

Theorem 1.6. *Suppose that $f \in C^{0,\beta}(\Omega; G_2)$, where G_2 is a step n_2 Carnot group, $\Omega \subset G_1$ is open, X_i is a left-invariant vector field belonging to the i -th layer and $0 < \beta \leq 1$. If $B(x_o, 2r) \subset \Omega$, then for all $0 < \varepsilon < r$ and the weight of ω_j is larger than the weight of X_i ,*

$$\begin{aligned} X_i(f_\varepsilon^* \omega_1^k) &\lesssim \varepsilon^{(i+1)\beta-i} \\ X_i(f_\varepsilon^* \omega_2^m) &\lesssim \varepsilon^{(i+1)\beta-i} \\ &\dots \\ X_i(f_\varepsilon^* \omega_{n_2}^m) &\lesssim \varepsilon^{(i+1)\beta-i} \end{aligned} \tag{1.3}$$

where $f_\varepsilon = f * \eta_\varepsilon = \int_{\mathbb{G}_{\mathbb{H}}} f(y) \eta_\varepsilon(y^{-1}x) dy$ and η_ε is the heat kernel on Carnot group G_1 .

Theorem 1.6 showcases the graded metric structure under the influence of the graded structure of Lie algebra structure even for non-differentiable case like Hölder maps. This greatly provide us a powerful tool to study the behaviors of Hölder lifts. Therefore, we have the the theorem below about the equivalent condition for the existence of Hölder lifts.

Theorem 1.7. *Let $U_1 \subset \mathbb{H}$ and $\tilde{U}_1 \subset \pi_1^{-1}(U_1) \subset \mathbb{G}_1$ be domains and $\text{rank}(\mathbb{G}_2) = \text{rank}(\mathbb{H}_2)$. For $\beta > \frac{1}{2}$, A Hölder map $f \in C^{0,\beta}$, $f : U_1 \rightarrow \mathbb{H}_2$ admits a Hölder lift $F \in C^{0,\beta}$, $F : \tilde{U}_1 \rightarrow \mathbb{G}_2$ if and only if*

$$\int_{f \circ \pi_1 \circ \tilde{\gamma}} \alpha = 0 \tag{1.4}$$

for all Lipschitz closed curves $\tilde{\gamma}$ on \mathbb{G}_1 . We define $\int_{\tilde{\gamma}} (f \circ \pi_1)^* \alpha := \lim_{\varepsilon \rightarrow 0} \int_{\tilde{\gamma}} (f_\varepsilon \circ \pi_1)^* \alpha$, where $f_\varepsilon = f * \eta_\varepsilon = \int_{\mathbb{G}_1} f(y) \eta_\varepsilon(y^{-1}x) dy$ and η_ε is the heat kernel on Carnot group \mathbb{G}_1 .

We generalize the definition of pullback of the differential form under β -Hölder map f . With it, we similarly define the lift of f with the form of integral. In the proof, we take advantage of the the Lie algebra graded-structure-preserving property for Hölder map to show the smoothness of Hölder lift F . To be specific, if

the in the horizontal layer the movement is approximately ε^β , the i -th layer would go $\varepsilon^{i\beta}$ (including the extension part $\exp(\int_\gamma (f \circ \pi_1)^* \alpha_2)$), which matches the metric structure of Carnot group.

We also show a criterion that is easy to check for the existence of Hölder lift.

Theorem 1.8. *Let $V_1 \rightarrow G_1 \rightarrow H$ and $V_2 \rightarrow G_2 \rightarrow H$ be central extensions, where $\mathfrak{h} = \mathfrak{h}^{[1]} \oplus \dots \oplus \mathfrak{h}^{[n]}$ and $V_1 = V_1^{[n+1]}$ and $V_2 = V_2^{[n+1]}$. Let U is a simply connected domain in H and $f : U \rightarrow H$ be a γ -Hölder map for $\beta > \frac{n}{n+1}$. Then f admits a β -Hölder lift F if and only if there is a linear map $\varphi : V_1 \rightarrow V_2$ and a V_2 -value 1-form λ with weight ≥ 2 such that the integral of $f^* \alpha_2$ on any horizontal curve equals to the integral of $\varphi \circ \alpha_1$ on the same curve, i.e.*

$$\varphi \circ \int_\gamma \alpha_1 = \int_\gamma f^* \alpha_2.$$

In theorem 1.8, because of lack of differential structure, we can just study under the graded structure. Although it is not as elegant as Lie algebra homomorphism, there is a special case which is easy to prove. If the central extension is just for the highest layer, we can decompose the lifting F into two part: $F(hk) = f(h)\Phi(k)$, where $h \in \mathbb{H}$, $k \in V_1$ and $\Phi : V_1 \rightarrow V_2$.

2 Preliminary

We recall the basic properties of Carnot groups. A Lie algebra \mathfrak{g} is stratified if it has a decomposition $\mathfrak{g} = \mathfrak{g}^{[1]} \oplus \dots \oplus \mathfrak{g}^{[s]}$ with $[\mathfrak{g}^{[1]}, \mathfrak{g}^{[k]}] = \mathfrak{g}^{[k+1]}$ for all $k \geq 1$, where we denote $\mathfrak{g}^{[k]} = \{0\}$ for $k \geq s+1$. A (sub-Riemannian) Carnot group is a simply connected nilpotent Lie group G with a stratified Lie algebra \mathfrak{g} equipped with an inner product on the horizontal layer $\mathfrak{g}^{[1]}$. For convenience, We will need an inner product not only on the horizontal layer $\mathfrak{g}^{[1]}$, but instead on the whole Lie algebra \mathfrak{g} . We always equip the Lie algebra of a Carnot group with an inner product for which the layers $\mathfrak{g}^{[1]} \oplus \dots \oplus \mathfrak{g}^{[s]}$ are pairwise orthogonal. The homogeneous dimension of G is $Q_G := \sum_{k=1}^s k \dim(\mathfrak{g}^{[k]})$. The (horizontal) rank of G is the dimension of the horizontal layer $\mathfrak{g}^{[1]}$.

Given $g \in G$, we denote by $L_g : G \rightarrow G$ the left-translation $L_g(h) = gh$. We identify $T_e G$ with \mathfrak{g} . An absolutely continuous curve $\gamma : [0, 1] \rightarrow G$ is horizontal if for almost every $t \in [0, 1]$ its left-trivialized derivative $(L_{\gamma(t)}^{-1})_* \dot{\gamma}(t)$ is contained in the horizontal layer $\mathfrak{g}^{[1]}$. The length of the horizontal curve γ is

$$\ell(\gamma) = \int_0^1 \left\| (L_{\gamma(t)}^{-1})_* \dot{\gamma}(t) \right\| dt$$

where the norm is the one induced by the inner product on $\mathfrak{g}^{[1]}$. The sub-Riemannian distance between two points $g, h \in G$ is

$$d(g, h) = \inf \{ \ell(\gamma) \mid \gamma : [0, 1] \rightarrow G \text{ horizontal}, \gamma(0) = g, \gamma(1) = h \}.$$

By construction, this distance is left-invariant. Carnot groups also admit a one-parameter group of automorphisms, known as the dilations $\delta_\lambda : G \rightarrow G$, $\lambda > 0$. The associated Lie algebra automorphisms, also denoted $\delta_\lambda : \mathfrak{g} \rightarrow \mathfrak{g}$, are the linear maps defined on each layer of the stratification as

$$\delta_\lambda(X) = \lambda^k X, \quad X \in \mathfrak{g}^{[k]}, \quad 1 \leq k \leq s.$$

The sub-Riemannian distance is 1-homogeneous with respect to these dilations, i.e., $d(\delta_\lambda g, \delta_\lambda h) = \lambda d(g, h)$.

If we want to study the properties of contact lift, we need to start with central extension and related concepts. Let \mathfrak{h} be a Lie algebra, V a vector space, and $\rho: \bigwedge^k \mathfrak{h} \rightarrow V$ a vector-valued k -form. Here, and in what follows, V is assumed to be a finite-dimensional vector space. The Lie algebra differential $d_0\rho$ of ρ is the vector-valued $(k+1)$ -form

$$\begin{aligned} d_0\rho(X_1, \dots, X_{k+1}) \\ = \sum_{i < j} (-1)^{i+j} \rho([X_i, X_j], X_1, \dots, \hat{X}_i, \dots, \hat{X}_j, \dots, X_{k+1}), \end{aligned}$$

where \hat{X}_i means that X_i is omitted from the list. We identify $\Lambda^1 \mathfrak{h} \simeq \mathfrak{h}$, so vector valued 1-forms are identified with linear maps $\mathfrak{h} \rightarrow V$. A k cocycle is a k -form whose Lie algebra differential vanishes.

Definition 2.1. Let \mathfrak{h} be a Lie algebra and $\rho: \bigwedge^2 \mathfrak{h} \rightarrow V$ a 2-cocycle with values in a vector space V . Denote the Lie bracket of \mathfrak{h} by $[\cdot, \cdot]_{\mathfrak{h}}$. The central extension of \mathfrak{h} by ρ is a Lie algebra \mathfrak{g} with a direct sum decomposition $\mathfrak{h} \oplus V$ equipped with the Lie bracket

$$[X + A, Y + B]_{\mathfrak{g}} = [X, Y]_{\mathfrak{h}} + \rho(X, Y), \quad \text{for } X, Y \in \mathfrak{h}, A, B \in V.$$

The direct sum decomposition $\mathfrak{g} = \mathfrak{h} \oplus V$ induces a natural inclusion $\iota_*: V \rightarrow \mathfrak{g}$ and a projection $\pi_*: \mathfrak{g} \rightarrow \mathfrak{h}$ which are Lie algebra homomorphisms. The central extension is denoted by

$$V \xrightarrow{\iota_*} \mathfrak{g} \xrightarrow{\pi_*} \mathfrak{h}.$$

If G and H are simply connected Lie groups with Lie algebras \mathfrak{g} and \mathfrak{h} , respectively, we also refer to the induced short exact sequence

$$V \xrightarrow{\iota} G \xrightarrow{\pi} H$$

as a central extension of H by ρ .

For the purpose of studying the functions that admit contact lift, we are supposed to define the central extension of Carnot groups.

Definition 2.2. A central extension $V \xrightarrow{\iota} G \xrightarrow{\pi} H$ by ρ is called stratified if the following properties hold.

- (i) The Lie algebras \mathfrak{h} and \mathfrak{g} are stratified and the vector space V is graded.
- (ii) The inclusion $\iota_*: V \rightarrow \mathfrak{g}$ and the projection $\pi_*: \mathfrak{g} \rightarrow \mathfrak{h}$ are graded with respect to the gradings from (i).

A stratified central extension $V \rightarrow G \xrightarrow{\iota} H$ by ρ is a central extension of Carnot groups if the following hold.

- (iii) The Lie groups G and H are Carnot groups and V is an inner product space.
- (iv) The inclusion $\iota_*: V \rightarrow \mathfrak{g}$ is an isometry and $\pi_*: \mathfrak{g} \rightarrow \mathfrak{h}$ is a submetry with respect to the inner product structures from (iii).

Remark 2.3. Consider Carnot groups G and H together with a graded homomorphism $\pi: G \rightarrow H$ for which $\pi_*: \mathfrak{g} \rightarrow \mathfrak{h}$ is a submetry and $\ker(\pi_*)$ is contained in the center of \mathfrak{g} .

Definition 2.4. Let X and Y be metric spaces and $0 < \gamma \leq 1$. A mapping $f: X \rightarrow Y$ is γ -Hölder continuous if

$$[f]_{\gamma} = [f]_{C^{0,\gamma}} := \sup \left\{ \frac{d_Y(f(x), f(y))}{d_X(x, y)^{\gamma}} : x \neq y \right\} < \infty.$$

The space of all γ -Hölder continuous mappings $f : X \rightarrow Y$ will be denoted by $C^{0,\gamma}(X; Y)$.

$C_b^{0,\gamma}(X; Y)$ is the space of bounded γ -Hölder continuous mappings i.e., the image of every mapping is contained in a ball in Y .

Note that $C_b^{0,\gamma}(X; \mathbb{R}^m)$ is a Banach space with respect to the norm

$$\|f\|_{C^{0,\gamma}} = \|f\|_\infty + [f]_\gamma.$$

$C_b^{0,\gamma}(X)$ is a real Banach space. Moreover if $f, g \in C_b^{0,\gamma}(X)$, then $fg \in C_b^{0,\gamma}(X)$ and $\|fg\|_{C^{0,\gamma}} \leq \|f\|_{C^{0,\gamma}} \|g\|_{C^{0,\gamma}}$.

3 Preservation of closed horizontal curves for Lipschitz contact lift

In this section, we connect the existence of Lipschitz contact lifts to a condition on closed horizontal curves.

For any horizontal curve, we can express its location in integral form on central extension of Carnot groups.(cf. [7], Lemma 3.5])

Lemma 3.1. *Let $V \xrightarrow{\iota} G \xrightarrow{\pi} H$ be a central extension of Carnot groups by ρ such that $\text{rank}(G) = \text{rank}(H)$. There exists a 1-form $\alpha \in \Omega^1(H; V)$ such that*

(i) $d\alpha = \rho$, and

(ii) if $\gamma : [0, 1] \rightarrow G$ is a horizontal curve starting from $\gamma(0) = e_G$, then

$$\log_G(\gamma(1)) = \log_H(\pi \circ \gamma(1)) + \int_{\pi \circ \gamma} \alpha \in \mathfrak{h} \oplus V = \mathfrak{g}.$$

Hence, we can judge if a curve on Carnot group H can be lifted to a larger Carnot group G . Given a Carnot group \mathbb{H} , a point $h \in \mathbb{H}$, and a domain $U \subset \mathbb{H}$, we denote by $\Gamma_{\text{LIP}}(h, U)$ the collection of closed Lipschitz curves $\gamma : [0, 1] \rightarrow U$ based at $\gamma(0) = h = \gamma(1)$.

For a central extension $V \rightarrow \mathbb{G} \rightarrow \mathbb{H}$ of Carnot groups by a 2-cocycle $\rho \in \Omega^2(\mathbb{H}; V)$, we denote by $\Gamma_{\text{LIP}}^\rho(h, U) \subset \Gamma_{\text{LIP}}(h, U)$ the subcollection which admit a horizontal lift to \mathbb{G} that is closed. Representing \mathbb{G} as a direct product $\mathbb{G} \simeq \hat{\mathbb{G}} \times W$, where $W \subset V$ is the horizontal component of the central extension $V \rightarrow \mathbb{G} \rightarrow \mathbb{H}$, the subcollection can be characterized as

$$\Gamma_{\text{LIP}}^\rho(h, U) = \left\{ \gamma \in \Gamma_{\text{LIP}}(h, U) : \int_\gamma \alpha = 0 \right\},$$

where $\alpha \in \Omega^1(\mathbb{H}; V)$ satisfies $d\alpha = \rho$. Note that the collection is independent of the specific potential since $\int_\gamma d\psi = 0$ for any closed curve $\gamma \in \Gamma_{\text{LIP}}(h, U)$ and any exact 1-form $d\psi \in \Omega^1(\mathbb{H}; V)$.

In fact, we can use the lifting of horizontal curves on H_2 to identify if a map admits a contact lift.

Lemma 3.2. *Let $U_1 \subset \mathbb{H}_1$ and $\tilde{U}_1 \subset \pi_1^{-1}(U_1) \subset \mathbb{G}_1$ be domains. A Lipschitz contact map $f : U_1 \rightarrow \mathbb{H}_2$ admits a Lipschitz contact lift $F : \tilde{U}_1 \rightarrow \mathbb{G}_2$ if and only if*

$$(f \circ \pi_1) \left(\Gamma_{\text{LIP}}(g, \tilde{U}_1) \right) \subset \Gamma_{\text{LIP}}^\rho(f(\pi_1(g)), \mathbb{H}_2) \quad (3.1)$$

for some $g \in \tilde{U}_1$.

Proof. If $F: \tilde{U}_1 \rightarrow G_2$ is a Lipschitz contact lift of $f: U_1 \rightarrow H_2$ and $\gamma \in \Gamma_{\text{LIP}}(g, \tilde{U}_1)$ is a closed horizontal curve based at an arbitrary $g \in \tilde{U}_1$, then $F \circ \gamma$ is a closed horizontal curve in G_2 based at $F(g)$ whose projection to H_2 is the closed horizontal curve $\pi_2 \circ F \circ \gamma = f \circ \pi_1 \circ \gamma$ based at $f(\pi_1(g))$. Thus (3.1) follows.

For the converse direction, we first observe that by Lemma 4.1 in [7], it suffices to consider the case $\text{rank}(G_2) = \text{rank}(H_2)$. Suppose that (3.1) holds for some $g \in \tilde{U}_1$. We will construct the lift F by a standard path lifting argument seen. We refer the reader to Section 8 in [1] for background on the endpoint map.

Fix a basepoint $p \in \pi_2^{-1}(f(\pi_1(g)))$. We construct a Lipschitz contact lift $F: \tilde{U}_1 \rightarrow G_2$ of f with $F(g) = p$. Recall that for every control $u \in L^2([0, 1], \mathfrak{g}_1^{[1]})$, there exists a corresponding trajectory $\gamma_u: [0, 1] \rightarrow G_1$ obtained as the unique solution of the Cauchy problem

$$\dot{\gamma}_u(t) = (L_{\gamma_u(t)})_* u(t), \quad \gamma_u(0) = g.$$

This defines the endpoint map

$$\text{End}_g: L^2([0, 1]; \mathfrak{g}_1^{[1]}) \rightarrow G_1, \quad \text{End}_g(u) = \gamma_u(1),$$

which by Proposition 8.5 in [1] is Lipschitz. We will also consider the two analogously defined endpoint maps

$$\text{End}_{f(\pi_1(g))}: L^2([0, 1]; \mathfrak{h}_2^{[1]}) \rightarrow H_2,$$

$$\text{End}_p: L^2([0, 1]; \mathfrak{g}_2^{[1]}) \rightarrow G_2.$$

We have $\mathfrak{h}_2^{[1]} = \mathfrak{g}_2^{[1]}$, so these latter two endpoint maps have the same domain. We note that $\text{End}_{f(\pi_1(g))} = \pi_2 \circ \text{End}_p$ by construction.

Let $W \subset L^2([0, 1]; \mathfrak{g}_1^{[1]})$ be the open subset of controls whose trajectories satisfy $\gamma_u([0, 1]) \subset \tilde{U}_1$. Since \tilde{U}_1 is a domain, $\text{End}_p(W) = \tilde{U}_1$. The Lipschitz contact map $f \circ \pi_1: \tilde{U}_1 \rightarrow H_2$ induces a bounded map $(f \circ \pi_1)_*: W \rightarrow L^2([0, 1]; \mathfrak{h}_2^{[1]})$ given by

$$((f \circ \pi_1)_* u)(t) = (L_{f \circ \pi_1 \circ \gamma_u(t)})_* \frac{d}{dt} (f \circ \pi_1 \circ \gamma_u(t)).$$

We define $\Phi := \text{End}_p \circ (f \circ \pi_1)_*: W \rightarrow G_2$. Assumption (3.1) implies that the value $\Phi(u)$ depends only on $\text{End}_g(u)$ for $u \in W$. Thus there exists a uniquely defined map $F: \tilde{U}_1 \rightarrow G_2$ for which $F \circ \text{End}_g|_W = \Phi$.

By a perturbation argument, we see that, for any $h \in \tilde{U}_1$, there exists a control $u \in W \cap \text{End}_g^{-1}(h)$ such that the differential $(d\text{End}_g)_u$ has full rank. So, by the implicit function theorem, every $h \in \tilde{U}_1$ has an open neighborhood $W \subset \tilde{U}_1$ such that a smooth right inverse $\Psi: V \rightarrow W$ of End_g exists. Therefore we have $F|_W = \Phi \circ \Psi$. Moreover, $\pi_2 \circ F = f \circ \pi_1$. Indeed, if $\gamma: [0, 1] \rightarrow \tilde{U}_1$ is a horizontal curve, then $F \circ \gamma$ is a horizontal lift of $f \circ \pi_1 \circ \gamma$ by the construction, the equality $\text{End}_f(\pi_1(g)) = \pi_2 \circ \text{End}_p$, and the assumption , and thus the lifting property follows. It also follows that F is contact.

If $\gamma: [0, 1] \rightarrow H_1$ is a horizontal curve starting from $\gamma(0) = e_{H_1}$, then

$$\tilde{\gamma}(1) = \gamma(1) \cdot \exp\left(\int_{\gamma} \alpha\right)$$

For the contact lift F of f , then

$$F(\tilde{\gamma}(1)) = f(\gamma(1)) \cdot \exp\left(\int_{\gamma} (f \circ \pi_1)^* \alpha_2\right)$$

By the assumption, we know $\alpha_2 \in \Omega^1(H_2; V_2)$, so locally there is a constant C such that $\|\alpha_2\| \leq C$, then $\|(f \circ \pi_1)_* \alpha_2\| \leq LC$, where L is the Lipschitz constant of f . Thus, F is Lipschitz contact lift of f . \square

4 Preservation of closed horizontal curves for Sobolev contact lift

In this section, we connect the existence of Sobolev contact lifts to a condition on closed horizontal curves.

We denote by $\Gamma_{W^{1,p}}(h, U)$ the collection of closed $W^{1,p}$ admissible curves $\gamma : [0, 1] \rightarrow U$ based at $\gamma(0) = h = \gamma(1)$.

We denote by $\Gamma_{W^{1,p}}^\rho(h, U) \subset \Gamma_{W^{1,p}}(h, U)$ the subcollection which admit a horizontal lift to \mathbb{G} that is closed. The subcollection can be characterized as

$$\Gamma_{W^{1,p}}^\rho(h, U) = \left\{ \gamma \in \Gamma_{W^{1,p}}(h, U) : \int_\gamma \alpha = 0 \right\}.$$

So, we can draw a similar conclusion as Lipschitz version just for Sobolev contact lift.

Lemma 4.1. *Let $U_1 \subset \mathbb{H}_1$ and $\tilde{U}_1 \subset \pi_1^{-1}(U_1) \subset \mathbb{G}_1$ be domains. A Sobolev contact map $f : U_1 \rightarrow \mathbb{H}_2$ admits a Sobolev contact lift $F : \tilde{U}_1 \rightarrow \mathbb{G}_2$ if and only if*

$$(f \circ \pi_1) \left(\Gamma_{LIP}(g, \tilde{U}_1) \right) \subset \Gamma_{W^{1,p}}^\rho(f(\pi_1(g)), \mathbb{H}_2) \text{ for } \mathcal{H}^{Q_1-1} - \text{a.e.} \quad (4.1)$$

for some $g \in \tilde{U}_1$, where $f, F \in W^{1,p}$ and $p > Q_1$, Q_1 is homogeneous dimension of H_1 .

Proof. For any $\gamma \in \Gamma_{LIP}(g, \tilde{U}_1)$, because $F \in W^{1,p}$ and $D_p F$ is a Lie algebra homomorphism, $F \circ \gamma$ is a horizontal curve. By $D_p F \in L^p(H_1)$, we can take a tubular neighbor U_r of γ with radius r , then

$$\int_{U_r} |D_p(f)|^p < \infty$$

f is differential on almost every curve and $f \circ \pi_1 \circ \gamma \in \Gamma_{W^{1,p}}$ for $\mathcal{H}^{Q_1-1} - \text{a.e.}$ Therefore we have $\int_\gamma (f \circ \pi_1)^* \alpha_2$ exists $\mathcal{H}^{Q_1-1} - \text{a.e.}$ Due to F is contact lift $\pi_2 \circ F \circ \gamma = f \circ \pi_1 \circ \gamma$, we have $\int_\gamma (f \circ \pi_1)^* \alpha_2 = 0$ for $\mathcal{H}^{Q_1-1} - \text{a.e.}$ Thus (4.1) follows.

For the converse direction, we first observe that $f \in \Gamma_{W^{1,p}}$, $p > Q_1$, so we can take f as Hölder continuous. And also because $L^p([0, 1]) \subset L^2([0, 1])$, we can find that the conditions for endpoint maps are all satisfied for $f \circ \pi_1 \circ \gamma$ $\mathcal{H}^{Q_1-1} - \text{a.e.}$ Therefore we can use the same method as the proof of Lipschitz contact lift to get a lift F of f . Obviously, F is continuous, since $\int_{\tilde{\gamma}} (f \circ \pi_1)^* \alpha_2 = 0$ for $\mathcal{H}^{Q_1-1} - \text{a.e.}$, we can take a Lipschitz $\tilde{\gamma}$ to define F without worrying about the existence of the limit of the integral

$$F(\tilde{\gamma}(1)) = f(\gamma(1)) \cdot \exp\left(\int_{\tilde{\gamma}} (f \circ \pi_1)^* \alpha_2\right),$$

where $\tilde{\gamma}(0) = e_G$.

The last step is to prove $F \in W^{1,p}$. We can assume $f(e) = e$, so $F(e) = e$. For

almost every Lipschitz curve $\tilde{\gamma}$, $\tilde{\gamma}(0) = e$

$$\begin{aligned}
& d_{CC}(F(\tilde{\gamma}(1)), e) \\
& \leq d_{CC}(e, \exp(\int_{\tilde{\gamma}} (f \circ \pi_1)^* \alpha_2)) \\
& + d_{CC}(\exp(\int_{\tilde{\gamma}} (f \circ \pi_1)^* \alpha_2), f(\gamma(1)) \cdot \exp(\int_{\tilde{\gamma}} (f \circ \pi_1)^* \alpha_2)) \\
& = d_{CC}(e, \exp(\int_{\tilde{\gamma}} (f \circ \pi_1)^* \alpha_2)) + d_{CC}(e, f(\gamma(1))) \\
& \leq \int_{\tilde{\gamma}} |(f \circ \pi_1)^* \alpha_2| + \int_{\tilde{\gamma}} |\nabla_H f| ds \\
& = \int_{\tilde{\gamma}} (|(f \circ \pi_1)^* \alpha_2| + |\nabla_H f|) ds,
\end{aligned}$$

so $|(f \circ \pi_1)^* \alpha_2| + |\nabla_H f|$ is the upper gradient of F and it belongs to L^p . Therefore $F \in W^{1,p}(G_1; G_2)$ \square

If we want to study the contact lift of quasiconformal maps we still need to know the fiber component of a sobolev lift and the decomposition of a Lipschitz or sobolev contact lift.

Lemma 4.2. *Let $V_1 \rightarrow G_1 \rightarrow H_1$ and $V_2 \rightarrow G_2 \rightarrow H_2$ be central extensions of Carnot groups such that $\text{rank}(G_2) = \text{rank}(H_2)$. Let $U_1 \subset H_1$ be a domain and $F : \pi_1^{-1}(U_1) \rightarrow G_2$ a Sobolev contact lift of a sobolev contact map $f : U_1 \rightarrow H_2$. Then there exists a Lie group homomorphism $\Phi : V_1 \rightarrow V_2$ such that $F(gk) = F(g)\Phi(k)$ for all $g \in \pi_1^{-1}(U_1)$ and $k \in V_1$.*

Proof. By the lift assumption, we have

$$\pi_2 \circ F(gk) = f \circ \pi_1(gk) = f \circ \pi_1(g) = \pi_2 \circ F(g).$$

Thus, for each $g \in \pi_1^{-1}(U_1)$ and $k \in V_1$, there exists some $\tilde{\Phi}(g, k) \in V_2$ such that $F(gk) = F(g)\tilde{\Phi}(g, k)$. We claim that $\tilde{\Phi}(g, k)$ is constant in $g \in \pi_1^{-1}(U_1)$. That is, there is a well defined map $\Phi : V_1 \rightarrow V_2$ such that $\tilde{\Phi}(g, k) = \Phi(k)$ for $g \in \pi_1^{-1}(U_1)$ and $k \in V_1$. By assumption, U_1 is connected, so it suffices to prove that $g \mapsto \tilde{\Phi}(g, k)$ is locally constant for fixed $k \in V_1$.

Let B' be a ball that is compactly contained in $\pi_1^{-1}(U_1)$. Let B be a ball with the same center and half the radius.

Let $g, h \in B$. Since the extensions are central, elements of V_1 commute with elements of G_1 , and similarly for V_2 . By left-invariance of the distance, we compute that

$$\begin{aligned}
d(\tilde{\Phi}(g, k), \tilde{\Phi}(h, k)) &= d(F(g)\tilde{\Phi}(g, k), F(g)\tilde{\Phi}(h, k)) \\
&\leq d(F(g)\tilde{\Phi}(g, k), F(h)\tilde{\Phi}(h, k)) \\
&\quad + d(F(h)\tilde{\Phi}(h, k), F(g)\tilde{\Phi}(h, k)) \\
&= d(F(gk), F(hk)) + d(F(h), F(g))
\end{aligned}$$

Because $F \in W^{1,p}$ and $|\nabla_H F| \in L^p$ is the upper gradient of F . Taking a geodesic curve connecting h and g , then $\gamma \cdot k$ is a geodesic curve connecting hk and gk . Therefore, we have

$$\begin{aligned}
& d(F(h), F(g)) + d(F(gk), F(hk)) \\
& \leq \int_{\gamma} |\nabla_H F| + \int_{\gamma \cdot k} |\nabla_H F| \\
& \leq \int_{\gamma} |\nabla_H F| + \int_{\gamma} |\nabla_H F| \circ r_k,
\end{aligned}$$

where r_k is the right multiplication by k . Therefore, the $\int_\gamma |\nabla_H F| + \int_\gamma |\nabla_H F| \circ r_k$ is the upper gradient of $\tilde{\Phi}(g, k)$, which means $\tilde{\Phi}(g, k) \in W^{1,p}$. By the equal rank assumption, V_2 does not contain any horizontal directions. Hence the only possible sobolev maps $B \rightarrow V_2$ are constants. Consequently, $g \mapsto \tilde{\Phi}(g, k)$ is locally constant for $k \in V_1$.

Next, we show that $\Phi : V_1 \rightarrow V_2$ is a group homomorphism. Let $k_1, k_2 \in V_1$ and let $g \in \pi_1^{-1}(U_1)$. Applying the definition of $\tilde{\Phi}$ with the pair $(g, k_1 k_2)$ shows

$$F(gk_1 k_2) = F(g)\Phi(k_1 k_2),$$

whereas applying the definition with the pairs (gk_1, k_2) and (g, k_1) yields

$$F(gk_1 k_2) = F(gk_1)\Phi(k_2) = F(g)\Phi(k_1)\Phi(k_2).$$

Cancelling out $F(g)$ implies that Φ is a group homomorphism.

Finally, we observe that $\Phi(k) = F(g)^{-1}F(gk)$ for any $g \in G_1$ and $k \in V_1$, so Φ is a $W^{1,p}$ map. \square

If we want to study the Pansu differential of a contact lift, the first thing is to study the Lie algebra homomorphism between central extension sequences.

Lemma 4.3. *Let $V_1 \rightarrow \mathfrak{g}_1 \rightarrow \mathfrak{h}_1$ and $V_2 \rightarrow \mathfrak{g}_2 \rightarrow \mathfrak{h}_2$ be central extensions by 2-cocycles ρ_1 and ρ_2 , respectively. Let $L : \mathfrak{h}_1 \rightarrow \mathfrak{h}_2$ be a Lie algebra homomorphism and let $\varphi : V_1 \rightarrow V_2$ be a linear map.*

(i) *There exists a Lie algebra homomorphism $\psi : \mathfrak{g}_1 \rightarrow \mathfrak{g}_2$ such that the diagram*

$$\begin{array}{ccccc} V_1 & \longrightarrow & \mathfrak{g}_1 & \xrightarrow{\pi_1} & \mathfrak{h}_1 \\ \varphi \downarrow & & \downarrow \psi & & \downarrow L \\ V_2 & \longrightarrow & \mathfrak{g}_2 & \xrightarrow{\pi_2} & \mathfrak{h}_2 \end{array}$$

commutes if and only if there exists a linear map $\mu : \mathfrak{h}_1 \rightarrow V_2$ such that $\varphi \circ \rho_1 - L^ \rho_2 = d_0 \mu$. When such a μ exists, a homomorphism ψ is given by*

$$\psi(X + Y) = L(X) + (\mu(X) + \varphi(Y)) \in \mathfrak{h}_2 \oplus V_2 = \mathfrak{g}_2 \quad (4.2)$$

for $X + Y \in \mathfrak{h}_1 \oplus V_1 = \mathfrak{g}_1$.

(ii) *When the central extensions are stratified and φ and L are graded, a graded homomorphism ψ is unique up to a graded linear map $\theta : \mathfrak{h}_1 \rightarrow V_2$ with $[\mathfrak{h}_1, \mathfrak{h}_1] \subset \ker \theta$. In particular, if $\mathfrak{h}_2^{[1]} = \mathfrak{g}_2^{[1]}$, then the graded homomorphism ψ is unique.*

Because Φ is Lie group homomorphism and V_1, V_2 are central, so Φ is a linear map.

$$\Phi(2k) = \Phi(k + k) = \Phi(k) + \Phi(k) = 2\Phi(k)$$

Let $g \in \pi_1^{-1}(U_1)$. Computing the Pansu derivative in a direction $h \in V_1$, we obtain

$$\begin{aligned} d_P F(g)h &= \lim_{\lambda \rightarrow 0^+} \delta_{1/\lambda} (F(g)^{-1}F(g\delta_\lambda h)) \\ &= \lim_{\lambda \rightarrow 0^+} \delta_{1/\lambda} (F(g)^{-1}F(g)\Phi(\delta_\lambda h)) \\ &= \lim_{\lambda \rightarrow 0^+} \delta_{1/\lambda} (\Phi(\delta_\lambda h)) =: \varphi h \end{aligned}$$

Since the projections π_1 and π_2 are homomorphisms, we have $\pi_2 \circ d_P F(g) = d_P f(\pi_1 g) \circ \pi_1$. This gives us the commutative diagram:

$$\begin{array}{ccccc} V_1 & \longrightarrow & G_1 & \xrightarrow{\pi_1} & H_1 \\ \varphi \downarrow & & \downarrow d_P F(g) & & \downarrow d_P f(\pi_1 g) \\ V_2 & \longrightarrow & G_2 & \xrightarrow{\pi_2} & H_2 \end{array}$$

By Lemma 4.3, such a diagram implies that the corresponding Lie algebra homomorphisms satisfy

$$\varphi \circ \rho_1 - d_P f(h)^* \rho_2 = d_0 \mu_h$$

for some linear map $\mu_h : \mathfrak{h}_1 \rightarrow V_2$. Thus, $\mu_h = \varphi \circ \alpha_1 - d_P f(h)^* \alpha_2$ and

$$D_P F = \begin{bmatrix} D_P f & 0 \\ \varphi \circ \alpha_1 - d_P f(h)^* \alpha_2 & \varphi \end{bmatrix} \quad (4.3)$$

With the result above, we know the Pansu differential for the lift of a quasiconformal contact map and get the lift is also a quasiconformal contact map when V_2 only has trivial horizontal part.

Lemma 4.4. *Let $U_1 \subset \mathbb{H}_1$ and $\tilde{U}_1 \subset \pi^{-1}(U_1) \subset \mathbb{G}_1$ be domains. When $\text{rank}(\mathbb{G}_2) = \text{rank}(\mathbb{H}_2)$, a quasiconformal contact map $f : U_1 \rightarrow \mathbb{H}_2$ admits a quasiconformal contact lift $F : \tilde{U}_1 \rightarrow \mathbb{G}_2$ if and only if this quasiconformal contact map $f : U_1 \rightarrow \mathbb{H}_2$ admits a sobolev contact lift $F : \tilde{U}_1 \rightarrow \mathbb{G}_2$.*

Proof. We know that a quasiconformal contact map is a W^{1,Q_1} map. Thus, we just need to prove the restriction of $D_P F$ to the horizontal layer is related to a quasiconformal map between Euclidean Spaces. Because of (4.3), the restriction of $D_P F$ to the horizontal layer only depends on the restriction of $D_P f$ to the horizontal layer. So, the sobolev contact lift F is also quasiconformal contact lift. \square

Theorem 4.5. *Suppose that $f_P^* \rho_2 = \varphi \circ \rho_1 + d_0 \omega$ for a graded linear map $\varphi : V_1 \rightarrow V_2$ and a 1-form $\omega \in \Omega^1(U_1; V_2)$. If the weight of ρ_2 is equal to or greater than the maximal weight of non-trivial 2-cocycles in the Rumin complex E_0 of \mathbb{H}_1 and U_1 is simply connected, then f admits a smooth contact lift $F : \mathbb{G}_1 \supset \pi_1^{-1}(U_1) \rightarrow \mathbb{G}_2$.*

Example 4.6. For Heisenberg group H_1

$$\begin{array}{ccc} R & \longrightarrow & H_1 \xrightarrow{\pi} \mathbb{R}^2 \\ & & \downarrow F \\ R & \longrightarrow & H_1 \xrightarrow{\pi} \mathbb{R}^2 \end{array}$$

We know that for Heisenberg group H_1 if $\varphi \circ \alpha_1 - d_P f(h)^* \alpha_2 = 0$ then there is a contact lift F of f . That is $\varphi = \det(Df)$ is a constant. For instance, $f(a, b) = (a + b^2, b)$ admits a smooth contact lift.

$$Df = \begin{bmatrix} 1 & 2b \\ 0 & 1 \end{bmatrix}$$

5 Rigidity of contact lift

In this chapter we find that all of the quasiconformal contact maps which admit contact lifts are bi-Lipschitz by the rigidity of the lift. Especially, for Carnot groups with rigid structure, the possible maps are very few.

Theorem 5.1. *Suppose that f is a K -quasiconformal contact map and it admits a contact lift F then f is bi-Lipschitz.*

Proof.

$$\begin{array}{ccc} V_1 & \longrightarrow & G_1 \xrightarrow{\pi} H_1 \\ \varphi \downarrow & & \downarrow F \\ V_1 & \longrightarrow & G_1 \xrightarrow{\pi} H_1 \end{array}$$

First of all, since Lemma 4.4 F is a quasiconformal map so that the dimensions of image of F must be the same as G_1 so as to φ is non-degenerate.

Let's begin with the easy condition when the central extension is for the highest layer, so for Case 1,2,3 we can directly compute by the restriction of φ .

Case 1: $V_1 = \mathbb{R}$ and $\mathfrak{h}_1 = \mathbb{R}^k$, so the central extension G_1 is a step 2 Carnot group. we know $V_1 = \text{span}\{\alpha\}$ and

$$\alpha = \sum_{i < j} a_{ij} dx_i \wedge dx_j$$

After using the following method below,

$$\begin{aligned} & dx_1 \wedge dx_2 + dx_1 \wedge dx_3 + dx_2 \wedge dx_3 \\ &= dx_1 \wedge (dx_2 + dx_3) + dx_2 \wedge (dx_2 + dx_3) \\ &= (dx_1 + dx_2) \wedge (dx_2 + dx_3) \end{aligned}$$

we can find a new basis of α like $\{dx_1 + dx_2, dx_2 + dx_3\}$ and note that as $\{d\tilde{x}_1, d\tilde{y}_2\}$. Thus, the α can be written as the form

$$\alpha = \sum_i b_i d\tilde{x}_i \wedge d\tilde{y}_i.$$

Because V_1 is the second layer, $\varphi \circ \alpha = \lambda \alpha$ and $f_P^* \alpha = \lambda \alpha$.

$$\sum_i b_i f_P^*(d\tilde{x}_i \wedge d\tilde{y}_i) = \sum_i b_i \lambda (d\tilde{x}_i \wedge d\tilde{y}_i)$$

Restricting the Pansu differential to horizontal layer, we can get $\|D_H f\| \approx |\lambda|^{\frac{1}{2}}$.

From the assumption of K -QC map f , we know f is bi-Lipschitz, since

$$|\lambda|^{\frac{1}{2}} K^{-1} \|X\| \leq |D_H f(X)| \leq |\lambda|^{\frac{1}{2}} K \|X\|$$

for horizontal vector X .

Case 2: If $V_1 = V_1^{[p+1]} = \text{span}\{\alpha\}$ and $\mathfrak{h}_1 = \mathfrak{h}_1^{[1]} \oplus \dots \oplus \mathfrak{h}_1^{[p]}$, for $p \geq 2$.

Due to $d_0 \rho = 0$,

$$\begin{aligned} & \rho([X_1, X_2], X_3) + \rho([X_2, X_3], X_1) + \rho([X_3, X_1], X_2) = 0 \\ & \rho(\mathfrak{h}_1^{[i]}, \mathfrak{h}_1^{[j]}) \subset \rho(\mathfrak{h}_1^{[i-1]}, \mathfrak{h}_1^{[j+1]}) \subset \dots \subset \rho(\mathfrak{h}_1^{[1]}, \mathfrak{h}_1^{[i+j-1]}) \end{aligned}$$

α has the form

$$\alpha = \sum_i b_i d\tilde{x}_i \wedge d\tilde{y}_i$$

where $d\tilde{x}_i \in (H_1^{[1]})^*$ and $d\tilde{y}_i \in (H_1^{[p]})^*$ with the weight $\text{wt}(d\tilde{x}_i) = -1$ and $\text{wt}(d\tilde{y}_i) = -p$.

Because V_1 is the highest layer, $\varphi \circ \alpha = \lambda \alpha$ and $f_P^* \alpha = \lambda \alpha$ so that we can get $\|D_H f\| \approx |\lambda|^{\frac{1}{p+1}}$.

Case 3: If $V_1 = V_1^{[p+1]} = \text{span}\{\alpha_1 \dots \alpha_j\}$ and $\mathfrak{h}_1 = \mathfrak{h}_1^{[1]} \oplus \dots \oplus \mathfrak{h}_1^{[p]}$ for $p \geq 2$.

So the same as Case 2, we can get $f_P^* \alpha = \varphi \circ \alpha$. When $D_P f$ is non-degenerate on $\mathfrak{h}_1^{[1]}$, $f_{P*}(\mathfrak{h}_1^{[1]}) = \mathfrak{h}_1^{[1]}$. Using all the vectors on $\mathfrak{h}_1^{[1]}$, they generate the whole Lie algebra \mathfrak{g}_1 , so $f_{P*}(\mathfrak{g}_1^{[1]}) = \mathfrak{g}_1^{[1]}$.

If $\|f_{P*}(X)\| \approx r \|X\|$ for horizontal vector X ,

$$\begin{aligned} \|f_{P*}([X_1, X_2])\| &= \|[f_{P*}(X_1), f_{P*}(X_2)]\| \\ &\leq 2 \|f_{P*}(X_1)\| \cdot \|f_{P*}(X_2)\| \\ &\lesssim 2r^2 \|X_1\| \cdot \|X_2\| \end{aligned}$$

where X_1 and X_2 are horizontal vectors. And also because f_{P*} is non-degenerate, there is a lower bound of $f_{P*}([\cdot, \cdot])$, combined with $D_P f$ commuting with dilation δ_r , therefore, if the horizontal layer is expanded by r times, then the k -th layer is expanded by r^k times. If $\|\varphi\| \approx \lambda$, then $\|D_H f\| \approx |\lambda|^{\frac{1}{p+1}}$.

Case 4: the central extension is not to add the highest layer, so the structure of $D_P F$ of the lifting F is not good enough. We need to estimate the determinant of $D_H f$ in a more general way which depends on the existence of the quasiconformal contact lift.

Because of Lemma 4.4, f admits a quasiconformal contact lift F and

$$D_P F|_{\mathfrak{g}_1^{[1]}} = \varphi|_{\mathfrak{g}_1^{[1]}} \subset \mathfrak{g}_1^{[1]}.$$

The homogeneous dimension of $\mathfrak{g}_1^{[1]}$ is Q' . If $\det(\varphi) = \lambda$, then $\|D_H f\| \approx \lambda^{\frac{1}{Q'}}$. \square

The result is not finished here, for some Carnot groups with rigid structure, we can even draw a better conclusion. For example, if F is a contact lift on quaternionic Heisenberg group $\mathbb{H}_{\mathbb{H}}^n$ of a map f on \mathbb{R}^{4n} , then $f : \mathbb{R}^{4n} \rightarrow \mathbb{R}^{4n}$ is a affine transformation.

Theorem 5.2. *Suppose f is a Sobolev continuous map on \mathbb{R}^{4n} which admits a contact lift F on $\mathbb{H}_{\mathbb{H}}^n$ and $D_P F$ is full-rank for every point, then f is an affine transformation.*

Proof. By assumption, F is a contact lift and its Pansu differential is

$$D_P F = \begin{bmatrix} Df & \\ & \varphi \end{bmatrix}$$

for some constant matrix φ .

$$\varphi = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

We define central extension as $\rho = (\rho_1, \rho_2, \rho_3)$,

$$\begin{aligned} \rho_1 &= \sum_{i=1}^n dx_i \wedge dy_i + dz_i \wedge dw_i \\ \rho_2 &= \sum_{i=1}^n dx_i \wedge dz_i + dw_i \wedge dy_i \\ \rho_3 &= \sum_{i=1}^n dx_i \wedge dw_i + dy_i \wedge dz_i. \end{aligned}$$

For any fixed point q , $D_P F(q)$ is a group homomorphism, because $D_P F$ is invertible for every point, we can define a map $\tilde{F} = F \circ D_P F(q)^{-1}$ with Pansu differential

$$D_P \tilde{F} = \begin{bmatrix} Df \circ (Df(q))^{-1} & \\ & id \end{bmatrix} \quad (5.1)$$

Obviously, \tilde{F} is also a contact lift, since $\tilde{f}^*(\alpha) = id \circ \alpha$ for constant map id , where $\tilde{f} = f \circ (Df(q))^{-1}$. Thus, $\tilde{F}_P^*(\rho) = \tilde{f}^*(\rho) = \rho$, which means \tilde{f} is a symplectic forms preserving map. This would greatly limit the possible choice of f .

The symplectic forms ρ_1, ρ_2, ρ_3 can be represent as the matrix J_1, J_2, J_3 in local coordinates.

$$J_1 = \begin{bmatrix} & I_n & \\ -I_n & & \\ & -I_n & I_n \end{bmatrix}$$

$$J_2 = \begin{bmatrix} & I_n & \\ -I_n & & \\ & -I_n & I_n \end{bmatrix}$$

$$J_3 = \begin{bmatrix} & & -I_n \\ & I_n & \\ I_n & & \end{bmatrix}$$

where satisfying $J_3 = J_1 J_2$. We denote $D\tilde{f} := A$. Due to map \tilde{f} preserving symplectic forms, we have

$$A^T J_i A = J_i \quad \text{for } i = 1, 2, 3.$$

and

$$\begin{aligned} A^T J_3 A &= J_3 \\ A^T (J_1 J_2) A &= J_1 J_2 \\ A^T (J_1 J_2) A &= A^T J_1 A A^T J_2 A \\ J_1 J_2 &= J_1 A A^T J_2 \\ I &= A A^T, \end{aligned}$$

since A is invertible. Thus, A is orthogonal.

$$\begin{aligned} A^T J_i A &= J_i \\ J_i A &= J_i A \quad \text{for } i = 1, 2, 3, \end{aligned}$$

which means $J_i \tilde{f}_* = \tilde{f}_* J_i$ and $\tilde{f}^*(\rho_i) = \rho_i$ for $i = 1, 2, 3$. Therefore, related to metric g on \mathbb{R}^{4n} ,

$$\begin{aligned} g(\tilde{f}_*(X), \tilde{f}_*(Y)) &= \rho_i(\tilde{f}_*(X), J_i \tilde{f}_*(Y)) \\ &= \rho_i(\tilde{f}_*(X), \tilde{f}_*(J_i Y)) \\ &= (\tilde{f}^* \rho_i)(X, J_i Y) \\ &= \rho_i(X, J_i Y) \\ &= g(X, Y). \end{aligned}$$

\tilde{f} is isometric on \mathbb{R}^{4n} . So, the only possible is \tilde{f} is an affine transformation as well as $f = \tilde{f} \circ Df(q)$.

□

Remark 5.3. If f is a quasiconformal map on \mathbb{R}^{4n} with a contact lift on $\mathbb{H}_{\mathbb{H}}^n$, then because of theorem 4.1 it's bi-Lipschitz and $D_P F$ is full-rank. Therefore, this kind of f would naturally satisfy the condition of theorem 4.2.

Remark 5.4. For quaternionic Heisenberg group $\mathbb{H}_{\mathbb{H}}^n$, its Lie group structure has more delicate symmetrical structure. To be specific, continue to use the notations from the theorem 4.2, $\rho_1^{2n} = \rho_2^{2n} = \rho_3^{2n} = 4^n dx_1 \wedge \cdots \wedge dx_n \wedge dy_1 \wedge \cdots \wedge dy_n \wedge dz_1 \wedge \cdots \wedge dz_n \wedge dw_1 \wedge \cdots \wedge dw_n$. From

$$\tilde{f}^*(\rho_1^{2n}) = \tilde{f}^*(\rho_2^{2n}) = \tilde{f}^*(\rho_3^{2n}),$$

we have

$$\tilde{f}^*(\rho_1)^{2n} = \tilde{f}^*(\rho_2)^{2n} = \tilde{f}^*(\rho_3)^{2n}$$

so that

$$a_{11}^{2n} + a_{12}^{2n} + a_{13}^{2n} = a_{21}^{2n} + a_{22}^{2n} + a_{23}^{2n} = a_{31}^{2n} + a_{32}^{2n} + a_{33}^{2n}.$$

However, if the structure of V in central extension is not rich enough, the contact lifts on the Carnot group are abundant.

Example 4.5.: Consider the real filiform Carnot group F^3 of step 3. The corresponding Lie algebra is given by

$$\mathfrak{f}^s = \text{span}(X_1, X_2) \oplus \text{span}(X_3) \oplus \text{span}(X_4),$$

where $[X_1, X_2] = X_3$ and $[X_1, X_3] = X_4$ are the only non-trivial relations. Consider the corresponding dual basis $\omega_1, \omega_2, \omega_3, \omega_4$.

$$\begin{aligned} X_1 &= \frac{\partial}{\partial x_1} \\ X_2 &= \frac{\partial}{\partial x_2} + x_1 \frac{\partial}{\partial x_3} + \frac{x_1^2}{2} \frac{\partial}{\partial x_4} \\ X_3 &= \frac{\partial}{\partial x_3} + x_1 \frac{\partial}{\partial x_4} \\ X_4 &= \frac{\partial}{\partial x_4} \end{aligned}$$

and

$$\begin{aligned} \omega_1 &= dx_1 \\ \omega_2 &= dx_2 \\ \omega_3 &= -x_1 dx_2 + dx_3 \\ \omega_4 &= \frac{x_1^2}{2} dx_2 - x_1 dx_3 + dx_4 \end{aligned}$$

(i) The easy way to find contact lifts is using central extension twice. The map $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ admits a contact lift F on Heisenberg group \mathbb{H}^1 if and only if f is a area-preserving map. If we want to lift F onto F^3 as $G : F^3 \rightarrow F^3$, its Pansu differential are supposed to be such form:

$$D_P G = \begin{bmatrix} a & b & & \\ 0 & \frac{1}{a^2} & & \\ & & \frac{1}{a} & \\ & & & 1 \end{bmatrix}$$

where a is any constant and for convenience we take $\varphi = id : \text{span}(X_4) \rightarrow \text{span}(X_4)$. Therefore, because b can be any continuous function, we can find plenty of maps f and F satisfying the condition.

(ii) The second way is more complicate. By applying the same method in [9] as Kor'anyi-Reimann theorem, we can use vector field to generate a one-parameter family of contact lift related to central extension $\mathbb{R} \rightarrow F^3 \rightarrow F^2$.

For vector field $V = p_1 X_1 + p_2 X_2 + p_3 X_3 + p_4 X_4$, as a generator of contact lift, it need to satisfy $\mathcal{L}_V \omega_4 = 0$ for a group homomorphism as initial condition. As the motion along V , every f_s would preserve ω_4 , which means it's a contact lift.

$$\begin{aligned} \mathcal{L}_V \omega_4 &= d(\iota_V \omega_4) + \iota_V (d\omega_4) \\ &= dp_4 + p_3 dx_1 + x_1 p_1 dx_2 - p_1 dx_3 \\ &= \left(\frac{\partial}{\partial x_1} p_4 + p_3 \right) dx_1 + \left(\frac{\partial}{\partial x_2} p_4 + x_1 p_1 \right) dx_2 + \left(\frac{\partial}{\partial x_3} p_4 - p_1 \right) dx_3 + \frac{\partial}{\partial x_4} p_4 dx_1 \end{aligned}$$

Therefore, the vector field satisfies

$$\begin{cases} p_1 = \frac{\partial}{\partial x_3} p_4 \\ x_1 p_1 = \frac{\partial}{\partial x_2} p_4 \\ p_3 = -\frac{\partial}{\partial x_1} p_4 \\ 0 = \frac{\partial}{\partial x_4} p_4 \end{cases} \quad (5.2)$$

and this kind of vector fields are abundant.

6 The approximation of contact lift on Heisenberg group

6.1 Sobolev contact lift

We can construct a family of new contact lifts in two ways, the first is to Constructe through mapped composition.

If f_1, f_2 admit contact lift F_1, F_2 , then $F_2 \circ F_1$ is the contact lift of $f_2 \circ f_1$, because we have the commutative diagram:

$$\begin{array}{ccccc} V_1 & \longrightarrow & G_1 & \xrightarrow{\pi_1} & H_1 \\ \Phi \downarrow & & \downarrow F_1 & & \downarrow f_1 \\ V_2 & \longrightarrow & G_2 & \xrightarrow{\pi_2} & H_2 \\ \Phi \downarrow & & \downarrow F_2 & & \downarrow f_2 \\ V_3 & \longrightarrow & G_3 & \xrightarrow{\pi_3} & H_3 \end{array}$$

and $F_2 \circ F_1(gk) = F_2(F_1(g)\Phi_1(k)) = F_2 \circ F_1(g) \cdot \Phi_2 \circ \Phi_1(k)$. It still holds under a limited number of composite operations.

The second is to find a family of smooth contact maps which adimts smooth contact lifts as the approximation. However, because the the space of Lie algebra homomorphism is not a vector space, we are not able to use convolution to construct Pansu differential smooth approximation functions. However, we can study the lifting F on Heisenberg group, due to f as the continuous function on \mathbb{R}^n .

Lemma 6.1. Suppose f and f_ε in $W^{1,p}(H_1; H_2)$ admit contact lift F and F_ε in $W^{1,p}(G_1; G_2)$, if $f_\varepsilon \rightarrow f$ in C^0 then $F_\varepsilon \rightarrow F$ in C^0 .

Proof. For Lipschitz curve $\tilde{\gamma}$,

$$\begin{aligned} & d(F_\varepsilon(\tilde{\gamma}(1)), F(\tilde{\gamma}(1))) \\ &= d(f(\gamma(1)) \cdot \exp(\int_\gamma (f \circ \pi_1)^* \alpha_2), f_\varepsilon(\gamma(1)) \cdot \exp(\int_\gamma (f_\varepsilon \circ \pi_1)^* \alpha_2)) \\ &\leq d(f(\gamma(1)), f_\varepsilon(\gamma(1))) + d(\exp(\int_\gamma (f \circ \pi_1)^* \alpha_2), \exp(\int_\gamma (f_\varepsilon \circ \pi_1)^* \alpha_2)) \\ &\rightarrow 0 \end{aligned} \quad (6.1)$$

□

For Heisenberg group H_n ,

$$\begin{array}{ccc} R & \longrightarrow & H_n \xrightarrow{\pi} \mathbb{R}^{2n} \\ & & \downarrow F \\ R & \longrightarrow & H_n \xrightarrow{\pi} \mathbb{R}^{2n} \end{array}$$

Taking a Lipschitz curve γ in \mathbb{R}^{2n}

$$\begin{aligned} \gamma^z(1) - \gamma^z(0) &= \sum_{i=a}^n \int_{\gamma} x_i dy_i - y_i dx_i \\ &= \sum_{i=a}^n \int_{D_i} dx_i \wedge dy_i, \end{aligned}$$

where D_i is the area in the projection of the closed curve to $x_i O y_i$ plane which connects γ and the line segment from 0 to $\gamma(1)$. If a sobolev map f admits a lift F , then $f_*(Z) = \lambda Z$ for some constant λ , which is equivalent to $f^*(\omega) = \lambda \omega$ where ω is the standard symplectic form on \mathbb{R}^{2n} . The Pansu differential of F is

$$D_P F = \begin{bmatrix} A & \\ & \lambda \end{bmatrix}$$

We can write the Lie algebra as follow:

$$\begin{aligned} [X, Y] &= X^T J Y \\ \lambda [X, Y] &= f_*([X, Y]) \\ &= [f_*(X), f_*(Y)] \\ &= [AX, AY] \\ &= X^T A^T J A Y \end{aligned}$$

where J is the standard symplectic matrix $J = \begin{bmatrix} 0 & I_n \\ -I_n & 0 \end{bmatrix}$. So, $\lambda J = A^T J A$.

$$\lambda^{2n} \det(J) = \det(A^T) \det(J) \det(A)$$

and $\lambda = (\det(A))^{1/n}$.

The next step is to construct a series of smooth g_ε which admit smooth contact lift G_ε and $g_\varepsilon \rightarrow f$ as $\varepsilon \rightarrow 0$. For continuous f , we can use convolution to find smooth functions f_ε and $f_\varepsilon \rightarrow f$ as $\varepsilon \rightarrow 0$, but for each f_ε it can not preserve symplectic form.

using the Moser's trick to construct geometric flow, we define the error terms

$$\eta_\varepsilon = f_\varepsilon^* \omega - \omega.$$

We want to find diffeomorphism $\Psi_\varepsilon : \mathbb{R}^{2n} \rightarrow \mathbb{R}^{2n}$, such that $g_\varepsilon = f_\varepsilon \circ \Psi_\varepsilon$ and $g_\varepsilon^* \omega = \Psi_\varepsilon^*(f_\varepsilon^* \omega) = \omega$. Therefore, $\Psi_\varepsilon^*(\omega + \eta_\varepsilon) = \Psi_\varepsilon^*(f_\varepsilon^* \omega) = \omega$.

Using vector field $X_{\varepsilon,t}$ define a flow $\phi_{\varepsilon,t}$, such that

$$\frac{d}{dt}(\phi_{\varepsilon,t}) = X_{\varepsilon,t} \circ \phi_{\varepsilon,t}, \quad \phi_{\varepsilon,0} = id \text{ and } \phi_{\varepsilon,1} = \Psi_\varepsilon$$

Considering $\omega_t = \omega + t\eta_\varepsilon$, $t \in [0, 1]$, then diffeomorphism $\phi_{\varepsilon,t}$ such that

$$\phi_{\varepsilon,t}^* \omega_t = \omega$$

We need to find $X_{\varepsilon,t}$ satisfying $\mathcal{L}_{X_{\varepsilon,t}}\omega_t = -\eta_\varepsilon$, then

$$\begin{aligned}\frac{d}{dt}(\phi_{\varepsilon,t}^*\omega_t) &= \phi_{\varepsilon,t}^*(\mathcal{L}_{X_{\varepsilon,t}}\omega_t + \frac{d}{dt}\omega_t) \\ &= \phi_{\varepsilon,t}^*(-\eta_\varepsilon + \eta_\varepsilon) \\ &= 0\end{aligned}$$

and

$$\Psi_\varepsilon^*(\omega + \eta_\varepsilon) = \phi_{\varepsilon,1}^*\omega_1 = \phi_{\varepsilon,0}^*\omega_0 = (id)^*\omega = \omega.$$

So, every g_ε admit a smooth contact lift and $g_\varepsilon \rightarrow f$ as $\varepsilon \rightarrow 0$. The rest is to prove the equation $\mathcal{L}_{X_{\varepsilon,t}}\omega_t = -\eta_\varepsilon$ is solvable.

Because of $d\omega_t = 0$,

$$\mathcal{L}_{X_{\varepsilon,t}}\omega_t = d(\iota_{X_{\varepsilon,t}}\omega_t) + \iota_{X_{\varepsilon,t}}(d\omega_t) = d(\iota_{X_{\varepsilon,t}}\omega_t),$$

it suffices to show $d(\iota_{X_{\varepsilon,t}}\omega_t) = -\eta_\varepsilon$ is solvable. For simply connected domain, closed from η_ε is exact. Thus, there is a 1-from $\alpha_{\varepsilon,t}$ such that $\alpha_{\varepsilon,t} = \iota_{X_{\varepsilon,t}}\omega_t$. For ε small enough, $\eta_\varepsilon = f_\varepsilon^*\omega - \omega$ is closed to 0 in L^p so that ω_t is closed to ω for any $t \in [0, 1]$, thus ω_t is a a.e. non-degenerated 2-form and $X \rightarrow \iota_X\omega_t$ is a isomorphism from TM to T^*M . Therefore, we can find $X_{\varepsilon,t}$ satisfying the condition in L^p .

Theorem 6.2. *For any sobolev map f on \mathbb{R}^{2n} and its contact lift F on \mathbb{H}^n , there are a serious of smooth g_ε which admit smooth contact lift G_ε , $g_\varepsilon \rightarrow f$ and $G_\varepsilon \rightarrow F$ as $\varepsilon \rightarrow 0$.*

Proof. The result follows by the proof above and Lemma 6.1. \square

6.2 Hölder lift

For Hölder lift, there is also a similar result about decomposion. At first, we need to clear the definition of Hölder lift. Without considering the differential structure, we just require the lifting F of f satisfying $f \circ \pi_1 = \pi_2 \circ F$ that is to say the following diagram commutes

$$\begin{array}{ccc} V_1 & \longrightarrow & G_1 & \xrightarrow{\pi_1} & H_1 \\ & & \downarrow F & & \downarrow f \\ V_2 & \longrightarrow & G_2 & \xrightarrow{\pi_2} & H_2 \end{array}$$

Before diving into the lifting, we are supposed to show a lemma for Hölder continuous maps between Carnot group, which establishes the anisotropic structure of Carnot groups.

For convenience, we just take step three Carnot group as an example to showcase the result, as for higher step Carnot groups the conclusion holds as well.

Let G be a step three Carnot group. Let d_1, \dots, d_r be a basis for $\mathfrak{g}_1, e_1, \dots, e_s$ a basis for \mathfrak{g}_2 , and f_1, \dots, f_t a basis for \mathfrak{g}_3 . We write

$$\begin{aligned}[d_i, d_j] &= \sum_{k=1}^s \alpha_k^{ij} e_k \\ [d_i, e_k] &= \sum_{m=1}^t \beta_m^{ik} f_m\end{aligned}$$

with all other bracket relations trivial. As in the step two case, we can identify G with \mathbb{R}^{r+s+t} equipped with the following operation via coordinates of the first kind:

$$(A_i, B_k, C_m) \star (a_i, b_k, c_m) = (\mathcal{A}_i, \mathcal{B}_k, \mathcal{C}_m)$$

where

$$\begin{aligned}\mathcal{A}_i &= A_i + a_i \\ \mathcal{B}_k &= B_k + b_k + \frac{1}{2} \sum_{i < j} \alpha_k^{ij} (A_i a_j - a_i A_j) \\ \mathcal{C}_m &= C_m + c_m + \frac{1}{2} \sum_{i,j} \beta_m^{ij} (A_i b_j - B_j a_i) + \frac{1}{12} \sum_{l,k} \sum_{i < j} (A_l - a_l) \alpha_k^{ij} (A_i a_j - a_i A_j) \beta_m^{lk}\end{aligned}$$

Observe $(A_i, B_k, C_m)^{-1} = (-A_i, -B_k, -C_m)$ and

$$(A_i, B_k, C_m)^{-1} \star (a_i, b_k, c_m) = (\tilde{\mathcal{A}}_i, \tilde{\mathcal{B}}_k, \tilde{\mathcal{C}}_m),$$

where

$$\begin{aligned}\tilde{\mathcal{A}}_i &= a_i - A_i \\ \tilde{\mathcal{B}}_k &= b_k - B_k - \frac{1}{2} \sum_{i < j} \alpha_k^{ij} (A_i a_j - a_i A_j) \\ \tilde{\mathcal{C}}_m &= c_m - C_m - \frac{1}{2} \sum_{i,j} \beta_m^{ij} (A_i b_j - B_j a_i) + \frac{1}{12} \sum_{l,k} \sum_{i < j} (A_l + a_l) \alpha_k^{ij} (A_i a_j - a_i A_j) \beta_m^{lk}\end{aligned}\tag{6.2}$$

Left-translating the canonical basis at the origin, we obtain the left-invariant vector fields

$$\begin{aligned}X^i &= \frac{\partial}{\partial A_i} - \frac{1}{2} \sum_{k=1}^s \sum_{i < j} \alpha_k^{ij} A_j \frac{\partial}{\partial B_k} \\ &\quad + \sum_{m=1}^t \left[-\frac{1}{2} \sum_{j=1}^s B_j \beta_m^{ij} + \frac{1}{12} \sum_{l=1}^r \sum_{k=1}^s A_l \left(\sum_{i < j} \alpha_k^{ij} A_j \right) \beta_m^{lk} \right] \frac{\partial}{\partial C_m} \\ Y^k &= \frac{\partial}{\partial B_k} + \sum_{m=1}^t \left(\frac{1}{2} \sum_{i=1}^r \beta_m^{ik} A_i \right) \frac{\partial}{\partial C_m} \\ Z^m &= \frac{\partial}{\partial C_m}\end{aligned}$$

It is clear that $\{X^i\}_i \cup \{Y^k\}_k \cup \{Z^m\}_m$ forms a basis for $\text{Lie}(\mathbb{R}^n, \star)$. Moreover, we have the expected step three stratification of $\text{Lie}(\mathbb{R}^n, \star)$:

$$\text{Lie}(\mathbb{R}^{r+s+t}, \star) = \langle X^i \rangle_{1 \leq i \leq r} \oplus \langle Y^k \rangle_{1 \leq k \leq s} \oplus \langle Z^m \rangle_{1 \leq m \leq t}$$

In fact, one can show using the Jacobi identity that the linear map $\varphi : \text{Lie}(\mathbb{R}^n, \star) \rightarrow \mathfrak{g}$ induced by

$$X^i \mapsto d_i, \quad Y^k \mapsto e_k, \quad Z^m \mapsto f_m$$

is a Lie algebra isomorphism. The contact forms are given by

$$\begin{aligned}\omega_1^k &:= dB_k - \frac{1}{2} \sum_{i=1}^r \sum_{i < j} \alpha_k^{ij} (A_i dA_j - A_j dA_i) \\ \omega_2^m &:= dC_m + \frac{1}{2} \sum_{i=1}^r \sum_{j=1}^s B_j \beta_m^{ij} dA_i - \frac{1}{12} \sum_{l=1}^r \sum_{k=1}^s \sum_{i < j} A_l \alpha_k^{ij} (A_i dA_j - A_j dA_i) \beta_m^{lk}\end{aligned}$$

We have

$$H(\mathbb{R}^{r+s+t}, \star) = \bigcap_{k=1}^s \ker \omega_1^k \cap \bigcap_{m=1}^t \ker \omega_2^m$$

so that a tangent vector v lies in $H_p(\mathbb{R}^{r+s+t}, \star)$ if and only if $(\omega_1^k)_p(v) = (\omega_2^m)_p(v) = 0$ for all k and m .

Every step n Carnot group has the relation that if $K \subset \mathbb{G}$ is compact, then there is a constant $C = C(K) \geq 1$ such that

$$C^{-1} |p - q| \leq d_{cc}(p, q) \leq C |p - q|^{1/n}\tag{6.3}$$

for all $p, q \in K$.

We can define the quasi-metric

$$d_K(p, q) = \sum_{i=1}^n \sum_{j=1}^{\dim(V_i)} |\pi_i^j(q^{-1}p)|^{\frac{1}{i}} \quad (6.4)$$

where π_i^j is the projection to the i -th layer and the j -th coordinate.

$$\begin{aligned} \pi_1^i(q^{-1}p) &= a_i - A_i \\ \pi_2^k(q^{-1}p) &:= \varphi_1^k(p, q) = b_k - B_k - \frac{1}{2} \sum_{i < j} \alpha_k^{ij} (A_i a_j - a_i A_j) \\ \pi_3^m(q^{-1}p) &:= \varphi_2^m(p, q) = c_m - C_m - \frac{1}{2} \sum_{i, j} \beta_m^{ij} (A_i b_j - B_j a_i) + \frac{1}{12} \sum_{l, k} \sum_{i < j} (A_l + a_l) \alpha_k^{ij} (A_i a_j - a_i A_j) \beta_m^{lk} \end{aligned}$$

It is worth to mention that d_K is comparable with C-C metric on Carnot group.

It is known that for Pansu differentiable map f the Pansu differential is a Lie algebra homomorphism. Thus, if the vector field X and contact form ω are not from the same layer, then $X(f_P^* \omega) = 0$. For instance, on Heisenberg group \mathbb{H}^1

$$\begin{aligned} D_P f &= \begin{bmatrix} X_1(f_P^* dx_1) & X_1(f_P^* dx_2) & X_1(f_P^* \omega_1) \\ X_2(f_P^* dx_1) & X_2(f_P^* dx_2) & X_2(f_P^* \omega_1) \\ Z(f_P^* dx_1) & Z(f_P^* dx_2) & Z(f_P^* \omega_1) \end{bmatrix} \\ &= \begin{bmatrix} X_1(f_P^* dx_1) & X_1(f_P^* dx_2) & 0 \\ X_2(f_P^* dx_1) & X_2(f_P^* dx_2) & 0 \\ 0 & 0 & Z(f_P^* \omega_1) \end{bmatrix} \end{aligned}$$

Although the Pansu differential of Hölder continuous maps do not exist, we can generalize the Pansu differential with the help of mollification.

Theorem 6.3. Suppose that $f \in C^{0, \beta}(\Omega; G_2)$, where G_2 is a step n_2 Carnot group, $\Omega \subset G_1$ is open, X is a left-invariant vector field belonging to the horizontal layer and $0 < \beta \leq 1$. If $B(x_o, 2r) \subset \Omega$, then for all $0 < \varepsilon < r$,

$$\begin{aligned} X_1(f_\varepsilon^* \omega_1^k) &\lesssim \varepsilon^{2\beta-1} \\ X_1(f_\varepsilon^* \omega_2^m) &\lesssim \varepsilon^{2\beta-1} \\ &\dots \\ X_1(f_\varepsilon^* \omega_{n_2}^m) &\lesssim \varepsilon^{2\beta-1} \end{aligned} \quad (6.5)$$

where $f_\varepsilon = f * \eta_\varepsilon = \int_{\mathbb{G}_1} f(y) \eta_\varepsilon(y^{-1}x) dy$ and η_ε is the heat kernel on Carnot group G_1 .

Proof. Recall that $\varphi(p, q)$ was defined in (6.4). Let $p \in B(x_o, r)$ and let $\varepsilon \in (0, r)$. In what follows we will identify $(f_\varepsilon^* \alpha)(p)$ with the vector (of equal length):

$$\begin{aligned} X(f_\varepsilon^* \omega_1^k)(p) &= X(f_\varepsilon^{B_k})(p) + \frac{1}{2} \alpha_k^{ij} \sum_{i < j} \left(f_\varepsilon^{A_j}(p) X(f_\varepsilon^{A_i})(p) - f_\varepsilon^{A_i}(p) X(f_\varepsilon^{A_j})(p) \right) = \\ &= \varepsilon^{-\nu_1-1} \int_{B_\varepsilon} \left(\left(f^{B_k}(z^{-1}p) - f_\varepsilon^{B_k}(p) + \frac{1}{2} \alpha_k^{ij} \sum_{i < j} (f_\varepsilon^{A_j}(p) f^{A_i}(z^{-1}p) - f_\varepsilon^{A_i}(p) f^{A_j}(z^{-1}p)) \right) X(\eta(\delta_{\varepsilon^{-1}}(z))) dz \right. \\ &= \varepsilon^{-\nu_1-1} \int_{B_\varepsilon} \varphi_k(f(z^{-1}p), f_\varepsilon(p)) X(\eta(\delta_{\varepsilon^{-1}}(z))) dz \\ &= \varepsilon^{-\nu_1-1} \int_{B_\varepsilon} \int_{B_\varepsilon} \varphi_k(f(z^{-1}p), f(z^{-1}w)) \eta_\varepsilon(w) X(\eta(\delta_{\varepsilon^{-1}}(z))) dw dz \end{aligned}$$

In the second equality we used the fact that $\int_{B_\varepsilon} f_\varepsilon^{t_k}(p) X(\eta(\delta_{\varepsilon^{-1}}(z))) dz = 0$. Easy verification of the last equality is left to the reader. Note that (6.4) yields

$$|\varphi_k(f(z^{-1}p), f(z^{-1}w))| \leq d_K(f(z^{-1}p), f(z^{-1}w))^2 \leq [f]_{\gamma, 2\varepsilon, B(x_o, 2r)}^2 (2\varepsilon)^{2\beta}$$

Therefore,

$$|X(f_\varepsilon^* \omega_1^k)(p)| \leq [f]_{\beta, 2\varepsilon, B(x_o, 2r)}^2 (2\varepsilon)^{2\beta} \varepsilon^{-1} \varepsilon^{-\nu_1} \int_{B_\varepsilon} |X(\eta(\delta_{\varepsilon^{-1}}(z)))| dz \lesssim \varepsilon^{2\beta-1} \quad (6.6)$$

The following method is enlightened by [8]. Similarly, to prove $|X(f_\varepsilon^* \omega_2^m)(p)| \lesssim \varepsilon^{2\beta-1}$ we would construct the equation have the form like

$$\varphi_2^m((a_i, b_k, c_m), (A_i, B_k, C_m)) = \tilde{\mathcal{C}}_m \\ = c_m - C_m - \frac{1}{2} \sum_{i,j} \beta_m^{ij} (A_i b_j - B_j a_i) + \frac{1}{12} \sum_{l,k} \sum_{i < j} (A_l + a_l) \alpha_k^{ij} (A_i a_j - a_i A_j) \beta_m^{lk}.$$

It is easy to find the following result.

$$\begin{aligned} & X(f_\varepsilon^* \omega_2^m) + X\left(f_\varepsilon^* \left(-\frac{1}{2} \sum_{i,j} \beta_m^{ij} A_i d B_j\right)\right) + X\left(f_\varepsilon^* \left(\frac{1}{12} \sum_{l,k} \sum_{i < j} A_l \alpha_k^{ij} (A_i d A_j - A_j d A_i) \beta_m^{lk}\right)\right) \\ &= \varepsilon^{-m-1} \int_{B_\varepsilon} \int_{B_\varepsilon} (f^{C_m}(z^{-1}p) - f^{C_m}(v^{-1}p)) \eta_\varepsilon(v) X(\eta(\delta_{\varepsilon^{-1}}(z))) dz dv \\ & - \frac{1}{2} \sum_{i,j} \beta_m^{ij} \varepsilon^{-m-1} \int_{B_\varepsilon} \int_{B_\varepsilon} \left(f_\varepsilon^{A_i}(v^{-1}p) f^{B_j}(z^{-1}p) - f_\varepsilon^{B_j}(v^{-1}p) f^{A_i}(z^{-1}p) \right) \\ & \quad \eta_\varepsilon(v) X(\eta(\delta_{\varepsilon^{-1}}(z))) dz dv \\ & + \frac{1}{12} \sum_{l,k} \sum_{i < j} \beta_m^{lk} \alpha_k^{ij} \int_{B_\varepsilon} \int_{B_\varepsilon} \int_{B_\varepsilon} (f^{A_l}(v^{-1}p) (f^{A_i}(v^{-1}p) f^{A_j}(z^{-1}p) - f^{A_j}(v^{-1}p) f^{A_i}(z^{-1}p)) \\ & \quad \eta_\varepsilon(w) \eta_\varepsilon(v) X(\eta(\delta_{\varepsilon^{-1}}(z))) dz dw dv \\ & + \frac{1}{12} \sum_{l,k} \sum_{i < j} \beta_m^{lk} \alpha_k^{ij} \int_{B_\varepsilon} \int_{B_\varepsilon} \int_{B_\varepsilon} (f^{A_l}(z^{-1}p) (f^{A_i}(v^{-1}p) f^{A_j}(z^{-1}p) - f^{A_j}(v^{-1}p) f^{A_i}(z^{-1}p)) \\ & \quad \eta_\varepsilon(v) \eta_\varepsilon(w) X(\eta(\delta_{\varepsilon^{-1}}(z))) dw dz dv \\ & := \mathbb{A} \end{aligned}$$

Since

$$\begin{aligned} |f_\varepsilon(p) f_\varepsilon(p) - (f(p) f(p))_\varepsilon| & \leq \int_{B_\varepsilon} \int_{B_\varepsilon} |f(z^{-1}p)| |f(v^{-1}p) - f(z^{-1}p)| \eta_\varepsilon(z) \eta_\varepsilon(v) dz dv \\ & \lesssim \int_{B_\varepsilon} \int_{B_\varepsilon} |f(z^{-1}p)| d_{cc}(f(v^{-1}p), f(z^{-1}p)) \eta_\varepsilon(z) \eta_\varepsilon(v) dz dv \\ & \leq \int_{B_\varepsilon} \int_{B_\varepsilon} \varepsilon^\beta |f(z^{-1}p)| \eta_\varepsilon(z) \eta_\varepsilon(v) dz dv \\ & \lesssim \varepsilon^\beta \end{aligned}$$

we have

$$\begin{aligned} & \left| \int_{B_\varepsilon} \int_{B_\varepsilon} \int_{B_\varepsilon} f^{A_l}(w^{-1}p) (f^{A_i}(v^{-1}p) f^{A_j}(z^{-1}p) - f^{A_j}(v^{-1}p) f^{A_i}(z^{-1}p)) \eta_\varepsilon(v) \eta_\varepsilon(w) X(\eta(\delta_{\varepsilon^{-1}}(z))) dz dw dv \right. \\ & \quad \left. - \int_{B_\varepsilon} \int_{B_\varepsilon} f^{A_l}(v^{-1}p) (f^{A_i}(v^{-1}p) f^{A_j}(z^{-1}p) - f^{A_j}(v^{-1}p) f^{A_i}(z^{-1}p)) \eta_\varepsilon(v) X(\eta(\delta_{\varepsilon^{-1}}(z))) dz dv \right| \\ & \lesssim \int_{B_\varepsilon} \int_{B_\varepsilon} \varepsilon^\beta \left| (f^{A_i}(v^{-1}p) f^{A_j}(z^{-1}p) - f^{A_j}(v^{-1}p) f^{A_i}(z^{-1}p)) \right| \eta_\varepsilon(v) X(\eta(\delta_{\varepsilon^{-1}}(z))) dz dv \end{aligned}$$

Thus, we have

$$\begin{aligned}
\mathbb{A} &= \varepsilon^{-m-1} \int_{B_\varepsilon} \int_{B_\varepsilon} \varphi_2^m(f(z^{-1}p), f(v^{-1}p)) \eta_\varepsilon(v) X(\eta(\delta_{\varepsilon^{-1}}(z))) dz dv \\
&\quad + \frac{1}{12} \sum_{l,k} \sum_{i < j} \beta_m^{lk} \alpha_k^{ij} \int_{B_\varepsilon} \int_{B_\varepsilon} \varepsilon^\beta (f^{A_i}(v^{-1}p) f^{A_j}(z^{-1}p) - f^{A_j}(v^{-1}p) f^{A_i}(z^{-1}p)) \eta_\varepsilon(v) X(\eta(\delta_{\varepsilon^{-1}}(z))) dz dv \\
&\lesssim \varepsilon^{-m-1} \int_{B_\varepsilon} \int_{B_\varepsilon} \varphi_2^m(f(z^{-1}p), f(v^{-1}p)) \eta_\varepsilon(v) X(\eta(\delta_{\varepsilon^{-1}}(z))) dz dv \\
&\quad + \frac{1}{6} \varepsilon^\beta \sum_{l,k} \beta_m^{lk} \int_{B_\varepsilon} \int_{B_\varepsilon} (|\varphi_1^k(f(z^{-1}p), f(v^{-1}p))| + |f^{B_k}(z^{-1}p) - f^{B_k}(v^{-1}p)|) \eta_\varepsilon(v) X(\eta(\delta_{\varepsilon^{-1}}(z))) dz dv \\
&\lesssim \varepsilon^{-m-1} \int_{B_\varepsilon} \int_{B_\varepsilon} \varphi_2^m(f(z^{-1}p), f(v^{-1}p)) \eta_\varepsilon(v) X(\eta(\delta_{\varepsilon^{-1}}(z))) dz dv \\
&\quad + \frac{1}{6} \varepsilon^\beta \sum_{l,k} \beta_m^{lk} \int_{B_\varepsilon} \int_{B_\varepsilon} (|\varphi_1^k(f(z^{-1}p), f(v^{-1}p))| + d_{cc}(f(z^{-1}p), f(v^{-1}p))) \eta_\varepsilon(v) X(\eta(\delta_{\varepsilon^{-1}}(z))) dz dv \\
&\lesssim \varepsilon^{-m-1} \int_{B_\varepsilon} \int_{B_\varepsilon} \varphi_2^m(f(z^{-1}p), f(v^{-1}p)) \eta_\varepsilon(v) X(\eta(\delta_{\varepsilon^{-1}}(z))) dz dv \\
&\quad + \varepsilon^{3\beta-1} + \varepsilon^{2\beta-1}
\end{aligned}$$

Since we have

$$|\varphi_2^m(f(z^{-1}p), f(v^{-1}p))| \lesssim d_{cc}(f(z^{-1}p), f(v^{-1}p))^3 \lesssim [f]_{\gamma, 2\varepsilon, B^m(x_o, 2r)}^3 (2\varepsilon)^{3\beta},$$

so

$$\begin{aligned}
&\left| X(f_\varepsilon^* \omega_2^m) + X\left(f_\varepsilon^* \left(-\frac{1}{2} \sum_{i,j} \beta_m^{ij} A_i dB_j\right)\right) + X\left(f_\varepsilon^* \left(\frac{1}{12} \sum_{l,k} \sum_{i < j} A_l \alpha_k^{ij} (A_i dA_j - A_j dA_i) \beta_m^{lk}\right)\right) \right| \\
&\leq \varepsilon^{-m-1} \int_{B_\varepsilon} \int_{B_\varepsilon} \int_{B_\varepsilon} |\varphi_2^m(f(z^{-1}p), f(v^{-1}p))| \eta_\varepsilon(v) \eta_\varepsilon(z) |X(\eta(\delta_{\varepsilon^{-1}}(z)))| dz dv + \varepsilon^{2\beta-1}
\end{aligned}$$

Taking use of $\|X(f_\varepsilon^* \omega_1^k)\|_{L^\infty(B^m(x_o, r))} \lesssim \varepsilon^{2\beta-1}$ we can get

$$\begin{aligned}
|X(f_\varepsilon^* \omega_2^m)| &\lesssim \varepsilon^{2\beta-1} + X\left(f_\varepsilon^* \left(-\frac{1}{2} \sum_{i,j} \beta_m^{ij} A_i dB_j\right)\right) + X\left(f_\varepsilon^* \left(\frac{1}{12} \sum_{l,k} \sum_{i < j} A_l \alpha_k^{ij} (A_i dA_j - A_j dA_i) \beta_m^{lk}\right)\right) \\
&\lesssim \varepsilon^{2\beta-1} + \left(\frac{5}{12} \sum_{l,k} \sum_{i < j} \beta_m^{lk} f_\varepsilon^{A(i)}(p) X f_\varepsilon^{B(j)}(p)\right) + \left(\frac{1}{12} \sum_{l,k} \sum_{i < j} \beta_m^{lk} f_\varepsilon^{A(i)}(p) \varepsilon^{2\beta-1}\right)
\end{aligned}$$

For $X(f_\varepsilon^{B(j)})(p)$,

$$X(f_\varepsilon^{B(j)}) \leq |X(f_\varepsilon^* \omega_1^k)(p)| + \left| X(f_\varepsilon^* \left(\frac{1}{2} \sum_{i < j} \alpha_k^{ij} (A_i dA_j - A_j dA_i)\right)) \right| \quad (6.7)$$

From (6.6), we know that $|X(f_\varepsilon^{B(j)})(p)| \lesssim \varepsilon^{2\beta-1}$

$$|f_\varepsilon^* \omega_2^m| \lesssim \varepsilon^{2\beta-1} + \varepsilon^{2\beta-1}$$

For higher step Carnot groups, we can similarly prove that $X_1(f_\varepsilon^* \omega_3^k) \lesssim \varepsilon^{2\beta-1}, \dots, X_1(f_\varepsilon^* \omega_{n_2}^m) \lesssim \varepsilon^{2\beta-1}$.

□

Remark 6.4. *The $\beta > \frac{1}{2}$ in theorem 6.3 is sharp. For $\Omega \hookrightarrow \mathbb{H}^n$ a domain on \mathbb{R}^m Hölder embedding into Heisenberg group, when $\beta < \frac{1}{2}$, we can construct a β -Hölder deformation with Hölder homotopic map to make the image $f(\Omega)$ to shrink to origin. We refer the reader to section 10 in [6].*

From the theorem 6.3, we know that for $f \in C^{0,\beta}(\Omega; G_2)$ and $1/2 < \beta \leq 1$ the pull-back of the left-invariant forms will decay to 0 at a rate proportional to $\varepsilon^{2\beta-1}$, when $i \geq 1$ for $X(f_\varepsilon^* \omega_i^k)$ which is one of the projections of $f_\varepsilon^* \omega_i^k$ to the horizontal layer. When considering the pansu differential for Lipschitz mappings, the pansu differential is a Lie algebra homomorphism so is a block diagonal matrix. Thus, we can get the similar result for Hölder mappings.

However, compared with the Pansu differential this result is weaker.

Theorem 6.5. *Suppose that $f \in C^{0,\beta}(\Omega; G_2)$, where G_2 is a step n_2 Carnot group, $\Omega \subset G_1$ is open, X_i is a left-invariant vector field belonging to the i -th layer and $0 < \beta \leq 1$. If $B(x_0, 2r) \subset \Omega$, then for all $0 < \varepsilon < r$ and the weight of ω_j is larger than the weight of X_i ,*

$$\begin{aligned} X_i(f_\varepsilon^* \omega_1^k) &\lesssim \varepsilon^{(i+1)\beta-i} \\ X_i(f_\varepsilon^* \omega_2^m) &\lesssim \varepsilon^{(i+1)\beta-i} \\ &\dots \\ X_i(f_\varepsilon^* \omega_{n_2}^m) &\lesssim \varepsilon^{(i+1)\beta-i} \end{aligned} \tag{6.8}$$

where $f_\varepsilon = f * \eta_\varepsilon = \int_{\mathbb{G}_\varepsilon} f(y) \eta_\varepsilon(y^{-1}x) dy$ and η_ε is the heat kernel on Carnot group G_1 .

Proof. For a vector field X_2 belongs to the second layer and X_1, X'_1 belong to the horizontal, for $j \geq 2$,

$$\begin{aligned} X_2(f_\varepsilon^* \omega_j^k) &= (f_\varepsilon^* \omega_j^k)[X_1, X'_1] \\ &= (d_0 f_\varepsilon^* \omega_j^k)(X_1, X'_1) \\ &= (f_\varepsilon^* d_0 \omega_j^k)(X_1, X'_1) \end{aligned}$$

By the definition of d_0 , the weight of $d_0 \omega_j^k$ is the same as ω_j^k , so it can be represented as the of a form on the horizontal layer and a contact form on the j -th layer under wedge product

$$d_0 \omega_j^k = \sum_i a_i dx_i \wedge \omega_{j-1}^i.$$

Thus,

$$\begin{aligned} X_2(f_\varepsilon^* \omega_j^k) &= (f_\varepsilon^* d_0 \omega_j^k)(X_1, X'_1) \\ &= \sum_i X_1 f_\varepsilon^*(a_i dx_i) X'_1 f_\varepsilon^*(\omega_{j-1}^i) \end{aligned}$$

By calculation,

$$\begin{aligned} X_1 f_\varepsilon &= \varepsilon^{Q-1} \int_{B_\varepsilon} (f(z^{-1}x) - f(x)) X_1(\eta(\delta_{\varepsilon^{-1}}(z))) dz \\ &\leq \varepsilon^{Q-1} \int_{B_\varepsilon} |f(z^{-1}x) - f(x)| |X_1(\eta(\delta_{\varepsilon^{-1}}(z)))| dz \\ &\leq \varepsilon^{Q-1} \int_{B_\varepsilon} d_{cc}(f(z^{-1}x), f(x)) |X_1(\eta(\delta_{\varepsilon^{-1}}(z)))| dz \\ &\lesssim \varepsilon^{\beta-1} \end{aligned}$$

and

$$\begin{aligned} X_2(f_\varepsilon^* \omega_j^k) &= \sum_i X_1 f_\varepsilon^*(a_i dx_i) X'_1 f_\varepsilon^*(\omega_{j-1}^i) \\ &\lesssim \varepsilon^{\beta-1} \cdot \varepsilon^{2\beta-1} \\ &= \varepsilon^{3\beta-2} \end{aligned}$$

For a vector field X_3 belongs to the third layer, for $j \geq 3$,

$$\begin{aligned} X_3(f_\varepsilon^* \omega_j^k) &= (f_\varepsilon^* \omega_j^k)[X_1, X_2] \\ &= (d_0 f_\varepsilon^* \omega_j^k)(X_1, X_2) \\ &= (f_\varepsilon^* d_0 \omega_j^k)(X_1, X_2) \end{aligned}$$

Because of

$$d_0 \omega_j^k = \sum_i a_i dx_i \wedge \omega_{j-1}^i.$$

Thus,

$$\begin{aligned} X_3(f_\varepsilon^* \omega_j^k) &= (f_\varepsilon^* d_0 \omega_j^k)(X_1, X_2) \\ &= \sum_i X_1 f_\varepsilon^*(a_i dx_i) X_2 f_\varepsilon^*(\omega_{j-1}^i) \\ &\lesssim \varepsilon^{\beta-1} \cdot \varepsilon^{3\beta-2} \\ &= \varepsilon^{4\beta-3} \end{aligned}$$

By induction, we have (6.8). \square

Therefore, as long as for β -Hölder continuous maps the parameter β is large enough, then we can make sure the general differential still exhibit properties similar to Lie algebra homomorphism.

The result above can be used to prove the property of Hölder contact lift.

Theorem 6.6. *Let $V_1 \rightarrow G_1 \rightarrow H_1$ and $V_2 \rightarrow G_2 \rightarrow H_2$ be central extensions of Carnot groups such that $\text{rank}(G_2) = \text{rank}(H_2)$. Let $U_1 \subset H_1$ be a domain and $F : \pi_1^{-1}(U_1) \rightarrow G_2$ a Hölder lift of a Hölder map $f : U_1 \rightarrow H_2$. Then there exists a Lie group homomorphism $\Phi : V_1 \rightarrow V_2$ such that $F(gk) = F(g)\Phi(k)$ for all $g \in \pi_1^{-1}(U_1)$ and $k \in V_1$.*

Proof. By the lift assumption, we have

$$\pi_2 \circ F(gk) = f \circ \pi_1(gk) = f \circ \pi_1(g) = \pi_2 \circ F(g).$$

Thus, for each $g \in \pi_1^{-1}(U_1)$ and $k \in V_1$, there exists some $\tilde{\Phi}(g, k) \in V_2$ such that $F(gk) = F(g)\tilde{\Phi}(g, k)$. We claim that $\tilde{\Phi}(g, k)$ is constant in $g \in \pi_1^{-1}(U_1)$. That is, there is a well defined map $\Phi : V_1 \rightarrow V_2$ such that $\tilde{\Phi}(g, k) = \Phi(k)$ for $g \in \pi_1^{-1}(U_1)$ and $k \in V_1$. By assumption, U_1 is connected, so it suffices to prove that $g \mapsto \tilde{\Phi}(g, k)$ is locally constant for fixed $k \in V_1$.

Let B' be a ball that is compactly contained in $\pi_1^{-1}(U_1)$. Let B be a ball with the same center and half the radius.

Let $g, h \in B$. Since the extensions are central, elements of V_1 commute with elements of G_1 , and similarly for V_2 . By left-invariance of the distance, we compute that

$$\begin{aligned} d(\tilde{\Phi}(g, k), \tilde{\Phi}(h, k)) &= d(F(g)\tilde{\Phi}(g, k), F(g)\tilde{\Phi}(h, k)) \\ &\leq d(F(g)\tilde{\Phi}(g, k), F(h)\tilde{\Phi}(h, k)) \\ &\quad + d(F(h)\tilde{\Phi}(h, k), F(g)\tilde{\Phi}(h, k)) \\ &= d(F(gk), F(hk)) + d(F(h), F(g)) \\ &\leq Ld(gk, hk)^\beta + Ld(h, g)^\beta \\ &= 2Ld(h, g)^\beta \end{aligned}$$

That means $\tilde{\Phi}(g, k) \in C^{0, \beta}$. By the equal rank assumption, V_2 does not contain any horizontal directions.

Claim: When $\beta > \frac{1}{2}$, the only possible Hölder maps $B \rightarrow V_2$ are constants.

By contradiction, assume a Hölder maps $h : B \rightarrow V_2$, for $\beta > \frac{1}{2}$, then there are two points p, q such that $h(p) \neq h(q)$. We take a family of smooth approximation h_ε of h as in theorem 6.3.

take a smooth horizontal curve $\alpha : [0, 1] \rightarrow G_1$ connecting p and q , $\alpha(0) = p$ and $\alpha(1) = q$. The tangent vector of α is

$$\alpha'(t) = \sum_{i=1}^{\dim(\mathfrak{g}_1^1)} a_i(t) X_i,$$

where the X_i are horizontal basis of \mathfrak{g}_1 .

$$\begin{aligned} h_{\varepsilon*}(\alpha'(t)) &= \sum_{i=1}^{\dim(\mathfrak{g}_1^1)} a_i(t) h_{\varepsilon*}(X_i) \\ &= \sum_{i=1}^{\dim(\mathfrak{g}_1^1)} a_i(t) \sum_{j=1}^{\dim(V_2)} X_i h_\varepsilon^*(\omega'_j) X'_j, \end{aligned}$$

where the X'_j and ω'_j are the vectors and dual vectors of V_2 . Therefore,

$$|h_{\varepsilon*}(\alpha'(t))| = \sum_{i=1}^{\dim(\mathfrak{g}_1^1)} \sum_{j=1}^{\dim(V_2)} |a_i(t)| |X_i h_\varepsilon^*(\omega'_j)| |X'_j| \lesssim \varepsilon^{2\beta-1}$$

This is because theorem 6.3.

Now, under Euclidean metric d ,

$$\begin{aligned} d(h(p), h(q)) &\leq d(h(p), h_\varepsilon(p)) + d(h_\varepsilon(p), h_\varepsilon(q)) + d(h_\varepsilon(q), h(q)) \\ &\leq d(h_\varepsilon(p), h_\varepsilon(q)) + 2C(\varepsilon) \\ &= 2C(\varepsilon) + \int_0^1 |h_{\varepsilon*}(\alpha'(t))| dt \\ &\lesssim 2C(\varepsilon) + \varepsilon^{2\beta-1} \rightarrow 0, \end{aligned}$$

where the second line is because $h_\varepsilon \rightrightarrows h$, then there is a constant $C(\varepsilon)$ such that $d(h(p), h_\varepsilon(p)) \leq C(\varepsilon)$ for all $p \in G_1$. Thus $h(p) = h(q)$, a contradiction.

This claim can also be written in the following sense, as a general sense of Pansu differential. If $\beta > 1/2$, by theorem 6.3, then for any horizontal vector field X

$$\pi_{g_2[i]}(\tilde{\Phi}_P)_*(X) = 0, \quad \text{for } i \geq 2$$

where $\pi_{g_2[i]}$ are the projection to the i -th layer. Although the Pansu differential of F does not exist, the limit still has a structure similar to Lie algebra homomorphism.

Consequently, $g \mapsto \tilde{\Phi}(g, k)$ is locally constant for $k \in V_1$.

Next, we show that $\Phi : V_1 \rightarrow V_2$ is a group homomorphism. Let $k_1, k_2 \in V_1$ and let $g \in \pi_1^{-1}(U_1)$. Applying the definition of $\tilde{\Phi}$ with the pair $(g, k_1 k_2)$ shows

$$F(gk_1 k_2) = F(g)\Phi(k_1 k_2),$$

whereas applying the definition with the pairs (gk_1, k_2) and (g, k_1) yields

$$F(gk_1 k_2) = F(gk_1)\Phi(k_2) = F(g)\Phi(k_1)\Phi(k_2).$$

Canceling out $F(g)$ implies that Φ is a group homomorphism.

Finally, we observe that $\Phi(k) = F(g)^{-1}F(gk)$ for any $g \in G_1$ and $k \in V_1$. \square

Especially, we can also use a family of smooth contact lift to approximate a Hölder lift when the case is on Heisenberger group H_1 .

Theorem 6.7. *Suppose that $f \in C^{0,\alpha}(\mathbb{R}^2; \mathbb{R}^2)$ is a Hölder map, for $\alpha > 1/2$, and it admits a lift $F \in C^{0,\alpha}(H_1; H_1)$, then there are a family of smooth contact maps G_ε on H_1 and $G_\varepsilon \rightarrow F$ as $\varepsilon \rightarrow 0$.*

Proof. For central extension

$$\begin{array}{ccccc} R & \longrightarrow & H_1 & \xrightarrow{\pi} & \mathbb{R}^2 \\ & & \downarrow F & & \downarrow f \\ R & \longrightarrow & H_1 & \xrightarrow{\pi} & \mathbb{R}^2 \end{array}$$

by the assumption $\alpha > 1/2$, the $\int_{f \circ \gamma} xdy - ydx$ exists for any Lipschitz curve γ in \mathbb{R}^2 .

$$\begin{aligned} \int_{f \circ \gamma} xdy - ydx &= \int_{\gamma} f^*(xdy - ydx) \\ &= \int_{\gamma} f_x df_y - f_y df_x \end{aligned}$$

By theorem 6.6, there is a Lie group homomorphism $\Phi : V_1 \rightarrow V_2$ such that $F(gk) = F(g)\Phi(k)$, then $\Phi(\gamma_z(1)) = \lambda\gamma_z(1)$ for some constant λ , where $\gamma(0) = 0$ and

$$\begin{aligned} \gamma_z(1) &= \int_{\gamma} xdy - ydx \\ &\stackrel{Stokes}{=} \int_D dS. \end{aligned}$$

Because γ is arbitrary, f maps any simply connected domain to a simply connected domain with an area of λ times. For convenience, We can assume $\lambda = 1$, so f is area preserving map. Defining measure:

$$\mu_f(D) := m(f(D)),$$

then μ_f is absolutely continuous related to Lebesgue measure on \mathbb{R}^2 . Thus, we can define determinant $\det(J_f) = 1$ almost everywhere.

For f , we can use convolution to find a family of smooth functions f_ε and $f_\varepsilon \rightarrow f$ as $\varepsilon \rightarrow 0$. By using the same method as in the proof of theorem 6.2, we can use diffeomorphism $\Psi_\varepsilon : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ to make every $g_\varepsilon = f_\varepsilon \circ \Psi_\varepsilon$ is area preserving. Therefore, every smooth g_ε admits a smooth contact lift G_ε and $G_\varepsilon \rightarrow F$ as $\varepsilon \rightarrow 0$. \square

Based on the result above, every Hölder lift on Heisenberg group H_1 , we can use a family of smooth contact lift to approximate. Conversely, we can use a family of smooth contact lift to construct a Hölder lift on Heisenberg group H_1 .

7 The existence of Hölder lift

In this section, we connect the existence of Hölder lifts to a condition on closed horizontal curves.

We can draw a similar conclusion as Lipschitz version just for Sobolev contact lift.

At first, we are supposed to generalize the integral for Hölder continuous function. The proposition 3.13, lemma 3.15 and corollary 3.16 in [6] tell us we can

generalize the the integral for Hölder condition. And also, it is easy to check the result can be used to any $\alpha, \beta > 0$ for $f \in C^{0,\alpha}([a, b])$ and $g \in C^{0,\beta}([a, b])$. So the following lemma holds.

Lemma 7.1. *Let $\alpha, \beta > 0$, $\alpha + \beta > 1$. If $f \in C^{0,\alpha}([a, b])$ and $g \in C^{0,\beta}([a, b])$, then*

$$\left| \int_a^b f(x)dg(x) \right| \leq C(\alpha, \beta, b-a) \|f\|_{C^{0,\alpha}} [g]_\beta.$$

If in addition $f(p) = 0$ for some $p \in [a, b]$, then

$$\left| \int_a^b f(x)dg(x) \right| \leq C(\alpha, \beta) |b-a|^{\alpha+\beta} [f]_\alpha [g]_\beta.$$

Let $0 < \alpha' \leq \alpha$, $0 < \beta' \leq \beta$ be such that $\alpha' + \beta' > 1$ and $\text{Lip}([a, b]) \ni f^\varepsilon \rightarrow f$ in $C_b^{0,\alpha'}([a, b])$ and $\text{Lip}([a, b]) \ni g^\varepsilon \rightarrow g$ in $C_b^{0,\beta'}([a, b])$, we define

$$\int_a^b f dg := \lim_{\varepsilon \rightarrow 0} \int_a^b f^\varepsilon dg^\varepsilon = \lim_{\varepsilon \rightarrow 0} \int_a^b f^\varepsilon(x) (g^\varepsilon)'(x) dx$$

It exists and it does not depend on the choice of the approximation.

Now, we can draw a similar conclusion for the existence of Hölder lift.

Theorem 7.2. *Let $U_1 \subset \mathbb{H}$ and $\tilde{U}_1 \subset \pi_1^{-1}(U_1) \subset \mathbb{G}_1$ be domains and $\text{rank}(\mathbb{G}_2) = \text{rank}(\mathbb{H}_2)$. For $\beta > \frac{1}{2}$, A Hölder map $f \in C^{0,\beta}$, $f : U_1 \rightarrow \mathbb{H}_2$ admits a Hölder lift $F \in C^{0,\beta}$, $F : \tilde{U}_1 \rightarrow \mathbb{G}_2$ if and only if*

$$\int_{f \circ \pi_1 \circ \tilde{\gamma}} \alpha = 0 \tag{7.1}$$

for all Lipschitz closed curves $\tilde{\gamma}$ on \mathbb{G}_1 . We define $\int_{\tilde{\gamma}} (f \circ \pi_1)^* \alpha := \lim_{\varepsilon \rightarrow 0} \int_{\tilde{\gamma}} (f_\varepsilon \circ \pi_1)^* \alpha$, where $f_\varepsilon = f * \eta_\varepsilon = \int_{\mathbb{G}_1} f(y) \eta_\varepsilon(y^{-1}x) dy$ and η_ε is the heat kernel on Carnot group \mathbb{G}_1 .

Proof. Taking any Lipschitz curve $\tilde{\gamma}$, the first step is to prove the limit of the integration exists. Because (6.3), all coordinate components of f are β -Hölder continuous functions. Since α does not extent the horizontal layer and $\beta > \frac{1}{2}$ and theorem 6.5, we know that the β -Hölder map F can still preserve the graded structure of Lie algebra. From [6] proposition 3.13, the limit of $\int_{\tilde{\gamma}} (f_\varepsilon \circ \pi_1)^* \alpha$ exists when $\varepsilon \rightarrow 0$. Thus, we define $\int_{\tilde{\gamma}} (f \circ \pi_1)^* \alpha := \lim_{\varepsilon \rightarrow 0} \int_{\tilde{\gamma}} (f_\varepsilon \circ \pi_1)^* \alpha$.

A case in point is Heisenberg group \mathbb{H}^n , where the $\alpha = dt - \sum_{i=1}^n x_i dy_i - y_i dx_i$ and $(f_\varepsilon \circ \pi_1)^* \alpha = \sum_{i=1}^n f_i^x df_i^y - f_i^y df_i^x$.

For step-3 Carnot group, we know that the space of β -Hölder continuous function is a Banach algebra and every coordinate component is β -Hölder continuous function, since

$$\begin{aligned} |f^{B_i}(x) - f^{B_i}(y)| &\lesssim |f(x) - f(y)| \\ &\leq C d_{cc}(f(x), f(y)) \\ &\lesssim d_{cc}(x, y)^\beta. \end{aligned}$$

where the constant $C = C(K)$ is from (6.3), for any $f(x), f(y) \neq \infty$ there is a compact set K such that $f(x), f(y) \in K$. Because $\text{rank}(\mathbb{G}_2) = \text{rank}(H)_2$, $X F_\varepsilon^*(\omega_2^m) \rightarrow 0$ for

$$\omega_2^m = dC_m + \frac{1}{2} \sum_{i=1}^r \sum_{j=1}^s B_j \beta_m^{ij} dA_i - \frac{1}{12} \sum_{l=1}^r \sum_{k=1}^s \sum_{i < j} A_l \alpha_k^{ij} (A_i dA_j - A_j dA_i) \beta_m^{lk}.$$

and horizontal vector field X . Therefore,

$$\begin{aligned} & \frac{1}{2} \sum_{i=1}^r \sum_{j=1}^s \beta_m^{ij} \int_a^b \tilde{\gamma}_j^B d\tilde{\gamma}_i^A - \frac{1}{12} \sum_{l=1}^r \sum_{k=1}^s \sum_{i < j} \alpha_k^{ij} \beta_m^{lk} \int_a^b \tilde{\gamma}_l^A (\tilde{\gamma}_i^A d\tilde{\gamma}_j^A - \tilde{\gamma}_j^A d\tilde{\gamma}_i^A) \\ &= \tilde{\gamma}^{C_m}(a) - \tilde{\gamma}^{C_m}(b) + \int_{\tilde{\gamma}} F^*(\omega_2^m) \\ &= \tilde{\gamma}^{C_m}(a) - \tilde{\gamma}^{C_m}(b) \end{aligned},$$

where the $\tilde{\gamma} = F \circ \gamma \in C^{0,\gamma}([0, 1]; \mathbb{G}_2)$ for Lipschitz curve γ .

For higher step situation, the way is the same.

Due to F is contact lift $\pi_2 \circ F \circ \tilde{\gamma} = f \circ \pi_1 \circ \tilde{\gamma}$, we have $\int_{\tilde{\gamma}} (f \circ \pi_1)^* \alpha_2 = 0$ when $\tilde{\gamma}$ is a closed Lipschitz curve.

For the converse direction, because $\int_{f \circ \pi_1 \circ \tilde{\gamma}} \alpha = 0$, we can define the lift F in the following sense.

If $\gamma : [0, 1] \rightarrow H_1$ is a horizontal curve starting from $\gamma(0) = e_{H_1}$, then

$$\tilde{\gamma}(1) = \gamma(1) \cdot \exp\left(\int_{\gamma} \alpha\right)$$

For the contact lift F of f , then

$$F(\tilde{\gamma}(1)) = f(\gamma(1)) \cdot \exp\left(\int_{\gamma} (f \circ \pi_1)^* \alpha_2\right)$$

Obviously, this F is well defined, since the value of F does not depend on the choice of curve γ .

The next step is to prove F is a γ -Hölder map. Let's take a Lipschitz curve $\tilde{\gamma}$ connecting e and $\tilde{\gamma}(1)$ and a Lipschitz curve $\hat{\gamma}$ connecting $\tilde{\gamma}(1) = \hat{\gamma}(0)$ and $\hat{\gamma}(\varepsilon)$, where $\hat{\gamma}$ is under arc length parameter.

$$\begin{aligned} & d_{CC}(F(\hat{\gamma}(0)), F(\hat{\gamma}(\varepsilon))) \\ & \leq d_{CC}(f(\pi_1(\hat{\gamma}(0))), f(\pi_1(\hat{\gamma}(\varepsilon)))) \\ & + d_{CC}(e, \exp\left(\int_{\hat{\gamma}} (f \circ \pi_1)^* \alpha_2\right)) \\ & \lesssim \varepsilon^\beta + d_{CC}(e, \exp\left(\int_{\hat{\gamma}} (f \circ \pi_1)^* \alpha_2\right)) \end{aligned}$$

Because

$$\int_{\hat{\gamma}} (f \circ \pi_1)^* \alpha_2 = \int_{(f \circ \pi_1 \circ \hat{\gamma}(0))^{-1} f \circ \pi_1 \circ \hat{\gamma}} (r_{f \circ \pi_1 \circ \hat{\gamma}(0)})^* \alpha_2$$

where the r is the right translation and $(f \circ \pi_1 \circ \hat{\gamma}(0))^{-1} f \circ \pi_1 \circ \hat{\gamma}$ is also a γ -Hölder curve, for convenience, we can assume $f \circ \pi_1 \circ \hat{\gamma}(0) = e$.

We take step 3 Carnot group as an example.

From (6.4),

$$\begin{aligned} |(f \circ \pi_1 \circ \hat{\gamma})^{B_i}(\varepsilon) - (f \circ \pi_1 \circ \hat{\gamma})^{B_i}(0)|^{\frac{1}{2}} & \leq d_K((f \circ \pi_1 \circ \hat{\gamma})(\varepsilon), (f \circ \pi_1 \circ \hat{\gamma})(0)) \\ & \approx d_{CC}(e, f \circ \pi_1 \circ \hat{\gamma}(\varepsilon)) \\ & \lesssim \varepsilon^\beta \end{aligned}$$

thus the coordinate component $(f \circ \pi_1 \circ \hat{\gamma})^{B_i}$ is a $\varepsilon^{2\beta}$ -Hölder function.

By the result form lemma 5.8 in [6] yields

$$\int_{\tilde{\gamma}} (f \circ \pi_1)^* \alpha \lesssim \varepsilon^{3\beta} [f \circ \pi_1 \circ \tilde{\gamma}]_{\gamma}^2$$

Thus,

$$\begin{aligned}
& d_{CC}(F(\hat{\gamma}(0)), F(\hat{\gamma}(\varepsilon))) \\
& \lesssim \varepsilon^\beta + d_{CC}(e, \exp(\int_{\hat{\gamma}} (f \circ \pi_1)^* \alpha_2)) \\
& \approx \varepsilon^\beta + \left| \int_{\hat{\gamma}} (f \circ \pi_1)^* \alpha \right|^{\frac{1}{3}} \\
& \lesssim \varepsilon^\beta + \varepsilon^\beta \\
& = 2\varepsilon^\beta
\end{aligned}$$

Therefore, the lift F is a β -Hölder map. For the higher step Carnot group, the way is the same. \square

In [7], theorem 1.1 and proposition 1.4 give decent results about the existence of contact lifts for differentiable maps. However, for Hölder map, the result about the existence of lift can just rely on the metric structure. Compared with [7], in this section we just consider a special case of central extension.

Theorem 7.3. *Let $V_1 \rightarrow G_1 \rightarrow H$ and $V_2 \rightarrow G_2 \rightarrow H$ be central extensions, where $\mathfrak{h} = \mathfrak{h}^{[1]} \oplus \dots \oplus \mathfrak{h}^{[n]}$ and $V_1 = V_1^{[n+1]}$ and $V_2 = V_2^{[n+1]}$. Let U is a simply connected domain in H and $f : U \rightarrow H$ be a γ -Hölder map for $\beta > \frac{n}{n+1}$. Then f admits a β -Hölder lift F if and only if there is a linear map $\varphi : V_1 \rightarrow V_2$ and a V_2 -value 1-form λ with weight ≥ 2 such that the integral of $f^* \alpha_2$ on any horizontal curve equals to the integral of $\varphi \circ \alpha_1$ on the same curve, i.e.*

$$\varphi \circ \int_{\gamma} \alpha_1 = \int_{\gamma} f^* \alpha_2.$$

such that for any horizontal curve.

$$\begin{array}{ccccc}
V_1 & \longrightarrow & G_1 & \xrightarrow{\pi_1} & H \\
\varphi \downarrow & & \downarrow F & & \downarrow f \\
V_2 & \longrightarrow & G_2 & \xrightarrow{\pi_2} & H
\end{array}$$

Proof. If f admits a β -Hölder lift F , thanks to theorem 6.6, there is a linear map $\varphi : V_1 \rightarrow V_2$. Because of $\beta > \frac{n}{n+1}$ and theorem 6.5, we know that the local β -Hölder map F can still preserve the graded structure of Lie algebra.

$$\begin{aligned}
F(hk) &= F(h)\Phi(k) \\
&= \pi_{V_2} \circ F(h) \cdot \pi_2 \circ F(h) \cdot \Phi(k) \\
&= \pi_2 \circ F(h) \cdot \Phi(k) \\
&= f \circ \pi_1(h) \cdot \Phi(k) \\
&= f(h)\Phi(k),
\end{aligned}$$

for $h \in H$ and $k \in V_1$. The second equation is because $\pi_{V_2} \circ F|_H : H \rightarrow V_2$ is a local β -Hölder map and the proof is the same as the claim in theorem 6.6. Thus,

$$\varphi \circ \int_{\gamma} \alpha_1 = \Phi(k) = \int_{\gamma} (f \circ \pi_1)^* \alpha_2,$$

where we take γ as the geodesic from e to k . From theorem 6.9, we know here the γ can be replaced by any Lipschitz curve.

For convenience, we can denote as the following form. There is a V_2 -value 1-form λ with weight ≥ 2 such that

$$\varphi \circ \alpha_1 = (f \circ \pi_1)^* \alpha_2 + \lambda.$$

The $(f \circ \pi_1)^* \alpha_2$ can be decomposed into two part: $f^* \alpha_2 : V_1 \rightarrow V_2$ and $\mu_h : H \rightarrow V_2$. For the $\varphi \circ \alpha_1$, it turns α_1 with weight $= n + 1$ to a form with weight $= n + 1$. For $f^* \alpha_2$, it turns α_2 with weight $= n + 1$ to a form with weight $= n + 1$. However, the $\mu_h = 0$, since it turns a 1-form with weight $< n + 1$ to a form with weight $= n + 1$.

Therefore,

$$\begin{aligned} \varphi \circ \alpha_1 &= f^* \alpha_2 + \mu_h + \lambda \\ &= f^* \alpha_2 + \lambda. \end{aligned}$$

and

$$\varphi \circ \int_{\gamma} \alpha_1 = \int_{\gamma} f^* \alpha_2.$$

For the converse direction, because distributionally $\varphi \circ \alpha_1 = f^* \alpha_2 + \lambda$ for any horizontal curve, locally for any fixed point p and its closed cubic neighbor $U(p)$

$$\int_{\tilde{\gamma}} f^* \alpha_2 = \varphi \circ \int_{\tilde{\gamma}} \alpha_1$$

where $\tilde{\gamma} \subset U(p)$. That means the value of F dose not depend on the choice of path in $U(p)$. We can divide the area into cubes $\{U(p)\}$. Since the domain U is simply connected, when considering the value of $\int_{\tilde{\gamma}} f^* \alpha_2$, any given horizontal curve $\tilde{\gamma}$ is equivalent to a curve near itself. Eventually, this curve can be moved to arbitrary horizontal curve. Therefore, for any closed Lipschitz curve $\hat{\gamma}$

$$\int_{\hat{\gamma}} f^* \alpha_2 = \varphi \circ \int_{\hat{\gamma}} \alpha_1 = 0.$$

Then f admits a β -Hölder lift F , since theorem 7.2. \square

Example 7.4. For Heisenberg group H_1 , We can take $f(x, y) = (ax + g(y), \frac{1}{a^2}y) : \mathbb{R}^2 \rightarrow \mathbb{R}^2$, where g is a Weierstrass $\frac{2}{3}$ -Hölder function

$$g(y) = \sum_{n=0}^{\infty} \frac{1}{9}^n \cos(27^n \pi y)$$

It is obvious that f admits a contact lift, since

$$\begin{aligned} f^*(dx \wedge dy) &= \lim_{\varepsilon} f_{\varepsilon}^*(dx \wedge dy) \\ &= \lim_{\varepsilon} f_{\varepsilon}^*(dx) \wedge f_{\varepsilon}^*(dy) \\ &= \lim_{\varepsilon} \left(\frac{\partial}{\partial x} f_{\varepsilon}^x dx + \frac{\partial}{\partial y} f_{\varepsilon}^x dy \right) \wedge \frac{1}{a^2} dy \\ &= \lim_{\varepsilon} \left(\frac{\partial}{\partial x} f^x * \eta_{\varepsilon} dx + \frac{\partial}{\partial y} f^x * \eta_{\varepsilon} dy \right) \wedge \frac{1}{a^2} dy \\ &= \lim_{\varepsilon} \frac{1}{a^2} \frac{\partial}{\partial x} f^x * \eta_{\varepsilon} dx \wedge dy \\ &= \lim_{\varepsilon} \frac{1}{a} dx \wedge dy \\ &= \frac{1}{a} dx \wedge dy. \end{aligned}$$

Therefore, with the help of theorem 7.3, we can find a $\frac{2}{3}$ -Hölder lift $F : H_1 \rightarrow H_1$.

The central extension is defined by $d_0\alpha = \rho = dx \wedge dy$ as

$$\begin{array}{ccc} R & \longrightarrow & H_1 \xrightarrow{\pi} \mathbb{R}^2 \\ & & \downarrow F \\ R & \longrightarrow & H_1 \xrightarrow{\pi} \mathbb{R}^2 \end{array}$$

If $\gamma : [0, 1] \rightarrow \mathbb{R}^2$ is a Lipschitz curve starting from $\gamma(0) = e$, then

$$\tilde{\gamma}(1) = \gamma(1) \cdot \exp\left(\int_{\gamma} \alpha\right)$$

and

$$F(\tilde{\gamma}(1)) = f(\gamma(1)) \cdot \exp\left(\int_{\gamma} (f \circ \pi)^* \alpha\right)$$

References

2019Agrachev

[1] Andrei Agrachev, Davide Barilari, and Ugo Boscain. *Endpoint Map and Exponential Map*, pages 246–294. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 2019.

Allcock1998

[2] D. Allcock. An isoperimetric inequality for the Heisenberg groups. *Geom. Funct. Anal.*, 8(2):219–233, 1998.

24aef8b54a84e703370bf

[3] {Zoltán M.} Balogh, Regula Hoefer-Isenegger, and {Jeremy T.} Tyson. Lifts of lipschitz maps and horizontal fractals in the heisenberg group. *Ergodic Theory and Dynamical Systems*, 26(3):621–651, June 2006.

CapognaTang1995

[4] Luca Capogna and Puqi Tang. Uniform domains and quasiconformal mappings on the Heisenberg group. *Manuscripta Math.*, 86(3):267–281, 1995.

FRANCHI2007152

[5] Bruno Franchi, Raul Serapioni, and Francesco Serra Cassano. Regular submanifolds, graphs and area formula in heisenberg groups. *Advances in Mathematics*, 211(1):152–203, 2007.

mappingsdifferential

[6] Piotr Hajłasz, Jacob Mirra, and Armin Schikorra. Hölder continuous mappings, differential forms and the heisenberg groups, 2025.

2025arXiv250814647H

[7] Eero Hakavuori, Susanna Heikkilä, and Toni Ikonen. Smooth contact lifts to central extensions of Carnot groups. *arXiv e-prints*, page arXiv:2508.14647, August 2025.

JUNG2017251

[8] Derek Jung. A variant of gromov’s problem on hölder equivalence of carnot groups. *Journal of Mathematical Analysis and Applications*, 456(1):251–273, 2017.

Koranyi1985

[9] A. Korányi and H. M. Reimann. Quasiconformal mappings on the heisenberg group. *Inventiones mathematicae*, 80(2):309–338, jun 1985.

book

[10] Steven Krantz. *Harmonic and Complex Analysis in Several Variables*. 01 2017.

Pansu1989MtriquesDC

[11] Pierre Pansu. Métriques de carnot-carthéodory et quasiisométries des espaces symétriques de rang un. *Annals of Mathematics*, 129:1–60, 1989.