

Maximum Diminished Sombor Index of Molecular Trees with a Perfect Matching*

Fei Guo^a, Fangxia Wang^b

^aCollege of Mathematics and System Sciences, Xinjiang University
Urumqi, Xinjiang 830046, P.R.China

^b School of Mathematical Science, Xinjiang Normal University
Urumqi 830054, P.R.China

Abstract The diminished Sombor index (DSO) of a graph G , introduced by Rajathagiri, is defined as

$$DSO(G) = \sum_{uv \in E} \frac{\sqrt{d_u^2 + d_v^2}}{d_u + d_v},$$

where d_u and d_v are the degrees of vertices u and v . A graph G is a molecular graph if $d_G(u) \leq 4$ for all $u \in V(G)$. In this paper, we examine the chemical applicability of the DSO index for predicting physicochemical properties of octane isomers. We also determine the maximum value of the diminished Sombor index among all molecular trees of order n with perfect matching and characterize all the corresponding extremal trees.

Keywords: Diminished Sombor index; Molecular graph; Perfect matching

Mathematics Subject Classification: 05C05; 05C07

1 Introduction

A single number, representing a chemical structure in graph-theoretical terms via the molecular graph, is called a topological index if it correlates with a molecular property. Topological indices are used to understand physicochemical properties of chemical compounds. By a molecular graph we understand a simple graph, where the vertices represent the carbon atoms and its edges represent the carbon-carbon bonds. The topological index, also known as a molecular descriptor, is

*The work was supported by the Doctoral Research Foundation of Xinjiang Normal University (No. XJNUZBS2446), the Natural Science Foundation of Xinjiang Uygur Autonomous Region under [Grant No. 2024D01B54] and the Tianchi Talent Training Program of Xinjiang Uygur Autonomous Region.

crucial in determining this link with no experimental procedures. This index is highly valuable in QSPR analysis, which contributes to computer-assisted drug design [3, 4]. Essentially, a topological index is a graph invariant that explains a molecular graph's structural features. Since Wiener's contribution [20], numerous types of indices have emerged in the literature relying on distinct parameters including degree, distance, eccentricity, spectrum, and so on. Among various classes of topological indices, the vertex-degree-based topological indices are perhaps the most extensively studied (see [9, 10] and references therein).

All graphs considered in this paper are finite, simple and undirected. We refer to [2] for undefined notation and terminology. The vertex and edge sets of G are denoted by $V(G)$ and $E(G)$, respectively. The *order* of G (the number of its vertices) is $|V(G)| = n$ and its *size* of G (the number of its edges) is $|E(G)| = m$. For a vertex $v \in V(G)$, the degree of v , denoted by $d_G(v)$, is the number of edges incident with v in G . An edge connecting two adjacent vertices u and v is denoted as $uv \in E$. Let $e_{i,j}$ be the number of edges of i -degree vertices and j -degree vertices in G . Let $E_{i,j}$ be the set of edges of i -degree vertices and j -degree vertices in G . A vertex is said to a *pendant vertex* if its degree is one. Analogously, an edge is said to a *pendant edge* if it is incident with a pendant vertex.

For a vertex $v \in V(G)$, $G - v$ is the graph obtained from G by deleting v and its incident edges. If $S \subseteq E(G)$, we use $G - S$ to denote the graph formed from G by removing the edges in S . Similarly, $G + S$ denotes the graph obtained from G by adding the edges in S . In particular, if $S = \{uv\}$, then $G - S$ and $G + S$ are simply denoted by $G - uv$ and $G + uv$, respectively. If M is a *matching*, the two ends of each edge of M are said to be matched under M , and each vertex incident with an edge of M is said to be covered by M . A *perfect matching* is one which covers every vertex of the graph.

Recall that an *acyclic graph* is one that contains no cycles. A connected acyclic graph is called a *tree*. A *molecular graph* is a graph with $d_G(u) \leq 4$ for all $u \in V(G)$. H. Y. Deng, Z. K. Tang and R. F. Wu [7] gave the sharp upper bound for the reduced Sombor index among all molecular trees of given order n . F. X. Wang and B. Wu [22] gave the maximum value of the reduced Sombor index among all molecular trees of order n with perfect matching and show that the maximum molecular trees of exponential reduced Sombor index. Du and Su [6] showed extremal results on bond incident degree indices of molecular trees with a fixed order and a fixed number of leaves. People may refer to [1, 5, 21, 23] for more relevant works.

Topological indices depending on end-vertex degrees of edges are called *vertex-degree-based topological indices* (VDB *topological indices* for short) [11]. During many years, scientists have been trying to improve the predictive power of the VDB topological index. The Sombor index and its variants belong to the class of vertex-degree-based (VDB) topological indices. The Sombor index, a important

index in chemical theory, is proposed in [8] and is defined as

$$SO(G) = \sum_{uv \in E} \sqrt{d_u^2 + d_v^2}.$$

Redžepović [17] studied chemical applicability of Sombor index. Specifically, the Sombor index was used to model entropy and enthalpy of vaporization of alkanes with satisfactory prediction potential, indicating that this topological index may be used successfully on modeling thermodynamic properties of compounds. The Sombor index has proven useful in QSPR and QSAR studies, contributing to its growing popularity in the literature [8, 12, 13]. Recently, Rajathagiri introduced the Sombor index's variants in [18]. F. Movahedi, I. Gutman, I. Redžepović and B. Furtula called the Sombor index's variants as Diminished Sombor index in [16]. This index is defined as follows:

$$DSO(G) = \sum_{uv \in E} \frac{\sqrt{d_u^2 + d_v^2}}{d_u + d_v}.$$

F. Movahedi, I. Gutman, I. Redžepović and B. Furtula [16] derived bounds for the *DSO* index and characterized the extremal graphs within the classes of trees, unicyclic graphs, and bicyclic graphs. They also presented numerical analyses regarding the structure-dependence of the *DSO* index and its potential applications in chemistry. In [14], the tricyclic graph of a specified order that attains the maximum *DSO* is identified, and its distinctive structural characteristics are examined. In [14], the relationships and inequalities between *DSO* and some classical topological indices are analyzed thoroughly, with characterizations of extremal graphs achieving equality conditions. Motivated by known results, in this paper, we will characterize molecular trees with a perfect matching attaining the maximum *DSO* index.

2 Chemical applicability

In order to evaluate the chemical relevance of the *DSO* index, we examined its statistical relationship with experimentally determined values for some physicochemical properties of octane isomers. The properties considered in this analysis are: density, melting point, boiling point, flash point, refractive index, critical volume, critical temperature, critical pressure, entropy. For each of the aforementioned properties, the Pearson correlation coefficient is computed to quantify the degree of linear association with the *DSO* index.




Figure 1. correlation between the DSO index and three properties.

In order to focus on the most statistically significant relationships, only those correlations with an absolute coefficient value $|corr| > 0.6$ were retained for further discussion. As illustrated in Figure 1, three properties satisfied this criterion: density, melting point and critical volume. These correlations suggest that the DSO index can predict some physicochemical properties of octane isomers.

3 The maximum DSO index of Molecular Trees with a Perfect Matching

In this section, we give the characterization for molecular trees which have maximum DSO index among all molecular trees of order n ($n \geq 12$) with a perfect matching. For molecular trees of $n \leq 10$ with a perfect matching, due to the order is small, it is easy to find the ones which has maximum DSO index by simple calculation, see Figure 2.




Figure 2. The corresponding molecular tree with the maximum *DSO* index of order n ($n \leq 10$).

Next, we give some useful lemmas and notations to describe properties of molecular trees which have maximum *DSO* index among all molecular trees of order n ($n \geq 12$) with a perfect matching.

Lemma 3.1. *Assume that a tree T has the maximum *DSO* index among all molecular trees of order n ($n \geq 12$) with a perfect matching. Let M be the perfect matching of T . If $e = uv \in M$, then e is a pendant edge of T .*

Proof. Suppose the result is not true. It follows that there must exist a vertex $u \in V(T)$ with $d_T(u) \geq 2$ such that for any $v \in N_T(u)$, $d_T(v) \geq 2$. We consider the following cases.

Case 1. $d_T(u) = 2$.

Let $v, u_1 \in N_T(u)$ and $uv \in M$. Let P be a maximal path which starts from u_1 and contains u_1u . Without loss of generality, suppose x is another end-point of P . Obviously, x is a pendant vertex. Let y be the neighbor of x . Since T has a perfect matching, then $d_T(y) = 2$. Let z be the another neighbor of y . Next, we distinguish the following two subcases.

Case 1.1. $z = v$.

Let $T' = T - uu_1 + yu_1$. Clearly, T' is also a molecular tree of order n with a perfect matching. In the following, we will obtain a contradiction by showing

$DSO(T') > DSO(T)$. Therefore

$$\begin{aligned}
& DSO(T') - DSO(T) \\
&= \frac{\sqrt{d_{T'}^2(u_1) + d_{T'}^2(y)}}{d_{T'}(u_1) + d_{T'}(y)} - \frac{\sqrt{d_T^2(u_1) + d_T^2(u)}}{d_T(u_1) + d_T(u)} + \frac{\sqrt{d_{T'}^2(y) + d_{T'}^2(x)}}{d_{T'}(y) + d_{T'}(x)} \\
&\quad - \frac{\sqrt{d_T^2(y) + d_T^2(x)}}{d_T(y) + d_T(x)} + \frac{\sqrt{d_{T'}^2(y) + d_{T'}^2(v)}}{d_{T'}(y) + d_{T'}(v)} - \frac{\sqrt{d_T^2(y) + d_T^2(v)}}{d_T(y) + d_T(v)} \\
&\quad + \frac{\sqrt{d_{T'}^2(u) + d_{T'}^2(v)}}{d_{T'}(u) + d_{T'}(v)} - \frac{\sqrt{d_T^2(u) + d_T^2(v)}}{d_T(u) + d_T(v)} \\
&= \frac{\sqrt{d_T^2(u_1) + 9}}{d_T(u_1) + 3} - \frac{\sqrt{d_T^2(u_1) + 4}}{d_T(u_1) + 2} + \frac{\sqrt{10}}{4} - \frac{\sqrt{5}}{3} + \frac{\sqrt{d_T^2(v) + 9}}{d_T(v) + 3} \\
&\quad - \frac{\sqrt{d_T^2(v) + 4}}{d_T(v) + 2} + \frac{\sqrt{d_T^2(v) + 1}}{d_T(v) + 1} - \frac{\sqrt{d_T^2(v) + 4}}{d_T(v) + 2}. \tag{1}
\end{aligned}$$

Since $2 \leq d_T(u_1) \leq 4$ and $2 \leq d_T(v) \leq 4$ in the last equation of (1), a simple calculation shows that the value of (1) become minimum when $d_T(u_1) = d_T(v) = 4$. Thus $DSO(T') - DSO(T) = \frac{10}{7} + \frac{\sqrt{10}}{4} - \frac{4\sqrt{5}}{3} + \frac{\sqrt{17}}{5} > 0$, contradicting the maximality of T .

Case 1.2. $z \neq v$.

Let $T' = T - uu_1 + yu_1$. Thus

$$\begin{aligned}
& DSO(T') - DSO(T) \\
&= \frac{\sqrt{d_{T'}^2(u_1) + d_{T'}^2(y)}}{d_{T'}(u_1) + d_{T'}(y)} - \frac{\sqrt{d_T^2(u_1) + d_T^2(u)}}{d_T(u_1) + d_T(u)} + \frac{\sqrt{d_{T'}^2(y) + d_{T'}^2(x)}}{d_{T'}(y) + d_{T'}(x)} \\
&\quad - \frac{\sqrt{d_T^2(y) + d_T^2(x)}}{d_T(y) + d_T(x)} + \frac{\sqrt{d_{T'}^2(y) + d_{T'}^2(z)}}{d_{T'}(y) + d_{T'}(z)} - \frac{\sqrt{d_T^2(y) + d_T^2(z)}}{d_T(y) + d_T(z)} \\
&\quad + \frac{\sqrt{d_{T'}^2(u) + d_{T'}^2(v)}}{d_{T'}(u) + d_{T'}(v)} - \frac{\sqrt{d_T^2(u) + d_T^2(v)}}{d_T(u) + d_T(v)} \\
&= \frac{\sqrt{d_T^2(u_1) + 9}}{d_T(u_1) + 3} - \frac{\sqrt{d_T^2(u_1) + 4}}{d_T(u_1) + 2} + \frac{\sqrt{10}}{4} - \frac{\sqrt{5}}{3} + \frac{\sqrt{d_T^2(z) + 9}}{d_T(z) + 3} \\
&\quad - \frac{\sqrt{d_T^2(z) + 4}}{d_T(z) + 2} + \frac{\sqrt{d_T^2(v) + 1}}{d_T(v) + 1} - \frac{\sqrt{d_T^2(v) + 4}}{d_T(v) + 2}. \tag{2}
\end{aligned}$$

Note that $2 \leq d_T(u_1) \leq 4$, $3 \leq d_T(v) \leq 4$ and $2 \leq d_T(z) \leq 4$. By checking through all possibilities, we can find the right side of (2) achieve its minimum

value at $d_T(u_1) = d_T(z) = 4$ and $d_T(v) = 2$. Thus $DSO(T') - DSO(T) = \frac{10}{7} + \frac{\sqrt{10}}{4} - \frac{2\sqrt{5}}{3} + \frac{\sqrt{2}}{2} > 0$, contradicting the maximality of T .

Case 2. $d_T(u) = 3$.

Let $v, u_1, u_2 \in N_T(u)$ and $uv \in M$. Let $d_T(v) \geq 3$. Otherwise, by Case 1, v is adjacent to a pendant vertex, a contradiction to $uv \in M$. Let P be a maximal path which starts from u and contains uv . Without loss of generality, suppose x is another end-point of P . Obviously, x is a pendant vertex. Let y be the neighbor of x . Since T has a perfect matching, then $d_T(y) = 2$. Let z be another neighbor of y . Next, we distinguish the following two subcases.

Case 2.1. $z = v$.

Let $T' = T - \{uu_1, uu_2\} + \{yu_1, yu_2\}$. Since $d_T(v) \geq 3$, we have

$$\begin{aligned}
& DSO(T') - DSO(T) \\
&= \frac{\sqrt{d_{T'}^2(u_1) + d_{T'}^2(y)}}{d_{T'}(u_1) + d_{T'}(y)} - \frac{\sqrt{d_T^2(u_1) + d_T^2(u)}}{d_T(u_1) + d_T(u)} + \frac{\sqrt{d_{T'}^2(y) + d_{T'}^2(u_2)}}{d_{T'}(y) + d_{T'}(u_2)} \\
&\quad - \frac{\sqrt{d_T^2(u_2) + d_T^2(u)}}{d_T(u_2) + d_T(u)} + \frac{\sqrt{d_{T'}^2(u) + d_{T'}^2(v)}}{d_{T'}(u) + d_{T'}(v)} - \frac{\sqrt{d_T^2(u) + d_T^2(v)}}{d_T(u) + d_T(v)} \\
&\quad + \frac{\sqrt{d_{T'}^2(y) + d_{T'}^2(x)}}{d_{T'}(y) + d_{T'}(x)} - \frac{\sqrt{d_T^2(y) + d_T^2(x)}}{d_T(y) + d_T(x)} + \frac{\sqrt{d_{T'}^2(v) + d_{T'}^2(y)}}{d_{T'}(v) + d_{T'}(y)} \\
&\quad - \frac{\sqrt{d_T^2(y) + d_T^2(v)}}{d_T(y) + d_T(v)} \\
&= \frac{\sqrt{d_T^2(u_1) + 16}}{d_T(u_1) + 4} - \frac{\sqrt{d_T^2(u_1) + 9}}{d_T(u_1) + 3} + \frac{\sqrt{17}}{5} - \frac{\sqrt{5}}{3} + \frac{\sqrt{d_T^2(v) + 16}}{d_T(v) + 4} \\
&\quad - \frac{\sqrt{d_T^2(v) + 4}}{d_T(v) + 2} + \frac{\sqrt{d_T^2(v) + 1}}{d_T(v) + 1} - \frac{\sqrt{d_T^2(v) + 9}}{d_T(v) + 3} + \frac{\sqrt{d_T^2(u_2) + 16}}{d_T(u_2) + 4} \\
&\quad - \frac{\sqrt{d_T^2(u_2) + 9}}{d_T(u_2) + 3}. \tag{3}
\end{aligned}$$

Note that $2 \leq d_T(u_1) \leq 4$, $3 \leq d_T(v) \leq 4$ and $2 \leq d_T(u_2) \leq 4$. By checking through all possibilities, we can find the right side of (3) achieve its minimum value at $d_T(u_1) = d_T(u_2) = 4$ and $d_T(v) = 2$. Thus $DSO(T') - DSO(T) = \frac{3\sqrt{2}}{2} + \frac{\sqrt{17}}{5} - \frac{10}{7} - \frac{\sqrt{5}}{3} - \frac{\sqrt{13}}{5} > 0$, contradicting the maximality of T .

Case 2.2. $z \neq v$.

Let $T' = T - \{uu_1, uu_2\} + \{yu_1, yu_2\}$. Since $d_T(v) \geq 3$, we have

$$\begin{aligned}
& DSO(T') - DSO(T) \\
&= \frac{\sqrt{d_{T'}^2(u_1) + d_{T'}^2(y)}}{d_{T'}(u_1) + d_{T'}(y)} - \frac{\sqrt{d_T^2(u_1) + d_T^2(u)}}{d_T(u_1) + d_T(u)} + \frac{\sqrt{d_{T'}^2(y) + d_{T'}^2(u_2)}}{d_{T'}(y) + d_{T'}(u_2)} \\
&\quad - \frac{\sqrt{d_T^2(u) + d_T^2(u_2)}}{d_T(u) + d_T(u_2)} + \frac{\sqrt{d_{T'}^2(u) + d_{T'}^2(v)}}{d_{T'}(u) + d_{T'}(v)} - \frac{\sqrt{d_T^2(u) + d_T^2(v)}}{d_T(u) + d_T(v)} \\
&\quad + \frac{\sqrt{d_{T'}^2(y) + d_{T'}^2(x)}}{d_{T'}(y) + d_{T'}(x)} - \frac{\sqrt{d_T^2(y) + d_T^2(x)}}{d_T(y) + d_T(x)} + \frac{\sqrt{d_{T'}^2(y) + d_{T'}^2(z)}}{d_{T'}(y) + d_{T'}(z)} \\
&\quad - \frac{\sqrt{d_T^2(y) + d_T^2(z)}}{d_T(y) + d_T(z)} \\
&= \frac{\sqrt{d_T^2(u_1) + 16}}{d_T(u_1) + 4} - \frac{\sqrt{d_T^2(u_1) + 9}}{d_T(u_1) + 3} + \frac{\sqrt{d_T^2(u_2) + 16}}{d_T(u_2) + 4} - \frac{\sqrt{d_T^2(u_2) + 9}}{d_T(u_2) + 3} \\
&\quad + \frac{\sqrt{17}}{5} - \frac{\sqrt{5}}{3} + \frac{\sqrt{d_T^2(z) + 16}}{d_T(z) + 4} - \frac{\sqrt{d_T^2(z) + 4}}{d_T(z) + 2} + \frac{\sqrt{d_T^2(v) + 1}}{d_T(v) + 1} \\
&\quad - \frac{\sqrt{d_T^2(v) + 9}}{d_T(v) + 3}. \tag{4}
\end{aligned}$$

Note that $2 \leq d_T(u_1) \leq 4$, $3 \leq d_T(v) \leq 4$, $2 \leq d_T(u_2) \leq 4$ and $2 \leq d_T(z) \leq 4$. By checking through all possibilities, we can find the right side of (4) achieve its minimum value at $d_T(u_1) = d_T(u_2) = d_T(z) = 4$ and $d_T(v) = 3$. Thus $DSO(T') - DSO(T) = \frac{\sqrt{10}+4\sqrt{2}}{4} - \frac{10}{7} + \frac{4\sqrt{17}}{5} - \frac{2\sqrt{5}}{3} > 0$, contradicting the maximality of T .

Case 3. $d_T(u) = 4$.

Let $v, u_1, u_2, u_3 \in N_T(u)$ and $uv \in M$. Let $d_T(v) = 4$. Otherwise, by Case 2, v is adjacent to a pendant vertex, a contradiction to $uv \in M$. Let P be a maximal path which starts from u and contains uv . Without loss of generality, suppose x_1 is another end-point of P . Obviously, x_1 is a pendant vertex. Let y_1 be the neighbor of x_1 . Since T has a perfect matching, we have $d_T(y_1) = 2$. Let z_1 be another neighbor of y_1 . Let P_1 be a maximal path which starts from u and contains uu_1 . Without loss of generality, suppose x_2 is another end-point of P_1 . Obviously, x_2 is a pendant vertex. Let y_2 be the neighbor of x_2 . Since T has a perfect matching, we have $d_T(y_2) = 2$. Let z_2 be another neighbor of y_2 . Next, we distinguish the following six subcases.

Case 3.1. $z_1 = v, y_2 = u_1$.

Let $T' = T - \{uu_1, uu_2, uu_3\} + \{y_1u_1, y_1u_3, u_1u_3\}$. Since $d_T(u_1) = d_T(y_2) = 2$ and $d_T(v) = 4$, we have

$$\begin{aligned}
& DSO(T') - DSO(T) \\
&= \frac{\sqrt{d_{T'}^2(u_1) + d_{T'}^2(y_1)}}{d_{T'}(u_1) + d_{T'}(y_1)} - \frac{\sqrt{d_T^2(u_1) + d_T^2(u)}}{d_T(u_1) + d_T(u)} + \frac{\sqrt{d_{T'}^2(u_2) + d_{T'}^2(u_1)}}{d_{T'}(u_2) + d_{T'}(u_1)} \\
&\quad - \frac{\sqrt{d_T^2(u) + d_T^2(u_2)}}{d_T(u) + d_T(u_2)} + \frac{\sqrt{d_{T'}^2(y_1) + d_{T'}^2(u_3)}}{d_{T'}(y_1) + d_{T'}(u_3)} - \frac{\sqrt{d_T^2(u_3) + d_T^2(u)}}{d_T(u_3) + d_T(u)} \\
&\quad + \frac{\sqrt{d_{T'}^2(u_1) + d_{T'}^2(x_2)}}{d_{T'}(u_1) + d_{T'}(x_2)} - \frac{\sqrt{d_T^2(u_1) + d_T^2(x_2)}}{d_T(u_1) + d_T(x_2)} + \frac{\sqrt{d_{T'}^2(u) + d_{T'}^2(v)}}{d_{T'}(u) + d_{T'}(v)} \\
&\quad - \frac{\sqrt{d_T^2(u) + d_T^2(v)}}{d_T(u) + d_T(v)} + \frac{\sqrt{d_{T'}^2(y_1) + d_{T'}^2(x_1)}}{d_{T'}(y_1) + d_{T'}(x_1)} - \frac{\sqrt{d_T^2(y_1) + d_T^2(x_1)}}{d_T(y_1) + d_T(x_1)} \\
&\quad + \frac{\sqrt{d_{T'}^2(v) + d_{T'}^2(y_1)}}{d_{T'}(v) + d_{T'}(y_1)} - \frac{\sqrt{d_T^2(v) + d_T^2(y_1)}}{d_T(v) + d_T(y_1)} \\
&= \frac{\sqrt{d_T^2(u_2) + 9}}{d_T(u_2) + 3} - \frac{\sqrt{d_T^2(u_2) + 16}}{d_T(u_2) + 4} + \frac{\sqrt{10}}{4} - \sqrt{5} + \frac{\sqrt{13}}{5} \\
&\quad - \frac{\sqrt{2}}{2} + \frac{2\sqrt{17}}{5}. \tag{5}
\end{aligned}$$

Since $2 \leq d_T(u_2) \leq 4$, by taking each possible value 2, 3, 4 into the last equation of (5), we find that the right side of (5) achieve its minimum at $d_T(u_2) = 2$. Thus $DSO(T') - DSO(T) = \frac{\sqrt{13}+2\sqrt{17}}{5} - \frac{5\sqrt{5}}{3} + \frac{\sqrt{10}}{4} + \frac{5}{7} > 0$, contradicting the maximality of T .

Case 3.2. $z_1 = v, z_2 = v$.

Let $T' = T - \{uu_1, uu_2, uu_3\} + \{y_1u_1, y_1u_3, y_2u_2\}$. Since $d_T(v) = 4$, we have

$$\begin{aligned}
& DSO(T') - DSO(T) \\
&= \frac{\sqrt{d_{T'}^2(y_2) + d_{T'}^2(u_2)}}{d_{T'}(y_2) + d_{T'}(u_2)} - \frac{\sqrt{d_T^2(u_2) + d_T^2(u)}}{d_T(u_2) + d_T(u)} + \frac{\sqrt{d_{T'}^2(y_1) + d_{T'}^2(u_1)}}{d_{T'}(y_1) + d_{T'}(u_1)} \\
&\quad - \frac{\sqrt{d_T^2(u) + d_T^2(u_1)}}{d_T(u) + d_T(u_1)} + \frac{\sqrt{d_{T'}^2(y_1) + d_{T'}^2(u_3)}}{d_{T'}(y_1) + d_{T'}(u_3)} - \frac{\sqrt{d_T^2(u_3) + d_T^2(u)}}{d_T(u_3) + d_T(u)} \\
&\quad + \frac{\sqrt{d_{T'}^2(y_2) + d_{T'}^2(x_2)}}{d_{T'}(y_2) + d_{T'}(x_2)} - \frac{\sqrt{d_T^2(y_2) + d_T^2(x_2)}}{d_T(y_2) + d_T(x_2)} + \frac{\sqrt{d_{T'}^2(u_1) + d_{T'}^2(y_2)}}{d_{T'}(u_1) + d_{T'}(y_2)} \\
&\quad - \frac{\sqrt{d_T^2(u_1) + d_T^2(y_2)}}{d_T(u_1) + d_T(y_2)} + \frac{\sqrt{d_{T'}^2(y_1) + d_{T'}^2(x_1)}}{d_{T'}(y_1) + d_{T'}(x_1)} - \frac{\sqrt{d_T^2(y_1) + d_T^2(x_1)}}{d_T(y_1) + d_T(x_1)} \\
&\quad + \frac{\sqrt{d_{T'}^2(v) + d_{T'}^2(y_1)}}{d_{T'}(v) + d_{T'}(y_1)} - \frac{\sqrt{d_T^2(v) + d_T^2(y_1)}}{d_T(v) + d_T(y_1)} + \frac{\sqrt{d_{T'}^2(v) + d_{T'}^2(u)}}{d_{T'}(v) + d_{T'}(u)} \\
&\quad - \frac{\sqrt{d_T^2(v) + d_T^2(u)}}{d_T(v) + d_T(u)} \\
&= \frac{\sqrt{d_T^2(u_2) + 9}}{d_T(u_2) + 3} - \frac{\sqrt{d_T^2(u_2) + 16}}{d_T(u_2) + 4} + \frac{\sqrt{d_T^2(u_1) + 9}}{d_T(u_1) + 3} - \frac{\sqrt{d_T^2(u_1) + 4}}{d_T(u_1) + 2} \\
&\quad + \frac{\sqrt{10}}{4} + \frac{2\sqrt{17}}{5} - \sqrt{5}. \tag{6}
\end{aligned}$$

Since $2 \leq d_T(u_2) \leq 4$ and $2 \leq d_T(u_1) \leq 4$ in the last equation of (6), a simple calculation shows that the value of (6) become minimum when $d_T(u_1) = 4$ and $d_T(u_2) = 2$. Thus $DSO(T') - DSO(T) = \frac{\sqrt{13}+2\sqrt{17}}{5} - \frac{5\sqrt{5}}{3} + \frac{\sqrt{10}}{4} + \frac{5}{7} > 0$, contradicting the maximality of T .

Case 3.3. $z_1 = v_2, z_2 \neq v$.

Let $T' = T - \{uv, uv_3, uv_4\} + \{y_1v, y_1v_4, y_2v_3\}$. Since $d_T(v_2) = 4$, we have

$$\begin{aligned}
& DSO(T') - DSO(T) \\
&= \frac{\sqrt{d_{T'}^2(y_2) + d_{T'}^2(u_2)}}{d_{T'}(y_2) + d_{T'}(u_2)} - \frac{\sqrt{d_T^2(u_2) + d_T^2(u)}}{d_T(u_2) + d_T(u)} + \frac{\sqrt{d_{T'}^2(y_1) + d_{T'}^2(u_1)}}{d_{T'}(y_1) + d_{T'}(u_1)} \\
&\quad - \frac{\sqrt{d_T^2(u) + d_T^2(u_1)}}{d_T(u) + d_T(u_1)} + \frac{\sqrt{d_{T'}^2(y_1) + d_{T'}^2(u_3)}}{d_{T'}(y_1) + d_{T'}(u_3)} - \frac{\sqrt{d_T^2(u_3) + d_T^2(u)}}{d_T(u_3) + d_T(u)} \\
&\quad + \frac{\sqrt{d_{T'}^2(y_2) + d_{T'}^2(x_2)}}{d_{T'}(y_2) + d_{T'}(x_2)} - \frac{\sqrt{d_T^2(y_2) + d_T^2(x_2)}}{d_T(y_2) + d_T(x_2)} + \frac{\sqrt{d_{T'}^2(z_2) + d_{T'}^2(y_2)}}{d_{T'}(z_2) + d_{T'}(y_2)} \\
&\quad - \frac{\sqrt{d_T^2(y_2) + d_T^2(z_2)}}{d_T(y_2) + d_T(z_2)} + \frac{\sqrt{d_{T'}^2(x_1) + d_{T'}^2(y_1)}}{d_{T'}(x_1) + d_{T'}(y_1)} - \frac{\sqrt{d_T^2(x_1) + d_T^2(y_1)}}{d_T(x_1) + d_T(y_1)} \\
&\quad + \frac{\sqrt{d_{T'}^2(y_1) + d_{T'}^2(v)}}{d_{T'}(y_1) + d_{T'}(v)} - \frac{\sqrt{d_T^2(y_1) + d_T^2(v)}}{d_T(y_1) + d_T(v)} + \frac{\sqrt{d_{T'}^2(u) + d_{T'}^2(v)}}{d_{T'}(u) + d_{T'}(v)} \\
&\quad - \frac{\sqrt{d_T^2(u) + d_T^2(v)}}{d_T(u) + d_T(v)} + \frac{\sqrt{d_{T'}^2(u_1) + d_{T'}^2(z_2)}}{d_{T'}(u_1) + d_{T'}(z_2)} - \frac{\sqrt{d_T^2(u_1) + d_T^2(z_2)}}{d_T(u_1) + d_T(z_2)} \\
&= \frac{\sqrt{d_T^2(u_2) + 9}}{d_T(u_2) + 3} - \frac{\sqrt{d_T^2(u_2) + 16}}{d_T(u_2) + 4} + \frac{\sqrt{d_T^2(z_2) + 9}}{d_T(z_2) + 3} - \frac{\sqrt{d_T^2(z_2) + 4}}{d_T(z_2) + 2} \\
&\quad + \frac{\sqrt{10}}{4} + \frac{2\sqrt{17}}{5} - \sqrt{5}. \tag{7}
\end{aligned}$$

Note that $2 \leq d_T(u_2) \leq 4$ and $3 \leq d_T(z_2) \leq 4$. By checking through all possibilities, we can find the right side of (7) achieve its minimum value at $d_T(u_2) = 2$ and $d_T(z_2) = 4$. Thus $DSO(T') - DSO(T) = \frac{\sqrt{13}+2\sqrt{17}}{5} - \frac{5\sqrt{5}}{3} + \frac{\sqrt{10}}{4} + \frac{5}{7} > 0$, contradicting the maximality of T .

Case 3.4. $z_1 \neq v, y_2 = u_1$.

Let $T' = T - \{uu_1, uu_2, uu_3\} + \{y_1u_1, y_1u_3, u_1u_2\}$. Since $d_T(u_1) = d_T(y_2) = 2$ and $d_T(v) = 4$, we have

$$\begin{aligned}
& DSO(T') - DSO(T) \\
&= \frac{\sqrt{d_{T'}^2(u_1) + d_{T'}^2(u_2)}}{d_{T'}(u_1) + d_{T'}(u_2)} - \frac{\sqrt{d_T^2(u_2) + d_T^2(u)}}{d_T(u_2) + d_T(u)} + \frac{\sqrt{d_{T'}^2(u_1) + d_{T'}^2(y_1)}}{d_{T'}(u_1) + d_{T'}(y_1)} \\
&\quad - \frac{\sqrt{d_T^2(u) + d_T^2(u_1)}}{d_T(u) + d_T(u_1)} + \frac{\sqrt{d_{T'}^2(u_3) + d_{T'}^2(y_1)}}{d_{T'}(y_1) + d_{T'}(u_3)} - \frac{\sqrt{d_T^2(u) + d_T^2(u_3)}}{d_T(u) + d_T(u_3)} \\
&\quad + \frac{\sqrt{d_{T'}^2(u_1) + d_{T'}^2(x_2)}}{d_{T'}(u_1) + d_{T'}(x_2)} - \frac{\sqrt{d_T^2(u_1) + d_T^2(x_2)}}{d_T(u_1) + d_T(x_2)} + \frac{\sqrt{d_{T'}^2(x_1) + d_{T'}^2(y_1)}}{d_{T'}(x_1) + d_{T'}(y_1)} \\
&\quad - \frac{\sqrt{d_T^2(x_1) + d_T^2(y_1)}}{d_T(x_1) + d_T(y_1)} + \frac{\sqrt{d_{T'}^2(y_1) + d_{T'}^2(z_1)}}{d_{T'}(y_1) + d_{T'}(z_1)} - \frac{\sqrt{d_T^2(y_1) + d_T^2(z_1)}}{d_T(y_1) + d_T(z_1)} \\
&\quad + \frac{\sqrt{d_{T'}^2(v) + d_{T'}^2(u)}}{d_{T'}(v) + d_{T'}(u)} - \frac{\sqrt{d_T^2(u) + d_T^2(v)}}{d_T(u) + d_T(v)} \\
&= \frac{\sqrt{d_T^2(u_2) + 9}}{d_T(u_2) + 3} - \frac{\sqrt{d_T^2(u_2) + 16}}{d_T(u_2) + 4} + \frac{\sqrt{d_T^2(z_1) + 16}}{d_T(z_1) + 4} - \frac{\sqrt{d_T^2(z_1) + 4}}{d_T(z_1) + 2} \\
&\quad + \frac{5}{7} - \sqrt{5} + \frac{\sqrt{10}}{4} - \frac{2\sqrt{17}}{5} - \frac{\sqrt{2}}{2}. \tag{8}
\end{aligned}$$

Note that $2 \leq d_T(u_2) \leq 4$ and $2 \leq d_T(z_1) \leq 4$. By checking through all possibilities, we can find the right side of (8) achieve its minimum value at $d_T(u_2) = 2$ and $d_T(z_1) = 4$. Thus $DSO(T') - DSO(T) = \frac{\sqrt{13}+2\sqrt{17}}{5} - \frac{5\sqrt{5}}{3} + \frac{\sqrt{10}}{4} + \frac{5}{7} > 0$, contradicting the maximality of T .

Case 3.5. $z_1 \neq v, z_2 = u_1$.

Let $T' = T - \{uu_1, uu_2, uu_3\} + \{y_1u_1, y_1u_3, y_2u_2\}$. Since $d_T(v) = 4$, we have

$$\begin{aligned}
& DSO(T') - DSO(T) \\
&= \frac{\sqrt{d_{T'}^2(u_1) + d_{T'}^2(y_1)}}{d_{T'}(u_1) + d_{T'}(y_1)} - \frac{\sqrt{d_T^2(u_1) + d_T^2(u)}}{d_T(u_1) + d_T(u)} + \frac{\sqrt{d_{T'}^2(y_2) + d_{T'}^2(u_2)}}{d_{T'}(y_2) + d_{T'}(u_2)} \\
&\quad - \frac{\sqrt{d_T^2(u) + d_T^2(u_2)}}{d_T(u) + d_T(u_2)} + \frac{\sqrt{d_{T'}^2(y_1) + d_{T'}^2(u_3)}}{d_{T'}(y_1) + d_{T'}(u_3)} - \frac{\sqrt{d_T^2(u_3) + d_T^2(u)}}{d_T(u_3) + d_T(u)} \\
&\quad + \frac{\sqrt{d_{T'}^2(y_2) + d_{T'}^2(x_2)}}{d_{T'}(y_2) + d_{T'}(x_2)} - \frac{\sqrt{d_T^2(y_2) + d_T^2(x_2)}}{d_T(y_2) + d_T(x_2)} + \frac{\sqrt{d_{T'}^2(u_1) + d_{T'}^2(y_2)}}{d_{T'}(u_1) + d_{T'}(y_2)} \\
&\quad - \frac{\sqrt{d_T^2(u_1) + d_T^2(y_2)}}{d_T(u_1) + d_T(y_2)} + \frac{\sqrt{d_{T'}^2(y_1) + d_{T'}^2(x_1)}}{d_{T'}(y_1) + d_{T'}(x_1)} - \frac{\sqrt{d_T^2(y_1) + d_T^2(x_1)}}{d_T(y_1) + d_T(x_1)} \\
&\quad + \frac{\sqrt{d_{T'}^2(y_1) + d_{T'}^2(z_1)}}{d_{T'}(y_1) + d_{T'}(z_1)} - \frac{\sqrt{d_T^2(y_1) + d_T^2(z_1)}}{d_T(y_1) + d_T(z_1)} + \frac{\sqrt{d_{T'}^2(v) + d_{T'}^2(u)}}{d_{T'}(v) + d_{T'}(u)} \\
&\quad - \frac{\sqrt{d_T^2(v) + d_T^2(u)}}{d_T(v) + d_T(u)} \\
&= \frac{\sqrt{d_T^2(u_2) + 9}}{d_T(u_2) + 3} - \frac{\sqrt{d_T^2(u_2) + 16}}{d_T(u_2) + 4} + \frac{\sqrt{d_T^2(u_1) + 9}}{d_T(u_1) + 3} + \frac{\sqrt{10}}{4} + \frac{2\sqrt{17}}{5} \\
&\quad - \frac{\sqrt{d_T^2(u_1) + 4}}{d_T(u_1) + 2} + \frac{\sqrt{d_T^2(z_1) + 16}}{d_T(z_1) + 4} - \frac{\sqrt{d_T^2(z_1) + 4}}{d_T(z_1) + 2} - \frac{2\sqrt{5}}{3} - \frac{\sqrt{2}}{2}. \tag{9}
\end{aligned}$$

Since $2 \leq d_T(u_2) \leq 4$, $2 \leq d_T(u_1) \leq 4$ and $2 \leq d_T(z_1) \leq 4$ in the last equation of (9), a simple calculation shows that the value of (9) become minimum when $d_T(u_1) = d_T(z_1) = 4$ and $d_T(u_2) = 2$. Thus $DSO(T') - DSO(T) = \frac{\sqrt{13}+2\sqrt{17}}{5} - \frac{5\sqrt{5}}{3} + \frac{\sqrt{10}}{4} + \frac{5}{7} > 0$, contradicting the maximality of T .

Case 3.6. $z_1 \neq v, z_2 \neq u_1$.

Let $T' = T - \{uu_1, uu_2, uu_3\} + \{y_1u_1, y_1u_3, y_2u_2\}$. Since $d_T(v_2) = 4$, we have

$$\begin{aligned}
& DSO(T') - DSO(T) \\
&= \frac{\sqrt{d_{T'}^2(u_2) + d_{T'}^2(y_2)}}{d_{T'}(u_2) + d_{T'}(y_2)} - \frac{\sqrt{d_T^2(u_2) + d_T^2(u)}}{d_T(u_2) + d_T(u)} + \frac{\sqrt{d_{T'}^2(y_1) + d_{T'}^2(u_1)}}{d_{T'}(y_1) + d_{T'}(u_1)} \\
&\quad - \frac{\sqrt{d_T^2(u) + d_T^2(u_1)}}{d_T(u) + d_T(u_1)} + \frac{\sqrt{d_{T'}^2(y_1) + d_{T'}^2(u_3)}}{d_{T'}(y_1) + d_{T'}(u_3)} - \frac{\sqrt{d_T^2(u_3) + d_T^2(u)}}{d_T(u_3) + d_T(u)} \\
&\quad + \frac{\sqrt{d_{T'}^2(y_2) + d_{T'}^2(x_2)}}{d_{T'}(y_2) + d_{T'}(x_2)} - \frac{\sqrt{d_T^2(y_2) + d_T^2(x_2)}}{d_T(y_2) + d_T(x_2)} + \frac{\sqrt{d_{T'}^2(z_2) + d_{T'}^2(y_2)}}{d_{T'}(z_2) + d_{T'}(y_2)} \\
&\quad - \frac{\sqrt{d_T^2(z_2) + d_T^2(y_2)}}{d_T(z_2) + d_T(y_2)} + \frac{\sqrt{d_{T'}^2(y_1) + d_{T'}^2(x_1)}}{d_{T'}(y_1) + d_{T'}(x_1)} - \frac{\sqrt{d_T^2(y_1) + d_T^2(x_1)}}{d_T(y_1) + d_T(x_1)} \\
&\quad + \frac{\sqrt{d_{T'}^2(y_1) + d_{T'}^2(z_1)}}{d_{T'}(y_1) + d_{T'}(z_1)} - \frac{\sqrt{d_T^2(y_1) + d_T^2(z_1)}}{d_T(y_1) + d_T(z_1)} + \frac{\sqrt{d_{T'}^2(v) + d_{T'}^2(u)}}{d_{T'}(v) + d_{T'}(u)} \\
&\quad - \frac{\sqrt{d_T^2(v) + d_T^2(u)}}{d_T(v) + d_T(u)} \\
&= \frac{\sqrt{d_T^2(u_2) + 9}}{d_T(u_2) + 3} - \frac{\sqrt{d_T^2(u_2) + 16}}{d_T(u_2) + 4} + \frac{\sqrt{d_T^2(z_2) + 9}}{d_T(z_2) + 3} + \frac{\sqrt{10}}{4} + \frac{2\sqrt{17}}{5} \\
&\quad - \frac{\sqrt{d_T^2(z_2) + 4}}{d_T(z_2) + 2} + \frac{\sqrt{d_T^2(z_1) + 16}}{d_T(z_1) + 4} - \frac{\sqrt{d_T^2(z_1) + 4}}{d_T(z_1) + 2} - \frac{2\sqrt{5}}{3} - \frac{\sqrt{2}}{2}. \tag{10}
\end{aligned}$$

Note that $2 \leq d_T(u_2) \leq 4$, $2 \leq d_T(z_1) \leq 4$ and $2 \leq d_T(u_2) \leq 4$. By checking through all possibilities, we can find the right side of (10) achieve its minimum value at $d_T(z_2) = d_T(z_1) = 4$ and $d_T(u_2) = 2$. Thus $DSO(T') - DSO(T) = \frac{\sqrt{13}+2\sqrt{17}}{5} - \frac{5\sqrt{5}}{3} + \frac{\sqrt{10}}{4} + \frac{5}{7} > 0$, contradicting the maximality of T .

The proof is completed. \square

Lemma 3.2. *Let T be a molecular tree which has maximum DOS index among all molecular trees of order n ($n \geq 12$) with a perfect matching M . If $d_T(u) = 3$, then $d_T(v) \in \{1, 4\}$ for any $v \in N_T(u)$.*

Proof. Suppose that $d_T(v) \in \{2, 3\}$. We consider two cases and will obtain a contradiction.

Case 1. $d_T(v) = 3$.

Let $N_T(u) \setminus \{v\} = \{u_1, u_2\}$ and $N_T(v) \setminus \{u\} = \{v_1, v_2\}$. Since T has a perfect matching, by Lemma 1 u is matched with one of its neighbor, say u_1 , v is matched with one of its neighbor, say v_1 . Thus $d_T(u_1) = 1$, $d_T(v_1) = 1$, $2 \leq d_T(u_2) \leq 4$ and $2 \leq d_T(v_2) \leq 4$. Let $T' = T - uu_2 + vu_2$. Clearly, T' is also a molecular tree of

order n with a perfect matching. In the following, we will obtain a contradiction by showing $DSO(T') > DSO(T)$.

$$\begin{aligned}
& DSO(T') - DSO(T) \\
&= \frac{\sqrt{d_{T'}^2(u_2) + d_{T'}^2(v)}}{d_{T'}(u_2) + d_{T'}(v)} - \frac{\sqrt{d_T^2(u_2) + d_T^2(u)}}{d_T(u_2) + d_T(u)} + \frac{\sqrt{d_{T'}^2(u) + d_{T'}^2(u_1)}}{d_{T'}(u) + d_{T'}(u_1)} \\
&\quad - \frac{\sqrt{d_T^2(u) + d_T^2(u_1)}}{d_T(u) + d_T(u_1)} + \frac{\sqrt{d_{T'}^2(u) + d_{T'}^2(v)}}{d_{T'}(u) + d_{T'}(v)} - \frac{\sqrt{d_T^2(u) + d_T^2(v)}}{d_T(u) + d_T(v)} \\
&\quad + \frac{\sqrt{d_{T'}^2(v) + d_{T'}^2(v_1)}}{d_{T'}(v) + d_{T'}(v_1)} - \frac{\sqrt{d_T^2(v) + d_T^2(v_1)}}{d_T(v) + d_T(v_1)} + \frac{\sqrt{d_{T'}^2(v) + d_{T'}^2(v_2)}}{d_{T'}(v) + d_{T'}(v_2)} \\
&\quad - \frac{\sqrt{d_T^2(v) + d_T^2(v_2)}}{d_T(v) + d_T(v_2)} \\
&= \frac{\sqrt{d_T^2(u_2) + 16}}{d_T(u_2) + 4} - \frac{\sqrt{d_T^2(u_2) + 9}}{d_T(u_2) + 3} + \frac{\sqrt{d_T^2(v_2) + 16}}{d_T(v_2) + 4} \\
&\quad - \frac{\sqrt{d_T^2(v_2) + 9}}{d_T(v_2) + 3} + \frac{2\sqrt{5}}{3} - \frac{\sqrt{10} + \sqrt{2}}{2} + \frac{\sqrt{17}}{5}. \tag{11}
\end{aligned}$$

Since $2 \leq d_T(u_2) \leq 4$ and $2 \leq d_T(v_2) \leq 4$ in the last equation of (11), a simple calculation shows that the value of (11) become minimum when $d_T(u_2) = d_T(v_2) = 4$. Thus $DSO(T') - DSO(T) = \frac{\sqrt{2}-\sqrt{10}}{2} - \frac{10}{7} + \frac{\sqrt{17}}{5} + \frac{2\sqrt{5}}{3} > 0$, contradicting the maximality of T .

Case 2. $d_T(v) = 2$

Let $N_T(u) \setminus \{v\} = \{u_1, u_2\}$ and $N_T(v) \setminus \{u\} = \{v_1\}$. Since T has a perfect matching, by Lemma 1 u is matched with one of its neighbor, say u_1 , v is matched with one of its neighbor, say v_1 . Thus $d_T(u_1) = 1$, $d_T(v_1) = 1$. In view of Case 1 and the assumption that $n \geq 12$, we have $d_T(u_2) = 4$. Let P be a maximal path of T which starts from u containing uu_2 . Without loss of generality, suppose x is another end-point of P . Obviously, x is a pendant vertex. Let y be the neighbor of x . Since T has a perfect matching, we have $d_T(y) = 2$. Let z be another neighbor of y . Since $n \geq 12$, we have $u_2 \neq z$.

Case 2.1. $d_T(z) = 3$.

Let $N_T(z) \setminus \{y\} = \{z_1, z_2\}$. Since T has a perfect matching, by Lemma 1 z is matched with one of its neighbor, say z_1 . Thus $d_T(z_1) = 1$. In view of Case 1 and $n \geq 12$, we can assume that $d_T(z_2) = 4$. Let $T' = T - uv + zv$. Clearly, T' is also a molecular tree of order n with a perfect matching. In the following, we will obtain a contradiction by showing $DSO(T') > DSO(T)$.

$$\begin{aligned}
& DSO(T') - DSO(T) \\
&= \frac{\sqrt{d_{T'}^2(v) + d_{T'}^2(z)}}{d_{T'}(v) + d_{T'}(z)} - \frac{\sqrt{d_T^2(v) + d_T^2(u)}}{d_T(v) + d_T(u)} + \frac{\sqrt{d_{T'}^2(u) + d_{T'}^2(u_1)}}{d_{T'}(u) + d_{T'}(u_1)} \\
&\quad - \frac{\sqrt{d_T^2(u) + d_T^2(u_1)}}{d_T(u) + d_T(u_1)} + \frac{\sqrt{d_{T'}^2(u) + d_{T'}^2(u_2)}}{d_{T'}(u) + d_{T'}(u_2)} - \frac{\sqrt{d_T^2(u) + d_T^2(u_2)}}{d_T(u) + d_T(u_2)} \\
&\quad + \frac{\sqrt{d_{T'}^2(z) + d_{T'}^2(z_1)}}{d_{T'}(z) + d_{T'}(z_1)} - \frac{\sqrt{d_T^2(z) + d_T^2(z_1)}}{d_T(z) + d_T(z_1)} + \frac{\sqrt{d_{T'}^2(z) + d_{T'}^2(y)}}{d_{T'}(z) + d_{T'}(y)} \\
&\quad - \frac{\sqrt{d_T^2(z) + d_T^2(y)}}{d_T(z) + d_T(y)} + \frac{\sqrt{d_{T'}^2(z_2) + d_{T'}^2(z)}}{d_{T'}(z_2) + d_{T'}(z)} - \frac{\sqrt{d_T^2(z_2) + d_T^2(z)}}{d_T(z_2) + d_T(z)} \\
&= \frac{4\sqrt{5}}{3} + \frac{\sqrt{17} - 2\sqrt{13}}{5} - \frac{10}{7} + \frac{\sqrt{2} - \sqrt{10}}{2} > 0.
\end{aligned} \tag{12}$$

Case 2.2 $d_T(z) = 4$.

Let $N_T(z) \setminus \{y\} = \{z_1, z_2, z_3\}$. Since T has a perfect matching, by Lemma 1 z is matched with one of its neighbor, say z_1 . Thus $d_T(z_1) = 1$ and $3 \leq d_T(z_2) \leq 4$. By the choice of P , we have $z_3 = 2$.

Case 2.2.1. $d_T(z) = 4, d_T(z_2) = 3$.

Let $N_T(z_2) \setminus \{z\} = \{z_4, z_5\}$. Since T has a perfect matching, by Lemma 1 z_2 is matched with one of its neighbor, say z_4 . Thus $d_T(z_4) = 1$. In view of Case 1, we may assume that $3 \leq d_T(z_5) \leq 4$. Let $T' = T - uv + vz_2$. Clearly, T' is a molecular tree of order n with a perfect matching. In the following, we will obtain a contradiction by showing $DSO(T') > DSO(T)$.

$$\begin{aligned}
& DSO(T') - DSO(T) \\
&= \frac{\sqrt{d_{T'}^2(z_2) + d_{T'}^2(v)}}{d_{T'}(z_2) + d_{T'}(v)} - \frac{\sqrt{d_T^2(v) + d_T^2(u)}}{d_T(v) + d_T(u)} + \frac{\sqrt{d_{T'}^2(u) + d_{T'}^2(u_1)}}{d_{T'}(u) + d_{T'}(u_1)} \\
&\quad - \frac{\sqrt{d_T^2(u) + d_T^2(u_1)}}{d_T(u) + d_T(u_1)} + \frac{\sqrt{d_{T'}^2(u) + d_{T'}^2(u_2)}}{d_{T'}(u) + d_{T'}(u_2)} - \frac{\sqrt{d_T^2(u) + d_T^2(u_2)}}{d_T(u) + d_T(u_2)} \\
&\quad + \frac{\sqrt{d_{T'}^2(z) + d_{T'}^2(z_2)}}{d_{T'}(z) + d_{T'}(z_2)} - \frac{\sqrt{d_T^2(z) + d_T^2(z_2)}}{d_T(z) + d_T(z_2)} + \frac{\sqrt{d_{T'}^2(z_2) + d_{T'}^2(z_4)}}{d_{T'}(z_2) + d_{T'}(z_4)} \\
&\quad - \frac{\sqrt{d_T^2(z_2) + d_T^2(z_4)}}{d_T(z_2) + d_T(z_4)} + \frac{\sqrt{d_{T'}^2(z_2) + d_{T'}^2(z_5)}}{d_{T'}(z_2) + d_{T'}(z_5)} - \frac{\sqrt{d_T^2(z_2) + d_T^2(z_5)}}{d_T(z_2) + d_T(z_5)} \\
&= \frac{\sqrt{d_T^2(z_5) + 16}}{d_T(z_5) + 4} - \frac{\sqrt{d_T^2(z_5) + 9}}{d_T(z_5) + 3} + \sqrt{5} - \frac{10}{7} + \frac{\sqrt{2} - \sqrt{10}}{2} \\
&\quad + \frac{\sqrt{17} - \sqrt{13}}{5}. \tag{13}
\end{aligned}$$

Since $3 \leq d_T(z_5) \leq 4$, by taking each possible value 3, 4 into the last equation of (12), we find that the right side of (12) achieve its minimum at $d_T(z_5) = 4$. Thus $DSO(T') - DSO(T) = \frac{2\sqrt{2}-\sqrt{10}}{2} - \frac{15}{7} + \sqrt{5} + \frac{\sqrt{17}-\sqrt{13}}{5} > 0$, contradicting the maximality of T .

Case 2.2.2. $d_T(z) = 4, d_T(z_2) = 4$.

Let $N_T(z_3) \setminus \{z\} = \{z_4\}$. Since T has a perfect matching, by Lemma 1 z_3 is matched with one of its neighbor, say z_4 . Thus $d_T(z_4) = 1$. Let $T' = T - \{zz_3, zy\} + \{vz_3, vy\}$. Clearly, T' is also a molecular tree of order n with a perfect matching. In the following, we will obtain a contradiction by showing $DSO(T') > DSO(T)$.

$$\begin{aligned}
& DSO(T') - DSO(T) \\
&= \frac{\sqrt{d_{T'}^2(v) + d_{T'}^2(y)}}{d_{T'}(v) + d_{T'}(y)} - \frac{\sqrt{d_T^2(z) + d_T^2(y)}}{d_T(z) + d_T(y)} + \frac{\sqrt{d_{T'}^2(v) + d_{T'}^2(z_3)}}{d_{T'}(v) + d_{T'}(z_3)} \\
&\quad - \frac{\sqrt{d_T^2(z) + d_T^2(z_3)}}{d_T(z) + d_T(z_3)} + \frac{\sqrt{d_{T'}^2(v) + d_{T'}^2(v_1)}}{d_{T'}(v) + d_{T'}(v_1)} - \frac{\sqrt{d_T^2(v) + d_T^2(v_1)}}{d_T(v) + d_T(v_1)} \\
&\quad + \frac{\sqrt{d_{T'}^2(v) + d_{T'}^2(u)}}{d_{T'}(v) + d_{T'}(u)} - \frac{\sqrt{d_T^2(v) + d_T^2(u)}}{d_T(v) + d_T(u)} + \frac{\sqrt{d_{T'}^2(z) + d_{T'}^2(z_1)}}{d_{T'}(z) + d_{T'}(z_1)} \\
&\quad - \frac{\sqrt{d_T^2(z) + d_T^2(z_1)}}{d_T(z) + d_T(z_1)} + \frac{\sqrt{d_{T'}^2(z) + d_{T'}^2(z_2)}}{d_{T'}(z) + d_{T'}(z_2)} - \frac{\sqrt{d_T^2(z) + d_T^2(z_2)}}{d_T(z) + d_T(z_2)} \\
&= \frac{\sqrt{5}}{3} - \frac{\sqrt{2}}{2} + \frac{5}{7} - \frac{\sqrt{13}}{5} > 0.
\end{aligned}$$

The proof is completed. \square

Lemma 3.3. *If a molecular tree T has the maximum DOS index among all molecular trees of order n ($n \geq 12$) with a perfect matching M , then $e_{4,4} \leq 2$ in T .*

Proof. We prove by contradiction. Suppose that $e_{4,4} \geq 3$. Let $u_1u_2, u_3u_4, u_5u_6 \in E_{4,4}$. Let P be a maximal path in T . Let x be an end of P . Obviously, x is a pendant vertex. Let y be the neighbor of x . Since T has a perfect matching, $d_T(y) = 2$. Let z be the neighbor of y other than x . Let $N_T(z) \setminus \{y\} = \{z_1, z_2, z_3\}$. Since T has a perfect matching, by Lemma 1 z is matched with one of its neighbor, say z_1 . Thus $d_T(z_1) = 1$. Since P is the maximal path, Lemma 1 and Lemma 2, we have $d_T(z_2) = 2$ and $3 \leq d_T(z_3) \leq 4$.

Case 1. $d_T(z_3) = 3$.

Let $N_T(z_3) \setminus \{z\} = \{z_4, z_5\}$. Since T has a perfect matching, by Lemma 1 z_3 is matched with one of its neighbor, say z_4 . Thus $d_T(z_4) = 1$. In view of Case 1 of Lemma 2, we may assume that $d_T(z_5) = 4$. Let $T' = T - \{u_1u_2, u_3u_4, u_5u_6, zy, zz_2, zz_3\} + \{u_1z, u_2z, u_3y, u_4y, u_5z_2, u_6z_2\}$. Therefore,

$$\begin{aligned}
& DSO(T') - DSO(T) \\
&= \frac{\sqrt{d_{T'}^2(u_1) + d_{T'}^2(z)}}{d_{T'}(u_1) + d_{T'}(z)} + \frac{\sqrt{d_{T'}^2(u_2) + d_{T'}^2(z)}}{d_{T'}(u_2) + d_{T'}(z)} + \frac{\sqrt{d_{T'}^2(u_3) + d_{T'}^2(y)}}{d_{T'}(u_3) + d_{T'}(y)} \\
&\quad + \frac{\sqrt{d_{T'}^2(y) + d_{T'}^2(u_4)}}{d_{T'}(y) + d_{T'}(u_4)} + \frac{\sqrt{d_{T'}^2(u_5) + d_{T'}^2(z_2)}}{d_{T'}(u_5) + d_{T'}(z_2)} + \frac{\sqrt{d_{T'}^2(u_6) + d_{T'}^2(z_2)}}{d_{T'}(u_6) + d_{T'}(z_2)} \\
&\quad - \frac{\sqrt{d_T^2(u_1) + d_T^2(u_2)}}{d_T(u_1) + d_T(u_2)} - \frac{\sqrt{d_T^2(u_3) + d_T^2(u_4)}}{d_T(u_3) + d_T(u_4)} - \frac{\sqrt{d_T^2(u_5) + d_T^2(u_6)}}{d_T(u_5) + d_T(u_6)} \\
&\quad - \frac{\sqrt{d_T^2(z) + d_T^2(y)}}{d_T(z) + d_T(y)} - \frac{\sqrt{d_T^2(z) + d_T^2(z_2)}}{d_T(z) + d_T(z_2)} - \frac{\sqrt{d_T^2(z) + d_T^2(z_3)}}{d_T(z) + d_T(z_3)} \\
&\quad + \frac{\sqrt{d_{T'}^2(z) + d_{T'}^2(z_1)}}{d_{T'}(z) + d_{T'}(z_1)} - \frac{\sqrt{d_T^2(z) + d_T^2(z_1)}}{d_T(z) + d_T(z_1)} + \frac{\sqrt{d_{T'}^2(z_3) + d_{T'}^2(z_4)}}{d_{T'}(z_3) + d_{T'}(z_4)} \\
&\quad - \frac{\sqrt{d_T^2(z_3) + d_T^2(z_4)}}{d_T(z_3) + d_T(z_4)} + \frac{\sqrt{d_{T'}^2(z_3) + d_{T'}^2(z_5)}}{d_{T'}(z_3) + d_{T'}(z_5)} - \frac{\sqrt{d_T^2(z_3) + d_T^2(z_5)}}{d_T(z_3) + d_T(z_5)} \\
&= \frac{\sqrt{20}}{7} - \frac{2\sqrt{5}}{3} + \frac{\sqrt{10} - 3\sqrt{2}}{2} - \frac{\sqrt{17}}{5} > 0.
\end{aligned}$$

Case 2. $d_T(z_3) = 4$.

Let $N_T(z_3) \setminus \{z\} = \{z_4, z_5, z_6\}$. Since T has a perfect matching, by Lemma 1 z_3 is matched with one of its neighbor, say z_4 . Thus $d_T(z_4) = 1$. $T' = T - \{u_1u_2, u_3u_4, zy, zz_2\} + \{u_1z_2, u_2z_2, u_3y, u_4y\}$.

$$\begin{aligned}
& DSO(T') - DSO(T) \\
&= \frac{\sqrt{d_{T'}^2(u_1) + d_{T'}^2(z_2)}}{d_{T'}(u_1) + d_{T'}(z_2)} + \frac{\sqrt{d_{T'}^2(u_2) + d_{T'}^2(z_2)}}{d_{T'}(u_2) + d_{T'}(z_2)} + \frac{\sqrt{d_{T'}^2(u_3) + d_{T'}^2(y)}}{d_{T'}(u_3) + d_{T'}(y)} \\
&\quad + \frac{\sqrt{d_{T'}^2(y) + d_{T'}^2(u_4)}}{d_{T'}(y) + d_{T'}(u_4)} - \frac{\sqrt{d_T^2(u_1) + d_T^2(u_2)}}{d_T(u_1) + d_T(u_2)} - \frac{\sqrt{d_T^2(u_3) + d_T^2(u_4)}}{d_T(u_3) + d_T(u_4)} \\
&\quad - \frac{\sqrt{d_T^2(z) + d_T^2(y)}}{d_T(z) + d_T(y)} - \frac{\sqrt{d_T^2(z) + d_T^2(z_2)}}{d_T(z) + d_T(z_2)} + \frac{\sqrt{d_{T'}^2(z) + d_{T'}^2(z_1)}}{d_{T'}(z) + d_{T'}(z_1)} \\
&\quad - \frac{\sqrt{d_T^2(z) + d_T^2(z_1)}}{d_T(z) + d_T(z_1)} + \frac{\sqrt{d_{T'}^2(z_3) + d_{T'}^2(z)}}{d_{T'}(z_3) + d_{T'}(z)} - \frac{\sqrt{d_T^2(z_3) + d_T^2(z)}}{d_T(z_3) + d_T(z)} \\
&= \frac{\sqrt{20}}{7} - \frac{2\sqrt{5}}{3} + \frac{\sqrt{10} - 3\sqrt{2}}{2} - \frac{\sqrt{17}}{5} > 0.
\end{aligned}$$

This complete the proof of this lemma. \square

Before presenting our main theorem, we define several classes of trees. Let $\mathcal{T}_{1,3}$ be the set of trees in which all vertices have degree 1 or 3. One can see that for any tree $T \in \mathcal{T}_{1,3}$ of order n , the number of pendant vertices is $\frac{n}{2} + 1$ and the number of vertices of degree 3 is $\frac{n}{2} - 1$. Three types of trees, denoted by $\mathcal{H}_i (i \in \{0, 1, 2\})$, are further obtained by $T \in \mathcal{T}_{1,3}$ inserting some vertices as follows:

$\mathcal{H}_0 = \{T : T \text{ is a tree obtained from } T' \in \mathcal{T}_{1,3} \text{ inserting exactly one vertex into each edge with both ends having degree 3}\}$.

$\mathcal{H}_1 = \{T : T \text{ is a tree obtained from } T' \in \mathcal{T}_{1,3} \text{ inserting exactly one vertex into all but one edge with both ends having degree 3}\}$.

$\mathcal{H}_2 = \{T : T \text{ is a tree obtained from } T' \in \mathcal{T}_{1,3} \text{ inserting exactly one vertex into all edges but two with both ends having degree 3}\}$. By the above definitions, if n

$$\text{is the order of a tree } T, \text{ then } n = \begin{cases} 3k + 7, & T \in \mathcal{H}_0 \\ 3k + 6, & T \in \mathcal{H}_1 \\ 3k + 8, & T \in \mathcal{H}_2, \end{cases}$$

where k is a nonnegative integer. Furthermore, let $\mathcal{G}_i = \{T : T \text{ is a tree obtained from } T' \in \mathcal{H}_i \text{ by adding } V(T') \text{ new vertices each of which joined to one vertex of } T'\}$ for each $i \in \{0, 1, 2\}$. Figure 3 shows some element in $\mathcal{G}_0, \mathcal{G}_1, \mathcal{G}_2$, respectively.




Figure 3. The tree with the maximum DSO index of molecular tree of order n ($n \geq 12$) with a perfect matching.

Theorem 3.4. *If T is a molecular tree of order n ($n \geq 12$) with perfect matching M , then*

$$DSO(T) \leq \begin{cases} \left(\frac{\sqrt{17}}{30} + \frac{\sqrt{5}}{9} + \frac{\sqrt{10}}{24} \cdot \frac{5}{21}\right)n - \frac{\sqrt{17}}{15} + \frac{10\sqrt{5}}{9} - \frac{10}{3} - \frac{40}{21}, & n - 12 \equiv 2 \pmod{6} \\ \left(\frac{\sqrt{17}}{30} + \frac{\sqrt{5}}{9} + \frac{\sqrt{10}}{24} \cdot \frac{5}{21}\right)n + \frac{4\sqrt{5}}{3} - \frac{\sqrt{10}}{2} - \frac{20}{7} - \frac{\sqrt{2}}{2}, & n - 12 \equiv 0 \pmod{6} \\ \left(\frac{\sqrt{17}}{30} + \frac{\sqrt{5}}{9} + \frac{\sqrt{10}}{24} \cdot \frac{5}{21}\right)n + \frac{\sqrt{17}}{15} + \frac{14\sqrt{5}}{9} - \frac{4\sqrt{10}}{6} - \frac{80}{21} - \sqrt{2}, & n - 12 \equiv 4 \pmod{6}. \end{cases}$$

For $n - 12 \equiv 2(\text{mod } 6)$, the equality holds if and only if $T \in \mathcal{G}_0$; for $n - 12 \equiv 0(\text{mod } 6)$, the equality holds if and only if $T \in \mathcal{G}_1$; for $n - 12 \equiv 4(\text{mod } 6)$, the equality holds if and only if $T \in \mathcal{G}_2$.

Proof. Let T^* be a molecular tree which has maximum DOS index with a perfect matching M . Let n_i be the number of vertices of degree i in T^* for each $i \in \{1, 2, 3, 4\}$. Clearly,

$$n_1 + n_2 + n_3 + n_4 = n \quad (14)$$

and

$$n_1 + 2n_2 + 3n_3 + 4n_4 = 2m = 2(n - 1). \quad (15)$$

By Lemma 1 and Lemma 2, we have

$$\begin{aligned} e_{1,2} + e_{1,3} + e_{1,4} + e_{2,4} + e_{3,4} + e_{4,4} &= n - 1 \\ 2n_2 + 3n_3 + n_4 + e_{4,4} &= n - 1. \end{aligned} \quad (16)$$

By Lemma 1, we have $n_1 = \frac{n}{2}$. Combining (14), (15) and (16), we have $n_3 = -2n_4 + \frac{n}{2} - 2$, $n_2 = n_4 + 2$, $e_{4,4} = 3n_4 - \frac{n}{2} + 1$.

Thus

$$\begin{aligned} DSO(T) &= \sum_{uv \in E_{1,2}} f(uv) + \sum_{uv \in E_{1,3}} f(uv) + \sum_{uv \in E_{1,4}} f(uv) + \sum_{uv \in E_{2,4}} f(uv) \\ &\quad + \sum_{uv \in E_{3,4}} f(uv) + \sum_{uv \in E_{4,4}} f(uv), \end{aligned}$$

where $f(uv) = \frac{\sqrt{d_G^2(u) + d_G^2(v)}}{d_G(u) + d_G(v)}$. We consider three cases in terms of n .

Case 1. $n = 6k + 14$, for an integer $k \geq 0$.

Combining $n = 6k + 14$, for an integer $k \geq 0$, we have $e_{4,4} = 3(n_4 - k) - 6$. By Lemma 3, we have $e_{4,4} = 0, 1, 2$. When $e_{4,4} = 1, 2$, it contradicts that k is an integer. Therefore, $e_{4,4} = 0$. Thus, $n_4 = k + 2$, $n_2 = k + 4$, $n_3 = k + 1$. By Lemma 1 and Lemma 2, $e_{2,4} = k + 4$, $e_{3,4} = 2k + 2$, $e_{1,2} = k + 4$, $e_{1,3} = k + 1$, $e_{1,4} = k + 2$. By the definition, $T^* \in \mathcal{G}_0$. Moreover,

$$\begin{aligned} DSO(T^*) &= \sum_{uv \in E_{1,2}} f(uv) + \sum_{uv \in E_{1,3}} f(uv) + \sum_{uv \in E_{1,4}} f(uv) + \sum_{uv \in E_{2,4}} f(uv) \\ &\quad + \sum_{uv \in E_{3,4}} f(uv) + \sum_{uv \in E_{4,4}} f(uv) \\ &= \left(\frac{2\sqrt{5}}{3} + \frac{\sqrt{10}}{4} + \frac{\sqrt{17}}{5} + \frac{10}{7} \right)k + \frac{\sqrt{5}}{3} + \frac{2\sqrt{17}}{5} + \frac{\sqrt{10}}{4} + \frac{10}{7} \\ &= \left(\frac{\sqrt{17}}{30} + \frac{\sqrt{5}}{9} + \frac{\sqrt{10}}{24} + \frac{5}{21} \right)n + \frac{10\sqrt{5}}{9} - \frac{\sqrt{17}}{15} - \frac{\sqrt{10}}{3} - \frac{40}{21}. \end{aligned}$$

Thus,

$$DSO(T) \leq \left(\frac{\sqrt{17}}{30} + \frac{\sqrt{5}}{9} + \frac{\sqrt{10}}{24} + \frac{5}{21} \right) n + \frac{10\sqrt{5}}{9} - \frac{\sqrt{17}}{15} - \frac{\sqrt{10}}{3} - \frac{40}{21}.$$

Case 2. $n = 6k + 12$, for an integer $k \geq 0$.

Combining $n = 6k + 12$, for an integer $k \geq 0$, we have $e_{4,4} = 3(n_4 - k) - 5$. By Lemma 3, we have $e_{4,4} = 0, 1, 2$. When $e_{4,4} = 0, 2$, it contradicts that k is an integer. Therefore, $e_{4,4} = 1$. Thus, $n_4 = k + 2, n_2 = k + 4, n_3 = k$. By Lemma 1 and Lemma 2, $e_{2,4} = k + 4, e_{3,4} = 2k, e_{1,2} = k + 4, e_{1,3} = k, e_{1,4} = k + 2$. By the definition, $T^* \in \mathcal{G}_1$. Moreover,

$$\begin{aligned} & DSO(T^*) \\ &= \sum_{uv \in E_{1,2}} f(uv) + \sum_{uv \in E_{1,3}} f(uv) + \sum_{uv \in E_{1,4}} f(uv) + \sum_{uv \in E_{2,4}} f(uv) \\ &\quad + \sum_{uv \in E_{3,4}} f(uv) + \sum_{uv \in E_{4,4}} f(uv) \\ &= \left(\frac{2\sqrt{5}}{3} + \frac{\sqrt{10}}{4} + \frac{\sqrt{17}}{5} + \frac{10}{7} \right) k + \frac{8\sqrt{5}}{3} + \frac{2\sqrt{17}}{5} + \frac{\sqrt{2}}{2} \\ &= \left(\frac{\sqrt{17}}{30} + \frac{\sqrt{5}}{9} + \frac{\sqrt{10}}{24} + \frac{5}{21} \right) n + \frac{4\sqrt{5}}{3} - \frac{\sqrt{10}}{2} - \frac{20}{7} + \frac{\sqrt{2}}{2}. \end{aligned}$$

Thus,

$$DSO(T) \leq \left(\frac{\sqrt{17}}{30} + \frac{\sqrt{5}}{9} + \frac{\sqrt{10}}{24} + \frac{5}{21} \right) n + \frac{4\sqrt{5}}{3} - \frac{\sqrt{10}}{2} - \frac{20}{7} + \frac{\sqrt{2}}{2}.$$

Case 3. $n = 6k + 16$, for an integer $k \geq 0$.

Combining $n = 6k + 16$, for an integer $k \geq 0$, we have $e_{4,4} = 3(n_4 - k) - 7$. By Lemma 3, we have $e_{4,4} = 0, 1, 2$. When $e_{4,4} = 0, 1$, it contradicts that k is an integer. Therefore, $e_{4,4} = 2$. Thus, $n_4 = k + 3, n_2 = k + 5, n_3 = k$. By Lemma 1 and Lemma 2, $e_{2,4} = k + 5, e_{3,4} = 2k, e_{1,2} = k + 5, e_{1,3} = k, e_{1,4} = k + 3$. By the definition, $T^* \in \mathcal{G}_2$. Moreover,

$$\begin{aligned} & DSO(T^*) \\ &= \sum_{uv \in E_{1,2}} f(uv) + \sum_{uv \in E_{1,3}} f(uv) + \sum_{uv \in E_{1,4}} f(uv) + \sum_{uv \in E_{2,4}} f(uv) \\ &\quad + \sum_{uv \in E_{3,4}} f(uv) + \sum_{uv \in E_{4,4}} f(uv) \\ &= \left(\frac{2\sqrt{5}}{3} + \frac{\sqrt{10}}{4} + \frac{\sqrt{17}}{5} + \frac{10}{7} \right) k + \frac{10\sqrt{5}}{3} + \frac{3\sqrt{17}}{5} + \sqrt{2} \\ &= \left(\frac{\sqrt{17}}{30} + \frac{\sqrt{5}}{9} + \frac{\sqrt{10}}{24} + \frac{5}{21} \right) n + \frac{\sqrt{17}}{15} + \frac{14\sqrt{5}}{9} - \frac{2\sqrt{10}}{3} - \frac{\sqrt{80}}{21} + \sqrt{2}. \end{aligned}$$

Thus,

$$DSO(T) \leq \left(\frac{\sqrt{17}}{30} + \frac{\sqrt{5}}{9} + \frac{\sqrt{10}}{24} + \frac{5}{21} \right) n + \frac{\sqrt{17}}{15} + \frac{14\sqrt{5}}{9} - \frac{2\sqrt{10}}{3} - \frac{\sqrt{80}}{21} + \sqrt{2}.$$

Summing up the above, we complete the proof. \square

Declarations of competing interest

The authors declare that have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- [1] A. Ali, B. Furtula, I. Redžepović, I. Gutman, Atom-bond sumconnectivity index, *J. Math. Chem.* **60**, 2081-2093 (2022)
- [2] J.A. Bondy, U.S.R. Murty, Graph Theory, Graduate Texts in Mathematics, vol. **244**, Springer, Heidelberg, (2008)
- [3] S.C. Basak, A.K. Bhattacharjee, Computational Approaches for the Design of Mosquito Repellent Chemicals. *Curr. Med. Chem.* **27**, 32-41 (2020)
- [4] S.C. Basak, M.G. Vracko, Parsimony Principle and its Proper use/ Application in Computer-assisted Drug Design and QSAR. *Curr. Comput. Aided Drug Des.* **16**, 1-5 (2020)
- [5] R. Cruz, I. Gutman, J. Rada, Sombor index of chemical graphs, *Appl. Math. Comput.* **399**, 126018 (2021)
- [6] J. W. Du, X. L. Su, On bond incident degree index of chemical trees with a fixed order and a fixed number of leaves, *Appl. Math. Comput.* **464**, 128390 (2024)
- [7] H. Y. Deng, Z. K. Tang, R. F. Wu, Molecular trees with extremal values of Sombor indices, *Int. J. Quantum Chem.* **121**, e26622 (2021)
- [8] I. Gutman, Geometric approach to degree-based topological indices: Sombor indices, *MATCH Commun. Math. Comput. Chem.* **86**, 11-16 (2021)
- [9] I. Gutman, Degree-based topological indices, *Croat. Chem. Acta* **86(4)**, 351-361 (2014)
- [10] I. Gutman, E. Milovanović, I. Milovanović, Beyond the Zagreb indices. *AKCE Int. J. Graphs Comb.* **17**, 74-85 (2018)

[11] I. Gutman, J. Tosović , Testing the quality of molecular structure descriptors: vertex-degree-based topological indices. *J. Serb. Chem. Soc.* **78**, 805-810 (2013)

[12] H. Liu, I. Gutman, L. You, Y. Huang, Sombor index: review of extremal results and bounds, *J. Math. Chem.* **60**, 771-798 (2022)

[13] F. Movahedi, M.H. Akhbari, Degree-based topological indices of the molecular structure of hyaluronic acid-methotrexate conjugates in cancer treatment, *Int. J. Quantum Chem.* vol.123 **7**, e27106 (2023)

[14] F. Movahedi, Maximal Diminished Sombor index of tricyclic graphs, *MATCH Commun. Math. Comput. Chem.* in press, (2025)

[15] F. Movahedi, Diminished Sombor index and its relationship with topological indices, submitted, (2025)

[16] F. Movahedi, I. Gutman, I. Redžepović, B. Furtula, Diminished Sombor index, *MATCH Commun. Math. Comput. Chem.* **95**, 141-162 (2026)

[17] I. Redžepović, Chemical applicability of Sombor indices, *J. Serb. Chem. Soc.* (2021)

[18] D. T. Rajathagiri, Enhanced mathematical models for the Sombor index: Reduced and co-Sombor index perspectives, *Data Anal. Artif. Intell.* **1(2)**, 215-228 (2021)

[19] R. Todeschini, V. Consonni, *Molecular Descriptors for Chemoinformatics* (Wiley-VCH, Weinheim, (2009)

[20] H. Wiener, Structural determination of paraffin boiling points. *J. Am. Chem. Soc.* **69**, 17-20 (1947)

[21] S. Wagner, H. Wang, G. Yu, Molecular graphs and the inverse Wiener index problem, *Discrete Appl. Math.* **157**, 1544-1554 (2009)

[22] F. X. Wang, B. Wu, The reduced Sombor index and the exponential reduced Sombor index of a molecular tree, *J. Math. Anal. Appl.* **515**, 126442 (2022)

[23] F. X. Wang, Y. Y. Zhang, X. Y. Ren, B. Zhang, Maximum Atom Bond Sum Connectivity Index of Molecular Trees with a Perfect Matching, *MATCH Commun. Math. Comput. Chem.* **92**, 653-669 (2024)