

A SEMICIRCLE LAW FOR THE NORMALIZED LAPLACIAN OF SPARSE RANDOM GRAPHS

YIMING CHEN, ZIJUN CHEN, AND YIZHE ZHU

ABSTRACT. We study the limiting spectral distribution of the normalized Laplacian \mathcal{L} of an Erdős-Rényi graph $G(n, p)$. To account for the presence of isolated vertices in the sparse regime, we define \mathcal{L} using the Moore-Penrose pseudoinverse of the degree matrix. Under this convention, we show that the empirical spectral distribution of a suitably normalized \mathcal{L} converges weakly in probability to the semicircle law whenever $np \rightarrow \infty$, thereby providing a rigorous justification of a prediction made in (Akara-pipattana and Evnin, 2023). Moreover, if $np > \log n + \omega(1)$, so that $G(n, p)$ has no isolated vertices with high probability, the same conclusion holds for the standard definition of \mathcal{L} . We further strengthen this result to almost sure convergence when $np = \Omega(\log n)$. Finally, we extend our approach to the Chung-Lu random graph model, where we establish a semicircle law for \mathcal{L} itself, improving upon (Chung, Lu, and Vu 2003), which obtained the semicircle law only for a proxy matrix.

1. INTRODUCTION

Spectral questions for random graphs lie at the interface of random matrix theory and random graph theory, and play a central role in a wide range of applications, including high-dimensional statistics [33] and theoretical computer science [29]. Given a graph, the eigenvalues and eigenvectors of its associated matrices control fundamental structural and algorithmic phenomena [21] such as expansion and connectivity, mixing and hitting times of random walks, and the performance of spectral algorithms for clustering and community detection. A canonical and extensively studied model is the Erdős-Rényi graph $G(n, p)$, whose adjacency matrix provides a prototypical example of a sparse random matrix. Over the past decades, global and local eigenvalue statistics, spectral norm bounds, and eigenvector properties for adjacency matrices have been developed across a broad range of sparsity regimes; see, e.g., [27, 40, 6, 3] and references therein.

In spectral graph theory [21], two particularly important operators are the (combinatorial) Laplacian $\mathbf{L} = \mathbf{D} - \mathbf{A}$, and the normalized Laplacian

$$\mathcal{L} = \mathbf{I} - \mathbf{D}^{-1/2} \mathbf{A} \mathbf{D}^{-1/2}, \quad (1)$$

where \mathbf{A} is the adjacency matrix and \mathbf{D} is the diagonal degree matrix. These matrices are ubiquitous in both theory and algorithms: \mathbf{L} is the basic object for cuts, connectivity, and electrical network interpretations [35], while \mathcal{L} (and the closely related random-walk matrix $\mathbf{D}^{-1} \mathbf{A}$) governs random-walk behavior and underpins spectral clustering [36]. Compared with adjacency matrices, however, the spectral theory of Laplacian-type matrices for random graphs is less complete, especially in regimes where degree fluctuations are significant. The normalization in (1) introduces a random and dependent reweighting through $\mathbf{D}^{-1/2}$, creating challenges that do not arise in the spectral analysis of \mathbf{A} .

Laplacian random matrices. For the Laplacian \mathbf{L} of an Erdős-Rényi graph, the empirical spectral distribution (after appropriate centering and scaling) has been shown to converge to the free convolution of a Gaussian distribution with the semicircle law as long as $np \rightarrow \infty$; see, e.g., [31, 25, 30, 14, 15, 13]. Beyond global limits, the behavior of extreme eigenvalues (or singular values) has also attracted considerable attention, with recent works establishing sharp asymptotics and fluctuation results in various regimes [12, 26, 18, 39]. Related deformed Laplacian matrices,

including the Bethe-Hessian, have been studied in the context of community detection in stochastic block models [37, 23, 38]. For directed sparse random graphs, limiting spectral laws and related phenomena have also been investigated; see [10].

The normalized Laplacian. A useful perspective on \mathcal{L} comes from random walks: writing

$$\mathcal{M} := \mathbf{D}^{-1/2} \mathbf{A} \mathbf{D}^{-1/2}, \quad \mathcal{L} = \mathbf{I} - \mathcal{M},$$

the matrix \mathcal{M} is a symmetric (reversible) Markov kernel obtained by symmetrizing the random-walk matrix $\mathbf{D}^{-1} \mathbf{A}$. Since the spectra of \mathcal{L} and \mathcal{M} have a deterministic relation, we will focus on the spectrum of \mathcal{M} below.

In dense regimes, since \mathbf{D} is concentrated around $(np)\mathbf{I}$, the semicircular law for \mathcal{M} is expected. When $np = \omega(\log n)$, Jiang [31] showed that the empirical spectral distribution of $\sqrt{np} \mathcal{M}$ converges almost surely to the semicircle law. In contrast, in the bounded expected degree regime, for a regularized version of \mathcal{M} , Chi [17] identified a non-degenerate symmetric limiting distribution characterized by return probabilities on Galton–Watson trees for $G(n, c/n)$.

In this paper, we bridge the gap between the constant expected-degree regime ($np = c$) and the supercritical regime ($np = \omega(\log n)$). We show that, as long as $np \rightarrow \infty$, the semicircle law for \mathcal{M} continues to hold, provided one adopts a convention for $\mathbf{D}^{-1/2}$ that remains well defined in the presence of isolated vertices (equivalently, by projecting away zero-degree coordinates). In view of the result of Chi [17], the condition $np \rightarrow \infty$ is both necessary and sufficient for the semicircle law to hold. The semicircle law for \mathcal{M} when $np \rightarrow \infty$ was also conjectured by Akara-pipattana and Evnin in [2, Section 5.2] based on predictions from the Fyodorov–Mirlin method, and we provide a rigorous justification.

Complementing asymptotic results, a substantial line of work establishes non-asymptotic extreme eigenvalue bounds of \mathcal{L} , motivated in part by stability guarantees for spectral clustering [32, 24, 16]; see also [28] for a sharp spectral-gap estimate restricted to the giant component of $G(n, p)$. Beyond random graph models, the limiting spectrum of the random walk matrix $\mathbf{D}^{-1} \mathbf{A}$ has also been considered for more general Hermitian [11, 8] or non-Hermitian random matrix \mathbf{A} [9].

The Chung–Lu model. Going beyond homogeneous Erdős–Rényi graphs, inhomogeneous random graphs $G(n, p_{ij})$, where edges (i, j) are present independently with probabilities p_{ij} , provide a flexible modeling framework that captures heterogeneous expected degrees and latent structure. A prominent special case is the rank-one (Chung–Lu) model [19, 22], which prescribes expected degrees via a factorized probability matrix. See Section 2.2 for a detailed description of the model. In this setting, Chung, Lu, and Vu [20] established a semicircle law for a matrix $\overline{\mathcal{M}} = \mathbb{E}[\mathbf{D}]^{-1/2} \mathbf{A} \mathbb{E}[\mathbf{D}]^{-1/2}$ approximating the normalized Laplacian, and further spectral estimates of \mathcal{L} were developed in [34]. Local spectral statistics for its adjacency matrix were investigated in [1].

We extend the semicircle law of the normalized Laplacian for $G(n, p)$ to the Chung–Lu model. In particular, we show that after normalization, a semicircle law holds for \mathcal{M} itself, rather than the proxy matrix $\overline{\mathcal{M}}$ studied in [20].

Beyond the rank-one setting, for general inhomogeneous random graphs, the limiting spectral distribution of \mathcal{M} need not be semicircular, even in dense regimes. If one further assumes that the variance profile of $\overline{\mathcal{M}}$ admits a graphon limit [43], the comparison argument developed in this paper can still be applied to characterize the corresponding limiting spectral distribution of \mathcal{L} .

Notation. Let $(\mathbb{R}, \mathcal{B})$ denote the measurable space, where \mathcal{B} is the Borel σ -algebra on \mathbb{R} . The indicator function is denoted by $\mathbf{1}(\cdot)$, and δ_x stands for the Dirac measure concentrated at $x \in \mathbb{R}^d$. The constant c may change from line to line. We also use the notation $[n] := \{1, 2, \dots, n\}$. For two sequences of nonnegative real numbers $\{a_n\}$ and $\{b_n\}$, we write $a_n = o(b_n)$ if $\lim_{n \rightarrow \infty} a_n/b_n = 0$, and $a_n = O(b_n)$ if $a_n \leq Cb_n$ for some $C > 0$ and all large n . We write $a_n \gg b_n$ if $a_n/b_n \rightarrow \infty$.

We denote by μ_{sc} the semicircle distribution supported on $[-2, 2]$, with density function

$$d\mu_{\text{sc}} = \frac{1}{2\pi} \sqrt{4 - x^2} \mathbf{1}\{|x| \leq 2\} dx. \quad (2)$$

Organization of the paper. The remainder of this paper is organized as follows. We state our main results for the normalized Laplacian of $G(n, p)$ and the Chung-Lu model in Section 2. Section 3 includes some basic facts about random matrix theory. We prove Theorems 1 and 2 in Section 4 and the proof of Theorem 3 is given in Section 5.

2. MAIN RESULTS

2.1. The Erdős-Rényi graph $G(n, p_n)$. Let $\mathbf{A}_n = (a_{ij}^{(n)})$ be the adjacency matrix of an Erdős-Rényi graph $G(n, p_n)$ such that $a_{ii}^{(n)} = 0, i \in [n]$. The normalized Laplacian matrix is defined by

$$\mathcal{L}_n = \mathbf{I}_n - \mathbf{D}_n^{-1/2} \mathbf{A}_n \mathbf{D}_n^{-1/2}, \quad (3)$$

where $\mathbf{D}_n = \text{diag}(\sum_{j \neq i} a_{ij}^{(n)})_{1 \leq i \leq n}$ is a diagonal matrix. If a graph has isolated vertices, \mathbf{D}_n is not invertible. Still, we can extend the definition of \mathbf{D}_n^{-1} using the Moore-Penrose pseudoinverse [41] by defining $\mathbf{D}_n^{-1}(i, i) = 0$ if $\mathbf{D}_n(i, i) = 0$. Under this convention, we show that the semicircle law holds when $np_n(1 - p_n) \rightarrow \infty$.

Theorem 1 (Semicircle law for \mathcal{L}_n). *Assume \mathbf{D}_n^{-1} is defined using the pseudo-inverse convention. Suppose that there exists a fixed constant $\varepsilon_0 > 0$ such that $\sup_n p_n \leq 1 - \varepsilon_0$ and*

$$np_n \rightarrow \infty \quad \text{as } n \rightarrow \infty.$$

Then the empirical spectral distribution of $\sqrt{\frac{np_n}{1-p_n}}(\mathbf{I}_n - \mathcal{L}_n)$ converges weakly to the semicircle law μ_{sc} defined in (2), in probability as $n \rightarrow \infty$.

Remark 1 (Optimality). *When $np_n = c$, [17, Theorem 5] showed that the limiting spectral distribution of $\mathbf{I}_n - \mathcal{L}_n$ is not semicircular. Consequently, the condition $np_n \rightarrow \infty$ is optimal for the semicircle law to hold, improving upon the stronger assumption $np_n = \omega(\log n)$ required in [31]. We note that Theorem 1 was also conjectured in [2, Section 5] based on predictions from the Fyodorov-Mirlin method.*

Since \mathbf{D}_n is invertible if and only if $G(n, p_n)$ has no isolated vertices, which occurs with high probability whenever $np_n > \log n + \omega(1)$ (see, e.g., [7, Theorem 7.3]), we obtain the following corollary of Theorem 1:

Corollary 1. *Assume that $\sup_n p_n < 1 - \varepsilon_0$ and there exists a sequence $c_n \rightarrow \infty$ such that*

$$np_n \geq \log n + c_n.$$

Then, with probability tending to one, the degree matrix \mathbf{D}_n is invertible. Moreover, the empirical spectral distribution of $\sqrt{\frac{np_n}{1-p_n}}(\mathbf{I}_n - \mathcal{L}_n)$ converges weakly in probability to the semicircle law as $n \rightarrow \infty$.

By further assuming that $np_n = \Omega(\log n)$, we can strengthen convergence in probability to almost sure convergence, as stated below:

Theorem 2. *Suppose that $\sup_n p_n \leq 1 - \varepsilon_0$, and $np_n \geq C \log n$ for some absolute constant C . Then, almost surely, the empirical spectral distribution of $\sqrt{\frac{np_n}{1-p_n}}(\mathbf{I}_n - \mathcal{L}_n)$ converges weakly to the semicircle law.*

A careful examination of the proof indicates that Theorem 2 holds for $C > 36$, although no attempt is made to optimize the numerical constant.

2.2. The Chung–Lu Model. The Chung–Lu model [19, 22] is a widely used random graph model for generating random graphs with a prescribed expected degree sequence. It can be viewed as a natural generalization of the Erdős–Rényi random graph. Let $n \in \mathbb{N}$ and let

$$\mathbf{w}_n = (w_{n,1}, w_{n,2}, \dots, w_{n,n})$$

be a sequence of positive weights. Define

$$\phi_n := \frac{1}{\sum_{i=1}^n w_{n,i}}.$$

The Chung–Lu random graph $G(n, \mathbf{w}_n)$ is an undirected random graph on the vertex set $[n] := \{1, 2, \dots, n\}$, in which edges between distinct vertices $i \neq j$ are placed independently with probabilities

$$\mathbb{P}(a_{ij}^{(n)} = 1) = p_{n,ij} := w_{n,i}w_{n,j}\phi_n, \quad \mathbb{P}(a_{ij}^{(n)} = 0) = 1 - p_{n,ij}.$$

Here $a_{ij}^{(n)}$ denotes the adjacency indicator between vertices i and j .

Let $\mathbf{A}_n = (a_{ij}^{(n)})_{1 \leq i,j \leq n}$ denote the adjacency matrix of $G(n, \mathbf{w}_n)$, and let $\mathbf{D}_n = \text{diag}(d_{n,1}, \dots, d_{n,n})$ be the corresponding degree matrix. We further denote by \mathbf{W}_n the diagonal matrix with diagonal entries $w_{n,i}$, representing the expected degrees of the vertices. A fundamental property of the Chung–Lu model is that the expected degree of vertex i satisfies

$$\mathbb{E}[d_{n,i}] = \sum_{j=1}^n p_{n,ij} \approx w_{n,i},$$

provided that

$$\max_{1 \leq i \leq n} w_{n,i}^2 < \sum_{j=1}^n w_{n,j},$$

which ensures that $p_{n,ij} \leq 1$ for all i, j . Consequently, the weight $w_{n,i}$ can be interpreted as the target expected degree of vertex i . As a special case, if $w_{n,i} \equiv np_n$ for all i , then $p_{n,ij} = p_n$, and the Chung–Lu model reduces to the classical Erdős–Rényi random graph $G(n, p_n)$.

We now state our main result under the Chung–Lu model.

Theorem 3 (Semicircle law for the Chung–Lu model). *Consider the Chung–Lu random graph $G(n, \mathbf{w}_n)$ defined above, and let*

$$w_{n,\min} := \min_{1 \leq i \leq n} w_{n,i}, \quad \bar{w}_n := \frac{1}{n} \sum_{i=1}^n w_{n,i}.$$

Assume that

$$w_{n,\min} \gg \sqrt{\bar{w}_n}. \tag{4}$$

Then the empirical spectral distribution of $\sqrt{\bar{w}_n}(\mathbf{I}_n - \mathcal{L}_n)$ converges weakly in probability to the semicircle law.

Remark 2 (Comparison with [20]). *Condition (4) implies that $w_{n,\min} \rightarrow \infty$ as $n \rightarrow \infty$. Under the same assumptions, Chung, Lu, and Vu [20, Theorem 6] established a semicircle law only for the proxy matrix $\mathbf{W}_n^{-1/2} \mathbf{A}_n \mathbf{W}_n^{-1/2}$. The present work closes this gap by establishing the semicircle law for the normalized Laplacian itself.*

3. PRELIMINARIES

Let \mathbf{M} be a symmetric matrix with eigenvalues $\lambda_1(\mathbf{M}) \geq \lambda_2(\mathbf{M}) \geq \dots \geq \lambda_n(\mathbf{M})$. The empirical spectral distribution (ESD) of \mathbf{M} , denoted by $\hat{\mu}(\mathbf{M})$, is defined as

$$\hat{\mu}(\mathbf{M}) = \frac{1}{n} \sum_{j=1}^n \delta_{\lambda_j(\mathbf{M})}.$$

If \mathbf{M} is random, then $\hat{\mu}(\mathbf{M})$ is a random probability measure on $(\mathbb{R}, \mathcal{B})$. We write $\hat{\mu}(\mathbf{M}_n) \Rightarrow \mu$ to denote that $\hat{\mu}(\mathbf{M}_n)$ converges weakly to a probability measure μ on $(\mathbb{R}, \mathcal{B})$.

Definition 1 (bounded Lipschitz metric). *Let d_{BL} denote the bounded Lipschitz metric*

$$d_{BL}(\mu, \nu) = \sup \left\{ \int f \, d\mu - \int f \, d\nu : \|f\|_\infty + \|f\|_L \leq 1 \right\},$$

where $\|f\|_\infty = \sup_x |f(x)|$, $\|f\|_L = \sup_{x \neq y} \frac{|f(x) - f(y)|}{|x - y|}$.

We will use the following comparison inequality to establish weak convergence of empirical spectral distributions:

Lemma 1. *For any symmetric matrices A, B , we have*

$$d_{BL}^2(\hat{\mu}(\mathbf{A}), \hat{\mu}(\mathbf{B})) \leq \frac{1}{n} \text{tr}((\mathbf{B} - \mathbf{A})^2).$$

Proof. For the spectral measures of $n \times n$ symmetric real matrices \mathbf{A}, \mathbf{B} we have

$$\begin{aligned} d_{BL}(\hat{\mu}(\mathbf{A}), \hat{\mu}(\mathbf{B})) &\leq \sup \left\{ \frac{1}{n} \sum_{j=1}^n |f(\lambda_j(\mathbf{A})) - f(\lambda_j(\mathbf{B}))| : \|f\|_L \leq 1 \right\} \\ &\leq \frac{1}{n} \sum_{j=1}^n |\lambda_j(\mathbf{A}) - \lambda_j(\mathbf{B})| \\ &\leq \frac{1}{\sqrt{n}} \sqrt{\sum_{j=1}^n |\lambda_j(\mathbf{A}) - \lambda_j(\mathbf{B})|^2} \leq \frac{1}{\sqrt{n}} \sqrt{\text{tr}((\mathbf{B} - \mathbf{A})^2)}, \end{aligned}$$

where the last inequality is from Lidskii's theorem (see, e.g., [4, Lemma 2.3]). \square

4. PROOF OF THEOREMS 1 AND 2

Before proceeding, we briefly outline the overall proof strategy. The proof is divided into several steps. From Ding and Jiang [25], the empirical spectral distribution of $\mathbf{A}_n / \sqrt{np_n(1-p_n)}$ converges weakly to the semicircle law when $np_n(1-p_n) \rightarrow \infty$. Therefore, it suffices to show that the bounded Lipschitz distance between the empirical spectral distribution under consideration, and that of $\mathbf{A}_n / \sqrt{np_n(1-p_n)}$ converges to zero almost surely as $n \rightarrow \infty$. A direct control of d_{BL} is technically challenging. To address this issue, we decompose the analysis into two parts: the expectation term and the deviation from its expectation. The expectation is handled in Lemma 2, where we derive a suitable tail estimate, while the fluctuation around the expectation is controlled in Lemma 3 by an improved bounded differences inequality.

Lemma 2 is first used to establish Theorem 1. Then, by combining Lemmas 2 and 3, we show that the bounded Lipschitz distance converges to zero almost surely, which in turn yields the desired spectral convergence and completes the proof of Theorem 2.

Set

$$\tilde{\mathcal{L}}_n = \mathbf{I}_n - \frac{1}{(n-1)p_n} \mathbf{A}_n, \quad u_n = (n-1)p_n, \quad d_{n,i} = \sum_{j \neq i} a_{ij}^{(n)}. \quad (5)$$

Note that

$$\begin{aligned} \tilde{\mathcal{L}}_n - \mathcal{L}_n &= -\frac{1}{(n-1)p_n} \mathbf{A}_n + \text{diag}(d_{n,1}^{-1/2}, \dots, d_{n,n}^{-1/2}) \mathbf{A}_n \text{diag}(d_{n,1}^{-1/2}, \dots, d_{n,n}^{-1/2}) \\ &= \left(\left(-\frac{1}{(n-1)p_n} + \frac{1}{\sqrt{d_{n,i}}\sqrt{d_{n,j}}} \right) a_{ij}^{(n)} \right)_{n \times n}. \end{aligned}$$

Lemma 2 (Approximation of $\frac{1}{n}\mathbb{E}(\text{tr}(\tilde{\mathcal{L}}_n - \mathcal{L}_n)^2)$). *Assume that the matrices \mathcal{L}_n and $\tilde{\mathcal{L}}_n$ are defined as (3) and (5). If $np_n \rightarrow \infty$, then*

$$\frac{1}{n}\mathbb{E}(\text{tr}(\tilde{\mathcal{L}}_n - \mathcal{L}_n)^2) = O(u_n^{-2}).$$

Lemma 3 (Concentration). *Assume that the matrices \mathcal{L}_n and $\tilde{\mathcal{L}}_n$ are defined as (3) and (5) and $np_n \geq C \log n$. Then, asymptotically almost surely,*

$$\frac{1}{n} \text{tr}(\tilde{\mathcal{L}}_n - \mathcal{L}_n)^2 - \frac{1}{n}\mathbb{E}(\text{tr}(\tilde{\mathcal{L}}_n - \mathcal{L}_n)^2) = o(u_n^{-1}).$$

4.1. Proof of Lemma 2. Through symmetry and the fact that $\{a_{ij}^{(n)}\}$ are i.i.d. random variables, we can derive that

$$\begin{aligned} \frac{1}{n}\mathbb{E}[\text{tr}(\tilde{\mathcal{L}}_n - \mathcal{L}_n)^2] &= \frac{1}{n}\mathbb{E}\left[\sum_{i \neq j}^n \left(-\frac{1}{u_n} + \frac{1}{\sqrt{d_{n,i}}\sqrt{d_{n,j}}}\right)^2 a_{ij}^{(n)}\right] \\ &= \frac{1}{n} \sum_{i \neq j}^n \mathbb{E}\left[\left(-\frac{1}{u_n} + \frac{1}{\sqrt{d_{n,i}}\sqrt{d_{n,j}}}\right)^2 a_{ij}^{(n)}\right] \\ &= \frac{1}{n} \cdot n(n-1)\mathbb{E}\left[\left(-\frac{1}{u_n} + \frac{1}{\sqrt{d_{n,1}}\sqrt{d_{n,2}}}\right)^2 a_{12}^{(n)}\right] \\ &= (n-1)\mathbb{E}\left[\left(-\frac{1}{u_n} + \frac{1}{\sqrt{d_{n,1}}\sqrt{d_{n,2}}}\right)^2 a_{12}^{(n)}\right]. \end{aligned} \quad (6)$$

Note that $\mathbb{P}(a_{12}^{(n)} = 1) = 1 - \mathbb{P}(a_{12}^{(n)} = 0) = p_n$, we get

$$\mathbb{E}\left[\left(-\frac{1}{u_n} + \frac{1}{\sqrt{d_{n,1}}\sqrt{d_{n,2}}}\right)^2 a_{12}^{(n)}\right] = p_n \mathbb{E}\left[\left(-\frac{1}{u_n} + \frac{1}{\sqrt{d_{n,1}}\sqrt{d_{n,2}}}\right)^2 | a_{12}^{(n)} = 1\right].$$

We briefly outline our approach for evaluating

$$\mathbb{E}\left[\left(-\frac{1}{u_n} + \frac{1}{\sqrt{d_{n,1}}\sqrt{d_{n,2}}}\right)^2 | a_{12}^{(n)} = 1\right].$$

We introduce an event E_1 under which both $d_{n,1}$ and $d_{n,2}$ remain close to u_n , so that the function $(x, y) \mapsto \left(\frac{1}{u_n} - \frac{1}{\sqrt{xy}}\right)^2$ can be locally approximated. The expectation is then decomposed into two parts corresponding to E_1 and its complement E_1^c . For E_1 , we use independence and moment estimates of binomial variables to show that the contribution is $O(u_n^{-3})$, while for E_1^c , a Chernoff

bound ensures that its probability is exponentially small. Combining these estimates yields the desired order of magnitude for the entire expectation.

Given that $a_{12}^{(n)} = 1$, let

$$d_{n,1} = 1 + Z_1 := 1 + \sum_{k \neq 1,2} a_{1k}^{(n)}, \quad d_{n,2} = 1 + Z_2 := 1 + \sum_{k \neq 1,2} a_{2k}^{(n)}.$$

So we have

$$Z_1 \sim \text{Bin}(n-2, p_n) \quad \text{and} \quad Z_2 \sim \text{Bin}(n-2, p_n),$$

where $\text{Bin}(n, p)$ denotes the binomial distribution with parameters n and p . Thus

$$\begin{aligned} \mathbb{E}(Z_1 + 1) &= \mathbb{E}(Z_2) = (n-2)p_n + 1 = u_n + o(u_n), \\ \mathbb{E}((Z_1 + 1)^2) &= \mathbb{E}((Z_2 + 1)^2) = 3(n-2)p_n + (n-2)(n-3)p_n^2 + 1 = u_n^2 + O(u_n). \end{aligned}$$

Let the event

$$E_1 := \{d_{n,1}, d_{n,2} \in [u_n/2, 3u_n/2]\}.$$

We first approximate

$$\mathbb{E}\left(\left(-\frac{1}{u_n} + \frac{1}{\sqrt{d_{n,1}}\sqrt{d_{n,2}}}\right)^2 \mathbf{1}(E_1) | a_{12}^{(n)} = 1\right).$$

Using the identity $|\frac{1}{\sqrt{x}} - \frac{1}{\sqrt{y}}| = \left|\frac{\sqrt{y} - \sqrt{x}}{\sqrt{xy}}\right| = \left|\frac{y-x}{\sqrt{xy}(\sqrt{x} + \sqrt{y})}\right| \leq \left|\frac{y-x}{y\sqrt{x}}\right|$ for $x = (Z_1 + 1)(Z_2 + 1)$, $y = u_n^2$, we obtain

$$\begin{aligned} \mathbb{E}\left(\left(-\frac{1}{u_n} + \frac{1}{\sqrt{d_{n,1}}\sqrt{d_{n,2}}}\right)^2 \mathbf{1}(E_1) | a_{12}^{(n)} = 1\right) &= \mathbb{E}\left(\left(-\frac{1}{u_n} + \frac{1}{\sqrt{Z_1 + 1}\sqrt{Z_2 + 1}}\right)^2 \mathbf{1}(E_1)\right) \\ &\leq \mathbb{E}\left(\left(\frac{(Z_1 + 1)(Z_2 + 1) - u_n^2}{u_n^2\sqrt{Z_1 + 1}\sqrt{Z_2 + 1}}\right)^2 \mathbf{1}(E_1)\right). \end{aligned}$$

By non-negativity, note that $((Z_1 + 1)(Z_2 + 1) - u_n^2)^2 \mathbf{1}(E_1) \leq ((Z_1 + 1)(Z_2 + 1) - u_n^2)^2$, and when event E_1 holds, $(Z_i + 1)^{-1} \leq 2/u_n$ for $i = 1, 2$. Since Z_1 and Z_2 are independent, we can therefore bound the expectation by

$$\begin{aligned} \mathbb{E}\left(\left(\frac{(Z_1 + 1)(Z_2 + 1) - u_n^2}{u_n^2\sqrt{Z_1 + 1}\sqrt{Z_2 + 1}}\right)^2 \mathbf{1}(E_1)\right) &\leq \mathbb{E}\left(\frac{4((Z_1 + 1)(Z_2 + 1) - u_n^2)^2}{u_n^6}\right) \\ &= \frac{4}{u_n^6} \cdot \mathbb{E}\left((Z_1 + 1)^2(Z_2 + 1)^2 - 2u_n^2(Z_1 + 1)(Z_2 + 1) + u_n^4\right) \\ &= \frac{4}{u_n^6} \cdot \left(\mathbb{E}\left((Z_1 + 1)^2\right)\mathbb{E}\left((Z_2 + 1)^2\right) - 2u_n^2\mathbb{E}(Z_1 + 1)\mathbb{E}(Z_2 + 1) + u_n^4\right) \\ &= \frac{4}{u_n^6}(u_n^4 + O(u_n^3) - 2u_n^4 + o(u_n^3) + u_n^4) \\ &= O(u_n^{-3}). \end{aligned} \tag{7}$$

Under the assumption $np_n \rightarrow \infty$, a direct application of Chernoff's bound yields

$$\mathbb{P}(E_1^c | a_{12} = 1) \leq 2 \exp(-u_n/12).$$

When event E_1^c occurs, observe that

$$\left|\frac{1}{u_n} - \frac{1}{\sqrt{d_{n,1}}\sqrt{d_{n,2}}}\right| \leq \left|\frac{1}{u_n}\right| + \left|\frac{1}{\sqrt{d_{n,1}}\sqrt{d_{n,2}}}\right| \leq 1 + 1 = 2,$$

where we use the pseudo-inverse convention of \mathbf{D}_n^{-1} .

Thus we have

$$\mathbb{E} \left(\left(-\frac{1}{u_n} + \frac{1}{\sqrt{d_{n,1}} \sqrt{d_{n,2}}} \right)^2 \mathbf{1}(E_1^C) | a_{12}^{(n)} = 1 \right) \leq 4\mathbb{P}(E_1^c) \leq 8 \exp(-u_n/12). \quad (8)$$

Since $u_n \rightarrow \infty$ as $n \rightarrow \infty$, it follows that

$$\exp(-u_n/12) = o(u_n^{-3}).$$

Combine (7) and (8), we get

$$\begin{aligned} \mathbb{E} \left(\left(-\frac{1}{u_n} + \frac{1}{\sqrt{d_{n,1}} \sqrt{d_{n,2}}} \right)^2 | a_{12}^{(n)} = 1 \right) &= \mathbb{E} \left(\left(-\frac{1}{u_n} + \frac{1}{\sqrt{d_{n,1}} \sqrt{d_{n,2}}} \right)^2 (\mathbf{1}(E_1) + \mathbf{1}(E_1^c)) | a_{12}^{(n)} = 1 \right) \\ &= O(u_n^{-3}). \end{aligned} \quad (9)$$

Substituting (9) into (6), we obtain

$$\frac{1}{n} \mathbb{E} \left[\text{tr} \left(\tilde{\mathcal{L}}_n - \mathcal{L}_n \right)^2 \right] = (n-1)p_n \cdot O(u_n^{-3}) = O(u_n^{-2}).$$

4.2. Proof of Theorem 1. We begin by comparing the empirical spectral distributions (ESDs) of \mathcal{L}_n and $\tilde{\mathcal{L}}_n$. Since the asymptotic spectral behavior of $\tilde{\mathcal{L}}_n$ has already been established in Ding and Jiang [25], it is enough to show that

$$d_{\text{BL}} \left(\hat{\mu}(\mathcal{L}_n), \hat{\mu}(\tilde{\mathcal{L}}_n) \right) \rightarrow 0 \quad \text{in probability as } n \rightarrow \infty.$$

By Lemma 2, we have

$$\mathbb{E} \left[d_{\text{BL}}^2 \left(\hat{\mu}(\tilde{\mathcal{L}}_n), \hat{\mu}(\mathcal{L}_n) \right) \right] \leq \frac{1}{n} \mathbb{E} \left[\text{tr} \left(\tilde{\mathcal{L}}_n - \mathcal{L}_n \right)^2 \right] = O(u_n^{-2}). \quad (10)$$

By Markov's inequality, (10) implies that the ESDs of \mathcal{L}_n and $\tilde{\mathcal{L}}_n$ are asymptotically indistinguishable in the d_{BL} metric. The same argument applies to scaled and shifted versions of \mathcal{L}_n and $\tilde{\mathcal{L}}_n$. Indeed, for any sequences of constants $\{\gamma_n\}$ and $\{\rho_n\}$,

$$\mathbb{E} \left[d_{\text{BL}}^2 \left(\hat{\mu}(\gamma_n \mathcal{L}_n + \rho_n \mathbf{I}_n), \hat{\mu}(\gamma_n \tilde{\mathcal{L}}_n + \rho_n \mathbf{I}_n) \right) \right] = O \left(\frac{\gamma_n^2}{u_n^2} \right).$$

Applying Markov's inequality yields

$$\mathbb{P} \left(d_{\text{BL}}^2 \left(\hat{\mu}(\gamma_n \mathcal{L}_n + \rho_n \mathbf{I}_n), \hat{\mu}(\gamma_n \tilde{\mathcal{L}}_n + \rho_n \mathbf{I}_n) \right) \geq \varepsilon \right) \leq c \frac{\gamma_n^2}{u_n^2 \varepsilon}.$$

Now choose

$$\gamma_n = -\rho_n = -\sqrt{np_n/(1-p_n)}. \quad (11)$$

Then

$$\frac{\gamma_n^2}{u_n^2} = O \left(\frac{1}{np_n(1-p_n)} \right) = o(1),$$

and hence

$$\hat{\mu}(\gamma_n \mathcal{L}_n + \rho_n \mathbf{I}_n) \Rightarrow \hat{\mu}(\gamma_n \tilde{\mathcal{L}}_n + \rho_n \mathbf{I}_n) \quad \text{in probability as } n \rightarrow \infty. \quad (12)$$

By the result of Ding and Jiang [25], we have

$$\frac{1}{n} \sum_{i=1}^n \mathbf{1} \left(\sqrt{\frac{np_n}{1-p_n}} \left(1 - \lambda_i(\tilde{\mathcal{L}}_n) \right) \leq x \right) \Rightarrow \mu_{\text{sc}}, \quad \text{a.s. as } n \rightarrow \infty.$$

With (11), for all $1 \leq i \leq n$,

$$\lambda_{n-i} \left(\gamma_n \tilde{\mathcal{L}}_n + \rho_n \mathbf{I}_n \right) = \sqrt{\frac{np_n}{1-p_n}} \left(1 - \lambda_i(\tilde{\mathcal{L}}_n) \right).$$

Therefore,

$$\hat{\mu}(\gamma_n \tilde{\mathcal{L}}_n + \rho_n \mathbf{I}_n) \implies \mu_{\text{sc}} \quad \text{a.s. as } n \rightarrow \infty. \quad (13)$$

Finally, combining (12) and (13) yields

$$\hat{\mu}(\gamma_n \mathcal{L}_n + \rho_n \mathbf{I}_n) \implies \mu_{\text{sc}} \quad \text{in probability as } n \rightarrow \infty,$$

which completes the proof.

4.3. Proof of Lemma 3. Firstly, we introduce the following result of Warnke [42].

Lemma 4 (Typical bounded differences inequality). *Let $X = (X_1, \dots, X_N)$ be a family of independent random variables with X_k taking values in a set Λ_k . Let $\Gamma \subseteq \prod_{j \in [N]} \Lambda_j$ be an event, and assume that the function $F : \prod_{j \in [N]} \Lambda_j \rightarrow \mathbb{R}$ satisfies the following typical Lipschitz condition:*

(TL) *There exist sequences $(c_k)_{k \in [N]}$ and $(c'_k)_{k \in [N]}$ with $c_k \leq c'_k$ such that whenever $x, \sigma_k(x) \in \prod_{j \in [N]} \Lambda_j$ differ only in the k -th coordinate, we have*

$$|F(x) - F(\sigma_k(x))| \leq \begin{cases} c_k, & \text{if } x \in \Gamma, \\ c'_k, & \text{otherwise.} \end{cases}$$

For any $(\beta_k)_{k \in [N]}$ with $\beta_k \in (0, 1]$, there exists an event $\mathcal{A} = \mathcal{A}(\Gamma, (\beta_k)_{k \in [N]})$ satisfying

$$\mathbb{P}(\mathcal{A}) \leq \sum_{k \in [N]} \beta_k^{-1} \mathbb{P}(X \notin \Gamma), \quad \neg \mathcal{A} \subseteq \Gamma, \quad (14)$$

such that, for $\mu = \mathbb{E}F(X)$, $\varepsilon_k = \beta_k(c'_k - c_k)$, and any $t \geq 0$,

$$\mathbb{P}(F(X) \geq \mu + t \text{ and } \neg \mathcal{A}) \leq \exp\left(-\frac{t^2}{2 \sum_{k \in [N]} (c_k + \varepsilon_k)^2}\right).$$

Proof of Lemma 3. Consider the function $f_n : \{0, 1\}^{n \times n} \rightarrow \mathbb{R}$ defined by

$$f_n(\{X_{ij}\}_{1 \leq i, j \leq n}) := \sum_{i \neq j}^n \left(u_n^{-1} - \left(\sum_{k \neq i} X_{ik} \sum_{k \neq j} X_{jk} \right)^{-1/2} \right)^2 X_{ij}, \quad (15)$$

which depends only on the adjacency variables $\{X_{ij}\}$ of the random graph. Then

$$F = \frac{1}{n} f_n = \frac{1}{n} \text{tr}(\tilde{\mathcal{L}}_n - \mathcal{L}_n)^2.$$

If we change only one coordinate (u, v) of $\{X_{ij}\}$ while keeping all others fixed, define

$$\sigma_{uv}(\{X_{ij}\}) := \begin{cases} 1 - X_{ij}, & \text{if } (i, j) = (u, v) \text{ or } (i, j) = (v, u), \\ X_{ij}, & \text{otherwise.} \end{cases}$$

Let $S_l = \sum_{k \neq l}^n X_{kl}$ and $S'_l = \sum_{k \neq l}^n \sigma_{uv}(\{X_{ij}\})_{kl}$. It is easy to verify that

$$|S_k - S'_k| = \begin{cases} 1, & \text{if } k = u \text{ or } k = v, \\ 0, & \text{otherwise.} \end{cases}$$

By definition of f_n in (15), we obtain

$$\begin{aligned}\delta_{uv}(\{X_{i,j}\}) &:= f_n(\{X_{i,j}\}) - f_n(\sigma_{uv}(\{X_{ij}\})) \\ &= 2 \sum_{k \neq u,v}^n \left[\left(u_n^{-1} - (S_u S_k)^{-1/2} \right)^2 - \left(u_n^{-1} - (S'_u S_k)^{-1/2} \right)^2 \right] X_{uk} \\ &\quad + 2 \sum_{k \neq u,v}^n \left[\left(u_n^{-1} - (S_v S_k)^{-1/2} \right)^2 - \left(u_n^{-1} - (S'_v S_k)^{-1/2} \right)^2 \right] X_{vk} \\ &\quad + 2 \left[\left(u_n^{-1} - (S_u S_v)^{-1/2} \right)^2 - \left(u_n^{-1} - (S'_u S'_v)^{-1/2} \right)^2 \right].\end{aligned}$$

Define the “*typical*” event

$$\Gamma := \left\{ d_{n,i} \in \left(\frac{u_n}{2}, \frac{3u_n}{2} \right) \text{ for all } 1 \leq i \leq n \right\}.$$

We now estimate the typical Lipschitz constants c_k .

Consider the auxiliary function $g(x, y) = \left(u_n^{-1} - \frac{1}{\sqrt{xy}} \right)^2$. A direct computation yields

$$\frac{\partial g}{\partial x} = 2 \left(u_n^{-1} - \frac{1}{\sqrt{xy}} \right) \left(-\frac{1}{2} x^{-3/2} y^{-1/2} \right) = \left(u_n^{-1} - \frac{1}{\sqrt{xy}} \right) x^{-3/2} y^{-1/2}.$$

For $x, y \in (u_n/2, 3u_n/2)$, we have $x^{-3/2} y^{-1/2} \leq 4u_n^{-2}$ and $|u_n^{-1} - x^{-1/2} y^{-1/2}| \leq u_n^{-1}$, hence $|\partial g / \partial x| \leq 4u_n^{-3}$. By symmetry, the same holds for $\partial g / \partial y$. Consequently, if both x and y change by at most 1, the function $g(x, y)$ changes by at most cu_n^{-3} .

When the event Γ holds, changing one coordinate (u, v) in $\{X_{ij}\}$ affects at most the degrees of vertices u and v . Recalling that $d_{n,i} = \sum_{j \neq i} a_{ij}$, we have

$$\begin{aligned}|\delta_{uv}(\mathbf{A}_n)| &\leq 2 \sum_{k \neq u,v}^n c u_n^{-3} a_{uk} + 2 \sum_{k \neq u,v}^n c u_n^{-3} a_{vk} + c u_n^{-3} \\ &\leq c u_n^{-3} (d_{n,u} + d_{n,v} + 1) \leq 3 c u_n^{-2},\end{aligned}$$

since $d_{n,i} \in [u_n/2, 3u_n/2]$ under Γ . Hence, for $F := n^{-1} f_n$, the *typical* Lipschitz constants satisfy

$$c_{uv} \leq \frac{C_1}{n u_n^2},$$

whenever the event Γ holds.

Outside the typical event Γ , we distinguish two cases.

First, if $S_u = 0$ or $S'_u = 0$, then necessarily $X_{uk} = 0$ for all $k \neq u, v$. In this case the corresponding contribution vanishes identically, namely,

$$\left| \sum_{k \neq u,v}^n \left[\left(u_n^{-1} - (S_u S_k)^{-1/2} \right)^2 - \left(u_n^{-1} - (S'_u S_k)^{-1/2} \right)^2 \right] X_{uk} \right| = 0.$$

Otherwise, assume that $S_u S'_u \neq 0$. Using the identities $x^2 - y^2 = (x+y)(x-y)$ and

$$x^{-1/2} - y^{-1/2} = \frac{x - y}{xy(x^{-1/2} + y^{-1/2})},$$

we obtain

$$\begin{aligned}
& \left| \sum_{k \neq u, v}^n \left[(u_n^{-1} - (S_u S_k)^{-1/2})^2 - (u_n^{-1} - (S'_u S_k)^{-1/2})^2 \right] X_{uk} \right| \\
& \leq \left| \sum_{k \neq u, v}^n \left(2u_n^{-1} - (S_u S_k)^{-1/2} - (S'_u S_k)^{-1/2} \right) \left((S_u S_k)^{-1/2} - (S'_u S_k)^{-1/2} \right) X_{uk} \right| \\
& \leq c \left| \sum_{k \neq u, v}^n \left((S_u S_k)^{-1/2} - (S'_u S_k)^{-1/2} \right) X_{uk} \right| \\
& \leq c \max_k S_k^{-1/2} \frac{|S_u - S'_u|}{S_u S'_u (S_u^{-1/2} + S'_u^{-1/2})} \sum_{k \neq u, v}^n X_{uk} \\
& \leq c \max_k S_k^{-1/2} \frac{S_u}{S_u S'_u (S_u^{-1/2} + S'_u^{-1/2})} \\
& = c \max_k S_k^{-1/2} \frac{1}{S'_u (S_u^{-1/2} + S'_u^{-1/2})}.
\end{aligned}$$

Under the pseudo-inverse convention, if there exists $1 \leq i \leq n$ such that $d_{n,i} = O(1)$, then the increment δ_{uv} may attain its maximal size, and we obtain the crude bound

$$\delta_{uv} = O(1).$$

Therefore, recall that $F := n^{-1} f_n$, we also have the crude *worst-case* bound

$$c'_{uv} \leq \frac{C_2}{n},$$

for some absolute constant $C_2 > 0$.

We choose the parameters uniformly over edges $k \leftrightarrow (u, v)$ by

$$\beta_k = \beta_n := u_n^{-2} \in (0, 1] \quad \text{for all sufficiently large } n,$$

so that $\varepsilon_k = \beta_n (c'_k - c_k) \leq C_2 n^{-1} u_n^{-2}$. Since $u_n \rightarrow \infty$ under the assumption $np_n \gtrsim \log n$, the typical constants dominate:

$$c_k + \varepsilon_k \leq (C_1 + C_2) n^{-1} u_n^{-2}.$$

Summing over the $N = \binom{n}{2}$ independent edges gives

$$\sum_k (c_k + \varepsilon_k)^2 \leq N \cdot \frac{(C_1 + C_2)^2}{n^2 u_n^4} = O(u_n^{-4}).$$

Therefore, by (14), we obtain the sub-Gaussian tail bound

$$\mathbb{P}(F - \mathbb{E}F \geq t \text{ and } \neg \mathcal{A}) \leq \exp(-c t^2 u_n^4). \quad (16)$$

for some absolute constant $c > 0$.

It remains to control $\mathbb{P}(\mathcal{A})$. Since $d_{n,i} \sim \text{Bin}(n-1, p_n)$ and $u_n = (n-1)p_n$, Chernoff's bound gives

$$\mathbb{P}(|d_{n,i} - u_n| \geq u_n/2) \leq 2e^{-u_n/12}.$$

By a union bound,

$$\mathbb{P}(\neg \Gamma) \leq 2ne^{-u_n/12}.$$

This also implies \mathbf{D}_n is invertible asymptotically almost surely. Using $\beta_n = u_n^{-2}$, we deduce from (14) that

$$\mathbb{P}(\mathcal{A}) \leq N \beta_n^{-1} \mathbb{P}(\neg \Gamma) \lesssim n^3 u_n^2 e^{-u_n/12}, \quad (17)$$

which is summable in n under $u_n \gtrsim \log n$ (equivalently, $np_n \gtrsim \log n$).

Finally, set $t_n = k/u_n$ with any fixed $k > 0$. By (16) and (17),

$$\sum_{n=1}^{\infty} \mathbb{P}(F - \mathbb{E}F \geq t_n) \leq \sum_{n=1}^{\infty} \mathbb{P}(F - \mathbb{E}F \geq t_n \text{ and } \neg \mathcal{A}) + \sum_{n=1}^{\infty} \mathbb{P}(\mathcal{A}) < \infty.$$

By the Borel–Cantelli lemma,

$$\frac{1}{n} \text{tr}(\tilde{\mathcal{L}}_n - \mathcal{L}_n)^2 - \frac{1}{n} \mathbb{E}[\text{tr}(\tilde{\mathcal{L}}_n - \mathcal{L}_n)^2] = o\left(\frac{1}{u_n}\right) \quad \text{a.s. as } n \rightarrow \infty,$$

which establishes the desired concentration result. \square

4.4. Proof of Theorem 2.

Proof. It suffices to show that

$$d_{\text{BL}}(\hat{\mu}(\mathcal{L}_n), \hat{\mu}(\tilde{\mathcal{L}}_n)) \rightarrow 0 \quad \text{a.s. as } n \rightarrow \infty.$$

By Lemmas 2 and 3, we have

$$\begin{aligned} d_{\text{BL}}^2(\hat{\mu}(\tilde{\mathcal{L}}_n), \hat{\mu}(\mathcal{L}_n)) &\leq \frac{1}{n} \text{tr}(\tilde{\mathcal{L}}_n - \mathcal{L}_n)^2 \\ &\leq \left| \frac{1}{n} \text{tr}(\tilde{\mathcal{L}}_n - \mathcal{L}_n)^2 - \frac{1}{n} \mathbb{E}[\text{tr}(\tilde{\mathcal{L}}_n - \mathcal{L}_n)^2] \right| + \frac{1}{n} \mathbb{E}[\text{tr}(\tilde{\mathcal{L}}_n - \mathcal{L}_n)^2]. \end{aligned}$$

Lemma 2 yields

$$\frac{1}{n} \mathbb{E}[\text{tr}(\tilde{\mathcal{L}}_n - \mathcal{L}_n)^2] = O(u_n^{-2}),$$

while Lemma 3 implies that the centered fluctuation term is $o(u_n^{-1})$ almost surely. Consequently,

$$d_{\text{BL}}^2(\hat{\mu}(\tilde{\mathcal{L}}_n), \hat{\mu}(\mathcal{L}_n)) = o(u_n^{-1}) \quad \text{a.s. as } n \rightarrow \infty. \quad (18)$$

(18) is also true under a linear scaling of $\{\gamma_n\}$ and $\{\rho_n\}$, we have

$$d_{\text{BL}}^2(\hat{\mu}(\gamma_n \mathcal{L}_n + \rho_n \mathbf{I}_n), \hat{\mu}(\gamma_n \tilde{\mathcal{L}}_n + \rho_n \mathbf{I}_n)) = o\left(\frac{\gamma_n^2}{u_n}\right) \quad \text{a.s. as } n \rightarrow \infty.$$

On the other hand, recall (13), we know that

$$\hat{\mu}(\gamma_n \tilde{\mathcal{L}}_n + \rho_n \mathbf{I}_n) \Rightarrow \mu_{\text{sc}} \quad \text{a.s. as } n \rightarrow \infty,$$

which is equivalent to

$$d_{\text{BL}}(\hat{\mu}(\gamma_n \tilde{\mathcal{L}}_n + \rho_n \mathbf{I}_n), \mu_{\text{sc}}) \rightarrow 0 \quad \text{a.s. as } n \rightarrow \infty.$$

Note that

$$\gamma_n = -\rho_n = -\sqrt{\frac{np_n}{1-p_n}}.$$

Under the assumption $np_n \geq C \log n$, we have $u_n \asymp np_n$ and hence

$$\frac{\gamma_n^2}{u_n} = O(1).$$

Therefore, by the triangle inequality for d_{BL} ,

$$\begin{aligned} d_{\text{BL}}(\hat{\mu}(\gamma_n \mathcal{L}_n + \rho_n \mathbf{I}_n), \mu_{\text{sc}}) &\leq d_{\text{BL}}(\hat{\mu}(\gamma_n \mathcal{L}_n + \rho_n \mathbf{I}_n), \hat{\mu}(\gamma_n \tilde{\mathcal{L}}_n + \rho_n \mathbf{I}_n)) \\ &\quad + d_{\text{BL}}(\hat{\mu}(\gamma_n \tilde{\mathcal{L}}_n + \rho_n \mathbf{I}_n), \mu_{\text{sc}}) \rightarrow 0 \quad \text{a.s. as } n \rightarrow \infty. \end{aligned}$$

This completes the proof of Theorem 2. \square

5. PROOF OF THEOREM 3

Proof. Define the normalized adjacency matrices

$$\mathbf{N}_n := \mathbf{D}_n^{-1/2} \mathbf{A}_n \mathbf{D}_n^{-1/2}, \quad \tilde{\mathbf{N}}_n := \mathbf{W}_n^{-1/2} \mathbf{A}_n \mathbf{W}_n^{-1/2}.$$

We proceed in two steps. First, we establish the semicircle law for the matrix $\tilde{\mathbf{N}}_n$. Second, we show that the same limiting law holds for \mathbf{N}_n by a perturbation analysis.

Recall that Chung, Lu, and Vu [20] proved that, under condition (4), the normalized matrix

$$\mathbf{C}_{\text{nor}} := \sqrt{\bar{w}_n} \mathbf{C}, \quad \mathbf{C} := \mathbf{W}_n^{-1/2} \mathbf{A}_n \mathbf{W}_n^{-1/2} - \phi_n W_n^{1/2} \mathbf{K} \mathbf{W}_n^{1/2},$$

has an empirical spectral distribution converging in probability to the semicircle distribution μ_{sc} .

Observe that

$$\tilde{\mathbf{N}}_n = \mathbf{W}_n^{-1/2} \mathbf{A}_n \mathbf{W}_n^{-1/2} = \mathbf{C} + \phi_n \mathbf{W}_n^{1/2} \mathbf{K} \mathbf{W}_n^{1/2} =: \mathbf{C} + \mathbf{R}_W,$$

where \mathbf{K} denotes the all-ones matrix. The matrix \mathbf{R}_W has rank one, since

$$\mathbf{W}_n^{1/2} \mathbf{K} \mathbf{W}_n^{1/2} = (\sqrt{w_{n,1}}, \dots, \sqrt{w_{n,n}})^\top (\sqrt{w_{n,1}}, \dots, \sqrt{w_{n,n}}).$$

Consequently, $\tilde{\mathbf{N}}_n$ differs from \mathbf{C} by a rank-one perturbation.

We now invoke the rank inequality due to Bai [5, Lemma 2.6].

Lemma 5 (Rank inequality). *Let \mathbf{A} and \mathbf{B} be two $n \times n$ Hermitian matrices. Then*

$$\|\hat{\mu}(\mathbf{A}) - \hat{\mu}(\mathbf{B})\|_\infty \leq \frac{1}{n} \text{rank}(\mathbf{A} - \mathbf{B}).$$

Applying Lemma 5 yields

$$\|\hat{\mu}(\tilde{\mathbf{N}}_n) - \hat{\mu}(\mathbf{C})\|_\infty \leq \frac{1}{n}, \tag{19}$$

which implies that $\tilde{\mathbf{N}}_n$ and \mathbf{C} have the same limiting spectral distribution.

Combining (19) observation with the semicircle law for \mathbf{C}_{nor} , we conclude that

$$\hat{\mu}(\sqrt{\bar{w}_n} \tilde{\mathbf{N}}_n) \implies \mu_{sc} \text{ in probability as } n \rightarrow \infty.$$

Lemma 6. *There exist absolute constants $c, c' > 0$ such that*

$$\frac{1}{n} \mathbb{E} \text{tr}((N_n - \tilde{N}_n)^2) \leq \frac{c}{\bar{w}_n w_{n,\min}} + c \bar{w}_n e^{-c' w_{n,\min}}.$$

In particular, since $w_{n,\min} \rightarrow \infty$, we have

$$\bar{w}_n \cdot \frac{1}{n} \mathbb{E} \text{tr}((N_n - \tilde{N}_n)^2) = O\left(\frac{1}{w_{n,\min}}\right) \rightarrow 0 \text{ in probability as } n \rightarrow \infty.$$

Proof. For $i \neq j$, the entries of \mathbf{N}_n and $\tilde{\mathbf{N}}_n$ are given by

$$(\mathbf{N}_n)_{ij} = \frac{a_{ij}^{(n)}}{\sqrt{d_{n,i} d_{n,j}}}, \quad (\tilde{\mathbf{N}}_n)_{ij} = \frac{a_{ij}^{(n)}}{\sqrt{w_{n,i} w_{n,j}}}.$$

Hence,

$$\tilde{\mathbf{N}}_n - \mathbf{N}_n = \left(\left(\frac{1}{\sqrt{w_{n,i} w_{n,j}}} - \frac{1}{\sqrt{d_{n,i} d_{n,j}}} \right) a_{ij}^{(n)} \right)_{1 \leq i,j \leq n}.$$

By symmetry and independence of the edge indicators,

$$\frac{1}{n} \mathbb{E} \text{tr}((\tilde{\mathbf{N}}_n - \mathbf{N}_n)^2) = \frac{1}{n} \sum_{i \neq j} \mathbb{E} \left[\left(\frac{1}{\sqrt{w_{n,i} w_{n,j}}} - \frac{1}{\sqrt{d_{n,i} d_{n,j}}} \right)^2 a_{ij}^{(n)} \right]. \tag{20}$$

Fix $i \neq j$. By conditioning on $a_{ij}^{(n)}$,

$$\mathbb{E} \left[\left(\frac{1}{\sqrt{d_{n,i}d_{n,j}}} - \frac{1}{\sqrt{w_{n,i}w_{n,j}}} \right)^2 a_{ij}^{(n)} \right] = p_{n,ij} \mathbb{E} \left[\left(\frac{1}{\sqrt{d_{n,i}d_{n,j}}} - \frac{1}{\sqrt{w_{n,i}w_{n,j}}} \right)^2 \mid a_{ij}^{(n)} = 1 \right].$$

Define the event

$$E_3 := \left\{ d_{n,i} \in [\frac{1}{2}w_{n,i}, \frac{3}{2}w_{n,i}], d_{n,j} \in [\frac{1}{2}w_{n,j}, \frac{3}{2}w_{n,j}] \right\}.$$

On E_3 , a Taylor expansion argument yields

$$\mathbb{E} \left[\left(\frac{1}{\sqrt{d_{n,i}d_{n,j}}} - \frac{1}{\sqrt{w_{n,i}w_{n,j}}} \right)^2 \mathbf{1}_{E_3} \mid a_{ij}^{(n)} = 1 \right] \leq c \left(\frac{1}{w_{n,i}^2 w_{n,j}} + \frac{1}{w_{n,j}^2 w_{n,i}} \right).$$

Since $d_{n,i}$ and $d_{n,j}$ are sums of independent Bernoulli random variables, by a Chernoff bound, we have

$$\mathbb{P}(E_3^c) \leq 2e^{-w_{n,i}/12} + 2e^{-w_{n,j}/12} \leq 4e^{-w_{n,\min}/12}.$$

On E_3^c , we use the crude bound and the pseudo-inverse convention to have

$$\left| \frac{1}{\sqrt{w_{n,i}w_{n,j}}} - \frac{1}{\sqrt{d_{n,i}d_{n,j}}} \right| \leq \frac{1}{\sqrt{w_{n,i}w_{n,j}}} + \frac{1}{\sqrt{d_{n,i}d_{n,j}}} \leq 2,$$

which implies

$$\mathbb{E} \left[\left(\frac{1}{\sqrt{d_{n,i}d_{n,j}}} - \frac{1}{\sqrt{w_{n,i}w_{n,j}}} \right)^2 \mathbf{1}_{E_3^c} \mid a_{ij}^{(n)} = 1 \right] \leq 16e^{-w_{n,\min}/12}.$$

Combining the two cases, we obtain

$$\mathbb{E} \left[\left(\frac{1}{\sqrt{d_{n,i}d_{n,j}}} - \frac{1}{\sqrt{w_{n,i}w_{n,j}}} \right)^2 \mid a_{ij}^{(n)} = 1 \right] \leq c \left(\frac{1}{w_{n,i}^2 w_{n,j}} + \frac{1}{w_{n,j}^2 w_{n,i}} \right) + 16e^{-w_{n,\min}/12}.$$

Multiplying by $p_{n,ij} = \phi_n w_{n,i} w_{n,j}$ gives

$$\mathbb{E} \left[\left(\frac{1}{\sqrt{d_{n,i}d_{n,j}}} - \frac{1}{\sqrt{w_{n,i}w_{n,j}}} \right)^2 a_{ij}^{(n)} \right] \leq c\phi_n \left(\frac{1}{w_{n,i}} + \frac{1}{w_{n,j}} \right) + 16p_{n,ij}e^{-w_{n,\min}/12}.$$

Summing over $i \neq j$ and using $\sum_{i,j} p_{n,ij} = n\bar{w}_n = 1/\phi_n$, we conclude from (20) that

$$\frac{1}{n} \mathbb{E} \text{tr} \left((\tilde{\mathbf{N}}_n - \mathbf{N}_n)^2 \right) \leq \frac{c}{\bar{w}_n w_{n,\min}} + c\bar{w}_n e^{-w_{n,\min}/12},$$

which completes the proof. \square

Recall that, for any sequences $\gamma_n \in \mathbb{R}$, we have

$$d_{BL}^2 \left(\hat{\mu}(\gamma_n \mathbf{N}_n), \hat{\mu}(\gamma_n \tilde{\mathbf{N}}_n) \right) \leq \gamma_n^2 \cdot \frac{1}{n} \text{tr} \left((\mathbf{N}_n - \tilde{\mathbf{N}}_n)^2 \right),$$

Taking $\gamma_n = \sqrt{\bar{w}_n}$, and using the expectation value and Lemma 6, we get:

$$\mathbb{E} \left[d_{BL}^2 \left(\hat{\mu}(\sqrt{\bar{w}_n} \mathbf{N}_n), \hat{\mu}(\sqrt{\bar{w}_n} \tilde{\mathbf{N}}_n) \right) \right] \leq \bar{w}_n \cdot \frac{1}{n} \mathbb{E} \text{tr} \left((\mathbf{N}_n - \tilde{\mathbf{N}}_n)^2 \right) = O \left(\frac{1}{w_{n,\min}} \right) \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

where we use the assumption $w_{n,\min} \gg \sqrt{\bar{w}_n}$. Thus, by Markov's inequality, we obtain

$$d_{BL} \left(\hat{\mu}(\sqrt{\bar{w}_n} \mathbf{N}_n), \hat{\mu}(\sqrt{\bar{w}_n} \tilde{\mathbf{N}}_n) \right) \rightarrow 0 \quad \text{in probability as } n \rightarrow \infty.$$

Thus, by (5) and the triangle inequality,

$$\begin{aligned}
 d_{BL}(\hat{\mu}(\sqrt{\bar{w}_n}\mathbf{N}_n), \mu_{sc}) &\leq d_{BL}\left(\hat{\mu}(\sqrt{\bar{w}_n}\mathbf{N}_n), \hat{\mu}(\sqrt{\bar{w}_n}\tilde{\mathbf{N}}_n)\right) \\
 &\quad + d_{BL}\left(\hat{\mu}(\sqrt{\bar{w}_n}\tilde{\mathbf{N}}_n), \mu_{sc}\right) \\
 &\longrightarrow 0 \quad \text{in probability as } n \rightarrow \infty.
 \end{aligned} \tag{21}$$

And (21) proves

$$\hat{\mu}(\sqrt{\bar{w}_n}\mathbf{D}_n^{-1/2}\mathbf{A}_n\mathbf{D}_n^{-1/2}) \Rightarrow \mu_{sc} \quad \text{in probability as } n \rightarrow \infty$$

as desired. \square

Acknowledgments. Y.Z. was partially supported by the Simons Grant MPS-TSM-00013944.

REFERENCES

- [1] Ben Adlam and Ziliang Che. Spectral statistics of sparse random graphs with a general degree distribution. *arXiv preprint arXiv:1509.03368*, 2015.
- [2] Pawat Akara-pipattana and Oleg Evnin. Random matrices with row constraints and eigenvalue distributions of graph laplacians. *Journal of Physics A: Mathematical and Theoretical*, 56(29):295001, 2023.
- [3] Johannes Alt, Raphaël Ducatez, and Antti Knowles. Extremal eigenvalues of critical Erdős–Rényi graphs. *The Annals of Probability*, 49(3):1347–1401, 2021.
- [4] Z. D. Bai. Methodologies in spectral analysis of large dimensional random matrices, a review. *Statistica Sinica*, 9:611–677, 1999.
- [5] Zhidong Bai and Jack W Silverstein. *Spectral analysis of large dimensional random matrices*. Springer, 2010.
- [6] Florent Benaych-Georges, Charles Bordenave, and Antti Knowles. Largest eigenvalues of sparse inhomogeneous erdős–rényi graphs. *The Annals of Probability*, 47(3):1653–1676, 2019.
- [7] Béla Bollobás. Random graphs. In *Modern graph theory*, pages 215–252. Springer, 2011.
- [8] Charles Bordenave, Pietro Caputo, and Djalil Chafaï. Spectrum of large random reversible markov chains: two examples. *ALEA: Latin American Journal of Probability and Mathematical Statistics*, 7:41–64, 2010.
- [9] Charles Bordenave, Pietro Caputo, and Djalil Chafaï. Circular law theorem for random markov matrices. *Probability Theory and Related Fields*, 152(3):751–779, 2012.
- [10] Charles Bordenave, Pietro Caputo, and Djalil Chafaï. Spectrum of markov generators on sparse random graphs. *Communications on Pure and Applied Mathematics*, 67(4):621–669, 2014.
- [11] Charles Bordenave, Pietro Caputo, and Djalil Chafaï. Spectrum of large random reversible markov chains: heavy-tailed weights on the complete graph. *The Annals of probability*, 39(4):1544–1590, 2011.
- [12] Andrew Campbell, Kyle Luh, Sean O’rourke, Santiago Arenas-Velilla, and Victor Pérez-Abreu. Extreme eigenvalues of laplacian random matrices with gaussian entries. *Electronic Journal of Probability*, 30:1–52, 2025.
- [13] Andrew Campbell and Sean O’Rourke. Spectrum of lévy–khintchine random laplacian matrices. *Journal of Theoretical Probability*, 37(1):933–973, 2024.
- [14] Arijit Chakrabarty, Rajat Subhra Hazra, Frank Den Hollander, and Matteo Sfragara. Spectra of adjacency and laplacian matrices of inhomogeneous erdős–rényi random graphs. *Random matrices: Theory and applications*, 10(01):2150009, 2021.
- [15] Anirban Chatterjee and Rajat Subhra Hazra. Spectral properties for the laplacian of a generalized wigner matrix. *Random Matrices: Theory and Applications*, 11(03):2250026, 2022.
- [16] Yiming Chen, Xuanang Hu, and Pengtao Li. On concentration inequality of the laplacian matrix of erdos–rényi graphs. *arXiv preprint arXiv:2502.02248*, 2025.
- [17] Zhiyi Chi. Random reversible markov matrices with tunable extremal eigenvalues. *Annals of applied probability*, 26(4):2257–2272, 2016.
- [18] Nicholas Christoffersen, Kyle Luh, Hoi H Nguyen, and Jingheng Wang. Eigenvalue gaps of the laplacian of random graphs. *arXiv preprint arXiv:2501.00234*, 2024.
- [19] Fan Chung and Linyuan Lu. The average distances in random graphs with given expected degrees. *Proceedings of the National Academy of Sciences*, 99(25):15879–15882, 2002.
- [20] Fan Chung, Linyuan Lu, and Van Vu. Spectra of random graphs with given expected degrees. *Proceedings of the National Academy of Sciences*, 100(11):6313–6318, 2003.
- [21] Fan RK Chung. *Spectral graph theory*, volume 92. American Mathematical Soc., 1997.
- [22] Fan RK Chung and Linyuan Lu. *Complex graphs and networks*. Number 107 in CBMS Regional Conference Series in Mathematics. American Mathematical Soc., 2006.

- [23] Lorenzo Dall'Amico, Romain Couillet, and Nicolas Tremblay. A unified framework for spectral clustering in sparse graphs. *Journal of Machine Learning Research*, 22(217):1–56, 2021.
- [24] Shaofeng Deng, Shuyang Ling, and Thomas Strohmer. Strong consistency, graph laplacians, and the stochastic block model. *Journal of Machine Learning Research*, 22(117):1–44, 2021.
- [25] Xue Ding and Tiefeng Jiang. Spectral distributions of adjacency and laplacian matrices of random graphs. *The annals of applied probability*, pages 2086–2117, 2010.
- [26] Raphael Ducatez and Renaud Rivier. Spectral gap and embedded trees for the laplacian of the erdős-rényi graph. *Electronic Journal of Probability*, 30:1–21, 2025.
- [27] László Erdos, Antti Knowles, Horng-Tzer Yau, and Jun Yin. Spectral statistics of erdős-rényi graphs i: Local semicircle law. *Annals of probability*, 41(3):2279–2375, 2013.
- [28] Christopher Hoffman, Matthew Kahle, and Elliot Paquette. Spectral gaps of random graphs and applications. *International Mathematics Research Notices*, 2021(11):8353–8404, 2021.
- [29] Shlomo Hoory, Nathan Linial, and Avi Wigderson. Expander graphs and their applications. *Bulletin of the American Mathematical Society*, 43(4):439–561, 2006.
- [30] Jiaoyang Huang and Benjamin Landon. Spectral statistics of sparse erdos-rényi graph laplacians. *Annales de l'Institut Henri Poincaré-Probabilités et Statistiques*, 56(1):120–154, 2020.
- [31] Tiefeng Jiang. Empirical distributions of laplacian matrices of large dilute random graphs. *Random Matrices: Theory and Applications*, 1(03):1250004, 2012.
- [32] Can M Le, Elizaveta Levina, and Roman Vershynin. Concentration and regularization of random graphs. *Random Structures & Algorithms*, 51(3):538–561, 2017.
- [33] Can M Le, Elizaveta Levina, and Roman Vershynin. Concentration of random graphs and application to community detection. In *Proceedings of the International Congress of Mathematicians: Rio de Janeiro 2018*, pages 2925–2943. World Scientific, 2018.
- [34] Linyuan Lu and Xing Peng. Spectra of edge-independent random graphs. *Electronic Journal of Combinatorics*, 20(4):Paper 27, 2013.
- [35] Russell Lyons and Yuval Peres. *Probability on trees and networks*, volume 42. Cambridge University Press, 2017.
- [36] Marina Meilă and Jianbo Shi. A random walks view of spectral segmentation. In *International workshop on artificial intelligence and statistics*, pages 203–208. PMLR, 2001.
- [37] Alaa Saade, Florent Krzakala, and Lenka Zdeborová. Spectral clustering of graphs with the bethe hessian. *Advances in neural information processing systems*, 27, 2014.
- [38] Ludovic Stephan and Yizhe Zhu. Community detection with the bethe-hessian. *arXiv preprint arXiv:2411.02835*, 2024.
- [39] Rajat Subhra Hazra, Frank den Hollander, and Maarten Markering. Large deviation principle for the norm of the laplacian matrix of inhomogeneous erdős-rényi random graphs. *Electronic Journal of Probability*, 30:1–22, 2025.
- [40] Linh V Tran, Van H Vu, and Ke Wang. Sparse random graphs: Eigenvalues and eigenvectors. *Random Structures & Algorithms*, 42(1):110–134, 2013.
- [41] Charles F Van Loan and G Golub. Matrix computations (johns hopkins studies in mathematical sciences). *Matrix Computations*, 5:32, 1996.
- [42] Lutz Warnke. On the method of typical bounded differences. *Combinatorics, Probability and Computing*, 25(2):269–299, 2016.
- [43] Yizhe Zhu. A graphon approach to limiting spectral distributions of wigner-type matrices. *Random Structures & Algorithms*, 56(1):251–279, 2020.

SCHOOL OF MATHEMATICAL SCIENCES, PEKING UNIVERSITY
Email address: ymchenmath@math.pku.edu.cn

SCHOOL OF MATHEMATICS, SHANDONG UNIVERSITY
Email address: czj4096@gmail.com

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA
Email address: yizhezhu@usc.edu