

Random Gradient-Free Optimization in Infinite Dimensional Spaces*

Caio Lins Peixoto[†]

Daniel Csillag[†]

Bernardo F. P. da Costa[†]

Yuri F. Saporito[†]

December 2025

Abstract

In this paper, we propose a random gradient-free method for optimization in infinite dimensional Hilbert spaces, applicable to functional optimization in diverse settings. Though such problems are often solved through finite-dimensional gradient descent over a parametrization of the functions, such as neural networks, an interesting alternative is to instead perform gradient descent directly in the function space by leveraging its Hilbert space structure, thus enabling provable guarantees and fast convergence. However, infinite-dimensional gradients are often hard to compute in practice, hindering the applicability of such methods. To overcome this limitation, our framework requires only the computation of directional derivatives and a pre-basis for the Hilbert space domain, i.e., a linearly-independent set whose span is dense in the Hilbert space. This fully resolves the tractability issue, as pre-bases are much more easily obtained than full orthonormal bases or reproducing kernels — which may not even exist — and individual directional derivatives can be easily computed using forward-mode scalar automatic differentiation. We showcase the use of our method to solve partial differential equations *à la* physics informed neural networks (PINNs), where it effectively enables provable convergence.

1 Introduction

Core to many applications is the solution of functional optimization problems; formally speaking, these are of the form

$$\arg \min_{h \in \mathcal{H}} \mathcal{R}(h), \quad (1)$$

where \mathcal{H} is a separable Hilbert space (e.g., L^2 space, or a Sobolev space), and $\mathcal{R} : \mathcal{H} \rightarrow \mathbb{R}$ is a Fréchet differentiable risk functional, often convex. Prominent examples include machine learning tasks such as supervised learning and generative modeling, optimization-based PDE solvers and broader inverse problems.

To solve problems like (1), practitioners typically introduce a finite-dimensional parametrization $\theta \mapsto h_\theta$ of the functions $h \in \mathcal{H}$ and then optimize over $\theta \in \mathbb{R}^D$ using some variant of gradient descent. This approach, however, is limited by the representation capacity of the chosen parametrization; moreover, because the parametrization is typically nonlinear (e.g., neural networks), even if \mathcal{R} is convex, the composition $\theta \mapsto \mathcal{R}(h_\theta)$ is most likely not, increasing the complexity of the resulting optimization problem. To overcome these issues, we perform gradient-based optimization *directly in the function space* \mathcal{H} , since its Hilbert space structure gives the existence of gradients as Riesz representers

*CLP was partially supported by CAPES through the CAPES/PICME master's scholarship and FGV EMAp during his work in this manuscript. YFS was supported by FAPERJ (Brasil) through the Jovem Cientista do Nosso Estado Program (E- 26/201.375/2022 (272760)) and by CNPq (Brasil) through the Productivity in Research Scholarship (306695/2021- 9).

[†]School of Applied Mathematics, Getulio Vargas Foundation, Rio de Janeiro, Brazil, caio.peixoto@fgv.br , daniel.csillag@fgv.br , bernardo.paulo@fgv.br , yuri.saporito@fgv.br .

of Fréchet derivatives. By doing so, we may retain convexity of the risk and also avoid the limited representation power of any specific parametrization.

However, gradient-based optimization in infinite-dimensional Hilbert spaces is challenging. For one, functional gradients themselves are often hard to compute exactly, being defined as solutions to variational problems involving the directional derivatives of the risk. Secondly, it is not generally clear how to represent infinite dimensional objects computationally. Thirdly, in the realm of stochastic gradient methods, controlling the variance of an infinite dimensional random element usually requires more care than that of a finite dimensional random vector.

In this paper, we adopt the functional optimization perspective, but we bypass exact gradient computation by taking inspiration from random gradient-free optimization methods. This is a class of algorithms that build stochastic gradient estimators based only on directional derivative computations, which are easily available through forward mode automatic differentiation. We adapt this framework to be applicable in infinite dimensions, yielding a general algorithm for minimizing Fréchet differentiable risk functionals over separable Hilbert spaces, with convergence guarantees in the convex case.

1.1 A motivating example: solving PDEs through gradient descent

Consider solving the following PDE with Dirichlet boundary conditions:

$$\begin{cases} L[u](\mathbf{x}) = 0 & \mathbf{x} \in \Omega, \\ u(\mathbf{x}) = f(\mathbf{x}) & \mathbf{x} \in \partial\Omega, \end{cases} \quad (2)$$

where $\Omega \subset \mathbb{R}^d$ is a bounded, open set with a smooth boundary and L is an ℓ -th order differential operator. For simplicity, in this section we will assume that L is a linear operator with bounded coefficients. Recently, a fast growing body of research [2, 15, 14, 19] has been dedicated to study deep learning algorithms that find approximate solutions to (2). These algorithms work by minimizing a functional of the form

$$\mathcal{R}(h) = \frac{1}{2} \|L[h]\|_{L^2(\Omega)}^2 + \frac{1}{2} \|h - f\|_{L^2(\partial\Omega)}^2, \quad (3)$$

where $h = h_\theta$ is taken to be a neural network with fixed architecture, parameterized by $\theta \in \mathbb{R}^D$ for some large $D \in \mathbb{N}$. The idea is to minimize $\mathcal{R}(h_\theta)$ over θ through variants of gradient descent in the parameter space.

We propose to see (3) as acting directly on functions, as opposed to a loss over finite-dimensional parameters. Our main motivation for doing so is preserving the convexity of the risk when L is linear.¹ This is also aligned with a recent surge in the *first optimize, then discretize* paradigm in the scientific ML literature, cf. [10, 11], which tries to emulate functional optimization algorithms when optimizing a parametric ansatz h_θ to solve (2). The functional perspective has also influenced recent works in statistical inverse problems [6] and nonparametric instrumental variable regression [5].

To proceed, we must first specify the domain \mathcal{H} of the risk \mathcal{R} . Due to the nature of problem (2), we take $\mathcal{H} = \mathbb{H}^\ell(\Omega)$, the Sobolev space of order ℓ and exponent 2 in Ω — that is, the space of all $h \in L^2(\Omega)$ whose weak derivatives exist up to order ℓ and lie in $L^2(\Omega)$, which we endow with the usual inner product $\langle h, h' \rangle_{\mathbb{H}^\ell(\Omega)} = \sum_j \langle D^{(j)}h, D^{(j)}h' \rangle_{L^2(\Omega)}$.² Thus, solving the PDE (2) becomes the optimization problem

$$\arg \min_{h \in \mathbb{H}^\ell(\Omega)} \mathcal{R}(h), \quad (4)$$

where \mathcal{R} is given by (3).

To solve this problem under the functional paradigm, the most direct approach would be to employ gradient descent in $\mathbb{H}^\ell(\Omega)$. By definition, if \mathcal{R} is Fréchet differentiable, the gradient $\nabla \mathcal{R}(h)$ is the unique element of $\mathbb{H}^\ell(\Omega)$ satisfying $\langle \nabla \mathcal{R}(h), v \rangle_{\mathbb{H}^\ell(\Omega)} = D\mathcal{R}(h; v)$ for all $v \in \mathbb{H}^\ell(\Omega)$, where

$$D\mathcal{R}(h; v) = \lim_{\delta \rightarrow 0} \frac{\mathcal{R}(h + \delta v) - \mathcal{R}(h)}{\delta} \quad (5)$$

¹If L is nonlinear we may lose convexity, but still obtain a problem that is simpler than what a non-linear parametrization would result in.

²We denote the j -th derivative of h by $D^{(j)}h$, which we interpret as a d^j -dimensional array of real numbers.

is the directional derivative of \mathcal{R} at h in the direction v . Thus, to obtain $\nabla\mathcal{R}(h)$ we first need to compute $D\mathcal{R}(h; v)$, which is straightforward due to the linearity of L :

$$D\mathcal{R}(h; v) = \int_{\Omega} L[h](\mathbf{x})L[v](\mathbf{x}) \, d\mathbf{x} + \int_{\partial\Omega} (h(\mathbf{x}) - f(\mathbf{x}))v(\mathbf{x}) \sigma(d\mathbf{x}),$$

where σ is the surface measure on $\partial\Omega$.

We can now see that the functional $D\mathcal{R}(h; \cdot)$ is linear and continuous in the $\mathbb{H}^\ell(\Omega)$ -norm. Furthermore, the map $h \mapsto D\mathcal{R}(h; \cdot)$, from $\mathbb{H}^\ell(\Omega)$ to its dual, is also continuous, implying that \mathcal{R} is Frechét differentiable [13]. Therefore, the gradient $\nabla\mathcal{R}(h)$ is well defined, and finding it is equivalent to solving the following problem:

Find $g \in \mathbb{H}^\ell(\Omega)$ such that

$$\sum_{j=0}^{\ell} \int_{\Omega} D^{(j)}g(\mathbf{x}) \cdot D^{(j)}v(\mathbf{x}) \, d\mathbf{x} = \int_{\Omega} L[h](\mathbf{x})L[v](\mathbf{x}) \, d\mathbf{x} + \int_{\partial\Omega} (h(\mathbf{x}) - f(\mathbf{x}))v(\mathbf{x}) \sigma(d\mathbf{x}), \quad (6)$$

for all $v \in \mathbb{H}^\ell(\Omega)$.

This is a variational problem for which no general closed form solution exists, creating the need to look beyond exact gradients in order to solve (4).

1.2 Background: random gradient-free methods

To avoid the computation of exact gradients, an existing branch of the literature constructs “random gradient-free methods,” which leverage stochastic gradient estimators based on computations of directional derivatives. Hence, though these methods are not derivative-free in a strict sense, they are gradient-free. So far, this approach has only been considered for *finite dimensional* optimization problems. In infinite dimensions, existing methods exhibit prohibitive limitations.

If $\mathcal{H} = \mathbb{R}^d$ for some $d \in \mathbb{N}$, then random gradient-free methods estimate the gradient $g \triangleq \nabla\mathcal{R}(h)$ as

$$\hat{g} = \frac{1}{M} \sum_{m=1}^M D\mathcal{R}(h; v_m) v_m, \quad (7)$$

where v_1, \dots, v_M are i.i.d. vectors in \mathbb{R}^d with zero mean and identity covariance matrix, cf. [12]. This is thus an unbiased estimator of g that, under additional assumptions on v , has finite second moment: $\mathbb{E}[\|\hat{g}\|^2] < +\infty$. For example, in the convenient case where $v_i \sim \mathcal{N}(0, \mathbf{I}_d)$, one may show that $\mathbb{E}[\|\hat{g}\|^2] = \|g\|^2 \left(\frac{M+d+1}{M}\right)$. This setup already exhibits a couple of issues that may arise when \mathcal{H} is high/infinite dimensional:

- (i) The expression for $\mathbb{E}[\|\hat{g}\|^2]$ presented above suggests taking the sample size M to be proportional to d . As $d \rightarrow +\infty$, this becomes infeasible.
- (ii) Many of the usual choices for the distribution of the v_i 's have no infinite-dimensional analogue: there is no standard Gaussian distribution [3], no uniform measure over the sphere [24], and no Rademacher distribution (since sequences of ± 1 's are not square-summable). These are distributions used, for example, in [8, 12, 20] for the finite dimensional scenario.
- (iii) More broadly, if a random element v in an infinite dimensional Hilbert space satisfies $\mathbb{E}[\|v\|^2] < +\infty$, then it necessarily has a compact covariance operator, which thus cannot be the identity — a necessary condition for the unbiasedness of \hat{g} in (7). In fact, since eigenvalues of compact operators concentrate around zero [18, Theorem 4.25], we have $\|T - I\| \geq 1$ for any compact operator $T : \mathcal{H} \rightarrow \mathcal{H}$.
- (iv) Generally speaking, it is difficult to design unbiased gradient estimators \hat{g} whose second moment is guaranteed to be finite for any gradient g .

In this paper, we resolve these issues by proposing a novel random gradient-free algorithm that is suitable for infinite-dimensional optimization. By first sampling a finite dimension K and then sampling random directions v_i from a specific K -dimensional subspace of \mathcal{H} , we are able to overcome issues (ii) and (iii); then, by choosing the sample size M proportional to the random dimension, we bypass issue (i). Finally, by introducing an appropriate preconditioning of the gradient descent process we are able to control the variance of our gradient estimates, thus resolving issue (iv).

Notation We denote the set of positive integers by \mathbb{N} . Given a real, separable Hilbert space $(\mathcal{H}, \langle \cdot, \cdot \rangle)$ and a finite dimension d , the elements of \mathcal{H} are denoted by regular font Latin letters f, g, h, \dots , whereas the elements of \mathbb{R}^d are denoted by boldface Latin letters $\mathbf{x}, \mathbf{y}, \mathbf{z}, \dots$. Regular font Greek letters will always denote scalars. For $h, h' \in \mathcal{H}$, we write $h \otimes h'$ to denote the rank one map $(h \otimes h')v = \langle h', v \rangle h$. If $\mathcal{E} = \{e_1, e_2, \dots\}$ is an orthonormal basis for \mathcal{H} , $\{\gamma_i\}_{i \in \mathbb{N}}$ is a positive, bounded sequence and³ $C = \sum_{i=1}^{\infty} \gamma_i (e_i \otimes e_i)$ is a diagonal operator, the powers C^s , for $s \in \mathbb{R}$, are defined through the usual functional calculus: $C^s = \sum_{i=1}^{\infty} (\gamma_i)^s (e_i \otimes e_i)$. These are possibly unbounded operators with restricted domains.

2 Hilbert Space Gradient-Free Descent

We assume to have access to a *pre-basis* $\mathcal{B} = \{b_1, b_2, \dots\} \subset \mathcal{H}$, i.e., a linearly independent subset of \mathcal{H} whose span is dense in \mathcal{H} . With \mathcal{B} in hand, we define the finite-dimensional subspaces $\mathcal{B}_k = \text{span}\{b_1, \dots, b_k\}$, for $k \in \mathbb{N}$. As a theoretical tool, it will also be useful to refer to the orthonormalization $\mathcal{E} = \{e_1, e_2, \dots\}$ of the pre-basis, defined through Gram-Schmidt. Since \mathcal{B} has dense span, \mathcal{E} is an actual orthonormal basis of \mathcal{H} . We emphasize that we do not assume an orthonormal basis to be available in closed form, rendering our approach applicable to spaces beyond the simple cases where such a basis is known.⁴

Fix a probability space $(\mathcal{S}, \mathcal{F}, \mathbb{P})$. At the n -th optimization step, our procedure starts by sampling a random finite dimension K_n from some distribution \mathbb{P}_K supported on an infinite subset of \mathbb{N} . Given $K_n = k$, it then performs the updates

$$h_{n+1} = h_n - \alpha_n \hat{g}_n(k) \quad \text{with} \quad \hat{g}_n(k) = \frac{\lambda_k}{M_k} \sum_{m=1}^{M_k} D\mathcal{R}(h_n; v_m) v_m, \quad (8)$$

where v_1, \dots, v_{M_k} are i.i.d. random directions sampled from a distribution $\mathbb{P}_{v|k}$ supported in $\mathcal{B}_k \subset \mathcal{H}$, $M_k \in \mathbb{N}$ denotes the sample-size for the stochastic gradient approximation, and $\lambda_k \in \mathbb{R}_+$ corresponds to a preconditioner on the gradient. We require $\{\lambda_i\}_{i \in \mathbb{N}}$ to be positive and bounded.

We seek to design a distribution the for v_i 's such that $\hat{g}_n(K_n)$ is an unbiased preconditioned gradient, i.e., letting $g_n \triangleq \nabla\mathcal{R}(h_n)$, we want the expectation of $\hat{g}_n(K_n)$ over the joint distribution of K_n and the v_i 's, conditional on h_1, \dots, h_n , to be Cg_n for a positive-definite operator $C : \mathcal{H} \rightarrow \mathcal{H}$, while ensuring finite second moment in an appropriate sense. This is straightforward in finite dimension, but challenging in infinite dimensional spaces. To this end, we construct the conditional distributions $\mathbb{P}_{v|k}$ so that their marginal \mathbb{P}_v over K_n has identity covariance; this guarantees preconditioned unbiasedness, and with additional conditions (to be derived in Theorem 2 and corollaries) will also ensure finite second moment.

To attain this, it will be useful to refer to the QR decomposition of quasi-matrices. Let B_k be the quasi-matrix $[b_1, \dots, b_k]$, i.e., the operator $B_k : \mathbb{R}^k \rightarrow \mathcal{H}$ given by

$$B_k = \sum_{i=1}^k b_i \otimes \mathbf{e}_i,$$

³Here and in similar situations where there is a series of operators, we define the series as the limit of the partial sums in the *strong operator topology*, not the uniform operator topology.

⁴For example, in our motivating application of optimization over Sobolev spaces, there are no explicitly-known orthonormal bases for general domains Ω and differentiability orders ℓ .

where $\{\mathbf{e}_1, \dots, \mathbf{e}_k\}$ is the canonical basis of \mathbb{R}^k . The QR-decomposition can be naturally extended to quasi-matrices, resulting in $B_k = Q_k \mathbf{R}_k$, where $Q_k = [e_1, \dots, e_k]$ is an orthogonal quasi-matrix and $\mathbf{R}_k \in \mathbb{R}^{k \times k}$ is an upper-triangular matrix [21].

Let $K \sim \mathbb{P}_K$ and define the tail probabilities $t_i = \mathbb{P}[K \geq i]$, as well as the matrix $\mathbf{T}_k = \text{diag}(t_1, \dots, t_k)$. Since \mathbb{P}_K has infinite support, we have $t_i > 0$ for all $i \in \mathbb{N}$. We then take $\mathbb{P}_{v|k}$ to be the distribution of

$$v = B_k \mathbf{R}_k^{-1} \mathbf{T}_k^{-\frac{1}{2}} \mathbf{z}, \quad \text{where } \mathbf{z} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_k). \quad (9)$$

With this choice, it follows:

Proposition 1. *The distributions $\mathbb{P}_{v|k}$ and \mathbb{P}_v both have zero mean and satisfy*

$$\text{Cov}[\mathbb{P}_{v|k}] = Q_k \mathbf{T}_k^{-1} Q_k^*, \quad \text{Cov}[\mathbb{P}_v] = I. \quad (10)$$

Consequently, $\mathbb{E}_n[\widehat{g}_n(K_n)] = C \nabla \mathcal{R}(h_n)$, where \mathbb{E}_n denotes the conditional expectation given h_1, \dots, h_n , and $C : \mathcal{H} \rightarrow \mathcal{H}$ is the diagonal operator

$$C = \sum_{i=1}^{\infty} \gamma_i (e_i \otimes e_i), \quad (11)$$

with $\gamma_i \triangleq \mathbb{E}[\lambda_K \mid K \geq i]$.

Proof. See Appendix A.1. □

Remark 1. *Distributions of v for which $\mathbb{E}[\|v\|^2] < +\infty$ are said to be of strong second order, and must have a compact covariance operator, cf. issue (iii) in Section 1. Since \mathcal{H} is infinite dimensional, the identity operator is not compact, implying that our proposed distribution \mathbb{P}_v is not of strong second order. It is, however, of weak second order, meaning that it satisfies $\mathbb{E}[\langle v, h \rangle^2] < +\infty$, for any $h \in \mathcal{H}$. Such distributions still have covariance operators, which do not need to be compact, cf. [1, 22].*

Leveraging this, we obtain Algorithm 1, which solves (1) by employing update (8) with the distribution $\mathbb{P}_{v|k}$ specified in (9). The analysis of our algorithm is inspired by that of preconditioned gradient and variable metric methods, see [4, Chapter 4] and the references therein. In this direction, a central object for the upcoming results is the (diagonal) inverse square-root operator $C^{-\frac{1}{2}} : C^{\frac{1}{2}}(\mathcal{H}) \rightarrow \mathcal{H}$. This operator is well defined because both C and $C^{\frac{1}{2}}$ are positive definite operators: we require $\lambda_i > 0$ for all i , implying that the diagonal of C is strictly positive. With $C^{-\frac{1}{2}}$ we can define the inner product $\langle h, h' \rangle_C \triangleq \langle C^{-\frac{1}{2}}h, C^{-\frac{1}{2}}h' \rangle$, for $h, h' \in C^{\frac{1}{2}}(\mathcal{H})$, as well as the norm $\|h\|_C^2 = \langle h, h \rangle_C$, which controls the behavior of our convergence bounds. It is straightforward to check that $C^{\frac{1}{2}}(\mathcal{H})$ contains the orthonormal basis \mathcal{E} , making $C^{-\frac{1}{2}}$ a densely-defined linear operator in \mathcal{H} .

Remark 2. *We note that C is the covariance operator of $\sqrt{\lambda_K}v$, for v given by (9). Though the distribution of $\sqrt{\lambda_K}v$ is not Gaussian, in the context of Gaussian measures the subspace $C^{\frac{1}{2}}(\mathcal{H})$ is called the Cameron-Martin space of the Gaussian with zero mean and covariance C , cf. [22].*

We can now state our first convergence bound. To simplify the notation we will write \widehat{g}_n instead of $\widehat{g}_n(K_n)$, leaving the dependence of \widehat{g}_n on K_n implicit.

Theorem 1. *Assume that \mathcal{R} is convex, and let h_1, \dots, h_N be the output sequence of Algorithm 1. Define the averaged estimate $\widehat{h} = (\sum_{n=1}^N \alpha_n h_n) / \sum_{n=1}^N \alpha_n$. Then, if $\mathbb{E}[\|\widehat{g}_n\|_C^2] < +\infty$ for all n , taking any $h \in C^{\frac{1}{2}}(\mathcal{H})$, we have:*

$$\mathbb{E}[\mathcal{R}(\widehat{h})] - \mathcal{R}(h) \leq \frac{\frac{1}{2}\|h - h_1\|_C^2}{\sum_{n=1}^N \alpha_n} + \frac{\frac{1}{2}\sum_{n=1}^N \alpha_n^2 \mathbb{E}[\|\widehat{g}_n\|_C^2]}{\sum_{n=1}^N \alpha_n}. \quad (12)$$

Proof. See Appendix A.2. □

Algorithm 1: Hilbert Space Gradient-Free Descent

Parameters: Distribution \mathbb{P}_K over \mathbb{N} , sequence of sample sizes $\{M_k\}_{k \in \mathbb{N}} \subset \mathbb{N}$, sequence of preconditioning parameters $\{\lambda_k\}_{k \in \mathbb{N}} \subset \mathbb{R}_+$, sequence of learning rates $\{\alpha_n\}_{n \in \mathbb{N}} \subset \mathbb{R}_+$, number of iterations $N \in \mathbb{N}$, starting point $h_1 \in C^{\frac{1}{2}}(\mathcal{H})$.

Input: Directional derivative $D\mathcal{R} : \mathcal{H} \times \mathcal{H} \rightarrow \mathbb{R}$, pre-basis $\mathcal{B} = \{b_1, b_2, \dots\}$, inner product $\langle \cdot, \cdot \rangle$.

Output: $h_1, \dots, h_{N+1} \in \mathcal{H}$.

for $1 \leq n \leq N$ **do**

$k \leftarrow$ sample from \mathbb{P}_K

$B_k \leftarrow [b_1, \dots, b_k]$

$\mathbf{T}_k \leftarrow \text{diag}(t_1, \dots, t_k)$

$\mathbf{R}_k \leftarrow \mathbf{R}$ matrix from QR decomposition of B_k $\text{// Can be precomputed}$

$\mathbf{z}_1, \dots, \mathbf{z}_{M_k} \leftarrow$ i.i.d. sample from $\mathcal{N}(\mathbf{0}, \mathbf{I}_k)$

$v_m \leftarrow B_k \mathbf{R}_k^{-1} \mathbf{T}_k^{-\frac{1}{2}} \mathbf{z}_m, 1 \leq m \leq M_k$

$\hat{g}_n \leftarrow \frac{\lambda_k}{M_k} \sum_{m=1}^{M_k} D\mathcal{R}(h_n; v_m) v_m$

$h_{n+1} \leftarrow h_n - \alpha_n \hat{g}_n$

end for

As is evident from Theorem 1, it is essential that the second moment term $\mathbb{E}[\|\hat{g}_n\|_C^2]$ be finite. By the tower property, a necessary condition would then be $\mathbb{E}_n[\|\hat{g}_n\|_C^2] \in L^1(\mathcal{S}, \mathcal{F}, \mathbb{P})$, where, we recall, \mathbb{E}_n denotes the conditional expectation given h_1, \dots, h_n . If \mathcal{H} were finite-dimensional, then usual choices for \mathbb{P}_v have finite moments of all orders, trivially ensuring a finite second moment for \hat{g}_n ; however, in infinite-dimensional spaces this is substantially harder, and is only possible due to (i) the particular design of our \mathbb{P}_v , and (ii) the introduction of the preconditioning λ_k , as we will now show.

Theorem 2. *Recall that $\gamma_i = \mathbb{E}[\lambda_K \mid K \geq i]$ and let $g_n \triangleq \nabla \mathcal{R}(h_n)$. The second moment of \hat{g}_n then satisfies*

$$\mathbb{E}_n[\|\hat{g}_n\|_C^2] = \mathbb{E}_n\left[\frac{\lambda_K^2}{M_K} \left(\sum_{i=1}^K \frac{1}{t_i \gamma_i}\right) \left(\sum_{i=1}^K \frac{\langle g_n, e_i \rangle^2}{t_i}\right)\right] + \mathbb{E}_n\left[\lambda_K^2 \left(1 + \frac{1}{M_K}\right) \left(\sum_{i=1}^K \frac{\langle g_n, e_i \rangle^2}{t_i^2 \gamma_i}\right)\right]. \quad (13)$$

Proof. See Appendix A.3. \square

Let us start by considering what happens when there is no preconditioning:

Corollary 1. *Let $\lambda_k \equiv 1$. Then, independent of the choice of M_k , a necessary condition for $\mathbb{E}_n[\|\hat{g}_n\|_C^2] < +\infty$ is*

$$\sum_{i=1}^{\infty} \frac{\langle g_n, e_i \rangle^2}{t_i} < +\infty. \quad (14)$$

Remark 3. *Note that if $\lambda_k \equiv 1$, then $C = I$ and $\|\cdot\|_C = \|\cdot\|$ meaning that, without preconditioning, we cannot guarantee $\mathbb{E}_n[\|\hat{g}_n\|^2] < +\infty$, for any $g_n \in \mathcal{H}$.*

Proof. From (13) it is immediate that

$$\mathbb{E}_n[\|\hat{g}_n\|_C^2] \geq \mathbb{E}_n\left[\lambda_K^2 \left(1 + \frac{1}{M_K}\right) \left(\sum_{i=1}^K \frac{\langle g_n, e_i \rangle^2}{t_i^2 \gamma_i}\right)\right] \geq \mathbb{E}_n\left[\sum_{i=1}^K \frac{\langle g_n, e_i \rangle^2}{t_i^2}\right],$$

where we have substituted $\lambda_k \equiv 1$. Now compute

$$\mathbb{E}_n\left[\sum_{i=1}^K \frac{\langle g_n, e_i \rangle^2}{t_i^2}\right] = \mathbb{E}_n\left[\sum_{i=1}^{\infty} \mathbf{1}\{K \geq i\} \frac{\langle g_n, e_i \rangle^2}{t_i^2}\right] = \sum_{i=1}^{\infty} \frac{\langle g_n, e_i \rangle^2}{t_i},$$

implying that $\sum_{i=1}^{\infty} \langle g_n, e_i \rangle^2 / t_i$ is a lower bound for $\mathbb{E}_n[\|\hat{g}_n\|_C^2]$. \square

Therefore, if $\lambda_k \equiv 1$, to obtain a useful bound in Theorem 1 we would need to assume (14) for every gradient g_n in Algorithm 1, which is rather restrictive. Fortunately, there is a better choice of λ_k and M_k which avoids this issue altogether. First, a useful lemma:

Lemma 1. *For any distribution \mathbb{P}_K over \mathbb{N} we have*

$$\frac{t_i}{2} \leq \mathbb{E}[t_K \mid K \geq i] \leq t_i, \quad \text{for all } i \in \mathbb{N}. \quad (15)$$

Proof. The upper bound is obvious, since $t_i = \mathbb{P}[K \geq i]$ is decreasing in i . To obtain the lower bound, expand the expectation and factor out a telescoping sum:

$$\begin{aligned} \mathbb{E}[t_K \mid K \geq i] &= \sum_{j=i}^{\infty} t_j \frac{\mathbb{P}[K=j]}{t_i} = \frac{1}{t_i} \sum_{j=i}^{\infty} t_j (t_j - t_{j+1}) \\ &= \frac{1}{t_i} \sum_{j=i}^{\infty} \frac{1}{2} (t_j^2 - t_{j+1}^2) + \frac{1}{2} (t_j - t_{j+1})^2 \\ &\geq \frac{1}{t_i} \sum_{j=i}^{\infty} \frac{1}{2} (t_j^2 - t_{j+1}^2) = \frac{t_i^2}{2t_i} = \frac{t_i}{2}, \end{aligned}$$

which follows from $t_i \rightarrow 0$ as $i \rightarrow +\infty$. \square

As an immediate consequence, we have the following second moment bound:

Corollary 2. *Choose $\lambda_k = t_k$ and $M_k = \lceil k/c \rceil$, for some $c > 0$. Then*

$$\mathbb{E}_n[\|\hat{g}_n\|_C^2] \leq 2(1+2c)\|g_n\|^2. \quad (16)$$

Proof. From (13) we can see that $\mathbb{E}_n[\|\hat{g}_n\|_C^2]$ is decreasing in M_k , so we can obtain an upper bound by substituting $M_k = k/c$. Then, by Lemma 1 we have

$$\begin{aligned} \mathbb{E}_n[\|\hat{g}_n\|_C^2] &\leq \mathbb{E}_n \left[\frac{ct_K^2}{K} \left(\sum_{i=1}^K \frac{2}{t_i^2} \right) \left(\sum_{i=1}^K \frac{\langle g_n, e_i \rangle^2}{t_i} \right) \right] + \mathbb{E}_n \left[t_K^2 \left(1 + \frac{c}{K} \right) \left(\sum_{i=1}^K \frac{2\langle g_n, e_i \rangle^2}{t_i^3} \right) \right] \\ &\leq 2c \mathbb{E}_n \left[\sum_{i=1}^K \frac{\langle g_n, e_i \rangle^2}{t_i} \right] + 2(1+c) \mathbb{E}_n \left[\sum_{i=1}^K \frac{\langle g_n, e_i \rangle^2}{t_i} \right] \\ &= 2(1+2c)\|g_n\|^2, \end{aligned}$$

where we have used the fact that $t_K \leq t_i$ if $i \leq K$. \square

Thus, combining Theorem 1 and Corollary 2, we obtain our main result:

Theorem 3. *Under the same setting as Theorem 1, take $\lambda_k = t_k$ and $M_k = \lceil k/c \rceil$ for some $c > 0$, and suppose that $\|g_n\|^2 \leq G^2$ almost surely, for all $1 \leq n \leq N$. Then, for any $h \in \mathcal{H}$,*

$$\mathbb{E}[\mathcal{R}(\hat{h})] - \mathcal{R}(h) \leq \frac{\frac{1}{2}\|h - h_1\|_C^2}{\sum_{n=1}^N \alpha_n} + (1+2c)G^2 \frac{\sum_{n=1}^N \alpha_n^2}{\sum_{n=1}^N \alpha_n}. \quad (17)$$

Proof. Immediate consequence of (12) and (16). \square

Theorem 3 implies that the learning rate α_n should be taken to satisfy the usual conditions

$$\sum_{n=1}^N \alpha_n \rightarrow +\infty \quad \text{and} \quad \frac{\sum_{n=1}^N \alpha_n^2}{\sum_{n=1}^N \alpha_n} \rightarrow 0 \quad (18)$$

as $N \rightarrow +\infty$. In fact, if we fix *a priori* the number of iterations N , then the following bound is optimal:

Corollary 3. *Under the setup of Theorem 3, take any $h \in C^{\frac{1}{2}}(\mathcal{H})$ and set*

$$\alpha_n \equiv \frac{\|h - h_1\|_C}{\sqrt{2(1+2c)G}\sqrt{N}}. \quad (19)$$

Then, the expected excess risk satisfies

$$\mathbb{E}[\mathcal{R}(\hat{h})] - \mathcal{R}(h) \leq \frac{\|h - h_1\|_C \sqrt{2(1+2c)G}}{\sqrt{N}}$$

and this bound is optimal.

Proof. See Appendix A.4. \square

In order to obtain the best possible bound in Theorem 3 or Corollary 3 with relation to $h \in C^{\frac{1}{2}}(\mathcal{H})$, we would like to take $h = h^* \in \arg \min_{h' \in \mathcal{H}} \mathcal{R}(h')$. However, if $h^* \notin C^{\frac{1}{2}}(\mathcal{H})$, then $\|h^* - h_1\|_C^2 = +\infty$, and the bound becomes vacuous. By Lemma 1, note that $h^* \in C^{\frac{1}{2}}(\mathcal{H})$ is equivalent to

$$\sum_{i=1}^{\infty} \frac{\langle h^*, e_i \rangle^2}{t_i} < +\infty. \quad (20)$$

Although there is no way to guarantee this holds *a priori* without additional knowledge about h^* , it is easy to show that, whatever h^* may be, there exists some distribution \mathbb{P}_K for which (20) holds.

Proposition 2. *For any $h^* \in \mathcal{H}$, there exists a distribution \mathbb{P}_K for which $t_i = \mathbb{P}_K([i, +\infty))$ satisfies (20). For this \mathbb{P}_K , we get $h^* \in C^{\frac{1}{2}}(\mathcal{H})$, implying $\|h^* - h_1\|_C^2 < +\infty$.*

Proof. Let $\theta_i = \langle h^*, e_i \rangle^2$, and set $A_i = \sum_{j \geq i} \theta_j$. Since $A_i \rightarrow 0$ as $i \rightarrow +\infty$, there exists a strictly increasing sequence $\{n_k\}_{k \in \mathbb{N}}$ such that $n_k \geq 2$ and $A_{n_k} \leq 2^{-k}$ for all $k \in \mathbb{N}$. Set $n_0 = 1$ and define $\{t_i\}_{i \in \mathbb{N}}$ by setting $t_i = 1/k$ for $n_{k-1} \leq i < n_k$. It is immediate that $t_1 = 1$, t_i is non-increasing and $t_i \rightarrow 0$ as $i \rightarrow +\infty$. Then

$$\sum_{i=1}^{\infty} \frac{\langle h^*, e_i \rangle^2}{t_i} = \sum_{i=1}^{\infty} \frac{\theta_i}{t_i} = \sum_{k=1}^{\infty} \sum_{i=n_{k-1}}^{n_k-1} k\theta_i \leq \sum_{k=1}^{\infty} kA_{n_{k-1}} \leq \sum_{k=1}^{\infty} k2^{-k+1} < +\infty.$$

Hence, defining \mathbb{P}_K by $\mathbb{P}_K(\{i\}) = t_i - t_{i+1}$, we get $h^* \in C^{\frac{1}{2}}(\mathcal{H})$. \square

Extensions Although we assume the distribution of \mathbf{z} given K to be exactly $\mathcal{N}(\mathbf{0}, \mathbf{I}_K)$, this is not necessary. Our proofs only rely on the first four moments of \mathbf{z} , therefore, all results remain valid if we swap $\mathcal{N}(\mathbf{0}, \mathbf{I}_K)$ by a distribution that matches all of its moments (including mixed ones) up to order four.

3 Application: solving PDEs with functional optimization

In this section, we show how our framework can be applied to tackle our motivating example of solving PDEs, presented in Section 1.1. To showcase our method's flexibility, we consider one linear PDE, the heat equation, and one non-linear PDE, an HJB equation. They result in convex and non-convex optimization problems, respectively.

3.1 Formal setup and parameters

Let $\Omega \subset \mathbb{R}^d$ be a bounded open set with a boundary smooth enough to admit a trace operator⁵ $\text{Tr} : \mathbb{H}^1(\Omega) \rightarrow L^2(\partial\Omega)$. For the sake of concreteness, we consider general PDEs with Dirichlet boundary

⁵We recall that the trace operator is the proper extension of the “restriction to the boundary” operator over continuous functions to Sobolev spaces. It exists, for example, if the boundary of Ω is Lipschitz [7, Theorem 1.5.1.3], in the sense that it is, locally and after possibly a change of coordinates, the graph of a Lipschitz function.

conditions:⁶

$$\begin{cases} L[u](\mathbf{x}) = 0 & \mathbf{x} \in \Omega, \\ u(\mathbf{x}) = f(\mathbf{x}) & \mathbf{x} \in \Lambda \subset \partial\Omega. \end{cases} \quad (21)$$

Here, the boundary condition is interpreted in the trace sense and L is a, possibly non-affine, ℓ -th order differential operator of the form

$$L[u](\mathbf{x}) = F(\mathbf{x}, u(\mathbf{x}), Du(\mathbf{x}), \dots, D^{(\ell)}u(\mathbf{x})),$$

with $F : \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d \times \dots \times \mathbb{R}^{d^\ell} \rightarrow \mathbb{R}$ being a given function. We assume that $L[u] \in L^2(\Omega)$ for all $u \in \mathbb{H}^\ell(\Omega)$. This is easily checked if L is an affine operator with bounded coefficients, in which case we further obtain convexity and Fréchet differentiability of the risk functional

$$\mathcal{R}(h) = \frac{1}{2} \|L[h]\|_{L^2(\Omega)}^2 + \frac{1}{2} \|(\text{Tr}[h] - f)\mathbf{1}_\Lambda\|_{L^2(\partial\Omega)}^2, \quad (22)$$

as in Equation (3). We will also consider the case of nonlinear L , in which this assumption may not hold, showing empirically that our algorithm still provides good approximations in more challenging scenarios.

To be applied in this setting, our framework requires a pre-basis $\mathcal{B} = \{b_i\}_{i \in \mathbb{N}}$ for the Sobolev space $\mathbb{H}^\ell(\Omega)$, where $\ell \in \mathbb{N}$. To obtain one, we will take advantage of the theory of Reproducing Kernel Hilbert Spaces (RKHSs) and their smoothness properties. Recall the Matérn kernel [16, Section 4.2], given by

$$\text{Matern}_{\nu, \eta}(\mathbf{x}, \mathbf{y}) = \frac{2^{1-\nu}}{\Gamma(\nu)} \left(\frac{\sqrt{2\nu} \|\mathbf{x} - \mathbf{y}\|}{\eta} \right)^\nu K_\nu \left(\frac{\sqrt{2\nu} \|\mathbf{x} - \mathbf{y}\|}{\eta} \right), \quad \mathbf{x}, \mathbf{y} \in \Omega, \quad (23)$$

where $\eta > 0$ is the bandwidth parameter, $\nu > 0$ is the smoothness parameter, Γ is the usual Γ -function and K_ν is a modified Bessel function. When $\nu - 1/2$ is a nonnegative integer, then $\text{Matern}_{\nu, \eta}$ admits a simple closed form as a product of exponential and polynomial terms [16].

The following proposition establishes that Matérn kernels centered at appropriately chosen centers \mathbf{x} form a pre-basis for $\mathbb{H}^\ell(\Omega)$.

Proposition 3. *Let $\Omega \subset \mathbb{R}^d$ be a bounded, open set with Lipschitz boundary, and let $\{\mathbf{x}_i\}_{i \in \mathbb{N}}$ be a dense subset of Ω . For $\ell \in \mathbb{N}$, take $\nu > 0$ such that $\nu + d/2$ is greater than or equal to ℓ . Then, for any $\eta > 0$, the set $\mathcal{B} = \{\text{Matern}_{\nu, \eta}(\mathbf{x}_i, \cdot)\}_{i \in \mathbb{N}}$ is a pre-basis for $\mathbb{H}^\ell(\Omega)$.*

Proof. The proof is given in Appendix A.5. □

So, as long as we have a countable set of points $\{\mathbf{x}_i\}_{i \in \mathbb{N}}$ that is dense in Ω (e.g., generated from quasi-random sampling at rational coordinates), the sequence of Matérn kernels centered at the \mathbf{x}_i is a suitable pre-basis. This holds for any differentiability order ℓ and any bounded domain Ω with Lipschitz boundary, even when the Sobolev space $\mathbb{H}^\ell(\Omega)$ is not an RKHS and/or the computation of an orthonormal basis becomes intractable.⁷

Implementation details In order to compute Sobolev inner products (needed for the QR decomposition of the quasi-matrices), as well as directional derivatives of the risk defined in (22), we need to compute multi-dimensional integrals. In our experiments, we estimate these integrals with Quasi-Monte Carlo using the Roberts sequence [17]. We chose this method for its scalability to higher dimensions, and superior performance compared to simple random sampling. Computation of the directional derivatives is done using scalar automatic differentiation, by writing $D\mathcal{R}(h; v) = \frac{d}{d\delta}\mathcal{R}(h + \delta v)|_{\delta=0}$.

⁶Other types of boundary conditions, such as Neumann or periodic boundary conditions, can be treated analogously.

⁷The computation of an orthonormal basis is highly dependent on both ℓ and Ω , and generally one is not known in closed form.




Figure 1: **Our method on the heat equation.** On the top row we visualize a heatmap of the solution $u(t, x)$, with $x \in [0, 2\pi]$ and $t \in [0, 1]$; on the bottom row, we visualize the solutions $u(t, x)$ at three points in time ($t = 0$, $t = 0.5$ and $t = 1$). As the theory suggests, our method with preconditioning effectively and provably finds the correct solution to the heat equation. Without preconditioning, the procedure quickly diverges.

3.2 Case I: Heat equation

As a first example, consider the heat equation

$$\begin{cases} \partial_t u(t, x) = \Delta u(t, x) & (t, x) \in (0, T) \times U, \\ u(t, x) = f(t, x) & (t, x) \in \Lambda, \end{cases} \quad (24)$$

for some open set $U \subset \mathbb{R}^d$ and boundary condition f over the parabolic boundary Λ of $(0, T) \times U$. When written in the form of Equation (2) we see that the corresponding differential operator $L = \partial_t - \Delta$ is linear, rendering the optimization problem convex. For the sake of concreteness, we shall consider the following special case:

$$\begin{cases} \partial_t u(t, x) = \Delta u(t, x) & t \in (0, T), x \in (0, 2\pi), \\ u(0, x) = \sin x & x \in (0, 2\pi), \\ u(t, 0) = u(t, 2\pi) = 0 & t \in (0, T). \end{cases} \quad (25)$$




Figure 2: **Convergence of our method on the heat equation.** The plots display (a) the evolution of the risk $\mathcal{R}(h_n)$ over iterations, and (b) the evolution of the distance from the estimates h_n to the optimal solution h^* in L^2 norm. Note how our algorithm with preconditioning steadily reduces both the risk and distance to the solution, whereas without preconditioning it quickly diverges.

This has a known unique solution given by $u(t, x) = e^{-t} \sin x$, allowing us to easily evaluate our procedure.

To solve Equation (25) we let $\Omega = (0, T) \times (0, 2\pi)$ and construct the risk functional $\mathcal{R} : \mathbb{H}^2(\Omega) \rightarrow \mathbb{R}$ given by

$$\begin{aligned} \mathcal{R}(h) &= \frac{1}{2} \|L[h]\|_{L^2(\Omega)}^2 + \frac{1}{2} \|h - f\|_{L^2(\partial\Omega)}^2 \\ &= \frac{1}{2} \int_0^T \int_0^{2\pi} [\partial_t h(t, x) - \Delta h(t, x)]^2 dx dt + \frac{1}{2} \int_0^{2\pi} [h(0, x) - \sin x]^2 dx \\ &\quad + \frac{1}{2} \int_0^T [h(t, 0) - 0]^2 dt + \frac{1}{2} \int_0^T [h(t, 2\pi) - 0]^2 dt. \end{aligned}$$

Note that, by the linearity of L and the discussion in Section 1.1, this is a well-defined, convex, Fréchet differentiable risk functional on $\mathbb{H}^2(\Omega)$, rendering our theory immediately applicable.

We take $T = 1$, $\mathbb{P}_K = \text{Poisson}(100)$ with sample sizes $M_k = \lceil k/2 \rceil$, initial position $h_1 \equiv 0$ and constant learning rates $\alpha_n \equiv 0.6$. For our pre-basis, we take $\eta = 20$ and sample centers $\{\mathbf{x}_i\}_{i \in \mathbb{N}} \subset \Omega$ with the Roberts quasi-random sequence [17]; this ensures that the centered kernels from Proposition 3 form a proper pre-basis.

Figure 1 illustrates our procedure in two settings: with preconditioning (i.e., $\lambda_k = t_k$), and without (i.e., $\lambda_k \equiv 1$). As we can see, under preconditioning we fall under the regime of Corollary 3, and our method converges to the true solution. However, without preconditioning, which is equivalent to running a naïve SGD method, our method quickly diverges. This is most likely caused by the high variance of the stochastic gradients, as noted in Corollary 1. In a similar vein, Figure 2 clearly displays the convergence of our preconditioned method to the minimizer, while also exhibiting the quick divergent behavior produced by vanilla stochastic gradients.

3.3 Case II: Hamilton-Jacobi-Bellman equation

For our second example, we consider an HJB equation, which is a type of nonlinear PDE with terminal boundary condition central to optimal control theory:

$$\begin{cases} \partial_t u(t, x) + \inf_{|c| \leq \bar{c}} H(x, \partial_x u(t, x), c) = 0, & (t, x) \in (0, T) \times (-\bar{x}, \bar{x}) \\ u(T, x) = Gx^2 & x \in (-\bar{x}, \bar{x}), \end{cases} \quad (26)$$

where $H(x, p, c) = Ax^2 + Bc^2 + p \cdot c$ is the Hamiltonian. The parameters A, B, G, \bar{c} and \bar{x} are positive constants. This particular HJB equation is satisfied by the value function of the following




Figure 3: **Our method on an HJB equation.** On the top row we visualize a heatmap of the solution $u(t, x)$, with $x \in [-3, +3]$ and $t \in [0, 5]$; on the middle row, we visualize the solutions $u(t, x)$ at three points in time ($t = 2/8$, $t = 2.5$ and $t = 5$). Finally, on the bottom row we visualize heatmaps of the optimal control arising from the corresponding solutions u . As we can see, our method finds the correct solution even without convexity of the risk functional.

linear-quadratic control problem:

$$\begin{aligned} \inf_{c \in L^\infty([t, T])} \quad & \int_t^T [Ax(s)^2 + Bc(s)^2] \, ds + Gx(T)^2, \\ \text{s.t.} \quad & \|c\|_\infty \leq \bar{c}, \\ & \dot{x} = c, \quad x(t) \in [-\bar{x}, \bar{x}]. \end{aligned} \tag{27}$$

We take $G = \sqrt{AB}$, so that (26) has a closed form solution, obtained by solving (27) and computing the cost attained by the optimal control $c^*(t, x)$.

To simplify the notation, we define

$$\tilde{H}(x, p) = \inf_{|c| \leq \bar{c}} H(x, p, c) = \begin{cases} Ax^2 - \frac{p^2}{4B}, & \text{if } |p| \leq 2B\bar{c}, \\ Ax^2 + B\bar{c}^2 - \bar{c}|p|, & \text{if } |p| > 2B\bar{c}, \end{cases} \tag{28}$$

and let the risk functional be given by

$$\mathcal{R}(h) = \frac{1}{2} \int_0^T \int_{-\bar{x}}^{\bar{x}} [\partial_t h(t, x) + \tilde{H}(x, \partial_x h(t, x))]^2 \, dx \, dt + \frac{1}{2} \int_{-\bar{x}}^{\bar{x}} [h(T, x) - Gx^2]^2 \, dx. \tag{29}$$

In this setting, we have:

Proposition 4. *The risk \mathcal{R} in (29) is a Fréchet differentiable, non-convex functional $\mathcal{R} : \mathbb{H}^1(\Omega) \rightarrow \mathbb{R}$, where $\Omega = (0, T) \times (-\bar{x}, \bar{x})$.*

Proof. See Appendix A.6. □

Nevertheless, the directional derivatives are still just as easily computable through automatic differentiation.

For this experiment we take $T = 5$, $A = B = 1$, $\bar{c} = 0.5$, $\bar{x} = 3$, $\mathbb{P}_K = \text{Poisson}(100)$ with sample sizes $M_k = \lceil k/1.5 \rceil$, initial position $h_1 \equiv 0$ and constant learning rates $\alpha_n \equiv 0.2$, and use the same Matérn-based pre-basis with $\eta = 20$ and centers sampled from the Roberts quasi-random sequence [17]. We furthermore use the proper preconditioning $\lambda_k = t_k$. Figure 3 shows the results of this experiment, featuring both the solution $u(t, x)$ of the differential equation and the resulting optimal control $c^*(t, x)$. We see that despite the apparent nonconvexity of the risk, our method still effectively converges to the correct solution, highlighting its versatility.

4 Conclusion

In this paper, we introduced a random gradient-free method for optimization in infinite-dimensional separable Hilbert spaces, avoiding the computational challenges common to functional gradient descent. By using only directional derivatives and a pre-basis, our approach offers a practical and broadly applicable framework for solving functional optimization problems. We believe that our work expands the applicability of functional gradient methods, particularly to settings with specialized losses and for which functional gradients would need to be computed accordingly. In the future it would be interesting to consider variants such as constrained, mirror and accelerated versions of our algorithm.

References

- [1] Charles R. Baker and Ian W. McKeague. “Compact covariance operators”. en. In: *Proceedings of the American Mathematical Society* 83.3 (Mar. 1981), pp. 590–590. ISSN: 0002-9939. DOI: 10.1090/S0002-9939-1981-0627699-7.
- [2] Salvatore Cuomo et al. “Scientific Machine Learning Through Physics-Informed Neural Networks: Where we are and What’s Next”. In: *Journal of Scientific Computing* 92 (2022).
- [3] Giuseppe Da Prato. *An introduction to infinite-dimensional analysis*. en. Universitext. Berlin: Springer, 2006. ISBN: 978-3-540-29020-9.
- [4] John Duchi. “Introductory lectures on stochastic optimization”. en. In: *IAS/Park City Mathematics Series*. Vol. 25. Providence, Rhode Island: American Mathematical Society, Nov. 2018, pp. 99–185. ISBN: 978-1-4704-3575-2. DOI: 10.1090/pcms/025/03. URL: <http://www.ams.org/pcms/025>.
- [5] Yuri Fonseca, Caio Peixoto, and Yuri Saporito. “Nonparametric Instrumental Variable Regression through Stochastic Approximate Gradients”. In: *The Thirty-eighth Annual Conference on Neural Information Processing Systems*. 2024.
- [6] Yuri Fonseca and Yuri Saporito. “Statistical Learning and Inverse Problems: A Stochastic Gradient Approach”. In: *Advances in Neural Information Processing Systems*. Ed. by Alice H. Oh et al. 2022.
- [7] Pierre Grisvard. *Elliptic problems in nonsmooth domains*. eng. Classics in applied mathematics. Philadelphia: Society for industrial and applied mathematics, 2011. ISBN: 978-1-61197-202-3.
- [8] David Kozak et al. “A stochastic subspace approach to gradient-free optimization in high dimensions”. en. In: *Computational Optimization and Applications* 79.2 (June 2021), pp. 339–368. ISSN: 0926-6003, 1573-2894. DOI: 10.1007/s10589-021-00271-w.
- [9] William McLean. *Strongly elliptic systems and boundary integral equations*. eng. 1. publ. Cambridge: Cambridge University Press, 2000. ISBN: 978-0-521-66375-5.
- [10] Johannes Müller and Marius Zeinhofer. “Achieving High Accuracy with PINNs via Energy Natural Gradient Descent”. In: *Proceedings of the 40th International Conference on Machine Learning*. Ed. by Andreas Krause et al. Vol. 202. Proceedings of Machine Learning Research. PMLR, July 2023, pp. 25471–25485.

- [11] Johannes Müller and Marius Zeinhofer. “Position: Optimization in SciML Should Employ the Function Space Geometry”. In: *Proceedings of the 41st International Conference on Machine Learning*. Ed. by Ruslan Salakhutdinov et al. Vol. 235. Proceedings of Machine Learning Research. PMLR, July 2024, pp. 36705–36722.
- [12] Yurii Nesterov and Vladimir Spokoiny. “Random Gradient-Free Minimization of Convex Functions”. en. In: *Foundations of Computational Mathematics* 17.2 (Apr. 2017), pp. 527–566. ISSN: 1615-3375, 1615-3383. DOI: 10.1007/s10208-015-9296-2.
- [13] Hemant Kumar Pathak. *An Introduction to Nonlinear Analysis and Fixed Point Theory*. Springer, 2018.
- [14] Maziar Raissi, Paris Perdikaris, and George Em Karniadakis. “Physics-informed neural networks: A deep learning framework for solving forward and inverse problems involving nonlinear partial differential equations”. In: *J. Comput. Phys.* 378 (2019), pp. 686–707.
- [15] Maziar Raissi et al. *Physics-Informed Neural Networks and Extensions*. 2024. arXiv: 2408.16806 [cs.LG].
- [16] Carl Edward Rasmussen and Christopher K. I. Williams. *Gaussian Processes for Machine Learning*. en. 3. print. Adaptive Computation and Machine Learning. Cambridge, Mass.: MIT Press, 2008. ISBN: 978-0-262-18253-9.
- [17] Martin Roberts. *The Unreasonable Effectiveness of Quasirandom Sequences*. 2018. URL: <https://extremelearning.com.au/unreasonable-effectiveness-of-quasirandom-sequences/> (visited on 12/11/2025).
- [18] W. Rudin. *Functional Analysis*. International series in pure and applied mathematics. Tata McGraw-Hill, 1974. ISBN: 978-0-07-099558-1.
- [19] Justin Sirignano and Konstantinos Spiliopoulos. “DGM: A deep learning algorithm for solving partial differential equations”. In: *Journal of Computational Physics* 375 (2018), pp. 1339–1364. ISSN: 0021-9991.
- [20] S. U. Stich, C. L. Müller, and B. Gärtner. “Optimization of Convex Functions with Random Pursuit”. In: *SIAM Journal on Optimization* 23.2 (2013), pp. 1284–1309. DOI: 10.1137/110853613. eprint: <https://doi.org/10.1137/110853613>.
- [21] L. N. Trefethen. “Householder triangularization of a quasimatrix”. en. In: *IMA Journal of Numerical Analysis* 30.4 (Oct. 2010), pp. 887–897. ISSN: 0272-4979, 1464-3642. DOI: 10.1093/imanum/drp018.
- [22] N. N. Vakhania and V. I. Tarieladze. “Covariance Operators of Probability Measures in Locally Convex Spaces”. en. In: *Theory of Probability & Its Applications* 23.1 (Dec. 1978), pp. 1–21. ISSN: 0040-585X, 1095-7219. DOI: 10.1137/1123001.
- [23] Holger Wendland. *Scattered Data Approximation*. 1st ed. Cambridge University Press, Dec. 2004. ISBN: 978-0-521-84335-5. DOI: 10.1017/CBO9780511617539.
- [24] Y Yamasaki. *Measures on Infinite Dimensional Spaces*. en. Vol. 5. Series in Pure Mathematics. WORLD SCIENTIFIC, July 1985. ISBN: 978-9971-978-52-5. DOI: 10.1142/0162.

A Proofs of selected results

A.1 Proof of Proposition 1

The fact that $\mathbb{E}[v | K] = 0$ is obvious from Equation (9), since $\mathbb{E}[\mathbf{z} | K] = \mathbf{0}$ and then

$$\mathbb{E}[v | K] = B_K \mathbf{R}_K^{-1} \mathbf{T}_K^{-\frac{1}{2}} \mathbb{E}[\mathbf{z} | K] = 0.$$

To show that $\mathbb{E}[v] = 0$, we need to show that $\mathbb{E}[\langle v, h \rangle] = 0$ for any $h \in \mathcal{H}$. This is easily obtained if we use the tower property of conditional expectations to compute $\mathbb{E}[\langle v, h \rangle] = \mathbb{E}[\mathbb{E}[\langle v, h \rangle | K]] = \mathbb{E}[\langle \mathbb{E}[v | K], h \rangle] = 0$. However, to make sure that the conditional expectations we are using actually exist and we can use the tower property, we must first ensure that $\langle v, h \rangle \in L^1(\mathcal{S}, \mathcal{F}, \mathbb{P})$. This is done in the following lemma:

Lemma 2. *For any $h \in \mathcal{H}$, $\langle v, h \rangle \in L^1(\mathcal{S}, \mathcal{F}, \mathbb{P})$.*

Proof. Since the random variable $|\langle v, h \rangle|$ is non-negative, its conditional expectation given K always exists, even if $|\langle v, h \rangle|$ is not integrable. Therefore, now we can use the tower property and write:

$$\mathbb{E}[|\langle v, h \rangle|] = \mathbb{E}[\mathbb{E}[|\langle v, h \rangle| | K]] \leq \mathbb{E} \left[\mathbb{E}[\langle v, h \rangle^2 | K]^{\frac{1}{2}} \right] = \mathbb{E} \left[\mathbb{E}[\langle Q_K \mathbf{T}_K^{-\frac{1}{2}} \mathbf{z}, x \rangle^2 | K]^{\frac{1}{2}} \right],$$

where we have used Jensen's inequality and the identity $v = Q_K \mathbf{T}_K^{-\frac{1}{2}} \mathbf{z}$, which is immediate from (9). Now note that, given K , we have

$$\langle Q_K \mathbf{T}_K^{-\frac{1}{2}} \mathbf{z}, h \rangle = \langle \mathbf{z}, \mathbf{T}_K^{-\frac{1}{2}} Q_K^* h \rangle \sim \mathcal{N}(0, \|\mathbf{T}_K^{-\frac{1}{2}} Q_K^* h\|^2).$$

This implies:

$$\begin{aligned} \mathbb{E} \left[\mathbb{E}[\langle Q_K \mathbf{T}_K^{-\frac{1}{2}} \mathbf{z}, h \rangle^2 | K]^{\frac{1}{2}} \right] &= \mathbb{E} \left[\|\mathbf{T}_K^{-\frac{1}{2}} Q_K^* h\| \right] \\ &\leq \mathbb{E} \left[\|\mathbf{T}_K^{-\frac{1}{2}} Q_K^* h\|^2 \right]^{\frac{1}{2}} = \mathbb{E} \left[\sum_{i=1}^K \frac{\langle h, e_i \rangle^2}{t_i} \right]^{\frac{1}{2}}. \end{aligned}$$

Now we rewrite the summation and apply Tonelli's Theorem to obtain

$$\begin{aligned} \mathbb{E} \left[\sum_{i=1}^K \frac{\langle h, e_i \rangle^2}{t_i} \right]^{\frac{1}{2}} &= \mathbb{E} \left[\sum_{i=1}^{\infty} \frac{\mathbf{1}\{K \geq i\} \langle h, e_i \rangle^2}{t_i} \right]^{\frac{1}{2}} \\ &= \left(\sum_{i=1}^{\infty} \mathbb{E} \left[\frac{\mathbf{1}\{K \geq i\} \langle h, e_i \rangle^2}{t_i} \right] \right)^{\frac{1}{2}} = \left(\sum_{i=1}^{\infty} \langle h, e_i \rangle^2 \right)^{\frac{1}{2}} = \|h\|. \end{aligned}$$

Thus, $\mathbb{E}[|\langle v, h \rangle|] \leq \|h\| < +\infty$. □

Since the distribution of v given K has zero mean and is finite dimensional, we have $\text{Cov}[v | K] = \mathbb{E}[v \otimes v | K]$. We expand this expression, using again that $v = Q_K \mathbf{T}_K^{-\frac{1}{2}} \mathbf{z}$:

$$\begin{aligned} \mathbb{E}[v \otimes v | K] &= \mathbb{E}[Q_K \mathbf{T}_K^{-\frac{1}{2}} \mathbf{z} \otimes Q_K \mathbf{T}_K^{-\frac{1}{2}} \mathbf{z} | K] \\ &= \mathbb{E}[Q_K \mathbf{T}_K^{-\frac{1}{2}} (\mathbf{z} \otimes \mathbf{z}) \mathbf{T}_K^{-\frac{1}{2}} Q_K^* | K] = Q_K \mathbf{T}_K^{-\frac{1}{2}} \mathbb{E}[\mathbf{z} \otimes \mathbf{z} | K] \mathbf{T}_K^{-\frac{1}{2}} Q_K^*. \end{aligned}$$

By the definition of \mathbf{z} in Equation (9), we have $\mathbb{E}[\mathbf{z} \otimes \mathbf{z} | K] = \mathbf{I}_K$. Thus:

$$\mathbb{E}[v \otimes v | K] = Q_K \mathbf{T}_K^{-1} Q_K^*.$$

This shows that $\text{Cov}[\mathbb{P}_{v|K}] = \text{Cov}[v | K] = Q_K \mathbf{T}_K^{-1} Q_K^*$.

To obtain $\text{Cov}[\mathbb{P}_v] = \text{Cov}[v] = I$, we must take $h, h' \in \mathcal{H}$ and show that $\mathbb{E}[\langle h, v \rangle \langle h', v \rangle] = \langle h, h' \rangle$. Once again, we would like to proceed by conditioning on K . The next lemma shows that this is possible:

Lemma 3. For any $h, h' \in \mathcal{H}$, we have $\langle h, v \rangle \langle h', v \rangle \in L^1(\mathcal{S}, \mathcal{F}, \mathbb{P})$.

Proof. By Cauchy-Schwarz in $L^2(\mathcal{S}, \mathcal{F}, \mathbb{P})$, we have

$$\mathbb{E}[|\langle h, v \rangle \langle h', v \rangle|] \leq \mathbb{E}[\langle h, v \rangle^2]^{\frac{1}{2}} \mathbb{E}[\langle h', v \rangle^2]^{\frac{1}{2}}.$$

By the arguments in the proof of Lemma 2 we know that

$$\mathbb{E}[\langle h', v \rangle^2] = \mathbb{E}[\mathbb{E}[\langle h', v \rangle^2 | K]] = \mathbb{E}[\|\mathbf{T}_K^{-\frac{1}{2}} Q_K^* h\|^2] = \|h\|^2,$$

and likewise for h' . Therefore, $\mathbb{E}[|\langle h, v \rangle \langle h', v \rangle|] \leq \|h\| \|h'\| < +\infty$. \square

Now, conditioning on K we obtain

$$\begin{aligned} \mathbb{E}[\langle h, v \rangle \langle h', v \rangle] &= \mathbb{E}[\mathbb{E}[\langle h, v \rangle \langle h', v \rangle | K]] \\ &= \mathbb{E}[\langle \text{Cov}[v | K] h, h' \rangle] = \mathbb{E}[\langle Q_K \mathbf{T}_K^{-1} Q_K^* h, h' \rangle] = \mathbb{E}[\langle \mathbf{T}_K^{-1} Q_K^* h, Q_K^* h' \rangle]. \end{aligned} \quad (30)$$

Since $Q_K = [e_1, \dots, e_K]$ and the e_i are orthonormal, we have

$$\langle \mathbf{T}_K^{-1} Q_K^* h, Q_K^* h' \rangle = \sum_{i=1}^K \frac{\langle h, e_i \rangle \langle h', e_i \rangle}{t_i} = \sum_{i=1}^{\infty} \frac{\mathbf{1}\{K \geq i\}}{t_i} \langle h, e_i \rangle \langle h', e_i \rangle. \quad (31)$$

Taking expectations on both sides and applying Fubini to bring out the sum, we get

$$\mathbb{E}[\langle h, v \rangle \langle h', v \rangle] = \sum_{i=1}^{\infty} \frac{\mathbb{E}[\mathbf{1}\{K \geq i\}]}{t_i} \langle h, e_i \rangle \langle h', e_i \rangle = \sum_{i=1}^{\infty} \langle h, e_i \rangle \langle h', e_i \rangle = \langle h, h' \rangle. \quad (32)$$

Therefore, $\text{Cov}[v] = I$. Finally, to show $\mathbb{E}_n[\widehat{g}_n(K)] = Cg_n$, we will take an arbitrary $h \in \mathcal{H}$ and show that $\mathbb{E}_n[\langle \widehat{g}_n(K), h \rangle] = \langle Cg_n, h \rangle$. The argument to show that $\langle \widehat{g}_n(K), h \rangle \in L^1(\mathcal{S}, \mathcal{F}, \mathbb{P})$ is very similar to the one presented in Lemma 3, so it will be omitted. Start by conditioning on K and recalling that $D\mathcal{R}(h_n, v_m) = \langle g_n, v_m \rangle$:

$$\begin{aligned} \mathbb{E}_n[\langle \widehat{g}_n(K), h \rangle] &= \mathbb{E}_n[\mathbb{E}_n[\langle \widehat{g}_n(K), h \rangle | K]] = \mathbb{E}_n\left[\frac{\lambda_K}{M_K} \sum_{m=1}^{M_K} \mathbb{E}_n[\langle g_n, v_m \rangle \langle x, v_m \rangle | K]\right] \\ &= \mathbb{E}_n[\lambda_K \mathbb{E}_n[\langle g_n, v \rangle \langle h, v \rangle | K]], \end{aligned}$$

where we have used that v_1, \dots, v_{M_K} are i.i.d. with the same distribution as v . By Equations (30) and (31) we know that

$$\mathbb{E}_n[\langle g_n, v \rangle \langle h, v \rangle | K] = \sum_{i=1}^{\infty} \frac{\mathbf{1}\{K \geq i\}}{t_i} \langle g_n, e_i \rangle \langle h, e_i \rangle. \quad (33)$$

Now, again by Fubini we obtain

$$\begin{aligned} \mathbb{E}_n[\langle \widehat{g}_n(K), h \rangle] &= \mathbb{E}_n\left[\sum_{i=1}^{\infty} \frac{\lambda_K \mathbf{1}\{K \geq i\}}{t_i} \langle g_n, e_i \rangle \langle h, e_i \rangle\right] \\ &= \sum_{i=1}^{\infty} \mathbb{E}\left[\frac{\lambda_K \mathbf{1}\{K \geq i\}}{t_i}\right] \langle g_n, e_i \rangle \langle h, e_i \rangle \\ &= \sum_{i=1}^{\infty} \mathbb{E}[\lambda_K | K \geq i] \langle g_n, e_i \rangle \langle h, e_i \rangle = \langle Cg_n, h \rangle, \end{aligned}$$

which finishes the proof.

A.2 Proof of Theorem 1

Let us start by recalling that $\widehat{g}_n \in \bigcup_{k=1}^{\infty} \mathcal{B}_k \subset C(\mathcal{H})$ almost surely. Thus, since $h_1 \in C(\mathcal{H})$ by assumption, we have $h_n \in C(\mathcal{H})$ for all $n = 1, \dots, N$. We emphasize that both \widehat{g}_n and h_n belong to $C(\mathcal{H})$ in addition to $C^{\frac{1}{2}}(\mathcal{H})$ because we will use the following property of $\langle \cdot, \cdot \rangle_C$: if $h \in C^{\frac{1}{2}}(\mathcal{H})$ and $h' \in C(\mathcal{H})$, we have $\langle h, h' \rangle_C = \langle h, C^{-1}h' \rangle$.

Set $g_n \triangleq \nabla \mathcal{R}(h_n)$ and expand $\frac{1}{2} \|h_{n+1} - h\|_C^2$:

$$\begin{aligned} \frac{1}{2} \|h_{n+1} - h\|_C^2 &= \frac{1}{2} \|h_n - \alpha_n \widehat{g}_n - h\|_C^2 \\ &= \frac{1}{2} \|h_n - h\|_C^2 + \frac{1}{2} \alpha_n^2 \|\widehat{g}_n\|_C^2 - \alpha_n \langle h_n - h, C^{-1} \widehat{g}_n \rangle \\ &= \frac{1}{2} \|h_n - h\|_C^2 + \frac{1}{2} \alpha_n^2 \|\widehat{g}_n\|_C^2 - \alpha_n \langle h_n - h, g_n \rangle \\ &\quad + \alpha_n \langle h_n - h, g_n - C^{-1} \widehat{g}_n \rangle. \end{aligned}$$

By the convexity of \mathcal{R} , we have $\mathcal{R}(h) - \mathcal{R}(h_n) \geq \langle h - h_n, g_n \rangle$, which gives us

$$-\alpha_n \langle h_n - h, g_n \rangle \leq -\alpha_n (\mathcal{R}(h_n) - \mathcal{R}(h)).$$

Hence, we obtain

$$\begin{aligned} \frac{1}{2} \|h_{n+1} - h\|_C^2 &\leq \frac{1}{2} \|h_n - h\|_C^2 + \frac{1}{2} \alpha_n^2 \|\widehat{g}_n\|_C^2 - \alpha_n (\mathcal{R}(h_n) - \mathcal{R}(h)) \\ &\quad + \alpha_n \langle \widehat{h}_n - h, g_n - C^{-1} \widehat{g}_n \rangle, \end{aligned}$$

which we rewrite as

$$\begin{aligned} \alpha_n (\mathcal{R}(h_n) - \mathcal{R}(h)) &\leq \frac{1}{2} \|h_n - h\|_C^2 + \frac{1}{2} \alpha_n^2 \|\widehat{g}_n\|_C^2 - \frac{1}{2} \|h_{n+1} - h\|_C^2 + \alpha_n \langle \widehat{h}_n - h, g_n - C^{-1} \widehat{g}_n \rangle. \end{aligned}$$

Summing this inequality for $n = 1, \dots, N$ we get

$$\sum_{n=1}^N \alpha_n (\mathcal{R}(h_n) - \mathcal{R}(h)) \leq \frac{1}{2} \|h_1 - h\|_C^2 + \frac{1}{2} \sum_{n=1}^N \alpha_n^2 \|\widehat{g}_n\|_C^2 + \sum_{n=1}^N \alpha_n \langle \widehat{h}_n - h, g_n - C^{-1} \widehat{g}_n \rangle.$$

Again by convexity, we have $\mathcal{R}(\widehat{h}) \leq (\sum_{n=1}^N \alpha_n \mathcal{R}(h_n)) / \sum_{n=1}^N \alpha_n$, implying

$$\mathcal{R}(\widehat{h}) - \mathcal{R}(h) \leq \frac{\frac{1}{2} \|h - h_1\|_C^2}{\sum_{n=1}^N \alpha_n} + \frac{\frac{1}{2} \sum_{n=1}^N \alpha_n^2 \|\widehat{g}_n\|_C^2}{\sum_{n=1}^N \alpha_n} + \frac{\sum_{n=1}^N \alpha_n \langle h_n - h, g_n - C^{-1} \widehat{g}_n \rangle}{\sum_{n=1}^N \alpha_n}.$$

Finally, we take expectations on both sides, hoping that the last term on the RHS will vanish. Although we will show this to be true, the reason is *not* because $\mathbb{E}[C^{-1} \widehat{g}_n] = C^{-1} \mathbb{E}[\widehat{g}_n]$. Recall that C^{-1} is a possibly unbounded operator, meaning that it does not necessarily commute with expectations, so we must proceed with care. The result we will use is the following:

Lemma 4. *If $\mathbb{E}_n[\|\widehat{g}_n\|_C^2] < +\infty$, then for any $h' \in C^{\frac{1}{2}}(\mathcal{H})$ we have*

$$\mathbb{E}_n[\langle h', C^{-1} \widehat{g}_n \rangle] = \langle h', g_n \rangle.$$

Proof. Analogously to the proof of Proposition 1, we will start by showing that $\langle h', C^{-1} \widehat{g}_n \rangle \in L^1(\mathcal{S}, \mathcal{F}, \mathbb{P}_n)$, where \mathbb{P}_n is the conditional measure given h_1, \dots, h_n . Expanding h' and \widehat{g}_n over the basis \mathcal{E} we get

$$\mathbb{E}_n[|\langle h', C^{-1} \widehat{g}_n \rangle|] = \mathbb{E}_n \left[\left| \sum_{i=1}^{\infty} \frac{\langle h', e_i \rangle \langle \widehat{g}_n, e_i \rangle}{\gamma_i} \right| \right] \leq \mathbb{E}_n \left[\sum_{i=1}^{\infty} \frac{|\langle h', e_i \rangle|}{\sqrt{\gamma_i}} \frac{|\langle \widehat{g}_n, e_i \rangle|}{\sqrt{\gamma_i}} \right].$$

Now, use Cauchy-Schwarz over the space of square-summable sequences to obtain

$$\sum_{i=1}^{\infty} \frac{|\langle h', e_i \rangle|}{\sqrt{\gamma_i}} \frac{|\langle \hat{g}_n, e_i \rangle|}{\sqrt{\gamma_i}} \leq \left(\sum_{i=1}^{\infty} \frac{\langle h', e_i \rangle^2}{\gamma_i} \right)^{\frac{1}{2}} \left(\sum_{i=1}^{\infty} \frac{\langle \hat{g}_n, e_i \rangle^2}{\gamma_i} \right)^{\frac{1}{2}} = \|h'\|_C \|\hat{g}_n\|_C.$$

Therefore,

$$\mathbb{E}_n[|\langle h', C^{-1}\hat{g}_n \rangle|] \leq \|h'\|_C \mathbb{E}_n[\|\hat{g}_n\|_C] \leq \|h'\|_C \mathbb{E}_n[\|\hat{g}_n\|_C^2]^{\frac{1}{2}} < +\infty,$$

by assumption. Thus, we can condition on K and use the fact that $\hat{g}_n \in \mathcal{B}_K$ to write

$$\begin{aligned} \mathbb{E}_n[\langle h', C^{-1}\hat{g}_n \rangle] &= \mathbb{E}_n \left[\mathbb{E}_n \left[\sum_{i=1}^{\infty} \frac{\langle h', e_i \rangle \langle \hat{g}_n, e_i \rangle}{\gamma_i} \mid K \right] \right] \\ &= \mathbb{E}_n \left[\mathbb{E}_n \left[\sum_{i=1}^K \frac{\langle h', e_i \rangle \langle \hat{g}_n, e_i \rangle}{\gamma_i} \mid K \right] \right] = \mathbb{E}_n \left[\sum_{i=1}^K \frac{\langle h', e_i \rangle \mathbb{E}_n[\langle \hat{g}_n, e_i \rangle \mid K]}{\gamma_i} \right]. \end{aligned}$$

Now recall that

$$\mathbb{E}_n[\langle \hat{g}_n, e_i \rangle \mid K] = \lambda_K \langle \mathbf{T}_K^{-1} Q_K^* g_n, Q_K^* e_i \rangle = \frac{\lambda_K \mathbf{1}\{K \geq i\}}{t_i} \langle g_n, e_i \rangle.$$

Thus, we have

$$\mathbb{E}_n[\langle h', C^{-1}\hat{g}_n \rangle] = \mathbb{E}_n \left[\sum_{i=1}^{\infty} \frac{\lambda_K \mathbf{1}\{K \geq i\}}{t_i \gamma_i} \langle h', e_i \rangle \langle g_n, e_i \rangle \right].$$

Using Fubini and the definition of γ_i as $\mathbb{E}[\lambda_K \mid K \geq i]$, we obtain

$$\mathbb{E}_n[\langle h', C^{-1}\hat{g}_n \rangle] = \sum_{i=1}^{\infty} \langle h', e_i \rangle \langle g_n, e_i \rangle = \langle h', g_n \rangle,$$

which finishes the proof of Lemma 4. \square

Going back to the proof of Theorem 1, since $h \in C^{\frac{1}{2}}(\mathcal{H})$ by assumption, we have $h_n - h \in C^{\frac{1}{2}}(\mathcal{H})$ for all n , letting us apply Lemma 4 to conclude that

$$\mathbb{E}[\langle h_n - h, g_n - C^{-1}\hat{g}_n \rangle] = \mathbb{E}[\mathbb{E}_n[\langle h_n - h, g_n - C^{-1}\hat{g}_n \rangle]] = 0.$$

A.3 Proof of Theorem 2

To simplify the notation in this proof, we will drop the subscript n and write \hat{g} instead of \hat{g}_n . Start by using the symmetry of $C^{-\frac{1}{2}}$, Tonelli and conditioning on K to obtain

$$\mathbb{E}[\|\hat{g}\|_C^2] = \mathbb{E} \left[\sum_{i=1}^{\infty} \langle C^{-\frac{1}{2}}\hat{g}, e_i \rangle^2 \right] = \sum_{i=1}^{\infty} \mathbb{E} \left[\frac{\lambda_K^2}{\mathbb{E}[\lambda_K \mid K \geq i]} \mathbb{E}[\langle \bar{g}, e_i \rangle^2 \mid K] \right], \quad (34)$$

where we define

$$\bar{g} \triangleq \lambda_K^{-1} \hat{g} = \frac{1}{M_K} \sum_{m=1}^M \langle g, v_m \rangle v_m. \quad (35)$$

Our goal now is to compute $\mathbb{E}[\langle \bar{g}, e_i \rangle^2 \mid K]$. Expanding the square and using the fact that v_1, \dots, v_{M_K} are i.i.d. given K , one may verify that

$$\mathbb{E}[\langle \bar{g}, e_i \rangle^2 \mid K] = \mathbb{E} \left[\left\langle \frac{1}{M_K} \sum_{m=1}^{M_K} \langle g, v_m \rangle v_m, e_i \right\rangle^2 \mid K \right]$$

$$= \frac{1}{M_K} \mathbb{E}[\langle g, v \rangle^2 \langle v, e_i \rangle^2 \mid K] + \left(1 - \frac{1}{M_K}\right) \frac{\mathbf{1}\{K \geq i\}}{t_i^2} \langle g, e_i \rangle^2. \quad (36)$$

To compute $\mathbb{E}[\langle g, e_i \rangle^2 \langle v, e_i \rangle^2 \mid K]$ we start by rewriting:

$$\begin{aligned} \langle g, e_i \rangle^2 \langle v, e_i \rangle^2 &= \langle (v \otimes v)(g \otimes g)(v \otimes v)e_i, e_i \rangle \\ &= \langle (Q\mathbf{T}^{-\frac{1}{2}}\mathbf{z} \otimes Q\mathbf{T}^{-\frac{1}{2}}\mathbf{z})(g \otimes g)(Q\mathbf{T}^{-\frac{1}{2}}\mathbf{z} \otimes Q\mathbf{T}^{-\frac{1}{2}}\mathbf{z})e_i, e_i \rangle \\ &= \langle Q\mathbf{T}^{-\frac{1}{2}}(\mathbf{z} \otimes \mathbf{z})(\mathbf{T}^{-\frac{1}{2}}Q^*g \otimes \mathbf{T}^{-\frac{1}{2}}Q^*g)(\mathbf{z} \otimes \mathbf{z})Q^*\mathbf{T}^{-\frac{1}{2}}e_i, e_i \rangle \\ &= \langle (\mathbf{z} \otimes \mathbf{z})\mathbf{L}(\mathbf{z} \otimes \mathbf{z})\mathbf{x}, \mathbf{x} \rangle_{\mathbb{R}^K}, \end{aligned}$$

where $\mathbf{L} = \mathbf{T}^{-\frac{1}{2}}Q^*g \otimes \mathbf{T}^{-\frac{1}{2}}Q^*g \in \mathbb{R}^{K \times K}$ and $\mathbf{x} = \mathbf{T}^{-\frac{1}{2}}Q^*e_i \in \mathbb{R}^K$. Hence,

$$\mathbb{E}[\langle g, e_i \rangle^2 \langle v, e_i \rangle^2 \mid K] = \langle \mathbb{E}[(\mathbf{z} \otimes \mathbf{z})\mathbf{L}(\mathbf{z} \otimes \mathbf{z}) \mid K] \mathbf{x}, \mathbf{x} \rangle.$$

Now, since $\mathbf{z} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_K)$ and \mathbf{L} is symmetric, one may show that the following fourth-moment identity holds:

$$\mathbb{E}[(\mathbf{z} \otimes \mathbf{z})\mathbf{L}(\mathbf{z} \otimes \mathbf{z}) \mid K] = (\text{tr } \mathbf{L})\mathbf{I}_K + 2\mathbf{L}.$$

This implies:

$$\mathbb{E}[\langle g, e_i \rangle^2 \langle v, e_i \rangle^2 \mid K] = \langle ((\text{tr } \mathbf{L})\mathbf{I}_K + 2\mathbf{L})\mathbf{x}, \mathbf{x} \rangle_{\mathbb{R}^K} = (\text{tr } \mathbf{L})\|\mathbf{x}\|_{\mathbb{R}^K}^2 + 2\langle \mathbf{L}\mathbf{x}, \mathbf{x} \rangle_{\mathbb{R}^K}.$$

Substituting \mathbf{L} and \mathbf{x} , we obtain

$$\mathbb{E}[\langle g, e_i \rangle^2 \langle v, e_i \rangle^2 \mid K] = \left(\sum_{j=1}^K \frac{\langle g, e_j \rangle^2}{t_j} \right) \frac{\mathbf{1}\{K \geq i\}}{t_i} + 2 \frac{\mathbf{1}\{K \geq i\}}{t_i^2} \langle g, e_i \rangle^2. \quad (37)$$

Substituting in (36), we find that

$$\mathbb{E}[\langle \bar{g}, e_i \rangle^2 \mid K] = \frac{1}{M_K} \left(\sum_{j=1}^K \frac{\langle g, e_j \rangle^2}{t_j} \right) \frac{\mathbf{1}\{K \geq i\}}{t_i} + \left(1 + \frac{1}{M_K}\right) \frac{\mathbf{1}\{K \geq i\}}{t_i^2} \langle g, e_i \rangle^2. \quad (38)$$

Now we simply substitute back in (34) and rearrange summations to obtain the desired result.

A.4 Proof of Corollary 3

Let $\psi : \mathbb{R}_+^N \rightarrow \mathbb{R}$ be given by the RHS of (17), that is:

$$\psi(\boldsymbol{\alpha}) = \frac{\frac{1}{2}\|h - h_1\|_C^2}{\|\boldsymbol{\alpha}\|_1} + (1 + 2c)G^2 \frac{\|\boldsymbol{\alpha}\|_2^2}{\|\boldsymbol{\alpha}\|_1},$$

where $\boldsymbol{\alpha} = (\alpha_1, \dots, \alpha_N)$ and $\|\boldsymbol{\alpha}\|_p$ denotes the p -th norm of $\boldsymbol{\alpha}$. The following facts about ψ are easily checked:

- (1) ψ is convex;
- (2) ψ is symmetric, in the sense that $\psi \circ \sigma = \psi$ for any coordinate permutation σ ;
- (3) ψ is coercive, meaning there exists $\tilde{c} > 0$ such that $\psi(\boldsymbol{\alpha}) \geq \tilde{c}\|\boldsymbol{\alpha}\|_1$, implying that $\psi(\boldsymbol{\alpha}) \rightarrow +\infty$ as $\|\boldsymbol{\alpha}\|_1 \rightarrow +\infty$.

Therefore, letting $S \subset \mathbb{R}_+^N$ be the set where ψ achieves its minimum, we know that S is a nonempty, convex set that satisfies $\sigma(S) \subset S$ for any coordinate permutation σ . Thus, S contains a point whose coordinates are all equal. Substituting $\alpha_n \equiv \alpha(N)$ in the RHS of Equation (17) and minimizing over $\alpha(N)$, we obtain (19).

A.5 Proof of Proposition 3

The fact that \mathcal{B} forms a linearly independent set follows directly from the positive-definiteness of the kernel $\text{Matern}_{\nu,\eta}$. To show \mathcal{B} has dense span, the main result we will use is the following theorem, which is an easy consequence of [16, Equation (4.15)], [23, Corollary 10.48] (see also the discussion before Corollary 11.33) and the extension theorem for fractional Sobolev spaces [9, Theorem A.4]:

Theorem 4. *Let $\text{Matern}_{\nu,\eta}$ be the Matérn kernel on $\Omega \subset \mathbb{R}^d$, with smoothness parameter ν , and let $s = \nu + d/2$. Then, the RKHS $\mathcal{H}_{\nu,\eta}$ generated by $\text{Matern}_{\nu,\eta}$ is equivalent to $\mathbb{H}^s(\Omega)$, meaning that $\mathcal{H}_{\nu,\eta} = \mathbb{H}^s(\Omega)$ as a set of functions and their norms are equivalent: there exist constants c_1 and c_2 such that*

$$c_1 \|h\|_{\mathbb{H}^s(\Omega)} \leq \|h\|_{\mathcal{H}_{\nu,\eta}} \leq c_2 \|h\|_{\mathbb{H}^s(\Omega)} \quad \text{for all } h \in \mathcal{H}_{\nu,\eta}.$$

Now we will prove Proposition 3. We start with some notation: If $(\mathcal{E}, \|\cdot\|_{\mathcal{E}})$ is a given normed space and $E \subset \mathcal{E}$, we denote by $\text{cl}(E; \mathcal{E})$ the closure of E with respect to $\|\cdot\|_{\mathcal{E}}$. In this notation, we want to show that, under the assumptions of Proposition 3, we have $\text{cl}(\text{span } \mathcal{B}; \mathbb{H}^\ell(\Omega)) = \mathbb{H}^\ell(\Omega)$. The idea will be to show that this closure actually contains a Sobolev space of order at least ℓ , which will be dense in $\mathbb{H}^\ell(\Omega)$. We divide the proof into four steps:

First, let $s = \nu + d/2$, which is greater or equal to ℓ , by assumption. We claim that $\text{cl}(\text{span } \mathcal{B}, \mathbb{H}^\ell(\Omega)) \supset \text{cl}(\text{span } \mathcal{B}, \mathbb{H}^s(\Omega))$. Indeed, since the inclusion $\mathbb{H}^s(\Omega) \hookrightarrow \mathbb{H}^\ell(\Omega)$ is continuous [9, Theorem 3.27], convergence in $\mathbb{H}^s(\Omega)$ -norm implies convergence in $\mathbb{H}^\ell(\Omega)$ -norm. Next, by Theorem 4, we have $\text{cl}(\text{span } \mathcal{B}, \mathbb{H}^s(\Omega)) = \text{cl}(\text{span } \mathcal{B}, \mathcal{H}_{\nu,\eta})$. Thirdly, let $\tilde{\mathcal{B}} = \{\Phi(\mathbf{x})\}_{\mathbf{x} \in \Omega}$. We then have $\text{cl}(\text{span } \mathcal{B}, \mathcal{H}_{\nu,\eta}) = \text{cl}(\text{span } \tilde{\mathcal{B}}, \mathcal{H}_{\nu,\eta}) = \mathcal{H}_{\nu,\eta}$. Indeed, for $\mathbf{x} \in \Omega$, let $\{\mathbf{y}_i\}_{i \in \mathbb{N}} \subset \{\mathbf{x}_i\}_{i \in \mathbb{N}}$ be such that $\mathbf{y}_i \rightarrow \mathbf{x}$ as $i \rightarrow +\infty$. Now compute:

$$\begin{aligned} & \|\text{Matern}_{\nu,\eta}(\mathbf{x}, \cdot) - \text{Matern}_{\nu,\eta}(\mathbf{y}_i, \cdot)\|_{\mathcal{H}_{\nu,\eta}}^2 \\ &= \langle \text{Matern}_{\nu,\eta}(\mathbf{x}, \cdot), \text{Matern}_{\nu,\eta}(\mathbf{x}, \cdot) \rangle + \langle \text{Matern}_{\nu,\eta}(\mathbf{y}_i, \cdot), \text{Matern}_{\nu,\eta}(\mathbf{y}_i, \cdot) \rangle \\ &\quad - 2 \langle \text{Matern}_{\nu,\eta}(\mathbf{x}, \cdot), \text{Matern}_{\nu,\eta}(\mathbf{y}_i, \cdot) \rangle \\ &= \text{Matern}_{\nu,\eta}(\mathbf{x}, \mathbf{x}) + \text{Matern}_{\nu,\eta}(\mathbf{y}_i, \mathbf{y}_i) - 2 \text{Matern}_{\nu,\eta}(\mathbf{x}, \mathbf{y}_i), \end{aligned}$$

where we have used the reproducing property of $\mathcal{H}_{\nu,\eta}$ in the last step. Now, note that $\text{Matern}_{\nu,\eta}$ is continuous in $\Omega \times \Omega$, which implies that $\lim \text{Matern}_{\nu,\eta}(\mathbf{y}_i, \mathbf{y}_i) = \lim \text{Matern}_{\nu,\eta}(\mathbf{x}, \mathbf{y}_i) = \text{Matern}_{\nu,\eta}(\mathbf{x}, \mathbf{x})$, thus $\|\text{Matern}_{\nu,\eta}(\mathbf{x}, \cdot) - \text{Matern}_{\nu,\eta}(\mathbf{y}_i, \cdot)\|_{\mathcal{H}_{\nu,\eta}}^2$ goes to zero as $i \rightarrow +\infty$. This proves the first equality. The second one is the known fact that an RKHS can be constructed as the closure of the span of the image of the feature map with respect to the RKHS norm, see [23, Section 10.2]. Finally, we have shown that $\mathbb{H}^s(\Omega) = \mathcal{H}_{\nu,\eta} \subset \text{cl}(\text{span } \mathcal{B}, \mathbb{H}^\ell(\Omega))$. Now, recall that since Ω is open, bounded and has a Lipschitz boundary, we know that $C^\infty(\bar{\Omega})$ is a dense subset of $\mathbb{H}^t(\Omega)$ for every $t \geq 0$ [9, Theorem 3.29]. Therefore, $\text{cl}(\mathbb{H}^s(\Omega), \mathbb{H}^\ell(\Omega)) \supset \text{cl}(C^\infty(\bar{\Omega}), \mathbb{H}^\ell(\Omega)) = \mathbb{H}^\ell(\Omega)$.

A.6 Proof of Proposition 4

Start by noting that

$$|\tilde{H}(x, p)| \leq x^2 + B\bar{c}^2 + \bar{c}|p| \quad (39)$$

for any $x, p \in \mathbb{R}$. Hence, if $h \in \mathbb{H}^1(\Omega)$, then the map $(t, x) \mapsto \tilde{H}(x, \partial_x h(t, x))$ is a member of $L^2(\Omega)$, implying that $\mathcal{R}(h)$ is well-defined for any $h \in \mathbb{H}^1(\Omega)$. Furthermore, it is easy to see that \tilde{H} is concave in p , implying that \mathcal{R} is not a convex function of h .

To show \mathcal{R} is Fréchet differentiable, we first note that the partial derivative of \tilde{H} with respect to p is given by

$$\partial_p \tilde{H}(x, p) = \begin{cases} -\frac{p}{2B}, & \text{if } |p| \leq 2B\bar{c}, \\ -\bar{c} \text{sign}(p), & \text{if } |p| > 2B\bar{c}, \end{cases}$$

where $\text{sign}(p)$ is the sign function, that is, $\text{sign}(p) = p/|p|$ for $p \neq 0$ and 0 for $p = 0$. Thus, $\partial_p \tilde{H}(x, p)$ is continuous and bounded by \bar{c} in absolute value. Now we take arbitrary $h, u \in \mathbb{H}^1(\Omega)$ and $\delta \in \mathbb{R}$ to compute

$$\mathcal{R}(h + \delta u)$$

$$\begin{aligned}
&= \frac{1}{2} \int_0^T \int_{-\bar{x}}^{\bar{x}} [\partial_t[h + \delta u](t, x) + \tilde{H}(x, \partial_x[h + \delta u](t, x))]^2 dx dt \\
&\quad + \frac{1}{2} \int_{-\bar{x}}^{\bar{x}} [(h + \delta u)(t, x) - Gx^2]^2 dx \\
&= \frac{1}{2} \int_0^T \int_{-\bar{x}}^{\bar{x}} [\partial_t[h + \delta u](t, x) + \tilde{H}(x, \partial_x h(t, x)) + \delta \partial_p \tilde{H}(x, p_\delta(t, x)) \partial_x u(t, x)]^2 dx dt \\
&\quad + \frac{1}{2} \int_{-\bar{x}}^{\bar{x}} [(h + \delta u)(t, x) - Gx^2]^2 dx,
\end{aligned}$$

where $p_\delta(t, x)$ lies between $\partial_x h(t, x)$ and $\partial_x[h + \delta u](t, x)$, and is due to the Mean Value Theorem. Now we expand:

$$\begin{aligned}
&\mathcal{R}(h + \delta u) \\
&= \mathcal{R}(h) + \delta \left\{ \int_0^T \int_{-\bar{x}}^{\bar{x}} [\partial_t h(t, x) + \tilde{H}(x, \partial_x h(t, x))] \right. \\
&\quad \cdot \left. [\partial_t u(t, x) + \partial_p \tilde{H}(x, p_\delta(t, x)) \partial_x u(t, x)] \right] dx dt \\
&\quad + \int_{-\bar{x}}^{\bar{x}} [h(t, x) - Gx^2] u(t, x) dx \Big\} \\
&\quad + \frac{\delta^2}{2} \left\{ \int_0^T \int_{-\bar{x}}^{\bar{x}} [\partial_t u(t, x) + \partial_p \tilde{H}(x, p_\delta(t, x)) \partial_x u(t, x)]^2 dx dt \right. \\
&\quad \left. + \int_{-\bar{x}}^{\bar{x}} u(t, x)^2 dx \right\},
\end{aligned}$$

obtaining:

$$\begin{aligned}
&\frac{\mathcal{R}(h + \delta u) - \mathcal{R}(h)}{\delta} \\
&= \int_0^T \int_{-\bar{x}}^{\bar{x}} [\partial_t h(t, x) + \tilde{H}(x, \partial_x h(t, x))] \\
&\quad \cdot [\partial_t u(t, x) + \partial_p \tilde{H}(x, p_\delta(t, x)) \partial_x u(t, x)] dx dt \\
&\quad + \int_{-\bar{x}}^{\bar{x}} [h(t, x) - Gx^2] u(t, x) dx \\
&\quad + \frac{\delta}{2} \left\{ \int_0^T \int_{-\bar{x}}^{\bar{x}} [\partial_p \tilde{H}(x, p_\delta(t, x)) \partial_x u(t, x)]^2 dx dt + \int_{-\bar{x}}^{\bar{x}} u(t, x)^2 dx \right\}.
\end{aligned}$$

Thus, since $\partial_p \tilde{H}$ is bounded in absolute value by 1 and also continuous, we can apply dominated convergence as we let $\delta \rightarrow 0$ to get

$$\begin{aligned}
&D\mathcal{R}(h; u) \\
&= \lim_{\delta \rightarrow 0} \frac{\mathcal{R}(h + \delta u) - \mathcal{R}(h)}{\delta} \\
&= \int_0^T \int_{-\bar{x}}^{\bar{x}} [\partial_t h(t, x) + \tilde{H}(x, \partial_x h(t, x))] \\
&\quad \cdot [\partial_t u(t, x) + \partial_p \tilde{H}(x, \partial_x h(t, x)) \partial_x u(t, x)] dx dt
\end{aligned}$$

$$+ \int_{-\bar{x}}^{\bar{x}} [h(t, x) - Gx^2]u(t, x) \, dx.$$

Again, since $\partial_p \tilde{H}$ is bounded and \tilde{H} satisfies (39), we conclude that $D\mathcal{R}(h; \cdot)$ is a bounded linear functional on $\mathbb{H}^1(\Omega)$ for any $h \in \mathbb{H}^1(\Omega)$. Furthermore, by using Cauchy-Schwarz as well as the fact that both \tilde{H} and $\partial_p \tilde{H}$ are Lipschitz continuous in p , one may show that the map $h \mapsto D(h; \cdot)$, from $\mathbb{H}^1(\Omega)$ to its dual, is also continuous. This implies the Fréchet differentiability of \mathcal{R} [13].