

REMARK ON A DETERMINANT INVOLVING PRIME NUMBERS

HUAN XIAO

ABSTRACT. Pausinger recently investigated a special determinant involving prime numbers. In this short note we point out that this type of determinants was already known in linear algebra and its computation is unrelated to prime numbers.

Let p_n be the n th prime number. Recently Pausinger [2] investigated a special determinant involving prime numbers. It is the determinant of the $n \times n$ matrix whose diagonal contains the first n prime numbers and all other elements are ones and thus

$$D_n = \begin{vmatrix} 2 & 1 & \cdots & 1 \\ 1 & 3 & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \cdots & p_n \end{vmatrix}.$$

It is the sequence A067549 of The On-Line Encyclopedia of Integer Sequences [1] and the first few values of D_n are 2, 5, 22, 140, 1448, 17856. Pausinger [2] relates this sequence to a counting problem and obtains a recurrence relation on D_n , see [2, Lemma 2].

Here in this short note we point out that this type of determinants like D_n was already known in linear algebra. Although at this moment I am unable to give a specific reference, the computation of D_n is standard and the value of it is well known. Indeed we have

Proposition 0.1. *Let $a_1 a_2 \cdots a_n \neq 0$, then*

$$d_n = \begin{vmatrix} 1 + a_1 & 1 & \cdots & 1 \\ 1 & 1 + a_2 & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \cdots & 1 + a_n \end{vmatrix} = \left(1 + \sum_{k=1}^n \frac{1}{a_k} \right) a_1 a_2 \cdots a_n.$$

Proof. This result is already known in linear algebra and for the completeness we give a proof here. Let r_i and c_i denote the i -th row and i -th column of the matrix in question. The first step is making the elementary transformations $r_i - r_1$ for $i = 2, \dots, n$, then we have

$$(0.1) \quad d_n = \begin{vmatrix} 1 + a_1 & 1 & \cdots & 1 \\ -a_1 & a_2 & & \\ \vdots & & \ddots & \\ -a_1 & & & a_n \end{vmatrix}.$$

Key words and phrases. determinant; prime numbers.

The second step is making the following elementary transformations for $i = 2, \dots, n$

$$c_1 + \frac{a_1}{a_i} c_i.$$

The resulting determinant is

$$(0.2) \quad d_n = \begin{vmatrix} b & 1 & \cdots & 1 \\ 0 & a_2 & & \\ \vdots & & \ddots & \\ 0 & & & a_n \end{vmatrix}$$

where

$$(0.3) \quad b = 1 + a_1 + a_1 \sum_{k=2}^n \frac{1}{a_k} = a_1 \left(1 + \sum_{k=1}^n \frac{1}{a_k} \right).$$

By (0.2) we have

$$(0.4) \quad d_n = \left(1 + \sum_{k=1}^n \frac{1}{a_k} \right) a_1 a_2 \cdots a_n.$$

□

Now taking $a_k = p_k - 1$ we have

Corollary 0.2.

$$(0.5) \quad D_n = \begin{vmatrix} 2 & 1 & \cdots & 1 \\ 1 & 3 & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \cdots & p_n \end{vmatrix} = \left(1 + \sum_{k=1}^n \frac{1}{p_k - 1} \right) \prod_{k=1}^n (p_k - 1).$$

REFERENCES

- [1] The Online Encyclopedia of Integer Sequences, <https://oeis.org/A067549>
- [2] Florian Pausinger, Counting appearances of integers in sets of arithmetic progressions, arXiv:2512.16358

SCHOOL OF ARTIFICIAL INTELLIGENCE, ZHUHAI CITY POLYTECHNIC, ZHUHAI, CHINA
Email address: xiaogo66@outlook.com