

AN ANDO-TYPE DILATION ON RIGHT LCM MONOIDS

BOYU LI AND MANSI SURYAWANSHI

ABSTRACT. We establish an Ando-type dilation theorem for a pair of commuting contractions together with a representation of a right LCM monoid via either the Cartesian or the free product. We prove that if each individual contraction together with the monoid representation has $*$ -regular dilation, then they can be dilated to commuting isometries and an isometric representation of the monoid. This extends an earlier result of Barik and Das.

1. INTRODUCTION

The study of dilation theory originated from the celebrated Sz.-Nagy dilation theorem [SN53, Sch55], which states that every contractive operator T on a Hilbert space \mathcal{H} admits an isometric dilation, which is isometry V on $\mathcal{K} \supset \mathcal{H}$ such that

$$P_{\mathcal{H}}V^k|_{\mathcal{H}} = T^k, \quad \text{for all } k \geq 0.$$

Extending Sz.-Nagy's dilation to the multivariate setting, Ando [And63] proved that a pair of commuting contractions can be dilated to commuting isometries. However, Ando's dilation fails to extend to three commuting contractions due to various counterexamples [Par70, Var74]. As a result, identifying additional conditions that guarantee the existence of isometric dilations has become an active and continuing line of research within multivariable operator theory.

In [Bre61], Brehmer first introduced the $*$ -regular condition for a family of commuting contractions, under which there exist commuting isometric dilations. Recently, Barik and Das [BD22] showed that for a family of commuting contractions $\{S_1, S_2, T_1, \dots, T_k\}$, suppose $\{S_i, T_1, \dots, T_k\}$ satisfies the Brehmer's $*$ -regular conditions for $i = 1, 2$, then this family of commuting contractions can be dilated to commuting isometries. This generalises earlier results of [GS97] for $k = 1$ and of [BDHS19] for a certain purity condition.

In the non-commutative setting, a family of contractions $\{T_1, \dots, T_k\}$ is called a row contraction if $\sum_{i=1}^k T_i T_i^* \leq I$. It is called a row isometry if each T_i is an isometry. It was shown in a series of papers ([Bun84, Fra82, Pop89]) that a row contraction can be dilated to a row isometry. This is now known as the Frazho-Bunce-Popescu dilation theorem.

The family of row isometries are particularly important in the study of operator algebras because of its close relationship to Cuntz algebras. In his seminal paper [Nic92], Nica introduced the notion of isometric Nica-covariant representations of quasi-lattice ordered semigroups. In the special case when the semigroup is the free semigroup \mathbb{F}_k^+ , an isometric Nica-covariant representation is determined by a family of row isometries. The C^* -algebra

Date: December 25, 2025.

2020 Mathematics Subject Classification. 47A20, 47D03, 47A13.

Key words and phrases. Ando's theorem, $*$ -regular dilation, minimal dilation.

Boyu Li was supported by an NSF grant (DMS-2350543), and he'd like to thank Kenneth Davidson for showing him the trick of proving Ando's dilation.

generated by isometric Nica-covariant representations is now known as the semigroup C^* -algebra, which has been generalized to more general semigroups [LS22, Li12]. This spurred a natural question about isometric dilations for contractive semigroup representations.

In [Li17], the first-named author observed a surprising connection between Brehmer's $*$ -regular dilation on commuting contractions and Frazho-Bunce-Popescu's isometric dilation on non-commuting row contractions. It turns out they are both special cases of dilations on the family of right-angled Artin monoids A_Γ^+ , also known as the graph product of \mathbb{N} . Building upon this discovery, one can extend Brehmer's notion of $*$ -regular dilation to representations of right-LCM monoids [Li19]. A contractive representation of a right-LCM monoid can be dilated to an isometric representation if it satisfies the generalized Brehmer's condition. Moreover, the minimal isometric $*$ -regular dilation is an isometric Nica-covariant co-extension.

Under this new perspective of $*$ -regular dilations, let us revisit the result of Barik and Das. The family $\{S_1, S_2, T_1, \dots, T_k\}$ can be viewed as a contractive representation of $\mathbb{N}^2 \times \mathbb{N}^k$, where S_1, S_2 determines a representation of S of \mathbb{N}^2 and $\{T_j\}$ determines a representation of T of \mathbb{N}^k . The main result in [BD22] can be restated as the following: suppose $S_i \times T$, as representations of $\mathbb{N} \times \mathbb{N}^k$ satisfy Brehmer's condition for each $i = 1, 2$, then one can find an isometric representation of $S \times T$ of $\mathbb{N}^2 \times \mathbb{N}^k$.

In this paper, we prove that we can replace \mathbb{N}^k by any right LCM monoid P , and the statement remains true. This establishes a new Ando-type dilation for a contractive representation of $\mathbb{N}^2 \times P$. We then considered the free product $\mathbb{N}^2 * P$ and obtained a similar Ando-type dilation result using the same technique. Our proof uses a trick of proving Ando's dilation theorem by Davidson and Katsoulis [DK11, Corollary 7.11].

2. PRELIMINARIES

A right LCM monoid P is a unital left-cancellative semigroup such that for all $p, q \in P$, $pP \cap qP$ is either \emptyset or rP for some $r \in P$. For any finite subset $U \subset P$ in a right LCM monoid, the right LCM condition implies that $\bigcap_{p \in U} pP$ is either \emptyset or rP for some $r \in P$. We use $r = \vee U$ to denote the latter case. Throughout this paper, we assume P is countable and discrete.

A contractive representation of a right LCM monoid P is a unital monoid homomorphism $T : P \rightarrow \mathcal{B}(\mathcal{H})$, where each $\|T(p)\| \leq 1$. The representation T is called isometric if each $T(p)$ is isometric. In addition, an isometric representation T is called *Nica-covariant* if for any $p, q \in P$,

$$T(p)T(p)^*T(q)T(q)^* = \begin{cases} T(r)T(r)^*, & \text{if } pP \cap qP = rP, \\ 0, & \text{otherwise.} \end{cases}$$

Isometric Nica-covariant representation was first introduced by Nica [Nic92], which plays a central role in the study of semigroup C^* -algebras [CL02, LR96, Li12, LS22]. Right LCM monoids form an important class of monoids in the study of semigroup C^* -algebras [ABLS19, aHNSY21, BLS17, BLS18, LL20, LL22, Sta15] due to their comparatively tractable Nica-covariant representations.

An isometric dilation for T is an isometric representation $V : P \rightarrow \mathcal{B}(\mathcal{K})$ on $\mathcal{K} \supset \mathcal{H}$, such that $T(p) = P_{\mathcal{H}}V(p)|_{\mathcal{H}}$ for all $p \in P$. The dilation V is called a *co-extension* of T if \mathcal{H}^\perp is invariant for V , or equivalently, \mathcal{H} is invariant for $V(p)^*$ for all $p \in P$.

The dilation V is called a *$*$ -regular dilation* if for any $p, q \in P$,

$$P_{\mathcal{H}}V(p)^*V(q)|_{\mathcal{H}} = \begin{cases} T(p^{-1}r)T(q^{-1}r)^*, & \text{if } pP \cap qP = rP, \\ 0, & \text{otherwise.} \end{cases}$$

The concept of $*$ -regular dilation originated from Brehmer's study [Bre61] on dilations of commuting contractions, which can be viewed as representations of \mathbb{N}^k . In a series of papers by the first-named author [Li16, Li17, Li19], the notion of $*$ -regular dilation was extended to general right LCM monoids, where it is shown that having $*$ -regular dilation is equivalent to the generalized Brehmer's condition [Li19, Theorem 3.9].

Notice that when V is a co-extension of T ,

$$P_{\mathcal{H}}V(a)V(b)^*|_{\mathcal{H}} = T(a)T(b)^*, \quad \text{for all } a, b \in P.$$

Therefore, if V is an isometric Nica-covariance co-extension,

$$\begin{aligned} P_{\mathcal{H}}V(p)^*V(q)|_{\mathcal{H}} &= \begin{cases} P_{\mathcal{H}}V(p^{-1}r)V(q^{-1}r)^*|_{\mathcal{H}}, & \text{if } pP \cap qP = rP, \\ 0, & \text{otherwise.} \end{cases} \\ &= \begin{cases} T(p^{-1}r)T(q^{-1}r)^*, & \text{if } pP \cap qP = rP, \\ 0, & \text{otherwise.} \end{cases} \end{aligned}$$

Hence, any isometric Nica-covariant co-extension must also be a $*$ -regular dilation.

A $*$ -regular dilation $V_0 : P \rightarrow \mathcal{B}(\mathcal{K}_0)$ is called minimal if $\mathcal{K}_0 = \overline{\text{span}}\{V_0(p)h : p \in P, h \in \mathcal{H}\}$. The minimal $*$ -regular dilation $V_0 : P \rightarrow \mathcal{B}(\mathcal{K}_0)$ is unique up to unitary equivalence. A key observation in [Li19, Theorem 3.8] is that the minimal $*$ -regular dilation $V_0 : P \rightarrow \mathcal{B}(\mathcal{K}_0)$ must be an isometric Nica-covariant co-extension.

This gives us a full characterization of isometric Nica-covariant co-extensions.

Lemma 2.1. *Let $V : P \rightarrow \mathcal{B}(\mathcal{L})$ be an isometric Nica-covariant co-extension of T . Then \mathcal{L} decomposes as $\mathcal{L} = \mathcal{K}_0 \oplus \mathcal{K}_1$ such that \mathcal{K}_i , $i = 0, 1$ reduces V . Moreover, $V|_{\mathcal{K}_0}$ is unitarily equivalent to the minimal isometric Nica-covariant dilation (hence $*$ -regular dilation) V_0 of T and $V|_{\mathcal{K}_1}$ is an isometric Nica-covariant representation.*

Proof. Given an isometric Nica-covariant co-extension $V : P \rightarrow \mathcal{B}(\mathcal{L})$ of T , define

$$\mathcal{K}_0 = \overline{\text{span}}\{V(p)h : p \in P, h \in \mathcal{H}\}.$$

It is clear that \mathcal{K}_0 is invariant for V . By the Nica-covariance condition, $V(p)^*V(q)h \in \mathcal{K}_0$ for all $p, q \in P$ and $h \in \mathcal{H}$, and hence \mathcal{K}_0 is also invariant for V^* . Therefore, \mathcal{K}_0 reduces V , and $\mathcal{L} = \mathcal{K}_0 \oplus \mathcal{K}_1$ for some Hilbert space \mathcal{K}_1 .

The restriction $V|_{\mathcal{K}_0}$ is an isometric Nica-covariant co-extension of T satisfying $\mathcal{K}_0 = \overline{\text{span}}\{V(p)h : p \in P, h \in \mathcal{H}\}$, and hence is minimal. Consequently, $V|_{\mathcal{K}_0}$ is a minimal $*$ -regular dilation of T , and by uniqueness of minimal $*$ -regular dilations, it is unitarily equivalent to V_0 . Therefore,

$$V \cong V_0 \oplus X,$$

where V_0 is the minimal $*$ -regular dilation of T and $X := V|_{\mathcal{K}_1}$ is an isometric Nica-covariant representation, since V is isometric and Nica-covariant. \square

Suppose P, Q are two right LCM monoids. We use $Q \times P$ to denote their Cartesian product, and $Q * P$ to denote their free product. Both $Q \times P$ and $Q * P$ are also right LCM monoids [FK09]. Given representations $S : Q \rightarrow \mathcal{B}(\mathcal{H})$ and $T : P \rightarrow \mathcal{B}(\mathcal{H})$, we define

$S \times T$ to be the representation of $Q \times P$ that maps (q, p) to $S(q)T(p)$. Since (q, e) and (e, p) commute in the Cartesian product, $S \times T$ defines a representation of $Q \times P$ if $S(q)$ commutes with $T(p)$ for all $q \in Q$ and $p \in P$. Similarly, define $S * T$ to be the representation of $Q * P$ that maps $q \in Q$ to $S(q)$ and $p \in P$ to $T(p)$. Since elements of Q and P do not need to commute in $Q * P$, $S * T$ is always a representation of $Q * P$.

For the rest of this paper, we focus on the case where Q is either \mathbb{N} or \mathbb{N}^2 . Let P be a right-LCM monoid and $T : P \rightarrow \mathcal{B}(\mathcal{H})$ be a representation. Suppose $S \in \mathcal{B}(\mathcal{H})$ commutes with all $T(p), p \in P$. One can treat S as a representation of \mathbb{N} by mapping each $k \geq 0$ to S^k . Since S commutes with T , one can define the representation $S \times T : \mathbb{N} \times P \rightarrow \mathcal{B}(\mathcal{H})$ by

$$S \times T(n, p) = S^n T(p).$$

Now suppose S_1, S_2 are two commuting contractions that both commute with T . Since S_1, S_2 commute, they define a representation $S_1 \times S_2$ of \mathbb{N}^2 by

$$(S_1 \times S_2)(n, m) = S_1^n S_2^m.$$

From here, we can similarly define the representation $(S_1 \times S_2) \times T$ of $\mathbb{N}^2 \times P$ by

$$(S_1 \times S_2) \times T(n, m, p) = S_1^n S_2^m T(p).$$

Recall, two operators A, B doubly commute if $AB = BA$ and $A^*B = BA^*$. Suppose B is a representation of P . We say A and B doubly commute if A and $B(p)$ doubly commute for each $p \in P$. The following Lemma is a well-known fact about isometric Nica-covariant representations of $\mathbb{N} \times P$.

Lemma 2.2. *Let $V \in \mathcal{B}(\mathcal{K})$ be an isometry and $W : P \rightarrow \mathcal{B}(\mathcal{K})$ be an isometric Nica-covariant representation of P . Then $V \times W$ is an isometric Nica-covariant representation of $\mathbb{N} \times P$ if and only if V and W doubly commute.*

Finally, for any contraction $S \in \mathcal{B}(\mathcal{H})$ and representation $T : P \rightarrow \mathcal{B}(\mathcal{H})$, one can always define $S * T$ to be the representation that maps the elements in P to $T(p)$ and the generator for \mathbb{N} to S . The following Lemma characterizes isometric Nica-covariant representations of $\mathbb{N} * P$.

Lemma 2.3. *Let $V \in \mathcal{B}(\mathcal{K})$ be an isometry and $W : P \rightarrow \mathcal{B}(\mathcal{K})$ be an isometric Nica-covariant representation of P . Let P_0 denote the set of all invertible elements in P . Then, $V * W$ is an isometric Nica-covariant representation of $\mathbb{N} * P$ if and only if V and $W(p)$ have orthogonal ranges for all $p \notin P_0$, or equivalently, $VV^* + W(p)W(p)^* \leq I$.*

3. MAIN RESULTS

The main goal of this paper is the following theorem.

Theorem 3.1. *Let P be a right LCM monoid and let $(S_1 \times S_2) \times T$ be a contractive representation of $\mathbb{N}^2 \times P$ on \mathcal{H} . Suppose that each $S_i \times T$ has a $*$ -regular dilation for $i = 1, 2$. Then there exists an isometric dilation $(V_1 \times V_2) \times W$ of $(S_1 \times S_2) \times T$. Moreover, for $i = 1, 2$, the representations $V_i \times W$ can be chosen to be isometric Nica-covariant co-extensions of $S_i \times T$.*

Proof. Let $A_i \times B_i$ be the minimal $*$ -regular dilations of $S_i \times T$, acting on \mathcal{M}_i , for $i = 1, 2$. By [Li19], $A_i \times B_i$ are isometric Nica-covariant co-extensions.

By restricting to P , the representations $B_i : P \rightarrow \mathcal{B}(\mathcal{M}_i)$ are isometric Nica-covariant co-extensions of T . Let $W_0 : P \rightarrow \mathcal{B}(\mathcal{M}_0)$ be the minimal isometric Nica-covariant dilation of T . By Lemma 2.1, for each $i = 1, 2$, there exists a reducing subspace $\mathcal{M}_i^{(0)} \subseteq \mathcal{M}_i$ such

that $B_i|_{\mathcal{M}_i^{(0)}} \cong W_0$. Replacing B_i by a unitarily equivalent representation, we may assume that $\mathcal{M}_i^{(0)} = \mathcal{M}_0$ and hence that

$$\mathcal{M}_i = \mathcal{M}_0 \oplus \mathcal{N}_i, \quad B_i = W_0 \oplus Z_i,$$

for some isometric representation $Z_i : P \rightarrow \mathcal{B}(\mathcal{N}_i)$. Since B_i is Nica-covariant, so is Z_i .

Now define $\mathcal{K} = \mathcal{M}_0 \oplus \mathcal{N}_1 \oplus \mathcal{N}_2$. Since $\mathcal{M}_i := \mathcal{M}_0 \oplus \mathcal{N}_i$, we may identify $\mathcal{K} \cong \mathcal{M}_1 \oplus \mathcal{N}_2 \cong \mathcal{M}_2 \oplus \mathcal{N}_1$. Define

$$\begin{aligned} V_1 &= A_1 \oplus I_{\mathcal{N}_2} \in \mathcal{B}(\mathcal{M}_1 \oplus \mathcal{N}_2) \cong \mathcal{B}(\mathcal{K}), \\ V_2 &= A_2 \oplus I_{\mathcal{N}_1} \in \mathcal{B}(\mathcal{M}_2 \oplus \mathcal{N}_1) \cong \mathcal{B}(\mathcal{K}), \\ W &= W_0 \oplus Z_1 \oplus Z_2 \in \mathcal{B}(\mathcal{M}_0 \oplus \mathcal{N}_1 \oplus \mathcal{N}_2) = \mathcal{B}(\mathcal{K}). \end{aligned}$$

Equivalently, V_1 acts as A_1 on $\mathcal{M}_0 \oplus \mathcal{N}_1$ and as the identity on \mathcal{N}_2 , while V_2 acts as A_2 on $\mathcal{M}_0 \oplus \mathcal{N}_2$ and as the identity on \mathcal{N}_1 . Since $W : P \rightarrow \mathcal{B}(\mathcal{K})$ is an isometric representation and satisfies $P_{\mathcal{H}}W(p)|_{\mathcal{H}} = P_{\mathcal{H}}W_0(p)|_{\mathcal{H}} = T(p)$, for all $p \in P$, it is an isometric dilation of T . Since $\mathcal{H} \subset \mathcal{M}_0$, V_i is an isometric dilation of S_i , for $i = 1, 2$. For $i \neq i'$, under the identification $\mathcal{K} \cong \mathcal{M}_i \oplus \mathcal{N}_{i'}$ we have

$$V_i \times W = (A_i \times B_i) \oplus (I_{\mathcal{N}_{i'}} \times Z_{i'}).$$

Since the identity representation $I_{\mathcal{N}_{i'}}$ doubly commutes with $Z_{i'}$ trivially, it follows that the product representation $I_{\mathcal{N}_{i'}} \times Z_{i'}$ satisfies the Nica covariance relations. As $A_i \times B_i$ is an isometric Nica-covariant co-extension of $S_i \times T$, and the summand $I_{\mathcal{N}_{i'}} \times Z_{i'}$ is isometric Nica-covariant representation. It follows that $V_i \times W$ is an isometric Nica-covariant co-extension of $S_i \times T$, for $i = 1, 2$.

Since $V_i \times W$ is Nica-covariant, V_i and W doubly commute by Lemma 2.2. Consequently, both V_1V_2 and V_2V_1 doubly commute with W . By Lemma 2.2, both $(V_1V_2) \times W$ and $(V_2V_1) \times W$ are isometric Nica-covariant representations of $\mathbb{N} \times P$. Since $V_i \times W$ are co-extensions of $S_i \times T$, \mathcal{H}^\perp is invariant for V_i and W and thus both $(V_1V_2) \times W$ and $(V_2V_1) \times W$ are isometric Nica-covariant co-extensions of $(S_2S_1) \times T = (S_1S_2) \times T$.

Again, by Lemma 2.1, we can decompose $\mathcal{K} = \mathcal{K}_1 \oplus \mathcal{K}_2$ into reducing subspaces, such that the restrictions of $(V_1V_2) \times W$ and $(V_2V_1) \times W$ on \mathcal{K}_1 are unitarily equivalent to the minimal isometric Nica-covariant dilation $M \times W_1$ for $(S_1S_2) \times T$, and $\mathcal{H} \subset \mathcal{K}_1$. The restrictions of $(V_1V_2) \times W$ and $(V_2V_1) \times W$ on \mathcal{K}_2 are some isometric Nica-covariant representation, and therefore, we may write

$$\begin{aligned} (V_1V_2) \times W &= (M \times W_1) \oplus (X \times W_2), \\ (V_2V_1) \times W &= (M \times W_1) \oplus (Y \times W_2), \end{aligned}$$

Here, X and Y are isometries acting on \mathcal{K}_2 , while $W_2 := W|_{\mathcal{K}_2}$. Moreover, $X \times W_2$ and $Y \times W_2$ are the restrictions of $(V_1V_2) \times W$ and $(V_2V_1) \times W$ on \mathcal{K}_2 .

Let $\mathcal{K}_2^\infty \cong \mathcal{K}_2 \otimes \ell^2(\mathbb{N})$ denote the infinite direct sum of \mathcal{K}_2 , and set $\mathcal{L} = \mathcal{K} \oplus \mathcal{K}_2^\infty \oplus \mathcal{K}_2^\infty$. Define operators on \mathcal{L} by

$$\begin{aligned} V'_1 &= V_1 \oplus X^\infty \oplus Y^\infty, \\ V'_2 &= V_2 \oplus I_{\mathcal{K}_2^\infty} \oplus I_{\mathcal{K}_2^\infty}, \\ W' &= W \oplus W_2^\infty \oplus W_2^\infty, \end{aligned}$$

where X^∞ (respectively, Y^∞) denotes the infinite ampliation of X (respectively, Y) acting on \mathcal{K}_2^∞ . Since $V_i \times W$ is an isometric Nica-covariant co-extension of $S_i \times T$, and since $X \times W_2$,

$Y \times W_2$, and $I_{\mathcal{K}_2} \times W_2$ are isometric Nica-covariant representations of $\mathbb{N} \times P$, it follows that $V'_i \times W'$ are isometric Nica-covariant representations of $\mathbb{N} \times P$. Moreover, as $\mathcal{H} \subset \mathcal{K} \subset \mathcal{L}$ and the additional summands in $V'_i \times W'$ act on subspaces orthogonal to \mathcal{H} , each $V'_i \times W'$ is an isometric Nica-covariant co-extension of $S_i \times T$.

With respect to the decomposition $\mathcal{L} = \mathcal{K}_1 \oplus \mathcal{K}_2 \oplus \mathcal{K}_2^\infty \oplus \mathcal{K}_2^\infty$, the operators $V'_1 V'_2$ and $V'_2 V'_1$ admit the block-diagonal forms

$$V'_1 V'_2 = (M \oplus X) \oplus X^\infty \oplus Y^\infty \quad \text{and} \quad V'_2 V'_1 = (M \oplus Y) \oplus X^\infty \oplus Y^\infty.$$

Let $F \in \mathcal{B}(\mathcal{K}_2 \oplus \mathcal{K}_2^\infty \oplus \mathcal{K}_2^\infty)$ be given by

$$F(k, \xi_1, \xi_2, \dots, \eta_1, \eta_2, \dots) = (\eta_1, k, \xi_1, \xi_2, \dots, \eta_2, \eta_3, \dots).$$

Define $U = I_{\mathcal{K}_1} \oplus F \in \mathcal{B}(\mathcal{L})$. Then a direct computation shows that,

$$U^*(V'_2 V'_1) U = V'_1 V'_2.$$

By construction, \mathcal{K}_1 reduces U and $U|_{\mathcal{K}_1} = I_{\mathcal{K}_1}$. Since $\mathcal{H} \subset \mathcal{K}_1$, it follows that \mathcal{H} also reduces U and $U|_{\mathcal{H}} = I_{\mathcal{H}}$.

Now define $\tilde{V}_1 := V'_1 U$, and $\tilde{V}_2 := U^* V'_2$. Then

$$\tilde{V}_1 \tilde{V}_2 = (V'_1 U)(U^* V'_2) = V'_1 V'_2 = U^*(V'_2 V'_1) U = \tilde{V}_2 \tilde{V}_1.$$

Hence, \tilde{V}_1 and \tilde{V}_2 commute. Moreover, since U is unitary and V'_1, V'_2 are isometries, it follows that \tilde{V}_1 and \tilde{V}_2 are also isometries. Since \tilde{V}_i and V'_i coincide on the reducing subspace \mathcal{K}_1 , \tilde{V}_i are isometric dilations of S_i for $i = 1, 2$.

Now, with respect to $\mathcal{L} = (\mathcal{K}_1 \oplus \mathcal{K}_2) \oplus \mathcal{K}_2^\infty \oplus \mathcal{K}_2^\infty$, we have

$$W' = (W_1 \oplus W_2) \oplus W_2^\infty \oplus W_2^\infty.$$

On $\mathcal{K}_1^\perp \cong \mathcal{K}_2 \oplus \mathcal{K}_2^\infty \oplus \mathcal{K}_2^\infty \cong \mathcal{K}_2^\infty$, $W'|_{\mathcal{K}_1^\perp} \cong W_2^\infty$, which obviously doubly commutes with the permutation operator F . On \mathcal{K}_1 , $U|_{\mathcal{K}_1} = I_{\mathcal{K}_1}$ also doubly commutes with $W'|_{\mathcal{K}_1}$. Therefore, U and W' doubly commute.

Now, V'_i and U all doubly commute with W' , and therefore, by Lemma 2.2, $\tilde{V}_i \times W'$ are isometric Nica-covariant representations. Each $\tilde{V}_i \times W'$ is a co-extension of $S_i \times T$ because it contains a direct summand of the minimal isometric Nica-covariant dilation $M \times W_1$. Moreover, since \tilde{V}_1 and \tilde{V}_2 commute, we have $(\tilde{V}_1 \times \tilde{V}_2) \times W'$ is an isometric dilation of $(S_1 \times S_2) \times T$. \square

Remark 3.2. We point out that in the simplest case, when $P = \{e\}$ and $T(e) = I$, Theorem 3.1 recovers Ando's dilation theorem: two commuting contractions S_1, S_2 admit commuting isometric dilations V_1, V_2 . In fact, the main technique used in the proof of Theorem 3.1 is inspired by the proof of Ando's theorem in [DK11, Corollary 7.11].

Example 3.3. When $P = \mathbb{N}^k$, a contractive representation $T : P \rightarrow \mathcal{B}(\mathcal{H})$ is uniquely determined by a family of k commuting contractions $\{T_1, \dots, T_k\}$. In this case, Theorem 3.1 asserts that if each $\{S_i, T_1, \dots, T_k\}$ admits a *regular dilation, then the commuting contractions $\{S_1, S_2, T_1, \dots, T_k\}$ admit commuting isometric dilations $\{V_1, V_2, W_1, \dots, W_k\}$. Moreover, for each $i = 1, 2$, the family $\{V_i, W_1, \dots, W_k\}$ can be chosen to be doubly commuting isometric co-extensions of $\{S_i, T_1, \dots, T_k\}$, since for isometric representations of \mathbb{N}^k , Nica-covariance is equivalent to double commutativity of the generators. This yields a short proof of the main result in [BD22, Theorem 3.9] by identifying $S_1 = T_1$, $S_2 = T_n$, and $(T_1, \dots, T_k) = (T_2, \dots, T_{n-1})$, (see also [BDHS19]).

As another application of Theorem 3.1, we can obtain a new Ando-type dilation on $\mathbb{N}^2 \times \mathbb{F}_k^+$.

Example 3.4. Let $P = \mathbb{F}_k^+$. A contractive representation T of P is determined by a family of k (not necessarily commuting) contractions $\{T_1, \dots, T_k\}$. Let S be a contraction that commutes with all T_j . Then $S \times T$ has a $*$ -regular dilation [Li17] if and only if $\{T_1, \dots, T_k\}$ is a row contraction and

$$I - SS^* - \sum_{j=1}^k T_j T_j^* + \sum_{j=1}^k S T_j T_j^* S^* \geq 0.$$

In this case, Theorem 3.1 yields the following. Suppose $\{T_1, \dots, T_k\}$ is a row contraction and S_1, S_2 are commuting contractions such that $S_i T_j = T_j S_i$ for all $i = 1, 2$ and all j . Assume that

$$I - S_i S_i^* - \sum_{j=1}^k T_j T_j^* + \sum_{j=1}^k S_i T_j T_j^* S_i^* \geq 0, \quad i = 1, 2.$$

Then S_1 and S_2 dilate to commuting isometries V_1 and V_2 , and $\{T_j\}$ dilates to a row isometry $\{W_j\}$. Moreover, for each $i = 1, 2$, V_i doubly commutes with each W_j .

We emphasize that one cannot, in general, expect V_1 and V_2 to be doubly commuting. Indeed, arranging this additional property would amount to producing a Nica-covariant (hence $*$ -regular) isometric dilation of $(S_1 \times S_2) \times T$, which is known to be equivalent to a stronger positivity condition in [Li17].

We now consider the right LCM monoid $\mathbb{N}^2 * P$. Using the same technique as in the proof of Theorem 3.1, we obtain the following result.

Theorem 3.5. *Let P be a discrete right-LCM monoid and let $(S_1 \times S_2) * T$ be a contractive representation of $\mathbb{N}^2 * P$ on a separable Hilbert space \mathcal{H} . Suppose $S_i * T$ have $*$ -regular dilation for $i = 1, 2$. Then there exists an isometric dilation $(V_1 \times V_2) * W$ of $(S_1 \times S_2) * T$. Moreover, $V_i * W$ can be chosen to be isometric Nica-covariant co-extensions.*

Proof. Let $A_i * B_i : \mathbb{N} * P \rightarrow \mathcal{B}(\mathcal{M}_i)$ denote the $*$ -regular, and hence minimal isometric Nica-covariant dilations of $S_i * T$ for $i = 1, 2$. For $i = 1, 2$, Lemma 2.1 guarantees a decomposition $\mathcal{M}_i = \mathcal{M}_0 \oplus \mathcal{N}_i$ into reducing subspaces such that, with respect to this decomposition,

$$B_i = W_0 \oplus Z_i,$$

where $W_0 : P \rightarrow \mathcal{B}(\mathcal{M}_0)$ is the minimal isometric Nica-covariant dilation of T , and $Z_i : P \rightarrow \mathcal{B}(\mathcal{N}_i)$ is an isometric Nica-covariant representation.

Define $\mathcal{K} = \mathcal{M}_0 \oplus \mathcal{N}_1 \oplus \mathcal{N}_2 \oplus \ell^2(P)^\infty$, where $\ell^2(P)^\infty$ denotes the infinite ampliation $\ell^2(P) \otimes \ell^2(\mathbb{N})$. Let

$$W = W_0 \oplus Z_1 \oplus Z_2 \oplus \lambda^\infty \quad \text{on } \mathcal{K}.$$

Here $\ell^2(P) = \overline{\text{span}}\{e_p : p \in P\}$ denotes the Hilbert space with canonical orthonormal basis $\{e_p\}_{p \in P}$, $\lambda : P \rightarrow \mathcal{B}(\ell^2(P))$ denotes the left regular representation defined by $\lambda(p)e_q = e_{pq}$, and λ^∞ denotes its infinite amplification. The representation λ is isometric and Nica-covariant.

Let $P_0 \subset P$ denote the set of invertible elements of P , and set $\mathcal{L}_0 = \ell^2(P_0)^\infty$. Since P is a monoid, $1 \in P_0$, and hence $P_0 \neq \emptyset$. We identify $\mathcal{L}_0 = \ell^2(P_0)^\infty$ with the closed subspace of $\ell^2(P)^\infty \cong \ell^2(P) \otimes \ell^2(\mathbb{N})$ spanned by $\{e_p \otimes \delta_n : p \in P_0, n \in \mathbb{N}\}$, where $\{\delta_n\}_{n \in \mathbb{N}}$ is the canonical orthonormal basis of $\ell^2(\mathbb{N})$. It is clear that \mathcal{L}_0 is an infinite-dimensional separable

Hilbert space. Therefore, we can pick isometries $C_i : \mathcal{N}_i \oplus \ell^2(P)^\infty \rightarrow \mathcal{L}_0 \subset \mathcal{N}_i \oplus \ell^2(P)^\infty$. Define

$$V_1 = A_1 \oplus C_2, \text{ and } V_2 = A_2 \oplus C_1.$$

Then V_1, V_2 are isometries on \mathcal{K} . Moreover, if $p \notin P_0$, then $pP \cap P_0 = \emptyset$, and hence

$$\text{ran } \lambda(p) = \overline{\text{span}}\{e_{pq} : q \in P\} \perp \ell^2(P_0).$$

Consequently, $\text{ran } \lambda(p)^\infty \perp \mathcal{L}_0$, and thus, one can verify that V_i and $W(p)$ have orthogonal ranges for all $p \notin P_0$. By Lemma 2.3, both $V_i * W$ are isometric Nica-covariant co-extensions of $S_i * T$.

From here, we can follow the same construction as in the proof of Theorem 3.1. One can check that both $(V_1 V_2) * W$ and $(V_2 V_1) * W$ are isometric Nica-covariant co-extensions of $(S_1 S_2) * T = (S_2 S_1) * T$. By the existence of a minimal isometric Nica-covariant dilation $M * W_1$ for $(S_1 S_2) * T$, we can decompose $\mathcal{K} = \mathcal{K}_1 \oplus \mathcal{K}_2$ using Lemma 2.1, under which

$$\begin{aligned} (V_1 V_2) * W &= (M * W_1) \oplus (X * W_2) \\ (V_2 V_1) * W &= (M * W_1) \oplus (Y * W_2) \end{aligned}$$

Define $\mathcal{K}_2^\infty \cong \mathcal{K}_2 \otimes \ell^2(\mathbb{N})$ and set $\mathcal{L} = \mathcal{K} \oplus \mathcal{K}_2^\infty \oplus \mathcal{K}_2^\infty$. Define operators on \mathcal{L} by

$$\begin{aligned} V'_1 &= V_1 \oplus X^\infty \oplus Y^\infty, \\ V'_2 &= V_2 \oplus I_{\mathcal{K}_2^\infty} \oplus I_{\mathcal{K}_2^\infty}, \\ W' &= W \oplus W_2^\infty \oplus W_2^\infty, \end{aligned}$$

where X^∞ (resp. Y^∞) denotes the infinite ampliation of X (resp. Y) acting on the corresponding copy of \mathcal{K}_2^∞ . By the same trick as in the proof of Theorem 3.1, there exists a unitary $U \in \mathcal{B}(\mathcal{L})$ such that

$$U^*(V'_2 V'_1) U = V'_1 V'_2.$$

Moreover, $U|_{\mathcal{K}_1} = I_{\mathcal{K}_1}$ and U doubly commutes with W' .

Now $\tilde{V}_1 = V'_1 U$ and $\tilde{V}_2 = U^* V'_2$ are commuting isometries and thus $(\tilde{V}_1 \times \tilde{V}_2) * W'$ is an isometric dilation of $(S_1 \times S_2) * T$. Moreover, since U and W' doubly commute, we have $\tilde{V}_i^* W'(p) = 0$ for all $p \notin P_0$. Therefore, both $\tilde{V}_i * W'$ are isometric Nica-covariant co-extensions of $S_i * T$. \square

Remark 3.6. We remark that the infinite amplification $\ell^2(P)^\infty$ in the proof is needed to make room for the ranges of the operators C_i , since each V_i is required to have range orthogonal to that of W .

Remark 3.7. Recall, from [Li19], $T : P \rightarrow \mathcal{B}(\mathcal{H})$ has $*$ -regular dilation if and only if for all finite $F \subset P$,

$$Z_F = \sum_{U \subset F} (-1)^{|U|} T_U T_U^* \geq 0.$$

Here,

$$T_U T_U^* = \begin{cases} T_r T_r^*, & \text{if } \cap_{p \in P} pP = rP, \\ 0, & \text{if } \cap_{p \in P} pP = \emptyset. \end{cases}$$

Now, for a contraction $S \in \mathcal{B}(\mathcal{H})$, to verify $S \times T$ and $S * T$ have $*$ -regular dilations on $\mathbb{N} \times P$ and $\mathbb{N} * P$, instead of considering all finite subsets F , the positivity of a much smaller collection of Z_F is needed. One can apply [Li19, Theorem 6.7] to obtain the following characterizations.

(1) For $S \times T$ on $\mathbb{N} \times P$, $S \times T$ has $*$ -regular dilation if for all finite $F \subset P$,

$$Z_F \geq 0, \text{ and } SZ_F S^* \leq Z_F.$$

(2) For $S * T$ on $\mathbb{N} * P$, $S * T$ has $*$ -regular dilation if for all finite $F \subset P$,

$$Z_F \geq 0, \text{ and } Z_F - SS^* \geq 0.$$

Given a simple graph Γ on n vertices $\{v_1, \dots, v_n\}$, recall the right-angled Artin monoid A_Γ^+ is the monoid

$$A_\Gamma^+ = \langle e_1, \dots, e_n : e_i e_j = e_j e_i \text{ if } v_i v_j \text{ is an edge in } \Gamma \rangle.$$

For a vertex v , use $\Gamma/\{v\}$ to denote the subgraph by removing v . Our main results establish the following Ando-type dilation theorem on a class of right-angled Artin monoids.

Example 3.8. Let Γ be a simple graph on n vertices where $v_1 v_2$ is an edge. Furthermore, suppose v_1, v_2 are either adjacent to all other vertices or not adjacent to any other vertices. Under this assumption, suppose a contractive representation $T : A_\Gamma^+ \rightarrow \mathcal{B}(\mathcal{H})$ satisfies that for $i = 1, 2$, $T|_{A_{\Gamma/\{v_i\}}^+}$ have $*$ -regular dilations (as characterized by [Li17, Li25]). Then, by Theorem 3.1 and 3.5, we know T must have an isometric co-extension V . Moreover, $V|_{A_{\Gamma/\{v_i\}}^+}$ are isometric Nica-covariant co-extensions for both $i = 1, 2$.

As a special case where the right-angled Artin monoid is

$$A_\Gamma^+ = \langle e_1, e_2, f_1, \dots, f_k : e_1 e_2 = e_2 e_1 \rangle \cong \mathbb{N}^2 * \mathbb{F}_k^+,$$

we obtain the following dilation theorem on $\mathbb{N}^2 * \mathbb{F}_k^+$.

Example 3.9. Consider the special case where $P = \mathbb{F}_k^+$. Let $\{S_1, S_2, T_1, \dots, T_k\}$ be contractions with $S_1 S_2 = S_2 S_1$. Assume that, for each $i = 1, 2$, the tuple (S_i, T_1, \dots, T_k) is a row contraction. Then, by Theorem 3.5, there exist row isometric dilations (V_i, W_1, \dots, W_k) of (S_i, T_1, \dots, T_k) such that V_1 and V_2 commute.

Comparing this new dilation result with other known results in the literature, the row contraction conditions are weaker than the $*$ -regular dilation conditions derived in [Li17]. While the existence of row isometric dilation for each pair (S_i, T) individually follows from Frazho-Bunce-Popescu's dilation of row contractions, it is not clear that W_1 and W_2 can be chosen to commute with each other. We would also like to point out that a result by Opěla [Opě06] states that the family $\{S_1, S_2, T_j : 1 \leq j \leq k\}$ has isometric dilations (where S_i are dilated to commuting isometries). However, the row isometric condition is not guaranteed.

REFERENCES

- [ABLS19] Zahra Afsar, Nathan Brownlowe, Nadia S. Larsen, and Nicolai Stammeier. Equilibrium states on right LCM semigroup C^* -algebras. *Int. Math. Res. Not. IMRN*, (6):1642–1698, 2019.
- [aHNSY21] Astrid an Huef, Brita Nucinkis, Camila F. Sehnem, and Dilian Yang. Nuclearity of semigroup C^* -algebras. *J. Funct. Anal.*, 280(2):Paper No. 108793, 46, 2021.
- [And63] T. Andô. On a pair of commutative contractions. *Acta Sci. Math. (Szeged)*, 24:88–90, 1963.
- [BD22] Sibaprasad Barik and B. Krishna Das. Isometric dilations of commuting contractions and Brehmer positivity. *Complex Anal. Oper. Theory*, 16(5):Paper No. 69, 25, 2022.
- [BDHS19] Sibaprasad Barik, B. Krishna Das, Kalpesh J. Haria, and Jaydeb Sarkar. Isometric dilations and von Neumann inequality for a class of tuples in the polydisc. *Trans. Amer. Math. Soc.*, 372(2):1429–1450, 2019.
- [BLS17] Nathan Brownlowe, Nadia S. Larsen, and Nicolai Stammeier. On C^* -algebras associated to right LCM semigroups. *Trans. Amer. Math. Soc.*, 369(1):31–68, 2017.

- [BLS18] Nathan Brownlowe, Nadia S. Larsen, and Nicolai Stammeier. C^* -algebras of algebraic dynamical systems and right LCM semigroups. *Indiana Univ. Math. J.*, 67(6):2453–2486, 2018.
- [Bre61] S. Brehmer. Über vetauschbare Kontraktionen des Hilbertschen Raumes. *Acta Sci. Math. Szeged*, 22:106–111, 1961.
- [Bun84] John W. Bunce. Models for n -tuples of noncommuting operators. *J. Funct. Anal.*, 57(1):21–30, 1984.
- [CL02] John Crisp and Marcelo Laca. On the Toeplitz algebras of right-angled and finite-type Artin groups. *J. Aust. Math. Soc.*, 72(2):223–245, 2002.
- [DK11] Kenneth R. Davidson and Elias G. Katsoulis. Dilatation theory, commutant lifting and semicrossed products. *Doc. Math.*, 16:781–868, 2011.
- [FK09] John Fountain and Mark Kambites. Graph products of right cancellative monoids. *J. Aust. Math. Soc.*, 87(2):227–252, 2009.
- [Fra82] Arthur E. Frazho. Models for noncommuting operators. *J. Funct. Anal.*, 48(1):1–11, 1982.
- [GS97] Dumitru Gaşpar and Nicolae Suciu. On the intertwinings of regular dilations. volume 66, pages 105–121. 1997. Volume dedicated to the memory of Włodzimierz Młak.
- [Li12] Xin Li. Semigroup C^* -algebras and amenability of semigroups. *J. Funct. Anal.*, 262(10):4302–4340, 2012.
- [Li16] Boyu Li. Regular representations of lattice ordered semigroups. *J. Operator Theory*, 76(1):33–56, 2016.
- [Li17] Boyu Li. Regular dilation on graph products of N . *J. Funct. Anal.*, 273(2):799–835, 2017.
- [Li19] Boyu Li. Regular dilation and Nica-covariant representation on right LCM semigroups. *Integral Equations Operator Theory*, 91(4):Paper No. 36, 35, 2019.
- [Li25] Boyu Li. Poisson transforms on right-angled artin monoids. *arXiv preprint arXiv:2504.00127*, 2025.
- [LL20] Marcelo Laca and Boyu Li. Amenability and functoriality of right-LCM semigroup C^* -algebras. *Proc. Amer. Math. Soc.*, 148(12):5209–5224, 2020.
- [LL22] Marcelo Laca and Boyu Li. Dilatation theory for right LCM semigroup dynamical systems. *J. Math. Anal. Appl.*, 505(2):Paper No. 125586, 37, 2022.
- [LR96] Marcelo Laca and Iain Raeburn. Semigroup crossed products and the Toeplitz algebras of non-abelian groups. *J. Funct. Anal.*, 139(2):415–440, 1996.
- [LS22] Marcelo Laca and Camila Sehnem. Toeplitz algebras of semigroups. *Trans. Amer. Math. Soc.*, 375(10):7443–7507, 2022.
- [Nic92] A. Nica. C^* -algebras generated by isometries and Wiener-Hopf operators. *J. Operator Theory*, 27(1):17–52, 1992.
- [Opě06] David Opěla. A generalization of Andô’s theorem and Parrott’s example. *Proc. Amer. Math. Soc.*, 134(9):2703–2710, 2006.
- [Par70] Stephen Parrott. Unitary dilations for commuting contractions. *Pacific J. Math.*, 34:481–490, 1970.
- [Pop89] Gelu Popescu. Isometric dilations for infinite sequences of noncommuting operators. *Trans. Amer. Math. Soc.*, 316(2):523–536, 1989.
- [Sch55] J. J. Schäffer. On unitary dilations of contractions. *Proc. Amer. Math. Soc.*, 6:322, 1955.
- [SN53] Béla Sz.-Nagy. Sur les contractions de l’espace de Hilbert. *Acta Sci. Math. (Szeged)*, 15:87–92, 1953.
- [Sta15] Charles Starling. Boundary quotients of C^* -algebras of right LCM semigroups. *J. Funct. Anal.*, 268(11):3326–3356, 2015.
- [Var74] N. Th. Varopoulos. On an inequality of von Neumann and an application of the metric theory of tensor products to operators theory. *J. Functional Analysis*, 16:83–100, 1974.

DEPARTMENT OF MATHEMATICAL SCIENCES, NEW MEXICO STATE UNIVERSITY, LAS CRUCES, NEW MEXICO, 88003, USA

Email address: boyuli@nmsu.edu

FACULTY OF MATHEMATICS, TECHNION – ISRAEL INSTITUTE OF TECHNOLOGY, HAIFA 3200003, ISRAEL

Email address: suryawanshi@campus.technion.ac.il