

ON NEW IDENTITIES OF JACOBI SUMS AND RELATED CYCLOTOMIC MATRICES

HAI-LIANG WU AND HAO PAN*

ABSTRACT. In this paper, using some arithmetic properties of Jacobi sums, we investigate some products involving Jacobi sums and reveal the connections between these products and certain cyclotomic matrices. In particular, as an application of our main results, we confirm a conjecture posed by Z.-W. Sun in 2019, and obtain a stronger result.

CONTENTS

| | |
|---|----|
| 1. Introduction | 2 |
| 1.1. Notation | 2 |
| 1.2. Background and motivation | 2 |
| 1.3. Main results | 4 |
| 1.4. Outline of this paper | 6 |
| 2. Preparations for the proofs of main results | 6 |
| 2.1. On a variant of the Gauss lemma | 6 |
| 2.2. Some known results on quadratic Gauss sums over $\mathbb{Z}/m\mathbb{Z}$ | 8 |
| 2.3. Circulant and almost circulant matrices | 9 |
| 2.4. Some necessary lemmas | 10 |
| 3. Proofs of Theorems 1.1–1.2 | 14 |
| 4. Proof of Corollary 1.1 | 16 |
| 5. Concluding Remarks | 20 |
| References | 21 |

Key words and phrases. Jacobi sums, cyclotomic matrices, finite fields.

2020 Mathematics Subject Classification. Primary 11L05, 15A15; Secondary 11R18, 12E20.

This research was supported by the Natural Science Foundation of China (Grant Nos. 12101321 and 12071208).

*Corresponding author.

1. INTRODUCTION

1.1. Notation. Throughout this paper, $q = p^f = 2n + 1$ is an odd prime power, where p is an odd prime and $f \in \mathbb{Z}^+ = \{1, 2, \dots\}$. The finite field with q elements is denoted by \mathbb{F}_q , and let $\mathbb{F}_q^\times = \mathbb{F}_q \setminus \{0\}$ be the multiplicative group of all nonzero elements over \mathbb{F}_q . Also,

$$\mathcal{S}_q = \{s_0 = 0, s_1 = 1, s_2, \dots, s_n\} = \{x^2 : x \in \mathbb{F}_q\}$$

denotes the set of all squares over \mathbb{F}_q .

Let $\widehat{\mathbb{F}_q^\times}$ denote the cyclic group of all multiplicative characters of \mathbb{F}_q . Fix a generator χ_q of $\widehat{\mathbb{F}_q^\times}$. Then, for any character $\chi_q^k \in \widehat{\mathbb{F}_q^\times}$, we define $\chi_q^k(0) = 0$. The quadratic character $\chi_q^{\pm n}$ is denoted by ϕ . Clearly,

$$\phi(x) = \begin{cases} 1 & \text{if } x \in \mathcal{S}_q \setminus \{0\}, \\ 0 & \text{if } x = 0, \\ -1 & \text{if } x \in \mathbb{F}_q \setminus \mathcal{S}_q. \end{cases}$$

For any $\chi_q^i, \chi_q^j \in \widehat{\mathbb{F}_q^\times}$, the Jacobi sum of χ_q^i and χ_q^j is defined by

$$J_q(\chi_q^i, \chi_q^j) := \sum_{x \in \mathbb{F}_q} \chi_q^i(x) \chi_q^j(1-x) \in \mathbb{Q}(\zeta_{q-1}),$$

where ζ_{q-1} is a primitive $(q-1)$ -th root of unity.

In addition, for any square matrix M over a field \mathbb{F} , we use $\det M$ to denote the determinant of M . Also, for any integers $a < b$, the symbol (a, b) denotes the set $\{a+1, \dots, b-1\}$, and the symbol $[a, b]$ denotes the set $\{a, a+1, \dots, b\}$.

1.2. Background and motivation. In 1827, Jacobi introduced the Jacobi sum in a letter to Gauss. Nowadays, Jacobi sums have extensive applications in number theory, and many well-known mathematicians made contributions to the theory of Jacobi sums. Readers may refer to [1] for detailed introduction on Jacobi sums.

In this paper, we mainly focus on some products involving Jacobi sums. We begin with a classical result due to Weil. Let $1 < m < q-1$ be a divisor of $q-1$ with $q-1 = lm$. Let

$$C_m := \{[x, y, z] \in \mathbb{P}^2(\mathbb{F}_q^{\text{alg}}) : x^m + y^m = z^m\}$$

be the projective Fermat curve defined over \mathbb{F}_q , where $\mathbb{F}_q^{\text{alg}}$ is an algebraic closure of \mathbb{F}_q , and $\mathbb{P}^2(\mathbb{F}_q^{\text{alg}})$ is the projective plane. Consider the zeta function of C_n defined by

$$\zeta_{C_m}(t) := \exp \left(\sum_{r \in \mathbb{Z}^+} \frac{N(q^r) \cdot t^r}{r} \right),$$

where $N(q^r)$ is the number of \mathbb{F}_{q^r} -rational points on C_m . As C_m is a nonsingular absolutely irreducible curve of genus $(m-1)(m-2)/2$, by the Weil theorem, there is an integral polynomial $P_m(t)$ with $\deg(P_m(t)) = (m-1)(m-2)$ such that

$$\zeta_{C_m}(t) = \frac{P_m(t)}{(1-t)(1-qt)}.$$

Weil proved that the integral polynomial $P_m(t)$ has close relations with certain product of Jacobi sums, that is,

$$P_m(t) = \prod_{\substack{i,j \in [1,m-1] \\ i+j \not\equiv 0 \pmod{m}}} (1 + J_q(\chi_q^{li}, \chi_q^{lj})t) \in \mathbb{Z}[t].$$

We next turn to Carlitz's result. Let $(\frac{\cdot}{p})$ be the Legendre symbol, where p is an odd prime. Carlitz [2] proved that the determinant of the matrix

$$(1.1) \quad C_p := \left[\left(\frac{i-j}{p} \right) \right]_{1 \leq i,j \leq p-1}$$

can be represented by a product of Jacobi sums, that is,

$$\det C_p = (-1)^{\frac{p-1}{2}} \cdot \prod_{k=1}^{p-1} J_p(\phi, \chi_p^k) = p^{\frac{p-3}{2}}.$$

Along this line, recently, the first author and Wang [14, Theorem 1.1] showed that if $q \equiv 3 \pmod{4}$ is a prime power, then

$$\det [\chi_q^r(s_i + s_j) + \chi_q^r(s_i - s_j)]_{1 \leq i,j \leq (q-1)/2} = \prod_{k=0}^{(q-3)/2} J_q(\chi_q^r, \chi_q^{2k})$$

for any $r \in [1, q-2]$.

It might be worth mentioning here that the above two results fully illustrate that the explicit values of certain products of Jacobi sums can be represented by the determinants of some cyclotomic matrices.

We next introduce some results due to Z.-W. Sun. Inspired by Carlitz's matrix C_p defined by (1.1), Z.-W. Sun [11] studied the matrix

$$S_p := \left[\left(\frac{i^2 + j^2}{p} \right) \right]_{1 \leq i, j \leq (p-1)/2},$$

and conjectured that $-\det S_p$ is a nonzero square of some integer whenever the prime $p \equiv 3 \pmod{4}$. This conjecture was later confirmed by Alekseyev and Krachun. When $p \equiv 1 \pmod{4}$, if we write $p = a^2 + 4b^2$ with $a, b \in \mathbb{Z}$ and $a \equiv 1 \pmod{4}$, then Cohen, Sun and Vsemirnov conjectured that $\det S_p/a$ is a nonzero square. The first author [13] proved this conjecture and revealed the connections between $\det S_p$ and the product

$$R_p(\chi_p) := \prod_{k \in (0, (p-1)/4)} (J_p(\phi, \chi_p^k) + J_p(\phi, \chi_p^{-k})),$$

which concerns the real part of the Jacobi sum $J_p(\phi, \chi_p^k)$.

Motivated by the above results, in this paper, we mainly investigate the product

$$(1.2) \quad I_q(\chi_q) := \prod_{k \in (0, n/2)} (J_q(\phi, \chi_q^k) - J_q(\phi, \chi_q^{-k})),$$

which involves the imaginary part of the Jacobi sum $J_q(\phi, \chi_q^k)$.

1.3. Main results. Now we state our two theorems concerning $I_q(\chi_q)$. Suppose first $q \equiv 3 \pmod{4}$ with $q = 2n + 1$. Note that the matrix

$$N_q(-1) := [\phi(s_i - s_j)]_{2 \leq i, j \leq n}$$

is a skew-symmetric matrix of even order with integer entries. Then by Stembridge's result [10, Proposition 2.2], $\det N_q(-1)$ is indeed a square of some integer. Our first theorem reveals the relationship between $I_q(\chi_q)$ and $\det N_q(-1)$.

Theorem 1.1. *Let $q = 2n + 1$ be an odd prime power with $q \equiv 3 \pmod{4}$, and let χ_q be a generator of $\widehat{\mathbb{F}_q^\times}$. Then*

$$x_q := \frac{I_q(\chi_q)}{\sqrt{(-1)^{\frac{n-1}{2}} \cdot n}} \in \mathbb{Z}.$$

Moreover, we have

$$x_q^2 = 2^{n-1} \cdot \det [\phi(s_i - s_j)]_{2 \leq i, j \leq n}.$$

We next turn to the case $q \equiv 1 \pmod{4}$. Inspired by Theorem 1.1, when $q \equiv 1 \pmod{4}$, one may expect that $I_q(\chi_q)$ has a close relationship with certain cyclotomic matrix involving ϕ . In fact, we will see that $I_q(\chi_q)$ is not only related to some cyclotomic matrix, but also to the number of \mathbb{F}_q -rational points on certain elliptic curve.

For any element $d \in \mathbb{F}_q^\times \setminus \mathcal{S}_q$, define the matrix

$$T_q(d) := [\phi(s_i + ds_j)]_{0 \leq i, j \leq n}.$$

Also, let X_d be the elliptic curve defined by the equation $y^2 = dx^3 + x$ over \mathbb{F}_q , and let

$$X_d(\mathbb{F}_q) = \{(x, y) \in \mathbb{F}_q \times \mathbb{F}_q : y^2 = dx^3 + x\} \cup \{\infty\}$$

be the set of all \mathbb{F}_q -rational points on X_d . Then it is known that

$$\#X_d(\mathbb{F}_q) = q + 1 - a_d(q),$$

where $\#S$ denotes the cardinality of a finite set S , and

$$(1.3) \quad a_d(q) = - \sum_{x \in \mathbb{F}_q} \phi(dx^3 + x)$$

is the trace of Frobenius map.

Let \mathbf{i} be the primitive 4-th root of unity with argument $\pi/2$. Now we state our second theorem.

Theorem 1.2. *Let $q = 2n + 1 \geq 7$ be an odd prime power with $q \equiv 1 \pmod{4}$, and χ_q be a generator of $\widehat{\mathbb{F}_q^\times}$. Then*

$$y_q := \mathbf{i}^{\frac{n-2}{2}} \cdot I_q(\chi_q) \cdot \sqrt{q-1} \in \mathbb{Z}.$$

Moreover, for any $d \in \mathbb{F}_q^\times \setminus \mathcal{S}_q$, we have

$$-a_d(q) \cdot y_q^2 = 2^n \cdot \det T_q(d).$$

In 2019, Z.-W. Sun [11, Conjecture 4.2(ii)] posed the following conjecture.

Conjecture 1.1. *Let $p \equiv 1 \pmod{4}$ be a prime with $p = c_p^2 + 4b_p^2$, where c_p and b_p are positive integers. Then for any $d \in \mathbb{F}_p^\times \setminus \mathcal{S}_p$, the number $|\det T_p(d)|/(2^{(p-1)/2}b_p)$ is a positive integral square not depending on d .*

As an application of Theorem 1.2, we can confirm this conjecture and obtain a stronger result.

Corollary 1.1. *Let $p \equiv 1 \pmod{4}$ be a prime and let notations be as above. Set*

$$z_p := \frac{y_p}{2^{(n-2)/2}}.$$

Then z_p is an integer such that

$$\left(\frac{z_p}{p} \right) = 1,$$

and

$$|\det T_p(d)|/(2^{(p-1)/2} b_p) = z_p^2.$$

1.4. Outline of this paper. In Section 2, we will prove some necessary lemmas. In particular, we will prove an important lemma, which can be viewed as a variant of the well-known Gauss lemma on quadratic residues. Our main results will be proved in Section 3 and Section 4 respectively. In Section 5, we shall briefly discuss a recent conjecture posed by Z.-W. Sun, which can be viewed as a refinement of Conjecture 1.1.

2. PREPARATIONS FOR THE PROOFS OF MAIN RESULTS

2.1. On a variant of the Gauss lemma. We first introduce some notations. For any real number y , let

$$\lfloor y \rfloor := \max\{u \in \mathbb{Z} : u \leq y\}.$$

For any positive integer m and any integer x , let $\{x\}_m$ denote the unique integer $r \in [0, m-1]$ with $r \equiv x \pmod{m}$.

Let p be an odd prime and $a \in \mathbb{Z}$ with $\gcd(a, p) = 1$. Then the famous Gauss lemma (cf. [4, p. 52]) says that

$$\left(\frac{a}{p} \right) = (-1)^{\#\{1 \leq x \leq \frac{p-1}{2} : \{ax\}_p > \frac{p}{2}\}}.$$

This result was later extended to Jacobi symbol by Jenkins. Let s be a positive odd integer and $a \in \mathbb{Z}$ with $\gcd(a, s) = 1$. Then Jenkins [5] proved that

$$(2.1) \quad \left(\frac{a}{s} \right) = (-1)^{\#\{1 \leq x \leq \frac{s-1}{2} : \{ax\}_s > \frac{s}{2}\}},$$

where $\left(\frac{\cdot}{s} \right)$ is the Jacobi symbol. Using this, it is easy to verify the following lemma.

Lemma 2.1. *Let s be a positive odd integer and $a \in \mathbb{Z}$ with $\gcd(a, s) = 1$. Then*

$$(-1)^K = \left(\frac{a}{s} \right),$$

where

$$K = \sum_{x=1}^{(s-1)/2} \left\lfloor \frac{2ax}{s} \right\rfloor.$$

The second author [8, Lemma 1] obtained the following result.

Lemma 2.2. *Let $m \geq 2$ be an integer and $a \in \mathbb{Z}$ with $\gcd(a, m) = 1$. Suppose that $1 \leq i < j \leq m-1$ and $\{ai\}_m > \{aj\}_m$. Then*

$$\left\lfloor \frac{aj}{m} \right\rfloor - \left\lfloor \frac{ai}{m} \right\rfloor - \left\lfloor \frac{a(j-i)}{m} \right\rfloor = 1.$$

We also need the following result.

Lemma 2.3. *Let $m \geq 2$ be an even integer and a be a positive integer with $\gcd(a, m) = 1$. Then*

$$\sum_{x=1}^{(a-1)/2} \left\lfloor \frac{2mx}{a} \right\rfloor + \sum_{y=1}^m \left\lfloor \frac{ay}{2m} \right\rfloor = \frac{(a-1)m}{2}.$$

Proof. By symmetry, one can verify that the sum

$$\sum_{x=1}^{(a-1)/2} \left\lfloor \frac{2mx}{a} \right\rfloor + \sum_{y=1}^m \left\lfloor \frac{ay}{2m} \right\rfloor$$

exactly counts the number of integral points of the set

$$\{(x, y) \in \mathbb{R} \times \mathbb{R} : 0 < x < a/2 \text{ and } 1 \leq y \leq m\}.$$

By this, our lemma holds immediately. \square

Now we are in a position to prove the following variant of Gauss lemma, which will play an important role in the proof of our second theorem.

Lemma 2.4. *Let $m \geq 2$ be an even integer and a be a positive integer with $\gcd(a, m) = 1$. Then*

$$(-1)^{\frac{(a-1)(m-2)}{4} + \#\{1 \leq x \leq \frac{m-2}{2} : \{ax\}_{2m} > m\}} = \left(\frac{2m}{a} \right).$$

Proof. Note that $\{am\}_{2m} = m$. By Lemma 2.2, one can verify that

$$\#\left\{1 \leq x \leq \frac{m-2}{2} : \{ax\}_{2m} > m\right\}$$

$$\begin{aligned}
&= \sum_{x=1}^{(m-2)/2} \left(\left\lfloor \frac{am}{2m} \right\rfloor - \left\lfloor \frac{ax}{2m} \right\rfloor - \left\lfloor \frac{a(m-x)}{2m} \right\rfloor \right) \\
(2.2) \quad &= \frac{(a-1)(m-2)}{4} + \left\lfloor \frac{a}{4} \right\rfloor - \sum_{y=1}^{m-1} \left\lfloor \frac{ay}{2m} \right\rfloor.
\end{aligned}$$

Meanwhile, by Lemma 2.3

$$(2.3) \quad \sum_{x=1}^{(a-1)/2} \left\lfloor \frac{2mx}{a} \right\rfloor + \sum_{y=1}^{m-1} \left\lfloor \frac{ay}{2m} \right\rfloor = \frac{(a-1)(m-1)}{2}.$$

Suppose first $a \equiv 1 \pmod{4}$. Combining (2.2) and (2.3) with Lemma 2.1, we obtain

$$\begin{aligned}
(-1)^{\#\{1 \leq x \leq \frac{m-2}{2}: \{ax\}_{2m} > m\}} &= (-1)^{\frac{a-1}{4} + \sum_{y=1}^{m-1} \left\lfloor \frac{ay}{2m} \right\rfloor} \\
&= (-1)^{\frac{a-1}{4} + \sum_{x=1}^{(a-1)/2} \left\lfloor \frac{2mx}{a} \right\rfloor} \\
&= \left(\frac{2m}{a} \right).
\end{aligned}$$

Now consider the case $a \equiv 3 \pmod{4}$. By (2.2), (2.3) and Lemma 2.1 again

$$\begin{aligned}
(-1)^{\#\{1 \leq x \leq \frac{m-2}{2}: \{ax\}_{2m} > m\}} &= (-1)^{\frac{m-2}{2} + \frac{a-3}{4} + \sum_{y=1}^{m-1} \left\lfloor \frac{ay}{2m} \right\rfloor} \\
&= (-1)^{\frac{m-2}{2} + \frac{a-3}{4} + 1 + \sum_{x=1}^{(a-1)/2} \left\lfloor \frac{2mx}{a} \right\rfloor} \\
&= (-1)^{\frac{m-2}{2}} \cdot \left(\frac{2m}{a} \right).
\end{aligned}$$

In view of the above, we have completed the proof. \square

2.2. Some known results on quadratic Gauss sums over $\mathbb{Z}/m\mathbb{Z}$. Let $p > 2$ be an odd prime. Gauss first determined the explicit value of the quadratic Gauss sum over $\mathbb{Z}/p\mathbb{Z}$, that is,

$$\sum_{x \in \mathbb{Z}/p\mathbb{Z}} \left(\frac{x}{p} \right) e^{2\pi i x/p} = \sum_{x \in \mathbb{Z}/p\mathbb{Z}} e^{2\pi i x^2/p} = \sqrt{(-1)^{\frac{p-1}{2}} p}.$$

The above result can be extended to the quadratic Gauss sum over $\mathbb{Z}/m\mathbb{Z}$, where m is an arbitrary positive integer (cf. [7, pp. 85-87]).

Lemma 2.5. *Let m be a positive integer. Then*

$$\sum_{x \in \mathbb{Z}/m\mathbb{Z}} e^{2\pi i x^2/m} = \begin{cases} 0 & \text{if } m \equiv 2 \pmod{4}, \\ (1+i)\sqrt{m} & \text{if } m \equiv 0 \pmod{4}, \\ \sqrt{(-1)^{\frac{m-1}{2}} m} & \text{if } m \equiv 1 \pmod{2}. \end{cases}$$

Moreover, if m is a positive odd integer, then

$$\sum_{x \in \mathbb{Z}/m\mathbb{Z}} e^{2\pi i a x^2/m} = \left(\frac{a}{m}\right) \cdot \sum_{x \in \mathbb{Z}/m\mathbb{Z}} e^{2\pi i x^2/m}$$

for any $a \in \mathbb{Z}$ with $\gcd(a, m) = 1$.

To state our next result, we introduce the Kronecker symbol, which is indeed an extension of Jacobi symbol. In fact, the Kronecker symbol is a function

$$\left(\frac{\bullet}{\bullet}\right) : (\mathbb{Z} \setminus \{0\}) \times (\mathbb{Z} \setminus \{0\}) \longrightarrow \{-1, 0, 1\}$$

defined by the following conditions:

- (i) $\left(\frac{b}{a_1}\right) \left(\frac{b}{a_2}\right) = \left(\frac{b}{a_1 a_2}\right)$ for any nonzero integers a_1, a_2, b ;
- (ii) If a is a positive odd integer, then $\left(\frac{b}{a}\right)$ coincides with the Jacobi symbol;
- (iii) For any nonzero integer b , we have $\left(\frac{b}{-1}\right) = \text{sign}(b) = b/|b|$;
- (iv) For any nonzero integer b , we have

$$\left(\frac{b}{2}\right) = \begin{cases} 0 & \text{if } b \equiv 0 \pmod{2}, \\ \left(\frac{2}{b}\right) & \text{if } b \equiv 1 \pmod{2}. \end{cases}$$

Now we state the following known result on character sums related to the Kronecker symbol (cf. [3, Corollary 2.1.47]).

Lemma 2.6. *Let K be a quadratic number field with discriminant D and let $m = |D|$. Then*

$$\sqrt{D} = \sum_{x \in (\mathbb{Z}/m\mathbb{Z}) \setminus \{0 \pmod{m}\}} \left(\frac{D}{x}\right) e^{2\pi i x/m}.$$

2.3. Circulant and almost circulant matrices. Let $m \geq 2$ be an integer and let

$$\mathbf{v} = (c_0, c_1, \dots, c_{m-1}) \in \mathbb{C}^m.$$

Then the *circulant matrix* $C(\mathbf{v})$ of the vector \mathbf{v} is an $m \times m$ matrix defined by

$$C(\mathbf{v}) := [c_{j-i}]_{0 \leq i, j \leq m-1},$$

where $c_x = c_{\{x\}_m}$ for any integer x . More precisely,

$$C(\mathbf{v}) = \begin{bmatrix} c_0 & c_1 & \cdots & c_{m-2} & c_{m-1} \\ c_{m-1} & c_0 & \cdots & c_{m-3} & c_{m-2} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ c_2 & c_3 & \cdots & c_0 & c_1 \\ c_1 & c_2 & \cdots & c_{m-1} & c_0 \end{bmatrix}.$$

Circulant matrices have extensively applications in number theory, combinatorics, as well as coding theory. Readers may refer to the survey paper [6] for more details on circulant matrices.

In 2025, the first author and Wang investigated the matrix

$$W(\mathbf{v}) := [c_{j-i}]_{1 \leq i, j \leq m-1},$$

which is obtained by deleting the first row and the first column of $C(\mathbf{v})$. The matrix $W(\mathbf{v})$ is called the *almost circulant matrix* of \mathbf{v} . For $W(\mathbf{v})$, the first author and Wang [14, Theorem 4.1] obtained the following result.

Lemma 2.7. *Let notations be as above and let $\lambda_1, \lambda_2, \dots, \lambda_m$ be exactly all the eigenvalues of the circulant matrix $C(\mathbf{v})$. Then*

$$\det W(\mathbf{v}) = \frac{1}{m} \sum_{l=1}^m \prod_{k \in [1, m] \setminus \{l\}} \lambda_k.$$

In particular, if $\lambda_m = 0$, then

$$\det W(\mathbf{v}) = \frac{1}{m} \lambda_1 \lambda_2 \cdots \lambda_{m-1}.$$

2.4. Some necessary lemmas. Here we introduce some lemmas, which will be used frequently in the proofs of main theorems. Recall that $q = 2n + 1$ is an odd prime power and χ_q is a generator of $\widehat{\mathbb{F}_q^\times}$. We begin with the following result.

Lemma 2.8. *Let notations be as above. Then*

$$J_q(\chi_q^i, \chi_q^j) = (-1)^i \cdot J_q(\chi_q^i, \chi_q^{-(i+j)})$$

for any integers i and j . In particular,

$$J_q(\phi, \chi_q^j) = \phi(-1) \cdot J_q(\phi, \chi_q^{n-j}).$$

Proof. By the definition of Jacobi sums, one can verify that

$$\begin{aligned}
(-1)^i \cdot J_q(\chi_q^i, \chi_q^{-(i+j)}) &= \sum_{x \in \mathbb{F}_q^\times} \chi_q^i(x-1) \chi_q^{i+j} \left(\frac{1}{x} \right) \\
&= \sum_{x \in \mathbb{F}_q^\times} \chi_q^i \left(1 - \frac{1}{x} \right) \chi_q^j \left(\frac{1}{x} \right) \\
&= \sum_{x \in \mathbb{F}_q^\times} \chi_q^i(1-x) \chi_q^j(x) \\
&= J_q(\chi_q^i, \chi_q^j).
\end{aligned}$$

This completes the proof. \square

Recall that

$$\mathcal{S}_q = \{s_0 = 0, s_1 = 1, s_2, \dots, s_n\} = \{x^2 : x \in \mathbb{F}_q\}.$$

Throughout out the remaining part of this paper, for $k = 1, 2, \dots, n$, we define

$$\lambda_k(d) := \sum_{s \in \mathcal{S}_q} \phi(1+ds) \chi_q^k(s).$$

Using these notations, we have the following result.

Lemma 2.9. *Let $d \in \mathbb{F}_q^\times$ and let the matrix*

$$M_q(d) := [\phi(s_i + ds_j)]_{1 \leq i, j \leq n}.$$

Then the following results hold.

- (i) $\lambda_1(d), \lambda_2(d), \dots, \lambda_n(d)$ are exactly all the eigenvalues of $M_q(d)$.
- (ii) Moreover, if $d \in \mathbb{F}_q^\times \setminus \mathcal{S}_q$, then for any $k = 1, 2, \dots, n$, we have

$$\chi_q^k(d) \cdot \lambda_k(d) = \frac{(-1)^k}{2} (J_q(\phi, \chi_q^k) - J_q(\phi, \chi_q^{-k})).$$

Proof. (i) For any $k \in [1, n]$, define the column vector

$$\mathbf{v}_k := (\chi_q^k(s_1), \chi_q^k(s_2), \dots, \chi_q^k(s_n))^T.$$

Then one can verify that

$$\begin{aligned}
\sum_{j \in [1, n]} \phi(s_i + ds_j) \chi_q^k(s_j) &= \sum_{j \in [1, n]} \phi(1 + ds_j/s_i) \chi_q^k(s_j/s_i) \chi_q^k(s_i) \\
&= \sum_{j \in [1, n]} \phi(1 + ds_j) \chi_q^k(s_j) \cdot \chi_q^k(s_i)
\end{aligned}$$

$$= \lambda_k(d) \cdot \chi_q^k(s_i).$$

This implies

$$M_q(d) \mathbf{v}_k = \lambda_k(d) \mathbf{v}_k$$

for any $k \in [1, n]$. As vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ are linearly independent over \mathbb{C} , the numbers $\lambda_1(d), \lambda_2(d), \dots, \lambda_n(d)$ are exactly all the eigenvalues of $M_q(d)$. Hence (i) holds.

(ii) Suppose now $d \in \mathbb{F}_q^\times \setminus \mathcal{S}_q$, i.e., $\phi(d) = -1$. Then the conjugate of the complex number $\chi_q^k(d) \cdot \lambda_k(d)$ is equal to

$$\begin{aligned} \overline{\chi_q^k(d) \cdot \lambda_k(d)} &= \sum_{s \in \mathcal{S}_q} \phi(1 + ds) \chi_q^{-k}(ds) \\ &= -\chi_q^k(d) \sum_{s \in \mathcal{S}_q} \phi(d + d^2 s) \chi_q^{-k}(d^2 s) \\ &= -\chi_q^k(d) \sum_{s \in \mathcal{S}_q} \phi(d + s) \chi_q^{-k}(s) \\ &= -\chi_q^k(d) \sum_{s \in \mathcal{S}_q \setminus \{0\}} \phi\left(d + \frac{1}{s}\right) \chi_q^k(s) \\ &= -\chi_q^k(d) \sum_{s \in \mathcal{S}_q \setminus \{0\}} \phi(1 + ds) \chi_q^k(s) \\ &= -\chi_q^k(d) \cdot \lambda_k(d). \end{aligned}$$

This implies that

$$\chi_q^k(d) \cdot \lambda_k \in \mathbb{R} \mathbf{i} = \{x \mathbf{i} : x \in \mathbb{R}\},$$

i.e., $\chi_q^k(d) \cdot \lambda_k$ is either 0 or a purely imaginary number. On the other hand, by (i), the numbers $\lambda_1(1), \lambda_2(1), \dots, \lambda_n(1)$ are precisely all the eigenvalues of the real symmetric matrix $M_q(1)$. Thus, $\lambda_k(1) \in \mathbb{R}$ for any $k \in [1, n]$.

As $d \in \mathbb{F}_q^\times \setminus \mathcal{S}_q$, we clearly have

$$\{ds : s \in \mathcal{S}_q \setminus \{0\}\} \cap \mathcal{S}_q = \emptyset,$$

and

$$\{ds : s \in \mathcal{S}_q \setminus \{0\}\} \cup \mathcal{S}_q = \mathbb{F}_q.$$

Using this, we obtain

$$\chi_q^k(d) \cdot \lambda_k(d) + \lambda_k(1) = \sum_{s \in \mathcal{S}_q} \phi(1 + ds) \chi_q^k(ds) + \sum_{s \in \mathcal{S}_q} \phi(1 + s) \chi_q^k(s)$$

$$\begin{aligned}
&= \sum_{x \in \mathbb{F}_q} \phi(1+x) \chi_q^k(x) \\
&= (-1)^k \cdot J_q(\phi, \chi_q^k).
\end{aligned}$$

Noting that $\chi_q^k(d) \cdot \lambda_k(d) \in \mathbb{R}i$ and $\lambda_k(1) \in \mathbb{R}$, we obtain

$$(2.4) \quad \chi_q^k(d) \cdot \lambda_k(d) = \frac{(-1)^k}{2} (J_q(\phi, \chi_q^k) - J_q(\phi, \chi_q^{-k}))$$

for any $k \in [1, n]$.

In view of the above, we have completed the proof. \square

For any $d \in \mathbb{F}_q^\times \setminus \mathcal{S}_q$, recall that the matrix

$$T_q(d) = [\phi(s_i + ds_j)]_{0 \leq i, j \leq n}.$$

We conclude this section with the following result.

Lemma 2.10. *Let $q = 2n + 1$ be an odd prime power with $q \equiv 1 \pmod{4}$. Then for any $d \in \mathbb{F}_q^\times \setminus \mathcal{S}_q$, the $n+1$ numbers $\pm\sqrt{-n}, \lambda_1(d), \lambda_2(d), \dots, \lambda_{n-1}(d)$ are exactly all the eigenvalues of $T_q(d)$.*

Proof. For $k = -1, 0, 1, \dots, n-1$, the column vector \mathbf{u}_k is defined by

$$\begin{cases} \mathbf{u}_k = ((-1)^{k+1} \cdot \sqrt{-n}, 1, 1, \dots, 1)^T & \text{if } k = -1, 0, \\ \mathbf{u}_k = (0, \chi_q^k(s_1), \chi_q^k(s_2), \dots, \chi_q^k(s_n))^T & \text{if } k \in [1, n-1]. \end{cases}$$

It is easy to see that the vectors $\mathbf{u}_{-1}, \mathbf{u}_0, \mathbf{u}_1, \dots, \mathbf{u}_{n-1}$ are linearly independent over \mathbb{C} .

For $k = -1, 0$, we clearly have

$$T_q(d)\mathbf{u}_k = (-1)^{k+1} \sqrt{-n} \mathbf{u}_k.$$

Also, with essentially the same method appeared in the proof of Lemma 2.9, we obtain

$$T_q(d)\mathbf{u}_k = \lambda_k(d)\mathbf{u}_k$$

for any $k \in [1, n-1]$. In view of the above, $\pm\sqrt{-n}, \lambda_1(d), \lambda_2(d), \dots, \lambda_{n-1}(d)$ are precisely all the eigenvalues of $T_q(d)$. This completes the proof. \square

3. PROOFS OF THEOREMS 1.1–1.2

Proofs of Theorems 1.1. Let $q \equiv 3 \pmod{4}$ be an odd prime power with $n = (q-1)/2$. We first prove that

$$x_q = \frac{I_q(\chi_q)}{\sqrt{(-1)^{\frac{n-1}{2}} \cdot n}} \in \mathbb{Q}.$$

Let $\sigma_a \in \text{Gal}(\mathbb{Q}(\zeta_{q-1})/\mathbb{Q})$ be an arbitrary automorphism with $a \in \mathbb{Z}$, $\gcd(a, q-1) = 1$ and $\sigma_a(\zeta_{q-1}) = \zeta_{q-1}^a$. For $j = 1, 2, 3, 4$, define the set

$$S_j(a) = \left\{ k \in (0, n/2) : \frac{j-1}{2}n < \{ka\}_{2n} < \frac{j}{2}n \right\}.$$

As $q \equiv 3 \pmod{4}$, applying Lemma 2.8, we obtain

$$J_q(\phi, \chi_q^k) = -1 \cdot J_q(\phi, \chi_q^{\pm n-k}).$$

By this, one can verify that

$$(3.1) \quad \sigma_a(I_q(\chi_q)) = (-1)^{\#S_2(a) \cup S_4(a)} \cdot I_q(\chi_q).$$

Note that

$$k_1a \not\equiv k_2a \pmod{n}$$

for any $k_1, k_2 \in (0, n/2)$ with $k_1 \neq k_2$. Thus,

$$S_2(a) \cup S_4(a) = \left\{ k \in (0, n/2) : \frac{n}{2} < \{ka\}_n < n \right\}.$$

Combining this with (2.1) and (3.1), we immediately obtain

$$(3.2) \quad \sigma_a(I_q(\chi_q)) = \left(\frac{a}{n}\right) \cdot I_q(\chi_q).$$

On the other hand, by Lemma 2.5

$$(3.3) \quad \sigma_a\left(\sqrt{(-1)^{\frac{n-1}{2}}n}\right) = \left(\frac{a}{n}\right) \cdot \sqrt{(-1)^{\frac{n-1}{2}}n}.$$

By (3.2) and (3.3), we see that

$$\sigma_a(x_q) = x_q$$

for any $\sigma_a \in \text{Gal}(\mathbb{Q}(\zeta_{q-1})/\mathbb{Q})$. Hence $x_q \in \mathbb{Q}$ by the Galois theory.

We next evaluate x_q^2 . Fix a generator g of cyclic group \mathbb{F}_q^\times and let $s_{j+1} = g^{2j}$ for $j = 0, 1, \dots, n-1$. Then the matrix

$$M_q(-1) = [\phi(s_i - s_j)]_{1 \leq i, j \leq n} = [\phi(1 - g^{2j-2i})]_{0 \leq i, j \leq n-1}$$

is indeed a circulant matrix $C(\mathbf{v})$ of the vector

$$\mathbf{v} = (\phi(1 - s_1), \phi(1 - s_2), \dots, \phi(1 - s_n)).$$

Hence

$$N_q(-1) = [\phi(s_i - s_j)]_{2 \leq i, j \leq n} = [\phi(1 - g^{2j-2i})]_{1 \leq i, j \leq n-1}$$

is the almost circulant matrix of \mathbf{v} . As $q \equiv 3 \pmod{4}$, we have $-1 \notin \mathcal{S}_q$. Note that $\lambda_n(-1) = 0$.

Applying Lemma 2.9 and Lemma 2.7, we obtain

$$\det N_q(-1) = \frac{1}{n} \prod_{k \in [1, n-1]} \lambda_k(-1) = \frac{1}{2^{n-1} \cdot n} \prod_{k \in [1, n-1]} (J_q(\phi, \chi_q^k) - J_q(\phi, \chi_q^{-k})).$$

Using Lemma 2.8 again, the above equality implies

$$(3.4) \quad 2^{n-1} \cdot \det N_q(-1) = x_q^2 \in \mathbb{Z}.$$

As $x_q \in \mathbb{Q}$, by (3.4) we see that $x_q \in \mathbb{Z}$.

In view of the above, we have completed the proof. \square

We next turn to the proof of our second theorem.

Proof of Theorem 1.2. Let $q = 2n + 1 \geq 7$ be an odd prime power with $q \equiv 1 \pmod{4}$. We first show that

$$y_q = \mathbf{i}^{\frac{n-2}{2}} \cdot I_q(\chi_q) \cdot \sqrt{q-1} \in \mathbb{Z}.$$

Let $\sigma_a \in \text{Gal}(\mathbb{Q}(\zeta_{q-1})/\mathbb{Q})$ be an arbitrary automorphism with $a \in [1, q-1]$, $\gcd(a, q-1) = 1$ and $\sigma_a(\zeta_{q-1}) = \zeta_{q-1}^a$. For any $j \in [1, 4]$, let

$$S_j(a) = \left\{ k \in (0, n/2) : \frac{j-1}{2}n < \{ka\}_{2n} < \frac{j}{2}n \right\}.$$

By Lemma 2.8 and $\phi(-1) = 1$, we have

$$J_q(\phi, \chi_q^k) = J_q(\phi, \chi_q^{n-k}).$$

By this and Lemma 2.4, one can verify that

$$(3.5) \quad \sigma_a \left(\mathbf{i}^{\frac{n-2}{2}} \cdot I_q(\chi_q) \right) = (-1)^{\frac{(a-1)(n-2)}{4} + \#S_3(a) \cup S_4(a)} \cdot \mathbf{i}^{\frac{n-2}{2}} \cdot I_q(\chi_q) = \left(\frac{q-1}{a} \right) \cdot \mathbf{i}^{\frac{n-2}{2}} \cdot I_q(\chi_q).$$

On the other hand, by Lemma 2.6 we see that

$$(3.6) \quad \sigma_a \left(\sqrt{q-1} \right) = \left(\frac{q-1}{a} \right) \cdot \sqrt{q-1}.$$

By (3.5) and (3.6), for any $\sigma_a \in \text{Gal}(\mathbb{Q}(\zeta_{q-1})/\mathbb{Q})$ we have

$$\sigma_a(y_q) = y_q,$$

and hence $y_q \in \mathbb{Q}$ by the Galois theory. As y_q is an algebraic integer, we further obtain $y_q \in \mathbb{Z}$.

We next consider y_q^2 . Recall that the matrix

$$T_d(q) = [\phi(s_i + ds_j)]_{0 \leq i, j \leq n}.$$

By Lemma 2.8, Lemma 2.9 and Lemma 2.10, one can verify that

$$\begin{aligned} \det T_q(d) &= n \cdot \lambda_{n/2}(d) \cdot \prod_{k \in [1, n-1] \setminus \{n/2\}} \lambda_k(d) \\ &= n \cdot \lambda_{n/2}(d) \cdot \prod_{k \in [1, n-1] \setminus \{n/2\}} \frac{(-1)^k \chi_q^{-k}(d)}{2} (J_q(\phi, \chi_q^k) - J_q(\phi, \chi_q^{-k})) \\ &= \frac{n \cdot \lambda_{n/2}(d)}{2^{n-2}} \cdot (-1)^{\frac{n-2}{2}} \cdot I_q(\chi_q)^2 \\ (3.7) \quad &= \frac{\lambda_{n/2}(d)}{2^{n-1}} \cdot y_q^2. \end{aligned}$$

Note also that

$$\begin{aligned} 2 \cdot \lambda_{n/2}(d) &= 2 \sum_{j \in [1, n]} \phi(1 + ds_j) \chi_q^{n/2}(s_j) \\ &= 2 \sum_{j \in [1, n]} \phi(1 + ds_j) \phi(\sqrt{s_j}) \\ &= \sum_{j \in [1, n]} \phi(1 + ds_j) \phi(\sqrt{s_j}) + \sum_{j \in [1, n]} \phi(1 + ds_j) \phi(-\sqrt{s_j}) \\ &= \sum_{x \in \mathbb{F}_q} \phi(1 + dx^2) \phi(x), \end{aligned}$$

where $\sqrt{s_j} \in \mathbb{F}_q$ such that $(\sqrt{s_j})^2 = s_j$. By this and (1.3) we obtain

$$(3.8) \quad a_d(q) = -2\lambda_{n/2}(d).$$

Now combining (3.7) with (3.8), we obtain

$$-1 \cdot 2^n \cdot \det T_q(d) = a_d(q) \cdot y_q^2.$$

In view of the above, we have completed the proof. \square

4. PROOF OF COROLLARY 1.1

Now we turn to the proof of our last result. We first introduce the following lemma.

Lemma 4.1. *Let $p \equiv 1 \pmod{4}$ be a prime and let*

$$B_p = \prod_{k=1}^{(p-1)/4} \binom{2k}{k}.$$

Then

$$\left(\frac{B_p}{p} \right) = \left(\frac{2}{p} \right).$$

Proof. By the definition of binomial coefficients, one can verify that

$$(4.1) \quad \left(\frac{B_p}{p} \right) = \prod_{k=1}^{(p-1)/4} \left(\frac{(2k)!}{p} \right) = \prod_{x=1}^{(p-1)/2} \left(\frac{x}{p} \right)^{\frac{p-1}{4}} \left(\frac{x}{p} \right)^{\lfloor \frac{x-1}{2} \rfloor} = (-1)^{\frac{p-1}{4}} \prod_{x=1}^{(p-1)/2} \left(\frac{x}{p} \right)^{\lfloor \frac{x-1}{2} \rfloor}.$$

For $i = 0, 1, 2, 3$, let

$$N_i(p) = \# \left\{ x \in (0, p/2) : \left(\frac{x}{p} \right) = -1 \text{ and } x \equiv i \pmod{4} \right\}.$$

As $p \equiv 1 \pmod{4}$, one can verify that

$$\begin{aligned} & N_0(p) + N_1(p) \\ &= \# \left\{ x \in (0, p/8) : \left(\frac{x}{p} \right) = -1 \right\} + \# \left\{ x \in (p/2, p) : \left(\frac{x}{p} \right) = -1 \text{ and } x \equiv 0 \pmod{4} \right\} \\ &= \# \left\{ x \in (0, p/8) : \left(\frac{x}{p} \right) = -1 \right\} + \# \left\{ x \in (p/8, p/4) : \left(\frac{x}{p} \right) = -1 \right\} \\ &= \# \left\{ x \in (0, p/4) : \left(\frac{x}{p} \right) = -1 \right\}, \end{aligned}$$

and that

$$N_0(p) + N_2(p) = \# \left\{ x \in (0, p/4) : \left(\frac{x}{p} \right) = - \left(\frac{2}{p} \right) \right\}.$$

This implies

$$(4.2) \quad N_1(p) + N_2(p) \equiv \frac{p-1}{4} \pmod{2}.$$

Using $p \equiv 1 \pmod{4}$ again, we obtain

$$(4.3) \quad N_0(p) + N_1(p) + N_2(p) + N_3(p) = \frac{p-1}{4}.$$

By (4.2) and (4.3), we see that

$$N_0(p) + N_3(p) \equiv 0 \pmod{2}.$$

By this and (4.1), we obtain

$$\left(\frac{B_p}{p}\right) = (-1)^{\frac{p-1}{4} + N_0(p) + N_3(p)} = (-1)^{\frac{p-1}{4}} = \left(\frac{2}{p}\right).$$

This completes the proof. \square

To state our next lemma, we need introduce the following notations. Let ζ_{p-1} be a primitive $(p-1)$ -th root of unity. Let the cyclotomic field $K = \mathbb{Q}(\zeta_{p-1})$ and let \mathcal{O}_K be the ring of all algebraic integers over K . Let \mathfrak{p} be a prime ideal of \mathcal{O}_K with $p \in \mathfrak{p}$. Then the Teichmüller character $\omega_{\mathfrak{p}}$ of \mathfrak{p} is defined by

$$\omega_{\mathfrak{p}}(x \bmod \mathfrak{p}) \equiv x \pmod{\mathfrak{p}}$$

for any $x \in \mathcal{O}_K$. As p splits totally over \mathcal{O}_K , we have

$$\mathcal{O}_K/\mathfrak{p} \cong \mathbb{F}_p.$$

Thus, if we identify $\mathcal{O}_K/\mathfrak{p}$ with \mathbb{F}_p , then it is easy to verify that $\omega_{\mathfrak{p}}$ is a generator of $\widehat{\mathbb{F}_p^\times}$.

We now state our next lemma (cf. [3, Proposition 3.6.4]).

Lemma 4.2. *Let notations be as above. Then for any integers $1 \leq i, j \leq p-2$,*

$$J_p(\omega_{\mathfrak{p}}^{-i}, \omega_{\mathfrak{p}}^{-j}) \equiv -\binom{i+j}{i} \pmod{\mathfrak{p}}.$$

In particular, if $i+j \geq p$, then

$$J_p(\omega_{\mathfrak{p}}^{-i}, \omega_{\mathfrak{p}}^{-j}) \equiv 0 \pmod{\mathfrak{p}}.$$

We also need the following known result (cf. [1, Theorem 2.1.4]).

Lemma 4.3. *Let $p > 2$ be an odd prime. Then for any nontrivial character $\psi \in \widehat{\mathbb{F}_p^\times}$,*

$$J_p(\phi, \psi) = \psi(4) \cdot J_p(\psi, \psi).$$

Now we are in a position to prove our last result.

Proof of Corollary 1.1. Let notations be as above and let $p \equiv 1 \pmod{4}$ be a prime with $p = 2n+1$. When $p = 5$, our result clearly holds.

Suppose now $p \geq 7$. As

$$\lambda_k(d) = \frac{(-1)^k \chi_p^{-k}(d)}{2} (J_p(\phi, \chi_q^k) - J_p(\phi, \chi_p^{-k}))$$

for any $k \in [1, n-1]$, the rational number

$$\frac{y_p}{2^{(n-2)/2}} = \mathbf{i}^{(n-2)/2} \cdot \sqrt{p-1} \cdot \prod_{k \in (0, n/2)} \frac{1}{2} (J_p(\phi, \chi_q^k) - J_p(\phi, \chi_p^{-k}))$$

is indeed an algebraic integer. Thus,

$$(4.4) \quad z_p = \frac{y_p}{2^{(n-2)/2}} \in \mathbb{Z}.$$

On the other hand, for any $d \in \mathbb{F}_p^\times \setminus \mathcal{S}_p$, by (3.8) one can verify that

$$\begin{aligned} (-1)^{n/2} \cdot J_p(\phi, \chi_p^{n/2}) &= \sum_{j \in [1, n]} \phi(1 + s_j) \chi_p^{n/2}(s_j) + \sum_{j \in [1, n]} \phi(1 + ds_j) \chi_p^{n/2}(ds_j) \\ &= \sum_{j \in [1, n]} \phi(1 + s_j) \phi(\sqrt{s_j}) \pm \mathbf{i} \sum_{j \in [1, n]} \phi(1 + ds_j) \phi(\sqrt{s_j}) \\ &= \lambda_{n/2}(1) \pm \mathbf{i} \cdot \lambda_{n/2}(d) \\ &= \lambda_{n/2}(1) \mp \mathbf{i} \cdot \frac{a_p(d)}{2}. \end{aligned}$$

Also, as $p \equiv 1 \pmod{4}$, we see that

$$\lambda_{n/2}(1) \equiv n-1 \equiv 1 \pmod{2}.$$

Hence, if we write $p = c_p^2 + 4b_p^2$ with $c_p, b_p \in \mathbb{Z}^+$, then

$$(4.5) \quad \pm a_p(d)/4 = b_p > 0.$$

Now applying Theorem 1.2, by (4.4) and (4.5) we see that

$$|\det T_d(p)| / (2^n \cdot b_p) = z_p^2$$

is a square of the integer z_p .

It remains to prove that the integer z_p is a quadratic residue modulo p . Let notations be as in Lemma 4.2. As $(\frac{z_p}{p})$ is independent of the choice of the generator χ_p , we now let $\chi_p = \omega_p$ be the Teichmüller character ω_p of \mathfrak{p} . Then by Lemma 4.3 and Lemma 4.2

$$\begin{aligned} y_p &\equiv \mathbf{i}^{(n-2)/2} \cdot \sqrt{p-1} \cdot \prod_{k \in (0, n/2)} (J_p(\phi, \chi_q^k) - J_p(\phi, \chi_p^{-k})) \\ &\equiv \mathbf{i}^{(n-2)/2} \cdot \sqrt{p-1} \cdot \prod_{k \in (0, n/2)} -J_p(\phi, \chi_p^{-k}) \\ &\equiv \mathbf{i}^{(n-2)/2} \cdot \sqrt{p-1} \cdot \prod_{k \in (0, n/2)} -\chi_p^{-k}(4) \cdot J_p(\chi_p^{-k}, \chi_p^{-k}) \end{aligned}$$

$$\equiv i^{(n-2)/2} \cdot \sqrt{p-1} \cdot \prod_{k \in (0, n/2)} \chi_p^{-k}(4) \cdot \binom{2k}{k} \pmod{\mathfrak{p}}.$$

By this and Lemma 4.1, one can verify that

$$\left(\frac{z_p}{p} \right) = \left(\frac{2}{p} \right)^{\frac{n-2}{2}} \left(\frac{y_p}{p} \right) = \left(\frac{2}{p} \right)^{\frac{n}{2}} \left(\frac{\sqrt{-1}}{p} \right)^{\frac{n}{2}} = 1.$$

In view of the above, we have completed the proof. \square

5. CONCLUDING REMARKS

Let notations be as above. Recently, Z.-W. Sun [12, Conjecture 6.1] posed the following conjecture, which is a refinement of Conjecture 1.1.

Conjecture 5.1. *Let $p \equiv 1 \pmod{4}$ be a prime. Then, there is an integer t_p with $(\frac{t_p}{p}) = 1$ such that*

$$\det T_p(d) = 2^{\frac{p-3}{2}} \cdot \left(\frac{p-1}{4} t_p \right)^2 \cdot \sum_{j=1}^{\frac{p-1}{2}} \left(\frac{x^3 + dx}{p} \right)$$

for any $d \in \mathbb{F}_p^\times \setminus \mathcal{S}_p$.

In this section, we briefly discuss this stronger conjecture. Using (4.5), we first observe that

$$\begin{aligned} 2 \sum_{j=1}^{(p-1)/2} \left(\frac{x^2 + d}{p} \right) \left(\frac{x}{p} \right) &= \sum_{x \in \mathbb{F}_p} \left(\frac{x^2 + d}{p} \right) \left(\frac{x}{p} \right) \\ &= \sum_{x \in \mathbb{F}_p} \left(\frac{(dx)^2 + d}{p} \right) \left(\frac{dx}{p} \right) \\ &= \sum_{x \in \mathbb{F}_p} \left(\frac{dx^2 + 1}{p} \right) \left(\frac{x}{p} \right) \\ &= -a_p(d) \\ &= \pm 2b_p. \end{aligned}$$

Thus, Conjecture 5.1 is equivalent to saying that

$$\frac{|\det T_p(d)|}{2^{(p-1)/2} \cdot b_p} = \left(\frac{p-1}{4} t_p \right)^2$$

for some integer t_p with $(\frac{t_p}{p}) = 1$. On the other hand, by the proof of Corollary 1.1, we have proven that

$$\frac{|\det T_p(d)|}{2^{(p-1)/2} \cdot b_p} = z_p^2,$$

where

$$z_p = \frac{i^{\frac{p-5}{4}} \cdot I_p(\chi_p) \cdot \sqrt{p-1}}{2^{(p-5)/4}} \in \mathbb{Z}$$

with $(\frac{z_p}{p}) = 1$.

In view of the above, we clearly see that Conjecture 5.1 holds if and only if the rational number $4z_p/(p-1)$ is an integer. However, we can not prove this result by the methods used in this paper.

REFERENCES

- [1] B. C. Berndt, R. J. Evans, K. S. Williams, *Gauss and Jacobi Sums*, Wiley, New York, 1998.
- [2] L. Carlitz, Some cyclotomic matrices, *Acta Arith.* 5 (1959), 293–308.
- [3] H. Cohen, *Number Theory, Vol. I. Tools and Diophantine Equations*, Graduate Texts in Math., 239, Springer, New York, 2007.
- [4] K. Ireland and M. Rosen, *A classical introduction to modern number theory*, 2nd Edition, Graduate Texts in Math., 84, Springer, New York, 1990.
- [5] M. Jenkins, Proof of an arithmetical theorem leading, by means of Gauss fourth demonstration of Legendre's law of reciprocity, to the extension of that law, *Proc. London Math. Soc.* 2 (1867) 29–32.
- [6] I. Kra, and S. R. Simanca, On circulant matrices, *Not. Am. Math. Soc.* 59 (2012), 368–377.
- [7] S. Lang, *Algebraic number theory*, 2nd Edition, Graduate Texts in Math., 110, Springer, New York, 1994.
- [8] H. Pan, A remark on Zolotera's theorem, preprint, arXiv:0601026, 2006.
- [9] J. H. Silverman, *The Arithmetic of Elliptic Curves*, 2nd ed., Springer, New York, 1990.
- [10] J. R. Stembridge, Nonintersecting paths, pfaffians and plane partitions, *Adv. in Math.* 83 (1990), 96–131.
- [11] Z.-W. Sun, On some determinants with Legendre symbol entries, *Finite Fields Appl.* 56 (2019), 285–307.
- [12] Z.-W. Sun, Some determinants involving quadratic residues modulo primes, *Frontiers Math.*, in press.
- [13] H.-L. Wu, Determinants concerning Legendre symbols. *C. R. Math. Acad. Sci. Paris* 359 (2021), 651–655.
- [14] H.-L. Wu and L.-Y. Wang, The Gross-Koblitz formula and almost circulant matrices related to Jacobi sums, *Finite Fields Appl.* 103 (2025), Article 102581.

(HAI-LIANG WU) SCHOOL OF SCIENCE, NANJING UNIVERSITY OF POSTS AND TELECOMMUNICATIONS, NANJING 210023, PEOPLE'S REPUBLIC OF CHINA

Email address: whl.math@smail.nju.edu.cn

(HAO PAN) SCHOOL OF APPLIED MATHEMATICS, NANJING UNIVERSITY OF FINANCE AND ECONOMICS,
NANJING 210046, PEOPLE'S REPUBLIC OF CHINA

Email address: haopan79@zoho.com