

Degenerate Vertex Cuts in Sparse Graphs

Thilo Hartel

Johannes Rauch

Dieter Rautenbach

Institute of Optimization and Operations Research

Ulm University, Ulm, Germany

thilo.hartel@uni-ulm.de

johannes.rauch@mailbox.org

dieter.rautenbach@uni-ulm.de

Abstract

For a non-negative integer k , a vertex cut in a graph is k -degenerate if it induces a k -degenerate subgraph. We show that a graph of order n at least $2k + 2$ without a k -degenerate cut has the size at least $\frac{1}{2} \left(k + \Omega(\sqrt{k}) \right) n$ and that a graph of order n at least 5 without a 2-degenerate cut has the size at least $\frac{27n-35}{10}$. For $k \geq 2$, we show that a connected graph G of order n at least $k + 6$ and size m at most $\frac{k+3}{2}n + \frac{k-1}{2}$ has a minimum k -degenerate cut.

1 Introduction

We consider finite, simple, and undirected graphs and use standard terminology. A graph G is k -degenerate for some non-negative integer k if every non-empty subgraph H of G has a vertex u of degree $d_H(u)$ at most k . In particular, a graph is 0-degenerate if it has no edges, and 1-degenerate if it is a forest. A set S of vertices of a graph G is a *cut of G* if $G - S$ is disconnected.

In the present paper we study the existence of k -degenerate cuts in sparse graphs, which are cuts inducing k -degenerate subgraphs. Answering a question of Caro, it was shown by Chen and Yu [6] that every graph of order n with less than $2n - 3$ edges has an *independent cut*, which is a 0-degenerate cut. This seminal result with its elegant inductive proof led to considerable further research, cf. [7, 10–12] and the references therein. At the 7th C5 Graph Theory Workshop (Kurort Rathen, 2003) [13], Atsushi Kaneko proposed the conjecture that every graph of order n with less than $3n - 6$ edges has a *forest cut*, which is a 1-degenerate cut. This conjecture was rediscovered by Chernyshev et al. [9] who verified it for planar graphs and showed that every graph of order n with less than $(11n - 18)/5$ edges has a forest cut. This bound was improved to $(9n - 15)/4$ by Botler et al. [5] and to $(19n - 28)/8$ for $n \geq 4$ by Bogdanov et al. [4] but the conjecture remains open. Bogdanov et al. also consider the existence of *bipartite cuts*, which are cutsets inducing a 2-colorable subgraph.

Chen and Yu's result and Kaneko's conjecture suggest the conjecture that every graph of sufficiently large order n with less than

$$(k+2)n - \binom{k+3}{2}$$

edges has a k -degenerate cut, which would be best possible in view of the join of K_{k+2} and \overline{K}_{n-k-2} . Unfortunately, this naive conjecture turns out to be quite false. Modifying a construction of Bessy et

al. [3] based on results of Axenovich et al. [2], Aubian et al. [1] construct graphs of arbitrarily large order n and less than

$$\frac{1}{2} \left(k + O \left(\sqrt{k \ln(k)} \right) \right) n$$

edges that do not have a k -degenerate cut or even a cut inducing a $(k+1)$ -colorable subgraph. Note that a graph G of order at least $k+2$ that does not have a k -degenerate cut has minimum degree at least $k+2$, because, otherwise, the neighborhood $N_G(u)$ of a vertex u of minimum degree is a k -degenerate cut. This trivial argument already implies that graphs of sufficiently large order n that do not have a k -degenerate cut have at least $\frac{1}{2}(k+2)n$ edges.

Our first contribution is the following result.

Theorem 1. *Let k be a positive integer. If G is a graph of order n at least $2k+2$ without a k -degenerate cut, then the size of G is at least $\frac{1}{2} \left(k + \frac{13\sqrt{k}}{190} - \frac{1}{38} \right) n$.*

The proof of Theorem 1 only considers vertices at distance at most 2 from small degree vertices, that is, it relies on a relatively local argument. Nevertheless, the results of Aubian et al. [1] imply that Theorem 1 is best possible up to a factor of order $\sqrt{\ln(k)}$ at the \sqrt{k} -term.

For $k=2$, we improve Theorem 1 as follows.

Theorem 2. *If G is a graph of order n at least 5 without a 2-degenerate cut, then the size of G is at least $\frac{27n-35}{10}$.*

The old and new results discussed so far concerned k -degenerate cuts of arbitrary order. Already Chen et al. [7] study under which conditions there are small independent cuts. In particular, for $s \leq 3$ and any positive integer n , they determine the largest value of m — up to a small additive error for $s=3$ — such that a graph of order n and size at most m has an independent cut of order at most s . Recently, Cheng et al. [8] consider the existence of minimum cuts that are k -degenerate for $k \in \{0, 1\}$. In particular, they show that every connected graph of order $n \geq 7$ and size at most $\lfloor \frac{3n}{2} \rfloor$ has an independent minimum cut, and that every connected graph of order $n \geq 7$ and size at most $2n$ has a minimum cut that induces a forest. In both cases, the bounds on the size are best possible.

Our third result concerns the existence of minimum cuts that are k -degenerate.

Theorem 3. *Let k be an integer at least 2. If G is a connected graph of order n at least $k+6$ and size m at most $\frac{k+3}{2}n + \frac{k-1}{2}$, then G has a minimum k -degenerate cut.*

Similarly as in [8], an explicit construction shows that Theorem 3 is best possible.

All proofs are given in the following section.

2 Proofs

A k -core in a graph G is a set C of vertices of G inducing a subgraph with no vertex of degree at most k , that is, the minimum degree $\delta(G[C])$ of the subgraph $G[C]$ of G induced by C is at least $k+1$. Clearly, every k -core contains at least $k+2$ vertices, the only k -core of order exactly $k+2$ is a clique, and a graph G is k -degenerate if and only if it contains no k -core.

We proceed to the proof of Theorem 1.

Proof of Theorem 1. Let k , G , and n be as in the statement. Since G has order at least $k+2$ and no k -degenerate cut, the minimum degree of G is at least $k+2$.

Claim 1. *If u is a vertex of G of degree at most $k + \frac{\sqrt{k}}{5}$, then u has least $k - \frac{2k}{25} - \frac{2\sqrt{k}}{5}$ neighbors of degree at least $k + \frac{\sqrt{k}}{5}$.*

Proof of Claim 1. Let u be a vertex of G of degree $d_G(u)$ at most $k + \frac{\sqrt{k}}{5}$. Let $R = V(G) \setminus N_G[u]$. Let X be a set of $\left\lfloor \frac{2\sqrt{k}}{5} \right\rfloor$ neighbors of u minimizing the number of edges between X and R . Let $Y = N_G(X) \setminus N_G[u]$ and $Z = N_G(u) \setminus X$. Since G has no k -degenerate cut and $n > 1 + d_G(u)$, there is a k -core C in G contained in $N_G(u)$.

If some vertex in X has at least $\frac{\sqrt{k}}{5}$ neighbors in R , then, by the choice of X , all vertices in Z have at least $\frac{\sqrt{k}}{5}$ neighbors in R . Now, each of the

$$|C \cap Z| \geq |C| - |X| \geq k + 2 - \left\lfloor \frac{2\sqrt{k}}{5} \right\rfloor$$

vertices in $C \cap Z$ has degree at least $k + \frac{\sqrt{k}}{5}$, which completes the proof of the claim. Hence, we may assume that every vertex in X has less than $\frac{\sqrt{k}}{5}$ neighbors in R , which implies

$$|Y| < \frac{\sqrt{k}}{5} \cdot |X| \leq \frac{\sqrt{k}}{5} \cdot \frac{2\sqrt{k}}{5} = \frac{2k}{25}.$$

If Z contains a set Z' of at least $k - \frac{2k}{25}$ vertices that have at least $\frac{\sqrt{k}}{5}$ neighbors in R , then each of the

$$\begin{aligned} |C \cap Z'| &\geq |C| - |N_G(u) \setminus Z'| \\ &= |C| - |N_G(u)| + |Z'| \\ &\geq (k + 2) - \left(k + \frac{\sqrt{k}}{5} \right) + \left(k - \frac{2k}{25} \right) \\ &> k - \frac{2k}{25} - \frac{2\sqrt{k}}{5} \end{aligned}$$

vertices in $C \cap Z'$ has degree at least $k + \frac{\sqrt{k}}{5}$, which completes the proof of the claim. Hence, we may assume that

$$Z \text{ contains less than } k - \frac{2k}{25} \text{ vertices that have at least } \frac{\sqrt{k}}{5} \text{ neighbors in } R. \quad (1)$$

Note that

$$|Z| = d_G(u) - |X| \leq k + \frac{\sqrt{k}}{5} - \left\lfloor \frac{2\sqrt{k}}{5} \right\rfloor \leq k - \frac{\sqrt{k}}{5} + 1.$$

Since G has no k -degenerate cut and

$$1 + |X| + |Y| + |Z| = 1 + d_G(u) + |Y| < 1 + k + \frac{\sqrt{k}}{5} + \frac{2k}{25} < n,$$

the set $Y \cup Z$ is a cut, and, hence, there is a k -core C' in G contained in $Y \cup Z$. Since every vertex in Z has at most $|Z| - 1 \leq k - \frac{\sqrt{k}}{5}$ neighbors in Z , every vertex in $C' \cap Z$ has at least $k + 1 - \left(k - \frac{\sqrt{k}}{5} \right) > \frac{\sqrt{k}}{5}$ neighbors in $Y \subseteq R$. Hence, the set Z contains at least

$$|C' \setminus Y| \geq |C'| - |Y| \geq k + 2 - \frac{2k}{25}$$

vertices that have at least $\frac{\sqrt{k}}{5}$ neighbors in R . This contradicts (1), which completes the proof of the claim. \square

Let V_ℓ be the set of vertices of G of degree at least $k + \frac{\sqrt{k}}{5}$ and let $V_s = V(G) \setminus V_\ell$. We use a discharging argument to obtain a lower bound on the degree sum of G . Initially, each vertex has charge equal to its degree. Now, each vertex in V_ℓ sends a charge of $\frac{5}{38\sqrt{k}}$ to its neighbors in V_s . By Claim 1, every vertex in V_s has at least $k - \frac{2k}{25} - \frac{2\sqrt{k}}{5} \geq \frac{13k}{25}$ neighbors in V_ℓ . Therefore a vertex in V_s receives from its large degree neighbors a total amount of charge of at least $\frac{5}{38\sqrt{k}} \cdot \frac{13k}{25} = \frac{13\sqrt{k}}{190}$, meaning that it has a final charge of at least $k + \frac{13\sqrt{k}}{190}$. On the other hand, a vertex in V_ℓ with degree d sends a total amount of charge of at most $d \cdot \frac{5}{38\sqrt{k}}$, implying that it has a final charge of at least

$$d \cdot \left(1 - \frac{5}{38\sqrt{k}}\right) \geq \left(k + \frac{\sqrt{k}}{5}\right) \cdot \left(1 - \frac{5}{38\sqrt{k}}\right) = k + \frac{13\sqrt{k}}{190} - \frac{1}{38}.$$

Summing the total final charge, we obtain

$$2m = \sum_{v \in V(G)} d_G(v) \geq \left(k + \frac{13\sqrt{k}}{190} - \frac{1}{38}\right) n,$$

which completes the proof. \square

For a vertex u in a graph G and a set U of vertices of G , let $d_U(u) = |U \cap N_G(u)|$ denote the number of neighbors of u in U .

Proof of Theorem 2. Suppose, for a contradiction, that G is a counterexample of minimum order n . In particular, the graph G has order $n \geq 5$, no 2-degenerate cut but less than $\frac{27n-35}{10}$ edges. If $n = 5$, then G has less than $\frac{27 \cdot 5 - 35}{10} = 10$ edges, which implies that G has two non-adjacent vertices, say u and v . Now, the set $V(G) \setminus \{u, v\}$ is a 2-degenerate cut, which is a contradiction. Hence, we have $n \geq 6$. If G has a cut S of order at most 4, then it follows that S has order exactly 4 and induces a K_4 . Let K be a component of $G - S$. Let $G_1 = G[K \cup S]$ and $G_2 = G - V(K)$. Clearly, the two graphs G_1 and G_2 have orders n_1 and n_2 , respectively, at least 5. If G_1 or G_2 has a 2-degenerate cut S' , then, since S is complete, the set S' is also a 2-degenerate cut in G , which is a contradiction. Hence G_1 and G_2 both have no 2-degenerate cut. By the choice of G , it follows that G_1 and G_2 have at least $\frac{27n_1-35}{10}$ and $\frac{27n_2-35}{10}$ edges, respectively. This implies that G has at least $\frac{27n_1-35}{10} + \frac{27n_2-35}{10} - 6 > \frac{27n-35}{10}$ edges, which is a contradiction. Hence, the vertex connectivity κ of G is at least 5. If $n \leq 17$, then the average degree of G is less than $\frac{27 \cdot 17 - 35}{5 \cdot 17} < 5$. This implies that the minimum degree, and, hence, also the vertex connectivity of G is at most 4, which is a contradiction. Hence, we have $n \geq 18$. Let u be a vertex of minimum degree in G . Since $5 \leq \kappa \leq d_G(u) < \frac{27 \cdot n - 35}{5 \cdot n} \leq \frac{27}{5}$, the degree of u is exactly 5.

Claim 2. $\sum_{v \in N_G(u)} d_G(v) \geq 29$.

Proof of Claim 2. Let $N_G(u) = \{v_1, v_2, v_3, v_4, v_5\}$. Let X be the set of vertices in $N_G(u)$ with at most one neighbor outside of $N_G(u)$. Let $\ell = |X|$. By symmetry, we may assume that $X = \{v_1, \dots, v_\ell\}$. If some vertex $v \in X$ has no neighbor outside of $N_G(u)$, then $N_G(u) \setminus \{v\}$ is a cut of order 4, which is a contradiction. If two vertices $v, v' \in X$ have the same neighbor w outside of $N_G(u)$, then $(N_G(u) \setminus \{v, v'\}) \cup \{w\}$ is a cut of order 4, which is a contradiction. Hence, the vertices v_1, \dots, v_ℓ have

distinct unique neighbors w_1, \dots, w_ℓ outside of $N_G[u]$, respectively. Since $N_G(u)$ is not a 2-degenerate cut, the set $N_G(u)$ contains a 2-core. We consider two cases.

Case 1. $G[N_G(u)]$ has minimum degree at least 3.

This implies that at least 8 edges have both endpoints in $N_G(u)$. If $|X| \geq 4$, then $\{v_1, v_2, w_3, w_4, v_5\}$ is a 2-degenerate cut, which is a contradiction. If $|X| \leq 2$, then

$$\sum_{v \in N_G(u)} d_G(v) \geq 5 + 2 \cdot 8 + 3 \cdot 2 + 2 = 29.$$

Hence, we may assume that $|X| = 3$. If v_1 is not adjacent to v_2 or v_4 , then $\{v_1, v_2, w_3, v_4, v_5\}$ is a 2-degenerate cut, which is a contradiction. By symmetry, this implies that $\{v_1, v_2, v_3\}$ is complete and that there are all possible 6 edges between $\{v_1, v_2, v_3\}$ and $\{v_4, v_5\}$. It follows that at least 9 edges have both endpoints in $N_G(u)$ and, hence,

$$\sum_{v \in N_G(u)} d_G(v) \geq 5 + 2 \cdot 9 + 2 \cdot 2 + 3 = 30.$$

Case 2. $G[N_G(u)]$ has minimum degree at most 2.

Since $N_G(u)$ contains a 2-core, we may assume, by symmetry, that v_1, v_2, v_3 , and v_4 induce a K_4 , and that v_5 has at most two neighbors in $N_G(u)$. Note that $v_5 \notin X$. If $d_{N_G(u)}(v_5) \geq \ell$, then

$$\begin{aligned} \sum_{v \in N_G(u)} d_G(v) &\geq 5 + (12 + 2 \cdot d_{N_G(u)}(v_5)) + 2(4 - \ell) + \ell + (4 - d_{N_G(u)}(v_5)) \\ &= 29 + d_{N_G(u)}(v_5) - \ell \geq 29. \end{aligned}$$

Hence, we may assume $d_{N_G(u)}(v_5) < \ell$. If $d_{N_G(u)}(v_5) = 1$, then $\ell \geq 2$ and $\{w_1, v_2, v_3, v_4, v_5\}$ is a 2-degenerate cut, which is a contradiction. If $d_{N_G(u)}(v_5) = 2$, then $\ell \geq 3$. By symmetry, we may assume that v_5 and v_1 are adjacent. Again, the set $\{w_1, v_2, v_3, v_4, v_5\}$ is 2-degenerate cut, which is a contradiction. If $d_{N_G(u)}(v_5) = 0$ and $\ell \geq 2$, then $\{w_1, v_2, v_3, v_4, v_5\}$ is a 2-degenerate cut, which is a contradiction. Hence, we may assume that $d_{N_G(u)}(v_5) = 0$ and $\ell = 1$. If v_2 is not adjacent to w_1 , then $\{w_1, v_2, v_3, v_4, v_5\}$ is a 2-degenerate cut, which is a contradiction. Hence, by symmetry, we may assume that w_1 is adjacent to v_2, v_3 , and v_4 . If v_2 has more than 2 neighbors outside of $N_G[u]$, the claim follows. Hence, by symmetry, we may assume that v_2, v_3 , and v_4 all have exactly one neighbor w_2, w_3 , and w_4 outside of $N_G[u] \cup \{w_1\}$, respectively. If $w_2 = w_3$, then $\{w_1, w_2, v_4, v_5\}$ is a cut of order 4, which is a contradiction. Hence, by symmetry, the vertices w_2, w_3 , and w_4 are distinct. Now, the set $\{w_1, w_2, v_3, v_4, v_5\}$ is a 2-degenerate cut, which is a contradiction. This completes the proof of the claim. \square

Let V_i be the set of vertices of G of degree i for $i \in \{5, 6, 7, 8\}$, and let V_9 be the set of vertices of G of degree at least 9. We use a discharging argument to obtain a lower bound on the degree sum of G . Initially, each vertex has charge equal to its degree. Then each vertex u in V_i for $i \in \{6, 7, 8, 9\}$ sends a charge of $\frac{i-5}{10}$ to each neighbor. By Claim 2, a vertex u in V_5 receives at least 4 times $\frac{1}{10}$ charge from its neighbors. This implies that the final charge of u is at least $\frac{27}{5}$. The final charge of a

vertex u in $V_6 \cup V_7 \cup V_8$ is at least $d_G(u) \left(1 - \frac{d_G(u)-5}{10}\right) \geq \frac{27}{5}$. The final charge of a vertex u in V_9 is at least $9 \left(1 - \frac{2}{5}\right) = \frac{27}{5}$. Summing the total final charge, we obtain $2m = \sum_{v \in V(G)} d_G(v) \geq \frac{27n}{5}$, which is a contradiction and completes the proof. \square

Our next goal is the proof of Theorem 3, which requires some preparatory results. Recall that every vertex in a minimum cut S of a graph G has a neighbor in each component of $G - S$. For a graph G , let $\kappa(G)$ denote the vertex connectivity of G .

Lemma 4. *Let k be an integer at least 2. If G is a connected graph of order n at least $k+6$ vertices, size m at most $\frac{k+3}{2}n + \frac{k-1}{2}$, and minimum degree exactly $k+3$, then G has a minimum k -degenerate cut.*

Proof. Let G be as in the statement. If $\kappa(G) \leq k+1$, then every minimum cut of G is k -degenerate. Hence, we may assume that $\kappa(G) \in \{k+2, k+3\}$. Let X be the set of all vertices of minimum degree in G . We have

$$(k+3)n + \sum_{u \in V(G) \setminus X} (d_G(u) - (k+3)) = 2m \leq (k+3)n + (k-1),$$

which implies

$$n - |X| \leq \sum_{u \in V(G) \setminus X} (d_G(u) - (k+3)) \leq k-1 \text{ and, hence,} \quad (2)$$

$$|X| \geq n - (k-1). \quad (3)$$

Let S be a minimum cut of G and suppose that S is not k -degenerate. Since $|S| \in \{k+2, k+3\}$, it follows from (3) that $|X \cap S| \geq |S| - (k-1) \geq 3$. Since S contains a k -core, there is a vertex v in $X \cap S$ such that v has at least $k+1$ neighbors in S and, hence, at most two neighbors outside of S . Since v has a neighbor in each component of $G - S$, the graph $G - S$ has exactly two components G_1 and G_2 . We may assume that $n(G_1) \geq n(G_2)$, which implies $n(G_1) \geq 2$ as $n \geq k+6$.

We consider distinct cases.

Case 1. $\kappa(G) = k+2$.

In this case, we have $G[S] = K_{k+2}$. Let $v \in X \cap S$. The vertex v has exactly one neighbor w in $V(G_1)$. If w has a neighbor v' in $X \cap S$ distinct from v , then w is the unique neighbor of v' in $V(G_1)$, which implies the contradiction that $(S \setminus \{v, v'\}) \cup \{w\}$ is a cut that is smaller than S . Since $X \cap S$ contains at least 3 vertices, the vertex w has a non-neighbor in S . It follows $(S \setminus \{v\}) \cup \{w\}$ is a minimum cut that is not complete, and, hence, k -degenerate cut, which completes the proof in this case.

Case 2. $\kappa(G) = k+3$ and S is a k -core.

In this case, we have $|X \cap S| \geq 4$. Let $v_1, v_2, v_3, v_4 \in X \cap S$. By the same argument as in Case 1, for each $i \in \{1, 2, 3, 4\}$, the vertex v_i has exactly one neighbor w_i in $V(G_1)$ and the vertex w_i has no neighbor in $(X \cap S) \setminus \{v_i\}$. In particular, the vertices w_1, w_2, w_3, w_4 are distinct, which implies

$n(G_1) \geq 4$. Counting the edges between $\{w_1, w_2, w_3, w_4\}$ and S , we obtain

$$\begin{aligned} d_S(w_1) + d_S(w_2) + d_S(w_3) + d_S(w_4) &\leq \sum_{v \in S} \left(d_G(v) - \underbrace{(k+1)}_{\leq d_S(v)} - \underbrace{1}_{\leq d_{V(G_2)}(v)} \right) \\ &= |S| + \sum_{v \in S \setminus X} \left(d_G(v) - (k+3) \right) \\ &\stackrel{(2)}{\leq} (k+3) + (k-1) = 2k+2. \end{aligned}$$

By symmetry, we may assume that $d_S(w_1), d_S(w_2) \leq k$, which implies that w_1 and w_2 both have at most $k-1$ neighbors in $S \setminus \{v_1, v_2\}$. Now, the set $(S \setminus \{v_1, v_2\}) \cup \{w_1, w_2\}$ is a minimum cut that is k -degenerate, which completes the proof in this case.

Case 3. $\kappa(G) = k+3$ and S is not a k -core.

Again, we have $|X \cap S| \geq 4$. Since S has order $k+3$, is not a k -core, but contains a k -core, there is a vertex u in S such that $d_S(u) \leq k$ and $S \setminus \{u\}$ is a clique. Since each vertex in $S \setminus \{u\}$ has a neighbor in $V(G_1)$, a neighbor in $V(G_2)$, and $k+1$ neighbors in $S \setminus \{u\}$, the at least 3 vertices in $(X \cap S) \setminus \{u\}$ are not adjacent to u . In particular, we have $d_S(u) \leq k-1$. Let $v \in (X \cap S) \setminus \{u\}$. As before, it follows that the vertex v has a unique neighbor w in $V(G_1)$, and w has no neighbor in $(X \cap S) \setminus \{u, v\}$. It follows that $(S \setminus \{v\}) \cup \{w\}$ is a minimum cut that is k -degenerate, which completes the proof in this final case. \square

We are now in a position to complete the proof of Theorem 3.

Proof of Theorem 3. Suppose, for a contradiction, that the theorem is false. Let G be a counterexample of minimum order n , that is, $n \geq k+6$, the size m of G is at most $\frac{k+3}{2}n + \frac{k-1}{2}$ but G has no minimum k -degenerate cut, which implies that $\kappa(G) \geq k+2$. The average degree $\frac{2m}{n}$ of G satisfies $\frac{2m}{n} \leq k+3 + \frac{k-1}{n} < k+4$, which implies $\delta(G) \leq k+3$. Lemma 4 implies $\delta(G) \leq k+2$. Since $\kappa(G) \leq \delta(G)$, we obtain $\kappa(G) = \delta(G) = k+2$. Let v be a vertex of minimum degree. The neighborhood $S = N_G(v)$ of v is a minimum cut, which implies $G[S] = K_{k+2}$.

First, suppose that $n = k+6$. Let $R = \{x, y, z\} = V(G) \setminus N_G[v]$. The number m' of edges incident with $\{x, y, z\}$ satisfies

$$m' \leq \frac{(k+3)(k+6)}{2} + \frac{k-1}{2} - \binom{k+3}{2} = \frac{5}{2}k + \frac{11}{2} \stackrel{k \geq 2}{<} 3k+5. \quad (4)$$

If $G[R]$ has at most one edge, then $m' \geq 3\delta(G) - 1 = 3(k+2) - 1$, which contradicts (4). If $G[R]$ contains at exactly two edges, say xy and xz , then (4) implies that x has degree $k+2$, and $N_G(x)$ is a minimum k -degenerate cut, which is a contradiction. It follows that R is complete. By (4), we may assume that $d_G(x) = d_G(y) = k+2$ and $d_G(z) \leq k+3$. Since $N_G(x)$ is not a k -degenerate cut, $N_G(x)$ is complete, which implies $N_G(x) \cap S = N_G(y) \cap S \subseteq N_G(z) \cap S$. Now, the set $N_G(z) \cap S$ is a cut of order at most $k+1$, which contradicts $\kappa(G) = k+2$. Hence, we may assume that $n > k+6$.

The graph $G' = G - v$ has order $n' = n-1 \geq k+6$ and size $m' = m - (k+2) < \frac{k+3}{2}(n-1)$. By the choice of G , the graph G' has a minimum k -degenerate cut S' . Since $|S'| \leq \kappa(G') \leq \delta(G') \leq k+2 = \kappa(G)$ and S is complete, the set S' is also a minimum k -degenerate cut in G . This final contradiction completes the proof. \square

We construct graphs showing that Theorem 3 is best possible. For integers $k \geq 2$ and $s \geq 3$, let G arise from the union of s disjoint $(k+2)$ -cliques C_0, \dots, C_{s-1} arranged in a cyclic order by

- adding sets M_0, \dots, M_{s-1} of edges, where M_i is a matching of size $k+2$ between C_i and C_{i+1} , and
- adding one further vertex u as well as all $k+2$ possible edges between u and C_0 .

See Figure 1 for an illustration. The only minimum cut of G is $N_G(u) = C_0$, which is not k -degenerate. The order of G is $n = (k+2)s+1$ and the size is $m = \left(\frac{k+3}{2}n + \frac{k-1}{2}\right) + 1$.




Figure 1: The constructed graph G for $k = 4$, $s = 12$, and a specific choice of the matchings M_i .

Acknowledgment The authors were partially supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – project number 545935699.

References

- [1] G. Aubian, M. Bonamy, R. Bourneuf, O. Fontaine, and L. Picasarri-Arrieta, On cuts of small chromatic number in sparse graphs, arXiv:2510.01791v1.
- [2] M. Axenovich, J.-S. Sereni, R. Snyder, and L. Weber, Bipartite independence number in graphs with bounded maximum degree, SIAM J. Discrete Math. **35** (2021), no. 2, 1136–1148.
- [3] S. Bessy, J. Rauch, D. Rautenbach, and U.S. Souza, Sparse vertex cutsets and the maximum degree, Electron. J. Combin. **32** (2025), no. 2, Paper No. 2.32, 11 pp..
- [4] I.I. Bogdanov, E. Neustroeva, G. Sokolov, A. Volostnov, N. Russkin, and V. Voronov, On forest and bipartite cuts in sparse graphs, arXiv:2505.16179.
- [5] F. Botler, Y.S. Couto, C.G. Fernandes, E.F. de Figueiredo, R. Gómez, V.F. dos Santos, and C.M. Sato, Extremal Problems on Forest Cuts and Acyclic Neighborhoods in Sparse Graphs, arXiv:2411.17885.

- [6] G. Chen and X. Yu, A note on fragile graphs, *Discrete Math.* **249** (2002), no. 1-3, 41–43.
- [7] G. Chen, R.J. Faudree, and M.S. Jacobson, Fragile graphs with small independent cuts, *J. Graph Theory* **41** (2002), no. 4, 327–341.
- [8] K. Cheng, Y. Tang, and X. Zhan, Sparse graphs with an independent or foresty minimum vertex cut, *Discrete Math.* **349** (2026), no. 1, Paper No. 114658, 6 pp..
- [9] V. Chernyshev, J. Rauch, and D. Rautenbach, Forest cuts in sparse graphs, *Discrete Math.* **348** (2025), no. 11, Paper No. 114594, 6 pp..
- [10] V. Chernyshev, J. Rauch, D. Rautenbach, and L. Redina, A faster algorithm for independent cut, *Theoret. Comput. Sci.* **1054** (2025), Paper No. 115542, 7 pp..
- [11] V.B. Le and F. Pfender, Extremal graphs having no stable cutsets, *Electron. J. Combin.* **20** (2013), no. 1, Paper 35, 7 pp..
- [12] J. Rauch and D. Rautenbach, Revisiting Extremal Graphs Having No Stable Cutsets, *Electron. J. Combin.* **32** (2025), no. 4, P4.25.
- [13] I. Schiermeyer, personal communication, November 19, 2024.