

A Sieve M-Estimator for Entropic Optimal Transport

Rami V. Tabri

Department of Econometrics and Business Statistics, Monash University,
Clayton, Victoria, Australia,
Email: rami.tabri@monash.edu.

January 5, 2026

Abstract

The entropically regularized optimal transport problem between probability measures on compact Euclidean subsets can be represented as an information projection with moment inequality constraints. This allows its Fenchel dual to be approximated by a sequence of convex, finite-dimensional problems using sieve methods, enabling tractable estimation of the primal value and dual optimizers from samples. Assuming only continuity of the cost function, I establish almost sure consistency of these estimators. I derive a finite-sample convergence rate for the primal value estimator, showing logarithmic dependence on sieve complexity, and quantify uncertainty for the dual optimal value estimator via matching stochastic bounds involving suprema of centered Gaussian processes. These results provide the first statistical guarantees for sieve-based estimators of entropic optimal transport, extending beyond the empirical Sinkhorn approach.

1 Introduction

Optimal transport (Villani, 2009) has emerged as a fundamental tool in modern statistics, supporting methodologies for distributional comparison, generative modeling, domain adaptation, and causal inference. Given two Borel probability spaces, $(\mathcal{X}, \mathcal{B}(\mathcal{X}), P_X)$ and $(\mathcal{Y}, \mathcal{B}(\mathcal{Y}), P_Y)$, with $\mathcal{X} \subset \mathbb{R}^{d_x}$ and $\mathcal{Y} \subset \mathbb{R}^{d_y}$, and a cost function $c : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R}_+$, the optimal transport problem is an infinite-dimensional linear optimization problem

$$\inf_{P \in \Pi(P_X, P_Y)} \int_{\mathcal{X} \times \mathcal{Y}} c(x, y) dP, \quad (1.1)$$

where the infimum is taken over the set $\Pi(P_X, P_Y)$ of couplings between P_X and P_Y . Central to its recent success in practice is the paradigm of entropic regularization, popularized by Cuturi (2013), enabling computational advances for data-rich applications in areas like machine learning or image processing (Peyré and Cuturi, 2019). For $\gamma > 0$, the entropically regularized optimal transport (EOT) problem is

$$\inf_{P \in \Pi(P_X, P_Y)} \int_{\mathcal{X} \times \mathcal{Y}} c(x, y) dP + \gamma^{-1} H(P | P_X \otimes P_Y), \quad (1.2)$$

where γ is the entropic regularization parameter, and $H(\cdot|P_X \otimes P_Y)$ denotes the relative entropy with respect to the product measure

$$H(P|P_X \otimes P_Y) = \begin{cases} \int_{\mathcal{X} \times \mathcal{Y}} \frac{dP}{d(P_X \otimes P_Y)} \log \left(\frac{dP}{d(P_X \otimes P_Y)} \right) d(P_X \otimes P_Y) & \text{if } P \ll P_X \otimes P_Y \\ \infty & \text{elsewhere.} \end{cases}$$

For fixed γ , the problem (1.2) yields a regularized optimal value, dual potentials, and a transport plan that approximate those of the problem (1.1), with large values of γ providing more accurate approximations (e.g., Léonard, 2012).

Entropic regularization enables scalable computation and smooths the geometry of the transport problem, and it is now routinely used in statistical practice via the empirical Sinkhorn divergence. Given samples, X_1, \dots, X_N i.i.d. from P_X and Y_1, \dots, Y_N i.i.d. from P_Y , and their empirical distributions, \hat{P}_X and \hat{P}_Y , respectively, the empirical Sinkhorn divergence solves the sample analogue of the EOT problem (1.2): it replaces the population distributions P_X and P_Y in the EOT problem with \hat{P}_X and \hat{P}_Y , respectively, and then applies the Sinkhorn algorithm to approximate the regularized optimal value, transport plan, and dual potentials. Despite its widespread use, relatively little is known about the statistical properties of the EOT problem beyond the narrow, though important, class of smooth cost functions.

Existing theory offers only partial answers: central limit theorems are available for quadratic costs and sub-Gaussian probability measures (e.g., del Barrio et al., 2023; Goldfeld et al., 2024), and more general cost functions have been analyzed only in discrete and semi-discrete settings (e.g., Bigot and Klein, 2019; Bercu and Bigot, 2021). Furthermore, sample-complexity bounds have been established under smooth costs and compactly supported measures (e.g., Genevay et al., 2019) and for quadratic costs with sub-Gaussian measures (e.g., Mena and Niles-Weed, 2019), ruling out many cost functions used in applications such as economics (Galichon, 2016; Chiappori and Salanié, 2016), imaging (Peyré and Cuturi, 2019), and machine learning (Arjovsky et al., 2017; Flamary and Courty, 2021). On the other hand, methods based on reproducing kernel Hilbert space (RKHS) parameterizations (Genevay et al., 2016) or neural networks (Seguy et al., 2018) are computationally flexible and accommodate general continuous cost functions, but impose continuity of the dual potentials as a structural assumption, motivated by optimization considerations. These methods lack statistical guarantees beyond consistency, which holds only under the correct specification of the potentials' continuity. As a result, the statistical behavior of EOT estimators under general continuous costs and general marginal distributions remains essentially unknown. This paper addresses this gap in the setting of compactly supported marginal distributions.

I develop a nonparametric estimation framework for the EOT value under minimal conditions: the cost function is continuous and the marginal distributions have compact support. The method is grounded in a representation of the EOT problem as an information projection under moment constraints—a classical perspective (e.g., Csiszár, 1975) that, to the best of my knowledge, has not previously been exploited for statistical estimation of entropic optimal transport. This formulation enables the use of sieve methods and empirical process theory to approximate the Fenchel dual of the information-projection problem. The resulting estimator is computationally tractable and applies to continuous cost functions with arbitrary compactly

supported marginal distributions, thereby extending statistical optimal transport to settings not covered by existing theory.

The present analysis develops a stochastic approximation scheme for the EOT problem (1.2) under the following minimal conditions.

Assumption 1. (i) \mathcal{X} and \mathcal{Y} are compact, (ii) $c : \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R}_+$ is continuous.

Under Assumption 1, the EOT problem (1.2) admits a unique solution, denoted by P_γ . Rewriting the objective as

$$\gamma^{-1} \log a_\gamma + \gamma^{-1} H(P \mid R_\gamma), \quad (1.3)$$

where the reference probability R_γ satisfies

$$\frac{dR_\gamma}{d(P_X \otimes P_Y)}(x, y) := \exp\{-\gamma c(x, y)\} / a_\gamma, \quad (x, y) \in \mathcal{X} \times \mathcal{Y}$$

with $a_\gamma := \int_{\mathcal{X} \times \mathcal{Y}} \exp\{-\gamma c(x, y)\} d(P_X \otimes P_Y)(x, y)$, reveals that P_γ is also the solution to the information projection (I -projection) problem

$$\min \{H(P \mid R_\gamma) : P \in \Pi(P_X, P_Y)\}. \quad (1.4)$$

The stochastic approximation scheme developed in this paper applies the *method of sieves* (Grenander, 1981) to the Fenchel dual of the I -projection problem (1.4). The construction rests on three main ideas:

- (a) reformulating the marginal constraints in (1.4) as moment inequality restrictions;
- (b) reparametrizing the Fenchel dual so that it admits finite-dimensional convex approximations; and
- (c) estimating these finite-dimensional programs via the *Sample Average Approximation* method (Shapiro et al., 2009).

Together, these steps yield a sieve M-estimator for the Fenchel dual problem. The approximation framework of Tabri (2025) provides a convergent sieve construction for implementing steps (a) and (b). The key insight behind step (a) is that cumulative distribution functions characterize probability measures supported on subsets of Euclidean space, allowing us to rewrite moment equality constraints as pairs of moment inequalities. This reveals that the resulting class of moment functions is uniformly bounded and Vapnik-Chervonenkis (VC), and hence suitably precompact for adapting the general results of Tabri (2025). Unlike existing approaches based on discretization or parametric approximation of dual potentials, this construction enforces the defining constraints of the EOT problem at the population level and uses the sample only to approximate expectations in the dual objective.

That framework reparametrizes the Fenchel dual so that the dual variable becomes a *Gelfand–Pettis integral* (Gelfand, 1936; Pettis, 1938): a weak vector-valued integral with respect to a positive Radon measure supported on the $L_1(R_\gamma)$ -closed convex hull of the moment functions defining the inequality constraints.

This representation is advantageous because it admits approximation by Riemann sums in the $L_1(R_\gamma)$ norm, forming the basis of the sieve construction in Tabri (2025). Specializing this scheme to the present setting produces a sequence of finite-dimensional convex stochastic programs with a common objective and expanding domains. Here, these domains are finite-dimensional spaces generated by selected moment functions, which remain VC because they consist of linear combinations of elements from a VC class.

To apply the SAA method, I begin by solving sample analogues of the sieve-based approximating programs. Because their domains are naturally unbounded, I restrict them in a way that still allows expansion with the sieve approximation. I justify this restriction by connecting the Fenchel dual problem's optimizers to Schrödinger potentials (Nutz and Wiesel, 2022), and I establish that these optimizers remain uniformly bounded under Assumption 1. With these restrictions, I show that when the sieve grows slowly enough relative to the sample size, a strong uniform law of large numbers (ULLN) holds for the corresponding empirical processes. This ULLN guarantees that optimizing the sample-based objective over the sieve is asymptotically equivalent to optimizing its expectation, the Fenchel dual objective. I provide the precise statement and proof of this ULLN in Appendix C. Notably, it accommodates sequences of function classes whose index sets evolve with sample size—a situation not addressed by existing ULLN results, to the best of my knowledge.

1.1 Summary of Contributions

The contributions of this paper are threefold.

1. I establish the almost sure convergence of the proposed stochastic approximation scheme. In particular, under random sampling from R_γ with sample size N ,

- (i) the optimal value of the dual problem is consistently estimated by the optimal values of a sequence of finite-dimensional SAA convex programs, and
- (ii) every weak-star accumulation point of the corresponding sequence of optimal solutions is an optimal solution of the dual problem,

as $N \rightarrow \infty$, with probability 1.

2. Using empirical process techniques, I derive a finite-sample rate in terms of mean-convergence for the estimator of the EOT value. The rate separates stochastic variation from sieve approximation error, yielding a nonparametric bias-variance trade-off. The stochastic component is controlled by Massart's finite class lemma (Massart, 2000), giving a bound $\sqrt{2 \log(n)/N}$, where n is the dimension of the approximating sieve function class. This yields rates slower than the classical \sqrt{N} benchmark—typical in nonparametric estimation—while exhibiting unusually mild dependence on model complexity due to the logarithmic factor.

3. I establish matching upper and lower stochastic bounds for the estimation error of the dual problem's optimal value, with leading terms given by the suprema of centered Gaussian processes indexed by the sieve class, divided by \sqrt{N} . These results follow from the Gaussian approximation of suprema of empirical processes in Chernozhukov et al. (2014b) and can be used to construct interval estimators for the EOT value. Importantly, all results require only continuity of the cost and compact supports, making the methodology applicable in settings for which no statistical theory currently exists.

1.2 Organization of the paper

Section 2 develops the moment-inequality representation of the EOT problem and establishes key structural properties of the associated Fenchel dual problem via a sequence of propositions. These results lay the analytical foundation for the statistical analysis. Section 3 introduces the sieve M-estimator and states the main result of the paper, Theorem 1, which establishes consistency, the convergence rate, and stochastic bounds. Section 4 discusses the scope of the main result, its implications for statistical optimal transport, and potential extensions. The Appendix collects proofs of the theorem and all propositions to streamline the exposition, as well as all technical lemmas, derivations, and a numerical illustration.

2 Moment Inequality Information Projection Problem and Approximation

Section 3 of Csiszár (1975) frames the I -projection problem (1.4) in terms of moment equality constraints. He argues that two probability measures on abstract spaces are equal if and only if they have equal moments for *all* integrable moment functions. Since I focus on the setup specified by Assumption 1, I can simplify this characterization. The constraint set $\Pi(P_X, P_Y)$ has the following characterization:

$$\{P \in \mathcal{P}(\mathcal{X} \times \mathcal{Y}) : P_1(A) = P_X(A) \forall A \in \mathcal{B}(\mathcal{X}) \text{ and } P_2(A) = P_Y(A) \forall A \in \mathcal{B}(\mathcal{Y})\}, \quad (2.1)$$

where $\mathcal{P}(\mathcal{X} \times \mathcal{Y})$ denotes the set of probability measures defined on the measurable space $(\mathcal{X} \times \mathcal{Y}, \mathcal{B}(\mathcal{X} \times \mathcal{Y}))$, and $\mathcal{B}(\mathcal{X} \times \mathcal{Y})$ is the Borel sigma-algebra of $\mathcal{X} \times \mathcal{Y}$. Under Assumption 1, the collection of sets $\mathcal{B}_X = \{A \in \mathcal{B}(\mathcal{X}) : A = \times_{i=1}^{d_x} (-\infty, x_i], x \in \mathcal{X}\}$ and $\mathcal{B}_Y = \{A \in \mathcal{B}(\mathcal{Y}) : A = \times_{i=1}^{d_y} (-\infty, y_i], y \in \mathcal{Y}\}$ generate the sigma-algebras $\mathcal{B}(\mathcal{X})$ and $\mathcal{B}(\mathcal{Y})$, respectively. Consequently, there is no loss of information in reformulating the constraints in (2.1) by replacing $\mathcal{B}(\mathcal{X})$ and $\mathcal{B}(\mathcal{Y})$ with \mathcal{B}_X and \mathcal{B}_Y , respectively.

Now this reformulation of the constraint set is connected to moment inequalities using the cumulative distribution functions of P_X and P_Y ,

$$F_X(x) := P_X\left(\times_{i=1}^{d_x} (-\infty, x_i]\right) \quad \text{and} \quad F_Y(y) := P_Y\left(\times_{i=1}^{d_y} (-\infty, y_i]\right),$$

which are defined on \mathcal{X} and \mathcal{Y} , respectively. To clarify, I can express the I -projection problem (1.4) as the following infinite-dimensional minimization problem,

$$\begin{aligned} \text{minimize} \quad m(p) &:= \begin{cases} \int_{\Omega} p \log(p) dR_{\gamma} & \text{if } p \geq 0, \\ \infty & \text{elsewhere,} \end{cases} \quad \int_{\Omega} p dR_{\gamma} = 1 \\ \text{subject to} \quad & \int_{\Omega} g_{x'} p dR_{\gamma} \leq 0 \text{ and } \int_{\Omega} -g_{x'} p dR_{\gamma} \leq 0 \quad \forall x' \in \mathcal{X}, \\ & \int_{\Omega} g_{y'} p dR_{\gamma} \leq 0 \text{ and } \int_{\Omega} -g_{y'} p dR_{\gamma} \leq 0 \quad \forall y' \in \mathcal{Y}, \\ & \text{and} \quad p \in L_1(R_{\gamma}), \end{aligned} \quad (2.2)$$

where $\Omega := \mathcal{X} \times \mathcal{Y}$, the moment functions are given by

$$g_{x'}(x, y) = F_X(x') - 1[x \preceq x'] \quad \text{and} \quad g_{y'}(x, y) = F_Y(y') - 1[y \preceq y'] , \quad (2.3)$$

with the notation " \preceq " means $x_i \leq x'_i$ and $y_j \leq y'_j$ for all $i = 1, \dots, d_x$ and $j = 1, \dots, d_y$, respectively, and

$$L_1(R_\gamma) := \left\{ h : \Omega \rightarrow \mathbb{R} : h \text{ is measurable } \mathcal{B}(\mathcal{X} \times \mathcal{Y})/\mathcal{B}(\mathbb{R}) \text{ and } \int_{\Omega} |h| dR_\gamma < \infty \right\} .$$

Let $\mathcal{V} := \{g_h, -g_h : h \in \mathcal{X} \cup \mathcal{Y}\}$. By Lemmas B.3 and B.4, this subset of $L_1(R_\gamma)$ is precompact and closed in the norm topology. These properties of \mathcal{V} are central to this section's results: existence and exponential family representation of the dual optimizers, and approximation. Towards that end, I formulate the I -projection's constraint set equivalently as

$$\mathcal{M} = \left\{ p \in L_1(R_\gamma) : m(p) < \infty, \int_{\Omega} vp dR_\gamma \leq 0 \forall v \in \mathcal{V} \right\} . \quad (2.4)$$

The positive conjugate cone of \mathcal{M} , using $L_0(R_\gamma)$ as the dual space, is thus defined as

$$\mathcal{M}^\oplus = \left\{ z \in L_0(R_\gamma) : \int_{\Omega} z p dR_\gamma \geq 0 \forall p \in \mathcal{M} \right\} .$$

In light of the form of \mathcal{M}^\oplus , I consider the dual optimization problem on the following domain

$$\mathcal{D} := \left\{ z \in \mathcal{M}^\oplus : z \in \overline{\text{co}}(\mathcal{V}) \cdot \alpha, \alpha \geq 0 \right\} , \quad (2.5)$$

where $\overline{\text{co}}(\mathcal{V})$ is the closed convex hull of \mathcal{V} in the $L_1(R_\gamma)$ -norm. In particular, the dual optimization problem is given by

$$\inf \left\{ \int_{\Omega} e^z dR_\gamma; z \in \mathcal{D} \right\} . \quad (2.6)$$

I have the following result on the existence and uniqueness of the solution to the dual problem (2.6), and its exponential representation.

Proposition 2.1. *Let the constraint set \mathcal{M} be given by (2.4), and suppose that Assumption 1 holds.*

1. $\arg \inf \left\{ \int_{\Omega} e^z dR_\gamma : z \in \mathcal{D} \right\} \neq \emptyset$ and the R_γ -density of P_γ has the following representation

$$p_\gamma = \frac{e^{z_{0,\gamma}}}{\int_{\Omega} e^{z_{0,\gamma}} dR_\gamma} \quad \text{where} \quad z_{0,\gamma} \equiv \arg \inf \left\{ \int_{\Omega} e^z dR_\gamma : z \in \mathcal{D} \right\} . \quad (2.7)$$

solves the I -projection problem (2.2).

2. $z_{0,\gamma} \in \overline{\text{span}_+(\mathcal{V})}$, where $\overline{\text{span}_+(\mathcal{V})}$ is the $L_1(R_\gamma)$ -norm closure of the positive linear span of \mathcal{V} .

Proof. See Appendix A.1. ■

Furthermore, duality holds:

$$m(p_\gamma) = -\log \left(\int_{\Omega} e^{z_0, \gamma} dR_\gamma \right). \quad (2.8)$$

I will now discuss the sieve approximation of the dual problem (2.6). Observe that the domain is infinite-dimensional because there are infinitely many inequality restrictions. Therefore, I approximate the dual problem and implement it numerically using a simulation-based procedure. The sieve approximation scheme is the one put forward by Tabri (2025). It approximates the dual problem using a sequence of finite-dimensional convex stochastic programs, and I use the SAA method in Section 3 to approximate these stochastic programs.

I begin by reparametrizing the dual problem (2.6) using the Gelfand–Pettis (G–F) integral. Because I work in $L_1(R_\gamma)$, I provide a self-contained definition and the minimal existence statement needed for this paper. This integral applies to *any* subset $J \subset L_1(R_\gamma)$ that is compact in the norm topology. Suppose ξ is a positive Radon measure on the measure space $(J, \mathcal{B}(J))$, where $\mathcal{B}(J)$ is the Borel sigma-algebra of J , and denote the topological dual of $L_1(R_\gamma)$ by $L_1(R_\gamma)^*$.

Definition 1 (Gelfand–Pettis integral in $L_1(R_\gamma)$). *Suppose that for every continuous linear functional $\Lambda \in L_1(R_\gamma)^*$, the scalar map $j \mapsto \Lambda(j)$ is ξ -integrable. If there exists $h \in L_1(R_\gamma)$ such that*

$$\Lambda(h) = \int_J \Lambda(j) d\xi(j) \quad \text{for all } \Lambda \in L_1(R_\gamma)^*,$$

then h is called the Gelfand–Pettis integral of the identity map on J with respect to ξ , and we write

$$\int_J j d\xi(j) := h.$$

As $L_1(R_\gamma)^*$ separates points on $L_1(R_\gamma)$, there is at most one such h that satisfies Definition 1. Thus, there is no uniqueness problem. The existence of h follows from an application of Theorem 3.27 of Rudin (1991), because (a) J is compact in the $L_1(R_\gamma)$ -norm, and (b) $L_1(R_\gamma)$ with its norm topology is a Fréchet space. As \mathcal{V} is compact in the norm topology of $L_1(R_\gamma)$, I can apply this definition with the choice $J = \mathcal{V}$.

In consequence, the dual problem (2.6) can be reparametrized in terms of Radon measures whose supported on \mathcal{V} . These measures are modeled as elements of the space $C(\mathcal{V})^*$ —the topological dual of the Banach space of continuous functions on \mathcal{V} , $C(\mathcal{V})$. The reparametrization arises from the following representation of elements in \mathcal{D} :

$$z \in \mathcal{D} \iff \exists \mu \in \mathcal{P} \text{ and } \alpha \geq 0 \text{ such that } z = \alpha \int_{\mathcal{V}} v d\mu(v). \quad (2.9)$$

where $\mathcal{P} \subset C(\mathcal{V})^*$ the set of Radon probability measures on \mathcal{V} , and $\int_{\mathcal{V}} v d\mu(v)$ the G–F integral. Now define $\Xi \subset C(\mathcal{V})^*$ as the set of all positive Radon measures on \mathcal{V} , and consider the following set

$$\Upsilon := \{\xi \in \Xi : \xi = \alpha \cdot \mu, \alpha \geq 0 \text{ and } \mu \in \mathcal{P}\}. \quad (2.10)$$

The dual problem (2.6) and its solution set can now be reparametrized as

$$\vartheta^* := \inf \left\{ \int_{\Omega} e^{\int_{\mathcal{V}} v \, d\xi(v)} \, dR_{\gamma} : \xi \in \Upsilon \right\}, \quad (2.11)$$

and $\mathcal{S}^* := \arg \inf \left\{ \int_{\Omega} e^{\int_{\mathcal{V}} v \, d\xi(v)} \, dR_{\gamma} : \xi \in \Upsilon \right\}$, respectively. I have the following result on the characterization of \mathcal{S}^* .

Corollary 2.1. *Suppose the conditions of Proposition 2.1 hold. Then it follows that*

$$\mathcal{S}^* = \left\{ \xi \in \Upsilon : z_{0,\gamma} = \int_{\mathcal{V}} v \, d\xi(v) \right\} \neq \emptyset.$$

Proof. See Appendix A.2. ■

The G-F integral representation of the solution $z_{0,\gamma}$ in (2.6) is $z_{0,\gamma} = \alpha_{0,\gamma} \int_{\mathcal{V}} v \, d\mu_{0,\gamma}(v)$. Since $z_{0,\gamma}$ is unique (up to equivalence class), it follows that $\alpha_{0,\gamma}$ is also unique, but that $\mu_{0,\gamma}$ may not be uniquely defined. This representation of $z_{0,\gamma}$ shows $\alpha_{0,\gamma}$ is also the total variation norm of the Radon measure $\xi_{0,\gamma} := \alpha_{0,\gamma} \cdot \mu_{0,\gamma}$.

Using Assumption 1, I can also obtain a representation of $z_{0,\gamma}$ in terms of Schrödinger potentials (Nutz and Wiesel, 2022), enabling me to deduce an upper bound on $\alpha_{0,\gamma}/\gamma$ via an application of Lemma 4.9 in Nutz (2022). The next result presents this bound.

Proposition 2.2. *Suppose Assumption 1 holds, and let $\|c\|_{\infty} = \sup_{(x,y) \in \mathcal{X} \times \mathcal{Y}} c(x, y)$. Then, $\alpha_{0,\gamma}/\gamma \leq \kappa \|c\|_{\infty} < \infty$, holds, where $\kappa := \frac{1}{\max\{1-F_X(\inf \mathcal{X}), 1-F_Y(\inf \mathcal{Y})\}}$.*

Proof. See Appendix A.3. ■

The sieve I consider is based on the approximation scheme put forward by Tabri (2025). His scheme is indexed by a sequence $\{\epsilon_{\ell}\}_{\ell \geq 1} \subset \mathbb{R}_{++}$ with $\epsilon_{\ell} \downarrow 0$ as $\ell \rightarrow \infty$. For each $\ell \in \mathbb{Z}_+$, let $U_{\ell} := \{z \in L_1(R_{\gamma}) : \|z\|_{L_1(R_{\gamma})} \leq \epsilon_{\ell}\}$. Then by Lemma A.2 of Tabri (2025), there corresponds a finite partition $\{E_{i,\ell}\}_{i=1}^{n_{\ell}}$ of \mathcal{V} with the property

$$\int_{\mathcal{V}} v \, d\mu(v) - \sum_{i=1}^{n_{\ell}} \mu(E_{i,\ell}) v_i \in U_{\ell} \quad \forall v_i \in E_{i,\ell}, \quad i = 1 \dots, n_{\ell}. \quad (2.12)$$

Remarkably, the only structure on the partitions that is required for this approximation to hold is that for each i : $v - v' \in U_{\ell}$ for all $v, v' \in E_{i,\ell}$. Consequently, the partitions depend only on ϵ_{ℓ} , R_{γ} and \mathcal{V} , and not μ . Of course, to satisfy the accuracy (2.12) the dimension of the discretization, n_{ℓ} , will diverge to ∞ as $\ell \rightarrow \infty$.

There are many partitions of \mathcal{V} that can be used in practice. By the proof of Lemma B.3, the set \mathcal{V} is VC subgraph; hence, there exists a finite partition of \mathcal{V} satisfying the accuracy (2.12) that does *not* depend on R_{γ} . Partitions of this sort are advantageous in practice since n_{ℓ} would then be independent of the regularizer γ . In particular, standard entropy bounds for uniformly bounded VC-subgraph classes (e.g., Theorem 2.6.7 in van der Vaart and Wellner, 1996) imply that, for each approximation level ϵ_{ℓ} , one can choose a partition

such that n_ℓ and the accuracy (2.12) satisfy $n_\ell \leq K \epsilon_\ell^{-\delta}$, for constants $K, \delta > 0$ depending only on the VC characteristics of \mathcal{V} and the uniform envelope (here, it is the constant function equal to 1), but not on the reference measure R_γ . Consequently, n_ℓ can always be taken to grow at most polynomially in ϵ_ℓ^{-1} , uniformly over the choice of the regularization parameter γ . Appendix D presents an example of such a partition.

The sieve approximation for the dual problem proceeds as follows. Fix a level $\ell \in \mathbb{Z}_+$ and choose an approximation tolerance $\epsilon_\ell > 0$. Let $n_\ell \in \mathbb{Z}_+$ and $\{E_{i,\ell}\}_{i=1}^{n_\ell}$ be a partition of \mathcal{V} satisfying the accuracy requirement (2.12). For each $i = 1, \dots, n_\ell$, select a representative $v_i \in E_{i,\ell}$ and let

$$G_\ell(\omega; \alpha, \mu) := \exp \left\{ \alpha \sum_{i=1}^{n_\ell} \mu_i v_i(\omega) \right\}, \quad \omega \in \Omega, (\alpha, \mu) \in \mathcal{C}_\ell, \quad (2.13)$$

where $\mathcal{C}_\ell := \left\{ (\alpha, \mu) \in \mathbb{R}_+^{n_\ell+1} : \sum_{i=1}^{n_\ell} \mu_i = 1 \right\}$. The finite-dimensional convex program associated with level ℓ is

$$\vartheta_\ell^* := \inf_{(\alpha, \mu) \in \mathcal{C}_\ell} \int_{\Omega} G_\ell(\omega; \alpha, \mu) dR_\gamma(\omega), \quad (2.14)$$

$$\mathcal{S}_\ell^* := \arg \inf_{(\alpha, \mu) \in \mathcal{C}_\ell} \int_{\Omega} G_\ell(\omega; \alpha, \mu) dR_\gamma(\omega). \quad (2.15)$$

I can also apply Theorem 3 of Tabri (2025) to establish convergence of the approximation scheme to the original problem (2.11) with the sequence of approximate solutions converging to a solution of the original problem, as $\ell \rightarrow \infty$. The next result formalizes this point.

Proposition 2.3. *Suppose Assumption 1 holds. Let $\bar{c} = \int_{\Omega} c(\omega) d(P_X \otimes P_Y)$ and let $\{\epsilon_\ell, U_\ell\}_{\ell \geq 1}$ be described as above. For each ℓ , let $\{E_{i,\ell}\}_{i=1}^{n_\ell}$ be a partition of \mathcal{V} such that, for each i , $v - v' \in U_\ell$ holds for all $v, v' \in E_{i,\ell}$. Then, for each ℓ , $\mathcal{S}_\ell^* \neq \emptyset$ for any $v_i \in E_{i,\ell}$ where $i = 1, \dots, n_\ell$. Furthermore, for each ℓ and $v_i \in E_{i,\ell}$ with $i = 1, \dots, n_\ell$, define the corresponding Radon measure $\xi_\ell = \alpha_{n_\ell} \sum_{i=1}^{n_\ell} \mu_{i,n_\ell} \delta_{v_i}$, where $(\alpha_{n_\ell}, \mu_{1,n_\ell}, \dots, \mu_{n_\ell,n_\ell}) \in \mathcal{S}_\ell^*$ and δ_{v_i} is the Dirac delta function at v_i for each i . Then the following statements hold.*

1. $\lim_{\ell \rightarrow \infty} \vartheta_\ell^* = \vartheta^*$,
2. *Convergence rate of $\{\log(\vartheta_\ell^*)/\gamma\}_{\ell \geq 1}$: $\gamma^{-1} |\log \vartheta_\ell^* - \log \vartheta^*| \leq \epsilon_\ell \kappa \|c\|_\infty e^{\gamma(\kappa \|c\|_\infty + \bar{c}) + \log a_\gamma}$ for all $\ell \in \mathbb{Z}_+$.*
3. *Every accumulation point of $\{\xi_\ell\}_{\ell \geq 1}$, in the weak-star topology of $C(\mathcal{V})^*$, is an element of \mathcal{S}^* .*

Proof. See Appendix A.4. ■

This section achieved two complementary objectives. First, it reformulated the entropically regularized optimal transport problem as an I -projection under an infinite collection of moment inequality restrictions and characterized its solution through a Fenchel dual representation involving a G-F integral. Second, it introduced a deterministic sieve approximation of this dual problem, replacing the infinite-dimensional

optimization with a sequence of finite-dimensional convex programs whose solutions converge to those of the original dual problem as the sieve resolution increases. These approximating programs provide a numerically tractable representation of the dual problem, but they still involve expectations with respect to the reference measure and are therefore not directly observable.

The following section builds on this deterministic approximation by introducing a sieve M-estimation framework. Specifically, the finite-dimensional convex programs constructed above are estimated using the SAA method, yielding estimators of both the dual optimal value and the corresponding dual variables. This two-step structure—deterministic sieve approximation followed by stochastic estimation—forms the basis for the consistency, rate, and inference results established in the remainder of the paper.

3 Sieve M-Estimation

As the approximating problem (2.14) is a stochastic program, I describe a procedure for computing ϑ_ℓ^* and elements of \mathcal{S}_ℓ^* using the SAA method. I also show the procedure converges to the Fenchel dual problem (2.11). Firstly, replace the constraint set \mathcal{C}_ℓ with $\bar{\mathcal{C}}_\ell := \{(\alpha, \mu) \in \mathcal{C}_\ell : \alpha \leq \gamma\kappa\|c\|_\infty\}$, where the upper bound is due to Proposition 2.2. Consequently, for each ℓ , the collection of functions

$$\mathcal{G}_\ell := \{G_\ell(\cdot, \alpha, \mu) : (\alpha, \mu) \in \bar{\mathcal{C}}_\ell\} \quad (3.1)$$

is a uniformly bounded class of functions, with common bound $e^{\gamma\kappa\|c\|_\infty}$. This property of \mathcal{G}_ℓ is useful for the ensuing analysis, as the level of the discretization's accuracy, ϵ_ℓ , must be coupled with the sample size through the complexity of this class of functions.

Next, replace the finite program (2.14) and its solution set \mathcal{S}_ℓ^* with

$$\begin{aligned} \vartheta_\ell &:= \inf \left\{ \int_{\Omega} G_\ell(\omega; \alpha, \mu) dR_\gamma(\omega) : (\alpha, \mu) \in \bar{\mathcal{C}}_\ell \right\} \quad \text{and} \\ S_\ell &:= \arg \min \left\{ \int_{\Omega} G_\ell(\omega; \alpha, \mu) dR_\gamma(\omega) : (\alpha, \mu) \in \bar{\mathcal{C}}_\ell \right\}, \end{aligned} \quad (3.2)$$

respectively. Now solving this optimization problem using the SAA method entails solving the sample-analogue of (3.2) with a simulated random sample from R_γ . Let $\omega^{N_\ell} := \{\omega_j, i \leq N_\ell\}$ be a random sample of size N_ℓ from R_γ . The SAA method solves

$$\hat{\vartheta}_\ell(\omega^{N_\ell}) := \inf \left\{ \frac{1}{N_\ell} \sum_{j=1}^{N_\ell} G_\ell(\omega_j, \alpha, \mu) : (\alpha, \mu) \in \bar{\mathcal{C}}_\ell \right\} \quad (3.3)$$

and approximates S_ℓ with $\hat{S}_\ell(\omega^{N_\ell}) := \arg \min \left\{ \frac{1}{N_\ell} \sum_{j=1}^{N_\ell} G_\ell(\omega_j, \alpha, \mu) : (\alpha, \mu) \in \bar{\mathcal{C}}_\ell \right\}$.

Remark 3.1 (Measurability). For each $\ell \in \mathbb{Z}_+$, the SAA objective function in (3.2) is defined on a common probability space $(\times_{i=1}^{N_\ell} \Omega, \times_{i=1}^{N_\ell} \mathcal{B}(\Omega), R_\gamma^{\otimes N_\ell})$, where $\mathcal{B}(\Omega)$ is the Borel sigma-algebra of Ω , and $R_\gamma^{\otimes N_\ell}$ is the N_ℓ -fold product of the probability measure R_γ . As the function $G_\ell(\omega; \alpha, \mu)$, defined in (2.13), is a

Carathéodory function, i.e., continuous in (α, μ) and measurable in ω , the SAA objective function is also such a function. In consequence, $\hat{\vartheta}_\ell(\omega^N)$ and $\hat{S}_\ell(\omega^N)$ are measurable. Furthermore, a particular optimal solution (α_ℓ, μ_ℓ) of the SAA problem is a measurable selection $(\alpha_\ell(\omega^{N_\ell}), \mu_\ell(\omega^{N_\ell})) \in \hat{S}_\ell(\omega^{N_\ell})$, and existence of such measurable selection is ensured by the Measurable Selection Theory (e.g., Theorem 7.34, Shapiro et al., 2009). This takes care of measurability questions.

In my setup, the sequence of SAA programs (3.3) comprise a nonparametric sieve estimation problem. A sufficient condition for its consistency, as the sample size N_ℓ increases with ℓ , for fixed γ , is that

$$\lim_{\ell \rightarrow \infty} N_\ell^{-1} \log n_\ell = 0. \quad (3.4)$$

This condition implies the Rademacher complexity, $\mathcal{R}_{N_\ell}(\mathcal{G}_\ell)$, described in Lemma C.1 and Corollary C.1, satisfies $\mathcal{R}_{N_\ell}(\mathcal{G}_\ell) = o(1)$ as $\ell \rightarrow \infty$. Consequently, I can establish that a uniform strong law of large numbers holds for the sequence of function classes $\{\mathcal{G}_\ell\}_{\ell \geq 1}$ —see Proposition C.1 in Appendix C. This large-sample result is key in the proof of my main result.

Theorem 1. *Suppose that Assumption 1 holds, let \mathcal{G}_ℓ be as given in (3.1) for each $\ell \in \mathbb{Z}_+$, and suppose that the limit (3.4) holds. For each ℓ , let ω^{N_ℓ} be a random sample from R_γ , and $\hat{\xi}_\ell(\omega^{N_\ell}) = \hat{\alpha}_\ell(\omega^{N_\ell}) \sum_{i=1}^{n_\ell} \hat{\mu}_{i,\ell}(\omega^{N_\ell}) \delta_{v_i}$ where $(\hat{\alpha}_\ell(\omega^{N_\ell}), \hat{\mu}_\ell(\omega^{N_\ell})) \in \hat{S}_\ell(\omega^{N_\ell})$. Then the following statements hold.*

1. *With probability 1,*

- (i) $\lim_{\ell \rightarrow \infty} \hat{\vartheta}_\ell = \vartheta^*$.
- (ii) *Every accumulation point of $\{\hat{\xi}_\ell\}_{\ell \geq 1}$, in the weak-star topology of $C(\bar{\mathcal{V}})^*$, is an element of \mathcal{S}^* .*

2. *Sample complexity: for each ℓ ,*

$$\gamma^{-1} E_{R_\gamma^{\otimes N_\ell}} \left[\left| \log \hat{\vartheta}_\ell - \log \vartheta^* \right| \right] \leq \max \left\{ \sqrt{\frac{2 \log n_\ell}{N_\ell}}, \epsilon_\ell \right\} 2\kappa \|c\|_\infty e^{\gamma 2\kappa \|c\|_\infty}.$$

3. *As $\ell \rightarrow \infty$,*

$$\begin{aligned} \vartheta^* &\geq \hat{\vartheta}_\ell - N_\ell^{-1/2} \sup_{g \in \mathcal{G}_\ell} B_\ell^l(g) - \epsilon_\ell \gamma \kappa \|c\|_\infty e^{\gamma \kappa \|c\|_\infty} + o_{R_\gamma}(N_\ell^{-1/2}), \\ \vartheta^* &\leq \hat{\vartheta}_\ell + N_\ell^{-1/2} \sup_{g \in \mathcal{G}_\ell^-} B_\ell^u(g) + o_{R_\gamma}(N_\ell^{-1/2}), \end{aligned}$$

where B_ℓ^l and B_ℓ^u are centered Gaussian processes indexed by \mathcal{G}_ℓ and $\mathcal{G}_\ell^- = -\mathcal{G}_\ell$ with common covariance kernel $\int_\Omega g(\omega)g'(\omega) dR_\gamma(\omega)$.

Proof. See Appendix A.5. ■

An immediate consequence of the result in Part 1(i) of Theorem 1 is on estimation of the EOT value (1.2). In the notation of Section 2, this value is given by $\gamma^{-1} (\log a_\gamma + m(p_\gamma))$, and with probability 1,

$$\lim_{\ell \rightarrow \infty} \gamma^{-1} \left(\log a_\gamma - \log \hat{\vartheta}_\ell \right) = \gamma^{-1} (\log a_\gamma - \log \vartheta^*) = \gamma^{-1} (\log a_\gamma + m(p_\gamma)),$$

by an application of the Continuous Mapping Theorem and optimal value duality (2.8).

Sections 2–3 together yield a two-stage construction: a deterministic sieve approximation of the Fenchel dual of the I -projection problem (2.6), followed by a stochastic approximation of the resulting finite-dimensional convex programs via the SAA method. Theorem 1 summarizes the statistical consequences of this construction, establishing (i) almost sure consistency for the estimators of the dual optimal value and the associated dual solutions, (ii) finite-sample mean-error bounds for the estimator of the primal optimal value, and (iii) stochastic upper and lower error bounds for the estimation error of the dual optimal value. The remainder of the paper discusses how these guarantees compare with the existing statistical optimal transport literature.

4 Discussion

This section contrasts the present moment-inequality-based approach with the empirical Sinkhorn divergence and with continuity-driven dual parametrizations, and it explains how the population-level constraint structure underlying the sieve M-estimator leads to different modes of convergence and different inferential tools. It also outlines extensions to entropic optimal transport problems with additional moment restrictions.

4.1 Other Approaches

A large body of work on approximating entropically regularized optimal transport focuses on computational formulations of the Fenchel–Rockafellar dual, often implemented via the Sinkhorn algorithm. In the general setting, the Fenchel–Rockafellar dual can be written as in Nutz and Wiesel (2022),

$$\sup_{f \in L_1(P_X), g \in L_1(P_Y)} \left(\int_{\mathcal{X}} f dP_X + \int_{\mathcal{Y}} g dP_Y - \frac{1}{\gamma} \left(\int_{\mathcal{X} \times \mathcal{Y}} e^{\gamma(f(x) + g(y) - c(x, y))} d(P_X \otimes P_Y) - 1 \right) \right),$$

where the functions f and g are called dual or Schrödinger potentials in the literature. Most practical algorithms do not use this L_1 -based convex-analytic dual. Instead, they require the potentials to be continuous and use parametrizations ensuring pointwise evaluation and stochastic-gradient updates are well defined. This category includes the RKHS-based sieve of Genevay et al. (2016) and the neural-network parametrizations of Seguy et al. (2018) and Arjovsky et al. (2017). The RKHS approach assumes the true potentials belong to a fixed RKHS—a strong modeling assumption that fails for many continuous functions on compact sets, as demonstrated by Theorem 1.1 of Steinwart (2024). Neural-network parametrizations avoid this restriction but still require continuity of the potentials as a structural assumption motivated by computational

considerations.

The dual framework developed in this paper fundamentally departs from continuity-based approaches. Here, the dual variables are finite Radon measures that act on a VC class of moment functions, placing them naturally in the dual space $C(\mathcal{V})^*$ with the weak-star topology. Unlike previous methods, this framework does not require continuity of the Schrödinger potentials. This choice aligns with the analytical structure of the entropic dual: regularization removes the pointwise constraint $f(x) + g(y) \leq c(x, y)$ found in unregularized optimal transport, so integrability—not continuity—is the minimal regularity ensured by the Schrödinger system. As a result, the sieve M-estimator developed here preserves the dual problem’s population-level structure without imposing unnecessary smoothness conditions.

A comparison with the empirical Sinkhorn divergence further clarifies the fundamental differences in how these estimators use data and the EOT constraint set. The sieve M-estimator enforces the EOT constraints at the population level by approximating the moment inequalities that define the Schrödinger system, guided by the geometry of the population-level feasible set of couplings. In contrast, the empirical Sinkhorn divergence replaces the true marginals P_X and P_Y with their empirical versions, solving a discrete EOT problem that enforces only empirical marginal constraints and derives its geometry entirely from the sampled data. As a result, the sieve M-estimator preserves the analytical structure of the population constraints and uses data solely to approximate expectations in the dual objective. In contrast, the empirical Sinkhorn divergence treats the empirical distributions as if they were the population distributions. This distinction has practical implications: the two estimators respond differently to sampling variability and to the geometry of the underlying distributions, and only the sieve estimator maintains the population constraint set inherent in the EOT problem.

These structural differences shape their statistical guarantees. Notably, only the empirical Sinkhorn divergence currently has established convergence rates or central limit theorems—and even these results hold only in special cases involving smooth or quadratic costs. In contrast, Theorem 1 shows that the sieve M-estimator achieves consistency under much weaker conditions and provides finite-sample mean error bounds and stochastic approximations valid for all continuous cost functions on compact supports. I discuss these results in detail below.

4.2 Part 1 of Theorem 1

The first part of Theorem 1 establishes the almost sure convergence of both the sieve optimal value estimator and the corresponding dual estimators, for any continuous cost function on compact supports. This significantly broadens the scope of statistical optimal transport beyond the traditional focus on smooth or quadratic costs and empirical Sinkhorn divergence. The main distinction between my dual consistency result and previous empirical Sinkhorn analyses lies in the mode and topology of convergence: my dual estimators converge almost surely in the weak-star topology of $C(\mathcal{V})^*$, whereas empirical Sinkhorn results typically establish convergence in mean with respect to the $L_2(\hat{P}_X \otimes \hat{P}_Y)$ norm, where \hat{P}_X and \hat{P}_Y are the empirical marginals (see, e.g., Lemma 4.8 in Section 4.2 of Chewi et al., 2025). The question of deriving convergence rates for my dual variable estimator remains open and is left for future work, though I briefly outline potential approaches below.

This difference in convergence mode has important practical implications. Weak-star almost sure convergence offers a strong qualitative guarantee: the estimated dual variables converge in the weak-star topology for almost every sample path, not just on average. This property is especially valuable for downstream tasks—such as sensitivity analysis, specification testing, or constructing confidence bounds for transport costs—where a stable dual representation is crucial. In contrast, mean L_2 convergence for empirical Sinkhorn duals depends on the empirical marginals and reflects approximation quality averaged over the sampled supports. While both types of convergence are informative, weak-star almost sure convergence uniquely captures how my dual variables utilize population-level information, rather than only the empirical support. This makes it particularly well suited for applications that rely on the dual structure inherent to the EOT problem.

A natural starting point for studying rates of convergence of my dual variable estimator is to endow the parameter space with a compatible metric. Recall that the dual variables lie in the set $\Upsilon_0 = \{\xi \in \Upsilon : \|\xi\|_{TV} \leq \gamma\kappa\|c\|_\infty\}$, where $\|\cdot\|_{TV}$ is the total variation norm on $C(\mathcal{V})^*$. As it is a norm-bounded subset of $C(\mathcal{V})^*$, by the Banach–Alaoglu theorem it must be weak-star compact, and the weak-star topology on it is metrizable. A convenient metric arises from the parametrization

$$\Phi : (0, \gamma\kappa\|c\|_\infty] \times \mathcal{P} \rightarrow \Upsilon_0, \quad \Phi(\alpha, \mu) = \alpha\mu,$$

where \mathcal{P} denotes the set of Radon probability measures on \mathcal{V} . Endow $(0, \gamma\kappa\|c\|_\infty]$ with the usual metric and \mathcal{P} with any metric that metrizes weak convergence (e.g., the bounded–Lipschitz or Prokhorov metric). Then Φ is a homeomorphism between the product space and $(\Upsilon_0, \text{weak-star})$, so the weak-star topology on Υ_0 may be described by a product metric. This makes it possible, at least in principle, to study rates of convergence of the dual estimators by analyzing the rates at which the scalar components $\hat{\alpha}_\ell$ and the probability-measure components $\hat{\mu}_\ell$ approach their population counterparts. A full development of such rates would require delicate stability properties of the dual problem and is therefore left for future research, but this parametrization clarifies how a metric-based analysis could proceed.

4.3 Part 2 of Theorem 1

4.3.1 Comparison to Theorem 3 in Genevay et al. (2019)

The second result of Theorem 1 provides a finite-sample rate on the approximation error incurred by the sieve M-estimator when estimating the optimal value of the EOT problem, namely

$$\gamma^{-1}(\log a_\gamma + \log \vartheta^*),$$

under Assumption 1. It is instructive to compare this bound with the in-mean convergence rate of the empirical Sinkhorn divergence established in Theorem 3 of Genevay et al. (2019). Their result applies when the cost function is infinitely differentiable and L -Lipschitz and when \mathcal{X} and \mathcal{Y} are bounded subsets of \mathbb{R}^d . In our notation, their discrepancy is bounded (up to constants) by

$$\frac{e^{\gamma(2L|\mathcal{X}| + \|c\|_\infty)}}{\sqrt{N}} \left(1 + \gamma^{\lfloor d/2 \rfloor}\right),$$

where $|\mathcal{X}|$ denotes the diameter of \mathcal{X} , and the constants depend only on $|\mathcal{X}|$, $|\mathcal{Y}|$, d , and $\|c^{(k)}\|_\infty$ for $k = 0, \dots, \lfloor d/2 \rfloor$.

There are several points of comparison and contrast.

(1) Scope of applicability. My rate applies to *all continuous* cost functions on compact supports, whereas theirs requires the substantially narrower class of infinitely differentiable, globally Lipschitz costs. Both bounds exhibit exponential dependence on γ . In the quadratic case $c(x, y) = \|x - y\|^2/2$, Mena and Niles-Weed (2019) show that this exponential dependence can be removed for the empirical Sinkhorn divergence. It is plausible that an analogous refinement may be possible for my sieve M-estimator, and I leave this investigation for future research.

(2) Dependence on dimension and regularization. In Genevay et al. (2019), the factor $(1 + \gamma^{\lfloor d/2 \rfloor})$ introduces a pronounced curse of dimensionality, especially for large γ . In contrast, my rate depends on dimension and regularization only through $\log n_\ell$, where n_ℓ is the size of the sieve. When the partition of \mathcal{V} is chosen independently of R_γ , this removes the $\gamma^{d/2}$ effect entirely. Even when the sieve is adapted to geometric or distributional features of P_X , P_Y , or R_γ , the influence of such features is substantially attenuated by the logarithm.

(3) Nature of the convergence rate. The rate obtained for the sieve estimator is nonparametric—it depends on $\max \left\{ \epsilon_\ell, \sqrt{2 \log(n_\ell) / N_\ell} \right\}$, and therefore may be slower than $N^{-1/2}$. Nevertheless, the numerical experiment in Appendix E shows that, for a moderate sample size and a well-chosen sieve, the proposed estimator can outperform the empirical Sinkhorn divergence in practice, despite its theoretically slower rate. This highlights the practical advantage of adapting the approximation to the structure of the I -projection rather than to the sample-induced discrete geometry.

4.3.2 Bias–variance trade-off in the rate

The rate established in Part 2 of Theorem 1 makes explicit the bias–variance trade-off inherent in the nonparametric estimation of $\gamma^{-1} \log \vartheta^*$. As is typical in nonparametric settings, the stochastic component of the error depends on the size and complexity of the sieve function class \mathcal{G}_ℓ . The term $\sqrt{2 \log(n_\ell) / N_\ell}$ increases with n_ℓ and, by Lemma S3.1, quantifies this complexity through a Rademacher-type bound. Enlarging \mathcal{G}_ℓ introduces greater variability into $\gamma^{-1} \log \hat{\vartheta}_\ell$, thereby inflating the variance, whereas the deterministic approximation error ϵ_ℓ represents a bias term that decreases as \mathcal{G}_ℓ becomes richer. Thus, reducing bias inevitably increases variance, reflecting the classical bias–variance trade-off.

In standard nonparametric estimation, the sample size is fixed and the optimal convergence rate is obtained by choosing a sieve dimension that balances the bias and variance terms. By contrast, when the EOT problem (1.2) is viewed as a stochastic optimization problem, the present framework first specifies the deterministic approximation error ϵ_ℓ , which in turn determines the size of \mathcal{G}_ℓ , and then selects a sample size that satisfies the rate condition (3.4). The optimal balance is achieved by equating the stochastic and deterministic errors:

$$\sqrt{\frac{2 \log n_\ell}{N_\ell}} = \epsilon_\ell \iff N_\ell = \frac{2 \log n_\ell}{\epsilon_\ell^2}. \quad (4.1)$$

A sample size of this order automatically satisfies the entropy condition (3.4) and yields the corresponding optimal rate of convergence for the sieve M-estimator.

Beyond clarifying the bias–variance trade-off, this calculation provides practically useful guidance on sample size selection. To the best of my knowledge, the Statistical Optimal Transport literature offers no explicit prescriptions for determining the number of samples required to achieve a desired accuracy level—whether for empirical Sinkhorn, RKHS-based methods, or neural-dual parametrizations. The expression $N_\ell = 2\epsilon_\ell^{-2} \log n_\ell$ gives a concrete, interpretable sample size that links the target accuracy ϵ_ℓ to the complexity of the sieve. Such guidance is especially valuable in applications where practitioners must tune both the regularization parameter γ and the resolution of the approximation scheme, yet currently have no principled tool for determining how large a sample is needed for a given precision.

4.4 Part 3 of Theorem 1

4.4.1 Comparison with Empirical Sinkhorn Divergence

It is instructive to contrast the stochastic bounds in Part 3 of Theorem 1 with recent results on the sample complexity of the empirical Sinkhorn divergence, such as Theorem 3 in Rigollet and Stromme (2024). The latter provides finite-sample concentration inequalities that control the estimation error in probability at a fixed sample size, for EOT value (1.2). By contrast, the present result yields an asymptotic stochastic approximation for the sieve M-estimator of the optimal value of the Fenchel dual problem, ϑ^* . Here, the estimation error is bounded above and below by suprema of Gaussian processes indexed by the sieve class and scaled by $N_\ell^{-1/2}$, up to a negligible remainder in probability.

These two types of results address complementary inferential questions. Concentration inequalities quantify tail behavior of the empirical Sinkhorn divergence viewed as a discretized transport problem. The Gaussian-process bounds derived here describe the full asymptotic fluctuation envelope of a population-level stochastic optimization problem, and are naturally suited to inference procedures such as confidence intervals and projection-based uncertainty quantification.

4.4.2 Confidence Interval for EOT value

The third part of Theorem 1 provides matching upper and lower stochastic inequalities for the estimation error of $\hat{\vartheta}_\ell$. These bounds naturally lead to the construction of an asymptotically valid confidence interval for ϑ^* , say $[b_\ell, b'_\ell]$. Because the map $\vartheta \mapsto \gamma^{-1}(\log a_\gamma - \log \vartheta)$ is strictly decreasing on $(0, \infty)$, this interval can be projected into an asymptotically valid confidence interval for the regularized optimal transport value $\gamma^{-1}(\log a_\gamma - \log \vartheta^*)$ via

$$\left[\gamma^{-1}(\log a_\gamma - \log b'_\ell), \quad \gamma^{-1}(\log a_\gamma - \log b_\ell) \right].$$

Let $q_{\delta, \ell}^{(l)}$ and $q_{\delta, \ell}^{(u)}$ denote the δ -quantiles of $\sup_{g \in \mathcal{G}_\ell} B_\ell^l(g)$ and $\sup_{g \in \mathcal{G}_\ell^-} B_\ell^u(g)$, respectively. At the pop-

ulation level, Part 3 of Theorem 1 suggests the asymptotic $(1 - \delta)$ confidence interval

$$\left[\hat{\vartheta}_\ell - N_\ell^{-1/2} q_{1-\delta/2,\ell}^{(l)} - \epsilon_\ell \gamma \kappa \|c\|_\infty e^{\gamma \kappa \|c\|_\infty}, \quad \hat{\vartheta}_\ell + N_\ell^{-1/2} q_{1-\delta/2,\ell}^{(u)} \right]$$

for ϑ^* . Since $\mathcal{G}_\ell^- = -\mathcal{G}_\ell$, the Gaussian processes B_ℓ^l and B_ℓ^u share the same covariance kernel and therefore have the same distribution, albeit indexed by different classes.

The lower bound includes a deterministic bias term $\epsilon_\ell \gamma \kappa \|c\|_\infty e^{\gamma \kappa \|c\|_\infty}$, which may be large for moderate or large values of γ . For practical implementation, it is therefore natural to consider the symmetric interval

$$\left[\hat{\vartheta}_\ell - N_\ell^{-1/2} q_{1-\delta/2,\ell}^{(l)}, \quad \hat{\vartheta}_\ell + N_\ell^{-1/2} q_{1-\delta/2,\ell}^{(u)} \right], \quad (4.2)$$

which omits this bias term. This modification is asymptotically valid under a standard undersmoothing condition: if $\sqrt{N_\ell} \epsilon_\ell = o(1)$, in addition to the rate condition (3.4) for fixed γ , then the omitted bias is negligible at the $N_\ell^{-1/2}$ scale.

The optimal sample size choice in (4.1) balances stochastic and approximation errors but does not satisfy this undersmoothing condition. More generally, the requirement $\sqrt{N_\ell} \epsilon_\ell = o(1)$ depends on how the sieve dimension n_ℓ grows with the approximation error ϵ_ℓ , which in turn is determined by the chosen partition of the moment class \mathcal{V} . Loosely speaking, selecting N_ℓ slightly larger than the bias–variance balance eliminates the approximation bias asymptotically. While a full analysis of optimal tuning is beyond the scope of this paper, the explicit form of the stochastic bounds provides clear guidance on how sample size and sieve complexity interact.

Finally, the quantiles $q_{1-\delta/2,\ell}^{(l)}$ and $q_{1-\delta/2,\ell}^{(u)}$ are unknown in practice and must be estimated from the data. Any procedure yielding consistent estimators of these quantiles as $\ell \rightarrow \infty$ —such as Gaussian approximation or multiplier bootstrap methods based on the empirical covariance structure of \mathcal{G}_ℓ in the spirit of Chernozhukov et al. (2014a)—leads to asymptotically valid confidence intervals for ϑ^* .

4.5 An Extension: Entropic Optimal Transport with Martingale Constraints

Recent work, such as Tang et al. (2025), has begun exploring entropically regularized martingale optimal transport problems from a computational perspective. These studies identify the existence, structure, and dual representations for entropic martingale transport, but leave open the question of statistical estimation under sampling.

The sieve M-estimation framework in this paper naturally extends to entropic martingale optimal transport. By formulating martingale and supermartingale constraints as conditional moment equalities or inequalities, and then converting them into (generally infinite) collections of unconditional moment restrictions using instrument functions, as in Tabri (2025), the resulting class of moment functions stays uniformly bounded and VC under Assumption 1, allowing the sieve approximation and stochastic optimization strategies presented here to handle these constraints and provide a principled estimation route.

However, to establish full statistical guarantees for constrained transport problems, we still need to develop additional tools. In particular, researchers have yet to find sharp bounds on the dual optimizers under general conditional moment restrictions—bounds that are crucial for controlling the stochastic program do-

mains and for deriving uniform laws of large numbers or Gaussian approximations. Advancing these aspects remains an important area for future research.

More broadly, this extension demonstrates the flexibility of viewing entropically regularized optimal transport as a stochastic optimization problem under moment restrictions. By focusing on population-level constraint sets rather than sample-induced discretizations, my framework creates new opportunities for statistical inference in constrained transport problems—areas previously explored mainly through analytical or computational methods.

4.6 Further Applications and Open Directions

The above discussion shows that the proposed sieve M-estimation framework relies on the moment-inequality structure of entropically regularized optimal transport, rather than on the analytic properties of dual potentials. This focus opens the door to a wide range of applications where entropic optimal transport yields a well-defined I -projection, even if we lack detailed regularity or representation results for the corresponding Schrödinger potentials.

Recent literature illustrates the variety of constraints we can express as moment restrictions in entropic optimal transport. Beyond classical marginal constraints, martingale and supermartingale constraints play a central role in model-independent finance and stochastic control (see, e.g., Beiglböck et al., 2013; Beiglböck and Juillet, 2016; Nutz and Stebegg, 2018, for supermartingale optimal transport; and Tang et al., 2025; Nutz and Wiesel, 2024, for entropic and Schrödinger bridge extensions). Convex order and dominance constraints can also be formulated as infinite families of linear moment inequalities indexed by convex test functions, which arise in risk aggregation and no-arbitrage pricing (Strassen, 1965; Rüsendorf, 1985; Beiglböck and Penkner, 2013). Related moment-based approaches appear also in causal and adapted transport (Lassalle, 2018; Backhoff-Veraguas and Pammer, 2020) and in Schrödinger bridge problems with structural or path-dependent constraints (Léonard, 2012). In many of these cases, one can establish entropic minimizers and dual representations using the results of Tabri (2025), but not sharp regularity results for the dual variables—such as continuity, boundedness, or smoothness.

My approach in this paper departs from traditional potential-based computational and analytical methods. Instead of parametrizing dual variables as functions and imposing continuity or smoothness for pointwise optimization, I operate directly on the population-level constraint set through moment inequalities. I treat dual variables as Radon measures acting on classes of moment functions and estimate them by approximating expectations in the Fenchel dual formulation. This method remains well defined even when Schrödinger potentials lack simple functional representations.

From this perspective, the lack of sharp results on dual potentials does not prevent statistical analysis of the corresponding entropic transport values. Instead, we need sufficient structural properties—such as boundedness of dual optimizers and VC complexity of the moment class—to control the stochastic approximation. Establishing these features for specific constrained entropic transport problems remains an important direction for future research.

Overall, this discussion highlights a conceptual shift: we can develop statistical optimal transport at the level of I -projections and moment restrictions, without requiring strong regularity assumptions on dual po-

tentials. This approach enables statistical inference across a wide range of constrained entropic transport problems, many of which researchers currently study primarily from variational or computational perspectives.

References

Alvarez-Mena, J. and O. Hernández-Lerma (2005). Convergence and approximation of optimization problems. *SIAM Journal on Optimization* 15(2), 527–539.

Arjovsky, M., S. Chintala, and L. Bottou (2017). Wasserstein generative adversarial networks. In *Proceedings of the 34th International Conference on Machine Learning*.

Backhoff-Veraguas, J. and G. Pammer (2020). Stability of martingale optimal transport and weak optimal transport. *Annals of Applied Probability* 30(2), 721–773.

Beiglböck, M., P. Henry-Labordère, and F. Penkner (2013). Model-independent bounds for option prices: A mass transport approach. *Finance and Stochastics* 17(3), 477–501.

Beiglböck, M. and N. Juillet (2016). On a problem of optimal transport under marginal martingale constraints. *The Annals of Probability* 44(1), 42 – 106.

Beiglböck, M. and F. Penkner (2013). A note on monotone martingale couplings. *Electronic Communications in Probability* 18, 1–11.

Bercu, B. and J. Bigot (2021). Asymptotic distribution and convergence rates of stochastic algorithms for entropic optimal transportation between probability measures. *The Annals of Statistics* 49(2), 968 – 987.

Bhattacharya, B. and R. L. Dykstra (1995). A General Duality Approach to I -Projections. *Journal of Statistical Planning and Inference* 47, 203–216.

Bigot, J. and T. Klein (2019). Central limit theorems for entropy-regularized optimal transport on finite spaces and statistical applications. *Electronic Journal of Statistics* 13(2), 5120–5150.

Chernozhukov, V., D. Chetverikov, and K. Kato (2014a). Anti-concentration and honest, adaptive confidence bands. *The Annals of Statistics* 42(5), 1787–1818.

Chernozhukov, V., D. Chetverikov, and K. Kato (2014b). Gaussian approximation of suprema of empirical processes. *The Annals of Statistics* 42(4), 1564–1597.

Chewi, S., J. Niles-Weed, and P. Rigollet (2025). *Statistical Optimal Transport*, Volume 2364 of *Lecture Notes in Mathematics*. Cham: Springer.

Chiappori, P.-A. and B. Salanié (2016). The econometrics of matching models. *Journal of Economic Literature* 54(3), 832–861.

Csiszár, I. (1975, 02). I -divergence geometry of probability distributions and minimization problems. *The Annals of Probability* 3(1), 146–158.

Cuturi, M. (2013). Sinkhorn distances: Lightspeed computation of optimal transport. In C. Burges, L. Bottou, M. Welling, Z. Ghahramani, and K. Weinberger (Eds.), *Advances in Neural Information Processing Systems*, Volume 26. Curran Associates, Inc.

del Barrio, E., A. González Sanz, J.-M. Loubes, and J. Niles-Weed (2023). An improved central limit

theorem and fast convergence rates for entropic transportation costs. *SIAM Journal on Mathematics of Data Science* 5(3), 639–669.

Flamary, R. and N. Courty (2021). *Optimal Transport for Machine Learning*. MIT Press. To appear.

Galichon, A. (2016). *Optimal Transport Methods in Economics*. Princeton University Press.

Gelfand, I. M. (1936). Sur un lemme de la theorie des espaces lineaires. *Communications de l'Institut des Sciences Mathématiques et mécaniques de l'Université de Kharkoff et la Société Mathématique de Kharkoff* 13, 35–40.

Genevay, A., M. Cuturi, G. Peyre, and F. Bach (2016). Stochastic optimization for large-scale optimal transport. *Neural Information Processing Systems*, 3440–3448.

Genevay, A., M. Cuturi, G. Peyré, and F. Bach (2019). Sample complexity of sinkhorn divergences. In *Advances in Neural Information Processing Systems*, Volume 32. Curran Associates, Inc.

Goldfeld, Z., K. Kato, G. Rioux, and R. Sadhu (2024). Statistical inference with regularized optimal transport. *Information and Inference: A Journal of the IMA* 13(1), 1–61.

Grenander, U. (1981). *Abstract Inference*. Wiley Series in Probability and Mathematical Statistics. John Wiley & Sons, Inc.

Lassalle, R. (2018). Causal transport plans and their monge–kantorovich problems. *Annales de l'Institut Henri Poincaré Probabilités et Statistiques* 54(1), 1–35.

Ledoux, M. and M. Talagrand (1991). *Probability in Banach Spaces: Isoperimetry and Processes*, Volume 23 of *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Springer.

Léonard (2012). From the Schrödinger problem to the Monge–Kantorovich problem. *Journal of Functional Analysis* 262(4), 1879–1920.

Luenberger, D. (1969). *Optimization by Vector Space Methods*. Series in Decision and Control. John Wiley & Sons, Inc.

Massart, P. (2000). About the constants in Talagrand’s concentration inequalities for empirical processes. *Annals of Probability* 28(2), 863–884.

Mena, G. and J. Niles-Weed (2019). Statistical bounds for entropic optimal transport: Sample complexity and the central limit theorem. In *Proceedings of the 33rd Conference on Neural Information Processing Systems (NeurIPS)*, pp. 1–12. Curran Associates, Inc.

Nutz, M. (2022). Introduction to entropic optimal transport. Revised lecture notes from courses given at Columbia University and Ecole Polytechnique.

Nutz, M. and F. Stebegg (2018). Canonical supermartingale couplings. *The Annals of Probability* 46(6), 3351 – 3398.

Nutz, M. and J. Wiesel (2022). Entropic optimal transport: convergence of potentials. *Probability Theory and Related Fields* 184, 401–424.

Nutz, M. and J. Wiesel (2024). Martingale schrödinger bridges. Preprint.

Pettis, B. J. (1938). On integration in vector spaces. *Transactions of The American Mathematical Society* 44, 70–74.

Peyré, G. and M. Cuturi (2019). *Computational Optimal Transport*, Volume 11 of *Foundations and Trends in Machine Learning*. Now Publishers.

Rigollet, P. and A. Stromme (2024). On the sample complexity of entropic optimal transport. *arXiv preprint arXiv:2402.XXXXX*. arXiv preprint.

Rudin, W. (1991). *Functional Analysis* (2 ed.). International Series in Pure and Applied Mathematics. McGraw-Hill.

Rüschendorf, L. (1985). The wasserstein distance and approximation theorems. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete* 70(1), 117–129.

Seguy, V., B. B. Damodaran, R. Flamary, N. Courty, A. Rolet, and M. Blondel (2018). Large-scale optimal transport and mapping estimation. In *International Conference on Learning Representations*.

Shapiro, A., D. Dentcheva, and A. Ruszczynski (2009). *Lectures on Stochastic Programming*. Society for Industrial and Applied Mathematics.

Steinwart, I. (2024). Reproducing kernel hilbert spaces cannot contain all continuous functions on a compact metric space. *Archiv der Mathematik* 122, 553–557.

Strassen, V. (1965). The existence of probability measures with given marginals. *Annals of Mathematical Statistics* 36(2), 423–439.

Tabri, R. V. (2025). The information projection in moment inequality models: Existence, dual representation, and approximation. mathematics of operations research. *Mathematics of Operations Research*.

Tang, X., M. Shavlovsky, H. Rahamanian, T. Xiao, and L. Ying (2025). Entropic optimal transport with martingale constraints. *arXiv preprint arXiv:2508.17641*.

van der Vaart, A. W. and J. Wellner (1996). *Weak Convergence and Empirical Processes* (First ed.). Springer Series in Statistics. Springer.

Villani, C. (2009). *Optimal Transport: Old and New*. Grundlehren der mathematischen Wissenschaften. Springer Berlin Heidelberg.

Wainwright, M. J. (2019). *High-Dimensional Statistics: A Non-Asymptotic Viewpoint*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press.

A Proofs of Results

A.1 Proposition 2.1

Proof. **Part 1.** We establish $\arg \inf \left\{ \int_{\Omega} e^z dR_{\gamma} : z \in \mathcal{D} \right\} \neq \emptyset$ through the existence of an element being a cluster point of the sequence $\{z_{0,n}\}_{n \geq 1}$, in the norm topology of $L_1(R_{\gamma})$, where

$$z_{0,n} = \arg \inf \left\{ \int_{\Omega} e^y dR_{\gamma} : z \in \mathcal{D}_n \right\} \quad \text{with} \quad \mathcal{D}_n = \{z \in \mathcal{D} : \alpha \leq \bar{\alpha}_n\}$$

for each n , and $\bar{\alpha}_n \nearrow \infty$ as $n \rightarrow \infty$. Lemma B.1 shows that I meet all of the conditions to apply Theorem 2.3 of Alvarez-Mena and Hernández-Lerma (2005), so that

$$\text{OL} \{ \{z_{0,n}\}_{n \geq 1} \} \subset \arg \inf \left\{ \int_{\Omega} e^z dR_{\gamma} : z \in \mathcal{D} \right\}, \quad (\text{A.1})$$

holds, where $\text{OL}\{\{z_{0,n}\}_{n \geq 1}\}$ denotes the outer limit of $\{z_{0,n}\}_{n \geq 1}$, which is also the set of cluster points of the sequence in the norm topology of $L_1(R_\gamma)$. Lemma B.2 establishes $\text{OL}\{\{z_{0,n}\}_{n \geq 1}\} \neq \emptyset$. Hence, $\arg \inf \{\int_\Omega e^z dR_\gamma : z \in \mathcal{D}\} \neq \emptyset$.

Now I shall establish the uniqueness a.s. $-R_\gamma$ of the minimizer. As any $v \in \mathcal{V}$ satisfies $\int_\Omega e^v dR_\gamma < \infty$, the minimizers cannot be where the objective function equals ∞ . Combining this implication with the strict convexity of the map $z \mapsto \int_\Omega e^z dR_\gamma$ on \mathcal{D} , implies that there is a unique minimizer (up to equivalence class). Let $\beta \in (0, 1)$ and $z_1, z_2 \in \mathcal{D}$ such that $z_1 \neq z_2$ holds as equivalence classes. Additionally, let $z_3 = \beta z_1 + (1 - \beta) z_2$. Then $\int_\Omega e^{z_3} dR_\gamma < \beta \int_\Omega e^{z_1} dR_\gamma + (1 - \beta) \int_\Omega e^{z_2} dR_\gamma$, holds, by the strict convexity of the exponential function. This establishes the strict convexity of the map, and hence, the set of minimizers $\arg \inf \{\int_\Omega e^z dR_\gamma : z \in \mathcal{D}\}$, is unique up to equivalence class.

Next, I develop the representation of the I -projection's R_γ -density. From the above arguments let $z_{0,\gamma} = \arg \inf \{\int_\Omega e^z dQ : z \in \mathcal{D}\}$. The set \mathcal{D} is convex, and the objective function $g(z) = \int_\Omega e^z dR_\gamma$ is Gâteaux differentiable, then by Theorem 2 on page 178 of Luenberger (1969), $\frac{d}{dt}g(z_{0,\gamma} + t(z - z_{0,\gamma}))|_{t=0} \geq 0$ $\forall z \in \mathcal{D}$, yielding $\int_\Omega (y - y_0) e^{y_0} dQ \geq 0 \forall y \in \mathcal{D}$. By choosing $y = cy_0$ first with $c > 1$ and then with $c < 1$ (since \mathcal{D} is also a cone), I obtain

$$\int_\Omega z_{0,\gamma} e^{z_{0,\gamma}} dR_\gamma = 0, \text{ and } \int_\Omega z e^{z_{0,\gamma}} dR_\gamma \geq 0 \forall z \in \mathcal{D}. \quad (\text{A.2})$$

Let $p_\gamma = \frac{e^{z_{0,\gamma}}}{\int_\Omega e^{z_{0,\gamma}} dR_\gamma}$, and note that the second part of (A.2) implies that $\int_\Omega v p_\gamma dR_\gamma \geq 0 \forall v \in \mathcal{V}$; hence, $p_\gamma \in \mathcal{M}$. Furthermore, $m(p_\gamma) + \log(\int_\Omega e^{z_{0,\gamma}} dR_\gamma) = \int_\Omega z_{0,\gamma} e^{z_{0,\gamma}} dR_\gamma = 0$, holds, by the first part of (A.2). Hence, by Theorem 2.2 of Bhattacharya and Dykstra (1995), p_γ solves the I -projection problem.

Part 2. The proof proceeds by the direct method. Firstly, observe that $\exists v \in \mathcal{V}$ such that $\int_\Omega v dR_\gamma > 0$; otherwise, the solution of the I -projection problem (2.2) would be R_γ . Hence, by Part (ii) of Theorem 1 in Tabri (2025), $z_{0,\gamma} \in \overline{\text{span}_+(B)}$, where $\overline{\text{span}_+(B)}$ is the $L_1(R_\gamma)$ -norm closure of the positive linear span of $B = \{v \in \mathcal{V} : \int_\Omega v dP_\gamma = 0\}$. Now $B = \mathcal{V}$ must hold, as the I -projection problem is a moment inequality formulation of the equality constraints that set the marginal distributions. Therefore, I must establish that $y_0 \in \overline{\text{span}_+(\mathcal{V})}$, holds. Since R_γ violates the moments inequality restrictions, $z_{0,\gamma} \in \mathcal{D}$ implies that $z_{0,\gamma} = \alpha_{0,\gamma} z'_{0,\gamma}$ with $\alpha_{0,\gamma} > 0$ and $z'_{0,\gamma} \in \overline{\text{co}}(\mathcal{V})$. The result of Lemma B.5 implies that there are only two cases to consider in establishing the desired result: (i) $z'_{0,\gamma} \in \text{ex}(\overline{\text{co}}(\mathcal{V}))$, and (ii) $z'_{0,\gamma} \notin \text{ex}(\overline{\text{co}}(\mathcal{V}))$, where $\text{ex}(\overline{\text{co}}(\mathcal{V}))$ denotes the set of extreme points of $\overline{\text{co}}(\mathcal{V})$.

Starting with case (i), since $\text{ex}(\overline{\text{co}}(\mathcal{V})) \subset \mathcal{V}$ also by Lemma B.5, it must be that $z'_{0,\gamma} \in \mathcal{V}$, and therefore, $z_{0,\gamma} \in \overline{\text{span}_+(\mathcal{V})}$. Next, consider case (ii): $z'_{0,\gamma} \notin \text{ex}(\overline{\text{co}}(\mathcal{V}))$. Then, $\exists n \in \mathbb{Z}_+, p_i > 0$ for each $i = 1, \dots, n$ such that $\sum_{i=1}^n p_i = 1$, for which $z_{0,\gamma} = \alpha_{0,\gamma} \sum_{i=1}^n p_i v_i$, where $\alpha_{0,\gamma} > 0$ and $\{v_1, \dots, v_n\} \subset \text{ex}(\overline{\text{co}}(\mathcal{V})) \subset \mathcal{V}$. Consequently, $z_{0,\gamma} \in \text{span}_+(\mathcal{V}) \subset \overline{\text{span}_+(\mathcal{V})}$. \blacksquare

A.2 Corollary 2.1

Proof. The proof proceeds by the direct method. I only need to establish the equality

$\mathcal{S}^* = \{\xi \in \Upsilon : z_{0,\gamma} = \int_{\mathcal{V}} v d\xi(v)\}$, as the non-emptiness trivially holds because $z_{0,\gamma}$ exists under the aforementioned conditions, and that $z_{0,\gamma}$ has a G-F integral representation (2.9). I start with the direction “ \subset ”.

Let $\xi' \in \mathcal{S}^*$, then we must show that $z_{0,\gamma} = \int_{\mathcal{V}} v d\xi'(v)$, holds. By optimality of ξ' , $\int_{\Omega} e^{\int_{\mathcal{V}} v d\xi'(v)} dR_{\gamma} \leq \int_{\Omega} e^{\int_{\mathcal{V}} v d\xi(v)} dR_{\gamma} \quad \forall \xi \in \Upsilon$, and that $\exists z' \in \mathcal{D}$ such that $z' = \int_{\mathcal{V}} v d\xi'(v)$, observe that

$$\int_{\Omega} e^{z'} dR_{\gamma} = \int_{\Omega} e^{\int_{\mathcal{V}} v d\xi'(v)} dR_{\gamma} \leq \int_{\Omega} e^{\int_{\mathcal{V}} v d\xi(v)} dR_{\gamma} \quad \forall \xi \in \Upsilon. \quad (\text{A.3})$$

Now by the G-F integral representation (2.9), I can rewrite (A.3) as

$$\int_{\Omega} e^{z'} dR_{\gamma} = \int_{\Omega} e^{\int_{\mathcal{V}} v d\xi'(v)} dR_{\gamma} \leq \int_{\Omega} e^z dR_{\gamma} \quad \forall z \in \mathcal{D}, \quad (\text{A.4})$$

and hence, $\int_{\Omega} e^{\int_{\mathcal{V}} v d\xi'(v)} dR_{\gamma} = \int_{\Omega} e^{z'} dR_{\gamma} = \int_{\Omega} e^{z_{0,\gamma}} dR_{\gamma}$ must hold by Proposition 2.1. Since $z_{0,\gamma}$ is unique (up to equivalence class) I must have $z_{0,\gamma} = z'$ a.s.- R_{γ} , and hence, $z_{0,\gamma} = \int_{\mathcal{V}} v d\xi'(v)$ a.s.- R_{γ} by the G-F integral representation (2.9). Therefore, $\xi' \in \{\xi \in \Upsilon : z_{0,\gamma} = \int_{\mathcal{V}} v d\xi(v)\}$.

Next, I consider the direction “ \supset ”. let $\xi' \in \{\xi \in \Upsilon : z_{0,\gamma} = \int_{\mathcal{V}} v d\xi(v)\}$, then we must show that $\xi' \in \mathcal{S}^*$. By the G-F integral representation (2.9) and the optimality of $z_{0,\gamma}$, observe that

$$\int_{\Omega} e^{\int_{\mathcal{V}} v d\xi'(v)} dR_{\gamma} = \int_{\Omega} e^{z_{0,\gamma}} dR_{\gamma} \leq \int_{\Omega} e^z dR_{\gamma} \quad \forall z \in \mathcal{D}. \quad (\text{A.5})$$

Now by the G-F integral representation (2.9), I can rewrite (A.5) as

$$\int_{\Omega} e^{\int_{\mathcal{V}} v d\xi'(v)} dR_{\gamma} \leq \int_{\Omega} e^{\int_{\mathcal{V}} v d\xi(v)} dR_{\gamma} \quad \forall \xi \in \Upsilon,$$

and hence, $\xi' \in \mathcal{S}^*$. ■

A.3 Proposition 2.2

Proof. The proof proceeds by the direct method. Under Assumption 1, the cost function, c , is bounded; i.e., $\|c\|_{\infty} < \infty$. The I -projection, P_{γ} , has density with respect to $P_X \otimes P_Y$ given by

$$\frac{dP_{\gamma}}{d(P_X \otimes P_Y)}(x, y) = \exp \{ \gamma (z_{0,\gamma}(x, y)/\gamma - c(x, y)) \} / (\vartheta^* a_{\gamma}). \quad (\text{A.6})$$

Under Assumption 1, I can apply Theorem 2.1 in Nutz (2022) to establish that this density is also that of the unique coupling $P \in \Pi(P_X, P_Y)$ having the form

$$\frac{dP_{\gamma}}{d(P_X \otimes P_Y)}(x, y) = \exp \{ \gamma (\varphi_{\gamma}(x) + \psi_{\gamma}(y) - c(x, y)) \} / (\vartheta^* a_{\gamma}), \quad (\text{A.7})$$

with $\int_{\mathcal{X}} \varphi_{\gamma}(x) dP_X = \int_{\mathcal{Y}} \psi_{\gamma}(y) dP_Y$, where $\varphi_{\gamma} \in L_1(P_X)$ and $\psi_{\gamma} \in L_1(P_Y)$ are the unique solution of the EOT's dual problem, given by

$$\sup_{f \in L_1(P_X), g \in L_1(P_Y)} \left(\int_{\mathcal{X}} f dP_X + \int_{\mathcal{Y}} g dP_Y - \frac{1}{\gamma} \left(\int_{\mathcal{X} \times \mathcal{Y}} e^{\gamma(f(x) + g(y) - c(x, y))} d(P_X \otimes P_Y) - 1 \right) \right).$$

Because $a_{\gamma} \leq 1$, holds, the above normalization implies $\int_{\mathcal{X}} \varphi_{\gamma}(x) dP_X, \int_{\mathcal{Y}} \psi_{\gamma}(y) dP_Y \geq 0$, under Assumption 1; see Remark 4.10 in Nutz (2022).

In consequence, I obtain $z_{0,\gamma}(x, y)/\gamma = \varphi_{\gamma}(x) + \psi_{\gamma}(y)$ upon equating the two forms of the density $\frac{dP_{\gamma}}{d(P_X \otimes P_Y)}$. Now because $z_{0,\gamma} = \alpha_{0,\gamma} \int_{\mathcal{V}} v d\mu_{0,\gamma}(v)$, integrating both sides with respect to $P_X \otimes P_Y$ yields, $\int_{\mathcal{X} \times \mathcal{Y}} z_{0,\gamma} d(P_X \otimes P_Y) = 0$, implying the desired normalization

$$0 = \int_{\mathcal{X} \times \mathcal{Y}} (\varphi(x) + \psi(y)) d(P_X \otimes P_Y) = \int_{\mathcal{X}} \varphi(x) dP_X + \int_{\mathcal{Y}} \psi(y) dP_Y. \quad (\text{A.8})$$

Note that $\int_{\mathcal{X} \times \mathcal{Y}} z_{0,\gamma} d(P_X \otimes P_Y) = 0$ follows from arguments in the proof of Part 2 of Proposition 2.1 that establish $z_{0,\gamma} = \alpha_{0,\gamma} z'_{0,\gamma}$ with $\alpha_{0,\gamma} > 0$ and $z'_{0,\gamma} \in \text{ex}(\overline{\text{co}}(\mathcal{V}))$ or $z'_{0,\gamma} \notin \text{ex}(\overline{\text{co}}(\mathcal{V}))$. Observe that $\int_{\mathcal{X} \times \mathcal{Y}} z'_{0,\gamma} d(P_X \otimes P_Y) = 0$, holds, in either case as the moment functions had the form described in (2.3).

Re-writing $z_{0,\gamma}/\gamma = \varphi_{\gamma} + \psi_{\gamma}$ as $(\alpha_{0,\gamma}/\gamma) \int_{\mathcal{V}} v d\mu_{0,\gamma}(v) = \varphi_{\gamma} + \psi_{\gamma}$, it follows that $\int_{\mathcal{V}} v d\mu_{0,\gamma}(v) = \varphi'_{\gamma} + \psi'_{\gamma}$, where $\varphi'_{\gamma} = (\gamma/\alpha_{0,\gamma})\varphi_{\gamma}$ and $\psi'_{\gamma} = (\gamma/\alpha_{0,\gamma})\psi_{\gamma}$. Now because $\int_{\mathcal{V}} v d\mu_{0,\gamma}(v) \in \overline{\text{co}}(\mathcal{V})$, I must have that $\varphi'_{\gamma}, \psi'_{\gamma} \in [-1, 1]$. Hence, by Lemma 4.9 in Nutz (2022), $(\alpha_{0,\gamma}/\gamma)\varphi'_{\gamma}(x) \leq \|c\|_{\infty}$ for each $x \in \mathcal{X}$. Then, for x such that $\varphi'_{\gamma}(x) > 0$, $(\alpha_{0,\gamma}/\gamma) \leq \|c\|_{\infty}/\varphi'_{\gamma}(x)$, and therefore, $(\alpha_{0,\gamma}/\gamma) \leq \inf \{\|c\|_{\infty}/\varphi'_{\gamma}(x) : \varphi'_{\gamma}(x) > 0\}$. Now observe that

$$\inf \{\|c\|_{\infty}/\varphi'_{\gamma}(x) : \varphi'_{\gamma}(x) > 0\} = \frac{\|c\|_{\infty}}{\sup_{\{x: \varphi'_{\gamma}(x) > 0\}} \varphi'_{\gamma}(x)},$$

and because φ'_{γ} must be a linear combination of functions of the form $x \mapsto F_X(x') - 1 [x \preceq x']$ or $x \mapsto 1 [x \preceq x'] - F_X(x')$, where the coefficients are positive and bounded from above by 1,

$$\frac{\|c\|_{\infty}}{\sup_{\{x: \varphi'_{\gamma}(x) > 0\}} \varphi'_{\gamma}(x)} \leq \frac{\|c\|_{\infty}}{1 - F_X(\inf \mathcal{X})}.$$

The same line of arguments holds if I also use ψ'_{γ} instead of φ'_{γ} , so that

$$\frac{\|c\|_{\infty}}{\sup_{\{x: \varphi'_{\gamma}(x) > 0\}} \varphi'_{\gamma}(x)} \leq \frac{\|c\|_{\infty}}{\max \{1 - F_X(\inf \mathcal{X}), 1 - F_Y(\inf \mathcal{Y})\}}$$

must hold. Hence $(\alpha_{0,\gamma}/\gamma) \leq \frac{\|c\|_{\infty}}{\max \{1 - F_X(\inf \mathcal{X}), 1 - F_Y(\inf \mathcal{Y})\}}$, as desired. ■

A.4 Proposition 2.3

Proof. The proof proceeds by the direct method.

Part 1. Let $\xi_{0,\ell} = \alpha_{0,\gamma} \sum_{i=1}^{n_\ell} \mu_{i,0,\ell} \delta_{v_i}$ such that $\mu_{i,0,\ell} = \mu_{0,\gamma}(E_{i,\ell})$ for each $i = 1, \dots, n_\ell$. By optimality of $\xi_{0,\gamma}$ and $(\alpha_\ell^*, \mu_\ell^*) \in \mathcal{S}_\ell^*$, I have the following inequalities

$$\vartheta^* = \int_{\Omega} e^{\int_{\mathcal{V}} v \, d\xi_{0,\ell}(v)} \, dR_\gamma \leq \vartheta_\ell^* = \int_{\Omega} G_\ell(\omega; \alpha_\ell^*, \mu_\ell^*) \, dR_\gamma \leq \int_{\Omega} e^{\int_{\mathcal{V}} v \, d\xi_{0,\ell}(v)} \, dR_\gamma. \quad (\text{A.9})$$

Then,

$$\left| \vartheta^* - \int_{\Omega} e^{\int_{\mathcal{V}} v \, d\xi_{0,\ell}(v)} \, dR_\gamma \right| = \left| \int_{\Omega} e^{\int_{\mathcal{V}} v \, d\xi_{0,\gamma}(v)} \, dR_\gamma - \int_{\Omega} e^{\int_{\mathcal{V}} v \, d\xi_{0,\ell}(v)} \, dR_\gamma \right| \quad (\text{A.10})$$

$$\begin{aligned} &\leq e^{\alpha_{0,\gamma}} \alpha_{0,\gamma} \left\| \int_{\mathcal{V}} v \, d\mu_{0,\gamma}(v) - \int_{\mathcal{V}} v \, d\mu_{0,\ell}(v) \right\|_{L_1(R_\gamma)} \\ &\leq \epsilon_\ell \gamma \kappa \|c\|_\infty e^{\gamma \kappa \|c\|_\infty}, \end{aligned} \quad (\text{A.11})$$

by applying the Mean Value Theorem to the discrepancy (A.10) with respect to the exponential function, and then using the result of Proposition 2.2 and the fact that $\int_{\mathcal{V}} v \, d\mu_{0,\gamma}(v), \int_{\mathcal{V}} v \, d\mu_{0,\ell}(v) \in [-1, 1]$ a.s.- R_γ , holds, to deduce the upper bound (A.11). This derivation holds for each ℓ , and hence,

$\lim_{\ell \rightarrow \infty} \int_{\Omega} e^{\int_{\mathcal{V}} v \, d\xi_{0,\ell}(v)} \, dR_\gamma = \vartheta^*$, because $\lim_{\ell \rightarrow \infty} \epsilon_\ell = 0$. Finally, from the inequalities (A.9), the above limit implies $\lim_{\ell \rightarrow \infty} \vartheta_\ell^* = \vartheta^*$.

Part 2. By the inequalities (A.9), observe that for each ℓ ,

$$\begin{aligned} |\log \vartheta_\ell^* - \log \vartheta^*| &= \log \vartheta_\ell^* - \log \vartheta^* = \log \left(1 + \frac{\vartheta_\ell^* - \vartheta^*}{\vartheta^*} \right) \leq \frac{\vartheta_\ell^* - \vartheta^*}{\vartheta^*} \\ &\leq \epsilon_\ell \kappa \gamma \|c\|_\infty e^{\gamma(\kappa \|c\|_\infty + \bar{c}) + \log a_\gamma}, \end{aligned}$$

where the second inequality arises from bounding ϑ^* from below by way of Lemma C.2.

Part 3. The proof follows steps identical to those in Part 2 of Theorem 4 in Tabri (2025). I present them here for completeness. Let ξ' be an accumulation point of $\{\xi_\ell^*\}_{\ell \geq 1}$ in the weak-star topology of $C(\mathcal{V})^*$. Therefore, there exists a subsequence $\{\xi_{\ell_h}^*\}_{h \geq 1}$ such that $\xi_{\ell_h}^* \xrightarrow{w^*} \xi'$. By observing that

$$\begin{aligned} \vartheta^* &= \int_{\Omega} e^{\int_{\mathcal{V}} v \, d\xi_{0,\gamma}(v)} \, dR_\gamma \leq \int_{\Omega} e^{\int_{\mathcal{V}} v \, d\xi_{\ell_h}(v)} \, dR_\gamma = \int_{\Omega} e^{\alpha_{\ell_h} \sum_{i=1}^{n_{\ell_h}} \mu_{i,\ell_h} v_i} \, dR_\gamma \\ &\leq \int_{\Omega} e^{\alpha_{0,\gamma} \sum_{i=1}^{n_{\ell_h}} \mu_0(E_{i,\ell_h}) v_i} \, dR_\gamma, \end{aligned}$$

holds, Part 1 of this proposition implies $\lim_{h \rightarrow \infty} \int_{\Omega} e^{\int_{\mathcal{V}} v \, d\xi_{\ell_h}} \, dR_\gamma = \int_{\Omega} e^{\int_{\mathcal{V}} v \, d\xi_{0,\gamma}(v)} \, dR_\gamma$. Now using the fact that the map $\xi \mapsto \int_{\mathcal{V}} v \, d\xi$ is continuous when $C(\mathcal{V})^*$ is given the weak-star topology and $L_1(R_\gamma)$ has the weak topology, it follows that $\int_{\mathcal{V}} v \, d\xi_{\ell_h} \xrightarrow{w} \int_{\mathcal{V}} v \, d\xi'$. Applying the Skorohod Representation Theorem, there exists a probability space $(\Omega', \mathcal{F}', Q')$ and real-valued measurable functions $\{z_h\}_{h \geq 1}$ and z on this probability space, such that

- (a) $\{z_h\}_{h \geq 1}$ and z have the same probability distributions as $\{e^{\int_{\mathcal{V}} v d\xi_{\ell_h}}\}_{h \geq 1}$ and $e^{\int_{\mathcal{V}} v d\xi'}$, respectively, and
- (b) $\lim_{h \rightarrow \infty} z_h(\omega') = z(\omega')$ a.s. $-Q'$.

One can set $\Omega' = [0, 1]$ with $\{z_h\}_{h \geq 1}$ and z being the quantile functions of $\{e^{\int_{\mathcal{V}} v d\xi_{\ell_h}}\}_{h \geq 1}$ and $e^{\int_{\mathcal{V}} v d\xi'}$, respectively. Consequently, $z_h(\omega'), z(\omega') \geq 0$ for all $\omega' \in [0, 1]$. Now we can use Fatou's Lemma to deduce the desired result:

$$\begin{aligned} \int_{\Omega} e^{\int_{\mathcal{V}} v d\xi'} dR_{\gamma} &= \int_{[0,1]} z dQ' \leq \liminf_{h \rightarrow \infty} \int_{[0,1]} z_h dQ' = \liminf_{h \rightarrow \infty} \int_{\Omega} e^{\int_{\mathcal{V}} v d\xi_{\ell_h}} dR_{\gamma} \\ &= \int_{\Omega} e^{\int_{\mathcal{V}} v d\xi_{0,\gamma}(v)} dR_{\gamma}. \end{aligned}$$

Therefore, $\xi' \in \arg \inf \left\{ \int_{\Omega} e^{\int_{\mathcal{V}} v d\xi(v)} dR_{\gamma} : \xi \in \Upsilon \right\}$ since $\xi_{0,\gamma}$ is an element of that set. This concludes the proof. \blacksquare

A.5 Proof of Theorem 1

Proof. The proof proceeds by the direct method. I firstly introduce some notation and set the stage to simplify the exposition. Recall that $\xi_{0,\gamma} = \alpha_{0,\gamma} \cdot \mu_{0,\gamma} \in \mathcal{S}^*$. Given ℓ , and hence, ϵ_{ℓ} , there corresponds a finite partition $\{E_{i,\ell}\}_{i=1}^{n_{\ell}}$ of \mathcal{V} with the accuracy (2.12), and let $v_i \in E_{i,\ell}$ for $i = 1, \dots, n_{\ell}$. Define $\xi_{0,\ell} = \alpha_{0,\gamma} \sum_{i=1}^{n_{\ell}} \mu_{i,0,\ell} \delta_{v_i}$, where $\mu_{i,0,\ell} = \mu_{0,\gamma}(E_{i,\ell})$ and δ_{v_i} is the Dirac delta function at v_i , for each $i = 1, \dots, n_{\ell}$. Now based on this partition of \mathcal{V} , let $(\alpha_{\ell}^*, \mu_{\ell}^*) \in \mathcal{S}_{\ell}^*$, $(\alpha_{\ell}, \mu_{\ell}) \in S_{\ell}$ and $(\hat{\alpha}_{\ell}, \hat{\mu}_{\ell}) \in \hat{S}_{\ell}$.

Note that for each ℓ , $\hat{S}_{\ell} \neq \emptyset$ for almost every sample realization of ω_{ℓ}^N under R_{γ} . The realizations of the sample $\omega^{N_{\ell}}$ where $\hat{S}_{\ell} = \emptyset$ arises is on a set of probability measure zero. This is because the Extreme Value Theorem applies for each realization of $\omega^{N_{\ell}}$, as $\bar{\mathcal{C}}_{\ell}$ is a nonempty compact subset of $\mathbb{R}^{n_{\ell}+1}$ and the objective function is a Carathéodory function – see Remark 3.1.

Part 1(i). Following the proof of Part (i) in Proposition 2.3, using the optimality of $\xi_{0,\gamma}$ and $(\alpha_{\ell}^*, \mu_{\ell}^*)$, I have the inequalities

$$\vartheta^* = \int_{\Omega} e^{\int_{\mathcal{V}} v d\xi_{0}(v)} dR_{\gamma} \leq \vartheta_{\ell}^* \leq \vartheta_{\ell} \leq \int_{\Omega} e^{\int_{\mathcal{V}} v d\xi_{0,\ell}(v)} dR_{\gamma}. \quad (\text{A.12})$$

Because $\lim_{\ell \rightarrow \infty} \int_{\Omega} e^{\int_{\mathcal{V}} v d\xi_{0,\ell}(v)} dR_{\gamma} = \vartheta^*$ holds by Part 1 in Proposition 2.3, by a sandwiching argument with the inequalities (A.12), it follows that $\lim_{\ell \rightarrow \infty} \vartheta_{\ell} = \lim_{\ell \rightarrow \infty} \vartheta_{\ell}^* = \vartheta^*$, must also hold. Now along realizations $\omega^{N_{\ell}}$ such that $\hat{S}_{\ell} \neq \emptyset$, I can add and subtract $\hat{\vartheta}_{\ell}$ and ϑ_{ℓ} in the center of the chain of inequalities (A.12) to obtain

$$\int_{\Omega} e^{\int_{\mathcal{V}} v d\xi_{0}(v)} dR_{\gamma} \leq \vartheta_{\ell}^* - \vartheta_{\ell} + \vartheta_{\ell} - \hat{\vartheta}_{\ell} + \hat{\vartheta}_{\ell} \leq \int_{\Omega} e^{\int_{\mathcal{V}} v d\xi_{0,\ell}(v)} dR_{\gamma}.$$

Hence, to obtain the desired result, I must establish that $\vartheta_{\ell} - \hat{\vartheta}_{\ell} = o_{a.s.}(1)$, holds.

Toward that end, note that $\vartheta_\ell - \hat{\vartheta}_\ell$ is given by

$$\int_{\Omega} G_\ell(\omega; \alpha_\ell, \mu_\ell) dR_\gamma - \frac{1}{N_\ell} \sum_{j=1} G_\ell(\omega_j; \hat{\alpha}_\ell, \hat{\mu}_\ell), \quad (\text{A.13})$$

I shall bound this term from above and from below by $o_{a.s}(1)$ random variables. The upper bound is

$$\int_{\Omega} G_\ell(\omega; \hat{\alpha}_\ell, \hat{\mu}_\ell) dR_\gamma - \frac{1}{N_\ell} \sum_{j=1} G_\ell(\omega_j; \hat{\alpha}_\ell, \hat{\mu}_\ell), \quad (\text{A.14})$$

as $\int_{\Omega} G_\ell(\omega; \hat{\alpha}_\ell, \hat{\mu}_\ell) dR_\gamma \geq \int_{\Omega} G_\ell(\omega; \alpha_\ell, \mu_\ell) dR_\gamma$. Now the absolute value of the term in (A.14) is bounded from above by

$$\sup_{(\alpha, \mu) \in \mathcal{C}_\ell} \left| \int_{\Omega} G_\ell(\omega; \alpha, \mu) dR_\gamma - \frac{1}{N_\ell} \sum_{j=1} G_\ell(\omega_j; \alpha, \mu) \right|, \quad (\text{A.15})$$

From Lemma C.1, and Corollary C.1, the Rademacher complexity of \mathcal{G}_ℓ , $\mathcal{R}_{N_\ell}(\mathcal{G}_\ell)$, satisfies $\mathcal{R}_{N_\ell}(\mathcal{G}_\ell) \leq \gamma \kappa \|c\|_\infty e^{\gamma \kappa \|c\|_\infty} \sqrt{\frac{2 \log n_\ell}{N_\ell}} \quad \forall \ell$, and hence, it vanishes under the limit condition (3.4). Therefore, by an application of by Proposition C.1, the term (A.15) must be $o_{a.s}(1)$. For the lower bound, observe that the term in (A.13) is bounded from below by

$$\int_{\Omega} G_\ell(\omega; \alpha_\ell, \mu_\ell) dR_\gamma - \frac{1}{N_\ell} \sum_{j=1} G_\ell(\omega_j; \alpha_\ell, \mu_\ell) \quad (\text{A.16})$$

as $\frac{1}{N_\ell} \sum_{j=1} G_\ell(\omega_j; \alpha_\ell, \mu_\ell) \geq \frac{1}{N_\ell} \sum_{j=1} G_\ell(\omega_j; \hat{\alpha}_\ell, \hat{\mu}_\ell)$. Now the absolute value of the term in (A.16) is bounded from above by the term in (A.15), so that it is also $o_{a.s}(1)$ by the same arguments. In consequence, $\vartheta^* = \int_{\Omega} e^{\int_{\mathcal{V}} v d\xi_0(v)} dR_\gamma \leq o(1) + o_{a.s}(1) + \hat{\vartheta}_\ell \leq \int_{\Omega} e^{\int_{\mathcal{V}} v d\xi_{0,\ell}(v)} dR_\gamma$, and taking the limit as $\ell \rightarrow \infty$ on all sides of these inequalities yields $\lim_{\ell \rightarrow \infty} \hat{\vartheta}_\ell = \vartheta^*$ a.s. $- R_\gamma$.

Part 1(ii). Suppose that $\{\hat{\xi}_{\ell_h}\}_{h \geq 1}$ converges to ξ' in the weak-star topology of $C(\mathcal{V})^*$ as $h \rightarrow \infty$, with probability 1. As the sequence $\{\hat{\vartheta}_{\ell_h}\}_{h \geq 1}$ is a subsequence of $\{\hat{\vartheta}_\ell\}_{\ell \geq 1}$, the result of Part 1(i) of this theorem implies $\lim_{h \rightarrow \infty} \hat{\vartheta}_{\ell_h} = \vartheta^*$ a.s. $- R_\gamma$. In consequence, I must establish the limit $\lim_{h \rightarrow \infty} \hat{\vartheta}_{\ell_h} = \int_{\Omega} e^{\int_{\mathcal{V}} v d\xi'} dR_\gamma$ with probability 1, to obtain the desired result, as it implies $\int_{\Omega} e^{\int_{\mathcal{V}} v d\xi'} dR_\gamma = \vartheta^*$, holds, and hence, $\xi' \in \mathcal{S}^*$.

By using the fact that the map $\xi \mapsto \int_{\mathcal{V}} v d\xi$ is continuous when $C(\mathcal{V})^*$ is given the weak-star topology and $L_1(R_\gamma)$ has the weak topology, it follows that $\left\{ \int_{\mathcal{V}} v d\hat{\xi}_{\ell_h} \right\}_{h \geq 1}$ converges to $\int_{\mathcal{V}} v d\xi'$ in the weak topology of $L_1(R_\gamma)$, with probability 1. I can apply the Skorohod Representation Theorem as follows: for almost every realization, there exists a probability space $(\Omega', \mathcal{F}', Q')$ and real-valued measurable functions $\{z_h\}_{h \geq 1}$ and z on this probability space, such that

- (a) $\{z_h\}_{h \geq 1}$ and z have the same probability distributions as $\{e^{\int_{\mathcal{V}} v d\hat{\xi}_{\ell_h}}\}_{h \geq 1}$ and $e^{\int_{\mathcal{V}} v d\xi'}$, respectively, and
- (b) $\lim_{h \rightarrow \infty} z_h(\omega') = z(\omega')$ a.s. $- Q'$.

One can set $\Omega' = [0, 1]$ with $\{z_h\}_{h \geq 1}$ and z being the quantile functions of $\{e^{\int_{\mathcal{V}} v d\hat{\xi}_{\ell_h}}\}_{h \geq 1}$ and $e^{\int_{\mathcal{V}} v d\xi'}$,

respectively. Consequently, $z_h(\omega'), z(\omega') \geq 0$ for all $\omega' \in [0, 1]$. Now applying Fatou's lemma on a per realization basis, for almost every realization,

$$\begin{aligned} \int_{\Omega} e^{\int_{\mathcal{V}} v \, d\xi'} \, dR_{\gamma} &= \int_{[0,1]} z \, dQ' \leq \liminf_{h \rightarrow \infty} \int_{[0,1]} z_h \, dQ' = \liminf_{h \rightarrow \infty} \int_{\Omega} e^{\int_{\mathcal{V}} v \, d\xi_{\ell_h}} \, dR_{\gamma} \\ &= \int_{\Omega} e^{\int_{\mathcal{V}} v \, d\xi_{0,\gamma}(v)} \, dR_{\gamma}. \end{aligned}$$

Therefore, $\xi' \in \arg \inf \left\{ \int_{\Omega} e^{\int_{\mathcal{V}} v \, d\xi(v)} \, dR_{\gamma} : \xi \in \Upsilon \right\}$ because $\xi_{0,\gamma}$ is an element of that set. This concludes the proof.

Part 2. By the triangle inequality, $|\log \hat{\vartheta}_{\ell} - \log \vartheta^*|$ is bounded from above by

$$|\log \hat{\vartheta}_{\ell} - \log \vartheta_{\ell}| + |\log \vartheta_{\ell} - \log \vartheta_{\ell}^*| + |\log \vartheta_{\ell}^* - \log \vartheta^*|.$$

The last two terms in the above display do not depend on the sample, and hence, are non-stochastic. I firstly focus on these non-stochastic terms. By Part 2 of Proposition 2.3,

$$|\log \vartheta_{\ell}^* - \log \vartheta^*| = \log \vartheta_{\ell}^* - \log \vartheta^* \leq \epsilon_{\ell} \gamma \kappa \|c\|_{\infty} e^{\gamma(\kappa\|c\|_{\infty} + \bar{c}) + \log a_{\gamma}}.$$

Furthermore, $|\log \vartheta_{\ell} - \log \vartheta_{\ell}^*| = \log \vartheta_{\ell}^* - \log \vartheta_{\ell} = \log \left(1 + \frac{\vartheta_{\ell}^* - \vartheta_{\ell}}{\vartheta_{\ell}} \right) \leq \frac{\vartheta_{\ell}^* - \vartheta_{\ell}}{\vartheta_{\ell}} \leq \frac{\vartheta_{\ell}^* - \vartheta^*}{\vartheta^*}$, because $\vartheta^* \leq \vartheta_{\ell}$ for each ℓ , and hence, by Part 2 of Proposition 2.3 $|\log \vartheta_{\ell} - \log \vartheta_{\ell}^*| \leq \epsilon_{\ell} \gamma \kappa \|c\|_{\infty} e^{\gamma(\kappa\|c\|_{\infty} + \bar{c}) + \log a_{\gamma}}$, as well.

I turn my focus now to bounding $E_{R_{\gamma}}^{\otimes N_{\ell}} [|\log \hat{\vartheta}_{\ell} - \log \vartheta_{\ell}|]$. Firstly, observe that by a first-order Taylor expansion of the logarithm function, $|\log \hat{\vartheta}_{\ell} - \log \vartheta_{\ell}| = \frac{|\hat{\vartheta}_{\ell} - \vartheta_{\ell}|}{\lambda \hat{\vartheta}_{\ell} + (1-\lambda) \vartheta_{\ell}}$, where $\lambda \in [0, 1]$. As $(\alpha, \mu) \in \bar{\mathcal{C}}_{\ell}$ and $v \in [-1, 1]$ for each sample and ℓ , it follows that $\hat{\vartheta}_{\ell} \geq e^{-\gamma \kappa \|c\|_{\infty}}$ for each sample and ℓ . Furthermore, $\vartheta_{\ell} \geq \vartheta^*$ for each ℓ by the inequality in (A.12), and $\vartheta^* \geq e^{-(\gamma \bar{c} + \log a_{\gamma})}$ by Lemma C.2. As $\kappa\|c\|_{\infty} \geq \bar{c} \geq \bar{c} + \gamma^{-1} \log a_{\gamma}$, $|\log \hat{\vartheta}_{\ell} - \log \vartheta_{\ell}| \leq e^{\gamma \kappa \|c\|_{\infty}} |\hat{\vartheta}_{\ell} - \vartheta_{\ell}|$ holds for each sample and ℓ , implying $E_{R_{\gamma}}^{\otimes N_{\ell}} [|\log \hat{\vartheta}_{\ell} - \log \vartheta_{\ell}|] \leq e^{\gamma \kappa \|c\|_{\infty}} E_{R_{\gamma}}^{\otimes N_{\ell}} [|\hat{\vartheta}_{\ell} - \vartheta_{\ell}|]$. Now,

$$\begin{aligned} e^{\gamma \kappa \|c\|_{\infty}} E_{R_{\gamma}}^{\otimes N_{\ell}} [|\hat{\vartheta}_{\ell} - \vartheta_{\ell}|] &\leq e^{\gamma \kappa \|c\|_{\infty}} E_{R_{\gamma}}^{\otimes N_{\ell}} \left[\sup_{(\alpha, \mu) \in \bar{\mathcal{C}}_{\ell}} \left| \frac{1}{N_{\ell}} \sum_{j=1}^{N_{\ell}} G_{\ell}(\omega_j, \alpha, \mu) - \int_{\Omega} G_{\ell}(\omega, \alpha, \mu) \right| \right] \\ &\leq e^{\gamma \kappa \|c\|_{\infty}} 2 \mathcal{R}_{N_{\ell}}(G_{\ell}) \leq 2 \gamma \kappa \|c\|_{\infty} \sqrt{\frac{2 \log n_{\ell}}{N_{\ell}}} e^{2 \gamma \kappa \|c\|_{\infty}}, \end{aligned}$$

with the second and third inequalities, holding, by applications of Lemma 2.3.1 in van der Vaart and Wellner (1996) (i.e. the Symmetrization Lemma), and Lemma C.1 and Corollary C.1, respectively. Therefore, $\gamma^{-1} E_{R_{\gamma}}^{\otimes N_{\ell}} [|\log \hat{\vartheta}_{\ell} - \log \vartheta_{\ell}|] \leq 2 \kappa \|c\|_{\infty} \sqrt{\frac{2 \log n_{\ell}}{N_{\ell}}} e^{2 \gamma \kappa \|c\|_{\infty}}$. Finally, combining these bounds from the

triangular inequality, I obtain

$$\begin{aligned}
\gamma^{-1} E_{R_\gamma}^{\otimes N_\ell} \left[\left| \log \hat{\vartheta}_\ell - \log \vartheta^* \right| \right] &\leq 2\kappa \|c\|_\infty \sqrt{\frac{2 \log n_\ell}{N_\ell}} e^{2\gamma\kappa\|c\|_\infty} + 2\epsilon_\ell \gamma \kappa \|c\|_\infty e^{\gamma(\kappa\|c\|_\infty + \bar{c}) + \log a_\gamma} \\
&\leq 2\kappa \|c\|_\infty \sqrt{\frac{2 \log n_\ell}{N_\ell}} e^{2\gamma\kappa\|c\|_\infty} + 2\epsilon_\ell \gamma \kappa \|c\|_\infty e^{\gamma 2\kappa\|c\|_\infty} \\
&\leq \max \left\{ \sqrt{\frac{2 \log n_\ell}{N_\ell}}, \epsilon_\ell \right\} 2\kappa \|c\|_\infty e^{\gamma 2\kappa\|c\|_\infty}.
\end{aligned}$$

Part 3. Starting with the lower bound, observe that

$$\sqrt{N_\ell}(\hat{\vartheta}_\ell - \vartheta^*) = \sqrt{N_\ell}(\hat{\vartheta}_\ell - \vartheta_\ell + \vartheta_\ell - \vartheta^*) \leq \sqrt{N_\ell}(\hat{\vartheta}_\ell - \vartheta_\ell) + \sqrt{N_\ell} \epsilon_\ell \gamma \kappa \|c\|_\infty e^{\gamma \kappa \|c\|_\infty}$$

holds by Part 2 of Proposition 2.3. I can re-write this inequality as

$$\begin{aligned}
\sqrt{N_\ell} \vartheta^* &\geq \sqrt{N_\ell} \hat{\vartheta}_\ell + \sqrt{N_\ell}(\vartheta_\ell - \hat{\vartheta}_\ell) - \sqrt{N_\ell} \epsilon_\ell \gamma \kappa \|c\|_\infty e^{\gamma \kappa \|c\|_\infty} \\
&= \sqrt{N_\ell} \hat{\vartheta}_\ell - \sqrt{N_\ell} \epsilon_\ell \gamma \kappa \|c\|_\infty e^{\kappa \gamma \|c\|_\infty} \\
&\quad + \sqrt{N_\ell} \left(\inf_{(\alpha, \mu) \in \overline{C}_\ell} E_{R_\gamma} [G_\ell(\alpha, \mu, \omega)] - \inf_{(\alpha, \mu) \in \overline{C}_\ell} \frac{1}{N_\ell} \sum_{j=1}^{N_\ell} G_\ell(\alpha, \mu, \omega_j) \right) \\
&\geq \sqrt{N_\ell} \hat{\vartheta}_\ell - \sqrt{N_\ell} \epsilon_\ell \gamma \kappa \|c\|_\infty e^{\kappa \gamma \|c\|_\infty} \\
&\quad + \sqrt{N_\ell} \inf_{(\alpha, \mu) \in \overline{C}_\ell} \left(E_{R_\gamma} [G_\ell(\alpha, \mu, \omega)] - \frac{1}{N_\ell} \sum_{j=1}^{N_\ell} G_\ell(\alpha, \mu, \omega_j) \right), \tag{A.17}
\end{aligned}$$

where the last inequality follows from the property of the infimum. Observe that the term (A.17) can be expressed as $-\sup_{g \in \mathcal{G}_\ell} \frac{1}{\sqrt{N_\ell}} \sum_{j=1}^{N_\ell} (g(\omega_j) - E_{R_\gamma}[g(\omega)])$. Now let $Z_\ell = \sup_{g \in \mathcal{G}_\ell} \frac{1}{\sqrt{N_\ell}} \sum_{j=1}^{N_\ell} (g(\omega_j) - E_{R_\gamma}[g(\omega)])$ for each ℓ , and by Lemma C.4 there exists $\tilde{Z}_\ell = \sup_{g \in \mathcal{G}_\ell} B_\ell^l g$, where B_ℓ^l is a centered Gaussian process indexed by \mathcal{G}_ℓ with covariance function $E_{R_\gamma}[B_\ell^l(g) B_\ell^l(g')] = \int_\Omega g(\omega) g'(\omega) dR_\gamma$, for $g, g' \in \mathcal{G}_\ell$, such that $|Z_\ell - \tilde{Z}_\ell| = O_{R_\gamma}(r_\ell)$ as $\ell \rightarrow \infty$ with $r_\ell = C \left[b K_{N_\ell} N_\ell^{-1/2} + (b\sigma)^{1/2} K_{N_\ell}^{3/4} N_\ell^{-1/4} + (b\sigma^2 K_{N_\ell}^2)^{1/3} N_\ell^{-1/6} \right]$, where $K_{N_\ell} = c v \max \left(\log N_\ell, \log \left(\frac{Ab}{\sigma} \right) \right)$, with $C, c > 0$ are constants *not* depending on N_ℓ , $b = \sigma = e^{\gamma \kappa \|c\|_\infty}$, $v = n_\ell + 2$, and $A = (K(n_\ell + 2))^{\frac{1}{n_\ell + 2}} 16e$ with K a universal constant.

In consequence, for each ℓ we have that $\sqrt{N_\ell} \vartheta^* \geq \sqrt{N_\ell} \hat{\vartheta}_\ell - \sqrt{N_\ell} \epsilon_\ell \gamma \kappa \|c\|_\infty e^{\kappa \gamma \|c\|_\infty} - Z_\ell + \tilde{Z}_\ell - \tilde{Z}_\ell$, holds; hence,

$$\vartheta^* \geq \hat{\vartheta}_\ell - \epsilon_\ell \gamma \kappa \|c\|_\infty e^{\kappa \gamma \|c\|_\infty} + O_{R_\gamma} \left(r_\ell / \sqrt{N_\ell} \right) - \tilde{Z}_\ell / \sqrt{N_\ell} \quad \text{as } \ell \rightarrow \infty.$$

where it is $|\tilde{Z}_\ell - Z_\ell| = O_{R_\gamma}(r_\ell / \sqrt{N_\ell})$ as $\ell \rightarrow \infty$. Finally, because $r_\ell \downarrow 0$ as $\ell \rightarrow \infty$, we must have $|\tilde{Z}_\ell - Z_\ell| = o_{R_\gamma}(1 / \sqrt{N_\ell})$ as $\ell \rightarrow \infty$.

Now I turn my focus to the upper bound. Starting with

$$\sqrt{N_\ell}(\hat{\vartheta}_\ell - \vartheta_\ell) = \sqrt{N_\ell} \left(\inf_{(\alpha, \mu) \in \overline{C}_\ell} \frac{1}{N_\ell} \sum_{j=1}^{N_\ell} G_\ell(\alpha, \mu, \omega_j) - \inf_{(\alpha, \mu) \in \overline{C}_\ell} E_{R_\gamma}[G_\ell(\alpha, \mu, \omega)] \right),$$

add and subtract $E_{R_\gamma}[G_\ell(\alpha, \mu, \omega)]$ under the sum in the right side of the above equality and using the lower bound property of the infimum of the sum, I obtain

$$\begin{aligned} \sqrt{N_\ell}(\hat{\vartheta}_\ell - \vartheta_\ell) &\geq \sqrt{N_\ell} \inf_{(\alpha, \mu) \in \overline{C}_\ell} \frac{1}{N_\ell} \sum_{j=1}^{N_\ell} (G_\ell(\alpha, \mu, \omega_j) - E_{R_\gamma}[G_\ell(\alpha, \mu, \omega)]) \\ &\quad + \sqrt{N_\ell} \inf_{(\alpha, \mu) \in \overline{C}_\ell} E_{R_\gamma}[G_\ell(\alpha, \mu, \omega)] - \sqrt{N_\ell} \inf_{(\alpha, \mu) \in \overline{C}_\ell} E_{R_\gamma}[G_\ell(\alpha, \mu, \omega)] \\ &= \sqrt{N_\ell} \inf_{(\alpha, \mu) \in \overline{C}_\ell} \frac{1}{N_\ell} \sum_{j=1}^{N_\ell} (G_\ell(\alpha, \mu, \omega_j) - E_{R_\gamma}[G_\ell(\alpha, \mu, \omega)]). \end{aligned}$$

Furthermore, $\vartheta^* \leq \vartheta_\ell$ for each ℓ , and hence,

$$\sqrt{N_\ell}(\hat{\vartheta}_\ell - \vartheta^*) \geq \sqrt{N_\ell} \inf_{(\alpha, \mu) \in \overline{C}_\ell} \frac{1}{N_\ell} \sum_{j=1}^{N_\ell} (G_\ell(\alpha, \mu, \omega_j) - E_{R_\gamma}[G_\ell(\alpha, \mu, \omega)]). \quad (\text{A.18})$$

The right side of (A.18) is equivalent to $-\sup_{g \in \mathcal{G}_\ell^-} \frac{1}{\sqrt{N_\ell}} \sum_{j=1}^{N_\ell} (g(\omega_j) - E_{R_\gamma}[g(\omega)])$, where $\mathcal{G}_\ell^- = -\mathcal{G}_\ell$. Now let $Z_\ell = \sup_{g \in \mathcal{G}_\ell^-} \frac{1}{\sqrt{N_\ell}} \sum_{j=1}^{N_\ell} (g(\omega_j) - E_{R_\gamma}[g(\omega)])$ for each ℓ . Because \mathcal{G}_ℓ^- must also be VC-subgraph by Part (iv) of Lemma 2.6.18 in van der Vaart and Wellner (1996), and that it must also be pointwise measurable, and has the same envelope function as that of \mathcal{G}_ℓ , I can apply Lemma C.4 to deduce the existence of $\tilde{Z}_\ell = \sup_{g \in \mathcal{G}_\ell^-} B_\ell^u g$, where B_ℓ^u is a centered Gaussian process indexed by \mathcal{G}_ℓ^- with covariance function $E_{R_\gamma}[B_\ell^u(g) B_\ell^u(g')] = \int_\Omega g(\omega) g'(\omega) dR_\gamma$, for $g, g' \in \mathcal{G}_\ell^-$, such that $|Z_\ell - \tilde{Z}_\ell| = O_{R_\gamma}(r_\ell)$ as $\ell \rightarrow \infty$ with r_ℓ as with the lower bound.

In consequence, $\sqrt{N_\ell}(\hat{\vartheta}_\ell - \vartheta^*) \geq (\tilde{Z}_\ell - Z_\ell) - \tilde{Z}_\ell \implies \vartheta^* \leq |\tilde{Z}_\ell - Z_\ell|/\sqrt{N_\ell} + \tilde{Z}_\ell/\sqrt{N_\ell} + \hat{\vartheta}_\ell$. Now noting that $|\tilde{Z}_\ell - Z_\ell|/\sqrt{N_\ell} = O_{R_\gamma}(r_\ell/\sqrt{N_\ell})$ as $\ell \rightarrow \infty$ and because $r_\ell \downarrow 0$ as $\ell \rightarrow \infty$, this term is in fact $o_{R_\gamma}(1/\sqrt{N_\ell})$ as $\ell \rightarrow \infty$. \blacksquare

B Technical Lemmas for Proposition 2.1

Lemma B.1. *Suppose the conditions of Proposition 2.1, hold. Furthermore, for each $n \in \mathbb{Z}_+$, let $z_{0,n} = \arg \inf \left\{ \int_\Omega e^z dR_\gamma : y \in \mathcal{D}_n \right\}$ where $\mathcal{D}_n = \{y \in \mathcal{D} : \alpha \leq \bar{\alpha}_n\}$, with $\bar{\alpha}_n \nearrow \infty$ as $n \rightarrow \infty$. Then*

$$OL\{\{z_{0,n}\}_{n \geq 1}\} \subset \arg \inf \left\{ \int_\Omega e^z dR_\gamma : z \in \mathcal{D} \right\}.$$

Proof. I will establish the three conditions of Assumption 2.1 in Alvarez-Mena and Hernández-Lerma (2005) hold in this example. Assumption 2.1(a) requires the set of limit points of the sequence $\{z_{0,n}\}_{n \geq 1}$ in

the norm topology of $L_1(R_\gamma)$, to be a subset of \mathcal{D} . We consider two cases: (i) the set of limit points is empty, and (ii) the set of limit points is non-empty. In case (i), Assumption 2.1(a) trivially holds since the empty set is a subset of every set. In case (ii), there exists a subsequence $\{z_{0,n_k}\}_{k \geq 1}$ of $\{z_{0,n}\}_{n \geq 1}$ that converges to a limit, z , and we must show $z \in \mathcal{D}$. Toward that end, first note that $z_{0,n_k} \in \mathcal{D}_{n_k}$ for each k means $z_{0,n_k} = \alpha_{0,n_k} z'_{0,n_k}$ where $\alpha_{0,n_k} \in [0, \bar{\alpha}_{n_k}]$ and $z'_{0,n_k} \in \overline{\text{co}}(\mathcal{V})$ for each k . Now since $z \in L_1(R_\gamma)$, $\exists L > 0$ such that $\alpha_{0,n_k} \leq L$ for all k , and hence, $\{z_{0,n_k}\}_{k \geq 1} \subset \mathcal{D}_{L'}$, where $L' = \min\{n_k : \bar{\alpha}_{n_k} \geq L\}$. Since $\mathcal{D}_{L'}$ is closed in the norm topology of $L_1(R_\gamma)$, we must have $z \in \mathcal{D}_{L'}$, and since $\mathcal{D}_{L'} \subset \mathcal{D}$, it follows that $z \in \mathcal{D}$, as desired.

Assumption 2.1(b) requires every subsequence $\{z_{0,n_k}\}_{k \geq 1}$ that converges to a limit, z , to satisfy

$$\liminf_{k \rightarrow \infty} \int_{\Omega} e^{z_{0,n_k}} dR_\gamma \geq \int_{\Omega} e^z dR_\gamma.$$

As with the proof of Assumption 2.1(a), since $z \in L_1(R_\gamma)$, $\exists L > 0$ such that $\alpha_{0,n_k} \leq L$ for all k . Then for some $\eta > 0$, $\int_{\Omega} e^{z_{0,n_k}} dR_\gamma \leq e^{\eta L}$ for all k , because $|z_{0,n_k}| \leq L$ for each k . Hence, the sequence $\{e^{z_{0,n_k}}\}_{k \geq 1}$ is uniformly integrable. Consequently, $\lim_{k \rightarrow \infty} \int_{\Omega} e^{z_{0,n_k}} dQ = \int_{\Omega} e^z dR_\gamma$, holds, implying the desired result.

Assumption 2.1(c) requires for each $z \in \mathcal{D}$, $\exists N \in \mathbb{N}$ and sequence $\{z_n\}_{n \geq 1}$ with $z_n \in \mathcal{D}_n$ for all $n \geq N$, and such that $z_n \xrightarrow{L_1(R_\gamma)} y$ and $\lim_{n \rightarrow \infty} \int_{\Omega} e^{z_n} dR_\gamma = \int_{\Omega} e^z dR_\gamma$. For $z \in \mathcal{D}$ means $z = \alpha z'$, and that $\exists N \in \mathbb{N}$ such that $\bar{\alpha}_n \geq \alpha$ for all $n \geq N$. Set $z_n = \alpha_n z'$ such that $\alpha_n \leq \bar{\alpha}_n$ and $\lim_{n \rightarrow \infty} \alpha_n = \alpha$. Observe that $z_n \xrightarrow{L_1(Q)} z$, holds, so that now we can repeat the same arguments in the previous paragraph to deduce that the sequence $\{e^{z_n}\}_{n \geq 1}$ is uniformly integrable, to conclude

$$\lim_{n \rightarrow \infty} \int_{\Omega} e^{z_n} dR_\gamma = \int_{\Omega} e^z dR_\gamma.$$

This concludes the proof. ■

Lemma B.2. *Suppose the conditions of Proposition 2.1, hold. Then $\text{OL}\{\{z_{0,n}\}_{n \geq 1}\} \neq \emptyset$.*

Proof. For each n , let $p_{0,n} = e^{z_{0,n}} / \int_{\Omega} e^{z_{0,n}} dR_\gamma$. I will now establish that $\{p_{0,n}\}_{n \geq 1} \subset \mathcal{M}$. The set \mathcal{D}_n is convex, and the objective function $g(y) = \int_{\Omega} e^y dR_\gamma$ is Gâteaux differentiable, then by Theorem 2 on page 178 of Luenberger (1969), $\frac{d}{dt}g(z_0 + t(z - z_0))|_{t=0} \geq 0 \ \forall z \in \mathcal{D}_n$, yielding $\int_{\Omega} (z - z_{0,n}) e^{z_{0,n}} dR_\gamma \geq 0 \ \forall z \in \mathcal{D}_n$. By choosing $z = cz_{0,n}$ first with $c > 1$ and then with $c < 1$ (since \mathcal{D}_n is also a cone), we obtain $\int_{\Omega} z_{0,n} e^{z_{0,n}} dR_\gamma = 0$ and $\int_{\Omega} z e^{z_{0,n}} dR_\gamma \geq 0 \ \forall z \in \mathcal{D}_n$. Since $\alpha_{0,n} \int_{\Omega} e^{z_{0,n}} dR_\gamma > 0$, these conditions are equivalent to

$$\int_{\Omega} \alpha'_{0,n} p_{0,n} dR_\gamma = 0 \quad \text{and} \quad \int_{\Omega} z' p_{0,n} dR_\gamma \geq 0 \ \forall a \in \mathcal{D}_n \tag{B.1}$$

where $z_{0,n} = \alpha_{0,n} z'_{0,n}$. Observe that $p_{0,n}$ is clearly a density function, so the main point to show is that it satisfies the moment inequality constraints. To show that it satisfies them, we use the fact that $v \in \overline{\text{co}}(\mathcal{V}) \ \forall v \in \mathcal{V}$. Now setting $z' = v$ and plugging this choice into the second condition in (B.1), we obtain

$\int_{\Omega} vp_{0,n} dR_{\gamma} \leq 0$, which holds for each $v \in \mathcal{V}$. Therefore, $p_{0,n} \in \mathcal{M}$. As n was arbitrary, $\{p_{0,n}\}_{n \geq 1} \subset \mathcal{M}$ must hold.

Now I will now establish that $\lim_{n \rightarrow \infty} \alpha_{0,n} = \infty$ cannot arise. For suppose that $\lim_{n \rightarrow \infty} \alpha_{0,n} = \infty$, holds, and consider a subsequence $\{z'_{0,n_k}\}_{k \geq 1}$ such that $z'_{0,n_k} \xrightarrow{L_1(R_{\gamma})} z'_{\infty} \in \overline{\text{co}}(\mathcal{V})$. The existence of such a subsequence is because $\{z'_{0,n_k}\}_{k \geq 1} \subset \overline{\text{co}}(\mathcal{V})$ and $\overline{\text{co}}(\mathcal{V})$ is compact in the norm topology of $L_1(R_{\gamma})$ (i.e., $L_1(R_{\gamma})$ is a Fréchet space and \mathcal{V} is precompact in the norm topology of $L_1(R_{\gamma})$ by Lemma B.3). Furthermore, note that $\lim_{n \rightarrow \infty} \int_{\Omega} e^{z_{0,n}} dR_{\gamma}$ exists since $\{\int_{\Omega} e^{z_{0,n}} dR_{\gamma}\}_{n \geq 1}$ is a non-increasing sequence of real numbers that is bounded from below by 1. Now, taking a further subsequence $\{z'_{0,n_{k_{\ell}}}\}_{\ell \geq 1}$ where $z'_{0,n_{k_{\ell}}} \rightarrow z'_{\infty}$ a.s.- R_{γ} ,

$$\begin{aligned} \int_{\Omega} e^{z_{0,n_{k_{\ell}}}} dR_{\gamma} &= \int_{\Omega} e^{z_{0,n_{k_{\ell}}}} \left(1 \left[e^{z_{0,n_{k_{\ell}}}} > \alpha_{0,n_{k_{\ell}}} \right] + 1 \left[e^{z_{0,n_{k_{\ell}}}} \leq \alpha_{0,n_{k_{\ell}}} \right] \right) dR_{\gamma} \\ &\geq \alpha_{0,n_{k_{\ell}}} R_{\gamma} \left(\omega \in \Omega : e^{z_{0,n_{k_{\ell}}}} > \alpha_{0,n_{k_{\ell}}} \right) \\ &= \alpha_{0,n_{k_{\ell}}} R_{\gamma} \left(\omega \in \Omega : z'_{0,n_{k_{\ell}}} > \frac{\log \alpha_{0,n_{k_{\ell}}}}{\alpha_{0,n_{k_{\ell}}}} \right). \end{aligned}$$

Now taking limits as $\ell \rightarrow \infty$ on both sides of these inequalities implies

$$\lim_{\ell \rightarrow \infty} R_{\gamma} \left(\omega \in \Omega : z'_{0,n_{k_{\ell}}} > \frac{\log \alpha_{0,n_{k_{\ell}}}}{\alpha_{0,n_{k_{\ell}}}} \right) = 0,$$

since we are assuming $\lim_{n \rightarrow \infty} \alpha_{0,n} = \infty$, which implies $\lim_{\ell \rightarrow \infty} \alpha_{0,n_{k_{\ell}}} = \infty$. And by the Bounded Convergence Theorem,

$$\begin{aligned} 0 &= \lim_{\ell \rightarrow \infty} R_{\gamma} \left(\omega \in \Omega : z'_{0,n_{k_{\ell}}} > \frac{\log \alpha_{0,n_{k_{\ell}}}}{\alpha_{0,n_{k_{\ell}}}} \right) = \int_{\Omega} \lim_{\ell \rightarrow \infty} 1 \left[\omega \in \Omega : z'_{0,n_{k_{\ell}}} > \frac{\log \alpha_{0,n_{k_{\ell}}}}{\alpha_{0,n_{k_{\ell}}}} \right] dR_{\gamma} \\ &= \int_{\Omega} 1 \left[\omega \in \Omega : y'_{\infty} > 0 \right] dQ = Q \left(\omega \in \Omega : y'_{\infty} > 0 \right). \end{aligned}$$

However, $z'_{\infty} \in \overline{\text{co}}(\mathcal{V})$ means there exists $\{v_{\gamma_i}, i = 1, \dots, m\} \subset \overline{\mathcal{V}} = \mathcal{V}$ and $\lambda_i > 0$ for $i = 1, \dots, m$ with $\sum_{i=1}^m \lambda_i = 1$, such that $z'_{\infty} = \sum_{i=1}^m \lambda_i v_{\gamma_i}$. In consequence,

$$0 = R_{\gamma} \left(\omega \in \Omega : z'_{\infty} > 0 \right) \geq (1 - \max_i \gamma_i) > 0,$$

yielding a contradiction.

Therefore, $\lim_{n \rightarrow \infty} \int_{\Omega} e^{z_{0,n}} dR_{\gamma} > 0$, holds. Note that this limit is finite, since $\int_{\Omega} e^{z_{0,n}} dR_{\gamma} = e^{-m(p_{0,n})} \leq e^{-m(p_{\gamma})} < \infty$ for every n , where p_{γ} is the I -projection onto \mathcal{M} . Hence, there exists a subsequence $\{\alpha_{0,n_k}\}_{k \geq 1}$ such that $\lim_{k \rightarrow \infty} \alpha_{0,n_k} \in \mathbb{R}_{++}$ (i.e., positive real numbers), and denote this limit by α_0 . Now taking a further subsequence, $\{z'_{0,n_{k_{\ell}}}\}_{\ell \geq 1}$, such that $z'_{0,n_{k_{\ell}}} \xrightarrow{L_1(R_{\gamma})} z'_{\infty} \in \overline{\text{co}}(\mathcal{V})$, observe that $z_{0,n_{k_{\ell}}} \xrightarrow{L_1(R_{\gamma})} \alpha_0 z'_{\infty} \in \text{OL} \{\{z_{0,n}\}_{n \geq 1}\}$. This concludes the proof. \blacksquare

Lemma B.3. Suppose Part (i) in Assumption 1 holds. For each $\gamma \in \mathbb{Z}_+$, the set \mathcal{V} is precompact in the norm

topology of $L_1(R_\gamma)$.

Proof. The proof proceeds by the direct method. The set \mathcal{V} can be expressed as the union of 4 sets, In particular, $\mathcal{V} = \bigcup_{i=1}^4 \mathcal{V}_i$, where

$$\begin{aligned}\mathcal{V}_1 &= \{(F_X(x') - 1[x \preceq x']) 1[y \in \mathcal{Y}], x' \in \mathcal{X}\}, \quad \mathcal{V}_2 = -\mathcal{V}_1 \\ \mathcal{V}_3 &= \{(F_Y(y') - 1[y \preceq y']) 1[x \in \mathcal{X}], y' \in \mathcal{Y}\}, \quad \mathcal{V}_4 = -\mathcal{V}_3.\end{aligned}$$

As norm precompactness in $L_1(R_\gamma)$ is preserved under finite unions of such sets, the proof requires that we establish these two properties for each of the aforementioned sets. Now because $\mathcal{V}_2 = -\mathcal{V}_1$ and $\mathcal{V}_4 = -\mathcal{V}_3$, and that \mathcal{V}_1 and \mathcal{V}_3 consist of the same type of elements with only the roles of the two marginals being reversed, it is sufficient to only show that \mathcal{V}_1 is precompact to obtain the desired result. Note that identical arguments would apply to establishing \mathcal{V}_3 being precompact, and the remaining sets are negatives of the previous two, which preserves the two properties.

I now establish that \mathcal{V}_1 is precompact in the norm topology of $L_1(R_\gamma)$. We know that the collection of sets $\{\{x \preceq x'\} : x' \in \mathcal{X}\}$ is a VC-class with index $d_x + 1$ – see, e.g., Example 2.6.1 in van der Vaart and Wellner (1996). This implies that the class of indicator functions $\{1[x \preceq x'] : x' \in \mathcal{X}\}$ is VC-subgraph in $\mathcal{X} \times \mathbb{R}$. By Part (iv) of Lemma 2.6.18 in van der Vaart and Wellner (1996), the collection $\{-1[x \preceq x'] : x' \in \mathcal{X}\}$ is also VC-subgraph in $\mathcal{X} \times \mathbb{R}$. Now since the map

$$-1[x \preceq x'] \mapsto F(x') - 1[x \preceq x']$$

is monotone, Part (viii) of Lemma 2.6.18 in van der Vaart and Wellner (1996) establishes that the collection \mathcal{V}_1 is also VC-subgraph in $\mathcal{X} \times \mathbb{R}$. Consequently, for every $\epsilon > 0$, Theorem 2.6.7 in van der Vaart and Wellner (1996) establishes the covering number of \mathcal{V}_1 in $L_1(R_\gamma)$ is finite, and hence, \mathcal{V}_1 is appropriately precompact. \blacksquare

Lemma B.4. *Suppose Part (i) in Assumption 1 holds. Then $\bar{\mathcal{V}} = \{g_\ell, -g_\ell : \ell \in \mathcal{X} \cup \mathcal{Y}\}$, where closure is in the norm topology of $L_1(R_\gamma)$.*

Proof. The proof proceeds by the direct method and uses the same representation of \mathcal{V} as in Lemma B.3. Starting with $\mathcal{V} = \bigcup_{i=1}^4 \mathcal{V}_i$, where

$$\begin{aligned}\mathcal{V}_1 &= \{(F_X(x') - 1[x \preceq x']) 1[y \in \mathcal{Y}], x' \in \mathcal{X}\}, \quad \mathcal{V}_2 = -\mathcal{V}_1 \\ \mathcal{V}_3 &= \{(F_Y(y') - 1[y \preceq y']) 1[x \in \mathcal{X}], y' \in \mathcal{Y}\}, \quad \mathcal{V}_4 = -\mathcal{V}_3,\end{aligned}$$

as norm closedness in $L_1(R_\gamma)$ is preserved under finite unions of such sets, the proof requires that we establish these two properties for each of the aforementioned sets. Now because $\mathcal{V}_2 = -\mathcal{V}_1$ and $\mathcal{V}_4 = -\mathcal{V}_3$, and that \mathcal{V}_1 and \mathcal{V}_3 consist of the same type of elements with only the roles of the two marginals being reversed, it is sufficient to only show that \mathcal{V}_1 is closed to obtain the desired result. Note that identical arguments would apply to establishing \mathcal{V}_3 is closed, and the remaining sets are negatives of the previous two, which preserves the two properties.

I now establish that the limit points of \mathcal{V}_1 , in the norm topology of $L_1(R_\gamma)$, have the same form of its elements. Consider an arbitrary sequence $\{v_n\}_{n \geq 1} \subset \mathcal{G}_1$, where

$$v_n(x, y) = (F_X(x'_n) - 1[x \preceq x'_n]) 1[y \in \mathcal{Y}] \quad n = 1, 2, \dots,$$

such that $v_n \xrightarrow{L_1(R_\gamma)} v$. To prove the desired result, we need to establish that

$$v(x, y) = (F_X(x') - 1[x \preceq x']) 1[y \in \mathcal{Y}]$$

for some $x' \in \mathcal{X}$. As norm-convergence in $L_1(R_\gamma)$ implies convergence in R_γ -measure, there exists a non-random increasing sequence of integers n_1, n_2, \dots , such that $\{v_{n_k}\}_{k \geq 1}$ converges to v a.s. $- R_\gamma$. That is,

$$v(x, y) = \lim_{k \rightarrow +\infty} v_{n_k}(x, y) \quad \text{a.s. } - R_\gamma. \quad (\text{B.2})$$

The limit (B.2) implies $\{x'_{n_k}\}_{k \geq 1} \subset \mathcal{X}$ holds. Since $\mathcal{X} \subset \mathbb{R}^{d_x}$ is compact, there exists a subsequence $\{x'_{n_{k_s}}\}_{s \geq 1}$ such that $\lim_{s \rightarrow +\infty} x'_{n_{k_s}} = x'^* \in \mathcal{X}$. If F_X is continuous at x'^* , then combining this conclusion with the limit (B.2) yields

$$v(x, y) = \lim_{s \rightarrow +\infty} v_{n_{k_s}}(x, y) = (F_X(x'^*) - 1[x \preceq x'^*]) 1[y \in \mathcal{Y}] \quad \text{a.s. } - R_\gamma,$$

as every subsequence of $\{v_{n_k}\}_{k \geq 1}$ converges to v a.s. $- R_\gamma$. Therefore, v has the appropriate form.

Now, we focus on the case where F_X is discontinuous at x'^* . The CDF F_X is monotonic, so it can have at most a countable number of points of discontinuity. With F_X discontinuous at x'^* , by following steps identical those above for the continuous case, we have

$$v(x, y) = \lim_{s \rightarrow +\infty} v_{n_{k_s}}(x, y) = (F_X(x'^*-) - 1[x \preceq x'^*]) 1[y \in \mathcal{Y}] \quad \text{a.s. } - R_\gamma,$$

where $\lim_{s \rightarrow +\infty} F_X(x'_{n_{k_s}}) = F_X(x'^* -)$ and $x'_{i, n_{k_s}} \uparrow x'^*_i$ for some component i at which F_X is discontinuous. \blacksquare

Lemma B.5. *Suppose Part (i) of Assumption 1 holds. The following statements hold: $\text{ex}(\overline{\text{co}}(\mathcal{V})) \neq \emptyset$, $\overline{\text{co}}(\text{ex}(\overline{\text{co}}(\mathcal{V}))) = \overline{\text{co}}(\mathcal{V})$, and $\text{ex}(\overline{\text{co}}(\mathcal{V})) \subset \mathcal{V}$.*

Proof. The proof proceeds by the direct method. By Lemmas B.3 and B.4, the set \mathcal{V} is compact in the $L_1(R_\gamma)$ -norm (as it is complete and totally bounded). Now, since $L_1(R_\gamma)$ is a Banach space, it is therefore a Fréchet space. This fact allows us to apply part (c) of Theorem 3.20 in Rudin (1991) to the set \mathcal{V} to deduce that $\overline{\text{co}}(\mathcal{V})$ is also compact in the same norm. Whence, I can apply the Krein-Milman Theorem (e.g., Theorem 3.23 in Rudin, 1991) to the set $\overline{\text{co}}(\mathcal{V})$ to deduce that its set of extreme points, $\text{ex}(\overline{\text{co}}(\mathcal{V}))$, is nonempty, and that $\overline{\text{co}}(\text{ex}(\overline{\text{co}}(\mathcal{V}))) = \overline{\text{co}}(\mathcal{V})$. Next, I can apply Milman's Theorem (e.g., Theorem 3.25 in Rudin, 1991) to establish $\text{ex}(\overline{\text{co}}(\mathcal{V})) \subset \mathcal{V}$. \blacksquare

C Technical Results for Theorem 1

Consider a class of functions \mathcal{F} of real-valued measurable functions defined on the probability space $(\mathcal{Z}, \mathcal{A}, P)$. The sample is denoted by Z_1, \dots, Z_N . Furthermore, let $\varepsilon_1, \dots, \varepsilon_N$ denotes a random sample of Rademacher variables that is independent Z_1, \dots, Z_N . The Rademacher complexity of the function class \mathcal{F} is defined as

$$\mathcal{R}_N(\mathcal{F}) = E_{(P_Z \otimes P_\varepsilon)^{\otimes N}} \left[\sup_{f \in \mathcal{F}} \left| \frac{1}{N} \sum_{i=1}^N \varepsilon_i f(Z_i) \right| \right]. \quad (\text{C.1})$$

Proposition C.1. *Consider a sequence $\{\mathcal{F}_\ell\}_{\ell \geq 1}$ where \mathcal{F}_ℓ is a class of real-valued measurable functions defined on the probability space $(\mathcal{Z}, \mathcal{A}, P)$, such that $\|f\|_{L_\infty(P)} \leq b \forall f \in \mathcal{F}_\ell$ for each ℓ . Furthermore, let $\{N_\ell\}_{\ell \geq 1}$ be a sequence of sample sizes such that $\lim_{\ell \rightarrow \infty} N_\ell = \infty$. If $\mathcal{R}_{N_\ell}(\mathcal{F}_\ell) = o(1)$ as $\ell \rightarrow \infty$, then $\sup_{f \in \mathcal{F}_\ell} \left| \frac{1}{N_\ell} \sum_{i=1}^{N_\ell} f(Z_i) - E(f(Z_i)) \right| \xrightarrow{a.s.} 0$.*

Proof. Fix $\epsilon > 0$ and define

$$A_{N_\ell}(\epsilon) := \left\{ \sup_{f \in \mathcal{F}_\ell} \left| \frac{1}{N_\ell} \sum_{i=1}^{N_\ell} f(Z_i) - Ef(Z_1) \right| > \epsilon \right\}, \quad B_m(\epsilon) := \bigcup_{N_\ell \geq m} A_{N_\ell}(\epsilon).$$

It suffices to show that $P(B_m(\epsilon)) \rightarrow 0$ as $m \rightarrow \infty$, since this implies

$$\sup_{f \in \mathcal{F}_\ell} \left| \frac{1}{N_\ell} \sum_{i=1}^{N_\ell} f(Z_i) - Ef(Z_1) \right| \rightarrow 0 \quad \text{almost surely.}$$

By assumption, $\mathcal{R}_{N_\ell}(\mathcal{F}_\ell) = o(1)$. Hence there exists ℓ_0 such that for all $\ell \geq \ell_0$,

$$2\mathcal{R}_{N_\ell}(\mathcal{F}_\ell) < \epsilon/2.$$

Let $\delta := \epsilon/2$. Then, for all $\ell \geq \ell_0$,

$$A_{N_\ell}(\epsilon) \subseteq \left\{ \sup_{f \in \mathcal{F}_\ell} \left| \frac{1}{N_\ell} \sum_{i=1}^{N_\ell} f(Z_i) - Ef(Z_1) \right| > 2\mathcal{R}_{N_\ell}(\mathcal{F}_\ell) + \delta \right\}.$$

By Theorem 4.10 of Wainwright (2019), since $\|f\|_{L_\infty(P)} \leq b$ for all $f \in \mathcal{F}_\ell$,

$$P \left(\sup_{f \in \mathcal{F}_\ell} \left| \frac{1}{N_\ell} \sum_{i=1}^{N_\ell} f(Z_i) - Ef(Z_1) \right| > 2\mathcal{R}_{N_\ell}(\mathcal{F}_\ell) + \delta \right) \leq \exp \left(-\frac{N_\ell \delta^2}{2b^2} \right).$$

Therefore, for all m sufficiently large,

$$P(B_m(\epsilon)) \leq \sum_{N_\ell \geq m} \exp \left(-\frac{N_\ell \delta^2}{2b^2} \right).$$

Since $N_\ell \rightarrow \infty$, the right-hand side is the tail of a convergent geometric series and hence converges to

zero as $m \rightarrow \infty$. This proves the claim. \blacksquare

Lemma C.1 (Expected Rademacher complexity of exp of convex mixtures). *Suppose Assumption 1 holds. Then for each sample size N_ℓ , the collection \mathcal{G}_ℓ in (3.1) satisfies*

$$\mathcal{R}_{N_\ell}(\mathcal{G}_\ell) \leq \gamma \kappa \|c\|_\infty e^{\gamma \kappa \|c\|_\infty} \mathcal{R}_{N_\ell}\left(\text{conv}\{v_1, \dots, v_{n_\ell}\}\right) = \gamma \kappa \|c\|_\infty e^{\gamma \kappa \|c\|_\infty} \mathcal{R}_{N_\ell}(\{v_1, \dots, v_{n_\ell}\}).$$

Proof. The proof proceeds by the direct method. To ease exposition, we introduce the following notation: $b = \gamma \kappa \|c\|_\infty$ and $\Delta_\ell = \{\mu \in \mathbb{R}_+^n : \sum_{i=1}^{n_\ell} \mu_i = 1\}$. Furthermore, fix a sample $S = (\omega_1, \dots, \omega_{N_\ell})$ and write $s_\mu := \sum_{i=1}^{n_\ell} \mu_i v_i$. Since $v_i \in [-1, 1]$ for each i and $\mu \in \Delta_\ell$, $s_\mu \in [-1, 1]$, hence for $\alpha \in [0, b]$, $\alpha s_\mu \in [-b, b]$.

(i) *Scaling.* Let $\mathbb{S} := \{s_\mu : \mu \in \Delta_\ell\}$ and $\mathcal{A} := \{\alpha s : \alpha \in [0, b], s \in \mathbb{S}\}$. For this fixed S ,

$$\mathfrak{R}_S(\mathcal{A}) = \mathbb{E}_{P_\epsilon^{\otimes N_\ell}} \left[\sup_{\alpha \in [0, b], s \in \mathbb{S}} \frac{\alpha}{N_\ell} \sum_{j=1}^{N_\ell} \epsilon_j s(\omega_j) \right] = b \mathfrak{R}_S(\mathbb{S}).$$

(ii) *Contraction.* Define $\phi(u) := e^u - 1$. On $[-b, b]$, ϕ is e^b -Lipschitz and $\phi(0) = 0$. Talagrand's contraction inequality (Ledoux and Talagrand, 1991, Theorem 4.12) yields

$$\mathfrak{R}_S(\{\phi \circ h : h \in \mathcal{A}\}) \leq e^b \mathfrak{R}_S(\mathcal{A}).$$

Since adding a constant does not change \mathfrak{R}_S , we have

$$\mathfrak{R}_S(\mathcal{G}_\ell) = \mathfrak{R}_S(\{e^{\alpha s}\}) = \mathfrak{R}_S(\{e^{\alpha s} - 1\}) \leq e^b \mathfrak{R}_S(\mathcal{A}) = b e^b \mathfrak{R}_S(\mathbb{S}).$$

(iii) *Convex hull invariance.* For fixed S ,

$$\sup_{s \in \text{conv}\{v_i\}} \sum_{j=1}^{N_\ell} \epsilon_j s(\omega_j) = \max_{1 \leq i \leq n_\ell} \sum_{j=1}^{N_\ell} \epsilon_j v_i(\omega_j),$$

so $\mathfrak{R}_S(\mathbb{S}) = \mathfrak{R}_S(\{v_1, \dots, v_{n_\ell}\})$. Combining (i)-(iii) gives, pointwise in S ,

$$\mathfrak{R}_S(\mathcal{G}_\ell) \leq b e^b \mathfrak{R}_S(\{v_1, \dots, v_{n_\ell}\}).$$

Taking expectation over $S \sim P^{\otimes N_\ell}$ yields the desired inequality for $\mathcal{R}_{N_\ell}(\mathcal{G}_\ell)$. \blacksquare

Corollary C.1 (Finite dictionary). *For any $N_\ell \geq 1$,*

$$\mathcal{R}_{N_\ell}(\{v_1, \dots, v_{n_\ell}\}) \leq \sqrt{\frac{2 \log n_\ell}{N_\ell}}.$$

Proof. The functions $v_i : \Omega \rightarrow [-1, 1]$, are bounded. Therefore, Massart's finite-class lemma (Massart, 2000) applied to the class $\{v_1, \dots, v_{n_\ell}\}$ and then averaged over $S = (\omega_1, \dots, \omega_{N_\ell})$, yields the result. \blacksquare

Lemma C.2. Suppose that Assumption 1 holds, and let $\bar{c} = \int_{\Omega} c(\omega) d(P_X \otimes P_Y)$. Then $\vartheta^* > e^{-(\gamma\bar{c} + \log a_{\gamma})}$.

Proof. The proof proceeds by the direct method. Under Assumption 1, I can apply Theorem 2.2 in Csiszár (1975) in our context with respect to the product measure $P_X \otimes P_Y$ to obtain $H(P_{\gamma}|R_{\gamma}) \leq H(P_X \otimes P_Y|R_{\gamma}) - H(P_X \otimes P_Y|P_{\gamma})$. Consequently, $H(P_{\gamma}|R_{\gamma}) < H(P_X \otimes P_Y|R_{\gamma})$, holds, since $H(P_X \otimes P_Y|P_{\gamma}) > 0$.

Noting that $H(P_X \otimes P_Y|R_{\gamma}) = \int_{\Omega} \log \left[\frac{dR_{\gamma}}{d(P_X \otimes P_Y)}(\omega) \right]^{-1} d(P_X \otimes P_Y) = \gamma\bar{c} + \log a_{\gamma}$, the above inequality becomes $H(P_{\gamma}|R_{\gamma}) < \gamma\bar{c} + \log a_{\gamma}$. Finally, I can combine this result with the fact that $H(P_{\gamma}|R_{\gamma}) = -\log(\vartheta^*)$ holds to deduce the desired result: $\vartheta^* > e^{-(\gamma\bar{c} + \log a_{\gamma})}$. \blacksquare

Lemma C.3 (VC-Dimension). Suppose that Assumption 1 holds. The collection \mathcal{G}_{ℓ} in (3.1) is a Vapnik-Chervonenkis (VC) class of functions with VC dimension V_{ℓ} bounded by $n_{\ell} + 2$, for each ℓ .

Proof. The proof proceeds by the direct method. For each $n_{\ell} \in \mathbb{Z}_+$, Lemma 2.6.15 in van der Vaart and Wellner (1996) establishes $\text{span}(v_1, \dots, v_{n_{\ell}})$ is a VC class of functions with VC-dimension bounded above by $n_{\ell} + 2$. Now because

$$\left\{ \alpha \sum_{i=1}^{n_{\ell}} \mu_i v_i : (\alpha, \mu) \in \mathcal{C}_{\ell} \right\} \subset \text{span}(v_1, \dots, v_{n_{\ell}}),$$

the class $\{\alpha \sum_{i=1}^{n_{\ell}} \mu_i v_i : (\alpha, \mu) \in \mathcal{C}_{\ell}\}$ must also be VC whose VC-dimension is also bounded from above by $n_{\ell} + 2$. I can apply Part (viii) of Lemma 2.6.18 in van der Vaart and Wellner (1996) to this last class with exponential function, which is strictly monotonic, to determine that the resulting class is also VC. Since the exponential function is strictly monotonic, and hence, one-to-one, the VC-dimension of this transformed class has the same VC-dimension. Finally, because $\mathcal{G}_{\ell} \subset \{\alpha \sum_{i=1}^{n_{\ell}} \mu_i v_i : (\alpha, \mu) \in \mathcal{C}_{\ell}\}$, it must be that $V_{\ell} \leq n_{\ell} + 2$. This concludes the proof. \blacksquare

Lemma C.4 (Gaussian Approximation). Let \mathcal{G}_{ℓ} be defined as in (3.1), and let $\omega_1, \dots, \omega_{N_{\ell}}$ be i.i.d. with law R_{γ} , and define the empirical process

$$\mathbb{G}_{N_{\ell}} g := \frac{1}{\sqrt{N_{\ell}}} \sum_{i=1}^{N_{\ell}} (g(\omega_i) - E_{R_{\gamma}}[g(\omega)]), \quad g \in \mathcal{G}_{\ell},$$

Consider the supremum statistic $Z_{\ell} = \sup_{g \in \mathcal{G}_{\ell}} \mathbb{G}_{N_{\ell}} g$. Let B_{ℓ} be a centered tight Gaussian process indexed by \mathcal{G}_{ℓ} with covariance function $E_{R_{\gamma}}[B_{\ell}(g) B_{\ell}(g')] = \int_{\Omega} g(\omega) g'(\omega) dR_{\gamma}$, for $g, g' \in \mathcal{G}_{\ell}$. Then there exists a random variable $\tilde{Z}_{N_{\ell}} \stackrel{d}{=} \sup_{g \in \mathcal{G}_{\ell}} B_{\ell}(g)$ such that $|Z_{\ell} - \tilde{Z}_{\ell}| = O_{R_{\gamma}}(r_{\ell})$, where

$$r_{\ell} := C \left[b K_{N_{\ell}} N_{\ell}^{-1/2} + (b\sigma)^{1/2} K_{N_{\ell}}^{3/4} N_{\ell}^{-1/4} + (b\sigma^2 K_{N_{\ell}}^2)^{1/3} N_{\ell}^{-1/6} \right],$$

$$K_{N_{\ell}} := c v \max \left(\log N_{\ell}, \log \left(\frac{Ab}{\sigma} \right) \right),$$

with $C, c > 0$ are constants not depending on N_{ℓ} , $b = \sigma = e^{\gamma\kappa\|c\|_{\infty}}$, $v = n_{\ell} + 2$, and $A = (K(n_{\ell} + 2))^{\frac{1}{n_{\ell} + 2}} 16e$ with K a universal constant.

Proof. The result follows by applying Corollary 2.2 in Chernozhukov et al. (2014b) in our setup. Towards that end, I shall verify the conditions for its application.

The class of functions \mathcal{G}_ℓ is trivially pointwise measurable, as it consists of functions that are indicator functions on upper sets plus a constant. By Lemma C.3, this class is also VC with VC-dimension bounded from above by $n_\ell + 2$, and it has the constant function, $\omega \mapsto e^{\kappa\gamma\|c\|_\infty}$, as its envelope function. This class of functions is also compatible with Definition 2.1 in Chernozhukov et al. (2014b): by Theorem 2.6.7 in van der Vaart and Wellner (1996) and Lemma C.3, I can specify v and A in their definition as $v = n_\ell + 2$ and $A = (K(n_\ell + 2))^{\frac{1}{n_\ell+2}} 16e$ with K a universal constant. Since the envelope function is a constant function, I can also specify b and σ in their corollary as $b = \sigma = e^{\kappa\gamma\|c\|_\infty}$. Finally, I can set $q = \infty$ in their corollary to obtain the desired result because \mathcal{G}_ℓ is uniformly bounded. \blacksquare

D Partition of \mathcal{V} Independent of R_γ

Lemma D.1. *Recall that \mathcal{V} is a uniformly bounded VC-subgraph class of measurable functions on the measurable space $(\Omega, \mathcal{B}(\mathcal{X} \times \mathcal{Y}))$, with envelope function, $F(\omega) = 1$ for all $\omega \in \Omega$. There exist constants $C, p > 0$, depending only on the VC characteristics of \mathcal{V} , such that the following holds.*

For every $\epsilon_\ell \in (0, 1)$ and every probability measure Q on $(\Omega, \mathcal{B}(\mathcal{X} \times \mathcal{Y}))$ there exists a finite subset $\{v_{1,\ell}, \dots, v_{n_\ell,\ell}\} \subset \mathcal{V}$ with

$$\sup_{v \in \mathcal{V}} \min_{1 \leq i \leq n_\ell} \|v - v_{i,\ell}\|_{L_1(Q)} \leq \epsilon_\ell \quad (\text{D.1})$$

and

$$n_\ell \leq C \epsilon_\ell^{-p}. \quad (\text{D.2})$$

Moreover, defining

$$E_{i,\ell} := \left\{ v \in \mathcal{V} : \|v - v_{i,\ell}\|_{L_1(Q)} = \min_{1 \leq j \leq n_\ell} \|v - v_{j,\ell}\|_{L_1(Q)} \right\}, \quad i = 1, \dots, n_\ell,$$

with any deterministic tie-breaking rule when the minimum is attained at multiple indices, yields a partition $\{E_{i,\ell}\}_{i=1}^{n_\ell}$ of \mathcal{V} such that

$$\sup_{v \in E_{i,\ell}} \|v - v_{i,\ell}\|_{L_1(Q)} \leq \epsilon_\ell, \quad i = 1, \dots, n_\ell. \quad (\text{D.3})$$

In particular, these conclusions hold uniformly in the choice of the reference measure $Q = R_\gamma$ for any $\gamma > 0$.

Proof. Since \mathcal{V} is a uniformly bounded VC–subgraph class with envelope F satisfying $\|F\|_\infty \leq 1$, standard entropy bounds for VC classes (see, e.g., Theorem 2.6.7 in van der Vaart and Wellner, 1996) imply that there exist constants $A, B > 0$ and an integer $v \geq 1$ (the VC index) such that, for all probability measures Q , all $\delta \in (0, 1)$ and all $r \in (0, 1]$,

$$\log N(\delta r, \mathcal{V}, L_2(Q)) \leq A v \log\left(\frac{B r}{\delta}\right). \quad (\text{D.4})$$

In particular, taking $r = \|F\|_{L_2(Q)} \leq 1$, I obtain

$$\log N(\delta, \mathcal{V}, L_2(Q)) \leq C_1 \log\left(\frac{C_2}{\delta}\right), \quad (\text{D.5})$$

for some constants $C_1, C_2 > 0$ depending only on A, B, v , but not on Q .

By uniform boundedness, $\|g\|_{L_1(Q)} \leq \|g\|_{L_2(Q)}$ for all $g \in \mathcal{V} - \mathcal{V}$ and all Q . Hence

$$N(\delta, \mathcal{V}, L_1(Q)) \leq N(\delta, \mathcal{V}, L_2(Q)), \quad (\text{D.6})$$

and (D.5) yields a corresponding $L_1(Q)$ entropy bound. In particular, for each $\epsilon_\ell \in (0, 1)$ and each Q , I may choose a minimal $L_1(Q)$ -covering of \mathcal{V} at radius $\epsilon_\ell/2$, that is, a finite set $\{v_{1,\ell}, \dots, v_{n_\ell,\ell}\} \subset \mathcal{V}$ such that

$$\mathcal{V} \subset \bigcup_{i=1}^{n_\ell} B_{L_1(Q)}(v_{i,\ell}, \epsilon_\ell/2),$$

and $n_\ell = N(\epsilon_\ell/2, \mathcal{V}, L_1(Q))$. By (D.6) and (D.5), there exist constants $C, p > 0$ depending only on the VC characteristics of \mathcal{V} such that

$$n_\ell \leq C \epsilon_\ell^{-p},$$

which is (D.2).

By the covering property, for every $v \in \mathcal{V}$ there exists $i \in \{1, \dots, n_\ell\}$ such that $\|v - v_{i,\ell}\|_{L_1(Q)} \leq \epsilon_\ell/2$. In particular,

$$\sup_{v \in \mathcal{V}} \min_{1 \leq i \leq n_\ell} \|v - v_{i,\ell}\|_{L_1(Q)} \leq \epsilon_\ell/2,$$

and hence (D.1) holds with ϵ_ℓ and a possibly enlarged constant C (absorbing the factor 2). Defining the sets $E_{i,\ell}$ by nearest-center assignment,

$$E_{i,\ell} := \left\{ v \in \mathcal{V} : \|v - v_{i,\ell}\|_{L_1(Q)} = \min_{1 \leq j \leq n_\ell} \|v - v_{j,\ell}\|_{L_1(Q)} \right\},$$

with deterministic tie-breaking, yields a partition $\{E_{i,\ell}\}_{i=1}^{n_\ell}$ of \mathcal{V} , and the covering property implies

$$\sup_{v \in E_{i,\ell}} \|v - v_{i,\ell}\|_{L_1(Q)} \leq \epsilon_\ell, \quad i = 1, \dots, n_\ell,$$

which is (D.3). The entropy bound (D.4) is uniform over all probability measures Q , so the constants C, p in (D.2) may be chosen independent of Q . In particular, the conclusions hold uniformly for $Q = R_\gamma$ with $\gamma > 0$. This proves the lemma. \blacksquare

E An Implementation and Numerical Illustration

This section presents a geometric implementation of the sieve M-estimation procedure, and illustrates its performance numerically relative to the empirical Sinkhorn divergence.

E.1 Implementation

The approximation scheme of Tabri (2025) requires a specification of $E_{i,\ell}$ for $i = 1, \dots, n_\ell$ satisfying the accuracy (2.12), which is a partition of the set of moment functions $\bar{\mathcal{V}}$. His scheme also allows the practitioner to select any set of $v_i \in E_{i,\ell}$ for $i = 1, \dots, n_\ell$, as the results of Proposition 2.3 are uniform in their choice. This section puts forward a specification of said partition and an approach to selecting the $v_i \in E_{i,N}$ based on the fact that $B := \{v \in \mathcal{V} : \int_{\Omega} v dP_\gamma = 0\}$ coincides with \mathcal{V} .

To elucidate, first note that $\mathcal{V} = \bigcup_{i=1}^4 \mathcal{V}_i$, where

$$\begin{aligned}\mathcal{V}_1 &= \{(F_X(x') - 1[x \preceq x']) 1[y \in \mathcal{Y}], x' \in \mathcal{X}\}, & \mathcal{V}_2 &= \{(1[x \preceq x'] - F_X(x')) 1[y \in \mathcal{Y}], x' \in \mathcal{X}\} \\ \mathcal{V}_3 &= \{(F_Y(y') - 1[y \preceq y']) 1[x \in \mathcal{X}], y' \in \mathcal{Y}\}, & \mathcal{V}_4 &= \{(1[y \preceq y'] - F_Y(y')) 1[x \in \mathcal{X}], y' \in \mathcal{Y}\}.\end{aligned}$$

In this notation, observe that $\mathcal{V}_2 = -\mathcal{V}_1$ and $\mathcal{V}_4 = -\mathcal{V}_3$. Now, given ϵ_ℓ , I can always find finite partitions of \mathcal{X} and \mathcal{Y} , given by $\{\mathcal{X}_j, j \leq n_x\}$ and $\{\mathcal{Y}_j, j \leq n_y\}$, respectively, such that

$$|F_X(x) - F_X(x')| \leq \epsilon_\ell/8 \quad \text{and} \quad |R_\gamma(x, \cdot) - R_{\gamma,X}(x')| \leq \epsilon_\ell/8 \quad \forall x, x' \in \mathcal{X}_j \quad \text{and} \quad (\text{E.1})$$

$$|F_Y(y) - F_Y(y')| \leq \epsilon_\ell/8 \quad \text{and} \quad |R_\gamma(Y(y), \cdot) - R_{\gamma,Y}(y')| \leq \epsilon_\ell/8 \quad \forall y, y' \in \mathcal{Y}_i, \quad (\text{E.2})$$

hold, for $j = 1, \dots, n_x$ and $i = 1, \dots, n_y$. The notation $R_{\gamma,X}(\cdot)$ and $R_{\gamma,Y}(\cdot)$ denote the distribution functions of the X -margin and Y -margin, respectively, of the joint distribution R_γ .

The partition has two parts, once for \mathcal{V}_1 and \mathcal{V}_2 , and one for \mathcal{V}_3 and \mathcal{V}_4 . Consider the following sets $E_{X,1,j} = \{(F_X(x') - 1[x \preceq x']) 1[y \in \mathcal{Y}], x' \in \mathcal{X}_j\}$, $E_{X,2,j} = \{(1[x \preceq x'] - F_X(x')) 1[y \in \mathcal{Y}], x' \in \mathcal{X}_j\}$, $E_{Y,1,i} = \{(F_Y(y') - 1[y \preceq y']) 1[x \in \mathcal{X}], y' \in \mathcal{Y}_i\}$, $E_{Y,2,i} = \{(1[y \preceq y'] - F_Y(y')) 1[x \in \mathcal{X}], y' \in \mathcal{Y}_i\}$, for $j = 1, \dots, n_x$ and $i = 1, \dots, n_y$. Then, I have partitions of the disjoint sets that comprise \mathcal{V} and satisfy the approximation's construction: $\mathcal{V}_1 = \bigcup_{j=1}^{n_x} E_{X,1,j}$ and $\mathcal{V}_2 = \bigcup_{j=1}^{n_x} E_{X,2,j}$, $\mathcal{V}_2 = \bigcup_{j=1}^{n_x} E_{X,2,j}$, $\mathcal{V}_3 = \bigcup_{i=1}^{n_y} E_{Y,1,i}$, and $\mathcal{V}_4 = \bigcup_{i=1}^{n_y} E_{Y,2,i}$. In consequence, $n_\ell = 2n_x + 2n_y$.

Now I shall make use of the fact that $B = \mathcal{V}$, holds, in selecting the functions v_i to implement a moment *equality* using two such inequalities. This can be done as follows: for each selected $v \in E_{X,1,j}$ and $v' \in E_{Y,1,i}$, select $-v \in E_{X,2,j}$ and $-v' \in E_{Y,2,i}$, and do this for $j = 1, \dots, n_x$ and $i = 1, \dots, n_y$. In consequence, I have reduced the number of choice variables in the approximating Fenchel dual programs by re-writing it to recognize moment equality constraints. In this case and using the result of Proposition 2.2, the problem (3.2) becomes

$$\inf \left\{ \int_{\Omega} G_\ell(\omega, \tau) dR_\gamma(\omega) : \tau \in [-1, 1]^{n_x+n_y}, \text{ and } \sum_{i=1}^{n_x+n_y} \tau_i \in [-1, 1] \right\} \quad \text{where} \quad (\text{E.3})$$

$$G_\ell(\omega, \tau) = \exp \left\{ \gamma \|c\|_\infty \sum_{i=1}^{n_x+n_y} \tau_i v_i(\omega) \right\}, \quad (\text{E.4})$$

reducing the number of choice variables from $2n_x + 2n_y$ to $n_x + n_y$ and constraining them to the box

$[-1, 1]^{n_x+n_y}$. With this formulation $n_\ell = n_x + n_y$, and function class

$$\left\{ G_\ell(\cdot, \tau), \tau \in [-1, 1]^{n_\ell} \text{ and } \sum_{i=1}^{n_\ell} \tau_i \in [-1, 1] \right\}$$

is still VC and uniformly bounded by $e^{\gamma\kappa\|c\|_\infty}$ for each ℓ , so that the results of Theorem 1 apply to the SAA version of (E.3).

In practice, we will consider large values of γ (e.g., $\gamma \geq 100$), which can create numerical challenges in solving the SAA version of (E.3),

$$\inf \left\{ \frac{1}{N_\ell} \sum_{j=1}^{N_\ell} G_\ell(\omega_j, \tau) : \tau \in [-1, 1]^{n_\ell} \text{ and } \sum_{i=1}^{n_\ell} \tau_i \in [-1, 1] \right\}, \quad (\text{E.5})$$

for a random sample of size N_ℓ from R_γ . A practical approach to addressing this challenge is to rescale the objective function in (E.5) by dividing it by $e^{\gamma\kappa\|c\|_\infty}$, and then to apply logarithms to quash the impact of large γ in the objective function (E.4), yielding

$$\inf \left\{ -\gamma\kappa\|c\|_\infty + \log \frac{1}{N_\ell} \sum_{j=1}^{N_\ell} G_\ell(\omega_j, \tau) : \tau \in [-1, 1]^{n_\ell} \text{ and } \sum_{i=1}^{n_\ell} \tau_i \in [-1, 1] \right\}, \quad (\text{E.6})$$

Because the logarithm is strictly positive monotonic transformation, the solutions of the problems (E.6) and (E.5) coincide. Furthermore, rescaling the objective function in this way implies

$$\left\{ e^{-\gamma\kappa\|c\|_\infty} G_\ell(\cdot, \tau), \tau \in [-1, 1]^{n_\ell} \text{ and } \sum_{i=1}^{n_\ell} \tau_i \in [-1, 1] \right\}$$

is still VC and now uniformly bounded by 1 for each ℓ . Hence, to calculate $\hat{\vartheta}_\ell$, I must first add $\gamma\kappa\|c\|_\infty$ to the optimal value (E.6) and then apply the exponential function to it. Finally, with this setup, I can implement the limit (3.4) as

$$\lim_{\ell \rightarrow \infty} \frac{\log n_\ell}{N_\ell} = 0. \quad (\text{E.7})$$

E.2 Numerical Illustration

This section considers a toy example that is useful for establishing proof of concept of my approximation scheme and its implementation. I specify $X \sim U[0, 1]$ and $Y \sim U[0, 2]$, and $c(x, y) = \frac{1}{2}(x - y)^2$. This specification satisfies the conditions of Assumption 1.

In this setting, the optimal transport value is the squared Wasserstein distance between P_X and P_Y (Villani, 2009), which is given by

$$W_2^2(P_X, P_Y) = \inf_{P \in \Pi(P_X, P_Y)} \frac{1}{2} \int_{\mathcal{X} \times \mathcal{Y}} (x - y)^2 dP. \quad (\text{E.8})$$

Table 1: Primitives

γ	κ	ϵ_ℓ	n_ℓ	N_ℓ
100	1	0.1	160	1015

The value and the solution of (E.8) can be calculated in closed-form. In particular, $W_2^2(P_X, P_Y) = 1/6 \approx 0.166$ and the solution is the singular distribution supported on $\{(x, y) : y = 2x\}$ having CDF

$$F_{X,Y}(x, y) = \begin{cases} 0, & x, y < 0, \\ \min\{x, y/2\}, & 0 \leq x \leq 1, 0 \leq y \leq 2, \\ 1, & x > 1, y > 2. \end{cases} \quad (\text{E.9})$$

The EOT value with $\gamma = 100$ is approximately 0.1846, and I have compared the finite-sample performance of the EOT value estimators based on the sieve M-estimator and empirical Sinkhorn divergence. I have studied their performance using 1000 Monte Carlo draws from R_γ to estimate the absolute deviation of their means from the target, 0.1846. I have set the sample size for both estimators to be the optimal one for the sieve-based estimator, given by display (4.1) in Section 4, and I repeat it here for convenience: $N_\ell = 2\epsilon_\ell^{-2} \log n_\ell$. The sieve M-estimator is $\hat{W}_2^2(P_X, P_Y) = -\gamma^{-1} \log \hat{\vartheta}_\ell + \gamma^{-1} \log a_\gamma$, where I have calculated $\hat{\vartheta}_\ell$ via the optimization problem (E.6). Table 1 reports the primitives of the optimization problem (E.6). Table 1 reports the primitives of the optimization problem (E.6). I have obtained the values 0.0254 and 0.0375 for the sieve M-estimator and the empirical Sinkhorn divergence, respectively. While this numerical result indicates the sieve M-estimator is less biased than its empirical Sinkhorn counterpart, it is more informative to report the boxplots of their MC estimates. The boxplots approximate the sampling distributions of estimators, and they are a powerful visual tool for illustrating finite-sample properties of the estimators.

Figure 1 reports the boxplots. Both estimators are centered below the OT benchmark, indicating a pronounced finite-sample downward bias in this high-regularization regime. This behavior contrasts with the population ordering, where the EOT value exceeds the unregularized OT cost, and highlights the impact of finite-sample effects and discretization error on the empirical objective. Relative to the empirical Sinkhorn divergence, the sieve M-estimator exhibits reduced variability and lower bias, suggesting improved stability at the theoretically prescribed sieve dimension and sample size.

The improved finite-sample performance of the sieve M-estimator relative to the empirical Sinkhorn estimator can be understood through the structure of their respective feasible sets. The population OT and EOT problems are characterized by exact marginal constraints, which may be equivalently expressed as an infinite collection of moment conditions. The sieve M-estimator replaces this infinite system by a finite, accuracy-controlled set of moment constraints indexed by ϵ_ℓ , thereby constructing a deterministic approximation to the population constraint set. As a result, even in finite samples, the sieve estimator




Figure 1: Boxplots of simulated estimates

optimizes over a feasible region that remains closely aligned with the population OT and EOT problems.

In contrast, the empirical Sinkhorn estimator enforces marginal constraints through empirical measures, leading to a feasible set that is random and subject to sampling fluctuations. These fluctuations introduce additional variability and bias into the estimated transport cost, particularly in finite samples. This distinction helps explain why, in the simulation results, the sieve M-estimator is both more concentrated and systematically closer to the population OT and EOT benchmark values than the empirical Sinkhorn estimator at the same sample size.