

The Burness-Giudici Conjecture on Primitive Groups of Lie-type with Rank One: Part (II)

HUYE CHEN

Guangxi University, Nanning 530004, P. R. China

SHAOFEI DU

Capital Normal University, Beijing 100048, P. R. China

WEICONG LI

Great Bay University, Dongguan 523000, P. R. China

Abstract

It was conjectured by Burness and Giudici that every primitive permutation group G containing some regular suborbits has the property that $\Gamma \cap \Gamma^g \neq \emptyset$ for any $g \in G$, where Γ is the union of all regular suborbits of G relative to α . We focused on proving the conjecture for all primitive groups G whose socle is a simple group of Lie-type of rank 1, that is, those with $\text{soc}(G) \in \{\text{PSL}(2, q), \text{PSU}(3, q), \text{Ree}(q), \text{Sz}(q)\}$. The case of $\text{soc}(G) = \text{PSL}(2, q)$ has been published in two papers. The case of $\text{soc}(G) = \text{PSU}(3, q)$ is divided into two parts: Part (I) addressed all primitive groups G with socle $\text{PSU}(3, q)$ whose point stabilizers contain $\text{PSO}(3, q)$ and based on it, this Part (II) will finish all cases of $\text{soc}(G) = \text{PSU}(3, q)$. The cases for $\text{soc}(G) \in \{\text{Ree}(q), \text{Sz}(q)\}$ will be treated in Part (III).

1 Introduction

Let $G \leq \text{Sym}(\Omega)$ be a permutation group. Then a subset Δ of Ω is called a *base* if its pointwise stabilizer therein is trivial. The *base size* of G on Ω , denoted by $b(G)$, is the minimal cardinality of a base for G . A *base size set* Δ for a finite permutation group $G \leq \text{Sym}(\Omega)$, is a base for G on the set Ω with $|\Delta| = b(G)$.

Recently, Burness and Giudici [3] introduced a graph, called the *Saxl graph* which is related to bases of a permutation group with $b(G) = 2$. Let $G \leq \text{Sym}(\Omega)$ be a permutation group on Ω with $b(G) = 2$. The vertex set of a Saxl graph $\Sigma(G)$ (denoted by Σ , simply) of G on Ω is just Ω and two vertices are adjacent if and only if they form a base size set.

Henceforth, assume that G acts transitively on Ω . Fix a point $\alpha \in \Omega$. The orbits of the stabilizer G_α are called the *suborbits* of G with respect to α , where $\{\alpha\}$ is said to be trivial. A pair α, β is a base for G if and only if G_α acts regularly on the suborbit containing β , given that $b(G) = 2$. Note that the Saxl graph $\Sigma(G)$ is vertex-transitive whenever G is a transitive permutation group on Ω . Moreover, the neighborhood $\Sigma_1(\alpha)$

Keywords base of permutation group, Saxl graph, simple group

E-mail addresses: chenhy280@gxu.edu.cn (H. Chen), dushf@mail.cnu.edu.cn (S. Du), liweicong@gbu.edu.cn (W. Li).

of α in $\Sigma(G)$ is the union Γ of the regular suborbits of G relative to α . In their paper [3], the authors investigated the valency, connectivity, hamiltonicity, and independence number of $\Sigma(G)$. They also proposed a conjecture, which can be stated as follows.

Conjecture 1.1 (BG-Conjecture) *Every finite primitive permutation group G with $b(G) = 2$ has the property that $\Gamma \cap \Gamma^g \neq \emptyset$ for any $g \in G$, where Γ is the union of all regular suborbits of G relative to a point α .*

To date, this conjecture is still open for simple groups of Lie type. Definitely, this is a huge and difficult topic. The first candidate may be those groups of rank one, that is $\text{soc}(G) \in \{\text{PSL}(2, q), \text{PSU}(3, q), \text{Ree}(q), \text{Sz}(q)\}$. The conjecture has been settled for groups with socle $\text{PSL}(2, q)$ in [4, 5]. Accordingly, we focus on the remaining three families. We shall finish them with three papers. The first one [6], had considered the case of $\text{soc}(G) = \text{PSU}(3, q)$ with a point stabilizer containing $\text{SO}(3, q)$. Based on it, this Part (II) will show the conjecture for all cases for $\text{soc}(G) = \text{PSU}(3, q)$ (see Therorem 1.2). Finally, Part (III)[7] will address the case of G with socle $\text{Ree}(q)$ and $\text{Sz}(q)$. For the relevant research background for this topic in details, see Part I [5]. Now we are ready to state the main result of this paper.

Theorem 1.2 *Let G be a primitive group with socle $\text{PSU}(3, q)$ and base size $b(G) = 2$, where $q = p^m \geq 5$ for a prime p . Then the BG-Conjecture holds for G , except for some small cases $G = \text{PSU}(3, q)\mathbb{Z}_{m_1}$ and $M = \text{SO}(3, q)\mathbb{Z}_{m_1}$ ($m_1 \mid 2m$ and $m_1 \neq 1$), where $\sqrt{q} \leq 17$ or 45 , if $3 \nmid (q+1)$ or $3 \mid (q+1)$, respectively.*

This paper is organized as follows. Following this introduction, Section 2 provides the necessary notation, basic definitions, and useful facts. The proof of Theorem 1.2 is presented in Section 3. With the exception of certain cases that are settled within Sections 3 and 7, the core of the proof is concluded to a family of groups. The detailed study of this family will be given in Sections 4-6, while general preparations are given in Section 4, and the two main subcases are analyzed in Sections 5 and 6, respectively.

2 Preliminaries

Let \mathbb{F}_q be a finite field of order $q = p^m$, where p is a prime and m is a positive integer. Let $\mathbb{F}_{q^2}^* = \langle \xi \rangle$ and $\mathbb{F}_q^* = \langle \theta \rangle$ be the cyclic multiplicative groups. The Frobenius automorphism $f \in \text{Aut}(\mathbb{F}_{q^2})$, defined as $f(\xi) = \xi^p$, has order $2m$.

Let V be the 3-dimensional unitary space of row vectors over \mathbb{F}_{q^2} , equipped with a Hermitian form (\cdot, \cdot) . We denote by $\text{GU}(3, q)$, $\text{SU}(3, q)$, $\text{PGU}(3, q)$ and $\text{PSU}(3, q)$ the general unitary, special unitary, projective general unitary, and projective special unitary groups, respectively. Note that $\text{PSU}(3, q)$ is simple for $q > 2$. We recall some known results for $T := \text{PSU}(3, q)$, whose order is given by $|T| = \frac{1}{d}(q^3 + 1)q^3(q^2 - 1)$ with $d := (3, q + 1)$. Throughout this paper, we let $\text{Syl}_p(G)$ denote the set of sylow

p -subgroups of a group G . Moreover, for any $a, b \in G$, we define the commutator as $[a, b] := aba^{-1}b^{-1}$.

(1) We choose a unitary basis w_1, w_2, w_3 whose Gram matrix is the antidiagonal matrix $J := [1, 1, 1]$ (see [8]). Using the automorphism $a \mapsto \bar{a} := a^q$ ($a \in \mathbb{F}_{q^2}$), we define $\text{GU}(3, q) := \{g \in \text{GL}(3, q^2) \mid gJ\bar{g}^T = J\}$. Let $\hat{X} := XZ/Z$ be the image of a subgroup X of $\text{GU}(3, q)$, where $Z := Z(\text{GU}(3, q))$ and use \hat{g} to denote the image of an element $g \in \text{GU}(3, q)$. In $\text{SU}(3, q)$, define

$$q(a, b) = \begin{pmatrix} 1 & a & b \\ 0 & 1 & -\bar{a} \\ 0 & 0 & 1 \end{pmatrix}, \quad h(k) = \begin{pmatrix} k^{-q} & 0 & 0 \\ 0 & k^{q-1} & 0 \\ 0 & 0 & k \end{pmatrix}, \quad \tau = \begin{pmatrix} 0 & 0 & 1 \\ 0 & -1 & 0 \\ 1 & 0 & 0 \end{pmatrix},$$

$$Q := \{\hat{q}(a, b) \mid a, b \in \mathbb{F}_{q^2}, b + \bar{b} + a\bar{a} = 0\}, \quad H := \{\hat{h}(k) \mid k \in \mathbb{F}_{q^2}^*\}.$$

Thus, Q is a Sylow p -subgroup of $T = \text{PSU}(3, q)$ of order q^3 ; and H is a cyclic subgroup of T of order $\frac{q^2-1}{d}$, with relations

$$\begin{aligned} \hat{q}(a, b)\hat{q}(a', b') &= \hat{q}(a + a', b + b' - a\bar{a}'), & \hat{\tau}\hat{h}(k)\hat{\tau} &= \hat{h}(k^{-q}), \\ \hat{h}(k)^{-1}\hat{q}(a, b)\hat{h}(k) &= \hat{q}(k^{2q-1}a, k^{q+1}b), & \hat{q}(a, b)^{-1} &= \hat{q}(-a, \bar{b}). \end{aligned} \quad (1)$$

Further, the center of Q , given by $Z(Q) =: Q_0 = \{\hat{q}(0, b) \mid b \in \mathbb{F}_{q^2}, b + \bar{b} = 0\}$, has order q .

(2) Let $\delta = [\mu, 1, 1]Z$, where $Z := Z(\text{GU}(3, q)) = \{[\mu, \mu, \mu] \mid \mu \in \mathbb{F}_{q^2}, \mu^{q+1} = 1\}$; here $[\mu, 1, 1]$ denotes the diagonal matrix with entries $(\mu, 1, 1)$, and $|\delta| = d$. Hence, $\text{PGU}(3, q) = T \rtimes \langle \delta \rangle$. The field automorphism of T induced by f is redenoted by f again so that $\text{Aut}(T) = T \rtimes \langle \delta, f \rangle$.

Proposition 2.1 [2, p.77,79]. *Let $T = \text{PSU}(3, q)$ and f the field automorphism. Then*

- (i) *Let $x \in \langle f \rangle$ such that x is of order a prime r . Then for any $g \in \text{PGU}(3, q)$, $|gx| = r$ if and only if $gx = x^{g_1}$ for some $g_1 \in \text{PGU}(3, q)$.*
- (ii) *Let $z \in \text{PGU}(3, q)$ be an element of prime order $r \neq p$. Then $z^T = z^{\text{PGU}(3, q)}$.*

Proposition 2.2 [9] *Let G be a transitive group on Ω and let $H = G_\alpha$ for some $\alpha \in \Omega$. Suppose that $K \leq G$ and at least one G -conjugate of K is contained in H . Suppose further that the set of G -conjugates of K which are contained in H form t conjugacy classes of H with representatives K_1, K_2, \dots, K_t . Then K fixes $\sum_{i=1}^t |\text{N}_G(K_i) : \text{N}_H(K_i)|$ points of Ω .*

Let $\text{fix}(x, G/M)$ denote the number of fixed points of x on $\Omega := [G : M]$ and let

$$\text{fpr}(x, G/M) = \frac{\text{fix}(x, G/M)}{|\Omega|}$$

be the fixed point ratio of x . The following proposition is modified from [3].

Proposition 2.3 Consider the transitive permutation representation of G on $\Omega := [G : M]$. Let $\check{Q}(G) := \frac{|M|}{|\Omega|} \left(\sum_{i=1}^k (|\langle x_i \rangle| - 1) \frac{\text{Fix}(\langle x_i \rangle)}{|\text{N}_M(\langle x_i \rangle)|} \right)$, where $\mathcal{P}^*(M) := \{\langle x_1 \rangle, \langle x_2 \rangle, \dots, \langle x_k \rangle\}$ is the set of representatives of conjugacy classes of subgroups of prime order in M and $\text{Fix}(\langle x_i \rangle)$ is the number of fixed points of $\langle x_i \rangle$ on Ω . Then any two vertices in $\Sigma(G)$ have a common neighbour, provided $\check{Q}(G) < \frac{1}{2}$.

Remark 2.4 Let G and G_0 be primitive groups on Ω with $b(G) = b(G_0) = 2$, where G_0 is a subgroup of G with a point stabilizer contained in that of G . If the BG-Conjecture holds for G acting on Ω , then the BG-Conjecture also holds for G_0 acting on Ω . The result follows from the inclusion $\Gamma(\alpha) \subseteq \Gamma_0(\alpha)$ of the respective sets of regular suborbits.

3 The proof of Theorem 1.2

Proof Let G be an almost simple group with socle $\text{soc}(G) = T = \text{PSU}(3, q)$, that is, $T \leq G \leq \text{Aut}(T)$. Consider the primitive permutation representations of G on the right cosets $[G : M]$ relative to maximal subgroups $\alpha = M$ of G , up to conjugacy. To prove Theorem 1.2, we shall find a common neighbor of α and α^g for any $g \in G$ in the Saxl graph.

Suppose that the group M is soluble. Then the conjecture is true by [4, Theorem 1.1]. So in what follows, we assume that M is insolvable. If $T \lesssim M$, then G/M_G is cyclic, where $M_G = \bigcap_{g \in G} M^g$. In this case, $b(G) = 1$, which is not our case. So we assume that $T \not\lesssim M$. The case when $q = 5$ may be verified by Magma directly and so assume that $q \geq 7$ below.

Let $M_0 := M \cap T$. Then by [2, Table 8.5 and Table 8.6], M_0 is a maximal subgroup of T and

$$M_0 \cong L \in \{\text{SO}(3, q), \text{SL}(2, q) \rtimes \mathbb{Z}_{\frac{q+1}{d}}, \text{PSU}(3, q').\mathbb{Z}_c, \text{PSL}(2, 7), \text{PSL}(2, 9)\},$$

where $q = q'^e$, $c := (\frac{q+1}{q'+1}, 3)$ and e is an odd prime. Note that $M_0 \cong \text{PSL}(2, 7)$ for $q = p \equiv 3, 5, 6 \pmod{7}$ with $q \neq 5$; and $M_0 \cong \text{PSL}(2, 9)$ for $q = p \equiv 11, 14 \pmod{15}$. These five cases will be dealt with separately.

(1) $M_0 \cong \text{SO}(3, q)$: This case has been dealt with in [6]. In there it was shown that the BG-Conjecture holds for G , except for some small cases: $G = \text{PSU}(3, q)\mathbb{Z}_{m_1}$ and $M = \text{SO}(3, q)\mathbb{Z}_{m_1}$ ($m_1 \mid 2m$ and $m_1 \neq 1$), where $\sqrt{q} \leq 17$ or 45 , if $3 \nmid (q+1)$ or $3 \mid (q+1)$, respectively.

(2) $M_0 \cong \text{SL}(2, q) \rtimes \mathbb{Z}_{\frac{q+1}{d}}$

Let G be an almost simple group with socle $\text{soc}(G) = T = \text{PSU}(3, q)$, M a maximal subgroup of G containing $M_0 = \text{SL}(2, q) \rtimes \mathbb{Z}_{\frac{q+1}{d}}$. In what follows, we shall show $b(G) \neq 2$.

As a subgroup of G , the action of T on $\Omega = [G : M]$ may be identified with its primitive right multiplication on cosets $\Omega_0 = [T : M_0]$. Clearly, there exist no regular suborbits of G on Ω , that is $b(G) \neq 2$, provided there exist no regular suborbits of T , that is $b(T) \neq 2$.

Now, the action of T on $\Omega_0 = [T : M_0]$ is equivalent to its action on the set \mathcal{P}_{non} of non-isotropic points in the unitary projective plane $\text{PG}(2, q^2)$. We shall show $T_{\langle \alpha \rangle} \cap T_{\langle \beta \rangle} \neq 1$ for any $\langle \alpha \rangle, \langle \beta \rangle \in \mathcal{P}_{non}$ so that $b(T) \neq 2$. Since T acts transitively on \mathcal{P}_{non} , we may fix $\alpha = (1, 0, 0)$ without loss of generality. For any $g \in \text{SU}(3, q)$, the defining condition $g\bar{g}^T = E$ (where E is the identity Gram matrix), when applied to the stabilizer $T_{\langle \alpha \rangle}$, implies that

$$T_{\langle \alpha \rangle} = \langle \hat{g} \mid g = \begin{pmatrix} a & 0 \\ 0 & A \end{pmatrix} \in \text{SU}(3, q), A \in \text{GU}(2, q), a|A| = 1 \rangle.$$

Now β is of form $(0, 0, 1)$, $(0, 1, c_1)$ or $(1, b, c_2)$ for some $b, c_1, c_2 \in \mathbb{F}_{q^2}$ with $c_1^{1+q} + 1 \neq 0$ and $1 + b^{1+q} + c_2^{1+q} \neq 0$. Since \hat{A} acts nonregularly on $\langle (0, 1) \rangle$, $\langle (1, c_1) \rangle$ and $\langle (b, c_2) \rangle$, we know $T_{\langle \alpha \rangle} \cap T_{\langle \beta \rangle} \neq 1$.

(3) $M_0 \cong \text{PSU}(3, q') \cdot \mathbb{Z}_c$, which will be done in Sections 4-6, see Theorem 4.1.

(4) $M_0 \cong \text{PSL}(2, 7)$, $\text{PSL}(2, 9)$, see Theorems 7.1 and 7.2 in Section 7.

In summary, Theorem 1.2 is proved. \square

4 $M_0 \cong \text{PSU}(3, q') \cdot (\frac{q+1}{q'+1}, 3)$

In this section, we mainly prove the following theorem.

Theorem 4.1 *Let G be a primitive group with slope $T = \text{PSU}(3, q)$ acting on a set Ω , such that a point stabilizer contains a subgroup $M_0 \cong \text{PSU}(3, q') \cdot (\frac{q+1}{q'+1}, 3)$. Here, $q = p^m$ and $q' = p^{m'}$ with $m = m'e$, where p is a prime and e is an odd prime. Suppose that $b(G) = 2$. Then the BG-Conjecture is true.*

Proof Let $c = (\frac{q+1}{q'+1}, 3)$, $T = \text{PSU}(3, q)$, and G be a group such that $T \leq G \leq \text{PGU}(3, q)$. Furthermore, let M be a maximal subgroup of G containing $M_0 = \text{PSU}(3, q') \cdot \mathbb{Z}_c$. As shown in Table 8.5 of [1], we have $T \leq G \leq T : \langle \delta^c, f \rangle$, where δ is a diagonal automorphism of T of order $d = (q+1, 3)$. By the definition of the unitary groups, with $q = p^m$ and $\gamma := f^m$, we have the conjugation relations $\delta^f = \delta^p$ and $\delta^\gamma = \delta^{-1}$.

Write $q' + 1 = 3k + t$, where $t \in \{0, 1, 2\}$ and k is a nonnegative integer. Then

$$q + 1 = (3k + t - 1)^e + 1 \equiv (t - 1)^e + 1 \equiv 0 \pmod{3}$$

if and only if $t = 0$, and so $(q' + 1, 3) = (q + 1, 3)$.

Since $\frac{q+1}{q'+1} = \frac{((q'+1)-1)^e+1}{q'+1} \equiv e \pmod{q'+1}$ and e is an odd prime, we get $c := (\frac{q+1}{q'+1}, 3) = 3$ if and only if $e = 3$ and $(q' + 1, 3) = 3$.

In the following two sections, we deal with two cases $c = 1$ and $e \geq 3$; and $c = e = 3$, separately. Therefore, Theorem 4.1 is a direct consequence of Theorems 5.3 and 6.3. \square

Combining Propositions 3.3.2, 3.3.3, 3.3.7, and Table B.4 in [2], we obtain the following lemma.

Lemma 4.2 *Let $T = \mathrm{PSU}(3, q)$ and $R = \mathrm{PGU}(3, q) = T \rtimes \langle \delta \rangle$, where $(3, q+1) = 3$. For any element $x \in R$ of prime order r , the R -conjugacy class of x is determined as follows:*

1. *If $r \neq 3$, then x is R -conjugate to an element of T .*
2. *If $r = 3$, then x is R -conjugate to one of the three diagonal matrices z_A , z_B , or $\delta = [\mu, 1, 1]Z$, whose centralizers in R are respectively $C_R(z_A) \cong (\mathbb{Z}_{q+1} \times \mathbb{Z}_{q+1}) \rtimes \mathbb{Z}_3$, $C_R(z_B) \cong \mathrm{SL}(2, q) \rtimes \mathbb{Z}_{q+1}$, and $C_R(\delta) \cong \mathbb{Z}_{q^2-q+1} \rtimes \mathbb{Z}_3$.*

In the remaining of this section, some preparations are given. To prove Theorem 4.1 via Proposition 2.3, we require information on the prime order subgroups of $\mathrm{PGU}(3, q)$.

Lemma 4.3 *Let $T = \mathrm{PSU}(3, q)$ and $R = \mathrm{PGU}(3, q)$. Then every cyclic subgroup of prime order r in T is T -conjugate to $\langle z \rangle$, where the canonical representation of z , $|N_T(\langle z \rangle)|$ and $|N_R(\langle z \rangle)|$ are given in Table 1.*

Proof Noting that $|T| = \frac{1}{d}(q^3+1)q^3(q^2-1)$, for any prime divisor r of $|T|$ the possible cases for r are $r \in \{2, 3, p\}$, or $r \mid (q-1)$, $r \mid (q+1)$ or $r \mid (q^2-q+1)$. From Table 2, which describes the conjugacy classes of elements in T [10], we obtain the following cases based on the value of r .

(1) $r = 2$ and p is odd:

Note that the centralizer of every involution is the stabilizer of a non-isotropic point. Since both T and R are transitive on the set of non-isotropic points, it follows that all subgroups isomorphic to \mathbb{Z}_2 are conjugate in the group. Moreover, for any such involution z_2 , we have $|N_T(\langle z_2 \rangle)| = q(q^2-1)(q+1)/d$ and $|N_R(\langle z_2 \rangle)| = q(q^2-1)(q+1)$.

(2) $r = p$:

From Table 2, we get that $|C_T(z_0)| = \frac{q^3(q+1)}{d}$ and there is only one conjugacy class subgroup $\langle z_0 \rangle$ of this form, which is presented in Table 1. Note that $N_T(\langle z_0 \rangle)/C_T(\langle z_0 \rangle) \lesssim \mathbb{Z}_{p-1}$. From Eq(1), we have $\hat{h}(k)^{-1}\hat{q}(0, 1)\hat{h}(k) = \hat{q}(0, k^{q+1})$ and $(\hat{q}(0, 1))^l = \hat{q}(0, l)$ for any $k \in \mathbb{F}_{q^2}$ and positive integer l . Therefore, we have $|N_T(\langle z_0 \rangle)| = \frac{q^3(q+1)(p-1)}{d}$. Furthermore, $|C_R(\langle z_0 \rangle)| = q^3(q+1)$ followed from Proposition 2.1. This implies that $|N_R(\langle z_0 \rangle)| = q^3(q+1)(p-1)$, as $N_R(\langle z_0 \rangle)/C_R(\langle z_0 \rangle) \lesssim \mathbb{Z}_{p-1}$.

Table 2 shows that there are d conjugacy classes of z'_l in T , each with $|C_T(\langle z'_l \rangle)| = q^2$, and that these d classes are fused into a single class under the action of R . This implies that $|C_R(\langle z'_l \rangle)| = q^2$. Note that we may, without loss of generality, choose one

of the elements z'_l to lie in $\mathrm{PGO}(3, q) \cong \mathrm{PGL}(2, q)$; denote this element by z'_0 . Since $\mathbb{Z}_{p-1} \lesssim \mathrm{N}_{\mathrm{PGO}(3, q)}(z'_0)$ and $\mathrm{N}_T(\langle z'_0 \rangle)/\mathrm{C}_T(\langle z'_0 \rangle) \lesssim \mathbb{Z}_{p-1}$, we have $\mathrm{N}_T(\langle z'_0 \rangle) = q^2(p-1)$. This implies that $\mathrm{N}_R(\langle z'_l \rangle) = q^2(p-1)$, for any l , since these elements are all conjugate in R . Consequently, we also have $\mathrm{N}_T(\langle z'_l \rangle) = q^2(p-1)$ for any l .

(3) $r = 3$ and $3 \mid q+1$:

As can be seen from Table 2, z_3^A exists only when $3 \mid q+1$, and z_3^B exists only when $9 \mid q+1$, where z_3^A and z_3^B are of the form presented in Table 1. And $\langle z_3^A \rangle$ is not conjugate to $\langle z_3^B \rangle$ in $\mathrm{PSU}(3, q)$. Moreover, for $3 \mid q+1$, there exist only these two conjugacy classes of elements of order $r = 3$ in $\mathrm{PSU}(3, q)$. By Table 2, $|\mathrm{C}_T(z_3^A)| = (q+1)^2$ and $|\mathrm{C}_T(z_3^B)| = \frac{q(q+1)^2(q-1)}{3}$. By Proposition 2.1, we get that $|T : \mathrm{C}_T(x)| = |R : \mathrm{C}_R(x)|$ for any $x \in T$ of order co-prime to p . Therefore, we get that $|\mathrm{C}_R(z_3^A)| = 3(q+1)^2$ and $|\mathrm{C}_R(z_3^B)| = q(q+1)^2(q-1)$.

Note that $\mathrm{N}_T(\langle z_3^A \rangle)/\mathrm{C}_T(\langle z_3^A \rangle) \lesssim \mathbb{Z}_2$. This implies that $|\mathrm{N}_T(\langle z_3^A \rangle)| = (q+1)^2s$, where $s = 1, 2$. Hence $(q+1)^2s \mid |\mathrm{N}_R(\langle z_3^A \rangle)|$. From $|\mathrm{C}_R(z_3^A)| = 3(q+1)^2$, we conclude that $3(q+1)^2 \mid |\mathrm{N}_R(\langle z_3^A \rangle)|$. Therefore, $3s(q+1)^2 \mid |\mathrm{N}_R(\langle z_3^A \rangle)|$. Hence, the order must be $|\mathrm{N}_R(\langle z_3^A \rangle)| = 3s(q+1)^2$ with $s = 1, 2$.

By analyzing the structure of the subgroup of $\mathrm{PSU}(3, q)$, we get that $\mathrm{C}_T(\langle z_3^B \rangle) \cong \mathrm{SL}(2, q) \rtimes \mathbb{Z}_{\frac{q+1}{d}}$ and $\mathrm{C}_R(\langle z_3^B \rangle) \cong \mathrm{SL}(2, q) \rtimes \mathbb{Z}_{q+1}$. Consequently, the centralizer $\mathrm{C}_T(\langle z_3^B \rangle)$ (resp. $\mathrm{C}_R(\langle z_3^B \rangle)$) is a maximal subgroup of T (resp. R). This implies that $|\mathrm{N}_T(\langle z_3^B \rangle)| = |\mathrm{C}_T(\langle z_3^B \rangle)| = \frac{q(q+1)^2(q-1)}{3}$ and $|\mathrm{N}_R(z_3^B)| = |\mathrm{C}_R(z_3^B)| = q(q+1)^2(q-1)$.

(4) $r \mid (q+1)$, $r \neq 2, 3$:

By Table 2, for $r \mid q+1$ and $r \neq 2, 3$, there exist only these two types of conjugate classes elements in $\mathrm{PSU}(3, q)$ of order r , with respective elements z_{q+1}^A and z_{q+1}^B . For the type z_{q+1}^B , there is only one conjugate classes subgroup $\langle z_{q+1}^B \rangle$ of this type. For the type z_{q+1}^A , there are at most $\frac{r-3}{2}$ (depending on the choices of $(1, a, b)$ such that $1+a+b \equiv 0 \pmod{r}$) conjugate classes subgroup $\langle z_{q+1}^A(a) \rangle$ of this type, depending on the choices of a . By Table 2, we have that $|\mathrm{C}_T(z_{q+1}^A(a))| = \frac{(q+1)^2}{d}$ and $|\mathrm{C}_T(z_{q+1}^B)| = \frac{q(q+1)^2(q-1)}{d}$. By Proposition 2.1, we conclude that $|\mathrm{C}_R(\langle z_{q+1}^A(a) \rangle)| = (q+1)^2$ and $|\mathrm{C}_R(\langle z_{q+1}^B \rangle)| = q(q+1)^2(q-1)$. It follows that $|\mathrm{N}_R(\langle z_{q+1}^B \rangle)| = q(q+1)^2(q-1)$, as a maximal subgroup of R has order $q(q+1)^2(q-1)$. And we can also get that $|\mathrm{N}_T(\langle z_{q+1}^B \rangle)| = \frac{q(q+1)^2(q-1)}{d}$, as a maximal subgroup of T has order $\frac{q(q+1)^2(q-1)}{d}$.

By analyzing the structure of group T , we get that $\mathrm{N}_T(\langle z_{q+1}^A \rangle) = \mathbb{Z}_{\frac{q+1}{d}} \times \mathbb{Z}_{q+1} \rtimes S$ where $S \lesssim S_3$ is a permutation matrix and S is determined by the choices of $(1, a, b)$ where $1+a+b \equiv 0 \pmod{r}$. So that $\mathrm{N}_R(\langle z_{q+1}^A \rangle) = \mathbb{Z}_{q+1} \times \mathbb{Z}_{q+1} \rtimes S$.

(5) $r \mid (q-1)$, $r \neq 2$:

Followed from Table 2, there is only one conjugate classes subgroup of order r , where $r \mid q-1$ and $r \neq 2$. Let $\langle z_{q-1} \rangle$ be a subgroup of $\mathbb{Z}_{\frac{q^2-1}{d}} \cong L (\leq T)$ that has prime

order r . Note that $N_T(L) \cong \mathbb{Z}_{\frac{q^2-1}{d}} \rtimes \mathbb{Z}_2$. Thus, we obtain $N_T(\langle z_{q-1} \rangle) = \mathbb{Z}_{\frac{q^2-1}{d}} \rtimes \mathbb{Z}_2$ from the structure of the subgroups of T . By Table 2, we have that $|C_T(\langle z_{q-1} \rangle)| = \frac{q^2-1}{d}$. It follows from Proposition 2.1 that $|C_R(\langle z_{q-1} \rangle)| = q^2 - 1$. Therefore, we have that $N_R(\langle z_{q-1} \rangle) = \mathbb{Z}_{q^2-1} \rtimes \mathbb{Z}_2$.

(6) $r \mid q^2 - q + 1$:

Let $\langle z_{q^2-q+1} \rangle$ be a character subgroup of $\mathbb{Z}_{\frac{q^2-q+1}{d}} \cong A$ ($\leq T$) with order r . Note that there is only one conjugate classes subgroup in $\mathrm{PSU}(3, q)$ of order $\frac{q^2-q+1}{d}$ and $N_T(A) = \mathbb{Z}_{\frac{q^2-q+1}{d}} \rtimes \mathbb{Z}_3$. Thus, we obtain $N_T(\langle z_{q^2-q+1} \rangle) \cong \mathbb{Z}_{\frac{q^2-q+1}{d}} \rtimes \mathbb{Z}_3$, which is maximal in T . By Table 2, we have that $|C_T(\langle z_{q^2-q+1} \rangle)| = \frac{q^2-q+1}{d}$. It follows from Proposition 2.1 that $|C_R(\langle z_{q^2-q+1} \rangle)| = q^2 - q + 1$. Therefore, we have that $N_R(\langle z_{q^2-q+1} \rangle) = \mathbb{Z}_{q^2-q+1} \rtimes \mathbb{Z}_3$. \square

For $q' = 2$, the group $\mathrm{PGU}(3, 2) \rtimes \langle f \rangle$ is soluble. In this case, the BG-Conjecture is established by [4]. For $e = 3$ and $q' = 3$, one may confirm the BG-Conjecture by Magma. From now on, we assume that either $e \geq 5$ and $q' \geq 3$ or $e = 3$ and $q' \geq 4$. We will address the cases where $(\frac{q+1}{q'+1}, 3)$ equals 1 and 3 in Sections 5 and 6, respectively.

5 $M_0 \cong \mathrm{PSU}(3, q').(\frac{q+1}{q'+1}, 3)$ with $(\frac{q+1}{q'+1}, 3) = 1$

Let $q = p^m$, $q' = p^{m'}$ for some odd prime e such that $m = m'e$. Assume that $c := (\frac{q+1}{q'+1}, 3) = 1$. Moreover, let $e \geq 5$ and $q' \geq 3$ or $e = 3$ and $q' \geq 4$. From the proof of Theorem 4.1, we get $(3, q+1) =: d = d' := (3, q'+1)$. Let

$$\begin{aligned} T &:= \mathrm{PSU}(3, q) \leq G_1 := \mathrm{PGU}(3, q) = T\langle \delta \rangle \leq G := \mathrm{PGU}(3, q) = G_1\langle f \rangle; \\ M_0 &:= \mathrm{PSU}(3, q') \leq M_1 := \mathrm{PGU}(3, q') = M_0\langle \delta \rangle \leq M := \mathrm{PGU}(3, q') : \langle f \rangle = M_1\langle f \rangle; \\ \Omega &= [G : M]. \end{aligned}$$

In what follows, we shall consider the action of G on $\Omega = [G : M]$ and prove $\check{Q}(G) < \frac{1}{2}$ so that the BG-Conjecture is true for G . By Remark 2.4, this implies that the BG-Conjecture is true for every group between T and G with the corresponding point-stabilizers.

First we determine the representatives $\langle z \rangle$ of conjugacy classes of subgroups of prime order in M , while these subgroups are devided into two subclasses: those are contained in M_1 or those are not. We shall deal with them in Lemmas 5.1 and 5.2, separately; measure $Q(\langle z \rangle)$ for these subgroups $\langle z \rangle$, where

$$Q(\langle z \rangle) := \frac{|M|}{|\Omega|} \frac{(|z| - 1)\mathrm{Fix}(\langle z \rangle)}{|N_M(\langle z \rangle)|};$$

and then prove $\check{Q}(G) < \frac{1}{2}$ by Proposition 2.3 so that Theorem 5.3 holds for the case $c = 1$.

Table 1: Subgroups of prime order r in $\mathrm{PSU}(3, q)$: conjugacy classes

r	p	p	2
representation	$z_0 = \begin{pmatrix} 1 & 0 & \epsilon \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$ $\mathrm{J}=[1,1,1]$	$z'_l = \begin{pmatrix} 1 & a' & b \\ 0 & 0 & -a'q \\ 0 & 1 & 1 \end{pmatrix}$ $\mathrm{J}=[1,1,1]$	$z_2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$ $\mathrm{J}=[1,1,1]$
$ \mathrm{C}_T(\langle z \rangle) $	$q^3(q+1)/d$	q^2	$q(q^2-1)(q+1)/d$
$ \mathrm{N}_T(\langle z \rangle) $	$q^3(q+1)(p-1)/d$	$q^2(p-1)$	$q(q^2-1)(q+1)/d$
$ \mathrm{C}_R(\langle z \rangle) $	$q^3(q+1)$	q^2	$q(q^2-1)(q+1)$
$ \mathrm{N}_R(\langle z \rangle) $	$q^3(q+1)(p-1)$	$q^2(p-1)$	$q(q^2-1)(q+1)$
Remark	$\epsilon^q + \epsilon = 0, \epsilon \in \mathbb{F}_{q^2}$; and $\epsilon = 1$ if $p = 2$	$\begin{aligned} 0 \leq l \leq d-1; \\ a = 3 \text{ and } b = -\frac{1}{3} \text{ if } d = 3; \\ z'_0 = z'_R = z'_2, \text{ if } d = 3 \end{aligned}$	$q \text{ odd}$
r	3	3	$r \mid q+1$
representation	$z_3^A = [1, \omega, \omega^2]$ $\mathrm{J}=[1,1,1]$	$z_3^B = [\rho^{\frac{q+1}{9}}, \rho^{\frac{q+1}{9}}, \rho^{\frac{-2(q+1)}{9}}]$ $\mathrm{J}=[1,1,1]$	$z_{q+1}^A(a) = [x, x^a, x^b]$ $\mathrm{J}=[1,1,1]$
$ \mathrm{C}_T(\langle z \rangle) $	$(q+1)^2$	$q(q+1)^2(q-1)/3$	$(q+1)^2/d$
$ \mathrm{N}_T(\langle z \rangle) $	$s(q+1)^2$	$q(q+1)^2(q-1)/3$	$s(q+1)^2/d$
$ \mathrm{C}_R(\langle z \rangle) $	$3(q+1)^2$	$q(q+1)^2(q-1)$	$(q+1)^2$
$ \mathrm{N}_R(\langle z \rangle) $	$3s(q+1)^2$	$q(q+1)^2(q-1)$	$s(q+1)^2$
Remark	$\begin{aligned} \omega \in \mathbb{F}_{q^2}, s=1,2 \\ \omega = 3, 3 \mid q+1 \end{aligned}$	$\begin{aligned} \rho \in \mathbb{F}_{q^2}; \\ \rho = q+1, 9 \mid q+1 \end{aligned}$	$\begin{aligned} x \in \mathbb{F}_{q^2}, 1+a+b \equiv 0 \pmod{r}; \\ \omega = r, r \mid q+1, s \mid 6; \\ 2 \leq a \leq \frac{q-1}{2}, \end{aligned}$
r	$r \mid q+1$	$r \mid q-1$	$r \mid q^2-q+1$
representation	$z_{q+1}^B := [x, x, x^{-2}]$ $\mathrm{J}=[1,1,1]$	$z_{q-1} := [\theta^{\frac{q-1}{r}}, 1, \theta^{-\frac{q-1}{r}}]$ $\mathrm{J}=[1,1,1]$	z_{q^2-q+1}
$ \mathrm{C}_T(\langle z \rangle) $	$q(q+1)^2(q-1)/d$	q^2-1/d	q^2-q+1/d
$ \mathrm{N}_T(\langle z \rangle) $	$q(q+1)^2(q-1)/d$	$2(q^2-1)/d$	$3(q^2-q+1)/d$
$ \mathrm{C}_R(\langle z \rangle) $	$q(q+1)^2(q-1)$	q^2-1	q^2-q+1
$ \mathrm{N}_R(\langle z \rangle) $	$q(q+1)^2(q-1)$	$2(q^2-1)$	$3(q^2-q+1)$
Remark	$\begin{aligned} x \in \mathbb{F}_{q^2}; \\ \omega = r, r \mid q+1 \end{aligned}$	$\mathbb{F}_q^* = \langle \theta \rangle$	*

Table 2: Elements of order greater than 2 in $\mathrm{PSU}(3, q)$: conjugacy classes

Label	number of classes	Representative g	Gram matrix \mathbf{J}	$ g $	$ C_T(g) $	parameters
\mathcal{C}_2	1	$\begin{pmatrix} 1 & 0 & \epsilon \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$	$\begin{bmatrix} 1, 1, 1 \\ 1, 1, 1 \end{bmatrix}$	p	$\frac{q^3(q+1)}{d}$	$\epsilon^q + \epsilon = 0, \epsilon \in \mathbb{F}_{q^2};$ and $\epsilon = 1$ if $p = 2$
$\mathcal{C}^{(l)}$	d	$\begin{pmatrix} 1 & a' & b \\ 0 & 1 & -a'^q \\ 0 & 0 & 1 \end{pmatrix}$	$\begin{bmatrix} 1, 1, 1 \\ 1, 1, 1 \end{bmatrix}$	p	q^2	$0 \leq l \leq d-1,$ $a'^{q+l} + b + b^q = 0, b \neq 0;$ $ a = 3$ and $b = -\frac{1}{2}$ if $d = 3$
$\mathcal{C}_4^{(k)}$	$\frac{q+1}{d} - 1$	$[\rho^k, \rho^k, \rho^{-2k}]$	$\begin{bmatrix} 1, 1, 1 \\ 1, 1, 1 \end{bmatrix}$	$ g \mid \frac{q+1}{k}$	$\frac{q(q+1)^2(q-1)}{d}$	$1 \leq k \leq \frac{q+1}{d} - 1;$ $\rho \in \mathbb{F}_{q^2}, \rho = q + 1$
\mathcal{C}'_6	$\frac{d-1}{2}$	$[1, \omega, \omega^2]$	$\begin{bmatrix} 1, 1, 1 \\ 1, 1, 1 \end{bmatrix}$	3	$(q+1)^2$	$\omega \in \mathbb{F}_{q^2};$ $ \omega = 3$
$\mathcal{C}_6^{(k,l,m)}$	$\frac{q^2-q+1-1}{6}$	$[\rho^k, \rho^l, \rho^m]$	$\begin{bmatrix} 1, 1, 1 \\ 1, 1, 1 \end{bmatrix}$	$ g \mid (q+1)$	$\frac{(q+1)^2}{d}$	$1 \leq k < l \leq \frac{q+1}{d};$ $l < m \leq q+1;$ $k + l + m \equiv 0 \pmod{q+1};$ $\rho \in \mathbb{F}_{q^2}, \rho = q + 1$
$\mathcal{C}_7^{(k)}$	$\frac{q^2-q+1-1}{2} - \frac{3-d}{2}$	$[\sigma^k, \rho^k, \sigma^{-qk}]$	$\begin{bmatrix} 1, 1, 1 \\ 1, 1, 1 \end{bmatrix}$	$ g \mid (q^2 - 1)$	$\frac{q^2-1}{d}$	$\rho, \sigma \in \mathbb{F}_{q^2};$ $ \rho = q + 1, \sigma = q^2 - 1;$ $1 \leq k \leq \frac{q^2-1}{d};$ $k \neq 0 \pmod{q-1};$ $\mathcal{C}_7^{(k)} = \mathcal{C}_7^{(-qk)}$
$\mathcal{C}_8^{(k)}$	$\frac{q^2-q+1-1}{3}$	*	*	$ g \mid q^2 - q + 1$	$\frac{q^2-q+1}{d}$	$\tau \in \mathbb{F}_{q^6}, \tau = q^2 - q + 1;$ $1 \leq k \leq \frac{q^2-q+1}{d} - 1;$ $\mathcal{C}_8^{(k)} = \mathcal{C}_8^{(-qk)} = \mathcal{C}_8^{(q^2k)}$

Lemma 5.1 Let $c = 1$ and either $e \geq 5$ and $q' \geq 3$ or $e = 3$ and $q' \geq 4$, where $q' = p^{m'}$. Then using the notations given in Table 1 and Lemma 4.2, every cyclic subgroup $\langle z \rangle$ of prime order r of M_1 is M -conjugate to one of the following groups.

(I) Cases either $e \geq 5$ and $q' \geq 3$ or $e = 3$ and $q' \geq 5$:

- (1) $r = 2$ and p is odd: $\langle z_2 \rangle$ with $Q(\langle z_2 \rangle) < \frac{1}{26}$.
- (2) $r = p$: $\langle z_0 \rangle$ with $Q(\langle z_0 \rangle) < \frac{1}{26}$; and $\langle z'_0 \rangle$ with $Q(\langle z'_0 \rangle) < \frac{1}{26}$.
- (3) $r = 3$ and $r \mid q'+1$: $\langle z_A \rangle$ with $Q(\langle z_A \rangle) < \frac{1}{26}$; $\langle z_B \rangle$ with $Q(\langle z_B \rangle) < \frac{1}{26}$; and $\langle \delta' \rangle \in M_1 \setminus M_0$ with $Q(\langle \delta' \rangle) < \frac{1}{26}$.
- (4) $r \mid q' - 1$ and $r \neq 2$: $\langle z_{q'-1} \rangle$ with $Q(\langle z_{q'-1} \rangle) < \frac{1}{26(q'-1)}$.
- (5) $r \mid q' + 1$ and $r \neq 2, 3$: $\langle z_{q'+1}^A(a) \rangle$ with $Q(\langle z_{q'+1}^A(a) \rangle) < \frac{1}{13(q'+1)^2}$; and $\langle z_{q'+1}^B \rangle$ with $Q(\langle z_{q'+1}^B \rangle) < \frac{1}{26(q'+1)}$.
- (6) $r \mid q'^2 - q' + 1$ with $r \neq 3$: $\langle z \rangle = \langle z_{q'^2-q'+1} \rangle$. If $(e, p, m') \notin \{(3, 2, 2), (3, 2, 4), (3, 3, 2)\}$, then $Q(\langle z_{q'^2-q'+1} \rangle) < \frac{1}{26(q'^2-q'+1)}$. Otherwise, there is a unique conjugacy class of subgroups of order r , and we have $Q(\langle z_{q'^2-q'+1} \rangle) < \frac{1}{26}$.

(II) Case $e = 3$ and $q' = 4$:

- (1) $r = 2$: $\langle z_0 \rangle$ with $Q(\langle z_0 \rangle) < 0.1 \times \frac{1}{26}$; and $\langle z'_0 \rangle$ with $Q(\langle z'_0 \rangle) < 0.85 \times \frac{1}{26}$.
- (2) $r = 3$: $\langle z_{q'-1} \rangle$ with $Q(\langle z_{q'-1} \rangle) < 0.02 \times \frac{1}{26}$.
- (3) $r = 5$: $\langle z_{q'+1}^A(a) \rangle$ with $Q(\langle z_{q'+1}^A(a) \rangle) < 0.75 \times \frac{1}{26}$; and $\langle z_{q'+1}^B \rangle$ with $Q(\langle z_{q'+1}^B \rangle) < 0.28 \times \frac{1}{26}$.
- (4) $r = 13$: $\langle z_{q'^2-q'+1} \rangle$ and $Q(\langle z_{q'^2-q'+1} \rangle) < 0.5 \times \frac{1}{26}$.

Proof Suppose $c = 1$ and either $e \geq 5$ and $q' \geq 3$ or $e = 3$ and $q' \geq 5$. The conclusion for $e = 3$ and $q' = 4$, that is (II) in the lemma, can be computed directly by inserting the exact value of e, q' in the proof of (I). So we assume either $e \geq 5$ and $q' \geq 3$ or $e = 3$ and $q' \geq 7$ (if $e = 3$, then $c = 1$ gives $q' \neq 5$). First we have

$$\begin{aligned} \frac{|M|}{|\Omega|} &= \frac{|M|^2}{|G|} = \frac{2m|\text{PGU}(3, q')|^2}{|\text{PGU}(3, q)|} = \frac{2mq'^6(q'^3+1)^2(q'^2-1)^2}{q^3(q^3+1)(q^2-1)} \\ &\leq \frac{2mq'^{16}}{q^8} \frac{(1+q'^{-3})^2}{(1-q'^{-2})} \frac{(1-q'^{-2})^2}{1+q'^{-3}} \leq \frac{2mq'^{16}}{q^8} \cdot \frac{13}{12} \cdot 1 \leq \frac{13mq'^{16}}{6q^8}. \end{aligned}$$

By Lemmas 4.2 and 4.3, we need to discuss the following cases, separately.

(1) Case $r = 2$ and p is odd: By Lemma 4.3, we may set $z = z_2$, while $\langle z_2 \rangle^f = \langle z_2 \rangle$. It follows from Proposition 2.2 and Table 1 that $\text{Fix}(\langle z_2 \rangle) = \frac{|\text{N}_G(\langle z_2 \rangle)|}{|\text{N}_M(\langle z_2 \rangle)|} = \frac{q(q^2-1)(q+1) \times 2m}{q'(q'^2-1)(q'+1) \times 2m} = \frac{q(q^2-1)(q+1)}{q'(q'^2-1)(q'+1)}$, so that

$$\begin{aligned} Q(\langle z_2 \rangle) &= \frac{|M|}{|\Omega|} \times \frac{(r-1) \times \text{Fix}(\langle z_2 \rangle)}{|\text{N}_M(\langle z_2 \rangle)|} = \frac{|M|}{|\Omega|} \frac{q(q^2-1)(q+1)}{2m[q'(q'^2-1)(q'+1)]^2} \\ &\leq \frac{13mq'^{16}}{6q^8} \frac{q^4(1+q'^{-1})}{2mq'^8(1-q'^{-2})^2} \leq \frac{793}{576q'^4(e-2)} \leq \frac{1}{26}. \end{aligned}$$

(2) Case $r = p$: By Lemma 4.3, we may set either $z = z_0$ or $z = z'_l$ where $0 \leq l \leq d' - 1$.

Suppose that $z = z_0$. Then $|\mathbf{N}_{G_1}(\langle z_0 \rangle)| = q^3(q+1)(p-1)$ and $|\mathbf{N}_{M_1}(\langle z_0 \rangle)| = q'^3(q'+1)(p-1)$. Now, we have

$$\begin{aligned} Q(\langle z_0 \rangle) &= \frac{|M|}{|\Omega|} \frac{(p-1)\text{Fix}(\langle z_0 \rangle)}{|\mathbf{N}_M(\langle z_0 \rangle)|} = \frac{|M|}{|\Omega|} \frac{(p-1)|\mathbf{N}_G(\langle z_0 \rangle)|}{|\mathbf{N}_M(\langle z_0 \rangle)|^2} \leq \frac{|M|}{|\Omega|} \frac{2m(p-1)q^3(q+1)(p-1)}{(q'^3(q'+1)(p-1))^2} \\ &\leq \frac{\frac{13m^2q'^{16}}{3q^8}q^4(1+q^{-1})}{q'^8} \leq \frac{3172m^2}{729q} \frac{1}{q'^{3e-8}} \leq \frac{1}{26}. \end{aligned}$$

For the case $e = 3$ and $q' = 4$, $|\mathbf{N}_G(\langle z_0 \rangle)| = 2m|\mathbf{N}_{G_1}(\langle z_0 \rangle)|$ and $|\mathbf{N}_M(\langle z_0 \rangle)| = 2m|\mathbf{N}_{M_1}(\langle z_0 \rangle)|$, as $p = 2$, where z_0 is of form in Table 1. So we get

$$Q(\langle z_0 \rangle) \leq \frac{|M|}{|\Omega|} \frac{(p-1)q^3(q+1)(p-1)}{(q'^3(q'+1)(p-1))^2 \times 2m} < 0.1 \times \frac{1}{26}.$$

Suppose that $z = z'_l$ where $0 \leq l \leq d' - 1$. If $d' = 3$, then three elements z'_l are conjugate to each other in M_1 and so set $z = z'_0$. This implies that

$$\begin{aligned} Q(\langle z'_0 \rangle) &= \frac{|M|}{|\Omega|} \times \frac{(p-1)\text{Fix}(\langle z'_0 \rangle)}{|\mathbf{N}_M(\langle z'_0 \rangle)|} = \frac{|M|}{|\Omega|} \frac{(p-1)|\mathbf{N}_G(\langle z'_0 \rangle)|}{|\mathbf{N}_M(\langle z'_0 \rangle)|^2} \leq \frac{13mq'^{16}}{6q^8} \frac{2m(p-1)q^2(p-1)}{(q'^2(p-1))^2} \leq \frac{13m^2}{3q} \frac{1}{q'^{5e-12}} \\ &\leq \frac{13}{81} \frac{1}{q'^{5e-12}} \leq \frac{1}{26}. \end{aligned}$$

(3) Case $r = 3$ and $r \mid q' + 1$: By Lemmas 4.2, we take $z = z_A, z_B$ and δ' . Hence, $\langle z \rangle^f = \langle z \rangle$.

$z = z_A$: $|\mathbf{N}_G(\langle z_A \rangle)| = 6ms(q+1)^2$ and $|\mathbf{N}_M(\langle z_A \rangle)| = 6ms'(q'+1)^2$, where $s, s' \in \{1, 2\}$. Hence,

$$Q(\langle z_A \rangle) = \frac{|M|}{|\Omega|} \frac{2\text{Fix}(\langle z_A \rangle)}{|\mathbf{N}_M(\langle z_A \rangle)|} = \frac{|M|}{|\Omega|} \frac{2|\mathbf{N}_G(\langle z_A \rangle)|}{|\mathbf{N}_M(\langle z_A \rangle)|^2} \leq \frac{13mq'^{16}}{6q^8} \frac{12ms(q+1)^2}{(6ms'(q'+1)^2)^2} \leq \frac{3172}{2187} \frac{1}{q'^{6(e-2)}} \leq \frac{1}{26}.$$

$z = z_B$: $|\mathbf{N}_G(\langle z_B \rangle)| = 2mq(q+1)^2(q-1)$ and $|\mathbf{N}_M(\langle z_B \rangle)| = 2mq'(q'+1)^2(q'-1)$. Therefore, we have

$$Q(\langle z_B \rangle) = \frac{|M|}{|\Omega|} \times \frac{2\text{Fix}(\langle z_B \rangle)}{|\mathbf{N}_M(\langle z_B \rangle)|} = \frac{|M|}{|\Omega|} \frac{2|\mathbf{N}_G(\langle z_B \rangle)|}{|\mathbf{N}_M(\langle z_B \rangle)|^2} \leq \frac{13mq'^{16}}{6q^8} \frac{4mq(q+1)^2(q-1)}{(2mq'(q'+1)^2(q'-1))^2} < \frac{5}{q'^{4(e-2)}} \leq \frac{1}{26}.$$

$z = \delta'$: Since $c = 1$, we get that $\delta' = \delta$. Then $|\mathbf{N}_G(\langle \delta' \rangle)| = |(\mathbb{Z}_{q^2-q+1} : \mathbb{Z}_3)| |\langle f \rangle| = 6m(q^2 - q + 1)$ and $|\mathbf{N}_M(\langle \delta' \rangle)| = 6m(q'^2 - q' + 1)$. Consequently, we have

$$Q(\langle \delta' \rangle) = \frac{|M|}{|\Omega|} \times \frac{2\text{Fix}(\langle \delta' \rangle)}{|\mathbf{N}_M(\langle \delta' \rangle)|} = \frac{|M|}{|\Omega|} \frac{2|\mathbf{N}_G(\langle \delta' \rangle)|}{|\mathbf{N}_M(\langle \delta' \rangle)|^2} \leq \frac{13mq'^{16}}{6q^8} \frac{12m(q^2 - q + 1)}{(6m(q'^2 - q' + 1))^2} \leq \frac{13}{8} \frac{1}{q'^{6(e-2)}} \leq \frac{1}{26}.$$

(4) Case $r \mid q' - 1$ and $r \neq 2$: By Lemma 4.3, we may set $z = z_{q'-1}$. Hence,

$$\begin{aligned} (q'-1)Q(\langle z_{q'-1} \rangle) &= (q'-1) \times \frac{|M|}{|\Omega|} \times \frac{(r-1)\text{Fix}(\langle z_{q'-1} \rangle)}{|\mathbf{N}_M(\langle z_{q'-1} \rangle)|} = (q'-1) \times \frac{|M|}{|\Omega|} \times \frac{(r-1)|\mathbf{N}_G(\langle z_{q'-1} \rangle)|}{|\mathbf{N}_M(\langle z_{q'-1} \rangle)|^2} \\ &= \frac{(q'-1)|M|}{|\Omega|} \times \frac{(r-1)2(q^2-1) \times 2m}{[2(q'^2-1) \times 2m]^2} \leq \frac{13mq'^{18}}{6q^8} \frac{q^2}{4mq'^2(1-q'^{-2})^2} \leq \frac{351}{512q'^{6e-16}} \leq \frac{1}{26}. \end{aligned}$$

(5) Case $r \mid q' + 1$ and $r \neq 2, 3$: By Lemma 4.3, we take $z = z_{q'+1}^A(a)$ and $z = z_{q'+1}^B$.

$z = z_{q'+1}^A(a)$: By the proof in Lemma 4.3, we get that if $z_{q'+1}^A(a_1)$ and $z_{q'+1}^A(a_2)$ are not conjugate in M_1 for some integers a_1 and a_2 , then they are also not conjugate in G_1 .

Since $\langle z_{q'+1}^A(a) \rangle^f = \langle z_{q'+1}^A(a) \rangle$, by Lemma 4.3, we have $|\mathrm{N}_G(\langle z_{q'+1}^A(a) \rangle)| = s(q+1)^2 \times 2m$ and $|\mathrm{N}_M(\langle z_{q'+1}^A(a) \rangle)| = s'(q'+1)^2 \times 2m$, where $s, s' \mid 6$. Therefore, we have

$$\begin{aligned} \frac{(q'+1)^2}{2} Q(\langle z_{q'+1}^A(a) \rangle) &= \frac{(q'+1)^2}{2} \times \frac{|M|}{|\Omega|} \times \frac{(r-1)\mathrm{Fix}(\langle z_{q'+1}^A(a) \rangle)}{|\mathrm{N}_M(\langle z_{q'+1}^A(a) \rangle)|} = \frac{(q'+1)^2}{2} \times \frac{|M|}{|\Omega|} \frac{(r-1)s(q+1)^2}{s'(q'+1)^2 s'(q'+1)^2 \times 2m} \\ &\leq \frac{13mq'^{16}}{12q^8} \frac{3(r-1)(q+1)^2}{m(q'+1)^2} \leq \frac{23q'^{15}}{7q^6} \leq \frac{1}{26} \quad (\text{as } r-1 \leq q'). \end{aligned}$$

$z = z_{q'+1}^B$: The number of conjugate classes of type $z_{q'+1}^B$ is only one. Since $\langle z_{q'+1}^B \rangle^f = \langle z_{q'+1}^B \rangle$, by Lemma 4.3, we have $|\mathrm{N}_G(\langle z_{q'+1}^B \rangle)| = q(q-1)(q+1)^2 \times 2m$ and $|\mathrm{N}_M(\langle z_{q'+1}^B \rangle)| = q'(q'-1)(q'+1)^2 \times 2m$. Noting $q' \geq 9$ and $r-1 \leq q'$ in this case, we have

$$\begin{aligned} (q'+1)Q(\langle z_{q'+1}^B \rangle) &= \frac{(q'+1)|M|}{|\Omega|} \times \frac{(r-1)|\mathrm{Fix}(z_{q'+1}^B)|}{|\mathrm{N}_M(\langle z_{q'+1}^B \rangle)|} = \frac{(q'+1)|M|}{|\Omega|} \times \frac{(r-1)q(q-1)(q+1)^2}{2m[q'(q'-1)(q'+1)^2]^2} \\ &\leq \frac{13mq'^{16}}{6q^8} \times \frac{(r-1)q^4(1+q^{-1})^2}{2mq'^7(1-q'^{-1})^2} \leq \frac{5q'^{10}}{2q^4} \leq \frac{1}{26}. \end{aligned}$$

(6) Case $r \mid q'^2 - q' + 1$ and $r \neq 3$: By Lemma 4.3, we may set $z = z_{q'^2-q'+1}$. Note that there is only one conjugate class subgroup with order r , and $\mathrm{Fix}(\langle z_{q'^2-q'+1} \rangle) = \frac{|\mathrm{N}_G(\langle z_{q'^2-q'+1} \rangle)|}{|\mathrm{N}_M(\langle z_{q'^2-q'+1} \rangle)|}$, $|\mathrm{N}_{G_1}(\langle z_{q'^2-q'+1} \rangle)| = 3(q^2 - q + 1)$ and $|\mathrm{N}_{M_1}(\langle z_{q'^2-q'+1} \rangle)| = 3(q'^2 - q' + 1)$ from Proposition 2.2 and Table 1. We now discuss the cases of $e = 3$ and $e \geq 5$ separately.

$e = 3$: Since $q = q'^3$ and $q'^2 - q' + 1 \mid q^2 - 1$, the element $z_{q'^2-q'+1}$ can be represented by a diagonal matrix of order r . Hence, we conclude that $\mathrm{Fix}(\langle z_{q'^2-q'+1} \rangle) = \frac{2m|\mathrm{N}_{G_1}(\langle z_{q'^2-q'+1} \rangle)|}{2m|\mathrm{N}_{M_1}(\langle z_{q'^2-q'+1} \rangle)|} = \frac{q^2 - q + 1}{q'^2 - q' + 1}$, due to $\langle z_{q'^2-q'+1} \rangle^f = \langle z_{q'^2-q'+1} \rangle$.

If $e = 3$ and $p = 2$, then $m' \neq 3$, as $c = 1$. For $(e, p, m') \in \{(3, 2, 2), (3, 2, 4), (3, 3, 2)\}$, there is only one conjugacy classes subgroup of order $r \mid q'^2 - q' + 1$, as $q'^2 - q' + 1$ is a prime. This implies that $Q(\langle z_{q'^2-q'+1} \rangle) = \frac{|M|}{|\Omega|} \frac{(r-1)\mathrm{Fix}(\langle z_{q'^2-q'+1} \rangle)}{|\mathrm{N}_M(\langle z_{q'^2-q'+1} \rangle)|} \leq \frac{|M|}{|\Omega|} \frac{(r-1)(q^2 - q + 1)}{3(q'^2 - q' + 1)^2} \leq \frac{1}{26}$, by inserting the these values of e, p, m' . And for other cases, we have

$$\begin{aligned} (q'^2 - q' + 1)Q(\langle z_{q'^2-q'+1} \rangle) &= \frac{(q'^2 - q' + 1)|M|}{|\Omega|} \times \frac{(r-1)\mathrm{Fix}(\langle z_{q'^2-q'+1} \rangle)}{|\mathrm{N}_M(\langle z_{q'^2-q'+1} \rangle)|} \\ &\leq \frac{|M|(q'^2 - q' + 1)}{|\Omega|} \frac{(r-1)(q^2 - q + 1)}{3(q'^2 - q' + 1)^2} \leq \frac{13mq'^{16}}{6q^8} \frac{(q^2 - q + 1)}{3} \leq \frac{13m'e}{18p^{6m'e-16m'}} \leq \frac{1}{26}. \end{aligned}$$

$e \geq 5$: We obtain that

$$\begin{aligned} (q'^2 - q' + 1)Q(\langle z_{q'^2-q'+1} \rangle) &= \frac{(q'^2 - q' + 1)|M|}{|\Omega|} \times \frac{(r-1)\mathrm{Fix}(\langle z_{q'^2-q'+1} \rangle)}{|\mathrm{N}_M(\langle z_{q'^2-q'+1} \rangle)|} \\ &\leq \frac{|M|(q'^2 - q' + 1)}{|\Omega|} \frac{(r-1)(q^2 - q + 1) \times 2m}{3(q'^2 - q' + 1)^2} \leq \frac{13mq'^{16}}{6q^8} \frac{(q^2 - q + 1) \times 2m}{3} \leq \frac{26m'^2e^2}{18p^{6m'e-16m'}} \leq \frac{1}{26}. \end{aligned} \quad \square$$

Lemma 5.2 *Let $c = 1$ and either $e \geq 5$ and $q' \geq 3$ or $e = 3$ and $q' \geq 4$. Then every cyclic subgroup $\langle z \rangle$ of prime order r of M not contained in M_1 is M -conjugate to one of the following groups.*

(I) *Cases either $e \geq 5$ and $q' \geq 3$ or $e = 3$ and $q' \geq 5$:*

$$(1) \quad r \notin \{e, 2\}: z = f^{\frac{2m}{r}} \text{ with } Q(\langle f^{\frac{2m}{r}} \rangle) < \frac{1}{26m'}.$$

$$(2) \quad r = 2: z = f^m \text{ with } Q(\langle f^m \rangle) < \frac{1}{26}.$$

(3) $r = e$:

- (3.1) $r \nmid |M_1| : z = f^{\frac{2m}{r}}$ with $Q(z) < \frac{1}{26}$;
- (3.2) $r = p : z = z_0 f^{\frac{2m}{r}}$ or $z'_0 f^{\frac{2m}{r}}$ with $Q(z_0 f^{\frac{2m}{r}}) + Q(z'_0 f^{\frac{2m}{r}}) < \frac{1}{26}$;
- (3.3) $r \mid q'+1$ and $r \neq 3 : z = f^{\frac{2m}{r}}$, $(z_{q'+1}^A(a))^i f^{\frac{2m}{r}}$ where $2 \leq a \leq \frac{r-1}{2}$ and $1 \leq i \leq r-1$,
or $(z_{q'+1}^B)^i f^{\frac{2m}{r}}$ with $1 \leq i \leq r-1$, and $Q(\langle f^{\frac{2m}{r}} \rangle) + \sum_{i=1}^{r-1} \sum_{a=2}^{\frac{r-1}{2}} Q(\langle (z_{q'+1}^A(a))^i f^{\frac{2m}{r}} \rangle) + \sum_{i=1}^{r-1} Q(\langle (z_{q'+1}^B)^i f^{\frac{2m}{r}} \rangle) < \frac{1}{26}$;
- (3.4) $r \mid q' - 1 : z_{q'-1}^i f^{\frac{2m}{r}}$ where $0 \leq i \leq r-1$, and $\sum_{i=0}^{r-1} Q(\langle (z_{q'-1})^i f^{\frac{2m}{r}} \rangle) < \frac{1}{26}$;
- (3.5) $r \mid (q'^2 - q' + 1) : z = z_{q'^2 - q' + 1}^i f^{\frac{2m}{r}}$, $0 \leq i \leq r-1$ and $\sum_{i=0}^{r-1} Q(\langle (z_{q'^2 - q' + 1})^i f^{\frac{2m}{r}} \rangle) < \frac{1}{26}$.

(II) Case $e = 3$ and $q' = 4$:

(1) $r = 2 : z = f^m$ with $Q(\langle f^m \rangle) < \frac{1}{20} \times \frac{1}{26}$.

(2) $r = 3 : z_{q'-1}^i f^{\frac{2m}{r}}$ where $0 \leq i \leq 2$, and $\sum_{i=0}^{r-1} Q(\langle (z_{q'-1})^i f^{\frac{2m}{r}} \rangle) < 4 \times \frac{1}{26}$.

Proof Let $u \in M \setminus M_1$. Then $z = u_1 f_1$ for some element $u_1 \in M_1$ and $f_1 \in \langle f \rangle$. Since $|z| = r$, we get $|f_1| = r$ and so we may set $z = u_1 f'$, where $f' := f^{\frac{2m}{r}}$. In what follows, we discuss three cases, separately. Similar to last lemma, we assume either $e \geq 5$ and $q' \geq 3$ or $e = 3$ and $q' \geq 7$, while the conclusion for $e = 3$ and $q' = 4$, that is (II) in the lemma, can be read from the proof of (I) directly.

(1) Case $r \notin \{e, 2\}$:

Suppose $r \notin \{e, 2\}$. Then by Proposition 2.1, we may set $u_1 = 1$ so that $z = f^{\frac{2m}{r}}$. Set $m = m'e = m_1re$ so that $\text{Aut}(\text{PSU}(3, q')) = \text{PGU}(3, q') \rtimes \langle f^e \rangle$. By Proposition 2.1, there is only one $\text{PGU}(3, q')$ -class $\langle f^{2m_1e} \rangle$, which is split into $(3, \frac{q'+1}{q'^{\frac{1}{r}}+1})$ distinct $\text{PSU}(3, q')$ -classes by [2, Proposition 3.3.12]. Since $N_M(\langle f^{\frac{2m}{r}} \rangle) = \text{PGU}(3, q'^{\frac{1}{r}}) \rtimes \langle f \rangle$ and $N_G(\langle f^{\frac{2m}{r}} \rangle) = \text{PGU}(3, q'^{\frac{1}{r}}) \rtimes \langle f \rangle$, we have

$$\begin{aligned} \text{Fix}(\langle f^{\frac{2m}{r}} \rangle) &= \frac{|\text{PGU}(3, q'^{\frac{1}{r}}) \rtimes \langle f \rangle|}{|\text{PGU}(3, q'^{\frac{1}{r}}) \rtimes \langle f \rangle|} = \frac{p^{3m_1e}(p^{3m_1e}+1)(p^{2m_1e}-1)}{p^{3m_1}(p^{3m_1}+1)(p^{2m_1}-1)}, \\ m'Q(\langle f^{\frac{2m}{r}} \rangle) &= \frac{m'|\Omega|}{|\Omega|} \frac{(r-1)\text{Fix}(\langle f^{\frac{2m}{r}} \rangle)}{|\text{N}_M(\langle f^{\frac{2m}{r}} \rangle)|} = \frac{m'|\Omega|}{|\Omega|} \frac{(r-1)q^{\frac{3}{r}}(q^{\frac{3}{r}}+1)(q^{\frac{2}{r}}-1)}{[q'^{\frac{3}{r}}(q'^{\frac{3}{r}}+1)(q'^{\frac{2}{r}}-1)]^2 \times 2m} \\ &\leq \frac{13m'mq'^{16}}{6q^8} \frac{2(r-1)q^{\frac{8}{r}}}{[q'^{\frac{8}{r}}(1-2^{-2})]^2 \times 2m} \leq \frac{104}{27q^{\frac{16e-38}{3}}} \leq \frac{1}{26} \text{ (as } r \geq 3, r-1 \leq q', m' \leq q' \text{ and } p \geq 2\text{).} \end{aligned}$$

(2) Case $r = 2$:

By Proposition 2.1 again, we take $z = f^m$. By [2, Proposition 3.3.15] we have $C_{M_1}(\langle f^m \rangle) \cong \text{PGO}(3, q')$ and $C_{G_1}(\langle f^m \rangle) \cong \text{PGO}(3, q)$. Hence, we have

$$\begin{aligned}\text{Fix}(\langle f^m \rangle) &= \frac{|\text{N}_G(\langle f^m \rangle)|}{|\text{N}_M(\langle f^m \rangle)|} = \frac{|\text{PGO}(3, q) \rtimes \langle f \rangle|}{|\text{PGO}(3, q') \rtimes \langle f \rangle|} = \frac{q(q^2-1)}{q'(q'^2-1)}, \\ Q(\langle f^m \rangle) &= \frac{|M|}{|\Omega|} \frac{\text{Fix}(\langle f^m \rangle)}{|\text{N}_M(\langle f^m \rangle)|} = \frac{|M|}{|\Omega|} \frac{q(q^2-1)}{2m[q'(q'^2-1)]^2} \leq \frac{13mq'^{16}}{6q^8} \frac{q^3}{2mq'^6(1-q'^{-2})^2} \leq \frac{351}{256q'^{5e-10}} \not\leq \frac{1}{26}.\end{aligned}$$

(3) Case $r = e$:

Suppose $r = e$. Then we may set $u = u_1 f^{\frac{2m}{e}} = u_1 f^{2m'}$, where $u_1 \in M_1$. Now we have $[x, f^{2m'}] = 1$ for any $x \in M_1$ and so $|u_1| = r$. Recalling $|M_1| = q'^3(q'^3+1)(q'^2-1)$, our prime r has the following possibilities: $r \nmid |M_1|$, $r = p$, $r \mid q' + 1$, $r \mid q' - 1$ and $r \mid q'^2 - q' + 1$, which are discussed separately.

(3.1) $r \nmid |M_1|$

Up to conjugate in M , set $z = f^{2m'}$. Hence, $\text{N}_G(\langle z \rangle) = \text{N}_M(\langle z \rangle) = M$ and

$$Q(\langle z \rangle) = \frac{|M|}{|\Omega|} \frac{(r-1) \times 1}{|M|} \leq \frac{13mq'^{16}}{6q^8} \frac{e}{2mq'^3(q'^3+1)(q'^2-1)} \leq \frac{39eq'^8}{32q^8} \not\leq \frac{1}{26} \text{ (as } r-1 \leq e\text{).}$$

(3.2) $r = p$

Since there are two M_1 -conjugacy classes of elements of order p , with the representatives z_0 and z'_0 , we set $\langle z \rangle = \langle f^{2m'} \rangle$, $\langle z_0 f^{2m'} \rangle$ and $\langle z'_0 f^{2m'} \rangle$. By Proposition 2.1, both of them are conjugate to $\langle f^{2m'} \rangle$ in G_1 . This implies that $|\text{N}_G(\langle z_0 f^{2m'} \rangle)| = |\text{N}_G(\langle z'_0 f^{2m'} \rangle)| = |\text{N}_G(\langle f^{2m'} \rangle)| = |M|$. By Lemma 4.3, we conclude that $|\text{C}_{M_1}(\langle z_0 \rangle)| = q'^3(q' + 1)$ and $|\text{C}_{M_1}(\langle z'_0 \rangle)| = q'^2$. It deduces that $|\text{N}_M(\langle z_0 f^{2m'} \rangle)| = |\text{C}_M(\langle z_0 \rangle)| \geq eq'^3(q' + 1)$ and $|\text{N}_M(\langle z'_0 f^{2m'} \rangle)| = |\text{C}_M(\langle z'_0 \rangle)| \geq eq'^2$. Therefore, we have

$$\begin{aligned}\text{Fix}(\langle z_0 f^{2m'} \rangle) &= \text{Fix}(\langle z'_0 f^{2m'} \rangle) = \text{Fix}(\langle f^{2m'} \rangle) \\ &= \frac{|\text{N}_G(\langle z_0 f^{2m'} \rangle)|}{|\text{N}_M(\langle z_0 f^{2m'} \rangle)|} + \frac{|\text{N}_G(\langle z'_0 f^{2m'} \rangle)|}{|\text{N}_M(\langle z'_0 f^{2m'} \rangle)|} + \frac{|\text{N}_G(\langle f^{2m'} \rangle)|}{|\text{N}_M(\langle f^{2m'} \rangle)|} \\ &\leq \frac{|M|}{eq'^3(q'+1)} + \frac{|M|}{eq'^2} + \frac{|M|}{|M|} = 2m'(q'^3+1)(q'-1)[1+q'(q'+1)]+1, \\ Q(\langle f^{2m'} \rangle) &= \frac{|M|}{|\Omega|} \frac{(p-1)\text{Fix}(\langle f' \rangle)}{|\text{N}_M(\langle f' \rangle)|} \leq \frac{|M|}{|\Omega|} \left(\frac{1+q'^2+q'}{q'^4} + \frac{p-1}{2mq'^6(q'^2-1)} \right) \leq \frac{|M|}{|\Omega|} \left(\frac{5}{3q'^2} + \frac{9}{16mq'^7} \right), \\ Q(\langle z_0 f^{2m'} \rangle) &= \frac{|M|}{|\Omega|} \frac{(p-1)\text{Fix}(\langle z_0 f' \rangle)}{|\text{N}_M(\langle z_0 f' \rangle)|} \leq \frac{|M|}{|\Omega|} \left(\frac{2m'(q'^2-q'+1)(1+q'^2+q')}{q'^2} + \frac{1}{q'^4} \right) \leq \frac{|M|}{|\Omega|} (10m'q'^2 + \frac{1}{q'^4}), \\ Q(\langle z'_0 f^{2m'} \rangle) &= \frac{|M|}{|\Omega|} \frac{(p-1)\text{Fix}(\langle z'_0 f' \rangle)}{|\text{N}_M(\langle z'_0 f' \rangle)|} \leq \frac{|M|}{|\Omega|} \left(\frac{2m'(q'^3+1)(d'+q'^2+q')}{q'} + \frac{1}{q'^2} \right) \leq \frac{|M|}{|\Omega|} \left(\frac{280m'q'^4}{81} + \frac{1}{q'^2} \right), \\ Q(\langle f' \rangle) + Q(\langle z_0 f' \rangle) + Q(\langle z'_0 f' \rangle) &\leq \frac{|M|}{|\Omega|} \left(\frac{8}{3q'^2} + \frac{1}{q'^4} + \frac{9}{16mq'^7} + 10m'q'^2 + \frac{280m'q'^4}{81} \right) \\ &\leq \frac{52m'pq'^{14}}{9q^8} + \frac{13m'pq'^{12}}{2q^8} + \frac{39q'^9}{32q^8} + \frac{65m'^2pq'^{18}}{3q^8} + \frac{1820m'^2pq'^{20}}{243q^8} \\ &\leq \frac{52q'^{15}}{9q^8} + \frac{13q'^{13}}{2q^8} + \frac{39q'^9}{32q^8} + \frac{65q'^{19}}{3q^8} + \frac{1820q'^{21}}{243q^8} \not\leq \frac{1}{26} \text{ (as } r = e = p \geq 3\text{).}\end{aligned}$$

(3.3) $r \mid q' + 1$

If $e = 3$, then $r = 3$ and so $3 \mid (q' + 1)$. Thus, $c = (\frac{q+1}{q'+1}, 3) = (q'^2 - q' + 1, 3) = 3$, a contradiction. Therefore, $e \geq 5$.

Let $z = u_0 f^{\frac{2m}{r}}$, where $u_0 \in M_1$. Since every subgroup of order r where $r \mid q' + 1$ of M_1 is conjugate to $\langle z_{q'+1}^A(a) \rangle$ where $2 \leq a \leq \frac{r-1}{2}$ or $\langle z_{q'+1}^B \rangle$, we may set $\langle z \rangle = \langle f^{\frac{2m}{r}} \rangle$, $\langle (z_{q'+1}^A(a))^i f^{\frac{2m}{r}} \rangle$ or $\langle (z_{q'+1}^B)^i f^{\frac{2m}{r}} \rangle$, where $1 \leq i \leq r-1$. By Proposition 2.1, all the elements $u_0 f^{\frac{2m}{r}}$ of order r in M are G_1 -conjugate to $f^{\frac{2m}{r}}$. Hence, $|\text{N}_G(\langle (z_{q'+1}^A(a))^i f' \rangle)| = |\text{N}_G(\langle (z_{q'+1}^B)^i f' \rangle)| = |\text{N}_G(\langle f' \rangle)| = |M|$. By Table 1, we get that

$$|\mathrm{N}_M(\langle\langle z_{q'+1}^A(a))^i f' \rangle)| = |\mathrm{C}_M(\langle\langle z_{q'+1}^A(a))^i \rangle)| = |\mathrm{C}_{M_1}(\langle\langle z_{q'+1}^A(a))^i \rangle)|e = (q'+1)^2e,$$

$$|\mathrm{N}_M(\langle\langle z_{q'+1}^B(a))^i f' \rangle)| = |\mathrm{C}_M(\langle\langle z_{q'+1}^B(a))^i \rangle)| = |\mathrm{C}_{M_1}(\langle\langle z_{q'+1}^B(a))^i \rangle)|e = q'(q'+1)^2(q'-1)e.$$

Now, we conclude that

$$\begin{aligned} \mathrm{Fix}(\langle f' \rangle) &= \mathrm{Fix}(\langle\langle z_{q'+1}^A(a))^i f' \rangle) = \mathrm{Fix}(\langle\langle z_{q'+1}^B(a))^i f' \rangle) \\ &\leq \frac{(r-3)(r-1)}{2} \frac{|\mathrm{N}_G(\langle\langle z_{q'+1}^A(a))^i f' \rangle)|}{|\mathrm{N}_M(\langle\langle z_{q'+1}^A(a))^i f' \rangle)|} + (r-1) \frac{|\mathrm{N}_G(\langle\langle z_{q'+1}^B(a))^i f' \rangle)|}{|\mathrm{N}_M(\langle\langle z_{q'+1}^B(a))^i f' \rangle)|} + \frac{|\mathrm{N}_G(\langle f' \rangle)|}{|\mathrm{N}_M(\langle f' \rangle)|} \\ &\leq \frac{(r-3)(r-1)}{2} \frac{|M|}{(q'+1)^2e} + (r-1) \frac{|M|}{q'(q'+1)^2(q'-1)e} + \frac{|M|}{|M|} \leq emq'^6 + 2mq'^4 + 1; \\ Q(\langle f' \rangle) &= \frac{|M|}{|\Omega|} \frac{(r-1)\mathrm{Fix}(\langle f' \rangle)}{|\mathrm{N}_M(\langle f' \rangle)|} \leq \frac{13mq'^{16}}{6q^8} \frac{r-1}{2mq'^3(q'^3+1)(q'^2-1)} (emq'^6 + 2mq'^4 + 1) \\ &\leq \frac{39e^2mq'^{14}}{32q^8} + \frac{39emq'^{12}}{16q^8} + \frac{39eq'^8}{32q^8}; \\ Q(\langle\langle z_{q'+1}^A(a))^i f' \rangle) &= \frac{|M|}{|\Omega|} \frac{(r-1)\mathrm{Fix}(\langle\langle z_{q'+1}^A(a))^i f' \rangle)}{|\mathrm{N}_M(\langle\langle z_{q'+1}^A(a))^i f' \rangle)|} \leq \frac{13mq'^{16}}{6q^8} \frac{r-1}{(q'+1)^2e} (emq'^6 + 2mq'^4 + 1) \\ &\leq \frac{13em^2q'^{20}}{6q^8} + \frac{13m^2q'^{18}}{3q^8} + \frac{13mq'^{14}}{6q^8}; \\ Q(\langle\langle z_{q'+1}^B(a))^i f' \rangle) &= \frac{|M|}{|\Omega|} \frac{(r-1)\mathrm{Fix}(\langle\langle z_{q'+1}^B(a))^i f' \rangle)}{|\mathrm{N}_M(\langle\langle z_{q'+1}^B(a))^i f' \rangle)|} \leq \frac{13mq'^{16}}{6q^8} \frac{r-1}{q'(q'+1)^2(q'-1)e} (emq'^6 + 2mq'^4 + 1) \\ &\leq \frac{13em^2q'^{18}}{4q^8} + \frac{13m^2q'^{16}}{2q^8} + \frac{13mq'^{12}}{4q^8}; \\ Q(\langle f' \rangle) + \sum_{i=1}^{r-1} \sum_{a=2}^{\frac{r-1}{2}} Q(\langle\langle z_{q'+1}^A(a))^i f' \rangle) + \sum_{i=1}^{r-1} Q(\langle\langle z_{q'+1}^B(a))^i f' \rangle) &\leq \frac{39e^2mq'^{14}}{32q^8} + \frac{39emq'^{12}}{16q^8} + \frac{39eq'^8}{32q^8} \\ &+ \frac{(r-1)(r-3)}{2} \left(\frac{13em^2q'^{20}}{6q^8} + \frac{13m^2q'^{18}}{3q^8} + \frac{13mq'^{14}}{6q^8} \right) + (r-1) \left(\frac{13em^2q'^{18}}{4q^8} + \frac{13m^2q'^{16}}{2q^8} + \frac{13mq'^{12}}{4q^8} \right) \\ &\leq \frac{221e^2mq'^{14}}{96q^8} + \frac{91emq'^{12}}{16q^8} + \frac{65e^2m^2q'^{18}}{12q^8} + \frac{13e^3m^2q'^{20}}{12q^8} + \frac{13em^2q'^{16}}{2q^8} + \frac{39eq'^8}{32q^8} \\ &\leq \frac{221 \times 5^3 q'^{14}}{96 \times 2^5 q^7} + \frac{91 \times 5^2 q'^{12}}{16 \times 2^5 q^7} + \frac{65 \times 5^4 q'^{18}}{12 \times 2^5 q^7} + \frac{13 \times 5^5 q'^{20}}{12 \times 2^5 q^7} + \frac{39 \times 5 q'^8}{2^6 q^7} + \frac{39 \times 5 q'^8}{32 \times 2^5 q^7} \not\leq \frac{1}{26}. \end{aligned}$$

(3.4) $r \mid q' - 1$

Let $z = u_0 f^{\frac{2m}{r}}$, where $u_0 \in M_1$. Since every subgroup of order r where $r \mid q' - 1$ of M_1 is conjugate to $\langle z_{q'-1} \rangle$, we set $\langle z \rangle = \langle z_{q'-1}^i f^{\frac{2m}{r}} \rangle$ where $0 \leq i \leq r-1$. By Proposition 2.1, all the elements $uf' \in M$ with order r are G_1 -conjugate to f' . Hence, $|\mathrm{N}_G(\langle z_{q'-1}^i f^{\frac{2m}{r}} \rangle)| = |\mathrm{N}_G(\langle f^{\frac{2m}{r}} \rangle)| = |M|$. By the arguments above and Table 1, we get that $|\mathrm{N}_M(\langle z_{q'-1}^i f' \rangle)| = |\mathrm{C}_M(\langle z_{q'-1} \rangle)| = 2e|\mathrm{C}_{M_1}(\langle z_{q'-1} \rangle)| = 2e(q'^2-1)$, as $z_{q'-1} \in \{h(k) \mid k \in \mathbb{F}_{q'}\}$. Hence,

$$\begin{aligned} \mathrm{Fix}(\langle f' \rangle) &= \mathrm{Fix}(\langle z_{q'-1}^i f' \rangle) \leq \frac{|\mathrm{N}_G(\langle f' \rangle)|}{|\mathrm{N}_M(\langle f' \rangle)|} + \sum_{i=1}^{r-1} \frac{|\mathrm{N}_G(\langle f' \rangle)|}{|\mathrm{N}_M(\langle z_{q'-1}^i f' \rangle)|} \\ &\leq \frac{|M|}{|M|} + (r-1) \frac{|M|}{2e(q'^2-1)} = 1 + (e-1) \frac{2mq'^3(q'^3+1)(q'^2-1)}{2e(q'^2-1)} \leq 1 + mq'^3(q'^3+1); \end{aligned}$$

$$\begin{aligned} Q(\langle f' \rangle) &= \frac{|M|}{|\Omega|} \frac{(r-1)\mathrm{Fix}(\langle f' \rangle)}{|\mathrm{N}_M(\langle f' \rangle)|} \leq \frac{13mq'^{16}}{6q^8} \frac{(e-1)[1+mq'^3(q'^3+1)]}{2mq'^3(q'^3+1)(q'^2-1)} \leq \frac{39eq'^8}{32q^8} + \frac{39meq'^{14}}{32q^8}; \\ Q(\langle z_{q'-1}^i f' \rangle) &= \frac{|M|}{|\Omega|} \frac{(r-1)\mathrm{Fix}(\langle z_{q'-1}^i f' \rangle)}{|\mathrm{N}_M(\langle z_{q'-1}^i f' \rangle)|} \leq \frac{13mq'^{16}}{6q^8} \frac{(r-1)[1+mq'^3(q'^3+1)]}{2e(q'^2-1)} \leq \frac{39mq'^{14}}{32q^8} + \frac{91m^2q'^{20}}{72q^8}; \\ \sum_{i=0}^{r-1} Q(\langle z_{q'-1}^i f' \rangle) &\leq \frac{39q'^9}{32q^8} + \frac{78emq'^{14}}{32q^8} + \frac{91em^2q'^{20}}{72q^8} \not\leq \frac{1}{26}. \end{aligned}$$

(3.5) $r \mid q'^2 - q' + 1$

If $e = 3$, then it follows from $1 = c = (\frac{q+1}{q'+1}, 3) = (q'^2 - q' + 1, 3)$ that $r = e \neq 3$. Therefore, $e \geq 5$.

Set $z = u_0 f^{\frac{2m}{r}}$, where $u_0 \in M_1$. Since M_1 has only one class of subgroups $\mathbb{Z}_{q'^2-q'+1}$, we may set $u_0 = z_{q'^2-q'+1}^i$, where $0 \leq i \leq r-1$. By Proposition 2.1, all these elements $z_{q'^2-q'+1}^i f^{2m'} \in M$ are G_1 -conjugate to f' . Hence, $|\mathrm{N}_G(\langle z_{q'^2-q'+1}^i f^{2m'} \rangle)| = |\mathrm{N}_G(\langle f^{2m'} \rangle)| = |M|$. By Table 1, we get that

$$|\mathrm{N}_M(\langle z_{q'^2-q'+1}^i f^{2m'} \rangle)| = |\mathrm{C}_M(\langle z_{q'^2-q'+1}^i \rangle)| = |\mathrm{C}_{M_1}(\langle z_{q'^2-q'+1}^i \rangle)|e = (q'^2 - q' + 1)e.$$

Therefore, we have

$$\begin{aligned} \mathrm{Fix}(\langle f^{2m'} \rangle) &= \mathrm{Fix}(\langle z_{q'^2-q'+1}^i f^{2m'} \rangle) \leq \frac{|\mathrm{N}_G(\langle f' \rangle)|}{|\mathrm{N}_M(\langle f' \rangle)|} + (r-1) \frac{|\mathrm{N}_G(\langle f^{2m'} \rangle)|}{|\mathrm{N}_M(\langle z_{q'^2-q'+1}^i f^{2m'} \rangle)|} \\ &\leq \frac{|M|}{|M|} + (r-1) \frac{|M|}{(q'^2-q'+1)e} = 1 + (e-1) \frac{2mq'^3(q'^3+1)(q'^2-1)}{(q'^2-q'+1)e} \leq 1 + 2mq'^3(q'+1)(q'^2-1), \\ Q(\langle f' \rangle) &= \frac{|M|}{|\Omega|} \frac{(r-1)|\mathrm{Fix}(\langle f' \rangle)|}{|\mathrm{N}_M(\langle f' \rangle)|} \leq \frac{13mq'^{16}}{6q^8} \frac{(r-1)[1+2mq'^3(q'+1)(q'^2-1)]}{|M|} \leq \frac{13mq'^{16}}{6q^8} \left(\frac{9}{16m'q^8} + \frac{3e}{2q'^2} \right) \\ &= \frac{39eq'^8}{32q^8} + \frac{13mq'^{14}}{4q^8}, \\ Q(\langle z_{q'^2-q'+1}^i f' \rangle) &= \frac{|M|}{|\Omega|} \frac{(r-1)|\mathrm{Fix}(\langle z_{q'^2-q'+1}^i f' \rangle)|}{|\mathrm{N}_M(\langle z_{q'^2-q'+1}^i f' \rangle)|} \leq \frac{13mq'^{16}}{6q^8} \frac{(r-1)[1+2mq'^3(q'+1)(q'^2-1)]}{(q'^2-q'+1)e} \\ &= \frac{13mq'^{16}}{6q^8} \left(\frac{1}{q'^2(1-q'^{-1})} + \frac{2mq'^4(1+q'^{-1})^2}{q'^2(1-q'^{-1})} \right) \leq \frac{13mq'^{14}}{4q^8} + \frac{104m^2q'^{18}}{9q^8}, \\ \sum_{i=0}^{r-1} Q(\langle z_{q'^2-q'+1}^i f' \rangle) &\leq \frac{39eq'^8}{32q^8} + \frac{13mq'^{14}}{4q^8} + (r-1) \left(\frac{13mq'^{14}}{4q^8} + \frac{104m^2q'^{18}}{9q^8} \right) \\ &\leq \frac{39eq'^8}{32q^8} + \frac{13emq'^{14}}{4q^8} + \frac{13emq'^{14}}{4q^8} + \frac{104em^2q'^{18}}{9q^8} \leq \frac{1}{26}. \end{aligned}$$

Theorem 5.3 *With $c = (\frac{q+1}{q'+1}, 3) = 1$, $e \geq 3$ an odd prime, $G = \mathrm{PGU}(3, q)$, and $M = \mathrm{PGU}(3, q') : \langle f \rangle$, the BG-Conjecture holds for the action of G on $[G : M]$.*

Proof Taking into account, all conjugate classes of subgroups $\langle z \rangle$ of prime order in M and an upper bound for the number $Q(\langle z \rangle)$ have been determined in Lemma 5.1 and Lemma 5.2, we are ready to show $\check{Q}(G) < \frac{1}{2}$, so that the BG-Conjecture is true.

Write $m' = r_1^{k_1} r_2^{k_2} \cdots r_l^{k_l}$ if $m' \neq 1$, where $2 \leq r_1 \leq r_2 \leq \cdots \leq r_l$ are distinct primes and $k_i \geq 1$ for all i . Further, let

- (a) $q' - 1 = 2^{\epsilon_2} t_1^{j_1} t_2^{j_2} \cdots t_o^{j_o}$, where $3 \leq t_1 \leq t_2 \leq \cdots \leq t_o$ are distinct primes;
- (b) $q' + 1 = 2^{\epsilon_1} 3^{\kappa_1} s_1^{i_1} s_2^{i_2} \cdots s_h^{i_h}$, where $5 \leq s_1 \leq s_2 \leq \cdots \leq s_h$ are distinct primes; and
- (c) $q'^2 - q' + 1 = 2^{\epsilon_3} 3^{\kappa_3} u_1^{\ell_1} u_2^{\ell_2} \cdots u_i^{\ell_i}$ where $5 \leq \ell_1 \leq \ell_2 \leq \cdots \leq \ell_i$ are distinct primes.

Furthermore, denote by $z_{q'-1,i}$ an element of order t_i in $\langle z_{q'-1} \rangle$; denote by $z_{q'+1,i}^A(a)$ and $z_{q'+1,i}^B$ elements of order s_i in $\langle z_{q'+1}^A(a) \rangle$ and $\langle z_{q'+1}^B \rangle$, respectively; denote by $z_{q'^2-q'+1,i}$ an element of order u_i in $\langle z_{q'^2-q'+1} \rangle$. Recall $\mathcal{P}^*(M) := \{\langle g_1 \rangle, \langle g_2 \rangle, \dots, \langle g_k \rangle\}$ is the set of representatives of conjugacy classes of subgroups of prime order in M .

Suppose that either $e \geq 5$ and $q' \geq 3$ or $e = 3$ and $q' \geq 7$. Then by Lemma 5.1.(I)

and Lemma 5.2.(I), we get

$$\begin{aligned}
\check{Q}(G) &= \frac{|\Omega|}{|\Omega|} \left(\sum_{i=1}^k (|\langle g_i \rangle| - 1) \frac{\text{Fix}(\langle g_i \rangle)}{|\text{N}_M(\langle g_i \rangle)|} \right) = \sum_{i=1}^k Q(\langle g_i \rangle) \\
&\leq (Q(\langle z_0 \rangle) + Q(\langle z'_0 \rangle) + Q(\langle z_2 \rangle) + Q(\langle z_A \rangle) + Q(\langle z_B \rangle) + Q(\langle \delta' \rangle)) \\
&\quad + \sum_{i=1}^o Q(\langle z_{q'-1,i} \rangle) + \sum_{i=1}^h \sum_{1 \leq a \leq \frac{s_i-3}{2}} Q(\langle z_{q'+1,i}^A(a) \rangle) + \sum_{i=1}^h Q(\langle z_{q'+1,i}^B \rangle) + \sum_{i=1}^i Q(\langle z_{q'^2-q'+1,i} \rangle) \\
&\quad + \sum_{r_i|m, r_i \neq 2, e} Q(\langle f^{\frac{2m}{r_i}} \rangle) + Q(\langle f^m \rangle) + \sum_{z \text{ is in Lemma 5.2(I)(3), } |z|=e} Q(\langle z \rangle) \\
&\leq \frac{6}{26} + (q' - 1)Q(\langle z_{q'-1} \rangle) + \frac{(q'+1)^2}{2}Q(\langle z_{q'+1}^A(a) \rangle) + (q' + 1)Q(\langle z_{q'+1}^B \rangle) \\
&\quad + (q'^2 - q' + 1)Q(\langle z_{q'^2-q'+1} \rangle) + m'Q(\langle f^{\frac{2m}{r}} \rangle) + \frac{1}{26} + \frac{1}{26} \\
&< \frac{6}{26} + 7 \cdot \frac{1}{26} = \frac{1}{2}.
\end{aligned}$$

Suppose that $e = 3$ and $q' = 4$. Then by Lemma 5.1.(II) and Lemma 5.2.(II), we have

$$\begin{aligned}
\check{Q}(G) &\leq (Q(\langle z_0 \rangle) + Q(\langle z'_0 \rangle) + Q(\langle z_{q'-1,1} \rangle) + Q(\langle z_{q'+1,1}^A(a) \rangle) + Q(\langle z_{q'+1,1}^B \rangle) \\
&\quad + Q(\langle z_{q'^2-q'+1,1} \rangle) + Q(\langle f^m \rangle) + \sum_{z \text{ is in Lemma 5.2(II)(3), } |z|=3} Q(\langle z \rangle) < \frac{1}{2},
\end{aligned}$$

as desired. \square

6 $M_0 \cong \text{PSU}(3, q').(\frac{q+1}{q'+1}, 3)$ with $(\frac{q+1}{q'+1}, 3) = 3$

Let $q = p^m$, $q' = p^{m'}$ for some odd prime e such that $m = m'e$. Assume that $c = (\frac{q+1}{q'+1}, 3) = 3$. From the proof of Lemma 4.1, we get $d := (3, q+1) = d' := (3, q'+1) = e = 3$. Let

$$\begin{aligned}
T &:= \text{PSU}(3, q) \leq G := \text{P}\Sigma\text{U}(3, q) = T\langle f \rangle; \\
M_0 &:= \text{PSU}(3, q') \leq M_1 := \text{PGU}(3, q') = M_0\langle \delta \rangle \leq M := \text{PGU}(3, q') : \langle f \rangle = M_1\langle f \rangle; \\
\Omega &= [G : M].
\end{aligned}$$

In what follows, we shall consider the action of G on Ω and prove $\hat{Q}(G) < \frac{1}{2}$, so that the BG-Conjecture is true for G . By Remark 2.4, the BG-Conjecture is true for every group G_1 between T and $\text{P}\Sigma\text{U}(3, q)$ with the corresponding point-stabilizers.

First we determine the representatives $\langle z \rangle$ of conjugacy classes of subgroups of prime order in M , while these subgroups are divided into two subclasses: those are contained M_1 or those not. We shall deal with them in Lemmas 6.1 and 6.2, separately; and also measure $Q(\langle z \rangle)$ for these subgroups $\langle z \rangle$, where

$$Q(\langle z \rangle) := \frac{|M|}{|\Omega|} \frac{(|z| - 1)\text{Fix}(\langle z \rangle)}{|\text{N}_M(\langle z \rangle)|};$$

and then prove the conjecture for this case in Theorem 6.3. Since $q' \neq 3, 4$ when $e = 3$, we may (and will) assume $q' \geq 5$ in the following analysis.

Lemma 6.1 *Let $c = e = 3$ and $q' \geq 5$. Then using the notations given in Table 1 and Lemma 4.2, every cyclic subgroup $\langle z \rangle$ of prime order r of M_1 is M conjugate to one of the following groups.*

- (1) $r = 2$ and p is odd: $\langle z_2 \rangle$ with $Q(\langle z_2 \rangle) < \frac{1}{26}$.
- (2) $r = p$: $\langle z_0 \rangle$ with $Q(\langle z_0 \rangle) < \frac{1}{26}$; and $\langle z'_0 \rangle$ with $Q(\langle z'_0 \rangle) < \frac{1}{26}$.
- (3) $r = 3$ and $r \mid q' + 1$: $\langle z \rangle = \langle z_A \rangle$, $\langle z_B \rangle$ or $\langle \delta' \rangle \in M_1 \setminus M_0$ with $Q(\langle z_A \rangle) + Q(\langle z_B \rangle) + Q(\langle \delta' \rangle) < \frac{3}{26}$.
- (4) $r \mid q' - 1$ and $r \neq 2$: $\langle z_{q'-1} \rangle$ with $Q(\langle z_{q'-1} \rangle) < \frac{1}{26(q'-1)}$.
- (5) $r \mid q' + 1$ and $r \neq 2, 3$: $\langle z_{q'+1}^A(a) \rangle$ with $Q(\langle z_{q'+1}^A(a) \rangle) < \frac{1}{13(q'+1)^2}$; and $\langle z_{q'+1}^B \rangle$ with $Q(\langle z_{q'+1}^B \rangle) < \frac{1}{26(q'+1)}$.
- (6) $r \mid q'^2 - q' + 1$ with $r \neq 3$: $\langle z_{q'^2-q'+1} \rangle$ and $Q(\langle z_{q'^2-q'+1} \rangle) < \frac{1}{26(q'^2-q'+1)}$.

Proof Suppose $c = 3$, $q' \geq 5$ and $q \geq 125$. Then

$$\frac{|M|}{|\Omega|} = \frac{|M|^2}{|G|} = \frac{2m|\text{PGU}(3, q')|^2}{|\text{PSU}(3, q)|} = \frac{6mq'^6(q'^3+1)^2(q'^2-1)^2}{q^3(q^3+1)(q^2-1)} \leq \frac{6mq'^{16}(1+q'^{-3})^2}{q^8(1-q^{-2})} < \frac{61m}{10q'^8}.$$

By Lemmas 4.2 and 4.3, we need to discuss the following cases, separately.

- (1) Case $r = 2$ and p is odd: By Lemma 4.3, we may set $z = z_2$, while $z_2^f = z_2$. Thus, $\text{Fix}(\langle z_2 \rangle) = \frac{|\text{N}_G(\langle z_2 \rangle)|}{|\text{N}_M(\langle z_2 \rangle)|} = \frac{2mq(q^2-1)(q+1)/3}{q'(q'^2-1)(q'+1) \times 2m} = \frac{q'^2(q'^2+q'+1)(q'^2-q'+1)^2}{3}$, and

$$Q(\langle z_2 \rangle) = \frac{|M|}{|\Omega|} \frac{(2-1)\text{Fix}(\langle z_2 \rangle)}{|\text{N}_M(\langle z_2 \rangle)|} = \frac{|M|}{|\Omega|} \frac{q'^2(q'^2+q'+1)(q'^2-q'+1)^2}{3 \times q'(q'^2-1)(q'+1) \times 2m} \leq \frac{61m}{10q'^8} \frac{q'^4 \frac{31}{25}}{6m \frac{24}{25}} < \frac{15}{11q'^4} < \frac{1}{26}.$$

- (2) Case $r = p$: By Lemma 4.3, we may set either $z = z_0$ or $z = z'_l$ where $0 \leq l \leq d' - 1$.

Suppose that $z = z_0$. Then $|\text{N}_T(\langle z_0 \rangle)| = q^3(q+1)(p-1)/3$ and $|\text{N}_{M_1}(\langle z_0 \rangle)| = q'^3(q'+1)(p-1)$. This implies that $|\text{N}_G(\langle z_0 \rangle)| \leq 2m|\text{N}_T(\langle z_0 \rangle)| = 2mq^3(q+1)(p-1)/3$, and $|\text{N}_M(\langle z_0 \rangle)| \geq e|\text{N}_{M_1}(\langle z_0 \rangle)| = eq'^3(q'+1)(p-1)$, as $f^{\frac{2m}{e}} \in \text{C}_G(M_1)$. Now, we conclude that. Thus,

$$Q(\langle z_0 \rangle) = \frac{|M|}{|\Omega|} \frac{(p-1)\text{Fix}(\langle z_0 \rangle)}{|\text{N}_M(\langle z_0 \rangle)|} = \frac{|M|}{|\Omega|} \frac{(p-1)|\text{N}_G(\langle z_0 \rangle)|}{|\text{N}_M(\langle z_0 \rangle)|^2} \leq \frac{|M|}{|\Omega|} \frac{2m(p-1)q^3(q+1)(p-1)}{3e^2q'^6(q'+1)^2(p-1)^2} < \frac{61m'^2}{15q'^4} < \frac{1}{26}.$$

Suppose that $z = z'_l$ where $0 \leq l \leq d' - 1$. Since these three elements z'_l are conjugate to each other in M_1 , we may set $z = z'_0$. Observe that $|\text{N}_G(\langle z'_0 \rangle)| \leq 2m|\text{N}_T(\langle z_0 \rangle)| = 2mq^2(p-1)$ and $|\text{N}_M(\langle z'_0 \rangle)| \geq e|\text{N}_{M_1}(\langle z'_0 \rangle)| = eq'^2(p-1)$. This implies that

$$Q(\langle z'_0 \rangle) = \frac{|M|}{|\Omega|} \frac{(p-1)\text{Fix}(\langle z'_0 \rangle)}{|\text{N}_M(\langle z'_0 \rangle)|} = \frac{|M|}{|\Omega|} \frac{(p-1)|\text{N}_G(\langle z'_0 \rangle)|}{|\text{N}_M(\langle z'_0 \rangle)|^2} \leq \frac{|M|}{|\Omega|} \frac{2m(p-1)q'^4}{q'^2(p-1) \times e^2} < \frac{61m}{10q'^8} \frac{2mq'^2}{e^2} = \frac{61m'^2}{5q'^6} < \frac{1}{26}.$$

- (3) Case $r = 3$ and $r \mid q' + 1$: By Lemma 4.2, we take $z = z_A, z_B$ and δ' . Since these are diagonal matrices, we have $\langle z \rangle^f = \langle z \rangle$. An it follows from Lemma 2.1 that there are two conjugacy classes subgroup in T of order 3, due to $9 \mid q + 1$.

$z = z_A$: $|\text{N}_G(\langle z_A \rangle)| = 2ms(q+1)^2$ and $|\text{N}_M(\langle z_A \rangle)| = 6ms'(q'+1)^2$, where $s, s' \in \{1, 2\}$. Hence,

$$\begin{aligned} Q(\langle z_A \rangle) &= \frac{|M|}{|\Omega|} \frac{(3-1)\text{Fix}(\langle z_A \rangle)}{|\text{N}_M(\langle z_A \rangle)|} = \frac{|M|}{|\Omega|} \frac{2|\text{N}_G(\langle z_A \rangle)|}{|\text{N}_{M_2}(\langle z_A \rangle)|^2} = \frac{|M|}{|\Omega|} \frac{2s(q'^2-q'+1)^2}{3s' \times 3s'(q'+1)^2 \times 2m} < \frac{61m}{10q'^8} \frac{2(q'^2-q'+1)^2}{9m(q'+1)^2} \\ &\leq \frac{61m}{10q'^8} \frac{2q'^2}{9m} = \frac{61}{45q'^6}. \end{aligned}$$

$z = z_B$ or $z = \delta'$: Since $c = 3$, we get that $\delta' \in \mathrm{PSU}(3, q)$. Obviously, $\langle \delta' \rangle$ is not M -conjugate to $\langle z_A \rangle$ and $\langle z_B \rangle$. But $\langle \delta' \rangle$ is T -conjugate to $\langle z_B \rangle$, where $\delta' \in [w^2, 1, 1]Z(\mathrm{GU}(3, q'))$ with $w \in \mathbb{F}_{q'^2}^*$ and $|w| = 3$.

Since $|\mathrm{N}_G(\langle z_B \rangle)| = |\mathrm{N}_G(\langle \delta' \rangle)| = q(q+1)(q^2-1)2m/3$, $|\mathrm{N}_M(\langle \delta' \rangle)| = 6m(q^2-q'+1)$ and $|\mathrm{N}_M(\langle z_B \rangle)| = 2mq'(q'+1)(q'^2-1)$, we have that

$$\begin{aligned} \mathrm{Fix}(\langle z_B \rangle) &= \mathrm{Fix}(\langle \delta' \rangle) = \frac{q(q+1)(q^2-1)2m}{3 \times q'(q'+1)(q'^2-1)2m} + \frac{q(q+1)(q^2-1)2m}{(q'^2-q'+1)18m} \\ &= \frac{1}{3}q'^2(q'^2-1'+1)(q'^2+q'+1)[q'^2-q'+1+\frac{1}{3}q'(q'+1)^2(q'-1)]; \\ Q(\langle z_B \rangle) &= \frac{|M|}{|\Omega|} \frac{2\mathrm{Fix}(\langle z_B \rangle)}{|\mathrm{N}_M(\langle z_B \rangle)|} = \frac{|M|}{|\Omega|} \frac{2}{3} \frac{q'^2(q'^2-q'+1)(q'^2+q'+1)[q'^2-q'+1+\frac{1}{3}q'(q'+1)^2(q'-1)]}{q'(q'+1)(q'^2-1)2m} \\ &\leq \frac{61m}{10q'^8} \frac{q'^3(q'^{-2}+q'^{-1}+1)q'^2[3q'^2-3q'+3+q'(q'+1)^2(q'-1)]}{9mq'^3(1-q'^{-2})} \\ &\leq \frac{61m}{10q'^8} \frac{\frac{31}{25}q'^2(q'^4+q'^3+2q'^2-4q'+3)}{9m\frac{24}{25}} \leq \frac{61m}{10q'^8} \frac{899q'^6}{5000m} < \frac{31}{28q'^2}; \\ Q(\langle \delta' \rangle) &= \frac{|M|}{|\Omega|} \frac{2\mathrm{Fix}(\langle \delta' \rangle)}{|\mathrm{N}_M(\langle \delta' \rangle)|} = \frac{|M|}{|\Omega|} \frac{2}{3} \frac{q'^2(q'^2-q'+1)(q'^2+q'+1)[q'^2-q'+1+\frac{1}{3}q'(q'+1)^2(q'-1)]}{3(q'^2-q'+1) \times 2m} \\ &\leq \frac{61m}{10q'^8} \frac{q'^8(q'^{-2}+q'^{-1}+1)(1+q'^{-1}+2q'^{-2}-4q'^{-3}+3q'^{-4})}{27m} < \frac{2}{5q'^2}. \end{aligned}$$

Consequently, we have $Q(\langle z_A \rangle) + Q(\langle z_B \rangle) + Q(\langle \delta' \rangle) < \frac{61}{45q^6} + \frac{31}{28q^2} + \frac{2}{5q^2} < \frac{3}{26}$.

(4) Case $r \mid q' - 1$ and $r \neq 2$: By Lemma 4.3, we may set $z = z_{q'-1}$. It follows that $|\mathrm{N}_G(\langle z_{q'-1} \rangle)| = 4m(q^2-1)/3$ and $|\mathrm{N}_M(\langle z_{q'-1} \rangle)| = 4m(q^2-1)$. Therefore, we have

$$\begin{aligned} (q'-1)Q(\langle z_{q'-1} \rangle) &= (q'-1) \frac{|M|}{|\Omega|} \frac{(r-1)\mathrm{Fix}(\langle z_{q'-1} \rangle)}{|\mathrm{N}_M(\langle z_{q'-1} \rangle)|} = (q'-1) \frac{|M|}{|\Omega|} \frac{(r-1)|\mathrm{N}_G(\langle z_{q'-1} \rangle)|}{|\mathrm{N}_M(\langle z_{q'-1} \rangle)|^2} \\ &= (q'-1) \frac{|M|}{|\Omega|} \frac{(r-1)\frac{4m(q^2-1)}{3}}{(4m(q'^2-1))^2} \leq \frac{61m}{10q'^8} \frac{(r-1)q'^5(q'^{-2}+q'^{-4}+1)}{12mq'^2(1-q'^{-2})} \leq \frac{61m}{10q'^8} \frac{(r-1)217q'^3}{2400m} < \frac{4}{7q'^4} \not\leq \frac{1}{26}, \end{aligned}$$

(5) Case $r \mid q' + 1$ and $r \neq 2, 3$: Since $r \notin 2, 3$ and $q' \geq 5$, it follows that $q' \geq 9$. By Lemma 4.3, we may take $z = z_{q'+1}^A(a)$ and $z = z_{q'+1}^B$, where $2 \leq a \leq \frac{r-1}{2}$. From the proof of Lemma 4.3, we get that these different M_1 -conjugate classes are also not conjugate in T .

$z = z_{q'+1}^A(a)$: It follows from Lemma 4.3 that $|\mathrm{N}_G(\langle z_{q'+1}^A(a) \rangle)| = \frac{(q+1)^2 \times 2m \times s}{3}$ and $|\mathrm{N}_M(\langle z_{q'+1}^A(a) \rangle)| = s'(q'+1)^2 \times 2m$, where s, s' are divisors of 6. Hence,

$$\begin{aligned} \frac{(q'+1)^2}{2}Q(\langle z_{q'+1}^A(a) \rangle) &= \frac{(q'+1)^2}{2} \frac{|M|}{|\Omega|} \frac{(r-1)\mathrm{Fix}(\langle z_{q'+1}^A(a) \rangle)}{|\mathrm{N}_M(\langle z_{q'+1}^A(a) \rangle)|} = \frac{(q'+1)^2}{2} \frac{|M|}{|\Omega|} \frac{(r-1)|\mathrm{N}_G(\langle z_{q'+1}^A(a) \rangle)|}{|\mathrm{N}_M(\langle z_{q'+1}^A(a) \rangle)|^2} \\ &= \frac{(q'+1)^2}{2} \frac{|M|}{|\Omega|} \frac{(r-1)s(q'^2-q'+1)^2}{3s'^2(q'+1)^2 \times 2m} \leq \frac{(q'+1)^2}{2} \frac{61m}{10q'^8} \frac{(r-1)(q'^2-q'+1)^2}{m(q'+1)^2} \leq \frac{61}{20q'^3} \not\leq \frac{1}{26}. \end{aligned}$$

$z = z_{q'+1}^B$: Applying Lemma 4.3, we obtain $|\mathrm{N}_G(\langle z_{q'+1}^B \rangle)| = \frac{q(q^2-1)(q+1) \times 2m}{3}$ and $|\mathrm{N}_M(\langle z_{q'+1}^B \rangle)| = q'(q^2-1)(q'+1) \times 2m$. Hence,

$$\begin{aligned} (q'+1)Q(\langle z_{q'+1}^B \rangle) &= (q'+1) \frac{|M|}{|\Omega|} \frac{(r-1)\mathrm{Fix}(\langle z_{q'+1}^B \rangle)}{|\mathrm{N}_M(\langle z_{q'+1}^B \rangle)|} = (q'+1) \frac{|M|}{|\Omega|} \frac{(r-1)|\mathrm{N}_G(\langle z_{q'+1}^B \rangle)|}{|\mathrm{N}_M(\langle z_{q'+1}^B \rangle)|^2} \\ &= (q'+1) \frac{|M|}{|\Omega|} \frac{(r-1)q'^2(q'^2-q'+1)^2(q'^2+q'+1)}{3 \times q'(q'^2-1)(q'+1) \times 2m} = \frac{|M|}{|\Omega|} \frac{(r-1)q'(q'^2-q'+1)^2(q'^2+q'+1)}{6m(q'^2-1)} \\ &\leq \frac{61m}{10q'^8} \frac{(r-1)q'^7(q'^{-2}+q'^{-1}+1)}{6m(1-q'^{-2})q'^2} \leq \frac{15}{11q'^2} \not\leq \frac{1}{26}, \end{aligned}$$

(6) Case $r \mid q'^2 - q' + 1$ and $r \neq 3$: Following the same arguments as in the proof of Lemma 5.1 for case (6), we can take $z = z_{q'^2-q'+1}$ as a diagonal matrix. Further,

by Lemma 4.3, we have that $|\mathrm{N}_G(\langle z_{q'^2-q'+1} \rangle)| = 2m(q^2 - q + 1)$, $|\mathrm{N}_M(\langle z_{q'^2-q'+1} \rangle)| = 6m(q'^2 - q' + 1)$. This follows that

$$(q'^2 - q' + 1)Q(\langle z_{q'^2-q'+1} \rangle) = (q'^2 - q' + 1) \frac{|M|}{|\Omega|} \frac{(r-1)(q'^6 - q'^3 + 1)}{6m(q'^2 - q' + 1) \times (q'^2 - q' + 1) \times 3} = \frac{|M|}{|\Omega|} \frac{(r-1)(q'^6 - q'^3 + 1)}{18m(q'^2 - q' + 1)} \\ \leq \frac{61m}{10q'^8} \frac{(r-1)q'^6}{18mq'^2(1-q'^{-1})} \leq \frac{61}{144q'^2} \nleq \frac{1}{26}. \quad \square$$

Lemma 6.2 *Let $c = e = 3$, $q = p^m$ and $q' = p^{m'} \geq 5$. Then every cyclic subgroup $\langle z \rangle$ of prime order r of M not contained in M_1 is M -conjugate to one of the following groups.*

- (1) $r \notin \{e, 2\}$: $z = f^{\frac{2m}{r}}$ with $Q(\langle f^{\frac{2m}{r}} \rangle) < \frac{1}{26m'}$.
- (2) $r = 2$: $z = f^m$ with $Q(\langle f^m \rangle) < \frac{1}{26}$.
- (3) $r = e$: $\langle f^{2m'} \rangle$, $z_A f^{2m'}$, $\langle z_B f^{2m'} \rangle$ and $\langle \delta' f^{2m'} \rangle$ with $Q(\langle f^{2m'} \rangle) + Q(\langle z_A f^{2m'} \rangle) + Q(\langle z_B f^{2m'} \rangle) + Q(\langle \delta' f^{2m'} \rangle) < \frac{1}{26}$.

Proof If $u \in M \setminus M_1$, then we can write $z = u_1 f_1$ for some element $u_1 \in M_1$ and $f_1 \in \langle f \rangle$. Since $|z| = r$, we get $|f_1| = r$ and so we may set $z = u_1 f'$, where $f' := f^{\frac{2m}{r}}$. In what follows, we discuss three cases, separately.

- (1) Case $r \notin \{2, e\}$:

Proposition 2.1 allows us to set $z = f^{\frac{2m}{r}}$, and there is only one M_1 -class $\langle f^{\frac{2m}{r}} \rangle$. We also get that $|\mathrm{N}_G(\langle f^{\frac{2m}{r}} \rangle)| = |\mathrm{PSU}(3, q'^{\frac{3}{r}}) \rtimes \langle f \rangle| = \frac{2m}{3}(q'^{\frac{3}{r}})^3((q'^{\frac{3}{r}})^3 + 1)((q'^{\frac{3}{r}})^2 - 1)$ and $|\mathrm{N}_M(\langle f^{\frac{2m}{r}} \rangle)| = |\mathrm{PGU}(3, q'^{\frac{1}{r}}) \rtimes \langle f \rangle| = 2m(q'^{\frac{1}{r}})^3((q'^{\frac{1}{r}})^3 + 1)((q'^{\frac{1}{r}})^2 - 1)$. Hence,

$$m'Q(\langle f^{\frac{2m}{r}} \rangle) = m' \frac{|M|}{|\Omega|} \frac{(r-1)\mathrm{Fix}(\langle f^{\frac{2m}{r}} \rangle)}{|\mathrm{N}_M(\langle f^{\frac{2m}{r}} \rangle)|} = m' \frac{|M|}{|\Omega|} \frac{(r-1)|\mathrm{N}_G(\langle f^{\frac{2m}{r}} \rangle)|}{|\mathrm{N}_M(\langle f^{\frac{2m}{r}} \rangle)|^2} \\ \leq \frac{m' |M| (r-1)q'^{\frac{16}{r}}(q'^{\frac{-9}{r}} + 1)(q'^{\frac{-4}{r}} + q'^{\frac{-2}{r}} + 1)}{6mq'^{\frac{8}{r}}(1-q'^{\frac{-2}{r}})} \leq \frac{61mm' (r-1)q'^{\frac{8}{5}}(p^{\frac{-9m'}{m'}} + 1)(p^{\frac{-4m'}{m'}} + p^{\frac{-2m'}{m'}} + 1)}{6m(1-p^{\frac{-2m'}{m'}})} \\ < \frac{9m'^2}{5q'^{\frac{32}{5}}} \nleq \frac{1}{26}.$$

- (2) Case $r = 2$:

By Proposition 2.1, we may take $z = f^m$. From [2, Proposition 3.3.15], we have $\mathrm{Fix}(\langle f^m \rangle) = \frac{|\mathrm{N}_G(\langle f^m \rangle)|}{|\mathrm{N}_M(\langle f^m \rangle)|} = \frac{|\mathrm{PSO}(3, q'^3)|2m}{|\mathrm{PGO}(3, q')|2m} = \frac{q'^3(q'^6 - 1)}{q'(q'^2 - 1)} = q'^2(q'^4 + q'^2 + 1)$. Consequently, we conclude that

$$Q(\langle f^m \rangle) = \frac{|M|}{|\Omega|} \frac{\mathrm{Fix}(\langle f^m \rangle)}{|\mathrm{N}_{M_2}(\langle f^m \rangle)|} = \frac{|M|}{|\Omega|} \frac{q'^2(q'^4 + q'^2 + 1)}{q'(q'^2 - 1) \times 2m} < \frac{4}{q'^5} \nleq \frac{1}{26}.$$

- (3) Case $r = e = d = d' = 3$:

Therefore, z can be expressed as $u_0 f^{2m'}$, where $u_0 \in M_1$. Since $[x, f^{2m'}] = 1$ for any $x \in M_1$, we have $|u_0| = r = 3$. Therefore, by Lemma 4.2, z must be of the form $f^{2m'}$, $z_A f^{2m'}$, $z_B f^{2m'}$ or $\delta' f^{2m'}$, where $\delta' \in M_1 \setminus M_0$.

By Proposition 2.1, we have that these elements are conjugate to each other in $\mathrm{PGU}(3, q)$. Now, we will prove that they are also conjugate to each other in T . Since

there exists $g \in \mathrm{PGU}(3, q)$ such that $f'^g = g^{-1}g^{f'^{-1}}f' = z_A f'$. It follows that $g = g^{f'^{-1}}(z_A)^{-1}$. Followed from [2, Lemma 2.1.1] that $x \in \mathrm{PSU}(3, q)$ if and only if $\det(x) = \lambda^3$ for some $\lambda \in \mathbb{F}_{q^2}^*$. Suppose that $\det(g) = \xi^k$, where $\mathbb{F}_{q^2}^* = \langle \xi \rangle$. Now, we have $\xi^k = \xi^{-kq'^2}\xi^{3l}$, where k, l are integers, which implies that $3 \mid k(1 + q'^2)$. It follows that $3 \mid k$, as $(q' + 1, 3) = 3$ and $(q'^2 + 1, 3) = 1$. Therefore, $g \in \mathrm{PSU}(3, q)$. Finally, by analogous arguments, the elements f' , $z_A f'$, $z_B f'$, and $\delta' f'$ are all conjugate to each other within $\mathrm{PSU}(3, q)$. Therefore, by Proposition 2.2, we have

$$\begin{aligned}
|\mathrm{Fix}(\langle f' \rangle)| &= |\mathrm{Fix}(\langle z_A f' \rangle)| = |\mathrm{Fix}(\langle z_B f' \rangle)| = |\mathrm{Fix}(\delta' f')| \\
&= \frac{|\mathrm{N}_G(\langle f' \rangle)|}{|\mathrm{N}_M(\langle f' \rangle)|} + \frac{|\mathrm{N}_G(\langle z_A f' \rangle)|}{|\mathrm{N}_M(\langle z_A f' \rangle)|} + \frac{|\mathrm{N}_G(\langle z_B f' \rangle)|}{|\mathrm{N}_M(\langle z_B f' \rangle)|} + \frac{|\mathrm{N}_G(\langle \delta' f' \rangle)|}{|\mathrm{N}_M(\langle \delta' f' \rangle)|} \\
&= \frac{|\mathrm{N}_G(\langle f' \rangle)|}{|\mathrm{N}_M(\langle f' \rangle)|} + \frac{|\mathrm{N}_G(\langle f' \rangle)|}{|\mathrm{C}_M(\langle z_A \rangle)|} + \frac{|\mathrm{N}_G(\langle f' \rangle)|}{|\mathrm{C}_M(\langle z_B \rangle)|} + \frac{|\mathrm{N}_G(\langle f' \rangle)|}{|\mathrm{C}_M(\langle \delta' \rangle)|} \\
&= \frac{|\mathrm{PGU}(3, q') : \langle f' \rangle|}{|\mathrm{PGU}(3, q') : \langle f' \rangle|} + \frac{|\mathrm{PGU}(3, q') : \langle f' \rangle|}{[3(q'+1)^2]r} + \frac{|\mathrm{PGU}(3, q') : \langle f' \rangle|}{[q'(q'+1)(q'^2-1)]r} + \frac{|\mathrm{PGU}(3, q') : \langle f' \rangle|}{[3(q'^2-q'+1)]r} \\
&= 1 + \frac{q'^3(q'^3+1)(q'^2-1) \times 2m}{(q'+1)^2 \times 3r} + \frac{q'^3(q'^3+1)(q'^2-1) \times 2m}{q'(q'+1)(q'^2-1)r} + \frac{q'^3(q'^3+1)(q'^2-1) \times 2m}{3(q'^2-q'+1)r} \\
&= 1 + 2m'q'^2(\frac{2}{3}q'^4 - \frac{1}{3}q'^3 + \frac{4}{3}q'^2 - \frac{5}{3}q' + 1) \leq \frac{4m'q'^6}{3}.
\end{aligned}$$

We therefore have

$$\begin{aligned}
Q(\langle f' \rangle) &= \frac{|M|}{|\Omega|} \frac{2\mathrm{Fix}(\langle f' \rangle)}{|\mathrm{N}_M(\langle f' \rangle)|} \leq \frac{|M|}{|\Omega|} \frac{2 \times \frac{4m'q'^6}{3}}{q'^3(q'^3+1)(q'^2-1) \times 2m} \leq \frac{|M|}{|\Omega|} \frac{25}{54q'^2}, \\
Q(\langle z_A f' \rangle) &= \frac{|M|}{|\Omega|} \frac{2\mathrm{Fix}(\langle z_A f' \rangle)}{|\mathrm{N}_M(\langle z_A f' \rangle)|} \leq \frac{|M|}{|\Omega|} \frac{2 \times \frac{4m'q'^6}{3}}{(q'+1)^2 \times 9} \leq \frac{|M|}{|\Omega|} \frac{8m'q'^4}{27}, \\
Q(\langle z_B f' \rangle) &= \frac{|M|}{|\Omega|} \frac{2\mathrm{Fix}(\langle z_B f' \rangle)}{|\mathrm{N}_M(\langle z_B f' \rangle)|} \leq \frac{|M|}{|\Omega|} \frac{2 \times \frac{4m'q'^6}{3}}{3q'(q'+1)(q'^2-1)} \leq \frac{|M|}{|\Omega|} \frac{25m'q'^2}{27}, \\
Q(\langle \delta' f' \rangle) &= \frac{|M|}{|\Omega|} \frac{2\mathrm{Fix}(\langle \delta' f' \rangle)}{|\mathrm{N}_M(\langle \delta' f' \rangle)|} \leq \frac{|M|}{|\Omega|} \frac{2 \times \frac{4m'q'^6}{3}}{9(q'^2-q'+1)} \leq \frac{|M|}{|\Omega|} \frac{10m'q'^4}{27}, \\
Q(\langle f' \rangle) + Q(\langle z_A f' \rangle) + Q(\langle z_B f' \rangle) + Q(\langle \delta' f' \rangle) &< \frac{61m}{10q^8}(\frac{25}{54q'^2} + \frac{18m'q'^4}{27} + \frac{25m'q'^2}{27}) \\
&\leq \frac{305m'}{36q^{10}} + \frac{61m'^2}{5q'^4} + \frac{305m'^2}{18q'^6} \not\leq \frac{1}{26}.
\end{aligned}$$

□

Theorem 6.3 *Let $c = e = 3$, $G = \mathrm{P\Sigma U}(3, q)$ and $M = \mathrm{PGU}(3, q') : \langle f \rangle$. Then the BG-Conjecture holds for G acting on $[G : M]$ by right multiplication.*

Proof Taking into account, all conjugate classes of subgroups $\langle z \rangle$ of prime order in M and an upper bound for the number $|Q(\langle z \rangle)|$ have been determined in Lemmas 6.1 and 6.2, we are ready to show $\check{Q}(G) < \frac{1}{2}$ so that the BG-Conjecture holds.

Write $m' = r_1^{k_1}r_2^{k_2} \cdots r_l^{k_l}$ if $m' \neq 1$, where $2 \leq r_1 \leq r_2 \leq \cdots \leq r_l$ are distinct primes and $k_i \geq 1$ for all i . Set $m_i = \frac{m}{r_i}$, $q_i = p^{m_i}$ and $f_i = f^{m_i}$. Further, let

- (a) $q' - 1 = 2^{\epsilon_2}t_1^{j_1}t_2^{j_2} \cdots t_o^{j_o}$, where $3 \leq t_1 \leq t_2 \leq \cdots \leq t_o$ are distinct primes;
- (b) $q' + 1 = 2^{\epsilon_1}3^{\kappa_1}s_1^{i_1}s_2^{i_2} \cdots s_h^{i_h}$, where $5 \leq s_1 \leq s_2 \leq \cdots \leq s_h$ are distinct primes; and
- (c) $q'^2 - q' + 1 = 2^{\epsilon_3}3^{\kappa_3}u_1^{\ell_1}u_2^{\ell_2} \cdots u_i^{\ell_i}$ where $5 \leq \ell_1 \leq \ell_2 \leq \cdots \leq \ell_i$ are distinct primes.

Furthermore, denote by $z_{q'-1,i}$ an element of order t_i in $\langle z_{q'-1} \rangle$; denote by $z_{q'+1,i}^A(a)$ and $z_{q'+1,i}^B$ elements of order s_i in $\langle z_{q'+1}^A(a) \rangle$ and $\langle z_{q'+1}^B \rangle$, respectively; denote by $z_{q'^2-q'+1,i}$ an element of order u_i in $\langle z_{q'^2-q'+1} \rangle$. Recall $\mathcal{P}^*(M) := \{\langle g_1 \rangle, \langle g_2 \rangle, \dots, \langle g_k \rangle\}$ is the set of representatives of conjugacy classes of subgroups of prime order in M .

By Lemmas 6.1 and 6.2, we have

$$\begin{aligned}
\check{Q}(G) &= \frac{|M|}{|\Omega|} \left(\sum_{i=1}^k (|\langle g_i \rangle| - 1) \frac{|\text{Fix}(\langle g_i \rangle)|}{|\text{N}_M(\langle g_i \rangle)|} \right) = \sum_{i=1}^k Q(\langle g_i \rangle) \\
&\leq (Q(\langle z_0 \rangle) + Q(\langle z'_0 \rangle) + Q(\langle z_2 \rangle) + Q(\langle z_3^A \rangle) + Q(\langle z_3^B \rangle) + Q(\langle \delta' \rangle)) \\
&+ \sum_{i=1}^o Q(\langle z_{q'-1,i} \rangle) + \sum_{i=1}^h \sum_{1 \leq a \leq \frac{s_i-3}{2}} Q(\langle z_{q'+1,i}^A(a) \rangle) + \sum_{i=1}^h Q(\langle z_{q'+1,i}^B \rangle) + \sum_{i=1}^i Q(\langle z_{q'^2-q'+1,i} \rangle) \\
&+ \sum_{r_i|m, r_i \neq 2, e} Q(\langle f^{\frac{2m}{r_i}} \rangle) + (Q(\langle f^{2m'} \rangle) + Q(\langle z_A f^{2m'} \rangle) + Q(\langle z_B f^{2m'} \rangle) + Q(\langle \delta' f^{2m'} \rangle)) + Q(\langle f^m \rangle) \\
&\leq \frac{6}{26} + (q' - 1)Q(\langle z_{q'-1} \rangle) + \frac{(q'+1)^2}{2}Q(\langle z_{q'+1}^A(a) \rangle) + (q' + 1)Q(\langle z_{q'+1}^B \rangle) \\
&+ (q'^2 - q' + 1)Q(\langle z_{q'^2-q'+1} \rangle) + m'Q(\langle f^{\frac{2m}{r}} \rangle) + \frac{1}{26} + \frac{1}{26} \\
&\leq \frac{6}{26} + 7 \cdot \frac{1}{26} < \frac{1}{2},
\end{aligned}$$

as desired. \square

7 $M_0 = \text{PSL}(2, 7)$ or $\text{PSL}(2, 9)$

Theorem 7.1 *Let $T = \text{PSU}(3, q)$ and $M_0 = \text{PSL}(2, 7)$ with $q = p \equiv 3, 5, 6 \pmod{7}$ and $q \neq 5$, so that $q = p \geq 13$. Let $T \leq G \leq \text{Aut}(\text{PSU}(3, q))$ and M a maximal subgroup of G containing M_0 . Then the BG-Conjecture holds for G acting on $[G : M]$ by right multiplication.*

Proof By checking [2, Tabel 8.6], we may set either $G = T$ and $M = M_0$; or $G = P\text{SU}(3, q)$ and $M = M_0 : \langle f \rangle \cong \text{PGL}(2, 7)$. By Remark 2.4, it suffices to show $\check{Q}(G) < \frac{1}{2}$ for the second case.

Set $\mathcal{P}^*(M)$ be the set of all the representatives of conjugacy classes of subgroups of prime order in M . Then

$$\mathcal{P}^*(M) = \{\langle x_1 \rangle, \langle x_2 \rangle, \langle x_3 \rangle, \langle x_4 \rangle\},$$

where $x_1, x_3, x_4 \in M_0$ and $x_2 \in M \setminus M_0$, $|x_1| = 2$, $|x_2| = 2$, $|x_3| = 3$ and $|x_4| = 7$. Note that $\langle x_1 \rangle$ and $\langle x_2 \rangle$ are also not conjugate to each other in G . Moreover, by Lemma 4.3 we have $\text{Fix}(\langle x_i \rangle) = \frac{|\text{N}_G(\langle x_i \rangle)|}{|\text{N}_M(\langle x_i \rangle)|}$, that is

$$\begin{aligned}
\text{Fix}(\langle x_1 \rangle) &= \frac{2 \times q(q^2-1)(q+1)/d}{16} = \frac{q(q^2-1)(q+1)}{8d}; & \text{Fix}(\langle x_2 \rangle) &= \frac{2|\text{SO}(3, q)|}{12} = \frac{q(q^2-1)}{6}; \\
\text{Fix}(\langle x_3 \rangle) &\leq \frac{2 \times q(q+1)(q^2-1)/d}{12} \leq \frac{7q^4}{39d}; & \text{Fix}(\langle x_4 \rangle) &\leq \frac{2 \times q(q+1)(q^2-1)/d}{42} \leq \frac{2q^4}{39d}.
\end{aligned}$$

Therefore,

$$\begin{aligned}
\check{Q}(G) &:= \frac{|M|}{|\Omega|} \left(\sum_{i=1}^4 (|\langle x_i \rangle| - 1) \frac{\text{Fix}(\langle x_i \rangle)}{|\text{N}_M(\langle x_i \rangle)|} \right) \\
&\leq \frac{336^2 \times d}{2 \times q^3(q^3+1)(q^2-1)} \left(\frac{q(q^2-1)(q+1)}{8d \times 16} + \frac{q(q^2-1)}{6 \times 12} + \frac{2 \times 7q^4}{39d \times 12} + \frac{6 \times 2q^4}{39d \times 42} \right) < \frac{1}{2}.
\end{aligned}$$

Theorem 7.2 Let $T = \mathrm{PSU}(3, q)$ and $M_0 = \mathrm{PSL}(2, 9)$ with $q = p \equiv 11, 14 \pmod{15}$. Let $T \leq G \leq \mathrm{Aut}(\mathrm{PSU}(3, q))$ and M is a maximal subgroup of G containing M_0 . Then the BG-Conjecture holds for G acting on $[G : M]$.

Proof By checking [2, Tabel 8.6] where $q \geq 7$, we get that either $G = T$ and $M = M_0$; or $G = \mathrm{P\Sigma U}(3, q) \cong T : \langle f \rangle$ and $M = M_0 : \langle f \rangle$, where f is given by $g^f = (g^{-1})^T$ for any $g \in T$ so that $M \cong \mathrm{PGL}(2, 9)$ (refer to [1, P.34]). By Remark 2.4, it suffices to show $\check{Q}(G) < \frac{1}{2}$ for the second case.

Set $\mathcal{P}^*(M)$ be the set of all the representatives of conjugacy classes of subgroups of prime order in M . Thus,

$$\mathcal{P}^*(M) = \{\langle x_1 \rangle, \langle x_2 \rangle, \langle x_3 \rangle, \langle x_4 \rangle\},$$

where $x_1, x_3, x_4 \in M_0$ and $x_2 \in M \setminus M_0$, $|x_1| = 2$, $|x_2| = 2$, $|x_3| = 3$ and $|x_4| = 5$. Note that $\langle x_1 \rangle$ and $\langle x_2 \rangle$ are also not conjugate to each other in G . Moreover, by Lemma 4.3 and the fact $q \geq 11$, we have

$$\begin{aligned} \mathrm{Fix}(\langle x_1 \rangle) &= \frac{2 \times q(q^2-1)(q+1)/d}{16} = \frac{q(q^2-1)(q+1)}{8d}; & \mathrm{Fix}(\langle x_2 \rangle) &= \frac{2|\mathrm{SO}(3,q)|}{20} = \frac{q(q^2-1)}{10}; \\ \mathrm{Fix}(\langle x_3 \rangle) &\leq \frac{2 \times q(q+1)(q^2-1)/d}{18} \leq \frac{4q^4}{33d}; & \mathrm{Fix}(\langle x_4 \rangle) &\leq \frac{2 \times q(q+1)(q^2-1)/d}{20} \leq \frac{6q^4}{55d}. \end{aligned}$$

Therefore,

$$\begin{aligned} \check{Q}(G) : &= \frac{|M|}{|\Omega|} \left(\sum_{i=1}^4 (|\langle x_i \rangle| - 1) \frac{\mathrm{Fix}(\langle x_i \rangle)}{|\mathrm{N}_M(\langle x_i \rangle)|} \right) \\ &\leq \frac{336^2 \times d}{2 \times q^3(q^3+1)(q^2-1)} \left(\frac{q(q^2-1)(q+1)}{8d \times 16} + \frac{q(q^2-1)}{10 \times 20} + \frac{2 \times 4q^4}{33d \times 18} + \frac{4 \times 6q^4}{55d \times 20} \right) < \frac{1}{2}. \end{aligned} \quad \square$$

Acknowledgements

The first author thanks the supports of the National Natural Science Foundation of China (12301446, 12571362). The second author thanks the supports of the National Natural Science Foundation of China (12471332). The third author thanks the supports of the National Natural Science Foundation of China (12301422).

References

- [1] J. N. Bray, D. F. Holt and C. M. Roney-Dougal, *The maximal subgroups of the low-dimensional finite classical groups*, London Mathematical Society Lecture Note Series, 407, Cambridge Univ. Press, Cambridge, 2013.
- [2] T. C. Burness and M. Giudici, *Classical groups, derangements and primes*, Australian Mathematical Society Lecture Series, 25, Cambridge Univ. Press, Cambridge, 2016.
- [3] T. C. Burness and M. Giudici, On the Saxl graph of a permutation group, *Math. Proc. Cambridge Philos. Soc.*, **168** (2020), 219–248.

- [4] T. C. Burness and H. Y. Huang, On the Saxl graphs of primitive groups with soluble stabilisers, *Algebr. Comb.*, **5** (2022), 1053–1087.
- [5] H. Chen and S. Du, On the Burness-Giudici conjecture, *Comm. Algebra*, **51** (2023), 5019–5045.
- [6] H. Chen, S. Du and W. Li, The Burness-Giudici Conjecture on Primitive Groups of Lie-type with Rank One: Part (I), arxiv.....
- [7] H. Chen and S. Du, The Burness-Giudici Conjecture on Primitive Groups of Lie-type with Rank One: Part (III), arxiv.....
- [8] B. Huppert, *Endliche Gruppen. I*, Die Grundlehren der mathematischen Wissenschaften, Band 134, Springer, Berlin-New York, 1967.
- [9] W. A. Manning, The order of primitive groups. III, *Trans. Amer. Math. Soc.*, **19** (1918), 127–142.
- [10] W. A. Simpson and J. S. Frame, The character tables for $\mathrm{SL}(3, q)$, $\mathrm{SU}(3, q^2)$, $\mathrm{PSL}(3, q)$, $\mathrm{PSU}(3, q^2)$, *Canadian J. Math.*, **25** (1973), 486–494.