

A GENERALIZATION OF THE "BROUWER – SCHAUDER – TYCHONOFF" FIXED-POINT THEOREM

RANJIT VOHRA

ABSTRACT. We prove a new fixed – point result for the image $\text{Im}(\varphi)$ of any continuous function φ from K to $K \times K$, where K is a compact convex subset of a Hausdorff locally convex space, provided that the projection of $\text{Im}(\varphi)$ to the first factor is onto, and a condition on the convex hull of $\text{Im}(\varphi)$ holds. A special case of our result is the Brouwer-Schauder-Tychonoff fixed-point theorem for continuous functions $f : K \rightarrow K$.

§ 1.

INTRODUCTION

We know, from the Brouwer-Schauder-Tychonoff theorem (see below for theorem statement), that any continuous function $f : K \rightarrow K$, where K is a compact convex subset of a Hausdorff locally convex space, admits a fixed point, meaning that the graph of f intersects the diagonal $\Delta \subset K \times K$. Intuitively, it seems plausible that any squiggly continuous image of K in $K \times K$, regardless of whether or not this image describes a function $f : K \rightarrow K$ in the target $K \times K$, should also intersect Δ . We make this intuition precise in our Theorem I below. Furthermore, since the Brouwer-Schauder-Tychonoff theorem can be regarded as a corollary of the Kakutani-Fan-Glicksberg ("KFG") fixed-point theorem for correspondences, we note (see Remark #2 in § 3 below) that the hypotheses of our Theorem I are not comparable to those of KFG.

Subject Classification: 46A99, 54A99, 91B50

§ 2.

REMINDERS

We recall the following two results.

BROUWER – SCHAUDER – TYCHONOFF FIXED-POINT THEOREM. Suppose:

- X is a Hausdorff locally convex space;
- K is a nonempty compact, convex subset of X ;
- $f : K \rightarrow K$ is a continuous function.

Then f has a fixed point.

Proof. See original papers [4], [11], [12]; and also 17.56, p.583 in [1].

KAKUTANI – FAN – GLICKSBERG FIXED-POINT THEOREM. Suppose:

- X is a Hausdorff locally convex space;
- K is a nonempty compact, convex subset of X ;
- $\Gamma : K \rightarrow K$ is a correspondence that has closed graph and nonempty, convex values.

Then Γ has a fixed point.

Proof. See original papers [10], [6], [7]; and also 17.55, p.583 in [1]

FACT: Every locally convex space (even if not Hausdorff) is locally connected, since each point has a local base of convex (hence connected) neighborhoods. In any locally connected space, each component of any open set is open. See e.g. p.199ff in [13]

§ 3.

THE MAIN RESULT

Remark #1. In the sequel we will see the following situation. There is a given function $\varphi : K \rightarrow K \times K$, $x \mapsto (\varphi_1(x), \varphi_2(x))$ such that φ_1 is onto. The image of φ , denoted $\text{Im}(\varphi)$, is a subset of $K \times K$ and can be regarded as the graph of the correspondence $\Psi : K \rightarrow K$, given by $t \mapsto \varphi_2(\{\varphi_1^{-1}(t)\})$. Thus $\text{Im}(\varphi) = \text{gr}(\Psi)$; where the correspondence Ψ is nonempty – valued (using that φ_1 is onto).

THEOREM I. Suppose:

- X is a Hausdorff locally convex space;
- K is a nonempty compact, convex subset of X ;
- $\varphi : K \rightarrow K \times K$, $x \mapsto (\varphi_1(x), \varphi_2(x))$ is a continuous function such that φ_1 is onto; & the correspondence $\Psi : K \rightarrow K$, $t \mapsto \varphi_2(\{\varphi_1^{-1}(t)\})$ has the following property: for each t in the domain of Ψ , $t \notin \Psi(t)$ implies $t \notin \text{co}(\Psi(t))$.

Then Ψ has a fixed point.

Proof. By Remark #1, the correspondence $\Psi : K \rightarrow K$, $t \mapsto \varphi_2(\{\varphi_1^{-1}(t)\})$ is nonempty – valued. Note also that $\text{gr}(\Psi)$ is connected, since $\text{gr}(\Psi) = \text{Im}(\varphi)$, and $\text{Im}(\varphi)$ is connected because it is the continuous image of the convex (hence connected) set K .

Now suppose, toward a contradiction, that the theorem hypotheses hold, and that Ψ has no fixed point, meaning that $\text{gr}(\Psi) \cap \Delta = \emptyset$, where Δ denotes the diagonal in the compact product $K \times K$. Then $t \notin \Psi(t)$ and indeed $t \notin \text{co}(\Psi(t))$ holds for all t in the domain K of Ψ , using the hypotheses on Ψ .

Put $\Psi'(t) := \text{co}(\Psi(t))$, and define a new correspondence $\Psi' : K \rightarrow K$ by $\Psi' := \bigcup_{t \in K} \text{co}(\Psi(t))$. We see that $\text{gr}(\Psi')$ has two important properties:

(P1) $\text{gr}(\Psi')$ is connected, since it is the union of the connected set $\text{gr}(\Psi)$ and the convex sets $\text{co}(\Psi(t))$ that are attached to points of $\text{gr}(\Psi)$.

(P2) $\text{gr}(\Psi') \cap \Delta = \emptyset$ since $t \notin \Psi'(t)$ holds for each $t \in K$, as noted above.

Since Δ is closed in $K \times K$ (using the Hausdorff hypothesis), we see from (P2) that $\text{gr}(\Psi')$ must belong to the open set $(K \times K) \setminus \Delta$. Moreover, since $\text{gr}(\Psi')$ is connected (P1), we know (see Fact in § 2) there is an open component \mathcal{U} of $(K \times K) \setminus \Delta$ such that $\text{gr}(\Psi') \subseteq \mathcal{U} \subseteq (K \times K) \setminus \Delta$.

The idea is to “fatten” Ψ' by adding open sets to its boundary $\partial\Psi'$, in order to produce a new correspondence Ψ^* that contains Ψ' and has open graph. Precisely, to each z of $\partial\Psi'$, we attach a basic open disk B_z (of the locally convex space $X \times X$). Thus for each t in K , and using notation $(B_z \mid t)$ for the restriction $B_z \cap (\{t\} \times K)$, we obtain:

$$\Psi^*(t) := \Psi'(t) \cup \left(\bigcup_{z \in \partial(\Psi'(t))} (B_z \mid t) \right) \quad (\dagger)$$

where each B_z is a basic open, convex disk of $X \times X$ that is centered at point z on the boundary of Ψ' , and is small enough that $\Psi^*(t) \subseteq \mathcal{U} \subseteq (K \times K) \setminus \Delta$.

We claim that $t \notin \text{co}(\Psi^*(t))$ for any $t \in K$. To see this, note that

$$\begin{aligned} \text{co}(\Psi^*(t)) &= \text{co} \left(\Psi'(t) \cup \left(\bigcup_{z \in \partial(\Psi'(t))} (B_z \mid t) \right) \right) && \text{using } (\dagger) \\ &= \Psi'(t) \cup \left(\bigcup_{z \in \partial(\Psi'(t))} (B_z \mid t) \right) \end{aligned}$$

as the expression on the last line is a convex set that (by choice of the disks B_z) does not contain t .

Put $\Psi^* := \bigcup_{t \in K} \Psi^*(t)$. Evidently Ψ^* is a nonempty – valued correspondence $K \rightarrow K$ with graph $\text{gr}(\Psi^*)$ that is open and connected, by construction. Thus we obtain the inclusions $\text{gr}(\Psi^*) \subseteq \mathcal{U} \subseteq (K \times K) \setminus \Delta$, and (as above) that $t \notin \text{co}(\Psi^*(t))$ holds for each t in the domain K of Ψ .

Let $\mathcal{H} : K \Rightarrow K$, $y \mapsto K \setminus (\Psi^*)^{-1}(y)$ be the “inverse complement correspondence”¹ of $\Psi^* : K \Rightarrow K$. Observe that Lemma 17.47 in [1]) applies to Ψ^* , and since $t \notin \text{co}(\Psi^*(t))$ for all $t \in K$, we see \mathcal{H} is a KKM correspondence. Furthermore (using that Ψ^* has open graph) Theorem 17.46 of [1] implies that there is some point x in the domain K of Ψ^* (and hence also of Ψ) that is empty-valued, contradicting that Ψ is nonempty-valued. We therefore conclude from this contradiction that our initial supposition $\text{gr}(\Psi) \cap \Delta = \emptyset$ is untenable, and that Ψ must have a fixed point. \square

COROLLARY. Suppose the function φ_1 is the identity map. Then $\Psi = \varphi_2$, whereby Ψ is a continuous function from K to K , and we recover the Brouwer-Schauder-Tychonoff fixed point theorem (see § 2) for Ψ .

Remark #2. Although the Brouwer – Schauder – Tychonoff fixed point theorem is a special case of both the Kakutani – Fan – Glicksberg theorem and of our Theorem I, the hypotheses of our theorem are different from those of KFG: neither implies or is implied by the other. The hypotheses of the FKG theorem require that the correspondence Γ has convex values, whereas our Theorem I has the weaker requirement that for each t in the domain of the correspondence Ψ , $t \notin \Psi(t)$ implies $t \notin \text{co}(\Psi(t))$. On the other hand, the correspondence Γ of the FKG theorem is not necessarily the image of a continuous function $\varphi : K \rightarrow K \times K$, so FKG is more general in this regard.

REFERENCES

- [1] C.D. Aliprantis, K.C. Border, Infinite Dimensional Analysis, 3rd ed., Springer, 2006.
- [2] M. Balaj, S. Muresan, Generalizations of the Fan-Browder Fixed Point Theorem & Minimax Inequalities, Archivum Mathematicum, Tomus 41 (2005), 399-407.
- [3] K.C. Border, Fixed point theorems with applications to economics and game theory, Cambridge University Press, 1985
- [4] L.E.J. Brouwer, 1912, Über Abbildung von Mannigfaltigkeiten, Mathematische Annalen 71:165 -168.
- [5] K. Fan, Fixed point and minimax theorems in locally convex topological spaces, Proc. Of the National Academy of Sciences, USA 38:121-126, (1952)
- [6] K. Fan, A generalization of Tychonoff’s fixed point theorem, Math. Ann. 142 (1962) 305–310.
- [7] I.L. Glicksberg, A further generalization of the Kakutani fixed point theorem with application to Nash equilibrium, Proc. Amer. Math. Soc. 3 (1952) 170–174.

¹ See e.g. p.578 in [1].

- [8] B.R. Halpern and G.M. Bergman, A Fixed Point Theorem for Inward and Outward Maps, *Trans. Amer. Math. Soc.*, 130, 353-358 (1968).
- [9] C.J. Himmelberg, Fixed points of compact multifunctions, *J. Math. Anal. Appl.* 38(1972), 205-207
- [10] S. Kakutani, A generalization of Brouwer's fixed point theorem, *Duke Mathematical Journal*, 8:457-459, 1941.
- [11] J. Schauder, 1930, Der Fixpunktsatz in Functionalraumen, *Studia Mathematica* 2; 171-180.
- [12] A. Tychonoff, 1935, Ein Fixpunktsatz, *Mathematische Annalen* 111:767-776.
- [13] S. Willard, *General Topology*, Addison Wesley, 1970 .

R. VOHRA

P.O. Box 240, Storrs CT 06268

rvohra@bridgew.edu