

Graph Limits via Quotients

Eitan Levin[†] and Venkat Chandrasekaran^{†,‡ *}

[†] Department of Computing and Mathematical Sciences

[‡] Department of Electrical Engineering
California Institute of Technology
Pasadena, CA 91125

December 30, 2025

Abstract

We present a new notion of limits of weighted directed graphs of growing size based on convergence of their random quotients. These limits are specified in terms of random exchangeable measures on the unit square. We call our limits *grapheurs* and show that these are dual to graphons in a precise sense. Grapheurs are well-suited to modeling hubs and connections between them in large graphs; previous notions of graph limits based on subgraph densities fail to adequately model such global structures as subgraphs are inherently local. Using our framework, we present an edge-based sampling approach for testing properties pertaining to hubs in large graphs. This method relies on an edge-based analog of the Szemerédi regularity lemma, whereby we show that sampling a small number of edges from a large graph approximately preserves its quotients. Finally, we observe that the random quotients of a graph are related to each other by equipartitions, and we conclude with a characterization of such random graph models.

Keywords: De Finetti, graphons, hubs, property testing, Szemerédi regularity

1 Introduction

Analyzing the structure of large graphs is a challenge that arises in numerous problem domains such as bioinformatics, network tomography, operations research, and the social sciences [1–5]. Graphons and other related notions of graph limits in the literature employ subgraph densities to summarize and compare large graphs. However, subgraph-based measures are inherently local and they fail to suitably account for global structures such as hubs and connections between them, which are commonly encountered in real-world networks [6–10]. For example, most subgraphs of a star, the prototypical example of a hub, do not contain any edges.

A *hub* is a vertex in a graph¹ with a significant fraction of the total edge weight incident to it. Our central thesis is that hubs in large graphs can be detected by their *quotients*. Formally, a quotient of size k of a graph on n vertices is defined by an ordered partition $f_{k,n} : [n] \rightarrow [k]$, which partitions the vertex set $[n]$ into k parts that are given by the fibers $f_{k,n}^{-1}(j)$ for $j \in [k]$. In particular, for a graph on n vertices with adjacency matrix $G \in \mathbb{R}_+^{n \times n}$, its quotient by the ordered partition $f_{k,n}$ is given by the graph $\rho(f_{k,n})G \in \mathbb{R}_+^{k \times k}$ on k vertices with edge weights:

$$[\rho(f_{k,n})G]_{i,j} = \sum_{\substack{v \in f_{k,n}^{-1}(i) \\ u \in f_{k,n}^{-1}(j)}} G_{v,u}. \quad (1)$$

*Email: {eitanl, venkatc}@caltech.edu

¹We consider directed graphs with possible self-loops and nonnegative edge weights.

Here we view $\rho(f_{k,n}) : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}^{k \times k}$ as a ‘linear representation’ of $f_{k,n}$. In words, the vertices of the quotient $\rho(f_{k,n})G$ correspond to the parts in the partition defined by $f_{k,n}$, and the edge weight between parts $i, j \in [k]$ is given by sums of all edge weights in G between vertices in parts $f_{k,n}^{-1}(i)$ and $f_{k,n}^{-1}(j)$. Thus, a quotient of G represents the edge weights between each pair of parts in a vertex partition.

As an illustration, we give in Figures 1–2 examples of small quotients of two networks. The first, depicted in Figure 1, is a traffic network representing the fraction of trips starting and ending at each zone of the Gold Coast, Australia [11]. The second, depicted in Figure 2, is a brain connectome, the network of connections between neurons of an adult male *C. elegans* roundworm [12, 13]. The structure of each quotient indicates some of the features of the underlying large network. As we will see in Section 4.3, the presence of hubs in the original network leads to nonuniform distribution of edge weight in their random quotients.




Figure 1: Traffic network for the Gold Coast, Australia, and an associated random quotient. The city is divided into 1068 zones, and the weight of an edge is the fraction of trips from a dataset with given start and end zones [11]. Here we plot the first 300 of these zones and their interconnections, displaying the width of an edge proportionally to its weight. We color the vertices according to their image in the quotient, and color and edge connecting two vertices with the same image by the color of that image. The nonuniformity in the edge weights of the quotient indicates the presence of hubs that we can see in the plot of the entire network.




Figure 2: Network of chemical connections among the 575 neurons of adult male *C. elegans* (a species of roundworm), and its random quotient. Edges are directed from pre-synaptic to post-synaptic cells, and their weight is the total number of EM serial sections of connectivity [12, 13]. The nonuniform distribution of self-loops and the unequal in-degrees and out-degrees among the four vertices in the quotient again indicate the presence of hubs.

In this paper, we define a new notion of convergence for sequences of growing-sized graphs based on convergence of their random quotients. In analogy with graphons [14, §1], we define suitable limit objects in terms of random measures for any quotient-convergent sequence of graphs. Finally, we show that finite (random) graphs sampled from our limit objects are quotient-convergent to the limit almost surely. The key concepts in our paper are dual to those in the literature on graphons, and we emphasize this duality throughout our development. In particular, we shall see that random quotients and random subgraphs, in terms of which graphon convergence is defined, are random linear maps which are adjoints of each other. Further, a dual analog of the cut metric plays a prominent role in our development, and we derive a dual Szemerédi lemma as a consequence of our results. We describe next these contributions in more detail.

1.1 Our Contributions

To set the stage, we define a random quotient of a graph on n vertices in terms of a uniformly random map $F_{k,n}: [n] \rightarrow [k]$. That is, each $i \in [n]$ gets sent independently to a uniform element $F_{k,n}(i) \in [k]$. Equivalently, this is a uniformly random ordered partition of $[n]$ into k parts, given by the sequence of (random) fibers $F_{k,n}^{-1}(j)$ for $j \in [k]$. We denote by $\Delta^{n \times n}$ the set of $n \times n$ matrices with nonnegative entries that sum to one.

Definition 1.1. *A sequence of weighted graphs $(G_n \in \Delta^{n \times n})_{n=1}^\infty$ with edge weights that sum to one quotient-converges if for each $k \in \mathbb{N}$ the sequence of random graphs $(\rho(F_{k,n})G_n \in \Delta^{k \times k})_{n=1}^\infty$ converges weakly. A general sequence of nonzero weighted graphs $(G_n \in \mathbb{R}_+^{n \times n})_n$ quotient-converges if $(G_n/\mathbb{1}^\top G_n \mathbb{1} \in \Delta^{n \times n})_n$ quotient-converges.*

Normalizing our graphs to have unit total edge weight does not change the structure of its hubs, which were defined above in terms of the *fraction* of edge weight incident on them. Unless specified otherwise, we will assume throughout that the total edge weight of our graphs equals one.

Note that $\rho(F_{n,k})^*G_n$ is the induced subgraph on k vertices sampled with replacement from the vertex set $[n]$ of G_n , in terms of which graphon convergence is defined [14, §9], giving a first sense in which our theory is dual to that of graphons. We remark that there is an alternative view of graphon limits via convergence of quotients [14, §12], but these latter quotients differ from ours. In particular, these latter quotients cannot detect hubs in a large graph, as we explain in Section 1.2 below.

1.1.1 Combinatorial Aspects of Quotient Convergence (Section 2)

As our first contribution, we analyze combinatorial aspects of quotient convergence in Section 2. Whereas previous notions of graph convergence were based on convergence of local “motifs” in the form of homomorphism densities (appropriately normalized), we show that our quotient convergence is equivalent to convergence of homomorphism *numbers*. More precisely, for a multigraph $H \in \mathbb{N}^{k \times k}$ possibly containing parallel edges and self-loops, and a general weighted graph $G \in \mathbb{R}^{n \times n}$ on n vertices, the homomorphism number and homomorphism density of H in G are given by [14, Eq. (5.3)]

$$\text{hom}(H; G) = \sum_{f: [k] \rightarrow [n]} \prod_{i,j=1}^k G_{f(i), f(j)}^{H_{i,j}}, \quad t(H; G) = \frac{1}{n^k} \text{hom}(H; G). \quad (2)$$

The terminology comes from the fact that when H and G are simple graphs, then $\text{hom}(H; G)$ is the number of graph homomorphisms from H to G and $t(H; G)$ counts the average number of times H occurs in a uniformly random subgraph of G . In particular, convergence of the homomorphism densities $t(H; \cdot)$ is used to define limits of dense graphs [15, 16].

In analogy, we define the *quotient density* $t_Q(H; G)$ of a multigraph $H \in \mathbb{N}^{k \times k}$ in a general weighted graph $G \in \mathbb{R}^{n \times n}$ as the average number of times H occurs in a uniformly random quotient of G :

$$t_Q(H; G) = \frac{1}{k^n} \sum_{f: [n] \rightarrow [k]} \prod_{i,j=1}^k (\rho(f)G)_{i,j}^{H_{i,j}}. \quad (3)$$

The role of homomorphism densities in graphons is played by quotient densities in our framework. We prove that a sequence of graphs is quotient-convergent if all of their quotient densities converge.

Theorem 1.2. *For a sequence of graphs $(G_n \in \Delta^{n \times n})_n$, the following are equivalent:*

1. *The sequence (G_n) is quotient-convergent.*
2. *For each multigraph $H \in \mathbb{N}^{k \times k}$, the sequence of quotient densities $(t_Q(H; G_n))_n$ converges.*
3. *For each multigraph $H \in \mathbb{N}^{k \times k}$, the sequence of homomorphism numbers $(\text{hom}(H; G_n))_n$ converges.*

The proof of this result is given in Section 2 based on expressing quotient densities in terms of (injective) homomorphism numbers. This expression involves a partial order on the collection of multigraphs induced by the quotient operation.

For simple graphs $(G_n \in \mathbb{S}_{\text{sim}}^n)_n$, i.e., undirected graphs with weights $(G_n)_{i,j} \in \{0, 1\}$ and with no self-loops so $\text{diag}(G_n) = 0$, we conclude that $(G_n/2|E(G_n)|)_n$ is quotient-convergent if and only if $\left(\frac{\text{hom}(H; G_n)}{|E(G_n)|^{|E(H)|}}\right)_n$ converges for all simple graphs $H \in \mathbb{S}_{\text{sim}}^k$ (see Corollary 2.7). Thus, quotient-convergence of simple graphs is equivalent to convergence of a homomorphism-density-like quantity, but one that is differently normalized than usual. To the best of our knowledge, such a normalization based on the number of edges has not appeared previously in the literature, as normalizations based on the numbers of vertices k and n are used (see Section 1.2). As a consequence, we will see below in our discussion of limit objects (Corollary 3.7) that quotient-convergent sequences of both dense and degree-bounded sequences of simple graphs converge to the same (uninteresting) limit. In fact, quotient limits of simple graphs can only detect stars, which are vertices on which a positive fraction of edges are incident. More broadly, the utility of our framework lies in uncovering such hub structure in weighted graphs.

1.1.2 Limit Objects for Quotient Convergence (Section 3)

Our second contribution consists of constructing analytic limit objects together with a notion of distance between them that allows us to compare graphs of different sizes, akin to graphons and the cut metric for dense graph sequences. Note that appending isolated vertices to a graph $G \in \Delta^{n \times n}$ or relabeling its vertices does not change the distribution of its quotients $\rho(F_{k,n})G_n$ for each $k \in \mathbb{N}$. Therefore, the limit object we associate to G must be the same as that associated to any other graph differing from it by extra isolated vertices and vertex relabeling. We describe our limit objects in terms of certain random exchangeable measures on the square, which we define next.

Definition 1.3 (Random exchangeable measures). *A random exchangeable probability measure on $[0, 1]^2$ is a random measure² \mathbf{M} satisfying $\mathbf{M} \stackrel{d}{=} \mathbf{M} \circ (\sigma, \sigma)$ for all (Lebesgue-)measure-preserving bijections $\sigma: [0, 1] \rightarrow [0, 1]$. We say that a sequence of random measures (\mathbf{M}_n) converges if it converges weakly viewed as a sequence of random variables taking values in the space of probability measures on $[0, 1]^2$ (see Section 1.3 for details).*

Any finite graph $G \in \Delta^{n \times n}$ can be associated with the random exchangeable measure:

$$\mathbf{M}_G = \sum_{i,j=1}^n G_{i,j} \delta_{(T_i, T_j)}, \quad \text{where } T_1, \dots, T_n \stackrel{iid}{\sim} \text{Unif}([0, 1]). \quad (4)$$

Note that relabeling the vertices of G or appending isolated vertices to it yields the same random measure in this way. More generally, Kallenberg showed that all random exchangeable measures on $[0, 1]^2$ have the form [17]:

$$\mathbf{M} = \sum_{i,j \in \mathbb{N}} E_{i,j} \delta_{(T_i, T_j)} + \sum_{i \in \mathbb{N}} [\sigma_i(\delta_{T_i} \otimes \lambda) + \varsigma_i(\lambda \otimes \delta_{T_i})] + \theta \lambda^2 + \vartheta \lambda_D, \quad (5)$$

where (i) λ is the uniform measure on $[0, 1]$, λ^2 is the uniform measure on $[0, 1]^2$, and λ_D is the uniform measure on the diagonal $\{(x, x) : x \in [0, 1]\}$; (ii) $T_1, T_2, \dots \stackrel{iid}{\sim} \text{Unif}([0, 1])$; and (iii) $E_{i,j}, \sigma_i, \varsigma_i, \theta, \vartheta \geq 0$ may potentially be random but must be independent of the (T_i) and must satisfy $\sum_{i,j} E_{i,j} + \sum_i (\sigma_i + \varsigma_i) + \theta + \vartheta = 1$ almost surely. Moreover, the collection of extremal random exchangeable measures, i.e., those that are not mixtures of other such random measures, are precisely the ones for which the coefficients $E, \sigma, \varsigma, \theta, \vartheta$ in (5) can be taken to be deterministic.

²We endow the space of probability measures on $[0, 1]^2$ with the weak topology and consider Borel random variables on it.

Definition 1.4. A grapheur (graph edge measure) is an extremal random exchangeable measure, i.e., one of the form (5) with deterministic coefficients $(E, \sigma, \varsigma, \theta, \vartheta)$. We denote the collection of grapheurs by

$$\mathcal{M} = \left\{ \mathbf{M} \text{ of the form (5)} : E \in \mathbb{R}_{\geq 0}^{N \times N}, \sigma, \varsigma \in \mathbb{R}_{\geq 0}^N, \theta, \vartheta \in \mathbb{R}_{\geq 0} \text{ s.t. } \sum_{i,j} E_{i,j} + \sum_i (\sigma_i + \varsigma_i) + \theta + \vartheta = 1 \right\}. \quad (6)$$

The space of grapheurs inherits the weak topology from the ambient space of random exchangeable measures.

The coefficients $(E, \sigma, \varsigma, \theta, \vartheta)$ describing a fixed grapheur $\mathbf{M} \in \mathcal{M}$ are essentially unique. Specifically, note that for any permutation $\pi: \mathbb{N} \rightarrow \mathbb{N}$, the coefficients $(E, \sigma, \varsigma, \theta, \vartheta)$ and $(\pi E \pi^\top, \pi \sigma, \pi \varsigma, \theta, \vartheta)$ define the same random exchangeable distribution via (5). Further, adding zero rows to E and corresponding zero entries to σ , or zero columns to E and corresponding zero entries to ς , does not change the resulting random measure. Kallenberg showed in [17, Prop. 3] that these are the only possible ambiguities in the coefficients. In particular, note that $\mathbf{M}_G \in \mathcal{M}$ is a grapheur for any finite graph G , and that $\mathbf{M}_G = \mathbf{M}_{G'}$ for two such graphs G, G' if and only if they differ by isolated vertices and vertex relabeling.

In analogy with the cut metric between graphons, we endow \mathcal{M} with the following metric between grapheurs that metrizes their convergence:

$$W_\square(\mathbf{M}_1, \mathbf{M}_2) = \inf_{\substack{\text{random } (\mathbf{M}'_1, \mathbf{M}'_2) \\ \mathbf{M}'_i \stackrel{d}{=} \mathbf{M}_i \text{ for } i=1,2}} \sup_{\substack{S, T \subseteq [0,1] \\ \text{intervals}}} \mathbb{E}|\mathbf{M}'_1(S \times T) - \mathbf{M}'_2(S \times T)|, \quad (7)$$

where the infimum is over couplings $(\mathbf{M}'_1, \mathbf{M}'_2)$ of the two random measures \mathbf{M}_1 and \mathbf{M}_2 . Here $\mathbf{M}'_i \stackrel{d}{=} \mathbf{M}_i$ means that the distribution of \mathbf{M}'_i is equal to that of \mathbf{M}_i , see Section 1.3 for more detail. We show that grapheurs are the limits of quotient-convergent sequences of graphs in the following precise sense.

Theorem 1.5. A sequence of grapheurs $(\mathbf{M}_n) \subseteq \mathcal{M}$ converges weakly to a grapheur $\mathbf{M} \in \mathcal{M}$ if and only if $\lim_n W_\square(\mathbf{M}_n, \mathbf{M}) = 0$. Furthermore, the following statements hold.

1. A sequence of graphs $(G_n \in \Delta^{n \times n})$ is quotient-convergent if and only if there is a grapheur $\mathbf{M} \in \mathcal{M}$ satisfying $\lim_n W_\square(\mathbf{M}, \mathbf{M}_{G_n}) = 0$.
2. Any sequence of graphs $(G_n \in \Delta^{n \times n})$ has a quotient-convergent subsequence.
3. For every grapheur $\mathbf{M} \in \mathcal{M}$ there is a sequence $(G_n \in \Delta^{n \times n})$ satisfying $\lim_n W_\square(\mathbf{M}, \mathbf{M}_{G_n}) = 0$.

In view of Theorem 1.5, we say that a sequence of graphs (G_n) converges to a grapheur $\mathbf{M} \in \mathcal{M}$ if $\lim_n \mathbf{M}_{G_n} = \mathbf{M}$ in \mathcal{M} . Theorem 1.5 states that any quotient-convergent sequence of graphs converges to a grapheur, and conversely each grapheur is the limit of a quotient-convergent sequence of graphs. Moreover, this theorem shows that the space of grapheurs endowed with the weak topology is compact, and that this compact topology is metrized by our W_\square metric (7).

To prove Theorem 1.5, we relate random quotients of finite graphs to their associated grapheurs (4). For each grapheur $\mathbf{M} \in \mathcal{M}$ and $k \in \mathbb{N}$, define the random graph $\mathbf{G}_k[\mathbf{M}] \in \Delta^{k \times k}$ whose entries are given by:

$$(\mathbf{G}_k[\mathbf{M}])_{i,j} = \mathbf{M}(I_i^{(n)} \times I_j^{(n)}) \quad (8)$$

for $i, j \in [n]$, where $I_i^{(n)} = [(i-1)/n, i/n]$ for $i \in [n]$.³ In words, we partition the square $[0, 1]^2$ into a $k \times k$ uniform grid, and let the entries of $\mathbf{G}_k[\mathbf{M}]$ be the (random) measure under \mathbf{M} of each square in this grid. We show in Section 3.1 that $\rho(F_{k,n})G \stackrel{d}{=} \mathbf{G}_k[\mathbf{M}_G]$ for any finite graph $G \in \Delta^{n \times n}$, and that a sequence of graphs (G_n) converges to the grapheur \mathbf{M} precisely when the random quotients $(\rho(F_{k,n})G_n)_k$ converge weakly to $\mathbf{G}_k[\mathbf{M}]$ for each $k \in \mathbb{N}$. Similarly, we can extend quotient densities to grapheurs by setting

$$t_Q(H; \mathbf{M}) = \mathbb{E} \prod_{i,j=1}^k (\mathbf{G}_k[\mathbf{M}])_{i,j}^{H_{i,j}} = \int \prod_{i,j \in [k]} \mu(I_i^{(k)} \times I_j^{(k)})^{H_{i,j}} d\mathbf{M}(\mu), \quad (9)$$

³Note that $\sum_{i,j} (\mathbf{G}_k[\mathbf{M}])_{i,j} = \mathbf{M}([0, 1]^2) = 1$ almost surely by (5).

for any multigraph $H \in \mathbb{N}^{k \times k}$, analogously to the extension of homomorphism densities to graphons [14, §7.2]. In terms of quotient densities, the sequence (G_n) converges to the grapheur \mathbf{M} if and only if $t_Q(H; G_n) \rightarrow t_Q(H; \mathbf{M})$ for all multigraphs H , see Section 3.1.

Having characterized our limit objects, we interpret in Section 3.2 each of the parameters $E, \sigma, \varsigma, \theta, \vartheta$ describing a grapheur (5) in terms of certain features of the large graphs converging to it. Informally, the array $E \in \mathbb{R}_{\geq 0}^{\mathbb{N} \times \mathbb{N}}$ specifies the connections between a discrete collection of hubs, with each entry corresponding to an edge with an asymptotically positive fraction of the total edge weight. The arrays $\sigma, \varsigma \in \mathbb{R}_{\geq 0}^{\mathbb{N}}$ specify directed stars centered at hubs, with each entry corresponding to the fraction of out-degree or in-degree of a hub from all the edges incident on it with asymptotically vanishing edge weights. Lastly, the parameters θ and ϑ specify the remaining edge weight that comes from all edges between vertices with asymptotically vanishing degrees and their self-loops. In particular, we say that a grapheur \mathbf{M} does not have hubs if $E = 0$ and $\sigma = \varsigma = 0$, and we say that a sequence (G_n) of graphs does not contain hubs asymptotically if it quotient-converges to such a grapheur.

Finally, we expand on the duality between graphons and grapheurs. We have seen previously that the sampling operators used to define each notion of graph limit, namely, those extracting random subgraphs and random quotients, respectively, are duals of each other. We further show in Section 3.3 that the space of graphons endowed with the cut metric is dual to the space \mathcal{M} endowed with the W_\square metric (7) in a precise sense.

Theorem 1.6 (Informal; see Theorem 3.10). *There is a random pairing $\langle \mathbf{M}, W \rangle$ associating a scalar random variable to each grapheur \mathbf{M} and graphon W satisfying the following properties. We have $\lim_n \mathbf{M}_n = \mathbf{M}$ in the W_\square metric if and only if $\langle \mathbf{M}_n, W_G \rangle \rightarrow \langle \mathbf{M}, W_G \rangle$ weakly for each step graphon W_G associated to a finite simple graph G . We also have $\lim_n W_n \rightarrow W$ in the cut metric if and only if $\langle \mathbf{M}_G, W_n \rangle \rightarrow \langle \mathbf{M}_G, W \rangle$ weakly for each grapheur \mathbf{M}_G associated to a finite undirected graph G with unit total edge weight and no self-loops.*

1.1.3 Edge Sampling and Property Testing (Section 4)

As our next contribution, we apply our framework to test properties of large graphs based on small summaries of them. Specifically, we focus on testing properties given by graph parameters that are continuous with respect to quotient convergence. While it would be natural to test such properties using random quotients, forming these requires access to the entire large graph, which limits their utility in practice. Motivated by this challenge, we develop an edge-based sampling perspective on quotient-convergence in Section 4.1. Specifically, we sample edges from a graph proportionally to their edge weights to form random graphs $\mathbf{G}^{(n)}$ with n edges, and extend this sampling procedure to any grapheur $\mathbf{M} \in \mathcal{M}$. We detail the general sampling procedure in Section 4.1.

We prove the following convergence results for the sequence of random graphs $(\mathbf{G}^{(n)})$ sampled from a grapheur \mathbf{M} back to \mathbf{M} , which are edge-based analogs of the second sampling lemma and Szemerédi's regularity lemma for graphons [14, Lemma 10.16, Lemma 9.15]. Here and throughout the paper, a superscript $G^{(n)}$ denotes a graph on n edges (and hence at most $2n$ vertices) while a subscript G_n denotes a graph on n vertices.

Theorem 1.7. *Fix any grapheur $\mathbf{M} \in \mathcal{M}$.*

(Edge sampling lemma) *For each $n \in \mathbb{N}$, let $\mathbf{G}^{(n)}$ be the random graph containing n edges sampled from \mathbf{M} as in Section 4. With probability at least $1 - e^{-2\epsilon^2}$, we have that $W_\square(\mathbf{M}_{\mathbf{G}^{(n)}}, \mathbf{M}) \leq \frac{174+\epsilon}{\sqrt{n}}$.*

(Edge-based Szemerédi regularity) *For any $\epsilon > 0$, there exists a graph $G^{(k(\epsilon))}$ on $k(\epsilon) = \lceil (174/\epsilon)^2 \rceil$ edges satisfying $W_\square(\mathbf{M}, \mathbf{M}_{G^{(k(\epsilon))}}) \leq \epsilon$.*

Theorem 1.7 allows us to test properties of large graphs by sampling a few of their edges at random. We formally define the graph parameters that can be tested using our framework in Section 4.3. These consist of graph parameters that are continuous with respect to quotient convergence, and include all polynomial graph parameters that do not change when appending isolated vertices to their inputs (Proposition 4.9). In particular, the graph parameter $f(G) = \|G\mathbf{1}\|_2^2 + \|G^\top \mathbf{1}\|_2^2$ is quotient-testable in our framework, and can be used to test for the presence of hubs in a growing sequence of graphs. Specifically, if (G_n) is a quotient-convergent sequence, we show in Example 4.11 that $f(G_n) \rightarrow 0$ precisely when (G_n) does not have hubs

asymptotically, i.e., it converges to a limiting grapheur (5) with $E = 0$ and $\sigma = \varsigma = 0$. We then describe an edge-based sampling procedure for testing this graph parameter. In contrast, this parameter cannot be tested in the graphons framework; see Section 4.3.

1.1.4 Equipartition-Consistent Random Graph Models (Section 5)

As our final contribution, we observe that random quotients $(G_k[M])_k$ of any grapheur M are related to each other by equipartitions of their vertex sets. Specifically, we observe that $\rho(d_{k,nk})G_{nk}[M] \stackrel{d}{=} G_k[M]$ for any equipartition $d_{k,nk}: [nk] \rightarrow [k]$, i.e., a map with equal-sized fibers. We call such sequences of random graphs *equipartition-consistent random graph models*, which play a role in our framework analogous to that played by consistent random graph models in graphon theory [18].

Definition 1.8 (Equipartition-consistent models). *A sequence of probability measures $(\mu_k \in \mathcal{P}(\Delta^{k \times k}))_{k \in \mathbb{N}}$ is an equipartition-consistent random graph model if $\rho(d_{k,nk})G_{nk} \stackrel{d}{=} G_k$ whenever $G_{nk} \sim \mu_{nk}$ and $G_k \sim \mu_k$ and for any equipartition $d_{k,nk}: [nk] \rightarrow [n]$.*

Further, we show that any equipartition-consistent random graph model is derived from a mixture of grapheurs.

Theorem 1.9. *For any equipartition-consistent random graph model $(\mu_k \in \mathcal{P}(\Delta^{k \times k}))_k$, there is a unique random exchangeable measure M on $[0, 1]^2$ satisfying $\mu_k = \text{Law}(G_k[M])$ for all $k \in \mathbb{N}$.*

Recall that grapheurs were defined as extremal random exchangeable measures in Definition 1.4. By Choquet's theorem, any random exchangeable measure is therefore a mixture of grapheurs. Finally, we show in Proposition 5.1 that the sequence $(G_k[M])$ of quotients of a grapheur M converge back to M in a certain sense.

1.2 Related Work

There are several well-studied notions of graph limits and random graph models in the literature. We review this body of work and discuss similarities and differences with the present paper.

Dense and degree-bounded graph limits. The earliest two types of graph limits pertain to sequences of simple graphs that are dense, whose limits are given by graphons [15, 16], and degree-bounded, whose limits are given by graphings [19–21]. We refer the reader to [14] for a survey of both limits. Both of these notions of graph limits are incomparable to ours, as both dense and degree-bounded sequences of simple graphs converge to the same limit in our framework, see Corollary 3.7. Further, dense graph limits can be characterized in terms of quotients [14, §12], called “right convergence” of graphons. However, these quotients are differently normalized compared to ours, and it is the set of quotients that converges in the graphon framework rather than their distribution as in our framework. Despite these distinctions, we highlight parallels between our development and the theory of graphons throughout our paper; in particular, grapheurs can be viewed as dual to graphons in a precise sense, see Section 3.3.

Edge-exchangeable random graphs. A class of random graph models closely related to the ones we obtain by sampling edges from a grapheur is called edge-exchangeable [22–26]. These random graphs are obtained by adding random edges to a fixed infinite vertex set (often with repetition, yielding multigraphs), and can yield both sparse and dense random graphs [26]. As explained in [27, Rmk. 4.4], every edge-exchangeable multigraph model can be obtained by sampling edges from a random exchangeable measure on $[0, 1]^2$. In fact, the random graphs we obtain by sampling edges in Section 4 are precisely these edge-exchangeable multigraphs, normalized to have unit edge weight, see Remark 4.2. From this perspective, convergence of grapheurs can be interpreted as convergence of the edge-exchangeable multigraph models associated to them, an interpretation we make precise in Proposition 4.1. Nevertheless, to our knowledge this notion of convergence of edge-exchangeable models has not been previously studied in the literature. In particular, the only metric between edge-exchangeable models we are aware of is an ℓ_1 distance between edge probabilities [23, Eq. (8)]. Convergence in this metric is too restrictive for our purposes, see Proposition 3.8. Further, our development of convergence via random quotients, which are not edge-exchangeable, allows us to take limits of general weighted graphs, to relate it to convergence of homomorphism densities and other graph parameters, and to test their values.

Graphex processes and limits. The closest line of work to ours is the study of graphex processes and their associated graph limits and random graph models [28–34]. Graphexes form a subset of random exchangeable measures on $\mathbb{R}_{\geq 0}^2$, and admit a characterization due to Kallenberg similarly to our grapheurs [17]. Graphexes give rise to random graph models obtained by sampling edges from them [29, 32, 35], analogously to our edge sampling procedure in Section 4.1. There are also several notions of limits for simple graphs and multigraphs, whereby such graphs converge to a limiting graphex [30, 33]. These notions of limits are conjectured to be inequivalent [34], and the space of graphexes with either topology is not compact. Some of these limits can be characterized by sampling random subgraphs (whose number of vertices is also random) [32], while others can be characterized by convergence of differently-normalized homomorphism numbers [30, §2.5] (the authors of [30] normalize $\text{hom}(H; G)$ by $(2|E(G)|)^{|V(H)|/2}$, whereas we normalize by $(2|E(G)|)^{|E(H)|}$). Graphex-based graph limits are again incomparable to ours. On the one hand, for a sequence of simple graphs our grapheur-based limits can only distinguish stars from a uniform component (Corollary 3.7), whereas graphex-based limits can capture more general limiting structures (see [34, §9] or [33]). On the other hand, grapheur-based graph limits are defined for general sequences of nonnegatively-weighted graphs, for which graphex-based limits have not been defined, allowing us to test properties pertaining to the distribution of edge weight in the graph (see Section 4.3). In fact, grapheurs are particularly well-suited for property testing, since the space of grapheurs we study here is compact, in contrast to the space of graphexes; to our knowledge, there is no characterization of properties that can be tested using graphex-based limits. Another consequence of the compactness of the space of grapheurs is that our notion of graph limits can be characterized in several equivalent ways: by convergence of random quotients, convergence of random edge-sampled graphs, and convergence of homomorphism numbers and related graph parameters.

1.3 Notation and Preliminaries

We denote by $\mathbb{N} = \{0, 1, 2, \dots\}$ the set of nonnegative integers. If $n \in \mathbb{N}$, then $[n] = \{1, \dots, n\}$. If $a \leq b$ are real numbers then $[a, b] = \{x \in \mathbb{R} : a \leq x \leq b\}$ and $[a, b) = [a, b] \setminus \{b\}$. We denote by λ the uniform measure on $[0, 1]$, by λ^2 the uniform measure on $[0, 1]^2$, and by λ_D the uniform measure on the diagonal $D = \{(x, x) : x \in [0, 1]\}$. We denote by $\mathbb{1}_n$ the vector of all-1's of length n , and we omit the subscript n when the size (possibly infinite) of the all-1's vector is clear from context. We denote by I_n the $n \times n$ identity matrix and by $\mathbb{1}_{n \times n} = \mathbb{1}_n \mathbb{1}_n^\top$ the $n \times n$ all-1's matrix.

We identify a (directed, weighted) graph with its adjacency matrix $G \in \mathbb{R}_{\geq 0}^{n \times n}$, where $G_{i,j}$ is the weight of the edge $i \rightarrow j$. The collection of all adjacency matrices of graphs with unit total edge weight is denoted by $\Delta^{n \times n}$. In this paper, all edge weights are nonnegative and sum to one unless stated otherwise. A graph is undirected if its adjacency matrix is symmetric. We denote the space of $n \times n$ symmetric matrices by \mathbb{S}^n , and the set of such matrices with nonnegative entries, which is the set of undirected graphs, by $\mathbb{S}_{\geq 0}^n$. An undirected graph G is simple if $\text{diag}(G) = 0$ and $G_{i,j} \in \{0, 1\}$ for all i, j , and we denote the set of simple graphs by $\mathbb{S}_{\text{sim}}^n$. A graph H is called a *multigraph* if it has integer edge weights $H \in \mathbb{N}^{k \times k}$, where we view $H_{i,j}$ as representing the number of parallel edges from i to j . Note that a simple graph is a multigraph. We denote by $|E(G)| = \mathbb{1}^\top G \mathbb{1}/2$ the number of edges in a *simple* graph G . If H is a multigraph, we write $\mathbb{1}^\top H \mathbb{1} = \|H\|_1$ to be the number of (directed) edges in H . We assume that multigraphs have no isolated vertices, defined as vertices with no self-loops and no edges incident on them. In other words, if $H \in \mathbb{N}^{k \times k}$ is a multigraph, then for every $i \in [k]$ there is a $j \in [k]$ such that $H_{i,j} > 0$ or $H_{j,i} > 0$. If $G \in \mathbb{R}_{\geq 0}^{n \times n}$ and $H \in \mathbb{N}^{k \times k}$ with $k \leq n$, we denote by $G^H = \prod_{i,j=1}^k G_{i,j}^{H_{i,j}}$ the monomial over $\mathbb{R}[G_{i,j}]$ defined by H . In this notation, we have $t_Q(H; G) = \mathbb{E}(\rho(F_{k,n})G)^H$ for a multigraph $H \in \mathbb{N}^{k \times k}$ and graph $G \in \Delta^{n \times n}$.

We denote by S_n the group of permutations of $[n]$, and by S_∞ the group of permutations of \mathbb{N} that fixes all but finitely-many integers. If $\pi \in \mathsf{S}_n$ and $G \in \mathbb{R}_{\geq 0}^{n \times n}$ is a graph, we denote by $\pi G \in \mathbb{R}_{\geq 0}^{n \times n}$ the permuted graph with entries $(\pi G)_{i,j} = G_{\pi^{-1}(i), \pi^{-1}(j)}$. This corresponds to relabeling the vertices of G according to π . Similarly, if $x \in \mathbb{R}^n$ is a vector we denote by $\pi x \in \mathbb{R}^n$ the permuted vector with entries $(\pi x)_i = x_{\pi^{-1}(i)}$. If $G \in \mathbb{R}_{\geq 0}^{\mathbb{N} \times \mathbb{N}}$ and $x \in \mathbb{R}^\mathbb{N}$ are infinite matrices and vectors, and $\pi \in \mathsf{S}_\infty$, we define πG and πx analogously.

If $\mu \in \mathcal{P}(S)$ is a probability measure on a compact Polish space S , we write $X \sim \mu$ and $\text{Law}(X) = \mu$ to denote a random variable distributed according to μ . If X, Y are two random variables on S , we write $X \stackrel{d}{=} Y$ to denote equality in distribution, i.e., the equality $\text{Law}(X) = \text{Law}(Y)$. If f is a function, we denote $\mathbb{E}_\mu f = \mathbb{E}_{X \sim \mu} f(X)$ its expectation with respect to μ . If (μ_n) is a sequence of measures in $\mathcal{P}(S)$, we say (μ_n)

converges weakly to μ and write $\mu_n \rightarrow \mu$ if $\mathbb{E}_{\mu_n} f \rightarrow \mathbb{E}_\mu f$ for all continuous f . All measures in this paper are supported in a single ambient compact set. In particular, all sequences of measures are tight.

A coupling of random variables X_1, \dots, X_k taking values in Polish spaces S_1, \dots, S_k , respectively, is a probability distribution $\Gamma \in \mathcal{P}(S_1 \times \dots \times S_k)$ such that if $(X'_1, \dots, X'_k) \sim \Gamma$ then $X'_i \stackrel{d}{=} X_i$ for all $i \in [k]$. In this paper we refer to a coupling of X_1, \dots, X_k simply by the tuple of jointly-distributed random variables (X'_1, \dots, X'_k) . If X_1, X_2 are random graphs taking values in $\mathbb{R}^{k \times k}$, we define the Wasserstein-1 distance between them by

$$W_1(X_1, X_2) = \inf_{\substack{\text{random } (X'_1, X'_2) \\ X'_i \stackrel{d}{=} X_i}} \mathbb{E} \|X'_1 - X'_2\|_1 = \sup_{\substack{f: \mathbb{R}^{k \times k} \rightarrow \mathbb{R} \\ \text{1-Lipschitz}}} |\mathbb{E} f(X_1) - \mathbb{E} f(X_2)|,$$

where the infimum is over couplings (X'_1, X'_2) of X_1 and X_2 , the norm $\|\cdot\|_1$ is the entrywise ℓ_1 norm, and the supremum is over 1-Lipschitz functions with respect to this norm. Note that a sequence of random graphs $(X_n) \subseteq \Delta^{k \times k}$ converges weakly to the random graph $X \in \Delta^{k \times k}$ if and only if $\lim_n W_1(X_n, X) = 0$ by [36, Thm. 6.9].

For a compact Polish space S , the space $\mathcal{P}(S)$ endowed with the weak topology is itself compact and Polish. If M is a random measure, that is, a random variable taking values in the Polish space $\mathcal{P}(S)$, we write $\mathbb{E}_M f$ for the scalar random variable obtained by integrating a deterministic function f against the random measure M . In particular, we write $M(B) = \mathbb{E}_M \mathbb{1}_B$ to denote the scalar random variable giving the (random) measure of a measurable set B , where $\mathbb{1}_B(x) = 1$ if $x \in B$ and zero otherwise. A sequence of random measures (M_n) converges weakly to M if $\text{Law}(M_n) \rightarrow \text{Law}(M)$ weakly, where these laws are probability measures over the space of probability measures. Equivalently, the sequence (M_n) converges weakly to M if the sequence of scalar random variables $\mathbb{E}_{M_n} f$ converges weakly to $\mathbb{E}_M f$ for all continuous functions f [37, Thm. A2.3].

2 Quotient Convergence: Global and Local Views

In this section we prove Theorem 1.2 relating quotient convergence to convergence of homomorphism numbers and quotient densities. Before presenting this proof, we give a few examples illustrating sequences of graphs that are quotient-convergent. As our first example, we show that the sequence of complete graphs is quotient-convergent.

Example 2.1 (Convergence of complete graphs). *To show that the sequence $(\mathbb{1}_{n \times n})_{n \in \mathbb{N}}$ of complete graphs converges, consider the associated normalized graphs $(G_n = \mathbb{1}_{n \times n}/n^2 \in \Delta^{n \times n})$ and observe that the random quotients are given as $\rho(F_{k,n})G_n = \frac{1}{n^2}NN^\top$ for a multinomial random variable $N = (N_1, \dots, N_k) \sim \text{Multinom}(n, k, \mathbb{1}_k/k)$. Since $\mathbb{E}\rho(F_{k,n})G_n = \frac{1}{k^2}\mathbb{1}_{k \times k} + O(1/n)$ and $\text{Var}[\rho(F_{k,n})G_n]_{i,j} = O(1/n)$ for all $i, j \in [k]$, we conclude that $\rho(F_{k,n})G_n$ converges weakly to the deterministic matrix $\frac{1}{k^2}\mathbb{1}_{k \times k}$ as $n \rightarrow \infty$. Since this holds for all $k \in \mathbb{N}$, we conclude that (G_n) is quotient-convergent. Note that the quotient densities and homomorphism numbers of this sequence likewise converge. Indeed, for any $H \in \mathbb{N}^{k \times k}$ we have $t_Q(H; G_n) = \mathbb{E}(\rho(F_{k,n})G_n)^H \xrightarrow{n \rightarrow \infty} (\frac{1}{k^2}\mathbb{1}_k\mathbb{1}_k^\top)^H = \frac{1}{k^2\|H\|_1}$. Similarly, $\text{hom}(H; G_n) = \frac{n^k}{n^2\|H\|_1}$, which converges to 1 if H is a disjoint union of edges and to 0 otherwise. An analogous set of arguments show that the sequence (I_n) of n self-loops is quotient-convergent.*

As our next example, we consider sequences of (directed) star graphs and we show that these are quotient-convergent as well. Note that stars are sparse graphs, as their edge density converges to zero. Nevertheless, we will see in Section 3 that growing-sized stars have nontrivial limits in our framework, whereas the limits of stars in the context of graphons yield trivial limiting objects. Similarly, stars are not degree-bounded, as the degree of the star center increases with the size of the star, hence they do not have a limit in the framework of graphings. Stars do have limits in the graphex sense, though they differ from ours, see Section 1.2.

Example 2.2 (Convergence of stars). *The sequence of directed star graphs all of whose edges point away from the center may be expressed as $(e_1\mathbb{1}_n^\top)$ for $e_1 = (1, 0, \dots, 0)^\top$. This sequence quotient-converges, since random quotients $\rho(F_{k,n})G_n$ of the associated normalized graphs $G_n = \frac{1}{n}e_1\mathbb{1}_n^\top$ are given by $\rho(F_{k,n})G_n = \frac{1}{n}e_1N^\top$ for a multinomial vector $N \sim \text{Multinom}(n, k, \mathbb{1}_k/k)$ and a uniformly random index $I \sim \text{Unif}([k])$. Since*

N/n weakly converges to the deterministic vector $\mathbb{1}_k/k$, we conclude that $\rho(F_{k,n})G_n$ converges weakly to the randomly-labeled star $\frac{1}{k}e_I\mathbb{1}_k^\top$, and hence that (G_n) is quotient-convergent. Similarly, the quotient densities and homomorphism numbers of this sequence of stars converge, since

$$\lim_{n \rightarrow \infty} t_Q(H; G_n) = \mathbb{E} \left(\frac{1}{k}e_I\mathbb{1}_k^\top \right)^H = \begin{cases} \frac{1}{k^{1+\sum_j m_j}} & \text{if } H = e_i m^\top \text{ for } m \in \mathbb{N}^k \text{ and } i \in [k], \\ 0 & \text{otherwise,} \end{cases}$$

and

$$\lim_{n \rightarrow \infty} \text{hom}(H; G_n) = \begin{cases} \lim_{n \rightarrow \infty} \frac{n^{k-1}}{n^{\sum_j m_j}} & \text{if } H = e_i m^\top, \\ 0 & \text{otherwise,} \end{cases} = \begin{cases} 1 & \text{if } H = e_i m^\top \text{ with } \sum_j m_j = k-1, \\ 0 & \text{otherwise.} \end{cases}$$

Analogous arguments show that the sequence of directed star graphs all of whose edges point towards the center quotient-converges, and so does the sequence of undirected stars.

As our final example of quotient-convergence, we consider an infinite weighted graph with edge weights summing to one and we show that any sequence of finite subgraphs obtained from this infinite graph is quotient-convergent.

Example 2.3 (Subgraphs of an infinite weighted graph). Suppose $G_\infty = (G_{i,j})_{i,j \in \mathbb{N}}$ is an infinite two-dimensional array with $G_{i,j} \geq 0$ and $\sum_{i,j} G_{i,j} = 1$, and consider the sequence of normalized truncations $G_n = \sum_{i,j=1}^n \frac{1}{G_{i,j}} (G_{i,j})_{i,j \in [n]}$. This sequence of subgraphs (G_n) is quotient-convergent. Indeed, let $F_k: \mathbb{N} \rightarrow [k]$ be a uniformly random map (whose images $F_k(i)$ are iid uniformly distributed over $[k]$), and consider the random graph $L_k = \rho(F_k)G_\infty$ whose entries are given by (1). We claim that the sequence $(\rho(F_{k,n})G_n)_n$ converges weakly to L_k . To this end, view G_n as an infinite array by zero-padding it, and note that $(\rho(F_k)G_n, \rho(F_k)G_\infty)$ is a coupling of $(\rho(F_{k,n})G_n, L_k)$. Using this coupling, we obtain the bound

$$W_1(\rho(F_{k,n})G_n, L_k) \leq \mathbb{E} \|\rho(F_k)(G_n - G_\infty)\|_1 \leq \|G_n - G_\infty\|_1 \xrightarrow{n \rightarrow \infty} 0,$$

where the second inequality follows from (1). Thus, we conclude that (G_n) is quotient-convergent, as claimed. Quotient densities and homomorphism numbers of this sequence again converge, with limits $\lim_n t_Q(H; G_n) = \mathbb{E} L_k^H$ and $\lim_n \text{hom}(H; G_n) = \sum_{f: [k] \rightarrow \mathbb{N}} \prod_{i,j=1}^k G_{f(i), f(j)}^{H_{i,j}}$, which is finite since $(G_{i,j})$ is summable.

Finally, we remark that convex combinations of quotient-convergent sequences also converge.

Example 2.4 (Combinations of graphs). Suppose (G_n) and (G'_n) are quotient-convergent sequences of graphs. Then for any $\theta \in [0, 1]$ the convex combination $(\theta G_n + (1 - \theta)G'_n)$ of these graphs is quotient-convergent as well. Indeed, for each $k \in \mathbb{N}$, let L_k and L'_k be the weak limits of $\rho(F_{k,n})G_n$ and $\rho(F_{k,n})G'_n$, respectively. For each $n \in \mathbb{N}$, let $(\tilde{L}_k, \tilde{F}_{k,n})$ and $(\tilde{L}'_k, \tilde{F}'_{k,n})$ be couplings of $(L_k, F_{k,n})$ and $(L'_k, F_{k,n})$, respectively, attaining the Wasserstein-1 distances $W_1(L_k, \rho(F_{k,n})G_n)$ and $W_1(L'_k, \rho(F_{k,n})G'_n)$. We define a coupling of $\theta L_k + (1 - \theta)L'_k$ and $\rho(F_{k,n})(\theta G_n + (1 - \theta)G'_n)$ as follows. Draw $F_{k,n}$ uniformly at random, and draw \tilde{L}_k from the coupling $(\tilde{L}_k, \tilde{F}_{k,n})$ conditioned on $\tilde{F}_{k,n} = F_{k,n}$. Draw \tilde{L}'_k similarly and independently. Then

$$\begin{aligned} W_1(\theta L_k + (1 - \theta)L'_k, \rho(F_{k,n})(\theta G_n + (1 - \theta)G'_n)) &\leq \mathbb{E} \|\theta \tilde{L}_k + (1 - \theta) \tilde{L}'_k - \rho(F_{k,n})(\theta G_n + (1 - \theta)G'_n)\|_1 \\ &\leq \theta W_1(L_k, \rho(F_{k,n})G_n) + (1 - \theta) W_1(L'_k, \rho(F_{k,n})G'_n) \xrightarrow{n \rightarrow \infty} 0. \end{aligned}$$

This proves the claim. An analogous argument shows that a convex combination of any finitely-many quotient-convergent sequences of graphs is also quotient-convergent.

Having seen a few examples of quotient convergence, we proceed to prove Theorem 1.2. We begin by showing the first part of that theorem on the equivalence between quotient convergence and convergence of quotient densities.

Proposition 2.5. A sequence of graphs $(G_n \in \Delta^{n \times n})$ is quotient-convergent if and only if $(t_Q(H; G_n))_n$ converges for each $H \in \mathbb{N}^{k \times k}$ and $k \in \mathbb{N}$.

Proof. By definition, the sequence $(G_n \in \Delta^{n \times n})$ is quotient-convergent when the sequence $(\rho(F_{k,n})G_n \in \Delta^{k \times k})_n$ converges weakly for each $k \in \mathbb{N}$. Since $\Delta^{k \times k}$ is compact, the latter sequence converges weakly if and only if all its associated sequences of moments converge. Now observe that the monomials in $\mathbb{R}[X_{i,j}]_{i,j \in [k]}$ all have the form $X^H = \prod_{i,j} X_{i,j}^{H_{i,j}}$ for some $H \in \mathbb{N}^{k \times k}$, and that the corresponding moment of $\rho(F_{k,n})G_n$ is precisely $\mathbb{E}(\rho(F_{k,n})G_n)^H = t_Q(H; G_n)$ by definition of t_Q in (3). This yields the desired conclusion. \square

This proves the first part of Theorem 1.2. To show the second part relating quotient convergence with convergence of homomorphism numbers, we express homomorphism numbers in terms of quotient densities and vice-versa. To this end, we begin by expressing quotient densities in terms of injective homomorphism numbers using [38, Prop. 6.8]. This expression is stated in terms of a partial order on multigraphs induced by quotients. Specifically, for two multigraphs $K \in \mathbb{N}^{n \times n}, H \in \mathbb{N}^{m \times m}$ we say that K is a *refinement* of H , denoted $K \leq_R H$, if H is a quotient of K , i.e., if there exists a surjective map $f: [n] \rightarrow [m]$ such that $H = \rho(f)K$. We further let $R_{K,H}$ be the number of such maps f . Note that $R_{H,H} = |\text{Aut}(H)|$ for any multigraph H . The poset of all multigraphs with the refinement order is a disjoint union of posets, one for each number of edges. The poset of all multigraphs with n edges has a minimal element, namely, the disjoint union of n edges, and a maximal element, namely, a single vertex with n self-loops. We illustrate the poset of all multigraphs with two edges in Figure 3.




Figure 3: Hasse diagram for the refinement poset consisting of multigraphs with two edges. Here there is an upward path from K to H precisely when $K \leq_R H$.

The quotient density $t_Q(H; G)$ can be expressed in terms of a linear combination of injective homomorphism numbers over all refinements of H [38, Prop. 6.8]:

$$t_Q(H; G) = \sum_{K \leq_R H} |V(H)|^{-|V(K)|} \frac{H!}{K!} \frac{R_{K,H}}{R_{K,K} R_{H,H}} \text{inj}(K; G), \quad (10)$$

where $H! = \prod_{i,j=1}^k (H_{i,j}!)$ and similarly for $K!$. Here $\text{inj}(K; G)$ is defined similarly to $\text{hom}(K; G)$ in (2) but with a sum only over injective maps. The expression (10) states that H is a subgraph of some quotient of G , i.e., $t_Q(H; G) > 0$, if and only if some refinement of H is a subgraph of G itself, i.e., $\text{inj}(K; G) > 0$.

We are now ready to prove the second part of Theorem 1.2. The following argument is a simple modification of the one from [38, Prop. 6.8].

Proposition 2.6. *A sequence of graphs $(G_n \in \Delta^{n \times n})$ is quotient-convergent if and only if $(\text{inj}(H; G_n))$ converges for each $H \in \mathbb{N}^{k \times k}$ and $k \in \mathbb{N}$, which is equivalent to $(\text{hom}(H; G_n))$ converging for each $H \in \mathbb{N}^{k \times k}$ and $k \in \mathbb{N}$.*

Proof. For each $m \in \mathbb{N}$, define the matrix M indexed by all multigraphs containing m edges whose entries are given by $M_{H,K} = |V(H)|^{-|V(K)|} \frac{H!}{K!} \frac{R_{K,H}}{R_{K,K} R_{H,H}}$ if $K \leq_R H$ and $M_{H,K} = 0$ otherwise. This is a lower-triangular matrix with respect to the refinement partial order, and has positive diagonal entries $M_{H,H} =$

$(|V(H)|^{|V(H)|} |\text{Aut}(H)|)^{-1}$, hence it is invertible. Thus, we can write

$$\text{inj}(H; G) = \sum_{H \leq_R K} (M^{-1})_{H,K} t_Q(K; G). \quad (11)$$

Based on (11) and (10), we conclude that $(t_Q(H; G_n))$ converges for all multigraphs H if and only if $(\text{inj}(H; G_n))$ converges for all multigraphs H . Injective and non-injective homomorphism numbers can be similarly related, with one of the quantities being expressed in terms of linear combinations of the other as in [14, §5.2.3]. For example, we have [38, Eq. (73)] that

$$\text{hom}(H; G) = \sum_{H \leq_R K} \frac{R_{H,K}}{R_{K,K}} \text{inj}(K; G), \quad (12)$$

where the sum is over multigraphs that H refines. This expression can be inverted as above to obtain injective homomorphism numbers in terms of non-injective homomorphism numbers. This allows us to conclude that convergence of $(\text{inj}(H; G_n))$ for all multigraphs H is equivalent to convergence of $(\text{hom}(H; G_n))$ for all multigraphs H . \square

Combined together, Propositions 2.5 and 2.6 complete the proof of Theorem 1.2. We illustrate the utility of these results next by expressing quotient convergence of simple graphs in terms of appropriately-normalized homomorphism numbers. The resulting expression allows us to conclude that all dense simple graphs are quotient-convergent; in fact, we prove that all the homomorphism numbers of dense simple graphs converge to the same limit (see also Corollary 3.7 below).

Corollary 2.7. *Let $(G_n \in \mathbb{S}_{\text{sim}}^n)$ be a sequence of simple graphs.*

(Convergence of simple graphs) *The sequence (G_n) is quotient-convergent if and only if $\left(\frac{\text{hom}(H; G_n)}{|E(G_n)|^{|E(H)|}}\right)$ converges for all simple and connected graphs $H \in \mathbb{S}_{\text{sim}}^k$.*

(Convergence of dense graphs) *Suppose $\liminf_{n \rightarrow \infty} \frac{|E(G_n)|}{n^2} > 0$. Then (G_n) is quotient-convergent. In fact, for a simple and connected $H \in \mathbb{S}_{\text{sim}}^k$, we have*

$$\lim_{n \rightarrow \infty} \frac{\text{hom}(H; G_n)}{|E(G_n)|^{|E(H)|}} = \begin{cases} 1 & \text{if } H = K_2, \\ 0 & \text{otherwise} \end{cases}$$

where K_2 denotes a graph with a single edge.

Proof. For the first claim, if $H \in \mathbb{N}^{k \times k}$ we define a strictly upper-triangular $\tilde{H} \in \{0, 1\}^{n \times n}$ by $\tilde{H}_{i,j} = 1$ if $i < j$ and either $H_{i,j} > 0$ or $H_{j,i} > 0$, and $\tilde{H}_{i,j} = 0$ otherwise. Then $\text{hom}(H; G_n) = \text{hom}(\tilde{H}, G_n)$ for all simple graphs G_n of all sizes. Therefore, we have

$$\text{hom}(H; \frac{G_n}{\mathbb{1}^\top G_n \mathbb{1}}) = \text{hom}(\tilde{H}; \frac{G_n}{\mathbb{1}^\top G_n \mathbb{1}}) \cdot (2|E(G_n)|)^{\mathbb{1}^\top (\tilde{H} - H) \mathbb{1}}.$$

Since $\mathbb{1}^\top \tilde{H} \mathbb{1} \leq \mathbb{1}^\top H \mathbb{1}$, we conclude that $\text{hom}(H; \frac{G_n}{\mathbb{1}^\top G_n \mathbb{1}})$ converges if $\text{hom}(\tilde{H}; \frac{G_n}{\mathbb{1}^\top G_n \mathbb{1}})$ converges, since either $|E(G_n)| \rightarrow \infty$ and $\text{hom}(H; \frac{G_n}{\mathbb{1}^\top G_n \mathbb{1}}) \rightarrow 0$ or $|E(G_n)|$ is constant for all large n . In turn, defining the simple graph $\tilde{H}_{\text{sym}} = (\tilde{H} + \tilde{H}^\top)/2$, we have

$$\text{hom}\left(\tilde{H}; \frac{G_n}{\mathbb{1}^\top G_n \mathbb{1}}\right) = \frac{\text{hom}(\tilde{H}_{\text{sym}}; G_n)}{(2|E(G_n)|)^{|E(\tilde{H}_{\text{sym}})|}}.$$

Thus, the sequence (G_n) quotient-converges if and only if $\left(\frac{\text{hom}(H; G_n)}{|E(G_n)|^{|E(H)|}}\right)_n$ converges for all simple H . To see that it suffices to consider simple and connected graphs, note that if $H = H_1 \sqcup \dots \sqcup H_\ell$ is a disjoint union of connected graphs, then $\frac{\text{hom}(H; G_n)}{|E(G_n)|^{|E(H)|}} = \prod_{i=1}^\ell \frac{\text{hom}(H_i; G_n)}{|E(G_n)|^{|E(H_i)|}}$. This proves the first claim.

For the second claim, if $H = K_2$ then $\text{hom}(H; G_n)/|E(G_n)| = 1$ by definition. Otherwise, suppose $\frac{|E(G_n)|}{n^2} \geq c > 0$ for all n . Note that $\text{hom}(H; G_n) \leq 2|E(G_n)| \cdot n^{|V(H)|-2}$, because the number of graph

homomorphisms $V(H) \rightarrow V(G_n)$ mapping all edges of H to edges of G_n is upper-bounded by the number of maps $V(H) \rightarrow V(G_n)$ that send one particular edge in H to an edge in G_n . Since H is connected and contains more than one edge, we have $|V(H)| < 2|E(H)|$, in which case

$$\frac{\hom(H; G_n)}{|E(G_n)|^{|E(H)|}} \leq \frac{2n^{|V(H)|-2}}{|E(G_n)|^{|E(H)|-1}} \leq \frac{2}{c^{|E(H)|-1}} \cdot n^{|V(H)|-2|E(H)|} \xrightarrow{n \rightarrow \infty} 0.$$

This proves the quotient-convergence of (G_n) by the first part of this corollary. \square

We have now seen several examples and characterizations of quotient-convergent sequences. However, so far it is not clear to what limits such sequences converge. In the next section, we derive limits for quotient-convergent graphs in the form of random exchangeable measures on $[0, 1]^2$.

3 Limits of Quotient-Convergent Graph Sequences

This section is devoted to characterizing limits of quotient-convergent graph sequences and understanding their properties. In Section 3.1, we prove Theorem 1.5 characterizing limits of quotient-convergent graph sequences via grpheurs. In Section 3.2 we interpret the different components of limiting grpheurs (5) in terms of combinatorial aspects of the sequence of graphs converging to them. In Section 3.3 we present a (random) duality pairing between grpheurs and graphons, and prove the formal version of Theorem 1.6 given in Theorem 3.10. Finally, Section 3.4 contains some tedious but straightforward proofs omitted from Section 3.1 for clarity of exposition.

3.1 Characterizing Limits via Grpheurs

To prove Theorem 1.5, we show that the random quotients of size k of a finite graph G are equal in distribution to the random graph $G_k[M_G]$ of size k obtained in (8) from the grpheur M_G associated in (4) to G .

Proposition 3.1. *For any $n, k \in \mathbb{N}$ and $G \in \Delta^{n \times n}$, we have $\rho(F_{k,n})G \stackrel{d}{=} G_k[M_G]$.*

Proof. Recall from (4) that $M_G = \sum_{i,j=1}^n G_{i,j} \delta_{(T_i, T_j)}$ where T_1, \dots, T_n are iid uniform on $[0, 1]$. With probability one, for each $i \in [n]$ there is a unique $j_i \in [k]$ such that $T_i \in I_{j_i}^{(k)}$. Moreover, the index j_i is uniformly distributed in $[k]$ by uniformity of T_i , and the indices j_1, \dots, j_n are independent of each other since the T_i are independent. Define the random map $F: [n] \rightarrow [k]$ sending $F(i) = j_i$ and observe that $F \stackrel{d}{=} F_{k,n}$ is uniformly random, since each $i \in [n]$ is mapped to an independent and uniformly random element of $[k]$. Further, note that

$$(G_k[M_G])_{i,j} = M_G(I_i^{(k)} \times I_j^{(k)}) = \sum_{u,v=1}^n G_{u,v} \mathbb{1}[T_u \in I_i^{(k)}, T_v \in I_j^{(k)}] = \sum_{\substack{u \in F^{-1}(i) \\ v \in F^{-1}(j)}} G_{u,v} = (\rho(F)G)_{i,j}.$$

Hence, we conclude that $G_k[M_G] = \rho(F)G \stackrel{d}{=} \rho(F_{k,n})G$, as claimed. \square

Consequently, we also have $t_Q(H; G) = t_Q(H; M_G)$, where the latter is given by (9). Proposition 3.1 implies that a sequence (G_n) is quotient-convergent if and only if the sequence of random graphs $(G_k[M_{G_n}])_n$ constructed from the associated grpheurs (M_{G_n}) converge weakly for each k . Since $\Delta^{k \times k}$ is compact, this is equivalent to convergence of $(G_k[M_{G_n}])_n$ in the Wasserstein-1 metric W_1 with respect to the entrywise ℓ_1 norm on $\mathbb{R}^{k \times k}$ for each k (see Section 1.3). Motivated by this observation, we relate our W_\square -metric (7) to this W_1 -metric.

Proposition 3.2. *For any two grpheurs $M, N \in \mathcal{M}$ and any $k \in \mathbb{N}$, we have*

$$W_1(G_k[M], G_k[N]) \leq k^2 W_\square(M, N) \quad \text{and} \quad W_\square(M, N) \leq W_1(G_k[M], G_k[N]) + \frac{4}{k}.$$

The proof is straightforward but somewhat technical, as it is based on approximating any rectangle appearing in our expression (7) for W_{\square} by squares in a uniform grid. We defer the proof to Section 3.4. As an immediate consequence, we conclude that a sequence of graphs (G_n) is quotient-convergent if and only if the associated sequence of grapheurs (M_{G_n}) is Cauchy in the W_{\square} -metric (7). We now observe that the metric space $(\mathcal{M}, W_{\square})$ is compact, because the metric W_{\square} metrizes weak convergence in \mathcal{M} and \mathcal{M} is weakly compact; see Section 1.3 for a review of weak convergence.

Proposition 3.3. *The space \mathcal{M} of grapheurs (6) is weakly compact. In particular, any Cauchy sequence $(M_n) \subseteq \mathcal{M}$ has a limit $M \in \mathcal{M}$. Moreover, the following statements are equivalent for $(M_n) \subseteq \mathcal{M}$ and $M \in \mathcal{M}$.*

1. *We have $\lim_n M_n = M$ weakly.*
2. *We have that $\lim_n W_{\square}(M_n, M) = 0$.*
3. *For each $k \in \mathbb{N}$, we have that $\lim_n G_k[M_n] = G_k[M]$ weakly.*
4. *For each multigraph $H \in \mathbb{N}^{k \times k}$, we have $\lim_n t_Q(H; M_n) = t_Q(H; M)$.*

In particular, we have $\lim_n W_{\square}(M_{G_n}, M) = 0$ for a sequence $(G_n \in \Delta^{n \times n})$ if and only if $\lim_n t_Q(H; G_n) = t_Q(H; M)$ for all multigraphs $H \in \mathbb{N}^{k \times k}$, as can be seen by combining Proposition 3.1 and Proposition 3.3(4). Once again, the proof is straightforward but involved, and we defer it to Section 3.4. Combining the preceding propositions, we are ready to prove Theorem 1.5.

Proof (Theorem 1.5). To prove the first statement, note that the sequence (G_n) is quotient-convergent if and only if $(G_k[M_{G_n}])$ is Cauchy in W_1 for each $k \in \mathbb{N}$ by Definition 1.1. By Proposition 3.2, this is equivalent to (M_{G_n}) being Cauchy in the W_{\square} -metric, as we remark previously. By Proposition 3.3, this is further equivalent to the existence of a limit $M \in \mathcal{M}$ for (M_{G_n}) in this metric, thus proving the first statement.

The second statement is a consequence of the compactness of \mathcal{M} , which follows from Proposition 3.3.

Finally, we prove the third statement by explicitly constructing a convergent sequence (G_n) of graphs for each grapheur $M \in \mathcal{M}$. Suppose M is given by (5) with $E, \sigma, \varsigma, \theta, \vartheta \geq 0$ and iid uniform $(T_i)_{i \in \mathbb{N}}$. Define the sequence of graphs $(G_n \in \Delta^{n \times n})$ with entries

$$(G_n)_{i,j} = \frac{1}{s_n} (E_{i,j} + \frac{1}{n} \sigma_i + \frac{1}{n} \varsigma_j + \frac{1}{n^2} \theta + \frac{1}{n} \vartheta \mathbb{1}_{i=j}), \quad s_n = \sum_{i,j=1}^n (E_{i,j} + (\sigma_i + \varsigma_j) + \theta + \vartheta), \quad (13)$$

where we note that $s_n \nearrow 1$ and we may assume $s_n > 0$ for all n by reordering the rows and columns of E, σ, ς if needed. Each summand in (13) quotient-converges to the corresponding summand in (5) as shown by combining Proposition 3.3 with the examples of Section 2. Moreover, by Example 2.4 the convex combination of these summands in (13) converges to the convex combination of the summands in (5), which is precisely the grapheur M . Thus, the third statement in Theorem 1.5 is proved. \square

In particular, the sequence of limits of random quotients of a convergent graph sequence fully determines its limit.

Corollary 3.4. *Two quotient-convergent sequences $(G_n^{(1)})$ and $(G_n^{(2)})$ have the same limit if and only if $\lim_n \rho(F_{k,n})G_n^{(1)} \stackrel{d}{=} \lim_n \rho(F_{k,n})G_n^{(2)}$ for all $k \in \mathbb{N}$.*

Proof. From Proposition 3.3, the two sequences have the same limit M if and only if $\lim_n \rho(F_{k,n})G_n^{(i)} \stackrel{d}{=} G_k[M]$ for both $i = 1, 2$. \square

We end this section by noting that valid polynomial inequalities in quotient densities precisely correspond to valid inequalities over grapheurs. We hope that grapheurs can be used to study such inequalities in future work, dually to the use of graphons in the study of inequalities in homomorphism densities [14, §16].

Corollary 3.5 (Inequalities in quotient densities). *Suppose H_1, \dots, H_k are multigraphs and $f: \mathbb{R}^k \rightarrow \mathbb{R}$ is a continuous function. We have $f(t_Q(H_1; G), \dots, t_Q(H_k; G)) \geq 0$ for all $G \in \Delta^{n \times n}$ and all $n \in \mathbb{N}$ if and only if $f(t_Q(H_1; M), \dots, t_Q(H_k; M)) \geq 0$ for all grapheurs $M \in \mathcal{M}$.*

Proof. Since $t_Q(H; G) = t_Q(H; \mathbf{M}_G)$, if $f(t_Q(H_1; \mathbf{M}), \dots, t_Q(H_k; \mathbf{M})) \geq 0$ for all grapheurs $\mathbf{M} \in \mathcal{M}$ then setting $\mathbf{M} = \mathbf{M}_G$ for $G \in \Delta^{n \times n}$ yields $f(t_Q(H_1; G), \dots, t_Q(H_k; G)) \geq 0$. Conversely, suppose $f(t_Q(H_1; G), \dots, t_Q(H_k; G)) \geq 0$ for all $G \in \Delta^{n \times n}$ and all $n \in \mathbb{N}$. For any $\mathbf{M} \in \mathcal{M}$, let (G_n) be a sequence of finite graphs that quotient-converges to \mathbf{M} . Then $t_Q(H; G_n) \rightarrow t_Q(H; \mathbf{M})$ for all multigraphs H , hence by continuity of f we have $f(t_Q(H_1; \mathbf{M}), \dots, t_Q(H_k; \mathbf{M})) = \lim_n f(t_Q(H_1; G_n), \dots, t_Q(H_k; G_n)) \geq 0$. \square

3.2 Structure of Grapheurs

Theorem 1.5 shows that limits of quotient-convergent sequences correspond to grapheurs in \mathcal{M} . Each such grapheur is described by a tuple $(E, \sigma, \varsigma, \theta, \vartheta)$ as in (5), and the goal of this section is to interpret each element of the tuple in terms of the combinatorics of the convergent sequence of graphs. Informally, the five parameters describing quotient limits correspond to the following properties.

- The array $E \in \mathbb{R}_{\geq 0}^{\mathbb{N} \times \mathbb{N}}$ describes a discrete collection of hubs and the fraction of edge weight between them. Specifically, each entry corresponds to an edge in a sequence of graphs that carries an asymptotically positive fraction of the total edge weight.
- The arrays $\sigma, \varsigma \in \mathbb{R}_{\geq 0}^{\mathbb{N}}$ specify directed stars. Specifically, each entry corresponds to the fraction of in-degree or out-degree, respectively, of a vertex that comes from all edges with asymptotically vanishing edge weights.
- The parameters θ and ϑ capture the remaining edge weight in the graph, coming from all edges between vertices with asymptotically vanishing degrees and their self-loops, respectively.

These properties can be seen in all the examples of Section 2. The following proposition proves these properties in general.

Proposition 3.6. *Suppose $(G_n \subseteq \Delta^{n \times n})$ quotient-converges to a grapheur \mathbf{M} that is described by the tuple $(E, \sigma, \varsigma, \theta, \vartheta)$ as in (5). View each G_n as an element of $\mathbb{R}^{\mathbb{N} \times \mathbb{N}}$ by zero-padding it with infinitely-many rows and columns. Then there exist permutations $\pi_n \in \mathsf{S}_\infty$ of \mathbb{N} fixing all but finitely-many integers such that $\lim_n \|E - \pi_n G_n\|_\infty = 0$, $\lim_n \|E\mathbf{1} + \sigma - \pi_n G_n \mathbf{1}\|_\infty = 0$, and $\lim_n \|E^\top \mathbf{1} + \varsigma - \pi_n G_n^\top \mathbf{1}\|_\infty = 0$. Furthermore, we have $\vartheta = \lim_n \text{Tr}(G_n) - \text{Tr}(E)$.*

Proposition 3.6 justifies the intuition outlined above: Up to a relabelling of the vertices of G_n , the entries of E are entrywise limits of the edge weights of G_n , the entries of σ, ς are limits of the remaining degrees of vertices in (G_n) after removing edges with positive limiting weights, and the parameters ϑ, θ capture the remaining edge weights in self loops and the rest of the edges, respectively. We prove Proposition 3.6 in Section 4.4 using an edge-sampling perspective on quotient convergence and a dual Szemerédi regularity lemma.

An immediate consequence of Proposition 3.6 is that all dense and degree-bounded sequences of simple graphs have the same limit.

Corollary 3.7. *Suppose $(G_n \in \mathbb{S}^n)$ is a sequence of undirected graphs such that the sequence $(G_n / \mathbf{1}^\top G_n \mathbf{1})$ quotient-converges.*

1. *The limit is described by $(E, \sigma, \varsigma, \theta, \vartheta)$ with $\sigma = \varsigma$ and $E^\top = E$.*
2. *If each G_n has no self-loops, then $\text{diag}(E) = \vartheta = 0$.*
3. *If $G_n \in \mathbb{S}_{\text{sim}}^n$ is a sequence of simple graphs with $|E(G_n)| \rightarrow \infty$, and is either dense or degree-bounded, then the sequence $(\frac{G_n}{2|E(G_n)|})$ quotient converges to the grapheur λ^2 .*

Proof. The first two claims follow directly from Proposition 3.6, since for any permutation $\pi_n \in \mathsf{S}_\infty$ the matrix $\pi_n G_n$ is, respectively, symmetric and has zero diagonal if G_n is symmetric and has no self-loops. For the third claim, since G_n is symmetric, we have $G_n^\top \mathbf{1} = G_n \mathbf{1} = \deg(G_n)$ is the vector of degrees of the vertices in G_n . If (G_n) is dense, then $\frac{\max_i \deg(G_n)_i}{2|E(G_n)|} \leq \frac{c}{n} \xrightarrow{n \rightarrow \infty} 0$ for some constant $c > 0$. If (G_n) is degree-bounded, with all degrees bounded by k , then $\frac{\max_i \deg(G_n)_i}{2|E(G_n)|} \leq \frac{k}{2|E(G_n)|} \xrightarrow{n \rightarrow \infty} 0$. In either case, Proposition 3.6 shows that $\lim_n \frac{G_n}{2|E(G_n)|} = \lambda^2$ since $\text{diag}(G_n) = 0$ for all n . \square

We end this section by giving a sufficient condition for the limit of a quotient-convergent sequence (G_n) to consist entirely of discrete hubs, i.e., such that $\sigma = \varsigma = 0$ and $\theta = \vartheta = 0$ in (5). In particular, this sufficient condition implies that the random quotients $(\rho(F_{k,n})G_n)$ and the grapheurs (M_{G_n}) converge not just weakly but also in total variation. Such total variation convergence was shown for graphs sampled both from graphons [14, Cor. 10.25] and from graphexes [32, §5], and was the primary mode of convergence studied for edge-exchangeable random graph models [23]. However, this condition is too restrictive for our purposes in this paper; for example, it does not apply to the sequence of stars in Example 2.2. In the following result, if μ, ν are two discrete measures on a space S then their total variation distance is $\text{dist}_{\text{TV}}(\mu, \nu) = \frac{1}{2} \sum_{x \in S} |\mu(\{x\}) - \nu(\{x\})|$.

Proposition 3.8. *Suppose $(G_n \in \Delta^{n \times n})$ and $E \in \mathbb{R}_{>0}^{N \times N}$ satisfy $\lim_n \|\pi_n G_n - E\|_1 = 0$ for some permutations $\pi_n \in S_n$. Then (G_n) is quotient-convergent to $M = \sum_{i,j} E_{i,j} \delta_{(T_i, T_j)}$.*

Proof. Draw iid uniform (T_i) and consider the coupling $M'_{G_n} = \sum_{i,j} (\pi_n G_n)_{i,j} \delta_{(T_i, T_j)}$, $M' = \sum_{i,j} E_{i,j} \delta_{(T_i, T_j)}$, using the same locations for both. Then with probability 1 we have $\text{dist}_{\text{TV}}(M', M'_{G_n}) \leq \|\pi_n G_n - E\|_1 / 2 \rightarrow 0$ since the atoms (T_i, T_j) are distinct almost surely. By (8), we then have $W_1(G_k[M], G_k[M_{G_n}]) \leq \mathbb{E} \|G_k[M'] - G_k[M'_{G_n}]\|_1 \leq k^2 \mathbb{E} \text{dist}_{\text{TV}}(M', M'_{G_n}) \rightarrow 0$ for each k , proving quotient convergence. \square

3.3 Duality between Grapheurs and Graphons

In this section, we characterize convergence of grapheurs in terms of graphons and vice-versa, exhibiting a facet of the duality between the two graph limit theories. Since graphon theory is most simply expressed in the undirected case (see [39, §8] for the directed case), we restrict ourselves to this setting in the present section.

We begin by recalling a few basic notions from graphon theory. Let $\widetilde{\mathcal{W}} = \{W: [0, 1]^2 \rightarrow [0, 1] \text{ measurable, symmetric}\}$ be the space of $[0, 1]$ -valued kernels. We define two kernels W and U to be equivalent, denoted $W \sim U$, if there exist (Lebesgue-)measure-preserving maps $\varphi, \psi: [0, 1] \rightarrow [0, 1]$ satisfying $W \circ (\varphi, \varphi) = U \circ (\psi, \psi)$. This is the weak isomorphism between kernels commonly defined in the dense graph limits literature [14, Cor. 10.35]. We denote by $\mathcal{W} = \widetilde{\mathcal{W}} / \sim$ the quotient space under this equivalence relation. Elements of \mathcal{W} are called graphons [14, §7.1]. A graphon $W \in \mathcal{W}$ defines random simple graphs of each size by setting $\mathbb{G}_k[W] = (\mathbb{B}_{T_i, T_j}^{(W)})_{i,j \in [k]}$ where $\mathbb{B}_{T_i, T_j}^{(W)} \sim \text{Ber}(W(T_i, T_j))$ are drawn independently for $i < j$ while $\mathbb{B}_{T_j, T_i}^{(W)} = \mathbb{B}_{T_i, T_j}^{(W)}$ for $i > j$ and $\mathbb{B}_{T_i, T_i}^{(W)} = 0$. Here $T_1, \dots, T_k \stackrel{iid}{\sim} \text{Unif}([0, 1])$ as before. A sequence $(W_n) \subseteq \mathcal{W}$ converges if for each k , the sequence of random graphs $(\mathbb{G}_k[W_n])_k$ converges weakly for each k .⁴ This is equivalent to convergence of homomorphism densities and to convergence in cut metric [14, §10]. Finally, a finite graph $G \in \mathbb{S}_{\text{sim}}^n$ defines a step graphon $W_G(x, y) = G_{i,j}$ if $x \in I_i^{(n)} \times I_j^{(n)}$ where $I_i^{(n)} = [(i-1)/n, i/n)$.

In our context, let $\mathcal{M}_{\text{sym}} = \{M \in \mathcal{M} : E = E^\top, \sigma = \varsigma, \text{diag}(E) = 0, \vartheta = 0, \theta \in \mathbb{R}_+\}$ be the collection of grapheurs corresponding to limits of undirected graphs with unit edge weight and no self-loops. For any $W \in \mathcal{W}$ and $M \in \mathcal{M}_{\text{sym}}$, define the random variable

$$\langle M, W \rangle = \sum_{i,j \in \mathbb{N}} E_{i,j} \mathbb{B}_{T_i, T_j}^{(W)} + 2 \sum_{i \in \mathbb{N}} \sigma_i \int_0^1 W(T_i, y) dy + \theta \int_{[0,1]^2} W(x, y) dx dy. \quad (14)$$

Note that this random variable exists and is bounded by $|\langle M, W \rangle| \leq 1$ because both $(E_{i,j})$ and (σ_i) are summable. Also note that $\langle M, W_G \rangle = \mathbb{E}_M W_G = \langle G, \mathbb{G}_k[M] \rangle$ if $G \in \mathbb{S}_{\text{sim}}^k$, since W_G takes values in $\{0, 1\}$ so $\mathbb{B}_{T_i, T_j}^{(W_G)} = W_G(T_i, T_j)$. Moreover, for finite graphs $G_1 \in \Delta^{n \times n}$, $G_2 \in \mathbb{S}_{\text{sim}}^m$, we have

$$\langle M_{G_1}, W_{G_2} \rangle = \langle \rho(F_{m,n})G_1, G_2 \rangle = \langle G_1, \rho(F_{m,n})^*G_2 \rangle,$$

which can equivalently be viewed as comparing a random quotient of G_1 to G_2 or comparing a random subgraph of G_2 to G_1 (see Section 1). We now prove that the random variable (14) only depends on the equivalence class of a graphon W .

⁴Graphon convergence is usually defined by total variation convergence of these random graphs, which is equivalent to their weak convergence since they are supported on the finite set $\mathbb{S}_{\text{sim}}^k$.

Proposition 3.9. *For any $W, U \in \widetilde{\mathcal{W}}$, if $W \sim U$ then $\langle \mathbf{M}, W \rangle \stackrel{d}{=} \langle \mathbf{M}, U \rangle$ for any $\mathbf{M} \in \mathcal{M}_{\text{sym}}$.*

Proof. Suppose $\mathbf{M} \in \mathcal{M}_{\text{sym}}$ is defined by $(E, \sigma = \varsigma, \theta, \vartheta = 0)$ as in (5). Since $W \sim U$, there are measure-preserving maps $\varphi, \psi: [0, 1] \rightarrow [0, 1]$ satisfying $W \circ (\varphi, \varphi) = U \circ (\psi, \psi)$. We now define a coupling of $\langle \mathbf{M}, W \rangle$ and $\langle \mathbf{M}, U \rangle$ under which the two are equal almost surely. Sample locations (T_i) iid uniformly on $[0, 1]$, and note that both $(\varphi(T_i)), (\psi(T_i)) \stackrel{iid}{\sim} \text{Unif}([0, 1])$ because φ and ψ are measure-preserving. Note that, conditioned on the locations (T_i) above, we have $\mathbf{B}_{\varphi(T_i), \varphi(T_j)}^{(W)} \stackrel{d}{=} \mathbf{B}_{\psi(T_i), \psi(T_j)}^{(U)}$, since they are both Bernoulli random variables with success probability $W(\varphi(T_i), \varphi(T_j)) = U(\psi(T_i), \psi(T_j))$. Letting $\mathbf{B}_{i,j}$ be a sample from the above distribution independently for each $i < j$, setting $\mathbf{B}_{i,j} = \mathbf{B}_{j,i}$ for $i > j$, and $\mathbf{B}_{i,i} = 0$ for all i , we define the coupling of $\langle \mathbf{M}, W \rangle$ and $\langle \mathbf{M}, U \rangle$ by using (14) with these common samples $(\mathbf{B}_{i,j})$ for both, and note that under this coupling they are equal almost surely. Thus, we have $\langle \mathbf{M}, W \rangle \stackrel{d}{=} \langle \mathbf{M}, U \rangle$ as desired. \square

Thus, the pairing (14) defines a pairing from $\mathcal{M}_{\text{sym}} \times \mathcal{W}$ to bounded scalar random variables. Importantly, we can characterize convergence in \mathcal{M}_{sym} in terms of graphons and characterize graphon convergence in \mathcal{W} in terms of grapheurs. The following is the formal version of Theorem 1.6 from Section 1.

Theorem 3.10. *We have $\lim_n \mathbf{M}_n = \mathbf{M}$ in \mathcal{M}_{sym} if and only if $\langle \mathbf{M}_n, W_G \rangle \rightarrow \langle \mathbf{M}, W_G \rangle$ weakly for all simple graphs G . We also have $\lim_n W_n \rightarrow W$ in \mathcal{W} if and only if $\langle \mathbf{M}_G, W_n \rangle \rightarrow \langle \mathbf{M}_G, W \rangle$ weakly for all undirected graphs G with unit edge weight and no self-loops.*

Proof. For the first claim, suppose $\langle \mathbf{M}_n, W_G \rangle \rightarrow \langle \mathbf{M}, W_G \rangle$ for all $G \in \mathbb{S}_{\text{sim}}^k$ and all $k \in \mathbb{N}$. Therefore, $\mathbb{E}_{\mathbf{M}_n} W_G \rightarrow \mathbb{E}_{\mathbf{M}} W_G$ for any such G . Taking linear combinations of such W_G , we get $\mathbb{E}_{\mathbf{M}_n} W \rightarrow \mathbb{E}_{\mathbf{M}} W$ for any $W: [0, 1]^2 \rightarrow \mathbb{R}$ that is piecewise-constant on the $k \times k$ uniform grid and equals zero on the diagonal for some $k \in \mathbb{N}$. For any continuous and symmetric $f: [0, 1]^2 \rightarrow \mathbb{R}$, define $f_k = \sum_{i,j=1}^k f_{i,j}^{(k)} \mathbb{1}_{I_i^{(k)} \times I_j^{(k)}}$ where if $i \neq j$ we set $f_{i,j}^{(k)} = f(\frac{i}{k} - \frac{1}{2k}, \frac{j}{k} - \frac{1}{2k})$ to the value of f at the center of the rectangle $I_i^{(k)} \times I_j^{(k)}$, and if $i = j$ we set $f_{i,i}^{(k)} = 0$. Since f is continuous on a compact set, it is uniformly continuous, so for any $\epsilon > 0$ we have $|f(x) - f_k(x)| \leq \epsilon$ for all $x \in [0, 1]^2 \setminus \bigcup_{i=1}^k (I_i^{(k)} \times I_i^{(k)})$. Therefore, for any grapheur $\mathbf{M}' \in \mathcal{M}_{\text{sym}}$ we have

$$\mathbb{E}_{\mathbf{M}'} |f - f_k| \leq \epsilon + \mathbb{E}_{\mathbf{M}'} \left(\bigcup_{i=1}^k I_i^{(k)} \times I_i^{(k)} \right) = \epsilon + \frac{1}{k},$$

for all large k since $\mathbb{E}_{\mathbf{M}'} = \lambda^2$ for $\mathbf{M}' \in \mathcal{M}_{\text{sym}}$. Thus, we have

$$W_1(\mathbb{E}_{\mathbf{M}_n} f, \mathbb{E}_{\mathbf{M}} f) \leq 2(\epsilon + k^{-1}) + W_1(\mathbb{E}_{\mathbf{M}_n} f_k, \mathbb{E}_{\mathbf{M}} f_k) \xrightarrow{n \rightarrow \infty} 2(\epsilon + k^{-1}),$$

since f_k is piecewise-constant on the $k \times k$ uniform grid and has zero values on its diagonal. We conclude by taking $k \rightarrow \infty$ that $\mathbb{E}_{\mathbf{M}_n} f \rightarrow \mathbb{E}_{\mathbf{M}} f$ weakly for any continuous $f: [0, 1]^2 \rightarrow \mathbb{R}$, and hence that $\mathbf{M}_n \rightarrow \mathbf{M}$ weakly. Conversely, suppose $\mathbf{M}_n \rightarrow \mathbf{M}$ weakly, and let $W = W_G$ for $G \in \mathbb{S}_{\text{sim}}^k$. Then

$$\langle \mathbf{M}_n, W_G \rangle = \mathbb{E}_{\mathbf{M}_n} W_G = \langle G, \mathbb{G}_k[\mathbf{M}_n] \rangle \rightarrow \langle G, \mathbb{G}_k[\mathbf{M}] \rangle = \mathbb{E}_{\mathbf{M}} W_G = \langle \mathbf{M}, W_G \rangle,$$

weakly, since $\mathbb{G}_k[\mathbf{M}_n] \rightarrow \mathbb{G}_k[\mathbf{M}]$ weakly for each $k \in \mathbb{N}$. This proves the first claim.

For the second claim, suppose $\langle \mathbf{M}_G, W_n \rangle \rightarrow \langle \mathbf{M}_G, W \rangle$ weakly for all graphs $G \in \mathbb{S}^k \cap \Delta^{k \times k}$ with $\text{diag}(G) = 0$, i.e., undirected weighted graphs with no self-loops and total edge weight of 1. We then have

$$\langle G, \mathbb{G}_k[W_n] \rangle = \langle \mathbf{M}_G, W_n \rangle \rightarrow \langle \mathbf{M}_G, W \rangle = \langle G, \mathbb{G}_k[W] \rangle,$$

weakly. Therefore, we get $\langle G, \mathbb{G}_k[W_n] \rangle \rightarrow \langle G, \mathbb{G}_k[W] \rangle$ for all $G \in \mathbb{S}^k$ by linearity and the fact that $\mathbb{G}_k[W_n]$ all have zero diagonal. This implies $\mathbb{G}_k[W_n] \rightarrow \mathbb{G}_k[W]$ weakly by Cramér–Wold [40, Cor. 6.5]. Since this holds for all $k \in \mathbb{N}$, we conclude that $W_n \rightarrow W$. Conversely, suppose $W_n \rightarrow W$. Then $\langle \mathbf{M}_G, W_n \rangle = \langle G, \mathbb{G}_k[W_n] \rangle \rightarrow \langle G, \mathbb{G}_k[W] \rangle = \langle \mathbf{M}_G, W \rangle$ for any $G \in \mathbb{S}^k \cap \Delta^{k \times k}$. \square

We conjecture that the above duality pairing is, in fact, bi-continuous.

Conjecture 3.11. *If $(\mathbf{M}_n) \subseteq \mathcal{M}_{\text{sym}}$ and $(W_n) \subseteq \mathcal{W}$ converge to \mathbf{M} and W , respectively, then $\langle \mathbf{M}_n, W_n \rangle \rightarrow \langle \mathbf{M}, W \rangle$ weakly.*

In particular, this conjecture implies that $M_n \rightarrow M$ if and only if $\langle M_n, W \rangle \rightarrow \langle M, W \rangle$ for all graphons $W \in \mathcal{W}$, and $W_n \rightarrow W$ if and only if $\langle M, W_n \rangle \rightarrow \langle M, W \rangle$ for all grapheurs $M \in \mathcal{M}_{\text{sym}}$, a strengthening of Theorem 3.10.

Theorem 3.10 exhibits a concrete, formal duality between quotient convergence and graphon convergence. This duality can be extended to general grapheurs and more general kernels on $[0, 1]^2$, but the extension of graphon convergence to this setting is more complicated [39, §8]. We therefore leave this and other extensions of this duality for future work.

3.4 Missing Proofs from Section 3

In this section, we give all the proofs deferred from Section 3.1 and Section 3.2, namely, Propositions 3.2 and 3.3.

Proof of Proposition 3.2

Many of the proofs in this section are based on standard approximations of rectangles, and continuous functions by step functions on a uniform grid and vice-versa. Such approximations allow us to compare the measures of these objects under our random exchangeable measures using the following lemma.

Lemma 3.12. *Fix any grapheur $M \in \mathcal{M}$.*

1. *There is $\theta \in [0, 1]$ satisfying $\mathbb{E}M = \theta\lambda^2 + (1 - \theta)\lambda_D$.*
2. *For any Borel set $B \subseteq [0, 1]^2$, we have $\mathbb{E}M(B) \leq \max\{\lambda^2(B), \lambda_D(B)\}$. In particular, if $\lambda^2(B) = \lambda_D(B) = 0$, then $M(B) = 0$ almost surely.*

Proof. For the first claim, since M is exchangeable we conclude that $\mu = \mathbb{E}M$ is a (deterministic) exchangeable measure on $[0, 1]^2$, meaning that $\mu \circ (\sigma, \sigma) = \mu$ for all (Lebesgue)-measure preserving bijections $\sigma: [0, 1] \rightarrow [0, 1]$. The space of such measures is two-dimensional and is spanned by λ^2, λ_D by [37, Thm. 9.12], so $\mu = \theta\lambda + \vartheta\lambda_D$ for some $\theta, \vartheta \in \mathbb{R}$. Since M is a probability measure almost surely, its mean μ is a probability measure as well, necessitating $\theta + \vartheta = 1$ and $\theta, \vartheta \geq 0$. This proves the first claim.

The second claim immediately follows from the first, noting in particular that if $\lambda^2(B) = \lambda_D(B) = 0$ then the scalar random variable $M(B)$ is nonnegative with a vanishing mean. \square

To prove Proposition 3.2, we also need to relate couplings of random measures to couplings of random graphs derived from them.

Lemma 3.13. *For any $M_1, M_2 \in \mathcal{M}$ and coupling (G_1, G_2) of $(G_k[M_1], G_k[M_2])$ for some $k \in \mathbb{N}$, there is a coupling (M'_1, M'_2) of (M_1, M_2) such that $(G_k[M'_1], G_k[M'_2]) \stackrel{d}{=} (G_1, G_2)$.*

Proof. We apply the disintegration theorem [41, Thm. A] to the continuous map $\pi_k: \mathcal{P}([0, 1]^2) \rightarrow \Delta^{k \times k}$ between compact metric spaces sending $\mu \mapsto (\mu(I_i^{(k)} \times I_j^{(k)}))_{i,j \in [k]}$. The disintegration theorem yields for each $M \in \mathcal{M}$ a Borel collection of probability measures $(\lambda_G^{(M)})_{G \in \Delta^{k \times k}}$ for each $M \in \mathcal{M}$ such that $\text{supp}(\lambda_G^{(M)}) \subseteq \pi_k^{-1}(G)$ and such that $\text{Law}(M) = \mu_k^{(M)} \otimes \lambda_G^{(M)}$ in the sense that

$$\mathbb{P}[M \in B] = \int_{\Delta^{k \times k}} \int_{\pi_k^{-1}(G)} \mathbb{1}_B(\mu) d\lambda_G^{(M)}(\mu) d\mu_k^{(M)}(s),$$

for any Borel $B \subseteq \mathcal{P}([0, 1]^2)$. The measure $\lambda_G^{(M)}$ is the distribution of M conditioned on $\pi_k(M) = G$.

Let $\gamma = \text{Law}((G_1, G_2))$. We construct the coupling $\Gamma = \text{Law}((M'_1, M'_2))$ explicitly by setting

$$\Gamma(B) = \int_{\Delta^{k \times k} \times \Delta^{k \times k}} \int_{\pi_k^{-1}(G_2)} \int_{\pi_k^{-1}(G_1)} \mathbb{1}_B(\mu, \nu) d\lambda_{G_1}^{(M)}(\mu) d\lambda_{G_2}^{(M)}(\nu) d\gamma(G_1, G_2).$$

for each Borel $B \subseteq \mathcal{P}([0, 1]^2) \times \mathcal{P}([0, 1]^2)$. This distribution corresponds to first sampling $(G_1, G_2) \sim \gamma$ and then sampling M'_1 and M'_2 conditioned on $G_k[M'_1] = G_1$ and $G_k[M'_2] = G_2$. It is routine to check that this is a coupling of (M_1, M_2) and that its pushforward under (π_k, π_k) is the given coupling γ of (G_1, G_2) , which is the claim we sought to prove. \square

We can now prove Proposition 3.2.

Proof (Proposition 3.2). Define the intervals $I_i^{(n)} = [(i-1)/n, i/n)$ for $i \in [n]$ and let (M', N') be a coupling of (M, N) attaining $W_{\square}(M, N)$ in (7). Such a coupling exists since the infimum in (7) consists of minimizing a lower-semicontinuous function of the coupling over the compact space of all possible couplings. For each $k \in \mathbb{N}$, we have

$$\begin{aligned} W_1(G_k[M], G_k[N]) &\leq \sum_{i,j=1}^k \mathbb{E} |M'(I_i^{(k)} \times I_j^{(k)}) - N'(I_i^{(k)} \times I_j^{(k)})| \\ &\leq k^2 W_{\square}(M, N), \end{aligned}$$

giving the first claimed inequality.

For the second inequality, consider arbitrary intervals $S = [a_1, a_2]$ and $T = [a_3, a_4]$ for $a, b, c, d \in [0, 1]$. For each $k \in \mathbb{N}$ and $j \in [4]$, let $i_j \in [k]$ be the unique index such that $a_j \in I_{i_j}^{(k)}$ (if $a_j = 1$ set $i_j = k$). Note that $S \subseteq \bigcup_{i=i_1}^{i_2} I_i^{(k)}$ and $|\bigcup_{i=i_1}^{i_2} I_i^{(k)} \setminus S| \leq 2/k$, and similarly for T . Therefore, we have

$$\lambda^2 \left(\bigcup_{i=i_1}^{i_2} \bigcup_{j=i_3}^{i_4} I_i^{(k)} \times I_j^{(k)} \setminus S \times T \right) \leq \frac{4}{k}, \quad \lambda_D \left(\bigcup_{i=i_1}^{i_2} \bigcup_{j=i_3}^{i_4} I_i^{(k)} \times I_j^{(k)} \setminus S \times T \right) \leq \frac{2\sqrt{2}}{k},$$

hence by Lemma 3.12 we have

$$\mathbb{E} |M'(S \times T) - N'(S \times T)| \leq \mathbb{E} \|G_k[M'] - G_k[N']\|_1 + \frac{4}{k}.$$

Taking the supremum over S, T , we get

$$W_{\square}(M', N') \leq \frac{4}{k} + \inf_{\substack{\text{couplings} \\ (M', N')}} \mathbb{E} \|G_k[M'] - G_k[N']\|_1 = \frac{4}{k} + W_1(G_k[M], G_k[N]),$$

where the last equality follows from Lemma 3.13. This is the second claimed inequality. \square

Proof of Proposition 3.3

Next, we turn to proving Proposition 3.3, which states that \mathcal{M} is weakly compact and that the metric W_{\square} metrizes this weak topology. To prove that \mathcal{M} is weakly compact, we rely on the following lemma and the characterization of \mathcal{M} from [37, §9] as the space of ergodic random exchangeable measures.

Lemma 3.14. *If (μ_n) is a weakly convergent sequence of probability distributions on a Polish space S with limit μ , and if $B \subseteq S$ is a measurable subset satisfying $\mu_n(B) = 1$ for all n , then $\mu(B) = 1$.*

Proof. By Urysohn's lemma, for any $\epsilon > 0$ we can find a subset $B_\epsilon \supseteq B$ with $\mu(B_\epsilon \setminus B) \leq \epsilon$ and a continuous function $f: S \rightarrow [0, 1]$ satisfying $f|_B = 1$ and $f|_{S \setminus B_\epsilon} = 0$. Since f is continuous, we have $\mathbb{E}_\mu f = \lim_n \mathbb{E}_{\mu_n} f \geq \lim_n \mu_n(S) = 1$. On the other hand, since $f \leq \mathbb{1}_{B_\epsilon \setminus B} + \mathbb{1}_B$ pointwise, we have $\mathbb{E}_\mu f \leq \mu(B) + \epsilon$. We conclude that $\mu(B) \geq 1$, and since μ is a probability measure, that $\mu(B) = 1$, as desired. \square

We are now to prove Proposition 3.3.

Proof (Proposition 3.3). We break down the proof into several steps.

Compactness of \mathcal{M} : The space $\mathcal{P}([0, 1]^2)$ of probability measures on $[0, 1]^2$ is compact in the weak topology. Therefore, the space $\mathcal{P}(\mathcal{P}([0, 1]^2))$ of random such measures is again compact in its own weak topology (see Section 1.3). Thus, it suffices to prove that $\mathcal{M} \subseteq \mathcal{P}(\mathcal{P}([0, 1]^2))$ is closed in this weak topology.

By [37, Prop. 9.1], the space of random exchangeable measures is precisely the collection of random measures on $[0, 1]^2$ invariant under the countable group \tilde{S}_∞ consisting of permutations of the intervals $(I_i^{(n)})_{i \in [n]}$ for all $n \in \mathbb{N}$. By [37, Lemma A1.2], we can characterize its extreme points by

$$\mathcal{M} = \left\{ M \text{ random exchangeable} : \mathbb{P}[M \in B] \in \{0, 1\} \text{ whenever } B \subseteq \mathcal{P}([0, 1]^2) \text{ is } \tilde{S}_\infty\text{-invariant} \right\}.$$

If $\mathbf{M}_n \rightarrow \mathbf{M}$ and $\text{Law}(\mathbf{M}_n)(B) \in \{0, 1\}$ for all n , then passing to a subsequence we may assume $\text{Law}(\mathbf{M}_n)(B) = 1$ for all n or $\text{Law}(\mathbf{M}_n)(B) = 0$ for all n . Applying Lemma 3.14 to either B or its complement, respectively, yields $\text{Law}(\mathbf{M})(B) \in \{0, 1\}$. We thus conclude that \mathcal{M} is weakly closed, and hence compact.

Equivalence of statements 2 and 3: The equivalence of statements 2 and 3 follows from Proposition 3.2, since $\lim_n G_k[\mathbf{M}_n] = G_k[\mathbf{M}]$ weakly for a given $k \in \mathbb{N}$ if and only if $\lim_n W_1(G_k[\mathbf{M}_n], G_k[\mathbf{M}]) = 0$.

Equivalence of statements 3 and 4: The equivalence of these two statements follows by the proof of Proposition 2.5. Specifically, the sequence $(G_k[\mathbf{M}_n])$ converges weakly to $G_k[\mathbf{M}]$ if and only if the moments of $(G_k[\mathbf{M}_n])$ converge to those of $G_k[\mathbf{M}]$, and these moments are precisely the quotient densities of grapheurs defined in (9).

Equivalence of statements 1 and 2: Endow \mathbb{R}^2 with the ℓ_∞ norm, endow $\mathcal{P}([0, 1]^2)$ with the Wasserstein-1 metric with respect to this norm, and endow \mathcal{M} with the Wasserstein-1 metric with respect to the Wasserstein-1 metric on $\mathcal{P}([0, 1]^2)$, which metrizes weak convergence of random measures [36, Thm. 6.9]. Explicitly, this metric is given by

$$\text{dist}_{W_1}(\mathbf{M}_1, \mathbf{M}_2) = \min_{\substack{\text{random } (\mathbf{M}'_1, \mathbf{M}'_2) \\ \mathbf{M}'_i \stackrel{d}{=} \mathbf{M}_i \text{ for } i=1,2}} \sup_{\substack{f: [0,1]^2 \rightarrow \mathbb{R} \\ f(0)=0 \\ \text{1-Lipschitz in } \ell_\infty}} |\mathbb{E}_{\mathbf{M}'_1} f - \mathbb{E}_{\mathbf{M}'_2} f|,$$

where the min over couplings is attained [36, Thm. 4.1]. It therefore suffices to prove that $\lim_n W_\square(\mathbf{M}_n, \mathbf{M}) = 0$ if and only if $\lim_n \text{dist}_{W_1}(\mathbf{M}_n, \mathbf{M}) = 0$. We proceed to prove this by standard approximations of continuous functions by step functions and vice-versa.

Suppose that $\lim_n \text{dist}_{W_1}(\mathbf{M}_n, \mathbf{M}) = 0$ and fix any $\epsilon > 0$. Let $(\mathbf{M}'_n, \mathbf{M}')$ be a coupling of $(\mathbf{M}_n, \mathbf{M})$ attaining dist_{W_1} . For any interval $S \subseteq [0, 1]$, define $S_\epsilon = (S + [-\epsilon, \epsilon]) \cap [0, 1]$. Note that

$$\lambda^2(S_\epsilon \times T_\epsilon \setminus S \times T) \leq 4\epsilon, \quad \lambda_D(S_\epsilon \times T_\epsilon \setminus S \times T) \leq 2\sqrt{2}\epsilon.$$

We can find an ϵ^{-1} -Lipschitz $f_\epsilon: [0, 1]^2 \rightarrow [0, 1]$ satisfying $f|_{S \times T} = 1$ and $f|_{[0,1]^2 \setminus S_\epsilon \times T_\epsilon} = 0$. For example, set $f_\epsilon(x) = \frac{\text{dist}(x, S_\epsilon \times T_\epsilon)}{\text{dist}(x, S \times T) + \text{dist}(x, S_\epsilon \times T_\epsilon)}$ where dist is with respect to the ℓ_∞ norm. For any $\mathbf{M} \in \mathcal{M}$, we have

$$\mathbb{E}|\mathbf{M}(S \times T) - \mathbb{E}_{\mathbf{M}} f_\epsilon| \leq \mathbb{E}\mathbf{M}(S_\epsilon \times T_\epsilon \setminus S \times T) \leq 4\epsilon,$$

by Lemma 3.12, hence

$$|\mathbb{E}\mathbf{M}'_n(S \times T) - \mathbb{E}\mathbf{M}'(S \times T)| \leq 8\epsilon + |\mathbb{E}_{\mathbf{M}'_n} f_\epsilon - \mathbb{E}_{\mathbf{M}'} f_\epsilon| \leq 8\epsilon + \frac{1}{\epsilon} \text{dist}_{W_1}(\mathbf{M}_n, \mathbf{M}).$$

We conclude that $W_\square(\mathbf{M}_n, \mathbf{M}) \leq 8\epsilon + \epsilon^{-1} \text{dist}_{W_1}(\mathbf{M}_n, \mathbf{M})$, so taking $n \rightarrow \infty$ and then $\epsilon \rightarrow 0$ we conclude that $\lim_n W_\square(\mathbf{M}_n, \mathbf{M}) = 0$.

Conversely, suppose $\lim_n W_\square(\mathbf{M}_n, \mathbf{M}) = 0$. Once again, we note that the infimum over couplings in (7) is attained, since it consists of minimizing a lower-semicontinuous function of the coupling over the compact space of all possible couplings. For each n , let $(\mathbf{M}'_n, \mathbf{M}_n)$ be a coupling attaining $W_\square(\mathbf{M}_n, \mathbf{M})$. For each $N \in \mathbb{N}$, define the intervals $I_i^{(N)} = [(i-1)/N, i/N]$ for $i \in \{1, \dots, N-1\}$ and $I_N^{(N)} = [1-1/N, 1]$. For any 1-Lipschitz $f: [0, 1] \rightarrow \mathbb{R}$ with $f(0) = 0$, let

$$f_N(x) = \sum_{i,j=1}^N f_{i,j}^{(N)} \mathbb{1}_{I_i^{(N)} \times I_j^{(N)}}(x),$$

where $f_{i,j}^{(N)} = f(\frac{i}{N} - \frac{1}{2N}, \frac{j}{N} - \frac{1}{2N})$ is the value of f on the center of the squares $I_i^{(N)} \times I_j^{(N)}$. Since f is 1-Lipschitz in ℓ_∞ and the diameter of these squares is $1/2N$ in ℓ_∞ , we have $|f(x) - f_N(x)| \leq \frac{1}{2N}$ for all $x \in [0, 1]^2$. Furthermore, since $f(0) = 0$ we have $|f_{i,j}^{(N)}| \leq 1$ for all i, j . Therefore, for any $\mathbf{M} \in \mathcal{M}$ we have

$$|\mathbb{E}_{\mathbf{M}} f - \mathbb{E}_{\mathbf{M}} f_N| \leq \frac{1}{2N},$$

and for our particular coupling $(\mathbf{M}', \mathbf{M}'_n)$, we have

$$|\mathbb{E}_{\mathbf{M}'} f - \mathbb{E}_{\mathbf{M}'_n} f| \leq \frac{1}{N} + |\mathbb{E}_{\mathbf{M}'} f_N - \mathbb{E}_{\mathbf{M}'_n} f_N| \leq \frac{1}{N} + \sum_{i,j=1}^N |\mathbf{M}'(I_i^{(N)} \times I_j^{(N)}) - \mathbf{M}'_n(I_i^{(N)} \times I_j^{(N)})|.$$

Since this inequality holds for any such function f , we conclude that

$$\text{dist}_{W_1}(\mathbf{M}_n, \mathbf{M}) \leq \frac{1}{N} + \sum_{i,j=1}^N \mathbb{E}|\mathbf{M}'(I_i^{(N)} \times I_j^{(N)}) - \mathbf{M}'_n(I_i^{(N)} \times I_j^{(N)})| \leq \frac{1}{N} + N^2 W_{\square}(\mathbf{M}_n, \mathbf{M}).$$

Letting $n \rightarrow \infty$ and then $N \rightarrow \infty$, we conclude that $\lim_n \text{dist}_{W_1}(\mathbf{M}_n, \mathbf{M}) = 0$. \square

4 Edge Sampling and Property Testing

In this section, we consider the problem of testing properties of large graphs by randomly sampling small summaries using our framework. For properties specified by graph parameters that are continuous with respect to quotient convergence, it would seem that random quotients are natural candidates for obtaining small summaries of large graphs. However, random quotients require access to the entire graph and are therefore impractical for large graphs. Moreover, we will see in Section 5 that random quotients do not concentrate very well, further obstructing property testing based on these summaries.

To remedy this situation, we present in this section an edge-based sampling procedure that yields a practical method for obtaining small summaries approximating large graphs of any size in the W_{\square} -metric with universal rates. This sampling procedure is based on an equivalent view of quotient convergence in terms of convergence of edge-exchangeable random graph models, the equivalence being furnished by the classical de Finetti's theorem. We describe this perspective on quotient convergence in Section 4.1 and an edge-based analog of the Szemerédi regularity lemma in Section 4.2. We leverage these results to present a practical approach for property testing, which we formally define and study in Section 4.3. We conclude in Section 4.4 by presenting proofs from Section 4.2 that are omitted from notational clarity as well as the proof of Proposition 3.6 from Section 3.2.

4.1 Edge Sampling View of Quotient Convergence

We begin by describing a procedure to obtain finite random graphs from a grapheur by sampling edges with replacement. We then reinterpret quotient convergence in terms of convergence of the resulting random edge-exchangeable graphs.

Given a grapheur \mathbf{M} , we can obtain a sequence of weighted random graphs $(\mathbf{G}^{(n)}[\mathbf{M}])_n$ where $\mathbf{G}^{(n)}[\mathbf{M}]$ has n edges as follows. We sample a measure $\mu \sim \mathbf{M}$ on $[0, 1]^2$, followed by sampling n edges $(x_1, y_1), \dots, (x_n, y_n) \stackrel{iid}{\sim} \mu$. Next we construct the weighted graph $\mathbf{G}^{(n)}[\mathbf{M}]$ on $N = |\{x_i\} \cup \{y_i\}|$ vertices by arbitrarily enumerating $\{t_1, \dots, t_N\} = \{x_i\} \cup \{y_i\}$ and setting the edge weights to:

$$(\mathbf{G}^{(n)}[\mathbf{M}])_{i,j} = |\{k \in [n] : (x_k, y_k) = (t_i, t_j)\}|/n$$

for $1 \leq i, j \leq N$. Note that $|V(\mathbf{G}^{(n)}[\mathbf{M}])| \leq 2n$.

If the grapheur \mathbf{M} is the limit of undirected graphs, we can further replace $\mathbf{G}^{(n)}[\mathbf{M}]$ by its symmetrization $\frac{\mathbf{G}^{(n)}[\mathbf{M}] + (\mathbf{G}^{(n)}[\mathbf{M}])^\top}{2}$ to obtain an undirected random graph; see Remark 4.6 below. When $\mathbf{M} = \mathbf{M}_G$ is the grapheur corresponding to a finite graph G as in (4), the above sampling procedure amounts to sampling with replacement the edges of G proportionally to their weight, and randomly labeling the resulting graph.

For a general grapheur \mathbf{M} described by parameters $(E, \sigma, \varsigma, \theta, \vartheta)$, edge sampling from \mathbf{M} as above corresponds to first sampling random locations $(T_i)_i$ and then sampling edges as follows.

- With probability $E_{i,j}$, sample the edge (T_i, T_j) .
- With probability σ_i , we sample a new vertex $T' \sim \text{Unif}([0, 1])$ and sample the edge (T_i, T') . With probability ς_i , we analogously sample (T', T_i) .

- With probability θ we sample two new vertices $T'_1, T'_2 \sim \text{Unif}([0, 1])$ and add the edge (T'_1, T'_2) . Similarly, with probability ϑ we add the edge (T'', T'') for a new location $T'' \sim \text{Unif}([0, 1])$.

After repeating the above procedure n times, we get a random exchangeable multigraph containing n edges (possibly with repetition), and normalize the resulting graph to have total edge weight one.

We now observe that convergence of edge-sampled graphs is equivalent to quotient convergence.

Proposition 4.1. *We have $\lim_n \mathbf{M}_n = \mathbf{M}$ in \mathcal{M} if and only if $(\mathbf{G}^{(k)}[\mathbf{M}_n])_n \subseteq \Delta^{2k \times 2k}$ converges weakly to $\mathbf{G}^{(k)}[\mathbf{M}]$ for all $k \in \mathbb{N}$.*

Proof. The grapheurs (\mathbf{M}_n) converge weakly if and only if the random exchangeable arrays

$$\begin{bmatrix} x_1^{(n)} & \cdots & x_k^{(n)} \\ y_1^{(n)} & \cdots & y_k^{(n)} \end{bmatrix} \in [0, 1]^{2 \times k}, \quad (15)$$

obtained by first sampling $\mu_n \sim \mathbf{M}_n$ and then sampling $(x_i, y_i) \stackrel{iid}{\sim} \mu_n$ converge weakly by [42, Thm. 1.1], since the distributions of (15) are precisely the finite-dimensional distributions of \mathbf{M}_n . The exchangeable array (15) is precisely the list of edges of the multigraph $k\mathbf{G}^{(k)}[\mathbf{M}_n]$ listed in arbitrary order after assigning its vertices iid labels in $\text{Unif}([0, 1])$. Formally, we have $\mathbf{M}_{\mathbf{G}^{(k)}[\mathbf{M}_n]} \stackrel{d}{=} \frac{1}{k} \sum_{i=1}^k \delta_{(x_i^{(n)}, y_i^{(n)})}$ for all $n, k \in \mathbb{N}$. Thus, the sequence (15) converges weakly for each $k \in \mathbb{N}$ precisely when the sequence of random graphs $(\mathbf{G}^{(k)}[\mathbf{M}_n])_n$ converges weakly. \square

In particular, a sequence of graphs $(G_n \in \Delta^{n \times n})$ quotient-converges precisely when the sequence of random graphs $(\mathbf{G}^{(k)}[\mathbf{M}_{G_n}])$ —obtained by independently sampling k edges from each G_n proportionally to their edge weights and randomly labeling the vertices—converges weakly for each $k \in \mathbb{N}$.

Remark 4.2 (Edge-exchangeable random graphs). *The random graphs we obtain by sampling edges from a grapheur appeared previously in the literature as edge-exchangeable random multigraph models [26]. As explained in [26, Rmk. 4.4], every edge-exchangeable random multigraph model is obtained from a random exchangeable measure on $[0, 1]^2$ as above after labelling the vertices by iid labels from $\text{Unif}([0, 1])$. Nevertheless, to our knowledge graph limits based on these edge-exchangeable random graphs as in Proposition 4.1 have not been previously studied in the literature, see also Section 1.2.*

As noted in [26], if we repeat the above edge sampling process infinitely-many times, each edge (T_i, T_j) corresponding to a positive edge weight $E_{i,j} > 0$ will be repeated infinitely-many times. In contrast, all the other edges sampled above will only appear once. In particular, the vertices T', T'_1, T'_2 sampled in this process will only appear in one edge. Such vertices are called “blips” in [26, 32], and the isolated edges and self-loops are called “dust”.

4.2 Edge-Based Szemerédi Regularity

We now provide bounds on the rate of convergence for $W_{\square}(\mathbf{M}_{\mathbf{G}^{(n)}[\mathbf{M}]}, \mathbf{M})$ to zero as n grows large, both in expectation and with high probability. In bounding the quantity $W_{\square}(\mathbf{M}_{\mathbf{G}^{(n)}[\mathbf{M}]}, \mathbf{M})$, it is important to distinguish between the randomness in the grapheur $\mathbf{M}_{\mathbf{G}^{(n)}[\mathbf{M}]}$ coming from the random atoms T_i and randomness coming from $\mathbf{G}^{(n)}[\mathbf{M}]$. In our next result, we fix a realization of $\mathbf{G}^{(n)}[\mathbf{M}]$, compute the distance $W_{\square}(\mathbf{M}_{\mathbf{G}^{(n)}[\mathbf{M}]}, \mathbf{M})$ depending on this realization (and hence random), and seek to bound both its expectation and its tails with respect to the randomness in $\mathbf{G}^{(n)}[\mathbf{M}]$.

In fact, our results provide convergence rates for the larger metric $\overline{W}_{\square}(\mathbf{M}, \mathbf{N})$ defined by

$$W_{\square}(\mathbf{M}, \mathbf{N}) \leq \overline{W}_{\square}(\mathbf{M}, \mathbf{N}) = \inf_{\substack{\text{random } (\mathbf{M}'_1, \mathbf{M}'_2) \\ \mathbf{M}'_i \stackrel{d}{=} \mathbf{M}_i \text{ for } i=1,2}} \mathbb{E} \left[\sup_{\substack{S, T \subseteq [0, 1] \\ \text{intervals}}} |\mathbf{M}'_1(S \times T) - \mathbf{M}'_2(S \times T)| \right],$$

where the expectation is taken with respect to the coupling $(\mathbf{M}'_1, \mathbf{M}'_2)$ of \mathbf{M}_1 and \mathbf{M}_2 . Note that \overline{W}_{\square} differs from W_{\square} in (7) by moving the supremum over axis-aligned rectangles into the expectation.

Theorem 4.3. For any graph $M \in \mathcal{M}$ and each $n \in \mathbb{N}$, let $G^{(n)}[M]$ be a random graph obtained by sampling n edges from M . Then

$$\mathbb{E}_{G^{(n)}[M]} [W_{\square}(M, M_{G^{(n)}[M]})] \leq \mathbb{E}_{G^{(n)}[M]} [\overline{W}_{\square}(M, M_{G^{(n)}[M]})] \leq \frac{174}{\sqrt{n}}.$$

Moreover, $W_{\square}(M, M_{G^{(n)}[M]}) \leq \overline{W}_{\square}(M, M_{G^{(n)}[M]}) \leq \frac{174+\epsilon}{\sqrt{n}}$ with probability at least $1 - \exp(-2\epsilon^2)$ over the random graph $G^{(n)}[M]$.

The proof of Theorem 4.3 relies on two key lemmas. The first involves the construction of a particular coupling of M , $M_{G^{(n)}[M]}$, and $G^{(n)}[M]$.

Lemma 4.4. There is a coupling $(M', M'_{G^{(n)}}, G^{(n)})$ of M , $M_{G^{(n)}[M]}$, and $G^{(n)}[M]$ satisfying the following two properties. First, in this coupling $M'_{G^{(n)}}$ is obtained by sampling $\mu \sim M'$, then sampling n iid points from μ , and finally forming the associated empirical measure. Second, conditioned on $G^{(n)} = G$, the pair $(M', M'_{G^{(n)}})$ is a coupling of M and M_G .

The second step in the proof of Theorem 4.3 involves using Vapnik-Chervonenkis theory to uniformly bound the discrepancy between a measure and its empirical sample on axis-aligned rectangles.

Lemma 4.5. For any measure $\mu \in \mathcal{P}([0, 1]^2)$, let $\mu_n = \frac{1}{n} \sum_{i=1}^n \delta_{X_i}$ be the random empirical measure obtained by sampling $X_1, \dots, X_n \stackrel{iid}{\sim} \mu$. Then

$$\mathbb{E}_{\mu_n} \sup_{\substack{S, T \subseteq [0, 1] \\ \text{intervals}}} |\mu(S \times T) - \mu_n(S \times T)| \leq \frac{174}{\sqrt{n}}.$$

The proofs of these two lemmas are straightforward but technical, and we defer them to Section 4.4 for clarity of exposition. Using the lemmas, we can now prove Theorem 4.3.

Proof (Theorem 4.3). Let $\mathcal{AAR} = \{[a, b] \times [c, d] : a, b, c, d \in [0, 1]\}$ be the class of axis-aligned rectangles contained in $[0, 1]^2$, and let $(M', M'_{G^{(n)}}, G^{(n)})$ be the coupling from Lemma 4.4. Define

$$f(G^{(n)}[M]) = f((x_1, y_1), \dots, (x_n, y_n)) = \overline{W}_{\square}(M_{G^{(n)}[M]}, M),$$

which is a function of the edges $(x_1, y_1), \dots, (x_n, y_n)$ sampled iid from $\mu \sim M$ to create $G^{(n)}[M]$. First note that

$$\mathbb{E}_{G^{(n)}[M]} f(G^{(n)}[M]) = \mathbb{E}_{G^{(n)}} f(G^{(n)}) \leq \mathbb{E}_{(M', M'_{G^{(n)}}, G^{(n)})} \sup_{R \in \mathcal{AAR}} |M'(R) - M'_{G^{(n)}}(R)| \leq \frac{174}{\sqrt{n}},$$

where the equality follows from $G^{(n)}[M] \stackrel{d}{=} G^{(n)}$, the first inequality follows by considering the coupling $(M', M'_{G^{(n)}})|G^{(n)}$ of M and $M_{G^{(n)}}$, and the last inequality follows by Lemma 4.5 and the fact that $M'_{G^{(n)}}$ is an empirical measure obtained from n iid samples from M' .

Second, note that f is $1/n$ -Lipschitz in each (x_i, y_i) . Indeed, if $G^{(n,i)}[M]$ differs from $G^{(n)}[M]$ only in the i th location (x_i, y_i) , and we glue the two couplings $(M'', M''_{G^{(n)}[M]})$ and $(M'', M''_{G^{(n,i)}[M]})$ attaining the corresponding \overline{W}_{\square} distances as in [36, §1], then

$$|f(G^{(n,i)}[M]) - f(G^{(n)}[M])| \leq \mathbb{E}_{(M''_{G^{(n)}[M]}, M''_{G^{(n,i)}[M]})} \sup_{R \in \mathcal{AAR}} |M''_{G^{(n)}[M]}(R) - M''_{G^{(n,i)}[M]}(R)| \leq \frac{1}{n},$$

since the only difference between $M''_{G^{(n,i)}[M]}$ and $M''_{G^{(n)}[M]}$ is a weight of $1/n$ placed at one of their atoms. Since $(x_1, y_1), \dots, (x_n, y_n)$ are sampled iid from a fixed measure μ (after it was sampled once from M), the bounded-difference concentration inequality [43, Cor. 2.21] implies that $f(G^{(n)}[M]) \leq \mathbb{E} f(G^{(n)}[M]) + \epsilon/\sqrt{n}$ with probability at least $1 - \exp(-2\epsilon^2)$, proving the theorem. \square

Using Theorem 4.3, we can now prove Theorem 1.7 from Section 1.

Proof (Theorem 1.7). The first claim follows from Theorem 4.3. The second claim follows from the fact that $\mathbb{E} W_{\square}(\mathbf{M}, \mathbf{M}_{G^{(n)}[\mathbf{M}]}) \leq \frac{174}{\sqrt{n}}$, which implies that some realization $G^{(n)}$ of $G^{(n)}[\mathbf{M}]$, satisfies $W_{\square}(\mathbf{M}, \mathbf{M}_{G^{(n)}}) \leq \frac{174}{\sqrt{n}}$. Setting $n = k(\epsilon) = \lceil (174/\epsilon)^2 \rceil$, we obtain the second claim. \square

We conclude this section by remarking that the above discussion applies to undirected random graphs sampled from symmetric graphheurs.

Remark 4.6 (Undirected graphs). *If \mathbf{M} is the limit of undirected graphs, then by Corollary 3.7 it is symmetric, i.e., we have $\mathbf{M}(S \times T) = \mathbf{M}(T \times S)$ for any measurable $S, T \subseteq [0, 1]$. In that case, we can sample undirected graphs from \mathbf{M} by first sampling $G^{(n)}[\mathbf{M}]$ as above and symmetrizing it to get $G_{\text{sym}}^{(n)}[\mathbf{M}] = \frac{G^{(n)}[\mathbf{M}] + (G^{(n)}[\mathbf{M}])^\top}{2}$. For any $S, T \subseteq [0, 1]$, we have*

$$\begin{aligned} |\mathbf{M}(S \times T) - \mathbf{M}_{G_{\text{sym}}^{(n)}[\mathbf{M}]}(S \times T)| &= \frac{1}{2} |\mathbf{M}(S \times T) + \mathbf{M}(T \times S) - \mathbf{M}_{G^{(n)}[\mathbf{M}]}(S \times T) - \mathbf{M}_{G^{(n)}[\mathbf{M}]}(T \times S)| \\ &\leq \max\{|\mathbf{M}(S \times T) - \mathbf{M}_{G^{(n)}[\mathbf{M}]}(S \times T)|, |\mathbf{M}(T \times S) - \mathbf{M}_{G^{(n)}[\mathbf{M}]}(T \times S)|\}, \end{aligned}$$

hence $W_{\square}(\mathbf{M}, \mathbf{M}_{G_{\text{sym}}^{(n)}[\mathbf{M}]}) \leq W_{\square}(\mathbf{M}, \mathbf{M}_{G^{(n)}[\mathbf{M}]})$ and similarly for \overline{W}_{\square} . Thus, all the results of this section apply to the undirected random graph $G_{\text{sym}}^{(n)}[\mathbf{M}]$.

4.3 Property Testing

We apply the preceding edge-based sampling procedure to test properties of large graphs based on small summaries. We begin by formally defining the class of graph parameters we can test using our framework. Here a *graph parameter* is a function $f: \bigsqcup_{n \in \mathbb{N}} \Delta^{n \times n} \rightarrow \mathbb{R}$ on graphs of all sizes that is permutation-invariant, in the sense that $f(\pi_n G_n) = f(G_n)$ for any permutation $\pi_n \in \mathsf{S}_n$ of the vertex labels.

Definition 4.7 (Quotient-testable graph parameters). *A quotient-testable graph parameter is a graph parameter $f: \bigsqcup_{n \in \mathbb{N}} \Delta^{n \times n} \rightarrow \mathbb{R}$ satisfying the following properties:*

(Isolate-indifference) $f(G_N) = f(G_n)$ whenever $G_N \in \Delta^{N \times N}$ differs from $G_n \in \Delta^{n \times n}$ by isolated vertices;

(Continuity) *The sequence $(f(G_n))$ converges whenever $(G_n \in \Delta^{n \times n})$ is quotient-convergent.*

We proceed to give several equivalent characterizations of quotient-testable parameters.

Proposition 4.8 (Testability as continuity). *The following are equivalent for a graph parameter f .*

1. *The parameter f is quotient-testable.*
2. *There is a function $\bar{f}: \mathcal{M} \rightarrow \mathbb{R}$ satisfying $\bar{f}(\mathbf{M}_G) = f(G)$ for all $G \in \bigsqcup_n \Delta^{n \times n}$. Such \bar{f} is then unique and continuous.*
3. *For any $\epsilon > 0$ there is a $\delta = \delta(\epsilon) > 0$ such that if $W_{\square}(\mathbf{M}_{G_n}, \mathbf{M}_{G_m}) \leq \delta$ then $|f(G_n) - f(G_m)| \leq \epsilon$ for any two graphs G_n, G_m of any two sizes.*

Proof. We prove these statements imply each other in order, with the last statement implying the first.

Suppose f is quotient-testable. Define $\mathbf{M}_{<\infty} = \{\mathbf{M}_G : G \in \Delta^{n \times n} \text{ for some } n\} \rightarrow \mathbb{R}$ to be the graphheurs in \mathbf{M} arising from finite graphs and define $\bar{f}: \mathbf{M}_{<\infty} \rightarrow \mathbb{R}$ by $\bar{f}(\mathbf{M}_G) = f(G)$. This is well-defined because f is permutation-invariant and isolate-indifferent, as $\mathbf{M}_G = \mathbf{M}_{G'}$ if and only if G and G' differ by isolated vertices and node relabeling [17, Prop. 3]. Further, \bar{f} is continuous by continuity of f , hence it extends uniquely to a continuous function on the closure \mathcal{M} of $\mathbf{M}_{<\infty}$. This proves statement 2.

Suppose f extends to a continuous $\bar{f}: \mathcal{M} \rightarrow \mathbb{R}$. Since \mathcal{M} is compact and this compact topology is metrized by the W_{\square} metric by Proposition 3.3, the function \bar{f} is uniformly continuous in the W_{\square} metric. This proves statement 3.

Finally, assume statement 3 holds. Since $W_{\square}(\mathbf{M}_{G_N}, \mathbf{M}_{G_n}) = 0$ if G_n and G_N differ by isolated vertices, we conclude that f is isolate-indifferent. Since quotient-convergent sequences $(G_n)_n$ are Cauchy in the W_{\square} -metric by Theorem 1.5, we conclude that $(f(G_n))_n \subseteq \mathbb{R}$ is Cauchy and hence converges. \square

As a first set of examples, quotient densities and homomorphism numbers are quotient-testable. More generally, we show next that a large class of polynomial graph parameters are testable, because all such polynomial parameters are functions of fixed-dimensional random quotients of their input graphs.

Proposition 4.9 (Polynomial parameters). *Suppose f is an isolate-indifferent graph parameter such that $f|_{\Delta^{n \times n}}$ is given by a polynomial function for each n , and that all these polynomials have degree $d \in \mathbb{N}$. Then f is quotient-testable.*

Further, a graph parameter f is quotient-testable if and only if for any $\epsilon > 0$ there exists a polynomial quotient-testable parameter p satisfying $|f(G) - p(G)| \leq \epsilon$ for all graphs G of all sizes.

Proof. For the first claim, it was shown in [38, Thm. 5.2] that there exists a polynomial q_k over $\mathbb{R}^{k \times k}$ satisfying $f(G_n) = \mathbb{E}q_k(\rho(F_{k,n})G_n)$ for all $G_n \in \Delta^{n \times n}$ and all $n \in \mathbb{N}$. Since q_k is L -Lipschitz on $\Delta^{k \times k}$ for some $L > 0$, we have by Proposition 3.2 that

$$\begin{aligned} |f(G_n) - f(G_m)| &= |\mathbb{E}q_k(\rho(F_{k,n})G_n) - \mathbb{E}q_k(\rho(F_{k,m})G_m)| \leq LW_1(\rho(F_{k,n})G_n, \rho(F_{k,m})G_m) \\ &\leq Lk^2W_{\square}(\mathbf{M}_{G_n}, \mathbf{M}_{G_m}), \end{aligned}$$

for any two graphs G_n, G_m of any two sizes. Thus, the function f is testable by Proposition 4.8(3), as can be seen by setting $\delta(\epsilon) = \frac{\epsilon}{Lk^2}$.

For the second claim, if for any $\epsilon > 0$ there is a testable parameter p satisfying $|f(G) - p(G)| \leq \epsilon$ for all G , then f is a uniform limit of W_{\square} -continuous functions and hence is W_{\square} -continuous itself. By Proposition 4.8, the parameter f is therefore testable. Conversely, if f is testable then it extends to a continuous $\bar{f}: \mathcal{M} \rightarrow \mathbb{R}$ by Proposition 4.8 again. Note that the collection of polynomial testable parameters is an algebra, contains constant functions, and separates points by Corollary 3.4. Since \mathcal{M} is compact, Stone–Weierstrass implies that \bar{f} is the uniform limit of polynomial testable parameters. \square

We combine the insights of Sections 4.1–4.2 along with the above characterization of quotient-testable parameters to show that these can indeed be tested by edge sampling.

Theorem 4.10. *Consider a graph parameter $f: \bigsqcup_n \Delta^{n \times n} \rightarrow \mathbb{R}$ whose restrictions $f|_{\Delta^{n \times n}}$ are continuous (in the usual topology on $\Delta^{n \times n}$) for each n . Then f is quotient-testable if and only if for every $\epsilon > 0$, there exists $k \in \mathbb{N}$ such that for any finite graph G of any size we have $|f(G) - f(G^{(k)}[\mathbf{M}_G])| \leq \epsilon$ with probability at least $1 - \epsilon$. In this case, we can take $k(\epsilon) = \left\lceil \left(\frac{174 + \sqrt{\log(\epsilon^{-1/2})}}{\delta(\epsilon)} \right)^2 \right\rceil$ where $\delta(\epsilon) > 0$ is as in Proposition 4.8(3).*

Proof. Suppose f is quotient-testable. For any $\epsilon > 0$, let $\delta(\epsilon)$ be as in Proposition 4.8(3) and set $k = k(\epsilon)$ as in the theorem statement. Then, noting that $\epsilon = e^{-2\tilde{\epsilon}^2}$ where $\tilde{\epsilon} = \sqrt{\log(\epsilon^{-1/2})}$, with probability at least $1 - \epsilon$ we have $W_{\square}(\mathbf{M}_G, \mathbf{M}_{G^{(k)}}) \leq \frac{174 + \tilde{\epsilon}}{\sqrt{k}} \leq \delta(\epsilon)$ and hence $|f(G) - f(G^{(k)}[\mathbf{M}_G])| \leq \epsilon$ by Theorem 4.3.

We turn to proving the converse. If G_1 and G_2 differ by isolated vertices, then $\mathbf{M}_{G_1} = \mathbf{M}_{G_2}$ and hence $G^{(k)}[\mathbf{M}_{G_1}] \stackrel{d}{=} G^{(k)}[\mathbf{M}_{G_2}]$. Therefore, for any $\epsilon > 0$ let $k \in \mathbb{N}$ be as in the theorem statement and let $G^{(k)}$ be a sample from this common distribution. Then with probability $1 - 2\epsilon$ we have $|f(G_1) - f(G^{(k)})| \leq \epsilon$ and $|f(G^{(k)}) - f(G_2)| \leq \epsilon$, so in particular there is a realization $G^{(k)}$ of $G^{(k)}$ satisfying both inequalities. We then have $|f(G_1) - f(G_2)| \leq |f(G_1) - f(G^{(k)})| + |f(G^{(k)}) - f(G_2)| \leq 2\epsilon$. Since $\epsilon > 0$ was arbitrary, we conclude that f is isolate-indifferent. A similar argument shows that $|f|$ is bounded. Indeed, since $|f|$ restricted to the compact set $\Delta^{2k \times 2k}$ is continuous, it is bounded by some $B_{2k} > 0$. For any other graph G of any size, some realization $G^{(k)}$ of $G^{(k)}[\mathbf{M}_G]$ satisfies $|f(G) - f(G^{(k)})| \leq \epsilon$, and hence $|f(G)| \leq B_{2k} + \epsilon$ for all graphs G . Denoting $B = \sup_G |f(G)|$, we then have $|f(G) - \mathbb{E}f(G^{(k)}[\mathbf{M}_G])| \leq \mathbb{E}|f(G) - f(G^{(k)}[\mathbf{M}_G])| \leq \epsilon(1 - \epsilon) + 2B\epsilon$. Finally, if (G_n) is quotient-convergent, then $(G^{(k)}[\mathbf{M}_{G_n}])_n$ is weakly-convergent, and by continuity of $f|_{\Delta^{2k \times 2k}}$, the sequence $(\mathbb{E}f(G^{(k)}[\mathbf{M}_{G_n}]))_n$ is convergent. Therefore, we have

$$|f(G_n) - f(G_m)| \leq 2\epsilon(1 - \epsilon + 2B) + |\mathbb{E}f(G^{(k)}[\mathbf{M}_{G_n}]) - \mathbb{E}f(G^{(k)}[\mathbf{M}_{G_m}])| \leq 3\epsilon(1 - \epsilon + 2B)$$

for all large n, m , showing that $(f(G_n))$ is convergent. \square

We remark that the graphon analog of the characterization of quotient-testable parameters in Theorem 4.10 was used as the definition of testable parameters in [16, Def. 2.11].

The upshot of this result is that a continuous graph parameter being quotient-testable is equivalent to the parameter being testable using edge sampling. In particular, Theorem 4.10 gives a simple recipe for estimating the value of a testable parameter $f(G)$ on an arbitrarily-large graph G : sample k edges iid from G proportionally to their edge weights, and compute the parameter $f(G^{(k)})$ of the resulting small graph $G^{(k)}$. The number of edges needed is given explicitly as $k = \left\lceil \left(\frac{174 + \sqrt{\log(\epsilon^{-1/2})}}{\delta(\epsilon)} \right)^2 \right\rceil$ in terms of the desired accuracy ϵ and the modulus of continuity $\delta(\epsilon)$ of f in the W_\square -metric. When f is L -Lipschitz continuous in the W_\square -metric, we can set $\delta(\epsilon) = \epsilon/L$.

Many graph parameters of interest are indeed Lipschitz-continuous, including all polynomial parameters as the proof of Proposition 4.9 demonstrates. Thus, the key quantity that is needed to bound the required number of edges to test the values of such parameters is their Lipschitz constants. We can explicitly bound the Lipschitz constants of quotient densities. Specifically, for a multigraph $H \in \mathbb{N}^{k \times k}$ we have

$$|t_Q(H; \mathbf{M}_1) - t_Q(H; \mathbf{M}_2)| = |\mathbb{E} \mathbf{G}_k[\mathbf{M}_1]^H - \mathbb{E} \mathbf{G}_k[\mathbf{M}_2]^H| \leq k^2 \|H\|_\infty W_\square(\mathbf{M}_1, \mathbf{M}_2),$$

since the polynomial $G \mapsto G^H$ is $\|H\|_\infty$ -Lipschitz with respect to the ℓ_1 norm on $\Delta^{k \times k}$. By using (10) and (12), we can similarly derive Lipschitz constants for homomorphism numbers. In the next example, we derive the Lipschitz constant of a particular graph parameter that can be used to test the presence of hubs.

Example 4.11 (Testing for hubs). *Consider the graph parameter*

$$f(G) = \text{hom}(S_2^{(\text{out})}; G) + \text{hom}(S_2^{(\text{in})}; G) = \|G \mathbf{1}\|_2^2 + \|G^\top \mathbf{1}\|_2^2,$$

where $S_2^{(\text{out})}, S_2^{(\text{in})} \in \mathbb{N}^{3 \times 3}$ are outward- and inward-pointing stars with two leaves, respectively. Define the polynomial q_2 on 2×2 matrices by

$$q_2(G) = (G_{1,1} + G_{1,2} - G_{2,1} - G_{2,2})^2 + (G_{1,1} + G_{2,1} - G_{1,2} - G_{2,2})^2,$$

and observe that $f(G_n) = \mathbb{E} q_2(\rho(F_{2,n})G_n)$ for any $G_n \in \Delta^{n \times n}$ and any $n \in \mathbb{N}$. Indeed, more generally for any grapheur \mathbf{M} with random locations (T_i) , let $R_i \stackrel{iid}{\sim} \text{Unif}(\{\pm 1\})$ be iid Rademacher random variables and note that

$$\begin{aligned} \mathbb{E} q_2(\mathbf{G}_2[\mathbf{M}]) &= \mathbb{E} \left[\sum_{i \in \mathbb{N}} (E \mathbf{1} + \sigma)_i (\mathbf{1}_{T_i < 1/2} - \mathbf{1}_{T_i > 1/2}) \right]^2 + \mathbb{E} \left[\sum_{i \in \mathbb{N}} (E^\top \mathbf{1} + \varsigma)_i (\mathbf{1}_{T_i < 1/2} - \mathbf{1}_{T_i > 1/2}) \right]^2 \\ &= \mathbb{E} \left[\sum_{i \in \mathbb{N}} (E \mathbf{1} + \sigma)_i R_i \right]^2 + \mathbb{E} \left[\sum_{i \in \mathbb{N}} (E^\top \mathbf{1} + \varsigma)_i R_i \right]^2 = \|E \mathbf{1} + \sigma\|_2^2 + \|E^\top \mathbf{1} + \varsigma\|_2^2, \end{aligned}$$

where $\mathbf{1}_{T_i < 1/2} = 1$ if $T_i < 1/2$ and zero otherwise and similarly for $\mathbf{1}_{T_i > 1/2}$, and we note that $(\mathbf{1}_{T_i < 1/2} - \mathbf{1}_{T_i > 1/2})_{i \in \mathbb{N}} \stackrel{d}{=} (R_i)_{i \in \mathbb{N}}$. We conclude that f can be extended to all grapheurs \mathbf{M} by $\bar{f}(\mathbf{M}) = \mathbb{E} q_2(\mathbf{G}_2[\mathbf{M}])$. This extension is Lipschitz-continuous. Explicitly, since $|\partial_{G_{i,j}} q_2(G)| = 2|G_{1,1} - G_{2,2}|$ if $i = j$ and $2|G_{i,j} - G_{j,i}|$ if $i \neq j$, we conclude that q_2 is 2-Lipschitz on $\Delta^{2 \times 2}$ with respect to $\|\cdot\|_1$, and hence that

$$|\bar{f}(\mathbf{M}_1) - \bar{f}(\mathbf{M}_2)| = |\mathbb{E} q_2(\mathbf{G}_2[\mathbf{M}_1]) - \mathbb{E} q_2(\mathbf{G}_2[\mathbf{M}_2])| \leq 2W_1(\mathbf{G}_2[\mathbf{M}_1], \mathbf{G}_2[\mathbf{M}_2]) \leq 8W_\square(\mathbf{M}_1, \mathbf{M}_2),$$

for any two grapheurs $\mathbf{M}_1, \mathbf{M}_2 \in \mathcal{M}$ by Proposition 3.2. Finally, observe that $\bar{f}(\mathbf{M}) = 0$ if and only if $E = 0$ and $\sigma = \varsigma = 0$, hence if and only if \mathbf{M} has no hubs. In particular, $W_\square(\mathbf{M}_G, \mathbf{M}) \geq f(G)/8$ for any grapheur \mathbf{M} with no hubs.

We can now use the above parameter to test for the existence of significant hubs in a graph. Specifically, suppose we are given an arbitrarily-large graph G , and we sample $k = \left\lceil 64 \left(\frac{174 + \sqrt{\log(\epsilon^{-1/2})}}{\epsilon} \right)^2 \right\rceil$ edges from G to obtain a graph $G^{(k)}$. Then with probability $1 - \epsilon$, we have $W_\square(\mathbf{M}_G, \mathbf{M}) \geq \frac{f(G^{(k)}) - \epsilon}{8}$ for any grapheur \mathbf{M} with no hubs. In this sense, large values of $f(G^{(k)})$ indicate the likely presence of significant hubs in G .

We observe that testing for hubs also has an appealing interpretation in terms of quotients. A graphneur \mathbf{M} has no hubs if and only if there is a $\theta \in [0, 1]$ satisfying $\mathbf{G}_m[\mathbf{M}] = \theta \frac{1}{m^2} \mathbf{1} \mathbf{1}^\top + (1 - \theta) \frac{1}{m} \mathbf{I}_m$ (a deterministic matrix) for all $m \in \mathbb{N}$. Thus, testing for hubs also amounts to testing whether the random quotients of a graph G deviate from a uniform distribution of edge weights, i.e., quotients of the above form. See Figures 1-2 for example. The parameter of Example 4.11 yields a lower bound on this deviation, as by Proposition 3.2 we have

$$\min_{\theta \in [0, 1]} \mathbb{E} \left\| \theta \frac{1}{m^2} \mathbf{1} \mathbf{1}^\top + (1 - \theta) \frac{1}{m} \mathbf{I}_m - \mathbf{G}_m[\mathbf{M}_G] \right\|_1 \geq \frac{f(G)}{8} - \frac{4}{m}.$$

For example, the values of the parameter f on the full graphs in Figures 1-2 are 0.0073 and 0.010, respectively. Estimating this parameter by sampling $k = 10^3$ edges, we obtain values of 0.0084 and 0.012, respectively. For comparison, the value of this parameter on an Erdős–Rényi random graph with edge probability 1/2 on 1,500 vertices is 0.001.

4.4 Missing Proofs from Section 4

In this section we give all the proofs missing from Section 4.

Proofs from Section 4.2

In this section, we prove the two lemmas used in Section 4.2 to prove Theorem 4.3. We begin by proving Lemma 4.4 by constructing an explicit coupling between \mathbf{M} , $\mathbf{M}_{\mathbf{G}^{(n)}[\mathbf{M}]}$, and $\mathbf{G}^{(n)}[\mathbf{M}]$ with the claimed properties.

Proof (Lemma 4.4). Sample $\mu \sim \mathbf{M}$ and set $\mathbf{M}' = \mu$. Next, sample $(x_i, y_i) \stackrel{iid}{\sim} \mu$, let $\{t_1, \dots, t_N\} = \{x_i\} \cup \{y_i\}$, form $\mathbf{G}^{(n)}$ as in Section 4.1, and set

$$\mathbf{M}'_{\mathbf{G}^{(n)}} = \frac{1}{n} \sum_{i=1}^n \delta_{(x_i, y_i)} = \sum_{i,j=1}^N (\mathbf{G}^{(n)})_{i,j} \delta_{(t_i, t_j)}. \quad (16)$$

We proceed to argue that this coupling satisfies all the claimed properties.

By construction, we have $\mathbf{M}' \stackrel{d}{=} \mathbf{M}$ and $\mathbf{G}^{(n)} \stackrel{d}{=} \mathbf{G}^{(n)}[\mathbf{M}]$. Next, because \mathbf{M} is exchangeable, we claim that $t_1, \dots, t_N \stackrel{iid}{\sim} \text{Unif}([0, 1])$. Indeed, $(x_i, y_i) \stackrel{d}{=} (\sigma(x_i), \sigma(y_i))$ for all $\sigma \in S_{[0,1]}$, hence $(x_i, y_i) \sim \theta \lambda^2 + (1 - \theta) \lambda D$. Therefore, for each i either $x_i = y_i \sim \text{Unif}([0, 1])$ or $x_i, y_i \stackrel{iid}{\sim} \lambda^2$. Combining this with the fact that the (x_i, y_i) are independent for different i , we obtain the claim. Therefore, $\mathbf{M}_{\mathbf{G}^{(n)}[\mathbf{M}]} \stackrel{d}{=} \frac{1}{n} \sum_{i=1}^n \delta_{(x_i, y_i)} = \mathbf{M}'_{\mathbf{G}^{(n)}}$, the latter being an empirical measure drawn from $\mu \sim \mathbf{M}$. Finally, we have $\mathbf{M}' | \mathbf{G}^{(n)} \stackrel{d}{=} \mathbf{M}$ and $\mathbf{M}'_{\mathbf{G}^{(n)}} | \mathbf{G}^{(n)} \stackrel{d}{=} \mathbf{M}_{\mathbf{G}^{(n)}}$ as can be seen from (16), as $\mathbf{G}^{(n)}$ only depends on the multiplicities of $\{(t_i, t_j)\}$ rather than their locations, and the multiplicities and locations of a random exchangeable measure are independent of each other [17, Thm. 2]. \square

We now turn to proving Lemma 4.5, combining standard results from VC theory.

Proof (Lemma 4.5). Fix any $\mu \in \mathcal{P}([0, 1]^2)$. By the proof of [43, Thm. 4.10], we have

$$\mathbb{E}_{\substack{\mu^{(n)} \\ \text{emp. meas.} \sim \mu}} \sup_{R \in \mathcal{AAR}} |\mu_n(R) - \mu(R)| \leq 2R_n(\mathcal{F}_{\mathcal{AAR}}),$$

where $R_n(\mathcal{F}) = \mathbb{E}_{\substack{X_i \stackrel{iid}{\sim} \mu \\ \varepsilon_i \stackrel{iid}{\sim} \text{Unif}(\{\pm 1\})}} \sup_{f \in \mathcal{F}} \left| \frac{1}{n} \sum_{i=1}^n \varepsilon_i f(X_i) \right|$ is the Rademacher complexity of a class of functions \mathcal{F} and we set $\mathcal{F}_{\mathcal{AAR}} = \{\mathbf{1}_R : R \in \mathcal{AAR}\}$. We further have by [43, Ex. 5.24] that

$$R_n(\mathcal{F}) \leq \frac{24}{\sqrt{n}} \int_0^{\text{diam}(\mathcal{F})} \mathbb{E} \sqrt{\log N(t; \mathcal{F}; \|\cdot\|_{\mu_n})} dt,$$

where $N(t; \mathcal{F}; \|\cdot\|_{\mu_n})$ is the t -covering number of \mathcal{F} in the (random) metric $\|f - g\|_{\mu_n} = \sqrt{\mathbb{E}_{\mu_n}(f - g)^2}$ defined by the empirical measure μ_n . If \mathcal{F} is class of $\{0, 1\}$ -valued functions, then $\text{diam}(\mathcal{F}) \leq 1$. Finally, we have by [44, Cor. 1] that

$$N(t; \mathcal{F}_{\mathcal{AAR}}; \|\cdot\|_{\mu}) \leq e(V+1) \left(\frac{2e}{t}\right)^V,$$

for any measure μ , where $V = \text{VC}(\mathcal{AAR}) = 4$ is the VC-dimension of the class of axis-aligned rectangles. Combining the above results, we conclude that

$$\mathbb{E}_{\substack{\mu^{(n)} \\ \text{emp. meas.} \sim \mu}} \sup_{R \in \mathcal{AAR}} |\mu_n(R) - \mu(R)| \leq \frac{48}{\sqrt{n}} \int_0^1 \sqrt{\log(5e) + 4 \log(2e/t)} dt \leq \frac{174}{\sqrt{n}},$$

where we numerically evaluated the last integral to 12 digits of accuracy to get $3.624000076562 \leq 3.625$. \square

Proof of Proposition 3.6 from Section 3.2

Using Theorem 4.3 and its proof, we can now prove Proposition 3.6 by comparing both the quotient-convergent sequence (G_n) and its limit \mathbf{M} to their edge-sampled random graphs.

Proof (Proposition 3.6). Since (G_n) quotient-converges to \mathbf{M} , we have $W_1(\mathbf{G}^{(k)}[\mathbf{M}_{G_n}], \mathbf{G}^{(k)}[\mathbf{M}]) \rightarrow 0$ as $n \rightarrow \infty$ for each $k \in \mathbb{N}$. Pick $k \geq (174/\epsilon)^2$, sample $(G_n^{(k)}, G_{\mathbf{M}}^{(k)})$ from a coupling of $\mathbf{G}^{(k)}[\mathbf{M}_{G_n}]$ and $\mathbf{G}^{(k)}[\mathbf{M}]$ attaining $W_1(\mathbf{G}^{(k)}[\mathbf{M}_{G_n}], \mathbf{G}^{(k)}[\mathbf{M}])$, and note that with probability $1 - 4e^{-2\epsilon^2} - \frac{W_1(\mathbf{G}^{(k)}[\mathbf{M}_{G_n}], \mathbf{G}^{(k)}[\mathbf{M}])}{\epsilon}$ we have

$$\sup_{R \in \mathcal{AAR}} |\mu_{G_n^{(k)}}(R) - \mu_{G_n}(R)| \leq \epsilon, \quad \sup_{R \in \mathcal{AAR}} |\mu_{G_{\mathbf{M}}^{(k)}}(R) - \mu_{\mathbf{M}}(R)| \leq \epsilon, \quad \text{and} \quad \|G_n^{(k)} - G_{\mathbf{M}}^{(k)}\|_1 \leq \epsilon, \quad (17)$$

for $(\mu_{G_{\mathbf{M}}^{(k)}}, \mu_{\mathbf{M}}) \sim (\mathbf{M}'_{G_{\mathbf{M}}^{(k)}}, \mathbf{M}')$ and $(\mu_{G_n^{(k)}}, \mu_{G_n}) \sim (\mathbf{M}'_{G_n^{(k)}}, \mathbf{M}'_{G_n})$ sampled from the coupling of Lemma 4.4 conditioned on $G_{\mathbf{M}}^{(k)}$ and $G_n^{(k)}$, respectively. Picking n large enough, the above occurs with positive probability, so we can find deterministic $G_n^{(k)}$ and $G_{\mathbf{M}}^{(k)}$ satisfying (17) for all large n . Explicitly, by Lemma 4.4 there are locations $t_i \in [0, 1]$ and a permutation $\pi_n \in \mathcal{S}_n$ satisfying

$$\mu_{G_n} = \sum_{i,j=1}^n (G_n)_{i,j} \delta_{(t_i, t_j)}, \quad \mu_{G_n^{(k)}} = \sum_{i,j=1}^n (\pi_n G_n^{(k)})_{i,j} \delta_{(t_i, t_j)}.$$

By considering rectangles in (17) centered at each of the atoms of μ_{G_n} and $\mu_{G_n^{(k)}}$, and around rows and columns of these two measures, we conclude that

$$\|G_n - \pi_n G_n^{(k)}\|_{\infty} \leq \epsilon, \quad \|G_n \mathbb{1} - \pi_n G_n^{(k)} \mathbb{1}\|_{\infty} \leq \epsilon, \quad \text{and} \quad \|G_n^{\top} \mathbb{1} - \pi_n (G_n^{(k)})^{\top} \mathbb{1}\|_{\infty} \leq \epsilon. \quad (18)$$

Similarly, we can find a permutation $\tilde{\pi}_M \in \mathcal{S}_M$ for large enough $M \in \mathbb{N}$ and atoms $(t_i)_{i \in \mathbb{N}}$ and $(t'_i)_{i=1}^{M-m}$ for some $m \leq M$ satisfying

$$\begin{aligned} \mu_{\mathbf{M}} &= \sum_{i,j \in \mathbb{N}} E_{i,j} \delta_{(t_i, t_j)} + \sum_{i \in \mathbb{N}} \sigma_i(\delta_{t_i} \otimes \lambda) + \sum_{i \in \mathbb{N}} \varsigma_i(\lambda \otimes \delta_{t_i}) + \theta \lambda^2 + \vartheta \lambda_D, \\ \mu_{G_{\mathbf{M}}^{(k)}} &= \sum_{i,j=1}^m (\tilde{\pi}_M G_{\mathbf{M}}^{(k)})_{i,j} \delta_{(t_i, t_j)} + \sum_{i=1}^m \sum_{j=m+1}^M (\tilde{\pi}_M G_{\mathbf{M}}^{(k)})_{i,j} \delta_{(t_i, t'_j)} + \sum_{i=m+1}^M \sum_{j=1}^m (\tilde{\pi}_M G_{\mathbf{M}}^{(k)})_{i,j} \delta_{(t'_i, t_j)}. \end{aligned}$$

Here we have separated the edges sampled from each summand in \mathbf{M} . Moreover, with probability 1 we have $(\tilde{\pi}_M G_{\mathbf{M}}^{(k)})_{i,j} \in \{0, 1/k\}$ if $i > m$ or $j > m$. Note that $k^{-1} \leq (\epsilon/174)^2 \leq \epsilon$ for all small $\epsilon > 0$. By considering similar rectangles in (17), we therefore

$$\|E - \tilde{\pi}_M G_{\mathbf{M}}^{(k)}\|_{\infty} \leq \epsilon, \quad \|E \mathbb{1} + \sigma - \tilde{\pi}_M G_{\mathbf{M}}^{(k)} \mathbb{1}\|_{\infty} \leq \epsilon, \quad \text{and} \quad \|E^{\top} \mathbb{1} + \varsigma - (\tilde{\pi}_M G_{\mathbf{M}}^{(k)})^{\top} \mathbb{1}\|_{\infty} \leq \epsilon. \quad (19)$$

Finally, since $\|G_n^{(k)} - G_{\mathbf{M}}^{(k)}\|_1 \leq \epsilon$, we conclude that

$$\|G_n^{(k)} - G_{\mathbf{M}}^{(k)}\|_{\infty} \leq \epsilon, \quad \|G_n^{(k)} \mathbf{1} - G_{\mathbf{M}}^{(k)} \mathbf{1}\|_{\infty} \leq \epsilon, \quad \text{and} \quad \|(G_n^{(k)})^{\top} \mathbf{1} - (G_{\mathbf{M}}^{(k)})^{\top} \mathbf{1}\|_{\infty} \leq \epsilon. \quad (20)$$

Combining (18), (19), and (20), we obtain

$$\|\tilde{\pi}_M \pi_n^{-1} G_n - E\|_{\infty} \leq 3\epsilon, \quad \|\tilde{\pi}_M \pi_n^{-1} G_n \mathbf{1} - (E \mathbf{1} + \sigma)\|_{\infty} \leq 3\epsilon, \quad \text{and} \quad \|\tilde{\pi}_M \pi_n^{-1} G_n^{\top} \mathbf{1} - (E^{\top} \mathbf{1} + \varsigma)\|_{\infty} \leq 3\epsilon.$$

Since $\epsilon > 0$ was arbitrary, our first three claims are proved. For the last claim, observe that $\lim_n \text{Tr}(G_n) = \lim_n \text{hom}([1], G_n)$ exists as it is the limit of the homomorphism numbers of a single self-loop. By Corollary 3.4 and the proof of Proposition 2.6, the limit of the homomorphism numbers is the same for any sequence of graphs quotient-converging to \mathbf{M} . By considering the sequence of graphs (13) derived from \mathbf{M} , we conclude that $\lim_n \text{Tr}(G_n) = \text{Tr}(E) + \vartheta$, proving the last claim. \square

5 Equipartition-Consistent Random Graph Models

Each grapheur \mathbf{M} yields a sequence of random graphs $(G_k[\mathbf{M}])$ via the construction (8). In Section 5.1, we show that these graphs are related to each other via equipartitions, and hence form an equipartition-consistent random graph model (Definition 1.8). We then prove that realizations of these random graphs $(G_k[\mathbf{M}])$ converge back to the grapheur \mathbf{M} in a suitable fashion. Finally, in Section 5.2 we prove Theorem 1.9, stating that any equipartition-consistent random graph model is a mixture of ones obtained from grapheurs as above.

5.1 Equipartition Consistency

We begin by arguing that $(G_k[\mathbf{M}])$ is equipartition-consistent. Note that $d_{k,nk} \circ F_{nk,m} \stackrel{d}{=} F_{k,m}$ for any $k, n, m \in \mathbb{N}$ by [38, Lemma 4.13], since each $i \in [m]$ is mapped independently by $F_{nk,m}$ to one of the fibers of $d_{k,nk}$ with the same probability. Therefore, for any $G \in \Delta^{n \times n}$ the sequence of distributions $(\text{Law}(\rho(F_{k,n})G_n))_k$ is indeed an equipartition-consistent model. If (G_m) is a sequence of finite graphs converging to a grapheur \mathbf{M} , then $\rho(F_{k,m})G_m \rightarrow G_k[\mathbf{M}]$ and $\rho(F_{nk,m})G_m \rightarrow G_{nk}[\mathbf{M}]$ weakly, hence

$$\text{Law}(\rho(d_{k,nk})G_{nk}[\mathbf{M}]) = \lim_m \text{Law}(\rho(d_{k,nk} \circ F_{nk,m})G_m) = \lim_m \text{Law}(\rho(F_{k,m})G_m) = \text{Law}(G_k[\mathbf{M}]),$$

showing that $(\text{Law}(G_k[\mathbf{M}]))_k$ is equipartition-consistent for any grapheur \mathbf{M} .

We now ask whether samples from an equipartition-consistent model $(G_k[\mathbf{M}])$ obtained from a grapheur $\mathbf{M} \in \mathcal{M}$ converge back to \mathbf{M} . The answer depends on how this sampling is performed.

Proposition 5.1. *Fix a grapheur $\mathbf{M} \in \mathcal{M}$ and consider the associated equipartition-consistent random graph model $(\mu_k = \text{Law}(G_k[\mathbf{M}]))_k$.*

1. *Draw $\mu \sim \mathbf{M}$ and let $G_k = (\mu(I_i^{(n)} \times I_j^{(n)}))_{i,j \in [k]}$ for each $k \in \mathbb{N}$. Then $G_k \sim \mu_k$ for all $k \in \mathbb{N}$, and (G_k) is quotient-convergent to \mathbf{M} almost surely.*
2. *Suppose $\mathbf{M} = \delta_{(T_1, T_2)}$ is the grapheur associated to a single edge. Draw $G_k \sim \mu_k$ independently of each other. Then $\limsup_k W_{\square}(\mathbf{M}_{G_k}, \mathbf{M}) \geq 1/4$ almost surely, so (G_k) does not converge to \mathbf{M} .*

Proof. For the first claim, fix any $\epsilon > 0$ and let $N \in \mathbb{N}$ be sufficiently large so that $\sum_{i>N \text{ or } j>N} E_{i,j} + \sum_{i>N} (\sigma_i + \varsigma_i) \leq \epsilon$. Drawing $\mu \sim \mathbf{M}$ amounts to sampling locations $T_i \stackrel{iid}{\sim} \text{Unif}([0, 1])$ in (5), and these samples are distinct almost surely. Therefore, there exists $K \geq N \in \mathbb{N}$ such that T_1, \dots, T_N lie in distinct intervals $(I_i^{(k)})_{i \in [k]}$ for all $k \geq K$. Therefore, for all $k \geq K$ we can find a permutation $\pi_k \in S_k$ such that $\pi_k G_k = \rho(\iota_{k,N})(E_{i,j} + \sigma_i/k + \varsigma_j/k)_{i,j \in [N]} + N_k + \theta \frac{1}{k^2} \mathbf{1}_k \mathbf{1}_k^{\top} + \vartheta \frac{1}{k} I_k$ where $\|N_k\|_1 \leq \epsilon$. Using the coupling $(\mathbf{M}'_{G_k}, \mathbf{M}')$ obtained by drawing random locations (T_i) , setting \mathbf{M}' to (5), and setting $\mathbf{M}'_{G_k} = \sum_{i,j=1}^N (\pi_k G_k)_{i,j} \delta_{(T_i, T_j)}$, we conclude that $\limsup_{k \rightarrow \infty} W_{\square}(\mathbf{M}_{G_k}, \mathbf{M}) \leq \epsilon$. Since $\epsilon > 0$ was arbitrary, we obtain the claimed convergence.

For the second claim, note that with probability $1/k$ we have $T_1, T_2 \in I_i^{(k)}$ for the same $i \in [k]$, in which case $G_k = \text{diag}(0, \dots, 1, \dots, 0)$ and $W_{\square}(\mathbf{M}_{G_k}, \mathbf{M}) \geq 1/4$ as can be seen by setting $S = [0, 1/2]$ and $T = [1/2, 1]$ in (7). Since these events are independent and the sum of their probabilities $\sum_k k^{-1}$ diverges, Borel–Cantelli implies that $\limsup_k W_{\square}(\mathbf{M}_{G_k}, \mathbf{M}) \geq 1/4$ almost surely. \square

In words, the quotients must be sampled in a manner that is related across different sizes for the samples to converge to the underlying grapheur; if we instead sample random quotients independently for each size, the resulting sequence almost surely does not converge to the underlying grapheur. In contrast, if we sample growing-sized random graphs from a graphon independently for different sizes, then the resulting sequence does converge back to the graphon almost surely as a consequence of Szemerédi's regularity lemma [14, Lemma 11.8]. Proposition 5.1(2) thus implies that the analog of Theorem 1.7 fails for random quotients. More precisely, for any rate of convergence $(R(k) \in \mathbb{R}_{\geq 0})_{n \in \mathbb{N}}$ with $R(k) \rightarrow 0$, we have $\sum_{k \in \mathbb{N}} \mathbb{P}[W_{\square}(\mathbf{M}, \mathbf{G}_k[\mathbf{M}]) > R(k)] = \infty$ by Borel–Cantelli. We leave the precise analysis of convergence rates of random quotients for future work.

5.2 Proof of Theorem 1.9

Our goal in this section is to prove Theorem 1.9, showing that every equipartition-consistent random graph model arises from a unique random exchangeable measure. We deduce the existence of the desired random measure from results on inverse limits of random histograms [45], and proving that this random measure is exchangeable proceeds via standard approximation arguments. To apply the results of [45], we require the following lemma on the sequence of means of an equipartition-consistent model.

Lemma 5.2. *For any equipartition-consistent random graph model (μ_k) , there exists $\theta \in [0, 1]$ satisfying $\mathbb{E}_{X \sim \mu_k} X = \theta \frac{1}{k^2} \mathbb{1}_k \mathbb{1}_k^\top + (1 - \theta) \frac{1}{k} I_k$ for all $k \in \mathbb{N}$.*

Proof. Note that $M_n = \mathbb{E}_{X \sim \mu_n} X \in \mathbb{S}^n$ is a deterministic matrix invariant under simultaneous permutations of its rows and columns, because μ_n is exchangeable. Any such matrix has the same on-diagonal and off-diagonal entries, hence there exist $\theta_n, \vartheta_n \in \mathbb{R}$ satisfying $M_n = \theta_n \frac{1}{n} \mathbb{1}_{n \times n} + \vartheta_n \frac{1}{n} I_n$. Since $\mathbb{1}^\top X \mathbb{1} = 1$ almost surely, we similarly have $\mathbb{1}^\top M_n \mathbb{1} = \theta_n + \vartheta_n = 1$. Also, since $X \geq 0$ almost surely, we must have $\theta_n \in [0, 1]$. Finally, since $\rho(d_{n,N}) X_N \stackrel{d}{=} X_n$ whenever $n|N$, we must have

$$M_n = \rho(d_{n,N}) M_N = \theta_N \frac{1}{n^2} \mathbb{1}_{n \times n} + (1 - \theta_N) \frac{1}{n} I_n.$$

Since $\frac{1}{n^2} \mathbb{1}_{n \times n}$ and $\frac{1}{n} I_n$ are linearly independent for all $n \geq 2$, we must have $\theta_N = \theta_n$ whenever $n|N$, and therefore for all $n, N \in \mathbb{N}$. This proves the claim. \square

Combining the above lemma with [45], we are ready to prove Theorem 1.9.

Proof (Theorem 1.9). Uniqueness is clear since squares of the form $I_i^{(k)} \times I_j^{(k)}$ generate the Borel σ -algebra on $[0, 1]^2$. To prove existence we appeal to [45, Thm. 1.1], which states that there exists a random probability measure \mathbf{M} on $[0, 1]^2$ (not guaranteed to be exchangeable) satisfying $\mathbf{M}(I_i^{(k)} \times I_j^{(k)}) \sim \mu_k$ for all k if there is a deterministic measure $\mu \in \mathcal{P}([0, 1]^2)$ satisfying $(\mu(I_i^{(k)} \times I_j^{(k)}))_{i,j \in [k]} = \mathbb{E}_{X \sim \mu_k} X$ for all $k \in \mathbb{N}$. Such a deterministic measure exists by Lemma 5.2, since

$$((\theta \lambda^2 + (1 - \theta) \lambda_D)(I_i^{(k)} \times I_j^{(k)}))_{i,j \in [k]} = \theta \frac{1}{k^2} \mathbb{1}_{k \times k} + (1 - \theta) \frac{1}{k} I_k.$$

We proceed to prove that the above \mathbf{M} is exchangeable. To do so, it suffices to prove that for any continuous function $f: [0, 1]^2 \rightarrow \mathbb{R}$ and any measure-preserving bijection σ , we have $\mathbb{E}_{\mathbf{M}} f \stackrel{d}{=} \mathbb{E}_{\mathbf{M}} f \circ (\sigma, \sigma)$. By [37, Prop. 9.1], it further suffices to prove this equality for $\sigma \in S_m$, a permutation of the m intervals $I_1^{(m)}, \dots, I_m^{(m)}$ acting by $\sigma((i-1)/m + x) = (\sigma(i)-1)/m + x$ for $x \in [0, 1/m)$ for $i \in [m]$.

For any continuous function $f: [0, 1]^2 \rightarrow \mathbb{R}$, let $f_n = \sum_{i,j=1}^n f_{i,j}^{(n)} \mathbb{1}_{I_i^{(n)} \times I_j^{(n)}}$ where $f_{i,j} = f(\frac{i}{n} - \frac{1}{2n}, \frac{j}{n} - \frac{1}{2n})$ is the value of f on the centers of the squares $I_i^{(n)} \times I_j^{(n)}$, whose diameter in ℓ_∞ is $1/2n$. Since f is continuous on a compact set, it is uniformly continuous, hence for any $\epsilon > 0$ we have $|f(x) - f_n(x)| \leq \epsilon$ for all $x \in [0, 1]^2$ and all large n . Note that

$$\mathbb{E}_{\mathbf{M}} f_n = \sum_{i,j=1}^n f_{i,j}^{(n)} \mathcal{M}(I_i^{(n)} \times I_j^{(n)}) = \sum_{i,j=1}^n f_{i,j}^{(n)} (\mathbf{G}_n[\mathbf{M}])_{i,j}.$$

Therefore, for any $m, k \in \mathbb{N}$ and $\sigma_m \in S_m$, we view σ_m as an interval permutation and as a permutation on $[mk]$ letters permuting consecutive intervals of length k , in which case we get

$$\mathbb{E}_M f_{mk} \circ (\sigma_m, \sigma_m) = \sum_{i,j=1}^m f_{i,j}^{(mk)} (G_{mk}^{(M)})_{\sigma^{-1}(i), \sigma^{-1}(j)} \stackrel{d}{=} \sum_{i,j=1}^m f_{i,j}^{(mk)} (G_{mk}^{(M)})_{i,j} = \mathbb{E}_M f_{mk}.$$

Since this holds for all $k \in \mathbb{N}$, we have

$$\begin{aligned} W_1(\mathbb{E}_M f, \mathbb{E}_M f \circ (\sigma_m, \sigma_m)) &\leq W_1(\mathbb{E}_M f, \mathbb{E}_M f_{mk}) + W_1(\mathbb{E}_M f_{mk}, \mathbb{E}_M f_{mk} \circ (\sigma_m, \sigma_m)) \\ &\quad + W_1(\mathbb{E}_M f_{mk} \circ (\sigma_m, \sigma_m), \mathbb{E}_M f \circ (\sigma_m, \sigma_m)) \leq 2\epsilon, \end{aligned}$$

for all large k , where the last inequality follows since M is supported on $[0, 1]^2 = \bigcup_{i,j} I_i^{(n)} \times I_j^{(n)}$. Since $\epsilon > 0$ was arbitrary, this proves the exchangeability of M . \square

6 Conclusions and Future Directions

We have introduced a new notion of limits of growing graphs based on convergence of their fixed-size random quotients. We showed that this notion of convergence can be equivalently characterized in terms of convergence of graph parameters such as homomorphism numbers and quotient densities. We characterized limits of quotient-convergent graph sequences via grapheurs, which are certain random exchangeable measures on $[0, 1]^2$. The limiting grapheur of a quotient-convergent sequence of graphs describes the asymptotic distribution of edge weight in the graphs. We then presented another equivalent view of quotient convergence via edge sampling, allowing us to prove a dual analog of Szemerédi's regularity lemma and to test properties of arbitrarily-large graphs by sampling edges from them. The number of edges needed is independent of the size of the graph, only depending on the desired accuracy and the Lipschitz constant of the graph parameter in question. Finally, we studied equipartition-consistent random graph models, showing that any such model corresponds to the sequence of random quotients associated to a mixture of grapheurs. We conclude by mentioning several questions suggested by our work.

(Optimal Szemerédi rate) Is the rate of $O(1/\sqrt{n})$ for approximating a grapheur by a graph on n edges optimal?

(Rates for random quotients) Is there a Szemerédi-type regularity lemma for random quotients, i.e., does $\mathbb{E}_{G_k[M]} W_{\square}(M_{G_k[M]}, M)$ converge to zero at a universal rate independent of M ? If not, how does this rate depend on M ?

(Extending graphon duality) Is Conjecture 3.11 on the bi-continuity of our duality pairing true? Can the duality with graphons in Section 3.3 be extended to limits of non-simple graphs? Can we relate the cut metric and our W_{\square} -metric via this duality?

(Inequalities in quotient densities) Corollary 3.5 shows that valid inequalities in quotient densities (or equivalently, homomorphism numbers) that hold for graphs of all sizes correspond to inequalities over grapheurs. Can we exploit the structure of the space of grapheurs to prove such inequalities, in analogy with the use of graphon theory to prove homomorphism density inequalities?

Acknowledgements

Eitan Levin and Venkat Chandrasekaran were supported in part by AFOSR grant FA9550-23-1-0070 and by NSF grant DMS-2502377.

References

[1] Mark Newman. *Networks*. Oxford University Press, 07 2018.

[2] Rui Castro, Mark Coates, Gang Liang, Robert Nowak, and Bin Yu. Network Tomography: Recent Developments. *Statistical Science*, 19(3):499 – 517, 2004.

[3] Shazia Tabassum, Fabiola S. F. Pereira, Sofia Fernandes, and João Gama. Social network analysis: An overview. *WIREs Data Mining and Knowledge Discovery*, 8(5):e1256, 2018.

[4] Matthew O. Jackson and Alison Watts. The evolution of social and economic networks. *Journal of Economic Theory*, 106(2):265–295, 2002.

[5] Danielle S Bassett and Olaf Sporns. Network neuroscience. *Nature neuroscience*, 20(3):353–364, 2017.

[6] Morton E O’Kelly. A geographer’s analysis of hub-and-spoke networks. *Journal of Transport Geography*, 6(3):171–186, 1998.

[7] Deborah L. Bryan and Morton E. O’Kelly. Hub-and-spoke networks in air transportation: An analytical review. *Journal of Regional Science*, 39(2):275–295, 1999.

[8] Jonathan D Power, Bradley L Schlaggar, Christina N Lessov-Schlaggar, and Steven E Petersen. Evidence for hubs in human functional brain networks. *Neuron*, 79(4):798–813, 2013.

[9] Xionglei He and Jianzhi Zhang. Why do hubs tend to be essential in protein networks? *PLOS Genetics*, 2(6):1–9, 06 2006.

[10] Martijn P Van den Heuvel and Olaf Sporns. Network hubs in the human brain. *Trends in cognitive sciences*, 17(12):683–696, 2013.

[11] Elizabeth Sall Ben Stabler, Hillel Bar-Gera. Transportation networks for research. <https://github.com/bstabler/TransportationNetworks>. Accessed: 2025-11-18.

[12] Steven J Cook, Travis A Jarrell, Christopher A Brittin, Yi Wang, Adam E Bloniarz, Maksim A Yakovlev, Ken CQ Nguyen, Leo T-H Tang, Emily A Bayer, Janet S Duerr, et al. Whole-animal connectomes of both *caenorhabditis elegans* sexes. *Nature*, 571(7763):63–71, 2019.

[13] Netzscheleuder. Complete *C. elegans* neurons (2019). https://networks.skewed.de/net/celegans_2019. Accessed: 2025-12-17.

[14] László Lovász. *Large networks and graph limits*, volume 60. American Mathematical Soc., 2012.

[15] László Lovász and Balázs Szegedy. Limits of dense graph sequences. *Journal of Combinatorial Theory, Series B*, 96(6):933–957, 2006.

[16] C. Borgs, J.T. Chayes, L. Lovász, V.T. Sós, and K. Vesztergombi. Convergent sequences of dense graphs I: Subgraph frequencies, metric properties and testing. *Advances in Mathematics*, 219(6):1801–1851, 2008.

[17] Olav Kallenberg. Exchangeable random measures in the plane. *Journal of Theoretical Probability*, 3(1):81–136, 1990.

[18] László Lovász and Balázs Szegedy. Random graphons and a weak positivstellensatz for graphs. *Journal of Graph Theory*, 70(2):214–225, 2012.

[19] Itai Benjamini and Oded Schramm. Recurrence of Distributional Limits of Finite Planar Graphs. *Electronic Journal of Probability*, 6(none):1 – 13, 2001.

[20] David Aldous and J Michael Steele. The objective method: probabilistic combinatorial optimization and local weak convergence. In *Probability on discrete structures*, pages 1–72. Springer, 2004.

[21] Gábor Elek. Note on limits of finite graphs. *Combinatorica*, 27(4):503–507, 2007.

[22] Harry Crane and Walter Dempsey. Edge exchangeable models for network data. *arXiv preprint arXiv:1603.04571*, 2016.

[23] Harry Crane and Walter Dempsey. Relational exchangeability. *Journal of Applied Probability*, 56(1):192–208, 2019.

[24] Diana Cai, Trevor Campbell, and Tamara Broderick. Edge-exchangeable graphs and sparsity. In D. Lee, M. Sugiyama, U. Luxburg, I. Guyon, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 29. Curran Associates, Inc., 2016.

[25] Trevor Campbell, Diana Cai, and Tamara Broderick. Exchangeable trait allocations. *Electronic Journal of Statistics*, 12(2):2290 – 2322, 2018.

[26] Svante Janson. On edge exchangeable random graphs. *Journal of statistical physics*, 173(3):448–484, 2018.

[27] Svante Janson. Graphons, cut norm and distance, couplings and rearrangements. *New York Journal of Mathematics*, 4, March 2013. NYJM Monographs.

[28] François Caron and Emily B. Fox. Sparse graphs using exchangeable random measures. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 79(5):1295–1366, 09 2017.

[29] Victor Veitch and Daniel M Roy. The class of random graphs arising from exchangeable random measures. *arXiv preprint arXiv:1512.03099*, 2015.

[30] Christian Borgs, Jennifer T. Chayes, Henry Cohn, and Nina Holden. Sparse exchangeable graphs and their limits via graphon processes. *Journal of Machine Learning Research*, 18(210):1–71, 2018.

[31] Christian Borgs, Jennifer T Chayes, Henry Cohn, and László Miklós Lovász. Identifiability for graphexes and the weak kernel metric. In *Building Bridges II: Mathematics of László Lovász*, pages 29–157. Springer, 2020.

[32] Christian Borgs, Jennifer T. Chayes, Henry Cohn, and Victor Veitch. Sampling perspectives on sparse exchangeable graphs. *The Annals of Probability*, 47(5):2754 – 2800, 2019.

[33] Christian Borgs, Jennifer T. Chayes, Souvik Dhara, and Subhabrata Sen. Limits of sparse configuration models and beyond: Graphexes and MultiGraphexes. *The Annals of Probability*, 49(6):2830 – 2873, 2021.

[34] Svante Janson. On convergence for graphexes. *European Journal of Combinatorics*, 104:103549, 2022.

[35] Victor Veitch and Daniel M. Roy. Sampling and estimation for (sparse) exchangeable graphs. *The Annals of Statistics*, 47(6):3274 – 3299, 2019.

[36] Cédric Villani et al. *Optimal transport: old and new*, volume 338. Springer, 2008.

[37] Olav Kallenberg et al. *Probabilistic symmetries and invariance principles*, volume 9. Springer, 2005.

[38] Eitan Levin and Venkat Chandrasekaran. Any-dimensional polynomial optimization via de Finetti theorems. *arXiv preprint arXiv:2507.15632*, 2025.

[39] Persi Diaconis and Svante Janson. Graph limits and exchangeable random graphs. *arXiv preprint arXiv:0712.2749*, 2007.

[40] Olav Kallenberg. *Foundations of modern probability*. Springer, 1997.

[41] Renata Possobon and Christian S. Rodrigues. Geometric properties of disintegration of measures. *Ergodic Theory and Dynamical Systems*, 45(5):1619–1648, 2025.

[42] Olav Kallenberg. Characterization and convergence of random measures and point processes. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete*, 27(1):9–21, 1973.

[43] Martin J Wainwright. *High-dimensional statistics: A non-asymptotic viewpoint*, volume 48. Cambridge university press, 2019.

- [44] David Haussler. Sphere packing numbers for subsets of the boolean n -cube with bounded Vapnik–Chervonenkis dimension. *Journal of Combinatorial Theory, Series A*, 69(2):217–232, 1995.
- [45] Peter Orbanz. Projective limit random probabilities on Polish spaces. *Electronic Journal of Statistics*, 5(none):1354 – 1373, 2011.