

ON HOPF IDEALS, INTEGRALITY, AND AUTOMORPHISMS OF QUANTUM GROUPS AT ROOTS OF 1

MATTHEW HARPER AND THOMAS KERLER

ABSTRACT. We consider skew-commutative subalgebras in Drinfeld-Jimbo quantum groups at a root of unity ζ generated by primitive power elements. We classify the centrality and commutativity of these skew-polynomial algebras depending on the Lie type and the order of ζ modulo 8. We describe Hopf ideals in the quantum group induced from ideals in these subalgebras, including the non-commutative cases.

Among these, we construct and analyze a family of Hopf ideals that depend on the choice of an element in the Weyl group. We show that they arise naturally both in the construction of (partial) R -matrices and as vanishing ideals of Bruhat subgroups. Specialization to the maximal element yields a rigorous construction of restricted quantum groups as pre-triangular Hopf algebras, independent of any choices.

Our treatment also includes even orders of ζ , non-simply laced Lie types, and minimal ground rings. Consequently, we extend some results of De Concini-Kac-Procesi, whose work focuses on odd orders of ζ , which forces the subalgebra to be strictly central, and complex ground fields. This includes the identification of the subalgebras for Lie types A_n and B_2 with the coordinate rings of associated algebraic groups in the commutative cases, even if ζ has even order. Our descriptions are computationally explicit and do not utilize Poisson structures.

As technical preparations, we discuss PBW bases over minimal rings, dependencies on choices of convex orderings, as well as various new constructions of, and relations among, automorphisms on quantum groups. The latter include formulae for the Garside element in the Lusztig-Artin group action and the family of Che-transformations.

CONTENTS

1. Introduction	3
1.1. Organization and Summary of Main Results	7
Acknowledgments	11
2. Root Systems and Weyl Groups	11
2.1. Root Systems and Weyl Reflections	12
2.2. Coxeter Systems, Reduced Words, and Bruhat Orders	13
2.3. Inversion Root Sets and Convex Orderings	14
2.4. Longest Elements, Involutions, and Complementary Words	18
2.5. Root Lattices and Special Elements	20
2.6. Φ -Symmetric Lattice Maps	21
2.7. Rank 2 Subsystems	23
3. Quantum Groups at Generic Parameter q	25
3.1. Quantization Rings, Quantum Numbers, Integral Subrings	25

3.2. Quantum Group Generators and Relations	27
3.3. Hopf Algebra Structures	28
4. Gradings and Automorphisms on Quantum Groups	29
4.1. Gradings	29
4.2. Che Transformations and Anti-Involutions	31
4.3. Lusztig's Action of Artin-Tits Monoid	33
4.4. More \mathcal{A}^+ Commutation Relations	35
5. Generators, Bases, and Integrality	37
5.1. Generators Associated to Words	38
5.2. Garside Automorphisms and Dictionary between Generators	41
5.3. Monomial Expressions, Exponent Sets, and Their Spans	43
5.4. Some Commutation Relations	47
5.5. Numberings, Boxes, and Refining Convex Orderings	51
5.6. PBW Bases for Box Orderings	53
5.7. Spanning Sets and Their Order Dependence	57
5.8. General Ordering Bases and Module Restrictions	60
6. Skew-Commutative Algebras at Roots of Unity	62
6.1. Quantum Numbers and Rings at Roots of Unity	62
6.2. Skew-Commutation Relations of Primitive Power Generators	66
6.3. Skew-Central Subalgebras and Monomial Bases	71
6.4. \mathfrak{U} -Invariance and Word Independence of \mathcal{Z}_w	75
6.5. \mathcal{Z} -Induced Ideals, \mathcal{G} -Invariance, and \mathcal{G} -Grading	81
7. Hopf Ideals for $U_q(\mathfrak{sl}_{n+1})$ from Solvable Linear Algebraic Groups	85
7.1. Basic Background and Elementary Examples	85
7.2. Standard Presentation of $U_q^{\geq 0}$ for A_n	91
7.3. Identification of $\mathcal{Z}_w^{\geq 0}$ as a (Semi) Classical Matrix Algebra	95
7.4. Ideals from Algebraic Subgroups of T_n and Bruhat Subgroups	99
8. The Structure of \mathcal{Z} and Hopf Ideals for Type B_2	105
8.1. Coalgebra Structure of $\mathcal{Z}_w^{\geq 0}$ for Type B_2	106
8.2. Proof of Proposition 8.1 for Non-simple Roots	108
8.3. The Borel Matrix Algebra on $SO(5, \mathbb{C})$	111
8.4. Isomorphisms between Hopf Algebras related to B_2	116
9. Hopf Ideals via Quasi- R -Matrices	119
9.1. Enhanced Tensor Structures	119
9.2. Completions and Quasi- R -Matrices for U_q	122
9.3. Pre-Triangular Structures for U_q	127
9.4. Maximal Hopf Ideals from Tanisaki-Lusztig Pairings	129
9.5. Truncated Quasi- R -Matrices and Coproduct Formulas	131
9.6. Quasi- R -Matrices for Restricted Quantum Algebras	137
Appendix A. Proof of Lemma 9.16	141
A.1. Case $A_{ij} = 2$	141

A.2. Case $A_{ij} = -1$	141
A.3. Case $A_{ij} = -2$	142
A.4. Case $A_{ij} = -3$	143
Appendix B. Proofs and Computations for Section 8.2	146
B.1. Proof of Proposition 8.3	146
B.2. A Summation Identity for the $c'_{k,p}$	148
Appendix C. Inexistence of Quantum Weyl Elements and Related Notes	149
C.1. An Example in Rank 1	152
Appendix D. Integral Form of Coalgebras	153
References	154

1. INTRODUCTION

One of the original motivations for this monograph was to provide accessible constructions and computational tools for quantum groups at roots of unity. Although initially aimed at practitioners engaged in newer developments in quantum topology, the topics and theorems presented here focus exclusively on the quantum algebra side, introducing several new results and perspectives. Keeping the intended audience in mind, algebraic background material will be selectively introduced.

Broadly speaking, a quantum group U is a Hopf algebra associated to a simple Lie algebra \mathfrak{g} over a ground ring Λ . In the original definitions by Drinfeld and Jimbo [Dri87, Jim85] Λ is naturally given as $\mathbb{Q}(q)$, where q is viewed as a formal “deformation parameter” relating U to the standard universal enveloping algebra of \mathfrak{g} . In many variants of the original definition Λ can be chosen as a subring of $\mathbb{Q}(q)$ or an extension of scalars of such a subring. Analogous to the universal envelope of \mathfrak{g} , any relevant version of U has a decomposition $U = U^+ U^0 U^-$ into subalgebras with generating sets typically of the form $\{E_i\}$, $\{K_i\}$, and $\{F_i\}$, respectively, where the indices enumerate a set of simple roots of \mathfrak{g} .

Quantum groups play a central role in low-dimensional and quantum topology thanks to their associated R -matrices, which give rise to solutions of the Yang-Baxter equation. The latter yield representations of braid groups and invariants of knots, links, and 3-manifolds. Early constructions of 3-manifold invariants and TQFTs, inspired by quantum field theories in physics, start from a Hopf algebra H [Hen96, Ker03], its representation theory [RT91], or, more generally, certain types of tensor categories [Tur16, KL01]. An important requirement for these constructions is that H is effectively finite-dimensional, to assure the existence of integrals, well-defined and finite Kirby colors, or certain coends (which are all essentially equivalent).

The so-called *small quantum groups* provide the quintessential examples of finite dimensional Hopf algebras in our context. Lusztig [Lus90b, Lus93] obtains these by first defining a divided power algebra $\mathcal{L}U$ version of U over $\Lambda = \mathbb{Z}[q, q^{-1}]$, and then extending this ring to $\mathbb{Z}[\zeta]$, where q is specialized to a primitive ℓ -th root of unity ζ . The small quantum group appears then as the subalgebra generated by the original $\{E_i, F_i, K_i\}$ in $\mathcal{L}U$. The construction is, thus, independent of further choices and may also be formulated as the kernel of a Frobenius homomorphism. An important caveat is that $\text{Spec}(U^0)$ is, of course, finite and discrete for these algebras, a feature that is not easily circumvented in this construction due to extra generators required in $\mathcal{L}U^0$.

More recently, invariants constructed from quantum groups with *continuous* $\text{Spec}(U^0)$ have gained considerable interest as they establish connections to classical topological invariants. Typically, parameters associated with $\text{Spec}(U^0)$ are identified with formal variables that appear in algebraic topology constructions. For generic q , one obtains R -matrix representations of braid groups for which the continuous weights of the K_i correspond to deck transformation parameters of homological representations [JK11, Ito15, Ang22, Mar22].

At roots of unity U admits finite-dimensional representations and a rigidity structure, while retaining continuity of $\text{Spec}(U^0)$. Invariants of knots and links are obtained by extending to modified quantum traces on the braid representations derived from R -matrices. In this setting of non-semisimple U and modified traces, the continuous parameters supplied by $\text{Spec}(U^0)$ often imply that the quantum invariants subsume classical invariants as special cases. Among the first such examples is the Alexander Polynomial, for which highest weights yield the indeterminate variable [Mur92, Mur93, Oht02]. For the more involved Akutsu-Deguchi-Ohtsuki (ADO) invariant [ADO92], the continuous weights have similar homological interpretations [Ito16, Ang24].

All the examples mentioned thus far assume the simplest Lie type $\mathfrak{g} = \mathfrak{sl}_2$. A higher rank example, leading to non-commutative properties of the knot invariant, has been studied in [Har]. In [CGPM14, BCGPM16] quantum groups at roots of unity (in rank one) are used to define infinitely graded categories from which 3-manifold invariants and TQFTs are constructed. These, in turn, extend both the Reshetikhin-Turaev (RT) invariants and the classical Turaev-Reidemeister torsion. The invariant depends on a 1-cohomology class $\omega \in H^1(M, \mathbb{C}/\mathbb{Z})$ whose coefficients stem again from the continuous spectrum of K . Specifically, the torsion map assigns the meridian homology generators of a surgery link to ζ^λ , where ζ is a fourth root of unity and λ a complex weight. Constructions for more general categories, such as those in [DR22], are, thus, likely candidates to admit similar connections to more involved classical geometric invariants.

The *restricted quantum groups* underlying these invariants may be characterized by the conditions that U^- and U^+ are finite-dimensional, that $U^0 = \Lambda[\{K_i^{\pm 1}\}]$, and that there is at least a quasi- R -matrix. Their construction typically starts from the specialization of the full (unrestricted) quantum to a root of unity ζ , sometimes also called the De Concini-Kac-Procesi (DCKP) algebra over $\mathbb{Q}(\zeta)$. The allowed restrictions to subrings Λ in $\mathbb{Q}(\zeta)$ heavily depend on renormalizations of generators and the condition that the algebra is stable under the Artin group action defined by Lusztig.

The restricted quantum group, for some given \mathfrak{g} and ζ , is then commonly written as the quotient of the DCKP algebra by an ideal generated by a collection of elements $X_\alpha = E_\alpha^{\ell_\alpha}$ and $Y_\alpha = F_\alpha^{\ell_\alpha}$, which we refer to as *primitive power generators*. Here α ranges over the set of positive roots Φ^+ and E_α is defined via the Artin group action, depending on the choice of a convex ordering of Φ^+ . The exponent ℓ_α is the order of $\zeta^{(\alpha|\alpha)}$, thus depending on the length of α .

Although it is clear that this definition yields continuous $\text{Spec}(U^0)$ as desired, it does not immediately imply that the constructed quotient is indeed a Hopf algebra with finite-dimensional U^\pm . To illustrate the involved subtleties, assume that \mathfrak{g} is of type B, C, or F and the order \mathbb{k} of ζ is a multiple of 4. In this case, there exists *no* (uniform) ℓ for which $\{E_\alpha^\ell, F_\alpha^\ell\}$ generates a Hopf ideal. Hence, the quotient fails to be a Hopf algebra for any ℓ .

A basic computation shows that the $X_i = X_{\alpha_i}$ are *skew-primitive* for simple roots α_i , meaning $\Delta(X_i) = X_i \otimes L_i + 1 \otimes X_i$ with $L_i = K_i^{\ell_i}$ group-like. One easily infers from this that the set $\{X_i, Y_i : i = 1, \dots, n\}$ indeed generates a Hopf ideal. However, the quotient algebra by this ideal yields *infinite* dimensional U^\pm in ranks greater than 1. Finally, for non-simple roots $\alpha \in \Phi^+ \setminus \Delta$, the X_α are far from skew-primitive. Consequently, the ideal generated by an arbitrary subset of these elements is generally also *not* a Hopf ideal.

Computationally explicit and self-contained proofs that the X_α and Y_α generate Hopf ideals for all Lie types and roots of unity tend to be difficult to locate in the existing literature. For the A_n Lie type one such treatment can be found in [BW04], relying on recursions that are considerably more complicated for other Lie types. For ζ of odd order \mathbb{k} , respective statements can be inferred indirectly from the differential methods developed in [DCKP92]. Our initial aim at the onset of this project has been the clarification of these points, which has led us to the exploration of numerous other directly related questions.

One of them is, which subsets of the X_α generate Hopf ideals. The answer we give here is in terms of the inversion sets $\mathcal{N}(s) = \Phi^+ \cap s(\Phi^-)$ (and unions thereof), where $s \in \mathcal{W}$ is an element of the Weyl group. The associated ideals $\hat{\mathcal{K}}(s)$ also occur naturally in the iterative construction of quasi- R -matrices, in the sense that intertwining relations for partial R -matrices hold on the (partial) quotients by the $\hat{\mathcal{K}}(s)$. Like the generators themselves, the $\hat{\mathcal{K}}(s)$ ideal *a priori* depends on the choice of a total convex ordering or, equivalently, a reduced word presentation of s . We show independence of the latter so that, in particular, the fully restricted quantum groups are also independent of these choices and admit unique quasi- R -matrices.

Another source of classical parameter spaces inherent to quantum groups at roots of unity and, thus, potential starting points for quantum invariants extending classical invariants, are Lie groups associated to central subalgebras. Assign to a connected, reductive algebraic group G a solvable group $G^\# < G \times G$ of the same dimension by choosing Borel subgroups $B^+ < G$ and $B^- < G$ for which $H = B^+ \cap B^-$ is a maximal torus in G and denote

$$G^\# = \ker(m \circ (\pi^- \times \pi^+) : B^- \times B^+ \rightarrow H) = \{(g, h) \in B^- \times B^+ : \pi^-(g) \cdot \pi^+(h) = 1\} , \quad (1.1)$$

where $\pi^\pm : B^\pm \rightarrow H$ are the usual abelianization maps. This group naturally appears in various contexts, such as the study of flag varieties, compactifications of algebraic groups, and Poisson Lie groups.

The algebra \mathcal{Z} generated by the X_α , Y_α , and L_i is in the center of the DCKP algebra for *odd* orders \mathbb{k} and is complementary to the generic Harish-Chandra center. In [DCKP92] De Concini, Kac, and Procesi establish an isomorphism of Poisson groups between $\text{Spec}(\mathcal{Z} \otimes \mathbb{C})$ and $G^\#$, or, equivalently, a Poisson Hopf isomorphism between $\mathcal{Z} \otimes \mathbb{C}$ and the coordinate ring of $G^\#$. The construction relies on differential Poisson structures that reduce the proofs to the comparison of skew-primitive elements for simple roots and, therefore, side-steps explicit assignments of the X_α generators for non-simple roots and comparisons of those coproducts.

For even \mathbb{k} , the algebra \mathcal{Z} is generally only a *skew-commutative* (commutative up to sign or unit factor) Hopf algebra. In [Tan16] Tanisaki defines the Frobenius center as the intersection of \mathcal{Z} with the center of U , which is generally no longer a Hopf subalgebra. He shows it is isomorphic to the coordinate ring of the quotient $G^\#/\mathcal{G}$ as an algebraic variety for a finite group \mathcal{G} , utilizing similar techniques as in [DCKP92].

In this paper, we provide Hopf algebra isomorphisms from $\mathcal{Z} \otimes \mathbb{C}$ to $\mathbb{C}[G^\#]$ with *explicit* assignments of *all* X_α generators for Lie type A_n with $\mathbb{k} \not\equiv 2 \pmod{4}$ and type B_2 for any \mathbb{k} . Our computations directly compare the coalgebra structures of \mathcal{Z} with those of $\mathbb{C}[\text{GL}(n, \mathbb{C})^\#]$ and $\mathbb{C}[\text{SO}(5, \mathbb{C})^\#]$, and do not rely on Poisson structures. The concrete forms of these identifications make them more suitable for applications in quantum topology.

The congruence restriction above for A_n is imposed solely to assure that \mathcal{Z} is commutative. For $\mathbb{k} \equiv 2 \pmod{4}$, the full \mathcal{Z} algebra is isomorphic to a non-commutative Artin-Schelter-Tate central extension [AST91] of classical matrix algebras. For this case, we suggest modifications of the Hopf algebra structures on $\mathbb{C}[G^\#]$ via \mathcal{G} weights. We provide a concrete description of \mathcal{G} and the isomorphism of $\mathbb{C}[G^\#/\mathcal{G}]$ with the Frobenius center, as proven abstractly in [Tan16].

Our calculations for the B_2 algebra illustrate further that, for \mathbb{k} a multiple of four, the group $G^\#$ needs to be replaced by $\check{G}^\#$. Here, the Langlands dual group \check{G} is the algebraic group associated to the coroot system $\check{\Phi}$, reflecting the fact that long and short roots are exchanged for the primitive power generators X_α . In both examples, we also show the ideals $\mathcal{K}(s)$ in \mathcal{Z} encountered above are isomorphically mapped to the vanishing ideals on the Bruhat subgroups. The latter are defined here as

$$B(s) = B \cap \dot{s}B\dot{s}^{-1}, \quad (1.2)$$

where \dot{s} is a lift from the geometric Weyl group to G .

An additional important feature of many quantum invariants is that they are valued in integral polynomial rings, since one may expect integer coefficients to relate to topological information. For example, RT-type invariants of homology spheres are valued in $\mathbb{Z}[\zeta]$ with coefficients giving rise to so-called finite type invariants of homology 3-spheres [Oht95, Oht96, Oht02],

which are, in turn, often related to geometric invariants. In a more direct approach, the Hennings invariant of a 3-manifold is valued in the ground ring of the Hopf algebra chosen for the construction, assuming adequate properties of (co)integrals [Hen96, GHPM22, GHPM23].

Among the different versions of quantum groups, we consider here the original Jimbo normalization with generators $\{E_i, F_i, K_i\}$ as well as elements obtained from these via the Artin group action. We write $U_{q,\mathbb{Q}}$ for the rational form of the quantum group with these generators. It is easy to see that the minimal ground ring for the Hopf subalgebra $U^{\geq 0}$ generated by $\{E_i, K_i\}$ must contain $\mathbb{Z}[q, q^{-1}, [\mathbf{e}]^{-1}]$. Here $[n]$ indicates the usual quantum number and $\mathbf{e} = 1$ for Lie types ADE, $\mathbf{e} = 2$ for types BCF, and $\mathbf{e} = 3$ for type G.

We prove that a PBW theorem holds for this minimal ring with respect to appropriate monomial bases. For the full quantum group with the described standard generators $\{E_i, F_i, K_i\}$ the ring over which the algebra is defined also includes $(q - q^{-1})^{-1}$. Minimal ground rings for quantum algebras with differently normalized generators are discussed as well.

Throughout this project, (anti)automorphisms of quantum groups are an important tool for generating and relating monomial basis elements. In addition to the Artin group action, symmetries are generated by numerous (anti)involutions as well as an infinite family of so-called *Che transformations* Ψ_u^h . The latter act on generators via multiplication with Cartan generators and scalars and appear in numerous calculations, particularly, relations and expressions involving the antipode.

We carefully organize the catalog of relations between all types of automorphisms, allowing us to derive explicit expressions for other automorphisms, most notably, the action of the Garside element with respect to the Artin group action. The study of these automorphisms serves to provide insights into symmetries of quantum invariants.

1.1. Organization and Summary of Main Results. Section 2 introduces basic background on spherical Artin and Coxeter groups as well as their actions on roots systems. In Sections 2.3, 2.4, and 2.5 we present numerous properties of inversion sets $\mathcal{N}(s)$ and discuss relations between reduced word expressions, weak Bruhat orders, as well as convex orderings of roots. We introduce root sums for inversion sets and a relation $v \rightsquigarrow w$ for reduced words, which is later used to describe involutions of basis elements. The notion of symmetric root lattice maps, discussed in Section 2.6, prepares the definition of Che transformations. We further compile formulae for rank 2 root systems in Section 2.7, as they will be frequently used in later proofs.

Section 3 covers the basic definitions of the Drinfeld-Jimbo quantum groups, which will be the focus of the paper as the most commonly used version. The section includes discussions of their various minimal ground rings, extensions of scalars, basic Hopf algebra structures, and pointedness. Specifically, the subalgebras $U_q^{\geq 0}$ generated by E_i and K_i are defined over Λ_{q,\mathbf{e}^*} , which is the extension of $\mathbb{Z}[q, q^{-1}]$ by $([\mathbf{e}]!)^{-1}$, where $[n]!$ is the usual quantum factorial and

$e \in \{1, 2, 3\}$ depends on the Lie type as above. The full algebra U_q is defined over Λ_{q,e^*} , which is obtained by further extending Λ_{q,e^*} by $(q - q^{-1})^{-1}$.

In addition to the standard integer root lattice grading of quantum groups, we introduce in Section 4.1 an independent $\mathbb{F}_2 = \mathbb{Z}/2\mathbb{Z}$ root lattice grading, which is useful for the derivation and structure analysis of the skew-commutative subalgebras at roots of unity. We assign to each Φ -symmetric lattice map h and homomorphism $u : \mathbb{Z}^\Delta \rightarrow \Lambda^*$ a Che automorphism Ψ_u^h in Section 4.2. We also introduce in that subsection (anti)involutions, such as Ω , \mathfrak{U} , Υ , Π , and Π , and list numerous relations between them, including a formula for the antipode.

Lusztig's action of the Artin group \mathcal{A} on U and its relations with the other automorphisms is explored in Section 4.3. Our convention differs slightly as we use instead the automorphisms $\Gamma_b = T_{\iota(b)}$, where $b \in \mathcal{A}$ and ι the inversion automorphism on \mathcal{A} .

In Section 5 we construct families of bases for U , starting in Section 5.1 with the definition of elements E_w and F_w assigned to any reduced word w of the underlying Weyl group. There we show that $E_w = E_k$ if the standard degree of E_w is a simple root α_k , extending the proof in [Lus90a] to all Lie types. As an application, we derive in Section 5.2 the formula

$$\Gamma_\partial = \mathcal{I} \circ \mathcal{P} \circ \mathfrak{U} \circ S$$

for the Lusztig-Artin action Γ_∂ of the Garside element $\partial \in \mathcal{A}$ in terms of the automorphisms from Section 4 as well as the involution \mathcal{I} given by symmetries of the Dynkin diagram. In the remainder of the subsection, we provide a dictionary between generators in Lusztig's and our convention in the rank 2 case via the \mathfrak{U} -involution. Aside from facilitating the translation of results and formulae, it serves to show that later definitions of ideals and quotients are independent of which of the two \mathcal{A} -actions is chosen.

Section 5.3 develops a formalism for PBW-style monomial elements $E_{w\bullet}^\psi$ and $E_{\bullet w}^\psi$, depending on reduced words w as well as an exponent functions $\psi : \mathcal{N}(w) \rightarrow \mathbb{N}_0$. We derive explicit formulae for the actions of automorphisms from Section 4 and introduce notations for spanning sets and spans. Various commutation relations between these monomials are gathered in Section 5.4 for later use.

In Sections 5.5 and 5.6 we adapt Lusztig's proof of a PBW basis for the divided power algebras from [Lus90a, Lus90b] to produce a PBW theorem for U_q^+ and U_q over respective minimal extensions of $\mathbb{Z}[q, q^{-1}]$ for one particular convex ordering. In Section 5.7 we state criteria for subsets of exponent functions from which one can deduce that their respective spans are independent of the choice of a reduced word expression for a given Weyl element. The results are combined in Section 5.8 to yield Proposition 5.17 and Theorem 5.18. They are restated in less formal terms as follows.

Theorem A. *Assume \mathfrak{g} to be of any simple Lie type and let z be any reduced word of maximal length. Then $\{E_{z\bullet}^\psi\}$ and $\{E_{z\bullet}^\psi \cdot K^\mu\}$ are Λ_{q,e^*} -bases of U_q^+ and $U_q^{\geq 0}$, respectively.*

Moreover, the set $\{E_{z\blacktriangleright}^\psi \cdot F_{z\blacktriangleright}^{\psi'} \cdot K^\mu\}$ is a $\Lambda_{q,\mathbf{e}^*}^*$ -basis of the Hopf subalgebra $U_q \subset U_{q,\mathbb{Q}}$.

The exponents are integer functions $\psi, \psi' \in \mathbb{N}_0^{\Phi^+}$ and $\mu \in \mathbb{Z}^\Delta$ on the positive and simple roots, respectively. Analogous statements hold for bases obtained from the monomials $E_{z\blacktriangleleft}^\psi$ defined with respect to the reverse direction of multiplication. We discuss a few less commonly used normalizations of generators that yield smaller ground rings.

In Section 6 we turn to the root of unity case, beginning in Section 6.1 with the introduction of various rings related to $\mathbb{Z}[\zeta] = \Lambda_\zeta$, notation for various orders, and formulae for multinomial coefficients. In Section 6.2 we derive skew-commutation relations involving the primitive power generators $X_w = E_w^{\ell_w}$, $Y_w = F_w^{\ell_w}$, and $L_i = K_i^{\ell_i}$ in terms of special functions $(\Phi^+)^2 \rightarrow \{\pm 1\}$. The discussion is extended in Section 6.3 to monomials, where we also introduce unital algebras such as \mathcal{Z}_w^+ , $\mathcal{Z}_w^{\geq 0}$, and \mathcal{Z}_w , generated over $\Lambda_{\zeta,\mathbf{e}^*}$ or $\Lambda_{\zeta,\mathbf{e}^*}^*$ by X_u , Y_u , and L^μ with $u \leq_R w$. We conclude the section with a full classification of Lie types and \mathbb{A} congruences for which these algebras are central as subalgebras of U_ζ and for which they are commutative. For $\square \in \{+, -, 0, \geq 0, \leq 0\}$ and a given reduced word w , they give rise to the ideal $\mathcal{K}(w)^\square$ in a maximal \mathcal{Z}_w^\square generated by the augmentation ideal A_w^+ of \mathcal{Z}_w^+ .

For each rank 2 type A_2 , B_2 , G_2 there are two possible convex orderings, each of which is represented by a maximal reduced word. In Section 6.4 we compute explicit formulae that express the X_w from one ordering in terms of those of the other ordering, with the exception of words of length 3 and 4 in type G_2 . Drawing on the results in previous section, the implied closure under the \mathfrak{G} -action entails that the algebras above do not depend on the word w representing an element s of the Weyl group, but only on s itself. Summarizing Theorem 6.19, Corollary 6.21, Theorem 6.22, and Corollary 6.26 this observation results in the following.

Theorem B. *Assume a Lie type different from G_2 and ζ of any order $\mathbb{A} \notin \{1, 2, \mathbf{e}, 2\mathbf{e}\}$. Suppose w is a reduced word and s the Weyl group element it represents. For $\square \in \{+, -, 0, \geq 0, \leq 0\}$ we have the following.*

- i) *The algebra \mathcal{Z}_w^\square depends only on s .*
- ii) *The maximal skew-commutative algebra \mathcal{Z}_w^\square is defined independently of the choice of ordering.*
- iii) *\mathcal{Z}_w is invariant under the Artin group action.*
- iv) *The augmentation ideal $\mathcal{K}(w)^\square$ in \mathcal{Z}_w^\square is only dependent on s .*

In Section 6.5 we consider the action of an elementary abelian 2-group \mathcal{G} on \mathcal{Z}_w and induce for any \mathcal{G} -stable ideal J in \mathcal{Z}_w a two-sided ideal \hat{J} in U_ζ . We introduce notations for special and extended cases of the resulting $\hat{\mathcal{K}}(s)$ and discuss various examples.

Sections 7 and 8 are devoted to the identifications of the \mathcal{Z}_w algebras with coordinate rings of algebraic groups in the basic A_n type as well as the doubly laced B_2 case. Section 7.1 collects background on algebraic groups and their coordinate rings, with emphasis on the correspondence between subgroups and Hopf ideals, and a discussion of relevant elementary examples.

Generators, commutation relations, and coproducts are computed for the quantum algebras for $\mathfrak{g} = \mathfrak{gl}_n$ and $\mathfrak{g} = \mathfrak{sl}_n$ in Section 7.2. Specializing to a root of unity of order ℓ , we identify in Section 7.3 the respective $\mathcal{Z}_\bullet^{\geq 0}$ algebra with the AST algebra over Λ_ζ [AST91]. For $\ell \not\equiv 2 \pmod{4}$ the extension is trivial, resulting in a Hopf algebra isomorphism to the coordinate ring of the group of upper triangular matrices over $\mathbb{k} = \mathbb{Q}(\zeta)$ or $\mathbb{k} = \mathbb{C}$. In Section 7.4 the full \mathcal{Z}_\bullet algebra for \mathfrak{gl}_n over \mathbb{C} is identified with the coordinate ring of $\mathrm{GL}(n, \mathbb{C})^\#$ and the augmentation ideals $\mathcal{K}(s)$ with vanishing ideals of Bruhat subgroups $B(s)$ defined in *loc. cit.* The section concludes with a discussion of potential interpretations of the skew-commutative case.

We start our treatment of the B_2 case with a complete description of the Hopf algebra structure of $\mathcal{Z}_\bullet^{\geq 0}$ in Section 8.1, deferring the detailed computations of the coproducts to Section 8.2 and Appendix B. The required replacement of the root system by the coroot system, as mentioned above, is achieved by the choice of an isomorphism between the B_2 and C_2 algebras.

In Section 8.3 we use a specific complex parametrization of $\mathrm{SO}(5, \mathbb{C})^{\geq 0}$ to derive formulae for the coalgebra structure of its coordinate ring. Using its Lie-Kolchin form, the vanishing ideals of the Bruhat subgroups are readily identified. Comparing the coalgebra structures obtained in Section 8.2 and 8.3, we obtain Hopf algebra isomorphisms between the central subalgebra and the coordinate ring in Section 8.4. The involved Hopf algebras fit into parametrized families of abstract weighted Hopf algebras, which are classified in the same sections.

The theorem below summarizes the main results from Lemma 7.18, Theorem 7.16, and Theorem 8.10. The field \mathbb{k} can be chosen as either $\mathbb{Q}(\zeta)$ or \mathbb{C} . The notation $G^\#$, B , and $B(s)$ for an algebraic group G is as in (1.1) and (1.2). The isomorphisms are given as explicit assignments of polynomial generators.

Theorem C. *Suppose \mathcal{Z}_\bullet is the full commutative subalgebra for $\mathfrak{g} = \mathfrak{gl}_n$ with $\ell \not\equiv 2 \pmod{4}$ or \mathfrak{g} of type B_2 for any ℓ . Let $G = \mathrm{GL}(n, \mathbb{k})$ or $G = \mathrm{SO}(5, \mathbb{k})$ respectively. Then*

- i) *There is a Hopf algebra isomorphism from $\mathcal{Z}_\bullet \otimes \mathbb{k}$ to $\mathbb{k}[G^\#]$, which restricts to an isomorphism between $\mathcal{Z}_\bullet^{\geq 0} \otimes \mathbb{k}$ and $\mathbb{k}[B]$.*
- ii) *The isomorphism maps the augmentation ideal $\mathcal{K}(s) \subseteq \mathcal{Z}_\bullet^{\geq 0}$ exactly to the vanishing ideal in $\mathbb{k}[B]$ of the Bruhat subgroup $B(s) < B$.*

Section 9 is mainly concerned with the construction of R -matrices and quasi-triangular structures. We review relevant algebra completions as well as Tanisaki's axioms for pre-triangular bialgebras in Section 9.1. The algebraic properties of elementary and partial quasi- R -matrices for generic q are discussed in Sections 9.2 and 9.3. We consider their relations with the Artin group action, introduce the Taniski automorphism Π for the pre-triangular structure, and define coefficients related to the Taniski-Lusztig pairings. Specialized at a root of unity, the latter is used in Section 9.4 to define the full quotient into the restricted quantum group, where maximal ideals are identified with null spaces.

The technical core result of Section 9.5 is a set of intertwining relations via truncated partial R -matrices between the coproduct $\Delta(x)$ and a twisted coproduct $\Delta^s(x)$, modulo ideals derived from the ideals $\widehat{\mathcal{K}}(w)$ introduced in Section 6. Here $\Delta^s(x)$ is the coproduct conjugated by Γ_s and the underlying quantum group $U_{\zeta,\ell}$ is now over the ring $\Lambda_{\zeta,\ell}^*$, which also includes $([\ell-1]!)^{-1}$. The rather involved computations for generators are deferred to Appendix A. An important consequence of these relations is that the two-sided ideals $\widehat{\mathcal{K}}(w)$ depend only on the Weyl elements represented by w and are Hopf for all cases, including G_2 . The following combines the statements in Theorem 9.20 and Proposition 9.21.

Theorem D. *Let ζ be of any order $\ell \notin \{1, 2, e, 2e\}$ and $U_{\zeta,\ell}$ of any simple Lie type. Suppose w is a reduced word representing a Weyl element s . Let $\widehat{\mathcal{K}}(w, w)$ be the two-sided ideal generated by all X_u and Y_u with $u \leq_R w$. Then*

- i) $\widehat{\mathcal{K}}(w, w)$ is a Hopf ideal depending only on s so that the quotient $U_{\zeta,\ell}^{[s]}$ by this ideal is a well-defined Hopf algebra over $\Lambda_{\zeta,\ell}^*$.
- ii) PBW bases of $U_{\zeta,\ell}^{[s]}$ are given by the set of exponents with $\psi(\alpha) < \ell_\alpha$ for all $\alpha \in \mathcal{N}(s)$.
- iii) For any reduced word u representing some $t \leq_R s$ there is a partial quasi- R -matrix $\mathcal{P}_{\blacktriangleleft u}^\star$ such that

$$\Delta^{[t]}(x) \cdot \mathcal{P}_{\blacktriangleleft u}^\star = \mathcal{P}_{\blacktriangleleft u}^\star \cdot \Delta(x),$$

where $x \in U_{\zeta,\ell}^{[s]}$ and $\Delta^{[t]}$ is the coproduct on $U_{\zeta,\ell}^{[s]}$ obtained by conjugation with Γ_t .

In Section 9.6 we consider the specialization to the maximal length element s_\bullet in the Weyl group, defining the restricted quantum group as $U_{\zeta,\ell}^{\text{rest}} = U_{\zeta,\ell}^{[s_\bullet]}$. Using our formula for the Garside element, we identify $\Delta^{[s_\bullet]}$ with the Tanisaki-Lusztig conjugate $\bar{\Delta}$ and show the maximal $\mathcal{P}_{\blacktriangleleft z}^\star$ is independent of the chosen word. A slightly less formal statement of Theorem 9.25 is as follows, which now encompasses also the G_2 case.

Theorem E. *For any Lie type and root of unity as in Theorem D, the triple $(U_{\zeta,\ell}^{\text{rest}}, \mathcal{P}^\star, \Pi)$ is a pre-triangular Hopf algebra over $\Lambda_{\zeta,\ell}^*$. Its construction is independent of any choices, it has the properties of a restricted quantum group, and it admits the actions of the Artin group and all involutions.*

Acknowledgments. MH was partially supported through the NSF-RTG grant #DMS-2135960.

2. ROOT SYSTEMS AND WEYL GROUPS

This section combines a general review of basic notions of root systems, Coxeter groups, and Artin groups with a discussion of several more specialized lemmas and concepts. The latter serve as technical tools in later sections, but are typically not treated in standard textbooks on Coxeter groups. Examples of such topics are various properties of inversion sets, complementary words, as well as Φ -symmetric lattice maps.

2.1. Root Systems and Weyl Reflections. Let Φ be an irreducible root system of rank n in some Euclidean space E . For a choice of simple roots $\Delta = \{\alpha_1, \dots, \alpha_n\}$ denote by Φ^+ and Φ^- the sets of positive and negative roots, with $\Phi = \Phi^+ \cup \Phi^-$. Although it is not dependent on the choice of Δ , we will frequently denote the root lattice of Φ in E by \mathbb{Z}^Δ .

The Weyl group \mathcal{W} is, by definition, generated by the reflections s_α of E along roots $\alpha \in \Phi$, which are expressed as $s_\alpha(x) = x - \langle \alpha, x \rangle \alpha$. The form $\langle \cdot, \cdot \rangle$ is linear in the second entry and gives rise to the Cartan matrix A associated to Φ via $A_{ij} = \langle \alpha_i, \alpha_j \rangle$. The convention for A chosen here conforms with that in [DCK90, Lus90b] but is transposed to that of textbooks such as [Bou02]. Clearly, \mathcal{W} also acts on \mathbb{Z}^Δ as an invariant subspace of E .

For a given Cartan matrix, we may find integers $d_i \in \{1, 2, 3\}$ that symmetrize the Cartan matrix in the sense that $C_{ij} = d_i A_{ij}$ is symmetric. Accordingly, we set $d_\alpha = d_i$ if α has the same length as α_i or, equivalently, if α is in the \mathcal{W} -orbit of α_i . We will use the following convention and notation,

$$\min_i \{d_i\} = 1 \quad \text{and} \quad e = \max_i \{d_i\} \in \{1, 2, 3\}. \quad (2.1)$$

Thus, in the simply laced cases ADE we have $e = 1$, meaning all $d_i = 1$. We have $e = 2$ for doubly laced Lie types BCF and $e = 3$ for type G. The renormalization leads to the definition of an inner form

$$(\cdot | \cdot) : \mathbb{Z}^\Delta \times \mathbb{Z}^\Delta \rightarrow \mathbb{Z} \quad (2.2)$$

by setting $(\alpha_i | \alpha_j) = C_{ij} = d_i A_{ij}$ for the basis roots vectors. It is readily verified that the Weyl group \mathcal{W} acts by isometries with respect to this form, meaning $(s(\alpha) | s(\beta)) = (\alpha | \beta)$ for any $s \in \mathcal{W}$ and $\alpha, \beta \in \mathbb{Z}^\Delta \supset \Phi$. So $(\cdot | \cdot)$ is a scalar multiple of the standard Euclidean inner product on E for an irreducible root system Φ . Specifically, it coincides with the canonical inner product for the realizations in the appendix of [Bou02] when the Lie type is one of ACDEG, and it is twice the canonical inner product given there if the Lie type is B or F. The integer from (2.1) may thus also be understood as the lower bound on inner products of positive roots,

$$e = \max_{i,j} \{-C_{ij}\} = \max_{i,j} \{-(\alpha_i | \alpha_j)\} = \max \{-(\alpha | \beta) : \alpha, \beta \in \Phi^+\}. \quad (2.3)$$

In these conventions we also find $(\alpha | \alpha) = 2d_\alpha$ by choosing some $s \in \mathcal{W}$ and $\alpha_i \in \Delta$ with $\alpha = s(\alpha_i)$ and, hence, $d_i = d_\alpha$. As usual, for a fixed root system, we will call α a *short* root if $d_\alpha = 1$ and, in the non-simply laced case, we say α is a *long* root if $d_\alpha = e > 1$. The initial integral pairing is recovered from $(\cdot | \cdot)$ via

$$\langle \cdot, \cdot \rangle : \Phi \times \Phi \rightarrow \mathbb{Z} \quad \text{with} \quad \langle \alpha, \beta \rangle = \frac{1}{d_\alpha} (\alpha | \beta) \in \mathbb{Z}. \quad (2.4)$$

For later notational convenience we also introduce

$$e^* = e + 1 \in \{2, 3, 4\}. \quad (2.5)$$

That is, $e^* = 2$ for Lie types ADE, $e^* = 3$ for the BCF types, and $e^* = 4$ for G_2 .

2.2. Coxeter Systems, Reduced Words, and Bruhat Orders. Recall that $(\mathcal{W}, \mathcal{S})$ forms a Coxeter system, where the set of generators $\mathcal{S} = \{s_1, \dots, s_n\}$ consists of the simple reflections $s_i = s_{a_i}$. The relations for a presentation of \mathcal{W} in generators \mathcal{S} are given by the involution $s_i^2 = 1$ and the Artin-Tits relations. They are given for each pair (i, j) with $i \neq j$ by

$$s_i s_j \dots = s_j s_i \dots \quad (2.6)$$

with m_{ij} reflections terms on each side with alternating indices. Here $m_{ij} = 2, 3, 4$, or 6 depending on whether the edge number $\max(|A_{ij}|, |A_{ji}|)$ is $0, 1, 2$, or 3 , respectively.

As usual, the length $l(s)$ of an element $s \in \mathcal{W}$ is the smallest number k such that $s = s_{i_1} \dots s_{i_k}$ can be expressed as a product of k simple reflections. A *reduced word* is a formal word expression $w_{i_1} \dots w_{i_k}$ in the letters w_1, \dots, w_n such that the respective product $s = s_{i_1} \dots s_{i_k}$ in \mathcal{W} is of minimal length $k = l(s)$. We use the separate notation \mathcal{W}^* for the set of reduced words, since several objects in the article will explicitly depend on words rather than just elements in \mathcal{W} .

An obvious surjection $r : \mathcal{W}^* \rightarrow \mathcal{W}$ is obtained by replacing each letter w_i by the respective s_i and mapping the empty word \emptyset to the identity element $1 \in \mathcal{W}$. We say $s \in \mathcal{W}$ is represented by a word $w \in \mathcal{W}^*$ if w maps to s so that, by definition, $l(s)$ is the number of letters in w . Denote by \mathcal{A} the *Artin group* associated to the same Coxeter data as \mathcal{W} . That is, \mathcal{A} shares the same generator set \mathcal{S} and the relations in (2.6) but lacks the relations $s_i^2 = 1$. Also denote by \mathcal{A}^+ the *Artin monoid*, defined as the subset of elements in \mathcal{A} that are a product of generators with positive powers.

Matsumoto's Theorem states that there is a well-defined map $s : \mathcal{W} \rightarrow \mathcal{A}^+ \subset \mathcal{A}$ such that $s \circ r : \mathcal{W}^* \rightarrow \mathcal{A}^+$ is given as the assignment of letters w_i to the respective Artin generators (see Part IV, §1.5 in [Bou02] or [Mat64]). The more practical formulation is as follows.

Corollary 2.1 (Matsumoto). *Suppose two words in \mathcal{W}^* represent the same element in \mathcal{W} . Then they are related by a sequence of Artin relations given in (2.6).*

Given a reduced word $w = w_{i_1} \dots w_{i_k} \in \mathcal{W}^*$, we write $w[a, b] = w_{i_a} \dots w_{i_b}$ for the subword between indices a and b with $1 \leq a \leq b \leq k$. Further, denote $w^* = w_{i_k} \dots w_{i_1}$ the word with letters multiplied in the opposite direction so that $r(w^*) = r(w)^{-1}$.

The *weak Bruhat orders* on \mathcal{W} and \mathcal{W}^* are defined in terms of smaller elements occurring as subwords. For our purposes, the most relevant is the weak right Bruhat order \leq_R . For two words $w, u \in \mathcal{W}^*$ we say $u \leq_R w$ if there exists $v \in \mathcal{W}^*$ such that $w = u \cdot v$ or, equivalently, if $u = w[1, j]$ for some j . Similarly, for two elements $s, t \in \mathcal{W}$ we have $s \leq_R t$ if t and s are represented by words w and u , respectively, with $u \leq_R w$.

To define the left weak Bruhat order, replace $u = w[j, k]$ in the above definition. If $s_\bullet \in \mathcal{W}$ is the unique longest element, then s_\bullet is also the unique maximal element for any of the Bruhat orders, including \leq_R (e.g., Proposition 3.1.2 in [BB05]).

Correspondingly, we write $\mathcal{W}_m^* = \tau^{-1}(s_\bullet)$ for the set of reduced words of maximal length. The set will be used to bijectively label all convex orders on Φ^+ . See, for example, [Sta84] for formulae for the size and rapid growth of $|\mathcal{W}_m^*|$.

Note that any reduced word $u \in \mathcal{W}^*$ can be written as the initial subword of some word in \mathcal{W}_m^* . Explicitly, let $s = \tau(u)$, set $s' = s^{-1}s_\bullet$, and choose any $u' \in \mathcal{W}^*$ representing s' . Since $s \leq_R s_\bullet$ we have $l(u') = l(s') = l(s_\bullet) - l(s) = l(s_\bullet) - l(u)$ by the same proposition in [BB05]. Now, $w = u \cdot u'$ represents s_\bullet and, since $l(u) + l(u') = l(s_\bullet)$, it is a reduced expression so that $w \in \mathcal{W}_m^*$.

Similarly, the element $\partial = \tau(s_\bullet) \in \mathcal{A}^+$ plays an important role as the fundamental element that renders $(\mathcal{A}^+, \partial)$ a Garside monoid (see, for example, Proposition 1.29 in [DDG⁺15]). In particular, Lemma 5.2 in [BS72] asserts that for all $b \in \mathcal{A}^+$ the commutation relation

$$\partial \cdot b = \sigma(b) \cdot \partial \quad (2.7)$$

holds, where σ is an involutive automorphism of \mathcal{A}^+ . Since σ needs to map generators to themselves, we may rewrite (2.7) as the relation $\partial \cdot w_i \cdot \partial^{-1} = w_{\sigma(i)}$ in \mathcal{A} , with σ redefined as an involution on $\{1, \dots, n\}$.

For example, for type A_n the Artin-Tits monoid is the set of positive braids B_n^+ in the braid group in n strands, and ∂ is the full positive half-twist. In this situation, relation (2.7) is readily visualized by moving a single positive neighbor crossing through this twist resulting in a crossing in reflected position, implying that $\sigma(w_i) = w_{n+1-i}$. For other types, the involution coincides with the respective ones in the Coxeter or Weyl group situation as in Section 2.4.

Note that relation (2.7) can also be derived directly from the presentations of s_\bullet for other Lie types given in Section 5.5 below, essentially following the methods for relative Garside elements developed in [ABI15].

2.3. Inversion Root Sets and Convex Orderings. In numerous calculations with words, it will be convenient to formally denote the index of the last letter as well as the word with the last letter deleted. That is, for a non-empty ($k \geq 1$) word

$$w = w_{i_1} \dots w_{i_k} \in \mathcal{W}^* \quad \text{set} \quad w^\flat = w_{i_1} \dots w_{i_{k-1}} \quad \text{and} \quad \tau(w) = i_k. \quad (2.8)$$

These components naturally enter the statement of the following lemma, which asserts an assignment of a positive root to any reduced word.

Lemma 2.2. *Suppose $w = w_{i_1} \dots w_{i_k} \in \mathcal{W}^*$ is a non-empty reduced word. Then we have*

$$\gamma(w) = w^\flat(\alpha_{\tau(w)}) = w_{i_1} \dots w_{i_{k-1}}(\alpha_{i_k}) \in \Phi^+,$$

where w acts on roots by its respective element in \mathcal{W} .

To see this, note that $s(\beta) \in \Phi^+$ if and only if $l(s \cdot s_\beta) > l(s)$ by Proposition 4.4.6 in [BB05], where s_β denotes the reflection along $\beta \in \Phi^+$. The lemma follows by specializing to $s = \tau(w^\flat)$

and $\beta = \alpha_{i_k}$. Next, define for any element $s \in \mathcal{W}$ the *inversion set* of s as

$$\mathcal{N}(s) = \{\alpha \in \Phi^+ : s^{-1}(\alpha) \in \Phi^-\} = \Phi^+ \cap s(\Phi^-). \quad (2.9)$$

For a type A_{n-1} root system realized as $\Phi^+ = \{\alpha_{i,j} = \epsilon_i - \epsilon_j : 1 \leq i < j \leq n\} \subset \mathbb{R}^n$ with $\mathcal{W} = S_n$, the symmetric group in n letters, $\mathcal{N}(s)$ is indeed given by the literal inversion set of the permutation s^{-1} . Proposition 3.6 in [Hil82] or Corollary 2 in Chapter VI §1.6 of [Bou02] show that for any reduced word $w \in \mathcal{W}^*$ as in (2.8) with $s = r(w) \in \mathcal{W}$ this set can be enumerated as

$$\mathcal{N}(s) = \{\beta_m = \gamma(w[1, m]) : 1 \leq m \leq k\} \quad \text{where } \beta_m = s_{i_1} \dots s_{i_{m-1}}(\alpha_{i_m}) \quad (2.10)$$

and $k = l(w) = l(s)$. Thus, any word representing s imposes a total ordering on $\mathcal{N}(s)$ given by $\beta_p \leq_w \beta_j$ if $p \leq j$, writing also $\beta_p <_w \beta_j$ if $p < j$. Clearly, $\mathcal{N}(1) = \emptyset$, $\mathcal{N}(s_i) = \{\alpha_i\}$, and $\mathcal{N}(w) = \{\gamma(u) : u \leq_R w\}$ if the notation is extended to words. Additional immediate consequences of the above are

$$l(s) = |\mathcal{N}(s)|, \quad s^{-1}(\mathcal{N}(s)) = -\mathcal{N}(s^{-1}), \quad \text{and} \quad \mathcal{N}(s) = \Phi^+. \quad (2.11)$$

We next explain how these sets determine the defect in length additivity for composites of Weyl group elements. In the first observation, we denote for an arbitrary subset $A \subseteq \Phi$ two subsets of Φ^+ as

$$A^+ = A \cap \Phi^+ \quad \text{and} \quad A^- = -(A \cap \Phi^-) = (-A) \cap \Phi^+,$$

so that $A = A^+ \sqcup -A^-$. Also write $B \setminus A$ for the complement of A in B .

Lemma 2.3. *Suppose $s = a \cdot b$ for $a, b \in \mathcal{W}$. Then $\mathcal{N}(a) \cap a(\mathcal{N}(b)) = \emptyset$ and*

$$\mathcal{N}(s) = a(\mathcal{N}(b))^+ \sqcup \mathcal{N}(a) \setminus a(\mathcal{N}(b))^- . \quad (2.12)$$

Moreover,

$$l(s) = l(a) + l(b) - 2\zeta(a, b) \quad \text{where} \quad \zeta(a, b) = |a(\mathcal{N}(b))^-| = |\mathcal{N}(b) \cap \mathcal{N}(a^{-1})|. \quad (2.13)$$

Proof. The disjointness statement is clear since $\mathcal{N}(b) \subseteq \Phi^+$ and $a^{-1}(\mathcal{N}(a)) \subseteq \Phi^-$. For (2.12) we may split $\mathcal{N}(s)$ according to the partition $\Phi = a(\Phi^+) \sqcup a(\Phi^-)$. Then $a(\Phi^+) \cap \mathcal{N}(s) = a(\Phi^+ \cap a^{-1}(\mathcal{N}(s))) = a(\Phi^+ \cap a^{-1}(\Phi^+) \cap b(\Phi^-)) = a(a^{-1}(\Phi^+) \cap \mathcal{N}(b)) = \Phi^+ \cap a(\mathcal{N}(b)) = a(\mathcal{N}(b))^+$.

Now, from $\Phi^+ = b(\Phi^+) \cap \Phi^+ \sqcup b(\Phi^-) \cap \Phi^+$ we see that $b(\Phi^+) \cap \Phi^+ = \Phi^+ \setminus \mathcal{N}(b)$ and, hence, $b(\Phi^-) \cap \Phi^+ = \Phi^- \setminus (-\mathcal{N}(b))$. Applying a to this we get $s(\Phi^-) \cap a(\Phi^-) = a(\Phi^-) \setminus (-a(\mathcal{N}(b)))$, and intersecting this with Φ^+ we find $\mathcal{N}(s) \cap a(\Phi^-) = \mathcal{N}(a) \setminus ((-a(\mathcal{N}(b))) \cap \Phi^+) = \mathcal{N}(a) \setminus a(\mathcal{N}(b))^-$.

Note that $-a(\mathcal{N}(b))^- = a(\mathcal{N}(b)) \cap \Phi^- = a(\mathcal{N}(b) \cap a^{-1}(\Phi^-)) = a(\mathcal{N}(b) \cap \mathcal{N}(a^{-1}))$ as $\mathcal{N}(b) \subseteq \Phi^+$. This readily implies the equality of the two expressions for $\zeta(a, b)$. Clearly, $|a(\mathcal{N}(b))^+| = |a(\mathcal{N}(b))| - |a(\mathcal{N}(b))^-| = |\mathcal{N}(b)| - \zeta(a, b) = l(b) - \zeta(a, b)$. Now, $a^{-1}(\mathcal{N}(b))^- = (-\mathcal{N}(b)) \cap a^{-1}(\Phi^+) = -(\mathcal{N}(b) \cap a^{-1}(\Phi^-)) = -(\mathcal{N}(b) \cap \mathcal{N}(a^{-1}))$. Thus, using also (2.11), $|\mathcal{N}(a) \setminus a(\mathcal{N}(b))^-| = |a^{-1}(\mathcal{N}(a)) \setminus a^{-1}(a(\mathcal{N}(b))^-)| = |(-\mathcal{N}(a^{-1})) \setminus (-(\mathcal{N}(b) \cap \mathcal{N}(a^{-1}))| = |\mathcal{N}(a^{-1})| - |\mathcal{N}(b) \cap \mathcal{N}(a^{-1})| = l(a^{-1}) - \zeta(a, b)$. Using $l(a^{-1}) = l(a)$ and adding these expressions yields the length defect equation in (2.13). \square

The following criteria for exact additivity and monotonicity are now easily derived from Lemma 2.3. For example, if $\mathcal{N}(a) \subseteq \mathcal{N}(s)$ we have $\zeta(a^{-1}, s) = |\mathcal{N}(s) \cap \mathcal{N}(a)| = |\mathcal{N}(a)| = l(a)$. So, for $b = a^{-1}s$ (2.13) implies $l(b) = l(a^{-1}) + l(s) - 2l(a) = l(s) - l(a)$ or $l(s) = l(a) + l(b)$. The latter and $s = a \cdot b$ now imply $a \leq_R s$ as claimed in (2.14).

Corollary 2.4. *For two elements $a, b \in \mathcal{W}$ the following conditions are equivalent.*

i) $l(a \cdot b) = l(a) + l(b)$	iii) $a(\mathcal{N}(b)) \subseteq \Phi^+$
ii) $\mathcal{N}(a \cdot b) = a(\mathcal{N}(b)) \sqcup \mathcal{N}(a)$	iv) $\mathcal{N}(b) \cap \mathcal{N}(a^{-1}) = \emptyset$

Moreover, we have for elements $a, b, s \in \mathcal{W}$ that

$$a \leq_R s \Leftrightarrow \mathcal{N}(a) \subseteq \mathcal{N}(s) \quad \text{and} \quad b \leq_L s \Leftrightarrow \mathcal{N}(b^{-1}) \subseteq \mathcal{N}(s^{-1}). \quad (2.14)$$

In particular, $\mathcal{N}(s) = \mathcal{N}(a)$ implies $s = a$.

The fact that the longest element of \mathcal{W} yields all of Φ^+ implies that any reduced word of maximal length $w \in \mathcal{W}_m^*$ imposes a total order \prec_w on the set of positive roots via $\Phi^+ = \mathcal{N}(w)$. Specifically, if we denote $\beta_j = \gamma(w[1, j])$ we have $\beta_1 \prec_w \beta_2 \prec_w \dots \prec_w \beta_L$ where $L = l(w)$. We write \leq_w if equality is included.

The result of Papi in [Pap94] states that the assignment $w \mapsto \leq_w$ establishes a bijection between elements in \mathcal{W}_m^* and the (total) *convex* orders on Φ^+ . Here convexity means that if for two roots $\alpha, \beta \in \Phi^+$ we have $\alpha \prec_w \beta$ and $\alpha + \beta \in \Phi^+$, then $\alpha \prec_w \alpha + \beta \prec_w \beta$.

Let $s = r(w)$ and observe that, since the β_j as above are distinct and $L = |\mathcal{N}(s)|$, we have that the map $[1, L] \rightarrow \mathcal{N}(s) : j \mapsto \beta_j$ is a bijection. Thus, to each $\alpha \in \mathcal{N}(s)$ we can assign a unique j_α such that $\alpha = \beta_{j_\alpha}$. It is convenient to also introduce the inverse map from the root set to words,

$$\mathcal{N}(s) \rightarrow \{u \in \mathcal{W}^* : u \leq_R w\} : \alpha \mapsto w[\alpha], \quad (2.15)$$

where we denote $w[\alpha] = w[1, j_\alpha]$ so that $\alpha = r(w[\alpha])$. In particular, if $w \in \mathcal{W}_m^*$ the correspondence (2.15) assigns to any positive root a reduced word $\leq_R w$.

For non-simply laced root systems (types BCFG) with $e > 1$ it is useful to distinguish the short and long roots portions of the various root sets. That is, we have $\Phi = \Phi_L \sqcup \Phi_S$ where $\Phi_L = \{\alpha \in \Phi : d_\alpha = e\}$ and $\Phi_S = \{\alpha \in \Phi : d_\alpha = 1\}$. Correspondingly, denote

$$\begin{aligned} \Phi_L^+ &= \Phi^+ \cap \Phi_L, & \Phi_L^- &= \Phi^- \cap \Phi_L, & \mathcal{N}(s)_L &= \mathcal{N}(s) \cap \Phi_L^+, \\ \Phi_S^+ &= \Phi^+ \cap \Phi_S, & \Phi_S^- &= \Phi^- \cap \Phi_S, & \mathcal{N}(s)_S &= \mathcal{N}(s) \cap \Phi_S^+. \end{aligned} \quad (2.16)$$

Since $s(\Phi_L) = \Phi_L$ and $s(\Phi_S) = \Phi_S$ for any $s \in \mathcal{W}$, we also have $\mathcal{N}(s)_L = s(\Phi_L^-) \cap \Phi_L^+$ and $\mathcal{N}(s)_S = s(\Phi_S^-) \cap \Phi_S^+$. Thus, for example, the relations between root sets in Lemma 2.3 will hold for the respective long root and short root portions separately. Furthermore, introduce the following subsets of the Weyl group.

$$\mathcal{W}_S = \{s \in \mathcal{W} : \mathcal{N}(s)_S = \emptyset\} \quad \text{and} \quad \mathcal{W}_L = \{s \in \mathcal{W} : \mathcal{N}(s)_L = \emptyset\} \quad (2.17)$$

Lemma 2.5. Suppose Φ is a non-simply laced root system with $e > 1$. Then \mathcal{W}_S and \mathcal{W}_L are the subgroups of \mathcal{W} generated by $\{s_i : d_i = e\}$ and $\{s_i : d_i = 1\}$ respectively.

Thus, for type B_n , we have that $\mathcal{W}_S \cong S_{n-1}$ is the canonical A_{n-1} -subgroup and $\mathcal{W}_S \cong \mathbb{F}_2$ the A_1 -subgroup. The assignments are reversed for C_n . For F_4 they are the respective A_2 -subgroups and for G_2 the respective A_1 -subgroups.

Proof. \mathcal{W}_S is readily identified with the stabilizer $\{s : s(\Phi_S^+) = \Phi_S^+\}$ and thus a subgroup of \mathcal{W} . If $d_i = e$ we also have $\mathcal{N}(s_i) = \{\alpha_i\}$ so that $\mathcal{N}(s_i)_S = \emptyset$ and, hence, $s_i \in \mathcal{W}_S$, implying that \mathcal{W}_S contains the subgroup generated by $\{s_i : d_i = e\}$. For the converse inclusion, assume $s = r(w) \in \mathcal{W}_S$ is given by a reduced expression $w = w_{i_1} \dots w_{i_k}$. Then $\beta_m = \gamma(w[1, m]) = t(\alpha_{i_m})$ where $t = r(w[1, m-1])$ so that β_m has the same length as α_{i_m} . Hence, $\mathcal{N}(s)_S = \emptyset$ if and only if $d_{i_1} = \dots = d_{i_k} = e$, implying that s is in the subgroup generated by s_i with $d_i = e$. The arguments for \mathcal{W}_L are analogous. \square

We will need the following observation about root decompositions later only in the A case. The elementary argument, though, extends to all classical types. The exceptional types are not discussed here.

Lemma 2.6. Let Φ be a root system of classical type ABCD. Assume that Φ^+ is equipped with any convex ordering \prec_w . Suppose $\beta = \beta_1 + \dots + \beta_k$ with $\beta, \beta_i \in \Phi^+$ for $i = 1, \dots, k$ with $k > 1$. Then

- i) There exists j for which $\beta - \beta_j \in \Phi^+$.
- ii) There exists r for which $\beta_r \prec_w \beta$.

Proof. Consider the same numbering of simple roots as in [Bou02, Plate I-IV] so that α_1 is opposite the special roots. The formulae in [Bou02] imply that every positive root is of the form $\beta_{a,b} = \sum_{s=a}^b \alpha_{c(s)}$, where $c : \{1, \dots, N\} \rightarrow \{1, \dots, n\}$ is a numbering sequence depending on Lie type, $n = |\Delta|$, and $1 \leq a \leq b \leq N$.

For $\beta = \sum_i c_i \alpha_i$ set $\text{ht}(\beta) = \sum_i c_i$ and $r(\beta) = \min\{i : c_i \neq 0\}$. The formulae $\text{ht}(\beta_{a,b}) = b - a + 1$ and $r(\beta_{a,b}) = \min\{c(a), c(b)\}$ are readily checked for all cases below. We use this notation to prove the statement *i*) for each Lie type separately.

For type A_n set $N = n$ and $c(i) = i$. Then any positive root is of the form $\beta = \beta_{a,b}$. Now, for β as in the lemma, at least one β_j must have non-zero coefficient for α_a and no β_i can have α_s with $s < a$ as a summand. Thus, we must have $r(\beta_j) = a$ for at least one j , which implies $\beta_j = \beta_{a,c}$ for some $c \leq b$ and, with $k > 1$, also $c < b$. Then $\beta - \beta_j = \beta_{a,b} - \beta_{a,c} = \beta_{c+1,b} \in \Phi^+$.

Next consider type C_n , setting $N = 2n - 1$ and $(c(1), \dots, c(N)) = (1, \dots, n-1, n, n-1, \dots, 1)$. It follows by inspection that all positive roots are of the form $\beta_{a,b}$ and all $\beta_{a,b} \in \Phi^+$. In fact, $(a, b) \mapsto \beta_{a,b}$ defines a bijection between Φ^+ and the set of pairs (a, b) for which $a \leq \min(n, b)$ and $c(a) \leq c(b)$. Let $\beta = \beta_{a,b}$ with these constraints so that $a = r(\beta)$. As before, we can pick β_j such

that $a = r(\beta_j)$ and, hence, $\beta_j = \beta_{a,c}$ for some $c < b$. Then, $\beta - \beta_j = \beta_{c+1,b}$ which is again in Φ^+ by previous remarks.

For B_n we set $N = 2n$ and $(c(1), \dots, c(N)) = (1, \dots, n-1, n, n, n-1, \dots, 1)$. As opposed to previous Lie types, not all $\beta_{a,b}$ are roots. Specifically, $\beta_{a,b} \notin \Phi^+$ if and only if $a \neq b$ but $c(a) = c(b)$. This case has to be excluded to establish a one-to-one correspondence, as in the C_n case. We may thus assume $\beta = \beta_{a,b}$ with $a = r(\beta) \leq n$ and $c(a) < c(b)$ since $k > 1$ implies $a \neq b$. Then, as before, we have $\beta_j = \beta_{a,c}$ for some $c < b$ so that $\gamma = \beta - \beta_j = \beta_{c+1,b}$. If either $c+1 = b$ or $c(c+1) \neq c(b)$ we are done since $\beta_{c+1,b} \in \Phi^+$. In case $c+1 < b$ and $c(c+1) = c(b)$ we have $\gamma = 2 \sum_{c < i \leq n} \alpha_i = \sum_{t \neq j} \beta_t$ so that there has to be an $s \neq j$ with $\beta_s = \beta_{c+1,d}$ with $d < b$. With $c(N+1-i) = c(i)$, $b = N - c$, and $d' = N - d$ we have $\beta_s = \beta_{d'+1,b}$. Now, $\beta - \beta_s = \beta_{a,d'}$ which is in Φ^+ since $c(a) < c(d')$.

For type D_n we may choose $N = 2n-2$ and $(c(1), \dots, c(N)) = (1, \dots, n-2, n-1, n, n-2, \dots, 1)$. A second sequence c' may be defined via $c'(i) = c(N+1-i) = \eta(c(i))$, where $\eta = (n-1, n)$ is the non-trivial involution of the Dynkin diagram. All positive roots are of the form $\beta_{a,b}$ or $\beta'_{a,b}$ defined via the respective sequences and $\beta_{c(a),c(b)} = \beta'_{a,b}$ if $\{a, b\} \cap \{n-1, n\} = \emptyset$. As for B_n , $\beta_{a,b} \notin \Phi^+$ if and only if $a \neq b$ and $c(a) = c(b)$. The argument now proceeds as in the D_n case, except that a subsequence may be chosen from either c or c' .

Item *ii*) follows now by induction in k . That is, if $\beta = \beta_1 + \dots + \beta_k$ and j such that $\beta' = \beta - \beta_j = \sum_{i \neq j} \beta_i \in \Phi^+$, convexity implies that either $\beta_j \prec_w \beta \prec_w \beta'$ or $\beta' \prec_w \beta \prec_w \beta_j$. In the former case we are done. In the latter, the induction hypothesis implies $\beta_i \prec_w \beta'$ for some i ($i \neq j$) so that, by transitivity, also $\beta_i \prec_w \beta$. \square

2.4. Longest Elements, Involutions, and Complementary Words. The longest element $s_\bullet \in \mathcal{W}$ is an involution by uniqueness. Its respective involutive action on the root system maps Φ^+ isometrically to Φ^- (e.g., Corollary 3 in [Bou02, VI.1.6]) and is thus of the form $s_\bullet(\alpha_i) = -\alpha_{\eta(i)}$ for an involution η of the Dynkin diagram associated to Φ .

As specified in Item XI of each plate in the appendix of [Bou02], η is the non-trivial involution for A_n , D_{2k+1} , and E_6 , and it is identity in all other cases. Mapping the \mathcal{A} -version of (2.7) to \mathcal{W} then implies that $\sigma = \eta$. We use the same notation η for the conjugation action of s_\bullet on \mathcal{W} . Since s_i is given by reflection along α_i we, thus, find

$$\eta(s_i) = s_{\eta(i)} \quad \text{where} \quad \eta(t) = s_\bullet t s_\bullet^{-1} = s_\bullet t s_\bullet. \quad (2.18)$$

We use the same notation for the obvious extension of η to the set of reduced words \mathcal{W}^* . We will frequently use the involution

$$w^\dagger = \eta(w)^* = \eta(w^*) \quad (2.19)$$

on \mathcal{W}^* , explicitly given as $(w_{i_1} \dots w_{i_N})^\dagger = w_{\eta(i_N)} \dots w_{\eta(i_1)}$ for reduced expressions. Clearly, $l(w^\dagger) = l(w)$ and $s^\dagger = \gamma(w^\dagger) = \eta(s)^{-1}$. A related involutive map on \mathcal{W} is given by right multiplication with the longest element by

$$t \mapsto t^\perp = t \cdot s_\bullet. \quad (2.20)$$

Lemma 2.7. *Suppose $t \in \mathcal{W}$ and t^\perp is as in (2.20). Then*

$i) \ t = t^{\perp\perp},$ $ii) \ s_\bullet \cdot t = \eta(t^\perp) = \eta(t)^\perp,$ $iii) \ l(t^\perp) = l(s_\bullet) - l(t),$	$iv) \ t \leq_R u \Leftrightarrow u^\perp \leq_R t^\perp,$ $v) \ \mathcal{N}(t^\perp) = \Phi^+ \setminus \mathcal{N}(t) = \Phi^+ \cap t(\Phi^+),$ $vi) \ t^\perp (\mathcal{N}(t^\dagger)) = \mathcal{N}(t).$
--	--

Proof. Items *i*, *ii*, and *iii* are immediate from the definitions. See also Proposition 2.3.2 in [BB05]. For Item *iv*) assume $t \leq_R u$, meaning $u = t \cdot a$ with $l(u) = l(t) + l(a)$. Using (2.18) this yields the decomposition $t^\perp = u^\perp \cdot \eta(a)^{-1}$. Further, using *iii*), $l(t^\perp) - l(s_\bullet) - l(t) = l(u^\perp) + l(a) = l(u^\perp) + l(\eta(a)^{-1})$. So, $u^\perp \leq_R t^\perp$. For Item *v*) note that $\mathcal{N}(t^\perp) = \Phi^+ \cap t^\perp(\Phi^-) = \Phi^+ \cap t(s_\bullet(\Phi^-)) = \Phi^+ \cap t(\Phi^+) = \Phi^+ \setminus \mathcal{N}(t)$. Finally, Item *vi*) follows from $t^\perp (\mathcal{N}(t^\dagger)) = ts_\bullet(\Phi^+ \cap t^\dagger(\Phi^-)) = t(s_\bullet(\Phi^+) \cap s_\bullet t^\dagger(\Phi^-)) = t(\Phi^- \cap t^{-1}s_\bullet(\Phi^-)) = t(\Phi^- \cap t^{-1}(\Phi^+)) = t(\Phi^-) \cap \Phi^+ = \mathcal{N}(t)$. \square

For the description of involutive automorphisms on quantum groups, the related notion of *complementary words* will be relevant. Given a non-empty reduced word $w \in \mathcal{W}^*$, we say that $v \in \mathcal{W}^*$ is complementary to w , writing $v \circ\circ w$, if the following two conditions hold.

$$v \circ\circ w \Leftrightarrow \tau(v) = \eta(\tau(w)) \quad \text{and} \quad \tau(v)^\perp = \tau(w^\dagger). \quad (2.21)$$

If we write $s = \tau(w^\dagger)$, $t = \tau(v^\dagger)$, $j = \tau(w)$, and $k = \tau(v)$ the condition in (2.21) can be rephrased more symmetrically as $k = \eta(j)$ and

$$t^{-1} \cdot s = q_j \quad \text{where} \quad q_j = s_\bullet \cdot s_j = s_{\eta(j)} \cdot s_\bullet. \quad (2.22)$$

Note, with $N = l(s_\bullet)$, these imply $l(w) + l(v) = N + 1$ or $l(t) + l(s) = N - 1$.

Lemma 2.8. *Suppose $v \circ\circ w$ for non-empty, reduced words $v, w \in \mathcal{W}^*$. Then*

$i) \ w \circ\circ v,$ $ii) \ \gamma(v) = \gamma(w),$	$iii) \ \mathcal{N}(w) \cap \mathcal{N}(v) = \{\gamma(w)\},$ $iv) \ \Phi^+ = \mathcal{N}(w) \cup \mathcal{N}(v).$
--	--

Moreover, if $z \in \mathcal{W}_m^*$ is a reduced word of maximal length $N = l(z)$ and $y = z^\dagger$, then

$$z[1, r] \circ\circ z[r, N]^\dagger = y[1, N + 1 - r]. \quad (2.23)$$

Proof. Symmetry in *i*) is immediate from (2.22) together with $q_j^{-1} = q_{\eta(j)}$ and $\eta^2 = \text{id}$. Note further that $q_j(\alpha_j) = s_\bullet(s_j(\alpha_j)) = s_\bullet(-\alpha_j) = \alpha_{\eta(j)}$. So, with notation as in (2.22), $\gamma(w) = s(\alpha_j) = t(q_j(\alpha_j)) = t(\alpha_{\eta(j)}) = \gamma(v)$. For *iii*) and *iv*) note that $\mathcal{N}(w) = \mathcal{N}(w^\dagger) \sqcup \{\gamma(w)\}$ by Corollary 2.4, and $\mathcal{N}(w^\dagger) = \Phi^+ \setminus \mathcal{N}(v)$ by (2.21) and Lemma 2.7 *v*).

For the last statement, let $z = w_{i_1} \dots w_{i_N}$ so that $y = w_{\eta(i_N)} \dots w_{\eta(1)}$. Denote further $w = z[1, r] = w_{i_1} \dots w_{i_r}$ and $v = z[r, N]^\dagger = \eta(w_{i_r} \dots w_{i_N})^* = w_{\eta(i_N)} \dots w_{\eta(i_r)} = y[1, N+1-r]$. Clearly, $\eta(\tau(w)) = \eta(i_r) = \tau(v)$, and $z = z[1, r-1]z[r, N] = w^\flat \eta(v^*)$. The latter implies $s_* = \tau(w^\flat) \tau(\eta(v)^*) = \tau(w^\flat) \tau(\eta(v))^{-1}$ and hence $\tau(w^\flat) = s_* \cdot \eta(\tau(v)) = \tau(v) \cdot s_* = \tau(v)^\perp$ as desired. \square

Combining Item *ii*) and (2.23) the following order reversal relation is an immediate consequence for the total ordering \prec_z on Φ^+ depending on a maximal length reduced word $z \in \mathcal{W}_m^*$ as introduced in Section 2.3.

Corollary 2.9. *Suppose $z \in \mathcal{W}_m^*$ and $y = z^\dagger$. Then $\alpha \prec_z \beta$ if and only if $\beta \prec_y \alpha$.*

Relation (2.23) provides explicit constructions of complementary words that imply a partitioning of Φ^+ into two totally ordered sets, which have only their maximal elements in common. That is, for a given $w \in \mathcal{W}^*$ find $z \in \mathcal{W}_m^*$ with $z \geq_R w$ and hence $w = z[1, r]$. Set then $v = y[N+1-r]$ with $y = z^\dagger$. Setting $\beta_i = \gamma(z[1, i])$ and $\mu_i = \gamma(y[1, i])$, the sets $\mathcal{N}(w)$ and $\mathcal{N}(v)$ are then given by the elements in respective maximal chains $\beta_1 \prec_z \beta_2 \prec_z \dots \prec_z \beta_r$ and $\mu_1 \prec_y \mu_2 \prec_y \dots \prec_y \mu_{N+1-r}$, with $\beta_r = \mu_{N+1-r}$ but $\beta_i \neq \mu_j$ otherwise.

For example, consider the reduced word $w = w_2 w_3 w_2 w_1$ in a type A_3 Coxeter system. A maximal length reduced word $z \in \mathcal{W}_m^*$ with $w \leq_R z$ is then given by $z = w_2 w_3 w_2 w_1 w_2 w_3$ so that $w = z[1, 4]$. With $\eta(1) = 3$ we find $y = \eta(z)^* = w_1 w_2 w_3 w_2 w_1 w_2$, which implies that $v = y[1, 3] = w_1 w_2 w_3$ is a complementary word $w \rightsquigarrow v$. Then the ordered elements of $\mathcal{N}(w)$ are $\alpha_2 \prec_z \alpha_2 + \alpha_3 \prec_z \alpha_3 \prec_z \alpha_1 + \alpha_2 + \alpha_3$ and those of $\mathcal{N}(v)$ are $\alpha_1 \prec_y \alpha_1 + \alpha_2 \prec_y \alpha_1 + \alpha_2 + \alpha_3$. Complementary words can also be constructed directly from the conditions in (2.22) by choosing a presentation for $t = \tau(w^\flat) q_j^{-1}$ and adjoining $w_{\tau(w)}$.

In the rank 2 case, complementary words are unique. A complete discussion of this situation is provided in Section 2.7 below.

2.5. Root Lattices and Special Elements. Let $J(s) \subseteq \{1, \dots, n\}$ be the set of indices $J(s) = \{i_1, \dots, i_M\}$ occurring in a reduced expression $s = s_{i_1} \dots s_{i_M}$. It is clear by either Matsumoto's theorem or properties of parabolic subgroups in Section 2.4 of [BB05] (or the lemma below) that $J(s)$ does not depend on the choice of this presentation.

Denote also by $\Xi(s) = \langle \mathcal{N}(s) \rangle_{\mathbb{Z}} \subset \mathbb{Z}^\Delta$ the sublattice of the root lattice generated by roots in the inversion set $\mathcal{N}(s)$. It admits the following simple description in terms of the set $J(s)$.

Lemma 2.10. *The subset $\{\alpha_i \in \Delta : i \in J(s)\}$ is a basis for $\Xi(s)$.*

Proof. Proceeding by induction in $l(s)$, we note that for $s = s_j$ we have $\Xi(s) = \langle \mathcal{N}(s_j) \rangle_{\mathbb{Z}} = \langle \{\alpha_j\} \rangle_{\mathbb{Z}} = \mathbb{Z} \alpha_j$ as desired. Suppose now $t = s_k \cdot s$ with $l(t) = l(s) + 1$ and assume the claim holds for s . By Corollary 2.4 we then have $\mathcal{N}(t) = s_k(\mathcal{N}(s)) \sqcup \{s_k\}$ and, hence, $\Xi(t) = s_k(\Xi(s)) + \mathbb{Z} \alpha_k$. By induction hypothesis $\Xi(s)$ is spanned by all α_i with $i \in J(s)$ so that the first summand is

spanned by all $s_k(\alpha_i) = \alpha_i - A_{ki}\alpha_k$ with $i \in J(s)$. Hence, $\Xi(t)$ is spanned as a lattice by all α_i with $i \in J(s) \cup \{k\} = J(t)$ as desired. \square

For later use in computations, we introduce the sum of roots $\theta(s)$ and sum of coroots $\check{\theta}(s)$ as elements of lattices in E , namely,

$$\theta(s) = \sum_{\alpha \in \mathcal{N}(s)} \alpha \in \mathbb{Z}^\Delta \quad \text{and} \quad \check{\theta}(s) = \sum_{\alpha \in \mathcal{N}(s)} \frac{1}{d_\alpha} \alpha = \sum_{\alpha \in \mathcal{N}(s)} \check{\alpha} \in \frac{1}{e} \mathbb{Z}^\Delta. \quad (2.24)$$

Here $\check{\alpha} = d_\alpha^{-1}\alpha$ are the usual coroots. Observe also that, by (2.4), the pairings $(\check{\theta}(s)|\beta)$ are integers for any $\beta \in \Phi$ and any $s \in \mathcal{W}$. Corollary 2.4 and \mathcal{W} -invariance of d_α readily imply that for elements $t_1, t_2 \in \mathcal{W}$ and $s = t_1 \cdot t_2$ with $l(s) = l(t_1) + l(t_2)$ we have

$$\theta(s) = \theta(t_1) + t_1(\theta(t_2)) \quad \text{and} \quad \check{\theta}(s) = \check{\theta}(t_1) + t_1(\check{\theta}(t_2)). \quad (2.25)$$

The following mirror relations for both roots sets and root sums are immediate from (2.9) and (2.24).

$$s^{-1}(\theta(s)) = -\theta(s^{-1}) \quad \text{and} \quad s^{-1}(\check{\theta}(s)) = -\check{\theta}(s^{-1}) \quad (2.26)$$

Finally, the vectors $\theta(s)$ and $\check{\theta}(s)$ can be expressed more directly using the usual half sums of positive roots or coroots via the formulae,

$$\begin{aligned} \theta(s) &= \rho - s(\rho) \quad \text{and} & \check{\theta}(s) &= \check{\rho} - s(\check{\rho}), \\ \text{where} \quad \rho &= \frac{1}{2} \sum_{\alpha \in \Phi^+} \alpha = \frac{1}{2} \theta(s_*) & \text{and} \quad \check{\rho} &= \frac{1}{2} \sum_{\alpha \in \Phi^+} \check{\alpha} = \frac{1}{2} \check{\theta}(s_*). \end{aligned} \quad (2.27)$$

The formula for $\theta(s)$ is proved in Proposition 21.15 in [Bum04] and the proof for the one for $\check{\theta}(s)$ is analogous using also invariance $d_{s(\alpha)} = d_\alpha$. Applying a simple reflection to (2.27), a standard argument shows that $(\rho|\alpha_i) = \frac{1}{2}(\alpha_i|\alpha_i) = d_i$ and $(\check{\rho}|\alpha_i) = 1$ for all $\alpha_i \in \Delta$.

The latter allows us to express the height function ht , given as $\text{ht}(\beta) = \sum_i b_i$ for a root vector $\beta = \sum_i b_i \alpha_i$, as $\text{ht}(\beta) = (\check{\rho}|\beta)$. Thus, together with (2.27) we can write

$$(\check{\theta}(s)|\beta) = \text{ht}(\beta) - \text{ht}(s^{-1}(\beta)) \in \mathbb{Z}. \quad (2.28)$$

Finally, we note that ρ is an element of the weight lattice Σ and that $\check{\rho} \in \frac{1}{e}\Sigma$.

2.6. Φ -Symmetric Lattice Maps. In preparation for the construction of a large family of automorphisms on quantum groups in Section 4.2, we generalize some classical notions of root systems to non-canonical integral inner products on E .

We say that a lattice homomorphism $h : \mathbb{Z}^\Delta \rightarrow \mathbb{Z}^\Delta$ is Φ -symmetric if $(h(\mu)|\nu) = (\mu|h(\nu))$ for all $\mu, \nu \in \mathbb{Z}^\Delta$, that is, h is symmetric with respect to an underlying Euclidean space E . Denote also $\mathbb{S}(\Phi)$ the space of such Φ -symmetric maps, which, thus, forms a sublattice of $\text{End}(\mathbb{Z}^\Delta)$ of rank $\binom{n+1}{2}$. Note that the quotient $\text{End}(\mathbb{Z}^\Delta)/\mathbb{S}(\Phi)$ will, generally, contain torsion.

Clearly, orthogonality of the Weyl group action implies that for any $s \in \mathcal{W}$ and Φ -symmetric map h also shs^{-1} is Φ -symmetric. Note that for any $\alpha \in \Phi$, the Weyl reflection $s_\alpha \in \mathbb{S}(\Phi)$ since $s_\alpha = s_\alpha^{-1}$. Hence, any integer combination of such reflections $\sum_\alpha c_\alpha s_\alpha$ is also Φ -symmetric. Generalizing the element ρ from (2.27), we define for a given Φ -symmetric lattice map h the vector ρ_h in E via the condition

$$(\rho_h|\alpha_i) = \frac{1}{2} (\alpha_i|h(\alpha_i)) . \quad (2.29)$$

While the concept of Φ -symmetric maps does not depend on the choice of a simple root system, the definition of ρ_h strongly depends on a choice of Δ . We, further, generalize formula (2.27) for the sum $\theta(s)$ above by setting for any $s \in \mathcal{W}$

$$\theta_h(s) = \rho_{shs^{-1}} - s(\rho_h) . \quad (2.30)$$

Several basic properties of these objects are summarized in the following lemma, where Σ is the weight lattice associated to Φ and $\text{id} : \mathbb{Z}^\Delta \rightarrow \mathbb{Z}^\Delta$ is the identity map, which is obviously Φ -symmetric.

Lemma 2.11. *For any Φ -symmetric lattice maps $h, k \in \mathbb{S}(\Phi)$, any root vector $\beta \in \mathbb{Z}^\Delta$, any Weyl group elements $s, t \in \mathcal{W}$, and any $m \in \mathbb{Z}$ the following hold:*

i) $\rho_{h+k} = \rho_h + \rho_k ,$ ii) $\rho_h \in \frac{1}{2}\Sigma ,$ iii) $\rho_{m \cdot \text{id}} = m \cdot \rho ,$ iv) $(\beta h(\beta)) - 2(\rho_h \beta) \in 2\mathbb{Z}$	v) $\theta_{h+k}(s) = \theta_h(s) + \theta_k(s) ,$ vi) $(\theta_h(s) \beta) \in \mathbb{Z} ,$ vii) $\theta_{m \cdot \text{id}}(s) = m \cdot \theta(s) ,$ viii) $\theta_h(ts) = \theta_{shs^{-1}}(t) + t(\theta_h(s)) .$
---	--

Proof. Items *i*), *iii*), *v*), *vii*), and *viii*) are immediate from the definitions in (2.29) and (2.30) as well as the formulae for $\theta(s)$ and ρ in (2.27). For the statement in *ii*) note that $(\rho_h|\check{\alpha}_i) = d_i^{-1}(\rho_h|\alpha_i) = \frac{1}{2}d_i^{-1}(\alpha_i|h(\alpha_i)) = \frac{1}{2}\sum_j d_i^{-1}(\alpha_i|\alpha_j)h_{ji} = \frac{1}{2}\sum_j A_{ij}h_{ji} \in \frac{1}{2}\mathbb{Z}$, where $\check{\alpha}_i = d_i^{-1}\alpha_i$ is the usual coroot, A_{ij} the Cartan data, and h_{ji} are the matrix coefficients of h in the Δ -basis.

For $\beta = \sum_i b_i \alpha_i$ and using symmetry of $n_{ij} = (\alpha_i|h(\alpha_j))$ the expression in *iv*) can be worked out as $2\sum_{i>j} b_i b_j n_{ij} + \sum_i (b_i^2 - b_i) n_{ii}$, which is clearly in $2\mathbb{Z}$. For Item *vi*), note that both terms $(s^{-1}(\beta)|h(s^{-1}(\beta))) - 2(\rho_h|s^{-1}(\beta)) = (\beta|(shs^{-1})(\beta)) - 2(s(\rho_h)|\beta)$ and $(\beta|(shs^{-1})(\beta)) - 2(\rho_{shs^{-1}}|\beta)$ are in $2\mathbb{Z}$ by *iv*). Thus, also their difference $2(\theta_h(s)|\beta)$ has to be in $2\mathbb{Z}$. \square

For an illustration of these notions, we briefly discuss the rank 2 root systems. We assume $A_{12} = e \in \{1, 2, 3\}$ depending on whether Φ is of type A_2 , B_2 , or G_2 , as well as $A_{21} = -1$. A \mathbb{Z} -basis for $\mathbb{S}(\Phi)$ for any type is given by $\{\text{id}, s_1, s_2\}$ where $s_i = s_{\alpha_i}$ are the usual reflections along the simple roots. Note that $s_{12} = s_{\alpha_1 + \alpha_2} = s_2 s_1 s_2 = (e-2)s_2 - s_1$. So, $\{\text{id}, s_1, s_{12}\}$ would be an alternate basis for $e \in \{1, 3\}$ but not for $e = 2$. It follows from elementary calculations that $\text{End}(\mathbb{Z}^\Delta)/\mathbb{S}(\Phi)$ is $\mathbb{Z} \oplus \mathbb{F}_2$ for $e = 2$ but only \mathbb{Z} for $e \in \{1, 3\}$.

Since ρ_h is additive in h it suffices to compute the vector for the basis elements $\{\text{id}, s_1, s_2\}$, for which we already know that $\rho_{\text{id}} = \rho$. The other two elements are computed as

$$\begin{aligned}\rho_{s_1} &= -\frac{1}{2}(1 + (\mathbf{e} - 1)(2\mathbf{e} - 3))\alpha_1 - \frac{1}{2}(\mathbf{e} - 1)^2\alpha_2, \\ \rho_{s_2} &= -(\mathbf{e} - 1)^2\alpha_1 - \frac{1}{2}(1 + (\mathbf{e} - 1)^2)\alpha_2.\end{aligned}$$

Thus, even in the $\mathbf{e} = 1$ (A_2) case we have $\rho_{s_i} = -\frac{1}{2}\alpha_i \notin \Sigma$, as opposed to ρ . Similarly, for the $\theta_h(s)$ it suffices to consider $h \in \{s_1, s_2\}$. The values $\theta_{s_i}(s_j)$ are explicitly computed below:

$$\begin{aligned}\theta_{s_1}(s_1) &= s_1(\alpha_1) = -\alpha_1, & \theta_{s_1}(s_2) &= s_1(\alpha_2) = \mathbf{e}\alpha_1 + \alpha_2, \\ \theta_{s_2}(s_2) &= s_2(\alpha_2) = -\alpha_2, & \theta_{s_1}(s_2) &= s_2(\alpha_1) + \binom{\mathbf{e}}{2}\alpha_1 = \frac{1}{2}(\mathbf{e}^2 - \mathbf{e} + 2)\alpha_1 + \mathbf{e}\alpha_2.\end{aligned}$$

From these one can obtain $\theta_{s_i}(s)$ using the recursion in Item *viii*) of Lemma 2.11 together with the explicit formulae $s_i s_j s_i = (\mathbf{e} - 2)s_i - s_j$ for $i \neq j$ for the adjoint action of \mathcal{W} on $\langle s_1, s_2 \rangle \subset \mathbb{S}(\Phi)$. In particular, we see that in this example $\theta_h(s) \in \Phi$ for all $s \in \mathcal{W}$ and $h \in \mathbb{S}(\Phi)$. Consider the case $\mathbf{e} = 1$ (A_2), for which $\theta_{s_j}(s_i) = s_i(\alpha_j) = s_j(\alpha_i)$, suppose $h = n_1 s_1 + n_2 s_2$, and set $\beta = n_1 \alpha_1 + n_2 \alpha_2$. The suggested computation then yields that

$$\theta_h(s) = s(\beta) \quad \text{if } l(s) \text{ is odd} \quad \text{and} \quad \theta_h(s) = 0 \quad \text{if } l(s) \text{ is even.}$$

We defer further discussion of Φ -symmetric maps to future work. Open questions include whether one can always find a subset $\mathcal{T} \subset \Phi^+$ such that $\{s_\alpha : \alpha \in \mathcal{T}\}$ is a \mathbb{Z} -basis for $\mathbb{S}(\Phi)$ and how to use this to efficiently compute the ρ_h and $\theta_h(s)$ in generality. Other questions are whether $\theta_h(s) \in \Phi$ also for higher ranks and what the possible torsion factors of $\text{End}(\mathbb{Z}^\Delta)/\mathbb{S}(\Phi)$ are.

2.7. Rank 2 Subsystems. As a simple but useful example, consider the Coxeter subsystem for the generator set $\{s_i, s_j\}$, given two distinct but fixed indices $i, j \in \{1, \dots, n\}$. The respective Weyl subgroup $\mathcal{W}_{(i,j)}$ is a dihedral group of order $2m_{ij} \in \{4, 6, 8, 12\}$. The longest element $s_{|ij|}$ is given by the expressions in (2.6) and, hence, of length $l(s_{|ij|}) = m_{ij}$. The element of maximal length has two word presentations given by

$$\mathcal{W}_{(i,j)}^* = \{z_{|ij|}, z_{|ji|}\} = \tau^{-1}(s_{|ij|}), \quad (2.31)$$

where the words with m_{ij} letters are $z_{|ij|} = w_i w_j \dots$ and $z_{|ji|} = w_j w_i \dots$. Thus, Matsumoto's Theorem says that two reduced $w, w' \in \mathcal{W}^*$ represent the same $s = \tau(w) = \tau(w')$ if w' is obtained from w by successively swapping sub words $z_{|ij|} \rightleftharpoons z_{|ji|}$.

For any element $s \in \mathcal{W}_{(i,j)}$ with $l(s) < m_{ij}$ there is exactly one reduced word in $\tau^{-1}(s) \subset \mathcal{W}_{(i,j)}^*$ representing s . To simplify notation here and in later applications, let $m = m_{ij}$ as well as

$$a_r = z_{|ij|}[1, r] = \begin{cases} (w_i w_j)^{\frac{r}{2}} & r \text{ even} \\ (w_i w_j)^{\frac{r-1}{2}} w_i & r \text{ odd} \end{cases} \quad b_r = z_{|ji|}[1, r] = \begin{cases} (w_j w_i)^{\frac{r}{2}} & r \text{ even} \\ (w_j w_i)^{\frac{r-1}{2}} w_j & r \text{ odd} \end{cases} \quad (2.32)$$

for $0 \leq r \leq m$, so, $a_m = z_{|ij|}$ and $b_m = z_{|ji|}$ and $l(a_r) = l(b_r) = r$. The a_r and b_r provide a distinct enumeration of elements in $\mathcal{W}^*_{(i,j)}$ with the exception of the empty word $a_0 = b_0 = \emptyset$. This organization of words provides a succinct way of expressing the complementary word relations introduced in Section 2.4.

Lemma 2.12. *With notation as above, (non-empty) complementary words are uniquely given for all finite Coxeter types by*

$$a_r \circ \circ b_{m+1-r} .$$

Proof. Note that for $m = 3$ we have $z_{|ij|}^* = z_{|ij|}$ and $\eta(z_{|ij|}) = z_{|ij|}$, whereas for $m \in \{4, 6\}$ we have $z_{|ij|}^* = z_{|ji|}$ and $\eta(z_{|ji|}) = z_{|ji|}$. Thus, $z_{|ij|}^\dagger = \eta(z_{|ij|}^*) = z_{|ij|}$ for all Lie types, which implies the statement by (2.23). Uniqueness is also clear since b_{m+1-r} is the only word matching the length constraint and the constraint on the final letter index. \square

In order to describe the total orders on rank 2 root systems implied by the two words $z_{|ij|}$ and $z_{|ji|}$ we choose α_i to be the shorter and α_j to be the longer root. Thus, in the conventions from Section 2.1, the Cartan data is given as $-A_{ji} = d_i = 1$ and $-A_{ij} = d_j = e \in \{1, 2, 3\}$ depending on Lie type. Following the definitions in Section 2.3, the maximal word $z_{|ij|}$ imposes a total order $\prec_{z_{|ij|}}$ with maximal chains given by the sequence of roots $\beta_r = \gamma(b_r) = \gamma(z_{|ij|}[1, r])$. Their explicit expressions in terms of simple roots α_i and α_j are as follows for Lie types A_2 , B_2 , and G_2 , respectively.

$$\begin{aligned} e = 1 \quad & \alpha_j \prec_{z_{|ji|}} \alpha_i + \alpha_j \prec_{z_{|ji|}} \alpha_i \\ e = 2 \quad & \alpha_j \prec_{z_{|ji|}} \alpha_i + \alpha_j \prec_{z_{|ji|}} 2\alpha_i + \alpha_j \prec_{z_{|ji|}} \alpha_i \\ e = 3 \quad & \alpha_j \prec_{z_{|ji|}} \alpha_i + \alpha_j \prec_{z_{|ji|}} 3\alpha_i + 2\alpha_j \prec_{z_{|ji|}} 2\alpha_i + \alpha_j \prec_{z_{|ji|}} 3\alpha_i + \alpha_j \prec_{z_{|ji|}} \alpha_i \end{aligned} \tag{2.33}$$

By Corollary 2.9 and the fact that $z_{|ij|} = z_{|ji|}^\dagger$ the total $\prec_{z_{|ij|}}$ order imposed by the other maximal reduced word $z_{|ij|}$ is exactly the opposite one.

Often questions for general Coxeter systems can be reduced to ones of rank 2 situations. This is illustrated in the following method of extracting from some $s \in \mathcal{W}$ a maximal element $r \leq_L s$ that belongs to such a subsystem. The proof here extends the argument used in the proof of Proposition 1.8 of [Lus90a] to also the non-simply laced Lie types.

Lemma 2.13. *Suppose $s \in \mathcal{W}$ and $1 \leq i \leq n$ are such that $l(s \cdot s_i) = l(s) + 1 \geq 2$. Then there exists $j \neq i, t \in \mathcal{W}$ and $y \in \mathcal{W}^*_{(i,j)}$, such that for $r = r(y)$ we have*

$$s = t \cdot r, \quad l(s) = l(t) + l(r), \quad 1 \leq l(r) < m, \quad t(\alpha_i), t(\alpha_j) \in \Phi^+, \quad \text{and} \quad \tau(y) = j.$$

Moreover, if $s(\alpha_i) = \alpha_k \in \Delta$, then $r \cdot s_i = s_{|ij|}$ so that $r(\alpha_i) \in \{\alpha_i, \alpha_j\}$ and, hence, either $t(\alpha_i) = \alpha_k$ or $t(\alpha_j) = \alpha_k$.

Proof. If $u \in \mathcal{W}^*$ with $r(u) = s$ we have by the first assumption that $u \cdot w_i$ is reduced as well. Thus, by Lemma 2.2, we have $\gamma(u \cdot w_i) = s(\alpha_i) \in \Phi^+$. Now let $j = \tau(u)$ so that $s = s' \cdot s_j$ with

$s' = \gamma(u^\flat)$. Therefore, $\gamma(u) = s'(\alpha_j) = -s'(s_j(\alpha_j)) = -s(\alpha_j) \in \Phi^+$ again by Lemma 2.2, which implies $s(\alpha_j) \in \Phi^-$ and, hence, $j \neq i$.

Next, let $t \in \mathcal{W}$ be an element of minimal length in the coset $s \cdot \mathcal{W}_{(ij)}$. Since $s \cdot s_j = \gamma(u^\flat)$ we have $l(s \cdot s_j) < l(s)$ so that $l(t) < l(s)$ and $l(r) \geq 1$. Minimality of t further implies that $l(t \cdot s_i) > l(t)$ and $l(t \cdot s_j) > l(t)$. Thus, if $x \in \mathcal{W}^*$ is a reduced word representing t also $x \cdot w_i$ and $x \cdot w_j$ are reduced, implying positivity of both $\gamma(x \cdot w_i) = t(\alpha_i)$, $\gamma(x \cdot w_j) = t(\alpha_j) \in \Phi^+$.

Since $\mathcal{N}(r) \subset \mathbb{N}_0\alpha_i + \mathbb{N}_0\alpha_j$ this also implies that $t(\mathcal{N}(r)) \subseteq \Phi^+$. By Corollary 2.4 we thus obtain that $l(s) = l(t) + l(r)$.

Observe further that $r(\alpha_i) = n_i\alpha_i + n_j\alpha_j$ with either both $n_i, n_j > 0$ if $r(\alpha_i) \in \Phi^+$ or both $n_i, n_j < 0$ if $r(\alpha_i) \in \Phi^-$. The latter case is not possible since it would imply $s(\alpha_i) = n_i t(\alpha_i) + n_j t(\alpha_j) \in \Phi^-$ by the previous observation. Now, $r(\alpha_i) \in \Phi^+$ implies that a reduced word representing r ends in w_j and that $l(r) < m$.

Finally, the condition $s(\alpha_i) = \alpha_k = n_i t(\alpha_i) + n_j t(\alpha_j)$ implies that either $t(\alpha_i) = \alpha_k$, with $(n_i, n_j) = (1, 0)$ and hence $r(\alpha_i) = \alpha_i$, or $t(\alpha_j) = \alpha_k$, with $(n_i, n_j) = (0, 1)$ and hence $r(\alpha_i) = \alpha_j$. Given that $l(r) \geq 1$ this now entails that $l(r) = m - 1$ so that $y \cdot w_i \in \mathcal{W}_{m(i,j)}^*$. \square

3. QUANTUM GROUPS AT GENERIC PARAMETER q

We recall the standard definitions of quantum groups as defined, for example, in [Jim85, CP94]. Various ground rings are introduced over which subalgebras and variants of the quantum groups can be defined. Throughout this section we fix a simple Lie algebra \mathfrak{g} with root system Φ and Cartan data A_{ij} as introduced in the previous section.

3.1. Quantization Rings, Quantum Numbers, Integral Subrings. In their original definitions, such as in [Dri87], quantum groups are obtained as deformations of the universal enveloping algebra of a Lie algebra \mathfrak{g} along a formal parameter \hbar . The natural ground ring in this context is the ring of formal power series $\mathbb{Q}[[\hbar]]$, which contains $q = e^\hbar = \sum_{n=0}^{\infty} \frac{1}{n!} \hbar^n$. The parameter identification yields a ring embedding $\mathbb{Q}(q)^1 \hookrightarrow \mathbb{Q}[[\hbar]]$, where $\mathbb{Q}(q)^1$ denotes the subring of rational functions without poles at $q = 1$. It extends to an embedding $\mathbb{Q}(q) \hookrightarrow \mathbb{Q}[[\hbar]]_\hbar$, where the localization $\mathbb{Q}[[\hbar]]_\hbar$ is given by formal Laurent series with finite numbers of negative powers. We continue our discussion with various frequently used quantities and formulae in $\mathbb{Z}[q, q^{-1}]$.

For a positive root $\alpha \in \Phi^+$ in the \mathcal{W} -orbit of $\alpha_i \in \Delta$, let $d_\alpha = d_i$ and $q_\alpha = q_i = q^{d_i}$. Given $a, b \in \mathbb{N}$, quantum numbers, factorials, and binomials are denoted with respect to this data as

$$[a]_\alpha = \frac{q_\alpha^a - q_\alpha^{-a}}{q_\alpha - q_\alpha^{-1}}, \quad [a]!_\alpha = \prod_{j=1}^a \frac{q_\alpha^j - q_\alpha^{-j}}{q_\alpha - q_\alpha^{-1}}, \quad \text{and} \quad \begin{bmatrix} a+b \\ a \end{bmatrix}_\alpha = \frac{[a+b]!_\alpha}{[a]!_\alpha [b]!_\alpha}, \quad (3.1)$$

all of which are valued in $\mathbb{Z}[q_\alpha, q_\alpha^{-1}] \subseteq \mathbb{Z}[q, q^{-1}]$. The subscript α is suppressed if $q_\alpha = q$, meaning $d_i = d_\alpha = 1$. We will frequently also use the abbreviation $[n]_i = [n]_{\alpha_i}$ for simple roots and

analogous notation for factorials and binomial coefficients. The binomial recursion below is easily verified and frequently used.

$$\begin{bmatrix} n+1 \\ t \end{bmatrix}_\alpha = q_\alpha^{\pm t} \begin{bmatrix} n \\ t \end{bmatrix}_\alpha + q_\alpha^{\mp(n+1-t)} \begin{bmatrix} n \\ t-1 \end{bmatrix}_\alpha \quad (3.2)$$

The q -binomial coefficients arise naturally in the respective binomial formula for q -commuting generators. Specifically, assume the algebra \mathcal{H}_q over $\mathbb{Z}[q, q^{-1}]$ is generated by U and V subject to the relation $VU = q^2UV$. Then the following is easily proved by induction,

$$(U + V)^a = \sum_{k=0}^a q^{-k(a-k)} \begin{bmatrix} a \\ k \end{bmatrix} U^k V^{a-k} = \sum_{k=0}^a q^{k(a-k)} \begin{bmatrix} a \\ k \end{bmatrix} V^k U^{a-k}. \quad (3.3)$$

Adjoining an inverse V^{-1} to \mathcal{H}_q and multiplying (3.3) by V^{-a} from the right one readily obtains, after the substitution $z = V^{-1}U$, the following variant of Gauss's formula,

$$\sum_{k=0}^a q^{-k(a+1)} \begin{bmatrix} a \\ k \end{bmatrix} z^k = \prod_{t=1}^a (q^{-2t}z + 1) = q^{-\binom{a+1}{2}} \prod_{t=1}^a (q^t + q^{-t}z). \quad (3.4)$$

For convenience in later computations we express ratios of factorials for different root lengths $d_j \in \{1, 2, 3\}$ as

$$\frac{[d_j k]!}{[k]!_j} = [d_j]^k \cdot [k : d_j]! \quad \text{and} \quad \frac{[k]!_j}{[k]!} = [d_j]^{-k} \cdot \{k : d_j\}, \quad (3.5)$$

where we use the following notation for elements in $\mathbb{Z}[q, q^{-1}]$,

$$[k : \mathbf{e}]! = \prod_{s=0}^{k-1} \prod_{t=1}^{\mathbf{e}-1} [s\mathbf{e} + t] \quad \text{and} \quad \{k : \mathbf{e}\} = \prod_{s=1}^k \frac{q^{\mathbf{e}s} - q^{-\mathbf{e}s}}{q^s - q^{-s}}. \quad (3.6)$$

Note that $[k : 1]! = 1 = \{k : 1\}$ for all $k \in \mathbb{N}$. On occasion, we will also use notation for quantum multinomial coefficients in $\mathbb{Z}[q, q^{-1}]$ given by

$$\begin{bmatrix} a_1 + a_2 + \dots + a_r \\ a_1, a_2, \dots, a_r \end{bmatrix}_\alpha = \frac{[a_1 + a_2 + \dots + a_r]!_\alpha}{[a_1]!_\alpha [a_2]!_\alpha \dots [a_r]!_\alpha} = \begin{bmatrix} a_1 + \dots + a_r \\ a_r \end{bmatrix}_\alpha \begin{bmatrix} a_1 + \dots + a_{r-1} \\ a_{r-1} \end{bmatrix}_\alpha \dots \begin{bmatrix} a_1 + a_2 \\ a_2 \end{bmatrix}_\alpha. \quad (3.7)$$

Concluding, we introduce the following notation for subrings of $\mathbb{Q}(q)$, over which various versions of quantum groups can be defined. Clearly, all of these are integral domains and have $\mathbb{Q}(q)$ as a common field of fractions. However, as opposed to $\mathbb{Q}(q)$, they have well-defined specializations to roots of unity of large enough orders.

$$\begin{aligned} \Lambda_q &= \mathbb{Z}[q, q^{-1}] & \Lambda_{q,n} &= \mathbb{Z}\left[q, q^{-1}, \frac{1}{[n-1]!}\right] \\ \Lambda_q^* &= \mathbb{Z}\left[q, q^{-1}, \frac{1}{q-q^{-1}}\right] & \Lambda_{q,n}^* &= \mathbb{Z}\left[q, q^{-1}, \frac{1}{q-q^{-1}}, \frac{1}{[n-1]!}\right] \end{aligned} \quad (3.8)$$

3.2. Quantum Group Generators and Relations. For Cartan data as above, the quantum universal enveloping algebra $U_{\hbar}(\mathfrak{g})$ is defined in [Dri87, CP94] as the algebra over $\mathbb{Q}[[\hbar]]$ topologically generated by elements E_i, F_i, H_i for $1 \leq i \leq n$ subject to relations

$$[H_i, H_j] = 0, \quad [E_i, F_j] = \delta_{ij} \frac{e^{d_i \hbar H_i} - e^{-d_i \hbar H_i}}{q_i - q_i^{-1}}, \quad (3.9)$$

$$[H_i, E_j] = A_{ij} E_j, \quad [H_i, F_j] = -A_{ij} F_j, \quad (3.10)$$

$$\sum_{r+s=1-A_{ij}} (-1)^s \begin{bmatrix} 1-A_{ij} \\ s \end{bmatrix}_i E_i^r E_j E_i^s = 0 \quad (\text{for } i \neq j), \quad (3.11)$$

$$\sum_{r+s=1-A_{ij}} (-1)^s \begin{bmatrix} 1-A_{ij} \\ s \end{bmatrix}_i F_i^r F_j F_i^s = 0 \quad (\text{for } i \neq j). \quad (3.12)$$

Although (3.9) and the other relations are well-defined over $\mathbb{Q}[[\hbar]]$ we extend the ground ring of $U_{\hbar}(\mathfrak{g})$ to $\mathbb{Q}[[\hbar]]_{\hbar}$ for convenience. If we denote as usual $K_i = e^{d_i \hbar H_i}$ the above imply the following relations:

$$K_i K_j = K_j K_i, \quad [E_i, F_j] = \delta_{ij} \frac{K_i - K_i^{-1}}{q_i - q_i^{-1}}, \quad (3.13)$$

$$K_i E_j K_i^{-1} = q_i^{A_{ij}} E_j = q^{(\alpha_i | \alpha_j)} E_j, \quad K_i F_j K_i^{-1} = q_i^{-A_{ij}} F_j = q^{-(\alpha_i | \alpha_j)} F_j. \quad (3.14)$$

Observing $(q_i - q_i^{-1}) = (q - q^{-1})[d_i]$, we note that all relations (3.11-3.14) are indeed over $\Lambda_{q, \mathbf{e}^*}$ where $\mathbf{e}^* \in \{2, 3, 4\}$ is as in (2.3) and we view $\Lambda_{q, \mathbf{e}^*} \subset \mathbb{Q}(q) \subset \mathbb{Q}[[\hbar]]_{\hbar}$. We thus define $U_q(\mathfrak{g})$ as the $\Lambda_{q, \mathbf{e}^*}$ -algebra generated by the elements $\{E_i, F_i, K_i : 1 \leq i \leq n\}$ and subject to relations (3.11-3.14). If \mathfrak{g} is a general simple Lie algebra or its type is clear from context, we will usually write U_q instead.

Additionally, we denote by U_q^+ the subalgebra generated by only the E_i , U_q^- the subalgebra generated by only the F_i , which may be considered as algebras over the subring $\Lambda_{q, \mathbf{e}^*}$ of $\Lambda_{q, \mathbf{e}^*}$. The algebra U_q^0 generated by only the K_i may be considered over Λ_q . Further, we set $U_q^{\geq 0} = U_q^+ U_q^0$ and $U_q^{\leq 0} = U_q^- U_q^0$ the respective subalgebras obtained from two sets of generators, which we also take to be defined over $\Lambda_{q, \mathbf{e}^*}$ by default.

Suppose $f : \Lambda_{q, \mathbf{e}^*} \rightarrow \Lambda$ or $f^* : \Lambda_{q, \mathbf{e}^*} \rightarrow \Lambda^*$ is a ring homomorphism where Λ or Λ^* is commutative with unit. We may define quantum algebras over ground rings Λ or Λ^* by respective extensions of scalars of the original algebras, viewed as $\Lambda_{q, \mathbf{e}^*}$ or $\Lambda_{q, \mathbf{e}^*}$ modules. Specifically, for any $\square \in \{+, -, 0, \geq 0, \leq 0\}$ we denote

$$U_{q, \Lambda^*} = f_!^* U_q = U_q \otimes_{\Lambda_{q, \mathbf{e}^*}} \Lambda^* \quad \text{and} \quad U_{q, \Lambda}^{\square} = f_! U_q^{\square} = U_q^{\square} \otimes_{\Lambda_{q, \mathbf{e}^*}} \Lambda. \quad (3.15)$$

The further abbreviated notation $U_{q, \mathbb{Q}}$ and $U_{q, \mathbb{Q}}^{\square}$ is used for algebras over $\mathbb{Q}(q)$ derived from the inclusions of $\Lambda_{q, \mathbf{e}^*}$ and $\Lambda_{q, \mathbf{e}^*}$ in $\mathbb{Q}(q)$. Similarly, for $m \geq \mathbf{e}^*$ we write $U_{q, m}$ and $U_{q, m}^{\square}$ for the

inclusions of Λ_{q,e^*} and Λ_{q,e^*} in $\Lambda_{q,m}$ and $\Lambda_{q,m}$, respectively. That is, the ground rings of $U_{q,m}$ and $U_{q,m}^\square$ contain also elements $\frac{1}{[m-1]!}$.

It is easy to see that the free abelian group algebra $\Lambda_q[\{K_i^{\pm 1}\}]$ embeds in the polynomial algebra $\mathbb{Q}[[\hbar]]_\hbar[\{H_i\}]$ via $K_i \mapsto e^{d_i \hbar H_i}$. Using the PBW results below and in [Ros89, CP94] it further follows that the respective algebra homomorphism $U_q \rightarrow U_\hbar(\mathfrak{g})$ is injective.

An important variant of quantum groups at roots of unity are Lusztig's divided power algebras $\mathcal{L}U$, see [Lus93, Lus90a, Lus90b]. They are defined as the Λ_q -subalgebras of $U_{q,\mathbb{Q}}$ generated by elements $E_i^{(k)} = \frac{1}{[k]!_i} E_i^k$ and $F_i^{(k)} = \frac{1}{[k]!_i} F_i^k$ as well as certain additional rational expressions in the K_i generators.

We denote the respective subalgebras by $\mathcal{L}U^+$, $\mathcal{L}U^-$, and $\mathcal{L}U^0$. Although they are not discussed explicitly in this paper, we will refer to these algebras in some of the proofs below as we adopt arguments from [Lus90a, Lus90b, Lus93].

3.3. Hopf Algebra Structures. Coalgebra structures on $U_\hbar(\mathfrak{g})$ and U_q are given by requiring K_i to be group-like, H_i to be primitive, and E_i and F_i to be *skew-primitives* via the co-relations

$$\Delta(E_i) = E_i \otimes K_i + 1 \otimes E_i \quad \text{and} \quad \Delta(F_i) = F_i \otimes 1 + K_i^{-1} \otimes F_i. \quad (3.16)$$

In this sense we have that the inclusions of the quantum algebras above, such as $U_q^{\geq 0} \hookrightarrow U_q \hookrightarrow U_\hbar(\mathfrak{g})$, are inclusions of Hopf algebras. A counit is given by $\varepsilon(E_i) = \varepsilon(F_i) = \varepsilon(H_i) = 0$ and $\varepsilon(K_i) = 1$. For later computations we also record here the coproducts for powers of generators, which easily follow from (3.14) and (3.16).

$$\Delta(E_i^m) = \sum_{s=0}^m q_i^{-s(m-s)} \begin{bmatrix} m \\ s \end{bmatrix}_i E_i^s \otimes K_i^s E_i^{m-s} \quad \Delta(F_i^m) = \sum_{s=0}^m q_i^{-s(m-s)} \begin{bmatrix} m \\ s \end{bmatrix}_i K_i^{-s} F_i^{m-s} \otimes F_i^s \quad (3.17)$$

The antipode acts on the Cartan generators as $S(K_i) = K_i^{-1}$ and $S(H_i) = -H_i$ and is given on the other generators as

$$S(E_i) = -E_i K_i^{-1} = -q_i^2 K_i^{-1} E_i \quad \text{and} \quad S(F_i) = -K_i F_i = -q_i^{-2} F_i K_i. \quad (3.18)$$

Again, it will be useful to also state the antipode of powers of generators.

$$S(E_i^m) = (-1)^m q_i^{-m(m-1)} E_i^m K_i^{-m} \quad S(F_i^m) = (-1)^m q_i^{m(m-1)} K_i^m F_i^m. \quad (3.19)$$

Clearly, $U_q^{\geq 0}$ and $U_q^{\leq 0}$ form Hopf subalgebras of U_q , and $U_q^0 = \Lambda_{q,e^*}[\{K_i^{\pm 1}\}]$ is isomorphic as a Hopf algebra to the group algebra of \mathbb{Z}^n .

Recall [Swe69, Mon93a, Rad12] that a coalgebra C over a field \mathbb{k} is called *pointed* if all its (co)simple subcoalgebras are 1-dimensional or, equivalently, if its coradical C_0 coincides with the subcoalgebra $\mathbb{k}[Gr(C)]$ of group-like elements. Pointed coalgebras and Hopf algebras have been extensively studied, though most results assume the ground ring to be a field.

Suppose \mathbb{k} is a field and $\vartheta : \Lambda_{q,e^*} \rightarrow \mathbb{k}$ a ring homomorphism. As noted in (3.15), this yields quantum algebras $U_{q,\mathbb{k}}$ and $U_{q,\mathbb{k}}^\square$ over the field \mathbb{k} for any $\square \in \{0, \geq 0, \leq 0\}$. Besides the case $\mathbb{k} = \mathbb{Q}(q)$ given by the obvious inclusions, the following result will also be of interest in the case of a cyclotomic field $\mathbb{k} = \mathbb{Q}(\zeta)$.

Proposition 3.1 ([Mon93b, Theorem 2.2]). *The Hopf algebras $U_{q,\mathbb{k}}$, $U_{q,\mathbb{k}}^{\geq 0}$, and $U_{q,\mathbb{k}}^{\leq 0}$ as defined above are pointed. Their common coradical is $U_{q,\mathbb{k}}^0$.*

The proof uses coalgebra filtrations $\{A_n\}$ of the quantum algebras, where $A_0 = U_{q,\mathbb{k}}^0$. Set $A_1 = A_0 \oplus \sum_i^\oplus E_i A_0 \oplus F_i A_0$ for $U_{q,\mathbb{k}}$ as well as $A_1 = A_0 \oplus \sum_i^\oplus E_i A_0$ for $U_{q,\mathbb{k}}^{\geq 0}$ and analogous for $U_{q,\mathbb{k}}^{\leq 0}$. The subspaces are then given as $A_n = (A_1)^n$, which clearly exhaust the entire algebra. Lemma 5.3.4 in [Mon93b], for example, now implies that C_0 is contained in and thus equal to $U_{q,\mathbb{k}}^0$. Pointedness now follows immediately from $C_0 = U_{q,\mathbb{k}}^0$, which is then also equal to the total subcoalgebra spanned by group-like elements.

4. GRADINGS AND AUTOMORPHISMS ON QUANTUM GROUPS

Several automorphisms and anti-automorphisms will play an important role in constructions throughout this paper. They include, besides the antipode, Lusztig's Artin group actions, involutions, and Che transformations. Additionally, we will refer to two types of gradings in several of our later the proofs. This section will collect basic properties of these automorphisms and gradings as well as establish various useful relations between them.

4.1. Gradings. Recall that a grading of an algebra A by a monoid G entails a decomposition $A = \bigoplus_{g \in G} A_g$ with $A_g \cdot A_h \subseteq A_{gh}$. We call elements $b \in A_g \setminus \{0\}$ for some $g \in G$ *homogeneous* with respect to such a given grading. If A is defined in terms of generators and relations, a grading on the free algebra of generators descends to A if the relations are homogeneous elements. Let G be a monoid and S any set. In the following discussion, we write G^S to mean interchangeably additive combinations $\sum_{s \in S} c_s s$ as well as maps $c : S \rightarrow G : s \mapsto c_s$.

A grading in $G = \mathbb{N}_0^\Delta$ with $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ on the algebra U_q^+ is given by defining $(U_q^+)_\mu$ as the subspace spanned by $E_{i_1} \dots E_{i_m}$ such that $\mu = \sum_{s=1}^m \alpha_{i_s} \in \mathbb{N}_0^\Delta$. It is clear that the relations in (3.11)-(3.14) are homogeneous with respect to this grading, which is therefore well-defined on U_q^+ . For a homogeneous element $b \in (U_q^+)_\mu$ we sometimes also write $w^+(b) = \mu$.

An analogous grading in \mathbb{N}_0^Δ is given on U_q^- by the function w^- from the set of homogeneous elements in U_q^- to \mathbb{N}_0^Δ . These extend to a \mathbb{Z}^Δ grading on U_q with $(U_q)_\mu$ spanned by elements $K_{j_1} \dots K_{j_k} E_{i_1}^{\epsilon_1} \dots E_{i_m}^{\epsilon_m}$ with $\mu = \sum_{s=1}^m \epsilon_s \alpha_{i_s}$ where $E_i^+ = E_i$ and $E_i^- = F_i$, observing that both sides of the relation for $[E_i, F_j]$ are in $(U_q)_{\alpha_i - \alpha_j}$. For a homogeneous element $a \in (U_q)_\mu$ with respect to this weight grading we also write $w(a) = \mu$. Thus, if $b \in U_q^+$ and $c \in U_q^-$ are homogeneous elements we have $w(bc) = w(cb) = w^+(b) - w^-(c)$, which readily implies, for example $(U_q^+)_\mu_1 \cdot (U_q^-)_\mu_2 \subseteq (U_q)_{\mu_1 - \mu_2}$.

For an element $v = \sum_i c_i \alpha_i \in \mathbb{Z}^\Delta$ in the root lattice, we denote a basis element of U_q^0 by

$$K^v = K_1^{c_1} \dots K_n^{c_n}. \quad (4.1)$$

Suppose $b \in U_q$ is homogeneous with $\mu = \text{w}(b) \in \mathbb{Z}^\Delta$. It is easy to derive the following commutation relation, where the inner form $(\cdot | \cdot)$ is naturally extended to a biadditive form on $\mathbb{Z}^\Delta \supset \Phi$,

$$K^v b = q^{(v|\text{w}(b))} b K^v. \quad (4.2)$$

Additional gradings in \mathbb{F}_2^Δ , where $\mathbb{F}_2 = \mathbb{Z}/2\mathbb{Z} = \{0, 1\}$, are artifacts of the $q \leftrightharpoons q^{-1}$ -symmetric definition of quantum numbers and Cartan elements for quantum groups. Two such gradings are defined on generators as follows:

$$\text{o}^+(E_i) = \alpha_i, \quad \text{o}^+(K_i) = \alpha_i, \quad \text{o}^+(F_i) = 0, \quad (4.3)$$

$$\text{o}^-(E_i) = 0, \quad \text{o}^-(K_i) = \alpha_i, \quad \text{o}^-(F_i) = \alpha_i. \quad (4.4)$$

As before, it is easy to check that relations (3.11)-(3.14) are homogeneous with respect to both o^+ and o^- , which are thus well-defined on U_q . For example, $\text{o}^+(E_i F_i) = \text{o}^+(F_i E_i) = \text{o}^+(K_i) = \text{o}^+(K_i^{-1}) = \alpha_i$ over \mathbb{F}_2 in the case of relation (3.13). We also note the following interactions of the antipode with these gradings, when evaluated on homogeneous elements:

$$\text{w} \circ S = \text{w}, \quad \text{o}^+ \circ S = \text{o}^-, \quad \text{o}^- \circ S = \text{o}^+. \quad (4.5)$$

Although the gradings are related by $\text{o}^+ - \text{o}^- = \text{w} \pmod{2}$, it is useful to denote them separately. Combining w with either o^+ or o^- yields a grading in $(\mathbb{Z} \times \mathbb{F}_2)^\Delta$ that provides a refinement of the usual weight decomposition, splitting summands further according to respective multiplicities of Cartan elements in the sense that $\text{o}^+(K^v) = v \pmod{2}$.

For a discussion of the behavior of the coproduct with respect to these gradings denote by U any of the Hopf algebras U_q , $U_q^{\geq 0}$ and $U_q^{\leq 0}$. For $\mu \in \mathbb{Z}^\Delta$ and $\eta \in \mathbb{F}_2^\Delta$ write also U_μ and U^η for the graded components for w and o^+ respectively. The algebra $U \otimes U$ thus assumes natural bi-gradings with $(U \otimes U)_{(\mu_1, \mu_2)} = U_{\mu_1} \otimes U_{\mu_2}$ and $(U \otimes U)^{(\eta_1, \eta_2)} = U^{\eta_1} \otimes U^{\eta_2}$. A simple verification on generators now yields

$$\Delta(U_\mu) \subseteq \sum_v U_{\mu-v} \otimes U_v \quad \text{and} \quad \Delta(U^\eta) \subseteq U \otimes U^\eta. \quad (4.6)$$

Finally, let $b \in (U_q^+)_\mu$ and $c \in (U_q^-)_{-\nu}$ be w -homogeneous elements with $\mu, \nu \in \mathbb{N}_0^\Delta$. It follows from an easy inductive verification on generators $E_{i_1} \dots E_{i_k}$ and $F_{j_1} \dots F_{j_m}$ that the $(\mu, 0)$ and $(0, \mu)$ graded components of $\Delta(b)$ as well as the $(-\nu, 0)$ and $(0, -\nu)$ graded components of $\Delta(c)$ have the form

$$\Delta(b) = b \otimes K^\mu + \dots + 1 \otimes b \quad \text{and} \quad \Delta(c) = c \otimes 1 + \dots + K^{-\nu} \otimes c. \quad (4.7)$$

4.2. Che Transformations and Anti-Involutions. A basic family of automorphisms can be obtained by multiplying the step generators by Cartan elements and units subject to certain consistency conditions. The main data characterizing such a *Che* (Ψ) *transformation* is a Φ -symmetric map $h \in \mathbb{S}(\Phi)$ as introduced in Section 2.6.

Additionally, assume a homomorphism $u : \mathbb{Z}^\Delta \rightarrow \Lambda^*$ from the root lattice to the multiplicative subgroup of units $\Lambda^* = \{\pm q^k : k \in \mathbb{Z}\}$ of the ground ring. We define a map Ψ_u^h on generators as follows:

$$\Psi_u^h(E_i) = u(\alpha_i)K^{h(\alpha_i)}E_i, \quad \Psi_u^h(F_i) = u(\alpha_i)^{-1}F_iK^{-h(\alpha_i)}, \quad \Psi_u^h(K_i) = K_i. \quad (4.8)$$

The next proposition establishes the existence and basic properties of these automorphisms. As before, U_q^\square may be U_q , $U_q^{\geq 0}$, or $U_q^{\leq 0}$ defined over the respective minimal ground rings. The q -exponent in the formula in *iii*) is integral by Lemma 2.11, so that the q -term is well-defined by *iv*) of the same lemma.

Proposition 4.1. *Let $h, k \in \mathbb{S}(\Phi)$ and $u, v \in \text{Hom}(\mathbb{Z}^\Delta, \Lambda^*)$ as above. Then the following hold.*

- i) *The map Ψ_u^h defined in (4.8) extends uniquely to an automorphism of U_q^\square .*
- ii) *$\Psi_u^h \circ \Psi_v^k = \Psi_{u \cdot v}^{h+k}$, and Ψ_1^0 is identity, where 1 is the constant map to 1 $\in \Lambda^*$.*
- iii) *If b is a homogeneous element with grading $\beta = \mathbf{w}(b)$ we have*

$$\Psi_u^h(b) = u(\beta)q^{-\frac{1}{2}((\beta|h(\beta))-2(\rho_h|\beta))}K^{h(\beta)}b.$$

- iv) *With b as above, the action on gradings is $\mathbf{w} \circ \Psi_u^h = \mathbf{w}$ and $\circ^+(\Psi_u^h(b)) = \circ^+(b) + h(\beta) \bmod 2$.*

Proof. Consider the algebra L_q^\square with the same generators as U_q^\square but subject only to the relations (3.14) in addition to those for $U_q^0 \cong \Lambda[\{K_i^{\pm 1}\}]$. L_q^\square admits the same gradings as U_q^\square , and commutation of the K_i in $L_q^0 \subset L_q^\square$ implies that Ψ_u^h is well-defined on L_q^\square . We next prove that *iii)* holds on L_q^\square by induction in the number of E_i and F_i generators in an expression for b . Since Ψ_u^h acts trivially on U_q^0 we may omit any additional K_i factors in b .

For $b = E_i$ and $\beta = \alpha_i$ the q -exponent is zero because of (2.29) so that the expression is the same as in (4.8). For $b = F_i$ and $\beta = -\alpha_i$ use $u(-\alpha_i) = u(\alpha_i)^{-1}$, $h(-\alpha_i) = -h(\alpha_i)$, and $(-\alpha_i|h(-\alpha_i)) - 2(\rho_h|-\alpha_i) = 2(\alpha_i|h(\alpha_i))$ again by (2.29).

For the induction step, we compute the formula for $E_i b$ with $\mathbf{w}(E_i b) = \alpha_i + \beta$ assuming the asserted expressions for E_i and b and using the fact that u and h are homomorphisms,

$$\begin{aligned} \Psi_u^h(E_i b) &= \Psi_u^h(E_i) \Psi_u^h(b) \\ &= u(\alpha_i)u(\beta)q^{-\frac{1}{2}((\alpha_i|h(\alpha_i))-2(\rho_h|\alpha_i))}q^{-\frac{1}{2}((\beta|h(\beta))-2(\rho_h|\beta))}K^{h(\alpha_i)}E_iK^{h(\beta)}b \\ &= u(\alpha_i + \beta)q^{-\frac{1}{2}((\alpha_i|h(\alpha_i))+(\beta|h(\beta))-2(\rho_h|\alpha_i+\beta))}q^{-(\alpha_i|h(\beta))}K^{h(\alpha_i)}K^{h(\beta)}E_i b \end{aligned}$$

$$\begin{aligned}
&= u(\alpha_i + \beta) q^{-\frac{1}{2}((\alpha_i|h(\alpha_i)) + 2(\alpha_i|h(\beta)) + (\beta|h(\beta)) - 2(\rho_h|\alpha_i + \beta))} K^{h(\alpha_i + \beta)} E_i b \\
&= u(\alpha_i + \beta) q^{-\frac{1}{2}((\alpha_i + \beta|h(\alpha_i + \beta)) - 2(\rho_h|\alpha_i + \beta))} K^{h(\alpha_i + \beta)} E_i b.
\end{aligned}$$

In the last step, we invoke the identity $(\alpha_i|h(\beta)) = (\beta|h(\alpha_i))$, which is equivalent to our assumption that h is Φ -symmetric. The respective computation for $F_i b$ is analogous, and the extension to Λ -linear combinations obvious.

Since $\Psi_u^h(b) = g_\beta b$ for a w -homogeneous elements b with g_β invertible, Ψ_u^h maps the two-sided ideal generated by b to itself. Thus, Ψ_u^h is well-defined on any quotient of L_q^\square obtained by dividing by w -homogeneous relations. Since all relations in (3.11)-(3.14) are indeed w -homogeneous, we conclude the statement in Item *i*). The statement in Item *ii*) is easily checked on the generators, and Item *iv*) is immediate from Item *iii*). \square

Given some $h \in \mathbb{S}(\Phi)$ and $\beta \in \mathbb{Z}^\Delta$, we know from Lemma 2.11 *ii*) that $2(\rho_h|\beta) \in \mathbb{Z}$. We can thus define a character $\varpi_h \in \text{Hom}(\mathbb{Z}^\Delta, \Lambda^\star)$ by

$$\varpi_h : \mathbb{Z}^\Delta \rightarrow \Lambda^\star : \beta \mapsto \varpi_h(\beta) = q^{2(\rho_h|\beta)}. \quad (4.9)$$

Note that $\varpi_h \cdot \varpi_k = \varpi_{h+k}$. We will also use the abbreviation $\varpi = \varpi_{id}$ so that $\varpi(\alpha_i) = q_i^2$.

In [Lus90b] Lusztig introduces two anti-involutions. The Cartan anti-involution $\Omega : U_q \rightarrow U_q^{\text{opp, cop}}$ is defined as the \mathbb{Z} -algebra homomorphism with the following actions on algebra generators and the indeterminate of the ground ring.

$$\Omega(E_i) = F_i, \quad \Omega(F_i) = E_i, \quad \Omega(K_i) = K_i^{-1}, \quad \Omega(q) = q^{-1}. \quad (4.10)$$

Secondly, let $\mathfrak{O} : U_q \rightarrow U_q^{\text{opp}}$ be the involutive $\Lambda_{q, \mathfrak{e}^*}^\star$ -algebra homomorphism defined on generators as follows.

$$\mathfrak{O}(E_i) = E_i, \quad \mathfrak{O}(F_i) = F_i, \quad \mathfrak{O}(K_i) = K_i^{-1}. \quad (4.11)$$

Clearly, \mathfrak{O} preserves both the w and \mathfrak{o}^\pm gradings, while $w \circ \Omega = -w$ and $\mathfrak{o}^+ \circ \Omega = \mathfrak{o}^-$. In addition to Lusztig's involutions, it will be useful to also introduce the conjugation automorphism $\Upsilon : U_q \rightarrow U_q^{\text{opp}}$ defined on generators by

$$\Upsilon(E_i) = E_i, \quad \Upsilon(F_i) = F_i, \quad \Upsilon(K_i) = K_i, \quad \Upsilon(q) = q^{-1}. \quad (4.12)$$

All three of the above involutions may also be viewed as involutive anti-homomorphisms. It follows easily by inspection on generators that Ω , \mathfrak{O} , and Υ commute with each other and, thus, implement an action of $(\mathbb{F}_2)^3$ on U_q .

For later use, we introduce notation for two composites of these. Firstly, $\Pi = \Upsilon \circ \Omega = \Omega \circ \Upsilon : U_q \rightarrow U_q$ is a $\Lambda_{q, \mathfrak{e}^*}^\star$ -algebra homomorphism given on generators by

$$\Pi(E_i) = F_i, \quad \Pi(F_i) = E_i, \quad \Pi(K_i) = K_i^{-1}. \quad (4.13)$$

Secondly, let $\Pi = \Upsilon \circ \mathfrak{U} = \mathfrak{U} \circ \Upsilon : U_q \rightarrow U_q$, which is a \mathbb{Z} -algebra homomorphism with non-trivial action on the ground ring given by

$$\Pi(E_i) = E_i, \quad \Pi(F_i) = F_i, \quad \Pi(K_i) = K_i^{-1}, \quad \Pi(q) = q^{-1}. \quad (4.14)$$

Clearly, the automorphisms \mathfrak{U} , Υ , and Π map each of the subalgebras U_q^\square with $\square \in \{0, +, -, \geq 0, \leq 0\}$ to itself, while the Ω and Π map these algebras to their respective opposites.

Additional commutation relations with the antipode and the Che transformations from (4.8) are immediate from checks on generators as well. More specifically, for a Φ -symmetric map $h \in \text{End}(\mathbb{Z}^\Delta)$ and a character $u \in \text{Hom}(\mathbb{Z}^\Delta, \Lambda^\star)$ we obtain the following commutations relations of these involutions with the Ψ_u^h transformations, where ϖ_h is as in (4.9).

$$\begin{aligned} \Omega \circ \Psi_u^h &= \Psi_u^h \circ \Omega & \mathfrak{U} \circ \Psi_u^h &= \Psi_{u \cdot \varpi_h}^{-h} \circ \mathfrak{U} \\ \Pi \circ \Psi_u^h &= \Psi_{u^{-1} \cdot \varpi_h^{-1}}^h \circ \Pi & \Upsilon \circ \Psi_u^h &= \Psi_{u^{-1} \cdot \varpi_h^{-1}}^h \circ \Upsilon \end{aligned} \quad (4.15)$$

Furthermore, the antipode can be expressed as the composite

$$S = \Psi_{\iota \cdot \varpi}^{-id} \circ \mathfrak{U} = \mathfrak{U} \circ \Psi_{\iota}^{id}, \quad \text{where } \iota(\beta) = (-1)^{\text{ht}(\beta)} = (-1)^{(\check{\rho}|\beta)}, \quad (4.16)$$

with ht and $\check{\rho}$ as discussed at the end of Section 2.3. Commutation relations of the antipode with the other automorphisms can now be derived directly from (4.16) and the commutation relations in (4.15). For example, S commutes with Ω , $S \circ \mathfrak{U} = \Psi_{\varpi}^{-2id} \circ \mathfrak{U} \circ S$, and $S^2 = \Psi_{\varpi}^0$.

Finally, note that \mathfrak{U} and Υ preserve the w and \mathfrak{o}^\pm gradings. For Ω we find

$$w \circ \Omega = -w, \quad \mathfrak{o}^+ \circ \Omega = \mathfrak{o}^-, \quad \text{and} \quad \mathfrak{o}^- \circ \Omega = \mathfrak{o}^+, \quad (4.17)$$

implying the same equations for Π .

4.3. Lusztig's Action of Artin-Tits Monoid. In [Lus90a, Lus90b, Lus93] Lusztig introduces for any Lie type \mathfrak{g} algebra automorphisms T_i on $U_{q,\mathbb{Q}}$ as well as on $\mathcal{L}U$ for $1 \leq i \leq n$ where n is the rank of \mathfrak{g} . Importantly, they fulfill the Artin-Tits relations (2.6) for the respective Cartan data.

In order to conform with conventions in the R -matrix constructions in [KR90] we will prefer to work with the inverses $\Gamma_i = T_i^{-1}$ of these automorphisms. Their explicit actions on the generators of U_q are as follows:

$$\Gamma_i(E_i) = -K_i^{-1}F_i, \quad \Gamma_i(F_i) = -E_iK_i, \quad \Gamma_i(K_j) = K_j K_i^{-A_{ij}}, \quad (4.18)$$

$$\Gamma_i^{-1}(E_i) = -F_iK_i, \quad \Gamma_i^{-1}(F_i) = -K_i^{-1}E_i, \quad \Gamma_i^{-1}(K_j) = K_j K_i^{-A_{ij}}, \quad (4.19)$$

and for $i \neq j$

$$\Gamma_i(E_j) = \sum_{r+s=-A_{ij}} \frac{(-1)^r q_i^{-s}}{[s]!_i [r]!_i} E_i^s E_j E_i^r, \quad \Gamma_i(F_j) = \sum_{r+s=-A_{ij}} \frac{(-1)^r q_i^s}{[r]!_i [s]!_i} F_i^r F_j F_i^s, \quad (4.20)$$

$$\Gamma_i^{-1}(E_j) = \sum_{r+s=-A_{ij}} \frac{(-1)^r q_i^{-s}}{[r]!_i [s]!_i} E_i^r E_j E_i^s, \quad \Gamma_i^{-1}(F_j) = \sum_{r+s=-A_{ij}} \frac{(-1)^r q_i^s}{[s]!_i [r]!_i} F_i^s F_j F_i^r. \quad (4.21)$$

Lemma 4.2. *The automorphisms Γ_i restrict to automorphisms on the algebra U_q over Λ_{q,\mathbf{e}^*} and fulfill the Artin-Tits relations (2.6) for all Lie types.*

The proof of the first assertion is immediate, observing that the denominators in the above expressions divide $[-A_{ij}]!_i$ and hence $[-d_i A_{ij}]!$ as well as $[\mathbf{e}]! = [\mathbf{e}^* - 1]!$. Thus, the Γ_i automorphisms are well-defined over the ground ring Λ_{q,\mathbf{e}^*} . Moreover, the Artin-Tits relations have been verified for the T_i automorphisms in [Lus90a, Lus90b, Lus93] and thus also hold for the $\Gamma_i = T_i^{-1}$.

An immediate consequence is that we can associate an automorphism to any Weyl group element. For each reduced word $w = w_{i_1} \dots w_{i_k} \in \mathcal{W}^*$, let $\Gamma_w = \Gamma_{i_1} \circ \dots \circ \Gamma_{i_k}$. By Lemma 2.1, the composed automorphism only depends on the Weyl group element $s = r(w)$ represented by w , allowing us to write $\Gamma_w = \Gamma_s$. Now, for $s, t \in \mathcal{W}$ the construction implies

$$\Gamma_{st} = \Gamma_s \circ \Gamma_t \quad \text{if } l(st) = l(s) + l(t). \quad (4.22)$$

That is, due to the length additivity condition $s \mapsto \Gamma_s$ is *not* a representation of \mathcal{W} . Instead, the above notation should be understood as the composite of the Matsumoto map $\mathfrak{s} : \mathcal{W} \rightarrow \mathcal{A}^+$ from Section 2.2 with the natural monoid homomorphism $\Gamma : \mathcal{A}^+ \rightarrow \text{Aut}(U_q)$ defined on the positive Artin-Tits monoid, which extends to an action $\Gamma : \mathcal{A} \rightarrow \text{Aut}(U_q)$ of the Artin group on U_q . Thus, depending on context, we allow subscripts of Γ to be in \mathcal{W} , \mathcal{W}^* , or \mathcal{A} . These notations are related, for example, by $\Gamma_w = \Gamma_{r(w)}$ if $w \in \mathcal{W}^*$ or $\Gamma_s = \Gamma_{\mathfrak{s}(s)}$ if $s \in \mathcal{W}$.

Observe that Ω commutes with all Γ_i and T_i . Also, for the \mathfrak{O} involution one readily infers $\mathfrak{O} \circ \Gamma_i \circ \mathfrak{O} = \Gamma_i^{-1} = T_i$, or, more generally,

$$\mathfrak{O} \circ \Gamma_s \circ \mathfrak{O} = (\Gamma_{s^{-1}})^{-1} = T_s \quad \forall s \in \mathcal{W}. \quad (4.23)$$

Here $T_s = T_{i_1} \circ \dots \circ T_{i_k}$ denotes the analogous automorphisms used by Lusztig to define generators associated to general positive roots. The relation in (4.23) will thus serve to relate our conventions of choosing generators to those in [Lus90a, Lus90b]. The second equality may also be written as $\Gamma_b = T_{\iota(b)}$ for an element $b \in \mathcal{A}$, where $\iota \in \text{Aut}(\mathcal{A})$ maps each generator of \mathcal{A} to its inverse. The action of the braid automorphisms on the U_q^0 basis elements is naturally given by

$$\Gamma_s(K^\nu) = K^{s(\nu)}. \quad (4.24)$$

Finally, for a homogeneous element $b \in U_q$ one readily works out the following equivariance relations for the gradings, where the action of $s \in \mathcal{W}$ is naturally extended to the root lattice $\mathbb{Z}^\Delta \supset \Phi$ as well as \mathbb{F}_2^Δ .

$$\mathfrak{w}(\Gamma_s(b)) = s(\mathfrak{w}(b)) \quad (4.25)$$

$$\mathfrak{o}^\pm(\Gamma_s(b)) = s(\mathfrak{o}^\pm(b)) \quad (4.26)$$

Clearly, (4.24), (4.25) and (4.26) hold also if the Γ_s are replaced by the original T_s automorphisms.

4.4. More \mathcal{A}^+ Commutation Relations. In this section, we provide commutation relations between the Γ_s automorphisms and the Che transformations Ψ_u^h , from which we also obtain respective commutation relations with the antipode. We denote the natural (left) actions of \mathcal{W} on elements $h \in \mathbb{S}(\Phi)$ and $u \in \text{Hom}(\mathbb{Z}^\Delta, \Lambda^\star)$ by $h^s = shs^{-1}$ and $s_*u = u \circ s^{-1}$, respectively. Additionally, define the character $\vartheta_s^h \in \text{Hom}(\mathbb{Z}^\Delta, \Lambda^\star)$ by

$$\vartheta_s^h(\beta) = q^{-(\theta_h(s)|\beta)}, \quad (4.27)$$

which is well-defined by Lemma 2.11 *vi*). The following basic relations involving this character are easily derived from statement *viii*) as well as the combination of (2.27) and (4.9):

$$\vartheta_{t \cdot s}^h = \vartheta_t^{h^s} \cdot t_*(\vartheta_s^h) \quad \text{and} \quad (\vartheta_s^h)^2 = (s_*\varpi_h) \cdot (\varpi_{h^s})^{-1}. \quad (4.28)$$

Proposition 4.3. *Suppose $h \in \mathbb{S}(\Phi)$, $u \in \text{Hom}(\mathbb{Z}^\Delta, \Lambda^\star)$, and $s \in \mathcal{W}$. Using notations and conventions as above we have*

$$\Gamma_s \circ \Psi_u^h = \Psi_{(s_*u) \cdot \vartheta_s^h}^{h^s} \circ \Gamma_s. \quad (4.29)$$

Proof. Since all maps in the relation are automorphisms it suffices to check (4.29) for the generators of U_q . Moreover, since the Ψ_u^h act trivially on the K_i 's and since Ω commutes with both Ψ_u^h and Γ_s it suffices to check (4.29) for only the E_i generators. Note further that for a homogeneous element b with $w(b) = \beta$ we have $\Psi_{s_*u}^0(\Gamma_s(b)) = (s_*u)(s(\beta))\Gamma_s(b) = u(\beta)\Gamma_s(b) = \Gamma_s(u(\beta)b) = \Gamma_s(\Psi_u^0(b))$, using $w(\Gamma_s(b)) = s(\beta)$ by (4.25), so that (4.29) holds for $h = 0$. Since $\Psi_u^h = \Psi_u^0 \circ \Psi_1^h$ we may, therefore, assume $u = 1$ in the following verification.

$$\begin{aligned} \Psi_{\vartheta_s^h}^{h^s}(\Gamma_s(E_i)) &= \vartheta_s^h(s(\alpha_i))q^{-\frac{1}{2}((s(\alpha_i)|h^s(s(\alpha_i)))-2(\rho_{h^s}|s(\alpha_i)))}K^{h^s(s(\alpha_i))}\Gamma_s(E_i) \\ &= q^{-(\theta_h(s)|s(\alpha_i))}q^{-\frac{1}{2}((s(\alpha_i)|s(h(\alpha_i)))-2(\rho_{shs^{-1}}|s(\alpha_i)))}K^{s(h(\alpha_i))}\Gamma_s(E_i) \\ &= q^{-(\rho_{shs^{-1}}|s(\alpha_i))-(s(\rho_h)|s(\alpha_i))}q^{-\frac{1}{2}((\alpha_i|h(\alpha_i))-2(\rho_{shs^{-1}}|s(\alpha_i)))}\Gamma_s(K^{h(\alpha_i)})\Gamma_s(E_i) \\ &= q^{-\frac{1}{2}((\alpha_i|h(\alpha_i))-2(s(\rho_h)|s(\alpha_i)))}\Gamma_s(K^{h(\alpha_i)})E_i \\ &= q^{-\frac{1}{2}((\alpha_i|h(\alpha_i))-2(\rho_h|\alpha_i))}\Gamma_s(\Psi_1^h(E_i)) = \Gamma_s \circ \Psi_1^h(E_i) \end{aligned}$$

In the first line, we use Proposition 4.1 *iii*) for $\beta = w(\Gamma_s(E_i)) = s(\alpha_i)$. In the second line, definitions are inserted, including $h^s(s(\alpha_i)) = shs^{-1}s(\alpha_i) = sh(\alpha_i)$. The third line invokes isometry of s and (4.24). In the fourth line, exponent terms are combined and canceled, and we use that Γ_s is an automorphism. In the last lines s -isometry, the definition for Ψ_1^h , and (2.29) are employed. \square

The action of the Artin-Tits monoid \mathcal{A}^+ on the set of Che transformations given by conjugation thus factors into an action of \mathcal{W} on $\mathbb{S}(\Phi) \times \text{Hom}(\mathbb{Z}^\Delta)$. The latter is defined as $s.(h, u) = (h^s, (s_*u) \cdot \vartheta_s^h)$ using (4.28). An immediate consequence is that any Che transformation commutes with the action of the *pure* Artin-Tits group on U_q , defined as the kernel of the map $\mathcal{A} \rightarrow \mathcal{W}$.

As an application, we also derive an explicit commutation relation between the Γ_s maps and the antipode. To this end, let $\kappa_s \in \text{Hom}(\mathbb{Z}^\Delta, \Lambda^\star)$ be the character defined by

$$\kappa_s(\beta) = (-1)^{(\check{\theta}(s)|\beta)} q^{-(\theta(s)|\beta)}, \quad (4.30)$$

with $\theta(s)$ and $\check{\theta}(s)$ as in (2.24) or (2.27). Relations (2.25) and (2.26) translate for $l(t \cdot s) = l(t) + l(s)$ into

$$\kappa_{t \cdot s} = \kappa_t \cdot t_*(\kappa_s) \quad \text{and} \quad s_*(\kappa_{s^{-1}}) = (\kappa_s)^{-1}. \quad (4.31)$$

Proposition 4.4. *For any $s \in \mathcal{W}$ we have*

$$S \circ (\Gamma_s)^{-1} = \Gamma_{(s^{-1})} \circ S \circ \Psi_{\kappa_s}^0 \quad \text{and} \quad (\Gamma_s)^{-1} \circ S \circ \Psi_{\kappa_s}^0 = S \circ \Gamma_{(s^{-1})}. \quad (4.32)$$

Proof. We first establish that $\kappa_s = (s_* \iota) \cdot \iota^{-1} \cdot \vartheta_s^{id}$ with ι as in (4.16). Using (2.27) and the fact that s is an isometry we compute

$$\begin{aligned} ((s_* \iota) \cdot \iota^{-1} \cdot \vartheta_s^{id})(\beta) &= \iota(s^{-1}(\beta)) \iota(\beta)^{-1} \vartheta_s^{id}(\beta) &= (-1)^{(\check{\rho}|s^{-1}(\beta))} (-1)^{-(\check{\rho}|\beta)} q^{-(\theta_{id}(s)|\beta)} \\ &= (-1)^{(s(\check{\rho})|\beta) - (\check{\rho}|\beta)} q^{-(\theta(s)|\beta)} &= (-1)^{(\check{\theta}(s)|\beta)} q^{-(\theta(s)|\beta)} &= \kappa_s(\beta). \end{aligned}$$

The following verification of the first identity invokes the expression in (4.16) for the antipode, the commutation rule from Proposition 4.3, the conjugation with \mathfrak{U} from (4.23), homomorphy from Proposition 4.1ii), and the above relation for κ_s .

$$\begin{aligned} S \circ (\Gamma_s)^{-1} &= \mathfrak{U} \circ \Psi_{\iota}^{id} \circ (\Gamma_s)^{-1} &= \mathfrak{U} \circ (\Gamma_s)^{-1} \circ \Psi_{(s_* \iota) \cdot \vartheta_s^{id}}^{id} &= \Gamma_{(s^{-1})} \circ \mathfrak{U} \circ \Psi_{(s_* \iota) \cdot \vartheta_s^{id}}^{id} \\ &= \Gamma_{(s^{-1})} \circ S \circ (\Psi_{\iota}^{id})^{-1} \circ \Psi_{(s_* \iota) \cdot \vartheta_s^{id}}^{id} &= \Gamma_{(s^{-1})} \circ S \circ \Psi_{\iota^{-1} \cdot (s_* \iota) \cdot \vartheta_s^{id}}^0 &= \Gamma_{(s^{-1})} \circ S \circ \Psi_{\kappa_s}^0 \end{aligned}$$

The second identity can be derived from the first by replacing s with s^{-1} and (4.31). Specifically, $S \circ (\Gamma_{(s^{-1})})^{-1} = \Gamma_s \circ S \circ \Psi_{\kappa_{(s^{-1})}}^0$ implies

$$(\Gamma_s)^{-1} \circ S = S \circ \Psi_{\kappa_{(s^{-1})}}^0 \circ \Gamma_{(s^{-1})} = S \circ \Psi_{(s^{-1})_* (\kappa_s^{-1})}^0 \circ \Gamma_{(s^{-1})} = S \circ \Gamma_{(s^{-1})} \circ \Psi_{(\kappa_s^{-1})}^0 = S \circ \Gamma_{(s^{-1})} \circ (\Psi_{\kappa_s}^0)^{-1},$$

and, hence, the second identity. \square

The character κ_s also appears in commutation relations of the Artin-Tits groups action with the conjugation anti-automorphism Υ .

Proposition 4.5. *For any element $s \in \mathcal{W}$ we have*

$$\Gamma_s \circ \Upsilon = \Upsilon \circ \Gamma_s \circ \Psi_{\kappa_{(s^{-1})}}^0 = \Psi_{\kappa_s}^0 \circ \Upsilon \circ \Gamma_s. \quad (4.33)$$

Proof. The second equality follows from $\Gamma_s \circ \Psi_{\kappa_{(s^{-1})}}^0 = \Psi_{s_* \kappa_{(s^{-1})}}^0 \circ \Gamma_s = \Psi_{\kappa_s^{-1}}^0 \circ \Gamma_s$ by (4.29) and (4.31) as well as (4.15). For $s = s_i$ equation (4.33) applied to a generator E_j reads

$$\begin{aligned} \Gamma_i(E_j) &= \Upsilon(\Gamma_i(\kappa_{s_i}(\alpha_j) E_j)) = \kappa_{s_i}(\alpha_j)^{-1} \Upsilon(\Gamma_i(E_j)) \\ &= (-1)^{(\check{\alpha}_i|\alpha_j)} q^{(\alpha_i|\alpha_j)} \Upsilon(\Gamma_i(E_j)) = (-1)^{A_{ij}} q_i^{A_{ij}} \Upsilon(\Gamma_i(E_j)). \end{aligned}$$

This is readily verified for both the $i = j$ and $i \neq j$ cases from (4.18), (4.20), and (4.12). Since Ω commutes with both Γ_i and Υ and, furthermore, inverts q the relation for $s = s_i$ also holds on F_i . As it is also trivially true on K_i we proved (4.33) for $s = s_i$. Suppose now (4.33) holds for two elements $s, t \in \mathcal{W}$ with $l(s), l(t) \geq 1$ and $l(ts) = l(t) + l(s)$. Then

$$\begin{aligned}\Gamma_{ts} \circ \Upsilon &= \Gamma_t \circ \Gamma_s \circ \Upsilon = \Gamma_t \circ \Psi_{\kappa_s}^0 \circ \Upsilon \circ \Gamma_s = \Psi_{t_*(\kappa_s)}^0 \circ \Gamma_t \circ \Upsilon \circ \Gamma_s \\ &= \Psi_{t_*(\kappa_s)}^0 \circ \Psi_{\kappa_t}^0 \circ \Upsilon \circ \Gamma_t \circ \Gamma_s = \Psi_{t_*(\kappa_s) \cdot \kappa_t}^0 \circ \Upsilon \circ \Gamma_{ts} = \Psi_{\kappa_{ts}}^0 \circ \Upsilon \circ \Gamma_{ts},\end{aligned}$$

where we used (4.29) and (4.31). The assertion now follows by induction. \square

5. GENERATORS, BASES, AND INTEGRALITY

Poincaré-Birkhoff-Witt (PBW) bases have been established for quantum deformations of Lie algebras in many settings. These typically consider quantum algebras defined over a field such as $\mathbb{Q}(q)$ or \mathbb{C} and assume specific fixed choices with respect to which the quantum analogs of the E_α operators are defined. See, for example, [Ros89, CP94, Jan96, Rin96, Lus93].

Lusztig describes the algebras $\mathcal{L}U^\square$ of divided powers as free modules over Λ_q in [Lus90a, Lus90b] for specific choices of E_α . In [Lus93] explains independence of some of these choices for respective bases over $\mathbb{Q}(q)$. Specifically, certain spanning sets $\mathfrak{B}_{w\bullet}^+$ of monomial expressions, to be defined below, will depend on the choice of a maximal reduced word $w \in \mathcal{W}_m^*$.

Corollary 40.2.2 in [Lus93] implies that $\mathfrak{B}_{w\bullet}^+$ is a basis of $U_{q,\mathbb{Q}}^+$ over the field $\mathbb{Q}(q)$ for all Lie types and all choices of w , which also implies that $\mathfrak{B}_{w\bullet}^+$ is a linearly independent set of generators for the Λ_{q,e^*} -module U_q^+ and that its span $\mathfrak{S}_{w\bullet}^+$ is a free Λ_{q,e^*} -module. The $\mathbb{Q}(q)$ -basis does, however, not exclude the possibility that $\mathcal{T}_w = U_q^+ / \mathfrak{S}_{w\bullet}^+$ is a non-zero torsion module and, thus, does not imply that $\mathfrak{B}_{w\bullet}^+$ is also a Λ_{q,e^*} -basis.

For versions of U_q^+ for classical Lie types, bases over Λ_q are obtained from iterated q -brackets in [Tak90] for specific lexicographic orderings. In the formalism and constructions of generators used there, however, it is difficult to describe quasi-triangular Hopf algebra structures or braid actions for reordering. Still, the example provided in Proposition 5.6 in the same paper illustrates the delicate dependence of integrality of bases on the chosen orders.

Specifically, Takeuchi shows for an algebra $\bar{\mathcal{B}}_3^\circ$, which can be roughly understood as a quotient of U_q^+ for Lie type B_3 , that the analog of the torsion module \mathcal{T}_w is indeed zero for a convex ordering but contains non-zero torsion summands such as $\mathbb{Z}[q, q^{-1}] / \langle q^3 - q^2 + 1 \rangle$ for a non-convexly ordered monomial basis. A careful treatment of orderings will thus have to be a crucial component in the derivation of integral PBW bases.

The plan of this section is to adapt arguments in [Lus90a, Lus90b] and related references to infer PBW bases that generate the algebras U_q^\pm and U_q as free Λ_{q,e^*} and Λ_{q,e^*}^* modules respectively. Generators E_w depending on reduced words $w \in \mathcal{W}^*$ are constructed in Section 5.1 and

some of their basic properties discussed. From these monomial expressions that, in addition, depend on exponent functions are introduced, their relations with the two anti-involutions Ω and \mathfrak{U} are explained in Section 5.3. A formalism for the spanning sets associated to the monomials is laid out in Section 5.7, where a general criterion for order independence is derived.

The relations between the box pre-orderings introduced by Lusztig, convex orderings, and words representing the longest elements in \mathcal{W} are explained in Section 5.5. Lusztig's proof of PBW bases is adapted to the U_q -algebras in Section 5.6 and the general theorem for integral PBW bases for general orders is stated in Section 5.8.

5.1. Generators Associated to Words. In a series of articles, Lusztig defines of a generator \tilde{E}_α for each $\alpha \in \Phi^+$, starting with a presentation $\alpha = s_{i_1} \dots s_{i_{k-1}}(\alpha_{i_k})$ obtained from a maximal reduced word. The simple root α_j is replaced by E_j and each s_i by an automorphism T_i described in Section 4.3. We will adopt and adjust the definition here by replacing the T_i by their inverses Γ_i and explicitly retain the dependence on words in the notation. That is, for a non-empty reduced word $w = w_{i_1} \dots w_{i_k}$ and using the notation in (2.8) we set

$$E_w = \Gamma_{w^\flat}(E_{\tau(w)}) = \Gamma_{i_1} \dots \Gamma_{i_{k-1}}(E_{i_k}) \quad \text{and} \quad F_w = \Gamma_{w^\flat}(F_{\tau(w)}). \quad (5.1)$$

As remarked in Section 4.3, we may write $E_w = \Gamma_s(E_j)$ where $s = \tau(w^\flat)$ and $j = \tau(w)$, implying that E_w only depends on the pair (s, j) . Further redundancies in the labeling are implied by Proposition 5.1 below, from which one can easily obtain examples of distinct pairs $(s, j) \neq (s', j')$ with $\Gamma_s(E_j) = \Gamma_{s'}(E_{j'})$. For computations with specific words $w = w_{i_1} \dots w_{i_k}$ it is often convenient to also use the index notation

$$E_w = E_{(i_1 \dots i_k)}. \quad (5.2)$$

Given a *fixed* reduced word $z \in \mathcal{W}_m^*$ of *maximal* length or, equivalently, a convex order \leq_w on Φ^+ , we will sometimes also use the abbreviated notation

$$E_\alpha = E_{z[\alpha]} \quad \text{and} \quad F_\alpha = F_{z[\alpha]} \quad (5.3)$$

for an $\alpha \in \Phi^+$ and with $z[\alpha]$ as defined in (2.15). On occasion, we may also use the analogous notation for the respectively defined Cartan element $K_w = \Gamma_{w^\flat}(K_{\tau(w)}) = K^\nu$, where K^ν is as defined above (4.2) with $\nu = \gamma(w)$. For most applications in this article the word labeling is, however, the preferred one as we often consider orderings depending on choices of words. We list next several immediate properties of these generators.

Suppose $w = u \cdot v$ for non-empty reduced words $u, v \in \mathcal{W}^*$ with $l(w) = l(u) + l(v)$ and let $s = \tau(u)$. Then (5.1) and (4.22) imply

$$E_w = \Gamma_s(E_v) \quad \text{and} \quad F_w = \Gamma_s(F_v). \quad (5.4)$$

Furthermore, commutation of the Γ_i with Ω yields the identity

$$\Omega(E_w) = F_w \quad \text{and} \quad \Omega(F_w) = E_w. \quad (5.5)$$

The generators from (5.1) as well as the Cartan element K^ν are homogeneous elements, for which gradings follow immediately from (4.25) and (4.26) with notation from Lemma 2.2.

$$\mathbf{w}(E_w) = \gamma(w) \quad \mathbf{w}(F_w) = -\gamma(w) \quad \mathbf{w}(K_\nu) = 0 \quad (5.6)$$

$$\mathbf{o}^+(E_w) = \gamma(w) \pmod{2} \quad \mathbf{o}^+(F_w) = 0 \quad \mathbf{o}^+(K_\nu) = \nu \pmod{2} \quad (5.7)$$

Extending the definition in (5.1) to non-reduced words leads to elements outside the respective Borel subalgebra, such as $E_{w_i \cdot w_i} = \Gamma_i(E_i) \notin U_q^+$. The proposition below states that this does not happen for reduced words. The statement is proved for simply laced Lie types ADE in [Lus90a]. In our proof below, we extend the arguments from [Lus90a] to all Lie types and the regular power quantum algebras U_q^\pm over Λ_{q,e^*} . The second assertion on the uniqueness of generators graded by simple roots will be used in later computation of braid automorphisms.

Proposition 5.1. *Suppose $w \in \mathcal{W}^*$ is any non-empty reduced word. Then $E_w \in U_q^+$ and $F_w \in U_q^-$.*

Moreover, if $\gamma(w) = \alpha_k \in \Delta$, then $E_w = E_k$ and $F_w = F_k$.

Proof. We first provide the proof of the statement for a rank 2 subalgebra $U_q^{(i,j)}$ generated by $\{E_i, E_j, F_i, F_j, K_i, K_j\}$ for some fixed pair of indices $i \neq j$. The notations $U_q^{(i,j)\pm}$ and minimal ground rings are chosen analogous to those for U_q . Omitting the trivial $A_1 \times A_1$ case, we thus consider the Lie types A_2 , $B_2 = C_2$, and G_2 . As in Section 39.2 of [Lus93] assume that α_i is the shorter root. So, $A_{ji} = -1$ and $A_{ij} = -e$ with $e \in \{1, 2, 3\}$, as well as $d_i = 1$ and $d_j = e$.

The respective sets of reduced words $\mathcal{W}^{*(i,j)}$ are given by alternating sequences of w_i and w_j of length at most $m_{ij} \in \{3, 4, 6\}$ have been described in Section 2.7 and subject to (2.6).

Note that the automorphisms Γ_i from (4.18) through (4.21) coincide with the $T'_{i,-1}$ operators defined in Section 37.1.3 of [Lus93]. There, Lusztig computes the E_z explicitly for every non-empty reduced word $z \in \mathcal{W}^{*(i,j)}$ in Section 39.2.2 for $e = 3$, in Section 39.2.3 for $e = 2$, and in Section 39.2.4 for $e = 1$. The (a) sequences in each section contain all words z with $\tau(z) = j$, that is, ending in the long root generator, and the (b) sequences correspond to words z starting in $\tau(z) = i$.

For $e = 1$ or 2 , each E_z is expressed as an element $x_{1,m;-1}$ or $x'_{1,m;-1}$ in the subalgebra generated by E_i and E_j over $\Lambda_{q,m+1}$. In the $e = 3$ case, additional elements \hat{x} and \hat{x}' are included, which are easily seen to be elements of the subalgebra over $\Lambda_{q,4}$. Hence, each E_z is in $U_q^{(ij)^\pm}$ defined over Λ_{q,e^*} as desired.

For a word $w \in \mathcal{W}^{*(i,j)}$ of maximal length $l(w) = m_{ij}$, the formulae in Section 39.2 in [Lus93] imply the identity $E_w = E_k$ where $k \in \{i, j\}$ such that $\gamma(w) = \alpha_k$. So, for example, we have $\Gamma_i \Gamma_j(E_i) = E_j$ in the $e = 1$ case and $\Gamma_i \Gamma_j \Gamma_i(E_j) = E_j$ in the $e = 2$ case. The same relations hold for i and j exchanged, as well as analogous relations for $e = 3$ according to the computations in [Lus93]. The respective relations for the F_i generators are obtained by application of the Ω involution, completing the proof in the rank 2 case.

The proof of the statement for higher ranks extends the induction argument in the length $l(w)$ in the proof of Proposition 1.8 of [Lus90a] to all Lie types. Let $s = \tau(w^\flat)$ and $i = \tau(w)$ so that $s(\alpha_i) = \gamma(w) \in \Phi^+$ by Lemma 2.2 and $E_w = \Gamma_s(E_i)$. Also $l(s \cdot s_i) = l(w) = l(w^\flat) + 1 = l(s) + 1$.

Following Lemma 2.13 we next write $s = t \cdot r$ with $r \in \mathcal{W}_{(i,j)}$. Since $l(s) = l(t) + l(r)$ we may infer from (4.22) that $\Gamma_s(E_i) = \Gamma_t \circ \Gamma_r(E_i) = \Gamma_t(E_z)$ where $E_z = \Gamma_r(E_i)$. Here $z = y \cdot w_i$ with $y \in \mathcal{W}_{(i,j)}^*$ and $r = \tau(y)$ as in Lemma 2.13. Now $l(s) = l(t) + l(r)$ and $l(s \cdot s_i) = l(s) + 1$ also imply $l(r \cdot s_i) = l(r) + 1$ so that z is indeed a reduced word. The statement above for the rank 2 case now implies that $E_z \in U_q^{(i,j)+}$ and, thus, can be written as a Λ_{q,e^*} combination of monomials in E_i and E_j .

Let now $x \in \mathcal{W}^*$ be a reduced word presenting $t = \tau(x)$. With $t(\alpha_i), t(\alpha_j) \in \Phi^+$ implied by Lemma 2.13, we also know that $u_i = x \cdot w_i$ and $u_j = x \cdot w_j$ are reduced words. Since $l(r) \geq 1$ we have $l(t) < l(s)$ and, hence, $l(u_i), l(u_j) < l(w)$. Thus, by induction hypothesis, we have that $E_{u_i} = \Gamma_t(E_i)$, $E_{u_j} = \Gamma_t(E_j) \in U_q^+$. Since E_z is a combination of monomials in E_i and E_j we conclude that $E_w = \Gamma_t(E_z)$ is the same combination of monomials in E_{u_i} and E_{u_j} . In particular, $E_w \in U_q^+$ as desired.

Suppose, further, that $\gamma(w) = s(\alpha_i) = \alpha_k \in \Delta$. By Lemma 2.13 we then have $r \cdot s_i = s_{|i,j|}$, meaning that $z = y \cdot w_i \in \mathcal{W}_{m(i,j)}^*$ and $r(\alpha_i) \in \{\alpha_i, \alpha_j\}$. The rank 2 results above assure that $E_z = E_m$ where $m \in \{i, j\}$ and $r(\alpha_i) = \alpha_m$. Also, $t(\alpha_m) = t(r(\alpha_i)) = s(\alpha_i) = \alpha_k$. The induction hypothesis, therefore, implies that $\Gamma_t(E_m) = E_k$ so that also $\Gamma_s(E_i) = \Gamma_t(\Gamma_r(E_i)) = \Gamma_t(E_m) = E_k$ as claimed.

The respective statements for the F_i follow by applications of the Ω involution. \square

An important special case of Proposition 5.1 are words $z \in \mathcal{W}_m^*$ of maximal length. If $j = \tau(z)$, recall from (2.22) that $q_j = \tau(z^\flat) = s_\bullet \cdot s_j = s_{\eta(j)} \cdot s_\bullet$, where η is the involution of the Dynkin diagrams as defined at the start of Section 2.4. As noted in the proof of Lemma 2.8, this yields $\gamma(z) = q_j(\alpha_j) = \alpha_{\eta(j)}$. Thus,

$$E_z = E_{\eta(\tau(z))} \quad \text{for all } z \in \mathcal{W}_m^*. \quad (5.8)$$

The next formulae summarize notations for two relevant renormalizations of a generator E_w for some word $w \in \mathcal{W}^*$ with $\alpha = \gamma(w)$. The quantities d_α , q_α , and $[n]_\alpha$ are as defined in (3.1) and $k \in \mathbb{N}$.

$$E_w^{(k)} = \frac{1}{[k]!_\alpha} E_w^k \quad (5.9)$$

$$\check{E}_w = (q_\alpha^{-1} - q_\alpha) E_w = (q^{-1} - q) [d_\alpha] E_w \quad (5.10)$$

The first extends Lusztig's *divided power generators*, already mentioned in Section 3.2 for simple roots, to all positive roots and *a priori* requires the ground ring to contain the reciprocals of all quantum numbers, such as for $\Lambda_{q,\infty}$ or $\mathbb{Q}(q)$.

The singularized generators \check{E}_w coincide, for appropriate ordering and up to signs, with the generators \overline{E}_β introduced in [DCP93, Sec 12.1]. In calculations, this renormalization often removes the $(q - q^{-1})$ denominators, such as in (3.13), and avoids the use of the extended ground rings such as Λ_q^* or $\Lambda_{q,n}^*$.

5.2. Garside Automorphisms and Dictionary between Generators. In this section, we aim to provide precise correspondences between Lusztig's and our definition of generators, which entails various useful identities. For a reduced word $w = w_{i_1} \dots w_{i_k}$, the T_i -analog of (5.1) is defined as

$$\check{E}_w = T_{w^\flat}(E_{\tau(w)}) = T_{i_1} \dots T_{i_{k-1}}(E_{i_k}), \quad (5.11)$$

with the analogous formula for \check{F}_w . As in (5.3), for a positive root $\alpha \in \Phi^+$, we set $\check{E}_\alpha = \check{E}_{z[\alpha]}$, matching the definition of generators in [Lus90a, Lus90b] for an adequately chosen reduced word $z \in \mathcal{W}_m^*$ of maximal length. They can be related to our convention using the \mathfrak{U} involution as well as the notion of complementary words defined in Section 2.4.

Proposition 5.2. *For non-empty reduced words $v, w \in \mathcal{W}^*$, assume that $w \rightsquigarrow v$. Then*

$$\check{E}_w = \mathfrak{U}(E_w) = E_v \quad \text{and} \quad \check{F}_w = \mathfrak{U}(F_w) = F_v.$$

Proof. The first equality is immediate from (4.23) expressed as $\mathfrak{U} \circ \Gamma_{i_1} \circ \dots \circ \Gamma_{i_{k-1}} = T_{i_1} \circ \dots \circ T_{i_{k-1}} \circ \mathfrak{U}$. For the second equality set $s = \tau(v^\flat)$, $t = \tau(w^\flat)$, and $j = \tau(v)$. As in (2.22) we have $t^{-1} \cdot s = q_j$ with $l(q_j) = l(t^{-1}) + l(s)$ so $\Gamma_{q_j} = \Gamma_{t^{-1}} \circ \Gamma_s$ by (4.22). With $\Gamma_{q_j}(E_j) = E_{\eta(j)}$ implied by (5.8) as well as (4.23), we thus compute $\mathfrak{U}(E_w) = \mathfrak{U}(\Gamma_t(E_{\eta(j)})) = \Gamma_{t^{-1}}^{-1}(\mathfrak{U}(E_{\eta(j)})) = \Gamma_{t^{-1}}^{-1}(E_{\eta(j)}) = \Gamma_{t^{-1}}^{-1}(\Gamma_{q_j}(E_j)) = \Gamma_s(E_j) = E_v$. The assertion for F_w follows again by application of Ω , which commutes with \mathfrak{U} . \square

As a second application of Proposition 5.1 we are able to explicitly compute the Lusztig automorphism Γ_∂ associated to the Garside element $\partial = \mathfrak{z}(s_\bullet) \in \mathcal{A}^+$ as in (2.7). In our conventions as explained in Section 4.3, we may also write $\Gamma_{s_\bullet} = \Gamma_\partial$, where $s_\bullet \in \mathcal{W}$ is the longest element.

The relation also involves the automorphism $\mathcal{I} : U_q \rightarrow U_q$ that implements the involution η on Dynkin diagrams defined at the beginning of Section 2.4. It is given on generators by

$$\mathcal{I}(E_i) = E_{\eta(i)}, \quad \mathcal{I}(F_i) = F_{\eta(i)}, \quad \mathcal{I}(K_i) = K_{\eta(i)}. \quad (5.12)$$

The maps $\iota, \varpi \in \text{Hom}(\mathbb{Z}^\Delta, \Lambda^*)$ are as in (4.9) and (4.16) and the remaining automorphisms are as defined in Section 4.

Corollary 5.3. *The Garside automorphism and its square are given by the following composites of elementary automorphisms:*

$$\Gamma_\partial = \mathcal{I} \circ \Pi \circ \Psi_\iota^{id} = \mathcal{I} \circ \Pi \circ \mathfrak{U} \circ S \quad \text{and} \quad \Gamma_\partial^2 = \Psi_{\varpi^{-1}}^{2id}. \quad (5.13)$$

Proof. As explained above, we express $\Gamma_{s_\bullet} = \Gamma_\partial$ in terms of the longest Weyl element. With the same notation as in (4.23) or the proof of Proposition 5.2 we have $s_\bullet = s_{\eta(i)} \cdot q_i$ with $l(s_\bullet) = l(q_i) + 1$

and $\Gamma_{q_i}(E_i) = E_{\eta(i)}$ for given $i \in \{1, \dots, n\}$. Hence,

$$\begin{aligned}\Gamma_{s_*}(E_i) &= \Gamma_{\eta(i)}(\Gamma_{q_i}(E_i)) = \Gamma_{\eta(i)}(E_{\eta(i)}) = -K_{\eta(i)}^{-1}F_{\eta(i)} = \\ &= \mathcal{I}(-K_i^{-1}F_i)) = \mathcal{I} \circ \Pi(-K_i E_i)) = \mathcal{I} \circ \Pi \circ \Upsilon_i^{id}(E_i),\end{aligned}\tag{5.14}$$

which proves the first identity (5.13) on E_i . The verification for F_i follows, as usual, by application of Ω to (5.14), and the automorphisms on both ends of (5.14) clearly map K^\vee to $K^{-\eta(\vee)}$, where η is the obvious extension to \mathbb{Z}^Δ . The formula for the square of the Garside automorphism is readily derived from the first identity and the commutation relations in (4.15). \square

Aside from \mathfrak{U} and Ω introduced in Section 4.2 we also record here the action of the conjugation automorphism Υ on the generators. The formula contains an additional factor in Λ^\star , which only depends on the root associated with the word label.

Lemma 5.4. *Let $w \in \mathcal{W}^*$ be a non-empty, reduced word and $\beta = \gamma(w)$ the associated root. Then*

$$\Upsilon(E_w) = (-1)^{\text{ht}(\beta)-1} q^{(\rho|\beta)-d_\beta} E_w \quad \text{and} \quad \Upsilon(F_w) = (-1)^{\text{ht}(\beta)-1} q^{-(\rho|\beta)+d_\beta} F_w. \tag{5.15}$$

Proof. Let $s = \gamma(w^\beta)$ and $i = \tau(w)$ so that $E_w = \Gamma_s(E_i)$ and $\beta = s(\alpha_i)$. Relation (4.33) implies $\Upsilon(E_w) = \Upsilon(\Gamma_s(E_i)) = (\Upsilon_{\kappa_s}^0)^{-1}(\Gamma_s(\Upsilon(E_i))) = (\kappa_s(\beta))^{-1}\Gamma_s(E_i) = (-1)^{(\check{\theta}(s)|\beta)}q^{(\theta(s)|\beta)}E_w$. Now, by (2.27), $(\theta(s)|\beta) = (\rho - s(\rho)|\beta) = (\rho|\beta) - (s(\rho)|s(\alpha_i)) = (\rho|\beta) - (\rho|\alpha_i) = (\rho|\beta) - d_i = (\rho|\beta) - d_\beta$ since d_β is \mathcal{W} -invariant. Also, (2.28) implies $(\check{\theta}(s)|\beta) = \text{ht}(\beta) - \text{ht}(\alpha_i) = \text{ht}(\beta) - 1$. The relation for F_w follows by application of Ω and (5.5). \square

In the remainder of this section, we discuss the generators and actions more explicitly in the rank 2 case. As in the proof of Proposition 5.1, let $U_{q^{(i,j)}}^+$ be the subalgebra generated by E_i and E_j for two indices i and j . We assume $A_{ji} = -1$, implying $d_i = 1$ and $-A_{ij} = d_j = \mathbf{e} \in \{1, 2, 3\}$, corresponding to Lie type A_2 , B_2 , or G_2 , respectively, for $U_{q^{(i,j)}}^+$. As before, let $m = m_{ij} \in \{3, 4, 6\}$ be the length of the longest element in $\mathcal{W}_{(i,j)}$.

For $r \in \{1, \dots, m\}$ let $a_r = w_i w_j \dots$ and $b_r = w_j w_i \dots$ be the two possible reduced words of length r in $\mathcal{W}_{(i,j)}^*$ as in (2.32). Combining Lemma 2.12 and Proposition 5.2 now yields the identities

$$\mathfrak{U}(E_{a_r}) = E_{b_{m-r+1}} = \tilde{E}_{a_r} \quad \text{and} \quad \mathfrak{U}(F_{a_r}) = F_{b_{m-r+1}} = \tilde{F}_{a_r}. \tag{5.16}$$

Since later proofs will refer to and invoke several commutation relations in [Lus90b], we provide here a dictionary from the notation used there for generators in a rank 2 system to our conventions. Specifically, in Section 5.1 of [Lus90b] Lusztig uses subscripts of the form 1 ... 12 ... 2 for the E -generators. Replacing $i \leftrightharpoons 1$ to index the short root and $j \leftrightharpoons 2$ for the (possibly) long root, the definition there can be summarized in the formula

$$\tilde{E}_{\underbrace{i \dots i}_{k} \underbrace{j \dots j}_{p}} = \tilde{E}_{z_{|ji|}[\alpha]} \quad \text{with} \quad \alpha = k\alpha_i + p\alpha_j \in \Phi^{+(ij)}. \tag{5.17}$$

That is, the frequency of i 's and j 's determined the coefficients of the root α in the root sublattice. This is identified with the unique word $b_r \leq_R z_{|ji|}$ as in (2.32) for which $\alpha = \gamma(b_r)$, with r

determined by the position of the root in the ordering in (2.33). The respective E -generator in our convention is then obtained via (5.16), which, in turn, can be expressed in the notation from (5.2).

For example, in the G_2 ($e = 3, m = 6$) case, \tilde{E}_{iiijj} indicates in [Lus90b] the generator associated to root $3\alpha_i + 2\alpha_j$. In the ordering for $z_{|j|}$ in (2.33) it appears in position $r = 3$ so that $\tilde{E}_{iiijj} = \tilde{E}_{b_3} = \tilde{E}_{w_j w_i w_j} = \tilde{E}_{(jij)} = T_j T_i(E_j)$. From (5.16) we next obtain that $\tilde{E}_{b_3} = E_{a_{6+1-3}} = E_{a_4} = E_{w_i w_j w_i w_j} = E_{(ijij)} = \Gamma_i \Gamma_j \Gamma_i(E_j)$. Similarly, the generators associated to simple roots are readily identified as

$$\tilde{E}_j = \tilde{E}_{w_j} = E_{z_{|j|}} = E_{\underbrace{(ij\dots)}_m} = E_j \quad \text{and} \quad \tilde{E}_i = \tilde{E}_{z_{|j|}} = \tilde{E}_{\underbrace{(ji\dots)}_m} = E_{(i)} = E_i. \quad (5.18)$$

The associated simple roots appear either as maximal or minimal elements in the two total orders on $\Phi^+_{(ij)}$.

The full dictionary translating between the notation in [Lus90b] and our conventions is as follows. Here, generators are ordered by the roots and $\prec_{z_{|j|}}$ as in (2.33), omitting the first and last terms given in (5.18) for even m . For $m = 3$ the identity is $\tilde{E}_i = E_i = E_{(iji)}$.

$$\begin{aligned} e = 1 \quad & \left\{ \begin{array}{l} \tilde{E}_{ij} = \tilde{E}_{(ji)} = E_{(ij)} \end{array} \right. \\ e = 2 \quad & \left\{ \begin{array}{l} \tilde{E}_{ij} = \tilde{E}_{(ji)} = E_{(iji)}, \quad \tilde{E}_{iij} = \tilde{E}_{(jij)} = E_{(ij)} \end{array} \right. \\ e = 3 \quad & \left\{ \begin{array}{l} \tilde{E}_{ij} = \tilde{E}_{(ji)} = E_{(ijiji)}, \quad \tilde{E}_{iiij} = \tilde{E}_{(jijij)} = E_{(ijij)}, \\ \tilde{E}_{ijj} = \tilde{E}_{(jijj)} = E_{(iji)}, \quad \tilde{E}_{iiij} = \tilde{E}_{(jijij)} = E_{(ij)} \end{array} \right. \end{aligned} \quad (5.19)$$

5.3. Monomial Expressions, Exponent Sets, and Their Spans. The PBW type bases of interest for us are given by monomials whose exponents are labeled by positive roots. To describe these more formally, we introduce *exponents sets*

$$\mathcal{E}(s) = \text{Maps}(\mathcal{N}(s), \mathbb{N}_0) = \mathbb{N}_0^{\mathcal{N}(s)} \quad \text{and} \quad \mathcal{E}^*(s) = \text{Maps}(\mathcal{N}(s), \mathbb{Z}) = \mathbb{Z}^{\mathcal{N}(s)}, \quad (5.20)$$

where the inversion sets $\mathcal{N}(s)$ are as in (2.9). Additionally, we use the more compact notation for the maximal sets

$$\mathcal{E}_\bullet = \mathcal{E}(s_\bullet) = \text{Maps}(\Phi^+, \mathbb{N}_0) = \mathbb{N}_0^{\Phi^+} \quad \text{and} \quad \mathcal{E}_\bullet^* = \mathcal{E}^*(s_\bullet) = \text{Maps}(\Phi^+, \mathbb{Z}). \quad (5.21)$$

Suppose $s = t_1 \cdot t_2$ with $l(s) = l(t_1) + l(t_2)$. Recall from Corollary 2.4 that $\mathcal{N}(s)$ is then the disjoint union of $\mathcal{N}(t_1)$ and $t_1(\mathcal{N}(t_2))$, which, correspondingly, implies the following bijections

of exponent sets:

$$\begin{aligned} \mathcal{E}(t_1) \times \mathcal{E}(t_2) \rightarrow \mathcal{E}(s) &: (\psi_1, \psi_2) \mapsto \psi = \psi_1 \sqcup (t_1^{-1})^*(\psi_2), \\ \mathcal{E}(s) \rightarrow \mathcal{E}(t_1) \times \mathcal{E}(t_2) &: \psi \mapsto (i_1^*\psi, i_2^*t_1^*(\psi)). \end{aligned} \quad (5.22)$$

Here, $s^*(\psi) = \psi \circ s$ denotes the usual pull-back, the disjoint union of maps indicates the respective definition on the two sets, and i_i^* is the restriction to $\mathcal{N}(s_i)$.

Suppose $w \in \mathcal{W}^*$, $s = \tau(w) \in \mathcal{W}$ with $L = l(w) = |\mathcal{N}(s)|$, and $\psi \in \mathcal{E}(s)$. Following (2.15) and the explanations in Section 2.3, we write $\beta_j = \gamma(w[1, j])$ for the ordered root sequence implied by w as well as $w^j = w[1, j] = w[\beta_j]$ for the words $\leq_R w$. Given these conventions, we define the following monomials, expressed in two equivalent notations and for each of two opposing directions of multiplication.

$$\begin{aligned} E_{w\blacktriangleright}^\psi &= E_{w^1}^{\psi(\gamma(w^1))} \dots E_{w^L}^{\psi(\gamma(w^L))} = E_{w[\beta_1]}^{\psi(\beta_1)} \dots E_{w[\beta_L]}^{\psi(\beta_L)} \\ E_{\blacktriangleleft w}^\psi &= E_{w^L}^{\psi(\gamma(w^L))} \dots E_{w^1}^{\psi(\gamma(w^1))} = E_{w[\beta_L]}^{\psi(\beta_L)} \dots E_{w[\beta_1]}^{\psi(\beta_1)} \end{aligned} \quad (5.23)$$

Generators $F_{w\blacktriangleright}^\psi$ and $F_{\blacktriangleleft w}^\psi$ are defined by the same formulae (5.23) with E replaced by F . Using that Ω is an anti-homomorphism we find from (5.5) the direction reversing relations

$$\Omega(E_{w\blacktriangleright}^\psi) = F_{\blacktriangleleft w}^\psi \quad \text{and} \quad \Omega(F_{w\blacktriangleright}^\psi) = E_{\blacktriangleleft w}^\psi. \quad (5.24)$$

Similarly, from Lemma 5.4 we find that the conjugation automorphism reverses the direction for the same type of monomial up to a unit factor. Specifically, if we denote $\vec{\psi} = \sum_\alpha \psi(\alpha) \alpha \in \mathbb{Z}^\Delta$, $d(\psi) = \sum_\alpha \psi(\alpha) d_\alpha \in \mathbb{N}_0$, and $n(\psi) = \sum_\alpha \psi(\alpha) \in \mathbb{N}_0$ the following formulae hold for the Υ anti-involution.

$$\begin{aligned} \Upsilon(E_{w\blacktriangleright}^\psi) &= (-1)^{\text{ht}(\vec{\psi}) - n(\psi)} q^{(\rho|\vec{\psi}) - d(\psi)} E_{\blacktriangleleft w}^\psi \\ \Upsilon(F_{w\blacktriangleright}^\psi) &= (-1)^{\text{ht}(\vec{\psi}) - n(\psi)} q^{-(\rho|\vec{\psi}) + d(\psi)} F_{\blacktriangleleft w}^\psi \end{aligned} \quad (5.25)$$

Suppose $w = u_1 \cdot u_2 \in \mathcal{W}^*$ with $l(w) = l(u_1) + l(u_2)$ and $s = \tau(w)$ and $t_i = \tau(u_i)$ for $i = 1, 2$. Moreover, assume that $\psi \in \mathcal{E}(s)$ is assigned to (ψ_1, ψ_2) in the correspondence in (5.22). Then the relation from (5.4) applied to each term of the monomial and the fact that the Γ_i are homomorphisms implies the following recursions for monomials:

$$E_{w\blacktriangleright}^\psi = E_{u_1\blacktriangleright}^{\psi_1} \cdot \Gamma_{t_1}(E_{u_2\blacktriangleright}^{\psi_2}) \quad \text{and} \quad E_{\blacktriangleleft w}^\psi = \Gamma_{t_1}(E_{\blacktriangleleft u_2}^{\psi_2}) \cdot E_{\blacktriangleleft u_1}^{\psi_1}. \quad (5.26)$$

Analogous monomial expressions $\tilde{E}_{w\blacktriangleright}^\psi$ and $\tilde{E}_{\blacktriangleleft w}^\psi$ are defined by replacing the $E_{w^j}^{\psi(\beta_j)}$ factors in (5.23) by Lusztig's $\tilde{E}_{w^j}^{\psi(\beta_j)}$ generators, with \tilde{E}_w as defined in (5.11). The next proposition extends the correspondence in Proposition 5.2 to these expressions.

Proposition 5.5. *For a non-empty $w \in \mathcal{W}^*$, let $t = \tau(w) \in \mathcal{W}$ and $r = t^\perp$ as in (2.20). Moreover, let $\psi \in \mathcal{E}(t)$ and $\phi = r_*\psi = \psi \circ r \in \mathcal{E}(t^\dagger)$. Then*

$$\mathfrak{O}(E_{w\blacktriangleright}^\psi) = \tilde{E}_{\blacktriangleleft w}^\psi = \Gamma_r \left(E_{w^\dagger\blacktriangleright}^\phi \right). \quad (5.27)$$

The analogous relation holds with directions flipped in all terms as well as with E generators replaced by F generators.

Proof. As noted in Section 2.2, we can find $u \in \mathcal{W}^*$ such that $z = w \cdot u$ is a reduced word of maximal length. So, with $N = l(z)$ and $L = l(w)$ we have $l(u) = N - L$. Moreover, $g = \nu(u) = t^{-1} \cdot s_* = s_* \cdot \eta(t)^{-1} = s_* \cdot t^\dagger$ and, hence, $\nu(u^\dagger) = g^\dagger = t \cdot s_* = t^\dagger = r$. Now, Lemma 2.7vi) implies $r(\mathcal{N}(t^\dagger)) = t^\dagger(\mathcal{N}(t^\dagger)) = \mathcal{N}(t)$ so that, indeed, r_* maps $\mathcal{E}(t)$ to $\mathcal{E}(t^\dagger)$.

Assume that $j \leq L$, and let $y = z^\dagger = u^\dagger \cdot w^\dagger$. Then we have $z[1, j] = w[1, j]$ and $y[1, N+1-j] = u^\dagger \cdot (w^\dagger[1, L+1-j])$. Thus, Lemma 2.8 implies $w[1, j] \rightsquigarrow u^\dagger \cdot (w^\dagger[1, L+1-j])$, which entails together with Proposition 5.2 and (5.4) that $\mathfrak{O}(E_{w[1,j]}) = \tilde{E}_{w[1,j]} = E_{u^\dagger \cdot (w^\dagger[1, L+1-j])} = \Gamma_{u^\dagger}(E_{w^\dagger[1, L+1-j]}) = \Gamma_r(E_{w^\dagger[1, L+1-j]})$. From Lemma 2.8 we also find that $\beta_j = \gamma(w[1, j]) = \gamma(u^\dagger \cdot (w^\dagger[1, L+1-j])) = r(\delta_{L+1-j})$ where $\delta_i = \gamma(w^\dagger[1, i])$. Thus,

$$\begin{aligned} \mathfrak{O}(E_{w^\dagger}^\psi) &= \mathfrak{O}\left(E_{w[1,1]}^{\psi(\beta_1)} \cdots E_{w[1,L]}^{\psi(\beta_L)}\right) &= \mathfrak{O}\left(E_{w[1,L]}^{\psi(\beta_L)}\right) \cdots \mathfrak{O}\left(E_{w[1,1]}^{\psi(\beta_1)}\right) & (5.28) \\ &= \Gamma_r\left(E_{w^\dagger[1,1]}^{\psi(\beta_L)}\right) \cdots \Gamma_r\left(E_{w^\dagger[1,L]}^{\psi(\beta_1)}\right) &= \Gamma_r\left(E_{w^\dagger[1,1]}^{\psi(r(\delta_1))}\right) \cdots \Gamma_r\left(E_{w^\dagger[1,L]}^{\psi(r(\delta_L))}\right) \\ &= \Gamma_r\left(E_{w^\dagger[1,1]}^{\phi(\delta_1)} \cdots E_{w^\dagger[1,L]}^{\phi(\delta_L)}\right) &= \Gamma_r(E_{w^\dagger}^\phi), \end{aligned}$$

which shows equality of the first and third term in (5.27). Furthermore, the last expression in the first line of (5.28) is readily identified with $\tilde{E}_{w[1,L]}^{\psi(\beta_L)} \cdots \tilde{E}_{w[1,1]}^{\psi(\beta_1)} = \tilde{E}_{w^\dagger}^\psi$. Identities for reversed directions follow analogously. The respective statements for F generators are obtained by application of Ω , using (5.24) as well as commutation with both \mathfrak{O} and Γ_t . \square

We will use this result mainly in the case of reduced words of maximal lengths, for which $t = s_*$ and $r = \text{id}$. That is, let $z \in \mathcal{W}_m^*$ and $\psi \in \mathcal{E}_*$ as in (5.21). Then (5.27) specializes to

$$\mathfrak{O}(E_{z^\dagger}^\psi) = \tilde{E}_{\mathfrak{z}^\dagger}^\psi = E_{z^\dagger}^\psi. \quad (5.29)$$

The respective identity for the opposite directions is readily derived from commutation of \mathfrak{O} with Υ . The variants with E 's replaced by F 's follows from commutation with Ω .

Further relations that directly relate (maximal) monomial expressions in the T and Γ conventions are provided by the composite automorphism $\Pi = \Upsilon \circ \mathfrak{O}$. For example, for z and ψ as above we find

$$\Pi(E_{z^\dagger}^\psi) = (-1)^{\text{ht}(\vec{\psi}) - n(\psi)} q^{(\rho|\vec{\psi}) - d(\psi)} \tilde{E}_{z^\dagger}^\psi = (-1)^{\text{ht}(\vec{\psi}) - n(\psi)} q^{(\rho|\vec{\psi}) - d(\psi)} E_{\mathfrak{z}^\dagger}^\psi \quad (5.30)$$

using the notation from (5.25). That is, up to unit factors, Π maps monomials in our convention to those in Lusztig's convention for the same choices of words and directions. Similarly, formulae for the antipode can be derived from those in (4.16), (5.29), and Proposition 4.1.

$$\begin{aligned} S(E_{z^\dagger}^\psi) &= (-1)^{\text{ht}(\vec{\psi})} q^{-\frac{1}{2}((\vec{\psi}|\vec{\psi}) - 2(\rho|\vec{\psi}))} E_{z^\dagger}^\psi K^{-\vec{\psi}} &= (-1)^{\text{ht}(\vec{\psi})} q^{\frac{1}{2}((\vec{\psi}|\vec{\psi}) + 2(\rho|\vec{\psi}))} K^{-\vec{\psi}} E_{z^\dagger}^\psi & (5.31) \\ S(F_{z^\dagger}^\psi) &= (-1)^{\text{ht}(\vec{\psi})} q^{-\frac{1}{2}((\vec{\psi}|\vec{\psi}) + 2(\rho|\vec{\psi}))} F_{z^\dagger}^\psi K^{\vec{\psi}} &= (-1)^{\text{ht}(\vec{\psi})} q^{\frac{1}{2}((\vec{\psi}|\vec{\psi}) - 2(\rho|\vec{\psi}))} K^{\vec{\psi}} F_{z^\dagger}^\psi \end{aligned}$$

The same equations hold for the basis elements with respect to the opposite directions, given the same is true for those in (5.29). For later reference, we also record the rank 2 case discussed in Section 2.7, for which $z_{|ij|}$ and $z_{|ji|} = z_{|ij|}^\dagger$ are maximal words:

$$\mathfrak{U}(E_{z_{|ij|}\blacktriangleright}^\psi) = \tilde{E}_{\blacktriangleleft z_{|ij|}}^\psi = E_{z_{|ji|}\blacktriangleright}^\psi. \quad (5.32)$$

Finally, we introduce notation for basis elements in $U_{q,\mathbb{Q}}$ given by divided power generators. Recall from (5.9) that for $k \in \mathbb{N}$ and $i = \tau(w)$ we have $E_w^{(k)} = \frac{1}{[k]!_i} E_w^k$. Given $w \in \mathcal{W}^*$, $L = l(w)$, $s = \tau(w) \in \mathcal{W}$, and $\psi \in \mathcal{E}(s)$, set, as before, $w^j = w[1, j]$, $\beta_j = \gamma(w^j)$, and $k_j = \psi(\beta_j)$. Set

$$E_{w\blacktriangleright}^{(\psi)} = E_{w^1}^{(k_1)} \cdots E_{w^L}^{(k_L)} = \frac{1}{[\psi]!} E_{w\blacktriangleright}^\psi \quad \text{and} \quad E_{\blacktriangleleft w}^{(\psi)} = E_{w^L}^{(k_L)} \cdots E_{w^1}^{(k_1)} = \frac{1}{[\psi]!} E_{\blacktriangleleft w}^\psi$$

(5.33)

where $[\psi]! = \prod_{j=1}^L [k_j]!_{d_j} = \prod_{\alpha \in \Phi^+} [\psi(\alpha)]!_\alpha$ with $d_j = \frac{1}{2} (\beta_j | \beta_j)$.

Here we wrote $[\psi]!$ as a product over Φ^+ rather than $\mathcal{N}(s)$, tacitly extending the exponent function as $\psi(\gamma) = 0$ for $\gamma \notin \mathcal{N}(s)$. With the same convention we introduce analogous notation for the singularized generators from (5.10) as

$$\check{E}_{w\blacktriangleright}^\psi = \check{E}_{w^1}^{k_1} \cdots \check{E}_{w^L}^{k_L} = \delta^\psi E_{w\blacktriangleright}^\psi \quad \text{and} \quad \check{E}_{\blacktriangleleft w}^\psi = \check{E}_{w^L}^{k_L} \cdots \check{E}_{w^1}^{k_1} = \delta^\psi E_{\blacktriangleleft w}^\psi$$

(5.34)

where $\delta^\psi = \delta^\psi(q) = \prod_{\alpha \in \Phi^+} (q_\alpha^{-1} - q_\alpha)^{\psi(\alpha)} = (q^{-1} - q)^{n(\psi)} \prod_{\alpha \in \Phi^+} [d_\alpha]^{\psi(\alpha)}$.

As usual, respective elements in U_q^- are defined by replacing all E 's by F 's. See also again [DCP93, Sec 12.1] for a variant of the $\check{E}_{w\blacktriangleright}^\psi$.

We conclude this section with notation for subsets of generators and their spans determined by subsets of exponent functions. Suppose $w \in \mathcal{W}^*$ with $s = \tau(w)$. We introduce the monomial spanning sets corresponding to a subset $S \subseteq \mathcal{E}(s)$ of exponents as follows:

$$\mathfrak{B}_{w\blacktriangleright}^{S,+} = \{E_{w\blacktriangleright}^\psi : \psi \in S\} \quad \text{and} \quad \mathfrak{B}_{\blacktriangleleft w}^{S,+} = \{E_{\blacktriangleleft w}^\psi : \psi \in S\}. \quad (5.35)$$

The sets $\mathfrak{B}_{w\blacktriangleright}^{S,-}$ and $\mathfrak{B}_{\blacktriangleleft w}^{S,-}$ are defined analogously with E 's replaced by F 's. The relation (5.24) then translates to the relations of sets

$$\mathfrak{B}_{w\blacktriangleright}^{S,-} = \Omega(\mathfrak{B}_{\blacktriangleleft w}^{S,+}) \quad \text{and} \quad \mathfrak{B}_{\blacktriangleleft w}^{S,-} = \Omega(\mathfrak{B}_{w\blacktriangleright}^{S,+}). \quad (5.36)$$

In the case that $S = \mathcal{E}(s)$, that is, if all exponent functions are included, we use the shorthand $\mathfrak{B}_{w\blacktriangleright}^{\pm} = \mathfrak{B}_{w\blacktriangleright}^{\mathcal{E}(s), \pm}$ and $\mathfrak{B}_{\blacktriangleleft w}^{\pm} = \mathfrak{B}_{\blacktriangleleft w}^{\mathcal{E}(s), \pm}$.

Recursion relations derived from (5.26) exist only in the special case when the exponent set S splits into a respective Cartesian product. More precisely, suppose $w = u_1 \cdot u_2$ and $s = t_1 \cdot t_2$ with $t_i = \tau(u_i)$ and $l(s) = l(t_1) + l(t_2)$. Assume that there are sets $S_i \subseteq \mathcal{E}(t_i)$ such that the bijection from (5.22) restricts to a bijection $S \rightarrow S_1 \times S_2$. In this case, (5.26) indeed implies

$$\mathfrak{B}_{w\blacktriangleright}^{S,+} = \mathfrak{B}_{u_1\blacktriangleright}^{S_1,+} \cdot \Gamma_{t_1}(\mathfrak{B}_{u_2\blacktriangleright}^{S_2,+}) \quad \text{and} \quad \mathfrak{B}_{\blacktriangleleft w}^{S,+} = \Gamma_{t_1}(\mathfrak{B}_{\blacktriangleleft u_2}^{S_2,+}) \cdot \mathfrak{B}_{\blacktriangleleft u_1}^{S_1,+}. \quad (5.37)$$

The above splitting property $S \cong S_1 \times S_2$ may be rephrased as the condition that $\psi \in S$ if and only if $i_1^*(\psi) \in i_1^*(S)$ and $i_2^*(i_1^*(\psi)) \in i_2^*(i_1^*(S))$, independently. Although very restrictive in general, the property applies to several useful situations below.

Assume now that Λ is an integral domain with a ring homomorphism $f: \Lambda_{q,\mathbf{e}^*} \rightarrow \Lambda$, and set $U_{q,\Lambda}^\pm = U_q^\pm \otimes_f \Lambda$ to be the respective quantum algebra over Λ . We denote the respective Λ -spans of the above sets in $U_{q,\Lambda}^\pm$ by

$$\mathfrak{S}_{w\blacktriangleright}^{S,\pm} = \langle \mathfrak{B}_{w\blacktriangleright}^{S,\pm} \rangle_\Lambda \quad \text{and} \quad \mathfrak{S}_{w\blacktriangleleft}^{S,\pm} = \langle \mathfrak{B}_{w\blacktriangleleft}^{S,\pm} \rangle_\Lambda. \quad (5.38)$$

As before, we write $\mathfrak{S}_{w\blacktriangleright}^\pm$ or $\mathfrak{S}_{w\blacktriangleleft}^\pm$ if $S = \mathcal{E}(s)$. Additional identities for the action of \mathfrak{O} on either basis sets or spans, such as $\mathfrak{O}(\mathfrak{B}_{w\blacktriangleright}^{S,+}) = \Gamma_r(\mathfrak{B}_{w\blacktriangleright}^{r^*(S),+})$, follow from Proposition 5.5.

5.4. Some Commutation Relations. We collect next various commutation relations among the E_w generators that are useful in establishing integral PBW bases and also support later computations of ideal properties. The most basic commutation relation for a pair of generators (A, B) is that they *skew-commute*, meaning that $BA = zAB$ where z is a unit in the ground ring. An important skew-commutation relation among the E_w generators is given by the following.

Lemma 5.6. *Suppose $w \in \mathcal{W}^*$ with reduced decomposition $w = u \cdot v$ for non-empty $u, v \in \mathcal{W}^*$. Denote $j = \tau(u)$, $\alpha = \gamma(u)$, and $\beta = \gamma(w)$. Assume further that w_j does not occur as a letter in v . Then*

$$E_w E_u = q^{(\alpha|\beta)} E_u E_w. \quad (5.39)$$

Proof. Since w_j is not a letter in v , E_v is an expression in $\{E_i\}_{i \neq j}$ and, hence, commutes with F_j . Also, by (5.6) and (4.2) we find $E_v K_j = q^{-(v|\alpha_j)} K_j E_v$ where $v = \gamma(v)$. Observe further that by (5.26) $E_w = \Gamma_u(E_v)$ and $E_u = \Gamma_{u^\flat}(E_j) = \Gamma_u(\Gamma_j^{-1}(E_j)) = \Gamma_u(-F_j K_j)$. Thus $E_w E_u = \Gamma_u(E_v(-F_j K_j)) = q^{-(v|\alpha_j)} \Gamma_u((-F_j K_j) E_v) = q^{-(\alpha_j|v)} E_u E_w$. Now, let $s = \tau(u)$ and $t = \tau(u^\flat)$ so that $s = t \cdot s_j$. Then $s(\alpha_j) = t(s_j(\alpha_j)) = -t(\alpha_j) = -\gamma(u) = -\alpha$. Moreover, we have $\beta = \gamma(u \cdot v) = s(\gamma(v)) = s(v)$. \mathcal{W} -invariance of the form now implies $(\alpha_j|v) = -(\alpha|\beta)$, from which we infer the assertion. \square

The most basic case is given by $v = w_k$, when $l(w) = l(u) + 1$ already implies the requirement $k \neq j$ of the lemma. It encompasses, for example, 2.3.(d3,d4) in [Lus90a] and several skew-commutation relations in Section 5.2 of [Lus90a]. The same references also contain strict commutation relations $E_\alpha E_\beta = E_\beta E_\alpha$ in special situations when $\alpha + \beta \notin \Phi^+$ and, hence, $(\alpha|\beta) = 0$. We will see, however, in the proof of Proposition 5.11 below that for non-simply laced Lie type B we may have weakly orthogonal pairs of roots $\alpha, \beta \in \Phi^+$, meaning $\alpha + \beta \in \Phi^+$ but $(\alpha|\beta) = 0$, for which the associated generators do not commute.

We next reformulate the commutation relations for divided and ordinary powers of simple root generators derived in (c), (i), and (a1) from Section 5.2 in [Lus90b] into our conventions and a more compact formalism. As in Section 2.7, assume $-A_{ji} = d_i = 1$ and $-A_{ij} = d_j = e \in \{1, 2, 3\}$ depending on Lie type. The translation utilizes the dictionary of generators from (5.19) as well as additional notation as follows.

To begin with, define for a root lattice vector $\vec{n} = M\alpha_i + N\alpha_j \in \mathbb{Z}^\Delta$ the set

$$S_e(\vec{n}) = \left\{ \psi \in \mathbb{N}_0^{\Phi^+} : \vec{\psi} = M\alpha_i + N\alpha_j \right\}, \quad (5.40)$$

where $\vec{\psi} = \sum_\alpha \psi(\alpha)\alpha \in \mathbb{Z}^\Delta$ as in Section 5.3. Note that the ordering of generators from right to left is the one determined by $z_{|ij|}$ as discussed in Section 2.7, opposite to the one displayed in (2.33) for $z_{|ji|}$. As before, set $\beta_s = \gamma(z_{|ij|}[1, s])$ so that $\beta_1 = \alpha_i$, $\beta_2 = s_i(\alpha_j) = e\alpha_i + \alpha_j$ and so forth.

For a given $\psi \in \mathbb{N}_0^{\Phi^+}$ define $k_s = \psi(\beta_s)$. So, explicitly, for $e = 1$ we have $k_1 = \psi(\alpha_i)$, $k_2 = \psi(\alpha_i + \alpha_j)$, and $k_3 = \psi(\alpha_j)$, and for $e = 2$ we have $k_1 = \psi(\alpha_i)$, $k_2 = \psi(2\alpha_i + \alpha_j)$, $k_3 = \psi(\alpha_i + \alpha_j)$ and $k_4 = \psi(\alpha_j)$. Moreover, if $e = 3$, we have $k_1 = \psi(\alpha_i)$, $k_2 = \psi(3\alpha_i + \alpha_j)$, $k_3 = \psi(2\alpha_i + \alpha_j)$, $k_4 = \psi(3\alpha_i + 2\alpha_j)$, $k_5 = \psi(\alpha_i + \alpha_j)$, and $k_6 = \psi(\alpha_j)$. The explicit forms of the associated root lattice vectors are thus

$$\vec{\psi} = \begin{cases} (k_1 + k_2)\alpha_i + (k_2 + k_3)\alpha_j & e = 1 \\ (k_1 + 2k_2 + k_3)\alpha_i + (k_2 + k_3 + k_4)\alpha_j & e = 2 \\ (k_1 + 3k_2 + 2k_3 + 3k_4 + k_5)\alpha_i + (k_2 + k_3 + 2k_4 + k_5 + k_6)\alpha_j & e = 3. \end{cases} \quad (5.41)$$

With these conventions we also define a quadratic function $f_e : \mathbb{N}_0^{\Phi^+} \rightarrow \mathbb{N}_0$ as

$$f_e(\psi) = \begin{cases} k_1k_3 + k_2 & e = 1 \\ 2k_4(k_1 + k_2) + k_3k_1 + 2k_2 + 2k_3 & e = 2 \\ 3k_6(k_1 + 2k_2 + k_3 + k_4) + k_5(2k_1 + 3k_2 + k_3) + 3k_4(k_1 + k_2) + k_3k_1 \\ \quad + 3k_2 + 4k_3 + 6k_4 + 3k_5 & e = 3. \end{cases} \quad (5.42)$$

Suppose $\hat{\chi}_i, \hat{\chi}_j \in \mathbb{N}_0^{\Phi^+}$ are the indicator functions for the simple roots α_i and α_j respectively. That is, (k_1, \dots, k_m) is equal to $(1, 0, \dots, 0)$ for the former and $(0, \dots, 0, 1)$ for the latter. The variation of f_e with respect to the simple root exponents can then be expressed as

$$\begin{aligned} f_e(\psi + s\hat{\chi}_j) &= f_e(\psi) + s\mathbf{e}(\psi(\alpha_j) + u_j(\vec{\psi})) \\ \text{and} \quad f_e(\psi + s\hat{\chi}_i) &= f_e(\psi) + s(\psi(\alpha_i) + u_i(\vec{\psi})). \end{aligned} \quad (5.43)$$

Here $u_i, u_j \in (\mathbb{Z}^\Delta)^*$ are defined for $\vec{\psi} = M\alpha_i + N\alpha_j$ as

$$u_j(\vec{\psi}) = M - N = \begin{cases} k_1 - k_3 & e = 1 \\ k_1 + k_2 - k_4 & e = 2 \\ k_1 + 2k_2 + k_3 + k_4 - k_6 & e = 3 \end{cases} \quad (5.44)$$

$$u_i(\vec{\psi}) = eN - M = \begin{cases} -k_1 + k_3 & e = 1 \\ -k_1 + k_3 + 2k_4 & e = 2 \\ k_1 + k_3 + 3k_4 + 2k_5 + 3k_6 & e = 3. \end{cases} \quad (5.45)$$

Using the basis elements from (5.33), the mentioned equations in [Lus90b] can thus be written in the following form, which applies to all Lie types.

$$E_i^{(M)} E_j^{(N)} = \sum_{\psi \in S_e(\vec{n})} q^{f_e(\psi)} E_{\vec{z}_{[ij]}}^{(\psi)} \quad (5.46)$$

For convenience, we provide the explicit forms of the basis elements

$$E_{\vec{z}_{[ij]}}^{(\psi)} = \begin{cases} E_j^{(k_3)} E_{(ij)}^{(k_2)} E_i^{(k_1)} & e = 1 \\ E_j^{(k_4)} E_{(iji)}^{(k_3)} E_{(ij)}^{(k_2)} E_i^{(k_1)} & e = 2 \\ E_j^{(k_6)} E_{(iji)j}^{(k_5)} E_{(iji)}^{(k_4)} E_{(iji)}^{(k_3)} E_{(ij)}^{(k_2)} E_i^{(k_1)} & e = 3. \end{cases} \quad (5.47)$$

Multiplying both sides of (5.46) by respective quantum factorials, we obtain the commutation relation of ordinary powers,

$$E_i^M E_j^N = \sum_{\psi \in S_e(\vec{n})} q^{f_e(\psi)} [\psi]_e E_{\vec{z}_{[ij]}}^{\psi}, \quad (5.48)$$

where all coefficients are in $\mathbb{Z}[q, q^{-1}]$. Explicitly, these are expressed in terms of quantum factorials and quantum multinomial coefficients as follows.

$$[\psi]_e = \begin{cases} [k_2]! \begin{bmatrix} k_1 + k_2 \\ k_2 \end{bmatrix} \begin{bmatrix} k_2 + k_3 \\ k_2 \end{bmatrix} & e = 1 \\ [2k_2]![k_3]!_j \begin{bmatrix} k_1 + 2k_2 + k_3 \\ k_1, 2k_2, k_3 \end{bmatrix} \begin{bmatrix} k_2 + k_3 + k_4 \\ k_2, k_3, k_4 \end{bmatrix}_j & e = 2 \\ [3k_2]![k_3]![k_3]!_j [3k_4]![k_4]!_j [k_5]!_j \begin{bmatrix} 2k_3 \\ k_3 \end{bmatrix} \begin{bmatrix} 2k_4 \\ k_4 \end{bmatrix}_j \\ \cdot \begin{bmatrix} k_1 + 3k_2 + 2k_3 + 3k_4 + k_5 \\ k_1, 3k_2, 2k_3, 3k_4, k_5 \end{bmatrix} \begin{bmatrix} k_2 + k_3 + 2k_4 + k_5 + k_6 \\ k_2, k_3, 2k_4, k_5, k_6 \end{bmatrix}_j & e = 3 \end{cases} \quad (5.49)$$

As usual, the j subscripts at the quantum numbers and quantum multinomial coefficients indicate that $q_j = q^e$ is used instead of $q_i = q$. The resummation formulae in the next lemma are an important tool for relating bases in later sections.

Lemma 5.7. Suppose $N, M \in \mathbb{N}_0$ and $\vec{n} = M\alpha_i + N\alpha_j$. Then, for an indeterminate z ,

$$\sum_{k=0}^N z^k q_j^{k(\alpha_j(\vec{n})-1)} E_j^{(k)} E_i^{(M)} E_j^{(N-k)} = \sum_{\phi \in S_e(\vec{n})} q^{f_e(\phi)} \prod_{t=1}^{\phi(\alpha_j)} (q_j^{-2t} z + 1) E_{\blacktriangleleft z_{|ij|}}^{(\phi)} \quad (5.50)$$

$$\sum_{k=0}^M z^k q_i^{k(\alpha_i(\vec{n})-1)} E_i^{(M-k)} E_j^{(N)} E_i^{(k)} = \sum_{\phi \in S_e(\vec{n})} q^{f_e(\phi)} \prod_{t=1}^{\phi(\alpha_i)} (q_i^{-2t} z + 1) E_{\blacktriangleleft z_{|ij|}}^{(\phi)}. \quad (5.51)$$

Proof. We show here only (5.50) since the proof for (5.51) is analogous. Set $w = zq_j^{\alpha_j(\vec{n})-1} = zq^{e(M-N-1)}$. The left side of (5.50) can be evaluated as

$$\begin{aligned} \sum_{k=0}^N w^k E_j^{(k)} E_i^{(M)} E_j^{(N-k)} &= \sum_{\substack{0 \leq k \leq N \\ \psi \in S_e(\vec{n}-k\alpha_j)}} w^k q^{f_e(\psi)} E_j^{(k)} E_{\blacktriangleleft z_{|ij|}}^{(\psi)} = \sum_{\substack{0 \leq k \leq N \\ \psi \in S_e(\vec{n}-k\alpha_j)}} w^k q^{f_e(\psi)} \begin{bmatrix} \psi(\alpha_j)+k \\ k \end{bmatrix}_j E_{\blacktriangleleft z_{|ij|}}^{(\psi+k\hat{\chi}_j)} \\ &= \sum_{\substack{0 \leq k \leq \phi(\alpha_j) \\ \phi \in S_e(\vec{n})}} w^k q^{f_e(\phi-k\hat{\chi}_j)} \begin{bmatrix} \phi(\alpha_j) \\ k \end{bmatrix}_j E_{\blacktriangleleft z_{|ij|}}^{(\phi)} = \sum_{\phi \in S_e(\vec{n})} q^{f_e(\phi)} \left(\sum_{k=0}^{\phi(\alpha_j)} w^k q^{-k\epsilon(\phi(\alpha_j)+M-N)} \begin{bmatrix} \phi(\alpha_j) \\ k \end{bmatrix}_j \right) E_{\blacktriangleleft z_{|ij|}}^{(\phi)}. \end{aligned}$$

Here we use (5.46) in the first equality with $\vec{n}-k\alpha_j = M\alpha_i + (N-k)\alpha_j$. The second step is immediate from the forms (5.42) and combining quantum factorials. In the third equality, we substitute $\phi = \psi + k\hat{\chi}_j$, summing instead over $\phi \in S_e(\vec{n})$ with $\phi(\alpha_j) \geq k$. The last step invokes (5.43) and rearranges summations. After resubstituting z in the last expression, we apply Gauss's formula from (3.3) with $a = \phi(\alpha_j)$ to arrive at the desired expression. \square

Specializing (5.50) to $z = -q_j^2$ and (5.51) to $z = -q_i^2$ yields the commutation relations in Section 5.5 of [Lus90b]. In this case, the product terms will be zero unless $\phi(\alpha_j) = k_m = 0$ or $\phi(\alpha_i) = k_1 = 0$, respectively. Assuming further $\alpha_j(\vec{n}) = M - N = 0$ for (5.50), it is immediate from (5.44) that $k_1 = \dots = k_{m-2} = 0$. Thus, the only exponent function ϕ' that occurs in the summation is the one that is N on $\gamma(a_{m-1}) = \alpha_i + \alpha_j$ and zero on all other roots, where a_j is as in (2.32). Observing also that $f_e(\phi') = \epsilon N$, the right side of (5.50) thus reduces to $q^{\epsilon N} E_{a_{m-1}}^{(N)}$.

Analogously, for (5.51) the constraint $\alpha_i(\vec{n}) = \epsilon N - M = 0$ implies $k_3 = \dots = k_m = 0$ so that the only remaining exponent function is the one which is N on $\gamma(a_2) = \epsilon \alpha_i + \alpha_j$ and zero on others. The right side of (5.50) thus becomes $q^{\epsilon N} E_{a_2}^{(N)}$. The resulting identities are summarized in the left column of the corollary below.

Corollary 5.8. The following identities hold:

$$E_{a_{m-1}}^{(N)} = \sum_{k=0}^N (-1)^{N-k} q_j^{-k} E_j^{(N-k)} E_i^{(N)} E_j^{(k)}, \quad E_{(ji)}^{(N)} = \sum_{k=0}^N (-1)^k q_j^{-(N-k)} E_j^{(N-k)} E_i^{(N)} E_j^{(k)}, \quad (5.52)$$

$$E_{(ij)}^{(N)} = \sum_{k=0}^{\epsilon N} (-1)^{\epsilon N-k} q^{-k} E_i^{(k)} E_j^{(N)} E_i^{(\epsilon N-k)}, \quad E_{b_{m-1}}^{(N)} = \sum_{k=0}^{\epsilon N} (-1)^k q^{-(\epsilon N-k)} E_i^{(k)} E_j^{(N)} E_i^{(\epsilon N-k)}. \quad (5.53)$$

The identities in the right column are obtained by applying \mathfrak{U} and using (5.16) as well as the relations $a_{m-1} \circ \circ b_2 = w_j w_i$ and $b_{m-1} \circ \circ a_2 = w_i w_j$ as in Lemma 2.12.

As already indicated, (5.52) corresponds to the first relation in Section 5.5 of [Lus90b]. For the change in conventions, note that $\tilde{E}_{ij} = \tilde{E}_{(ji)} = E_{a_{m-1}}$ by (5.19). Similarly, the right identity of (5.53) is the translation of the second expression in [Lus90b, Section 5.5]. For this, the generators \tilde{E}_{ij} , \tilde{E}_{iij} , and \tilde{E}_{iiij} listed in [Lus90b] for $e = 1, 2$, and 3 , respectively, coincide in all cases with $\tilde{E}_{b_{m-1}} = E_{(ij)}$ via (5.19).

We thus have expansions into products of generators for simple roots of divided power generators $E_w^{(N)}$ for all words, with exception of the words $a_3 \circ \circ b_4$ and $a_4 \circ \circ b_3$ in the G_2 -case. Yet, unlike (5.48), equations (5.52) and (5.53) cannot be written as expansions of ordinary powers over $\mathbb{Z}[q, q^{-1}]$.

5.5. Numberings, Boxes, and Refining Convex Orderings. The integral PBW bases derived in [Lus90a, Lus90b] for the divided power algebra $\mathcal{L}U^+$ require orderings on Φ^+ that are compatible with a specific pre-order \precsim on Φ^+ , defined in Section 4.3 of [Lus90b]. As we adapt the proofs there to the situation of regular powers, we review this pre-order next and establish various relations to convex orderings and maximal words.

Lusztig's construction of \precsim starts by imposing a *good* numbering on the set Δ of simple roots. For a given indexing $\Delta = \{\alpha_1, \dots, \alpha_n\}$, write $\Phi^+[1, j]$ for the respective subroot system generated by $\{\alpha_1, \dots, \alpha_j\}$. Assuming that each $\Phi^+[1, j]$ is connected, set $L_j = \Phi^+[1, j] - \{\rho_j\}$, where ρ_j is the highest root in $\Phi^+[1, j]$. The numbering is called good if for any root $\beta \in L_j$ the coefficient of α_j is at most 1.

The numberings in appendix of [Bou02] are good with the exception of types B and F. For the latter two, the reverse numberings given by $i \leftrightharpoons l+1-i$ are good. For type A any numbering is good, for C also the reverse of [Bou02] is good, and for type D any numbering coinciding with the one in [Bou02] in the highest index as well as the reverse of the one in [Bou02] are good.

For a given good numbering, any positive root β can be uniquely written as $\beta = b_j \alpha_j + b_{j-1} \alpha_{j-1} + \dots + b_1 \alpha_1$ with $b_j > 0$. Lusztig introduces maps $g : \Phi^+ \rightarrow \{1, \dots, n\}$ and $c : \Phi^+ \rightarrow \{1, 2\}$ given by $g(\beta) = j$ and $c(\beta) = c_j$. It is clear from the good numbering construction that $c(\beta) = 2$ implies that $\beta = \rho_j$ in $\Phi^+[1, j]$. Lusztig further defines $h' : \Phi^+ \rightarrow \mathbb{Q}$ by $h'(\beta) = c(\beta)^{-1} \text{ht}(\beta)$, where $\text{ht}(\beta) = \sum_i b_i$ is the height of the root with respect to Δ .

The pre-order is now defined by writing $\alpha \precsim \beta$ if $g(\alpha) \geq g(\beta)$ and $h'(\alpha) \leq h'(\beta)$. The equivalence classes $R_{m,k} = \{\beta \in \Phi^+ : g(\beta) = m, h'(\beta) = k\}$ are called *boxes* in [Lus90b]. We record the following two observations.

Lemma 5.9. *Suppose $\alpha, \beta \in \Phi^+$ ($\alpha \neq \beta$) belong to the same box. Then none of $\pm \alpha \pm \beta$ are roots.*

Proof. Given $m = g(\alpha) = g(\beta)$ we have $\alpha, \beta \in M = \Phi^+[1, m]$, where M itself is a root system with highest root ρ , so that it suffices to show $\pm \alpha \pm \beta \notin M$. Assume first $c(\alpha) = c(\beta) = 1$. Then

$\text{ht}(\alpha - \beta) = \text{ht}(\alpha) - \text{ht}(\beta) = h'(\alpha) - h'(\beta) = 0 = \text{ht}(\beta - \alpha)$, which is not possible. For $\gamma = \alpha + \beta$ we have $c(\gamma) = 2$ and $\text{ht}(\gamma) = 2k$. If $\gamma \in M$ the former implies that $\gamma = \rho$ and that M is of type E_8 , F_4 , or G_2 . In all three cases $\text{ht}(\rho)$ is odd (29, 11, or 5), leading to a contradiction. Now, $c(\alpha) = 2$ implies $\alpha = \rho$ and hence $k = h'(\alpha) \in \{14.5, 5.5, 2.5\}$ so that there is no other root in the same box. \square

An immediate corollary is that any two roots in the same box are (strongly) orthogonal. In [Lus90a, Lus90b] Lusztig goes on to consider total orders refining \precsim . With good numberings as above, we notice the following.

Lemma 5.10. *Any total order on Φ^+ refining \precsim is convex and thus of the form \preceq_w for some $w \in \mathcal{W}_m^*$.*

Proof. Let $\alpha, \beta, \gamma = \alpha + \beta \in \Phi^+$ and suppose $\alpha \in R_{m,k}$ and $\beta \in R_{m',k'}$. Set $t = c(\alpha)c(\gamma)^{-1}$ and $s = c(\beta)c(\gamma)^{-1}$ so that $h'(\gamma) = t \cdot k + s \cdot k'$, using that $\text{ht}(\gamma) = \text{ht}(\alpha) + \text{ht}(\beta)$. Write also $\mu \nsim \nu$ to indicate two roots are not in the same box.

Suppose now $<$ is a total order refining \precsim and assume $\alpha < \beta$. Since $\gamma \in \Phi^+$, Lemma 5.9 implies $\alpha \nsim \beta$. Hence, either we have $m > m'$ or we have $m = m'$ and $k < k'$. The former case implies $g(\gamma) = m$, which means $\gamma \precsim \beta$ and $\beta \nsim \gamma$ and, hence, $\gamma < \beta$. Moreover, with $c(\gamma) = c(\alpha)$, we find $t = 1$ so that $h'(\gamma) = k + s \cdot k' > k$. Hence, $\gamma \gtrsim \alpha$ and $\gamma \nsim \alpha$ so that $\alpha < \gamma$.

In the second case we have $g(\alpha) = g(\beta) = g(\gamma)$ and $c(\gamma) = c(\alpha) + c(\beta)$ so that $s = (1 - t)$. Thus $h'(\gamma) = t \cdot k + (1 - t) \cdot k'$ and $k < h'(\gamma) < k'$ since $0 < t < 1$. We, therefore, obtain $\alpha < \gamma < \beta$ in this case as well and, hence, convexity of the total ordering. As shown in [Pap94] this implies that $<$ needs to be of the form \preceq_w as desired. \square

Note, for Lie types A , B , and G each box contains no more than one element. Thus, there is only one $w \in \mathcal{W}_m^*$ for which the total order \preceq_w refines \precsim . Specifically, the chosen good numbering for A_n is realized by the maximal word $w_{A_n} = a_1 a_2 \dots a_n$, where $a_k = w_n w_{n-1} \dots w_k$. Also, for B_n we have $w_{B_n} = b_n \dots b_1$, where $b_k = w_k w_{k-1} \dots w_2 w_1 w_2 \dots w_{k-1} w_k$, and for G_2 that $w_{G_2} = (w_2 w_1)^3$.

For C_n with the Bourbaki numbering one longest word choice refining \precsim is given as a product $w = u \cdot v$. Here $u = \alpha_0 \beta_0 \alpha_1 \beta_1 \dots \beta_1 \alpha_1 \beta_0 \alpha_0$ is a palindromic expression in $\alpha_j = w_n w_{n-2} \dots w_{n-2j}$ and $\beta_j = w_{n-1} w_{n-3} \dots w_{n-2j-1}$ centered around the longest α_k or β_k , and v is a longest word for A_{n-1} . The reduced word

$$w_4 w_3 w_2 w_3 w_1 w_4 w_2 w_3 w_4 w_2 w_3 w_1 w_2 w_3 w_4 w_3 w_2 w_3 w_1 w_2 w_3 w_1 w_2 w_1$$

is compatible with the pre-order for F_4 using the reverse numbering of that in [Bou02]. The words for the simply laced D and E Lie types are similarly worked out with additional choices within boxes.

5.6. PBW Bases for Box Orderings. The sole focus of this section is the adaption of the proofs in [Lus90a, Lus90b] to yields a PBW theorem for regular power generators instead of divided power generators. The main change from the original proof is that the involved commutation and recursion relations for generators are valid only over Λ_{q,e^*} rather than Λ_q as in [Lus90a, Lus90b]. The second difference is merely concerning conventions, namely the alternate choice of generators E_w defined via the automorphisms Γ_i in Section 5.1 rather than the \tilde{E}_w used in [Lus90a, Lus90b]. Thanks to the relation in (5.29) it is not difficult to move between these two settings using the \mathfrak{U} involution, while also replacing the root ordering for z by the one for z^\dagger . The form (5.25) for the action of the Υ involutions implies, further, that bases with generators multiplied in opposing directions have the same spans.

Although the inductive argument for a PBW theorem itself is unchanged from [Lus90a, Lus90b], we outline the main steps again in the proof below. We also adapt the formulae to fit our conventions, even as the mentioned automorphisms and symmetries would afford us the flexibility to retain those in [Lus90a, Lus90b]. We freely use the notation for monomial sets and spans introduced at the end of Section 5.3, with a focus on the minimal ground ring Λ_{q,e^*} and the maximal exponent set $S = \mathcal{E}(s_\bullet)$.

Proposition 5.11. *Suppose $w \in \mathcal{W}_m^*$ is such that \preceq_w refines \preceq . Then U_q^+ is free a Λ_{q,e^*} -module with basis given by either the generators in $\mathfrak{B}_{w\blacktriangleright}^+$ or those in $\mathfrak{B}_{\blacktriangleleft w}^+$ as defined in (5.35). The analogous statement holds for U_q^- and the sets $\mathfrak{B}_{w\blacktriangleright}^-$ and $\mathfrak{B}_{\blacktriangleleft w}^-$.*

Proof. Recall that, by Proposition 5.1, the Λ_{q,e^*} -module $\mathfrak{S}_{w\blacktriangleright}^+$, generated by elements in $\mathfrak{B}_{w\blacktriangleright}^+$, is contained in U_q^+ . Since the elements in $\mathfrak{B}_{w\blacktriangleright}^+$ form a basis of $U_{q,\mathbb{Q}}$ over the field $\mathbb{Q}(q)$, see Corollary 40.2.2 in [Lus93], we also know that they form a basis of $\mathfrak{S}_{w\blacktriangleright}^+$ as a Λ_{q,e^*} -module and that $U_q^+/\mathfrak{S}_{w\blacktriangleright}^+$ is at most a torsion module. Since $1 \in \mathfrak{S}_{w\blacktriangleright}^+$ is a cyclic vector for the left or right regular action of U_q^+ on itself, either $U_q \cdot \mathfrak{S}_{w\blacktriangleright}^+ \subseteq \mathfrak{S}_{w\blacktriangleright}^+$ or $\mathfrak{S}_{w\blacktriangleright}^+ \cdot U_q \subseteq \mathfrak{S}_{w\blacktriangleright}^+$ would imply $U_q = \mathfrak{S}_{w\blacktriangleright}^+$. Thus, it remains to show that $\mathfrak{S}_{w\blacktriangleright}^+ \cdot E_i \subseteq \mathfrak{S}_{w\blacktriangleright}^+$ for all i or $E_i \cdot \mathfrak{S}_{w\blacktriangleright}^+ \subseteq \mathfrak{S}_{w\blacktriangleright}^+$ for all i .

The inductive proof in the simply laced case can be obtained by modifying the arguments in Section 2 of [Lus90a]. Specifically, instead of the spaces in 2.10 we consider analogous spaces defined with regular rather than divided powers. For example, $X_{m,k}$ is replaced by $X_{m,k}^*$, defined as the Λ_{q,e^*} -algebra of U_q^+ generated by $E_{w[\alpha]}$ (as in (2.15)) with $\alpha \in R_{m,k}$ as well as E_j with $j < m$. Similarly, $Y_{m,k}^*$ is the free Λ_{q,e^*} -submodule generated by $\mathfrak{B}_{u\blacktriangleright}^+$ where $u \leq_R w$ is the maximal word with $\mathcal{N}(u) \cap R_{m,k}^+ = \emptyset$. Statement (b) of Lemma 2.11 in [Lus90a] is then changed to $Y_{m,k}^* \cdot X_{m,k}^* \subseteq Y_{m',k'}^* \cdot X_{m',k'}^*$ if $R_{m',k'}^+$ is the next smaller box in the \preceq order after $R_{m,k}^+$ with integer k' .

In the adjustment of the proof, we first redefine the generators ξ_t as ordinary powers $E_{\alpha_t}^{N_t}$. The notions of distinguished and non-distinguished generators as well as bad or good pairs are adapted to ordinary powers in the same manner. The commutation argument invokes equations (d2)-(d5) in §2.3 of [Lus90a] that are imposed on the algebra V^+ defined there. The isomorphism $V^+ \cong \mathcal{L}U^+$ in Section 4.7 implies that these relations also hold in $U_{q,\mathbb{Q}}^+$ for the divided powers of

the \ddot{E}_α generators. Analogous equations can be readily derived from these for ordinary powers of the E_α generators, as defined in (5.1) and (5.3). For example, for α and α' fulfilling (e1) or (e2), equation (d3) in [Lus90a] specializes for $N = M = 1$ to $\ddot{E}_{\alpha'} \ddot{E}_\alpha = q \ddot{E}_{\alpha+\alpha'} + q \ddot{E}_\alpha \ddot{E}_{\alpha'}$. Applying \mathfrak{U} with $\mathfrak{U}(\ddot{E}_\alpha) = E_\alpha$ we obtain $E_{\alpha'} E_\alpha = (-1) E_{\alpha+\alpha'} + q^{-1} E_\alpha E_{\alpha'}$. Analogs for (d2), (d4), and (d5) are also obtained by application of \mathfrak{U} and multiplication of factorials. The desired analog of (d3) is then obtained from these relations by the same induction,

$$E_{\alpha'}^N E_\alpha^M = \sum_{j=0}^{\min(N,M)} (-1)^j q^{-(N-j)(M-j)} [j]! \begin{bmatrix} N \\ j \end{bmatrix} \begin{bmatrix} M \\ j \end{bmatrix} E_\alpha^{M-j} E_{\alpha+\alpha'}^j E_{\alpha'}^{N-j}.$$

Clearly, all coefficients are in $\Lambda_q = \Lambda_{q,2}$. The same process is applied to (b) in Proposition 2.7 to express $E_{\alpha_i}^N E_{\beta_0}^M$ as Λ_q -combinations of respective ordinary power generators with reversed cyclic order of indices, using also that $\psi_0 = 1$ and $\begin{bmatrix} N \\ s \end{bmatrix} \begin{bmatrix} M \\ s \end{bmatrix} \in \Lambda_q$. The other equations in Proposition 2.7 and 2.8 are modified, as before, by application of \mathfrak{U} and multiplication of factorials.

All other parts of the proof of Lemma 2.11 in [Lus90a] can be adapted verbatim. As in Proposition 2.12(b) the inclusion can be iterated to infer $Y_{n,1}^* \cdot X_{n,1}^* \subseteq Y_{1,1}^* \cdot X_{1,1}^*$, which is exactly the inclusion $U_q^+ \subseteq \mathbf{B}^+$.

As noted in [Lus90b] for the case of divided powers, the proof for the non-simply laced cases follows the same type of inductive argument. Commutation relations of all divided power generators in the rank 2 case $\mathbf{B}_2 = \mathbf{C}_2$ are explicitly worked out in Sections 5.2 and 5.3 of [Lus90b]. As with the simply laced cases, it is straightforward to derive relations for $e = 2$ and $e = 3$ in Section 5.2 for the opposite orders of generators and, hence, after application of \mathfrak{U} , respective relations for our convention of generators. From these, analogs of the equations in Section 5.3 can be readily derived for our generators by rescaling arguments. The coefficient will thus differ only by units in Λ_{q,e^*} .

Multiplication of the analogs of equations (d) through (i) in the same section of [Lus90b] by respective quantum factorials readily yields commutation relations for ordinary powers with coefficients in $\Lambda_{q,3}$. The variant of (g) for our generators is, for example, as follows,

$$E_\gamma^k E_\delta^{k'} = \sum_{s=0}^{\min(k,k')} \frac{1}{[2]^s} q^{2s(k+k')-3s^2} (q - q^{-1})^s [s]_2! \begin{bmatrix} k \\ s \end{bmatrix}_2 \begin{bmatrix} k' \\ s \end{bmatrix}_2 E_\delta^{k'-s} E_\beta^{2s} E_\gamma^{k-s}. \quad (5.54)$$

Here $\delta = \alpha_2$, $\gamma = \alpha_2 + 2\alpha_1$, and $\beta = \alpha_1 + \alpha_2$ with α_2 the long root. The coefficients are all in $\Lambda_{q,3}$ but not in Λ_q due to the additional $\frac{1}{[2]}$ factors. The respective variant for (h) is

$$E_1^k E_\tau^{k'} = \sum_{s=0}^{\min(k,k')} (-1)^s [2]^s q^{s(k+k')-\frac{1}{2}s(7s+3)} [s]! \begin{bmatrix} k \\ s \end{bmatrix} \begin{bmatrix} k' \\ s \end{bmatrix} E_\tau^{k'-s} E_{\tau+\alpha_1}^s E_1^{k-s}, \quad (5.55)$$

where $\tau = \alpha_2 + \alpha_1$. Using respective multinomial coefficients and $v = \alpha_2$ we also reexpress equation (i) in [Lus90b] as

$$E_1^k E_v^{k'} = \sum_{\substack{r,s,t,u \geq 0 \\ r+s+t=k' \\ s+2t+u=k}} (-1)^s q^{-2r(u+t)+us} [s]_2 [2t]! \begin{bmatrix} k \\ s, 2t, u \end{bmatrix}_2 \begin{bmatrix} k' \\ r, s, t \end{bmatrix}_2 E_v^r E_{v+\alpha_1}^s E_{v+2\alpha_1}^t E_1^u. \quad (5.56)$$

It is immediate that the set of relations allows reordering of any monomial over $\Lambda_{q,3}$ into the order on the positive roots of B_2 given by $w_2 w_1 w_2 w_1$. Explicit relations for G_2 are given in Section 5.4 of [Lus90b] as well. Although more complicated, all of them can be multiplied by respective factorials and transformed into relations for our generators in the same manner, except that powers of $\frac{1}{[3]}$ occur in addition to those of $\frac{1}{[2]}$.

The higher rank BCF cases follow as in the simply laced case from commutation relations obtained by application of Artin automorphisms to the rank 2 relations as well as those for the respective A type subalgebras. In the above ordering for the B case Lemma 5.6 readily implies that generators from consecutive boxes $R_{m,k}^+$ and $R_{m,k+1}^+$ skew-commute. Consider next required commutation relations between elements E_τ with $\tau \in R_{m,k}^+$ and generators E_j for $j < m$. From Proposition 5.1 it follows that $\Gamma_{d_m}(E_j) = E_j$ for $d_m = b_n \dots b_{m+1} \in \mathcal{W}^*$, reducing the arguments to the case $m = n$ and, hence, $E_\tau = E_v$ with $v \leq_R b_m$.

In all cases in which $\tau + \alpha_j \notin \Phi^+$ it is then readily checked that E_τ and E_j and, hence, all their powers commute up to signs. For $\tau = \alpha_m + \dots + \alpha_{j+1}$ when $j > 1$ we invoke the known commutation relations for $A_{n-1} \subset B_n$. Applying $\Gamma_{w_m \dots w_3}$ to (5.55) we see that the relation holds more generally for $v = \alpha_m + \dots + \alpha_2$, yielding the needed relation between E_v and E_1 . Similarly, applying $\Gamma_{w_m \dots w_3}$ to (5.55) extends the relation to the case $\tau = \alpha_m + \dots + \alpha_1$. Note that in this case $(\tau|\alpha_1) = 0$, even as $\tau + \alpha_1 \in \Phi^+$.

Finally, for $j > 1$ and $\tau = \alpha_m + \dots + \alpha_j + 2\alpha_{j-1} + \dots + 2\alpha_1$ we find $E_j E_\tau = q^{-2} E_\tau E_j - E_{\tau+\alpha_j}$, $E_{\tau+\alpha_j} E_j = q^{-2} E_j E_{\tau+\alpha_j}$, and $E_{\tau+\alpha_j} E_\tau = q^2 E_\tau E_{\tau+\alpha_j}$, which implies E_τ and E_j fulfill the A_2 Serre relations with q replaced by q^2 . This, again, implies the usual A type commutation relations, only with q replaced by q^2 .

The treatment for the C and F cases uses the same arguments and analogous computations. Indeed, choosing the numbering opposite to that of [Bou02] is also good for C, leading to essentially the same relations as in the B case. The F_4 can be treated explicitly as an extension of B_3 or C_3 . This completes the proof for the basis $\mathfrak{B}_{w\bullet}^+$ for one direction of multiplication.

It follows now immediately from (5.25) that $\mathfrak{B}_{w\bullet}^+$ provides a basis of U_q^+ as well, since Υ maps elements from $\mathfrak{B}_{w\bullet}^+$ to those of $\mathfrak{B}_{w\bullet}^+$ up to unit factors. Similarly, (5.24) implies the respective statements for U_q^- with basis elements mapped to each other by Ω . \square

As noted in the introductory remarks, we could have also followed the Lusztig's conventions with the respectively renormalized commutation relations in the \tilde{E}_w . This would have yielded $\tilde{E}_{w\bullet}^\psi$ as a Λ_{q,e^*} -basis for U_q^+ for w compatible with the box ordering chosen in [Lus90a, Lus90b].

Since U_q^+ is invariant under \mathfrak{U} , we have from (5.29) that $E_{\blacktriangleleft w^\dagger}^\psi$ is also a basis. So, after application Υ , we find that $E_{w^\dagger \blacktriangleright}^\psi$ is a basis as well. For the general PBW theorem, it suffices to establish a PBW statement for only one ordering, so that the replacement of w by w^\dagger does not change the proof of Propositions 5.17 or Theorem 5.18 below.

Corollary 5.12. *For $w \in \mathcal{W}_m^*$ as in Proposition 5.11 the sets $\mathfrak{B}_{w \blacktriangleright}^\pm$ and $\mathfrak{B}_{\blacktriangleleft w}^\pm$ are bases of U_q^- as a free $\Lambda_{q, \mathbf{e}^*}$ -module.*

We conclude with another useful consequence of Proposition 5.11, which allows us to identify subalgebras generated by a subset of generators as quantum algebras themselves.

Corollary 5.13. *For any pair of distinct indices $i, j \in \{1, \dots, n\}$, let $U_q^{(i,j)+}$ be the rank 2 quantum algebra with respectively restricted Cartan data. Then the canonical algebra homomorphism $U_q^{(i,j)+} \rightarrow U_q^+$ is well-defined and injective, and its image is the subalgebra of U_q^+ generated by E_i and E_j .*

Proof. Surjectivity on the subalgebra generated by E_i and E_j is obvious. Similarly, it is clear the map is well-defined, since the relations of $U_q^{(i,j)+}$ are also fulfilled in U_q^+ . For G_2 (or any rank 2 case) the statement is also obvious. To show injectivity in the $A_1 \times A_1$, A_2 , and $B_2 = C_2$ cases, we note that either numbering for each of these is a good numbering and can thus be chosen to inherit the order of the numbering imposed by U_q^+ .

Since also the height and lead coefficients of positive roots are the same, the pre-order \precsim of Φ^+ for U_q^+ restricts to the pre-order of positive root system $\Phi^{(i,j)+}$ of $U_q^{(i,j)+}$. This implies, by Proposition 5.11, that we have a basis for $U_q^{(i,j)+}$ that is mapped to basis elements of U_q^+ , implying injectivity. Indeed, $U_q^{(i,j)+}$ is a direct summand of U_q^+ as a $\Lambda_{q, \mathbf{e}^*}$ -module. \square

We conclude our discussion of the existence of bases with a brief outline of other approaches. As mentioned previously, Lusztig proves in [Lus90a, Lus90b, Lus93] that the monomials $E_{w \blacktriangleright}^{(\psi)}$ from (5.33) define a PBW basis of his divided power algebra $\mathcal{L}U^+$ over $\Lambda_q = \mathbb{Z}[q, q^{-1}]$. It can be correspondingly extended to a basis of all of $\mathcal{L}U$ over Λ_q at the expense of introducing additional generators in $\mathcal{L}U^0$ given by rational expressions in the K_i and q .

In [DCP93, Sec 12], De Concini and Procesi define the algebra ${}_sU$ to be the smallest subalgebra of $U_q \otimes \mathbb{k}(q)$ over $\mathbb{k}[q, q^{-1}]$ that contains the singularized generators \check{E}_i and \check{F}_i from (5.10) and is stable under the Artin group action generated by the Γ_i . They offer a proof that the monomials $\check{E}_{w \blacktriangleright}^\psi K^\mu \check{F}_{w \blacktriangleright}^\chi$ from (5.34) form a Λ_q -basis of ${}_sU$. Their argument is based on recursions formulae over \mathbb{C} in [LS91] and proves that recursion coefficients are indeed in $\mathbb{k}[q, q^{-1}]$ for an appropriate \mathbb{k} .

The fact that the $\check{E}_{\blacktriangleleft w}^\psi$ are a basis of ${}_sU^+$ may also be inferred from the Lusztig-Tanisaki pairing discussed in Section 9.4 and the PBW theorem for $\mathcal{L}U^+$ in [Lus90a, Lus90b, Lus93]. Comparing the coefficients in (5.33), (5.34), and (9.11) the pairing in (9.25) can be rewritten as

$$\langle \cdot, \cdot \rangle_z : \mathcal{L}U^- \times {}_sU^+ \rightarrow \Lambda_q \quad \text{with} \quad \langle F_{\blacktriangleleft w}^{(\psi)}, \check{E}_{\blacktriangleleft z}^\phi \rangle = \delta_{\psi, \phi} \cdot (-1)^{n(\psi)} \cdot q^{-\sum_\alpha d_\alpha \binom{\psi(\alpha)}{2}} \quad (5.57)$$

Since the pairings for $\psi = \phi$ are units in Λ_q the span of the $\check{E}_{\blacktriangleleft z}^\phi$ is the exact Λ_q -dual to $\mathcal{L}U^-$. The product and the q -braided coproduct $\bar{\Delta}$ defined by Lusztig [Lus93] on $\mathcal{L}U^-$ are then dual to respective structures on this span. This now identifies the latter with ${}_sU^+$ so that the basis result for ${}_sU^{\geq 0}$ is essentially the dual equivalent of Lusztig's basis result on $\mathcal{L}U^{\leq 0}$.

However, as algebras over Λ_q or Λ_{q,e^*} , ${}_sU^+$ and U_q^+ exhibit rather different behaviors. This becomes especially apparent in the $q \mapsto 1$ specialization, realized by the obvious ring maps $\Lambda_q \rightarrow \mathbb{Z}$ or $\Lambda_{q,e^*} \rightarrow \mathbb{Z}[\mathbf{e}^{-1}]$. For example, for Lie type A_n one easily sees from Proposition 7.7 that the specialization ${}_sU^+ \otimes \mathbb{Z}$ is the *commutative* polynomial algebra in the $E_{i,j}$, while the specialization $U_q^+ \otimes \mathbb{Z}$ is the universal envelope $U(\mathfrak{n})$, where \mathfrak{n} is the Lie algebra of the strictly upper triangular matrices.

Another variant is to consider renormalized generators $\mathring{E}_w = [d_w]E_w = \Gamma_{w^\flat}(\mathring{E}_i)$ with $i = \tau(w)$ and $\mathring{E}_i = [d_i]E_i$. The respective renormalization of the formulae (5.54, 5.55, 5.56) then yields expansion coefficients entirely in Λ_q . We, thus, expect that there is a Hopf algebra $\mathring{U}_q^{\geq 0}$ over Λ_q with a PBW basis over this ring, even for the non-simply laced Lie types.

5.7. Spanning Sets and Their Order Dependence. For any subset of exponent functions $\psi \in \mathcal{E}(s)$ for some $s \in \mathcal{W}$ one can consider the submodule spanned by the respective monomials $E_{w\blacktriangleright}^\psi$, where $s = r(w)$. Numerous subalgebras and ideals we will consider later can be defined as such spans. In this section we extract sufficient conditions on a set of exponent functions which imply that the respective free submodule is independent of the choice of a convex root ordering or, equivalently, dependent only on the Weyl element $s = r(w)$.

Suppose $s \in \mathcal{W}$. We say (a, b) with $a, b \in \mathcal{W}$ is a *relator pair* for s if

$$s = a \cdot s_{|ij|} \cdot b \quad \text{with} \quad l(s) = l(a) + l(s_{|ij|}) + l(b) \quad (5.58)$$

for the longest element $s_{|ij|}$ is some rank 2 subsystem $\Phi_{(ij)}$ of Φ . Observe now that Corollary 2.4 provides us with a bijection of inversion sets given as

$$\mathcal{O}_{(a,b)} = \text{id} \sqcup a \sqcup a \cdot s_{|ij|} : \mathcal{N}(a) \sqcup \mathcal{N}(s_{|ij|}) \sqcup \mathcal{N}(b) \longrightarrow \mathcal{N}(s). \quad (5.59)$$

Correspondingly, as in (5.22), we obtain the bijection of exponent sets

$$\mathcal{O}_{(a,b)}^* : \mathcal{E}(s) \rightarrow \mathcal{E}(a) \times \mathcal{E}(s_{|ij|}) \times \mathcal{E}(b). \quad (5.60)$$

Given a subset $S \subseteq \mathcal{E}(s)$ we abbreviate its image in the Cartesian product as $S_{(a,b)} = \mathcal{O}_{(a,b)}^*(S)$.

The following discussion assumes a general ground ring Λ with an extension of scalars function $f : \Lambda_{q,e^*} \rightarrow \Lambda$. Note, that Proposition 5.11 implies, that both $\mathfrak{B}_{z_{|ij}\blacktriangleright}^+$ and $\mathfrak{B}_{z_{|ji}\blacktriangleright}^+$ are bases of $U_{q,\Lambda^{(i,j)}}^+ = U_q^{(i,j)} \otimes_f \Lambda$. Recall from Corollary 5.13, that the latter may be viewed as the subalgebra generated by E_i and E_j . We, thus, can define coefficients $\mathfrak{v}_{\varphi'}^\varphi \in \Lambda$ for any $\varphi, \varphi' \in \mathcal{E}(s_{|ij|})$ by

$$\mathfrak{O}(E_{z_{|ij}\blacktriangleright}^\varphi) = E_{z_{|ij}\blacktriangleright}^\varphi = \sum_{\varphi'} \mathfrak{v}_{\varphi'}^\varphi \cdot E_{z_{|ij}\blacktriangleright}^{\varphi'}. \quad (5.61)$$

From these we define the following subsets of $\mathcal{E}(s_{|ij|})$ for any $\varphi \in \mathcal{E}(s_{|ij|})$ and $Q \subseteq \mathcal{E}(s_{|ij|})$:

$$\mathfrak{U}_{\{ij\}}(\varphi) = \{\varphi' \in \mathcal{E}(s_{|ij|}) : \mathfrak{U}_{\varphi'} \neq 0\} \quad \text{and} \quad \mathfrak{U}_{\{ij\}}(Q) = \bigcup_{\varphi \in Q} \mathfrak{U}_{\{ij\}}(\varphi). \quad (5.62)$$

Note that, since \mathfrak{U} is an involution, we have $\mathfrak{U}_{\{ij\}}(Q) = \mathfrak{U}_{\{ji\}}(Q)$. Commutation of \mathfrak{U} with Υ and (5.25) imply that the sets are the same if the oppositely ordered basis elements are used instead. Commutation with Ω shows the same for the respective bases of U_q^- . The sets may, however, depend on the choice of the ground ring as f may map some non-zero coefficients in Λ_{q,e^*} to zero in Λ .

For given $s \in \mathcal{W}$ we say an exponent set $S \subseteq \mathcal{E}(s)$ is *relator stable* if for any relator pair (a, b) for s we have the implication

$$(\chi, \varphi, \xi) \in S_{(a,b)} \quad \text{and} \quad \varphi' \in \mathfrak{U}_{\{ij\}}(\varphi) \quad \Rightarrow \quad (\chi, \varphi', \xi) \in S_{(a,b)}. \quad (5.63)$$

Defining $U_{\chi, \xi}^{S,a,b} \subseteq U_{q,\Lambda^{(i,j)}}^+$ as the Λ -span of the set $\{E_{z_{|ij|}\blacktriangleright}^\varphi : (\chi, \varphi, \xi) \in S_{(a,b)}\}$, it is clear from (5.61) that this condition may be equivalently restated as the requirement that $\mathfrak{U}(U_{\chi, \xi}^{S,a,b}) \subseteq U_{\chi, \xi}^{S,a,b}$ for all relator pairs (a, b) and $(\chi, \xi) \in \mathcal{E}(a) \times \mathcal{E}(b)$.

Note that we only need to consider cases when $m_{ij} = l(s_{|ij|}) \in \{3, 4, 6\}$ since the condition is void for $m_{ij} = 2$. Indeed, in the latter case $\Phi_{(ij)} = \{\alpha_i, \alpha_j\}$ and E_i and E_j commute so that $E_{z_{|ij|}\blacktriangleright}^\varphi = E_{z_{|ji|}\blacktriangleright}^\varphi$. Thus $\mathfrak{U}_{\{ij\}}(\varphi) = \{\varphi\}$, rendering (5.63) trivially true.

Equipped with this terminology, we state the following criterion for word independence of spanning sets, which may be viewed as a reformulation of Matsumoto's equivalence as a property of exponent sets.

Proposition 5.14. *Let $w, \tilde{w} \in \mathcal{W}^*$ be two reduced words representing the same element $s = r(w) = r(\tilde{w})$ in \mathcal{W} . Assume further that $S \subseteq \mathcal{E}(s)$ is a relator stable exponent set for given ground ring Λ . Then we have for the Λ -spans $\mathfrak{S}_w^{S,+} = \mathfrak{S}_{\tilde{w}}^{S,+}$. Further, if the elements of $\mathfrak{B}_w^{S,+}$ are linearly independent, then so are those of $\mathfrak{B}_{\tilde{w}}^{S,+}$.*

The analogous assertions hold for the $\mathfrak{B}_w^{S,-}$ and $\mathfrak{B}_w^{S,\pm}$ and their spans.

Proof. By Matsumoto's Theorem (see Corollary 2.1) the exists a sequence of reduced words, starting with w and ending with \tilde{w} , each of which is obtained from the previous one by replacing a substring of the form $z_{|ij|}$ by $z_{|ji|}$, vice versa. By transitivity of the assertions, it suffices to consider the case in which \tilde{w} is obtained from w by one such substitution. That is, $w = u \cdot z_{|ij|} \cdot v$ and $\tilde{w} = u \cdot z_{|ji|} \cdot v$ as reduced decompositions. Thus, with $a = r(u)$ and $b = r(v)$, we find $s = a \cdot s_{|ij|} \cdot b$ and $l(s) = l(a) + m_{ij} + l(b)$. In particular, (a, b) is a relator pair for s .

Suppose now $\psi \in S$ so that $E_{w\blacktriangleright}^\psi$ is a spanning element for $\mathfrak{S}_w^{S,+}$ and let $(\chi, \varphi, \xi) = \mathcal{C}_{(a,b)}^*(\psi)$. Applying (5.26) twice, we find $E_{w\blacktriangleright}^\psi = E_{u\blacktriangleright}^\chi \cdot \Gamma_a(E_{z_{|ij|}\blacktriangleright}^\varphi) \cdot \Gamma_{a \cdot s_{|ij|}}(E_{v\blacktriangleright}^\xi)$. Expanding $E_{z_{|ij|}\blacktriangleright}^\varphi$ as in (5.61) into a sum of $E_{z_{|ji|}\blacktriangleright}^{\varphi'}$, each term can then be rewritten as some $E_{\tilde{w}\blacktriangleright}^\psi$ using again (5.26). More

precisely, writing $\psi(\varphi') = (\mathcal{Q}_{(a,b)}^*)^{-1}(\chi, \varphi', \xi)$ for fixed χ and ξ we find

$$E_{w\blacktriangleright}^\psi = E_{w\blacktriangleright}^{\psi(\varphi)} = \sum_{\varphi'} \mathfrak{v}_{\varphi'}^\varphi \cdot E_{\tilde{w}\blacktriangleright}^{\psi(\varphi')} . \quad (5.64)$$

Condition (5.63) finally ensures that all basis elements on the right side are in $\mathfrak{B}_{\tilde{w}\blacktriangleright}^{S,+}$ whenever $\mathfrak{v}_{\varphi'}^\varphi \neq 0$. Thus $E_{w\blacktriangleright}^\psi \in \mathfrak{S}_{\tilde{w}\blacktriangleright}^{S,+}$ and, therefore, $\mathfrak{S}_{w\blacktriangleright}^{S,+} \subseteq \mathfrak{S}_{\tilde{w}\blacktriangleright}^{S,+}$. The opposite inclusion is implied by \mathfrak{V} -symmetry.

Now, for fixed χ and ξ , (5.64) yields a transformation of the spanning elements $B_w^{(\chi, \xi)} = \{E_{w\blacktriangleright}^\psi : \psi(\varphi) \in S\}$ to those of $B_{\tilde{w}}^{(\chi, \xi)}$ via the restricted involution expressed by $\mathfrak{v}_{\varphi'}^\varphi$. Thus, $B_w^{(\chi, \xi)}$ is a basis if and only if $B_{\tilde{w}}^{(\chi, \xi)}$ is a basis, proving the second statement. Similarly, applications of Υ and Ω yield the assertions for the other types of spans. \square

As an example, we may consider the following basic type of exponent sets. Given two sets $V_1, V_e \subseteq \mathbb{N}_0$ of non-negative integers and $s \in \mathcal{W}$, define

$$V^{[s]} = \{\psi \in \mathcal{E}(s) : \psi(\alpha) \in V_{d_\alpha}, \forall \alpha \in \mathcal{N}(s)\} \quad \text{and} \quad V^{[ij]} = V^{[s_{ij}]} . \quad (5.65)$$

Suppose that $s = r \cdot t$ with $l(s) = l(r) + l(t)$. Observe that the bijection $\mathcal{E}(s) \rightarrow \mathcal{E}(r) \times \mathcal{E}(t)$ from (5.22) restricts to a bijection $V^{[s]} \rightarrow V^{[r]} \times V^{[t]}$. Conversely, it is not hard to show that any family of exponent sets $\{V^{[s]} : s \in \mathcal{W}\}$ with this *splitting property* has to be of the form in (5.65). For convenience, we also introduce notation for the complements

$$\bar{V}^{[s]} = \mathcal{E}(s) \setminus V^{[s]} = \{\psi \in \mathcal{E}(s) : \exists \alpha \in \mathcal{N}(s) \text{ with } \psi(\alpha) \notin V_{d_\alpha}\} , \quad (5.66)$$

and, analogously, $\bar{V}^{[ij]} = \bar{V}^{[s_{ij}]} .$

Corollary 5.15.

Let $w, \tilde{w} \in \mathcal{W}^*$ with $s = r(w) = r(\tilde{w})$ and $\{V^{[s]}\}$ be a family of exponent sets as in (5.65).

- i) Suppose \mathfrak{V} maps the Λ -span $\mathfrak{S}_{z_{ij}\blacktriangleright}^{V^{[ij]},+}$ to itself for all $1 \leq i, j \leq n$. Then we have for the Λ -spans $\mathfrak{S}_{w\blacktriangleright}^{V^{[s]},+} = \mathfrak{S}_{\tilde{w}\blacktriangleright}^{V^{[s]},+}$. Moreover, $\mathfrak{B}_{w\blacktriangleright}^{V^{[s]},+}$ is linearly independent if and only if $\mathfrak{B}_{\tilde{w}\blacktriangleright}^{V^{[s]},+}$ is.
- ii) Suppose \mathfrak{V} maps the Λ -span $\mathfrak{S}_{z_{ij}\blacktriangleright}^{\bar{V}^{[ij]},+}$ to itself for all $1 \leq i, j \leq n$. Then we have for the Λ -spans $\mathfrak{S}_{w\blacktriangleright}^{\bar{V}^{[s]},+} = \mathfrak{S}_{\tilde{w}\blacktriangleright}^{\bar{V}^{[s]},+}$. Moreover, $\mathfrak{B}_{w\blacktriangleright}^{\bar{V}^{[s]},+}$ is linearly independent if and only if $\mathfrak{B}_{\tilde{w}\blacktriangleright}^{\bar{V}^{[s]},+}$ is.

The analogous statements hold for opposite directions and respective bases of $U_{q,\Lambda}^-$.

Proof. Suppose (a, b) is a relator pair for s . Then the aforementioned splitting property implies that $V_{(a,b)}^{[s]} = V^{[a]} \times V^{[ij]} \times V^{[b]}$. So, in the case of i), (5.63) reduces to the localized condition that $\varphi \in V^{[ij]}$ and $\mathfrak{v}_{\varphi'}^\varphi \neq 0$ implies $\varphi' \in V^{[ij]}$, which is equivalent to the condition that \mathfrak{V} preserves $\mathfrak{S}_{z_{ij}\blacktriangleright}^{V^{[ij]},+}$. Alternatively, this can be inferred from the remark following in (5.63), observing that $U_{\chi, \xi}^{S,a,b} = \mathfrak{S}_{z_{ij}\blacktriangleright}^{V^{[ij]},+}$ whenever $(\chi, \xi) \in V^{[a]} \times V^{[b]}$ and zero otherwise.

For the complementary statement in *ii)*, note that

$$\bar{V}_{(a,b)}^{[s]} = \bar{V}^{[a]} \times \mathcal{E}(s_{|ij|}) \times \mathcal{E}(b) \cup \mathcal{E}(a) \times \bar{V}^{[s_{|ij|}]} \times \mathcal{E}(b) \cup \mathcal{E}(a) \times \mathcal{E}(s_{|ij|}) \times \bar{V}^{[b]}.$$

If (χ, φ, ξ) is either in the first or third set, the condition (5.63) is trivially true since \mathfrak{O} preserves $U_{\chi, \xi}^{S,a,b} = U_{q,\Lambda^{(i,j)}}^+$. If (χ, φ, ξ) is in the second set the condition (5.63) is, again, equivalent to \mathfrak{O} -stability of $U_{\chi, \xi}^{S,a,b} = \mathfrak{S}_{z_{|ij|}}^{\bar{V}^{[ij]}, +}$. \square

Observe, finally, that Corollary 5.15 can be slightly strengthened by noting that only pairs (i, j) need to be considered for which there is a reduced decomposition $s = a \cdot s_{|ij|} \cdot b$.

5.8. General Ordering Bases and Module Restrictions. The aim of this section is to extend the special bases from Proposition 5.11 to bases for any convex ordering and the entire U_q algebra. We will, further, consider extensions of scalars to ground rings Λ or Λ' as in (3.15) for given homomorphisms $f : \Lambda_{q,e^*} \rightarrow \Lambda$ and $f' : \Lambda'_{q,e^*} \rightarrow \Lambda'$. Since the associated functor $-\otimes_{\Lambda_{q,e^*}} \Lambda$ maps free modules to free modules, any Λ_{q,e^*} -basis of U_q^\square is also a Λ -basis for $U_{q,\Lambda}^\square$, with $\square \in \{+, -, 0, \geq 0, \leq 0\}$. Similar considerations apply to $U_{q,\Lambda'}$.

We begin by adapting Rosso's arguments for the tensor decomposition of the full quantum algebra over a field to our situation for rings Λ or Λ' . Lemma 2 in [Ros88] proves that $U_{q,\Lambda}^0$ is isomorphic to $\Lambda[\{K_i^\pm\}] \cong \Lambda[\mathbb{Z}^n]$ and admits a Λ -basis is given by

$$\mathfrak{B}^0 = \{K^\nu : \nu \in \mathbb{Z}^\Delta\} \quad (5.67)$$

for Λ a field. The argument there does not require the latter assumption so that the statement holds for any integral domain Λ as a ground ring. We next consider multiplication maps, such as the following.

$$\begin{aligned} U_{q,\Lambda}^0 \otimes_\Lambda U_{q,\Lambda}^+ &\rightarrow U_{q,\Lambda}^{\geq 0} \\ U_{q,\Lambda'}^+ \otimes_{\Lambda'} U_{q,\Lambda'}^0 \otimes_{\Lambda'} U_{q,\Lambda'}^- &\rightarrow U_{q,\Lambda'} \end{aligned} \quad (5.68)$$

Obvious additional maps are given by permuting the tensor factors as well as swapping the algebras $U_q^+ \leftrightharpoons U_q^-$ and $U_q^{\geq 0} \leftrightharpoons U_q^{\leq 0}$.

To show surjectivity of these maps, it suffices to show that their respective images are preserved under the left regular action of the respective target algebra, which is immediate from the commutation relations (3.13) and (3.13). For example, applying E_k to a general element in the image of the first type of map yields with $E_k(K^\nu E_{i_1} \dots E_{i_N}) = q^{-(\nu|a_k)}(K^\nu E_k E_{i_1} \dots E_{i_N})$ again an element in the image. Similarly, for the second type of maps, $F_k(E_{i_1} \dots E_{i_N} K^\nu F_{j_1} \dots F_{j_M})$ can be expressed as a Λ'_q -combination of summands of the form $E_{i_1} \dots E_{i_N} K^\nu F_k F_{j_1} \dots F_{j_M}$ and $E_{i_1} \dots \widehat{E_{i_s}} \dots E_{i_N} K^\mu F_{j_1} \dots F_{j_M}$, where $\mu \in \mathbb{Z}^\Delta$ and the hat denotes omission whenever $i_s = k$.

The proofs of Proposition 1 and 2 in [Ros88] for injectivity assume that the quantum algebra is defined over a field. As before, a straightforward inspection of the arguments there immediately shows that the only assumptions needed are the existence of a free basis over the ground ring

as well as a grading-preserving coalgebra structure. The former is assured by Proposition 5.11 and Corollary 5.12 above. The latter is clearly also defined over the respective integral domains, and its grading properties are listed in (4.6) and (4.7). These observations are summarized next.

Lemma 5.16 ([Ros88]). *The multiplication maps from (5.68) and their obvious variants are isomorphisms of free Λ or Λ' modules.*

We next combine results from Sections 5.7 and 5.6 to obtain PBW bases for the $U_{q,\Lambda}^\pm$ algebras for general orderings.

Proposition 5.17. *Suppose $w \in \mathcal{W}_m^*$ is any reduced word of maximal length. Then both $\mathfrak{B}_{w\blacktriangleright}^+$ and $\mathfrak{B}_{\blacktriangleleft w}^+$ are Λ -bases for $U_{q,\Lambda}^+$. Similarly, $\mathfrak{B}_{w\blacktriangleright}^-$ and $\mathfrak{B}_{\blacktriangleleft w}^-$ are Λ -bases for $U_{q,\Lambda}^-$.*

Proof. Proposition 5.11 ensures that there is at least one reduced word $w^* \in \tau^{-1}(s_\bullet)$ for which $\mathfrak{B}_{w^*\blacktriangleright}^+$ and $\mathfrak{B}_{\blacktriangleleft w^*}^+$ are Λ_{q,e^*} -bases of U_q^+ and hence also Λ -bases of $U_{q,\Lambda}^+$. Suppose now $w \in \mathcal{W}_m^*$ is another presentation of s_\bullet .

The full exponent set $S = \mathcal{E}(s_\bullet) = \mathbb{N}_0^{\Phi^+}$ is clearly of the form (5.65) with $V_d = \mathbb{N}_0$. Moreover, as already noted in the proof of Corollary 5.15, we have for the full Λ -span $\mathfrak{S}_{z_{ij}\blacktriangleright}^+ = \mathfrak{S}_{z_{ij}\blacktriangleright}^+ = U_{q,\Lambda}^{(i,j)+}$. Thus, the second condition of Corollary 5.15 is also fulfilled, which implies that $\mathfrak{S}_{w^*\blacktriangleright}^+ = \mathfrak{S}_{w\blacktriangleright}^+$. Thus $\mathfrak{B}_{w\blacktriangleright}^+$ also spans $U_{q,\Lambda}^+$ and is linearly independent. The second case for U_q^- follows readily from Corollary 5.12 in the same manner. \square

Clearly, Lemma 5.16 implies that, given bases for the tensor factors on the left sides of (5.68), their products as spanning sets yield bases for the quantum algebras on the right side. This, combined with Proposition 5.17 and the observation for U_q^0 in (5.67), now yields the following general construction of PBW bases.

Theorem 5.18. *Let $w, u \in \mathcal{W}_m^*$ be any reduced words of maximal length and Λ, Λ' rings as in (3.15). Then any of the spanning sets $\mathfrak{B}_{w\blacktriangleright}^+ \cdot \mathfrak{B}^0$, $\mathfrak{B}^0 \cdot \mathfrak{B}_{w\blacktriangleright}^+$, $\mathfrak{B}_{\blacktriangleleft w}^+ \cdot \mathfrak{B}^0$, or $\mathfrak{B}^0 \cdot \mathfrak{B}_{\blacktriangleleft w}^+$ is a Λ -basis for $U_{q,\Lambda}^{\geq 0}$.*

The analogous statement holds for $U_{q,\Lambda}^{\leq 0}$. Moreover, the spanning set $\mathfrak{B}_{w\blacktriangleright}^+ \cdot \mathfrak{B}^0 \cdot \mathfrak{B}_{u\blacktriangleright}^-$ and any reordering of the factor sets as well as any direction reversals (24 versions in total) is a Λ' -basis for $U_{q,\Lambda'}$.

One elementary but useful application of this theorem arises when subspaces constructed over fields need to be pulled back to submodules over the other rings. More concretely, suppose \mathfrak{B} is any Λ -basis as above for one of the algebras $U_{q,\Lambda}^\square$. Assume that Λ is an integral domain and \mathbb{F} its field of fractions so that we may view $U_{q,\Lambda}^\square$ as a subalgebra of $U_{q,\mathbb{F}}^\square$.

Suppose now $\mathcal{I} \subset \mathcal{B}$ is a subset of basis elements and denote $I = \langle \mathcal{I} \rangle$ its Λ -span in $U_{q,\Lambda}^\square$. Similarly, let $I_{\mathbb{F}} \cong I \otimes \mathbb{F}$ be the respective \mathbb{F} -span in $U_{q,\mathbb{F}}^\square$. Since I is a direct summand, we then immediately have

$$I = I_{\mathbb{F}} \cap U_{q,\Lambda}^\square. \quad (5.69)$$

6. SKEW-COMMUTATIVE ALGEBRAS AT ROOTS OF UNITY

In this section, we turn to the specialization $q \mapsto \zeta$ of a quantum group to a primitive \mathbb{k} -th root of unity ζ . The main focus of our discussion are the primitive power generators $X_\alpha = E_\alpha^{\ell_\alpha}$, where ℓ_α is the order of $\zeta^{(\alpha|\alpha)}$. For odd \mathbb{k} , the algebra \mathcal{Z}^+ generated by these elements lies in the center of U_ζ and has been studied, for example, in [DCK90, DCP93].

Our treatment here includes, in addition, *all* even $\mathbb{k} > 2e^*$. In many of these cases, the generators X_α commute only up to signs with other elements in U_ζ . Depending on Lie type and the congruence of \mathbb{k} modulo 8, the algebra \mathcal{Z}^+ may be central, commutative but not central, or neither. Besides a classification of these cases, we also prove word independence of respective partial subalgebras \mathcal{Z}_w^+ (with one exception) and invariance under automorphisms. Various types of implied and induced ideals are discussed as well.

6.1. Quantum Numbers and Rings at Roots of Unity. Assume a root system with integers $d_i \in \{1, 2, 3\}$ as in Section 2.1 such that $1 = \min_i \{d_i\}$ and $e = \max_i \{d_i\}$. Fix an integer $\mathbb{k} \in \mathbb{N}$ with

$$\mathbb{k} \notin \{1, 2, e, 2e\}. \quad (6.1)$$

The ring $\mathbb{Z}[\zeta]$ of cyclotomic integers may be understood as the quotient of either $\mathbb{Z}[q]$ or $\mathbb{Z}[q, q^{-1}]$ by the principal ideal $J_\mathbb{k} = (\Phi_\mathbb{k}(q))$ of the \mathbb{k} -th cyclotomic polynomial. Equivalently, we may view $\mathbb{Z}[\zeta]$ as the ring of integers of the cyclotomic field $\mathbb{Q}(\zeta)$, considered as a subfield of \mathbb{C} for a given a choice of a primitive \mathbb{k} -th root of unity $\zeta \in \mathbb{C}$. That is, the kernel of the map $\mathbb{Z}[q] \rightarrow \mathbb{C} : q \mapsto \zeta$ is precisely $J_\mathbb{k}$ (see, for example, [Was82] for details). We use the same notation as in (3.1) for the images of the quantum numbers in $\mathbb{Z}[\zeta]$.

Denote by $\ell = \mathbb{k}/\gcd(\mathbb{k}, 2)$ the order of ζ^2 . For integers d_i as above write $\zeta_i = \zeta^{d_i}$ and denote by $\mathbb{k}_i = \mathbb{k}/\gcd(\mathbb{k}, d_i)$ the order of ζ_i . Similarly, we set $\ell_i = \ell/\gcd(\ell, d_i) = \mathbb{k}/\gcd(\mathbb{k}, 2d_i) = \mathbb{k}_i/\gcd(\mathbb{k}_i, 2)$ to be the order of $\zeta_i^2 = \zeta^{2d_i} = \zeta^{(\alpha_i|\alpha_i)}$. Note also that for short roots we have $\ell_i = \ell$ since, by definition, $d_i = 1$.

Condition (6.1) is indeed equivalent to requiring $\zeta_i^2 \neq 1$ for all $i \in \{1, \dots, n\}$. This ensures that the factor $(\zeta_i - \zeta_i^{-1})^{-1}$ occurring in (3.13) exists in $\mathbb{Q}(\zeta)$ so that the specialization $q \mapsto \zeta$ for quantum groups is well-defined. Similarly, (6.1) implies that $m = \ell_i$ is precisely the smallest positive integer for which $[m]_i = 0$.

The following additional powers of ζ will frequently appear in subsequent formulae.

$$\xi_i = \zeta_i^{\ell_i} = \zeta^{d_i \ell_i} = \zeta^{\text{lcm}(d_i, \ell)} \quad \xi_i = \xi_i^{\ell_i} = \zeta_i^{\ell_i^2} \quad \xi_i = \zeta_i^{\binom{\ell_i}{2}} \quad (6.2)$$

We note that $\xi_i, \xi_i \in \{+1, -1\}$. Specifically, $\xi_i = -1$ if and only if \mathbb{k}_i is even, and $\xi_i = -1$ if and only if $\mathbb{k}_i \equiv 2 \pmod{4}$. Moreover, ξ_i is a primitive fourth root of unity ($\xi_i^2 = -1$) if $\mathbb{k}_i \equiv 0 \pmod{4}$, we have $\xi_i = -1$ if $\mathbb{k}_i \equiv 6 \pmod{8}$, and $\xi_i = 1$ in all other cases. These relations also imply the identity $\xi_i^2 = (-1)^{\ell_i-1}$ in all cases.

As for generic q , if $d_i = 1$, we will often drop the i subscript from the roots of unity appearing in (6.2) as well as the quantum numbers. In any of these, we may replace any subscript i by a subscript α if $\alpha \in \Phi^+$ is in the \mathcal{W} -orbit of α_i . So, for example, $\ell_\alpha = \ell_i$, $\mathbb{k}_\alpha = \mathbb{k}_i$, $\zeta_\alpha = \zeta_i$, and $\xi_\alpha = \xi_i$. Analogously, for a reduced word $w \in \mathcal{W}^*$ we will write $\ell_w = \ell_i$, $\zeta_w = \zeta_i$, $\xi_w = \xi_i$, and so forth if $i = \tau(w)$ or, equivalently, $\gamma(w)$ is in the \mathcal{W} -orbit of α_i .

The commutation relations in (3.13) for the standard definition of quantum groups also contain denominators $(q_i - q_i^{-1})^{-1}$, which *a priori* map only to $\mathbb{Q}(\zeta)$ but not $\mathbb{Z}[\zeta]$. A slightly sharper statement asserts that the image $(\zeta_i - \zeta_i^{-1})^{-1}$ is an element of the smaller ring $\mathbb{Z}[\zeta, \frac{1}{\ell_i}] \subset \mathbb{Q}(\zeta)$. This follows readily from the following formula, in which $r = \ell / \gcd(m, \ell)$ is the order of ζ^{2m} :

$$(\zeta^m - \zeta^{-m})^{-1} = -\frac{1}{r} \sum_{j=0}^{r-2} (r-j-1) \zeta^{m(2j+1)} \in \mathbb{Z}[\zeta, \frac{1}{r}] \subset \mathbb{Q}(\zeta). \quad (6.3)$$

For non-prime orders \mathbb{k} , smaller denominators may be chosen as, for example, $(\zeta - \zeta^{-1})^{-1} \in \mathbb{Z}[\zeta, \frac{1}{3}]$ if $\ell = \mathbb{k} = 9$. The use of singularized generators as in (5.10) would avoid these denominators.

Aside from vanishing quantum numbers, the specialization also leads to additional units. For example, if r and ℓ_i are coprime, then $[r]_i$ is a unit in $\mathbb{Z}[\zeta]$. More generally, for any $m \in \mathbb{N}$ and $d = \gcd(m, \ell_i)$ we have the factorization $[m]_i = [d]_i \cdot u$, where u is a (cyclotomic) unit in $\mathbb{Z}[\zeta]$.

From (6.3) we also find that $([m]_i)^{-1} \in \mathbb{Z}[\zeta, \frac{1}{s}]$, where $s = \ell_i/d \in \mathbb{Z}$ is the order of $\zeta_i^{2m} = \zeta^{2d_i m}$. The next identity expresses the last non-vanishing factorial in terms of the order ℓ_i and the element $(\zeta_i - \zeta_i^{-1})$.

$$D_i = [\ell_i - 1]_i! = \ell_i \cdot \xi_i^{-1} \cdot (\zeta_i^{-1} - \zeta_i)^{-\ell_i+1} \quad (6.4)$$

For an integer $n > 1$, we introduce the following notation for specializations of the rings in (3.8), which will be used frequently in later sections. They are well-defined as subrings of $\mathbb{Q}(\zeta)$ if and only if $n \leq \ell$, which we will thus assume.

$$\begin{aligned} \Lambda_\zeta &= \mathbb{Z}[\zeta] & \Lambda_{\zeta,n} &= \mathbb{Z}\left[\zeta, \frac{1}{[n-1]!}\right] \\ \Lambda_\zeta^* &= \mathbb{Z}\left[\zeta, \frac{1}{\zeta - \zeta^{-1}}\right] & \Lambda_{\zeta,n}^* &= \mathbb{Z}\left[\zeta, \frac{1}{\zeta - \zeta^{-1}}, \frac{1}{[n-1]!}\right] \end{aligned} \quad (6.5)$$

The minimal rings Λ_{q,e^*} and Λ_{q,e^*}^* over which U_q^\pm and U_q are defined specialize to Λ_{ζ,e^*} and Λ_{ζ,e^*}^* , respectively, if and only if $\ell \geq e^*$, where $e^* = e + 1 \in \{2, 3, 4\}$ is as in (2.3) depending on Lie type. As a condition on \mathbb{k} , the bound on ℓ is equivalent to requiring (6.1) and, additionally, $\mathbb{k} \neq 4$ whenever $e = 3$. The latter exclusion refers to the G_2 quantum group, which is well-defined for $\mathbb{k} = 4$ but for which the braid automorphisms from Section 4.3 are not defined.

Previous remarks about cyclotomic units imply that $\Lambda_\zeta = \Lambda_{\zeta,n}$ and $\Lambda_\zeta^* = \Lambda_{\zeta,n}^*$ if j and ℓ are coprime ($\gcd(\ell, j) = 1$) for all $1 \leq j \leq n$. This is, of course, the case if ℓ itself is a prime or if $n = 1$.

Consequently, we have $\Lambda_{\zeta, \mathbf{e}^*} = \Lambda_\zeta$ and $\Lambda_{\zeta, \mathbf{e}^*}^* = \Lambda_\zeta^*$ for types ADE and all $\ell > 1$, for types BCF if ℓ is odd, and for type G₂ if $\gcd(\ell, 6) = 1$.

As before, the relation $(\zeta_i - \zeta_i^{-1}) = [d_i](\zeta - \zeta^{-1})$ implies that $(\zeta_i - \zeta_i^{-1})^{-1} \in \Lambda_{\zeta, \mathbf{e}^*}^*$. The definition of an R -matrix in our setting will require all non-zero factorials to appear as denominators, thus requiring $n = \ell$. It is clear that all rings listed in (6.5) are subrings of $\Lambda_{\zeta, \ell}^*$. In fact, equations (6.3) and (6.4) imply the equality of rings

$$\Lambda_{\zeta, \ell}^* = \mathbb{Z}[\zeta, \frac{1}{\ell}].$$

The evaluation of quantum binomial coefficients at a root of unity also leads to vanishing patterns and factorizations of classical binomial coefficients that will be important later. In order to describe the former, we introduce functions $c_i : \mathbb{N}_0 \times \mathbb{N}_0 \rightarrow \{0, 1\}$ depending on ℓ_i as follows. For two integers $u, v \in \mathbb{N}_0$, write $u = a \cdot \ell_i + r$ and $v = b \cdot \ell_i + s$ with remainders $r, s \in \{0, \dots, \ell_i - 1\}$. Define then

$$c_i(u, v) = \left\lfloor \frac{u+v}{\ell_i} \right\rfloor - \left\lfloor \frac{u}{\ell_i} \right\rfloor - \left\lfloor \frac{v}{\ell_i} \right\rfloor = c_i(r, s) = \begin{cases} 1 & \text{for } r+s \geq \ell_i \\ 0 & \text{for } r+s < \ell_i. \end{cases} \quad (6.6)$$

Here $\lfloor x \rfloor$ is the floor function, which yields the greatest integer less than or equal to x . The expression clearly only depends on the classes $\bar{u}, \bar{v} \in \mathbb{Z}/\ell_i \mathbb{Z}$ and, as a map on these, is the standard 2-cocycle in $Z^2(\mathbb{Z}/\ell_i \mathbb{Z})$ generating $H^2(\mathbb{Z}/\ell_i \mathbb{Z})$.

Lemma 6.1. *With integers $u, v, r, s \in \mathbb{N}_0$ as above, the image of the quantum binomial coefficient in $\mathbb{Z}[\zeta]$ is given by the formula*

$$\left[\begin{matrix} u+v \\ u \end{matrix} \right]_i = \xi_i^{(as+br)} \cdot \xi_i^{ab} \cdot \left[\begin{matrix} r+s \\ r \end{matrix} \right]_i \cdot \binom{a+b}{a}, \quad (6.7)$$

$$\text{so that} \quad \left[\begin{matrix} u+v \\ u \end{matrix} \right]_i = 0 \quad \iff \quad c_i(u, v) \neq 0.$$

Both the quantum binomial coefficient in r and s at ζ_i and the classical binomial coefficient in a and b are elements in $\mathbb{Z}[\zeta]$ and the powers of ξ and $\bar{\xi}$ are signs ± 1 . The vanishing criterion follows from inspection of the quantum binomial term in r and s .

Importantly, this implies that if $u+v$ is a multiple of ℓ_i , then the expression is non-zero only if also both u and v are multiples of ℓ_i as well. In this case, $r = s = 0$ and the quantum binomial is given by the respective classical binomial up to a sign. A main ingredient in the derivation of (6.7) is that the quantum factorial for u in $\mathbb{Z}[q, q^{-1}]$ is divisible by $[\ell_i]^a$ and can thus be evaluated at ζ_i . More precisely, we have for $\ell_i > 1$ that

$$\left. \frac{[u]_{q_i}!}{[\ell_i]_{q_i}^a} \right|_{q_i \rightarrow \zeta_i} = \xi_i^{ar} \cdot \xi_i^{\binom{a}{2}} \cdot [r]_{\zeta_i}! \cdot a! \cdot \mathcal{D}_i^a, \quad (6.8)$$

where the vertical line indicates the map to $\mathbb{Z}[\zeta]$. We will later use the following basic consequence of Lemma 6.1 for the multinomial coefficients from (3.7).

Corollary 6.2. *Suppose $a_1, \dots, a_r \in \mathbb{N}_0$ such that $a_1 + \dots + a_r = \ell_i$. Then*

$$\left[\begin{matrix} a_1 + \dots + a_r \\ a_1, \dots, a_r \end{matrix} \right]_i = \begin{cases} 1 & \text{if } \exists s : a_s = \ell_i \text{ and } a_t = 0 \text{ for } t \neq s \\ 0 & \text{otherwise} \end{cases}$$

We will distinguish two types of specializations of quantum groups to roots of unity, one given by the minimal choice $n = e^*$ and the other by the maximal choice $n = \ell$. The notation is as follows, where $\square \in \{\geq 0, \leq 0, +, -\}$ and the indicated extensions of scalars are with respect to the evaluation maps $q \mapsto \zeta$ between the rings in (3.8) and those in (6.5).

$$\begin{aligned} U_\zeta^\square &= U_q^\square \otimes \Lambda_{\zeta, e^*} & U_\zeta &= U_q \otimes \Lambda_{\zeta, e^*}^\star & U_{\zeta, \mathbb{Q}}^\square &= U_q^\square \otimes \mathbb{Q}(\zeta) \\ U_{\zeta, \ell}^\square &= U_q^\square \otimes \Lambda_{\zeta, \ell} & U_{\zeta, \ell} &= U_q \otimes \Lambda_{\zeta, \ell}^\star \end{aligned} \tag{6.9}$$

Many properties established for quantum groups at generic q in previous sections extend without changes to these quantum groups at roots of unity.

Corollary 6.3. *The following statements hold.*

- i) *The algebras defined in (6.9) are free modules over their respective ground rings, with bases given as in Theorem 5.18.*
- ii) *The natural inclusions of algebras in (3.8) induce respective embeddings for the algebras in (6.9).*
- iii) *All gradings and automorphisms defined in Section 4 factor into well-defined gradings and automorphisms for the algebras in (6.9).*

The first assertion is obvious, since all modules are free. Similarly, the existence of compatible free bases for the natural inclusions assures that the induced maps are also injective, mapping basis elements to each other. The last assertion is clear as well, since all automorphisms are, in particular, module isomorphisms or module anti-isomorphisms with respect to the conjugations on the ground rings.

In later discussions, we will consider root lattice vectors $\ell\alpha = \ell_\alpha \cdot \alpha$ for a given root $\alpha \in \Phi$ in a connected root system Φ of finite type and fixed $\ell > 1$. Let $e = \max\{d_i\}_i \in \{1, 2, 3\}$ as in Section 2.1. If e and ℓ are coprime, then all $\ell_\alpha = \ell$ so that $\ell\alpha = \ell \cdot \alpha$. Conversely, if $e > 1$ and divides ℓ , we have $\ell_\alpha = d_\alpha^{-1}\ell$ so that $\ell\alpha = \ell \cdot \check{\alpha}$ in our convention for coroots.

The collection $\Phi_\ell = \{\ell\alpha = \ell_\alpha \cdot \alpha : \alpha \in \Phi\}$ thus has the following properties.

Lemma 6.4. *We have $\Phi_\ell = \ell \cdot \Phi$ if $\gcd(e, \ell) = 1$ and $\Phi_\ell = \ell \cdot \check{\Phi}$ if $\gcd(e, \ell) > 1$, where $\check{\Phi}$ denotes the coroot system. Thus Φ_ℓ is itself a root system that is canonically isomorphic to either Φ or $\check{\Phi}$.*

The Weyl group \mathcal{W} of Φ preserves Φ_ℓ and its action coincides with that of the intrinsic Weyl group of Φ_ℓ in the sense that $s_{\ell\alpha}(\ell\beta) = \ell(s_\alpha(\beta)) = s_\alpha(\ell\beta)$. Moreover, the set $\Delta_\ell = \{\ell_i\alpha_i\}_i$ is a basis of Φ_ℓ .

The assertions are immediate from the preceding discussion, the fact that uniform scaling of a root system does not change its isomorphism class, and well-known properties of coroot systems.

6.2. Skew-Commutation Relations of Primitive Power Generators. In Lemma 3.1 of [DCK90] De Concini and Kac observe that ℓ -th powers of E_α and F_α generators commute with other generators of U_ζ up to unit factors. In this section, we will sharpen this result to include the ℓ_α -th powers, where $\ell_\alpha = \ell / \gcd(\ell, d_\alpha)$. This extension is required for our later discussion since for $\ell_\alpha < \ell$, certain subsets of $E_\alpha^{\ell_\alpha}$ elements will generate Hopf ideals, while the E_α^ℓ elements generally do not give rise to Hopf ideals.

We provide here an explicit and conceptual proof of the commutation relations based on the ϕ^\pm -gradings introduced in Section 4.1 and their \mathcal{W} -covariance properties. Moreover, statements will be in terms of word-dependent generators, which allows us to study independence of the relations of root orderings.

The function κ , defined in (6.10), keeps track of the signs in commutation relations. Here $\xi_\alpha = \zeta_\alpha^{\ell_\alpha} = \zeta^{d_\alpha\ell_\alpha} \in \{+1, -1\}$ is as in (6.2) and $\check{\alpha} = d_\alpha^{-1}\alpha$ denotes the coroot, implying the last equality below.

$$\kappa : \Phi \times \mathbb{F}_2^\Delta \rightarrow \{+1, -1\} \quad (\alpha, \bar{\mu}) \mapsto \kappa(\alpha, \bar{\mu}) = \zeta^{\ell_\alpha(\alpha|\bar{\mu})} = \xi_\alpha^{(\check{\alpha}|\bar{\mu})}. \quad (6.10)$$

Note, the respective map $\Phi \times \mathbb{Z}^\Delta \rightarrow \{+1, -1\}$, given by the same formula for $(\alpha, \mu) \in \Phi \times \mathbb{Z}^\Delta$, is a homomorphism in the μ argument onto the group $\{+1, -1\} \cong \mathbb{Z}/2\mathbb{Z}$. The map thus factors into $\Phi \times \mathbb{F}_2^\Delta$ as implied. It is also immediate from $\ell_\alpha = \ell_{s(\alpha)}$ that κ is \mathcal{W} -bivariant in the sense that $\kappa(s(\alpha), s(\bar{\mu})) = \kappa(\alpha, \bar{\mu})$ for all $s \in \mathcal{W}$.

Note that $\kappa(\alpha, \bar{\mu}) = 1$ for all α and $\bar{\mu}$ if ℓ is odd since in this case $\xi_\alpha = 1$. The next lemma discusses several cases for even ℓ .

Lemma 6.5. *Let $\ell = 2\ell$. Suppose ℓ is odd or Φ is of oddly laced Lie type ADEG. Then $\kappa(\alpha, \bar{\mu}) = \varepsilon(\check{\alpha}, \bar{\mu})$ for all $\alpha \in \Phi$ and $\bar{\mu} \in \mathbb{F}_2^\Delta$, where ε is the bihomomorphism*

$$\varepsilon : \mathbb{F}_2^\Delta \times \mathbb{F}_2^\Delta \longrightarrow \{+1, -1\} : (\bar{v}, \bar{\mu}) \mapsto \varepsilon(\bar{v}, \bar{\mu}) = (-1)^{(\bar{v}|\bar{\mu})}, \quad (6.11)$$

and $\check{\alpha}$ is the image of α under the canonical map $\Phi \rightarrow \mathbb{Z}^\Delta \rightarrow \mathbb{F}_2^\Delta$.

If ℓ is even ($\ell \equiv 0 \pmod{4}$) and Φ is of Lie type B_n, C_n , or F_4 with $n \geq 2$, then

$$\kappa(\alpha, \bar{\mu}) = (-1)^{(\check{\alpha}|\bar{\mu})}, \quad (6.12)$$

which is not additive in the first argument and, thus, does not extend to a bihomomorphic lattice map.

The assertion for oddly laced types follows from the observations that in these cases $\xi_\alpha = -1$ and $(\check{\alpha}|\bar{\mu}) \equiv (\alpha|\bar{\mu}) \pmod{2}$. Moreover, for types B, C and F and odd ℓ one verifies $\xi_\alpha = (-1)^{d_\alpha}$. Note that for B the form $(\cdot|\cdot)$ is always even so that $\kappa = \varepsilon = 1$, given odd ℓ .

For the doubly laced types BCF but with even ℓ , equation (6.12) follows from the fact that $\xi_\alpha = -1$ for all $\alpha \in \Phi^+$.

Failure of additivity for even ℓ is readily verified for type B_2 , which occurs as a subsystem in the listed Lie types. If α_1 is the short and α_2 the long root (so, $d_1 = 1$ and $d_2 = 2$) one finds that $\kappa(\alpha_1, \bar{\alpha}_1) = 1$, $\kappa(\alpha_2, \bar{\alpha}_1) = -1$, and $\kappa(\alpha_1 + \alpha_2, \bar{\alpha}_1) = 1$ as $\alpha_1 + \alpha_2$ is also a short root. So, clearly, $\kappa(\alpha_1, \bar{\alpha}_1)\kappa(\alpha_2, \bar{\alpha}_1) \neq \kappa(\alpha_1 + \alpha_2, \bar{\alpha}_1)$, obstructing an extension as a lattice map to $\mathbb{Z}^\Delta \supset \Phi^+$.

Since κ will be more frequently used in combination with the \circ^\pm -gradings from (4.3) and (4.4) it is convenient to introduce the following functions $\hat{\kappa}(\cdot, \cdot)^+$ and $\hat{\kappa}(\cdot, \cdot)^-$. Here $w \in \mathcal{W}^*$ is a (non-empty) reduced word and b is a \circ^\pm -homogeneous element in either U_q or U_ζ :

$$\hat{\kappa}(w, b)^\pm = \kappa(\gamma(w), \circ^\pm(b)). \quad (6.13)$$

For w and b as above, suppose $w_i \cdot w$ is also a reduced word. Using $\gamma(w_i \cdot w) = s_i(\gamma(w))$, (4.26), and \mathcal{W} -bivariance of $\kappa(\cdot, \cdot)$ we infer that for any i ,

$$\hat{\kappa}(w_i \cdot w, \Gamma_i(b))^\pm = \hat{\kappa}(w, b)^\pm. \quad (6.14)$$

For a given non-empty, reduced word $w \in \mathcal{W}^*$, $\ell_w = \ell_{\tau(w)} = \ell_{\gamma(w)}$, and root $\alpha \in \Phi$ we now introduce the notation for the respective *primitive power generators*:

$$X_w = E_w^{\ell_w}, \quad Y_w = F_w^{\ell_w}, \quad \text{and} \quad L_\alpha = K^{(\ell_\alpha \cdot \alpha)}, \quad (6.15)$$

where K^μ is as in (4.1) and $\ell_\alpha \cdot \alpha$ indicates the integer multiple of the root vector. That is, $L_i = L_{\alpha_i} = K_i^{\ell_i}$. For $\mu \in \mathbb{Z}^\Delta$, $\mu_i = \mu(\alpha_i)$, and $\eta = \sum_i \mu_i \ell_i \alpha_i$ set

$$L^\mu = L_1^{\mu_1} \cdot \dots \cdot L_n^{\mu_n} = K^\eta. \quad (6.16)$$

The w -grading in \mathbb{Z}^Δ from Section 4.1 on a generator for $\alpha = \gamma(w)$ is then given by

$$\mathbb{w}(X_w) = \ell_\alpha \cdot \alpha = \frac{\ell}{\gcd(\ell, d_\alpha)} \cdot \alpha = \begin{cases} \ell \cdot \alpha & \text{if } \gcd(\ell, e) = 1 \\ \ell \cdot \check{\alpha} & \text{if } \gcd(\ell, e) = e. \end{cases} \quad (6.17)$$

That is, if e and ℓ are coprime the generators are graded by the lattice of $\ell \cdot \Phi \cong \Phi$. However, if e divides ℓ the grading is given by the ℓ -scaled (but equivalent) version of the lattice of the coroot system $\check{\Phi}$. So, for example, if ℓ is even and Φ is of type B_n , then the X_w generators admit a natural grading in the root lattice for C_n .

Recall that in Lusztig's construction of the divided power algebras the elements X_w and Y_w are mapped to zero in the specialization from a Λ_q -algebra to a Λ_ζ -algebra. Since the Γ_j are

algebra automorphisms and the ℓ_α are \mathcal{W} -invariant, the definition in (5.1) readily translates into

$$X_w = \Gamma_{w^\flat}(X_{\tau(w)}) = \Gamma_u(X_v) \quad \text{and} \quad Y_w = \Gamma_{w^\flat}(Y_{\tau(w)}) = \Gamma_u(Y_v), \quad (6.18)$$

assuming $w = u \cdot v$ is reduced and $l(v) \geq 1$. Similarly, (4.24), (6.15), and the same \mathcal{W} -invariance imply

$$\Gamma_s(L_\alpha) = L_{s(\alpha)} \quad \text{for all } \alpha \in \Phi \text{ and } s \in \mathcal{W}. \quad (6.19)$$

Various other relations and properties for the E_w and F_w generators from Section 5.1 are easily extended to properties of the X_w and Y_w elements in the same manner. Remarkably, this includes the following identity that holds for all \mathbb{K} , due to the identity $\zeta^2 = (-1)^{\ell_i-1}$ mentioned in Section 6.1:

$$\Gamma_i(X_i) = -L_i^{-1}Y_i = -Y_iL_i^{-1} \quad \text{and} \quad \Gamma_i(Y_i) = -L_iX_i = -X_iL_i. \quad (6.20)$$

The following commutation relations for a rank 2 subsystem in U_q^+ are obtained from equations (c) and (i) in Section 5.3 and equations (a2) and (a6) in Section 5.4 of [Lus90b]. As in Section 2.7 above, assume for a pair of indices (i, j) that $-A_{ji} = d_i = 1$ and $-A_{ij} = d_j = e \in \{1, 2, 3\}$. The index notation for generators is as defined in (5.2) or, specialized to the rank 2 setting, in (5.19), where also translations to Lusztig's conventions are explained.

$$e = 1 \quad \left\{ \begin{array}{l} E_i^k E_j = q^{-k} E_j E_i^k + q^{-1} [k]_i E_{(ij)} E_i^{k-1} \end{array} \right. \quad (6.21)$$

$$e = 2 \quad \left\{ \begin{array}{l} E_j^k E_i = q^{-2k} E_i E_j^k + q^{-2} [k]_j E_{(ji)} E_j^{k-1} \\ E_i^k E_j = q^{2k} E_j E_i^k - q^{(k-1)} [k]_i E_i^{k-1} E_{(ji)} - q^{2(k-1)} [k]_i [k-1]_i E_i^{k-2} E_{(jij)} \end{array} \right. \quad (6.22)$$

$$e = 3 \quad \left\{ \begin{array}{l} E_j^k E_i = q^{-3k} E_i E_j^k + q^{-3} [k]_j E_{(ji)} E_j^{k-1} \\ E_i^k E_j = q^{3k} E_j E_i^k - q^{(k-1)} [k]_i E_i^{k-1} E_{(ji)} - q^{2(k-1)} [k]_i [k-1]_i E_i^{k-2} E_{(jiji)} \\ \quad - q^{3(k-1)} [k]_i [k-1]_i [k-2]_i E_i^{k-3} E_{(ijiji)} \end{array} \right. \quad (6.23)$$

Since all equations in (6.21) are well-defined equations in U_q^+ over Λ_{q,e^*} they descend to the same equations in U_ζ^+ or U_ζ via (6.9), upon specializing $q \mapsto \zeta$ with the previous assumptions in the order of ζ . If we set $k = \ell_i$ in the equations with an E_i^k -term and $k = \ell_j$ in equations with an E_j^k -term, only the first term on the right side of each equation remains, since $[\ell_i]_i = [\ell_j]_j = 0$ by the restriction in (6.1). The resulting relations can be summarized in the following lemma, where $X_s = X_{\alpha_s} = E_s^{\ell_s}$.

Lemma 6.6. *For all $r, s \in \{1, \dots, n\}$ we have $X_s E_r = \zeta^{\ell_s d_s A_{sr}} E_r X_s$.*

Proof. If $r \neq s$ and $A_{rs} \neq 0$ the ζ power is easily verified to be identical to the respective lead coefficient in the formulae in (6.21) for $(s, r) = (i, j)$ and $(s, r) = (j, i)$. If $A_{rs} = 0$, the relation is clear since E_s and E_r strictly commute to begin with. If $r = s$ we find $\zeta^{\ell_s d_s A_{ss}} = \zeta^{\ell_s d_s 2} = 1$ by definition and as desired. \square

Note, further, that the relations in (3.13) and (3.14) entail commutation relations $E_s^k K_r = q^{-k(\alpha_s|\alpha_r)} K_r E_s$ and $[E_s^k, F_r] = \delta_{sr}(q_s - q_s^{-1})^{-1} [k]_s E_s^{k-1} (q_s^{k-1} K_s - q_s^{-(k-1)} K_s^{-1})$ in U_q , which specialize for $k = \ell_i$ and $q \mapsto \zeta$ to

$$X_s F_r = F_r X_s \quad \text{and} \quad X_s K_r = \zeta^{\ell_s(\alpha_s|\alpha_r)} K_r X_s. \quad (6.24)$$

Proposition 6.7. *Let $w \in \mathcal{W}^*$ be a non-empty, reduced word and b a \circ -homogeneous element. Then*

$$X_w b = \hat{K}(w, b)^+ \cdot b X_w \quad \text{and} \quad Y_w b = \hat{K}(w, b)^- \cdot b Y_w.$$

Each relation holds in U_ζ as well as each U_ζ^\square with $\square \in \{+, -, \geq 0, \leq 0\}$, provided the elements in the relation belong to the respective algebra.

Proof. We begin by considering the relation for $X_s = X_{w_s}$ in the full algebra U_ζ . Since $\hat{K}(w_s, E_r)^+ = K(\gamma(w_s), \circ^+(E_r)) = K(\alpha_s, \alpha_r) = \zeta^{\ell_s(\alpha_s|\alpha_r)} = \zeta^{\ell_s d_s A_{sr}}$, Lemma 6.6 implies the desired relation for $b = E_r$. The calculation for $\hat{K}(w_s, K_r)^+$ is exactly the same so that, by the second identity in (6.24), the assertion also holds for $b = K_r$. Further, the first identity in (6.24) together with $\hat{K}(w_s, F_r)^+ = K(\gamma(w_s), \circ^+(F_r)) = K(\alpha_s, 0) = 1$ shows the commutation formula for $b = F_r$.

Next, observe that $\hat{K}(w, bc)^+ = \hat{K}(w, b)^+ \hat{K}(w, c)^+$ for \circ -homogeneous elements b and c because K is a homomorphism in the second argument and \circ is a grading. Thus, we can establish $X_s b = \hat{K}(w, b)^+ \cdot b X_s$ for any $b = b_1 \dots b_N$ with $b_j \in \{E_r, F_r, K_r\}$ by iteration and, by additivity, further extend the relation to all homogeneous $b \in U_\zeta$.

The proof for general $w \in \mathcal{W}^*$ follows now by induction in $l(w)$. Suppose w and $w_i \cdot w$ are reduced words so that $\Gamma_i(X_w) = X_{w_i \cdot w}$ by (6.18). Assume further the commutation relation holds for w and any homogeneous element in U_ζ . Applying the automorphism Γ_i to $X_w c = \hat{K}(w, c)^+ \cdot c X_w$ then yields $X_{w_i \cdot w} b = \hat{K}(w, c)^+ \cdot b X_{w_i \cdot w}$, where $b = \Gamma_i(c)$. From (6.14) we find that $\hat{K}(w, c)^+ = \hat{K}(w_i \cdot w, b)^+$. Since Γ_i is a bijection among homogeneous elements, this shows that the relation also holds for $w_i \cdot w$ and, hence, for all reduced words.

Note that (5.5) implies $\Omega(X_w) = Y_w$, and we have $\hat{K}(w, \Omega(b))^+ = \hat{K}(w, b)^-$ by (4.17). From this, the second relation with Y_w is inferred by applying Ω to the first, using also that the $\hat{K}(w, b)^\pm \in \{+1, -1\}$. Finally, the relations in U_ζ imply the same relations in the embedded algebras U_ζ^\square via Corollary 6.3.ii). \square

Define also $\hat{K}(\alpha, b)^\circ = K(\alpha, \mathbf{w}(b)) = K(\alpha, \circ^+(b))K(\alpha, \circ^-(b))$ for a root $\alpha \in \Phi$ and a fully homogeneous element b , where the second equality results from $\mathbf{w} = \circ^+ - \circ^-$, noted in Section 4.1. Equation (4.2) now immediately implies the following commutation relations for generators defined in

(6.15) and (6.16):

$$L_\alpha b = \hat{K}(\alpha, b)^\circ \cdot b L_\alpha \quad \text{and} \quad L^\mu b = \hat{K}(\mu, b)^\circ \cdot b L^\mu. \quad (6.25)$$

Commutation relations among the primitive power generators themselves are readily derived from Proposition 6.7:

$$X_v X_w = \hat{\pi}(v, w) X_w X_v, \quad Y_v Y_w = \hat{\pi}(v, w) Y_w Y_v, \quad \text{and} \quad X_v Y_w = Y_w X_v \quad (6.26)$$

where

$$\hat{\pi}(v, w) = \pi(\gamma(v), \gamma(w)) \quad \text{with} \quad \pi(\alpha, \beta) = \zeta^{\ell_\alpha \ell_\beta (\alpha|\beta)} \in \{+1, -1\}. \quad (6.27)$$

For the generators L_α as in (6.15) we also derive from $K^\eta E_w = q^{(\eta|\gamma(w))} E_w K^\eta$ that

$$L_\alpha X = \pi(\alpha, \gamma(w)) X L_\alpha \quad \text{and} \quad L_\alpha Y = \pi(\alpha, \gamma(w)) Y L_\alpha. \quad (6.28)$$

We conclude this section with criteria for strict commutativity of a primitive power generator X_v with $\alpha = \gamma(v)$ for non-empty $v \in \mathcal{W}^*$. As before, we assume that the root system is indecomposable of rank ≥ 2 and denote by κ the order of ζ .

Lemma 6.8. *The power generator X_v is central in U_ζ^+ if and only if (at least) one of the following conditions holds.*

- i) κ is odd.
- ii) $\kappa \equiv 2 \pmod{4}$, $d_\alpha = 2$ (that is, the Lie type is BCF and α is a long root).
- iii) The Lie type is B and α is a short root (that is, $d_\alpha = 1$).

In these cases X_v also commutes with all elements in U_ζ . The analogous statement holds for Y_v and L_α .

Note that the first condition includes all Lie types and was already observed in [DCK90]. See also the remarks following Proposition 6.12 below. The third condition applies to all κ and it is the only allowed situation if $\kappa \equiv 0 \pmod{4}$. The proof consists of checking for which α , Lie types, and orders κ we have that $\kappa(\alpha, \bar{\mu}) = 1$ for all $\bar{\mu}$. As a weaker condition, one may consider commutation of X_v with only the other primitive power generators.

Lemma 6.9. *X_v commutes with all X_w if and only if (at least) one of the following conditions holds.*

- i) κ is odd or $\kappa \equiv 0 \pmod{8}$.
- ii) $\kappa \equiv 2 \pmod{4}$ and the Lie type is B.
- iii) $\kappa \equiv 2 \pmod{4}$, the Lie type is CF, and α is a long root ($d_\alpha = 2$).
- iv) $\kappa \equiv 4 \pmod{8}$ and the Lie type is ACDEG.
- v) $\kappa \equiv 4 \pmod{8}$, the Lie type is F, and α is a short root.
- vi) κ is arbitrary, the Lie type is B, and α is a short root.

In these cases X_v also commutes with all Y_v and L_μ . The analogous statement holds for Y_v and L_α .

6.3. Skew-Central Subalgebras and Monomial Bases. The X_w and Y_w elements introduced in Section 6.2 generate almost polynomial subalgebras whose indeterminate variables commute up to signs. This section establishes the basic properties of these algebras and classifies when they are central, commutative but not central, or neither. We begin with definitions of the monomial elements in analogy to those introduced in Section 5.3.

Let $\mathbf{l} \in \mathbb{N}_0^\Phi$ be the length function $\mathbf{l}(\alpha) = \ell_\alpha$ and, for $\psi \in \mathbb{N}_0^A$ with $A \subseteq \Phi$, denote by $\mathbf{l} \cdot \psi$ the point-wise product $(\mathbf{l} \cdot \psi)(\alpha) = \ell_\alpha \psi(\alpha) \in \mathbb{N}_0$. For a non-empty, reduced word $w \in \mathcal{W}^*$, $\psi \in \mathcal{E}(s)$ with $s = r(w)$, and with conventions as for (5.20), set

$$X_{w\blacktriangleright}^\psi = E_{w\blacktriangleright}^{\mathbf{l} \cdot \psi} = X_{w[\beta_1]}^{\psi(\beta_1)} \dots X_{w[\beta_L]}^{\psi(\beta_L)} = E_{w[\beta_1]}^{\ell_{\beta_1} \psi(\beta_1)} \dots E_{w[\beta_L]}^{\ell_{\beta_L} \psi(\beta_L)}. \quad (6.29)$$

Analogously, we denote $X_{w\blacktriangleleft}^\psi = E_{w\blacktriangleleft}^{\mathbf{l} \cdot \psi}$, $Y_{w\blacktriangleright}^\psi = F_{w\blacktriangleright}^{\mathbf{l} \cdot \psi}$, and $Y_{w\blacktriangleleft}^\psi = F_{w\blacktriangleleft}^{\mathbf{l} \cdot \psi}$. Suppose next that $w, u \in \mathcal{W}^*$ are reduced words with associated Weyl group elements $s = r(w)$ and $t = r(u)$. Further, let $\psi, \phi \in \mathcal{E}(s)$ and $\chi \in \mathcal{E}(t)$ be respective exponent maps. Using the root sets and orderings explained in Section 2.3 we then define bihomomorphisms valued in $\{+1, -1\}$ as

$$\hat{\zeta}(\psi, \chi)_{w,u} = \prod_{\substack{\alpha \in \mathcal{N}(s) \\ \beta \in \mathcal{N}(t)}} \pi(\alpha, \beta)^{\psi(\alpha)\chi(\beta)} \quad \text{and} \quad \zeta(\psi, \phi)_w = \prod_{\substack{\alpha, \beta \in \mathcal{N}(s) \\ \alpha \prec_w \beta}} \pi(\alpha, \beta)^{\psi(\beta)\phi(\alpha)}, \quad (6.30)$$

for which we observe that $\zeta(\psi, \phi)_w \zeta(\phi, \psi)_w = \hat{\zeta}(\psi, \phi)_{w,w}$.

The commutation relations from (6.26) readily generalize to

$$X_w^\psi X_u^\chi = \hat{\zeta}(\psi, \chi)_{w,u} X_u^\chi X_w^\psi, \quad Y_w^\psi Y_u^\chi = \hat{\zeta}(\psi, \chi)_{w,u} Y_u^\chi Y_w^\psi, \quad \text{and} \quad Y_w^\psi X_u^\chi = X_u^\chi Y_w^\psi. \quad (6.31)$$

Here, X_w^ψ and Y_w^ψ denotes a monomial with any fixed direction of multiplication. That is, we may replace X_w^ψ in each equation by either $X_{w\blacktriangleright}^\psi$ or $X_{w\blacktriangleleft}^\psi$ and similarly for Y_w^ψ . Commutation relations for a fixed reduced word are given by

$$X_{w\blacktriangleright}^\psi X_{w\blacktriangleright}^\phi = \zeta(\psi, \phi)_w X_{w\blacktriangleright}^{\psi+\phi}, \quad X_{w\blacktriangleleft}^\psi X_{w\blacktriangleleft}^\phi = \zeta(\phi, \psi)_w X_{w\blacktriangleleft}^{\psi+\phi}, \quad \text{and} \quad X_{w\blacktriangleright}^\psi = \zeta(\psi, \psi)_w X_{w\blacktriangleleft}^\psi. \quad (6.32)$$

The same equations hold if all X 's are replaced by Y 's. We also record formulae for the involutions from Section 4.2 given by

$$\Omega(X_{w\blacktriangleright}^\psi) = Y_{w\blacktriangleleft}^\psi = \zeta(\psi, \psi)_w Y_{w\blacktriangleright}^\psi \quad \text{and} \quad \mathfrak{O}(X_{w\blacktriangleright}^\psi) = \Gamma_r(X_{w\blacktriangleright}^{\phi_{\psi}}). \quad (6.33)$$

The first is immediate from (5.24) and the second follows from Proposition 5.5 with elements r and ϕ as stated there. In either formula we use that \mathcal{W} preserves root lengths.

For a non-empty, reduced word $w \in \mathcal{W}^*$ we denote by $\mathcal{Z}_w^+ \subseteq U_\zeta^+$ and $\mathcal{Z}_w^- \subseteq U_\zeta^-$ the unital subalgebras generated by the sets $\{X_v : v \leq_R w\} \cup \{1\}$ and $\{Y_v : v \leq_R w\} \cup \{1\}$, respectively. Further, let \mathcal{Z}_s^0 be the unital subalgebra generated by 1 and all $L_\alpha^{\pm 1}$ with $\alpha \in \mathcal{N}(s)$. For a given $s \in \mathcal{W}$ denote also the ℓ -scaled exponent set

$$(\mathbf{l}, s) = \mathbf{l} \cdot \mathcal{E}(s) = \{\phi : \phi(\alpha) \in \ell_\alpha \mathbb{N}_0\}. \quad (6.34)$$

In the notation from (5.35) we thus have $\mathfrak{B}_{w\blacktriangleright}^{(\mathfrak{l},s),+} = \{X_{w\blacktriangleright}^\psi : \psi \in \mathcal{E}(s)\}$ and analogous identities for $\mathfrak{B}_{w\blacktriangleleft}^{(\mathfrak{l},s),+}$, $\mathfrak{B}_{w\blacktriangleright}^{(\mathfrak{l},s),-}$, and $\mathfrak{B}_{w\blacktriangleleft}^{(\mathfrak{l},s),-}$. The ground ring Λ in the following result is normally assumed to be $\Lambda_{\zeta, \mathbf{e}^*}$ for \mathcal{Z}_w^\pm or Λ_ζ for \mathcal{Z}_s^0 but may be extended to any $\Lambda_{\zeta,n}$ with $n \leq \ell$ or $\mathbb{Q}(\zeta)$. We are also reusing the notation for spans $\mathfrak{B}_{w\blacktriangleright}^{S,\pm}$ from (5.38) for these ground rings Λ .

Proposition 6.10. *Suppose $w \in \mathcal{W}^*$ and $s = r(w)$. Then both $\mathfrak{B}_{w\blacktriangleright}^{(\mathfrak{l},s),+}$ and $\mathfrak{B}_{w\blacktriangleleft}^{(\mathfrak{l},s),+}$ are Λ -bases for \mathcal{Z}_w^+ . In particular, $\mathcal{Z}_w^+ = \mathfrak{G}_{w\blacktriangleright}^{(\mathfrak{l},s),+} = \mathfrak{G}_{w\blacktriangleleft}^{(\mathfrak{l},s),+}$.*

The subalgebra \mathcal{Z}_w^+ is isomorphic to the abstract Λ -algebra with 1, freely generated by elements $X_{w\blacktriangleright}^\psi$ and subject to the first set of relations in (6.32). The analogous statements for \mathcal{Z}_w^- apply.

The subalgebra \mathcal{Z}_s^0 is generated by $L_i^{\pm 1}$ with $i \in J(s)$ as in Section 2.3. It is thus isomorphic to the group algebra $\Lambda [\{L_i^\pm : i \in J(s)\}] \cong \Lambda [\Xi(s)]$.

Proof. The relations in (6.32) imply that for any $v \in \mathcal{W}^*$ with $v \leq_R w$ we have $X_v \cdot \mathfrak{G}_{w\blacktriangleright}^{(\mathfrak{l},s),+} \subseteq \mathfrak{G}_{w\blacktriangleright}^{(\mathfrak{l},s),+}$, writing $X_v = X_{w\blacktriangleright}^{\chi_\alpha}$ with χ_α the indicator function for $\alpha = \gamma(v)$. Thus $\mathcal{Z}_w^+ = \mathcal{Z}_w^+ \cdot 1 \subseteq \mathfrak{G}_{w\blacktriangleright}^{(\mathfrak{l},s),+}$. From the definition in (6.29), it is further clear that 1 is a cyclic element of $\mathfrak{G}_{w\blacktriangleright}^{(\mathfrak{l},s),+}$ as a \mathcal{Z}_w^+ -module and hence $\mathcal{Z}_w^+ = \mathfrak{G}_{w\blacktriangleright}^{(\mathfrak{l},s),+}$ as desired. Linear independence established in Section 5 then completes the proof of the first claim. Independence under reversal of the direction of multiplication is immediate from (6.32).

It is clear that the set of generators $X_{w\blacktriangleright}^\psi$ of an abstractly defined algebra with relations as in (6.32) is also a basis. Thus, the canonical map from this algebra onto \mathcal{Z}_w^+ is an isomorphism of Λ -modules and, hence, also an algebra isomorphism.

By its definition, \mathcal{Z}_s^0 is the Λ -span of all K^ν with ν in the root sublattice generated by all $\ell\alpha = \ell_\alpha \cdot \alpha$ with $\alpha \in \mathcal{N}(s)$. Following Lemma 6.4, we consider the root system Φ_ℓ with the same Weyl group action. Since, clearly, $s(\alpha) \in \Phi^-$ iff $s(\ell\alpha) \in \Phi^-_\ell$ the inversion set $\mathcal{N}(s)_\ell$ with respect to Φ_ℓ is given precisely by $\{\ell\alpha : \alpha \in \mathcal{N}(s)\}$. By Lemma 2.10 the root sublattice generated by this set is spanned by the simple roots of Φ_ℓ with indices in $J(s)$. By Lemma 6.4 these are $\ell\alpha_i = \ell_i\alpha_i$, yielding the $L_i = K^{\ell\alpha_i}$ as free generators. \square

The next observations are consequences of (6.33), particularly, in the case when the word in the \mathfrak{U} formula is maximal so that $r = 1$. The stated equivariance under the antipode follows from (4.16).

Corollary 6.11. *Suppose $w \in \mathcal{W}^*$ is a reduced word and $z \in \mathcal{W}_m^*$ one of maximal length. Then $\Omega(\mathcal{Z}_s^0) = \mathfrak{U}(\mathcal{Z}_s^0) = S(\mathcal{Z}_s^0) = \mathcal{Z}_s^0$ as well as*

$$\Omega(\mathcal{Z}_w^\pm) = \mathcal{Z}_w^\mp, \quad \mathfrak{U}(\mathcal{Z}_z^\pm) = \mathcal{Z}_{z^\dagger}^\pm, \quad \text{and} \quad S(\mathcal{Z}_z^0) = \mathcal{Z}_{z^\dagger}^0. \quad (6.35)$$

Proposition 6.7 further implies the following relations between $X_{w\bullet}^\psi$ and $E_{w\bullet}^\rho$ elements. Here w is a reduced word and $\psi, \rho \in \mathcal{E}(s)$ with $s = \tau(w)$, and κ from (6.10) is lifted to $\Phi \times \mathbb{Z}^\Delta$:

$$X_{w\bullet}^\psi \cdot E_{w\bullet}^\rho = \vec{\kappa}(\psi, \rho)_w \cdot E_{w\bullet}^{\mathbf{l} \cdot \psi + \rho} \quad \text{and} \quad E_{w\bullet}^\rho \cdot X_{w\bullet}^\psi = \overleftarrow{\kappa}(\psi, \rho)_w \cdot E_{w\bullet}^{\mathbf{l} \cdot \psi + \rho}$$

where $\vec{\kappa}(\psi, \rho)_w = \prod_{\substack{\alpha, \beta \in \mathcal{N}(s) \\ \alpha \prec_w \beta}} K(\alpha, \beta)^{\psi(\beta)\rho(\alpha)}$ and $\overleftarrow{\kappa}(\psi, \rho)_w = \prod_{\substack{\alpha, \beta \in \mathcal{N}(s) \\ \alpha \prec_w \beta}} K(\alpha, \beta)^{\psi(\alpha)\rho(\beta)}$. (6.36)

Using division with remainder pointwise for each α , any $\phi \in \mathcal{E}(s)$ can be uniquely written as $\phi = \mathbf{l} \cdot \psi + \rho$ with $\rho \in [0, \mathbf{l}]_s$, where

$$[0, \mathbf{l}]_s = \{\psi \in \mathcal{E}(s) : \psi(\alpha) < \ell_\alpha, \forall \alpha \in \mathcal{N}(s)\} \quad \text{and} \quad [0, \mathbf{l}]_* = [0, \mathbf{l}]_s. \quad (6.37)$$

It is now immediate that the multiplication map below is an isomorphism of Λ -modules, mapping pairs of PBW basis elements to another PBW basis element up to a sign.

$$m_w^+ : \mathcal{Z}_w^+ \otimes_\Lambda \mathfrak{S}_{w\bullet}^{[0, \mathbf{l}]_s, +} \longrightarrow \mathfrak{S}_{w\bullet}^+ \quad (6.38)$$

Viewing $\mathfrak{S}_{w\bullet}^+$ as a \mathcal{Z}_w^+ -module with its left regular action given by the signed shifts in (6.36), this implies that $\mathfrak{S}_{w\bullet}^+$ splits into directed summands, each of which has $E_{w\bullet}^\rho$ for some $\rho \in [0, \mathbf{l}]_s$ as cyclic element.

Note also that $\mathfrak{S}_{w\bullet}^{[0, \mathbf{l}]_s, +}$ generally depends explicitly on the choice of the reduced word w rather than just the respective Weyl element $s = \tau(w)$. For example, for Lie type A_2 and $\mathbf{k} = 4$ both $w = w_1 w_2 w_1$ and $\bar{w} = w_2 w_1 w_2$ represent the maximal element in \mathcal{W} . Given that $\ell = 2$, the components of $\mathfrak{S}_{w\bullet}^{[0, \mathbf{l}]_s, +}$ and $\mathfrak{S}_{\bar{w}\bullet}^{[0, \mathbf{l}]_s, +}$ with maximal w -grading of $2\alpha_1 + 2\alpha_2$ are $\Lambda E_{w\bullet}^1$ and $\Lambda E_{\bar{w}\bullet}^1$ respectively, where $\mathbf{l}(\alpha) = 1$ for all $\alpha \in \Phi^+$. With $E_{w\bullet}^1 = E_1 E_{(12)} E_2$ and $E_{\bar{w}\bullet}^1 = E_2 E_{(21)} E_1 = E_{w\bullet}^1 + E_{(12)}^2 - 2\zeta^{-1} E_1^2 E_2^2$, these are clearly different spaces. We will, however, prove below that \mathcal{Z}_w^+ only depends on $s = \tau(w)$ for almost all Lie types.

Recall from Theorem 5.18 and Corollary 6.3 that $\mathfrak{S}_{w\bullet}^+ = U_\zeta^+$ for a reduced word $w \in \mathcal{W}_m^*$ of maximal length. Variations of (6.36) and (6.38) are readily derived with reversed directions or replacing E 's by F 's and X 's by Y 's.

Similar to the full quantum groups, we introduce combined subalgebras such as

$$\mathcal{Z}_w^{\geq 0} = \mathcal{Z}_w^+ \cdot \mathcal{Z}_s^0, \quad \mathcal{Z}_w^{\leq 0} = \mathcal{Z}_w^- \cdot \mathcal{Z}_s^0, \quad \text{and} \quad \mathcal{Z}_w = \mathcal{Z}_w^+ \cdot \mathcal{Z}_w^- \cdot \mathcal{Z}_s^0, \quad (6.39)$$

where $s = \tau(w)$. Analogous statements about PBW bases and identifications with signed polynomial algebras hold for each of these subalgebras. Since all elements in \mathcal{Z}_w^+ commute with all elements in \mathcal{Z}_w^- we do not require the denominators $(\zeta - \zeta^{-1})^{-1}$ so that each \mathcal{Z}_w is well-defined over $\Lambda_{\zeta, \mathbf{e}^*}$. We next record the precise criteria for these subalgebras to be strictly central or strictly commutative.

Proposition 6.12. *Suppose $w \in \mathcal{W}^*$ is non-empty, $s = \tau(w)$, and $\square \in \{ -, +, \geq 0\}$. The algebra \mathcal{Z}_w^+ is central in U_q^\square if and only if (at least) one of the following conditions holds:*

- i) \mathbb{K} is odd (any Lie type).
- ii) $\mathbb{K} \not\equiv 0 \pmod{4}$ and the Lie type is B.
- iii) $\mathbb{K} \equiv 2 \pmod{4}$, the Lie type is CF, and $s \in \mathcal{W}_S$.
- iv) Lie type is B and $s \in \mathcal{W}_L$ (any \mathbb{K}).

Thus, if $w \in \mathcal{W}_m^*$, then \mathcal{Z}_w^+ is central if and only if (at least) one of i) and ii) hold. Analogous statements hold for \mathcal{Z}_w^\square .

The subgroups \mathcal{W}_L and \mathcal{W}_S are as defined in (2.17). The one referred to in items iii) and iv) are of order 2 for the B, C types and of order 6 for F_4 by Lemma 2.5. The assertions in Proposition 6.12 are readily derived from Lemma 6.8. We note that the last statement for maximal words is commensurable with Corollary 3.1 of [DCK90]. The elements considered there in part (b) correspond to our X_w if $\alpha = \gamma(w)$ is a short root but to our X_w^2 if α is a long root in case \mathbb{K} is a multiple of 4. Requiring centrality of X_w rather than X_w^2 then excludes this case.

In order to formulate criteria for commutativity of the subalgebras, we say that an element $s \in \mathcal{W}$ is *mod-2 discrete* if

$$(\check{\alpha}|\beta) \equiv 0 \pmod{2} \quad \forall \alpha, \beta \in \mathcal{N}(s) \text{ with } d_\alpha \geq d_\beta. \quad (6.40)$$

For example, for Lie type A_5 and $s = s_1 s_3 s_6$ we have $\mathcal{N}(s) = \{\alpha_1, \alpha_3, \alpha_6\}$, which are all orthogonal so that s is mod-2 discrete. For non-simply laced types we will also call an element $s \in \mathcal{W}$ *mod-2 long discrete* if condition (6.40) is required only for pairs $\alpha, \beta \in \mathcal{N}(s)_L$. The notion of *mod-2 short discrete* is analogous.

Proposition 6.13. *Suppose $w \in \mathcal{W}^*$ is non-empty and $s = r(w)$. The subalgebra \mathcal{Z}_w^+ is commutative if and only if (at least) one of the following conditions holds:*

- i) s is mod-2 discrete (any Lie type and \mathbb{K}).
- ii) \mathbb{K} is odd or $\mathbb{K} \equiv 0 \pmod{8}$ (any Lie type).
- iii) $\mathbb{K} \equiv 0 \pmod{4}$ and the Lie type is ADEG.
- iv) $\mathbb{K} \equiv 2 \pmod{4}$, the Lie type is BCF, and s is mod-2 short discrete.
- v) $\mathbb{K} \equiv 4 \pmod{8}$, the Lie type is BCF and s is mod-2 long discrete.

The identical statement holds for $\mathcal{Z}_w^{\geq 0}$.

The extension from \mathcal{Z}_w^+ to $\mathcal{Z}_w^{\geq 0}$ holds since, by (6.28), the sign occurring in the commutation of X_α with X_β (with respect to some order) is the same as the one in the commutation of X_α with L_β . Note also that for type A_1 case i) holds for any s .

For the maximal case, observe that the longest element $s_\bullet \in \mathcal{W}$ is never mod-2 discrete for any Lie type of rank ≥ 2 . In the non-simply laced case we have that $s_\bullet \in \mathcal{W}$ is mod-2 short discrete if and only if the Lie type is B. It is mod-2 long discrete if and only if the Lie type is C.

This yields the following characterization of maximal commutative subalgebras, which can also be derived from Lemma 6.9.

Corollary 6.14. *Let $z \in \mathcal{W}_m^*$ be a reduced word of maximal length. Then \mathcal{Z}_z^+ is commutative if and only if (at least) one of the following applies:*

- i) $\mathbb{K} \equiv 0, 1, 3, 5, \text{ or } 7 \pmod{8}$ (any Lie type).
- ii) $\mathbb{K} \equiv 2 \text{ or } 6 \pmod{8}$ and the Lie type is B .
- iii) $\mathbb{K} \equiv 4 \pmod{8}$ and the Lie type is ACDEG.
- iv) The Lie type is A_1 (any \mathbb{K}).

The identical statement holds for $\mathcal{Z}_z^{\geq 0}$.

Remark 6.15. Corollary 6.14 implies that \mathcal{Z}_z^+ in type $B_2 = C_2$ is commutative for any \mathbb{K} and any $z \in \mathcal{W}_m^*$. However, it is central only if \mathbb{K} is not a multiple of 4. \diamond

It is immediate from Proposition 6.10 that, if \mathcal{Z}_w^+ is strictly commutative, it is isomorphic to a free polynomial algebra, assigning each X_v with $v \leq_R w$ to an indeterminate x_α , where $\alpha = \gamma(v) \in \mathcal{N}(s)$ and $s = \tau(w)$. That is,

$$\mathcal{Z}_w^+ \cong \Lambda \left[\{x_\alpha : \alpha \in \mathcal{N}(s)\} \right]. \quad (6.41)$$

Comparing the criteria in Propositions 6.12 and Corollary 6.14, we note that there are various situations in which the subalgebras are commutative but not central. Given a maximal $w \in \mathcal{W}_m^*$, \mathcal{Z}_w^+ has this property if $\mathbb{K} \equiv 0 \pmod{8}$ for any Lie type as well as if $\mathbb{K} \equiv 4 \pmod{8}$ and the Lie type is neither B nor F . So, for example, in the simply laced cases (ADE) and $\mathbb{K} \equiv 0 \pmod{4}$ we have that \mathcal{Z}_w^+ is commutative as in (6.41) but for any $b \in U_q$ with $\mu = \phi^+(b)$ and $\alpha = \gamma(v)$ we have

$$X_v b = (-1)^{(\alpha|\mu)} b X_v. \quad (6.42)$$

6.4. \mathfrak{U} -Invariance and Word Independence of \mathcal{Z}_w . The definition of the algebras \mathcal{Z}_w^\square in Section 6.3 *a priori* depends on a choice of a word $w \in \mathcal{W}^*$. In this section, we show for Lie types different from G_2 that \mathcal{Z}_w^\square depends only on the respective Weyl element $s = \tau(w)$, employing the techniques from Section 5.7.

We begin with the derivation of formulae for the action of \mathfrak{U} on generators in U_q , starting with the generic case and coefficients in $\mathbb{Q}(q)$. We continue here the conventions from Section 5.4, with i the index of a short root ($d_i = 1$) and j the index of a (possibly) long root ($d_j = e$). Observe that specializing the left side of (5.50) to $z = -1$ and $u_j(\vec{n}) = 0$ yields the expression for $E_{(ji)}^{(N)}$ in (5.52) multiplied by $(-1)^N$. Similarly, for $z = -1$ and $u_i(\vec{n}) = 0$, the left side of (5.51) yields the expression for $E_{b_{m-1}}^{(N)}$ in (5.52) multiplied by $(-1)^{eN}$. The products over the respective $(1 - q_\alpha^{-2t}) = q_\alpha^{-t}(q_\alpha - q_\alpha^{-1})[t]_\alpha$ terms are then expressed as factorials, resulting in the following

formulae.

$$E_{(ji)}^{(N)} = (-1)^N \sum_{\substack{\phi \in S_{\mathbf{e}}(\vec{\sigma}) \\ k_m = \phi(\alpha_j)}} q^{f_{\mathbf{e}}(\phi)} (q_j - q_j^{-1})^{k_m} q_j^{-\binom{k_m+1}{2}} [k_m]!_j E_{\zeta_{|ij|}}^{(\phi)} \quad \vec{\sigma} = N s_j(\alpha_i) = N \alpha_i + N \alpha_j \quad (6.43)$$

$$E_{b_{m-1}}^{(N)} = (-1)^{\mathbf{e}N} \sum_{\substack{\phi \in S_{\mathbf{e}}(\vec{\tau}) \\ k_1 = \phi(\alpha_i)}} q^{f_{\mathbf{e}}(\phi)} (q - q^{-1})^{k_1} q^{-\binom{k_1+1}{2}} [k_1]!_j E_{\zeta_{|ij|}}^{(\phi)} \quad \vec{\tau} = N s_i(\alpha_j) = \mathbf{e}N \alpha_i + N \alpha_j \quad (6.44)$$

Multiplying these expressions by $[N]!$ and $[N]!_j$, respectively, we obtain expressions for ordinary powers in terms of elements $\langle \phi, i \rangle_{\mathbf{e}}$ and $\langle \phi, j \rangle_{\mathbf{e}}$, obtained by combining q -factorials.

$$E_{(ji)}^N = (-1)^N \sum_{\substack{\phi \in S_{\mathbf{e}}(\vec{\sigma}) \\ k_m = \phi(\alpha_j)}} q^{f_{\mathbf{e}}(\phi)} (q - q^{-1})^{k_m} q_j^{-\binom{k_m+1}{2}} \langle \phi, j \rangle_{\mathbf{e}} E_{\zeta_{|ij|}}^{\phi} \quad (6.45)$$

$$E_{b_{m-1}}^N = (-1)^{\mathbf{e}N} \sum_{\substack{\phi \in S_{\mathbf{e}}(\vec{\tau}) \\ k_1 = \phi(\alpha_i)}} q^{f_{\mathbf{e}}(\phi)} (q - q^{-1})^{k_1} q^{-\binom{k_1+1}{2}} \langle \phi, i \rangle_{\mathbf{e}} E_{\zeta_{|ij|}}^{\phi} \quad (6.46)$$

In the case of (6.45) we have $u_j(\vec{\sigma}) = 0$, which implies $k_3 = k_1$ for $\mathbf{e} = 1$, $k_4 = k_1 + k_2$ for $\mathbf{e} = 2$, and $k_6 = k_1 + 2k_2 + k_3 + k_4$ for $\mathbf{e} = 3$. The ratios of factorials can readily be worked out as follows, using the elements in (3.6) and identities in (3.5).

$$\begin{aligned} \langle \phi, j \rangle_1 &= \begin{bmatrix} k_1 + k_2 \\ k_1 \end{bmatrix} & \langle \phi, j \rangle_2 &= [2]^{k_2+k_4} [k_2 : 2]!_j \cdot \begin{bmatrix} k_1 + 2k_2 + k_3 \\ k_1, 2k_2, k_3 \end{bmatrix} \\ \langle \phi, j \rangle_3 &= [3]^{k_2+k_4+k_6} \cdot [k_2 : 3]!_j \cdot [k_4 : 3]!_j \cdot [k_3]!_j \cdot \begin{bmatrix} 2k_3 \\ k_3 \end{bmatrix} \cdot \begin{bmatrix} k_1 + 3k_2 + 2k_3 + 3k_4 + k_5 \\ k_1, 3k_2, 2k_3, 3k_4, k_5 \end{bmatrix} \end{aligned} \quad (6.47)$$

The multinomial coefficients are as defined in (3.7). Similarly, for (6.45), the condition $u_i(\vec{\tau}) = 0$ implies $k_1 = k_3$ for $\mathbf{e} = 1$, $k_1 = k_3 + 2k_4$ for $\mathbf{e} = 2$, and $k_1 = k_3 + 3k_4 + 2k_5 + 3k_6$ for $\mathbf{e} = 3$. As before, with notation from (3.6) and (3.5), we find analogous formulae.

$$\begin{aligned} \langle \phi, i \rangle_1 &= \begin{bmatrix} k_2 + k_3 \\ k_3 \end{bmatrix} & \langle \phi, i \rangle_2 &= [2]^{-k_3} \cdot \{k_3 : 2\} \cdot \begin{bmatrix} k_2 + k_3 + k_4 \\ k_2, k_3, k_4 \end{bmatrix}_j \\ \langle \phi, i \rangle_3 &= [3]^{-(k_3+k_5)} \cdot \{k_3 : 3\} \cdot \{k_5 : 3\} \cdot [k_4]!_j \cdot \begin{bmatrix} 2k_4 \\ k_4 \end{bmatrix}_j \cdot \begin{bmatrix} k_2 + k_3 + 2k_4 + k_5 + k_6 \\ k_2, k_3, 2k_4, k_5, k_6 \end{bmatrix}_j \end{aligned} \quad (6.48)$$

Clearly, all coefficients in (6.47) and (6.48) lie in Λ_{q,\mathbf{e}^*} as in (3.8). It is easy to check that in a corresponding relation for rescaled generators $\dot{E}_{\alpha} = [d_{\alpha}]E_{\alpha}$, already mentioned at the end of Section 5.6, the factors of the form $[\mathbf{e}]^{-k}$ drop out so that the coefficients would be entirely in Λ_q .

In either case, we may specialize the identity to a root of unity ζ , assuming that $\ell > \mathbf{e}$ where ℓ is the order of ζ^2 . This allows us, in particular, to derive expansions for $X_{(ji)} = E_{(ji)}^{\ell}$ and $X_{b_{m-1}} = E_{b_{m-1}}^{\ell_j}$ in terms of the $E_{\zeta_{|ij|}}^{(\phi)}$ basis elements. The number of non-zero terms will depend

in either case on whether $\ell = \ell_j$ or $\ell > \ell_j$, that is, whether $\gcd(\ell, \mathbf{e}) = 1$ or not. To keep subsequent formulae more compact, it is convenient to introduce

$$\mathbf{d} = \gcd(\ell, \mathbf{e}) \in \{1, \mathbf{e}\} \quad (6.49)$$

for given ℓ and \mathbf{e} . Using the Kronecker notation, we can then express the evaluation of the factorial ratio from (3.6) at $q = \zeta$ as

$$\{\ell_j : \mathbf{e}\} = \delta_{\mathbf{d}, 1} \cdot \mathbf{e} \cdot \zeta^{-(\mathbf{e}-1)(\ell+1)/2}, \quad (6.50)$$

implying that $\{\ell_j : \mathbf{e}\} = 0$ if $\ell_j < \ell$. The other factorial ratio from (3.6) can be evaluated at $q = \zeta$ using (6.4).

Lemma 6.16. *Suppose $\ell = \mathbf{e} \cdot \ell_j$. Then*

$$[\ell_j : \mathbf{e}]^{\mathbf{d}} = \mathbf{e} \cdot \zeta^{-\binom{\mathbf{e}}{2}\ell_j^2} \cdot (\zeta^{-1} - \zeta)^{-(\mathbf{e}-1)\ell_j}.$$

Another quantity that frequently appears in the computations below is the expression

$$\nabla_j := (\zeta_j - \zeta_j^{-1})^{\ell_j} = [\mathbf{e}]^{\ell_j} (\zeta - \zeta^{-1})^{\ell_j}. \quad (6.51)$$

This is the singularization factor of X_j in the sense of (5.10), as we introduce the analogous notation $\check{X}_j = \check{E}_j^{\ell_j} = (-1)^{\ell_j} \nabla_j E_j^{\ell_j}$. Notice also that $\nabla_j^{\mathbf{d}} = \nabla_i \cdot [\mathbf{e}]^{\ell}$.

The next two propositions consider the specializations of (6.45) and (6.46) to a root of unity ζ with $N = \ell$ and $N = \ell_j$, respectively. These yield explicit expressions for $X_{(ji)} = E_{(ji)}^{\ell}$ and $X_{b_{m-1}} = E_{b_{m-1}}^{\ell_j}$.

Proposition 6.17. *We have $X_{(ji)} \in \mathcal{Z}_{z_{[ij]}}^+$ as a ring over $\mathbb{Z}[\zeta]$, given by*

$$(-1)^\ell X_{(ji)} = \zeta_j^\ell X_{a_{m-1}} + \zeta_j^{\binom{\ell}{2}} \nabla_j^{\mathbf{d}} X_j^{\mathbf{d}} X_i + (1 - \delta_{\mathbf{d}, 1}) \cdot \mathbf{e} \cdot (\zeta - \zeta^{-1})^{-\ell} \cdot T_1$$

where $T_1 = \begin{cases} (-\zeta)^{\ell_j} \nabla_j^2 X_j X_{(ij)} & \mathbf{e} = 2 \\ (-\zeta_j)^{-\ell_j^2} \nabla_j^3 X_j^2 X_{(ij)} + \zeta_j^{-\binom{\ell_j+1}{2}} \nabla_j^2 X_j X_{(iji)} & \mathbf{e} = 3. \end{cases} \quad (6.52)$

The computation starts with the observation that, by Corollary 6.2, we only need to consider cases in which one of the terms in the large multinomial coefficient for each $(\phi, j) \mathbb{D}_{\mathbf{e}}$ in (6.47) is equal to ℓ and all the other k_s with $1 \leq s \leq m-1$ are zero. The value of k_m is then determined via $u_j(\phi) = 0$ and (5.44) from the non-zero k_s .

The first term in (6.52) results from the case $k_{m-1} = \ell$ and all other $k_s = 0$. The second term accounts for the case $k_1 = k_m = \ell$ and all other $k_s = 0$. Note also that $2k_3 = \ell$ for $\mathbf{e} = 3$ will still lead to $(\phi, j) \mathbb{D}_{\mathbf{e}} = 0$ due to the extra binomial term.

All other cases require that \mathbf{e} divides ℓ , meaning $\mathbf{d} = \mathbf{e} > 1$ and $\ell = \mathbf{e}\ell_j$. For $\mathbf{e} = 2$ the only remaining situation is then $k_2 = k_4 = \ell_j$ and $k_1 = k_3 = 0$. For $\mathbf{e} = 3$ we have two cases, the first

of which is $k_2 = \ell_j$, $k_6 = 2\ell_j$ and $k_1 = k_3 = k_4 = k_5 = 0$. The second is given by $k_4 = \ell_j$, $k_6 = \ell_j$ and $k_1 = k_2 = k_3 = k_5 = 0$.

The coefficients for each of these values of k_s are then worked out using Lemma 6.16. Next, we provide the expressions for $X_{b_{m-1}}$, where we use the additional notation

$$\mathbf{n} = \frac{\mathbf{e}}{\mathbf{d}} \quad \text{so that} \quad \mathbf{n} \cdot \ell = \mathbf{e} \cdot \ell_j. \quad (6.53)$$

Note that the third term in (6.54) is non-zero only if $\mathbf{e} > 1$ and $\gcd(\ell, \mathbf{e}) = 1$. As remarked in Section 6.1 this condition implies that $[\mathbf{e}]$ is a unit, meaning $[\mathbf{e}]^{-1} \in \mathbb{Z}[\zeta]$. Analogous to (6.51) we also write $\nabla_i = (\zeta - \zeta^{-1})^\ell$.

Proposition 6.18. *We have $X_{b_{m-1}} \in \mathcal{Z}_{z_{ij}}^+$ as a ring over $\mathbb{Z}[\zeta]$, given by*

$$(-1)^{\mathbf{e}\ell_j} X_{b_{m-1}} = \zeta_j^{\ell_j} X_{(ij)} + \zeta^{\binom{n-\ell}{2}} \nabla_i^{\mathbf{n}} X_j X_i^{\mathbf{n}} + \delta_{\mathbf{d},1} \cdot (1 - \delta_{\mathbf{e},1}) \cdot \mathbf{e} \cdot [\mathbf{e}]^{-\ell} \cdot T_2$$

where $T_2 = \begin{cases} \zeta^\ell \nabla_i X_{(iji)} X_i & \mathbf{e} = 2 \\ \zeta^{-\binom{\ell}{2}} \nabla_i X_{(iji)} X_i + \zeta^{\ell(\ell+1)} \nabla_i^2 X_{(ijiji)} X_i^2 & \mathbf{e} = 3. \end{cases} \quad (6.54)$

As for Proposition 6.17, the only contributing terms are those for which exactly one of the integers in the multinomial expressions of (6.48) is ℓ_j and all others are zero, once we set $N = \ell_j$ and $q = \zeta$ in (6.46). The first term in (6.54) accounts for the case when $k_2 = \ell_j$, implying by $u_i = 0$ that all other $k_s = 0$. The second term corresponds to the case $k_m = \ell_j$, which entails $k_1 = \mathbf{e}\ell_j = \mathbf{n}\ell$ and all remaining $k_s = 0$.

The last terms for $\mathbf{e} > 1$ all involve coefficients $\{\ell_j : \mathbf{e}\}$, which, by (6.50), are non-zero only if $\mathbf{d} = 1$ and hence $\ell = \ell_j$. The extra term for $\mathbf{e} = 2$ corresponds to $k_3 = k_1 = \ell_j = \ell$ and all other indices zero. As before, for $\mathbf{e} = 3$ we only need to consider the cases $k_3 = \ell_j = \ell$ and $k_5 = \ell_j = \ell$, which imply $k_1 = \ell$ and $k_1 = 2\ell$, respectively.

Comparing Propositions 6.17 and 6.18, we note that additional terms and exponents occur in the non-simply laced cases in exactly the opposite situations with respect to whether ℓ and \mathbf{e} are coprime or not. In addition, a term by term matching, up to coefficients, can also be achieved by switching an X_{a_j} by an $X_{a_{m+1-j}}$. This can be thought of replacing the underlying root system by a corresponding coroot system as in (6.17), a phenomenon that will resurface in Section 8.1.

We discuss a normalization of the generators X_α in Appendix D for which the above formulae are integral. The main application of (6.52) and (6.54) word independence of the skew-central subalgebras.

Theorem 6.19. *Suppose Φ is not of Lie type G_2 . Assume $w, u \in \mathcal{W}^*$ are reduced words for the same Weyl element $s = r(w) = r(u)$. Then $\mathcal{Z}_w^+ = \mathcal{Z}_u^+$ as w -graded subalgebras in U_ζ^+ over $\Lambda_{\zeta, \mathbf{e}^*}$.*

Proof. By Proposition 6.10 \mathcal{Z}_w^+ is as a $\Lambda_{\zeta, \mathbf{e}^*}$ -module given by $\mathfrak{S}_{\mathbf{w}}^{(\mathbf{l}, s), +}$. We note that (\mathbf{l}, s) , as defined in (6.34), is of the form prescribed in (5.65) by choosing $V_d = \ell_d \cdot \mathbb{N}_0$ with $\ell_d = \ell / \gcd(d, \ell)$.

Following Corollary 5.15 it thus suffices to show that \mathfrak{U} maps $\mathcal{Z}_{z_{ijj}}^+ = \mathfrak{S}_{\zeta_{ijj}}^{(\mathfrak{l}, s_{ijj}), +}$ to itself for any pair (i, j) that occurs for relator pairs. Now, $\mathfrak{S}_{\zeta_{ijj}}^{(\mathfrak{l}, s), +}$ is spanned by monomials in the skew-commuting generators X_{a_s} . Note also that by (5.32) and (5.16) and with words as in (2.32) we also have

$$\mathfrak{U}(X_{a_s}) = X_{b_{m+1-s}}. \quad (6.55)$$

Thus \mathfrak{U} -invariance of the space $\mathfrak{S}_{\zeta_{ijj}}^{(\mathfrak{l}, s), +}$ is given by showing that it contains all X_{b_k} . This is obvious for $X_{b_1} = X_j$ and $X_{b_m} = X_i$, implied, for example, by (5.18). For $X_{b_2} = X_{(ji)}$ this is verified in Proposition 6.17 and for $X_{b_{m-1}}$ in Proposition 6.18. Thus, if $m \in \{3, 4\}$ all cases are covered.

By construction any \mathcal{Z}_w^+ inherits the w -grading from U_ζ since it admits a w -graded basis. The graded components of \mathcal{Z}_w^+ and \mathcal{Z}_u^+ thus also have to coincide. \square

Missing from the proof for the G_2 Lie type are computations that show also X_{b_3} and X_{b_4} can be written as a polynomial expression in the $\{X_{a_i}\}$ only, meaning they are in $\mathfrak{S}_{\zeta_{ijj}}^{(\mathfrak{l}, s_{ijj}), +}$. We leave this as a question, which appears to be open, at least for even \mathfrak{k} .

Conjecture 6.20. *Theorem 6.19 extends also to the G_2 case for all $\mathfrak{k} > 6$.*

For Lie types different from G_2 , we may thus unambiguously write \mathcal{Z}_s^+ for an element in $s \in \mathcal{W}$. In the G_2 case, \mathcal{Z}_s^+ is, of course, also well-defined for $l(s) \leq 5$, since there is only one reduced expression for such s . For the longest element s_\bullet , there are exactly two reduced expressions. So, the only missing equality between subalgebras addressed by this conjecture is the one between $\mathcal{Z}_{z_{ijj}}^+$ and $\mathcal{Z}_{z_{ijj}}^+$.

Assuming the other Lie types, we denote the maximal subalgebras associated to the unique longest element $s_\bullet \in \mathcal{W}$ as

$$\mathcal{Z}_\bullet^\pm = \mathcal{Z}_{s_\bullet}^\pm, \quad \mathcal{Z}_\bullet^{\geq 0} = \mathcal{Z}_\bullet^+ \cdot \mathcal{Z}_\bullet^0, \quad \mathcal{Z}_\bullet^{\leq 0} = \mathcal{Z}_\bullet^- \cdot \mathcal{Z}_\bullet^0, \quad \text{and} \quad \mathcal{Z}_\bullet = \mathcal{Z}_\bullet^+ \cdot \mathcal{Z}_\bullet^0 \cdot \mathcal{Z}_\bullet^-, \quad (6.56)$$

where \mathcal{Z}_\bullet^0 is the full subalgebra generated by the L_i . Also immediate from the definition is that

$$\mathcal{Z}_t^+ \subseteq \mathcal{Z}_s^+ \quad \forall s, t \in \mathcal{W} \text{ with } t \leq_R s. \quad (6.57)$$

Analogous statements hold for the \mathcal{Z}_s^\square algebras with $\square \in \{+, -, \geq 0, \leq 0\}$. Note also that Corollary 6.35 and Theorem 6.19 imply for these Lie types

$$\Omega(\mathcal{Z}_s^\pm) = \mathcal{Z}_s^\mp, \quad \mathfrak{U}(\mathcal{Z}_\bullet^\pm) = \mathcal{Z}_\bullet^\pm, \quad \text{and} \quad S(\mathcal{Z}_\bullet^{\geq 0}) = \mathcal{Z}_\bullet^{\geq 0}. \quad (6.58)$$

Combining this with \mathfrak{U} -correspondence with Lusztig's conventions from Proposition 5.5 we make the following observations.

Corollary 6.21. *The maximal algebras in (6.56) remain unchanged if the Γ_i are replaced by T_i in the definition of the generators E_w and X_w .*

We conclude this section with a proof that the full maximal algebra is invariant under the actions of Lusztig's T_i automorphisms. It uses the explicit expression of the Garside element Γ_{s_\bullet} from Section 5.2.

Theorem 6.22. *Assuming Lie types other than G_2 and any $\mathbb{k} > 2e$ (or $\mathbb{k} = 3$ for $e = 2$), the full algebra \mathcal{Z}_\bullet over Λ_{ζ, e^*}^* is invariant under the Artin group action generated by the Γ_i .*

Proof. Invariance of \mathcal{Z}_\bullet^0 under Γ_i follows immediately from (6.19) and Proposition 6.10. It thus suffices to show $\Gamma_i(\mathcal{Z}_\bullet^+) \subset \mathcal{Z}_\bullet$ for each fixed $i \in \{1, \dots, n\}$, which implies $\Gamma_i(\mathcal{Z}_\bullet^-) \subset \mathcal{Z}_\bullet$ via (6.58) and commutation of Ω with Γ_i .

Let $u \in \mathcal{W}^*$ be a reduced word for $q_{\eta(i)} = s_i \cdot s_\bullet = s_\bullet \cdot s_{\eta(i)}$ as in (2.22) so that $z = w_i \cdot u$ and $w = u \cdot w_{\eta(i)}$ are maximal reduced words $z, w \in \mathcal{W}_m^*$. By Theorem 6.19 we can view $\mathcal{Z}_\bullet^+ = \mathcal{Z}_w^+$, as generated by X_v with $v \leq_R w = u \cdot w_{\eta(i)}$. The latter generating set splits into the union $\{X_v : v \leq_R u\} \sqcup \{X_w\}$. If $v \leq_R u$ we have that $w_i \cdot v$ is reduced since $w_i \cdot u$ is reduced. Hence $\Gamma_i(X_v) = X_{w_i \cdot v} \in \mathcal{Z}_\bullet^+ \subset \mathcal{Z}_\bullet$ by (5.4) and (6.18).

Now, by definition, $X_w = \Gamma_u(X_{\eta(i)})$ so that $\Gamma_i(X_w) = \Gamma_i \circ \Gamma_u(X_{\eta(i)}) = \Gamma_z(X_{\eta(i)}) = \Gamma_{s_\bullet}(X_{\eta(i)})$ since $z = w_i \cdot u$ is reduced and maximal. The formula (5.14) in the proof of Corollary 5.3 then implies that $\Gamma_i(X_w) = \Gamma_{s_\bullet}(E_{\eta(i)}^{\ell_i}) = (-K_i^{-1}F_i)^{\ell_i} = -L_i^{-1}Y_i \in \mathcal{Z}_\bullet^{\leq 0} \subset \mathcal{Z}_\bullet$. Thus, Γ_i maps all generators of \mathcal{Z}_w^+ to \mathcal{Z}_\bullet as desired. \square

The action of the Artin groups on the maximal skew-commutative algebras is rather non-trivial. We leave it here as an open question whether \mathcal{A} acts effectively or faithfully, as suggested by the identifications with actions on Lie groups discussed in Sections 7.4 and 8.4.

For illustration, we gather below explicit formulae for the Lie type A_2 and odd \mathbb{k} , with $\nabla = (\zeta - \zeta^{-1})^\ell$. The action of Γ_1 on the Cartan algebra, given by $\Gamma_1(L_1) = L_1^{-1}$ and $\Gamma_1(L_2) = L_2L_1^{-1}$, factors through the quotient $\mathcal{A} = B_3 \rightarrow \mathcal{W} = S_3$. The remaining expressions are as follows.

$$\begin{aligned} \Gamma_1(X_1) &= -L_1^{-1}Y_1 & \Gamma_1(X_2) &= X_{(12)} & \Gamma_1(X_{(12)}) &= -X_2 + \nabla L_1^{-1}X_{(12)}Y_1 \\ \Gamma_1(Y_1) &= -L_1X_1 & \Gamma_1(Y_2) &= Y_{(12)} & \Gamma_1(Y_{(12)}) &= -Y_2 - \nabla L_1Y_{(12)}X_1 \end{aligned} \tag{6.59}$$

Those in the first two columns are immediate from (6.20) and (6.18). For those in the third column, we invoke the identity $X_{(12)} + X_{(21)} + \nabla X_1X_2 = 0$ with $\nabla = (\zeta - \zeta^{-1})^\ell$, which is obtained by specializing Propositions 6.17. We also use the identity $\Gamma_1(X_{(21)}) = X_2$, which is implied by Propositions 5.1. The action on $Y_{(12)}$ follows via application of Ω .

It is now not hard to find many infinite families of braids in B_3 for which the action in (6.59) on generators produces polynomials whose (naïve) degrees monotonously increase with the crossing numbers of the braids.

6.5. \mathcal{Z} -Induced Ideals, \mathcal{G} -Invariance, and \mathcal{G} -Grading. In this section, we describe the general process of obtaining ideals for the full quantum groups from ideals of the skew-central subalgebras. To keep the exposition more readable, we will often suppress the choice of a maximal word $z \in \mathcal{W}_m^*$ in the notation of the maximal algebra $\mathcal{Z}_\bullet^\square = \mathcal{Z}_z^\square$. By Theorem 6.19 this is unambiguous for Lie types other than G_2 . In the case G_2 it is tacitly understood that there may be (at most) two or four choices, depending on \square .

As skew-polynomial algebras, the $\mathcal{Z}_\bullet^\square$ have a rich ideal structure. Ideals in $\mathcal{Z}_\bullet^\square$ extend to ideals in U_ζ^\square provided they are commensurable with the action of an elementary abelian 2-group stemming from commutation relations as in Proposition 6.7.

More formally, the latter group is defined as $\mathcal{G} = \mathbb{F}_2^\Delta \times \mathbb{F}_2^\Delta$, acting on the algebra \mathcal{Z}_w for any $w \in \mathcal{W}^*$. An element $(\bar{\mu}, \bar{v}) \in \mathcal{G}$ acts on generators in \mathcal{Z}_w associated to some $v \leq_R w$ by multiplication with ± 1 via the formulae

$$(\bar{\mu}, \bar{v})_v X_v = \kappa(\gamma(v), \bar{\mu}) X_v, \quad (\bar{\mu}, \bar{v})_v Y_v = \kappa(\gamma(v), \bar{v}) Y_v, \quad \text{and} \quad (\bar{\mu}, \bar{v})_v L_i = \kappa(\alpha_i, \bar{\mu} + \bar{v}) L_i. \quad (6.60)$$

Since the $\{X_v, Y_v, L_i : v \leq_R w, 1 \leq i \leq n\}$ are independent (sign-commuting) generators of \mathcal{Z}_w , the formulae in (6.60) uniquely extend to a \mathcal{G} -action on \mathcal{Z}_w as algebra automorphisms. The action clearly preserves all \mathcal{Z}_w^\square subalgebras and specializes to a \mathbb{F}_2^Δ -action on either \mathcal{Z}_w^+ or \mathcal{Z}_w^- as one of the components acts trivially.

The commutation relations from Proposition 6.7 and (6.25) can now be rephrased more compactly using the \mathcal{G} -action. Specifically, for a fully homogeneous element $b \in U_q$ and $W \in \mathcal{Z}_w$ we find

$$W \cdot b = b \cdot \mathfrak{o}^*(b)_v W, \quad \text{where} \quad \mathfrak{o}^*(b) = (\mathfrak{o}^+(b), \mathfrak{o}^-(b)) \in \mathcal{G}, \quad (6.61)$$

by straightforward verification on the generators. Here we use also $w = \mathfrak{o}^+ + \mathfrak{o}^- \pmod{2}$ and the extension to general elements via standard action properties. We say that a subspace $A \subseteq \mathcal{Z}_w^\square$ is \mathcal{G} -invariant if $g_v x \in A$ for all $g \in \mathcal{G}$ and $x \in A$.

Lemma 6.23. *The \mathcal{G} -action on $\mathcal{Z}_\bullet^\square$ does not depend on a choice of $w \in \mathcal{W}_m^*$.*

If $W \in \mathcal{Z}_\bullet = \mathcal{Z}_{s_\bullet}$ and $w, z \in \mathcal{W}_m^*$ represent s_\bullet denote by $(\bar{\mu}, \bar{v})_v^w$ and $(\bar{\mu}, \bar{v})_v^z$ the respective actions defined via (6.60). Equation (6.61) then implies $b \cdot \mathfrak{o}^*(b)_v^z W = b \cdot \mathfrak{o}^*(b)_v^w W$ for homogeneous elements b . Choosing respective PBW basis elements one may then infer that the actions need to be the same. Another immediate consequence of (6.61) is as follows.

Lemma 6.24. *If J is a \mathcal{G} -invariant left or right ideal in $\mathcal{Z}_\bullet^\square$, then J is also a two-sided ideal.*

For example, suppose J is left sided, $W \in J$, and $b \in \mathcal{Z}_\bullet^+$ is homogeneous. Then $W \cdot b = b \cdot W'$ where $W' = \mathfrak{o}^*(b)_v W \in J$ by \mathcal{G} -invariance. Thus $W \cdot b = b \cdot W' \in J$ since J is a left ideal and, hence, also a right ideal. It is not difficult to find counterexamples to Lemma 6.24 if the condition of \mathcal{G} -invariance is dropped. In the next proposition, we consider *algebra* ideals only, disregarding any coalgebra structures.

Proposition 6.25. Suppose J is a \mathcal{G} -invariant (any-sided) ideal in $\mathcal{Z}_\bullet^\square$. Then $\hat{J} = J \cdot U_\zeta^\square$ is a two-sided ideal in U_ζ^\square . Moreover, the multiplication map from (6.38) induces an isomorphism of $\mathcal{Z}_\bullet^\square$ -modules,

$$(\mathcal{Z}_\bullet^\square / J) \otimes_\Lambda \mathfrak{S}_{z\blacktriangleright}^{[0, \mathbf{l}], +} \longrightarrow U_\zeta^\square / \hat{J} : ([z], b) \mapsto [z \cdot b], \quad (6.62)$$

where $[.]$ denotes the class of an element in the respective quotient algebra and $z \in \mathcal{W}_m^*$ is a reduced word of maximal length.

Proof. \hat{J} is a right ideal by definition. To see it is also a left ideal, write general elements $B \in U_\zeta^\square$ and $C \in J$ as $B = \sum_s b_s$ and $C = \sum_t W_t c_t$ where $b_s, c_t \in U_\zeta^\square$ are fully homogeneous and $W_t \in J$. (6.61) then implies $B \cdot C = \sum_{s,t} \phi^*(b_s)_{\blacktriangleright} (W_t) b_s c_t$. By \mathcal{G} -invariance of J we have $\phi^*(b_s)_{\blacktriangleright} (W_t) \in J$ so that $B \cdot C \in \hat{J}$. Thus $U_\zeta^\square \cdot J \subseteq \hat{J}$, which implies $U_\zeta^\square \cdot \hat{J} = U_\zeta^\square \cdot J \cdot U_\zeta^\square \subseteq \hat{J} \cdot U_\zeta^\square \subseteq \hat{J}$ as desired.

For the second assertion, it suffices to show that $m_z^+ (J \otimes_\Lambda \mathfrak{S}_{z\blacktriangleright}^{[0, \mathbf{l}], +}) = \hat{J}$. The inclusion of the image in \hat{J} is clear from the ideal property. Surjectivity of the restriction of m_z^+ from $J \otimes_\Lambda \mathfrak{S}_{z\blacktriangleright}^{[0, \mathbf{l}], +}$ to \hat{J} is immediate from the commutative diagram below, where the top horizontal arrow is an isomorphism and the two vertical arrows are surjections.

$$\begin{array}{ccc} J \otimes_\Lambda \mathcal{Z}_\bullet^\square \otimes_\Lambda \mathfrak{S}_{z\blacktriangleright}^{[0, \mathbf{l}], +} & \xrightarrow{\text{id}_J \otimes m_z^+} & J \otimes_\Lambda U_\zeta^\square \\ \downarrow m_J \otimes_\Lambda \text{id} & & \downarrow m_J \\ J \otimes_\Lambda \mathfrak{S}_{z\blacktriangleright}^{[0, \mathbf{l}], +} & \xrightarrow{m_z^+} & \hat{J} \end{array}$$

□

An important special family of \mathcal{G} -invariant ideals are those generated by augmentation ideals of the subalgebra \mathcal{Z}_w^+ for some reduced word $w \in \mathcal{W}$ and $\square = +, \square = \geq 0$, or \square omitted for the full algebra. They are defined as

$$\mathcal{K}(w)^\square = (A_w^+) = A_w^+ \cdot \mathcal{Z}_\bullet^\square \quad \text{where} \quad A_w^+ = \ker(\varepsilon : \mathcal{Z}_w^+ \rightarrow \Lambda). \quad (6.63)$$

It is clear from the PBW bases that $\mathcal{K}(w)^+$ is identical to the ideal generated by $\{X_v : \emptyset \neq v \leq_R w\}$. Assume some $z \in \mathcal{W}_m^*$ with $w \leq_R z$ and let $s = r(w)$. It is then not hard to see that, as a free Λ -module, $\mathcal{K}(w)^+$ is spanned by all $X_{z\blacktriangleright}^\psi$ in $\mathfrak{B}_{z\blacktriangleright}^{(\mathbf{l}, s), +}$ for which the restriction of ψ to the inversion set $\mathcal{N}(s) \subseteq \Phi^+$ is non-zero.

For a more concise description of the ideals in the full quantum algebras, we introduce abbreviated notation for exponent subsets complementary to the $[0, \mathbf{l}]_s$ from (6.37), namely,

$$[\mathbf{l}, \infty)^s = \{\psi \in \mathcal{E}(s) : \exists \alpha \in \mathcal{N}(s) \text{ for which } \psi(\alpha) \geq \ell_\alpha\} \quad \text{and} \quad [\mathbf{l}, \infty)^s = [\mathbf{l}, \infty)^{s^*}. \quad (6.64)$$

It is clear from the isomorphism in (6.38) and with notation as in (5.38) that the associated ideals in U_ζ^+ may also be written as the two-sided ideals generated by the spans corresponding to these exponent sets. That is,

$$\hat{\mathcal{K}}(w)^+ = U_\zeta^+ \cdot \mathfrak{S}_{w\blacktriangleright}^{[\mathbf{l}, \infty)^s, +} = U_\zeta^+ \cdot \mathfrak{S}_{\blacktriangleleft w}^{[\mathbf{l}, \infty)^s, +}. \quad (6.65)$$

Observe that for a maximal word $z \in \mathcal{W}_m^*$ we have $\mathcal{K}(z)^+ = A_z^+$, as defined in (6.63), and $\mathcal{K}(z)^{\geq 0} = A_z^+ \cdot \mathcal{Z}^0$. Correspondingly, (6.65) reduces to $\hat{\mathcal{K}}(z)^+ = \mathfrak{S}_{z\bullet}^{[l,\infty)^*,+}$.

Note that since the automorphisms from Corollary 6.11 preserve the counit, the analogs of the identities in (6.35) also hold for the respective ideals. Applied to the induced ideals in the quantum algebras one finds

$$\Omega(\hat{\mathcal{K}}(w)^\pm) = \hat{\mathcal{K}}(w)^\mp, \quad \mathfrak{O}(\hat{\mathcal{K}}(z)^\pm) = \hat{\mathcal{K}}(z^\dagger)^\pm, \quad \text{and} \quad S(\hat{\mathcal{K}}(z)^{\geq 0(\leq 0)}) = \hat{\mathcal{K}}(z^\dagger)^{\geq 0(\leq 0)}. \quad (6.66)$$

Similarly, Theorem 6.19 and the fact that the definition in (6.63) does not refer to choices of bases imply the next remark.

Corollary 6.26. *Assume the Lie type is not G_2 , as before, and $s = r(w) = r(w')$ for two reduced words $w, w' \in \mathcal{W}^*$. Then $\mathcal{K}(w)^\square = \mathcal{K}(w')^\square$.*

We may thus write $\mathcal{K}(s)^\square$ for Lie types different from G_2 . We then adopt the abbreviated notation for a maximal reduced word $z \in \mathcal{W}_m^*$

$$\mathcal{K}_\bullet^\square = \mathcal{K}(z)^\square = \mathcal{K}(s_\bullet)^\square \quad \text{and} \quad \hat{\mathcal{K}}_\bullet^\square = \hat{\mathcal{K}}(z)^\square = \hat{\mathcal{K}}(s_\bullet)^\square, \quad (6.67)$$

where the first equalities also apply to the G_2 case (for which $\mathcal{K}_\bullet^\square$ may be ambiguous). Furthermore, \mathcal{A} -invariance of \mathcal{Z}_\bullet , as asserted in Theorem 6.22, the recursion (6.19), and the basis generators for $\mathcal{K}(s)$ imply the following recursion for ideals.

Lemma 6.27. *Suppose $\mathcal{K}(s)$ is the ideal of the full algebra \mathcal{Z}_\bullet , $l(s \cdot t) = l(s) + l(t)$ for $s, t \in \mathcal{W}$, and Lie type $\neq G_2$. Then*

$$\mathcal{K}(s \cdot t) = \mathcal{K}(s) + \Gamma_s(\mathcal{K}(t)). \quad (6.68)$$

Analogous augmentation ideals $\mathcal{Z}_\bullet^{\leq 0}$ and \mathcal{Z}_\bullet^- are defined in using $A_s^- = \ker(\varepsilon : \mathcal{Z}_s^- \rightarrow \Lambda)$ as generators. For a pair $s, t \in \mathcal{W}$, combined ideals in the full algebra \mathcal{Z}_\bullet are given as

$$\mathcal{K}(s, t) = (A_s^+, A_t^-) = (A_s^+ + A_t^-)\mathcal{Z}_\bullet. \quad (6.69)$$

Observe also that $A_1^\pm = 0$ and than (6.58) implies $\Omega(A_s^+) = A_s^-$, which yields the identity

$$\mathcal{K}(s) = \mathcal{K}(s, 1) = \Omega(\mathcal{K}(1, s)). \quad (6.70)$$

For later convenience, we introduce compact notation for the ideals in \mathcal{Z}_\bullet corresponding to the longest element $s_\bullet \in \mathcal{W}$ as

$$\mathcal{K}_{\bullet\bullet} = \mathcal{K}(s_\bullet, 1), \quad \mathcal{K}_{\bullet\bullet} = \mathcal{K}(1, s_\bullet), \quad \text{and} \quad \mathcal{K}_{\bullet\bullet} = \mathcal{K}(s_\bullet, s_\bullet). \quad (6.71)$$

Also for Lie types different from G_2 (or keeping in mind the aforementioned possible ambiguity for G_2) we use the suggested notation from Proposition 6.25 for the entailed ideals in U_ζ such as, for example,

$$\hat{\mathcal{K}}(s), \quad \hat{\mathcal{K}}(s, t), \quad \hat{\mathcal{K}}_{\bullet\bullet} = \hat{\mathcal{K}}(s_\bullet, 1), \quad \text{or} \quad \hat{\mathcal{K}}_{\bullet\bullet} = \hat{\mathcal{K}}(s_\bullet, s_\bullet). \quad (6.72)$$

More generally, any subset $M \subset \mathfrak{B}_{w\blacktriangleright}^{(\mathfrak{l},s),+}$ generates a \mathcal{G} -invariant ideal $J = (M)$ of \mathcal{Z}_w^+ since all monomials in $\mathfrak{B}_{w\blacktriangleright}^{(\mathfrak{l},s),+}$ are homogeneous in the sense that $g \triangleright X_{w\blacktriangleright}^\psi = \pm X_{w\blacktriangleright}^\psi$ for all $g \in \mathcal{G}$. So, if $M = (X_{w\blacktriangleright}^{\chi_1}, \dots, X_{w\blacktriangleright}^{\chi_K})$ a PBW basis for \mathcal{Z}_w^+ / M is given by the set of all $X_{w\blacktriangleright}^\psi$ for which $\psi \not\geq \chi_i$ for any $i = 1, \dots, K$. The partial order used here is defined as $\psi \geq \chi$ iff $\psi(\alpha) \geq \chi(\alpha)$ for all $\alpha \in \mathcal{N}(w)$. Consequently, a PBW basis of U_ζ^+ / \widehat{M} is given by the set of all $E_{w\blacktriangleright}^\phi$ for which $\phi \not\geq \mathfrak{l} \cdot \chi_i$ for any $i = 1, \dots, K$.

We briefly elaborate here on the caution expressed in the introduction to this monograph regarding quotients by only the X_i for simple roots. Let M_Δ be ideal in \mathcal{Z}_z^+ generated by the set $\{X_i : i = 1, \dots, n\} = \{X_v : \gamma(v) \in \Delta, v \leq_R z\}$, see Proposition 5.1. Considering only the positive part, a PBW basis of $U_\zeta^+ / \widehat{M}_\Delta$ is given by $X_{w\blacktriangleright}^\psi$ with $\psi(\alpha_i) < \ell_i$ on simple roots but ψ unbounded on $\Phi^+ \setminus \Delta$, so that $U_\zeta^+ / \widehat{M}_\Delta$ is of *infinite* rank if $n > 1$.

Even more concretely, for Lie type A_2 and $\mathfrak{k} = 2\ell$, the algebra obtained by imposing only relations $E_1^\ell = 0$ and $E_2^\ell = 0$ (but not $E_{(12)}^\ell = 0$) has PBW basis $\{E_1^{i_1} E_{(12)}^j E_2^{i_2} : 0 \leq i_1, i_2 < \ell$ and $j \in \mathbb{N}_0\}$. The observations for the full quantum groups are analogous, leading to algebras without the desired properties of a restricted quantum group.

We mention also a natural \mathcal{G} -grading z^* on any of the \mathcal{Z}_w^+ algebras, which provides a dual picture to \mathcal{G} -invariance and which is analogous to \mathfrak{o}^* on U_q . Given $\alpha = \gamma(v)$ we set $z^*(X_v) = (\bar{\alpha}, 0)$ and $z^*(Y_v) = (0, \bar{\alpha})$ where $\bar{\alpha} \in \mathbb{F}_2^\Delta$ is the image of the canonical lattice map $\Phi^+ \rightarrow \mathbb{F}_2^\Delta$ that assigns $\alpha_i \mapsto \alpha_i$. Similarly, we set $z^*(L^\mu) = (\bar{\mu}, \bar{\mu})$ for $\mu \in \mathbb{Z}^\Delta$ and L as in (6.16). By Proposition 6.10 the grading extends uniquely to all of \mathcal{Z}_w^+ , yielding a decomposition

$$\mathcal{Z}_w^+ = \bigoplus_{g \in \mathcal{G}} (\mathcal{Z}_w^+)_g. \quad (6.73)$$

A pairing of \mathcal{G} with itself is now given by the symmetric bihomomorphism

$$v : \mathcal{G} \times \mathcal{G} \longrightarrow \{+1, -1\} \quad \text{with} \quad v((\bar{\mu}, \bar{v}), (\bar{\mu}', \bar{v}')) = \varepsilon(\bar{\mu}, \bar{\mu}') \cdot \varepsilon(\bar{v}, \bar{v}'), \quad (6.74)$$

where ε is as defined in (6.11).

Suppose now that $\mathfrak{k} = 2\ell$ and that Φ is of oddly laced Lie type (ADEG). Then Lemma 6.5 implies that (6.60) can be expressed more compactly. Namely, for any $g, h \in \mathcal{G}$ and $W \in (\mathcal{Z}_w^+)_g$

$$h \triangleright W = v(g, h) \cdot W. \quad (6.75)$$

Assume, further, that the Lie type is A_{2k}, E_6, E_8 , or G_2 . For each of these, ε and, hence, v are non-degenerate pairings of respective \mathbb{F}_2 lattices. So, in these cases a submodule $J \subseteq \mathcal{Z}_w^+$ is \mathcal{G} -invariant if and only if it respects the decomposition in (6.73), meaning that for $J_g = J \cap (\mathcal{Z}_w^+)_g$ we have

$$J = \bigoplus_{g \in \mathcal{G}} J_g. \quad (6.76)$$

The decomposition in (6.73) can be described in more concrete terms under further assumptions. Aside from the restriction to $\{A_{2k}, E_6, E_8, G_2\}$, suppose further ℓ is even ($\mathfrak{k} \equiv 0 \pmod{4}$)

and $w \in \mathcal{W}_m^*$ is maximal so that, by Corollary 6.14, \mathcal{Z}_w^\square can be identified with a polynomial algebra as in (6.41). There is a natural epimorphism from the former with indeterminates labeled by all positive roots to the polynomial algebra with indeterminates given by only the simple roots.

$$\Lambda [\{x_\alpha : \alpha \in \Phi^+\}] \rightarrow \Lambda [z_1, \dots, z_n] : x_\alpha \mapsto \prod_i z_i^{c_i} \quad \text{for } \alpha = \sum_i c_i \alpha_i.$$

Given $\mu \in \mathbb{F}_2^\Delta$ we say that a polynomial $f \in \Lambda [z_1, \dots, z_n]$ has parity μ if

$$f(z_1, \dots, z_{i-1}, -z_i, z_{i+1}, \dots, z_n) = (-1)^{\mu_i} f(z_1, \dots, z_n) \quad \forall i = 1, \dots, n.$$

Thus, a polynomial $p \in \Lambda [\{x_\alpha : \alpha \in \Phi^+\}]$ represents an element in $(\mathcal{Z}_w^+)_{{\bar{\mu}}, 0}$ with $\bar{\mu} = (\mu_1, \dots, \mu_n)$ precisely if it is mapped to a polynomial in $\Lambda [z_1, \dots, z_n]$ of parity μ , in which case we assign the parity $\bar{\mu}$ also to p and well as the respective element in \mathcal{Z}_w^+ .

Thus, for example for type A_2 , the element $X_{(12)}^3 + X_1^5 X_2$ is of parity $(1, 1)$ in $(\mathcal{Z}_w^+)_{((1, 1), 0)}$, while the element $X_1 + X_2$ is not of any fixed parity since $X_1 \in (\mathcal{Z}_w^+)_{((1, 0), 0)}$ but $X_2 \in (\mathcal{Z}_w^+)_{((0, 1), 0)}$. Thus, $(X_{(12)}^3 + X_1^5 X_2)$ is a \mathcal{G} -invariant ideal of \mathcal{Z}_w^+ but $(X_1 + X_2)$ is not. The smallest \mathcal{G} -invariant ideal the latter is contained in is $(X_1 + X_2, 2X_1)$.

Any finite collection of elements with fixed parity in \mathcal{Z}_w^+ thus generates a \mathcal{G} -invariant ideal. For a converse statement, suppose \mathbb{L} is a Noetherian integral domain in which 2 is a unit, suppose $\jmath : \Lambda \rightarrow \mathbb{L}$ is a ring epimorphism, and let $\mathcal{Z}_{w, \mathbb{L}}^+ = \mathcal{Z}_w^+ \otimes_{\mathbb{Z}} \mathbb{L}$. Then, in fact, all \mathcal{G} -invariant ideals of $\mathcal{Z}_{w, \mathbb{L}}^+$ are given as ideals generated by a finite collection of elements of fixed parity.

7. HOPF IDEALS FOR $U_\zeta(\mathfrak{sl}_{n+1})$ FROM SOLVABLE LINEAR ALGEBRAIC GROUPS

The A_n case is computationally more accessible than other Lie types, allowing explicit descriptions of the correspondence between the skew-central subalgebras discussed in the previous chapter and the (semi) classical coordinate rings of Borel subgroups. We start by providing basic background on linear algebraic groups and their coordinate rings in Section 7.1.

In Section 7.2 we construct specific generators of $U_\zeta(\mathfrak{g})$ for both $\mathfrak{g} = \mathfrak{sl}_{n+1}$ and $\mathfrak{g} = \mathfrak{gl}_{n+1}$. They are used in Section 7.3 to establish $\mathcal{Z}_\zeta^{\geq 0}$ as a Hopf subalgebra with explicit forms for the coproduct. The latter are used for identifications with centrally extended quantum matrix algebras, whose complexifications for $\mathbb{k} \not\equiv 2 \pmod{4}$, coincide with the coordinate rings of upper triangular matrices.

Recall from Section 6.3 that for $\mathbb{k} \equiv 0 \pmod{4}$ the algebra $\mathcal{Z}_\zeta^{\geq 0}$ is commutative but *not* central in $U_\zeta(\mathfrak{g})$. The remainder and Section 7.4 discuss the construction of Hopf ideals from subgroups and the relation of the $\mathcal{K}(s)^{\geq 0}$ ideals from (6.63) to respective Bruhat subgroups.

7.1. Basic Background and Elementary Examples. We provide here some basic vocabulary and elementary examples as preparation and motivation for the correspondences between Hopf ideals and algebraic subgroups discussed in Sections 7 and 8. Most of the material reviewed here can be found, for example, in the early chapters of [Spr09] and [Mil17].

Suppose G is any algebraic set and M some ring of functions on G . Further, assume H is any subset of G and I some ideal in M . We denote the *vanishing ideal* of H in M as well as the *vanishing locus* of I in G as

$$\mathcal{I}(H) = \{f \in M : f(h) = 0 \forall h \in H\} \quad \text{and} \quad \mathcal{V}(I) = \{g \in G : f(g) = 0 \forall f \in I\}, \quad (7.1)$$

respectively. We, trivially, have inclusions $H \subseteq \mathcal{V}(\mathcal{I}(H))$ and $I \subseteq \mathcal{I}(\mathcal{V}(I))$. Suppose \mathbb{k} is an algebraically closed field, $G = \mathbb{k}^m$, and $M = \mathbb{k}[x_1, \dots, x_m]$ the ring of polynomials viewed as functions on G . In this case, Hilbert's Nullstellensatz and the definition of the Zariski topology imply that

$$\mathcal{I}(\mathcal{V}(I)) = \sqrt{I} \quad \text{and} \quad \mathcal{V}(\mathcal{I}(H)) = \text{Zcl}(H), \quad (7.2)$$

where \sqrt{I} denotes the radical of I and $\text{Zcl}(H)$ the usual Zariski closure, establishing a one-to-one correspondence between Zariski closed subsets and radical ideals. This correspondence generalizes from \mathbb{k}^n to any algebraic set G .

Recall that a linear algebraic group is an affine algebraic set $G \subseteq \mathbb{k}^N$ with group operations that are morphisms of varieties. The coordinate ring on G , customarily denoted by $\mathbb{k}[G] = \mathbb{k}[x_1, \dots, x_N]/\mathcal{I}(G)$, admits for each $g \in G$ a well-defined algebraic evaluation map $e_g : \mathbb{k}[G] \rightarrow \mathbb{k}$ with $e_g([p]) = p(g)$ where p is a polynomial in $\{x_i\}$.

Importantly, these admit a natural Hopf algebra structure, see for example [Spr09, Ch 2], [Mil17, Ch 3]. The coproduct $\Delta : \mathbb{k}[G] \rightarrow \mathbb{k}[G \times G] \cong \mathbb{k}[G] \otimes \mathbb{k}[G]$ is the dual of the product morphism on G such that $(e_a \otimes e_b) \circ \Delta = e_{ab}$ or, more informally, $\Delta([p])(a, b) = p(ab)$. An antipode is obtained similarly from inversion on G .

As a central example, consider $\mathbb{k}[\text{GL}(n, \mathbb{k})] \cong \mathbb{k}[\{x_{ij}\}, \tau]/\mathcal{I}$, where $\mathcal{I} = (\tau \cdot \det(x) - 1)$. The presentation implies that any element $[p] \in \mathbb{k}[\text{GL}(n, \mathbb{k})]$ can be represented by a polynomial $p \in \mathbb{k}[\{x_{ij}\}, \tau]$ in $N = n^2 + 1$ indeterminates. The coalgebra structure is readily worked out as

$$\Delta([x_{ij}]) = \sum_{k=1}^n [x_{ik}] \otimes [x_{kj}] \quad \text{and} \quad \Delta([\tau]) = [\tau] \otimes [\tau] \quad (7.3)$$

and it is immediate that \mathcal{I} is a Hopf ideal.

As usual, for a Hopf algebra A , denote the subgroup of *group-like* elements as $\text{Gr}(A) = \{a \in A : \Delta(a) = a \otimes a\}$. Applied to a coordinate ring $A = \mathbb{k}[G]$ this group is naturally identified with the algebraic character group $\text{Spec}_m(G) = \text{Hom}_{\text{alg}}(G, \mathbb{k}^\times) = \text{Gr}(\mathbb{k}[G])$ consisting of group homomorphisms $\sigma : G \rightarrow \mathbb{k}^\times$ that are also morphisms of algebraic sets. For algebraically closed \mathbb{k} with $\text{char}(\mathbb{k}) = 0$ the character group of $\text{GL}(n, \mathbb{k})$ is infinite cyclic generated by the determinant function. That is,

$$\text{Spec}_m(\text{GL}(n, \mathbb{k})) = \mathbb{Z}_{(\det)}. \quad (7.4)$$

Closely related is the fact that $\text{GL}(n, \mathbb{k})$ is linearly reductive or, equivalently, that $\mathbb{k}[\text{GL}(n, \mathbb{k})]$ is cosemisimple for such \mathbb{k} .

Next, let $\delta : \mathbb{k}[\mathrm{GL}(n, \mathbb{k})] \rightarrow \mathbb{N}_0$ be the function assigning the minimal degree of representatives $\delta(f) = \min\{\deg(p) : f = [p]\}$, where $\deg(p)$ denotes the total degree of p as a polynomial in the $n^2 + 1$ variables. Observe that the finite-dimensional linear subspaces $D_m = \{f \in \mathbb{k}[\mathrm{GL}(n, \mathbb{k})] : \delta(f) \leq m\}$ form an ascending chain of subcoalgebras. Indeed, if $f = [p]$ and p is of minimal degree $d = \delta(f)$, it is clear from the formulae in (7.3) that $\Delta(f)$ can be written as a summation of pure tensors of monomials, each of which with degree $\leq d$.

It is a basic fact that for any linear algebraic group H over \mathbb{k} there exists a (Zariski) closed monomorphism $j_H : H \hookrightarrow \mathrm{GL}(n, \mathbb{k})$ of algebraic groups for some n , e.g. [Spr09, 2.3.7]. Given such an embedding, we may define an ascending sequence of finite-dimensional linear spaces $D_m(H) = j_H^*(D_m)$, each of which is clearly a subcoalgebra of $\mathbb{k}[H]$.

The next lemma establishes a one-to-one correspondence between Hopf ideals and (Zariski) closed algebraic subgroups in characteristic zero. It is easy to find examples in positive characteristic for which *ii*) does not hold.

Lemma 7.1. *Let G be a linear algebraic group over an algebraically closed \mathbb{k} with $\mathrm{char}(\mathbb{k}) = 0$.*

- i) *If $J \subset \mathbb{k}[G]$ is a Hopf ideal, then $H = \mathcal{V}(J)$ is a closed algebraic subgroup in G .*
- ii) *Further, J is a radical ideal so that $J = \mathcal{I}(H) = \mathcal{I}(\mathcal{V}(J))$.*
- iii) *If $H \leq G$ is any subgroup, then $\mathcal{I}(H)$ is a Hopf ideal in $\mathbb{k}[G]$.*
- iv) *In this case $\mathcal{I}(H) = \mathcal{I}(\mathrm{Zcl}(H))$.*
- v) *If I and J are Hopf ideals in $\mathbb{k}[G]$, then $\mathcal{V}(I + J) = \mathcal{V}(I) \cap \mathcal{V}(J)$.*
- vi) *If H and K are closed algebraic subgroups of G , then $\mathcal{I}(H \cap K) = \mathcal{I}(H) + \mathcal{I}(K)$.*

Proof. For *i*) note that $a \in \mathcal{V}(J)$ iff $J \subseteq \ker(e_a)$. So, for $a, b \in \mathcal{V}(J)$ and J a bi-ideal we have $e_{ab}(J) = e_a \otimes e_b(\Delta(J)) \subseteq e_a \otimes e_b(J \otimes \mathbb{k}[G] + \mathbb{k}[G] \otimes J) = \{0\}$ and hence $ab \in \mathcal{V}(J)$. The argument for the antipode is analogous. Given that $\mathrm{char}(\mathbb{k}) = 0$ and using the fact that $\mathbb{k}[G]/J$ is a commutative Hopf algebra, Theorem 13.1.2 in [Swe69] states that $\mathbb{k}[G]/J$ is a reduced \mathbb{k} -algebra. This, in turn, means that $J = \sqrt{J}$ is a radical ideal. The statements in *ii*) follow now from (7.2) as $J = \mathcal{I}(\mathcal{V}(J)) = \mathcal{I}(H)$.

For *iii*) use the chain of finite-dimensional coalgebras $D_0 \subset D_1 \subset \dots$ with $\mathbb{k}[G] = \bigcup_j D_j$ described above. Set $\mathcal{I}_m = \mathcal{I}(H) \cap D_m$, let $D_m = \mathcal{I}_m \oplus R_m$ for some complementary space, and pick a basis $\{g_v\}$ for R_m with $\dim(R_m) = M$. Let $\vec{g} : G \rightarrow \mathbb{k}^M$ be given by $\vec{g}(a) = (g_1(a), \dots, g_M(a))$. Thus, if $\vec{v}^\top \vec{g}(a) = 0$ for all $a \in H$ then $\vec{v}^\top \vec{g} \in \mathcal{I}(H)$ and hence $\vec{v} = 0$ by linear independence. Given $f \in \mathcal{I}_m$, there are now unique coefficients $b_{v,\mu} \in \mathbb{k}$ such that $\Delta(f) = \sum_{v,\mu} b_{v,\mu} g_v \otimes g_\mu + T$ with $T \in \mathcal{I}_m \otimes D_m + D_m \otimes \mathcal{I}_m$. Evaluating at (a, b) for $a, b \in H$ so that $ab \in H$ we have $0 = f(ab) = e_a \otimes e_b \Delta(f) = \sum_{v,\mu} b_{v,\mu} g_v(a) g_\mu(b) + e_a \otimes e_b(T) = \vec{g}(a)^\top B \vec{g}(b)$, where $B = (b_{v,\mu})$ and $\mathcal{I}_m \subseteq \ker(e_a)$.

As $\vec{g}(a)^\top B \vec{g}(b) = 0$ holds for all $a, b \in H$, the previous remarks imply $B = 0$ and, hence, $\Delta(f) = T \in \mathcal{I}_m \otimes D_m + D_m \otimes \mathcal{I}_m \subseteq \mathcal{I}(H) \otimes \mathbb{k}[G] + \mathbb{k}[G] \otimes \mathcal{I}(H)$, proving that $\mathcal{I}(H)$ is a bi-ideal.

Invariance under the antipode is clear from $S(f)(a) = f(a^{-1}) = 0$ if $a \in H$ and $f \in \mathcal{I}(H)$. Finally, combining *iii*) with *ii*) and (7.2) we find $\mathcal{I}(H) = \mathcal{I}(\mathcal{V}(\mathcal{I}(H))) = \mathcal{I}(\text{Zcl}(H))$.

The last two statements follow from the standard correspondence between algebraic sets and radical ideals. The only additional observation needed is that the sum of two Hopf ideals is again a Hopf ideal so that by *ii*) $\mathcal{I}(H \cap K) = \sqrt{(\mathcal{I}(H) + \mathcal{I}(K))} = \mathcal{I}(H) + \mathcal{I}(K)$. \square

The most elementary example is the additive group \mathbb{k}_a , for which $\mathbb{k}[\mathbb{k}_a] = \mathbb{k}[x]$ with $\Delta(x) = x \otimes 1 + 1 \otimes x$ and $S(x) = -x$. The vanishing locus of any polynomial in $\mathbb{k}[x]$ is finite so that a non-trivial Hopf ideal needs to correspond to a finite subgroup of \mathbb{k}_a . In $\text{char}(\mathbb{k}) = 0$ the only such subgroup is $\{0\}$. Thus, by Lemma 7.1, the only proper Hopf ideal of $\mathbb{k}[x]$ is the augmentation ideal $(x) = \mathcal{I}(\{0\})$. In characteristic $p > 0$ we, additionally, have that the principal ideal (x^p) generated by x^p is a Hopf ideal.

For the multiplicative group $\mathbb{k}_m = (\mathbb{k}^\times, \cdot)$, we have $\mathbb{k}[\mathbb{k}_m] = \mathbb{k}[\lambda, \lambda^{-1}] = \mathbb{k}[\lambda, \tau]/(\lambda\tau - 1)$ with $\Delta(\lambda) = \lambda \otimes \lambda$. Recall any finite subgroup of \mathbb{k}_m is cyclic and, thus, of the form $Z_r = \langle \xi \rangle$ where ξ a primitive r -th root of unity. We denote $Z_1 = \{1\}$ and use the convention $Z_0 = \mathbb{k}^\times$. The Z_r are algebraic subgroups with vanishing ideal $F_r = \mathcal{I}(Z_r) = (\lambda^r - 1)$ so that $\{F_r : r \in \mathbb{N}\}$ is the complete set of non-trivial Hopf ideals of $\mathbb{k}[\lambda, \lambda^{-1}]$.

A somewhat more involved example is the semidirect product $P = \mathbb{k}^\times \ltimes \mathbb{k}$ with group product given by $(\lambda, x)(\lambda', x') = (\lambda\lambda', x\lambda' + x')$. It may be viewed as the closed algebraic subgroup of the upper triangular matrices in GL_2 with 1 in the upper left entry. Alternatively, P may be viewed as $\text{PGL}_2^{\geq 0}$, given by the image of the upper triangular matrices in the projective linear group PGL_2 . Denote abelian algebraic subgroups as $\bar{Z}_r = Z_r \times \{0\}$ and $U = \{1\} \times \mathbb{k} = [P, P] \cong \mathbb{k}_a$.

Lemma 7.2. *Any closed algebraic subgroup of P is of the form $g\bar{Z}_r g^{-1}$ or $\bar{Z}_r U$ with $g \in U$ and $r \in \mathbb{N}_0$.*

Proof. Suppose H is an algebraic subgroup in P . Let $\sigma : P \rightarrow \mathbb{k}^\times$ be the abelianization map and note that $S = \sigma(H) \leq \mathbb{k}^\times$ is an abelian (*a priori* only constructible) subgroup. Suppose $\rho \in S \setminus \{1\}$ is a non-trivial element, meaning $(\rho, y) \in H$ for some $y \in \mathbb{k}$. It is then easy to check that $g(\rho, 0)g^{-1} = (\rho, y)$ for $g = (1, (\rho - 1)^{-1}y) \in U$.

Note that $H \cap U = \ker(\sigma|_H)$ is an algebraic subgroup of \mathbb{k}_a and, hence, either $\mathbb{1} = (1, 0)$ or U . If $H \cap U = \{\mathbb{1}\}$ the map $\sigma : H \rightarrow S$ is a group isomorphism so that H is abelian. If $S \neq \{1\}$ there are, by the previous remark, $g \in U$ and $\rho \in \mathbb{k}^\times$ with $\rho \neq 1$ and $g(\rho, 0)g^{-1} \in H$. Since H is abelian it has to be in the centralizer of $g\bar{Z}_0 g^{-1} = C(g(\rho, 0)g^{-1}) \cong \mathbb{k}_m$. Thus, $H = g\bar{Z}_r g^{-1}$ for some $r \in \mathbb{N}$. If $S = \{1\}$ clearly $H = \{\mathbb{1}\}$.

Finally, assume $H \cap U = U$ and $S \neq \{1\}$. By the observation above we have that $(\rho, y) \in H$ implies $(\rho, 0) \in H$ as now $g \in H$. Hence, $H = S \ltimes \mathbb{k}$ so that $S \cong H \cap \bar{Z}_0$ is an algebraic group. It follows that $H = K_r \ltimes \mathbb{k} = \bar{Z}_r U$. If also $S = \{1\}$, clearly $H = U = \bar{Z}_1 U$. \square

For a clearer formal distinction between coordinates of and functions on G denote by $\hat{t}, \hat{x} : G \rightarrow \mathbb{k}$ the projections given by $\hat{t}(g) = \lambda$ and $\hat{x}(g) = x$ if $g = (\lambda, x)$. Similar to the previous

examples, we have $\mathbb{k}[\mathrm{PGL}_2^{\geq 0}] \cong \mathbb{k}[P] = \mathbb{k}[\hat{\lambda}, \hat{\lambda}^{-1}, \hat{x}]$ as a commutative algebra. The coproducts are easily worked out from the product rule on P as

$$\Delta(\hat{\lambda}) = \hat{\lambda} \otimes \hat{\lambda} \quad \text{and} \quad \Delta(\hat{x}) = \hat{x} \otimes \hat{\lambda} + 1 \otimes \hat{x}, \quad (7.5)$$

with $S(\hat{\lambda}) = \hat{\lambda}^{-1}$ and $S(\hat{x}) = -\hat{x}\hat{\lambda}^{-1}$.

Assuming $\mathrm{char}(\mathbb{k}) = 0$ and using Lemma 7.1, we obtain a complete list of Hopf ideals for $\mathbb{k}[P]$ as the vanishing ideals of the subgroups classified in Lemma 7.2. For example, for $g = (1, v)$ with $v \in \mathbb{k}$ we find $J_v = \mathcal{I}(g\bar{Z}_0g^{-1}) = (\hat{x} - v(\hat{\lambda} - 1))$ and for any $r \in \mathbb{N}$ we have $F_r = \mathcal{I}(\bar{Z}_rU) = (\hat{\lambda}^r - 1)$. For the remaining subgroups, the respective Hopf ideals are no longer principal, since for $r \in \mathbb{N}$ $\mathcal{I}(g\bar{Z}_r g^{-1}) = \mathcal{I}(g\bar{Z}_0 g^{-1} \cap \bar{Z}_r U) = F_r + J_v = (\hat{\lambda}^r - 1, \hat{x} - v(\hat{\lambda} - 1))$. Note, finally, that the augmentation ideal is given by $\ker(\varepsilon) = F_1 + J_0 = J_v + J_\mu$ for $v \neq \mu$ and that $F_s + F_r = F_{\gcd(s,r)}$, providing a full description of the additive semigroup of Hopf ideals of $\mathbb{k}[P]$.

The discussion of $\mathbb{k}[P]$ now directly applies to quantum groups U_ζ or $U_\zeta^{\geq 0}$ at a root of unity ζ . For a simple root α_i , denote by $\mathcal{Z}_i^{\geq 0}$ the subalgebra generated by $L_i = K_i^{\ell_i}$ and $X_i = E_i^{\ell_i}$.

Corollary 7.3. *The algebra $\mathcal{Z}_i^{\geq 0}$ is a commutative Hopf subalgebra of $U_\zeta^{(\geq 0)}$.*

The semigroup of possible Hopf ideals of $\mathcal{Z}_i^{\geq 0} \otimes \mathbb{C}$ is given by the principal ideals $\ddot{F}_r = (L_i^r - 1)$ with $r \in \mathbb{N}$ and $\ddot{J}_v = (X_i - v(L_i - 1))$ with $v \in \mathbb{C}$ as well as the sums of ideals $\ddot{F}_r + \ddot{J}_v$.

Proof. Note first $X_i L_i = \zeta_i^{2\ell_i^2} L_i X_i = L_i X_i$ by definition of ζ_i and ℓ_i . So, $\mathcal{Z}_i^{\geq 0}$ is commutative, as also stated in Corollary 6.14. Clearly, L_i is group-like since K_i is. Moreover, we find from (3.17) and Corollary 6.2 the coproduct formula

$$\Delta(X_i) = \sum_{s=0}^{\ell_i} \begin{bmatrix} \ell_i \\ s \end{bmatrix}_i \zeta_i^{n(\ell_i-s)} E_i^s \otimes E_i^{\ell_i-s} K_i^s = E_i^{\ell_i} \otimes K_i^{\ell_i} + 1 \otimes E_i^{\ell_i} = X_i \otimes L_i + 1 \otimes X_i. \quad (7.6)$$

Thus, $\mathcal{Z}_i^{\geq 0} \otimes \mathbb{C} \cong \mathbb{C}[P]$ as Hopf algebras over \mathbb{C} . The claim now follows from the previously discussed classification of Hopf ideals of $\mathbb{C}[P]$. \square

Non-trivial Hopf ideals in $\mathcal{Z}_i^{\geq 0}$ for a general ground ring Λ containing Λ_ζ can be found as preimages of the extension of scalars map $\mathcal{Z}_i^{\geq 0} \rightarrow \mathcal{Z}_i^{\geq 0} \otimes \mathbb{C}$, provided that $v \in \Lambda$. All of these are readily seen to be free and complemented Λ -submodules of $\mathcal{Z}_i^{\geq 0}$.

The classification in Corollary 7.3 underscores the need to choose the minimal exponents $\ell_i = \ell / \gcd(d_i, \ell)$ in the definition of the X_i generators. Specifically, suppose α_i is a long root in the doubly laced Lie type case, meaning $d_i = 2$, and ℓ is even. Then $\ell = 2\ell_i$ so that $E_i^\ell = X_i^2$, generating an ideal $(E_i^\ell) = (X_i^2)$ that is not even radical and hence not Hopf. The latter is also obvious from the coproduct

$$\Delta(X_i^2) = X_i^2 \otimes L_i^2 + 2X_i \otimes L_i X_i + 1 \otimes X_i^2 \notin (X_i^2) \otimes U_\zeta + U_\zeta \otimes (X_i^2).$$

Since restricted groups are occasionally defined in the literature with such generators we add the above observation as a more formal remark, which has an obvious analogue for G_2 .

Remark 7.4. Assume $U_\zeta^{(\geq 0)}$ is of Lie type B_n , C_n , or F_4 with $n \geq 2$ as well as $\ell \equiv 0 \pmod{4}$. Then the ideal generated by $\{E_\alpha^\ell : \alpha \in R\}$ is **not** a Hopf ideal if $R \subseteq \Phi^+$ contains a long root. \diamond

Recall that any reductive algebraic subgroups over an algebraically closed field admits a Borel subgroup, unique up to conjugation. The fundamental example of such a Borel subgroup is the subgroup $T_n < \mathrm{GL}(n, \mathbb{k})$ of upper triangular matrices with non-zero diagonal entries. Denote also by D_m the subgroup of diagonal matrices and U_n the subgroup of upper triangular matrices with 1's on the diagonal so that $T_n = D_m \ltimes U_n$. As algebraic sets, $U_n \cong \mathbb{k}^{\binom{n}{2}}$ and $D_m \cong (\mathbb{k}^\times)^n$, where the latter is realized as either $\{(x_{1,1}, \dots, x_{n,n}, \tau) \in \mathbb{k}^n \times \mathbb{k} : x_{1,1} \dots x_{n,n} \tau = 1\}$ or, equivalently, as $(D_1)^n$. The former is commensurable with the embedding in $\mathrm{GL}(n, \mathbb{k})$, while the latter yields the coordinate rings as Laurent polynomials.

The coordinate ring is, thus, given as $\mathbb{k}[T_n] = \mathbb{k}[\{\hat{x}_{i,j} : i \leq j\}, \{\hat{x}_{i,i}^{-1}\}]$ so that the set of functions that are polynomials in off-diagonal entries and Laurent polynomials in the diagonal entries. As before, we notationally distinguish coordinates and coordinate functions. The group of group-like elements in $\mathbb{k}[T_n]$, corresponding to algebraic characters on T_n , are given

$$\mathrm{Spec}_m(T_n) \cong \mathbb{Z}^n \quad \text{with generators } \{\hat{x}_{i,i} : i = 1, \dots, n\}. \quad (7.7)$$

The counit is simply $\epsilon(\hat{x}_{i,j}) = \delta_{ij}$. The coproduct and antipode for the remaining generators $i < j$ are readily derived from (7.3) as

$$\Delta(\hat{x}_{i,j}) = \sum_{i \leq k \leq j} \hat{x}_{i,k} \otimes \hat{x}_{k,j} \quad \text{and} \quad S(\hat{x}_{i,j}) = \sum_{\substack{1 \leq m \leq j-i \\ i=k_0 < k_1 < \dots < k_m=j}} (-1)^m \hat{x}_{i,i}^{-1} \hat{x}_{i,k_1} \hat{x}_{k_1,k_1}^{-1} \hat{x}_{k_1,k_2} \dots \hat{x}_{k_{m-1},j} \hat{x}_{j,j}^{-1}. \quad (7.8)$$

We finally mention the notion of a *cosolvable* Hopf algebra H following [MW98]. In the case of a commutative Hopf algebra H , it starts from a finite ascending series of $\mathbb{k}1 = H_0 \subset H_1 \subset \dots \subset H_n = H$ Hopf subalgebras. From the augmentation ideals $H_s^\circ = H_s \cap \ker(\epsilon)$ one obtains for each s a Hopf ideal $H_{s-1}^\circ H_s$ in H_s and, thus, a Hopf algebra $\bar{H}_s = H_s / H_{s-1}^\circ H_s$, called the s -th factor of the series. H is said to be cosolvable if all of the factors are cocommutative for some such series.

In the case $H = \mathbb{k}[T_n]$, these properties can be easily verified if we define H_s as the polynomial subalgebra generated by $\{\hat{x}_{i,j} : j - i < s\}$ and 1. It is immediate from (7.8) that each H_s defines a Hopf subalgebra. Moreover, it is clear that $\bar{H}_1 = \mathbb{k}[D_m]$, generated by group-like elements. For $s > 1$, one readily verifies that $\bar{H}_s \cong \mathbb{k}[\{\bar{x}_{i,j} : i - j = s - 1\}]$ is a polynomial algebra, for which all generators are strictly primitive. Hence, each H_s is cocommutative and $\mathbb{k}[T_n]$ is cosolvable.

Recall, further, that by the Lie-Kolchin Theorem every closed, *connected*, and *solvable* linear algebraic group B is conjugate to a subgroup of T_n , see [Spr09, 6.3.1.]. The images of the H_s under restriction and conjugation map $\mathbb{k}[T_n] \rightarrow \mathbb{k}[B]$ give rise to a respective series with the same properties as the one for $\mathbb{k}[T_n]$.

Corollary 7.5. Suppose B is a closed, connected, solvable algebraic subgroup of $\mathrm{GL}(n, \mathbb{k})$ over an algebraically closed field \mathbb{k} . Then $\mathbb{k}[B]$ is cosolvable in the sense of [MW98].

In particular, the coordinate ring of $\mathbb{k}[\mathrm{GL}(n, \mathbb{k})]$ is fundamentally different from that of a (connected) Borel subgroup. The former is cosemisimple and thus not cosolvable and its character group in (7.4) also differs from that in (7.7).

7.2. Standard Presentation of $U_q^{\geq 0}$ for A_n . We provide next detailed relations and coproduct formulae for generators E_w of $U_q^{\geq 0}(\mathfrak{sl}_{n+1})$ with respect to a particularly convenient choice of a maximal word. It is given by the conjugate $z_n = w_{A_n}^\dagger$ of the one mentioned in Section 5.5 and inductively defined as

$$z_n = z_{n-1} c_{n,1} \quad \text{with} \quad c_{n,k} = w_n \dots w_{n+1-k} \quad \text{so that} \quad c_{n,k}^\dagger = w_n \dots w_{n+2-k} = c_{n,k-1}. \quad (7.9)$$

Observe that z_m is the word of maximal length in the canonically embedded A_m subsystem in A_n for $n > m$ with $l(z_m) = \binom{m+1}{2}$. Denote by $\bar{z}_n = \gamma(z_n)$ and $\bar{c}_{n,k} = \gamma(c_{n,k})$ the respective elements in the Weyl group \mathcal{W} . In the standard Euclidean realization of the A_n root system in $\mathbb{R}^{n+1} = \langle \epsilon_1, \dots, \epsilon_{n+1} \rangle$, simple roots are given as $\alpha_i = \epsilon_i - \epsilon_{i+1}$ and \mathcal{W} acts as the group of permutations in $n+1$ letters via $s(\epsilon_i) = \epsilon_{s(i)}$, with generators $s_i = (i, i+1)$. Specifically, $\bar{c}_{n,k} = (n+1, n, \dots, n+1-k)$ is a cyclic permutation and $\bar{z}_m(\epsilon_i) = \epsilon_{m+2-i}$ for $1 \leq i \leq m+1$ and $\bar{z}_m(\epsilon_i) = \epsilon_i$ for $i > m+1$.

Suppose $N = \binom{m}{2} + k$ for $1 \leq k \leq m$. We have then $z_n[1, N] = z_{m-1} c_{m,k}$ for the first N letters in z_n . The N -th term in the ordering sequence is now computed as

$$\begin{aligned} \beta_N &= \gamma(z_n[1, N]) = \gamma(z_{m-1} c_{m,k}) = \bar{z}_{m-1} \bar{c}_{m,k-1}(\alpha_{m+1-k}) \\ &= \bar{z}_{m-1}(m+1 \dots m+2-k)(\epsilon_{m+1-k} - \epsilon_{m+2-k}) \\ &= \bar{z}_{m-1}(\epsilon_{m+1-k} - \epsilon_{m+1}) = \epsilon_k - \epsilon_{m+1} = \alpha_k + \dots + \alpha_m. \end{aligned} \quad (7.10)$$

Hence, the ordering \prec_z defined in Section 2.3 is simply the lexicographic ordering. Namely, for $i < j$ and $r < s$ we have

$$\epsilon_i - \epsilon_j \prec_z \epsilon_r - \epsilon_s \quad \text{iff} \quad j < s \quad \text{or} \quad (j = s \text{ and } i < r). \quad (7.11)$$

Consider next for $1 \leq k \leq m$ the reduced words

$$b_{m,k} = w_k \dots w_m = c_{m,m+1-k}^* \quad \text{so that} \quad b_{m,k}^\dagger = w_k \dots w_{m-1} = b_{m-1,k}. \quad (7.12)$$

As before, we denote the respective element in the Weyl group as $\bar{b}_{m,k} = (k, \dots, m, m+1)$ and compute

$$\gamma(b_{m,k}) = \bar{b}_{m-1,k}(\alpha_m) = (k, \dots, m)(\epsilon_m - \epsilon_{m+1}) = \epsilon_k - \epsilon_{m+1}. \quad (7.13)$$

Note next that $\bar{b}_{m,k}$ is in the A_m subsystem and thus $\bar{b}_{m,k} \leq_R \bar{z}_m$. Thus, there exists some $\bar{u}_{m,k}$ in the Weyl group of the subsystem such that

$$\bar{z}_m = \bar{b}_{m,k} \cdot \bar{u}_{m,k} \quad \text{with} \quad l(\bar{z}_m) = l(\bar{b}_{m,k}) + l(\bar{u}_{m,k}). \quad (7.14)$$

In fact, it is easy to verify that a representative reduced word is given by $u_{m,k} = z_{m-1} \cdot c_{m,k-1}$. The calculations in (7.10) and (7.13) show that

$$\bar{b}_{m-1,k}(\alpha_m) = \bar{z}_{m-1} \bar{c}_{m,k-1}(\alpha_{m+1-k}) = \bar{b}_{m-1,k} \bar{u}_{m-1,k} \bar{c}_{m,k-1}(\alpha_{m+1-k}).$$

Together with length comparisons we find

$$\begin{aligned} \alpha_m &= \bar{u}_{m-1,k} \bar{c}_{m,k-1} (\alpha_{m+1-k}) \\ \text{and} \quad l(z_{m-1} c_{m,k-1}) &= l(b_{m-1,k}) + l(u_{m-1,k}) + l(c_{m,k-1}). \end{aligned} \tag{7.15}$$

For $1 \leq i < j \leq n+1$, denote now the generators of U_q^+ as follows.

$$E_{i,j} = E_{z_n[\epsilon_i - \epsilon_j]} = E_{z_n[1,N]} \quad \text{with } N = \binom{j-1}{2} + i \tag{7.16}$$

Proposition 7.6. *For a given $j \in \{2, \dots, n+1\}$, the elements in (7.16) are determined by the identity $E_{j-1,j} = E_{j-1}$ and the recursive relation*

$$E_{i,j} = \Gamma_i(E_{i+1,j}) = q^{-1} E_i E_{i+1,j} - E_{i+1,j} E_i \quad \text{for } i < j-1. \tag{7.17}$$

Proof. Set $m = j-1$ and $k = i$. We have $\gamma(z_n[1,N]) = \epsilon_i - \epsilon_j$ for $N = \binom{j-1}{2} + i$ by (7.10). Thus, by definition, $E_{z_n[1,N]} = \Gamma_{z_{j-2} c_{j-1,i-1}}(E_{j-i}) = \Gamma_{b_{j-2,i} u_{j-2,i} c_{j-1,i-1}}(E_{j-i}) = \Gamma_{b_{j-2,i}}(\Gamma_{u_{j-2,i} c_{j-1,i-1}}(E_{j-i}))$ since the Γ only depend on the Weyl element and $l(b_{j-2,i} u_{j-2,i} c_{j-1,i-1}) = l(b_{j-2,i}) + l(u_{j-2,i} c_{j-1,i-1})$ by (7.15).

Note, further, that by (7.15) we have that the weight of $E_{u_{j-2,i} c_{j-1,i}} = \Gamma_{u_{j-2,i} c_{j-1,i-1}}(E_{j-i})$ is α_{j-1} . It, therefore, follows from Proposition 5.1 that $E_{u_{j-2,i} c_{j-1,i}} = E_{j-1}$. Plugging this into the previous expression for $E_{z_n[1,N]}$ we find

$$E_{i,j} = \Gamma_{b_{j-2,i}}(E_{j-1}) = \Gamma_i \dots \Gamma_{j-2}(E_{j-1}) = \Gamma_i(E_{i+1,j}). \tag{7.18}$$

In particular, $E_{j-1,j} = E_{j-1}$. The recursion in (7.17) follows by induction, decreasing in i . Assuming the identity for $i+1$, one can express $E_{i+1,j}$ by E_{i+1} and $E_{i+2,j}$. Since the latter is in the subalgebra $\langle E_{i+2}, \dots, E_{j-1} \rangle$ generated by $\{E_{i+2}, \dots, E_{j-1}\}$, Γ_i acts as identity on $E_{i+2,j}$ and E_i commutes with $E_{i+2,j}$. The resulting expression in E_i , E_{i+1} , and $E_{i+2,j}$ can now be worked out and identified with the left side of (7.17). \square

A full set of commutation relations as well as nested subalgebras of \mathbf{A} -types are derived in the next proposition. Most of these can be found in other references, with the possible exception of the relation in Item *iii*) below.

Proposition 7.7. *For $1 \leq i < s < j < t \leq n+1$ the following commutation relations hold.*

$$\begin{array}{ll} i) \quad E_{i,t} = q^{-1} E_{i,j} E_{j,t} - E_{j,t} E_{i,j} & iv) \quad E_{i,s} E_{j,t} = E_{j,t} E_{i,s} \\ ii) \quad E_{i,t} E_{s,j} = E_{s,j} E_{i,t} & v) \quad E_{i,t} E_{i,j} = q E_{i,j} E_{i,t} \\ iii) \quad E_{i,j} E_{s,t} - E_{s,t} E_{i,j} = -(q - q^{-1}) E_{s,j} E_{i,t} & vi) \quad E_{j,t} E_{i,t} = q E_{i,t} E_{j,t} \end{array}$$

More generally, suppose $\hat{j} = (j_1, \dots, j_{m+1})$ is a sequence with $1 \leq j_1 < j_2 < \dots < j_{m+1} \leq n+1$. Then there exists an injective algebra homomorphism $\phi_{\hat{j}} : U_q^+(\mathfrak{sl}_{m+1}) \rightarrow U_q^+(\mathfrak{sl}_{n+1})$ with

$$\phi_{\hat{j}}(E_{r,s}) = E_{j_r, j_s}. \tag{7.19}$$

Proof. The relation in *i*) is obtained by applying $\Gamma_{b_{i-1,k}} = \Gamma_k \dots \Gamma_{i-1}$ to (7.17), using that $\Gamma_{b_{i-1,k}}$ acts trivially on $E_{i+1,j} \in \langle E_{i+1}, \dots, E_{j-1} \rangle$, and substituting indices. Item *iv*) is clear, since $E_{i,s} \in \langle E_i, \dots, E_{s-1} \rangle$ and $E_{j,t} \in \langle E_j, \dots, E_{t-1} \rangle$ are in algebras for which the generators of one commute with the generators of the other.

Next note that, by (7.18), we can write $E_{i,j} = E_{w_{ij}}$, where $w_{ij} = w_i \dots w_{j-1}$. Clearly, $w_{it} = w_{ij}w_{jt}$ with $l(w_{it}) = l(w_{ij}) + l(w_{jt})$. Further, the index of $\tau(w_{ij}) = j-1$ does not occur w_{jt} . We also have $\alpha = \gamma(w_{ij}) = \epsilon_i - \epsilon_j$ and $\beta = \gamma(w_{it}) = \epsilon_i - \epsilon_t$, so that $(\alpha|\beta) = 1$. The relation in *v*) follows now directly from Lemma 5.6, with $w = w_{it}$, $u = w_{ij}$, and $v = w_{jt}$.

Alternatively, we find from (7.10) and (7.16) that $E_{it} = E_{z_{t-2}c_{t-1,i}}$. From (7.9) we have $c_{t-1,j} = c_{t-1,i}v$, where $v = w_{t-1-i} \dots w_{t-j}$. So, if we set $w = z_{t-2}c_{t-1,j}$ and $u = z_{t-2}c_{t-1,i}$ we find $w = uv$ with $l(w) = l(u) + l(v)$. As before $\tau(u) = t-j$ does not occur as an index in v . Moreover, $\alpha = \gamma(u) = \epsilon_i - \epsilon_t$ and $\beta = \gamma(w) = \epsilon_j - \epsilon_t$, so that $(\alpha|\beta) = 1$. Lemma 5.6, therefore, implies the relation in Item *vi*).

Combining relation *i*) with *v*) as well as combining relation *i*) with *vi*) we obtain respective Serre relations

$$E_{ij}^2 E_{jt} - [2]E_{ij}E_{jt}E_{ij} + E_{jt}E_{ij}^2 = 0 \quad \text{and} \quad E_{jt}^2 E_{ij} - [2]E_{jt}E_{ij}E_{jt} + E_{ij}E_{jt}^2 = 0.$$

Together with the relation in *v*) we observe that, for a sequence $j_1 < \dots < j_{m+1}$ as above, the elements $\tilde{E}_r = E_{j_r, j_{r+1}}$ fulfill the relations of the standard generators of $U_q^+(\mathfrak{sl}_{m+1})$. Hence, we have a well-defined algebra homomorphism $\phi_{\hat{j}}$ that maps the E_r generator in $U_q^+(\mathfrak{sl}_{m+1})$ to \tilde{E}_r in $U_q^+(\mathfrak{sl}_{n+1})$. Setting $(i, j, t) = (j_k, j_{k+1}, j_m)$ and denoting $\tilde{E}_{k,m} = E_{j_k, j_m}$ relation *v*) can be written as $\tilde{E}_{k,m} = q^{-1}\tilde{E}_k\tilde{E}_{k+1,m} - \tilde{E}_{k+1,m}\tilde{E}_k$. Since this is the same recursion (7.17) for the generators in $U_q^+(\mathfrak{sl}_m)$ and $\phi_{\hat{j}}$ is a homomorphism, (7.19) follows inductively. This also means that the PBW basis elements given by ordered monomials in the $E_{i,j}$ in $U_q^+(\mathfrak{sl}_{m+1})$ are mapped to elements of standard PBW basis of $U_q^+(\mathfrak{sl}_{n+1})$, which, in turn, implies that $\phi_{\hat{j}}$ is injective.

Given the $A_m \hookrightarrow A_n$ inclusions, it now suffices for the remaining relations to consider the A_3 case, using $\phi_{\hat{j}}$ with $\hat{j} = (j_1, j_2, j_3, j_4) = (i, s, j, t)$. The relation in *ii*) is then implied by $E_{1,4}E_2 = E_2E_{1,4}$. To see this, observe $E_{1,4} = \Gamma_1\Gamma_2(E_3)$ and $E_2 = \Gamma_1\Gamma_2(E_1)$ as in Proposition 5.1. Thus, $[E_{1,4}, E_2] = [\Gamma_1\Gamma_2(E_3), \Gamma_1\Gamma_2(E_1)] = \Gamma_1\Gamma_2([E_1, E_3]) = 0$ as desired.

To derive the last relation *iii*), we set $\theta_2 = \Gamma_2^{-1}(E_2) = -F_2K_2$ and compute from the relations in Section 3.2 the following equations in the full quantum group U_q .

$$\theta_2 E_i = q^{-1} E_i \theta_2 \quad \text{for } i \in \{1, 3\} \quad \text{and} \quad \theta_2 E_2 = \frac{q^2}{q-q^{-1}} (K_2^2 - 1) + q^2 E_2 \theta_2$$

Letting $c = \frac{q^2}{q-q^{-1}}$ and $i, j \in \{1, 3\}$, these entail commutators

$$\begin{aligned} [\theta_2, E_i E_2 E_j] &= c(qK_2^2 - q^{-1})E_i E_j, & [\theta_2, E_2 E_i E_j] &= c(K_2^2 - 1)E_i E_j, \\ [\theta_2, E_i E_j E_2] &= c(q^2 K_2^2 - q^{-2})E_i E_j. \end{aligned} \tag{7.20}$$

Using the braid relations for the Γ_i as well as $\Gamma_2\Gamma_3(E_2) = E_3$ we, further, compute

$$\begin{aligned}\Gamma_2^{-1}\Gamma_1^{-1}\Gamma_2(E_3) &= \Gamma_1\Gamma_2^{-1}\Gamma_1^{-1}(E_3) = \Gamma_1\Gamma_2^{-1}(E_3) = \Gamma_1\Gamma_3(E_2) \\ &= q^{-2}E_3E_1E_2 - q^{-1}(E_3E_2E_1 + E_1E_2E_3) + E_2E_1E_3.\end{aligned}\tag{7.21}$$

Combining (7.20) and (7.21) we arrive at

$$[\Gamma_2^{-1}(E_2), \Gamma_2^{-1}\Gamma_1^{-1}\Gamma_2(E_3)] = -(q - q^{-1})E_1E_3,$$

which yields, after application of $\Gamma_1\Gamma_2$ and using $\Gamma_1\Gamma_2(E_1) = E_2 = E_{2,3}$, the desired form

$$[E_{1,3}, E_{2,4}] = [\Gamma_1(E_2), \Gamma_2(E_3)] = -(q - q^{-1})\Gamma_1\Gamma_2(E_1)\Gamma_1\Gamma_2(E_3) = -(q - q^{-1})E_{2,3}E_{1,4}.$$

□

To express the coproducts of these elements we introduce notations $K_{i,j} = K_i \dots K_{j-1}$ and $E_{i,i} = E_{j,j} = 1$.

Lemma 7.8. *For $i < j$, the coproduct of $E_{i,j}$ is*

$$\Delta(E_{i,j}) = \sum_{k=i}^j c_k E_{i,k} \otimes K_{i,k} E_{k,j} \tag{7.22}$$

where $c_i = c_j = 1$ and $c_k = -(q - q^{-1})$ for $i < k < j$.

Proof. The formula is immediate for the coproduct of E_{j-1} when $i = j - 1$. The assertion for $i < j - 1$ is then obtained by descending induction in i , using the recursion (7.17). For $T_k^{i,j} = E_{i,k} \otimes K_{i,k} E_{k,j}$ a basic calculation yields

$$q^{-1}\Delta(E_i)T_k^{i+1,j} - T_k^{i+1,j}\Delta(E_i) = \begin{cases} T_k^{i,j} & \text{for } k > i + 1 \\ -(q - q^{-1})T_{i+1}^{i,j} + T_i^{i,j} & \text{for } k = i + 1. \end{cases}$$

Assuming (7.22) for $\Delta(E_{i+1,j})$ we then use this computation to derive (7.22) for the coproduct as $\Delta(E_{i,j}) = q^{-1}\Delta(E_i)\Delta(E_{i+1,j}) - \Delta(E_{i+1,j})\Delta(E_i)$. □

The coproduct for powers of these generators is derived from commutation relations for the elements $T_k^{i,j}$ defined in the proof above. First, observe that the identities *v*) and *vi*) in Proposition 7.7 imply that

$$T_r^{i,j}T_s^{i,j} = q^{-2}T_s^{i,j}T_r^{i,j} \quad \text{for all } i \leq r < s \leq j. \tag{7.23}$$

Thus, if $D_m = \sum_{k=i}^m c_k T_k^{i,j}$ we find from (7.23) that also $D_{m-1}T_m^{i,j} = q^{-2}T_m^{i,j}D_{m-1}$. We may now write $D_m = V + U$ where $V = D_{m-1}$ and $U = c_m T_m^{i,j}$. Applying (3.3), this allows us to infer $D_m^N = \sum_r q^{r(N-r)} \begin{bmatrix} N \\ r \end{bmatrix} D_{m-1}^{N-r} (c_m T_m^{i,j})^r$. Iterating this formula, we arrive at the following

expression,

$$\Delta(E_{i,j}^N) = \sum_{\substack{r_i, \dots, r_j \geq 0 \\ r_i + \dots + r_j = N}} q^{\sum_{i \leq s < t \leq j} r_s r_t} (q^{-1} - q)^{\sum_{i < s < j} r_s} \begin{bmatrix} N \\ r_i, \dots, r_j \end{bmatrix} (T_i^{i,j})^{r_i} \dots (T_j^{i,j})^{r_j}. \quad (7.24)$$

7.3. Identification of $\mathcal{Z}_\bullet^{\geq 0}$ as a (Semi) Classical Matrix Algebra. We consider next the specialization to a primitive ℓ -th root of unity ζ , with ℓ the order of ζ^2 . In the simply laced setting all $\ell_\alpha = \ell$. Thus, for the root $\alpha = \alpha_i + \dots + \alpha_{j-1} = \epsilon_i - \epsilon_j$, we have $X_\alpha = X_{i,j} = E_{i,j}^\ell$ with respect to the ordering given by (7.9). We extend this notation to $X_{i,i} = 1$. Write also $L_{i,j} = L^\alpha = K_{i,j}^\ell$ as in (6.16) and define $\mathcal{Z}_\bullet^{\geq 0} = \mathcal{Z}_{s_\bullet}^{\geq 0}$ as in (6.39) and (6.56).

The coproducts of the X_α have been computed in [BW04, (2.24)] for a 2-parameter version $U_{r,s}(\mathfrak{sl}_n)$, which coincides with U_ζ if $r = s^{-1}$ is a respective root of unity. The results there imply that the collection of all X_α generate a Hopf ideals, leading to a well-defined (2-parameter) restricted quantum group. A general form for the coproduct for other cases can also be found in [DCKP92, Sect. 5.5], though the stated formulae are modulo higher ideal terms and no explicit coefficients are offered. A similar version of the next lemma can be found in [BW04].

Lemma 7.9. *For $1 \leq i < j \leq n + 1$ and with notation as above, we have*

$$\Delta(X_{i,j}) = \sum_{k=i}^j b_k X_{i,k} \otimes L_{i,k} X_{k,j}, \quad (7.25)$$

where $b_k = (\zeta^{-1} - \zeta)^\ell \zeta^{\binom{\ell}{2}}$ for $i < k < j$ and $b_i = b_j = 1$.

In particular, $\mathcal{Z}_\bullet^{\geq 0}$ is a Hopf subalgebra of $U_\zeta^{\geq 0}$.

Proof. Setting $N = \ell$ in (7.24) and with Corollary 6.2 we are left with a summation over terms in which one index $r_k = \ell$ and all other indices are zero. For $i < k < j$ we, further, note that $(T_k^{i,j})^\ell = (E_{i,k})^\ell \otimes (K_{i,k} E_{k,j})^\ell = \zeta^{\binom{\ell}{2}} E_{i,k}^\ell \otimes K_{i,k}^\ell E_{k,j}^\ell$ since $K_{i,k} E_{k,j} = \zeta^{-1} E_{k,j} K_{i,k}$. The resulting formula for the coproduct in (7.25) as well as $\Delta(L_{i,j}) = L_{i,j} \otimes L_{i,j}$ contain only generators of $\mathcal{Z}_\bullet^{\geq 0}$, so that $\mathcal{Z}_\bullet^{\geq 0}$ is a sub-bialgebra.

As noted in the proof of Theorem 6.19 in Section 6.4, $\mathcal{Z}_\bullet^{\geq 0}$ is also invariant under \mathfrak{U} . From Item *iv*) in Proposition 4.1 we also know that $\Psi_u^{id}(X_{i,j}) = \lambda K^{\ell(\epsilon_i - \epsilon_j)} X_{i,j} = \lambda L_{i,j} X_{i,j}$, where $\lambda = \pm \zeta^a$ for some $a \in \mathbb{Z}$. Relation (4.16) now implies that $\mathcal{Z}_\bullet^{\geq 0}$ is also invariant under the antipode S . \square

A normalization of generators for which the coproduct structure is defined over \mathbb{Z} is discussed in Appendix D. The fact that the full skew-central subalgebra forms a Hopf subalgebra allows us now to prove that the ideals defined in Section 6.5 are indeed Hopf ideals.

Theorem 7.10. *For any $s \in \mathcal{W}$ the augmentation ideal $\mathcal{K}(s)^{\geq 0}$ as in (6.63) is a Hopf algebra ideal in $\mathcal{Z}_\bullet^{\geq 0}$.*

Proof. Since $s \leq_R s_\bullet$ there are reduced words $w, b \in \mathcal{W}^*$ with $z = w \cdot b \in \mathcal{W}_m^*$, $s_\bullet = \tau(z)$, and $s = \tau(w)$. As in Section 2.3, this imposes a total convex ordering \prec_z on Φ^+ . As in the remarks

following (6.63), word independence of Theorem 6.19 implies that $\mathcal{K}(s)^{\geq 0}$ is the ideal generated by all X_v with $\emptyset \neq v \leq_R w$. Given the fixed $z \in \mathcal{W}_m^*$, we may relabel generators $X_\alpha = X_{z[\alpha]}$ analogous to (5.3). Thus, with $\gamma = \gamma(w)$ and this notation, $\mathcal{K}(s)^{\geq 0}$ is the ideal generated by X_β with $\beta \leq_z \gamma$.

Next, note that, by Proposition 6.10, the monomials $L^\mu X_{z\bullet}^\psi = \pm L^\mu \prod_{\alpha \in \Phi^+} X_\alpha^{\psi(\alpha)}$ are a basis for $\mathcal{Z}_\bullet^{\geq 0}$. Since, by Lemma 7.9, $\mathcal{K}_\bullet^{\geq 0}$ is a Hopf algebra over Λ , we can write $\Delta(X_\beta)$ with $\beta \in \Phi^+$ as a Λ -linear combination of terms of the form $L^\nu X_{\beta_1} \dots X_{\beta_s} \otimes L^\mu X_{\beta_{s+1}} \dots X_{\beta_k}$, where the sequence of roots $\beta_i \in \Phi^+$ may have repetitions.

By w -grading, we must have $\beta = \sum_i \beta_i$. Lemma 2.6 now implies that for at least one j we have $\beta_j \leq_z \beta$. Thus, if $\beta \leq_z \gamma$ also $\beta_j \leq_z \gamma$ and, hence, that $X_{\beta_j} \in \mathcal{K}(s)^{\geq 0}$. Therefore, any of these terms is in $\mathcal{K}(s)^{\geq 0} \otimes \mathcal{Z}_\bullet^{\geq 0} + \mathcal{Z}_\bullet^{\geq 0} \otimes \mathcal{K}(s)^{\geq 0}$, which, thus, contains $\Delta(X_\beta)$ for every generator X_β of $\mathcal{K}(s)^{\geq 0}$ (with $\beta \leq_z \gamma$) and, consequently, also $\Delta(\mathcal{K}(s)^{\geq 0})$. \square

The proof of Theorem 7.10 extends to other classical Lie types for which $\mathcal{Z}_\bullet^{\geq 0}$ can be shown to be a Hopf subalgebra, as well as other Lie types to which Lemma 2.6 can be generalized.

The $\mathcal{Z}_w^{\geq 0}$ subalgebras, however, will generally *not* form sub-bialgebras. For example, for type A_2 and $w = w_1 w_2 = z_2[1, 2]$ we have that \mathcal{Z}_w^+ is the subalgebra generated by $X_{1,3}$ and $X_{2,3}$. Yet, $\Delta(X_{1,3})$ will also contain a $X_{1,2}$ factor.

Conversely, call a subset $C \subseteq \Phi^+$ a *lower set* if $\mu + \nu \in C$ for $\mu, \nu \in \Phi^+$ implies $\mu, \nu \in C$. The coproduct in (7.25) implies that $\langle X_\alpha, L_\alpha : \alpha \in C \rangle$ is a Hopf subalgebra for any lower set C , where the generators are defined with respect to the maximal word from (7.9). For a given root $\beta = \epsilon_i - \epsilon_j = \alpha_i + \dots + \alpha_{j-1}$, let C_β be the set of all positive roots $\leq \beta$ in the usual partial order, which is just the root subsystem $\{\epsilon_s - \epsilon_r : i \leq s < r \leq j\}$. It is clear that a subset C has the above property if and only if it is a union of (possibly intersecting) sets C_β .

One may, further, ask for which $s \in \mathcal{W}$, both, the algebra $\mathcal{Z}_s^{\geq 0}$ is a Hopf subalgebra so that $\mathcal{K}(s)^{\geq 0}$ is a Hopf ideal. In the A_n case with $\mathcal{W} = S_{n+1}$ it is not difficult to see that the root set $\mathcal{N}(s)$ for $\mathcal{Z}_s^{\geq 0}$ is a lower set if it is the *disjoint* union of sets C_β . Particularly, the lower set property for $\mathcal{N}(s)$ implies that if (i, j) with $i < j$ is an inversion pair for $s \in S_{n+1}$ then any (a, b) with $i \leq a < b \leq j$ also needs to be an inversion pair. Suppose that $[i, j]$ is a maximal interval with $\epsilon_i - \epsilon_j \in \mathcal{N}(s)$. It is then not hard to see that s then maps the intervals $[1, i-1]$, $[i, j]$, and $[j+1, n]$ respectively to themselves and that it acts on $[i, j]$ via the longest (full inversion) element of S_{j-1+1} .

Thus, by induction, s must be the product of longest elements on disjoint intervals $a_s = [i_s, j_s]$, with $j_s < i_{s+1}$. The associated subalgebra $\mathcal{Z}_s^{\geq 0}$ is then identified with $\mathcal{Z}_\bullet^{\geq 0}(a_1) \otimes \dots \otimes \mathcal{Z}_\bullet^{\geq 0}(a_m)$, where each $\mathcal{Z}_\bullet^{\geq 0}(a_s)$ is the subalgebra generated X_α and L_α with α in the respective $A_{j_s-i_s+1}$ subsystem. Requiring the augmentation ideals $\mathcal{K}(s)^{\geq 0}$ to derive from Hopf algebras, thus, leaves us with only these obvious examples.

In the remainder of this section, we discuss identifications of the Hopf algebra $\mathcal{Z}_\bullet^{\geq 0}$ as in Lemma 7.9 with (semi) classical Hopf algebra constructions that are (mostly) independent of a choice of roots of unity ζ . These are more conveniently worked out for the quantum group associated to \mathfrak{gl}_{n+1} , which contains the standard one for \mathfrak{sl}_{n+1} .

The generators of $U_q(\mathfrak{gl}_{n+1})$ are given by the usual E_i and F_i for $1 \leq i \leq n$ as well as $P_i^{\pm 1}$ for $1 \leq i \leq n+1$. The elements $P_i^{\pm 1}$ are required to be group like and to commute with each other. Relations (3.13) and (3.16) are replaced by ones in which we substitute K_i by $P_i P_{i+1}^{-1}$. The conjugation relations in (3.14) are replaced by

$$P_i E_j P_i^{-1} = q^{(\epsilon_i | \alpha_j)} E_j = q^{\delta_{i,j} - \delta_{i,j+1}} E_j \quad \text{and} \quad P_i F_j P_i^{-1} = q^{-(\epsilon_i | \alpha_j)} F_j = q^{-\delta_{i,j} + \delta_{i,j+1}} F_j, \quad (7.26)$$

and the Serre relations (3.11) and (3.12) remain unchanged.

A \mathbb{Z} -grading \mathfrak{p} on $U_q(\mathfrak{gl}_{n+1})$ is provided by setting $\mathfrak{p}(P_i) = 1$ and $\mathfrak{p}(E_i) = \mathfrak{p}(F_i) = 0$ for all $i = 1, \dots, n$. The original algebra $U_q(\mathfrak{sl}_{n+1})$ is then recovered as the 0-graded subalgebra of $U_q(\mathfrak{gl}_{n+1})$ with respect to \mathfrak{p} . The inclusion map of $U_q(\mathfrak{sl}_{n+1})$ into $U_q(\mathfrak{gl}_{n+1})$ is now given by $K_i \mapsto P_i P_{i+1}^{-1}$ on the Cartan subalgebra and identity on all other generators.

The specialization of $U_q(\mathfrak{gl}_{n+1})$ to a root of unity ζ is commensurable with this inclusion. The analogous skew-central subalgebra $\tilde{\mathcal{Z}}_\bullet^{\geq 0}$ is generated by $Q_i = P_i^\ell$ as well as the $X_{i,j}$ as before. The \mathfrak{p} -grading extends to $\tilde{\mathcal{Z}}_\bullet^{\geq 0}$ with $\mathfrak{p}(Q_i) = \ell$.

The ground ring Λ may be any extension of $\Lambda_\zeta = \mathbb{Z}[\zeta]$, such as the ones from (6.5). If $(\zeta - \zeta^{-1})^{-1}$ is not in Λ , we denote by $\tilde{\mathcal{Z}}_\bullet^{\geq 0}$ the subalgebra generated by $Q_i = P_i^\ell$ and $(\zeta - \zeta^{-1})^\ell X_{i,j}$. Thus, for example for $\Lambda = \Lambda_\zeta$, the quotient $\tilde{\mathcal{Z}}_\bullet^{\geq 0}/\tilde{\mathcal{Z}}_\bullet^{\geq 0}$ is a torsion module for Λ . Conversely, if the ground ring extends Λ_ζ we have $\tilde{\mathcal{Z}}_\bullet^{\geq 0} = \tilde{\mathcal{Z}}_\bullet^{\geq 0}$.

As for the generic case, the original subalgebra $\mathcal{Z}_\bullet^{\geq 0}$ is identified with the zero-graded component of $\tilde{\mathcal{Z}}_\bullet^{\geq 0}$ with respect to \mathfrak{p} . Analogously, we write $\mathcal{Z}_\bullet^{\geq 0}$ for the subalgebra generated by the L_j and $(\zeta - \zeta^{-1})^\ell X_{i,j}$, or, equivalently, the zero-graded part of $\tilde{\mathcal{Z}}_\bullet^{\geq 0}$.

We next consider a smaller variant of the extended quantum matrix algebra introduced by Artin, Schelter, and Tate in [AST91]. Suppose $c : \{1, \dots, n\}^2 \rightarrow \Lambda^\times$ are coefficients for which $c_{(i,j)} = c_{(j,i)}^{-1}$ and $c_{(i,i)} = 1$. Define $\mathcal{O}(T_n, \Lambda, c)$ to be the unital algebra over Λ with generators $R_{i,j}$ and relations

$$R_{i,j} R_{s,t} = \frac{c_{(i,s)}}{c_{(j,t)}} R_{s,t} R_{i,j} \quad \text{and} \quad R_{i,i}^{-1} R_{i,i} = 1 = R_{i,i} R_{i,i}^{-1} \quad (7.27)$$

for all valid indices. This algebra differs from the one in [AST91] in that we only use the coefficients on the subgroup T_n of upper triangular matrices in $GL(n, \mathbb{C})$, and that an extra parameter there is $\lambda = 1$. The algebra $\mathcal{O}(T_n, \Lambda, c)$ may also be viewed as a central extension of the polynomial algebra $\Lambda[\{x_{i,j}, x_{i,i}^{-1}\}_{i \leq j}]$, in which the c expression features as the respective 2-cocycle.

The bialgebra portion of the following result is a straightforward adaption of Theorems 1 and 2 in [AST91]. As opposed to the situation in [AST91], an antipode is assured by the invertibility

of the diagonal elements and does not require a determinant relation. Its explicit form below is readily verified by direct computation.

Proposition 7.11. *The algebra $\mathcal{O}(T_n, \Lambda, c)$ is a free Λ module with basis given by monomials $\prod_{i \leq j} R_{i,j}^{n_{i,j}}$, where $n_{i,j} \in \mathbb{Z}$ and $n_{i,j} \in \mathbb{N}_0$ for $1 \leq i < j \leq n$ (for some fixed ordering).*

Further, $\mathcal{O}(T_n, c, \Lambda)$ admits the structure of a Hopf algebra over Λ with coproduct defined by

$$\Delta(R_{i,j}) = \sum_{k=i}^j R_{i,k} \otimes R_{k,j}. \quad (7.28)$$

The counit is $\varepsilon(R_{i,j}) = \delta_{i,j}$. The unique antipode is given by $S(R_{i,i}) = R_{i,i}^{-1}$ and, for $j > i$, by

$$S(R_{i,j}) = \sum_{\substack{1 \leq m \leq j-i \\ i=s_0 < s_1 < \dots < s_m=j}} (-1)^m R_{s_0,s_0}^{-1} R_{s_0,s_1} R_{s_1,s_1}^{-1} R_{s_1,s_2} \dots R_{s_{m-1},s_{m-1}}^{-1} R_{s_{m-1},s_m} R_{s_m,s_m}^{-1}.$$

Two special cases of fully symmetric and anti symmetric coefficient sets are given by

$$s(i,j) = 1 \quad \text{and} \quad a(i,j) = -(-1)^{\delta_{ij}}. \quad (7.29)$$

We will also abbreviate $\mathcal{O}(T_n, \Lambda) = \mathcal{O}(T_n, s, \Lambda)$ for the classical matrix algebra. Note also, that the coefficients in (7.29) are units in \mathbb{Z} and the structure coefficients in Proposition 7.11 are integers as well. Thus, the matrix algebras can be defined over \mathbb{Z} so that

$$\mathcal{O}(T_n, \Lambda) = \mathcal{O}(T_n, \mathbb{Z}) \otimes \Lambda \quad \text{and} \quad \mathcal{O}(T_n, a, \Lambda) = \mathcal{O}(T_n, a, \mathbb{Z}) \otimes \Lambda. \quad (7.30)$$

The algebra $\mathcal{O}(T_n, c, \Lambda)$ admits many integer gradings. One of them, given on generators as $\mathfrak{n}(R_{i,j}) = \epsilon_j - \epsilon_i$, extends to a Hopf algebra grading. Another is the total polynomial degree, given by $\mathfrak{q}(R_{i,j}) = 1$ and $\mathfrak{q}(R_{i,i}^{-1}) = -1$, which does not extend to a Hopf algebra grading. Nevertheless, it is not hard to see that the zero graded subalgebra, spanned by monomials M with $\mathfrak{q}(M) = 0$, forms indeed a Hopf subalgebra, which we denote as $\mathcal{O}_o(T_n, c, \Lambda)$. Similarly, we abbreviate $\mathcal{O}_o(T_n, \Lambda) = \mathcal{O}_o(T_n, s, \Lambda)$.

The following identifications of the skew-central subalgebras assume that Λ contains Λ_ζ , where ζ is a primitive ℓ -th root of unity.

Theorem 7.12. *Let $\tilde{\mathcal{Z}}_{\bullet, \bullet}^{\geq 0}$ and $\mathcal{Z}_{\bullet, \bullet}^{\geq 0}$ be the subalgebra of $U_\zeta(\mathfrak{gl}_n)$ and $U_\zeta(\mathfrak{sl}_n)$ as defined above. There exist isomorphisms of Hopf algebras over Λ as follows.*

$$\begin{aligned} \tilde{\mathcal{Z}}_{\bullet, \bullet}^{\geq 0} &\cong \mathcal{O}(T_n, \Lambda) & \text{and} & \quad \mathcal{Z}_{\bullet, \bullet}^{\geq 0} \cong \mathcal{O}_o(T_n, \Lambda) & \quad \text{if } \ell \not\equiv 2 \pmod{4} \\ \tilde{\mathcal{Z}}_{\bullet, \bullet}^{\geq 0} &\cong \mathcal{O}(T_n, a, \Lambda) & \text{and} & \quad \mathcal{Z}_{\bullet, \bullet}^{\geq 0} \cong \mathcal{O}_o(T_n, a, \Lambda) & \quad \text{if } \ell \equiv 2 \pmod{4} \end{aligned} \quad (7.31)$$

Proof. Set $b = (\zeta^{-1} - \zeta)^\ell \zeta^{\binom{\ell}{2}}$ and consider the assignments of generators of $\mathcal{O}(T_n, c, \Lambda)$ to generators of $\tilde{\mathcal{Z}}_{\bullet, \bullet}^{\geq 0}$ given by

$$R_{i,j} \mapsto b Q_i^{-1} X_{i,j} \quad \text{for } i < j \quad \text{and} \quad R_{i,i} \mapsto Q_i^{-1}. \quad (7.32)$$

Using $Q_i Q_j^{-1} = P_i^\ell P_j^{-\ell} = K_{i,j}^\ell = L_{i,j}$ and the coproduct formula in (7.25) we find

$$\begin{aligned}\Delta(bQ_i^{-1}X_{i,j}) &= b(Q_i^{-1} \otimes Q_i^{-1}) \left(1 \otimes Q_i Q_i^{-1} X_{i,j} + \sum_{i < k < j} bX_{i,k} \otimes Q_i Q_k^{-1} X_{k,j} + X_{i,j} \otimes Q_i Q_j^{-1} \right) \\ &= Q_i^{-1} \otimes bQ_i^{-1} X_{i,j} + \sum_{i < k < j} bQ_i^{-1} X_{i,k} \otimes bQ_k^{-1} X_{k,j} + bQ_i^{-1} X_{i,j} \otimes Q_j^{-1}.\end{aligned}$$

That is, the assignment preserves the corelation of generators from (7.28).

The commutation relations 6.26 specialize to $X_\alpha X_\beta = \xi^{(\alpha|\beta)} X_\beta X_\alpha$ in the simply laced case. Here $\xi = \zeta^{\ell^2}$, as in (6.2), is $\xi = -1$ if $\ell \equiv 2 \pmod{4}$ and $\xi = +1$ otherwise. From (7.26) we also have $Q_i X_\alpha = \xi^{(\epsilon_i|\alpha)} X_\alpha Q_i$. Thus, $Q_i^{-1} X_\alpha Q_s^{-1} X_\beta = \xi^{(\alpha|\epsilon_s) + (\epsilon_i|\beta) + (\alpha|\beta)} Q_s^{-1} X_\beta Q_i^{-1} X_\alpha$. Setting $\alpha = \epsilon_i - \epsilon_j$ and $\beta = \epsilon_s - \epsilon_t$ with $i \leq j$ and $s \leq t$ we find the exponent is $\delta_{i,s} + \delta_{j,t} = (1 + \delta_{i,s}) + (1 + \delta_{j,t}) \pmod{2}$.

Thus $(Q_i^{-1} X_{i,j})(Q_s^{-1} X_{s,t}) = \frac{c(i,s)}{c(j,t)} (Q_s^{-1} X_{s,t})(Q_i^{-1} X_{i,j})$, where $c(i,j) = \xi^{1+\delta_{i,j}}$. Clearly, for $\xi = 1$ this matches the commutation relations in $\mathcal{O}(T_n, \mathfrak{s}, \Lambda)$ and for $\xi = -1$ the ones for $\mathcal{O}(T_n, \mathfrak{a}, \Lambda)$. Consequently, the assignment in (7.32) extends to an algebra homomorphism, with the previous observation, also to a bialgebra homomorphism, and, using uniqueness of antipodes, to a Hopf algebra homomorphism.

It is clear that this homomorphism maps PBW bases to each other, yielding an isomorphism of Hopf algebras and, thus, proving the statements for the $U_\zeta(\mathfrak{gl}_n)$ subalgebras. The respective statements for the $U_\zeta(\mathfrak{sl}_n)$ are inferred from the fact that the assignment in (7.32) maps a generator with \mathfrak{q} -grading of 1 to an element with \mathfrak{p} -grading $-\ell$. Therefore, the isomorphism restricts to an isomorphism of the respective zero-graded Hopf subalgebras. \square

Theorem 7.12 together with (7.30) implies that in the skew-central subalgebras are determined by Hopf algebras that are *independent* of the choice of a root of unity, except for the mod 4-parity of its order.

Conjecture 7.13. *For given root system Φ , denote by B be the Borel subgroup of the respective complex, simply connected simple algebraic group and by \check{B} the Borel subgroup so associated to the coroot system $\check{\Phi}$. Let further $\check{\mathcal{Z}}_{\bullet}^{\geq 0}$ be the analogous skew-central subalgebra in an extension of $U_\zeta^{\geq 0}$ for a primitive ℓ -th root of unity ζ .*

- i) $\check{\mathcal{Z}}_{\bullet}^{\geq 0}$ is also a Hopf subalgebra for any ℓ , isomorphic to a centrally extended matrix algebra, generalizing the definition of [AST91] and associated to either B or \check{B} or a finite quotient of these.
- ii) This extended matrix algebra depends up to isomorphism only on parity of ℓ modulo 4e.

7.4. Ideals from Algebraic Subgroups of T_n and Bruhat Subgroups. In this section we focus entirely on type A_{n-1} algebras and the case $\ell \not\equiv 2 \pmod{4}$, for which $\mathcal{Z}_{\bullet}^{\geq 0}$ is commutative. In this setting, Theorem 7.12 yields an isomorphism between commutative rings, whose scalars we can be further extended to a field $\mathbb{k} = \mathbb{Q}(\zeta)$ or $\mathbb{k} = \mathbb{C}$, assuming an embedded ground ring $\Lambda \hookrightarrow \mathbb{k}$.

As before, let $T_n < \mathrm{GL}(n, \mathbb{k})$ be the algebraic group of upper triangular matrices over \mathbb{k} . Denote also $\mathrm{PT}_n < \mathrm{PGL}(n, \mathbb{k})$ the respective subgroup of the projective linear group. We will use the following identifications with coordinate rings.

Corollary 7.14. *Assume $\mathbb{k} \not\equiv 2 \pmod{4}$ and Lie types \mathfrak{gl}_n or \mathfrak{sl}_n as above. Then the isomorphisms from Theorem 7.12 composed with the assignment $R_{i,j} \mapsto \hat{x}_{i,j}$ yield isomorphisms of Hopf algebras*

$$\tilde{\mathcal{Z}}_{\bullet}^{\geq 0} \otimes \mathbb{k} \cong \mathbb{k}[T_n] \quad \text{and} \quad \mathcal{Z}_{\bullet}^{\geq 0} \otimes \mathbb{k} \cong \mathbb{k}[\mathrm{PT}_n].$$

Proof. Clearly, $\tilde{\mathcal{Z}}_{\bullet}^{\geq 0} \otimes \mathbb{k} = \tilde{\mathcal{Z}}_{\bullet}^{\geq 0} \otimes \mathbb{k}$ and $\mathcal{Z}_{\bullet}^{\geq 0} \otimes \mathbb{k} = \mathcal{Z}_{\bullet}^{\geq 0} \otimes \mathbb{k}$. It is also easy to see that $\mathcal{O}(T_n, \mathbb{k}) \cong \mathbb{k}[T_n]$ as commutative Hopf algebras by comparing coproducts in (7.8) and (7.28). On the diagonal the assignment extends uniquely to $R_{i,i}^{-1} \mapsto \hat{x}_{i,i}^{-1}$. Recall that PT_n is defined as the quotient T_n/C by the center $C = \mathbb{k}^\times \cdot 1$ so that $\mathbb{k}[\mathrm{PT}_n]$ embeds in $\mathbb{k}[T_n]$. Functions in $\mathbb{k}[\mathrm{PT}_n]$ are those constant on each C -coset. For given $\lambda \in \mathbb{k}^\times$, $g \in T_n$, and $\hat{x}^f = \prod_{i \leq j} \hat{x}_{i,j}^{f_{ij}}$ a general monomial, one readily verifies that $\hat{x}^f(\lambda g) = \lambda^{\sum_{i \leq j} f_{ij}} \cdot \hat{x}^f(g)$. Thus $\hat{x}^f \in \mathbb{k}[\mathrm{PT}_n]$ iff $\sum_{i \leq j} f_{ij} = 0$. Since $\mathfrak{q}(\prod_{i \leq j} R_{i,j}^{f_{ij}}) = \sum_{i \leq j} f_{ij}$, the isomorphism $\mathbb{k}[T_n] \rightarrow \mathcal{O}(T_n, \mathbb{k})$ maps $\mathbb{k}[\mathrm{PT}_n]$ precisely to $\mathcal{O}_o(T_n, \mathbb{k})$, which is thus identified with $\mathcal{Z}_{\bullet}^{\geq 0} \otimes \mathbb{k}$. \square

Note that the coalgebra structure for $\mathbb{k}[\mathrm{PT}_2] \cong \mathbb{k}[\hat{\lambda}, \hat{\lambda}^{-1}, \hat{x}]$ is given by (7.5), whereas the coordinate ring of the Borel part in the 2-fold cover $\widetilde{\mathrm{PT}}_2 < \mathrm{SL}(2, \mathbb{C})$ is of the form $\mathbb{k}[\hat{\chi}, \hat{\chi}^{-1}, \hat{x}]$ with coproduct $\Delta(\hat{x}) = \hat{x} \otimes \hat{\chi} + \chi^{-1} \otimes \hat{x}$. Clearly, the two groups and their respective coordinate rings are *not* isomorphic. To avoid these types of subtleties, we will confine our discussion below mostly to the case $T_n < \mathrm{GL}(n, \mathbb{C})$. Corollary 7.14 permits us now to defined large families of Hopf ideals via the composition of maps

$$\mathcal{h} : \mathcal{Z}_{\bullet}^{\geq 0} \hookrightarrow \tilde{\mathcal{Z}}_{\bullet}^{\geq 0} \rightarrow \tilde{\mathcal{Z}}_{\bullet}^{\geq 0} \otimes \mathbb{C} \xrightarrow{\cong} \mathbb{C}[T_n]. \quad (7.33)$$

Lemma 7.15. *Assume Lie type A_{n-1} , $\Lambda \hookrightarrow \mathbb{C}$ and $\mathbb{k} \not\equiv 2 \pmod{4}$ as above. Let $H \leq T_n$ be any subgroup of the complex subgroup of upper triangular matrices. Then*

$$\mathcal{A}(H) = \{z \in \mathcal{Z}_{\bullet}^{\geq 0} : \mathcal{h}(z)(g) = 0 \ \forall g \in H\} \quad (7.34)$$

is a Hopf ideal in $\mathcal{Z}_{\bullet}^{\geq 0}$. Moreover, $\mathcal{A}(H)$ depends only on the (Zariski) closure of H and for two closed algebraic subgroups $H, K \leq T_n$ we have $\mathcal{A}(H \cap K) = \mathcal{A}(H) + \mathcal{A}(K)$.

Proof. The definition in (7.34) can be rephrased as $\mathcal{A}(H) = \mathcal{h}^{-1}(\mathcal{I}(H))$, with notation for the vanishing ideal of H in $\mathbb{C}[T_n]$ as in Section 7.1. Since \mathcal{h} is a Hopf algebra homomorphism as a composite of such, the preimage of a Hopf ideal is again a Hopf ideal. The other properties follow directly from Lemma 7.1. \square

Subgroups of $P \leq T_2$ were discussed and classified already in Corollary 7.3. More general examples of algebraic subgroups may be structured using $T_n = D_m \ltimes U_n$, where D_m and U_n are as in Section 7.1. Algebraic subgroups $A \leq D_m$ are given as products of finite groups and copies of \mathbb{C}^\times , yielding further abelian subgroups $A' = gAg^{-1}$ for any $g \in U_n$. Any such A' acts

on U_n by conjugation, and any algebraic subgroup $V \leq U_n$ that is A' -stable yields an algebraic subgroup $A'V \leq T_n$. Respective Hopf ideals in $\mathcal{Z}_*^{\geq 0}$ for $AU_n = A'U_n$ are generated by expressions of the form $\prod_i L_i^{k_i} - 1$, while the ideals for general $A'V$ are typically generated also by linear expressions in the coordinate functions.

More relevant for our purposes are algebraic subgroups closely related to the Bruhat decomposition $G = \bigsqcup_{s \in \mathcal{W}} C(s)$ with Bruhat cells $C(s) = B \dot{s} B$ defined as double cosets. Here G is a connected complex reductive algebraic group, $B \leq G$ a Borel subgroup, and $\dot{s} \in N(T)$ a representative of $s \in \mathcal{W} \cong N(T)/T$, where $T \leq B$ is a maximal torus of G and $N(T)$ its normalizer. We will call the following the *Bruhat subgroup* associated to an element $s \in \mathcal{W}$.

$$B(s) = B \cap \dot{s}B\dot{s}^{-1}. \quad (7.35)$$

To explain the relation to the Bruhat decomposition, let $U \leq B$ be the usual unipotent group such that B is a semidirect product of $B = D \ltimes U$ with the Cartan subgroup $D \geq T$ (in the sense of $(\mathbb{C}^\times)^n \geq (S^1)^n$). Analogous to (7.35), set $U(s) = U \cap \dot{s}U\dot{s}^{-1} = B(s) \cap U$, which is generated by subgroups $\{U_\alpha : \alpha \in \mathcal{N}(s^\perp)\}$, where s^\perp is as in (2.20). For each $s \in \mathcal{W}$, the map $U(s^\perp) \rightarrow C(s)/B : u \mapsto [u\dot{s}]$ is a bijection, thus relating the collection $\{U(s)\}$ of subgroups to the Bruhat partition of the coset space G/B . Note that, $U(s^\perp) = U \cap \dot{s}U^-\dot{s}^{-1}$, where $U^- = \dot{s}_* U \dot{s}_*^{-1}$ is the respective group associated to negative roots. For details see, for example, [Bum04, Ch30], [Spr09, §8.3], or [Con17, §10.3,N].

The augmentation ideals $\mathcal{K}(s)$, defined in (6.63), are next identified with the vanishing ideals on the Bruhat subgroups. Here, the latter are defined with respect to $G = \mathrm{GL}(n, \mathbb{C})$, $B = T_n$, and $U = U_n$ as above.

Theorem 7.16. *For Lie type A, $\mathbb{k} \neq 2 \pmod{4}$, and any $s \in \mathcal{W}$ we have*

$$\mathcal{A}(B(s)) = \mathcal{K}(s)^{\geq 0}.$$

Proof. Note first that for $s \in \mathcal{W} = S_n$ the representative \dot{s} is given by the usual permutation matrix, acting on the canonical basis by $\dot{s}e_i = e_{s(i)}$. Suppose $L \in T_n$, meaning that $L_{i,i} \neq 0$ and $L_{i,j} = 0$ for $i > j$. The additional condition $L \in \dot{s}B\dot{s}^{-1}$ is then equivalent to $L_{i,j} = 0$ if $i < j$ but $s^{-1}(i) > s^{-1}(j)$. Thus, with $\alpha = \epsilon_i - \epsilon_j \in \Phi^+$, the latter is equivalent to $s^{-1}(\alpha) = \epsilon_{s^{-1}(i)} - \epsilon_{s^{-1}(j)} \in \Phi^-$ and, hence, $\alpha \in \mathcal{N}(s)$. It is thus clear that a polynomial in $\hat{x}_{i,j}$ and $\hat{x}_{i,i}^{-1}$ is zero on $B(s)$ if and only if it is in the ideal generated by $\{\hat{x}_{i,j} : \epsilon_i - \epsilon_j \in \mathcal{N}(s)\}$. The \mathbb{k} -preimage of this set is then the ideal generated by $\{X_\alpha : \alpha \in \mathcal{N}(s)\}$, which is precisely $\mathcal{K}(s)^{\geq 0}$. \square

It is important to keep in mind that the definition of $\mathcal{A}(H)$ in Lemma 7.33 depends on the isomorphisms in Theorem 7.12. These, in turn, depend on the choice of the ordering in (7.11) which determine the $E_{i,j}$ and thus $X_{i,j}$ generators. The equality in Theorem 7.16 implies now shared properties of the identified ideals.

Corollary 7.17. *The following statements hold.*

- i) $\mathcal{A}(H)$ does not depend on a choice of ordering or isomorphism if H is a Bruhat subgroup.
- ii) If $\mathbb{k} \not\equiv 2 \pmod{4}$, then $\widehat{\mathcal{K}}(s)^{\geq 0}$ is a Hopf ideal in $U_\zeta^{\geq 0}(\mathfrak{sl}_n)$ and $U_\zeta(\mathfrak{sl}_n)$ for any $s \in \mathcal{W}$.

The first statement is immediate from word-independence of $\mathcal{K}(s)$, which itself a consequence of Theorem 6.19 and the definition in (6.63).

For the second assertion recall the notation $\widehat{J} = J \cdot U_\zeta^\square$ from Proposition 6.25, extending ideals in the subalgebras $\mathcal{Z}_\bullet^{\geq 0}$ to ideals in the respective quantum algebras. \mathcal{G} -invariance trivially holds in the commutative case. It is clear that, if $\mathcal{Z}_\bullet^{\geq 0}$ is a Hopf subalgebra and J a Hopf ideal in $\mathcal{Z}_\bullet^{\geq 0}$, then \widehat{J} is a Hopf ideal in the quantum algebra as well. By Theorem 7.16 and Lemma 7.15 this is the case for $J = \mathcal{K}(s)^{\geq 0}$ for any $s \in \mathcal{W}$.

A natural question related to the first item in Corollary 7.17 asks, for which closed algebraic subgroups $C \leq B$ the resulting ideal $\mathcal{A}(C)$ can be constructed without the choice of a particular ordering. The observation in Item ii) of the corollary will be proved for all Lie types in Theorem 9.20 below.

The full $\tilde{\mathcal{Z}}_\bullet = \tilde{\mathcal{Z}}_\bullet^{\geq 0} \cdot \tilde{\mathcal{Z}}_\bullet^{\leq 0}$ subalgebra naturally also forms a Hopf subalgebra, which, over \mathbb{C} , can also be identified with the coordinate ring of the algebraic group $W_n = \mathrm{GL}(n, \mathbb{C})^\#$ as defined in (1.1). One Borel algebra $B^+ = T_b$ is given by the usual upper triangular matrices and the opposite $B^- = T_n^-$ by the lower triangular ones. The algebraic maximal torus $D_n = T_n^- \cap T_n$ is, hence, given by the diagonal matrices in $\mathrm{GL}(n, \mathbb{C})$. Rephrasing definition (1.1) in this notation, we have

$$W_n = \ker(T_n^- \times T_n \longrightarrow D_n \times D_n \longrightarrow D_n).$$

The construction also implies $W_n \cong D_n \ltimes (U_n^- \times U_n)$, where U_n^- and U_n are the respective unipotent subgroups. The next result extends Corollary 7.14. Although the existence of a Hopf algebra isomorphism is already implied in [DCKP92] for odd \mathbb{k} , we construct the map here explicitly via assignments of the polynomial generators of the form $R_{i,j}^\pm \mapsto \hat{x}_{i,j}^\pm$ and also include the case $\mathbb{k} \equiv 0 \pmod{4}$.

Lemma 7.18. *Suppose ζ is a \mathbb{k} -th root of unity and $U_\zeta = U_\zeta(\mathfrak{gl}_n)$ as defined above.*

- i) $\tilde{\mathcal{Z}}_\bullet$ is a Hopf subalgebra of U_ζ .
- ii) For $\mathbb{k} \not\equiv 2 \pmod{4}$ there is an isomorphism of Hopf algebras $\tilde{\mathcal{Z}}_\bullet \otimes \mathbb{C} \cong \mathbb{C}[W_n]$.
- iii) Moreover, for such \mathbb{k} , any closed algebraic subgroup of W_n defines a Hopf ideal in $\tilde{\mathcal{Z}}_\bullet$ and, hence, a Hopf ideal of U_ζ .

Proof. Recall that the involution $\Pi : U_\zeta \rightarrow U_\zeta^{\mathrm{cop}}$ from (4.13) maps $U_\zeta^{\leq 0(\geq 0)}$ to $U_\zeta^{\geq 0(\leq 0)}$ with the opposite coproduct. In particular, since $\tilde{\mathcal{Z}}_\bullet^{\geq 0}$ is a Hopf subalgebra, then so is $\tilde{\mathcal{Z}}_\bullet^{\leq 0} = \Pi(\tilde{\mathcal{Z}}_\bullet^{\geq 0})$. The product map $\tilde{\mathcal{Z}}_\bullet^{\leq 0} \otimes \tilde{\mathcal{Z}}_\bullet^{\geq 0} \rightarrow \tilde{\mathcal{Z}}_\bullet$ is a surjective Hopf algebra homomorphism with kernel given by the Hopf ideal $J = \sum_i (L_i^{-1} \otimes L_i - 1)$. Composing this with $\Pi \otimes \mathrm{id}$, we obtain a surjection $\tilde{\mathcal{Z}}_\bullet^{\geq 0, \mathrm{cop}} \otimes \tilde{\mathcal{Z}}_\bullet^{\geq 0} \rightarrow \tilde{\mathcal{Z}}_\bullet$ with kernel $J' = \sum_i (L_i \otimes L_i - 1)$. Over \mathbb{C} the isomorphism from Corollary 7.14

allows us to replace $\tilde{\mathcal{Z}}_{\bullet}^{\geq 0}$ with $\mathbb{C}[T_n]$ and $\tilde{\mathcal{Z}}_{\bullet}^{\geq 0, \text{cop}}$ with $\mathbb{C}[T_n^{\text{opp}}]$, where T_n^{opp} is naturally identified with T_n^- via transposition.

The kernel of the resulting epimorphism $\mathbb{C}[T_n^- \times T_n] = \mathbb{C}[T_n^-] \otimes \mathbb{C}[T_n] \rightarrow \tilde{\mathcal{Z}}_{\bullet}$ is now the ideal $J'' = \sum_i ((\hat{x}_{i,i}, \hat{x}_{i,i}) - 1) = \sum_i (\hat{x}_{i,i} \otimes \hat{x}_{i,i} - 1)$. The subgroup $\mathcal{V}(J'')$ now consists of $(a, b) \in T_n^- \times T_n$ such that $\hat{x}_{i,i}(a)\hat{x}_{i,i}(b) = 1$ for all $i = 1, \dots, n$. The set of these pairs is now exactly the kernel W_n defined above. Lemma 7.1 then implies $J'' = \mathcal{I}(W_n)$ and thus $\mathbb{C}[W_n] \cong \tilde{\mathcal{Z}}_{\bullet}$. The last statement is now also immediate from Lemma 7.1. \square

Consider the natural center action of \mathbb{C}^{\times} on W_n , given by $\lambda.(b^-, b^+) = (\lambda^{-1}b^-, \lambda b^+)$ with $b^- \in T_n^-$ and $b \in T_n$. The respective \mathbb{C}^{\times} action on the coordinate ring of W_n over \mathbb{C} yields a direct decomposition $\mathbb{C}[W_n] = \bigoplus_{k \in \mathbb{Z}} \mathbb{C}[W_n]_k$ with $f \in \mathbb{C}[W_n]_k$ characterized by $\lambda.f = \lambda^k \cdot f$. It is easy to check that this decomposition coincides with the one imposed by the \mathfrak{p} -grading introduced in Section 7.3.

Thus, for $\mathfrak{k} \not\equiv 2 \pmod{4}$, the isomorphism in Lemma 7.18 restricts to an isomorphism of Hopf algebras between $\mathcal{Z}_{\bullet} \otimes \mathbb{C}$ and $\mathbb{C}[W_n]_0$, the \mathbb{C}^{\times} -invariant part of the coordinate ring, which, in turn, may be identified with $\mathbb{C}[\text{PGL}(n, \mathbb{C})^{\#}]$.

The Artin group $\mathcal{A} = B_n$ action on $U_q(\mathfrak{sl}_n)$ generated by the Γ_i naturally extends to an action on $U_q(\mathfrak{gl}_n)$ by $\Gamma_s(P_i) = P_{s(i)}$, where $i \mapsto s(i)$ is the usual permutation for $s \in \mathcal{W} = S_n$. Lemma 7.18 now implies an action of B_n by algebra automorphisms of $\mathbb{C}[W_n]$. Invoking the standard correspondence between algebra homomorphisms of coordinate rings and morphisms of algebraic varieties, this implies an action of B_n on W_n .

Corollary 7.19. *There exists a non-trivial action of the braid group B_n on n strands by algebraic automorphisms on the algebraic group W_n , commensurable with Lusztig's B_n -action on the quantum group of Lie type A_{n-1} .*

It is, worth contrasting the correspondence in Lemma 7.18 for the central subalgebras with that of the traditional correspondences for the original quantum groups. For the latter $U_q^{\geq 0}$ and $U_q^{\leq 0}$ are associated with the opposing Borel subgroups and $U_q = U_q^{\geq 0} \cdot U_q^{\leq 0}$ with the original simple or reductive Lie group.

However, for the central subalgebras, the associated Lie group W_n is a solvable group and thus fundamentally different from $\text{GL}(n, \mathbb{C})$. Correspondingly, the coordinate ring $\mathbb{C}[W_n] \cong \tilde{\mathcal{Z}}_{\bullet}$ is cosolvable in the sense of Corollary 7.5, whereas $\mathbb{C}[\text{GL}(n, \mathbb{C})]$ is cosemisimple since $\text{GL}(n, \mathbb{C})$ is reductive. Similarly, the of group algebraic characters $\text{Spec}_{\mathfrak{m}}(W_n) \cong \text{Gr}(\tilde{\mathcal{Z}}_{\bullet}) \cong \mathbb{Z}^n$ differs from the one in (7.4).

Lemma 7.18 allows us to associate, for example, the Hopf ideal $\mathcal{I}((K \times H) \cap W_n)$ to any pair of subgroups $H \leq T_n$ and $K \leq T_n^-$, which then define Hopf ideals $\mathcal{A}(K, H)$ in \mathcal{Z}_{\bullet} as before. These may then be reexpressed as $\mathcal{A}(K, H) = \mathcal{A}(K, T_n) + \mathcal{A}(T_n^-, H) = (\mathcal{A}(K)^{\leq 0} + \mathcal{A}(H)^{\geq 0}) \cdot \mathcal{Z}_{\bullet}$. Specialized to the Bruhat subgroups this yields, together with Theorem 7.16 and a respective definition of a

Bruhat subgroup $B(t)^- < T_n^-$, the identity of ideals

$$\mathcal{A}(B(t)^-, B(s)) = (\mathcal{K}(t)^{\leq 0} + \mathcal{K}(s)^{\geq 0}) \cdot \mathcal{Z}. \quad \text{for any } s, t \in \mathcal{W}. \quad (7.36)$$

Concluding our discussion of the A_{n-1} algebras, we briefly address the case $\mathcal{K} \equiv 2 \pmod{4}$, when $\xi = -1$, as well as an approach to the next simply laced Lie types.

We first provide a concrete description of Tanisaki's isomorphism from [Tan16] for the subgroup $T_n < \mathrm{GL}(n, \mathbb{C})$. Let $I_s \in T_n$ be the diagonal matrix with $(I_s)_{s,s} = -1$ and $(I_s)_{i,i} = 1$ for $i \neq s$. For any $1 \leq s \leq t \leq n$ define the map $\tau_{s,t} : T_n \rightarrow T_n$ by $\tau_{s,t}(b) = I_s \cdot b \cdot I_t$. Clearly, all $\tau_{s,t}$ are of order 2 and commute with each other, generating an elementary abelian 2-group \mathcal{G} of isomorphisms of T_n as an algebraic variety. Obviously, none of the $\tau_{s,t}$ are a group homomorphism of T_n and the quotient variety T_n/\mathcal{G} retains no group structure either.

Lemma 7.20. *Assume $\mathcal{K} \equiv 2 \pmod{4}$ and Lie type A_{n-1} . Let $\mathcal{Z}_{\mathrm{Fr}}^{\geq 0}$ be the intersection of $\tilde{\mathcal{Z}}_{\bullet}^{\geq 0} \otimes \mathbb{C}$ with the center of $U_{\zeta, \mathbb{C}}$. Then there is a weight-preserving algebraic isomorphism of commutative algebras,*

$$\mathcal{Z}_{\mathrm{Fr}}^{\geq 0} \cong \mathbb{C}[T_n]^{\mathcal{G}} \cong \mathbb{C}[T_n/\mathcal{G}].$$

Proof. It follows from the commutation relations in Section 6.2 that $\mathcal{Z}_{\mathrm{Fr}}^{\geq 0}$ is identical with the center of $\tilde{\mathcal{Z}}_{\bullet}^{\geq 0} \otimes \mathbb{C} \cong \mathcal{O}(T_n, \mathfrak{a}, \mathbb{C})$ itself. Denote $R^f = \prod_{i \leq j} R_{i,j}^{f_{i,j}}$ for some fixed ordering and $\hat{x}^f = \prod_{i \leq j} \hat{x}_{i,j}^{f_{i,j}}$ as before. Recall that $R_{i,j} R_{s,t} = (-1)^{\delta_{i,s} + \delta_{j,t}} R_{s,t} R_{i,j}$ and observe that $\tau_{s,t}^*(\hat{x}_{i,j}) = (-1)^{\delta_{i,s} + \delta_{j,t}} \hat{x}_{i,j}$. It is clear from these formulae that R^f is central if and only if \hat{x}^f is \mathcal{G} -stable. Write $M \subseteq \mathbb{Z}^{\binom{n+1}{2}}$ for the set of exponents $f = (f_{i,j})$ for which this is the case, namely, those for which $\sum_{s \leq j} f_{s,j} + \sum_{i \leq t} f_{i,t} \equiv 0 \pmod{2}$ for all $s \leq t$. Depending on the chosen ordering, it now follows from commutativity that one can then choose a quadratic form ϑ on $M \otimes \mathbb{Z}$ such that for $\tilde{R}^f = (\sqrt{-1})^{\vartheta(f)} R^f$ we have $\tilde{R}^f \cdot \tilde{R}^g = \tilde{R}^{f+g}$. Thus, $\hat{x}^f \mapsto \tilde{R}^f$ defines an isomorphism of algebras over \mathbb{C} . \square

As an elementary example, it is easy to see that $\mathbb{C}[T_2]^{\mathcal{G}}$ corresponds to the subalgebra generated by $R_{1,1}^2, R_{1,2}^2$, and $R_{2,2} R_{1,1}^{-1}$, which is clearly not a Hopf subalgebra. Also, we have $\mathcal{G} \cong \mathbb{F}_2 \times \mathbb{F}_2$ acting component-wise on $T_2 \cong (\mathbb{C}^\times)^2 \times \mathbb{C}$ by $(x_{11}, x_{22}) \mapsto (-x_{11}, -x_{22})$ and $x_{12} \mapsto -x_{12}$, respectively. So, T_2/\mathcal{G} has no obvious group structure.

As an alternative approach to finding an interpretation of $\mathcal{O}(T_n, \mathfrak{a}, \mathbb{C})$ and thus $\tilde{\mathcal{Z}}_{\bullet}^{\geq 0} \otimes \mathbb{C}$ we may seek to retain the coordinate ring $\mathbb{C}[T_n]$ as a target set, but modify the multiplication and comultiplication maps as weighted averages with respect to some \mathcal{G} -action. We propose the following form of a possible Hopf algebra structure.

Conjecture 7.21. *There exist modified multiplication \star and comultiplication $\ddot{\Delta}$ on $\mathbb{C}[T_n]$ of the form*

$$\phi \star \psi = \sum_{\sigma, \tau} \mu(\sigma, \tau) \tau^*(\phi) \sigma^*(\psi) \quad \text{and} \quad \ddot{\Delta}(\phi)(a, b) = \sum_{\sigma, \tau} \eta(\sigma, \tau) \phi(\sigma(a) \tau(b)), \quad (7.37)$$

such that $(\mathbb{C}[B_n], \star, \ddot{\Delta})$ is isomorphic to $\mathcal{O}(T_n, \mathfrak{a}, \mathbb{C})$. Here summations are over algebraic involutions $\sigma : T_n \rightarrow T_n$ with $\sigma^*(\hat{x}_{i,j}) = \pm \hat{x}_{i,j}$ and the weights μ and η take values in $\mathbb{Z}[\frac{1}{2}]$.

Again, the involutions are not required to be homomorphisms. In this form, the definition of Hopf ideals as vanishing ideals on subgroups then requires that these subgroups are preserved under the groups of involutions. This is indeed the case for the Bruhat subgroups. We mention that versions of the specific quantum matrix algebras $\mathcal{O}(T_n, \mathfrak{a}, \Lambda)$ are recovered also in Proposition 2 of [Cli94] as images of a Frobenius map.

Overall, however, the skew matrix algebras $\mathcal{O}(T_n, \mathfrak{a}, \Lambda)$ are scarcely studied in the literature despite their natural occurrences, lacking any semi-classical interpretations one might expect. Following [AST91] the algebra $\mathcal{O}(T_n, \mathfrak{a}, \Lambda)$ is constructable from the skew coordinate algebra generated by $\{x_i : i = 1, \dots, n\}$ subject to relations $x_i x_j = -x_j x_i$ if $i \neq j$. Particularly, the generators x_i remain of infinite order and the algebra is fundamentally distinct from finite-dimensional exterior or Clifford algebras. There, thus, appear to be no obvious identifications of the skew matrix algebras with function algebras on super or spin groups.

Among the simply laced Lie types, D_n has several well understood relations to A type algebras. Clearly, it contains an A_{n-2} algebra and we, conjecturally, expect there to be a quotient of the respective algebras from the A_{2n-1} onto the D_n . The derivations of explicit coproduct formulae as those in (7.22) and (7.25), however, are considerably more challenging.

Kharchenko [Kha15] provides explicit coproduct formulae for the generic D_n algebra with coefficients over Λ_q reminiscent of the ones in (7.22) for type A . Computations of powers to arrive at the analogues of (7.25), however, are substantially more complicated since two types of basis generators are involved. They are defined via recursions as the one in (7.17) but in opposing directions, and neither subset of generators spans a subcoalgebra.

8. THE STRUCTURE OF \mathcal{Z} AND HOPF IDEALS FOR TYPE B_2

The accessible computational tools for \mathfrak{gl}_n allowed us to verify the claims in Conjectures 7.13 and 8.11 computationally for all A_{n-1} algebras. As additional evidence for these conjectures, we prove them in this section also for the simplest non-simply laced Lie type $B_2 = C_2$. The descriptions of isomorphisms and subspaces will be in very explicit terms, while computations are much more involved. The latter are assured to play a central role in extensions to other doubly-laced Lie types, as they illustrate additional methods and principles for the construction of isomorphisms.

Throughout this section, we assume that the order of ζ is $\mathbb{k} > 4$ or $\mathbb{k} = 3$, but impose no other restrictions on \mathbb{k} . As pointed out in Remark 6.15 for B_2 , the subalgebra $\mathcal{Z}_\bullet^{\geq 0}$ is commutative for all \mathbb{k} , though central in U_ζ only if \mathbb{k} is not a multiple of 4.

In Section 8.1 we describe the explicit Hopf algebra structure of $\mathcal{Z}_\bullet^{\geq 0}$, deferring the computation of coproducts for non-simple roots to Section 8.2. The fact that \mathcal{Z} is a Hopf subalgebra

may be indirectly inferred also from [DCKP92] for odd \mathbb{k} , though no explicit formulae or computations are given there, leaving open also the case of even roots of unity. The Hopf algebra structure of the Borel coordinate ring $\mathbb{C}[\mathrm{SO}_5^{\geq 0}]$ is explicitly determined in Section 8.3. A similarly explicit isomorphism of $\mathbb{C}[\mathrm{SO}_5^{\geq 0}]$ with $\mathcal{Z}_\bullet^{\geq 0} \otimes \mathbb{C}$ is established in Section 8.4.

8.1. Coalgebra Structure of $\mathcal{Z}_\bullet^{\geq 0}$ for Type B_2 . This section provides the full list of coproduct formulae for the generators of $\mathcal{Z}_\bullet^{\geq 0}$ for Lie type B_2 , preceded by a review of essential notation and conventions.

As in Section 2.7, we denote by α_i the short root of B_2 and by α_j the long root. Thus, $s_j(\alpha_i) = \alpha_i + \alpha_j$ is the second short root and $s_i(\alpha_j) = 2\alpha_i + \alpha_j$ the second long root. The default ground ring for the generic B_2 algebra is $\Lambda = \Lambda_{q,3} = \mathbb{Z}[q^{\pm 1}, [2]^{-1}]$ as defined in (3.8).

Recall the formulae for the standard generators with respect to $z_{[ij]}$ as defined in (5.19), which follow readily from the automorphisms in (4.20) and (4.21), as well as their w -gradings in \mathbb{Z}^Δ .

$$E_{(iji)} = \check{E}_{(ji)} = \Gamma_j^{-1}(E_i) = q^{-2}E_iE_j - E_jE_i \quad w(E_{(iji)}) = \alpha_i + \alpha_j \quad (8.1)$$

$$E_{(ij)} = \Gamma_i(E_j) = \frac{1}{[2]}(E_iE_{(iji)} - E_{(iji)}E_i) \quad w(E_{(ij)}) = 2\alpha_i + \alpha_j \quad (8.2)$$

From the coproducts for E_i and E_j , as in (3.16), the following expressions for the coproducts of the other two generators are easily derived. Here \check{E}_i and \check{E}_j are singularized generators as defined in (5.10).

$$\Delta(E_{(iji)}) = E_{(iji)} \otimes K_iK_j - E_i \otimes K_i\check{E}_j + 1 \otimes E_{(iji)} \quad (8.3)$$

$$\Delta(E_{(ij)}) = E_{(ij)} \otimes K_i^2K_j + q\check{E}_i^2 \otimes K_i^2E_j - \check{E}_i \otimes K_iE_{(iji)} + 1 \otimes E_{(ij)} \quad (8.4)$$

Choose next a \mathbb{k} -th primitive root of unity ζ with \mathbb{k} as above and ℓ the order of ζ^2 . We also reemploy the notation from (6.49) and (6.53) with $(\mathfrak{d}, \mathfrak{n}) = (1, 2)$ if ℓ is odd and $(\mathfrak{d}, \mathfrak{n}) = (2, 1)$ if ℓ is even.

The terms occurring in the coproduct formulae for the primitive power generators $X_w = E_w^{\ell_w}$ depend on the parity of ℓ as they are differently constrained by the w -grading. More precisely, for $\alpha = \gamma(w)$, the observation in (6.17) implies that $\ell^{-1}w(X_w)$ is given by α if ℓ is odd and the respective coroot $\check{\alpha}$ if ℓ is even. Recall that the coroot system $\check{\Phi}_{B_n}$ is isomorphic to the root system Φ_{C_n} . For $n = 2$ the latter is isomorphic to Φ_{B_2} , yielding a combined root system isomorphism $\ell : \check{\Phi}_{B_2} \rightarrow \Phi_{B_2}$.

The composite $\varphi(\alpha) = \ell(\check{\alpha})$ then defines an involution in $\Phi_{B_2}^+$ that exchanges long with short roots and which is given by $\varphi(\alpha_i) = \alpha_j$, $\varphi(\alpha_j) = \alpha_i$, $\varphi(\alpha_i + \alpha_j) = 2\alpha_i + \alpha_j$, and $\varphi(2\alpha_i + \alpha_j) = \alpha_i + \alpha_j$. Thus, if we define $u(X_w) = \alpha$ for odd ℓ and $u(X_w) = \varphi(\alpha)$ if ℓ is even, we obtain a grading on $\mathcal{Z}_\bullet^{\geq 0}$ that is isomorphic to the induced w -grading, but, for either parity, conveniently valued in $\Phi_{B_2}^+$. As such, the coproduct needs to preserve the u -grading.

For more convenient reference, we list in the table below the \mathfrak{u} -weights for the primitive power generators for words less than $z_{|ij|}$ together with the special constants.

	$\mathfrak{u}(X_i)$	$\mathfrak{u}(X_j)$	$\mathfrak{u}(X_{(ij)})$	$\mathfrak{u}(X_{(iji)})$	\mathfrak{d}	\mathfrak{n}
ℓ odd	α_i	α_j	$2\alpha_i + \alpha_j$	$\alpha_i + \alpha_j$	1	2
ℓ even	α_j	α_i	$\alpha_i + \alpha_j$	$2\alpha_i + \alpha_j$	2	1

(8.5)

We state next the main result of this section, assuming notation and conventions as above as well as the notation ∇_j from (6.51).

Proposition 8.1. *The coproducts of X_w for $\emptyset \neq w \leq_R z_{|ij|}$ are as follows:*

$$\Delta(X_i) = X_i \otimes L_i + 1 \otimes X_i, \quad \Delta(X_j) = X_j \otimes L_j + 1 \otimes X_j, \quad (8.6)$$

$$\begin{aligned} \Delta(X_{(iji)}) &= 1 \otimes X_{(iji)} - \nabla_j^{\mathfrak{d}} X_i \otimes L_i X_j^{\mathfrak{d}} + X_{(iji)} \otimes L_i L_j^{\mathfrak{d}} \\ &\quad - \delta_{\mathfrak{d},2} \cdot 2 \cdot [2]^{\ell} (-\zeta)^{\ell_j} X_{(ij)} \otimes L_i L_j X_j, \end{aligned} \quad (8.7)$$

$$\begin{aligned} \Delta(X_{(ij)}) &= 1 \otimes X_{(ij)} + \zeta^{\binom{2^{\ell_j}}{2}} \nabla_i^{\mathfrak{n}} X_i^{\mathfrak{n}} \otimes L_i^{\mathfrak{n}} X_j + X_{(ij)} \otimes L_i^{\mathfrak{n}} L_j \\ &\quad - \delta_{\mathfrak{n},2} \cdot 2 \cdot \zeta^{\ell} [2]^{-\ell} \nabla_i \cdot X_i \otimes L_i X_{(iji)}. \end{aligned} \quad (8.8)$$

As usual, $\delta_{a,b}$ is 1 if $a = b$ and 0 is otherwise. In cases when $\delta_{\mathfrak{d},2}$ or $\delta_{\mathfrak{n},2}$ are zero, the respective terms are indeed not consistent with the \mathfrak{u} -grading. For example, if ℓ is even and $\mathfrak{n} = 1$ the \mathfrak{u} -grading of $X_i \otimes X_{(iji)}$ is $(\alpha_j) + (2\alpha_i + \alpha_j) = 2(\alpha_i + \alpha_j)$, while $\mathfrak{u}(X_{(ij)}) = \alpha_i + \alpha_j$.

The expressions in (8.6) are restatements of (7.6). The explicit proofs of (8.8) and (8.7) are postponed to the following section. The formulae in Proposition 8.1 also involve only elements in $\mathcal{Z}_*^{\geq 0}$, implying the latter is a sub-bialgebra.

Explicit expressions for the antipode can be derived from the axioms, applied as $0 = \epsilon(X_{(iji)}) = \text{mo}(\text{id} \otimes S) \circ \Delta(X_{(iji)})$ and $0 = \epsilon(X_{(ij)}) = \text{mo}(S \otimes \text{id}) \circ \Delta(X_{(ij)})$.

$$S(X_{(ij)}) L_i^{\mathfrak{n}} L_j = -X_{(ij)} - (-1)^{\mathfrak{n}} \zeta^{\binom{2^{\ell_j}}{2}} \nabla_i^{\mathfrak{n}} \cdot X_i^{\mathfrak{n}} X_j - \delta_{\mathfrak{n},2} \cdot 2 \cdot \zeta^{\ell} [2]^{-\ell} \nabla_i \cdot X_i X_{(iji)} \quad (8.9)$$

$$S(X_{(iji)}) L_i L_j^{\mathfrak{d}} = -X_{(iji)} + (-1)^{\mathfrak{d}} \nabla_j^{\mathfrak{d}} X_i X_j^{\mathfrak{d}} - \delta_{\mathfrak{d},2} \cdot 2 \cdot [2]^{\ell} (-\zeta)^{\ell_j} X_{(ij)} X_j \quad (8.10)$$

Proposition 8.1 and the above formulae thus provide further evidence for the first parts of Conjectures 7.13 and 8.11.

Corollary 8.2. *For Lie type B_2 and any \mathfrak{K} as above, $\mathcal{Z}_*^{\geq 0}$ is a Hopf subalgebra over $\Lambda_{\zeta,3}$.*

We remark that the Hopf algebra structure computed above provides an alternative proof for the statement of Propositions 6.17 and 6.18 for $\mathfrak{e} = 2$. Specifically, recall from (4.16) that the antipode is related to the \mathfrak{O} -involution via $\Psi_{(\iota \cdot \varpi)^{-1}}^{\text{id}} \circ S = \mathfrak{O}$. As $\Psi_{(\iota \cdot \varpi)^{-1}}^{\text{id}}(X_{(iji)}) = (-1)^{\ell+1} X_{(iji)} L_i L_j^{\mathfrak{d}}$

and $\Psi_{(\iota \cdot \varpi)^{-1}}^{id}(X_{(ij)}) = (-1)^\ell X_{(ij)} L_i^{\mathfrak{n}} L_j$, applying $\Psi_{(\iota \cdot \varpi)^{-1}}^{id}$ to (8.9) and (8.10) yields the relations below.

$$(-1)^\ell \mathfrak{O}(X_{(ij)}) = -X_{(ij)} - (-1)^{\mathfrak{n}} \zeta^{\binom{2\ell}{2}} \nabla_i^{\mathfrak{n}} X_i^{\mathfrak{n}} X_j - \delta_{\mathfrak{n},2} \cdot 2 \cdot \zeta^\ell [2]^{-\ell} \nabla_i \cdot X_i X_{(iji)} \quad (8.11)$$

$$(-1)^\ell \mathfrak{O}(X_{(iji)}) = X_{(iji)} - (-1)^{\mathfrak{d}} \nabla_j^{\mathfrak{d}} X_i^{\mathfrak{d}} X_j^{\mathfrak{d}} + \delta_{\mathfrak{d},2} \cdot 2 \cdot [2]^\ell (-\zeta)^{\ell_j} X_{(ij)} X_j \quad (8.12)$$

Specializing relation (6.55) to $s = 2$ and $s = 3$ and writing the explicit words as in (2.32) yields $\mathfrak{O}(X_{(ij)}) = X_{b_{m-1}} = X_{(jij)}$ and $\mathfrak{O}(X_{(iji)}) = X_{b_2} = X_{(ji)}$. Comparison of coefficients then yields (6.54) and (6.52) for $\mathbf{e} = 2$.

See also Appendix D for an integral form of the coalgebra structure in Proposition 8.1.

8.2. Proof of Proposition 8.1 for Non-simple Roots. The proof of the formulae (8.8) and (8.7) relies on a generalized quantum multinomial formula. It involves a special type of quantum number, which we introduce first. For $k, t \in \mathbb{N}_0$, define $c_{k,t} \in \Lambda_q$ recursively in t by

$$c_{k,t} = \sum_{i=0}^k c_{i,t-1} q^{-(2t-2+i)} [2t-1+i], \quad (8.13)$$

with initial conditions $c_{k,0} = 1$. For $k, t > 0$ one readily derives the relations

$$c_{k,t} = c_{k-1,t} + c_{k,t-1} q^{-(2t-2+k)} [2t+k-1] \quad \text{and} \quad c_{0,t} = q^{-t(t-1)} [2t-1]!! , \quad (8.14)$$

denoting as usual the quantum double factorial $[a]!! = [a][a-2] \cdots [a-2\lfloor(a-1)/2\rfloor]$. The following identity is a generalization of [Lus90a, Lemma 1.6], where it applies only to the simply laced situation.

Proposition 8.3. *Suppose A, B, C, D satisfy*

$$DA = q^{-2}AD + B, \quad CA = q^{-2}AC, \quad BA = q^{-4}AB, \quad DC = q^{-2}CD, \quad DB = q^{-4}BD \quad CB = BC.$$

Then $(A + C + D)^n$ is equal to

$$\sum_{s=0}^n \sum_{t=0}^{\min(s,n-s)} \sum_{k=0}^{n-s-t} q^{-(s+t+k)(n-s-t-k)-(s-t)(2t+k)} \begin{bmatrix} n \\ s+t+k \end{bmatrix} \begin{bmatrix} s+t+k \\ s-t \end{bmatrix} c_{k,t} A^{s-t} B^t C^k D^{n-s-t-k}.$$

The technical proof of this proposition is given in Appendix B. For the following computations at generic q we consider the quantum algebra $U_q^{\geq 0}$ over the ground ring $\Lambda_{q,3}$ for Lie type B_2 . Additional commutation relations follow, for example, from Lemma 5.6 with $E_{(iji)} = E_j$:

$$E_{(iji)} E_{(ij)} = q^2 E_{(ij)} E_{(iji)}, \quad E_{(ij)} E_i = q^2 E_i E_{(ij)}, \quad \text{and} \quad E_j E_{(iji)} = q^2 E_{(iji)} E_j. \quad (8.15)$$

We first compute the coproduct of the generator for the non-simple short root.

Proof of Equation (8.7). We begin with the computation of $\Delta(E_{(iji)})^n$ for generic q . Proposition 8.3 is applied by setting $C = 0$ as well as

$$A = E_{(iji)} \otimes K_i K_j, \quad D = D_1 + D_2 \quad \text{with} \quad D_1 = (q^{-2} - q^2) E_i \otimes K_i E_j \quad \text{and} \quad D_2 = 1 \otimes E_{(iji)}, \quad (8.16)$$

so that $\Delta(\check{E}_{(iji)}) = A + D$. One easily checks that $D_2 D_1 = q^2 D_1 D_2$. We obtain identities

$$D^m = \sum_{u=0}^m q^{u(m-u)} \begin{bmatrix} m \\ u \end{bmatrix} D_1^u D_2^{m-u} \quad \text{and} \quad B = q^{-2}[2](q^{-2} - q^2) E_{(ij)} \otimes K_i^2 K_j E_j, \quad (8.17)$$

where the former now follows from (3.3) and the latter from $B = DA - q^{-2}AD$, as stipulated in Proposition 8.3. The other required commutation relations among A , B , and D are derived from (8.1) and (8.15). Applying the proposition to the case $C = 0$ forces $k = 0$ in the summation, which yields, together with (8.17) and (8.14), the following expression.

$$\begin{aligned} \Delta(E_{(iji)}^n) = (A + D)^n &= \sum_{s=0}^n \sum_{t=0}^{\min(s,n-s)} q^{-(s+t)(n-s-t)-(s-t)(2t)} \begin{bmatrix} n \\ s+t \end{bmatrix} \begin{bmatrix} s+t \\ s-t \end{bmatrix} c_{0,t} A^{s-t} B^t D^{n-s-t} \\ &= \sum_{s=0}^n \sum_{t=0}^{\min(s,n-s)} \sum_{u=0}^{n-s-t} q^{e(n,s,t,u)} [2t-1]!! \begin{bmatrix} n \\ s+t \end{bmatrix} \begin{bmatrix} s+t \\ s-t \end{bmatrix} \begin{bmatrix} n-s-t \\ u \end{bmatrix} A^{s-t} B^t D_1^u D_2^{n-s-t-u}, \end{aligned} \quad (8.18)$$

$$\text{where } e(n, s, t, u) = -ns - nt + s^2 + 2t^2 + t + un - us - ut - u^2.$$

Next, we specialize q to a primitive \mathbb{k} -th root of unity ζ with $\mathbb{k} > 4$ or $\mathbb{k} = 3$. We further set $n = \ell$ with ℓ the order of ζ^2 . The product of the three binomial coefficients in (8.18) is readily identified with a multinomial coefficient as in (3.7) for $r = 4$, $a_1 = \ell - s - t - u$, $a_2 = s - t$, $a_3 = 2t$, and $a_4 = u$. Corollary 6.2 now implies that the only contributing non-zero terms are given by the following combinations of indices.

$$(s, t, u) \in \{(0, 0, 0), (0, 0, \ell), (\ell, 0, 0), (\ell/2, \ell/2, 0)\}.$$

Of course, the term $(\ell/2, \ell/2, 0)$ only contributes when ℓ is even, meaning $\mathfrak{d} = 2$. Summing over these four index-tuples gives

$$\begin{aligned} \Delta(X_{(iji)}) &= D_2^\ell + D_1^\ell + A^\ell + \delta_{\mathfrak{d},2} \cdot \zeta_j^{-\binom{\ell}{2}} [\ell-1]!! \cdot B^{\ell/2} \\ &= 1 \otimes E_{(iji)}^\ell + (\zeta^{-2} - \zeta^2)^\ell \zeta_j^{\binom{\ell}{2}} \cdot E_i^\ell \otimes K_i^\ell E_j^\ell + E_{(iji)}^\ell \otimes K_i^\ell K_j^\ell \\ &\quad - \delta_{\mathfrak{d},2} \cdot ([2](\zeta^{-2} - \zeta^2))^{\ell/2} \zeta_j^{-\binom{\ell}{2}} [\ell_j : 2]!! \cdot E_{(ij)}^{\ell_j} \otimes K_i^{\ell_j} K_j^{\ell_j} E_j^{\ell_j} \\ &= 1 \otimes X_{(iji)} - \nabla_j^\mathfrak{d} \cdot X_i \otimes L_i X_j^\mathfrak{d} + X_{(iji)} \otimes L_i L_j^\mathfrak{d} \\ &\quad - \delta_{\mathfrak{d},2} \cdot 2 \cdot (-\zeta)^{\ell_j} [2]^\ell \cdot X_{(ij)} \otimes L_i L_j X_j. \end{aligned}$$

The last step uses Lemma 6.16 to simplify coefficients, yielding (8.7) as the final expression. \square

The proof of the coproduct formula in the case of the non-simple long root is more involved. The required commutation relations are more naturally expressed in terms of $q_j = q^2$ rather than $q_i = q$. Denoting by $\text{sq} : \Lambda_q \rightarrow \Lambda_q$ the homomorphism that maps q to q^2 (that is, q_i to q_j) we define $c'_{k,t} = \text{sq}(c_{k,t})$.

Proof of Equation (8.8). As before, we start with the computation of $\Delta(\check{E}_{(ij)}^n)$ for generic q . The main tool for this proof is the identity in Proposition 8.3 with q replaced by q_j , obtained by applying sq to all coefficients, using $\text{sq}\left(\begin{bmatrix} a \\ b \end{bmatrix}\right) = \begin{bmatrix} a \\ b \end{bmatrix}_j$. We begin by defining summands

$$A' = q(q - q^{-1})^2 E_i^2 \otimes K_i^2 E_j, \quad C' = -(q - q^{-1}) E_i \otimes K_i E_{(iji)},$$

$$D' = 1 \otimes E_{(ij)}, \quad G = E_{(ij)} \otimes K_i^2 K_j,$$

so that

$$\Delta(E_{(ij)}) = A' + C' + D' + G \quad \text{by (8.4).}$$

Commutation relations between these follow again from (8.2) and (8.15). We find

$$GA' = q_j^2 A'G, \quad GC' = q_j^2 C'G, \quad \text{and} \quad GD' = q_j^2 D'G,$$

$$\text{so that} \quad G(A' + C' + D') = q_j^2 (A' + C' + D')G \quad (8.19)$$

$$\text{as well as} \quad C'A' = q_j^{-2} A'C' \quad \text{and} \quad D'C' = q_j^{-2} C'D'. \quad (8.20)$$

Similarly, we compute from previous identities

$$B' = D'A' - q_j^{-2} A'D' = -q^{-2} \frac{(q-q^{-1})^3}{q+q^{-1}} E_i^2 \otimes K_i^2 E_{(iji)}^2 = -q^{-2} \frac{q-q^{-1}}{q+q^{-1}} (C')^2, \quad (8.21)$$

which entails the additional commutation relations

$$B'A' = q_j^{-4} A'B', \quad D'B' = q_j^{-4} B'D', \quad \text{and} \quad C'B' = B'C'. \quad (8.22)$$

In the computation below we first apply the binomial formula (3.3) using (8.19). The second step invokes Proposition 8.3, as we verified the requirements for A' , B' , C' , and D' in (8.21), (8.20), and (8.22) for q_j in place of q .

$$\begin{aligned} \Delta(E_{(ij)}^n) &= ((A' + C' + D') + G)^n = \sum_{m=0}^n \begin{bmatrix} n \\ m \end{bmatrix}_j q_j^{m(n-m)} (A' + C' + D')^{n-m} G^m \\ &= \sum_{\substack{m,s,t,u \geq 0, \\ n \geq m+s+t+u}} q_j^{c(n,m,s,t,u)} \begin{bmatrix} n \\ m \end{bmatrix}_j \begin{bmatrix} n-m \\ s+t+u \end{bmatrix}_j \begin{bmatrix} s+t+u \\ s-t \end{bmatrix}_j c'_{u,t} \\ &\quad \cdot (A')^{s-t} (B')^t (C')^u (D')^{n-m-s-t-u} G^m, \end{aligned} \quad (8.23)$$

$$\text{with} \quad c(n, m, s, t, u) = m(n-m) - (s+t+u)(n-m-s-t-u) - (s-t)(2t+u).$$

Next, we specialize q to the root of unity ζ as before and set $n = \ell_j$. The product of the three binomial coefficients in (8.23) is again a multinomial coefficient for $r = 4$, now with $a_1 = m$, $a_2 = s - t$, $a_3 = 2t + u$, and $a_4 = \ell_j - m - s - t - u$. Corollary 6.2 then confines the summation to the following tuples of indices.

$$(m, s, t, u) \in \{(0, 0, 0, 0), (0, \ell_j, 0, 0), (\ell_j, 0, 0, 0)\} \cup \{(0, t, t, \ell_j - 2t) : 0 \leq t \leq \lfloor \ell_j/2 \rfloor\}$$

Note that with $\mathfrak{c} = (a_1 + a_4)(a_2 + a_3) + a_1 a_4 - a_2 a_3$ the q -exponential in (8.23) is 1 in all cases. Observe further that on the subset indexed by t the operator term is

$$(A')^{s-t}(B')^t(C')^u(D')^{n-m-s-t-u}G^m = (B')^t(C')^{\ell_j-2t} = \left(-\zeta^{-2}\frac{\zeta-\zeta^{-1}}{\zeta+\zeta^{-1}}\right)^t (C')^{\ell_j}.$$

The summation over the non-zero terms, thus, gives

$$\begin{aligned} \Delta(X_{(ij)}) &= (D')^{\ell_j} + (A')^{\ell_j} + G^{\ell_j} + \sum_{t=0}^{\lfloor \ell_j/2 \rfloor} c'_{\ell_j-2t,t} \left(-\frac{\zeta-\zeta^{-1}}{\zeta^2[2]}\right)^t (C')^{\ell_j} \\ &= 1 \otimes E_{(ij)}^{\ell_j} + \zeta^{\ell_j}(q-q^{-1})^{2\ell_j} E_i^{2\ell_j} \otimes \zeta_j^{2\binom{\ell_j}{2}} K_i^{2\ell_j} E_j^{\ell_j} + E_{(ij)}^{\ell_j} \otimes K_i^{2\ell_j} K_j^{\ell_j} \\ &\quad + \zeta^{-\binom{\ell_j}{2}}[2]^{-\ell_j} \{\ell_j:2\} \cdot (-1)^{\ell_j} (\zeta-\zeta^{-1})^{\ell_j} E_i^{\ell_j} \otimes K_i^{\ell_j} E_{(iji)}^{\ell_j} \\ &= 1 \otimes X_{(ij)} + \zeta^{\binom{\ell_j}{2}} \nabla_i^{\mathfrak{n}} X_i^{\mathfrak{n}} \otimes L_i^{\mathfrak{n}} X_j + X_{(ij)} \otimes L_i^{\mathfrak{n}} L_j \\ &\quad - \delta_{\mathfrak{n},2} \cdot 2 \cdot \zeta^{\ell_j}[2]^{-\ell_j} \nabla_i \cdot X_i \otimes L_i X_{(iji)}. \end{aligned}$$

In the last step we used Lemma B.3. Recall from (6.50) that $\{\ell_j:2\} = \delta_{\mathfrak{d},1} \cdot 2 \cdot \zeta^{-\binom{\ell_j+1}{2}}$ is nonzero only if $\ell = \ell_j$, or equivalently if $\mathfrak{n} = 2$. In this case, $E_i^{\ell_j} = X_i$ and $E_{(iji)}^{\ell_j} = X_{(iji)}$ which now gives the desired expression. \square

We note that similar coproduct formulae appear in [BDR02], where the authors compute liftings of the Nichols algebra of a Yetter-Drinfeld module of type B_2 . Indeed, the q -binomial theorem in [BDR02, Sec 4] is the special case of our Proposition 8.3 for B a multiple of C^2 .

8.3. The Borel Matrix Algebra on $\mathrm{SO}(5, \mathbb{C})$. We adopt here the conventions from [GW09, §2.1.2] for the definition of the *complex* orthogonal group. The basis for the symmetric form in odd dimensions contains two isotropic subbases besides an additional vector. The explicit definition in this form is

$$\begin{aligned} \mathrm{SO}(5, \mathbb{C}) &= \{A \in \mathrm{SL}(5, \mathbb{C}) : A^T J A = J\} \quad \text{for} \quad J = \begin{bmatrix} 0 & & 0 & & I \\ & \ddots & & & \\ & & 0 & & \\ & & & \ddots & \\ & & & & 0 & 0 \end{bmatrix}, \quad (8.24) \\ \text{where} \quad I &= \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad \text{and} \quad \mathbb{0} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}. \end{aligned}$$

Following [GW09, §2.1.2], a Cartan subgroup H is given by diagonal matrices $\mathrm{diag}[\lambda, \mu, 1, \mu^{-1}, \lambda^{-1}]$ with $\lambda, \mu \in \mathbb{C}^\times$. For an element $h = \mathrm{diag}[h_1, h_2, 0, -h_2, -h_1] \in \mathfrak{h}$ in the respective Cartan Lie subalgebra, a set of simple roots in \mathfrak{h}^* is then given by $\alpha_1(h) = h_2$ and $\alpha_2(h) = h_1 - h_2$, where α_1 is short and α_2 the long root for B_2 , implying $A_{12} = -2$ and $A_{21} = -1$ for the Cartan data defined in Section 2.

These choices realize the Lie-Kolchin Theorem, meaning, the entailed Borel subgroup $\mathrm{SO}_5^{\geq 0} = \mathrm{SO}(5, \mathbb{C}) \cap \mathrm{T}_5$ is precisely given by the upper triangular matrices in $\mathrm{SO}(5, \mathbb{C})$. It is then straightforward to work out the following expression for an isomorphism $\phi : (\mathbb{C}^\times)^2 \times \mathbb{C}^4 \rightarrow \mathrm{SO}_5^{\geq 0}$ of algebraic sets.

$$\phi \begin{pmatrix} \lambda & a & c \\ \mu & b & d \end{pmatrix} = \begin{bmatrix} F & \vdots & -Fv & \vdots & F(S - \frac{1}{2}vv^\top)I \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 1 & & (Iv)^\top \\ \cdots & \cdots & 0 & & \cdots \\ 0 & & 0 & & (IF^\top I)^{-1} \end{bmatrix} \quad \text{where} \quad F = \begin{bmatrix} \lambda & b \\ 0 & \mu \end{bmatrix},$$

$$S = \begin{bmatrix} 0 & d \\ -d & 0 \end{bmatrix}, \text{ and } v = \begin{bmatrix} c \\ a \end{bmatrix}. \quad (8.25)$$

In these coordinates, the group or matrix multiplication of two elements in $\mathrm{SO}_5^{\geq 0}$ is given by

$$\phi \begin{pmatrix} \lambda_1 & a_1 & c_1 \\ \mu_1 & b_1 & d_1 \end{pmatrix} \cdot \phi \begin{pmatrix} \lambda_2 & a_2 & c_2 \\ \mu_2 & b_2 & d_2 \end{pmatrix} = \phi \begin{pmatrix} \lambda_3 & a_3 & c_3 \\ \mu_3 & b_3 & d_3 \end{pmatrix},$$

where

$$\begin{aligned} \lambda_3 &= \lambda_1 \lambda_2 & \mu_3 &= \mu_1 \mu_2 & a_3 &= \mu_2^{-1} a_1 + a_2 \\ b_3 &= \mu_2 b_1 + \lambda_1 b_2 & c_3 &= \lambda_2^{-1} c_1 + c_2 - \lambda_2^{-1} \mu_2^{-1} a_1 b_2 \\ d_3 &= \lambda_2^{-1} \mu_2^{-1} d_1 + d_2 + \frac{1}{2} (\mu_2^{-1} a_1 c_2 - \lambda_2^{-1} c_1 a_2 + \lambda_2^{-1} \mu_2^{-1} a_1 a_2 b_2). \end{aligned} \quad (8.26)$$

The Cartan subgroup SO_5^0 is then the ϕ -image of the subspace $(\mathbb{C}^\times)^2 \times \{(0, 0, 0, 0)\}$, that is, the matrices with $a = b = c = d = 0$. Similarly, the respective unipotent subgroup SO_5^+ consists of matrices with $\lambda = \mu = 1$, meaning the ϕ -image of $\{(1, 1)\} \times \mathbb{C}^4$.

Note that the inverse map $\phi^{-1} : \mathrm{SO}_5^{\geq 0} \rightarrow (\mathbb{C}^\times)^2 \times \mathbb{C}^4$ can be obtained as the restriction of a polynomial map on $\mathrm{GL}(5, \mathbb{C})$. Specifically, $\lambda, \lambda^{-1}, \mu, \mu^{-1}, a, b$, and c are simply given as the restriction of some $\hat{x}_{i,j}$, and the expression for the d coordinate is easily worked out from these. Thus, $(\mathbb{C}^\times)^2 \times \mathbb{C}^4$ equipped with the multiplication operation from (8.26) is isomorphic to $\mathrm{SO}_5^{\geq 0}$ as an algebraic group.

Correspondingly, we introduce the generating functions as the respective coordinate projections $\mathrm{SO}_5^{\geq 0} \rightarrow (\mathbb{C}^\times)^2 \times \mathbb{C}^4 \rightarrow \mathbb{C}$ for this parametrization, denoting by $\hat{\gamma}$ the function corresponding to a coordinate labeled by γ . So, for example, $\hat{\lambda}(G) = (\phi^{-1}(G))_{1,1}$ or $\hat{d}(G) = (\phi^{-1}(G))_{2,3}$ with $G \in \mathrm{SO}_5^{\geq 0}$. Thus, as a commutative algebra, $\mathbb{C}[\mathrm{SO}_5^{\geq 0}]$ is naturally identified with $\mathbb{C}[\hat{\lambda}^{\pm 1}, \hat{\mu}^{\pm 1}, \hat{a}, \hat{b}, \hat{c}, \hat{d}]$.

Recall from Section 7.1 that the coalgebra structure on $\mathbb{C}[\mathrm{SO}_5^{\geq 0}]$ is defined by $\Delta(f)(a, b) = f(ab)$. The respective coproduct formulae for the generators follow immediately from the expressions in (8.26). For example, if $G_3 = G_1 G_2$ then $\hat{a}(G_3) = a_3 = \mu_2^{-1} a_1 + a_2 = \hat{a}(G_1) \hat{\mu}^{-1}(G_2) + \hat{a}(G_2) = (\hat{a} \otimes \hat{\mu}^{-1} + 1 \otimes \hat{a})(G_1, G_2)$. The complete coalgebra structure is, thus, given as follows.

$$\begin{aligned}
\Delta(\hat{\lambda}) &= \hat{\lambda} \otimes \hat{\lambda} & \Delta(\hat{\mu}) &= \hat{\mu} \otimes \hat{\mu} & \Delta(\hat{a}) &= \hat{a} \otimes \hat{\mu}^{-1} + 1 \otimes \hat{a} \\
\Delta(\hat{b}) &= \hat{b} \otimes \hat{\mu} + \hat{\lambda} \otimes \hat{b} & \Delta(\hat{c}) &= \hat{c} \otimes \hat{\lambda}^{-1} + 1 \otimes \hat{c} - \hat{a} \otimes \hat{\lambda}^{-1} \hat{\mu}^{-1} \hat{b} \\
\Delta(\hat{d}) &= \hat{d} \otimes \hat{\lambda}^{-1} \hat{\mu}^{-1} + 1 \otimes \hat{d} + \frac{1}{2} \hat{a} \otimes \hat{\mu}^{-1} \hat{c} - \frac{1}{2} \hat{c} \otimes \hat{\lambda}^{-1} \hat{a} + \frac{1}{2} \hat{a} \otimes \hat{\lambda}^{-1} \hat{\mu}^{-1} \hat{a} \hat{b}
\end{aligned} \tag{8.27}$$

An antipode is defined by $S(f)(L) = f(L^{-1})$ but can, alternatively, also be computed iteratively from the antipode axiom and the corelations in (8.27). Additionally, a \mathbb{Z}^Δ grading w is defined on $\mathbb{C}[\mathrm{SO}_5^{\geq 0}]$ by assigning to each of the generators a positive roots as follows.

$$w(\hat{\lambda}) = w(\hat{\mu}) = 0, \quad w(\hat{a}) = \alpha_1, \quad w(\hat{b}) = \alpha_2, \quad w(\hat{c}) = \alpha_1 + \alpha_2, \quad \text{and} \quad w(\hat{d}) = 2\alpha_1 + \alpha_2 \tag{8.28}$$

We summarize our computations next.

Proposition 8.4. *The coordinate ring $\mathbb{C}[\mathrm{SO}_5^{\geq 0}]$ is isomorphic, as a w -graded Hopf algebra, to the complex polynomial algebra $\mathbb{C}[\hat{\lambda}^{\pm 1}, \hat{\mu}^{\pm 1}, \hat{a}, \hat{b}, \hat{c}, \hat{d}]$ equipped with the corelations given by (8.27).*

The fact that the abstract polynomial algebra is isomorphic to $\mathbb{C}[\mathrm{SO}_5^{\geq 0}]$ is clear from the fact that monomials $\{\hat{\lambda}^{m_1} \hat{\mu}^{m_2} \hat{a}^{k_1} \hat{b}^{k_2} \hat{c}^{k_3} \hat{d}^{k_4} : m_i \in \mathbb{Z}, k_i \in \mathbb{N}_0\}$ form a basis for both the abstract algebra and for $\mathbb{C}[\mathrm{SO}_5^{\geq 0}]$. The former is well-defined as a coalgebra since the invertible generators are group-like. It is also easy to check that the corelations in (8.27) preserve the grading from (8.28).

Recall that the Weyl group \mathcal{W} for B_2 is the dihedral group of order 8, generated by reflections s_1 and s_2 along $\alpha_1 = \epsilon_2$ and $\alpha_2 = \epsilon_1 - \epsilon_2$, respectively, so that $s_1(\alpha_2) = 2\alpha_1 + \alpha_2$ and $s_2(\alpha_1) = \alpha_1 + \alpha_2$. On the Cartan algebra for $\mathrm{SO}(5, \mathbb{C})$, these reflections correspond to $(h_1, h_2) \mapsto (h_1, -h_2)$ and $(h_1, h_2) \mapsto (h_2, h_1)$ for s_1 and s_2 , respectively.

The lifted maps on the Cartan subgroup SO_5^0 in the form given in (8.25) are implemented by conjugation with the following representatives of s_1 and s_2 as elements of the geometric Weyl group in the normalizer of SO_5^0 in $\mathrm{SO}(5, \mathbb{C})$.

$$\dot{s}_1 = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 1 & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix} \quad \text{and} \quad \dot{s}_2 = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 1 & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix}. \tag{8.29}$$

Representatives $\dot{s} \in N(\mathrm{SO}_5^0)$ for other elements $s \in \mathcal{W}$ are easily worked out from those in (8.29). This allows us to define the parabolic and unipotent Bruhat subgroups for $\mathrm{SO}(5, \mathbb{C})$.

$$\mathrm{SO}_5^{\geq 0}(s) = \mathrm{SO}_5^{\geq 0} \cap \dot{s} \mathrm{SO}_5^{\geq 0} \dot{s}^{-1} \quad \text{and} \quad \mathrm{SO}_5^+(s) = \mathrm{SO}_5^+ \cap \dot{s} \mathrm{SO}_5^+ \dot{s}^{-1} \tag{8.30}$$

Clearly, $\mathrm{SO}_5^{\geq 0}(s) \cong \mathrm{SO}_5^0 \ltimes \mathrm{SO}_5^+(s)$. As before, we define for any $s \in \mathcal{W}$ a vanishing ideal for the Bruhat subgroup as

$$\mathcal{I}_5^{\mathrm{SO}}(s) = \{f \in \mathbb{C}[\mathrm{SO}_5^{\geq 0}] : f(h) = 0 \ \forall h \in \mathrm{SO}_5^{\geq 0}(s)\}, \quad (8.31)$$

which is a Hopf ideal by Lemma 7.1. For $s \in \mathcal{W}$ we also introduce the ideal in $\mathbb{C}[\mathrm{SO}_5^{\geq 0}]$ generated by subsets of coordinate functions of $\mathbb{C}[\mathrm{SO}_5^{\geq 0}]$ that have grading in $\mathcal{N}(s)$. That is,

$$\mathcal{J}(s) = (G(s)) \quad \text{where} \quad G(s) = \{\hat{y} \in \{\hat{a}, \hat{b}, \hat{c}, \hat{d}\} : \mathbf{w}(\hat{y}) \in \mathcal{N}(s)\}. \quad (8.32)$$

For reference and convenience, we list all the non-trivial cases here explicitly.

$$\begin{array}{lll} \mathcal{J}(s_1) = (\hat{a}) & \mathcal{J}(s_1 s_2) = (\hat{a}, \hat{d}) & \mathcal{J}(s_1 s_2 s_1) = (\hat{a}, \hat{c}, \hat{d}) \\ \mathcal{J}(s_2) = (\hat{b}) & \mathcal{J}(s_2 s_1) = (\hat{b}, \hat{c}) & \mathcal{J}(s_2 s_1 s_2) = (\hat{b}, \hat{c}, \hat{d}) \end{array} \quad (8.33)$$

The following can be verified immediately from the coproduct formulae in (8.27) in each of the cases listed in (8.33).

Lemma 8.5. *For any $s \in \mathcal{W}$ the ideal $\mathcal{J}(s)$ is a Hopf ideal.*

The two types of Hopf ideals are identified next.

Proposition 8.6. *For any $s \in \mathcal{W}$ we have $\mathcal{J}(s) = \mathcal{I}_5^{\mathrm{SO}}(s)$.*

Proof. First, observe that the elements in (8.29) also represent elements of the Weyl group of the ambient $\mathrm{GL}(5, \mathbb{C})$, which we identify with S_5 . More precisely, consider the Cayley embedding $\iota : \mathcal{W} \hookrightarrow S_5$ defined by $\iota(s_1) = (2\ 4)$ and $\iota(s_2) = (1\ 2)(4\ 5)$, mapping the longest element $s_\bullet = (s_1 s_2)^2$ in \mathcal{W} to the longest element $\iota(s_\bullet) = (1\ 5)(2\ 4)$ in S_5 . Thus, if $\pi = \iota(s)$, we have that \dot{s} is given by the ordinary permutation matrix $\bar{\pi}$ for π in $\mathrm{GL}(5, \mathbb{C})$ up to sign changes of the 1-entries.

Next, observe that, because of $\mathrm{SO}_5^{\geq 0} = T_5 \cap \mathrm{SO}(5, \mathbb{C})$, we have $\mathrm{SO}_5^{\geq 0}(s) = \mathrm{SO}_5^{\geq 0} \cap \dot{s} \mathrm{SO}_5^{\geq 0} \dot{s}^{-1} = T_5 \cap \dot{s} T_5 \dot{s}^{-1} \cap \mathrm{SO}(5, \mathbb{C}) = T_5 \cap \bar{\pi} T_5 \bar{\pi}^{-1} \cap \mathrm{SO}(5, \mathbb{C})$. Hence,

$$\mathrm{SO}_5^{\geq 0}(s) = B(\iota(s)) \cap \mathrm{SO}(5, \mathbb{C}) \quad \text{and} \quad \mathrm{SO}_5^+(s) = U(\iota(s)) \cap \mathrm{SO}(5, \mathbb{C}),$$

with analogous relations for the unipotent parts. For $\pi \in S_5$ let

$$\mathrm{Inv}(\pi) = \{(i, j) \in \mathbb{N}^2 : 1 \leq i < j \leq 5 \text{ and } \pi^{-1}(i) > \pi^{-1}(j)\}.$$

Recall from the proof of Theorem 7.16 that $L \in B_5(\pi)$ iff $L_{i,j} = 0$ whenever $(i, j) \in \mathrm{Inv}(\pi)$.

Thus, for a given $s \in \mathcal{W}$ and setting $N(s) = \mathrm{Inv}(\iota(s))$, any $(i, j) \in N(s)$ imposes a constraint on the parameters in $(\mathbb{C}^\times)^2 \times \mathbb{C}^4$ in the parametrization of $\mathrm{SO}_5^{\geq 0}$, given in (8.25) by setting the (i, j) coefficient to zero. The corresponding constraints $\phi(y)_{i,j} = 0$ on elements $y \in (\mathbb{C}^\times)^2 \times \mathbb{C}^4$ then define a subvariety in $(\mathbb{C}^\times)^2 \times \mathbb{C}^4$, for which the vanishing ideal can be expressed in terms of the generating coordinate functions.

We apply this method next to verify all six non-trivial cases above. One can readily work out $N(s_1) = \{(2, 3), (2, 4), (3, 4)\}$ and $N(s_2) = \{(1, 2), (4, 5)\}$ for the generators. Setting the respective

matrix coefficients to zero yields the constraint $a = 0$ for the former and $b = 0$ for the latter. The vanishing ideals are thus (\hat{a}) and (\hat{b}) , respectively.

For the first length 2 case, we have $N(s_1 s_2) = \{(1, 4), (2, 3), (2, 4), (2, 5), (3, 4)\}$, which imposes constraints that are equivalent to the condition $a = d = 0$. Similarly, we find $N(s_2 s_1) = \{(1, 2), (1, 3), (1, 5), (3, 5), (4, 5)\}$ for the second case, yielding $b = c = 0$. Finally, the length 3 constraints are given by the following data.

$$\begin{aligned} N(s_2 s_1 s_2) &= \{(1, 2), (1, 3), (1, 4), (1, 5), (2, 5), (3, 5), (4, 5)\} & \Rightarrow b = c = d = 0 \\ N(s_1 s_2 s_1) &= \{(1, 3), (1, 4), (1, 5), (2, 3), (2, 4), (2, 5), (3, 4), (3, 5)\} & \Rightarrow a = c = d = 0, \end{aligned}$$

which implies the remaining identifications of ideals. \square

In order to better match the Hopf algebra structure of $\mathbb{C}[\mathrm{SO}_5^{>0}]$ with its quantum group counterpart, it is convenient to introduce a modified set of generators. For a given parameter t , we define them by the following polynomial expressions.

$$\begin{aligned} \bar{g} &= \hat{\lambda}^{-1} \hat{\mu} & \bar{a} &= \hat{a} & \bar{c}_t &= \hat{c} + t \hat{\lambda}^{-1} \hat{a} \hat{b} \\ \bar{h} &= \hat{\mu}^{-1} & \bar{b} &= \hat{\lambda}^{-1} \hat{b} & \bar{d}_t &= \hat{d} + (\frac{1}{2} - t) \hat{a} \hat{c} - \frac{1}{2} t^2 \hat{\lambda}^{-1} \hat{a}^2 \hat{b} \end{aligned} \quad (8.34)$$

It is easy to check that the generators in $\{\hat{\lambda}^{\pm 1}, \hat{\mu}^{\pm 1}, \hat{a}, \hat{b}, \hat{c}, \hat{d}\}$ are, conversely, given as polynomial expressions in the $\{\bar{g}^{\pm 1}, \bar{h}^{\pm 1}, \bar{a}, \bar{b}, \bar{c}_t, \bar{d}_t\}$, which implies an isomorphism between the corresponding polynomial algebras for each parameter t . The assignment is, further, grading preserving in the sense that $w(\bar{a}) = \alpha_1$, $w(\bar{b}) = \alpha_2$, $w(\bar{c}_t) = \alpha_1 + \alpha_2$, and $w(\bar{d}_t) = 2\alpha_1 + \alpha_2$.

Next, observe that the ideals in (8.33) are unchanged if the new generators are used instead. That is, we have, for example, $\mathcal{J}(s_1 s_2) = (\bar{a}, \bar{d}_t)$ and $\mathcal{J}(s_2 s_1 s_2) = (\bar{a}, \bar{c}_t, \bar{d}_t)$. The coproducts for the generators in (8.34) are determined from those in (8.27) via an elementary but somewhat lengthy computation.

$$\begin{aligned} \Delta(\bar{g}) &= \bar{g} \otimes \bar{g} & \Delta(\bar{h}) &= \bar{h} \otimes \bar{h} & \Delta(\bar{a}) &= \bar{a} \otimes \bar{h} + 1 \otimes \bar{a} \\ \Delta(\bar{b}) &= \bar{b} \otimes \bar{g} + 1 \otimes \bar{b} & \Delta(\bar{c}_t) &= \bar{c}_t \otimes \bar{h} \bar{g} + 1 \otimes \bar{c}_t + t \bar{b} \otimes \bar{g} \bar{a} - (1 - t) \bar{a} \otimes \bar{h} \bar{b} \\ \Delta(\bar{d}_t) &= \bar{d}_t \otimes \bar{h}^2 \bar{g} + 1 \otimes \bar{d}_t + (1 - t) \bar{a} \otimes \bar{h} \bar{c}_t - \frac{1}{2} (1 - t)^2 \bar{a}^2 \otimes \bar{h}^2 \bar{b} - t \bar{c}_t \otimes \bar{h} \bar{g} \bar{a} - \frac{1}{2} t^2 \bar{b} \otimes \bar{g} \bar{a}^2 \end{aligned} \quad (8.35)$$

An important feature of this form is that in each of the tensor factors only one type of non-Cartan generator occurs, avoiding mixed terms such as $\bar{a} \otimes \bar{a} \bar{b}$ or $\bar{a} \bar{c}_t \otimes 1$. The two relevant parameter choices are $t = 1$ and $t = 0$, for which terms in the coproducts of \bar{c}_t and \bar{d}_t vanish.

The proposition below summarizes our findings, relevant to the identifications with quantum algebras. Here, it is useful to introduce for a parameter t the free abstract polynomial algebra \mathcal{M}_t with generators $\{\bar{g}^{\pm 1}, \bar{h}^{\pm 1}, \bar{a}, \bar{b}, \bar{c}_t, \bar{d}_t\}$ with the usual commutation and inverse relation. The algebra \mathcal{M}_t is equipped with the coalgebra structure defined in (8.35), which uniquely extends to a Hopf algebra structure.

Proposition 8.7. *The following statements hold for any choice of t .*

- i) $\mathbb{C}[\mathrm{SO}_5^{\geq 0}]$ is isomorphic to \mathcal{M}_t as a \mathbb{Z}^Δ -graded Hopf algebra.
- ii) Any subgroup of $\mathrm{SO}_5^{\geq 0}$ defines a Hopf ideal in \mathcal{M}_t .
- iii) The Hopf ideal corresponding to a Bruhat subgroup $\mathrm{SO}_5^{\geq 0}(s)$ coincides with the ideal in \mathcal{M}_t generated by the set $\{\bar{\gamma} \in \{\bar{a}, \bar{b}, \bar{c}_t, \bar{d}_t\} : \mathbf{w}(\bar{\gamma}) \in \mathcal{N}(s)\}$.

The isomorphism in *i*), of course, implies that the \mathcal{M}_t are isomorphic to each other for different t . The proof of the statement in *ii*) is the same as the one for Lemma 7.15. As noted before, Item *i*) follows readily from Proposition 8.6 and inspection of the change of generators in (8.34).

8.4. Isomorphisms between Hopf Algebras related to B_2 . The weight grading of either $\mathcal{Z}_\bullet^{\geq 0}$ or $\mathbb{C}[\mathrm{SO}_5^{\geq 0}]$ implies a sequence of (commutative) Hopf subalgebras as in the definition of cosolvability at the end of Section 7.1. This facilitates classifications of weight preserving coalgebra structures, as they can be viewed as successive extensions of each other. We develop here a restricted classification of coalgebra structures associated to the B_2 root system, for which the employed methods are likely to generalize to much larger classes of graded, cosolvable Hopf algebras.

The underlying associative algebra is, as before, of the form $\mathbb{C}[\tilde{g}^{\pm 1}, \tilde{h}^{\pm 1}, \tilde{a}, \tilde{b}, \tilde{c}, \tilde{d}]$ with weights $\mathbf{w}(\tilde{g}) = \mathbf{w}(\tilde{h}) = 0$, $\mathbf{w}(\tilde{a}) = \alpha_1$, $\mathbf{w}(\tilde{b}) = \alpha_2$, $\mathbf{w}(\tilde{c}) = \alpha_1 + \alpha_2$, and $\mathbf{w}(\tilde{d}) = 2\alpha_1 + \alpha_2$. Aside from preserving the \mathbf{w} -grading, we require \tilde{a} and \tilde{b} to be skew-primitive with respect to group-likes $(1, \tilde{h})$ and $(1, \tilde{g})$.

We further constrain the allowable coproducts by requiring lead coefficients to be 1 and that the group-likes in the second tensor-factor are determined by the weight of the first tensor-factor as, for example, in (4.7). Moreover, as discussed in (8.35), we omit terms with mixed monomials in tensor-factors as these do not occur or can be removed in the previous examples. In this restricted setting we find six allowable terms besides the lead terms, with coefficients given by six parameters (u, v, x, y, r, s) as follows.

$$\begin{aligned} \Delta(\tilde{g}) &= \tilde{g} \otimes \tilde{g} & \Delta(\tilde{h}) &= \tilde{h} \otimes \tilde{h} & \Delta(\tilde{a}) &= \tilde{a} \otimes \tilde{h} + 1 \otimes \tilde{a} & \Delta(\tilde{b}) &= \tilde{b} \otimes \tilde{g} + 1 \otimes \tilde{b} \\ \Delta(\tilde{c}) &= \tilde{c} \otimes \tilde{h}\tilde{g} + 1 \otimes \tilde{c} + u \cdot \tilde{a} \otimes \tilde{h}\tilde{b} + v \cdot \tilde{b} \otimes \tilde{g}\tilde{a} & & & & & & (8.36) \\ \Delta(\tilde{d}) &= \tilde{d} \otimes \tilde{h}^2\tilde{g} + 1 \otimes \tilde{d} + x \cdot \tilde{a} \otimes \tilde{h}\tilde{c} + y \cdot \tilde{c} \otimes \tilde{h}\tilde{g}\tilde{a} + r \cdot \tilde{a}^2 \otimes \tilde{h}^2\tilde{b} + s \cdot \tilde{b} \otimes \tilde{g}\tilde{a}^2 & & & & & & \end{aligned}$$

The axiom of coassociativity now imposes additional conditions on these parameters. It is easy to see that for the subalgebra in which \tilde{d} is omitted as a generator, the corelations in (8.36) define a coalgebra structure for any choices of u and v . Testing coassociativity for \tilde{d} yields non-trivial constraints only for the comparisons of the $\tilde{a} \otimes \tilde{h}\tilde{a} \otimes \tilde{h}^2\tilde{b}$, $\tilde{a} \otimes \tilde{h}\tilde{b} \otimes \tilde{h}\tilde{g}\tilde{a}$, and $\tilde{b} \otimes \tilde{g}\tilde{a} \otimes \tilde{h}\tilde{g}\tilde{a}$ terms. The existence of an antipode for any bialgebra structure is immediate.

Lemma 8.8. *The formulae in (8.36) define a Hopf algebra structure on $\mathbb{C}[\tilde{g}^{\pm 1}, \tilde{h}^{\pm 1}, \tilde{a}, \tilde{b}, \tilde{c}, \tilde{d}]$ iff*

$$vx = uy, \quad s = \frac{1}{2}vy, \quad \text{and} \quad r = \frac{1}{2}ux. \quad (8.37)$$

In particular, the parameters r and s are determined by the four others. We, thus, denote by $\mathcal{M}(u, v, x, y)$ the Hopf algebra defined by the coproducts in (8.36) subject to the parameter constraints in (8.37). The obvious generator to generator map yields the following isomorphism with the Hopf algebra $\text{SO}_5^{\geq 0}$ as presented in (8.35).

$$\mathcal{M}_t \cong \mathcal{M}(t-1, t, 1-t, -t) \quad (8.38)$$

We next determine isomorphism classes within this family of Hopf algebras. We require an isomorphism to be polynomial, weight-preserving, as well as preserving the general structure in (8.36). It is an easy exercise to show that such a transformation must be the identity on group-likes and maps the subring $\mathbb{C}[\tilde{a}, \tilde{b}, \tilde{c}, \tilde{d}]$ of non-group-like generators to itself. Hence, an isomorphism $\varphi : \mathcal{M}(u, v, x, y) \rightarrow \mathcal{M}(u', v', x', y')$ necessarily acts on the simple root generators as $\varphi(\tilde{a}) = v_1\tilde{a}$ and $\varphi(\tilde{b}) = v_2\tilde{b}$ for $v_1, v_2 \in \mathbb{C}^\times$. Moreover, for any such pair (v_1, v_2) , there is an automorphism of the same $\mathcal{M}(u, v, x, y)$ to itself. So, without loss of generality, we may assume φ is identity on the Hopf subalgebra $\mathcal{M}^0 = \mathbb{C}[\tilde{h}^{\pm 1}, \tilde{g}^{\pm 1}, \tilde{a}, \tilde{b}]$.

It follows that $\varphi(\tilde{c})$ is a \mathbb{C} -linear combination of \tilde{c} and $\tilde{a}\tilde{b}$, and $\varphi(\tilde{d})$ is a \mathbb{C} -linear combination of \tilde{d} , $\tilde{a}\tilde{c}$, and $\tilde{a}^2\tilde{b}$. The two diagonal coefficients need to be non-zero in order for φ to be invertible. Two additional conditions on the five coefficients of φ stem from the requirement to preserve the form in (8.36) and the relations in (8.37), each depending also on the parameters (u, v, x, y) of the source algebra.

One can then work out that an isomorphism $\varphi : \mathcal{M}(u, v, x, y) \rightarrow \mathcal{M}(u', v', x', y')$ exists iff $(u', v') = \sigma(u, v) + \eta(1, 1)$ and $(x', y') = \mu(x, y) + \tau(1, 1)$ for parameters $\sigma, \mu \in \mathbb{C}^\times$ and $\eta, \tau \in \mathbb{C}$ with $\mu\eta(x - y) = \sigma\tau(u - v)$. This now classifies Hopf algebras of the above form.

Proposition 8.9. *There exist weight-preserving isomorphisms of Hopf algebras as follows.*

$$\mathcal{M}(u, v, x, y) \cong \begin{cases} \mathcal{M}(1, 0, 1, 0) & \text{if } u \neq v \text{ and } x \neq y \\ \mathcal{M}(1, 0, 0, 0) & \text{if } u \neq v \text{ and } x = y \\ \mathcal{M}(0, 0, 1, 0) & \text{if } u = v \text{ and } x \neq y \\ \mathcal{M}(0, 0, 0, 0) & \text{if } u = v \text{ and } x = y \end{cases}$$

Furthermore, there are no weight-preserving isomorphisms of the required form between any of the four listed special cases.

Observe that $(u, v) = (0, 0)$ means that \tilde{c} is skew-primitive and $(x, y) = (0, 0)$ that \tilde{d} is skew-primitive. Given \mathcal{M}^0 as before, denote also by \mathcal{M}^1 the Hopf subalgebra with additional generator \tilde{c} and $\mathcal{M}^2 = \mathcal{M}(u, v, x, y)$ the full algebra. The four cases above may then be characterized by whether the extension from \mathcal{M}^0 to \mathcal{M}^1 as well as the one \mathcal{M}^1 to \mathcal{M}^2 is trivial (in the sense that only an independent skew-primitive is added) or non-trivial.

Note, finally, that $u = v \neq 0$ implies by (8.37) that $x = y$. So, the third case in Proposition 8.9 can occur only if $u = v = 0$. Analogously, the second case implies $x = y = 0$. We refer to the first version as the *generic* case. Clearly, $\mathbb{C}[\mathrm{SO}_5^{\geq 0}]$ is an example by (8.38) and Proposition 8.7.

The classification immediately entails the identifications of the Hopf algebras and Hopf ideals summarized below. For an element $s \in \mathcal{W}$, we denote here by $\check{s} \in \mathcal{W}$ the action of s on the coroot system $\check{\Phi}^+$ conjugated by the identification of $\check{\Phi}^+$ with Φ^+ discussed in the beginning of Section 8.1. So, $s \mapsto \check{s}$ is the group homomorphism given by $\check{s}_1 = s_2$ and $\check{s}_2 = s_1$.

Theorem 8.10. *Suppose $U_\zeta^{\geq 0}$ is of Lie type $\mathrm{B}_2 = \mathrm{C}_2$ with ζ a root of unity of any order $\mathbb{k} > 4$ or $\mathbb{k} = 3$.*

- i) $\mathcal{Z}_\bullet^{\geq 0} \otimes \mathbb{C}$ is isomorphic as a Hopf algebra to $\mathbb{C}[\mathrm{SO}_5^{\geq 0}]$.
- ii) Any subgroup $H < \mathrm{SO}_5^{\geq 0}$ yields Hopf ideals $\mathcal{A}(H)$ and $\widehat{\mathcal{A}}(H)$ of $\mathcal{Z}_\bullet^{\geq 0}$ and $U_\zeta^{\geq 0}$, respectively.
- iii) For ℓ odd, $\mathcal{A}(\mathrm{SO}_5^{\geq 0}(s)) = \mathcal{K}(s)^{\geq 0}$ and, for ℓ even, $\mathcal{A}(\mathrm{SO}_5^{\geq 0}(\check{s})) = \mathcal{K}(s)^{\geq 0}$ for any $s \in \mathcal{W}$.

Proof. Remark 6.15 implies that $\mathcal{Z}_\bullet^{\geq 0}$ is commutative for any root of unity and, thus, isomorphic to the polynomial ring $\mathbb{C}[L_i^{\pm 1}, L_j^{\pm 1}, \{X_\alpha : \alpha \in \Phi^+\}]$. We next note that the coalgebra structure from Proposition 8.1 is indeed of the form given in (8.36).

Specifically, if ℓ is odd ($\mathfrak{n} = 2$ and $\mathfrak{d} = 1$) assign $\tilde{h} \mapsto L_i$, $\tilde{g} \mapsto L_j$, $\tilde{a} \mapsto X_i$, $\tilde{b} \mapsto X_j$, $\tilde{c} \mapsto X_{(ij)}$, and $\tilde{d} \mapsto X_{(ij)}$. Comparison shows that the coalgebra structure coincides with the one of $\mathcal{M}(u, 0, x, 0)$, where $u \neq 0 \neq x$ are given by the coefficients in (8.8) and (8.7). Thus, $\mathcal{Z}_\bullet^{\geq 0}$ coincides with the generic case in Proposition 8.9 and is, therefore, isomorphic to $\mathbb{C}[\mathrm{SO}_5^{\geq 0}]$.

Conforming with the \mathfrak{u} -grading from (8.5), the assignments for even ℓ ($\mathfrak{n} = 1$ and $\mathfrak{d} = 2$) are $\tilde{h} \mapsto L_j$, $\tilde{g} \mapsto L_i$, $\tilde{a} \mapsto X_j$, $\tilde{b} \mapsto X_i$, $\tilde{c} \mapsto X_{(ij)}$, and $\tilde{d} \mapsto X_{(iji)}$. This yields a Hopf algebra isomorphism from $\mathbb{C}[\mathrm{SO}_5^{\geq 0}]$ with $\mathcal{Z}_\bullet^{\geq 0}$ via the generic form $\mathcal{M}(0, v, 0, y)$ with $v \neq 0 \neq y$.

The construction of Hopf ideals is the same as in Lemma (7.34). The identifications of the ideals associated to the Bruhat groups follow from the construction of \mathfrak{u} -grading preserving isomorphisms and may also be checked from the explicit forms in (8.33). \square

Comparing (8.38) and the isomorphisms described in the above proof, we observe that for odd ℓ the non-zero terms occurring in the coproducts for $\mathcal{Z}_\bullet^{\geq 0}$ are the same as for \mathcal{M}_0 , and for even ℓ they coincide with those of \mathcal{M}_1 . Specifically,

$$\text{odd } \ell: \quad u = -\nabla_j \quad v = 0 \quad x = -2\zeta^\ell [2]^{-\ell} \nabla_i \quad y = 0 \quad (8.39)$$

$$\text{even } \ell: \quad u = 0 \quad v = \zeta^{\binom{\ell}{2}} \nabla_i \quad x = 0 \quad y = -2[2]^\ell (-\zeta)^\ell \quad (8.40)$$

Thus, in conclusion, we can provide explicit grading preserving maps given only by scalings of generators.

Theorem 8.10 confirms Conjecture 7.13 in the B_2 case and illustrates the need to switch to the coroot system for certain \mathbb{k} congruences. Furthermore, it is not difficult to construct and prove an analog statement of Lemma 7.18 for a respective subgroup $W(\mathrm{B}_2)$ of $\mathrm{SO}_5^{\leq 0} \times \mathrm{SO}_5^{\geq 0}$. As

for Corollary 7.19, Theorem 6.22 now implies an action of the finite-type dihedral Artin group $\mathcal{A} \cong \langle a, b \mid [a^2, b] \rangle$ on $W(\mathbf{B}_2)$.

We conclude our discussions of special cases in Section 7 and 8 with a list of conjectures, generalizing our findings for \mathbf{A}_n and \mathbf{B}_2 . As already mentioned above, the existence of an isomorphism for odd \mathbb{k} in items *i*) and *iii*) below has been proved in [DCKP92] via additional Poisson structures and not in the suggested explicit form. For a given simple Lie type Φ , denote by G the associated simply connected algebraic group and B a Borel algebra over \mathbb{C} . Denote also \check{G} and \check{B} the dual groups associated with the coroot lattice, as well as $G^\#$ and $\check{G}^\#$ as in (1.1).

Conjecture 8.11.

- i) In all the \mathbb{k} -congruence cases for which subalgebras are strictly commutative (Corollary 6.14), the algebra $\mathcal{Z}_\bullet^{\geq 0} \otimes \mathbb{C}$ is isomorphic to $\mathbb{C}[\check{B}]$, where \check{B} is B or \check{B} or a finite covering quotient (isogeny) of these. The isomorphism map can be chosen as an explicit assignment of generators X_w (with $w \leq_R z$ for some $z \in \mathcal{W}_\mathbf{m}^*$) to weighted polynomials of matrix elements.
- ii) For any $s \in \mathcal{W}$, the vanishing ideal of the Bruhat group $\check{B}(s)$, as defined in (7.35), is mapped exactly to the $\mathcal{K}(s)^{\geq 0} \otimes \mathbb{C}$ ideal.
- iii) The isomorphism from i) extends to an isomorphism from $\mathcal{Z}_\bullet \otimes \mathbb{C}$ to a respective coordinate ring of $G^\#$ or $\check{G}^\#$ or a respective finite quotient.
- iv) Lusztig's action of the Artin group \mathcal{A} on $\mathcal{Z}_\bullet \otimes \mathbb{C}$ from Theorem 6.22 coincides with the induced action of \mathcal{A} on $G^\#$ in [DCKP92, Sec 7.5] also for the cases when \mathbb{k} is even and $\mathcal{Z}_\bullet^{\geq 0}$ commutative.

9. HOPF IDEALS VIA QUASI- R -MATRICES

R -matrices are the primary motivation for studying quantum groups, as they give rise to solutions to Yang Baxter equations and, more generally, braided tensor categories. Their construction for quantum groups at roots of unity is one of the goals of this section.

As another application, we use here the R -matrix formalism in an iterative argument, showing that the $\hat{\mathcal{K}}(s)$ ideals are Hopf ideals. Their quotients, therefore, produce a family of well-defined Hopf algebras $U_{\zeta, \epsilon}^{[s]}$. The methods introduced and developed in this section also include truncated quasi- R -matrices, Tanisaki's formalism of pre-triangular Hopf algebras, and the Tanisaki-Lusztig pairings.

9.1. Enhanced Tensor Structures. In this section, we provide some general constructions relevant to defining R -matrices in the generic setting. The first helps to make sense of the infinite sums used in the formulae for R -matrices using certain inverse limits.

Generally, if A is an algebra over a ring Λ and I^n a descending sequence of *two-sided* ideals, $\varprojlim A/I^n$ admits a canonical algebra structure as an inverse system in the category of algebras.

For left (or right) ideals A^n , however, the spaces A/A^n are not algebras so that there is no obvious algebra structure on their limit. This can be remedied by an additional assumption, roughly stating that the A^n are right ideals up to a shift that is locally bounded.

Specifically, let $A = \bigoplus_{n \in \mathbb{N}_0} A_n$ and assume there is a function $\chi : \bigcup_n A_n \rightarrow \mathbb{N}_0$ such that

$$\text{for all } n, m \in \mathbb{N}_0, x \in A_n, y \in A_m \text{ we have } xy \in A^{m+\max(0, n-\chi(y))}, \quad (9.1)$$

where $A^n = \bigoplus_{k \geq n} A_k$. Denote $\hat{A} = \varprojlim_n A/A^n$. We shall call χ the *local bound* of A .

Clearly, by (9.1), the A^n are left (but not necessarily right) ideals, their intersection is zero, and we have the standard embedding $A \hookrightarrow \hat{A}$ as a Λ -module. For a second algebra B of this form with local bound $\gamma : \bigcup_n B_n \rightarrow \mathbb{N}_0$ one may consider degree preserving homomorphisms $f : A \rightarrow B$ in the sense that $f(A_n) \subseteq B_n$ as well as a grading of the tensor algebra $A \otimes B$ with components $(A \otimes B)_n = \bigoplus_s A_s \otimes B_{n-s}$.

The assumption of a decomposition into components A_n can be circumvented with further work, but fits our applications to quantum groups.

Lemma 9.1. *Let A, B be algebras and $f : A \rightarrow B$ a homomorphism, all as above.*

- i) *\hat{A} admits a unique algebra structure for which A is a subalgebra.*
- ii) *There is a unique algebra homomorphism $\hat{f} : \hat{A} \rightarrow \hat{B}$ that restricts to f on A .*
- iii) *$A \otimes B$ has the property (9.1) with local bound determined by $\eta(y \otimes y') = \chi(y) + \gamma(y')$. The respective completion canonically contains $\hat{A} \otimes \hat{B}$ as a subalgebra and gives rise to an associative completion $\hat{\otimes}$ of the tensor product as $\widehat{A \otimes B} = \hat{A} \hat{\otimes} \hat{B}$.*
- iv) *If A admits a bialgebra structure with coproduct $\Delta(A_n) \subseteq (A \otimes A)_n$, then \hat{A} admits a canonical bialgebra structure with respect to the completed tensor, in the sense that $\hat{\Delta} : \hat{A} \rightarrow \hat{A} \hat{\otimes} \hat{A}$.*

Proof. By definition, an element of the inverse limit \hat{A} is uniquely presented by a sequence $(x_0, x_1, \dots) \in \prod_{n \geq 0} A_n$, which is more suggestively written as a formal sum $\sum_{k \geq 0} x_k$. The submodule A is given by sequences with only finitely many non-zero terms. For two elements $\sum_n x_n, \sum_m y_m \in \hat{A}$, (9.1) implies that we can write

$$x_n y_m = \sum_{k \geq m, k + \chi(b_m) \geq n + m} z_k^{n,m} \quad \text{for some } z_k^{n,m} \in A_k.$$

For fixed k , the bounds imply $m \leq k$ and $n \leq k + r_k$ with $r_k = \max_{0 \leq m \leq k} \chi(b_m) - m$. Thus, $x_n y_m$ has a component in A_k only for a finite number of pairs (n, m) so that $(xy)_k = \sum_{n, m} z_k^{n,m}$ is well-defined. It is clear that the so defined product yields an algebra structure as proposed in Item i). Item ii) is then also immediate.

Suppose $x \in A_i, x' \in B_j, y \in A_s$, and $y' \in B_t$. Then the component of $(x \otimes x')(y \otimes y')$ = $xy \otimes x'y'$ in $(A \otimes B)_k$ is, by (9.1), applied to each tensor factor, zero if $k \leq (s + \max(0, i - \chi(y)) + (t + \max(0, j - \gamma(y')) \leq s + t + \max(0, (i + j - (\chi(y) + \gamma(y'))))$, which implies the claim for pure component

tensors. For a general tensor in $A_i \otimes B_j$, η may be defined as the smallest number d such that the tensor can be written as a sum of pure tensors, each of which with η -value less than d . For a general element in $(A \otimes B)_n$, take the maximum over the $A_s \otimes B_{n-s}$ graded components.

The embedding of $\hat{A} \otimes \hat{B}$ into $\hat{A} \hat{\otimes} \hat{B}$ is obvious, and standard properties of $\hat{\otimes}$ are readily checked. Item *iv*) is now a consequence of *ii*) and *iii*). \square

For an A -module V set $V_n = \{v \in V : A^n \cdot v = 0\}$, yielding an ascending chain of subspaces $0 = V_0 \subseteq V_1 \subseteq V_2 \subseteq \dots$. We say V is *locally finite* if $V = \bigcup_n V_n$. We collect several basic observations next.

Lemma 9.2. *Suppose V and W are locally finite A -modules as above.*

- i) *If $y \in A_k$ and $v \in V_n$ then $y \cdot v \in V_{n-k+\chi(y)}$.*
- ii) *The action of A extends to an action of \hat{A} on V .*
- iii) *The action of $A \otimes A$ on $V \otimes W$ is locally finite and, thus, extends to an action of $\hat{A} \hat{\otimes} \hat{A}$.*
- iv) *If A has a bialgebra structure as above, the locally finite modules form a tensor category.*

Proof. Item *i*) follows readily from the definitions, Item *ii*) holds since only finitely many terms are non-zero in any $\sum_k a_k v$, Item *iii*) is a consequence of $(A \otimes A)^{n+m}(V_n \otimes W_m) = 0$, and Item *iv*) is immediate from Item *iii*). \square

Observe that the category noted in the last statement does *not* need to be rigid. In fact, the locally finite condition above emulates the respective property for Verma modules (or, more generally, modules in category \mathcal{O}) for which standard duals do not exist. Thus, at the level of algebras, we generally do not expect the antipode on a Hopf algebra A to extend to the limit \hat{A} .

The second ingredient required to define a braided structure from formal series are Tanisaki's axioms [Tan92] that aim to reexpress the ill-defined diagonal parts of an R -matrix in terms of certain automorphisms.

Definition 9.3. *Let A be a bialgebra with structure as in (9.1), $\mathcal{F} \in \hat{A} \hat{\otimes} \hat{A}$ invertible, and f a component-preserving algebra automorphism of $A \otimes A$. We say (A, \mathcal{F}, f) is a pre-triangular bialgebra if the following hold:*

$$\begin{aligned} f_{12}(\mathcal{F}_{13})\mathcal{F}_{12} &= (\text{id} \otimes \Delta)(\mathcal{F}), & f_{23}(\mathcal{F}_{13})\mathcal{F}_{23} &= (\Delta \otimes \text{id})(\mathcal{F}), \\ \text{and} \quad \mathcal{F}\Delta(x) &= (f \circ \Delta^{\text{opp}}(x))\mathcal{F}, & \text{for all } x \in A. \end{aligned} \tag{9.2}$$

The notation in (9.2) omits the $\hat{\cdot}$ accent on automorphisms and coproducts. Since f is assumed to preserve components, such an extension to $\hat{A} \hat{\otimes} \hat{A}$ exists. Similarly, one checks that, for example, $\hat{\Delta} \hat{\otimes} \text{id} : \hat{A} \hat{\otimes}^2 \rightarrow \hat{A} \hat{\otimes}^3$ is well-defined. The subscripts for \mathcal{F} indicate the usual canonical inclusions extended to limits $\hat{A} \hat{\otimes}^2 \hookrightarrow \hat{A} \hat{\otimes}^3$ and the subscripts on the automorphism mean, for example, $f_{23} = \text{id} \hat{\otimes} \hat{f}$.

Assume a pre-triangular bialgebra A with properties as in (9.1) as well as a tensor (sub)category \mathcal{M} of locally finite modules. We say (A, \mathcal{F}, f) is *implementable* in \mathcal{M} if for any two modules $V, W \in \mathcal{M}$ there is an isomorphism $D_{V,W} : V \otimes W \rightarrow V \otimes W$ such that for all $V, W, Z \in \mathcal{M}$ the following hold.

$$\begin{aligned} f(c) &= D_{V,W}^{-1} c D_{V,W} \quad \text{for all } c \in A \otimes A \text{ acting on } V \otimes W \\ D_{V \otimes W, Z} &= (T \otimes \text{id})(\text{id} \otimes D_{V,Z})(T \otimes \text{id})(\text{id} \otimes D_{W,Z}) \\ D_{V,W \otimes Z} &= (T \otimes \text{id})(\text{id} \otimes D_{V,Z})(T \otimes \text{id})(D_{V,W} \otimes \text{id}) \end{aligned} \quad (9.3)$$

Here T denotes the ordinary transposition of tensor factors. We suppress notation for the representation map $A \rightarrow \text{End}(V)$ or $A \otimes A \rightarrow \text{End}(V \otimes W)$ whenever the respective actions are clear. A routine verification of categorical triangle (hexagon) relations yields the next observation, which is central to many applications of quantum groups.

Corollary 9.4. *If (A, \mathcal{F}, f) is implementable in \mathcal{M} , then the natural isomorphism*

$$R(V, W) = TD_{V,W}\mathcal{F} : V \otimes W \rightarrow W \otimes V$$

defines a braiding on \mathcal{M} .

An immediate consequence is the existence of representations of the braid group B_n on $V^{\otimes n}$. The axioms in [Tan92] assume also relations $f_{23} \circ f_{13}(\mathcal{F}_{12}) = \mathcal{F}_{12}$ and $f_{13} \circ f_{23}(\mathcal{F}_{23}) = \mathcal{F}_{23}$ in addition to those in (9.2) but omit the relations for $D_{V \otimes W, Z}$ and $D_{V,W \otimes Z}$ in (9.3). This leads to the same braid representations without the categorical context.

9.2. Completions and Quasi- R -Matrices for U_q . We will construct the quasi- R -matrices for $U_{q,\mathbb{Q}} = U_q \otimes \mathbb{Q}(q)$ from simple quasi- R -matrices, *a priori* given by the formal expressions

$$\mathcal{P}_i = \sum_{s=0}^{\infty} \frac{q_i^{s(s-1)/2}}{[s]!_i} (q_i - q_i^{-1})^s E_i^s \otimes F_i^s = \sum_{s=0}^{\infty} q_i^{s(s-1)/2} (-1)^s \check{E}_i^s \otimes F_i^{(s)} \quad \in \hat{U}_{q,\mathbb{Q}} \hat{\otimes} \hat{U}_{q,\mathbb{Q}} \quad (9.4)$$

for any $i = 1, \dots, n$. The renormalized generators in the second expression are as in (5.9) and (5.10). Several limits $\hat{U}_{q,\mathbb{Q}}$ and completions $\hat{\otimes}$ may be considered to provide a rigorous meaning to (9.4). The ones relevant here start from the \mathbb{N}_0 -gradings w^+ and w^- on U_q^+ and U_q^- introduced in Section 4.1. The height function from Section 2.5 extends to a homomorphism $ht : \mathbb{N}_0^\Delta \rightarrow \mathbb{N}_0$, which, thus, entails respective integer-gradings $ht \circ w^\pm$. We denote the corresponding components as

$$U_{q,\mathbb{Q}}^{k,+} = \bigoplus_{ht(\mu)=k} (U_{q,\mathbb{Q}}^+)_\mu^{w^+} \quad \text{and} \quad U_{q,\mathbb{Q}}^{k,-} = \bigoplus_{ht(\mu)=k} (U_{q,\mathbb{Q}}^-)_\mu^{w^-}. \quad (9.5)$$

Also set $U_{q,\mathbb{Q}}^{\geq k, \pm} = \bigoplus_{s \geq k} U_{q,\mathbb{Q}}^{s, \pm}$. The completion suggested, for example, in [CP94] fits the construction of Section 9.1 with

$$A_n = U_{q,\mathbb{Q}} \cdot U_{q,\mathbb{Q}}^{n,+} \quad \text{and} \quad A^n = U_{q,\mathbb{Q}} \cdot U_{q,\mathbb{Q}}^{\geq n,+}. \quad (9.6)$$

Note that, by the PBW theorems, $A_n = \bigoplus_{t \in \mathbb{N}_0} A_{t,n}$, where $A_{t,n} = U_{q,\mathbb{Q}}^{t,-} U_{q,\mathbb{Q}}^0 U_{q,\mathbb{Q}}^{n,+}$. A function $\chi : A_n \rightarrow \mathbb{N}_0$ is now defined by setting $\chi(y) = m$ to be the smallest m such that $y \in \bigoplus_{t \leq m} A_{t,n}$.

Lemma 9.5. *The A_n and χ , as defined above, fulfill (9.1). Thus, $\hat{U}_{q,\mathbb{Q}} = \varprojlim U_{q,\mathbb{Q}}/A^n$ is a bialgebra with $\hat{\otimes}$ as in Lemma 9.1.*

Proof. Set $B_{r,t} = U_{q,\mathbb{Q}}^{r,+} \cdot U_{q,\mathbb{Q}}^{t,-}$. We first observe that $B_{r,t} \subseteq \bigoplus_{j=0}^{\min(r,t)} A_{t-j,r-j}$. This follows by induction on expressions $E_{u_1} \dots E_{u_r} F_{v_1} \dots F_{v_t}$ that span $B_{r,t}$, using the basic commutation relations in (3.13) and (3.14). Multiplying this inclusion from the left with $U_{q,\mathbb{Q}}^0$ and from the right with $U_{q,\mathbb{Q}}^0 \cdot U_{q,\mathbb{Q}}^{m,+}$ we find $A_r \cdot A_{t,m} \subseteq \bigoplus_{j=0}^{\min(t,r)} U_{q,\mathbb{Q}}^0 \cdot A_{t-j,r-j} \cdot U_{q,\mathbb{Q}}^{m,+} = \bigoplus_{j=0}^{\min(t,r)} A_{r+m-j} \subseteq A^{r+m-\min(t,r)} = A^{m+\min(r-t,0)}$, which implies (9.1). Since the coproduct on $U_{q,\mathbb{Q}}^{\geq 0}$ preserves the w^+ -grading, the additional assumption in Lemma 9.1 holds as well, leading to a well-defined bialgebra structure on the limit with respect to the completion $\hat{\otimes}$. \square

The locally finite representations for this completion encompass all BGG-Category \mathcal{O} modules as in [AM15]. This includes infinite-dimensional Verma modules, for which the entailed braid representations are of interest, as they relate to geometrically constructed ones [JK11].

To see that the antipode does not extend to the limit, consider the formal element $x = \sum_{m \geq 0} F^m E^m$ in $\hat{U}_q(\mathfrak{sl}_2)$. From (3.18) we find $S(F^m E^m) = E^m F^m = \sum_{j=0}^m c_j^m F^j E^j$ with $c_j^m \in U_q^0$. In particular, $c_0^m = [m]! \prod_{s=0}^{m-1} \frac{Kq^{-s} - K^{-1}q^s}{q - q^{-1}}$ leading to ill-defined polynomials of infinite order in the 0-component of $S(x)$.

We do, however, have a well-defined extension of the antipode to the respective limits of Borel subalgebras $\hat{U}_{q,\mathbb{Q}}^{\geq 0}$ and $\hat{U}_{q,\mathbb{Q}}^{\leq 0}$.

As an alternative to the above limit, we may consider, for example, the two-sided ideals $I^{n,+} = \langle U_{q,\mathbb{Q}}^{n,+} \rangle$ generated by elements in $U_{q,\mathbb{Q}}^{n,+}$. It is not hard to see that $\Delta(I^{n,+}) \subseteq \bigoplus_k I^{n-k,+} \otimes I^{k,+}$. Thus, the inverse limit $\tilde{U}_{q,\mathbb{Q}} = \varprojlim U_q/I^{n,+}$ yields again a bialgebra structure and tensor-completion for which (9.4) is well-defined.

As opposed to the previous situation, the algebra structure of $\tilde{U}_{q,\mathbb{Q}}$ here follows trivially from the quotients, which are algebras. Moreover, the antipode is well-defined on the limit. However, the set of $U_{q,\mathbb{Q}}$ -modules to which $\tilde{U}_{q,\mathbb{Q}}$ extends is much smaller. Specifically, for any cyclic submodule W there needs to exist some n such that $U_{q,\mathbb{Q}}^{n,+}$ vanishes on all of W . Thus, the only allowed highest weight modules are finite-dimensional, which excludes, for example, all Verma modules.

Using (3.4) with $z = -q^2$ one can check by direct computation that the following expression is an inverse in $\hat{U}_{q,\mathbb{Q}} \hat{\otimes} \hat{U}_{q,\mathbb{Q}}$.

$$\mathcal{P}_i^{-1} = \sum_{s=0}^{\infty} \frac{q_i^{-s(s-1)/2}}{[s]!_i} (-1)^s (q_i - q_i^{-1})^s E_i^s \otimes F_i^s \quad (9.7)$$

Note that the braid automorphisms from Sections 4.3 also do *not* extend to the full completion $\hat{U}_{q,\mathbb{Q}}$. Specifically, in rank 1, $\Gamma_1(x)$ is ill-defined on the same element $x = \sum_{m \geq 0} F^m E^m$ for the same reason as for the anitpode above. The automorphisms are, however, well-defined as linear operators on certain linear subspaces in $\hat{U}_{q,\mathbb{Q}}$, whose construction we outline next.

In the notation of Section 5.3, observe that for fixed $z \in \mathcal{W}_m^*$ and $k \in \mathbb{N}_0$, there are only finitely many $\psi \in \mathcal{E}_*$ with $n(\psi) = \mathbf{w}(E_{\blacktriangleleft z}^\psi) = k$. Thus, an element in $\hat{U}_{q,\mathbb{Q}}$ can be written uniquely as a formal sum $\sum_{\psi \in \mathcal{E}_*} R_\psi E_{\blacktriangleleft z}^\psi$ with $R_\psi \in U_{q,\mathbb{Q}}^{\leq 0}$. Similarly, an element in $\hat{U}_{q,\mathbb{Q}} \hat{\otimes} \hat{U}_{q,\mathbb{Q}}$ is formally given by $\sum_{\phi, \psi \in \mathcal{E}_*} R_{\phi, \psi} (E_{\blacktriangleleft z}^\phi \otimes E_{\blacktriangleleft z}^\psi)$ with $R_{\phi, \psi} \in (U_{q,\mathbb{Q}}^{\leq 0})^{\otimes 2}$.

As in (5.38), denote by $\mathfrak{S}_{\blacktriangleleft w}^-$ the span of $F_{\blacktriangleleft w}^\psi$ for a (not necessarily maximal) reduced word $w \in \mathcal{W}^*$, $s = r(w)$, and $\psi \in \mathcal{E}(s)$. Next, for $s = r(w)$, define the linear subspace of formal sums

$$\mathring{\mathcal{L}}_{\blacktriangleleft w} = \left\{ \sum_{\psi \in \mathcal{E}(s)} R_\psi E_{\blacktriangleleft w}^\psi : R_\psi \in \mathfrak{S}_{\blacktriangleleft w}^- \cdot U_{q,\mathbb{Q}}^0 \right\} \subseteq \hat{U}_{q,\mathbb{Q}}. \quad (9.8)$$

These subspaces now generate the $U_{q,\mathbb{Q}}$ -sub-bimodules $\hat{\mathcal{L}}_{\blacktriangleleft w} = U_{q,\mathbb{Q}} \cdot \mathring{\mathcal{L}}_{\blacktriangleleft w} \cdot U_{q,\mathbb{Q}}$. Analogously, the subspaces $\mathring{\mathcal{L}}_{\blacktriangleleft w}^2 = \mathring{\mathcal{L}}_{\blacktriangleleft w} \hat{\otimes} \mathring{\mathcal{L}}_{\blacktriangleleft w}$ are defined from formal sums $\sum_{\phi, \psi \in \mathcal{E}(s)} R_{\phi, \psi} (E_{\blacktriangleleft w}^\phi \otimes E_{\blacktriangleleft w}^\psi)$ with $R_{\phi, \psi} \in (\mathfrak{S}_{\blacktriangleleft w}^- \cdot U_{q,\mathbb{Q}}^0)^{\otimes 2}$, generating the $U_{q,\mathbb{Q}}^{\otimes 2}$ -sub-bimodules $\hat{\mathcal{L}}_{\blacktriangleleft w}^2 = \hat{\mathcal{L}}_{\blacktriangleleft w} \hat{\otimes} \hat{\mathcal{L}}_{\blacktriangleleft w}$.

The next lemma follows by applying (5.37) or (5.26) to the limiting expressions.

Lemma 9.6. *Suppose $u_1, u_2 \in \mathcal{W}^*$ are reduced words such that $w = u_1 \cdot u_2$ is reduced. Let $t_i = r(u_i)$ and $s = t_1 \cdot t_2 = r(w)$ so that $l(s) = l(t_1) + l(t_2)$. Then Γ_{t_1} extends to a linear map on $\mathring{\mathcal{L}}_{\blacktriangleleft u_2}$ as well as $\hat{\mathcal{L}}_{\blacktriangleleft u_2}$ with images $\Gamma_{t_1}(\hat{\mathcal{L}}_{\blacktriangleleft u_2}) \subseteq \hat{\mathcal{L}}_{\blacktriangleleft w}$ and $\Gamma_{t_1}(\mathring{\mathcal{L}}_{\blacktriangleleft u_2}) \subseteq \mathring{\mathcal{L}}_{\blacktriangleleft w}$ in $\hat{U}_{q,\mathbb{Q}}$.*

Analogously, $\Gamma_{t_1}^{\otimes 2}$ extends to a linear map on $\mathring{\mathcal{L}}_{\blacktriangleleft u_2}^2$ and $\hat{\mathcal{L}}_{\blacktriangleleft u_2}^2$ with images $\Gamma_{t_1}^{\otimes 2}(\hat{\mathcal{L}}_{\blacktriangleleft u_2}^2) \subseteq \hat{\mathcal{L}}_{\blacktriangleleft w}^2$ and $\Gamma_{t_1}^{\otimes 2}(\mathring{\mathcal{L}}_{\blacktriangleleft u_2}^2) \subseteq \mathring{\mathcal{L}}_{\blacktriangleleft w}^2$ in $\hat{U}_{q,\mathbb{Q}}^{\hat{\otimes} 2}$.

Clearly, $\mathcal{P}_i, \mathcal{P}_i^{-1} \in \mathring{\mathcal{L}}_{\blacktriangleleft w_i}^2$ with w_i a word of length one. Suppose $w \in \mathcal{W}^*$ with $r(w) = i$, so that $w = w^\flat \cdot w_i$. Then Lemma 9.6 allows us to define an *elementary* quasi- R -matrix associated to the reduced word w as

$$\mathcal{P}_w = \Gamma_{w^\flat} \otimes \Gamma_{w^\flat}(\mathcal{P}_i) = \sum_{j=0}^{\infty} \frac{q_i^{j(j-1)/2}}{[j]!_i} (q_i - q_i^{-1})^j E_w^j \otimes F_w^j. \quad (9.9)$$

Following the conventions of (2.15) and Section 2.3, we write $\beta^j = \gamma(w[1, j])$ and $w^j = w[1, j] = w[\beta_j]$. Assume that $L = l(w)$. In the notation of Section 5.3, we define the *partial* quasi- R -matrix associated to w by the following products in $\hat{U}_{q,\mathbb{Q}} \hat{\otimes} \hat{U}_{q,\mathbb{Q}}$.

$$\mathcal{P}_{\blacktriangleleft w} = \mathcal{P}_{w^L} \dots \mathcal{P}_{w^1} = \mathcal{P}_{w[\beta_L]} \dots \mathcal{P}_{w[\beta_1]} = \sum_{\psi \in \mathcal{E}(s)} \mathbf{c}_\psi(q)^{-1} \cdot E_{\blacktriangleleft w}^\psi \otimes F_{\blacktriangleleft w}^\psi \in \mathring{\mathcal{L}}_{\blacktriangleleft w}^2 \quad (9.10)$$

Here $s = r(w)$ and the coefficients in $\mathbb{Q}(q)$ are given by

$$\begin{aligned} \mathbf{c}_{\psi}(q) &= \frac{[\psi]!}{\sigma_{\psi}(q)} \in \Lambda_{q,e^*} \quad \text{where} \quad [\psi]! = \prod_{\alpha \in \Phi^+} [\psi(\alpha)]!_{\alpha} \\ \text{and} \quad \sigma_{\psi}(q) &= \prod_{\alpha \in \Phi^+} q_{\alpha}^{\psi(\alpha)(\psi(\alpha)-1)/2} (q_{\alpha} - q_{\alpha}^{-1})^{\psi(\alpha)} = (-1)^{n(\psi)} \cdot \delta^{\psi}(q) \cdot q^{\sum_{\alpha} d_{\alpha} \binom{\psi(\alpha)}{2}}. \end{aligned} \quad (9.11)$$

The elements $[\psi]!$ and $\delta^{\psi}(q)$ are as defined in (5.33) and (5.34), respectively. Thus, in the product expressions, we consider all positive roots rather than just those in $\mathcal{N}(s)$ and assume $\psi \in \mathcal{E}(s)$ is extended to all of Φ^+ by setting ψ to zero outside of $\mathcal{N}(s)$.

The inclusion in Λ_{q,e^*} is obvious from the definitions, and the formal sum expression in (9.10) is immediate from the definitions in (5.23). It also implies the indicated inclusion into $\mathring{\mathcal{L}}_{\blacktriangleleft w}^2$.

Let $w = u_1 \cdot u_2 \in \mathcal{W}^*$ and $t_i = r(u_i)$ as above so that $l(s) = l(t_1) + l(t_2)$ for $s = t_1 t_2 = r(w)$. Lemma 9.6 now implies that $\Gamma_{t_1}^{\otimes 2}(\mathcal{P}_{\blacktriangleleft u_2})$ is well-defined and given by applying $\Gamma_{t_1}^{\otimes 2}$ to individual summands. Since the q_{α} depend only on the \mathcal{W} -orbit of α , one readily checks that $\mathbf{c}_{s^*(\psi)}(q) = \mathbf{c}_{\psi}(q)$. In the situation where ψ and (ψ_1, ψ_2) are related as in (5.22), we infer from this that $\mathbf{c}_{\psi}(q) = \mathbf{c}_{\psi_1}(q) \mathbf{c}_{(t_1^{-1})^*(\psi_2)}(q) = \mathbf{c}_{\psi_1}(q) \mathbf{c}_{\psi_2}(q)$. Together with (5.26) we then find the following well-defined recursion.

$$\mathcal{P}_{\blacktriangleleft w} = \Gamma_{t_1}^{\otimes 2} \left(\mathcal{P}_{\blacktriangleleft u_2} \right) \cdot \mathcal{P}_{\blacktriangleleft u_1} \in \mathring{\mathcal{L}}_{\blacktriangleleft w}^2 \quad (9.12)$$

A central property of the $\mathcal{P}_{\blacktriangleleft w}$ elements is that they intertwine the standard coproduct Δ with Δ^{Γ_w} , the coproduct conjugated by the braid automorphism for the same word.

Proposition 9.7. *Suppose $w \in \mathcal{W}^*$ and $x \in U_{q,\mathbb{Q}}$. Then the following relation holds in $\mathring{\mathcal{L}}_{\blacktriangleleft w}^2$,*

$$\mathcal{P}_{\blacktriangleleft w} \cdot \Delta(\Gamma_w(x)) = \Gamma_w^{\otimes 2}(\Delta(x)) \cdot \mathcal{P}_{\blacktriangleleft w}. \quad (9.13)$$

Proof. The proof proceeds by induction on the length of w . For a length one word $w = w_i$ we have $\mathcal{P}_{\blacktriangleleft w_i} = \mathcal{P}_i$, reducing (9.13) to

$$\mathcal{P}_i \cdot \Delta(\Gamma_i(x)) = \Gamma_i^{\otimes 2}(\Delta(x)) \cdot \mathcal{P}_i.$$

This case is verified, for example, by taking a formal limit $\ell_i \rightarrow \infty$ for the computations in Appendix A. It also appears in some equivalent variations in [LS90, Prop.2.4.1], [KR90, pg.429], or [AJS94, App.C.4].

Assume next that (9.13) holds for $w \in \mathcal{W}$ and that $w \cdot w_i \in \mathcal{W}$ is reduced. From (9.12) we obtain $\mathcal{P}_{\blacktriangleleft w \cdot w_i} = \Gamma_w^{\otimes 2}(\mathcal{P}_i) \cdot \mathcal{P}_{\blacktriangleleft w}$ as a well-defined relation in $\mathring{\mathcal{L}}_{\blacktriangleleft w \cdot w_i}$. The expressions and calculations below are now well-defined on $\hat{\mathcal{L}}_{\blacktriangleleft w \cdot w_i}$ by Lemma 9.6. They use both the induction hypothesis and the base case relations.

$$\begin{aligned} \Gamma_{w \cdot w_i}^{\otimes 2}(\Delta(x)) \mathcal{P}_{\blacktriangleleft w \cdot w_i} &= \Gamma_w^{\otimes 2} \circ \Gamma_i^{\otimes 2}(\Delta(x)) \cdot \Gamma_w^{\otimes 2}(\mathcal{P}_i) \cdot \mathcal{P}_{\blacktriangleleft w} &= \Gamma_w^{\otimes 2} \left(\Gamma_i^{\otimes 2}(\Delta(x)) \cdot \mathcal{P}_i \right) \cdot \mathcal{P}_{\blacktriangleleft w} \\ &= \Gamma_w^{\otimes 2} \left(\mathcal{P}_i \cdot \Delta(\Gamma_i(x)) \right) \cdot \mathcal{P}_{\blacktriangleleft w} &= \Gamma_w^{\otimes 2}(\mathcal{P}_i) \cdot \Gamma_w^{\otimes 2} \left(\Delta(\Gamma_i(x)) \right) \cdot \mathcal{P}_{\blacktriangleleft w} \end{aligned}$$

$$= \Gamma_w^{\otimes 2}(\mathcal{P}_i) \cdot \mathcal{P}_{\blacktriangleleft w} \cdot \Delta(\Gamma_w(\Gamma_i(x))) \quad = \mathcal{P}_{\blacktriangleleft w \cdot w_i} \cdot \Delta(\Gamma_{w \cdot w_i}(x))$$

This shows the assertion for $w \cdot w_i$, completing the proof. \square

An identity similar to (9.13) can be found in [CP94, Prop 8.3.7] in the special case of $x = E_i$ with $w \cdot w_i$ reduced. The derivation there also involves certain quantum Weyl elements, which generally exist only in representations that admit a compatible \mathcal{W} -action and, hence, limiting the statement to an identity between linear maps on such representations. We, therefore, abstain from the use of techniques using Weyl elements here. See, Appendix C for further details.

A *full* quasi- R -matrix is associated to any reduced word of maximal length. The first step in verifying independence of the choice of word is an intertwining relation with a twisted coproduct defined on generators by

$$\bar{\Delta}(E_i) = E_i \otimes K_i^{-1} + 1 \otimes E_i, \quad \bar{\Delta}(F_i) = F_i \otimes 1 + K_i \otimes F_i, \quad \text{and} \quad \bar{\Delta}(K_i) = K_i \otimes K_i. \quad (9.14)$$

It is easy to check that $\bar{\Delta} \circ \Pi = \Pi^{\otimes 2} \circ \Delta$, where Π is as in (4.13). The latter involution coincides with the bar involution in [Lus93, 3.1.12] so that $\bar{\Delta}$ coincides with the alternate coproduct defined in [Lus93, 4.1.1]. The next result uses the fact that, for maximal length words, the coproduct is conjugated by the Garside automorphism, for which we derived an explicit form in Section 5.2.

Lemma 9.8. *Suppose $z \in \mathcal{W}_m^*$ is a reduced word of maximal length and $x \in U_{q,\mathbb{Q}}$. Then*

$$\mathcal{P}_{\blacktriangleleft z} \cdot \Delta(x) = \bar{\Delta}(x) \cdot \mathcal{P}_{\blacktriangleleft z}. \quad (9.15)$$

Proof. Recall from Corollary 5.3 that $\Gamma_{s_\bullet} = \mathcal{I} \circ \Pi \circ \mathcal{U} \circ S$ for the longest element $s_\bullet \in \mathcal{W}$. It is easy to check that $\Pi \circ \mathcal{U} = \Pi \circ \Omega$ for the commuting involutions defined in Section 4.2. Thus, $\Gamma_{s_\bullet} = \Pi \circ Q$, where $Q = \mathcal{I} \circ \Omega \circ S$. Recall that both $\Omega, S : U_q \rightarrow U_q^{\text{opp, cop}}$ are anti-coalgebra anti-automorphisms and \mathcal{I} , defined in (5.12), is a Hopf algebra automorphism. Thus, Q intertwines the coalgebra structure $Q^{\otimes 2}(\Delta(y)) = \Delta(Q(y))$. Given that $s_\bullet = \tau(z)$, Proposition 9.7 now implies

$$\begin{aligned} \mathcal{P}_{\blacktriangleleft z} \cdot \Delta(x) &= \Gamma_{s_\bullet}^{\otimes 2} \left(\Delta(\Gamma_{s_\bullet}^{-1}(x)) \right) \cdot \mathcal{P}_{\blacktriangleleft z} &= \Pi^{\otimes 2} \circ Q^{\otimes 2} \left(\Delta(Q^{-1} \circ \Pi^{-1}(x)) \right) \cdot \mathcal{P}_{\blacktriangleleft z} \\ &= \Pi^{\otimes 2} \left(\Delta(\Pi^{-1}(x)) \right) \cdot \mathcal{P}_{\blacktriangleleft z} &= \bar{\Delta}(x) \cdot \mathcal{P}_{\blacktriangleleft z}. \end{aligned} \quad \square$$

In [Lus93, Thm.4.1.2(a)] Lusztig shows that an element Θ in some completion of $\bigoplus_v (U_{q,\mathbb{Q}}^-)_v \otimes (U_{q,\mathbb{Q}}^+)_v$ is uniquely determined by the relation $\Theta \cdot \bar{\Delta}^{\text{opp}}(y) = \Delta^{\text{opp}}(y) \cdot \Theta$ (since [Lus93] uses the opposite coproduct of ours). The required assumptions can be readily checked for $\Theta = (\mathcal{P}_{\blacktriangleleft z}^{-1})^{\text{opp}}$ using the general form as in (9.10), the commutation (9.15), and noting that the completions align. From the uniqueness of solutions to (9.15) we infer word independence for quasi- R -matrices associated to reduced words of maximal length.

Corollary 9.9. *There exists $\mathcal{P}_\bullet \in \hat{U}_{q,\mathbb{Q}}^{\otimes 2}$ such that $\mathcal{P}_\bullet = \mathcal{P}_{\blacktriangleleft z}$ for all $z \in \mathcal{W}_m^*$. Moreover, $\Pi^{\otimes 2}(\mathcal{P}_\bullet) = \mathcal{P}_\bullet^{-1}$.*

9.3. Pre-Triangular Structures for U_q . Since the coefficients in (9.9) may be evaluated at $q = 1$, the element \mathcal{P}_\bullet has a well-defined image $\mathcal{P}_\bullet^\hbar$ in the power series completion $U_\hbar(\mathfrak{g}) \hat{\otimes} U_\hbar(\mathfrak{g})$ over $\mathbb{Q}[[\hbar]]$ as in Section 3.2. Originally, universal R -matrices were constructed in this setting by Drinfeld [Dri87] as products $\mathcal{R}^\hbar = D \cdot \mathcal{P}_\bullet^\hbar$ in $U_\hbar(\mathfrak{g})^{\hat{\otimes} 2}$. The “diagonal” term is given by

$$D = \exp \left(\hbar \sum_{i,j=1}^n (B^{-1})_{ij} H_i \otimes H_j \right), \quad (9.16)$$

where $B_{ij} = A_{ij} d_j^{-1} = (\check{\alpha}_i | \check{\alpha}_j) = C_{ij} d_i^{-1} d_j^{-1} \in \mathbf{e}^{-1} \mathbb{Z}$ with notation as in Section 2.1. Explicit formulae are obtained in [Ros89] for the A_n case and [KR90] for general Lie types, see also [CP94]. Tanisaki’s axioms in [Tan92] or (9.2) above are motivated by the fact that D cannot be defined in $U_q^{\otimes 2}$, but conjugation by D can be expressed as an automorphism on $\hat{U}_{q,\mathbb{Q}}^{\hat{\otimes} 2}$.

Specifically, if x and y are homogeneously graded elements with $\mathbf{w}(x) = \mu, \mathbf{w}(y) = \nu \in \mathbb{Z}^\Delta$, we have $[H_i, x] = (\mu | \check{\alpha}_i)x$ and $[H_j, y] = (\nu | \check{\alpha}_j)y$. From this we obtain $\sum_{i,j} (B^{-1})_{ij} [H_i \otimes H_j, x \otimes y] = x \otimes (H_\mu \cdot y) + (x \cdot H_\nu) \otimes y$, where $H_\mu = \sum_i \mu_i d_i H_i$ if $\mu = \sum_i \mu_i \alpha_i$. Exponentiation $t \mapsto e^{-\hbar t}$ of the implied adjoint and regular actions on $x \otimes y$ together with the notation from (4.1) then yields the identities below.

$$\Pi(x \otimes y) = D^{-1}(x \otimes y)D = xK^{-\nu} \otimes K^{-\mu}y = q^{-(\mu|\nu)}xK^{-\nu} \otimes yK^{-\mu} \quad (9.17)$$

Thus, although the Tanisaki automorphism is initially defined as $\Pi = \text{Ad}(D^{-1})$ on $U_\hbar(\mathfrak{g})^{\hat{\otimes} 2}$, it clearly maps the subalgebras $U_q^{\otimes 2}$ and $\hat{U}_{q,\mathbb{Q}}^{\hat{\otimes} 2}$ to themselves. The form in (9.16) can often still be defined, at least up to scalars, on tensor products of certain modules of $U_q^{\otimes 2}$, implying solutions to the Yang Baxter equations or solutions to (9.3).

For example, for generic highest weight representations V and W , the maps $D_{V,W}$ can be designed starting from generalized Verma modules. For a tuple $\kappa = (\kappa_1, \dots, \kappa_n) \in \mathbf{u}^\Delta$ with \mathbf{u} a group of units in the ground ring, the *generic Verma module* $V(\kappa)$ is defined as the induced representation from the one-dimensional $U_q^{\geq 0}$ representation, given by $K_i v_\kappa = \kappa_i v_\kappa$ and $E_i v_\kappa = 0$ on a basis vector v_κ . As usual, there is a linear isomorphism $U_q^- \rightarrow V(\kappa) : x \mapsto x \cdot v$.

Assume now for homogeneous $x, y \in U_q^-$ that $-\mathbf{w}(x) = \mathbf{w}^-(x) = \mu$ and $-\mathbf{w}(y) = \mathbf{w}^-(y) = \nu$ and write $\kappa^\mu = \kappa_1^{\mu_1} \dots \kappa_n^{\mu_n}$ for $\mu \in \mathbb{Z}^\Delta$. Then linear maps on tensor products of generic Verma modules are well-defined by the formula below. They are unique solutions to the equation $D^{-1}cD = \Pi(c)$ of module maps (with Π as in (9.17)) up to an overall scalar factor.

$$D_{V(\kappa), V(\lambda)}((x \cdot v_\kappa) \otimes (y \cdot v_\lambda)) = q^{(\mu|\nu)} \kappa^{-\nu} \lambda^{-\mu} ((x \cdot v_\kappa) \otimes (y \cdot v_\lambda)) \quad (9.18)$$

It is not hard to check that the respective isomorphisms $V(\kappa) \otimes V(\lambda) \rightarrow V(\lambda) \otimes V(\kappa)$, as defined in Corollary 9.4, obey the Yang Baxter Equation and, thus, produce a representation of the colored braid groupoids over $\mathbb{Q}(q)[\{\kappa_i^{\pm 1}, \lambda_i^{\pm 1}, \dots\}]$.

The categorical triangle relations may be realized in a more general setting, in which we consider a commutative ring \mathcal{E} with $\text{char}(\mathcal{E}) = 0$ and an extension of $(\cdot | \cdot) : \mathcal{E}^\Delta \times \mathcal{E}^\Delta \rightarrow \mathcal{E}$ for

the form from (2.2). Elements of the group ring $\mathbb{Z}[\mathcal{E}]$ may be written formally as q^x with $x \in \mathcal{E}$. Denoting $\mathbb{Q}(\mathcal{E})$ as the field of fractions, we have $\mathbb{Q}(\mathbb{Z}) = \mathbb{Q}(q)$. As a tensor category, one may then consider representations of U_q on \mathcal{E}^Δ -graded vector spaces $V = \bigoplus_\gamma V_\gamma$ subject to the condition that $K_i v = q^{(\alpha_i|\gamma)} v$ for $v \in V_\gamma$ and $\gamma \in \mathcal{E}^\Delta$. For two graded representations V and W , define then $D_{V,W}$ to be $q^{(\gamma|\eta)}$ on the $V_\gamma \otimes W_\eta$ component.

The relations (9.3) and (9.17) are now easily verified on such modules, yielding a braided tensor category \mathcal{C} over $\mathbb{Q}(\mathcal{E})$. As an abelian category the latter naturally splits in a sum $\bigoplus_\rho \mathcal{C}_\rho$ with $\rho \in \mathcal{E}^1 = \mathcal{E}/\mathbb{Z}$ with the obvious \mathcal{E}^1 -graded behavior for tensors.

The main result for the generic case is summarized in the proposition below. It has been stated at various levels of rigor and formality in numerous other places in the literature. We adapt here results from [Lus93] to our conventions, which itself is following similar calculations as those in [Tan92, Theorem 4.3.3].

Proposition 9.10. *The triple $(\hat{U}_{q,\mathbb{Q}}, \mathcal{P}_\bullet, \Pi)$ is a pre-triangular bialgebra.*

Proof. A straightforward verification on generators shows that $\bar{\Delta} = \Pi \circ \Delta^{\text{opp}}$. Lemma 9.8 then implies the last relation in (9.2).

The argument preceding Corollary 9.9 identifies Lusztig's universal element with our quasi- R -matrix via $\Theta = (\mathcal{P}_\bullet^{-1})^{\text{opp}}$ or, equivalently, $\bar{\Theta}^{\text{opp}} = \mathcal{P}_\bullet$ so that, restricted to weight components, we have for any $v \in \mathbb{Z}^\Delta$

$$\bar{\Theta}_v^{\text{opp}} = (\mathcal{P}_\bullet)_v = \sum_{\psi : \vec{\psi} = v} \mathbf{c}_\psi(q)^{-1} \cdot E_{\blacktriangleleft z}^\psi \otimes F_{\blacktriangleleft z}^\psi,$$

with $\vec{\psi}$ as defined in Section 5.3 and $z \in \mathcal{W}_m^*$. Exchanging Lusztig's conventions with ours, namely Δ with Δ^{opp} and \tilde{K}_{-v} by K^{-v} , the formula in [Lus93, pg 38] reads

$$(\Delta^{\text{opp}} \otimes 1) \bar{\Theta}_\mu = \sum_{v' + v'' = \mu} \bar{\Theta}_{v'}^{13} (1 \otimes K^{-v'} \otimes 1) \bar{\Theta}_{v''}^{23}.$$

Applying the (13) permutation to the tensor factors and substituting the \mathcal{P}_\bullet components above we arrive at

$$(1 \otimes \Delta) \mathcal{P}_{\bullet,\mu} = \sum_{v' + v'' = \mu} \mathcal{P}_{\bullet,v'}^{13} (1 \otimes K^{-v'} \otimes 1) \mathcal{P}_{\bullet,v''}^{12} = \sum_{v' + v'' = \mu} \Pi_{12}(\mathcal{P}_{\bullet,v'}^{13}) \mathcal{P}_{\bullet,v''}^{12}, \quad (9.19)$$

using formula (9.17) for Π . Summation, in the sense of the inverse limits, then yields the first identity in (9.2). The second follows analogously. \square

A left inverse of \mathcal{P}_\bullet may be constructed by applying the map $a \otimes b \otimes c \mapsto a \otimes S^{-1}(c)b$ to the first relation in (9.19). Using the antipode axiom for Δ^{opp} , the expression in (9.10) for a maximal word $z \in \mathcal{W}_m^*$, the formulae for the antipode on basis elements in (5.31), as well as notations from (5.21) and (2.19) we obtain that

$$\tilde{\mathcal{P}}_{\blacktriangleleft z} = \sum_{\psi \in \mathcal{E}_\bullet} (-1)^{\text{ht}(\vec{\psi})} q^{-\frac{1}{2}((\vec{\psi}|\vec{\psi}) - 2(\rho|\vec{\psi}))} \mathbf{c}_\psi(q)^{-1} \cdot E_{\blacktriangleleft z}^\psi \otimes F_{\blacktriangleleft z^\dagger}^\psi \in \mathring{\mathcal{L}}_{\blacktriangleleft z}^2 \quad (9.20)$$

is a left inverse of $\mathcal{P}_{\blacktriangleleft z}$ by summation over all graded components in the limit. A similar calculation, applying $a \otimes b \otimes c \mapsto a \otimes c S^{-1}(b)$ and the automorphism $x \otimes y \mapsto x \otimes y K^\nu$ for $w(x) = \nu$, shows that $\tilde{\mathcal{P}}_{\blacktriangleleft z}$ is also a right inverse of $\mathcal{P}_{\blacktriangleleft z}$.

Applying $a \otimes b \otimes c \mapsto a S(b) \otimes c$ and $a \otimes b \otimes c \mapsto S(a)b \otimes c$ to the second relation in (9.19) shows that $\tilde{\mathcal{P}}_{\blacktriangleleft z^\dagger}$ is also a two-sided inverse of $\mathcal{P}_{\blacktriangleleft z}$ so that

$$\mathcal{P}_{\blacktriangleleft z}^{-1} = \tilde{\mathcal{P}}_{\blacktriangleleft z} = \tilde{\mathcal{P}}_{\blacktriangleleft z^\dagger} = \mathcal{P}_{\blacktriangleleft z^\dagger}^{-1}. \quad (9.21)$$

From this identity together with (5.31), (9.17), and word independence one readily derives

$$S \hat{\otimes} S(\mathcal{P}_\bullet) = \Pi^{-1}(\mathcal{P}_\bullet). \quad (9.22)$$

It is useful to express (9.19) also as a relation of (co)multiplication coefficients. Suppose $z \in \mathcal{W}_m^*$ is a reduced word of maximal length. The PBW theorems from Section 5, the fact that U_q^+ is an algebra over Λ_{q,e^*} , and $U_q^{<0}$ a coalgebra over Λ_q then imply that there are elements $\mathbf{m}_\chi^{\psi,\phi}(q) \in \Lambda_{q,e^*}$ and $\mathbf{d}_{\phi,\psi}^\chi(q) \in \Lambda_q$ with $\psi, \phi, \chi \in \mathbb{N}_0^{\Phi^+}$ such that

$$E_{\blacktriangleleft z}^\psi E_{\blacktriangleleft z}^\phi = \sum_\chi \mathbf{m}_\chi^{\psi,\phi}(q) \cdot E_{\blacktriangleleft z}^\chi \quad \text{and} \quad \Delta(F_{\blacktriangleleft z}^\chi) = \sum_{\phi,\psi} \mathbf{d}_{\phi,\psi}^\chi(q) \cdot K^{-\vec{\psi}} F_{\blacktriangleleft z}^\phi \otimes F_{\blacktriangleleft z}^\psi. \quad (9.23)$$

The summations are over only finitely many elements in $\mathbb{N}_0^{\Phi^+}$ since they are subject to the grading constraint $\vec{\chi} = \vec{\phi} + \vec{\psi}$. It is clear that the only allowed terms for $\Delta(F_{\blacktriangleleft z}^\chi)$ are those appearing in the sum (9.23). Linear independence of the elements $E_{\blacktriangleleft z}^\phi \otimes K^{-\vec{\psi}} F_{\blacktriangleleft z}^\phi \otimes F_{\blacktriangleleft z}^\psi$ now implies that equation (9.19) is, indeed, *equivalent* to the condition that

$$\mathbf{c}_\chi(q) \mathbf{m}_\chi^{\psi,\phi}(q) = \mathbf{c}_\psi(q) \mathbf{c}_\phi(q) \mathbf{d}_{\phi,\psi}^\chi(q) \quad \text{for all } \chi, \phi, \psi \in \mathbb{N}_0^{\Phi^+}. \quad (9.24)$$

An analogous equation of (co) multiplication coefficients, equivalent to the second Tanisaki relation in (9.19), can be derived in the same manner.

9.4. Maximal Hopf Ideals from Tanisaki-Lusztig Pairings. Yet another viewpoint of the same axioms in (9.19) is derived from the pairings used in [Lus93, Tan92]. Fix some $z \in \mathcal{W}_m^*$ and observe that, by (9.11), we may define a pairing of U_q^+ with U_q^- over Λ_{q,e^*} as follows using PBW bases.

$$\langle \cdot, \cdot \rangle_z : U_q^- \times U_q^+ \rightarrow \Lambda_{q,e^*} \quad \text{with} \quad \langle F_{\blacktriangleleft z}^\psi, E_{\blacktriangleleft z}^\phi \rangle_z = \delta_{\psi,\phi} \cdot \mathbf{c}_\psi(q). \quad (9.25)$$

Note that (9.25) is dual to (9.10) in the sense that $(\langle F_{\blacktriangleleft z}^\psi, \cdot \rangle_z \hat{\otimes} \text{id})(\mathcal{P}_{\blacktriangleleft z}) = F_{\blacktriangleleft z}^\psi$. Word independence and linearity of this relation imply that $(\langle f, \cdot \rangle_z \otimes \text{id})(\mathcal{P}_\bullet) = f$ for all $f \in U_q^-$. This, in turn, shows that the pairing in (9.25) is independent of z as well.

Define a linear isomorphisms $\Pi_R : A^{\otimes 2} \rightarrow A^{\otimes 2}$ by $\Pi_R(x \otimes y) = x \otimes K^{-\mu} y$, where x is homogeneous with $w(x) = \mu$ and A is any graded subalgebra that contains U_q^0 . The form of the coproduct in (9.23) now implies that we have a well-defined map

$$\hat{\Delta} = \Pi_R \circ \Delta^{\text{opp}} : U_q^- \rightarrow U_q^- \otimes U_q^-, \quad (9.26)$$

which is a homomorphism with respect to the braided product structure. The latter is given on homogeneous elements in $(U_q^-)^{\otimes 2}$ by $(x \otimes y)(x' \otimes y') = q^{-(\mu|\nu)}xx' \otimes yy'$, where $\mu = \mathbf{w}(y)$ and $\nu = \mathbf{w}(x')$. One easily checks that $\hat{\Delta}$ is coassociative.

In this language, the first relation in (9.2) as well as the coefficient relation in (9.24) are equivalent to the requirement that $\hat{\Delta}$ is adjoint to the multiplication in U_q^+ with respect to the pairing in (9.25). That is,

$$\langle \hat{\Delta}(a), b \otimes c \rangle = \langle a, bc \rangle \quad \text{for all } a \in U_q^- \quad \text{and} \quad b, c \in U_q^+. \quad (9.27)$$

As before, analogous statements hold for the second Tanisaki relation.

Consider next the specialization to a primitive ℓ -th root of unity ζ such that $\ell > \mathbf{e}$. As in (6.9), denote for $\mathbf{e} + 1 = \mathbf{e}^* \leq m \leq \ell$ the algebras

$$U_{\zeta,m} = U_q \otimes \Lambda_{\zeta,m}^* \quad \text{and} \quad U_{\zeta,m}^\square = U_q^\square \otimes \Lambda_{\zeta,m} \quad \text{if } \square \in \{\geq 0, \leq 0, +, -\}. \quad (9.28)$$

The form (9.25) clearly specializes to a form $\langle \cdot, \cdot \rangle_{\zeta,m} : U_{\zeta,m}^- \times U_{\zeta,m}^+ \rightarrow \Lambda_{\zeta,m}^*$ but will be degenerate. Denote by $N_{\zeta,m}^\pm \subset U_{\zeta,m}^\pm$ the respective null spaces.

Lemma 9.11. *With notation as in (6.72) and (6.65) we have $N_{\zeta,m}^\pm = \hat{\mathcal{K}}_\bullet^\pm \otimes \Lambda_{\zeta,m}$. In particular, $\hat{\mathcal{K}}_\bullet^\pm$ is uniquely defined for all Lie types.*

Proof. Observe that, with the exponent set from (6.64),

$$\mathbf{c}_\psi(\zeta) = 0 \quad \text{if and only if} \quad \psi \in [\mathbf{l}, \infty)^*.$$

The claim is now immediate from the relation in (6.65). \square

Recall from Section 6.5 the remaining ambiguity for the G_2 type, which is resolved here via the word independence of the pairing. As before, we will suppress the ground ring from our notation $N_{\zeta,m}^\pm = \hat{\mathcal{K}}_\bullet^\pm$ whenever it is clear from the context.

Proposition 9.12. *The ideals $\hat{\mathcal{K}}_\bullet^{\leq 0} \subset U_{\zeta,m}^{\leq 0}$ and $\hat{\mathcal{K}}_\bullet^{\geq 0} \subset U_{\zeta,m}^{\geq 0}$ are Hopf ideals, independent of choices of reduced words of maximal length.*

Proof. The identity (9.27) implies that the left null space has to be a co-ideal with respect to $\hat{\Delta}$. Together with Lemma 9.11 we thus obtain

$$\hat{\Delta}(\hat{\mathcal{K}}_\bullet^-) \subseteq \hat{\mathcal{K}}_\bullet^- \otimes U_{\zeta,m}^- + U_{\zeta,m}^- \otimes \hat{\mathcal{K}}_\bullet^-.$$

With $U_{\zeta,m}^{\leq 0} = U_{\zeta,m}^0 \cdot U_{\zeta,m}^-$ and $\hat{\mathcal{K}}_\bullet^{\leq 0} = U_{\zeta,m}^0 \cdot \hat{\mathcal{K}}_\bullet^-$ it is easy to check that Π_R maps $\hat{\mathcal{K}}_\bullet^{\leq 0} \otimes U_{\zeta,m}^{\leq 0}$ as well as $U_{\zeta,m}^{\leq 0} \otimes \hat{\mathcal{K}}_\bullet^{\leq 0}$ to itself. The co-ideal equation for $\hat{\Delta}$ and (9.26) now imply the respective co-ideal condition for the regular coproduct with respect to $\hat{\mathcal{K}}_\bullet^{\leq 0}$. Word independence of $\hat{\mathcal{K}}_\bullet^{\leq 0}$ from Lemma 9.11 and relation (6.66) now entails that $\hat{\mathcal{K}}_\bullet^{\leq 0}$ is invariant under the antipode and, hence, a Hopf ideal. Finally, recall that $\Omega(\hat{\mathcal{K}}_\bullet^{\leq 0}) \subseteq \hat{\mathcal{K}}_\bullet^{\geq 0}$, $\Omega(U_{\zeta,m}^{\leq 0}) \subseteq U_{\zeta,m}^{\geq 0}$, $\Omega^{\otimes 2} \circ \Delta = \Delta^{\text{opp}} \circ \Omega$, and $\Omega \circ S = S \circ \Omega$. From these relations one readily derives that $\hat{\mathcal{K}}_\bullet^{\geq 0}$ is also a Hopf ideal. \square

We may consider now the full maxiaml ideal $\hat{\mathcal{K}}_{\bullet\bullet} = (\hat{\mathcal{K}}_{\bullet}^{\geq 0} + \hat{\mathcal{K}}_{\bullet}^{\leq 0}) \cdot U_{\zeta, m}$ for the full algebra $U_{\zeta, m}$ over $\Lambda_{\zeta, m}^*$, which has all desired properties.

Corollary 9.13. *Suppose $e^* \leq m \leq \ell$.*

- i) *For any Lie type and root of unity with ζ as in (6.1), $\hat{\mathcal{K}}_{\bullet\bullet}$ is a Hopf ideal in $U_{\zeta, m}$.*
- ii) *$\hat{\mathcal{K}}_{\bullet\bullet}$ is equal to the ideal generated by the set $\{X_w, Y_{w'} : \emptyset \neq w \leq_R z, \emptyset \neq w' \leq_R z'\}$ for any choice of $z, z' \in \mathcal{W}_m^*$.*
- iii) *$\hat{\mathcal{K}}_{\bullet\bullet}$ is stable under all automorphisms discussed in Section 4.*

Proof. The first part is immediate from Proposition 9.12. Invariance under the automorphisms Ω , \mathfrak{U} , and S follows from (6.66) and word independence. Gradings and identities, such as (5.15) and $\Pi = \Upsilon \circ \Omega$, imply invariance for all others in Section 4.2. \square

For the Artin generators Γ_i we may repeat the proof of Theorem 6.22 nearly verbatim. Specifically, for the choice of w therein, it is shown that for all $v \leq_R w$ the element $\Gamma_i(X_v)$ is again some power generator and, hence, in $\hat{\mathcal{K}}_{\bullet\bullet}$ by word independence. \square

Thus, if we define the *restricted quantum groups* by the quotient

$$U_{\zeta, m}^{\text{rest}} = U_{\zeta, m} / \hat{\mathcal{K}}_{\bullet\bullet}, \quad (9.29)$$

then Corollary 9.13 implies that $U_{\zeta, m}^{\text{rest}}$ is indeed a Hopf algebra, which is independent of any choices made in its definition. The construction of a full poset of ideals under $\hat{\mathcal{K}}_{\bullet\bullet}$ and respective quasi- R -matrices will be addressed in the next sections with different techniques.

9.5. Truncated Quasi- R -Matrices and Coproduct Formulas. In this section we develop a calculus of quasi- R -matrices at roots of unity analogous to the one in Section 9.2. To avoid singularities, the summations defining these R -matrices need to be truncated so that the corresponding identities hold only up to certain ideals.

It is, therefore, useful to extend the ideals in U_{ζ}^{\pm} defined (6.65) to the full algebra $U_{\zeta, m}$ over $\Lambda_{\zeta, m}^*$ for a given primitive ℓ -th root of unity ζ and $e^* \leq m \leq \ell$. For any pair of reduced words $u, w \in \mathcal{W}^*$, we write

$$\hat{\mathcal{K}}(u, w)_m = (\hat{\mathcal{K}}(u)^+ + \hat{\mathcal{K}}(w)^-) \cdot U_{\zeta, m} = \mathcal{K}(u, w) \cdot U_{\zeta, m} = \hat{\mathcal{K}}(u, \emptyset)_m + \hat{\mathcal{K}}(\emptyset, w)_m, \quad (9.30)$$

which may, alternatively, be viewed as the two-sided ideal in $U_{\zeta, m}$ generated by the set $\{X_a, Y_b : \emptyset \neq a, b \leq_R u\}$. Correspondingly, for reduced words $a, b \in \mathcal{W}^*$ for which $a \cdot b$ is also reduced, we infer that, for any Lie type,

$$\hat{\mathcal{K}}(a \cdot b, \emptyset)_m = \hat{\mathcal{K}}(a, \emptyset)_m + \Gamma_a(\hat{\mathcal{K}}(b, \emptyset)_m) \quad \text{and} \quad \Omega(\hat{\mathcal{K}}(a, \emptyset)_m) = \hat{\mathcal{K}}(\emptyset, a)_m \quad (9.31)$$

hold as well as the respective identities with flipped arguments and summations over ideals. A useful symmetry for various tensor identities is

$$\Omega^\# = (1 \ 2) \circ \Omega^{\otimes 2} : U_{\zeta, \ell}^{\otimes 2} \rightarrow U_{\zeta, \ell}^{\otimes 2} : x \otimes y \mapsto \Omega^\#(x \otimes y) = \Omega(y) \otimes \Omega(x), \quad (9.32)$$

which is an anti-linear anti-involution by the properties in (4.10).

To ensure minimal truncations, we consider R -matrices defined over the maximal ground rings $\Lambda_{\zeta, \ell}$ or $\Lambda_{\zeta, \ell}^*$, observing that $[s]!_i \neq 0$ whenever $s < \ell_i$. We begin with the definition of the *simple truncated* quasi- R -matrix associated to a given $\alpha_i \in \Delta$,

$$\mathcal{P}_i^\bullet = \sum_{s=0}^{\ell_i-1} \frac{\zeta_i^{s(s-1)/2}}{[s]!_i} (\zeta_i - \zeta_i^{-1})^s \cdot E_i^s \otimes F_i^s \in U_{\zeta, \ell}^+ \otimes U_{\zeta, \ell}^-, \quad (9.33)$$

which is well-defined over $\Lambda_{\zeta, \ell}$. Analogous to the generic case we, further, denote the conjugate R -matrix as

$$\bar{\mathcal{P}}_i^\bullet = \Omega^\#(\mathcal{P}_i^\bullet) = \sum_{s=0}^{\ell_i-1} \frac{\zeta_i^{-s(s-1)/2}}{[s]!_i} (-1)^s (\zeta_i - \zeta_i^{-1})^s \cdot E_i^s \otimes F_i^s \in U_{\zeta, \ell}^+ \otimes U_{\zeta, \ell}^-, \quad (9.34)$$

where $\Omega^\#$ is as in (9.32). To express these as inverses, we introduce for any $w \in \mathcal{W}^*$ the ideals

$$\mathcal{N}_o(w) = (\hat{\mathcal{K}}(w)^+ \otimes \hat{\mathcal{K}}(w)^-) \otimes \Lambda_{\zeta, \ell} = (\hat{\mathcal{K}}(w, \emptyset)_\ell \otimes \hat{\mathcal{K}}(\emptyset, w)_\ell) \cap (U_{\zeta, \ell}^+ \otimes U_{\zeta, \ell}^-), \quad (9.35)$$

generated by all $X_a \otimes Y_b$ with $\emptyset \neq a, b \leq_R w$. In calculations throughout this section, we also employ the standard notation $x \equiv y \pmod{V}$ to mean $x - y \in V$ for a given submodule V .

Lemma 9.14 (cf. [Oht02, Lemma A.1]). *For all $\alpha_i \in \Delta$ we have*

$$\bar{\mathcal{P}}_i^\bullet \cdot \mathcal{P}_i^\bullet \equiv \mathcal{P}_i^\bullet \cdot \bar{\mathcal{P}}_i^\bullet \equiv 1 \otimes 1 \pmod{\mathcal{N}_o(w_i)}.$$

Proof. We start with a direct computation and resummation.

$$\begin{aligned} \mathcal{P}_i^\bullet \cdot \bar{\mathcal{P}}_i^\bullet &= \sum_{r,n=0}^{\ell_i-1} \frac{\zeta_i^{r(r-1)/2-n(n-1)/2}}{[r]!_i [n]!_i} (-1)^n (\zeta_i - \zeta_i^{-1})^{r+n} E_i^{r+n} \otimes F_i^{r+n} \\ &= \sum_{k=0}^{2\ell_i-2} \rho_k \cdot \zeta_i^{-k(k-1)/2} (-1)^k (\zeta_i - \zeta_i^{-1})^k E_i^k \otimes F_i^k, \\ \text{where } \rho_k &= \sum_{r=\max(0, k-(\ell_i-1))}^{\min(k, \ell_i-1)} \frac{1}{[r]!_i [k-r]!_i} (-1)^r \zeta_i^{r(k-1)}. \end{aligned}$$

For $0 < k < \ell_i$, we compute

$$\begin{aligned} \rho_k &= \sum_{r=0}^k \frac{1}{[r]!_i [k-r]!_i} (-1)^r \zeta_i^{r(k-1)} = \frac{1}{[k]!} \sum_{r=0}^k \begin{bmatrix} k \\ r \end{bmatrix}_i (-1)^r \zeta_i^{r(k-1)} \\ &= \frac{1}{[k]!} \prod_{t=1}^k (-\zeta_i^{2(t-1)} + 1) = 0, \end{aligned}$$

where we used (3.4) with substitutions $a \mapsto k$, $k \mapsto r$, $q \mapsto \zeta_i^{-1}$, and $z \mapsto -\zeta_i^{-2}$. Observing that $\rho_0 = 1$, $X_i = E_i^{\ell_i}$, and $Y_i = F_i^{\ell_i}$ we now find

$$\mathcal{P}_i^\bullet \cdot \bar{\mathcal{P}}_i^\bullet = 1 \otimes 1 + (X_i \otimes Y_i) \sum_{k=\ell_i}^{2\ell_i-2} \rho_k \cdot \zeta_i^{-k(k-1)/2} (-1)^k (\zeta_i - \zeta_i^{-1})^k E_i^{k-\ell_i} \otimes F_i^{k-\ell_i}, \quad (9.36)$$

for which the latter term is clearly in $\mathcal{N}_o(w_i)$. The verification for $\bar{\mathcal{P}}_i^\bullet \cdot \mathcal{P}_i^\bullet$ is nearly identical. \square

Analogous to the generic case and using definitions from (5.11), we now define the *elementary* truncated quasi- R -matrices associated to a reduced word $w \in \mathcal{W}^*$ as

$$\begin{aligned} \mathcal{P}_w^\bullet &= \Gamma_{w^b}^{\otimes 2} (\mathcal{P}_{\tau(w)}^\bullet) = \sum_{s=0}^{\ell_w-1} \frac{\zeta_w^{s(s-1)/2}}{[s]!_w} (\zeta_w - \zeta_w^{-1})^s E_w^s \otimes F_w^s \\ \text{and} \quad \bar{\mathcal{P}}_w^\bullet &= \Gamma_{w^b}^{\otimes 2} (\bar{\mathcal{P}}_{\tau(w)}^\bullet) = \sum_{s=0}^{\ell_w-1} \frac{\zeta_w^{-s(s-1)/2}}{[s]!_w} (-1)^s (\zeta_w - \zeta_w^{-1})^s E_w^s \otimes F_w^s, \end{aligned} \quad (9.37)$$

with notations as in (2.8) and $\zeta_w = \zeta_{\tau(w)} = \zeta_{\gamma(w)}$. The following assertion is readily derived by applying $\Gamma_{w^b}^{\otimes 2}$ to (9.36) with $i = \tau(w)$, resulting in the same equation with i replaced by w .

Corollary 9.15. *For all $w \in \mathcal{W}^*$ we have*

$$\bar{\mathcal{P}}_w^\bullet \cdot \mathcal{P}_w^\bullet \equiv \mathcal{P}_w^\bullet \cdot \bar{\mathcal{P}}_w^\bullet \equiv 1 \otimes 1 \quad \text{mod } \mathcal{N}_o(w).$$

We define, similarly, the *partial* truncated quasi- R -matrices associated to some $w \in \mathcal{W}^*$ as

$$\begin{aligned} \mathcal{P}_{\blacktriangleleft w}^\bullet &= \mathcal{P}_{w^L}^\bullet \dots \mathcal{P}_{w^1}^\bullet = \sum_{\psi \in [0, \mathbf{I}]_s} \mathbf{c}_\psi(\zeta)^{-1} \cdot E_{\blacktriangleleft w}^\psi \otimes F_{\blacktriangleleft w}^\psi \\ \text{and} \quad \bar{\mathcal{P}}_{w^\blacktriangleright}^\bullet &= \bar{\mathcal{P}}_{w^1}^\bullet \dots \bar{\mathcal{P}}_{w^L}^\bullet = \sum_{\psi \in [0, \mathbf{I}]_s} \mathbf{c}_\psi(\zeta^{-1})^{-1} \cdot E_{w^\blacktriangleright}^\psi \otimes F_{w^\blacktriangleright}^\psi, \end{aligned} \quad (9.38)$$

with notation as in (9.10), $L = l(w)$, and $[0, \mathbf{I}]_s = [0, \mathbf{I}]_s \cap \mathcal{E}(s)$ as in (6.37). The identities below, with $a, b \in \mathcal{W}^*$ such that $a \cdot b$ is reduced, are straightforward from Corollary 9.15, the mentioned properties of $\Omega^\#$, and (5.26).

$$\begin{aligned} \bar{\mathcal{P}}_{w^\blacktriangleright}^\bullet &= \Omega^\#(\mathcal{P}_{\blacktriangleleft w}^\bullet) & \bar{\mathcal{P}}_{w^\blacktriangleright}^\bullet \cdot \mathcal{P}_{\blacktriangleleft w}^\bullet &\equiv 1^{\otimes 2} \equiv \bar{\mathcal{P}}_{w^\blacktriangleright}^\bullet \cdot \mathcal{P}_{\blacktriangleleft w}^\bullet \quad \text{mod } \mathcal{N}_o(w) \\ \mathcal{P}_{\blacktriangleleft a \cdot b}^\bullet &= \Gamma_a^{\otimes 2}(\mathcal{P}_{\blacktriangleleft b}^\bullet) \cdot \mathcal{P}_{\blacktriangleleft a}^\bullet & \bar{\mathcal{P}}_{a \cdot b^\blacktriangleright}^\bullet &= \bar{\mathcal{P}}_{a^\blacktriangleright}^\bullet \cdot \Gamma_a^{\otimes 2}(\bar{\mathcal{P}}_{b^\blacktriangleright}^\bullet) \end{aligned} \quad (9.39)$$

The relevant ideals for the intertwining relations of the truncated R -matrices are given for each $w \in \mathcal{W}^*$ as

$$\mathcal{N}(w) = \hat{\mathcal{K}}(w, \emptyset)_\ell \otimes U_{\zeta, \ell} + U_{\zeta, \ell} \otimes \hat{\mathcal{K}}(\emptyset, w)_\ell, \quad (9.40)$$

which is the ideal in $U_{\zeta, \ell}^{\otimes 2}$ generated by all $X_a \otimes 1$ and $1 \otimes Y_b$ with $\emptyset \neq a, b \leq_R w$. Assuming $w \cdot u$ is reduced for $w, u \in \mathcal{W}^*$, one easily verifies the identities

$$\mathcal{N}_o(w) \subset \mathcal{N}(w), \quad \mathcal{N}(w \cdot u) = \Gamma_w^{\otimes 2}(\mathcal{N}(u)) + \mathcal{N}(w), \quad \text{and} \quad \Omega^\#(\mathcal{N}(w)) = \mathcal{N}(w). \quad (9.41)$$

The truncated analog of Proposition 9.7 for a simple reflection w_i and simple generators is now stated as follows.

Lemma 9.16. *Let Φ be of any Lie type and $1 \leq i, j \leq n$. Then*

$$\Delta \circ \Gamma_i(E_j) \equiv \bar{\mathcal{P}}_i^\bullet \cdot \Gamma_i^{\otimes 2} \circ \Delta(E_j) \cdot \mathcal{P}_i^\bullet \quad \text{mod } \mathcal{N}(w_i). \quad (9.42)$$

The proof of this lemma is provided in Appendix A by explicit computations of both sides of (9.42), depending on the Cartan matrix entries A_{ij} and A_{ji} . The properties of automorphisms and ideals developed thus far allow us to extend this identity to all words and generators.

Proposition 9.17. *Let Φ be of any Lie type, $w \in \mathcal{W}^*$, and $x \in U_{\zeta, \ell}$. Then*

$$\Delta \circ \Gamma_w(x) \equiv \bar{\mathcal{P}}_{w\bullet}^\bullet \cdot \Gamma_w^{\otimes 2} \circ \Delta(x) \cdot \mathcal{P}_{\bullet w}^\bullet \quad \text{mod } \mathcal{N}(w).$$

Proof. We start with the case $w = w_i$ for which $\Gamma_w = \Gamma_i$ and $\mathcal{P}_{\bullet w_i}^\bullet = \mathcal{P}_i^\bullet$. Denote the quotient algebra $\mathbf{U}(w) = U_{\zeta, \ell}^{\otimes 2}/\mathcal{N}(w)$ and note that by Corollary 9.15 and (9.41) the images of $\bar{\mathcal{P}}_i^\bullet$ and \mathcal{P}_i^\bullet in $\mathbf{U}(w_i)$ are inverse to each other. The maps from $U_{\zeta, \ell}$ to $\mathbf{U}(w_i)$ given by $x \mapsto [\Delta \circ \Gamma_i(x)]$ and $x \mapsto [\bar{\mathcal{P}}_i^\bullet \cdot \Gamma_i^{\otimes 2} \circ \Delta(x) \cdot \mathcal{P}_i^\bullet]$ are thus algebra homomorphisms. Since, by Proposition 9.16, they coincide on generators, they have to be equal on all of $U_{\zeta, \ell}^+$.

Note next that $\Delta \circ \Gamma_i$ and $\Gamma_i^{\otimes 2} \circ \Delta$ coincide on any K^\vee and, further, that $K^\vee \otimes K^\vee$ commutes with \mathcal{P}_i^\bullet . This implies that the two homomorphisms also coincide on $U_{\zeta, \ell}^0$ and, hence, that the identity above holds for $x \in U_{\zeta, \ell}^{\geq 0}$ and $w = w_i$.

Recall that, with $\Omega^\#$ as in (9.32), $\Omega^\#(\Delta(x)) = \Delta(\Omega(x))$ and $\Omega^\#$ commutes with $\Gamma_i^{\otimes 2}$. It readily follows from the fact that $\Omega^\#$ is an anti-linear anti-involution, (4.10), and (9.34) that $\Omega^\#(\mathcal{P}_i^\bullet) = \bar{\mathcal{P}}_i^\bullet$. Applying $\Omega^\#$ to the identity in Proposition 9.16 for some $x \in U_{\zeta, \ell}^{\geq 0}$, these observations now imply the same relation for $\Omega(x) \in U_{\zeta, \ell}^{\geq 0}$, using also that $\mathcal{N}(w)$ is $\Omega^\#$ -invariant by (9.41). Hence, the relation holds for any $x \in U_{\zeta, \ell}$, completing the proof for $w = w_i$.

The extension to general words is similar to the one in Corollary 9.15. We proceed by induction on $L = l(w)$, starting from the base case already proven above. For a reduced word $u \in \mathcal{W}^*$ let $w = u^\flat$ and $i = \tau(u)$ so that $l(u) > l(w)$. Assuming the assertion for w , we first observe

$$\Delta \circ \Gamma_u(x) = \Delta \circ \Gamma_w(\Gamma_i(x)) \equiv \bar{\mathcal{P}}_{w\bullet}^\bullet \cdot \Gamma_w^{\otimes 2} \circ \Delta(\Gamma_i(x)) \mathcal{P}_{\bullet w}^\bullet \quad \text{mod } \mathcal{N}(w).$$

Applying the relation for w_i to the $\Delta(\Gamma_i(x))$ term, we find

$$\begin{aligned} \Delta \circ \Gamma_u(x) &\in \bar{\mathcal{P}}_{w\bullet}^\bullet \cdot \Gamma_w^{\otimes 2} \left(\bar{\mathcal{P}}_i^\bullet \cdot \Gamma_i^{\otimes 2}(\Delta(x)) \cdot \mathcal{P}_i^\bullet + \mathcal{N}(w_i) \right) \mathcal{P}_{\bullet w}^\bullet + \mathcal{N}(w) \\ &\subseteq \bar{\mathcal{P}}_{w\bullet}^\bullet \cdot \Gamma_w^{\otimes 2}(\bar{\mathcal{P}}_i^\bullet) \cdot \Gamma_{w \cdot w_i}^{\otimes 2}(\Delta(x)) \cdot \Gamma_w^{\otimes 2}(\mathcal{P}_{w_i}^\bullet) \cdot \mathcal{P}_{\bullet w}^\bullet + \bar{\mathcal{P}}_{w\bullet}^\bullet \cdot \Gamma_w^{\otimes 2}(\mathcal{N}(w_i)) \cdot \mathcal{P}_{\bullet w}^\bullet + \mathcal{N}(w) \\ &\subseteq \bar{\mathcal{P}}_{w \cdot w_i \bullet}^\bullet \cdot \Gamma_{w \cdot w_i}^{\otimes 2}(\Delta(x)) \cdot \mathcal{P}_{\bullet w \cdot w_i}^\bullet + \Gamma_w^{\otimes 2}(\mathcal{N}(w_i)) + \mathcal{N}(w) \\ &= \bar{\mathcal{P}}_{u\bullet}^\bullet \cdot \Gamma_u^{\otimes 2}(\Delta(x)) \cdot \mathcal{P}_{\bullet u}^\bullet + \mathcal{N}(u). \end{aligned}$$

The step from the second to the third line uses (9.39) for the first term and the fact that $\Gamma_w^{\otimes 2}(\mathcal{N}(w_i))$ is itself a two-sided ideal for the second term. The last step uses the iteration of ideals from (9.41) and $u = w \cdot w_i$. Taking the inclusion modulo $\mathcal{N}(u)$ yields the claim. \square

Recall from the proof of Lemma 9.8 that $\bar{\Delta}(x) = \Gamma_{s_*}^{\otimes 2}(\Delta(\Gamma_{s_*}^{-1}(x)))$ for the twisted coproduct defined in (9.14). The relation clearly descends to $U_{\zeta, \ell}$. Applying, Proposition 9.17 to a reduced word of maximal length now yields the following analog of Lemma 9.8.

Corollary 9.18. *Suppose $z \in \mathcal{W}_m^*$ and $x \in U_{\zeta, \ell}$. Then*

$$\bar{\Delta}(x) \cdot \mathcal{P}_{\blacktriangleleft z}^\bullet \equiv \mathcal{P}_{\blacktriangleleft z}^\bullet \cdot \Delta(x) \quad \text{mod } \mathcal{N}(z).$$

The iteration of the coproduct with truncated quasi- R -matrices provided in the above proposition allows us to establish further Hopf ideals. It is immediate from the form in (9.40) that

$$\mathcal{N}(u) \subseteq \mathcal{N}_*(u) = \hat{\mathcal{K}}(u, u)_\ell \otimes U_{\zeta, \ell} + U_{\zeta, \ell} \otimes \hat{\mathcal{K}}(u, u)_\ell, \quad (9.43)$$

which suggests that we first consider these diagonal ideals.

Proposition 9.19. *For every $u \in \mathcal{W}^*$ the ideal $\hat{\mathcal{K}}(u, u)_m$ is a Hopf ideal in $U_{\zeta, m}$.*

Proof. We start by showing that the $\hat{\mathcal{K}}(u, u)_\ell$ are bi-ideals. Given their definition, this means verifying that $\Delta(X_v), \Delta(Y_v) \in \mathcal{N}_*(u)$ for all $\emptyset \neq v \leq_R w$. Since $\mathcal{N}_*(v) \subseteq \mathcal{N}_*(u)$ and using Ω -symmetry as before, it suffices to check $\Delta(X_v) \in \mathcal{N}_*(v)$ for any reduced word v . Setting $w = v^\flat$ and $i = \tau(v)$ and using (7.6) we first observe that

$$\begin{aligned} \Gamma_w^{\otimes 2} \circ \Delta(X_i) &= \Gamma_w^{\otimes 2}(X_i \otimes L_i + 1 \otimes X_i) = X_{w \cdot w_i} \otimes L_{w \cdot w_i} + 1 \otimes X_{w \cdot w_i} \\ &= X_v \otimes L_v + 1 \otimes X_v \in \mathcal{N}_*(v). \end{aligned}$$

Using this, Proposition 9.17, the coproduct from (7.6), the fact that $\mathcal{N}_*(v)$ is a two-sided ideal in $U_{\zeta, \ell}^{\otimes 2}$, and $\mathcal{N}(w) \subseteq \mathcal{N}_*(v)$ we compute

$$\begin{aligned} \Delta(X_v) &= \Delta(X_{w \cdot w_i}) = \Delta \circ \Gamma_w(X_i) \in \bar{\mathcal{P}}_{w \blacktriangleright}^\bullet \cdot \Gamma_w^{\otimes 2} \circ \Delta(X_i) \cdot \mathcal{P}_{\blacktriangleleft w}^\bullet + \mathcal{N}(w) \\ &\subseteq \bar{\mathcal{P}}_{w \blacktriangleright}^\bullet(\mathcal{N}_*(v)) \mathcal{P}_{\blacktriangleleft w}^\bullet + \mathcal{N}(w) \subseteq \mathcal{N}_*(v), \end{aligned}$$

so that $\hat{\mathcal{K}}(u, u)_\ell$ is indeed a bi-ideal.

We now invoke a result of Nichols, which states that any bi-ideal in a pointed Hopf algebra is necessarily a Hopf ideal. See [Nic78, Thm 1(v)]. By Proposition 3.1, $U_{q, \mathbb{k}} \cong U_{\zeta, \mathbb{Q}} = U_{\zeta, \ell} \otimes \mathbb{Q}(\zeta)$ is pointed for $\mathbb{k} = \mathbb{Q}(\zeta)$ via the obvious map $\vartheta : \Lambda_{q, \mathbb{e}^*}^\bullet \rightarrow \mathbb{Q}(\zeta)$. Since $\hat{\mathcal{K}}(u, u)_\mathbb{Q} = \hat{\mathcal{K}}(u, u)_\ell \otimes \mathbb{Q}(\zeta)$ is a bi-ideal, Nichols's result implies that $\hat{\mathcal{K}}(u, u)_\mathbb{Q}$ is indeed a Hopf ideal, meaning it is preserved under the antipode S .

Finally, by (6.65) and (9.30), the ideals $\hat{\mathcal{K}}(u, v)_m$ are the free $\Lambda_{\zeta, m}^\bullet$ -submodules spanned by a subset of elements of a PBW basis of $U_{\zeta, m}$, implying $\hat{\mathcal{K}}(u, v)_m = \hat{\mathcal{K}}(u, v)_\mathbb{Q} \cap U_{\zeta, m}$. Since the intersection of a Hopf subalgebra and Hopf ideal is again a Hopf ideal, we arrive at the claim for $\hat{\mathcal{K}}(u, u)_m$. \square

We next seek to extend the observation above to the related families of ideals defined in previous sections. The two-sided ideals $\hat{\mathcal{K}}(w)^+$ defined in (6.65) are not Hopf ideals since U_{ζ, e^*}^+ is not a bialgebra, nor are they even Hopf subalgebras. We, thus, consider the extension to the Hopf algebra $U_{\zeta, e^*}^{\geq 0}$ by including the Cartan generators, and defining

$$\hat{\mathcal{K}}(w)^{\geq 0} = \hat{\mathcal{K}}(w)^+ \cdot U_{\zeta, e^*}^{\geq 0} \quad \text{with } \Lambda_{\zeta, e^*}\text{-basis } \{b \cdot K^\nu \mid \nu \in \mathbb{Z}^\Delta, b \in \mathfrak{B}_{z\blacktriangleright}^{i^*([l, \infty)^s), +}\}. \quad (9.44)$$

Here, $z \in \mathcal{W}_m^*$ with $w \leq_R z$, $s = r(w)$, and $\iota : \mathcal{N}(s) \hookrightarrow \Phi^+$. The definition of $\hat{\mathcal{K}}(w)^{\leq 0}$ is analogous and the $\hat{\mathcal{K}}(u, v)_m$ are obtained from the $\hat{\mathcal{K}}(w)^{\geq 0, \leq 0}$ in the same way as in (9.30).

The statement in Item *i*) in Theorem 9.20 already follows, independently, from the results in Sections 7 and 8 for the respective Lie types A_n and B_2 , where the ideals are explicitly identified with vanishing ideals of Bruhat subgroups.

Word independence, asserted in Item *iii*), is already implied by Theorem 6.19 for Lie types different from G_2 . The use of Nicols's result for Hopf ideals in Proposition 9.19 allows us to extend word independence also to the G_2 case for the ideals in the full algebras.

Theorem 9.20. *Suppose Φ is of any Lie type and $e^* \leq m \leq \ell$. Assume further $u, v, u', v' \in \mathcal{W}^*$ are reduced words with $s = r(u) = r(u')$ and $t = r(v) = r(v')$.*

i) The ideals $\hat{\mathcal{K}}(u)^{\geq 0}$ and $\hat{\mathcal{K}}(u)^{\leq 0}$ are Hopf ideals in $U_{\zeta, e^}^{\geq 0}$ and $U_{\zeta, e^*}^{\leq 0}$, respectively.*

ii) The ideal $\hat{\mathcal{K}}(u, v)_m$ is a Hopf ideal in $U_{\zeta, m}$.

iii) The ideals depend only on the elements represented in \mathcal{W} .

That is, $\hat{\mathcal{K}}(u)^{\geq 0} = \hat{\mathcal{K}}(u')^{\geq 0}$, $\hat{\mathcal{K}}(u)^{\leq 0} = \hat{\mathcal{K}}(u')^{\leq 0}$, and $\hat{\mathcal{K}}(u, v)_m = \hat{\mathcal{K}}(u', v')_m$.

Proof. The same reasoning as in the proof of Proposition 9.19 applies to show that $\hat{\mathcal{K}}(u)^{\geq 0} = \hat{\mathcal{K}}(u, u)_m \cap U_{\zeta, e^*}^{\geq 0}$, by restricting both the basis elements defined with respect to u to those in (9.44) and reducing the ground ring to Λ_{ζ, e^*} . As before, since $U_{\zeta, e^*}^{\geq 0}$ is a Hopf subalgebra it follows from Proposition 9.19 that $\hat{\mathcal{K}}(u)^{\geq 0}$ is a Hopf ideal. The argument for $\hat{\mathcal{K}}(u)^{\leq 0}$ is identical, and the assertion in Item *ii*) is now immediate from the definition in (9.30).

The statement in Item *i*) implies, in particular, that for any $w \in \mathcal{W}^*$, $\hat{\mathcal{K}}(w)^{\geq 0}$ is invariant under the antipode. The formulae in (4.16) and the fact that the Ψ_u^h preserves any subspace of $U_{\zeta, \ell}^{\geq 0}$ with a w -graded decomposition implies that \mathfrak{O} maps $\hat{\mathcal{K}}(w)^{\geq 0}$ to itself. The decomposition into a positive and Cartan part now implies $\mathfrak{O}(\hat{\mathcal{K}}(w)^+) = \hat{\mathcal{K}}(w)^+$.

Relation (6.65) shows that word independence of $\hat{\mathcal{K}}(w)^+$ is implied by word independence of $\mathfrak{S}_{w\blacktriangleright}^{[l, \infty)^s, +}$. To prove the latter we first observe that $V^{[s]} = [0, l]_s$ is of the split form (5.65) with $V_d = \{0, 1, \dots, \ell_d - 1\}$. The exponent set $\bar{V}^{[s]} = [l, \infty)^s$ is defined as the complement and thus of the required in the case *ii*) of Corollary 5.15.

As in the remark following (6.65), we have $\hat{\mathcal{K}}(z_{[ij]})^+ = \mathfrak{S}_{z_{[ij]}\blacktriangleright}^{[l, \infty)^s, +} = \mathfrak{S}_{z_{[ij]}\blacktriangleright}^{\bar{V}^{[ij]}, +}$ in $U_{\zeta, ij}^+$. By the previous argument we have that $\hat{\mathcal{K}}(z_{[ij]})^+$ is stable under \mathfrak{O} so that the second assumption of

Corollary 5.15 is also fulfilled. It follows that that $\hat{\mathcal{K}}(w)^+$ is independent of the choice of w . The respective statements for the other ideals are now immediate from the identities in (9.30) and (9.44). \square

Theorem 9.20 implies, in particular, that the ideals $\hat{\mathcal{K}}_{\bullet\bullet}$, $\hat{\mathcal{K}}_{\bullet\bullet}$, and $\hat{\mathcal{K}}_{\bullet\bullet}$ defined in (6.72) are indeed independent of choices of maximal reduced words as well, including the G_2 case.

The intertwining relations derived above may also be understood in terms of partial Hopf algebra quotients. For any $s \in \mathcal{W}$ and $w \in \mathcal{W}^*$ with $s = r(w)$ define

$$\Delta^s = \Gamma_s^{\otimes 2} \circ \Delta \circ \Gamma_s^{-1} \quad \text{and} \quad U_{\zeta, \ell}^{[s]} = U_{\zeta, \ell} / \hat{\mathcal{K}}(w, w)_\ell. \quad (9.45)$$

As with any automorphism, the conjugate Δ^s defines an alternative Hopf algebra structure with antipode S^{Γ_s} . Theorem 9.20 implies that $U_{\zeta, \ell}^{[s]}$ is a Hopf algebra defined independent of the choice of the presentation w of s .

Proposition 9.21. *Suppose $t = r(u)$ as above and $t \leq_R s$ for $s \in \mathcal{W}$. Let $\mathcal{P}_{\bullet u}^\star = [\mathcal{P}_{\bullet u}^\bullet]$ be the image of the partial quasi- R -matrix in $(U_{\zeta, \ell}^{[s]})^{\otimes 2}$. Then Δ^t descends to a coproduct $\Delta^{[t]}$ on $U_{\zeta, \ell}^{[s]}$, $\mathcal{P}_{\bullet u}^\star$ is invertible, and*

$$\Delta^{[t]}(x) \cdot \mathcal{P}_{\bullet u}^\star = \mathcal{P}_{\bullet u}^\star \cdot \Delta(x) \quad \text{for all } x \in U_{\zeta, \ell}^{[s]}.$$

Proof. Recall first from (9.41) and (9.43) that $\mathcal{N}_\circ(u) \subset \mathcal{N}(u) \subset \mathcal{N}_*(u) = \mathcal{N}_*(t) \subseteq \mathcal{N}_*(s)$. Thus, equations in Corollary 9.15 and Proposition 9.17 also hold modulo $\mathcal{N}_*(t)$. The former shows that $\mathcal{P}_{\bullet u}^\star$ is invertible. The latter can be rewritten as

$$\Delta^t(x) \equiv \mathcal{P}_{\bullet u}^\bullet \cdot \Delta(x) \cdot \bar{\mathcal{P}}_{\bullet u}^\bullet \quad \text{mod } \mathcal{N}_*(s).$$

If $x \in \hat{\mathcal{K}}(u, u)_\ell$, Proposition 9.19 implies that $\Delta(x) \in \mathcal{N}_*(s)$. Thus, the right-hand side and, hence, also $\Delta^t(x)$ is in $\mathcal{N}_*(s)$. It follows that Δ^t descends to a coproduct $\Delta^{[t]}$ on $U_{\zeta, \ell}^{[s]}$ for which the claimed identity now clearly holds. \square

It is important to keep in mind, though, that the automorphisms Γ_s themselves generally do not factor through these Hopf algebra quotients.

9.6. Quasi- R -Matrices for Restricted Quantum Algebras. In this section, we return to the maximal ideal case and collect previous observations for the construction of unique quasi- R -matrices for the restricted quantum algebras. The defining relations in tensor products of $U_{\zeta, \ell}$ will generally hold only up to ideals, which we introduce first. For $a \in \mathbb{N}$, let $\mathcal{N}_{\bullet\bullet}^a$ be the ideal in $U_{\zeta, \ell}^{\otimes a}$ generated by all elements $1^{\otimes(j-1)} \otimes X_v \otimes 1^{\otimes(a-j)}$ and $1^{\otimes(j-1)} \otimes Y_v \otimes 1^{\otimes(a-j)}$ for all $\emptyset \neq v \leq_R z$ and $j = 1, \dots, a$ for some $z \in \mathcal{W}_m^*$. Theorem 9.20 implies the word independent presentation

$$\mathcal{N}_{\bullet\bullet}^a = \hat{\mathcal{K}}_{\bullet\bullet} \otimes U_{\zeta, \ell}^{\otimes(a-1)} + U_{\zeta, \ell} \otimes \hat{\mathcal{K}}_{\bullet\bullet} \otimes U_{\zeta, \ell}^{\otimes(a-2)} + \dots + U_{\zeta, \ell}^{\otimes(a-1)} \otimes \hat{\mathcal{K}}_{\bullet\bullet}, \quad (9.46)$$

where $\hat{\mathcal{K}}_{\bullet\bullet}$ is as in (6.72). Since, for example, $\mathcal{N}(z) \subseteq \mathcal{N}_{\bullet\bullet}^2$, Corollary 9.18 implies

$$\bar{\Delta}(x) \cdot \mathcal{P}_{\bullet z}^\bullet \equiv \mathcal{P}_{\bullet z}^\bullet \cdot \Delta(x) \quad \text{mod } \mathcal{N}_{\bullet\bullet}^2, \quad (9.47)$$

for any $x \in U_{\zeta, \ell}$ and $\mathcal{P}_{\blacktriangleleft z}^*$ as in (9.37). The remainders in these ideals have been explicitly computed in Appendix A. Likewise, the Drinfeld-Tanisaki relations from (9.2) hold for the truncated R -matrices also only up to remainder terms in these ideals. For example, in the rank one case with $\mathcal{P}^* = \sum_{s=0}^{\ell-1} \mathbf{c}_s(\zeta)^{-1} E^s \otimes F^s$ we have

$$\Pi_{12}(\mathcal{P}_{13}^*) \cdot \mathcal{P}_{12}^* - (\text{id} \otimes \Delta)(\mathcal{P}^*) = (X \otimes 1 \otimes 1) \cdot \sum_{\substack{0 \leq s, t \leq \ell \\ s+t \geq \ell}} (\mathbf{c}_s(\zeta) \mathbf{c}_t(\zeta))^{-1} E^{t+s-\ell} \otimes K^{-s} F^t \otimes F^s,$$

which is clearly in $\mathcal{N}_{\bullet\bullet}^3$, but not zero. The general case follows similarly.

Proposition 9.22. *For a given maximal reduced word $z \in \mathcal{W}_m^*$ and any Lie type denote $\mathcal{P}^* = \mathcal{P}_{\blacktriangleleft z}^*$.*

$$\text{Then} \quad (\text{id} \otimes \Delta)(\mathcal{P}^*) \equiv \Pi_{12}(\mathcal{P}_{13}^*) \cdot \mathcal{P}_{12}^* \quad \text{mod } \mathcal{N}_{\bullet\bullet}^3$$

$$\text{and} \quad (\Delta \otimes \text{id})(\mathcal{P}^*) \equiv \Pi_{23}(\mathcal{P}_{13}^*) \cdot \mathcal{P}_{23}^* \quad \text{mod } \mathcal{N}_{\bullet\bullet}^3.$$

Proof. The first identity is straightforwardly verified by juxtaposing the explicit expressions for both sides using the expressions in (9.38) and (9.23), all of which specialize to $q \mapsto \zeta$.

$$\begin{aligned} (\text{id} \otimes \Delta)(\mathcal{P}^*) &= \sum_{\substack{\chi \in [0, \mathfrak{l}] \\ \phi, \psi \in \mathcal{E}}} \mathbf{c}_\chi(\zeta)^{-1} \mathbf{d}_{\phi, \psi}^\chi(\zeta) \cdot E_{\blacktriangleleft z}^\chi \otimes K^{-\bar{\psi}} F_{\blacktriangleleft z}^\phi \otimes F_{\blacktriangleleft z}^\psi \\ \Pi_{12}(\mathcal{P}_{13}^*) \cdot \mathcal{P}_{12}^* &= \sum_{\substack{\chi \in \mathcal{E} \\ \phi, \psi \in [0, \mathfrak{l}]}} \mathbf{c}_\phi(\zeta)^{-1} \mathbf{c}_\psi(\zeta)^{-1} \mathbf{m}_\chi^{\phi, \psi}(\zeta) \cdot E_{\blacktriangleleft z}^\chi \otimes K^{-\bar{\psi}} F_{\blacktriangleleft z}^\phi \otimes F_{\blacktriangleleft z}^\psi \end{aligned}$$

Recall here that $\mathbf{c}_\phi(\zeta) \neq 0$ iff $\phi \in [0, \mathfrak{l}]$, and observe that (9.24) has a well-defined specialization of q to ζ . We infer from this that for terms for which all three exponents χ, ϕ , and ψ are in $[0, \mathfrak{l}]$, the respective coefficients indeed coincide. The difference between the equations above is, thus, a sum of terms, each of which has at least one exponent in the complement $[\mathfrak{l}, \infty)^*$. The monomial element associated to this exponent is then, by Corollary 9.13, in $\hat{\mathcal{K}}_{\bullet\bullet}$ so that the full tensor term is in $\mathcal{N}_{\bullet\bullet}^3$.

This implies the first congruence above. The second identity follows analogously. An alternative argument can be designed by immediately passing to the quotient $U_{\zeta, \ell}^{\text{rest}}$ from (9.29) and observing that the pairing from (9.25) factors into a non-degenerate pairing between the positive and negative parts of $U_{\zeta, \ell}^{\text{rest}}$ upon the specialization $q \mapsto \zeta$. The image of \mathcal{P}^* in $(U_{\zeta, \ell}^{\text{rest}})^{\otimes 2}$ is then the element dual to this form. The duality relation (9.27) also factors into the form on $U_{\zeta, \ell}^{\text{rest}}$ and is equivalent to the Tanisaki relation on $U_{\zeta, \ell}^{\text{rest}}$. \square

Indeed, combining Proposition 9.22 and (9.47) implies that, for any maximal reduced word $z \in \mathcal{W}_m^*$, the triple $(U_{\zeta, \ell}^{\text{rest}}, [\mathcal{P}_{\blacktriangleleft z}^*], \Pi)$ is a pre-triangular Hopf algebra in the sense of (9.3), where $[\mathcal{P}_{\blacktriangleleft z}^*]$ denotes the image of $\mathcal{P}_{\blacktriangleleft z}^*$ in $(U_{\zeta, \ell}^{\text{rest}})^{\otimes 2}$. The remaining question is the word independence of $[\mathcal{P}_{\blacktriangleleft z}^*]$ or $\mathcal{P}_{\blacktriangleleft z}^*$ modulo $\mathcal{N}_{\bullet\bullet}^2$.

At this point, we have the tools to prove this claim in several different ways. One is to reconsider the pairing from (9.25), which is independent of a choice of $z \in \mathcal{W}_m^*$ over $\Lambda_{q,e}^*$ and, thus, also over $\Lambda_{\zeta,e}^*$ as a non-degenerate pairing of the positive and negative parts of $U_{\zeta,\ell}^{\text{rest}}$. Since the element $[\mathcal{P}_{\zeta,z}^*]$ is dual to this pairing it is also word independent. A second approach is to adapt the uniqueness proof of [Lus93, Thm.4.1.2(a)] to $U_{\zeta,\ell}^{\text{rest}}$ and use it directly to show the analog for Corollary 9.9 for $U_{\zeta,\ell}^{\text{rest}}$. We offer here a third option that exhibits further symmetries of the quasi- R -matrices.

Lemma 9.23. *Suppose $z \in \mathcal{W}_m^*$. Then $\mathcal{P}_{z^\dagger}^* = \mathfrak{U} \otimes \mathfrak{U}(\mathcal{P}_{z}^*) \equiv \mathcal{P}_{z}^* \pmod{\mathcal{N}_{\bullet\bullet}^2}$.*

Proof. Clearly, the automorphisms Π , S , and \mathfrak{U} preserve the maximal ideals and are, hence, well-defined on $U_{\zeta,\ell}^{\text{rest}}$. The Tanisaki relations from Proposition 9.22, therefore, hold as equalities in the restricted quantum group for the image $\mathcal{P}_z^* = [\mathcal{P}_{z}^*] \in (U_{\zeta,\ell}^{\text{rest}})^{\otimes 2}$. There, we may now repeat the arguments in (9.20) to show that $\tilde{\mathcal{P}}_z^* = [\tilde{\mathcal{P}}_{z}^*]$ and $\tilde{\mathcal{P}}_{z^\dagger}^*$ are both two-sided inverses of \mathcal{P}_z^* , where $\tilde{\mathcal{P}}_{z}^*$ denotes the truncated version of the element from (9.20).

Setting $\mathbf{f}(\mu) = (-1)^{\text{ht}(\mu)} \zeta^{-\frac{1}{2}((\mu|\mu)-2(\rho|\mu))}$ for $\mu \in \mathbb{Z}^\Delta$, we may re-express the implied equality $\tilde{\mathcal{P}}_z^* = \tilde{\mathcal{P}}_{z^\dagger}^*$ as the following congruence,

$$\sum_{\psi \in [0,1)} \mathbf{f}(\vec{\psi}) \mathbf{c}_\psi(\zeta)^{-1} \cdot E_{\zeta,z}^\psi \otimes F_{\zeta,z^\dagger}^\psi = \sum_{\psi \in [0,1)} \mathbf{f}(\vec{\psi}) \mathbf{c}_\psi(\zeta)^{-1} \cdot E_{\zeta,z^\dagger}^\psi \otimes F_{\zeta,z}^\psi \pmod{\mathcal{N}_{\bullet\bullet}^2}.$$

Applying to this the map $x \otimes y \mapsto \mathbf{f}(\mu)^{-1} x \otimes y$ for $\mathbf{w}(x) = \mu$ as well as $\mathfrak{U} \otimes \text{id}$, yields the claimed identity. \square

Corollary 9.24. *The element $\mathcal{P}^* = [\mathcal{P}_{z}^*] \in (U_{\zeta,\ell}^{\text{rest}})^{\otimes 2}$ is independent of the choice of $z \in \mathcal{W}_m^*$.*

Proof. By Matsumoto's Theorem, it suffices to show $\mathcal{P}_{z}^* \equiv \mathcal{P}_{w}^* \pmod{\mathcal{N}_{\bullet\bullet}^2}$ for maximal reduced words of the form $z = a \cdot z_{|ij|} \cdot b$ and $w = a \cdot z_{|ji|} \cdot b$. Following (9.39) we have

$$\mathcal{P}_{z}^* = \Gamma_{a \cdot z_{|ij|}}^{\otimes 2}(\mathcal{P}_{z_{|ij|}}^*) \cdot \Gamma_a^{\otimes 2}(\mathcal{P}_{z_{|ij|}}^*) \cdot \mathcal{P}_{a}^*,$$

and the analogous expression for \mathcal{P}_{w}^* . By Lemma 9.23 the difference between $\mathcal{P}_{z_{|ij|}}^*$ and $\mathcal{P}_{z_{|ji|}}^* = \mathfrak{U} \otimes \mathfrak{U}(\mathcal{P}_{z_{|ij|}}^*)$ is sum of terms of the form $(X_c \otimes 1)A$ and $(1 \otimes Y_c)B$ with $c \leq_R z_{|ij|}$. Since for these $a \cdot c$ is also reduced, we have $\Gamma_a(X_c) = X_{a \cdot c}$ and $\Gamma_a(Y_c) = Y_{a \cdot c}$. It follows from their above decompositions that the difference of \mathcal{P}_{z}^* and \mathcal{P}_{w}^* is a sum of terms of the form $(X_{a \cdot c} \otimes 1)A'$ and $(1 \otimes Y_{a \cdot c})B'$, which is, therefore, in $\mathcal{N}_{\bullet\bullet}^2$ by word independence of the ideal. \square

The findings of this section together with Corollary 9.13 can now be summarized as follows.

Theorem 9.25. *For any Lie type and root of unity ζ of order ℓ as in (6.1), the triple $(U_{\zeta,\ell}^{\text{rest}}, \mathcal{P}^*, \Pi)$ is a pre-triangular Hopf algebra over $\Lambda_{\zeta,\ell}^*$, independent of choices of maximal reduced words (or convex orderings) used in its construction.*

Moreover, all (anti)automorphisms defined in Section 4.1, including the Artin group action, factor into well-defined (anti)automorphisms on $U_{\zeta, \ell}^{\text{rest}}$. The construction is, thus, also independent of chosen directions of multiplications and independent of whether the Γ_i or T_i are used in the definition of generators.

The construction of braided tensor categories follows along the same steps as described in Section 9.3, where Σ is now allowed to have positive characteristic.

APPENDIX A. PROOF OF LEMMA 9.16

In this section, we prove Lemma 9.16. It suffices to show that equation (9.42) holds for each choice of $A_{ij} \in \{2, 0, -1, -2, -3\}$. The proof is carried out in a series of computations, with a subsection dedicated to each choice of A_{ij} . In most cases, we follow the same sequence of steps.

Proof. The $A_{ij} = 0$ case is obvious because \mathcal{P}_i^* commutes with $\Delta(E_j)$ and since the action $\Gamma_i(E_j) = E_j$ is trivial.

In the remaining cases, we make use of the fact that the coefficients $a_s = \frac{\zeta_i^{s(s-1)/2}}{[s]_i!}(\zeta_i - \zeta_i^{-1})^s$ of $\mathcal{P}_i^* = \sum_{s=0}^{\ell_i-1} a_s \cdot E_i^s \otimes F_i^s$ satisfy the recursion relation

$$a_{s+1} = \frac{\zeta_i^s(\zeta_i - \zeta_i^{-1})}{[s+1]_i} a_s. \quad (\text{A.1})$$

A.1. Case $A_{ij} = 2$. In this case, we write $i = j$. Using (4.19), equation (9.42) then reduces to

$$-K_i^{-1}F_i \otimes K_i^{-1} - K_i^{-2} \otimes K_i^{-1}F_i \equiv \bar{\mathcal{P}}_i^* \cdot (-K_i^{-1}F_i \otimes K_i^{-1} - 1 \otimes K_i^{-1}F_i) \cdot \mathcal{P}_i^* \quad \text{mod } \mathcal{N}(w_i).$$

We will use the commutation relation

$$[F_i, E_i^s] = -[s]_i \frac{\zeta_i^{s-1} K_i - \zeta_i^{-(s-1)} K_i^{-1}}{\zeta_i - \zeta_i^{-1}} E_i^{s-1},$$

which follows from a straightforward induction. Applying these commutators to each summand of the R -matrix we find the relations

$$\begin{aligned} (K_i^{-1}F_i \otimes K_i^{-1})\mathcal{P}_i^* &= \mathcal{P}_i^*(K_i^{-1}F_i \otimes K_i^{-1}) - \sum_{s=0}^{\ell_i-2} a_s(E_i^s \otimes F_i^s) ((\zeta_i^{2s} - K_i^{-2}) \otimes K_i^{-1}F_i) \\ &= \mathcal{P}_i^*(K_i^{-1}F_i \otimes K_i^{-1} + K_i^{-2} \otimes K_i^{-1}F_i) - \sum_{s=0}^{\ell_i-2} a_s \zeta_i^{2s} (E_i^s \otimes F_i^s) (1 \otimes K_i^{-1}F_i), \\ (1 \otimes K_i^{-1}F_i)\mathcal{P}_i^* &= \sum_{s=0}^{\ell_i-1} a_s \zeta_i^{2s} (E_i^s \otimes F_i^s) (1 \otimes K_i^{-1}F_i). \end{aligned}$$

Combining terms now yields

$$(-K_i^{-1}F_i \otimes K_i^{-1} - 1 \otimes K_i^{-1}F_i)\mathcal{P}_i^* = \mathcal{P}_i^*(-K_i^{-1}F_i \otimes K_i^{-1} - K_i^{-2} \otimes K_i^{-1}F_i) + a_{\ell_i-1} q^{2\ell_i} E_i^{\ell_i-1} K_i^{-2} \otimes F_i^{\ell_i} K_i^{-1},$$

which implies the desired equality modulo the ideal $\mathcal{N}(w_i)$.

A.2. Case $A_{ij} = -1$. In this case $E_{(ij)} = -E_j E_i + \zeta^{-1} E_i E_j$. We use the following relations from [Lus90b, Section 5.3].

$$E_{(ij)} E_i = \zeta E_i E_{(ij)}, \quad [E_{(ij)}, F_i^s] = -\zeta^{1-s} [s]_i F_i^{s-1} K_i E_j,$$

with the latter relation obtained by applying $\Gamma_i \circ \mathfrak{U}$ to Lusztig's equation (c) with $k' = 1$. These give the following relations between \mathcal{P}_i^* and terms in $\Gamma_i^{\otimes 2} \circ \Delta(E_j)$.

$$\begin{aligned}
(E_{(ij)} \otimes K_i K_j) \mathcal{P}_i^* &= \mathcal{P}_i^* (E_{(ij)} \otimes K_i K_j), \\
(1 \otimes E_{(ij)}) \mathcal{P}_i^* &= \mathcal{P}_i^* (1 \otimes E_{(ij)}) - (\zeta - \zeta^{-1}) \sum_{s=0}^{\ell_i-2} a_s (E_i^s \otimes F_i^s) (E_i \otimes K_i E_j), \\
\bar{\mathcal{P}}_i^* \cdot \Gamma_i^{\otimes 2} \circ \Delta(E_j) \mathcal{P}_i^* &= \bar{\mathcal{P}}_i^* (E_{(ij)} \otimes K_i K_j + 1 \otimes E_{(ij)}) \mathcal{P}_i^* \\
&= E_{(ij)} \otimes K_i K_j + 1 \otimes E_{(ij)} - (\zeta - \zeta^{-1}) E_i \otimes K_i E_j \\
&\quad + \bar{\mathcal{P}}_i^* (\zeta - \zeta^{-1}) a_{\ell_i-1} (E_i^{\ell_i} \otimes F_i^{\ell_i-1} K_i E_j) \\
&\equiv E_{(ij)} \otimes K_i K_j + 1 \otimes E_{(ij)} - (\zeta - \zeta^{-1}) E_i \otimes K_i E_j \quad \text{mod } \mathcal{N}(w_i).
\end{aligned}$$

This expression agrees with the coproduct of $E_{(ij)}$ computed below and, therefore, verifies the claim in this case.

$$\begin{aligned}
\Delta \circ \Gamma_i(E_j) &= \begin{cases} -(E_j E_i \otimes K_i K_j + E_j \otimes K_j E_i + E_i \otimes E_j K_i + 1 \otimes E_j E_i) \\ + \zeta^{-1} (E_i E_j \otimes K_i K_j + E_j \otimes E_i K_j + E_i \otimes K_i E_j + 1 \otimes E_i E_j) \end{cases} \\
&= E_{(ij)} \otimes K_i K_j + 1 \otimes E_{(ij)} - (\zeta - \zeta^{-1}) E_i \otimes K_i E_j
\end{aligned}$$

A.3. Case $A_{ij} = -2$. The following equations are obtained from

$$E_{(ij)} = \zeta^{-2} E_i^{(2)} E_j - \zeta^{-1} E_i E_j E_i + E_j E_i^{(2)}$$

as well as the ones listed in [Lus90b, Section 5.3].

$$\begin{aligned}
E_{(ij)} E_i &= \zeta^2 E_i E_{(ij)} \\
[E_{(ij)}, F_i^s] &= -[s] \zeta^{1-s} F_i^{s-1} K_i E_{(ij)} + [s][s-1] \zeta^{-2s+4} F_i^{s-2} K_i^2 E_j
\end{aligned}$$

The latter equation is obtained by applying $\Gamma_i \circ \mathfrak{U}$ to Lusztig's equation (i). We apply these next to compute the commutator of \mathcal{P}_i^* with $\Gamma^{\otimes 2} \circ \Delta(E_j)$.

$$\begin{aligned}
(E_{(ij)} \otimes K_i^2 K_j) \mathcal{P}_i^* &= \mathcal{P}_i^* (E_{(ij)} \otimes K_i^2 K_j), \\
(1 \otimes E_{(ij)}) \mathcal{P}_i^* &= \mathcal{P}_i^* (1 \otimes E_{(ij)}) - (\zeta - \zeta^{-1}) \sum_{s=0}^{\ell_i-2} a_s (E_i^s \otimes F_i^s) (E_i \otimes K_i E_{(ij)}) \\
&\quad + \zeta (\zeta - \zeta^{-1})^2 \sum_{s=0}^{\ell_i-3} a_s (E_i^s \otimes F_i^s) (E_i^2 \otimes K_i^2 E_j).
\end{aligned}$$

We infer from these relations that $\bar{\mathcal{P}}_i^\bullet \cdot \Gamma_i^{\otimes 2} \circ \Delta(E_j) \mathcal{P}_i^\bullet$ equals

$$\begin{aligned} & E_{(ij)} \otimes K_i^2 K_j + 1 \otimes E_{(ij)} - (\zeta - \zeta^{-1}) E_i \otimes K_i E_{(iji)} + \zeta(\zeta - \zeta^{-1})^2 E_i^2 \otimes K_i^2 E_j \\ & + \bar{\mathcal{P}}_i^\bullet \left(a_{\ell-1} (\zeta - \zeta^{-1}) E_i^\ell \otimes F_i^{\ell-1} K_i E_{(iji)} - \zeta(\zeta - \zeta^{-1})^2 \sum_{s=\ell-2}^{\ell-1} a_s E_i^{s+2} \otimes F_i^s K_i^2 E_j \right) \\ & \equiv E_{(ij)} \otimes K_i^2 K_j + 1 \otimes E_{(ij)} - (\zeta - \zeta^{-1}) E_i \otimes K_i E_{(iji)} + \zeta(\zeta - \zeta^{-1})^2 E_i^2 \otimes K_i^2 E_j \quad \bmod \mathcal{N}(w_i). \end{aligned}$$

Since

$$\begin{aligned} \Delta \circ \Gamma_i(E_j) &= \begin{cases} \zeta^{-2} \left(E_i^{(2)} E_j \otimes K_i^2 K_j + \zeta E_i E_j \otimes E_i K_i K_j + E_j \otimes E_i^{(2)} K_j \right. \\ \left. + E_i^{(2)} \otimes K_i^2 E_j + \zeta_i E_i \otimes E_i K_i E_j + 1 \otimes E_i^{(2)} E_j \right) \\ - \zeta^{-1} \left(E_i E_j E_i \otimes K_i^2 K_j + E_i E_j \otimes K_i K_j E_i + E_i^2 \otimes K_i E_j K_i + E_j E_i \otimes E_i K_i K_j \right. \\ \left. + E_i \otimes K_i E_j E_i + E_j \otimes E_i K_j E_i + E_i \otimes E_i E_j K_i + 1 \otimes E_i E_j E_i \right) \\ + \left(E_j E_i^{(2)} \otimes K_j K_i^2 + \zeta E_j E_i \otimes K_j E_i K_i + E_j \otimes K_j E_i^{(2)} \right. \\ \left. + E_i^{(2)} \otimes E_j K_i^2 + \zeta E_i \otimes E_j E_i K_i + 1 \otimes E_j E_i^{(2)} \right) \end{cases} \\ &= E_{(ij)} \otimes K_j K_i^2 + 1 \otimes E_{(ij)} - (\zeta - \zeta^{-1}) E_i \otimes (K_i(\zeta^{-2} E_i E_j - E_j E_i)) + \zeta(\zeta - \zeta^{-1})^2 E_i^2 \otimes K_i^2 E_j \\ &= E_{(ij)} \otimes K_i^2 K_j + 1 \otimes E_{(ij)} - (\zeta - \zeta^{-1}) E_i \otimes K_i E_{(iji)} + \zeta(\zeta - \zeta^{-1})^2 E_i^2 \otimes K_i^2 E_j, \end{aligned}$$

this proves the case for $A_{ij} = -2$.

A.4. Case $A_{ij} = -3$. In this case,

$$E_{(ij)} = \Gamma_i(E_j) = \zeta^{-3} E_i^{(3)} E_j - \zeta^{-2} E_i^{(2)} E_j E_i + \zeta^{-1} E_i E_j E_i^{(2)} - E_j E_i^{(3)} = (T_{w_j w_i})^2(E_j).$$

Below, we refer to Lusztig's equations (a1) and (a3) in [Lus90b, Section 5.4].

$$\begin{aligned} E_{(ij)} E_i &= \zeta^3 E_i E_{(ij)} \\ [E_{(ij)}, F_i^s] &= -[s] \zeta^{1-s} F_i^{s-1} K_i E_{(iji)} + [s]! \zeta^{4-2n} F_i^{(s-2)} K_i^2 E_{(ijiji)} - \zeta^{9-3n} [s]! F_i^{(s-3)} K_i^3 E_j \end{aligned}$$

These relations yield expressions for conjugation of $\Gamma_i^{\otimes 2} \circ \Delta(E_j) = E_{(ij)} \otimes K_i^3 K_j + 1 \otimes E_{(ij)}$ by \mathcal{P}_i^\bullet via the following computation.

$$\begin{aligned} (E_{(ij)} \otimes K_i^3 K_j) \mathcal{P}_i^\bullet &= \mathcal{P}_i^\bullet (E_{(ij)} \otimes K_i^3 K_j), \\ (1 \otimes E_{(ij)}) \mathcal{P}_i^\bullet &= \mathcal{P}_i^\bullet (1 \otimes E_{(ij)}) - (\zeta - \zeta^{-1}) \sum_{s=0}^{\ell-2} a_s (E_i^s \otimes F_i^s) (E_i \otimes K_i E_{(iji)}) \\ &+ \zeta(\zeta - \zeta^{-1})^2 \sum_{s=0}^{\ell-3} a_s (E_i^s \otimes F_i^s) (E_i^2 \otimes K_i^2 E_{(ijiji)}) \\ &- \zeta^3 (\zeta - \zeta^{-1})^3 \sum_{s=0}^{\ell-4} a_s (E_i^s \otimes F_i^s) (E_i^3 \otimes K_i^3 E_j), \end{aligned}$$

$$\begin{aligned}
\bar{\mathcal{P}}_i^\bullet \cdot \Gamma_i^{\otimes 2} \circ \Delta(E_j) \mathcal{P}_i^\bullet &= E_{(ij)} \otimes K_i^3 K_j + 1 \otimes E_{(ij)} - (\zeta - \zeta^{-1}) E_i \otimes K_i E_{(iji)} \\
&\quad + \zeta(\zeta - \zeta^{-1})^2 E_i^2 \otimes K_i^2 E_{(ijiji)} - \zeta^3(\zeta - \zeta^{-1})^3 E_i^3 \otimes K_i^3 E_j \\
&\quad + \bar{\mathcal{P}}_i^\bullet \left(\begin{array}{l} (\zeta - \zeta^{-1}) a_{\ell-1} E_i^\ell \otimes F_i^{\ell-1} K_i E_{(iji)} \\ - \zeta(\zeta - \zeta^{-1})^2 \sum_{s=\ell-2}^{\ell-1} a_s (E_i^{s+2} \otimes F_i^s K_i^2 E_{(ijiji)}) \\ + \zeta^3(\zeta - \zeta^{-1})^3 \sum_{s=\ell-3}^{\ell-1} a_s (E_i^{s+3} \otimes F_i^s K_i^3 E_j) \end{array} \right) \\
&\equiv E_{(ij)} \otimes K_i^3 K_j + 1 \otimes E_{(ij)} - (\zeta - \zeta^{-1}) E_i \otimes K_i E_{(iji)} \\
&\quad + \zeta(\zeta - \zeta^{-1})^2 E_i^2 \otimes K_i^2 E_{(ijiji)} - \zeta^3(\zeta - \zeta^{-1})^3 E_i^3 \otimes K_i^3 E_j \quad \text{mod } \mathcal{N}(w_i).
\end{aligned}$$

It remains to verify the coproduct identity

$$\begin{aligned}
\Delta \circ \Gamma_i(E_j) = \Delta(E_{(ij)}) &= E_{(ij)} \otimes K_j K_i^3 + 1 \otimes E_{(ij)} - (\zeta - \zeta^{-1}) E_i \otimes K_i E_{(iji)} \\
&\quad + \zeta(\zeta - \zeta^{-1})^2 E_i^2 \otimes K_i^2 E_{(ijiji)} - \zeta^3(\zeta - \zeta^{-1})^3 E_i^3 \otimes K_i^3 E_j.
\end{aligned}$$

We begin by expanding the coproduct for the divided powers expressions in $\Delta(E_{(ij)})$:

$$\begin{aligned}
&\zeta^{-3} \left(\begin{array}{l} E_i^{(3)} E_j \otimes K_i^3 K_j + \zeta^2 E_i^{(2)} E_j \otimes E_i K_i^2 K_j + \zeta^2 E_i E_j \otimes E_i^{(2)} K_i K_j + E_j \otimes E_i^{(3)} K_j \\ + E_i^{(3)} \otimes K_i^3 E_j + \zeta^2 E_i^{(2)} \otimes E_i K_i^2 E_j + \zeta^2 E_i \otimes E_i^{(2)} K_i E_j + 1 \otimes E_i^{(3)} E_j \end{array} \right) \\
&- \left(\begin{array}{l} \zeta^{-2} \left(\begin{array}{l} E_i^{(2)} E_j \otimes K_i^2 K_j + \zeta E_i E_j \otimes E_i K_i K_j + E_j \otimes E_i^{(2)} K_j \\ + E_i^{(2)} \otimes K_i^2 E_j + \zeta E_i \otimes E_i K_i E_j + 1 \otimes E_i^{(2)} E_j \end{array} \right) \\ - \frac{\zeta^{-1}}{[2]} \left(\begin{array}{l} E_i E_j E_i \otimes K_i^2 K_j + E_i E_j \otimes K_i K_j E_i + E_i^2 \otimes K_i E_j K_i + E_j E_i \otimes E_i K_i K_j \\ + E_i \otimes K_i E_j E_i + E_j \otimes E_i K_j E_i + E_i \otimes E_i E_j K_i + 1 \otimes E_i E_j E_i \end{array} \right) \\ + \frac{1}{[3]} \left(\begin{array}{l} E_j E_i^{(2)} \otimes K_j K_i^2 + \zeta E_j E_i \otimes K_j E_i K_i + E_j \otimes K_j E_i^{(2)} \\ + E_i^{(2)} \otimes E_j K_i^2 + \zeta E_i \otimes E_j E_i K_i + 1 \otimes E_j E_i^{(2)} \end{array} \right) \end{array} \right) \Delta(E_i)
\end{aligned}$$

$$= E_{(ij)} \otimes K_j K_i^3 + 1 \otimes E_{(ij)}$$

$$\begin{aligned}
& + \zeta^{-3} \left(\begin{array}{l} \zeta^2 E_i^{(2)} E_j \otimes E_i K_i^2 K_j + \zeta^2 E_i E_j \otimes E_i^{(2)} K_i K_j + E_j \otimes E_i^{(3)} K_j \\ + E_i^{(3)} \otimes K_i^3 E_j + \zeta^2 E_i^{(2)} \otimes E_i K_i^2 E_j + \zeta^2 E_i \otimes E_i^{(2)} K_i E_j \end{array} \right) \\
& + \left(\begin{array}{l} -\zeta^{-2} E_i^{(2)} E_j \otimes K_i^2 K_j E_i - \zeta^{-2} E_i \otimes E_i^{(2)} E_j K_i + \frac{\zeta^{-1}}{[2]} E_i E_j E_i \otimes K_i^2 K_j E_i \\ + \frac{\zeta^{-1}}{[2]} E_i \otimes E_i E_j E_i K_i - \frac{1}{[3]} E_j E_i^{(2)} \otimes K_j K_i^2 E_i - \frac{1}{[3]} E_i \otimes E_j E_i^{(2)} K_i \end{array} \right) \\
& - \left(\begin{array}{l} \zeta^{-2} \left(\begin{array}{l} \zeta E_i E_j \otimes E_i K_i K_j + E_j \otimes E_i^{(2)} K_j \\ + E_i^{(2)} \otimes K_i^2 E_j + \zeta E_i \otimes E_i K_i E_j \end{array} \right) \\ - \frac{\zeta^{-1}}{[2]} \left(\begin{array}{l} E_i E_j \otimes K_i K_j E_i + E_i^2 \otimes K_i E_j K_i + E_j E_i \otimes E_i K_i K_j \\ + E_i \otimes K_i E_j E_i + E_j \otimes E_i K_j E_i + E_i \otimes E_i E_j K_i \end{array} \right) \\ + \frac{1}{[3]} \left(\begin{array}{l} \zeta E_j E_i \otimes K_j E_i K_i + E_j \otimes K_j E_i^{(2)} \\ + E_i^{(2)} \otimes E_j K_i^2 + \zeta E_i \otimes E_j E_i K_i \end{array} \right) \end{array} \right) \Delta(E_i) \\
& = E_{(ij)} \otimes K_j K_i^3 + 1 \otimes E_{(ij)} \\
& + \zeta^{-3} \left(\begin{array}{l} \zeta E_i^{(2)} E_j \otimes K_i^2 K_j E_i + \zeta^4 E_i E_j \otimes K_i K_j E_i^{(2)} + \zeta^9 E_j \otimes K_j E_i^{(3)} \\ + E_i^{(3)} \otimes K_i^3 E_j + \zeta^{-2} E_i^{(2)} \otimes K_i^2 E_i E_j + \zeta^{-2} E_i \otimes K_i E_i^{(2)} E_j \end{array} \right) \\
& + \left(\begin{array}{l} -\zeta^{-2} E_i^{(2)} E_j \otimes K_i^2 K_j E_i - \zeta^{-3} E_i \otimes K_i E_i^{(2)} E_j + \frac{\zeta^{-1}}{[2]} E_i E_j E_i \otimes K_i^2 K_j E_i \\ + \frac{\zeta^{-2}}{[2]} E_i \otimes K_i E_i E_j E_i - \frac{1}{[3]} E_j E_i^{(2)} \otimes K_j K_i^2 E_i - \frac{\zeta^{-1}}{[3]} E_i \otimes K_i E_j E_i^{(2)} \end{array} \right) \\
& - \left(\begin{array}{l} \frac{\zeta}{[2]} E_i E_j \otimes K_i K_j E_i + \frac{\zeta^6}{[3]} E_j \otimes K_i E_i^{(2)} + \frac{\zeta^6 - \zeta^4 - \zeta^2 + \zeta^{-2} + \zeta^{-4}}{[3]} E_i^{(2)} \otimes K_i^2 E_j \\ + \frac{\zeta^{-4} + \zeta^{-2} - 1}{[2]} E_i \otimes K_i E_i E_j - \frac{\zeta^2}{[2][3]} E_j E_i \otimes K_i K_j E_i + \frac{\zeta^3 - \zeta^{-1} - \zeta^{-3}}{[2][3]} E_i \otimes K_i E_j E_i \end{array} \right) \Delta(E_i) \\
& = E_{(ij)} \otimes K_j K_i^3 + 1 \otimes E_{(ij)} - \zeta^{-4} (\zeta - \zeta^{-1}) E_i \otimes K_i E_i^{(2)} E_j \\
& + \zeta^{-3} \left(\begin{array}{l} \zeta^4 E_i E_j \otimes K_i K_j E_i^{(2)} + \zeta^9 E_j \otimes K_j E_i^{(3)} \\ + E_i^{(3)} \otimes K_i^3 E_j + \zeta^{-2} E_i^{(2)} \otimes K_i^2 E_i E_j \end{array} \right) + \left(\begin{array}{l} \frac{\zeta^{-1}}{[2]} E_i E_j E_i \otimes K_i^2 K_j E_i + \frac{\zeta^{-2}}{[2]} E_i \otimes K_i E_i E_j E_i \\ - \frac{1}{[3]} E_j E_i^{(2)} \otimes K_j K_i^2 E_i - \frac{\zeta^{-1}}{[3]} E_i \otimes K_i E_j E_i^{(2)} \end{array} \right) \\
& - \left(\begin{array}{l} \frac{\zeta^{-1}}{[2]} E_i E_j E_i \otimes K_i^2 K_j E_i + \frac{\zeta}{[2]} E_i E_j \otimes K_i K_j E_i^2 + \frac{\zeta^6}{[2][3]} E_j \otimes K_i E_i^3 \\ + \frac{\zeta^9 - \zeta^7 - \zeta^5 + \zeta + \zeta^{-1}}{[2][3]} E_i^3 \otimes K_i^3 E_j + \zeta (\zeta - \zeta^{-1}) E_i^2 \otimes K_i^2 E_j E_i + \frac{\zeta^{-3} + \zeta^{-1} - \zeta}{[2]} E_i^2 \otimes K_i^2 E_i E_j \\ + \frac{\zeta^{-4} + \zeta^{-2} - 1}{[2]} E_i \otimes K_i E_i E_j E_i - \frac{1}{[2][3]} E_j E_i^2 \otimes K_i^2 K_j E_i + \frac{\zeta^3 - \zeta^{-1} - \zeta^{-3}}{[2][3]} E_i \otimes K_i E_j E_i^2 \end{array} \right)
\end{aligned}$$

Further collapsing these expressions, we obtain

$$\begin{aligned}
\Delta(E_{(ij)}) &= E_{(ij)} \otimes K_j K_i^3 + 1 \otimes E_{(ij)} - \zeta^{-4}(\zeta - \zeta^{-1})E_i \otimes K_i E_i^{(2)} E_j - (\zeta - \zeta^{-1})E_i \otimes K_i E_j E_i^{(2)} \\
&\quad + \zeta^{-2}(\zeta - \zeta^{-1})E_i \otimes K_i E_i E_j E_i - \zeta(\zeta - \zeta^{-1})^2 E_i^2 \otimes K_i^2 E_j E_i \\
&\quad - \zeta^3(\zeta - \zeta^{-1})^3 E_i^3 \otimes K_i^3 E_j - \zeta^{-2}(\zeta - \zeta^{-1})^2 E_i^2 \otimes K_i^2 E_i E_j \\
&= E_{(ij)} \otimes K_j K_i^3 + 1 \otimes E_{(ij)} - (\zeta - \zeta^{-1})E_i \otimes K_i(\zeta^{-4}E_i^{(2)} E_j - \zeta^{-2}E_i E_j E_i + E_j E_i^{(2)}) \\
&\quad + \zeta(\zeta - \zeta^{-1})^2 E_i^2 \otimes K_i^2(\zeta^{-3}E_i E_j - E_j E_i) - \zeta^3(\zeta - \zeta^{-1})^3 E_i^3 \otimes K_i^3 E_j \\
&= E_{(ij)} \otimes K_j K_i^3 + 1 \otimes E_{(ij)} - (\zeta - \zeta^{-1})E_i \otimes K_i E_{(iji)} \\
&\quad + \zeta(\zeta - \zeta^{-1})^2 E_i^2 \otimes K_i^2 E_{(iji)} - \zeta^3(\zeta - \zeta^{-1})^3 E_i^3 \otimes K_i^3 E_j.
\end{aligned}$$

This now proves $\Delta \circ \Gamma_i(E_j) \equiv \bar{\mathcal{P}}_i^\bullet \cdot \Gamma_i^{\otimes 2} \circ \Delta(E_j) \cdot \mathcal{P}_i^\bullet \pmod{\mathcal{N}(w_i)}$. \square

APPENDIX B. PROOFS AND COMPUTATIONS FOR SECTION 8.2

We supply here the computational proofs of Proposition 8.3 and Lemma B.3 used in Sections 8.1 and 8.2.

B.1. Proof of Proposition 8.3. We begin with a variation of the recursive identity used to define the coefficients $c_{k,t}$ in (8.13) that applies more directly to the desired proof. To this end we introduce notation for the coefficients actually appearing in the formula of Proposition 8.3, namely,

$$a_{s,t,k}^n = q^{-(s+t+k)(n-s-t-k)-(s-t)(2t+k)} \begin{bmatrix} n \\ s+t+k \end{bmatrix} \begin{bmatrix} s+t+k \\ s-t \end{bmatrix} c_{k,t}, \quad (\text{B.1})$$

for $s \geq t \geq 0$, $k \geq 0$, and $n \geq s+t+k$. We further have $a_{0,0,0}^0 = 1$ and use the convention that $a_{s,t,k}^n = 0$ whenever any one of the index bounds is exceeded. That is, whenever

$$s, t, k < 0, \quad s > n, \quad t > \min(s, n-s), \quad \text{or} \quad k > n-s-t. \quad (\text{B.2})$$

Lemma B.1. *With definitions and conventions as above, we have the recursion*

$$a_{s,t,k}^{n+1} = q^{-2(n+1-s+t)} a_{s-1,t,k}^n + q^{-3(n+1-s-t-k)} [n+2-s-t-k] a_{s-1,t-1,k}^n + q^{-2(n+1-s-t-k)} a_{s,t,k-1}^n + a_{s,t,k}^n.$$

Proof. The following straightforward verification makes repeated use of the q -binomial formula from (3.2). To shorten some formulae, we use $u = s+t+k$.

$$\begin{aligned}
a_{s,t,k}^{n+1} &= q^{-u(n+1-u)-(s-t)(2t+k)} \begin{bmatrix} n+1 \\ u \end{bmatrix} \begin{bmatrix} u \\ s-t \end{bmatrix} c_{k,t} \\
&= q^{-u(n+1-u)-(s-t)(2t+k)} \left(q^u \begin{bmatrix} n \\ u \end{bmatrix} + q^{-(n+1-u)} \begin{bmatrix} n \\ u-1 \end{bmatrix} \right) \begin{bmatrix} u \\ s-t \end{bmatrix} c_{k,t}
\end{aligned}$$

$$\begin{aligned}
&= a_{s,t,k}^n + q^{-(u+1)(n+1-u)-(s-t)(2t+k)} \begin{bmatrix} n \\ u-1 \end{bmatrix} \left(q^{s-t} \begin{bmatrix} u-1 \\ s-t \end{bmatrix} + q^{-(2t+k)} \begin{bmatrix} u-1 \\ s-1-t \end{bmatrix} \right) c_{k,t} \\
&= a_{s,t,k}^n + q^{-2(n+1-s+t)} a_{s-1,t,k}^n \\
&\quad + q^{-(u+1)(n+1-u)-(s-t)(2t+k-1)} \begin{bmatrix} n \\ u-1 \end{bmatrix} \begin{bmatrix} u-1 \\ s-t \end{bmatrix} (c_{k-1,t} + q^{-(2t+k-2)} [2t+k-1] c_{k,t-1}) \\
&= a_{s,t,k}^n + q^{-2(n+1-s+t)} a_{s-1,t,k}^n + q^{-2(n+1-u)} a_{s,t,k-1}^n \\
&\quad + q^{-(u+1)(n+1-u)-(s-t+1)(2t+k-1)+1} [n+2-u] \begin{bmatrix} n \\ u-2 \end{bmatrix} \begin{bmatrix} u-2 \\ s-t \end{bmatrix} c_{k,t-1} \\
&= a_{s,t,k}^n + q^{-2(n+1-s+t)} a_{s-1,t,k}^n + q^{-2(n+1-u)} a_{s,t,k-1}^n + q^{-3(n+1-u)} [n+2-u] a_{s-1,t-1,k}^n.
\end{aligned}$$

□

Observe that we may also consider the $a_{s,t,k}^n$ to be defined via the recursion in Lemma B.1 and the mentioned boundary conditions. The following identity is proved in [Lus90a, Lemma 1.6] for the special case $v = 1$.

Lemma B.2. *Suppose A, B, D satisfy*

$$DA = v^2 AD + B, \quad BA = q^{-2} v^2 AB, \quad \text{and} \quad DB = q^{-2} v^2 BD.$$

Then $D^m A = v^{2m} AD^m + (q^{-1} v^2)^{m-1} [m] BD^{m-1}$ and

$$D^m A^n = \sum_{i=0}^{\min(m,n)} v^{2mn-i(i+1)} q^{-i(n+m-2i)-i(i-1)/2} \begin{bmatrix} n \\ i \end{bmatrix} \begin{bmatrix} m \\ i \end{bmatrix} [i]! A^{n-i} B^i D^{m-i},$$

where quantum numbers and coefficients are with respect to q .

Proof. Each of the two claims of the lemma is proved by induction. The base case of the former is the assumption $DA = v^2 AD + B$. For the induction step, we compute

$$\begin{aligned}
D^{m+1} A &= D(v^{2m} AD^m + (q^{-1} v^2)^{m-1} [m] BD^{m-1}) \\
&= v^{2(m+1)} AD^{m+1} + v^{2m} BD^m + (q^{-2} v^2)(q^{-1} v^2)^{m-1} [m] BD^m \\
&= v^{2(m+1)} AD^{m+1} + (q^{-1} v^2)^m [m+1] BD^m,
\end{aligned}$$

where the last equality uses $1 + q^{-(m+1)} [m] = q^{-m} [m+1]$. The second statement is now derived by induction in n using the first identity as the base case. It amounts to repeated use of (3.2) and resummations. □

Note that the q -binomials disappear in the formula of Lemma B.2 if it is rewritten in terms of divided power generators $A^{(n)} = \frac{1}{[n]!} A^n$. That is, we find instead

$$D^{(m)} A^{(n)} = \sum_{i=0}^{\min(m,n)} v^{2mn-i(i+1)} q^{i(n+m-2i)+i(i-1)/2} A^{(n-i)} B^{(i)} D^{(m-i)}.$$

With these preparations we proceed to the main goal of this section.

Proof of Proposition 8.3. Assume A , B , C , and D fulfill the commutation relations stated in the proposition. The assertion is trivial for $n = 0$ and easily worked out for $n = 1$. Assume, thus, that the formula hold for $(A + C + D)^n$, with coefficients $a_{s,t,k}^n$ as in (B.1). We compute

$$\begin{aligned}
(A + C + D)^{n+1} &= \sum_{s=0}^n \sum_{t=0}^{\min(s, n-s)} \sum_{k=0}^{n-s-t} a_{s,t,k}^n A^{s-t} B^t C^k D^{n-s-t-k} (A + C + D) \\
&= \sum_{s=0}^n \sum_{t=0}^{\min(s, n-s)} \sum_{k=0}^{n-s-t} a_{s,t,k}^n q^{-2(n-s+t)} A^{s-t+1} B^t C^k D^{n-s-t-k} \\
&\quad + a_{s,t,k}^n q^{-3(n-1-s-t-k)} [n-s-t-k] A^{s-t} B^{t+1} C^k D^{n-s-t-k-1} \\
&\quad + a_{s,t,k}^n q^{-2(n-s-t-k)} A^{s-t} B^t C^{k+1} D^{n-s-t-k} \\
&\quad + a_{s,t,k}^n A^{s-t} B^t C^k D^{n+1-s-t-k} \\
&= \sum_{s=1}^{n+1} \sum_{t=0}^{\min(s-1, n+1-s)} \sum_{k=0}^{n+1-s-t} a_{s-1,t,k}^n q^{-2(n+1-s+t)} A^{s-t} B^t C^k D^{n+1-s-t-k} \\
&\quad + \sum_{s=1}^{n+1} \sum_{t=1}^{\min(s, n+1-s)} \sum_{k=0}^{n+1-s-t} a_{s-1,t-1,k}^n q^{-3(n+1-s-t-k)} [n+2-s-t-k] A^{s-t} B^t C^k D^{n+1-s-t-k} \\
&\quad + \sum_{s=0}^n \sum_{t=0}^{\min(s, n-s)} \sum_{k=1}^{n+1-s-t} a_{s,t,k-1}^n q^{-2(n+1-s-t-k)} A^{s-t} B^t C^k D^{n+1-s-t-k} \\
&\quad + \sum_{s=0}^n \sum_{t=0}^{\min(s, n-s)} \sum_{k=0}^{n-s-t} a_{s,t,k}^n A^{s-t} B^t C^k D^{n+1-s-t-k} \\
&= \sum_{s=0}^{n+1} \sum_{t=0}^{\min(s, n+1-s)} \sum_{k=0}^{n+1-s-t} a_{s,t,k}^{n+1} A^{s-t} B^t C^k D^{n+1-s-t-k}.
\end{aligned}$$

In the first step, we employ the original commutation relations as well as Lemma B.2. In the second step, we shift summation indices and make use of the boundary conditions on the $a_{s,t,k}^n$. The last equation then results from Lemma B.1, completing the induction step. \square

B.2. A Summation Identity for the $c'_{k,p}$. As before, let $c_{k,p}$ be the coefficients defined recursively in (8.13) and $c'_{k,p} = \text{sq}(c_{k,p})$ the element in Λ_q with q replaced by q^2 as explained in Section 8.2.

Lemma B.3. *For all $n \in \mathbb{N}_0$, the following equality holds in $\Lambda_{q,3}$,*

$$\sum_{p=0}^{\lfloor n/2 \rfloor} \left(-\frac{q - q^{-1}}{q^2[2]} \right)^p c'_{n-2p,p} = q^{-\binom{n}{2}} [2]^{-n} \{n:2\}.$$

Proof. Recall that $\{n:2\} = \prod_{i=1}^n (q^i + q^{-i})$ under the specialization of $e = 2$ in (3.6). We prove that both expressions in the proposed equality are sequences in n with the same initial values and

satisfy the same recursion relation. Denote the sides of the desired equality by

$$A_n = \sum_{p=0}^{\lfloor n/2 \rfloor} \phi_p c'_{n-2p,p}, \quad \text{where } \phi_p = \left(-\frac{q-q^{-1}}{q^2[2]} \right)^p, \text{ and}$$

$$B_n = q^{-\binom{n}{2}} \psi_n, \quad \text{where } \psi_n = [2]^{-n} \{n : 2\}.$$

Observe that $A_0 = B_0 = A_1 = B_1 = 1$. The recursion for A_n is given below. Note that if n is even, $c'_{n-2(\lfloor n/2 \rfloor - 1),p} = 0$ and $n/2 - 1 = \lfloor (n-1)/2 \rfloor$. Whereas if n is odd, $\lfloor n/2 \rfloor = \lfloor (n-1)/2 \rfloor$ and $c'_{n-2(\lfloor n/2 \rfloor - 1),p} \neq 0$.

$$\begin{aligned} A_n &= \sum_{p=0}^{\lfloor n/2 \rfloor} \phi_p c'_{n-2p,p} = \sum_{p=0}^{\lfloor n/2 \rfloor} \phi_p (c'_{n-2p-1,p} + c'_{n-2p,p-1} q^{-2(n-2)} [n-1]_{q^2}) \\ &= A_{n-1} + q^{-2(n-2)} [n-1]_{q^2} \sum_{p=0}^{\lfloor n/2 \rfloor - 1} \phi_{p+1} c'_{n-2p-2,p} \\ &= A_{n-1} - q^{-2(n-2)} [n-1]_{q^2} \left(\frac{q-q^{-1}}{q^2[2]} \right) A_{n-2} \\ &= A_{n-1} - q^{-2(n-1)} [n-1]_{q^2} \frac{q-q^{-1}}{[2]} A_{n-2} \end{aligned}$$

Hence $A_n - A_{n-1} = -q^{-2(n-1)} [n-1]_{q^2} \frac{q-q^{-1}}{[2]} A_{n-2}$. An identical recursion for B_n is found by computing

$$\begin{aligned} B_n - B_{n-1} &= \psi_{n-2} [2]^{-1} (q^{n-1} + q^{-(n-1)}) \left(q^{-\binom{n}{2}} (q^n + q^{-n}) [2]^{-1} - q^{-\binom{n-1}{2}} \right) \\ &= \psi_{n-2} [2]^{-1} (q^{n-1} + q^{-(n-1)}) q^{-\binom{n-2}{2}} (q^{-2n+3} (q^n + q^{-n}) [2]^{-1} - q^{-n+2}) \\ &= B_{n-2} \frac{[n-1]_{q^2}}{[2][n-1]} ((q^{-n+3} + q^{-3n+3}) - (q^{-n+3} + q^{-n+1})) \\ &= -q^{-2(n-1)} \frac{[n-1]_{q^2}}{[2]} (q - q^{-1}) B_{n-2}. \end{aligned}$$

It follows that $A_n = B_n$ by strong induction, which is the identity claimed in the lemma. \square

APPENDIX C. INEXISTENCE OF QUANTUM WEYL ELEMENTS AND RELATED NOTES

In the finite-dimensional representation theory of classical, simple Lie algebras, Weyl elements are formal elements that are instrumental in the derivation of various dimension and character formulae. For a simply connected simple Lie group G with maximal torus $T \subset G$ the geometric Weyl group \mathcal{W} , defined as $N(T)/T$, is acting on \mathfrak{h} , the Lie algebra of T , via conjugation by representatives in the normalizer $N(T)$. The action of a standard generator $s_i \in \mathcal{W}$ is given by $S_i = \text{Ad}(\omega_i)$, where $\omega_i = \exp(E_i) \exp(-F_i) \exp(E_i) \in N(T)$. (see, for example, [Hum78]). The elements ω_i also appear in the theory of Chevalley groups and Tits systems.

The \mathcal{W} -action extends naturally to \mathfrak{g} and hence to $U(\mathfrak{g})$. There, the generators S_i coincide with the T_i or Γ_i actions on $U_{\hbar}(\mathfrak{g})$ from Section 4.3 in the specialization $\hbar \rightarrow 0$. On finite-dimensional highest weight representations V of the corresponding Lie algebra \mathfrak{g} the w_i , as elements of $\text{End}(V)$, are well-defined since E_i and F_i act nilpotently on V .

Quantum Weyl elements attempt to define analogs of w_i for certain versions of quantum groups, which implement the T_i or Γ_i actions in the same manner by conjugation. Constructions of such elements are described in [Soi90, LS90, KR90] and [CP94, 8.2]. In essence, these approaches map a quantum group U into a completion of $\bigoplus_{\lambda} \text{End}(V_{\lambda})$, where λ ranges over dominant highest weights so that each V_{λ} naturally carries a \mathcal{W} -action. The w_i elements are then used to derive various formulae for R -matrices and \mathcal{A} -actions.

For relevant versions of quantum groups at roots of unity, however, quantum Weyl elements implementing the T_i -actions *almost never* exist for U_{ζ} and related quantum algebras, even if formulated in a very weak sense. The criteria discussed in this section illustrate that such implementations exist only in very exceptional types of representations.

Consider a representation $\rho : U_{\zeta, \mathbb{Q}} \rightarrow \text{End}(V)$ on a vector space V over \mathbb{C} give an embedding $\mathbb{Q}(\zeta) \subset \mathbb{C}$. Let \mathfrak{z} be the center of $U_{\zeta, \mathbb{Q}}$ and $\mathcal{Z} = \mathcal{Z}_{\bullet} \otimes \mathbb{C}$ the full subalgebra described in Section 6.3. With primes indicating commutants, we have inclusions $\text{im}(\rho) \subseteq \text{im}(\rho)'' \subseteq (\rho(\mathfrak{z}))' \subseteq (\rho(\mathfrak{z} \cap \mathcal{Z}))' =: C_{\rho}$ of subalgebras of $\text{End}(V)$. We say that a representation (ρ, V) is *weakly \mathcal{A} -equivariant* if there exist invertible operators $\mathsf{M}_i \in C_{\rho}$ such that $\rho(\Gamma_i(x)) = \mathsf{M}_i \rho(x) \mathsf{M}_i^{-1}$ for all $i = 1, \dots, n$ and $x \in U_{\zeta, \mathbb{Q}}$. The existence of quantum Weyl elements $w_i \in U_{\zeta, \mathbb{Q}}$ would immediately imply this property for *all* representations by setting $\mathsf{M}_i = \rho(w_i) \in \text{im}(\rho)$ and is, thus, a far stronger requirement.

For the assertions and counterexamples that follow, assume \mathbb{k} is odd and exclude Lie types B and G. Fix a maximal word $z \in \mathcal{W}_{\mathfrak{m}}^*$ and denote respective generators by E_{α} in the sense of (5.3). Similar results hold for other Lie types and even \mathbb{k} but are technically more involved.

For a given representation, (ρ, V) use the shorthand $\bar{A} = \rho(A)$ for $A \in U_{\zeta, \mathbb{Q}}$ as well as $F_{\alpha} = E_{-\alpha}$ and $Y_{\alpha} = X_{-\alpha}$ for $\alpha \in \Phi^+$. Let also $\Lambda_{\mathbb{k}} = \mathbb{Z}^{\Delta}/\mathbb{k}\mathbb{Z}^{\Delta} \cong (\mathbb{F}_{\mathbb{k}})^{\Delta}$ be the weight space mod \mathbb{k} , generated by the fundamental weights ϖ_i defined via $(\varpi_i | \check{\alpha}_j) = \delta_{ij}$. Assuming (ρ, V) is weakly \mathcal{A} -equivariant, denote by \mathbb{M} the subgroup of $\text{GL}(V) \cap C_{\rho}$ generated by M_i . Moreover, for a reduced word $w = w_{i_1} \dots w_{i_k}$, write $\mathsf{M}_w = \mathsf{M}_{i_1} \cdot \dots \cdot \mathsf{M}_{i_k}$. Recall also the augmentation ideals $\mathcal{K}_{\bullet}^{\square}$ and $\hat{\mathcal{K}}_{\bullet\bullet}$ as defined in (6.67) and (6.72) of Section 6.5.

Proposition C.1. *Suppose (ρ, V) is weakly \mathcal{A} -equivariant with $\{\mathsf{M}_i\}$ and \mathbb{M} as above, \mathbb{k} odd, and the Lie type is neither B nor G. Then*

- i) *We have decompositions $V = \bigoplus_{\lambda \in \Lambda_{\mathbb{k}}} V_{\lambda}$, where $K_i v = \zeta^{(\lambda | \alpha_i)} v$ for $v \in V_{\lambda}$. Moreover, $\bar{E}_{\alpha}(V_{\lambda}) \subseteq V_{\lambda+\alpha}$ and $\bar{F}_{\alpha}(V_{\lambda}) \subseteq V_{\lambda-\alpha}$.*
- ii) *For any $w \in \mathcal{W}^*$ we have $\mathsf{M}_w(V_{\lambda}) = V_{s(\lambda)}$ where $s = \gamma(w)$.*

- iii) For any pair $\alpha, \beta \in \Phi^+$ with $d_\alpha = d_\beta$ there exists an $\mathbf{M} \in \mathbb{M}$ such that $\mathbf{M}\bar{E}_\alpha\mathbf{M}^{-1} = \bar{E}_\beta$. Similarly, \bar{F}_α and \bar{F}_β are conjugate by some element in \mathbb{M} , as are \bar{F}_α and $-\bar{E}_\beta\bar{K}^\beta$.
- iv) There are operators $\bar{Z}_1, \bar{Z}_e \in \mathbb{M}' \cap (\text{im}(\rho))' \subset \text{End}(V)$ such that $\bar{X}_\alpha = -\bar{Y}_\alpha = \bar{Z}_1$ for all short roots $\alpha \in \Phi^+$ (that is, $d_\alpha = 1$) and $\bar{X}_\alpha = -\bar{Y}_\alpha = \bar{Z}_e$ if $\alpha \in \Phi^+$ is a long root ($d_\alpha = e$).
- v) Suppose V is cyclic with a cyclic vector $\mathbf{v} \in V$ such that $U_{\zeta, \mathbb{Q}}^+ \mathbf{v} = 0$. Then $\hat{\mathcal{K}}_{\bullet\bullet} \subseteq \ker(\rho)$ so that (ρ, V) factors into a representation of a respective small quantum group.

Proof. Note first that the definition of weak \mathcal{A} -equivariance implies that for any $a \in \mathcal{Z} \cap \mathfrak{z}$ we have $\rho(\Gamma_s(a)) = \rho(a)$. Now, since \mathbb{k} is odd we have $L_i = K_i^\mathbb{k} \in \mathcal{Z} \cap \mathfrak{z}$, which implies with $\Gamma_j(L_i) = L_i L_j^{-A_{ji}}$ that $\bar{L}_j^{-A_{ji}} = 1 \in \text{End}(V)$ for all $i, j = \{1, \dots, n\}$. For Lie types different from \mathbf{B} we have $\text{gcd}(A_{j1}, \dots, A_{jn}) = 1$ for any j so that, indeed, $\bar{L}_j = \bar{K}_j^\mathbb{k} = 1$ for all j . Over a field containing ζ , this implies a decomposition into common eigenspaces of the \bar{K}_j with eigenvalues of \bar{K}_j given by some powers ζ^{λ_i} with $\lambda_i \in \mathbb{F}_\mathbb{k}$. Now, for $\lambda = \sum_i \lambda_i \varpi_i$ we have $\lambda_i = (\lambda | \check{\alpha}_i) \equiv (\lambda | \alpha_i)$ mod \mathbb{k} since \mathbb{k} is coprime to all d_i if the Lie type \mathbf{G} is also excluded, proving the decomposition in Item *i*). The other claims in Item *i*) as well as Item *ii*) are immediate from commutation relations with the K_j and the actions of the Γ_i on the K_j .

For Item *iii*) write $E_\alpha = \Gamma_s(E_i)$ and $E_\beta = \Gamma_t(E_j)$ for some $s, t \in \mathbb{W}$ and $i, j \in \{1, \dots, n\}$ with $d_i = d_j$. So, $\bar{E}_\alpha = \mathbf{M}_w \bar{E}_i \mathbf{M}_w^{-1}$ and $\bar{E}_\beta = \mathbf{M}_u \bar{E}_j \mathbf{M}_u^{-1}$ with $s = \nu(w)$ and $t = \nu(u)$. Since α_i and α_j have the same length, there is some $r \in \mathbb{W}$ such that $r(\alpha_i) = \alpha_j$. From Proposition 5.1 we then find that $\Gamma_r(E_i) = E_j$ and, hence, $\mathbf{M}_z \bar{E}_i \mathbf{M}_z^{-1} = \bar{E}_j$ where $r = \nu(z)$. The claim now follows for $\mathbf{M} = \mathbf{M}_u \mathbf{M}_z \mathbf{M}_w^{-1} \in \mathbb{M}$. The remaining assertions follow from analogous arguments.

Since all $X_\alpha, Y_\alpha \in \mathcal{Z} \cap \mathfrak{z}$ by Proposition 6.12, we infer that $\bar{X}_\alpha, \bar{Y}_\alpha \in \mathbb{M}' \cap (\text{im}(\rho))'$ for all $\alpha \in \Phi^+$. Item *iii*) now implies that $\bar{X}_\alpha = \bar{X}_\beta$ and $\bar{Y}_\alpha = \bar{Y}_\beta$ whenever $d_\alpha = d_\beta$. Next, $\bar{L}_i = 1$ and (6.20) also imply $\bar{X}_i = -\bar{Y}_i$ for all i so that also $\bar{X}_\alpha = -\bar{Y}_\beta$ whenever $d_\alpha = d_\beta$, proving the claim in Item *iv*).

For a cyclic vector $\mathbf{v} \in V$ as in Item *v*), note that, by Item *iv*), any $\bar{Z} \in \{\bar{Z}_1, \bar{Z}_e\}$ is given as $\bar{Z} = \bar{X}_i$ for some i so that $\bar{Z}\mathbf{v} = \bar{X}_i\mathbf{v} = 0$. Given that any $\bar{A} \in \text{im}(\rho)$ commutes with \bar{Z} , we also find $\bar{Z}\bar{A}\mathbf{v} = \bar{A}\bar{Z}\mathbf{v} = 0$ and hence $\bar{Z} = 0$ by the cyclicity assumption. Item *iv*) now implies that all $\bar{X}_\alpha = \bar{Y}_\alpha = 0$. Thus, $X_\alpha = E_\alpha^{\ell_\alpha}$, $Y_\alpha = F_\alpha^{\ell_\alpha}$, and $K_i^{\ell_i} - 1 = L_i - 1$ are all in the kernel of ρ , which then factors into the small quantum group. \square

In the remainder of this section, we provide several examples of representations that are not weakly \mathcal{A} -equivariant using the criteria from the above proposition.

Denote by V_κ the weight space on which each K_i acts as $\kappa_i \mathbb{1}$ for an n -tuple $\kappa = (\kappa_1, \dots, \kappa_n) \in \mathbb{t} = (\mathbb{C}^\times)^n$. For any $\kappa \in \mathbb{t}$, a representation with $V_\kappa \neq 0$ can be easily constructed, for example, as a highest weight representation with highest weight κ . The latter can also be chosen to be finite-dimensional and irreducible by dividing out maximal submodules as usual. The condition in Item *i*) thus restricts the set of possible weights from the infinite group \mathbb{t} to the finite subgroup

$\mathbb{S} = \{\kappa \in \mathbb{t} : \kappa_i^k = 1 \forall i\}$ of tuples of roots of unity. In this sense, even the weak \mathcal{A} -equivariance condition forces a restriction to a null set.

Requiring $\bar{L}_i = 1$ still leaves large families of finite-dimensional representations that are not weakly \mathcal{A} -equivariant. Assuming that k is odd, these are readily constructed by choosing *any* ideal $J \subsetneq \mathcal{K}_\bullet^- \subsetneq \mathcal{Z}^- \cong \mathbb{C}[\{Y_\alpha : \alpha \in \Phi^+\}]$ such that \mathcal{Z}^-/J is of finite rank $m > 1$. Since \mathcal{Z} is central, any weight $\kappa \in \mathbb{S}$ defines a one-dimensional representation of $B_J = J \cdot U_{\zeta, \mathbb{k}}^{\geq 0}$ with $K_i \mathbf{v} = \kappa_i \mathbf{v}$ and $E_i \mathbf{v} = b \mathbf{v} = 0$ for all $i \in \{1, \dots, n\}$ and all $b \in J$. The induced module $U_{\zeta, \mathbb{k}} \otimes_{B_J} \mathbf{v}$ is now finite-dimensional with $\bar{L}_i = 1$, but violates Item [v](#)) in the proposition above and is, thus, not \mathcal{A} -equivariant. There are many choices for J for which the \bar{Y}_α are not zero and may in fact even be invertible.

C.1. An Example in Rank 1. Even among representations that factor through the small quantum groups, most are still not weakly \mathcal{A} -equivariant. This can already be seen in the rank 1 case $\mathfrak{g} = \mathfrak{sl}_2$ and $k = 3$ for which $E^3 = F^3 = 0$ and $K^3 = 1$.

Denote by V_0 , V_- , and V_+ the eigenspaces of K for eigenvalues 1, $\zeta^2 = \zeta^{-1}$, and $\zeta^{-2} = \zeta^1$, respectively. We consider a special family of representations in which V_+ and V_- are isomorphic as vector spaces to a common space V_* so that, after conjugation by respective isomorphisms, the restrictions $E : V_- \rightarrow V_+$ and $F : V_+ \rightarrow V_-$ may be viewed as linear endomorphisms of V_* . Further specializing our example, we assume that these two endomorphisms are the identity on V_* . The restrictions of E and F to other eigenspaces are then denoted as $E_-, F_+ : V_0 \rightarrow V_*$ and $E_+, F_- : V_* \rightarrow V_0$. They form a representation of the small quantum group iff

$$\begin{aligned} E_+ E_- &= F_- F_+ = 0 & \text{and} & \quad E_+ F_+ = F_- E_- & \quad \text{on } V_0 \\ \text{and} \quad E_- E_+ &= F_+ F_- = E_- F_- = F_+ E_+ = 0 & & & \quad \text{on } V_*. \end{aligned} \tag{C.1}$$

This type of reduction of representations of $U_\zeta(\mathfrak{sl}_2)$ to a quiver with relations as in (C.1) extends to all roots of unity as shown, for example, in [\[Ker89\]](#). The representation given by the E_\pm and F_\pm as above is \mathcal{A} -equivariant if and only if there are isomorphisms $W_0 \in \text{GL}(V_0)$ and $W_* \in \text{GL}(V_*)$ such that

$$W_0 E_+ W_*^{-1} = F_-, \quad W_* E_- W_0^{-1} = -F_+, \quad W_* F_+ W_0^{-1} = E_-, \quad \text{and} \quad W_0 F_- W_*^{-1} = -E_+ \tag{C.2}$$

implying, in particular, $\text{rank}(E_+) = \text{rank}(F_-)$ and $\text{rank}(E_-) = \text{rank}(F_+)$.

It is not hard to construct maps that fulfill (C.1) but violate (C.2). For example, setting $E_+ = F_- = 0$ automatically solves all equations in (C.1) so that any choice of maps $E_-, F_+ : V_0 \rightarrow V_*$ produces a representation. The residual representation theory is, thus, equivalent to that of the (2) Kronecker quiver, which is tame and for which the indecomposable modules are classified [\[Ben91, Theorem 4.3.2\]](#).

Kronecker's classification includes a generic family in which $\dim(V_0) = \dim(V_*)$ and one of the maps can be chosen as the identity matrix and the other an indecomposable rational form. For example, if $\mathbb{k} = \mathbb{C}$ the latter implies an indecomposable Jordan block $J = \kappa \mathbb{1} + N$, where N

is maximally nilpotent. It is not hard to check that this representation is weakly \mathcal{A} -equivariant if and only if $\kappa^2 = -1$, again singling out a finite set from an infinite parameter space.

For the second family in the classification, the dimensions of V_0 and V_* differ by one and there is exactly one representation for each such choice of dimensions. Bases $\{v_i^0\}$ of V_0 and $\{v_i^*\}$ of V_* may be chosen such that $E_- v_i^0 = v_i^*$ and $F_+ v_i^0 = v_{i+1}^*$ or $F_+ v_i^0 = v_{i-1}^*$. These special cases are indeed weakly \mathcal{A} -equivariant. The W_0 and W_* may be chosen as anti-diagonal matrices with alternating +1 and -1 entries.

Note that if we require representations to lift to a larger (unrolled) quantum group that contains the element H as in (3.10), the generic family is excluded unless $\kappa = 0$ (which does not yield a weakly \mathcal{A} -equivariant module due to the rank condition). In this situation, however, the notion of weak \mathcal{A} -equivariance on the larger algebra implies also $-\bar{H} = M\bar{H}M^{-1}$ since $\Gamma(H) = -H$, implying symmetry of H -weights around 0. The latter is easily broken by shifts $\bar{H} \mapsto \bar{H} + m\hbar\mathbb{1}$, producing infinite families of lifts of a representation that are not weakly \mathcal{A} -equivariant.

For irreducible representations arising from sufficiently small dominant highest weights (in a respective principal affine Weyl alcove) one may expect weak \mathcal{A} -equivariance to apply based on the original constructions of quantum Weyl elements. In the $U_\zeta(\mathfrak{sl}_2)$ situation above, this corresponds to the trivial solution of (C.1) and (C.2) in which all maps are zero.

Finally, we note that the standard projective representation in the special $U_\zeta(\mathfrak{sl}_2)$ case is in fact weakly \mathcal{A} -equivariant. In terms of the maps above it is given in a suitable basis by matrices

$$E_+ = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \quad E_- = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \quad F_+ = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \quad \text{and} \quad F_- = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}.$$

The conditions (C.2) are then solved, for example, by $W_0 = \mathbb{1}$ and $W_* = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$.

APPENDIX D. INTEGRAL FORM OF COALGEBRAS

We introduce an additional normalization \hat{X}_α of the De Concini-Procesi singularized power generators \check{X}_α , which entails integral coproduct expressions in our formulae for types A_n and B_2 . Notice from (8.7) that the singularized generator $\check{X}_{(iji)}$ has a coproduct over the Gauss integers $\mathbb{Z}[\sqrt{-1}]$ when ℓ is even. Thus, the integral coordinate rings arise as Hopf subalgebras of $U_\zeta(\mathfrak{g})$ in these types. These integral forms also support simplified computations in future work.

In type A_n , we gave in Lemma 7.9 a formula for the coproduct of generators $X_{i,j}$

$$\Delta(X_{i,j}) = \sum_{k=i}^j b_k X_{i,k} \otimes L_{i,k} X_{k,j}, \tag{D.1}$$

where $b_k = (\zeta^{-1} - \zeta)^\ell \zeta^{\binom{\ell}{2}}$ for $i < k < j$ and $b_i = b_j = 1$. Replacing all instances of X_α with the singular generator \check{X}_α removes the coefficient $(\zeta^{-1} - \zeta)^\ell$ from b_k . If we set $\hat{X}_\alpha = \zeta^{\binom{\ell}{2}} \check{X}_\alpha = (\zeta^{-1} - \zeta)^\ell \zeta^{\binom{\ell}{2}} X_\alpha$, then we obtain a coproduct formula in which all nonzero coefficients are 1.

$$\Delta(\hat{X}_{i,j}) = \sum_{k=i}^j \hat{X}_{i,k} \otimes L_{i,k} \hat{X}_{k,j} \quad (\text{D.2})$$

More generally, set $\hat{E}_i^k = \zeta_i^{\binom{k}{2}} \check{E}_i^k$, then write $\hat{X}_i = \hat{E}_i^{\ell_i}$. The coproduct formulae of Proposition 8.1 have simpler presentations using the rescaled singularized generators

$$\begin{aligned} \Delta(\hat{X}_{(iji)}) &= 1 \otimes \hat{X}_{(iji)} + \hat{X}_i \otimes L_i \hat{X}_j^\mathbf{d} + \hat{X}_{(iji)} \otimes L_i L_j^\mathbf{d}, \\ &+ \delta_{\mathbf{d},2} \cdot 2 \cdot (-1)^{\ell_j} \hat{X}_{(ij)} \otimes L_i L_j \hat{X}_j \end{aligned} \quad (\text{D.3})$$

$$\begin{aligned} \Delta(\hat{X}_{(ij)}) &= 1 \otimes \hat{X}_{(ij)} + \hat{X}_i^\mathbf{n} \otimes L_i^\mathbf{n} \hat{X}_j + \hat{X}_{(ij)} \otimes L_i^\mathbf{n} L_j \\ &+ \delta_{\mathbf{n},2} \cdot 2 \cdot \zeta^{\binom{\ell+1}{2}} \hat{X}_i \otimes L_i \hat{X}_{(iji)}. \end{aligned} \quad (\text{D.4})$$

These follow from the observations $\hat{X}_j^\mathbf{d} = \zeta_j^{\binom{\ell}{2}} \check{X}_j^\mathbf{d} = (-1)^{\ell+1} \check{X}_j^\mathbf{d}$ and $\hat{X}_i^\mathbf{n} = \zeta^{\binom{2\ell_j}{2}} \check{X}_i^\mathbf{n}$. Note also that $\delta_{\mathbf{d},2} \zeta^{\binom{\ell}{2}} = \delta_{\mathbf{d},2} (-\zeta)^{-\ell_j}$, $\delta_{\mathbf{n},2} \zeta^{\binom{\ell}{2}} = 1$, $\delta_{\mathbf{n},2} \zeta^{\binom{\ell}{2}+\ell_j} = \delta_{\mathbf{n},2} \zeta^{\binom{\ell+1}{2}} = \delta_{\mathbf{n},2} \zeta^{\binom{\ell+1}{2}}$ are all valued in $\{-1, 0, 1\}$.

The automorphism \mathfrak{O} is given for certain power generators in (6.52) and (6.54). Here, we regard it as a factor of the antipode, which takes values over the same ring as the coproduct. We deduce integrality for the \hat{X}_α from the respective equations.

$$\begin{aligned} (-1)^\ell \hat{X}_{(ji)} &= \zeta_j^\ell \hat{X}_{a_{m-1}} + (-1)^\ell \hat{X}_j^\mathbf{d} \hat{X}_i + (1 - \delta_{\mathbf{d},1}) \cdot \mathbf{e} \cdot (-1)^\ell \cdot T_1 \\ \text{where} \quad T_1 &= \begin{cases} (-1)^{\ell_j+1} \hat{X}_j \hat{X}_{(ij)} & \mathbf{e} = 2 \\ \zeta_j^{2\binom{\ell_j}{2}} \hat{X}_j^2 \hat{X}_{(ij)} + \hat{X}_j \hat{X}_{(iji)} & \mathbf{e} = 3, \end{cases} \end{aligned} \quad (\text{D.5})$$

$$\begin{aligned} (-1)^{\mathbf{e}\ell_j} \hat{X}_{b_{m-1}} &= \zeta_j^{\ell_j} \hat{X}_{(ij)} + (-1)^{\mathbf{n}\cdot\ell} \hat{X}_j \hat{X}_i^\mathbf{n} + \delta_{\mathbf{d},1} \cdot (1 - \delta_{\mathbf{e},1}) \cdot \mathbf{e} \cdot (-1)^\ell \cdot T_2 \\ \text{where} \quad T_2 &= \begin{cases} \zeta^\ell \hat{X}_{(iji)} \hat{X}_i & \mathbf{e} = 2 \\ \hat{X}_{(iji)} \hat{X}_i + (-1)^{\ell+1} \hat{X}_{(iji)} \hat{X}_i^2 & \mathbf{e} = 3. \end{cases} \end{aligned} \quad (\text{D.6})$$

The integrality of these expressions suggests that the normalized generators above produce integral coproducts for all Lie types and roots of unity.

Conjecture D.1. *Let $z \in \mathcal{W}_m^*$. The set of \hat{X}_w for $w \leq_R z$ generate a \mathbb{Z} -Hopf subalgebra of $U_\zeta(\mathfrak{g})$.*

REFERENCES

[ABI15] Usman Ali, Barbu Berceanu, and Zaffar Iqbal. Relative Garside elements of Artin monoids. *Rev. Roumaine Math. Pures Appl.*, 60(3):267–279, 2015.

[ADO92] Yasuhiro Akutsu, Tetsuo Deguchi, and Tomotada Ohtsuki. Invariants of colored links. *J. Knot Theory Ramifications*, 1(2):161–184, 1992.

[AJS94] Henning Haahr Andersen, Jens Carsten Jantzen, and Wolfgang Soergel. Representations of quantum groups at a p th root of unity and of semisimple groups in characteristic p : independence of p . *Astérisque*, 220:321, 1994.

[AM15] Henning Haahr Andersen and Volodymyr Mazorchuk. Category \mathcal{O} for quantum groups. *J. Eur. Math. Soc. (JEMS)*, 17(2):405–431, 2015.

[Ang22] Cristina Ana-Maria Anghel. A topological model for the coloured Jones polynomials. *Selecta Math. (N.S.)*, 28(3):Paper No. 63, 50, 2022.

[Ang24] Cristina Ana-Maria Anghel. ADO invariants directly from partial traces of homological representations. *New York J. Math.*, 30:481–501, 2024.

[AST91] Michael Artin, William Schelter, and John Tate. Quantum deformations of GL_n . *Comm. Pure Appl. Math.*, 44(8-9):879–895, 1991.

[BB05] Anders Björner and Francesco Brenti. *Combinatorics of Coxeter groups*, volume 231 of *Graduate Texts in Mathematics*. Springer, New York, 2005.

[BCGPM16] Christian Blanchet, Francesco Costantino, Nathan Geer, and Bertrand Patureau-Mirand. Non-semi-simple TQFTs, Reidemeister torsion and Kashaev’s invariants. *Adv. Math.*, 301:1–78, 2016.

[BDR02] Margaret Beattie, Sorin Dăscălescu, and Șerban Raianu. Lifting of Nichols algebras of type B_2 . *Israel J. Math.*, 132:1–28, 2002. With an appendix by the authors and Ian Rutherford.

[Ben91] David John Benson. *Representations and cohomology. I*, volume 30 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1991. Basic representation theory of finite groups and associative algebras.

[Bou02] Nicolas Bourbaki. *Lie groups and Lie algebras. Chapters 4–6*. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 2002. Translated from the 1968 French original by Andrew Pressley.

[BS72] Egbert Brieskorn and Kyoji Saito. Artin-Gruppen und Coxeter-Gruppen. *Invent. Math.*, 17:245–271, 1972.

[Bum04] Daniel Bump. *Lie groups*, volume 225 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2004.

[BW04] Georgia Benkart and Sarah Witherspoon. Restricted two-parameter quantum groups. In *Representations of finite dimensional algebras and related topics in Lie theory and geometry*, volume 40 of *Fields Inst. Commun.*, pages 293–318. Amer. Math. Soc., Providence, RI, 2004.

[CGPM14] Francesco Costantino, Nathan Geer, and Bertrand Patureau-Mirand. Quantum invariants of 3-manifolds via link surgery presentations and non-semi-simple categories. *J. Topol.*, 7(4):1005–1053, 2014.

[Cli94] Gerald Cliff. A tensor product theorem for quantum linear groups at even roots of unity. *J. Algebra*, 165(3):566–575, 1994.

[Con17] Brian Conrad. Algebraic groups II. AMS Open Math Notes, 2017. [OMN:201701.110663](#).

[CP94] Vyjayanthi Chari and Andrew Pressley. *A guide to quantum groups*. Cambridge University Press, Cambridge, 1994.

[DCK90] Corrado De Concini and Victor G. Kac. Representations of quantum groups at roots of 1. In *Operator algebras, unitary representations, enveloping algebras, and invariant theory (Paris, 1989)*, volume 92 of *Progr. Math.*, pages 471–506. Birkhäuser Boston, Boston, MA, 1990.

[DCKP92] Corrado De Concini, Victor G. Kac, and Claudio Procesi. Quantum coadjoint action. *J. Amer. Math. Soc.*, 5(1):151–189, 1992.

[DCP93] Corrado De Concini and Claudio Procesi. Quantum groups. In *D-modules, representation theory, and quantum groups (Venice, 1992)*, volume 1565 of *Lecture Notes in Math.*, pages 31–140. Springer, Berlin, 1993.

[DDG⁺15] Patrick Dehornoy, Francois Digne, Eddy Godelle, Daan Krammer, and Jean Michel. *Foundations of Garside theory*, volume 22 of *EMS Tracts in Mathematics*. European Mathematical Society (EMS), Zürich, 2015.

- [DR22] Marco De Renzi. Non-semisimple extended topological quantum field theories. *Mem. Amer. Math. Soc.*, 277(1364):v+161, 2022.
- [Dri87] Vladimir G. Drinfeld. Quantum groups. In *Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Berkeley, Calif., 1986)*, pages 798–820. Amer. Math. Soc., Providence, RI, 1987.
- [GHPM22] Nathan Geer, Ngoc Phu Ha, and Bertrand Patureau-Mirand. Modified graded Hennings invariants from unrolled quantum groups and modified integral. *J. Pure Appl. Algebra*, 226(3):Paper No. 106815, 46, 2022.
- [GHPM23] Nathan Geer, Ngoc Phu Ha, and Bertrand Patureau-Mirand. Modified symmetrized integral in G -coalgebras. *J. Knot Theory Ramifications*, 32(14):Paper No. 2350082, 18, 2023.
- [GW09] Roe Goodman and Nolan R. Wallach. *Symmetry, representations, and invariants*, volume 255 of *Graduate Texts in Mathematics*. Springer, Dordrecht, 2009.
- [Har] Matthew Harper. A non-abelian generalization of the Alexander Polynomial from Quantum \mathfrak{sl}_3 . Preprint 2020, [arXiv:2008.06983](https://arxiv.org/abs/2008.06983).
- [Hen96] Mark Hennings. Invariants of links and 3-manifolds obtained from Hopf algebras. *J. London Math. Soc.* (2), 54(3):594–624, 1996.
- [Hil82] Howard Hiller. *Geometry of Coxeter groups*, volume 54 of *Research Notes in Mathematics*. Pitman (Advanced Publishing Program), Boston, Mass.-London, 1982.
- [Hum78] James E. Humphreys. *Introduction to Lie algebras and representation theory*, volume 9 of *Graduate Texts in Mathematics*. Springer-Verlag, New York-Berlin, 1978. Second printing, revised.
- [Ito15] Tetsuya Ito. Reading the dual Garside length of braids from homological and quantum representations. *Comm. Math. Phys.*, 335(1):345–367, 2015.
- [Ito16] Tetsuya Ito. A homological representation formula of colored Alexander invariants. *Adv. Math.*, 289:142–160, 2016.
- [Jan96] Jens Carsten Jantzen. *Lectures on quantum groups*, volume 6 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 1996.
- [Jim85] Michio Jimbo. A q -difference analogue of $U(\mathfrak{g})$ and the Yang-Baxter equation. *Lett. Math. Phys.*, 10(1):63–69, 1985.
- [JK11] Craig Jackson and Thomas Kerler. The Lawrence-Krammer-Bigelow representations of the braid groups via $U_q(\mathfrak{sl}_2)$. *Adv. Math.*, 228(3):1689–1717, 2011.
- [Ker89] Thomas Kerler. *Darstellungen der Quantengruppen und Anwendungen*. Diploma thesis, ETH Zürich, 1989.
- [Ker03] Thomas Kerler. Towards an algebraic characterization of 3-dimensional cobordisms. In *Diagrammatic morphisms and applications (San Francisco, CA, 2000)*, volume 318 of *Contemp. Math.*, pages 141–173. Amer. Math. Soc., Providence, RI, 2003.
- [Kha15] Vladislav K. Kharchenko. Explicit coproduct formula for quantum groups of infinite series. *Israel J. Math.*, 208(1):13–43, 2015.
- [KL01] Thomas Kerler and Volodymyr V. Lyubashenko. *Non-semisimple topological quantum field theories for 3-manifolds with corners*, volume 1765 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2001.
- [KR90] Anatoliĭ N. Kirillov and Nikolai Yu. Reshetikhin. q -Weyl group and a multiplicative formula for universal R -matrices. *Comm. Math. Phys.*, 134(2):421–431, 1990.
- [LS90] Sergei Z. Levendorskiĭ and Yan S. Soibelman. Some applications of the quantum Weyl groups. *J. Geom. Phys.*, 7(2):241–254, 1990.
- [LS91] Serge Levendorskiĭ and Yan Soibelman. Algebras of functions on compact quantum groups, Schubert cells and quantum tori. *Comm. Math. Phys.*, 139(1):141–170, 1991.
- [Lus90a] George Lusztig. Finite-dimensional Hopf algebras arising from quantized universal enveloping algebra. *J. Amer. Math. Soc.*, 3(1):257–296, 1990.
- [Lus90b] George Lusztig. Quantum groups at roots of 1. *Geom. Dedicata*, 35(1-3):89–113, 1990.
- [Lus93] George Lusztig. *Introduction to quantum groups*. Birkhäuser Boston Inc., Boston, MA, 1993.

[Mar22] Jules Martel. A homological model for $U_q\mathfrak{sl}_2$ Verma modules and their braid representations. *Geom. Topol.*, 26(3):1225–1289, 2022.

[Mat64] Hideya Matsumoto. Générateurs et relations des groupes de Weyl généralisés. *C. R. Acad. Sci. Paris*, 258:3419–3422, 1964.

[Mil17] James Stuart Milne. *Algebraic groups*, volume 170 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2017. The theory of group schemes of finite type over a field.

[Mon93a] Susan Montgomery. *Hopf algebras and their actions on rings*, volume 82 of *CBMS Regional Conference Series in Mathematics*. American Mathematical Society, Providence, RI, 1993.

[Mon93b] Susan Montgomery. Some remarks on filtrations of Hopf algebras. *Comm. Algebra*, 21(3):999–1007, 1993.

[Mur92] Jun Murakami. The multi-variable Alexander polynomial and a one-parameter family of representations of $U_q(\mathfrak{sl}(2, \mathbb{C}))$ at $q^2 = -1$. In *Quantum groups (Leningrad, 1990)*, volume 1510 of *Lecture Notes in Math.*, pages 350–353. Springer, Berlin, 1992.

[Mur93] Jun Murakami. A state model for the multivariable Alexander polynomial. *Pacific J. Math.*, 157(1):109–135, 1993.

[MW98] Susan Montgomery and Sarah J. Witherspoon. Irreducible representations of crossed products. *J. Pure Appl. Algebra*, 129(3):315–326, 1998.

[Nic78] Warren D. Nichols. Quotients of Hopf algebras. *Comm. Algebra*, 6(17):1789–1800, 1978.

[Oht95] Tomotada Ohtsuki. A polynomial invariant of integral homology 3-spheres. *Math. Proc. Cambridge Philos. Soc.*, 117(1):83–112, 1995.

[Oht96] Tomotada Ohtsuki. Finite type invariants of integral homology 3-spheres. *J. Knot Theory Ramifications*, 5(1):101–115, 1996.

[Oht02] Tomotada Ohtsuki. *Quantum invariants*, volume 29 of *Series on Knots and Everything*. World Scientific Publishing Co., Inc., River Edge, NJ, 2002. A study of knots, 3-manifolds, and their sets.

[Pap94] Paolo Papi. A characterization of a special ordering in a root system. *Proc. Amer. Math. Soc.*, 120(3):661–665, 1994.

[Rad12] David E. Radford. *Hopf algebras*, volume 49 of *Series on Knots and Everything*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2012.

[Rin96] Claus Michael Ringel. PBW-bases of quantum groups. *J. Reine Angew. Math.*, 470:51–88, 1996.

[Ros88] Marc Rosso. Finite-dimensional representations of the quantum analog of the enveloping algebra of a complex simple Lie algebra. *Comm. Math. Phys.*, 117(4):581–593, 1988.

[Ros89] Marc Rosso. An analogue of P.B.W. theorem and the universal R -matrix for $U_h\mathrm{sl}(N+1)$. *Comm. Math. Phys.*, 124(2):307–318, 1989.

[RT91] Nikolai Yu. Reshetikhin and Vladimir G. Turaev. Invariants of 3-manifolds via link polynomials and quantum groups. *Invent. Math.*, 103(3):547–597, 1991.

[Soi90] Yan S. Soibelman. Gel'fand-Naimark-Segal states and Weyl groups for the quantum group $\mathrm{SU}(n)$. *Funktional. Anal. i Prilozhen.*, 24(3):92–93, 1990.

[Spr09] Tonny A. Springer. *Linear algebraic groups*. Modern Birkhäuser Classics. Birkhäuser Boston, Inc., Boston, MA, second edition, 2009.

[Sta84] Richard P. Stanley. On the number of reduced decompositions of elements of Coxeter groups. *European J. Combin.*, 5(4):359–372, 1984.

[Swe69] Moss E. Sweedler. *Hopf algebras*. Mathematics Lecture Note Series. W. A. Benjamin, Inc., New York, 1969.

[Tak90] Mitsuhiro Takeuchi. The q -bracket product and quantum enveloping algebras of classical types. *J. Math. Soc. Japan*, 42(4):605–629, 1990.

[Tan92] Toshiyuki Tanisaki. Killing forms, Harish-Chandra isomorphisms, and universal R -matrices for quantum algebras. In *Infinite analysis, Part A, B (Kyoto, 1991)*, volume 16 of *Adv. Ser. Math. Phys.*, pages 941–961. World Sci. Publ., River Edge, NJ, 1992.

- [Tan16] Toshiyuki Tanisaki. The center of a quantized enveloping algebra at an even root of unity. *Osaka J. Math.*, 53(1):47–81, 2016.
- [Tur16] Vladimir G. Turaev. *Quantum invariants of knots and 3-manifolds*, volume 18 of *De Gruyter Studies in Mathematics*. De Gruyter, Berlin, third edition, 2016.
- [Was82] Lawrence C. Washington. *Introduction to cyclotomic fields*, volume 83 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1982.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, RIVERSIDE
DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY

Email address: mrhmath@proton.me

DEPARTMENT OF MATHEMATICS, THE OHIO STATE UNIVERSITY

Email address: kerler.2@osu.edu