

**ASYMPTOTICS OF THE SHIFTED FINITE DIFFERENCES OF THE  
OVERPARTITION FUNCTION AND A PROBLEM OF  
WANG–XIE–ZHANG**

GARGI MUKHERJEE

**ABSTRACT.** Let  $\bar{p}(n)$  denote the overpartition function, and for  $j \in \mathbb{N}$ ,  $\Delta_j^r$  denote the  $r$ -fold applications of the shifted difference operator  $\Delta_j$  defined by  $\Delta_j(a)(n) := a(n) - a(n-j)$ . The main goal of this paper is to derive an asymptotic expansion of  $\Delta_j^r(\bar{p})(n)$  with an effective error bound which subsequently gives an answer to a problem of Wang, Xie, and Zhang. In order to get the asymptotics of  $\Delta_j^r(\bar{p})(n)$ , we derive an asymptotic expansion of the shifted overpartition function  $\bar{p}(n+k)$  for any integer  $k \neq 0$ .

1. INTRODUCTION

An overpartition of a positive integer  $n$ , introduced by Corteel and Lovejoy [6], is a non-increasing sequence of positive integers whose sum is  $n$  in which the first occurrence of a number may be overlined,  $\bar{p}(n)$  denotes the number of overpartitions of  $n$ , and  $\bar{p}(0) := 1$ . For instance, there are 8 overpartitions of 3 enumerated by

$$3, \bar{3}, 2+1, \bar{2}+1, 2+\bar{1}, \bar{2}+\bar{1}, 1+1+1, \bar{1}+1+1.$$

Analogous to the celebrated Hardy-Ramanujan-Rademacher formula for the partition function  $p(n)$  (see [13, 24]), Zuckerman's [27] formula for  $\bar{p}(n)$  states that

$$\bar{p}(n) = \frac{1}{2\pi} \sum_{\substack{k=1 \\ 2 \nmid k}}^{\infty} \sqrt{k} \sum_{\substack{h=0 \\ (h,k)=1}}^{k-1} \frac{\omega(h,k)^2}{\omega(2h,k)} e^{-\frac{2\pi i nh}{k}} \frac{d}{dn} \left( \frac{\sinh \frac{\pi \sqrt{n}}{k}}{\sqrt{n}} \right), \quad (1.1)$$

where

$$\omega(h,k) = \exp \left( \pi i \sum_{r=1}^{k-1} \frac{r}{k} \left( \frac{hr}{k} - \left\lfloor \frac{hr}{k} \right\rfloor - \frac{1}{2} \right) \right)$$

for  $(h,k) \in \mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 1}$ . Similar to the work done by Lehmer [16] for  $p(n)$ , Engel [8] estimated an error term for  $\bar{p}(n)$  as

$$\bar{p}(n) = \frac{1}{2\pi} \sum_{\substack{k=1 \\ 2 \nmid k}}^N \sqrt{k} \sum_{\substack{h=0 \\ (h,k)=1}}^{k-1} \frac{\omega(h,k)^2}{\omega(2h,k)} e^{-\frac{2\pi i nh}{k}} \frac{d}{dn} \left( \frac{\sinh \frac{\pi \sqrt{n}}{k}}{\sqrt{n}} \right) + R_2(n, N), \quad (1.2)$$

---

2020 *Mathematics Subject Classification.* 05A16, 05A20, 11P82.

*Key words and phrases.* Overpartitions, Finite differences, Asymptotic expansion.

with

$$|R_2(n, N)| < \frac{N^{5/2}}{\pi n^{3/2}} \sinh\left(\frac{\pi\sqrt{n}}{N}\right). \quad (1.3)$$

Let  $\Delta$  be the forward difference operator defined by  $\Delta(a)(n) := a(n) - a(n - 1)$  for a sequence  $(a(n))_{n \geq 0}$  and  $\Delta^r$  be the  $r$ -fold applications of  $\Delta$  defined as  $\Delta^r a(n) := \Delta(\Delta^{r-1} a(n))$  for  $r \in \mathbb{N}$ . In 1977, Good [10] proposed a problem on  $\Delta^r(p)(n)$  which states that for  $r \geq 3$ ,  $(\Delta^r(p)(n))_{n \geq 1}$  is a sequence of alternating terms for a certain range  $1 \leq n \leq n_0(r)$  and then become positive. Using Hardy-Ramanujan-Rademacher [13, 24] formula for  $p(n)$ , Gupta proved that  $\Delta^r(p)(n) > 0$  for all  $n > n_0(r)$  but without any explicit estimate for  $n_0(r)$  and conjectured that  $n_0(r) \sim r^3$  as  $r \rightarrow \infty$ . Odlyzko [22] settled the Good-Gupta conjecture by proving that  $(-1)^n \Delta^r(p)(n) \geq 0$  for all  $1 \leq n < n_0(r)$  and  $\Delta^r(p)(n) \geq 0$  for  $n \geq n_0(r)$  with  $n_0(r) \sim \frac{6}{\pi^2} r^2 (\log r)^2$  as  $r \rightarrow \infty$ . Remarkably, an exact equation for  $n_0(r)$  was given by Knessl and Keller [15] which subsequently refined the asymptotic estimation of  $n(r)$  by Odlyzko mentioned before. Later Almkvist [1] obtained an exact formula for  $\Delta^r(p)(n)$ . Recently, Gomez, Males, and Rolen [9] considered a generalized version of the difference operator  $\Delta$  and defined  $j$ -shifted difference by  $\Delta_j(a)(n) := a(n) - a(n - j)$  (note that, for  $j = 1$ , we retrieve the classical  $\Delta$  operator) and obtained an asymptotic expansion (up to first 4 terms) of  $\Delta^r(p)(n)$  along with an explicit estimation of the error term (cf. [9, Theorem 1.1]) and consequently proved that  $\Delta_j^2(p)(n) > 0$  for all  $n > 16j^2 + 1/24$  (cf. [9, Theorem 1.2]). A combinatorial proof of  $\Delta_j^2 p(n) > 0$  is given independently by Gomez, Males, and Rolen [9] and Banerjee [4]. Banerjee [2, Section 7.2] studied the asymptotic expansion of  $\Delta_j^r(p)(n)$  but without any estimation of the error bound. Regarding the logarithmic variation of finite differences of the partition function, denoted by  $(-1)^{r-1} \Delta^r(\log p)(n)$ , the positivity phenomena similar to the difference of partition function, first studied in detail by Chen, Wang, and Xie [5, Theorem 3.1 and Theorem 4.1] and settled a conjecture of DeSalvo and Pak [7, Conjecture 4.3]. Banerjee [3, Theorem 4.9] obtained an asymptotic expansion for  $(-1)^{r-1} \Delta^r(\log p)(n)$ .

In context of the overpartition function, Wang, Xie, and Zhang [25] studied the positivity of the higher order differences  $\Delta^r(\bar{p})(n)$  (cf. [25, Theorems 2.1 and 2.3]) and its logarithmic variation  $(-1)^{r-1} \Delta^r(\log \bar{p})(n)$  (cf. [25, Theorem 3.1 and Theorem 4.1]). The author [20] settled a problem of Wang, Xie, and Zhang [25, Problem 3.4] by proving an asymptotic lower bound of  $(-1)^{r-1} \Delta^r(\log \bar{p})(n)$ . A combinatorial proof of  $\Delta_j^2(\bar{p})(n) > 0$  is given in [4]. The key motivation of this paper arises from the following problem of Wang, Xie, and Zhang [25, Problem 2.4]. Before we state the problem, let us state the result on  $\Delta^r(\bar{p})(n)$  by Wang, Xie, and Zhang.

**Theorem 1.1.** [25, Theorem 2.1 and Theorem 2.3] *For  $r \geq 1$ , there exists a positive integer  $n(r)$  such that for all  $n \geq n(r)$ ,*

$$0 < \Delta^r(\bar{p})(n) < 2^{r-3} \left(1 - 2^{-\frac{3}{2}}\right) \zeta\left(\frac{3}{2}\right) \frac{e^{\pi\sqrt{n+r}}}{n+r}, \quad (1.4)$$

where  $\zeta$  is the Riemann zeta function.

This result led Wang, Xie, and Zhang to propose the following problem.

**Problem 1.2.** [25, Problem 2.4] *Find a sharp lower bound of  $\Delta^r(\bar{p})(n)$ .*

In other words, it is equivalent to find the asymptotic growth of  $\Delta^r(\bar{p})(n)$  as  $n \rightarrow \infty$ .

In attempting to give an affirmative answer to this question, we first derive an asymptotic expansion of  $\bar{p}(n+k)$  for any  $k \in \mathbb{Z} \setminus \{0\}$  in Theorem 1.3. Using Theorem 1.3, we obtain the asymptotic expansion of the  $r$ -fold applications of the  $j$ -shifted differences of the overpartition function in Theorem 1.4 which in turn gives a sharper lower and upper bounds for  $\Delta_j^r(\bar{p})(n-j)$ .

Before we state the main results, for  $(N, j, r) \in \mathbb{N}^3$  and  $k \in \mathbb{Z} \setminus \{0\}$ , let us define the following constants depending on parameters  $N, j, r, k$  which will be used in stating the main results.

$$N(m) := \begin{cases} 9, & \text{if } m = 1, \\ 10m \log m - m \log \log m, & \text{if } m \geq 2, \end{cases} \quad (1.5)$$

$$N_0(m) := \left\lceil \left( \frac{N(m)}{\pi} \right)^2 \right\rceil, \quad (1.6)$$

$$N_1(m, k) := \min_{k \in \mathbb{Z} \setminus \{0\}} \{N_0(m) - k\}, \quad (1.7)$$

$$n(N, k) := \begin{cases} \max \left\{ N_0(N+1), 4k, k^2, \left\lceil \frac{k^2 \pi^2}{4} \right\rceil \right\}, & \text{if } k > 0, \\ \max \left\{ N_0(N+1), 4|k| \right\}, & \text{if } k < 0, \end{cases} \quad (1.8)$$

$$\tilde{E}_1(N, k) := \frac{2|k|^{\frac{N}{2}} \sinh(\pi \sqrt{|k|})}{\pi^{3/2}} \left( 1 + \sqrt{2(N+1)} \right), \quad (1.9)$$

$$\tilde{E}_2(N, k) := \frac{2|k|^{\frac{N}{2}} \cosh(\pi \sqrt{|k|}) \sqrt{2(N+2)}}{\pi^{3/2}}, \quad (1.10)$$

$$\bar{E}_1(N, k) := \tilde{E}_1(N, k) + \tilde{E}_2(N, k), \quad (1.11)$$

$$\hat{E}_2(N) = \hat{E}_2(N, k) := \begin{cases} \frac{1+\pi}{\pi^{N+1}}, & \text{if } k > 0, \\ \frac{2^{\frac{N+3}{2}}}{\pi^{N+1}}, & \text{if } k < 0, \end{cases} \quad (1.12)$$

$$E(N, k) := \bar{E}_1(N, k) + \hat{E}_2(N), \quad (1.13)$$

$$n(N, r, j) := \max_{0 \leq m \leq r} \{n(N, -(m+1)j)\} \quad (1.14)$$

$$E(N, r, j) := \sum_{m=0}^r \binom{r}{m} E(N, -(m+1)j). \quad (1.15)$$

**Theorem 1.3.** *Let  $n(N, k)$ ,  $E(N, k)$  be as in (1.8) and (1.13) respectively.. For  $N \in \mathbb{N}$ ,  $k \in \mathbb{Z} \setminus \{0\}$ , and  $n \geq n(N, k)$ , we have*

$$\bar{p}(n+k) = \frac{e^{\pi \sqrt{n}}}{8n} \left( \sum_{t=0}^N \frac{A_k(t)}{n^{\frac{t}{2}}} + O_{\leq E(N, k)} \left( n^{-\frac{N+1}{2}} \right) \right), \quad (1.16)$$

where

$$A_k(t) = \left(\frac{k}{2}\right)^t \sum_{\ell=0}^{\lfloor \frac{t+1}{2} \rfloor} \frac{(-1)^\ell (t+1-\ell)}{(t+1-2\ell)!} \binom{t+1}{\ell} \frac{\pi^{t-2\ell}}{k^\ell}. \quad (1.17)$$

**Theorem 1.4.** Let  $n(N, r, j)$  and  $E(N, r, j)$  be as defined in (1.14) and (1.15) respectively. For all  $j, r \in \mathbb{N}$ ,  $N \in \mathbb{Z}_{\geq r}$  and  $n \geq n(N, r, j)$  we have

$$\Delta_j^r(\bar{p})(n-j) = \frac{e^{\pi\sqrt{n}}}{8n} \left( \sum_{t=r}^N \frac{A_j(t, r)}{n^{\frac{t}{2}}} + O_{\leq E(N, r, j)}(n^{-\frac{N+1}{2}}) \right), \quad (1.18)$$

where

$$A_j(t, r) = \left(-\frac{1}{2}\right)^t \sum_{\ell=0}^{\lfloor \frac{t+1}{2} \rfloor} \binom{t+1}{\ell} \frac{j^{t-\ell} \pi^{t-2\ell} (t+1-\ell)}{(t+1-2\ell)!} \sum_{m=0}^r (-1)^m \binom{r}{m} (m+1)^{t-\ell}. \quad (1.19)$$

**Remark 1.5.** In particular, we see that

$$A_j(r, r) = \left(\frac{\pi j}{2}\right)^r. \quad (1.20)$$

From Theorem 1.4, we have the following asymptotic growth of  $\Delta_j^r(\bar{p})(n)$ .

**Corollary 1.6.** We have

$$\Delta_j^r(\bar{p})(n) \sim \left(\frac{\pi j}{2}\right)^r \frac{e^{\pi\sqrt{n}}}{8 n^{\frac{r}{2}+1}} \quad \text{as } n \rightarrow \infty.$$

Setting  $j = 1$  in Corollary 1.6, we get the asymptotic major term of  $\Delta^r(\bar{p})(n)$  stated below.

**Corollary 1.7.** We have

$$\Delta^r(\bar{p})(n) \sim \left(\frac{\pi}{2}\right)^r \frac{e^{\pi\sqrt{n}}}{8 n^{\frac{r}{2}+1}} \quad \text{as } n \rightarrow \infty.$$

The paper is organized in the following way. In Section 2, we present some preparatory lemmas in deriving asymptotic expansion of the shifted overpartition function which will be helpful to get error estimations carried out in Section 3 that finally lead to the proof of Theorem 1.3 and applying Theorem 1.3, subsequently, we will present the proof of Theorem 1.4. Finally, we conclude the paper with raising some open questions in Section 4.

## 2. PRELIMINARY LEMMAS

Following the sketch done in [19, Lemma 2.1.], we have the following lemma.

**Lemma 2.1.** Let  $N(m)$  be as defined in (1.5). Then for any integer  $m \geq 1$ ,

$$10x^m e^{-\frac{x}{2}} < 1, \quad \text{for } x \geq N(m).$$

*Proof.* For  $m = 1$ , it is clear that  $N(m) = 9$ . For  $m \geq 2$ , define  $f(x) := \frac{x}{2} - m \log x - \log 10$ . Now,  $f'(x) = \frac{1}{2} - \frac{m}{x} > 0$  implies that  $f(x)$  is increasing for  $x > 2m$ . In order to show  $f(x) > 0$  for all  $x \geq N(m)$ , it is enough to show  $f(x) > f(N(m)) > 0$  for all  $x \geq N(m) > 2m$ ; i.e., we first show that (i)  $N(m) > 2m$  and then we show (ii)  $f(N(m)) > 0$ . It is easy to verify that  $N(m) > 2m$ , that is,  $10m \log m - m \log \log m > 2m$  for  $m \geq 2$ . Next, in order to prove  $f(N(m)) > 0$ , it is equivalent to show that

$$5m \log m - \frac{m \log \log m}{2} - m \log(10m \log m - m \log \log m) - \log 10 > 0,$$

which in turn, amounts to prove

$$\log m > \frac{2m+1}{8m} \log 10 + \frac{3}{8} \log \log m + \frac{1}{8} \log \left(1 - \frac{\log \log m}{10 \log m}\right).$$

Since  $\frac{2m+1}{8m} > \frac{1}{4}$  and  $\log \left(1 - \frac{\log \log m}{10 \log m}\right) < 0$  for  $m \geq 3$ , we have  $m > 10^{\frac{1}{4}} \log m^{\frac{3}{8}}$  which is equivalent to the inequality stated before.  $\square$

Set  $\bar{\mu}(n) := \pi\sqrt{n}$ .

**Lemma 2.2.** Let  $N_0(m)$  be as defined in (1.6). Then for all  $k \in \mathbb{Z} \setminus \{0\}$ ,  $m \geq 2$  and  $n \geq N_0(m)$ ,

$$\bar{p}(n+k) = \frac{e^{\bar{\mu}(n+k)}}{8(n+k)} \left(1 - \frac{1}{\bar{\mu}(n+k)} + O_{\leq 1}(\bar{\mu}(n+k)^{-m})\right). \quad (2.1)$$

*Proof.* From [17, equation (3.5)], we have

$$\bar{p}(n) = \frac{e^{\bar{\mu}(n)}}{8n} \left(1 - \frac{1}{\bar{\mu}(n)} + \tilde{T}(n)\right)$$

with

$$\tilde{T}(n) = \left(1 + \frac{1}{\bar{\mu}(n)}\right) e^{-2\bar{\mu}(n)} + \frac{8n}{e^{\bar{\mu}(n)}} R_2(n, 2),$$

and  $R_2(n, 2)$  is the error term of ((1.2)), given in ((1.3)). By [17, equation (3.6)] we have

$$|\tilde{T}(n)| < 10e^{-\frac{1}{2}\bar{\mu}(n)}.$$

Using Lemma 2.1 we get for all  $m \geq 2$  and  $\bar{\mu}(n) \geq N(m)$ ; i.e, for  $n \geq N_0(m)$ ,

$$\bar{p}(n) = \frac{e^{\bar{\mu}(n)}}{8n} \left(1 - \frac{1}{\bar{\mu}(n)} + O_{\leq 1}(\bar{\mu}(n)^{-m})\right). \quad (2.2)$$

Thus applying (2.2) with  $n \mapsto n+k$  and following (1.7), we have for all  $k \in \mathbb{Z} \setminus \{0\}$ ,  $m \geq 2$  and  $n \geq N_1(m, k)$ ,

$$\bar{p}(n+k) = \frac{e^{\bar{\mu}(n+k)}}{8(n+k)} \left(1 - \frac{1}{\bar{\mu}(n+k)} + O_{\leq 1}(\bar{\mu}(n+k)^{-m})\right).$$

This finishes the proof of the lemma.  $\square$

To derive the coefficient sequence arising from the Taylor expansion of the asymptotic main term  $\frac{e^{\bar{\mu}(n+k)}}{8(n+k)} \left(1 - \frac{1}{\bar{\mu}(n+k)}\right)$  after extracting out the factor  $\frac{e^{\pi\sqrt{n}}}{8n}$ , we closely follow the methodology carried out in [23, Proposition 4.4].

**Lemma 2.3.** *For  $k \in \mathbb{Z} \setminus \{0\}$ , we have*

$$\frac{e^{\bar{\mu}(n+k)}}{8(n+k)} \left(1 - \frac{1}{\bar{\mu}(n+k)}\right) = \frac{e^{\pi\sqrt{n}}}{8n} \sum_{t \geq 0} \frac{A_k(t)}{n^{\frac{t}{2}}},$$

with

$$A_k(t) := \left(\frac{\pi k}{2}\right)^t \sum_{\ell=0}^{\lfloor \frac{t+1}{2} \rfloor} \left(-\frac{1}{\pi^2 k}\right)^\ell \frac{(t-\ell+1)}{(t+1-2\ell)!} \binom{t+1}{\ell}.$$

*Proof.* We first estimate  $\frac{e^{\bar{\mu}(n+k)}}{8(n+k)} \left(1 - \frac{1}{\bar{\mu}(n+k)}\right)$  up to some truncation point say  $N \in \mathbb{N}$ .

Setting  $z := \frac{1}{\sqrt{n}}$  and  $c := \pi$ , we have

$$\begin{aligned} \frac{e^{\bar{\mu}(n+k)}}{8(n+k)} \left(1 - \frac{1}{\bar{\mu}(n+k)}\right) &= \frac{e^{\pi\sqrt{n}}}{8n} \left( \frac{e^{\pi\sqrt{n}(\sqrt{1+\frac{k}{n}}-1)}}{(1+\frac{k}{n})} \left(1 - \frac{1}{\pi\sqrt{n}\sqrt{1+\frac{k}{n}}}\right) \right) \\ &= \frac{e^{\pi\sqrt{n}}}{8n} \left[ e^{\pi\sqrt{n}(\sqrt{1+\frac{k}{n}}-1)} \left( \frac{1}{1+\frac{k}{n}} - \frac{1}{\pi\sqrt{n}(1+\frac{k}{n})^{3/2}} \right) \right] \\ &= \frac{e^{\pi\sqrt{n}}}{8n} \left[ \exp\left(\frac{c}{z}(\sqrt{1+kz^2}-1)\right) \left( \frac{1}{1+kz^2} - \frac{1}{\frac{c}{z}(1+kz^2)^{3/2}} \right) \right]. \end{aligned} \quad (2.3)$$

Let  $\alpha := c\sqrt{n}$ ,  $x := kz^2 = \frac{k}{n}$ . Considering the left hand side of (2.3) as a formal power series in  $x$ , we get

$$\exp(\alpha(\sqrt{x+1}-1)) \left( \frac{1}{1+x} - \frac{1}{\alpha(1+x)^{3/2}} \right) = \sum_{j=0}^{\infty} \xi_j(\alpha) x^j. \quad (2.4)$$

Integrating (2.4) twice with respect to  $x$  yields

$$\xi_j(\alpha) = (j+1)(j+2)[x^{j+2}] \frac{4}{\alpha^2} \sum_{\ell=0}^{\infty} \frac{\alpha^\ell}{\ell!} (\sqrt{1+x}-1)^\ell.$$

Let  $y := \sqrt{1+x}$  and  $\mathcal{B}_{-1}(x)$  be the generalized binomial series (see [11, page 203]). Then for all  $m \in \mathbb{C}$  with  $m \notin \mathbb{Z}_{\geq 0}$ , it follows that

$$\left(\frac{1+y}{2}\right)^m = \left(\frac{1+\sqrt{1+x}}{2}\right)^m = \mathcal{B}_{-1}\left(\frac{x}{4}\right)^m = \sum_{j=0}^{\infty} \frac{m}{m-j} \binom{m-j}{j} \frac{x^j}{4^j}. \quad (2.5)$$

Therefore,

$$\xi_j(\alpha) = (j+1)(j+2) \sum_{\ell=0}^{\infty} \left(\frac{\alpha}{2}\right)^{\ell-2} \frac{1}{\ell!} [x^{j+2-\ell}] \left(\frac{1+y}{2}\right)^{-\ell}$$

$$\begin{aligned}
&= (j+1)(j+2) \sum_{\ell=0}^{j+1} \left(\frac{\alpha}{2}\right)^{j-\ell} \frac{1}{(j+2-\ell)!} [x^\ell] \left(\frac{1+y}{2}\right)^{-(j+2-\ell)} \\
&= (j+1) \sum_{\ell=0}^{j+1} \left(\frac{\alpha}{2}\right)^{j-\ell} \frac{1}{(j+2-\ell)!} \frac{j+2-\ell}{j+2} \binom{-j-2}{\ell} \frac{1}{4^\ell} \\
&= \sum_{\ell=0}^{j+1} \frac{(j+1)\alpha^{j-\ell}(-1)^\ell}{2^{j+\ell}(j+\ell+1)!} \binom{j+\ell+1}{\ell},
\end{aligned}$$

where in the second step, we have used (2.5) along with the fact that

$$\left(\frac{1+y}{2}\right)^{-(j+2-\ell)} = 0 \quad \text{for } \ell \geq j+2 \quad \text{as } \binom{m-j-1}{j-1} = 0 \text{ for } \left\lfloor \frac{m}{2} \right\rfloor + 1 \leq j \leq m-1.$$

Since, in (2.4),  $\xi_j(\alpha)x^j$  contains the term  $\alpha^{j-\ell}x^j = \pi^{j-\ell}k^j n^{-\frac{j+\ell}{2}}$ , thus we have

$$A_k(t) = \sum_{\substack{j+\ell=t \\ \ell \leq j+1}} \frac{(-1)^\ell(j+1)\pi^{j-\ell}k^j}{2^{j+\ell}(j+1-\ell)!} \binom{j+\ell+1}{\ell} = \sum_{\ell=0}^{\lfloor \frac{t+1}{2} \rfloor} \frac{(-1)^\ell(t-\ell+1)\pi^{t-2\ell}k^{t-\ell}}{2^t(t+1-2\ell)!} \binom{t+1}{\ell},$$

as  $j+\ell=t$  and  $\ell \leq j+1$ , we have  $t-j \leq j+1 \Leftrightarrow j \geq \frac{t-1}{2}$  which implies  $\ell \leq t-j \leq \frac{t+1}{2}$  and this finally concludes the proof.  $\square$

### 3. PROOF OF THEOREM 1.3 AND THEOREM 1.4

Before we present the proof of Theorem 1.3, we first need to estimate an error bound of  $\sum_{t \geq N+1} \frac{A_k(t)}{n^{\frac{t}{2}}}$ .

**Lemma 3.1.** *Let  $n(N, k)$  be as in (1.8) and  $\bar{E}_1(N, k)$  be as in (1.11). Then for  $k \neq 0$ , and  $n \geq n(N, k)$ , we have*

$$\sum_{t \geq N+1} \frac{A_k(t)}{n^{\frac{t}{2}}} = O_{\leq \bar{E}_1(N, k)} \left( n^{-\frac{N+1}{2}} \right).$$

*Proof.* From Lemma 2.3, it follows that for  $k > 0$ ,

$$A_k(2t) = \left(-\frac{k}{4}\right)^t \sum_{\ell=0}^t (-1)^\ell \binom{2t+1}{t-\ell} (t+\ell+1) \frac{(\pi\sqrt{k})^{2\ell}}{(2\ell+1)!}, \quad (3.1)$$

$$A_k(2t+1) = \frac{k^t(-1)^{t+1}}{2^{2t+1}\pi} \sum_{\ell=0}^{t+1} (-1)^\ell \binom{2t+2}{t-\ell+1} (t+\ell+1) \frac{(\pi\sqrt{k})^{2\ell}}{(2\ell)!}. \quad (3.2)$$

For  $t \in \mathbb{N}$ , define  $c(t, \ell) := \binom{2t+1}{t-\ell} (t+\ell+1)$  and we see that

$$\frac{c(t, \ell)}{c(t, 0)} = \frac{\prod_{j=1}^\ell (t-j+1)}{\prod_{j=1}^\ell (t+j)} \leq 1,$$

which implies that  $c(t, \ell) \leq c(t, 0)$  for all  $0 \leq \ell \leq t$ . Consequently, using (3.1) and the fact that  $\binom{2t}{t} \leq \frac{4^t}{\sqrt{\pi t}}$ , we have

$$|A_k(2t)| \leq \frac{1}{\pi\sqrt{k}} \binom{k}{4}^t \binom{2t+1}{t} (t+1) \left| \sum_{\ell=0}^t \frac{(-1)^\ell (\pi\sqrt{k})^{2\ell+1}}{(2\ell+1)!} \right| \leq \frac{k^{t-1/2} (2t+1)}{\pi^{3/2} \sqrt{t}} \sinh(\pi\sqrt{k}),$$

and so

$$\begin{aligned} \sum_{t \geq \frac{N+1}{2}} \frac{|A_k(2t)|}{n^t} &\leq \frac{k^{-\frac{1}{2}} \sinh(\pi\sqrt{k})}{\pi^{3/2}} \sum_{t \geq \frac{N+1}{2}} \frac{2t+1}{\sqrt{t}} \frac{k^t}{n^t} \\ &\leq \frac{k^{\frac{N}{2}} \sinh(\pi\sqrt{k})}{\pi^{3/2}} n^{-\frac{N+1}{2}} \sum_{t \geq 0} \left( 2\sqrt{\frac{N+1}{2}} \sqrt{1 + \frac{2t}{N+1}} + \frac{1}{\sqrt{t + \frac{N+1}{2}}} \right) \left( \frac{k}{n} \right)^t \\ &\leq \frac{k^{\frac{N}{2}} \sinh(\pi\sqrt{k})}{\pi^{3/2}} n^{-\frac{N+1}{2}} \sum_{t \geq 0} \left( 2\sqrt{\frac{N+1}{2}} \sqrt{1+t} + \frac{1}{\sqrt{t+1}} \right) \left( \frac{k}{n} \right)^t \\ &\leq \frac{k^{\frac{N}{2}} \sinh(\pi\sqrt{k})}{\pi^{3/2}} n^{-\frac{N+1}{2}} \left( 1 + \sqrt{2(N+1)} \right) \sum_{t \geq 0} \left( \frac{2k}{n} \right)^t \\ &\leq \left( \frac{2k^{\frac{N}{2}} \sinh(\pi\sqrt{k})}{\pi^{3/2}} \left( 1 + \sqrt{2(N+1)} \right) \right) n^{-\frac{N+1}{2}} = \tilde{E}_1(N, k) n^{-\frac{N+1}{2}}, \end{aligned} \quad (3.3)$$

using the fact  $\sqrt{t+1} \leq 2^t$  for  $t > 0$  in the fourth step and  $n \geq n(N, k) \geq 4k$  in the second last step and (1.9) in the final step.

Similarly, we estimate

$$|A_k(2t+1)| \leq \frac{k^t (t+1)}{4^t 2\pi} \binom{2t+2}{t+1} \cosh(\pi\sqrt{k}) \leq \frac{2k^t \sqrt{t+1}}{\pi^{3/2}} \cosh(\pi\sqrt{k}),$$

and consequently using (3.2), (1.10), and  $n \geq n(N, k) \geq 4k$ , it follows that

$$\begin{aligned} \sum_{t \geq \frac{N}{2}} \frac{|A_k(2t+1)|}{n^{t+\frac{1}{2}}} &\leq n^{-\frac{1}{2}} \frac{2 \cosh(\pi\sqrt{k})}{\pi^{3/2}} \sum_{t \geq \frac{N}{2}} \sqrt{t+1} \frac{k^t}{n^t} \\ &\leq \frac{2k^{N/2} \cosh(\pi\sqrt{k})}{\pi^{3/2}} n^{-\frac{N+1}{2}} \sum_{t \geq 0} \frac{k^t}{n^t} \sqrt{t + \frac{N}{2} + 1} \leq \tilde{E}_2(N, k) n^{-\frac{N+1}{2}}. \end{aligned} \quad (3.4)$$

Combining (3.3) and (3.4), we get

$$\begin{aligned} \left| \sum_{t \geq N+1} \frac{A_k(t)}{n^{\frac{t}{2}}} \right| &= \left| \sum_{t \geq \frac{N+1}{2}} \frac{A_k(2t)}{n^t} + \sum_{t \geq \frac{N}{2}} \frac{A_k(2t+1)}{n^{t+\frac{1}{2}}} \right| \leq \sum_{t \geq \frac{N+1}{2}} \frac{|A_k(2t)|}{n^t} + \sum_{t \geq \frac{N}{2}} \frac{|A_k(2t+1)|}{n^{t+\frac{1}{2}}} \\ &= \overline{E}_1(N, k) n^{-\frac{N+1}{2}}. \end{aligned} \quad (3.5)$$

This concludes the proof of for the case  $k > 0$ . Next, we proceed to the case where  $k < 0$ .

Let  $k = -m$ ,  $m \in \mathbb{Z}_{>0}$ ; i.e,  $m = |k|$ . Then we have

$$A_{-m}(t) = \frac{(-m)^t}{2^t} \sum_{\ell=0}^{\lfloor \frac{t+1}{2} \rfloor} \frac{(t+1-\ell)}{(t+1-2\ell)!} \binom{t+1}{\ell} \frac{\pi^{t-2\ell}}{m^\ell}.$$

In a similar way like in the case  $k \in \mathbb{Z}_{>0}$ , here we have the following bounds

$$|A_{-m}(2t+1)| \leq \frac{2m^t \sqrt{t+1}}{\pi^{3/2}} \cosh(\pi\sqrt{m}), \quad |A_{-m}(2t)| \leq \frac{m^{t-\frac{1}{2}} (2t+1)}{\pi^{3/2} \sqrt{t}} \sinh(\pi\sqrt{m}),$$

which consequently implies that for  $n \geq n(N, k)$

$$\sum_{t \geq \frac{N}{2}} \frac{|A_k(2t+1)|}{n^{t+\frac{1}{2}}} \leq \tilde{E}_2(N, k) n^{-\frac{N+1}{2}} \quad \text{and} \quad \sum_{t \geq \frac{N+1}{2}} \frac{|A_k(2t)|}{n^t} \leq \tilde{E}_1(N, k) n^{-\frac{N+1}{2}}.$$

Analogous to the estimation done in (3.5), in this case, we have also

$$\left| \sum_{t \geq N+1} \frac{A_k(t)}{n^{\frac{t}{2}}} \right| \leq \bar{E}_1(N, k) n^{-\frac{N+1}{2}},$$

which concludes the proof.  $\square$

**Lemma 3.2.** *Let  $n(N, k)$  be as in (1.8) and  $\hat{E}_1(N, k)$  be as in (1.12). Then for  $k \neq 0$ , and  $n \geq n(N, k)$ , we have*

$$\frac{e^{\pi\sqrt{n+k}}}{8(n+k)} \bar{\mu}(n+k)^{-(N+1)} = \frac{e^{\pi\sqrt{n}}}{8n} O_{\leq \hat{E}_2(N)} \left( n^{-\frac{N+1}{2}} \right).$$

*Proof.* Before we move on to estimate the error term, we note down the following well known Bernoulli's inequality

$$(1+x)^r \leq 1 + rx \quad \text{for } 0 \leq r \leq 1 \quad \text{and} \quad x \geq -1. \quad (3.6)$$

For  $k > 0$  and  $n \geq n(N, k) \geq \max\{k^2, \frac{k^2\pi^2}{4}\}$ , and using the fact that  $e^x < 1+2x$  for  $0 < x < 1$ , we bound the error term as

$$\begin{aligned} \frac{e^{\pi\sqrt{n+k}}}{8(n+k)} \bar{\mu}(n+k)^{-(N+1)} &= \frac{e^{\pi\sqrt{n}}}{8n} \frac{e^{\pi\sqrt{n}(\sqrt{1+\frac{k}{n}}-1)}}{\pi^{N+1} n^{\frac{N+1}{2}}} \frac{1}{(1+\frac{k}{n})^{\frac{N+3}{2}}} \leq \frac{e^{\pi\sqrt{n}}}{8n} \frac{e^{\pi\sqrt{n}(\sqrt{1+\frac{k}{n}}-1)}}{\pi^{N+1} n^{\frac{N+1}{2}}} \\ &\leq \frac{e^{\pi\sqrt{n}}}{8n} \frac{e^{\frac{k\pi}{2\sqrt{n}}}}{\pi^{N+1} n^{\frac{N+1}{2}}} \leq \frac{e^{\pi\sqrt{n}}}{8n} \left( \frac{1+\pi}{\pi^{N+1}} \right) n^{-\frac{N+1}{2}} = \hat{E}_2(N) n^{-\frac{N+1}{2}}, \end{aligned}$$

using (3.6) in the third step and for  $k < 0$ , we estimate in the following way

$$\begin{aligned} \frac{e^{\pi\sqrt{n-k}}}{8(n-k)} \bar{\mu}(n-k)^{-(N+1)} &= \frac{e^{\pi\sqrt{n}}}{8n} \frac{e^{\pi\sqrt{n}(\sqrt{1-\frac{k}{n}}-1)}}{\pi^{N+1} n^{-\frac{N+1}{2}}} \frac{1}{(1-\frac{k}{n})^{\frac{N+3}{2}}} \leq \frac{e^{\pi\sqrt{n}}}{8n} \frac{2^{\frac{N+3}{2}} e^{\pi\sqrt{n}(\sqrt{1-\frac{k}{n}}-1)}}{\pi^{N+1} n^{-\frac{N+1}{2}}} \\ &\leq \frac{e^{\pi\sqrt{n}}}{8n} \frac{2^{\frac{N+3}{2}} e^{-\frac{k\pi}{2\sqrt{n}}}}{\pi^{N+1} n^{-\frac{N+1}{2}}} \leq \frac{e^{\pi\sqrt{n}}}{8n} \hat{E}_2(N) n^{-\frac{N+1}{2}}, \end{aligned}$$

using (3.6) in the third step. From the above two estimates done separately for both cases for  $k$ , we conclude the proof.  $\square$

*Proof of Theorem 1.3* Applying Lemmas 2.3, 3.1, and 3.2 to (2.2) with (1.13), we get for  $n \geq n(N, k)$ ,

$$\bar{p}(n+k) = \frac{e^{\pi\sqrt{n}}}{8n} \left( \sum_{t=0}^N \frac{A_k(t)}{n^{\frac{t}{2}}} + O_{\leq E(N, k)} \left( n^{-\frac{N+1}{2}} \right) \right).$$

This finishes the proof.  $\square$

Before we proceed to present the proof of Theorem 1.4, we recall the definition of Stirling number. Adopting the notation from [11], here  $\{n\}_m$  denotes the Stirling number of second kind which counts the number of ways to partition a set of  $n$  elements into  $m$  nonempty subsets. Here we state three elementary facts about  $\{n\}_m$ . From [11, Table 264 and equation (6.19)], we have

$$\{n\}_m = 0 \text{ for } m > n \text{ and } m! \{n\}_m = \sum_{k=0}^m \binom{m}{k} k^n (-1)^{m-k}. \quad (3.7)$$

Using Theorem 1.3, in this section, we are now ready to prove Theorem 1.4.

*Proof of Theorem 1.4* Following the definition of the operator  $\Delta_j^r$ , it follows that

$$\Delta_j^r(\bar{p})(n-j) = \sum_{m=0}^r (-1)^m \binom{r}{m} \bar{p}(n-m.j-j) = \sum_{m=0}^r (-1)^m \binom{r}{m} \bar{p}(n-(m+1)j),$$

and therefore, with  $k \mapsto -(m+1)j$ , by Theorem 1.3, we have for  $n \geq \max_{0 \leq m \leq r} \{n(N, -(m+1)j)\}$ ,

$$\begin{aligned} \Delta_j^r(\bar{p})(n-j) &= \sum_{\substack{m=0 \\ m \equiv 0 \pmod{2}}}^r \binom{r}{m} \bar{p}(n-(m+1)j) - \sum_{\substack{m=0 \\ m \equiv 1 \pmod{2}}}^r \binom{r}{m} \bar{p}(n-(m+1)j) \\ &\leq \frac{e^{\pi\sqrt{n}}}{8n} \sum_{\substack{m=0 \\ m \equiv 0 \pmod{2}}}^r \binom{r}{m} \left( \sum_{t=0}^N \frac{A_{-(m+1).j}(t)}{n^{\frac{t}{2}}} + \frac{E(N, -(m+1)j)}{n^{\frac{N+1}{2}}} \right) \\ &\quad - \frac{e^{\pi\sqrt{n}}}{8n} \sum_{\substack{m=0 \\ m \equiv 1 \pmod{2}}}^r \binom{r}{m} \left( \sum_{t=0}^N \frac{A_{-(m+1).j}(t)}{n^{\frac{t}{2}}} - \frac{E(N, -(m+1)j)}{n^{\frac{N+1}{2}}} \right) \\ &= \frac{e^{\pi\sqrt{n}}}{8n} \left( \sum_{m=0}^r (-1)^m \binom{r}{m} \sum_{t=0}^N \frac{A_{-(m+1).j}(t)}{n^{\frac{t}{2}}} + \sum_{m=0}^r \binom{r}{m} \frac{E(N, -(m+1)j)}{n^{\frac{N+1}{2}}} \right) \\ &= \frac{e^{\pi\sqrt{n}}}{8n} \left( \sum_{m=0}^r (-1)^m \binom{r}{m} \sum_{t=0}^N \frac{A_{-(m+1).j}(t)}{n^{\frac{t}{2}}} + \frac{E(N, r, j)}{n^{\frac{N+1}{2}}} \right). \end{aligned} \quad (3.8)$$

Similarly, it follows that

$$\Delta_j^r(\bar{p})(n-j) \geq \frac{e^{\pi\sqrt{n}}}{8n} \left( \sum_{m=0}^r (-1)^m \binom{r}{m} \sum_{t=0}^N \frac{A_{-(m+1).j}(t)}{n^{\frac{t}{2}}} - \frac{E(N, r, j)}{n^{\frac{N+1}{2}}} \right). \quad (3.9)$$

Hence, combining (3.8) and (3.9), for  $n \geq n(N, r, j)$ , we get

$$\Delta_j^r(\bar{p})(n-j) = \frac{e^{\pi\sqrt{n}}}{8n} \left( \sum_{m=0}^r (-1)^m \binom{r}{m} \sum_{t=0}^N \frac{A_{-(m+1),j}(t)}{n^{\frac{t}{2}}} + O_{\leq E(N,r,j)}\left(n^{-\frac{N+1}{2}}\right) \right). \quad (3.10)$$

Next, following (1.17) with  $k \mapsto -(m+1)j$ , we simplify

$$\begin{aligned} & \sum_{m=0}^r (-1)^m \binom{r}{m} \sum_{t=0}^N \frac{A_{-(m+1),j}(t)}{n^{\frac{t}{2}}} \\ &= \sum_{m=0}^r (-1)^m \binom{r}{m} \sum_{t=0}^N \left( \frac{-(m+1)j}{2\sqrt{n}} \right)^t \sum_{\ell=0}^{\lfloor \frac{t+1}{2} \rfloor} \frac{(t+1-\ell)}{(t+1-2\ell)!} \binom{t+1}{\ell} \frac{\pi^{t-2\ell}}{((m+1)j)^\ell} \\ &= \sum_{t=0}^N \left( \left( -\frac{1}{2} \right)^t \sum_{\ell=0}^{\lfloor \frac{t+1}{2} \rfloor} \binom{t+1}{\ell} \frac{j^{t-\ell} \pi^{t-2\ell} (t+1-\ell)}{(t+1-2\ell)!} \sum_{m=0}^r (-1)^m \binom{r}{m} (m+1)^{t-\ell} \right) \frac{1}{n^{\frac{t}{2}}} \\ &= \sum_{t=0}^N \frac{A_j(t, r)}{n^{t/2}}. \end{aligned} \quad (3.11)$$

We observe that for  $0 \leq t \leq r-1$ ,

$$\begin{aligned} \sum_{m=0}^r (-1)^m \binom{r}{m} (m+1)^{t-\ell} &= \sum_{m=0}^r (-1)^m \binom{r}{m} \sum_{s=0}^{t-\ell} \binom{t-\ell}{s} m^s \\ &= (-1)^r r! \sum_{s=0}^{t-\ell} \binom{t-\ell}{s} \left\{ \begin{matrix} s \\ r \end{matrix} \right\}, \end{aligned} \quad (3.12)$$

using (3.7) in the final step. As  $\ell \geq 0$ , so for  $s \leq t-\ell < r$ , using (3.7), we have

$$\sum_{s=0}^{t-\ell} \binom{t-\ell}{s} \left\{ \begin{matrix} s \\ r \end{matrix} \right\} = 0,$$

Consequently, (3.11) reduces to

$$\sum_{m=0}^r (-1)^m \binom{r}{m} \sum_{t=0}^N \frac{A_{-(m+1),j}(t)}{n^{\frac{t}{2}}} = \sum_{t=r}^N \frac{A_j(t, r)}{n^{t/2}}.$$

Hence, with (3.10), we finish the proof of (1.18).

Next, to prove (1.20), from (3.12), we see that for  $t = r$  and  $r \in \mathbb{N}$ ,

$$\sum_{m=0}^r (-1)^m \binom{r}{m} (m+1)^{r-\ell} = \begin{cases} 0, & \text{if } \ell \geq 1 \text{ and } \ell = 0, 0 \leq s < r, \\ (-1)^r r!, & \text{if } (\ell, s) = (0, r). \end{cases}$$

Hence, with the only choice  $(\ell, t) = (0, r)$ , from (3.11), it follows that

$$A_j(r, r) = (-1)^r r! \left( -\frac{1}{2} \right)^r \frac{j^r \pi^r (r+1)}{(r+1)!} = \left( \frac{\pi j}{2} \right)^r.$$

This concludes the proof of the theorem.  $\square$

#### 4. CONCLUSION

We conclude this paper by raising the following questions with a brief background on each of the problem.

(1) Following up the work of Odlyzko [22] on the positivity of  $\Delta^r(p)(n)$  for  $n \geq n(r)$  with  $n(r) \sim \frac{6}{\pi^2} r^2 (\log r)^2$  as  $r \rightarrow \infty$ , we ask the following:

**Problem 4.1.** *What is the asymptotic growth of  $n(r)$  as  $r \rightarrow \infty$  such that for all  $n \geq n(r)$ ,  $\Delta^r(\bar{p})(n) \geq 0$ ? Can one obtain an exact (and/or simplified) formula for such  $n(r)$  as in Almkvist, Knessl and Keller's work [1, 15] for the overpartition function?*

(2) Let  $PL(n)$  denotes the plane partition function introduced by MacMahon [18], and the generating function is given by

$$\sum_{n=0}^{\infty} PL(n)q^n = \prod_{n=1}^{\infty} \frac{1}{(1-q^n)^n}.$$

Knessl [14] studied the positivity of  $\Delta^r(PL)(n)$ , showed that  $\Delta^r(PL)(n) > 0$  for  $n \geq n(r)$  with

$$n(r) \sim \frac{2}{3} (3\zeta(3))^{-\frac{1}{2}} (r \log r)^{\frac{3}{2}} \text{ as } r \rightarrow \infty.$$

Recently, thanks to the seminal work of Ono, Pujahari and Rolen [21] on the asymptotic expansion of  $PL(n)$ , a substantial modification of Wright's formula [26], we have an explicit estimate of the error bound for the asymptotic expansion of  $PL(n)$ . We raise the following problem for  $PL(n)$ .

**Problem 4.2.** *For  $r \in \mathbb{N}$ , what is the asymptotic growth of  $\Delta_j^r(PL)(n)$  as  $n \rightarrow \infty$ ?*

#### Acknowledgements

The author would like to thank the anonymous referee for his/her comments that improved the paper substantially. This work has been carried out as a postdoctoral fellow at school of mathematical sciences, National Institute of Science Education and Research, Bhubaneswar. The author would like to thank the institute for its hospitality and support.

#### REFERENCES

- [1] G. Almkvist, *On the differences of the partition function*, Acta Arith. **61** (1992), 173–181.
- [2] K. Banerjee, *Invariants of the quartic binary form and proofs of Chen's conjectures on inequalities for the partition function and the Andrews' spt function*, Res. Math. Sci. **12** (2025), <https://doi.org/10.1007/s40687-024-00485-4>.
- [3] K. Banerjee, *A unified framework to prove multiplicative inequalities for the partition function*, Adv. in Appl. Math. **152** (2024), p. 102590.
- [4] K. Banerjee, *Positivity of the second shifted difference of partitions and overpartitions: a combinatorial approach*, *Enumer. Comb. Appl.* **3:2** (2023), article #S2R12.
- [5] W. Y. C. Chen, L. Wang, and G. Xie, *Finite differences of the logarithm of the partition function*, *Math. Comp.* **85** (2016), 825–847.

- [6] S. Corteel and J. Lovejoy, *Overpartitions*, Trans. Amer. Math. Soc. **356** (2004), 1623–1635.
- [7] S. DeSalvo and I. Pak, *Log-concavity of the partition function*, Ramanujan J. **38** (2015), 61–73.
- [8] B. Engel, *Log-concavity of the overpartition function*, Ramanujan J. **43** (2017), 229–241.
- [9] K. Gomez, J. Males, and L. Rolen, *The second shifted difference of partitions and its applications*, Bull. Aust. Math. Soc. **107** (2023), 66–78.
- [10] I. J. Good, *The differences of the partition function*, Problem 6137, Amer. Math. Monthly **84** (1977), p. 141.
- [11] R. L. Graham, D. E. Knuth, and O. Patashnik, *Concrete Mathematics*, Addison Wesley, Reading MA, 2nd edition, 1994.
- [12] H. Gupta, *Finite differences of the partition function*, Math. Comp. **32** (1978), 1241–1243.
- [13] G. H. Hardy and S. Ramanujan, *Asymptotic formulae in combinatory analysis*, Proc. Lond. Math. Soc. (3) **2** (1918), 75–115.
- [14] C. Knobell, *Asymptotic behavior of high-order differences of the plane partition function*, Discrete Math. **126** (1994), 179–193.
- [15] C. Knobell and J. B. Keller, *Asymptotic behavior of high-order differences of the partition function*, Comm. Pure Appl. Math. **44** (1991), 1033–1045.
- [16] D. H. Lehmer, *On the remainders and convergence of the series for the partition function*, Trans. Amer. Math. Soc. **46** (1939), 362–373.
- [17] E. Y. S. Liu and H. W. J. Zhang, *Inequalities for the overpartition function*, Ramanujan J. **54** (2021), 485–509.
- [18] P. A. MacMahon, *Combinatory analysis*, Vol. I, II, Dover Publ., Mineola, New York, 2004.
- [19] G. Mukherjee, H. W. J. Zhang, and Y. Zhong, *Higher order log-concavity of the overpartition function and its consequences*, Proc. Edinb. Math. Soc. (2) **66** (2023), 164–181.
- [20] G. Mukherjee, *Inequalities for higher order differences of the logarithm of the overpartition function and a problem of Wang–Xie–Zhang*, Res. Number Theory **9**, 9 (2023).
- [21] K. Ono, S. Pujahari, and L. Rolen, *Turán inequalities for the plane partition function*, Adv. Math. **409** (2022), p. 108692.
- [22] A. M. Odlyzko, *Differences of the partition function*, Acta Arith. **49** (1988), 237–254.
- [23] C. O’Sullivan, *Detailed asymptotic expansions for partitions into powers*, Int. J. Number Theory **9** (2023), 2163–2196.
- [24] H. Rademacher, *On the expansion of the partition function in a series*, Ann. of Math. (2) **44** (1943), 416–422.
- [25] L. X. W. Wang, G. Y. B. Xie, and A. Q. Zhang, *Finite difference of the overpartition function*, Adv. in Appl. Math. **92** (2018), 51–72.
- [26] E. Wright, *Asymptotic partition formulae I: plane partitions*, Q. J. Math. **1** (1931), 177–189.
- [27] H. S. Zuckerman, *On the coefficients of certain modular forms belonging to subgroups of the modular group*, Trans. Amer. Math. Soc. **45** (1939), 298–321.

SCHOOL OF MATHEMATICAL SCIENCES, NATIONAL INSTITUTE OF SCIENCE EDUCATION AND RESEARCH, BHBANESWAR, AN OCC OF HOMI BHABHA NATIONAL INSTITUTE, VIA- JATNI, KHURDA, ODISHA- 752050, INDIA

*Email address:* `gargi@niser.ac.in`