

# Many-electron characterizations of higher-charge superconductors

Zi-Hao Dong and Yi Zhang\*

International Center for Quantum Materials, School of Physics, Peking University, Beijing, 100871, China

(Dated: January 1, 2026)

The theoretical understanding of conventional superconductivity as the phonon-assisted formation and condensation of two-electron Cooper pairs is a significant triumph in condensed matter physics. Here, we propose many-electron characterizations of higher-charge superconductivity with Cooper pairs consisting of more than two electrons, by implementing translation symmetrization on parent pair-density-wave-ordered states. In particular, we demonstrate many-electron constructions with vanishing charge-2e sectors, but with sharp signatures in charge-4e or charge-6e expectation values instead. Such characterizations are consistent with previous phenomenology of vestigial order and Ginzburg-Landau theory, yet, instead of point-group-symmetry presumptions, we show that momentum conservation is both vital and sufficient. Our study thus offers a novel, general, and microscopic route to understand and characterize higher-charge superconductivity, including non-trivial experimental signatures such as fractional magnetic flux and period in interferometry, as well as localized Cooper pairs at lattice topological defects.

*Introduction.*— Superconductors naturally exhibit zero electrical resistance and the Meissner effect below a critical temperature [1–3]. Discovered by Onnes in 1911, the origin of superconductivity remained a microscopic mystery, despite the phenomenological success of its Ginzburg-Landau theory [4], until the BCS theory was proposed in 1957, which explains the dissipation-less current through the formation and condensation of Cooper pairs - two electrons bound together via lattice-mediated attractive interactions [5, 6]. Such Cooper pairs, when generalized to finite momenta, give rise to pair density waves (PDWs), whose superconducting order parameter breaks translation symmetries and varies periodically in space [7–15]. Signatures of such exotic phases have been established in recent experiments [16–20], including those involving the layered Kagomé material  $\text{CsV}_3\text{Sb}_5$  [18, 19].

Higher-charge superconductors can emerge as vestigial orders of an incommensurate PDW, which may be (partially) melted by disorder, thermal fluctuations, quantum fluctuations, or quasi-crystal structures [9, 21–25]. Subsequently, the broken continuous translation symmetry is restored, and only the  $U(1)$  gauge symmetry (charge conservation) breaking is retained. In a Ginzburg-Landau theory framework, these exotic superconducting phases reflect higher orders of the PDW order parameter fields, characterized by multiplets of Cooper pairs, such as with charges of 4e or 6e [9, 21–23, 26–31]. The corresponding fractional magnetic flux quanta, aside from alternatives such as multi-component or time-reversal-symmetry-breaking superconductors [32–35], have been potentially observed in quantum-interference experiments [36]. While theoretical mechanisms have been explored via the Hubbard models [37–40], vestigial fluctuations [41–43], and geometric frustration [44–46], a comprehensive, microscopic many-electron characterization remains lacking, limiting our description and understanding of these fascinating quantum phenomena and correlated systems.

Candidate wavefunctions play a crucial role in understanding exotic quantum matters. A landmark example is the Laughlin wavefunctions, which successfully characterize fractional quantum Hall liquids and encode the fractional statistics of topological order [47–49]. In the realm of quantum spin liquids, Gutzwiller-projected wavefunctions derived from slave-particle approaches offer a powerful variational ansatz [50–53], supporting an effective field theory for universal behaviors such as long-range entanglement and emergent gauge structures. Overall, these wavefunctions serve not only as trial states for variational studies [54–62] but also as conceptual bridges linking microscopic models to emergent quantum phases. Furthermore, the restoration of translation symmetry via projection is a well-established operation for recovering well-defined quantum numbers in variational studies across quantum chemistry [63], nuclear physics [64, 65], and condensed matter physics [66, 67].

In this work, we propose and study candidate many-electron quantum states of higher-charge superconductors, based on translation-symmetrizing parent PDW states. For the simplest scenario of a charge-4e superconducting state rooted from a stripe PDW in two dimensions, the resulting states display vanishing single-Cooper-pair sectors, yet finite expectation values over certain charge-4e Cooper pairs, e.g.,  $\langle c_{\mathbf{k},\uparrow}^\dagger c_{-\mathbf{k}+\mathbf{Q},\downarrow}^\dagger c_{\mathbf{k}',\uparrow}^\dagger c_{-\mathbf{k}'-\mathbf{Q},\downarrow}^\dagger \rangle$ , a composite of two Cooper pairs with momentum  $\pm\mathbf{Q}$ , respectively. Generalizations to even higher-charge states, such as charge-6e, and to higher dimensions, are also straightforward. The formalism is consistent with the physical picture of vortex binding between the PDW order parameter fields in previous vestigial order and classical Ginzburg-Landau theory studies. Unlike previous presumptions and beliefs, however, we demonstrate that while momentum conservation between the PDW order parameter fields is crucial, the lattice geometry and point-group symmetry are less dominant, which we further elaborate in a perturbation theory.

Furthermore, we examine higher-charge superconduc-

\* frankzhangyi@pku.edu.com

tors with respect to an interferometry geometry and observe apparent periodicity in the threading magnetic fluxes associated with the proposed higher-charge Cooper pairs, which differs significantly from that of the parent charge-2e PDW. Interestingly, we find that such higher-charge superconductors interact nontrivially with certain topological defects appearing in the form of edge dislocations [68] of the underlying lattices, with Burgers vectors compatible with the parent PDW wave vectors, allowing localized single Cooper pairs to re-emerge. Also, such charge-4e superconductivity naturally offers non-trivial pairing symmetry and exchange statistics resembling those of a topological  $p+ip$  superconductor [69–74], yet without the necessity of broken time-reversal symmetry. These discoveries offer smoking-gun signatures and novel characterizing probes for higher-charge superconductivity and its parent PDW phases.

*Many-electron states for higher-charge superconductors.*— For simplicity, we first start with a tight-binding model with a (mean-field) stripe PDW in two dimensions (2D) as our parent Hamiltonian:

$$\hat{H}_{\text{PDW}} = -t \sum_{\langle \mathbf{r}, \mathbf{r}' \rangle, s} c_{\mathbf{r}', s}^\dagger c_{\mathbf{r}, s} + c_{\mathbf{r}, s}^\dagger c_{\mathbf{r}', s} - \mu \sum_{\mathbf{r}, s} c_{\mathbf{r}, s}^\dagger c_{\mathbf{r}, s} + V \sum_{\mathbf{r}} \cos(Qx) c_{\mathbf{r}, \uparrow}^\dagger c_{\mathbf{r}, \downarrow} + \text{h.c.}, \quad (1)$$

where  $s = \uparrow, \downarrow$  labels the electron spin and  $\mathbf{r} = (x, y)$  labels the sites on a  $L \times L$  square lattice under periodic boundary conditions (PBCs), and we set  $L = 200$  unless noted otherwise;  $t = 1$  is the hopping amplitude as well as our unit of energy;  $V$  describes the strength of a stripe PDW with wave vector  $\mathbf{Q} = Q\hat{x}$  along the  $x$ -direction; see illustration in Fig. 1(a). Ideally, our choice of  $Q$  should be incommensurate; however, such  $Q$  would obstruct the PBC and later, the translation symmetrization in numerical models, where we are compelled to employ  $Q = 2\pi q/p$  with a large  $p \sim O(L)$ , whose distinction with an incommensurate  $Q$  becomes negligible for a sufficiently large  $L$ . Through a particle-hole transformation  $a_{\mathbf{r}, \downarrow} = c_{\mathbf{r}, \downarrow}^\dagger$  and diagonalization of  $\hat{H}_{\text{PDW}}$ , we obtain the ground state  $|\Psi\rangle = \prod_{\epsilon_n < 0} \sum_{\mathbf{r}} [\psi_n(\mathbf{r}, \uparrow) c_{\mathbf{r}, \uparrow}^\dagger + \psi_n(\mathbf{r}, \downarrow) a_{\mathbf{r}, \downarrow}^\dagger] |\emptyset\rangle$ , where  $|\emptyset\rangle$  is the vacuum state of  $c_\uparrow$  and  $a_\downarrow$  (full state of  $c_\downarrow$ ). On top of the electron Fermi sea dictated by the dispersion  $\epsilon_{\mathbf{k}} = -2t(\cos k_x + \cos k_y) - \mu$ , the PDW introduces coupling between  $c_{\mathbf{k}, \uparrow}^\dagger$  and  $c_{-\mathbf{k} \mp \mathbf{Q}, \downarrow}^\dagger$  ( $a_{\mathbf{k} \mp \mathbf{Q}, \downarrow}$ ), introducing a momentum transfer of  $\pm \mathbf{Q}$ , respectively; such pairing is most active around selected hot spots on the otherwise sharp Fermi surface, as shown in Fig. 1(b).

Next, we introduce a translation symmetrization  $P_T$ , where  $P_T = 1 + T_x + T_x^2 + \dots + T_x^{L-1}$  and  $T_x$  is the translation operator of one lattice spacing along the  $\hat{x}$  direction. The subsequent projected wavefunction,  $P_T|\Psi\rangle = T_x(P_T|\Psi\rangle)$ , is translation invariant, just as the vestigial order after the (partial) melting of the initial PDW. Interestingly, unlike the parent PDW state  $|\Psi\rangle$  (Fig. 2(b) [75]), the Cooper-pair expectation values,  $\langle c_{\mathbf{r}, \downarrow}^\dagger c_{\mathbf{r}', \uparrow}^\dagger \rangle$ , vanish universally for arbitrary  $\mathbf{r}$  and  $\mathbf{r}'$ , showing that the single




FIG. 1. (a) Illustration of the model in Eq. 1 on a two-dimensional lattice (blue dots). In contrast to the parent state with a stripe PDW (left), the many-electron state following translation symmetrization exhibits vanishing 2e Cooper-pair density  $\Delta$ . (b) A schematic diagram illustration of the Fermi sea and the  $\pm \mathbf{Q}$  momentum transfers induced due to the PDW. The hot spots (purple crosses) are located at the intersections of the original and shifted Fermi surfaces. Correspondingly, the momentum-space electron density  $\langle c_{\mathbf{k}, s}^\dagger c_{\mathbf{k}, s} \rangle$  of the model in Eq. 1 clearly reveals its Fermi sea and the hot spots. Here,  $V = 0.1$ ,  $Q = 1.76$ , and  $\mu = -1.0$ .




FIG. 2. (a) Heat map of the four-point correlation functions  $|\langle c_{\mathbf{k}, x}^\dagger c_{\mathbf{k} + \mathbf{Q}, x}^\dagger c_{-\mathbf{k}, x}^\dagger c_{-\mathbf{k} - \mathbf{Q}, x} \rangle|$  of the charge-4e superconducting state  $P_T|\Psi\rangle$  show a sharp peak at  $(k_x, k'_x) = (Q/2, -Q/2)$ , which corresponds to the hot spots on the Fermi surface in Fig. 1(b); we have fixed  $k_y = k'_y = 1.71$  for simpler display. (b) While the charge-2e Cooper-pair density  $\Delta(\mathbf{k}) = \langle c_{\mathbf{k}}^\dagger c_{-\mathbf{k} + \mathbf{Q}}^\dagger \rangle$  flourishes in the parent PDW state  $|\Psi\rangle$ , (c) it is completely suppressed in the higher-charge superconducting state  $P_T|\Psi\rangle$ .

2e Cooper-pair sector is fully absent in  $P_T|\Psi\rangle$ ; see Fig. 2(c). Instead, the minimum charge difference is 4e - two Cooper pairs, as shown by the finite expectation value of  $\langle c_{\mathbf{k}, \uparrow}^\dagger c_{\mathbf{k} + \mathbf{Q}, \downarrow}^\dagger c_{\mathbf{k}', \uparrow}^\dagger c_{\mathbf{k}' - \mathbf{Q}, \downarrow}^\dagger \rangle$  in Fig. 2(a). We detail our method for evaluating such expectation values in the Supplemental Material [76]. Such a contrast between two-electrons and four-electrons terms already indicates that  $P_T|\Psi\rangle$  deviates significantly from a simple Hartree-Fock state, whose correlations always follow straightforward factorizations of Wick's theorem.

The Gutzwiller projection of spinful electrons offers quantum spin-liquid wavefunctions for variational studies and universal characterizations [51, 54–62, 77–80], by

retaining the physically relevant spinon degrees of freedom while removing the chargon degrees of freedom. Effectively, the projection binds the spin-up electrons and spin-down holes as an auxiliary (gauge) field. Likewise, a stripe PDW consists of two order parameter fields  $\Delta_{\mathbf{Q}}(\mathbf{r}), \Delta_{-\mathbf{Q}}(\mathbf{r}) \in \mathbb{C}$ , corresponding to its  $\pm \mathbf{Q}$  components, whose anti-binding  $\Delta_{\mathbf{Q}}^*(\mathbf{r})\Delta_{-\mathbf{Q}}(\mathbf{r})$  and binding  $\Delta_{\mathbf{Q}}(\mathbf{r})\Delta_{-\mathbf{Q}}(\mathbf{r})$  are associated with the CDW and the charge-4e superconductivity, breaking translation symmetry and charge conservation, respectively - the translation symmetrization projects out the former and retains the latter degrees of freedom. Phenomenologically, the projection  $P_T$  represents the proliferating vortices binding the two order parameter fields as proposed in previous Ginzburg-Landau theories [9, 21–23, 26, 27, 31].

Intuitively, we may understand the above unusual properties of  $P_T|\Psi\rangle$  through the perturbation theory for relatively small  $V$ . The zeroth-order state  $|\Psi_0\rangle$ , the electron Fermi sea of the first line in Eq. 1, is fully translation-symmetric with a zero total momentum, thus remains intact upon  $P_T$ . Following first-order perturbation, the contributions to the state,  $|\Psi_1\rangle \propto V \sum_{\mathbf{k}} (c_{\mathbf{k},\uparrow}^\dagger c_{-\mathbf{k}\pm\mathbf{Q},\downarrow}^\dagger + \text{h.c.}) |\Psi_0\rangle$ , introduce charge-2e Cooper pairs, yet alter the total momentum by  $\pm \mathbf{Q}$ , and subsequently, become projected out by  $P_T$ . It is only until the second order that receives a nontrivial contribution  $|\Psi_2\rangle \propto V^2 \sum_{\mathbf{k},\mathbf{k}'} c_{\mathbf{k},\uparrow}^\dagger c_{-\mathbf{k}\pm\mathbf{Q},\downarrow}^\dagger \cdot c_{\mathbf{k}',\uparrow}^\dagger c_{-\mathbf{k}'\mp\mathbf{Q},\downarrow}^\dagger |\Psi_0\rangle$ , corresponding to sequential scatterings from the PDW with opposing thus mutually canceling momentum transfers, which manages to conserve its total momentum and survive the projection  $P_T$ . Indeed, our numerical results indicate that  $P_T|\Psi\rangle$ 's expectation values of charge-4e Cooper pairs are most significant at momenta  $\pm \mathbf{Q}$ :  $\langle c_{\mathbf{k},\uparrow}^\dagger c_{-\mathbf{k}+\mathbf{Q},\downarrow}^\dagger c_{\mathbf{k}',\uparrow}^\dagger c_{-\mathbf{k}'-\mathbf{Q},\downarrow}^\dagger \rangle$ , with amplitudes quadratic in  $V$ , see Fig. 3(g).

Importantly, while previous Ginzburg-Landau theory analysis is somewhat reliant on point-group symmetries, such as inversion symmetry for charge-4e superconductivity [9, 21, 23, 26, 27],  $C_3$  rotation symmetry for charge-6e superconductivity [22, 31], and so on, our legitimate quantum many-electron states explicitly suggest that momentum conservation alone should be sufficient for their advent. For instance, we may generalize  $P_T|\Psi\rangle$  to cases without inversion symmetry, e.g., either a non-reciprocal PDW  $V_{\mathbf{Q}} \sum_{\mathbf{k}} c_{\mathbf{k},\uparrow}^\dagger c_{-\mathbf{k}+\mathbf{Q},\downarrow}^\dagger + V_{\mathbf{Q}'} \sum_{\mathbf{k}} c_{\mathbf{k},\uparrow}^\dagger c_{-\mathbf{k}+\mathbf{Q}',\downarrow}^\dagger + \text{h.c.}$  ( $\mathbf{Q}' = -\mathbf{Q}$ ), or more fundamentally, an asymmetric Fermi sea  $|\Psi_0\rangle$ . Note that in the latter case, the translation symmetrization should be adjusted as  $P_T = 1 + T_x e^{iK} + T_x^2 e^{i2K} + \dots + T_x^{L-1} e^{iK(L-1)}$  for a Fermi sea with a finite total momentum  $\mathbf{K}$ ; for simplicity, here we consider  $\mathbf{K} = K\hat{x}$ . Consistent with the aforementioned suppression of charge-2e sectors, Figs. 3(a)(b) demonstrate that a finite charge-4e signal is clearly evident in both scenarios. Further, the charge-4e amplitude at  $\langle c_{\mathbf{k},\uparrow}^\dagger c_{-\mathbf{k}+\mathbf{Q},\downarrow}^\dagger c_{\mathbf{k}',\uparrow}^\dagger c_{-\mathbf{k}'-\mathbf{Q},\downarrow}^\dagger \rangle$ , which we dub the charge-4e order parameter  $V_{4e}$  and summarize in Fig. 3(g), shows a clear linear dependence on both  $V_{\mathbf{Q}}$  and  $V_{-\mathbf{Q}}$ :  $V_{4e} \propto$




FIG. 3. The correlations  $|\langle c_{\mathbf{k}}^\dagger c_{-\mathbf{k}+\mathbf{Q}}^\dagger c_{\mathbf{k}'}^\dagger c_{-\mathbf{k}'-\mathbf{Q}}^\dagger \rangle|$  of the  $P_T|\Psi\rangle$  states with (a) asymmetric pairing amplitude  $V_{\mathbf{Q}} \neq V_{-\mathbf{Q}}$ , (b) asymmetric Fermi sea with non-zero total momentum  $\mathbf{K}$ , and (c) nonreciprocal PDW momentum  $\mathbf{Q}' \neq -\mathbf{Q}$ , suggests that the charge-4e superconductivity may exists as long as momentum conserves despite broken inversion symmetry. (d) The electron density  $\langle c_{\mathbf{k},s}^\dagger c_{\mathbf{k},s} \rangle$  of a charge-6e superconductor displays hot spots on its Fermi surface. The corresponding  $P_T|\Psi\rangle$  exhibits vanishing charge-2e and (e) charge-4e expectation values, yet (f) a sharp signal in charge-6e correlations (we have fixed  $k_y = k'_y = k''_y = 2.0$ ). (g) The charge-4e order parameter  $V_{4e}$  scales proportionally to both  $V_{\mathbf{Q}}$  and  $V_{-\mathbf{Q}}$ ; (h) similarly, the charge-6e order parameter scales tri-linearly to each of the three parent PDWs' amplitudes. Here, we set  $L = 1000$ .

$V_{\mathbf{Q}} V_{-\mathbf{Q}}$ , consistent with our perturbative analysis and irrespective of broken inversion symmetry. On the contrary, all charge-4e expectation values  $\langle c_{\mathbf{r},\uparrow}^\dagger c_{\mathbf{r}',\downarrow}^\dagger c_{\mathbf{r}'',\uparrow}^\dagger c_{\mathbf{r}'',\downarrow}^\dagger \rangle$  vanish in  $P_T|\Psi\rangle$  in the absence of momentum conservation, e.g., the momentum components  $\mathbf{Q}' \neq -\mathbf{Q}$  within the PDW, as we summarize in Figs. 3(c)(g). These results and conclusions also extend to relatively large values of  $V_{\pm\mathbf{Q}}$ , as we discuss further in the Supplemental Material [76].

Further, we can generalize such a construction to higher-charge superconductivity, e.g., charge-6e Cooper pairs with three PDW momentum components,  $\mathbf{Q}_1$ ,  $\mathbf{Q}_2$ , and  $\mathbf{Q}_3$ , which conserves total momentum  $\mathbf{Q}_1 + \mathbf{Q}_2 + \mathbf{Q}_3 = 0$ . For instance, we employ the following PDW term on top of the Fermi sea (first line) in Eq. 1:

$$H_{\text{PDW}} = \sum_{\mathbf{r}} V(\mathbf{r}) c_{\mathbf{r},\uparrow}^\dagger c_{\mathbf{r},\downarrow}^\dagger + \text{h.c.},$$

$$V(\mathbf{r}) = V_{\mathbf{Q}} e^{iQx} + V_{-\mathbf{Q}/2} e^{-iQx/2}. \quad (2)$$




FIG. 4. (a) An illustration of our interferometer setup  $P_T|\Psi(\phi)\rangle$ : translation-symmetrizing the parent PDW model in Eq. 1 with a PDW in the  $x$ -direction and a magnetic flux  $\phi$  in the  $y$ -direction. (b) Unlike the parent PDW system  $|\Psi(\phi)\rangle$  with a  $\pi$  periodicity in  $\phi$  corresponding to conventional charge-2e Cooper pairs, the susceptibility  $\mathcal{L}(\phi)$  in Eq. 3 of  $P_T|\Psi(\phi)\rangle$  exhibits an unmistakable charge-4e signature:  $\pi/2$  period with respect to the threaded magnetic flux  $\phi$ . Here, our system size is  $L = 1000$ .

Following translation symmetrization, the resulting state  $P_T|\Psi\rangle$  witnesses vanishing charge-2e and charge-4e correlations, yet a clear charge-6e pairing signature ( $\langle c_{\mathbf{k},\uparrow}^\dagger c_{-\mathbf{k}+\mathbf{Q},\downarrow}^\dagger c_{\mathbf{k}',\uparrow}^\dagger c_{-\mathbf{k}',-\mathbf{Q}/2,\downarrow}^\dagger c_{\mathbf{k}'',\uparrow}^\dagger c_{-\mathbf{k}''-\mathbf{Q}/2,\downarrow}^\dagger \rangle$ ), which scales as  $V^3$ , or proportionally to the amplitude of each of the participating PDWs. These results are presented in Figs. 3(d)-(f)(h). We note that potential signatures on sequential or coexisting charge-4e and charge-6e orderings observed in  $\text{CsV}_3\text{Sb}_5$  materials [36] are consistent with the previously hosting PDW patterns [18, 19], which conserve the total momentum under both charge-4e and charge-6e mechanisms. Generalizations to higher dimensions or other PDW scenarios, such as the checkerboard order, are straightforward, albeit a corresponding translation symmetrization  $P_T$  may need to be introduced.

*Phenomena of higher-charge superconductivity.*— Our microscopic many-electron models of higher-charge superconductors have opened new opportunities to understand and discover their novel properties. For instance, we may probe their exotic magnetic-flux quantization by measuring the periodicity of their physical properties in an interferometer geometry. In particular, we may introduce to the parent PDW model in Eq. 1 a magnetic flux along the  $y$ -direction through a phase  $\phi$  across the boundary along the  $x$ -direction; see our setup geometry in Fig. 4(a). We neglect the Zeeman effect on the electron spins and the magnetic flux vortices for simplicity.

After the translation symmetrization, fully compatible with the magnetic flux, we evaluate the susceptibility of the charge-4e Cooper pair:

$$\mathcal{L}(\phi) = \frac{\partial \langle c_{\mathbf{k},\uparrow}^\dagger c_{-\mathbf{k}+\mathbf{Q},\downarrow}^\dagger c_{\mathbf{k}',\uparrow}^\dagger c_{-\mathbf{k}',-\mathbf{Q},\downarrow}^\dagger \rangle|_{P_T|\Psi(\phi)\rangle}}{\partial V}, \quad (3)$$

for the subsequent many-electron state  $P_T|\Psi(\phi)\rangle$ , which we summarize in Fig. 4(b). Indeed,  $\mathcal{L}(\phi)$  exhibits clear,




FIG. 5. (a) Charge-2e Cooper pairs re-emerge locally at topological defects of the underlying lattice when the Burgers vector  $\mathbf{B}$  aligns with the parent PDW wave vector  $\pm\mathbf{Q}$ , yet (b) no such modes are visible when  $\mathbf{B}$  and  $\mathbf{Q}$  are vertical. The upper panels show schematic illustrations of the lattices with a pair of topological defects and the PDWs before translation symmetrization  $P_T$ . The dashed lines illustrate the local hopping connectivity, which becomes 1-to-2 on the edge-dislocation sites. The lower panels are the charge-2e expectation values  $\langle c_{\mathbf{r},\uparrow}^\dagger c_{\mathbf{r},\downarrow}^\dagger \rangle$  for such  $P_T|\Psi\rangle$  with various distances  $l$  between the two topological defects. Here, we have  $L = 50$ .

distinctive  $\pi/2$  periodicity in magnetic fluxes and offers strong evidence for its charge-4e fundamentals. We attribute the slight deviations from a perfect 4e-period to residue contributions of the electrons, especially the remaining Fermi surface, and the finite-size effects; please refer to the Supplemental Material for additional results and generalizations [76]. This is in sharp contrast with its parent PDW state, where the  $\pi$  magnetic-flux period plays a dominant role in the pairing susceptibility contributed by charge-2e Cooper pairs.

In addition, our formalism may reveal new features of higher-charge superconducting states: while the charge-2e Cooper-pair channel is initially gapped out, localized Cooper pairs may re-emerge at certain topological defects, as long as their Burgers vector  $\mathbf{B}$  is compatible, i.e., parallel or anti-parallel, with the parent PDW wave vector  $\mathbf{Q}$ . Intuitively, a topological defect may serve as a source or drain of momentum along its Burgers vector [76], thus assisting charge-2e Cooper pairs originating from single PDW scatterings to conserve momentum and exist locally. We note that a topological defect and its Burgers vector - essentially a topological invariant - remains robust upon perturbations; likewise, such localized Cooper pairs also remain relatively stable. When and only when two topological defects with opposing Burgers vectors encounter and annihilate, their respective local charge-2e Cooper pairs with  $\pm\mathbf{Q}$  momentum merge into the charge-4e Cooper pair condensate.

To demonstrate such localized charge-2e Cooper pairs, we introduce a pair of edge dislocations with opposing Burgers vectors  $\pm\mathbf{B}$  into the 2D lattice hosting the parent Hamiltonian in Eq. 1, see Fig. 5 upper panels. In particular, we insert  $l$  additional lattice sites in the bulk, thus separating the two edge dislocations by a distance  $l$ . Except in the vicinity of the topological defects, the system remains effectively translation-invariant, and we can generalize our translation symmetrization  $P_T$  accordingly, see further details in Ref. [76].

As a result, Fig. 5 lower panels display the charge-2e Cooper pair density  $\langle c_{\mathbf{r},\uparrow}^\dagger c_{\mathbf{r},\downarrow}^\dagger \rangle$  for the many-electron state  $P_T|\Psi\rangle$ . When the Burgers vector  $\mathbf{B}$  is perpendicular to the PDW wave vector  $\mathbf{Q}$ , charge-2e Cooper pairs remain fully absent. In contrast, when  $\mathbf{B}$  is parallel to  $\mathbf{Q}$ , clear charge-2e pairing signals emerge at the dislocations, while everywhere else in the lattice remains oblivious to the topological defects and minimally charge-4e. As we gradually decrease the separation  $l$  between the two dislocations, the localized charge-2e Cooper pairs approach and eventually annihilate each other, restoring a charge-4e superconducting state without topological defects. These unusual localized-Cooper-pair phenomena are fully consistent with our theoretical expectations, and offer smoking-gun signatures for identifying charge-4e superconductivity as well as parent PDW in candidate materials, e.g.,  $\text{CsV}_3\text{Sb}_5$ , via scanning tunneling spectroscopy experiments.

*Discussion.*— We have established many-electron characterizations of higher-charge superconductivity by translation-symmetrizing parent PDW systems, consistent with the Ginzburg-Landau theory of vestigial orders. In particular, with simple stripe PDWs, we clearly demonstrate charge-4e and charge-6e superconductivity, with vanishing charge-2e Cooper-pair sectors. Interestingly, momentum conservation, rather than stricter point-group symmetries, is sufficient for their establish-

ment. The obtained higher-charge superconductors exhibit consistent magnetic flux quantization and periodicity in their interferometry. We also show that charge-2e Cooper pairs may re-emerge at topological defects of the underlying lattice, if their Burgers vector and the parent PDW wave vector are compatible. In addition, unrestricted by the presence of time reversal symmetry, the spin-up (or spin-down) electrons of the above charge-4e superconductors offer a pairing and exchange symmetry that resembles a nodeless  $p+ip$ -wave superconductor, and thus offer a novel arena for nontrivial topological phenomena and applications, as we discuss further in the Supplemental Material [76]. Overall, our formalism offers unprecedented opportunities to understand and identify these exotic quantum matters.

Finally, generalizations of the current formalism to higher dimensions and other density-wave scenarios are straightforward. Interestingly, applications to spin density waves and charge density waves may offer a novel perspective and method for the physics of intertwined orders [81–84] in strongly correlated systems [85, 86]. There, in addition to the universal behaviors of the superconducting phases that we have mainly focused on, details such as the parent density wave’s pattern, strength, and wave vector may impact their overall energetic competitiveness. Besides, as in the topological theories of gapped quantum spin liquids, it will be crucial to investigate the physics of such Cooper-pair-binding vortices, based on particular Fermi seas entirely gapped out by the parent density waves and thus clean from residual contributions.

*Acknowledgment:* We acknowledge Fa Wang, Jian Wang, and Hong-Hao Tu for helpful discussions and support from the National Key R&D Program of China (Grant No.2021YFA1401900 & No.2022YFA1403700), the National Natural Science Foundation of China (Grants No.12174008 & No.92270102), and Shanghai Municipal Science and Technology Project (Grant No.25LZ2601100).

---

[1] K. Onnes, The resistance of pure mercury at helium temperatures, *Commun. Phys. Lab. Univ. Leiden*, b **120** (1911).

[2] W. Meissner and R. Ochsenfeld, Ein neuer effekt bei eintritt der supraleitfähigkeit, *Naturwissenschaften* **21**, 787 (1933).

[3] D. Van Delft and P. Kes, The discovery of superconductivity, *Physics today* **63**, 38 (2010).

[4] V. L. Ginzburg and L. D. Landau, On the theory of superconductivity, in *On superconductivity and superfluidity: a scientific autobiography* (Springer, 2009) pp. 113–137.

[5] L. N. Cooper, Bound electron pairs in a degenerate fermi gas, *Physical Review* **104**, 1189 (1956).

[6] J. Bardeen, L. N. Cooper, and J. R. Schrieffer, Theory of superconductivity, *Physical review* **108**, 1175 (1957).

[7] P. Fulde and R. A. Ferrell, Superconductivity in a strong spin-exchange field, *Physical Review* **135**, A550 (1964).

[8] A. Larkin and Y. N. Ovchinnikov, Nonuniform state of superconductors, *Soviet Physics-JETP* **20**, 762 (1965).

[9] D. Agterberg and H. Tsunetsugu, Dislocations and vortices in pair-density-wave superconductors, *Nature Physics* **4**, 639 (2008).

[10] E. Berg, E. Fradkin, S. A. Kivelson, and J. M. Tranquada, Striped superconductors: how spin, charge and superconducting orders intertwine in the cuprates, *New Journal of Physics* **11**, 115004 (2009).

[11] P. A. Lee, Amperean pairing and the pseudogap phase of cuprate superconductors, *Physical Review X* **4**, 031017 (2014).

[12] E. Fradkin, S. A. Kivelson, and J. M. Tranquada, Colloquium: Theory of intertwined orders in high temperature superconductors, *Reviews of Modern Physics* **87**, 457 (2015).

[13] D. F. Agterberg, J. S. Davis, S. D. Edkins, E. Fradkin, D. J. Van Harlingen, S. A. Kivelson, P. A. Lee, L. Radzihovsky, J. M. Tranquada, and Y. Wang, The physics

of pair-density waves: cuprate superconductors and beyond, *Annual Review of Condensed Matter Physics* **11**, 231 (2020).

[14] Y.-M. Wu, P. A. Nosov, A. A. Patel, and S. Raghu, Pair density wave order from electron repulsion, *Physical Review Letters* **130**, 026001 (2023).

[15] N. S. Ticea, S. Raghu, and Y.-M. Wu, Pair density wave order in multiband systems, *Physical Review B* **110**, 094515 (2024).

[16] M. Hamidian, S. D. Edkins, S. H. Joo, A. Kostin, H. Eisaki, S. Uchida, M. Lawler, E.-A. Kim, A. P. Mackenzie, K. Fujita, *et al.*, Detection of a cooper-pair density wave in  $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_8 + x$ , *Nature* **532**, 343 (2016).

[17] X. Liu, Y. X. Chong, R. Sharma, and J. S. Davis, Discovery of a cooper-pair density wave state in a transition-metal dichalcogenide, *Science* **372**, 1447 (2021).

[18] H. Chen, H. Yang, B. Hu, Z. Zhao, J. Yuan, Y. Xing, G. Qian, Z. Huang, G. Li, Y. Ye, *et al.*, Roton pair density wave in a strong-coupling kagome superconductor, *Nature* **599**, 222 (2021).

[19] H. Zhao, H. Li, B. R. Ortiz, S. M. Teicher, T. Park, M. Ye, Z. Wang, L. Balents, S. D. Wilson, and I. Zeljkovic, Cascade of correlated electron states in the kagome superconductor  $\text{CsV}_3\text{Sb}_5$ , *Nature* **599**, 216 (2021).

[20] Q. Gu, J. P. Carroll, S. Wang, S. Ran, C. Broyles, H. Siddiquee, N. P. Butch, S. R. Saha, J. Paglione, J. S. Davis, *et al.*, Detection of a pair density wave state in  $\text{UTe}_2$ , *Nature* **618**, 921 (2023).

[21] E. Berg, E. Fradkin, and S. A. Kivelson, Charge-4e superconductivity from pair-density-wave order in certain high-temperature superconductors, *Nature Physics* **5**, 830 (2009).

[22] D. Agterberg, M. Geracie, and H. Tsunetsugu, Conventional and charge-six superfluids from melting hexagonal fulde-ferrell-larkin-ovchinnikov phases in two dimensions, *Physical Review B—Condensed Matter and Materials Physics* **84**, 014513 (2011).

[23] S.-K. Jian, Y. Huang, and H. Yao, Charge-4e superconductivity from nematic superconductors in two and three dimensions, *Physical review letters* **127**, 227001 (2021).

[24] L. Nie, G. Tarjus, and S. A. Kivelson, Quenched disorder and vestigial nematicity in the pseudogap regime of the cuprates, *Proceedings of the National Academy of Sciences* **111**, 7980 (2014).

[25] Y.-B. Liu, J. Zhou, and F. Yang, Nematic superconductivity and its critical vestigial phases in the quasicrystal, *Physical Review Letters* **133**, 136002 (2024).

[26] E. V. Herland, E. Babaev, and A. Sudbø, Phase transitions in a three dimensional  $\text{U}(1) \times \text{U}(1)$  lattice london superconductor: Metallic superfluid and charge-4e superconducting states, *Physical Review B—Condensed Matter and Materials Physics* **82**, 134511 (2010).

[27] R. M. Fernandes and L. Fu, Charge-4e superconductivity from multicomponent nematic pairing: Application to twisted bilayer graphene, *Physical review letters* **127**, 047001 (2021).

[28] J. H. Han and P. A. Lee, Understanding resistance oscillation in the  $\text{CsV}_3\text{Sb}_5$  superconductor, *Physical Review B* **106**, 184515 (2022).

[29] C. M. Varma and Z. Wang, Extended superconducting fluctuation region and 6e and 4e flux quantization in a kagome compound with a normal state of 3Q order, *Physical Review B* **108**, 214516 (2023).

[30] L.-F. Zhang, Z. Wang, and X. Hu, Higgs-leggett mechanism for the elusive  $\phi_0/3 = hc/6e$  oscillation in little-parks setup of kagome superconductor  $\text{CsV}_3\text{Sb}_5$ , *Communications Physics* **7**, 210 (2024).

[31] T.-Y. Lin, F.-F. Song, and G.-M. Zhang, Theory of the charge-6e condensed phase in kagome-lattice superconductors, *Physical Review B* **111**, 054508 (2025).

[32] V. B. Geshkenbein, A. I. Larkin, and A. Barone, Vortices with half magnetic flux quanta in “heavy-fermion”superconductors, *Physical Review B* **36**, 235 (1987).

[33] M. Sigrist, T. Rice, and K. Ueda, Low-field magnetic response of complex superconductors, *Physical review letters* **63**, 1727 (1989).

[34] H.-Y. Kee, Y. B. Kim, and K. Maki, Half-quantum vortex and d-soliton in  $\text{Sr}_2\text{RuO}_4$ , *arXiv preprint cond-mat/0005510* (2000).

[35] M. A. Rampp and J. Schmalian, Integer and fraction- alized vortex lattices and off-diagonal long-range order, *Journal of Physics Communications* **6**, 055013 (2022).

[36] J. Ge, P. Wang, Y. Xing, Q. Yin, A. Wang, J. Shen, H. Lei, Z. Wang, and J. Wang, Charge-4e and charge-6e flux quantization and higher charge superconductivity in kagome superconductor ring devices, *Physical Review X* **14**, 021025 (2024).

[37] Y.-F. Jiang, Z.-X. Li, S. A. Kivelson, and H. Yao, Charge-4e superconductors: A majorana quantum monte carlo study, *Physical Review B* **95**, 241103 (2017).

[38] N. V. Gnezdilov and Y. Wang, Solvable model for a charge-4e superconductor, *Physical Review B* **106**, 094508 (2022).

[39] M. O. Soldini, M. H. Fischer, and T. Neupert, Charge-4e superconductivity in a hubbard model, *Physical Review B* **109**, 214509 (2024).

[40] P. Li, K. Jiang, and J. Hu, Charge 4e superconductor: A wavefunction approach, *Science Bulletin* **69**, 2328 (2024).

[41] Y.-B. Liu, J. Zhou, C. Wu, and F. Yang, Charge-4e superconductivity and chiral metal in 45-twisted bilayer cuprates and related bilayers, *Nature Communications* **14**, 7926 (2023).

[42] Y.-M. Wu and Y. Wang, d-wave charge-4e superconductivity from fluctuating pair density waves, *npj Quantum Materials* **9**, 66 (2024).

[43] F.-F. Song and G.-M. Zhang, Phase coherence of charge-6e superconductors with a frustrated kagome xy antiferromagnet, *Chinese Physics Letters* **42**, 037401 (2025).

[44] S. Zhou and Z. Wang, Chern fermi pocket, topological pair density wave, and charge-4e and charge-6e superconductivity in kagomé superconductors, *Nature Communications* **13**, 7288 (2022).

[45] Z. Pan, C. Lu, F. Yang, and C. Wu, Frustrated superconductivity and sextetting order, *Science China Physics, Mechanics & Astronomy* **67**, 287412 (2024).

[46] M. Zeng, L.-H. Hu, H.-Y. Hu, Y.-Z. You, and C. Wu, High-order time-reversal symmetry breaking normal state, *Science China Physics, Mechanics & Astronomy* **67**, 237411 (2024).

[47] R. B. Laughlin, Anomalous quantum hall effect: An incompressible quantum fluid with fractionally charged excitations, *Physical Review Letters* **50**, 1395 (1983).

[48] D. Arovas, J. R. Schrieffer, and F. Wilczek, Fractional statistics and the quantum hall effect, *Physical review letters* **53**, 722 (1984).

[49] J. K. Jain, Composite-fermion approach for the fractional quantum hall effect, *Physical review letters* **63**, 199 (1989).

[50] M. C. Gutzwiller, Effect of correlation on the ferromagnetism of transition metals, *Physical Review Letters* **10**, 159 (1963).

[51] X.-G. Wen, Mean-field theory of spin-liquid states with finite energy gap and topological orders, *Physical Review B* **44**, 2664 (1991).

[52] P. A. Lee, N. Nagaosa, and X.-G. Wen, Doping a mott insulator: Physics of high-temperature superconductivity, *Reviews of modern physics* **78**, 17 (2006).

[53] Y. Zhou, K. Kanoda, and T.-K. Ng, Quantum spin liquid states, *Reviews of Modern Physics* **89**, 025003 (2017).

[54] H. Yokoyama and H. Shiba, Variational monte-carlo studies of hubbard model. I, *Journal of the Physical Society of Japan* **56**, 1490 (1987).

[55] H. Yokoyama and H. Shiba, Variational monte-carlo studies of hubbard model. II, *Journal of the Physical Society of Japan* **56**, 3582 (1987).

[56] C. Gros, Physics of projected wavefunctions, *Annals of Physics* **189**, 53 (1989).

[57] A. Paramekanti, M. Randeria, and N. Trivedi, Projected wave functions and high temperature superconductivity, *Physical review letters* **87**, 217002 (2001).

[58] A. Himeda, T. Kato, and M. Ogata, Stripe states with spatially oscillating d-wave superconductivity in the two-dimensional  $t - t' - J$  model, *Physical review letters* **88**, 117001 (2002).

[59] O. I. Motrunich, Variational study of triangular lattice spin-1/2 model with ring exchanges and spin liquid state in  $\kappa$ -(ET)<sub>2</sub>Cu<sub>2</sub>(CN)<sub>3</sub>, *Physical Review B—Condensed Matter and Materials Physics* **72**, 045105 (2005).

[60] B. Edegger, V. N. Muthukumar, and C. Gros, Gutzwiller-rvb theory of high-temperature superconductivity: Results from renormalized mean-field theory and variational monte carlo calculations, *Advances in Physics* **56**, 927 (2007).

[61] Y. Iqbal, F. Becca, and D. Poilblanc, Projected wave function study of Z2 spin liquids on the kagome lattice for the spin-1/2 quantum heisenberg antiferromagnet, *Physical Review B—Condensed Matter and Materials Physics* **84**, 020407 (2011).

[62] Y. Iqbal, F. Becca, S. Sorella, and D. Poilblanc, Gapless spin-liquid phase in the kagome spin-1/2 heisenberg antiferromagnet, *Physical Review B—Condensed Matter and Materials Physics* **87**, 060405 (2013).

[63] C. A. Jiménez-Hoyos, T. M. Henderson, T. Tsuchimochi, and G. E. Scuseria, Projected hartree–fock theory, *The Journal of chemical physics* **136** (2012).

[64] T. Nikšić, D. Vretenar, and P. Ring, Beyond the relativistic mean-field approximation. II. configuration mixing of mean-field wave functions projected on angular momentum and particle number, *Physical Review C—Nuclear Physics* **74**, 064309 (2006).

[65] T. Sogo, R. Lazauskas, G. Röpke, and P. Schuck, Critical temperature for *alpha*-particle condensation within a momentum-projected mean-field approach, *Physical Review C* **79**, 051301 (2009).

[66] T. Mizusaki and M. Imada, Quantum-number projection in the path-integral renormalization group method, *Physical Review B* **69**, 125110 (2004).

[67] T. Heitmann and J. Schnack, Combined use of translational and spin-rotational invariance for spin systems, *Physical Review B* **99**, 134405 (2019).

[68] N. D. Mermin, The topological theory of defects in ordered media, *Reviews of Modern Physics* **51**, 591 (1979).

[69] T. Rice and M. Sigrist, Sr<sub>2</sub>RuO<sub>4</sub>: an electronic analogue of 3he?, *Journal of Physics: Condensed Matter* **7**, L643 (1995).

[70] N. Read and D. Green, Paired states of fermions in two dimensions with breaking of parity and time-reversal symmetries and the fractional quantum hall effect, *Physical Review B* **61**, 10267 (2000).

[71] A. P. Mackenzie and Y. Maeno, The superconductivity of Sr<sub>2</sub>RuO<sub>4</sub> and the physics of spin-triplet pairing, *Reviews of Modern Physics* **75**, 657 (2003).

[72] G. M. Luke, Y. Fudamoto, K. Kojima, M. Larkin, J. Mermin, B. Nachumi, Y. Uemura, Y. Maeno, Z. Mao, Y. Mori, *et al.*, Time-reversal symmetry-breaking superconductivity in Sr<sub>2</sub>RuO<sub>4</sub>, *Nature* **394**, 558 (1998).

[73] K. Nelson, Z. Mao, Y. Maeno, and Y. Liu, Odd-parity superconductivity in Sr<sub>2</sub>RuO<sub>4</sub>, *Science* **306**, 1151 (2004).

[74] C. Kallin and J. Berlinsky, Chiral superconductors, *Reports on Progress in Physics* **79**, 054502 (2016).

[75] We note that the PDW has a finite charge-2e Cooper-pair density in most places, yet the average of states is different from the average of expectation values.

[76] Please refer to the supplemental material for further details.

[77] Y. Zhang, T. Grover, and A. Vishwanath, Entanglement entropy of critical spin liquids, *Physical review letters* **107**, 067202 (2011).

[78] Y. Zhang, T. Grover, A. Turner, M. Oshikawa, and A. Vishwanath, Quasiparticle statistics and braiding from ground-state entanglement, *Physical Review B—Condensed Matter and Materials Physics* **85**, 235151 (2012).

[79] T. Grover, Y. Zhang, and A. Vishwanath, Entanglement entropy as a portal to the physics of quantum spin liquids, *New Journal of Physics* **15**, 025002 (2013).

[80] Y. Zhang and A. Vishwanath, Establishing non-abelian topological order in gutzwiller-projected chern insulators via entanglement entropy and modular s-matrix, *Physical Review B—Condensed Matter and Materials Physics* **87**, 161113 (2013).

[81] J. S. Davis and D.-H. Lee, Concepts relating magnetic interactions, intertwined electronic orders, and strongly correlated superconductivity, *Proceedings of the National Academy of Sciences* **110**, 17623 (2013).

[82] Y. Wang and A. Chubukov, Charge-density-wave order with momentum  $(2Q, 0)$  and  $(0, 2Q)$  within the spin-fermion model: Continuous and discrete symmetry breaking, preemptive composite order, and relation to pseudogap in hole-doped cuprates, *Physical Review B* **90**, 035149 (2014).

[83] B. Keimer, S. A. Kivelson, M. R. Norman, S. Uchida, and J. Zaanen, From quantum matter to high-temperature superconductivity in copper oxides, *Nature* **518**, 179 (2015).

[84] R. M. Fernandes, P. P. Orth, and J. Schmalian, Intertwined vestigial order in quantum materials: Nematicity and beyond, *Annual Review of Condensed Matter Physics* **10**, 133 (2019).

[85] G. Grüner, The dynamics of charge-density waves, *Reviews of modern physics* **60**, 1129 (1988).

[86] G. Grüner, The dynamics of spin-density waves, *Reviews of modern physics* **66**, 1 (1994).