

A REGULARITY THEORY FOR SECOND-ORDER PARABOLIC PARTIAL DIFFERENTIAL EQUATIONS IN WEIGHTED MIXED NORM SOBOLEV-ZYGMUND SPACES

JAE-HWAN CHOI AND JUNHEE RYU

ABSTRACT. We develop an optimal regularity theory for parabolic partial differential equations in weighted mixed norm Sobolev-Zygmund spaces. The results extend the classical Schauder estimates to coefficients that are merely measurable in time and to the critical case of integer-order regularity. In addition, nonzero initial data are treated in the optimal trace space via a sharp trace theorem.

1. INTRODUCTION AND MAIN RESULTS

1.1. Goal and setting. In this paper, we study the existence, uniqueness, and regularity of solutions to the second-order parabolic partial differential equation (PDE)

$$\partial_t u(t, x) = \sum_{i,j=1}^d a_{ij}(t, x) D_{x^i x^j} u(t, x) + \sum_{i=1}^d b_i(t, x) D_{x^i} u(t, x) + c(t, x) u(t, x) + f(t, x), \quad (1.1)$$

posed on $(0, T) \times \mathbb{R}^d$, with non-zero initial data. Our framework is based on the weighted mixed norm Sobolev-Zygmund space $L_p((0, T), w dt; \Lambda^\gamma(\mathbb{R}^d))$, where $\Lambda^\gamma(\mathbb{R}^d)$ is the Hölder-Zygmund space of order $\gamma \in (0, \infty)$ and the weight w belongs to the Muckenhoupt class $A_p(\mathbb{R})$. For $p \in (1, \infty]$, the mixed norm is given by

$$\|u\|_{L_p((0, T), w dt; \Lambda^\gamma(\mathbb{R}^d))} := \begin{cases} \left(\int_0^T \|u(t, \cdot)\|_{\Lambda^\gamma(\mathbb{R}^d)}^p w(t) dt \right)^{1/p}, & p \in (1, \infty), \\ \operatorname{ess\,sup}_{t \in (0, T)} \|u(t, \cdot)\|_{\Lambda^\gamma(\mathbb{R}^d)}, & p = \infty. \end{cases}$$

We begin by recalling the definition of Hölder-Zygmund spaces and Muckenhoupt's weight class.

Definition 1.1 (Hölder-Zygmund space). Let $\gamma \in (0, \infty)$ and $p \in [1, \infty]$. The space $\Lambda_p^\gamma(\mathbb{R}^d)$ is defined by

$$\Lambda_p^\gamma(\mathbb{R}^d) := \{f \in L_\infty(\mathbb{R}^d) : \|f\|_{\Lambda_p^\gamma(\mathbb{R}^d)} < \infty\}.$$

Here the norm $\|f\|_{\Lambda_p^\gamma(\mathbb{R}^d)}$ is given by

$$\|f\|_{\Lambda_p^\gamma(\mathbb{R}^d)} := \|f\|_{L_\infty(\mathbb{R}^d)} + \|f\|_{\dot{\Lambda}_p^\gamma(\mathbb{R}^d)},$$

where

$$\|f\|_{\dot{\Lambda}_p^\gamma(\mathbb{R}^d)} := \begin{cases} \left(\int_{\mathbb{R}^d} \|\mathcal{D}_h^{[\gamma]} f\|_{L_\infty(\mathbb{R}^d)}^p \frac{dh}{|h|^{d+p\gamma}} \right)^{1/p}, & p \in [1, \infty), \\ \sup_{h \in \mathbb{R}^d} \frac{\|\mathcal{D}_h^{[\gamma]} f\|_{L_\infty(\mathbb{R}^d)}}{|h|^\gamma}, & p = \infty. \end{cases}$$

2020 *Mathematics Subject Classification.* 35K15, 35B65, 26A16, 35D35.

Key words and phrases. Parabolic equations with variable coefficients, Mixed norm Sobolev-Zygmund estimates, Partial Schauder theory.

J.-H. Choi was supported by a KIAS Individual Grant (MG102701) at Korea Institute for Advanced Study.

J. Ryu was supported by a KIAS Individual Grant (MG101501) at Korea Institute for Advanced Study.

Here $\mathcal{D}_h f(x) := f(x + h) - f(x)$, $\mathcal{D}_h^n f(x) = \mathcal{D}_h^{n-1}(\mathcal{D}_h f)(x)$, and $[\gamma]^-$ is the smallest integer strictly larger than γ . We denote $\Lambda^\gamma(\mathbb{R}^d) := \Lambda_\infty^\gamma(\mathbb{R}^d)$ and $\dot{\Lambda}^\gamma(\mathbb{R}^d) := \dot{\Lambda}_\infty^\gamma(\mathbb{R}^d)$.

Definition 1.2 (Muckenhoupt's class). For $p \in (1, \infty)$, let $A_p = A_p(\mathbb{R})$ be the class of all nonnegative and locally integrable functions $w : \mathbb{R} \rightarrow [0, \infty)$ satisfying

$$[w]_{A_p} := \sup_{r>0, t \in \mathbb{R}} \left(\int_{t-r}^{t+r} w(s) ds \right) \times \left(\int_{t-r}^{t+r} w(s)^{-\frac{1}{p-1}} ds \right)^{p-1} < \infty.$$

1.2. Motivation and backgrounds. The regularity theory for linear parabolic equations of the form (1.1) has been extensively studied. A classical cornerstone is the *Schauder theory* which roughly states that

$$a_{ij}, b_i, c, f \in C_{t,x}^{\alpha/2, \alpha}, \quad \alpha \in (0, 1) \implies u \in C_{t,x}^{1+\alpha/2, 2+\alpha}.$$

However, the classical Schauder theory faces two fundamental limitations:

1. it does not apply when the coefficients a_{ij}, b_i , and c are merely *measurable in time*;
2. it breaks down at the *critical index* $\alpha = 1$.

To overcome the first limitation, Brandt [5] introduced the *partial Schauder theory*, in which regularity is measured in mixed norm spaces $L_\infty(\mathbb{R}, dt; C^\alpha(\mathbb{R}^d))$, thereby accommodating time-measurable coefficients while preserving spatial Hölder continuity. Lorenzi [20] and Krylov [17, 18] subsequently generalized Brandt's approach to the mixed norm spaces $L_p(\mathbb{R}, dt; C^\alpha(\mathbb{R}^d))$ for $p \in (1, \infty]$. Following their work, several further extensions have been developed. Boccia and Krylov [4] extended the theory to higher-order parabolic systems. Stinga and Torrea [22] proved weighted Sobolev-Schauder estimates for second-order parabolic equations in the mixed norm setting

$$L_p(\mathbb{R}, w(t) dt; C^\alpha(\mathbb{R}^d)), \quad w \in A_p(\mathbb{R}).$$

Despite these advances, classical Hölder spaces remain inadequate in the critical case of *integer* regularity $\alpha = 1$, where the Schauder estimate is known to fail; see, for instance, [14, Chapter 2.2].

A natural way to address the second limitation is to work within the Hölder-Zygmund spaces Λ^γ . These spaces coincide with Hölder spaces for non-integer orders, $\Lambda^\gamma = C^\gamma$ for $\gamma \notin \mathbb{N}$, but become strictly larger for integer orders (see Proposition 2.2):

$$C^\gamma \subsetneq \Lambda^\gamma, \quad \gamma \in \mathbb{N}.$$

Within this framework, Kim [15] established regularity estimates in $L_p((0, T), dt; \Lambda^\alpha(\mathbb{R}^d))$, but only for equations with zero initial data. This assumption is likewise adopted in the results of Krylov [17, 18], Boccia-Krylov [4], and Stinga-Torrea [22]. Recently, the first author of the present paper [7] resolved this limitation by developing a regularity theory in the weighted mixed norm spaces

$$L_p((0, T), w(t) dt; \Lambda^\gamma(\mathbb{R}^d)), \quad w \in A_p(\mathbb{R}), \quad p \in (1, \infty], \quad (1.2)$$

allowing for nonzero initial data through the use of the optimal trace theorem.

While the aforementioned results focus on time-dependent coefficients, the corresponding theory for space-time dependent coefficients within the Zygmund framework has not been fully established. The extension to variable coefficients is motivated not only by theoretical completeness but also by its relevance to nonlinear parabolic PDEs. A pertinent example is the two-dimensional incompressible Navier-Stokes equation in vorticity form

$$\partial_t \omega + u \cdot \nabla_x \omega = \Delta_x \omega, \quad u = \nabla_x^\perp \Delta_x^{-1} \omega. \quad (1.3)$$

It is well-known that for initial vorticity $\omega_0 \in L_1(\mathbb{R}^2) \cap L_\infty(\mathbb{R}^2)$, there exists a unique global weak solution satisfying $\omega \in L_\infty((0, T); L_1(\mathbb{R}^2) \cap L_\infty(\mathbb{R}^2))$, as established by Ben-Artzi and Brezis [2, 6]. Moreover, Ben-Artzi [2] also proved

$$\partial_t D_x^k \omega(t, \cdot), \quad D_x^k \omega(t, \cdot) \in L_\infty(\mathbb{R}^2), \quad \forall k \in \mathbb{N}, t > 0.$$

We note that this regularization is recovered via the partial Schauder theory. Treating the velocity field u as a transport coefficient that is bounded in time yet regular in space motivates the study of

linear theories in such mixed norm classes. Such a perspective provides a useful analytical tool for linearized models arising from nonlinear PDEs.

Recently, Wei et al. [24] studied parabolic PDEs with coefficients depending on both time and space, and obtained regularity estimates for solutions in Lebesgue-Hölder-Dini spaces. Their results, however, are derived under zero initial data and unweighted time norms, and do not extend to Zygmund regularity, weighted mixed norm settings, or the delicate *integer*-order case. Consequently, despite the well-developed theory for time-dependent coefficients, the Zygmund theory for fully variable coefficients $a_{ij}(t, x)$ remains largely open. Motivated by this gap, we study equation (1.1) in the weighted mixed norm space (1.2), thereby extending the existing partial Schauder and Zygmund regularity theories to parabolic PDEs with variable coefficients, including the integer-order regime $\gamma \in \mathbb{N}$.

We also refer to several works that investigate related regularity questions in different settings, such as those based on Zygmund spaces or the partial Schauder theory; see *e.g.*, [11, 12, 13, 19, 21].

1.3. Main results. We begin by introducing the weighted mixed norm Sobolev-Zygmund spaces that serve as the natural setting for our solutions. These spaces combine spatial Zygmund regularity with time-dependent L_p integrability under the Muckenhoupt weights, which is essential for treating parabolic PDEs whose coefficients are merely measurable in time.

Definition 1.3 (Solution space). Let $\gamma \in (0, \infty)$, $p \in (1, \infty]$, and $w \in A_p(\mathbb{R})$. We put $w \equiv 1$ when $p = \infty$.

(i) For $0 < S < T < \infty$, we denote

$$\Lambda_{p,w}^\gamma(S, T) := \begin{cases} L_p((S, T), w \, dt; \Lambda^\gamma(\mathbb{R}^d)), & \text{if } p \in (1, \infty), \\ L_\infty((S, T), dt; \Lambda^\gamma(\mathbb{R}^d)), & \text{if } p = \infty, \end{cases}$$

equipped with the norm defined by

$$\|u\|_{\Lambda_{p,w}^\gamma(S, T)} := \left(\int_S^T \|u(t, \cdot)\|_{\Lambda^\gamma(\mathbb{R}^d)}^p w(t) dt \right)^{1/p} 1_{p \in (1, \infty)} + \operatorname{ess\,sup}_{t \in (S, T)} \|u(t, \cdot)\|_{\Lambda^\gamma(\mathbb{R}^d)} 1_{p=\infty}.$$

When $S = 0$, we simply denote $\Lambda_{p,w}^\gamma(T) := \Lambda_{p,w}^\gamma(0, T)$.

(ii) We define $\Lambda_p^{\gamma,w}(\mathbb{R}^d)$ to be the set of all continuous functions u defined on \mathbb{R}^d satisfying

$$\|u\|_{\Lambda_p^{\gamma,w}(\mathbb{R}^d)} := \|u\|_{L_\infty(\mathbb{R}^d)} + \|u\|_{\dot{\Lambda}_p^{\gamma,w}(\mathbb{R}^d)} < \infty,$$

where

$$\|u\|_{\dot{\Lambda}_p^{\gamma,w}(\mathbb{R}^d)} := \left(\int_{\mathbb{R}^d} \frac{\|\mathcal{D}_h^{[\gamma]} u\|_{L_\infty(\mathbb{R}^d)}^p}{W(|h|^2)^{-1}} \frac{dh}{|h|^{d+p\gamma}} \right)^{1/p} 1_{p \in (1, \infty)} + \|u\|_{\dot{\Lambda}^\gamma(\mathbb{R}^d)} 1_{p=\infty},$$

and $W(\lambda) := \int_0^\lambda w(s) ds$.

(iii) The space $\mathbf{H}_{p,w}^{\gamma+2}(S, T)$ is the set of all $u \in \Lambda_{p,w}^{\gamma+2}(S, T)$ for which there exist $u_0 \in \Lambda_p^{\gamma+2,w}(\mathbb{R}^d)$ and $f \in \Lambda_{p,w}^\gamma(S, T)$ such that

$$u(t, x) = u_0(x) + \int_S^t f(s, x) ds$$

for all $(t, x) \in [S, T] \times \mathbb{R}^d$. We also define

$$\|u\|_{\mathbf{H}_{p,w}^{\gamma+2}(S, T)} := \|u\|_{\Lambda_{p,w}^{\gamma+2}(S, T)} + \|u_0\|_{\Lambda_p^{\gamma+2,w}(\mathbb{R}^d)} + \|f\|_{\Lambda_{p,w}^\gamma(S, T)}.$$

Remark 1.4. It can be verified directly that the pair $(u_0, f) \in \Lambda_p^{\gamma+2,w}(\mathbb{R}^d) \times \Lambda_{p,w}^\gamma(S, T)$ associated with u is unique. Hence, we may unambiguously write $f = \partial_t u$.

We now state the assumptions on the coefficients and present the main result. The conditions below ensure uniform ellipticity of the operator and spatial Λ^γ -regularity of the coefficients.

Assumption 1.5 (γ). There exist positive constants ν and K satisfying the following conditions:

(1) For all $(t, x) \in (0, \infty) \times \mathbb{R}^d$,

$$\nu|\xi|^2 \leq \sum_{i,j=1}^d a_{ij}(t, x)\xi_i\xi_j, \quad \forall \xi \in \mathbb{R}^d. \quad (1.4)$$

(2) The coefficients a_{ij} , b_i , and c are measurable, and satisfy

$$\sum_{i,j=1}^d \|a_{ij}\|_{L_\infty((0,\infty);\Lambda^\gamma(\mathbb{R}^d))} + \sum_{i=1}^d \|b_i\|_{L_\infty((0,\infty);\Lambda^\gamma(\mathbb{R}^d))} + \|c\|_{L_\infty((0,\infty);\Lambda^\gamma(\mathbb{R}^d))} \leq K.$$

Theorem 1.6. Let $T \in (0, \infty)$, $p \in (1, \infty]$, $\gamma > 0$, and let $w \in A_p(\mathbb{R})$ with $[w]_{A_p} \leq K_0$ (put $w \equiv 1$ and $K_0 = 1$ when $p = \infty$). Suppose that Assumption 1.5 (γ) holds. Then, for any pair $(u_0, f) \in \Lambda_p^{\gamma+2,w}(\mathbb{R}^d) \times \Lambda_{p,w}^\gamma(T)$, there exists a unique solution $u \in \mathbf{H}_{p,w}^{\gamma+2}(T)$ to (1.1) in $(0, T) \times \mathbb{R}^d$, with the initial condition $u(0, x) = u_0(x)$. Moreover, the following estimate holds:

$$\|u\|_{\mathbf{H}_{p,w}^{\gamma+2}(T)} \leq N(\|u_0\|_{\Lambda_p^{\gamma+2,w}(\mathbb{R}^d)} + \|f\|_{\Lambda_{p,w}^\gamma(T)}), \quad (1.5)$$

where $N = N(d, \gamma, K_0, K, \nu, T, p)$.

2. PRELIMINARIES ON HÖLDER-ZYGMUND SPACES

2.1. Characterizations of Hölder-Zygmund Spaces. In this subsection, we recall several equivalent descriptions of the Hölder-Zygmund spaces $\Lambda_p^\gamma(\mathbb{R}^d)$.

Proposition 2.1. Let $\gamma \in (0, \infty)$ and $p \in (1, \infty]$.

(i) (Equivalence with Besov spaces) Let $\{\Delta_j\}_{j \in \mathbb{Z}}$ be a standard Littlewood-Paley projection operators defined by

$$\Delta_j f := \psi_j * f, \quad \mathcal{F}[\psi_j](\xi) := \mathcal{F}[\psi](2^{-j}\xi), \quad \text{supp}(\mathcal{F}[\psi]) = \{\xi \in \mathbb{R}^d : 1/2 \leq |\xi| \leq 2\},$$

with $\mathcal{F}[\psi] \geq 0$ and $\sum_{j \in \mathbb{Z}} \mathcal{F}[\psi](2^{-j}\xi) = 1$ for $\xi \neq 0$. Then

$$\|f\|_{\dot{\Lambda}_p^\gamma(\mathbb{R}^d)} \simeq \|f\|_{\dot{B}_{\infty,p}^\gamma(\mathbb{R}^d)}$$

where

$$\|f\|_{\dot{B}_{\infty,p}^\gamma(\mathbb{R}^d)} := \begin{cases} \left(\sum_{j \in \mathbb{Z}} 2^{j\gamma p} \|\Delta_j f\|_{L_\infty(\mathbb{R}^d)}^p \right)^{1/p}, & \text{if } p \in (1, \infty), \\ \sup_{j \in \mathbb{Z}} 2^{j\gamma} \|\Delta_j f\|_{L_\infty(\mathbb{R}^d)}, & \text{if } p = \infty. \end{cases}$$

(ii) (Characterization via derivatives) If $\gamma = n + \delta$, $n \in \mathbb{N} \cup \{0\}$, and $\delta \in (0, 1]$, then

$$\|f\|_{\Lambda_p^\gamma(\mathbb{R}^d)} \simeq \sum_{|\alpha| \leq n} \|D^\alpha f\|_{L_\infty(\mathbb{R}^d)} + \sum_{|\beta|=n} \|D^\beta f\|_{\dot{\Lambda}_p^\delta(\mathbb{R}^d)}.$$

The equivalence constants depend only on d , γ , and p .

Proof. The result (i) is well-known, e.g., see [8, Theorem 1.5, Theorem 1.6].

For (ii), iterating elementary difference estimates, we have

$$\|\mathcal{D}_h^{[\gamma]} f\|_{L_\infty(\mathbb{R}^d)} \leq |h|^n \sum_{|\alpha|=n} \|\mathcal{D}_h^{[\delta]} D^\alpha f\|_{L_\infty(\mathbb{R}^d)}.$$

Therefore,

$$\|f\|_{\dot{\Lambda}_p^\gamma(\mathbb{R}^d)} \leq N(n, p) \sum_{|\alpha|=n} \|D^\alpha f\|_{\dot{\Lambda}_p^\delta(\mathbb{R}^d)},$$

which proves the “ \leq ” direction.

For the converse inequality, note that for any multi-index α ,

$$\|\Delta_j D^\alpha f\|_{L_\infty(\mathbb{R}^d)} \leq N 2^{j|\alpha|} \|\Delta_j f\|_{L_\infty(\mathbb{R}^d)}. \quad (2.6)$$

Hence,

$$\begin{aligned} \|D^\alpha f\|_{L_\infty(\mathbb{R}^d)} &\leq N \sum_{j \in \mathbb{Z}} 2^{j|\alpha|} \|\Delta_j f\|_{L_\infty(\mathbb{R}^d)} \\ &\leq N \left(\|f\|_{L_\infty(\mathbb{R}^d)} + \sum_{j=1}^{\infty} 2^{j|\alpha|} \|\Delta_j f\|_{L_\infty(\mathbb{R}^d)} \right), \end{aligned} \quad (2.7)$$

For $|\alpha| \leq n$ and $p \in (1, \infty)$, Hölder's inequality gives

$$\begin{aligned} \sum_{j=1}^{\infty} 2^{j|\alpha|} \|\Delta_j f\|_{L_\infty(\mathbb{R}^d)} &\leq \left(\sum_{j=1}^{\infty} 2^{j\gamma p} \|\Delta_j f\|_{L_\infty(\mathbb{R}^d)}^p \right)^{1/p} \left(\sum_{j=1}^{\infty} 2^{j(|\alpha|-\gamma)p'} \right)^{1/p'} \\ &\leq N \left(\sum_{j=1}^{\infty} 2^{j\gamma p} \|\Delta_j f\|_{L_\infty(\mathbb{R}^d)}^p \right)^{1/p} \leq N \|f\|_{\dot{B}_{\infty,p}^\gamma(\mathbb{R}^d)}, \end{aligned} \quad (2.8)$$

where $p' = p/(p-1)$. Combining (2.7), (2.8), and (i), we obtain the desired bound for $p \in (1, \infty)$. When $p = \infty$, we instead use the trivial bound

$$\sum_{j=1}^{\infty} 2^{j|\alpha|} \|\Delta_j f\|_{L_\infty} \leq \sup_{j \in \mathbb{Z}} (2^{j\gamma} \|\Delta_j f\|_{L_\infty}) \sum_{j=1}^{\infty} 2^{j(|\alpha|-\gamma)} \leq N \|f\|_{\dot{B}_{\infty,\infty}^\gamma(\mathbb{R}^d)}.$$

Thus, for all $p \in (1, \infty]$,

$$\|D^\alpha f\|_{L_\infty(\mathbb{R}^d)} \leq N \|f\|_{\Lambda_p^\gamma(\mathbb{R}^d)}, \quad \forall |\alpha| \leq n.$$

Finally, by (i) and (2.6), for $|\beta| = n$ we have

$$\|D^\beta f\|_{\dot{\Lambda}_p^\delta(\mathbb{R}^d)} \simeq \left(\sum_{j \in \mathbb{Z}} 2^{j\delta p} \|\Delta_j D^\beta f\|_{L_\infty(\mathbb{R}^d)}^p \right)^{1/p} \leq N \left(\sum_{j \in \mathbb{Z}} 2^{j\gamma p} \|\Delta_j f\|_{L_\infty(\mathbb{R}^d)}^p \right)^{1/p}$$

with the obvious modification when $p = \infty$ (replace ℓ_p by ℓ_∞). Summing over all $|\beta| = n$ completes the proof. \square

If $p = \infty$ and $\gamma > 0$ is a non-integer, then the Zygmund space $\Lambda^\gamma(\mathbb{R}^d)$ coincides with the classical Hölder space $C^\gamma(\mathbb{R}^d)$. At integer orders $\gamma \in \mathbb{N}$, however, this identification fails. The next result states the precise inclusions between the corresponding spaces.

Proposition 2.2. *For $n \in \mathbb{N}$,*

$$C^n(\mathbb{R}^d) \subsetneq C^{n-1,1}(\mathbb{R}^d) \subsetneq \Lambda^n(\mathbb{R}^d) \subsetneq C_{\log}^{n-1,1}(\mathbb{R}^d),$$

where

$$\|f\|_{C_{\log}^{n-1,1}(\mathbb{R}^d)} := \sum_{|\alpha| \leq n-1} \|D^\alpha f\|_{L_\infty(\mathbb{R}^d)} + \sum_{|\beta|=n-1} \sup_{|x-y|<1} \frac{|D^\beta f(x) - D^\beta f(y)|}{|x-y| \log_2(2/|x-y|)}.$$

Proof. The strict inclusion $C^n(\mathbb{R}^d) \subsetneq C^{n-1,1}(\mathbb{R}^d)$ is well known, so we focus on the remaining inclusions. By Proposition 2.1-(ii),

$$\|f\|_{\Lambda^n(\mathbb{R}^d)} \simeq \sum_{|\beta|=n-1} \|D^\beta f\|_{\Lambda^1(\mathbb{R}^d)} + 1_{n \geq 2} \sum_{|\alpha| \leq n-2} \|D^\alpha f\|_{L_\infty(\mathbb{R}^d)}.$$

Thus, it suffices to establish the result for $n = 1$, namely

$$C^{0,1}(\mathbb{R}^d) \subsetneq \Lambda^1(\mathbb{R}^d) \subsetneq C_{\log}^{0,1}(\mathbb{R}^d). \quad (2.9)$$

1. $C^{0,1}(\mathbb{R}^d) \subsetneq \Lambda^1(\mathbb{R}^d)$.

Let $\zeta \in C_c^\infty(\mathbb{R}^d)$ be a nonnegative cut-off function near zero, and $f(x) := x^1 \log_2(2/|x|)$. Then for $|h| < 1$,

$$|f(h) - f(0)| = |h^1| \log_2(2/|h|),$$

which implies that $f\zeta \notin C^{0,1}(\mathbb{R}^d)$. However, $f\zeta \in \Lambda^1(\mathbb{R}^d)$, which easily follows from

$$|f(h) + f(-h) - 2f(0)| = 0 \leq |h|.$$

2. $\Lambda^1(\mathbb{R}^d) \subsetneq C_{\log}^{0,1}(\mathbb{R}^d)$.

The proof of $\Lambda^1(\mathbb{R}^d) \subset C_{\log}^{0,1}(\mathbb{R}^d)$ is provided in [1, Proposition 2.107], but the strictness is not. For the sake of completeness, we provide the proof and a counterexample. For every $m \in \mathbb{N}$,

$$\mathcal{D}_h f = -\frac{1}{2} \sum_{k=0}^{m-1} 2^{-k} \mathcal{D}_{2^k h}^2 f + 2^{-m} \mathcal{D}_{2^m h} f. \quad (2.10)$$

Taking L^∞ -norms gives

$$\begin{aligned} \|\mathcal{D}_h f\|_{L^\infty(\mathbb{R}^d)} &\leq \frac{1}{2} \sum_{k=0}^{m-1} 2^{-k} \|\mathcal{D}_{2^k h}^2 f\|_{L^\infty(\mathbb{R}^d)} + 2^{-m} \|\mathcal{D}_{2^m h} f\|_{L^\infty(\mathbb{R}^d)} \\ &\leq \frac{1}{2} \sum_{k=0}^{m-1} 2^{-k} (\|f\|_{\dot{\Lambda}^1(\mathbb{R}^d)} 2^k |h|) + 2^{1-m} \|f\|_{L^\infty(\mathbb{R}^d)} \\ &= \frac{m|h|}{2} \|f\|_{\dot{\Lambda}^1(\mathbb{R}^d)} + 2^{1-m} \|f\|_{L^\infty(\mathbb{R}^d)}. \end{aligned}$$

For $|h| < 1$ choose $m = \lfloor \log_2(2/|h|) \rfloor$ so that $m \leq \log_2(2/|h|)$ and $2^{-m} \leq |h|$. Then

$$\|\mathcal{D}_h f\|_{L^\infty(\mathbb{R}^d)} \leq |h| \log_2 \left(\frac{2}{|h|} \right) \|f\|_{\dot{\Lambda}^1(\mathbb{R}^d)} + 2|h| \|f\|_{L^\infty(\mathbb{R}^d)} \leq 3 \|f\|_{\Lambda^1(\mathbb{R}^d)} |h| \log_2 \left(\frac{2}{|h|} \right),$$

since $\log_2(2/|h|) \geq 1$ for $|h| < 1$. Taking the supremum over $|h| < 1$ yields

$$\sup_{|x-y| < 1} \frac{|f(x) - f(y)|}{|x-y| \log_2(2/|x-y|)} \leq 2 \|f\|_{\Lambda^1(\mathbb{R}^d)}. \quad (2.11)$$

To show $\Lambda^1(\mathbb{R}^d) \neq C_{\log}^{0,1}(\mathbb{R}^d)$, we put $g(x) := |x| \log_2(2/|x|)$. Then

$$|g(h) - g(-h) - 2g(0)| = 2|h| \log_2(2/|h|),$$

which implies that $g\zeta \notin \Lambda^1(\mathbb{R}^d)$. However, $g\zeta \in C_{\log}^{0,1}(\mathbb{R}^d)$. The proposition is proved. \square

2.2. Properties of Hölder-Zygmund Spaces. We next record several lemmas used throughout the proof. They provide

- tool for localization via partition of unity (Lemma 2.3),
- interpolation between Zygmund spaces (Lemma 2.4),
- estimates for products of functions in Zygmund classes (Lemma 2.5).

All of these will be employed in Step 1 of the proof of Theorem 3.1.

The following partition of unity lemma is a modified version of [16, Lemma 4.1.1], and we include the proof for completeness.

Lemma 2.3 (Partition of unity). *Let $\gamma > 0$, and let $\zeta \in C_c^\infty(\mathbb{R}^d)$ be a cut-off function such that $\zeta(x) = 1$ for $|x| \leq 1$, $\zeta(x) = 0$ for $|x| \geq 2$, and $0 \leq \zeta \leq 1$. Then for any $R > 0$, $y \in \mathbb{R}^d$, and $u \in \Lambda^\gamma(\mathbb{R}^d)$, we have*

$$\|u\|_{\Lambda^\gamma(\mathbb{R}^d)} \leq N_0 \sup_{z \in \mathbb{R}^d} \|u \zeta_R^z\|_{\dot{\Lambda}^\gamma(\mathbb{R}^d)} + N_1 \sup_{z \in \mathbb{R}^d} \|u \zeta_R^z\|_{L^\infty(\mathbb{R}^d)}, \quad (2.12)$$

where $\zeta_R^z(x) := \zeta\left(\frac{x-z}{R}\right)$, $N_0 = N_0(d, \gamma, \zeta)$, and $N_1 = N_1(d, \gamma, R, \zeta)$.

Proof. We first claim that

$$\|u\|_{\dot{\Lambda}^\gamma(\mathbb{R}^d)} \leq N(\gamma, R) \|u\|_{L_\infty(\mathbb{R}^d)} + \sup_{y \in \mathbb{R}^d} \sup_{|h| \leq R/[\gamma]^-} \frac{\|\mathcal{D}_h^{[\gamma]^-} u\|_{L_\infty(B_R(y))}}{|h|^\gamma}. \quad (2.13)$$

Indeed, for any $h, y \in \mathbb{R}^d$ and $x \in B_R(y)$,

$$\begin{aligned} \frac{|\mathcal{D}_h^{[\gamma]^-} u(x)|}{|h|^\gamma} &= \frac{|\mathcal{D}_h^{[\gamma]^-} u(x)|}{|h|^\gamma} \mathbf{1}_{|h| \leq \frac{R}{[\gamma]^-}} + \frac{|\mathcal{D}_h^{[\gamma]^-} u(x)|}{|h|^\gamma} \mathbf{1}_{|h| > \frac{R}{[\gamma]^-}} \\ &\leq \sup_{|h| \leq \frac{R}{[\gamma]^-}} \frac{\|\mathcal{D}_h^{[\gamma]^-} u\|_{L_\infty(B_R(y))}}{|h|^\gamma} + N(\gamma, R) \|u\|_{L_\infty(\mathbb{R}^d)}. \end{aligned}$$

Next, fix $y \in \mathbb{R}^d$. For $x \in B_{2R}(y)$, we have the identity

$$u(x) = cR^{-d} \int_{|z-y| \leq 4R} u(z) \zeta \left(\frac{x-z}{R} \right) dz = cR^{-d} \int_{|z-y| \leq 4R} u(z) \zeta_R^z(x) dz, \quad (2.14)$$

where $c^{-1} := \int_{\mathbb{R}^d} \zeta(z) dz$. Due to (2.14),

$$\|u\|_{L_\infty(\mathbb{R}^d)} \leq N \sup_{z \in \mathbb{R}^d} \|u \zeta_R^z\|_{L_\infty(\mathbb{R}^d)}. \quad (2.15)$$

For any natural number $k \leq [\gamma]^-$ and $|h| \leq R/[\gamma]^-$, we have for $x \in B_R(y)$, $x + kh \in B_{2R}(y)$. Hence, by (2.14),

$$\begin{aligned} u(x + kh) &= cR^{-d} \int_{|z-y| \leq 4R} u(z + kh) \zeta \left(\frac{x + kh - z}{R} \right) dz \\ &= cR^{-d} \int_{|z-y| \leq 4R} u(z + kh) \zeta_R^z(x + kh) dz. \end{aligned}$$

Therefore,

$$\mathcal{D}_h^{[\gamma]^-} u(x) = cR^{-d} \int_{|z-y| \leq 4R} \mathcal{D}_h^{[\gamma]^-} (u \zeta_R^z)(x) dz, \quad \forall x \in B_R(y).$$

Taking the supremum over $x \in B_R(y)$ yields

$$\frac{\|\mathcal{D}_h^{[\gamma]^-} u\|_{L_\infty(B_R(y))}}{|h|^\gamma} \leq cR^{-d} \int_{|z-y| \leq 4R} \frac{\|\mathcal{D}_h^{[\gamma]^-} (u \zeta_R^z)\|_{L_\infty(\mathbb{R}^d)}}{|h|^\gamma} dz \leq N \|u \zeta_R^z\|_{\dot{\Lambda}^\gamma(\mathbb{R}^d)}. \quad (2.16)$$

Combining (2.13), (2.15), and (2.16) completes the proof. \square

Lemma 2.4 (Interpolation inequality). *Let $\gamma \in (0, \infty)$ and $f \in \Lambda^\gamma(\mathbb{R}^d)$. Then for $\theta \in (0, 1)$ and $\varepsilon > 0$,*

$$\|f\|_{\dot{\Lambda}^{\theta\gamma}(\mathbb{R}^d)} \leq N\varepsilon \|f\|_{\dot{\Lambda}^\gamma(\mathbb{R}^d)} + N\varepsilon^{-\frac{\theta}{1-\theta}} \|f\|_{L_\infty(\mathbb{R}^d)}, \quad \forall \varepsilon > 0,$$

where N is independent of f and ε .

Proof. By [3, Theorem 6.4.5] and Proposition 2.1,

$$\Lambda^{\theta\gamma}(\mathbb{R}^d) = (B_{\infty,\infty}^0(\mathbb{R}^d), \Lambda^\gamma(\mathbb{R}^d))_{\theta,\infty},$$

where $B_{\infty,\infty}^0(\mathbb{R}^d)$ denotes the Besov space of order 0. By [23, (6) in Section 2.4.1],

$$\|f\|_{\Lambda^{\theta\gamma}(\mathbb{R}^d)} \simeq \|f\|_{(B_{\infty,\infty}^0(\mathbb{R}^d), \Lambda^\gamma(\mathbb{R}^d))_{\theta,\infty}} \leq 2 \|f\|_{B_{\infty,\infty}^0(\mathbb{R}^d)}^{1-\theta} \|f\|_{\Lambda^\gamma(\mathbb{R}^d)}^\theta. \quad (2.17)$$

Let

$$S_0 f(x) = \int_{\mathbb{R}^d} f(x-y) \Phi(y) dy, \quad \Delta_j f(x) = \int_{\mathbb{R}^d} f(x-y) \psi_j(y) dy,$$

where $\Phi := \sum_{j \leq 0} \psi_j$. Then we obtain

$$\|f\|_{B_{\infty, \infty}^0(\mathbb{R}^d)} := \|S_0 f\|_{L^\infty(\mathbb{R}^d)} + \sup_{j \in \mathbb{N}} \|\Delta_j f\|_{L^\infty(\mathbb{R}^d)} \leq C_\psi \|f\|_{L^\infty(\mathbb{R}^d)}, \quad (2.18)$$

where

$$C_\psi := \|\Phi\|_{L^1(\mathbb{R}^d)} + \sup_{j \in \mathbb{Z}} \|\psi_j\|_{L^1(\mathbb{R}^d)} < \infty.$$

For any $c, \eta > 0$

$$\|f(c \cdot)\|_{\dot{\Lambda}^\eta(\mathbb{R}^d)} = c^\eta \|f\|_{\dot{\Lambda}^\eta(\mathbb{R}^d)}. \quad (2.19)$$

Combining (2.17)–(2.19) yields

$$\begin{aligned} \|f(c \cdot)\|_{\Lambda^{\theta\gamma}(\mathbb{R}^d)} &= \|f\|_{L^\infty(\mathbb{R}^d)} + c^{\theta\gamma} \|f\|_{\dot{\Lambda}^{\theta\gamma}(\mathbb{R}^d)} \\ &\leq N \|f\|_{L^\infty(\mathbb{R}^d)}^{1-\theta} (\|f\|_{L^\infty(\mathbb{R}^d)} + c^\gamma \|f\|_{\dot{\Lambda}^\gamma(\mathbb{R}^d)})^\theta \\ &\leq N \|f\|_{L^\infty(\mathbb{R}^d)} + N c^{\theta\gamma} \|f\|_{L^\infty(\mathbb{R}^d)}^{1-\theta} \|f\|_{\dot{\Lambda}^\gamma(\mathbb{R}^d)}^\theta, \end{aligned} \quad (2.20)$$

where N is independent of c . Dividing both sides by $c^{\theta\gamma}$ and letting $c \rightarrow \infty$, we obtain

$$\|f\|_{\dot{\Lambda}^{\theta\gamma}(\mathbb{R}^d)} \leq N \|f\|_{L^\infty(\mathbb{R}^d)}^{1-\theta} \|f\|_{\dot{\Lambda}^\gamma(\mathbb{R}^d)}^\theta.$$

Applying Young's inequality gives the desired estimate. \square

Lemma 2.5. *Let $\gamma \in (0, \infty)$ and $f, g \in \Lambda^\gamma(\mathbb{R}^d)$. Then $fg \in \Lambda^\gamma(\mathbb{R}^d)$, and*

$$\begin{aligned} \|fg\|_{\dot{\Lambda}^\gamma(\mathbb{R}^d)} &\leq \|f\|_{L^\infty(\mathbb{R}^d)} \|g\|_{\dot{\Lambda}^\gamma(\mathbb{R}^d)} + \sum_{j=1}^{[\gamma]-1} \binom{[\gamma]-1}{j} \|f\|_{\dot{\Lambda}^j(\mathbb{R}^d)} \|g\|_{\dot{\Lambda}^{\gamma-j}(\mathbb{R}^d)} \\ &\quad + \|f\|_{\dot{\Lambda}^\gamma(\mathbb{R}^d)} \|g\|_{L^\infty(\mathbb{R}^d)}. \end{aligned} \quad (2.21)$$

Proof. The inequality (2.21) can be directly obtained from

$$\mathcal{D}_h^{[\gamma]-} (fg)(x) = \sum_{j=0}^{[\gamma]-} \binom{[\gamma]-1}{j} \mathcal{D}_h^j f(x + ([\gamma]-j)h) \mathcal{D}_h^{[\gamma]-j} g(x).$$

The lemma is proved. \square

3. PROOF OF THEOREM 1.6

3.1. Proof of Theorem 1.6 with zero initial data. In this section, we consider the Cauchy problem

$$\partial_t u(t, x) = \sum_{i,j=1}^d a_{ij}(t, x) D_{ij} u(t, x) + \sum_{i=1}^d b_i(t, x) D_i u + c(t, x) u(t, x) + f(t, x), \quad (3.22)$$

subject to the zero initial condition $u(0, \cdot) = 0$.

Theorem 3.1. *Let $T \in (0, \infty)$, $p \in (1, \infty]$, $\gamma > 0$, and let $w \in A_p(\mathbb{R})$ with $[w]_{A_p} \leq K_0$ (put $w \equiv 1$ and $K_0 = 1$ when $p = \infty$). Suppose that Assumption 1.5 (γ) holds. Then, for any $f \in \Lambda_{p,w}^\gamma(T)$, there exists a unique solution $u \in \mathbf{H}_{p,w}^{\gamma+2}(T)$ to the equation (3.22) in $(0, T) \times \mathbb{R}^d$, with zero initial condition $u(0, x) = 0$. Moreover, the following estimate holds:*

$$\|u\|_{\mathbf{H}_{p,w}^{\gamma+2}(T)} \leq N \|f\|_{\Lambda_{p,w}^\gamma(T)},$$

where $N = N(d, \gamma, K_0, \nu, T, p)$.

The next two lemmas play crucial roles in the proof of a priori estimate.

Lemma 3.2. *Let $\eta > 0$, $p \in (1, \infty]$, and $T > 0$. Suppose that*

- $a = (a_{ij}(t))_{i,j}$ are independent of x .
- $a = (a_{ij}(t))_{i,j}$ is bounded and satisfies (1.4).

- $b_i = c = 0$ for all $i = 1, \dots, d$.

If u is a solution to (3.22) with $f \in \mathbf{A}_{p,w}^\eta(T)$, then for each $t \in (0, T)$,

$$\|u(t, \cdot)\|_{\Lambda^{\eta+2}(\mathbb{R}^d)} \leq N(T) \mathcal{M}_t (\|f(*, \cdot)\|_{\Lambda^\eta(\mathbb{R}^d)} 1_{(0,T)}(*)) (t), \quad (3.23)$$

where

$$\mathcal{M}_t f(t) := \sup_{r>0} \int_{t-r}^{t+r} f(s) ds.$$

In particular, $N(T)$ is increasing in T .

Proof. This is a direct consequence of [7, Lemma 2.6] with $\phi(\lambda) = \lambda^\eta$, $\gamma = 2$, and $\phi_\gamma(\lambda) = \lambda^{\eta+2}$. \square

Lemma 3.3. Let $\gamma \in (0, \infty)$, $p \in (1, \infty]$, and $w \in A_p$ such that $[w]_{A_p} \leq K_0$ (put $w \equiv 1$ and $K_0 = 1$ when $p = \infty$). Suppose that $u \in \mathbf{A}_{p,w}^\gamma(a, b)$ satisfies $u(a, \cdot) = 0$. Then we have

$$\|u\|_{L_p((a,b), w dt; L_\infty(\mathbb{R}^d))} \leq N(b-a) \|\partial_t u\|_{L_p((a,b), w dt; L_\infty(\mathbb{R}^d))}, \quad (3.24)$$

where $N = N(d, p, K_0)$.

Proof. By the fundamental theorem of calculus,

$$|u(t, x)| \leq \int_a^t |\partial_t u(s, x)| ds = (t-a) \int_a^t |\partial_t u(s, x)| ds.$$

Consequently,

$$\|u(t, \cdot)\|_{L_\infty(\mathbb{R}^d)} \leq 2(t-a) \mathcal{M}_t (\|\partial_t u(*, \cdot)\|_{L_\infty(\mathbb{R}^d)} 1_{(a,b)}(*)) (t), \quad \forall t \in (a, b).$$

Applying the weighted Hardy–Littlewood maximal function theorem (see, e.g., [10, Theorem 2.2]) yields (3.24). This completes the proof. \square

We now prove the a priori estimate for the case with zero initial data.

Lemma 3.4 (A priori estimate). Let $\gamma \in (0, \infty)$, $p \in (1, \infty]$, $T > 0$, and $w \in A_p$ with $[w]_{A_p} \leq K_0$ (put $w \equiv 1$ and $K_0 = 1$ when $p = \infty$). Suppose that $f \in \mathbf{A}_{p,w}^\gamma(T)$, and $u \in \mathbf{H}_{p,w}^{\gamma+2}(T)$ is a solution to (3.22). Then

$$\|\partial_t u\|_{\mathbf{A}_{p,w}^\gamma(T)} + \|u\|_{\mathbf{A}_{p,w}^{\gamma+2}(T)} \leq N \|f\|_{\mathbf{A}_{p,w}^\gamma(T)}, \quad (3.25)$$

where $N = N(d, \nu, K, p, K_0, T)$.

Proof. We divide the proof into two steps.

Step 1. We first establish

$$\|\partial_t u\|_{\mathbf{A}_{p,w}^\gamma(T)} + \|u\|_{\mathbf{A}_{p,w}^{\gamma+2}(T)} \leq N \left(\|f\|_{\mathbf{A}_{p,w}^\gamma(T)} + \|u\|_{L_p((0,T), w dt; L_\infty(\mathbb{R}^d))} \right), \quad (3.26)$$

where $N = N(d, \nu, K, K_0, p, T)$.

Since u satisfies the equation (3.22), we have

$$\begin{aligned} \|\partial_t u\|_{\mathbf{A}_{p,w}^\gamma(T)} &= \left\| \sum_{i,j=1}^d a_{ij} D_{ij} u + \sum_{i=1}^d b_i D_i u + c u + f \right\|_{\mathbf{A}_{p,w}^\gamma(T)} \\ &\leq N \left(\|u\|_{\mathbf{A}_{p,w}^{\gamma+2}(T)} + \|f\|_{\mathbf{A}_{p,w}^\gamma(T)} \right). \end{aligned}$$

Thus, in order to obtain (3.26), it remains to estimate $\|u\|_{\mathbf{A}_{p,w}^{\gamma+2}(T)}$ in terms of the right-hand side.

To achieve this, it is enough to deal with the case $b_i \equiv c \equiv 0$. Indeed, once the required estimate holds under this assumption, we have

$$\begin{aligned} \|\partial_t u\|_{\Lambda_{p,w}^\gamma(T)} + \|u\|_{\Lambda_{p,w}^{\gamma+2}(T)} &\leq N \left\| f + \sum_{i=1}^d b_i D_i u + c u \right\|_{\Lambda_{p,w}^\gamma(T)} \\ &\leq N \left(\|f\|_{\Lambda_{p,w}^\gamma(T)} + \|u\|_{\Lambda_{p,w}^{\gamma+1}(T)} \right). \end{aligned}$$

We now estimate the remaining $\|u\|_{\Lambda_{p,w}^{\gamma+1}(T)}$ term. By Lemma 2.4,

$$\|u\|_{\Lambda_{p,w}^{\gamma+1}(T)} \leq N\varepsilon \|u\|_{\Lambda_{p,w}^{\gamma+2}(T)} + N(\varepsilon) \|u\|_{L_p((0,T), w \, dt; L_\infty(\mathbb{R}^d))}. \quad (3.27)$$

Choosing $\varepsilon > 0$ sufficiently small allows the first term on the right-hand side to be absorbed into the left-hand side of the estimate, thereby yielding (3.26). Therefore, we assume that $f \in \Lambda_{p,w}^\gamma(T)$, and $u \in \mathbf{H}_{p,w}^{\gamma+2}(T)$ is a solution to (3.22) with $b_i = c = 0$ and zero initial condition.

We first consider $\gamma \in (0, 1]$. Let $\zeta \in C_c^\infty(\mathbb{R}^d)$ such that $\zeta = 1$ on B_1 , $\zeta = 0$ on $\mathbb{R}^d \setminus B_2$, and $0 \leq \zeta \leq 1$. For $R > 0$ and $y \in \mathbb{R}^d$, define $\zeta_R^y(x) := \zeta((x - y)/R)$ and set $v^y := u\zeta_R^y$. Then v^y satisfies

$$\begin{aligned} \partial_t v^y &= \sum_{i,j=1}^d a_{ij} D_{ij} v^y + \zeta_R^y f - \sum_{i,j=1}^d (a_{ij} + a_{ji}) D_i u D_j \zeta_R^y - \sum_{i,j=1}^d a_{ij} u D_{ij} \zeta_R^y \\ &= \sum_{i,j=1}^d a_{ij}(t, y) D_{ij} v^y + F, \end{aligned}$$

where

$$\begin{aligned} F &:= \sum_{i,j=1}^d (a_{ij} - a_{ij}(t, y)) D_{ij} v^y + \zeta_R^y f - \sum_{i,j=1}^d (a_{ij} + a_{ji}) D_i u D_j \zeta_R^y - \sum_{i,j=1}^d a_{ij} u D_{ij} \zeta_R^y \\ &=: \sum_{i,j=1}^d (a_{ij} - a_{ij}(t, y)) D_{ij} v^y + G. \end{aligned}$$

Due to $\Lambda^\gamma = C^\gamma$ if $\gamma \in (0, 1)$, and (2.11), $\|a_{ij}(t, \cdot) - a_{ij}(t, y)\|_{L_\infty(B_{2R}(y))} \leq N A_\gamma(R)$, where $N = N(\nu)$ and $A_\gamma(R) := R^\gamma 1_{\gamma \in (0,1)} + R \log_2(2/R) 1_{\gamma=1}$. Using Lemma 2.5, Proposition 2.1-(ii) and Lemma 2.4 with $\varepsilon = A_\gamma(R)$, we have

$$\begin{aligned} &\left\| \sum_{i,j=1}^d (a_{ij}(t, \cdot) - a_{ij}(t, y)) D_{ij} v^y(t, \cdot) \right\|_{\Lambda^\gamma(\mathbb{R}^d)} \\ &\leq N(d) \sum_{i,j=1}^d \| (a_{ij}(t, \cdot) - a_{ij}(t, y)) \|_{L_\infty(B_{2R}(y))} \| D_x^2 v^y(t, \cdot) \|_{\Lambda^\gamma(\mathbb{R}^d)} \\ &\quad + N(d) \sum_{i,j=1}^d \| (a_{ij}(t, \cdot) - a_{ij}(t, y)) \|_{\Lambda^\gamma(\mathbb{R}^d)} \| D_x^2 v^y(t, \cdot) \|_{L_\infty(\mathbb{R}^d)} \\ &\leq N A_\gamma(R) \| v^y(t, \cdot) \|_{\Lambda^{\gamma+2}(\mathbb{R}^d)} + N \| v^y(t, \cdot) \|_{\Lambda^2(\mathbb{R}^d)} \\ &\leq N A_\gamma(R) \| v^y(t, \cdot) \|_{\Lambda^{\gamma+2}(\mathbb{R}^d)} + N(d, \nu, \gamma, R) \| u(t, \cdot) \|_{L_\infty(\mathbb{R}^d)}. \end{aligned} \quad (3.28)$$

Due to (2.19), for $n \in \mathbb{N} \cup \{0\}$

$$\| D_x^n \zeta_R^y \|_{\Lambda^\gamma(\mathbb{R}^d)} = R^{-n} \| D_x^n \zeta \|_{L_\infty(\mathbb{R}^d)} + R^{-n-\gamma} \| D_x^n \zeta \|_{\dot{\Lambda}^\gamma(\mathbb{R}^d)}. \quad (3.29)$$

For G , Lemma 2.5 and (3.29) imply

$$\begin{aligned} \|G(t, \cdot)\|_{\Lambda^\gamma(\mathbb{R}^d)} &\leq N \|\zeta_R^y\|_{\Lambda^\gamma(\mathbb{R}^d)} \|f(t, \cdot)\|_{\Lambda^\gamma(\mathbb{R}^d)} \\ &\quad + N(\|D_x \zeta_R^y\|_{\Lambda^\gamma(\mathbb{R}^d)} + \|D_{xx} \zeta_R^y\|_{\Lambda^\gamma(\mathbb{R}^d)}) \|u(t, \cdot)\|_{\Lambda^{\gamma+1}(\mathbb{R}^d)} \\ &\leq N(R) \|f(t, \cdot)\|_{\Lambda^\gamma(\mathbb{R}^d)} + N(R) \|u(t, \cdot)\|_{\Lambda^{\gamma+1}(\mathbb{R}^d)}, \end{aligned} \quad (3.30)$$

where $N(R)$ is independent of y . Combining (3.28), (3.30), and Lemma 3.2 gives

$$\begin{aligned} &\|v^y(t, \cdot)\|_{\Lambda^{\gamma+2}(\mathbb{R}^d)} \\ &\leq N \mathcal{M}_t (\|F(*, \cdot)\|_{\Lambda^\gamma(\mathbb{R}^d)} 1_{(0,T)}(*)) (t) \\ &\leq N \mathcal{M}_t (\|(a_{ij} - a_{ij}(*, y)) D_{ij} v^y(*, \cdot)\|_{\Lambda^\gamma(\mathbb{R}^d)} 1_{(0,T)}(*)) (t) \\ &\quad + N \mathcal{M}_t (\|G(*, \cdot)\|_{\Lambda^\gamma(\mathbb{R}^d)} 1_{(0,T)}(*)) (t) \\ &\leq N A_\gamma(R) \mathcal{M}_t (\|v^y(*, \cdot)\|_{\Lambda^{\gamma+2}(\mathbb{R}^d)} 1_{(0,T)}(*)) (t) \\ &\quad + N(R) \mathcal{M}_t ((\|u(*, \cdot)\|_{\Lambda^{\gamma+1}(\mathbb{R}^d)} + \|f(*, \cdot)\|_{\Lambda^\gamma(\mathbb{R}^d)} + \|u(*, \cdot)\|_{L_\infty(\mathbb{R}^d)}) 1_{(0,T)}(*)) (t). \end{aligned} \quad (3.31)$$

Since the constants in (3.31) are independent of y , we have

$$\begin{aligned} &\sup_{y \in \mathbb{R}^d} \|v^y(t, \cdot)\|_{\Lambda^{\gamma+2}(\mathbb{R}^d)} \\ &\leq N A_\gamma(R) \mathcal{M}_t \left(\sup_{y \in \mathbb{R}^d} \|v^y(*, \cdot)\|_{\Lambda^{\gamma+2}(\mathbb{R}^d)} 1_{(0,T)}(*) \right) (t) \\ &\quad + N(R) \mathcal{M}_t ((\|u(*, \cdot)\|_{\Lambda^{\gamma+1}(\mathbb{R}^d)} + \|f(*, \cdot)\|_{\Lambda^\gamma(\mathbb{R}^d)} + \|u(*, \cdot)\|_{L_\infty(\mathbb{R}^d)}) 1_{(0,T)}(*)) (t) \end{aligned}$$

By the Hardy-Littlewood maximal function theorem, when $p \in (1, \infty)$,

$$\begin{aligned} &\int_0^T \sup_{y \in \mathbb{R}^d} \|v^y(t, \cdot)\|_{\Lambda^{\gamma+2}(\mathbb{R}^d)}^p w(t) dt \\ &\leq N A_\gamma(R) \int_0^T \sup_{y \in \mathbb{R}^d} \|v^y(t, \cdot)\|_{\Lambda^{\gamma+2}(\mathbb{R}^d)}^p w(t) dt \\ &\quad + N(R) \left(\|u\|_{\mathbf{A}_{p,w}^{\gamma+1}(T)} + \|f\|_{\mathbf{A}_{p,w}^\gamma(T)} + \|u\|_{L_p((0,T), w dt; L_\infty(\mathbb{R}^d))} \right). \end{aligned}$$

By taking sufficiently small $R = R_0 \in (0, 1)$ such that $N A_\gamma(R_0) < 1$ (N is independent of R), we have

$$\begin{aligned} &\int_0^T \sup_{y \in \mathbb{R}^d} \|v^y(t, \cdot)\|_{\Lambda^{\gamma+2}(\mathbb{R}^d)} w(t) dt \\ &\leq N(R_0) \left(\|u\|_{\mathbf{A}_{p,w}^{\gamma+1}(T)} + \|f\|_{\mathbf{A}_{p,w}^\gamma(T)} + \|u\|_{L_p((0,T), w dt; L_\infty(\mathbb{R}^d))} \right). \end{aligned} \quad (3.32)$$

Since by Lemma 2.3,

$$\begin{aligned} \|u\|_{\mathbf{A}_{p,w}^{\gamma+2}(T)}^p &= \int_0^T \|u(t, \cdot)\|_{\Lambda^{\gamma+2}(\mathbb{R}^d)}^p w(t) dt \\ &\leq N(R_0) \int_0^T \sup_{y \in \mathbb{R}^d} \|v^y(t, \cdot)\|_{\Lambda^{\gamma+2}(\mathbb{R}^d)}^p w(t) dt \end{aligned}$$

one can deduce that

$$\|u\|_{\mathbf{A}_{p,w}^{\gamma+2}(T)} \leq N(R_0) \left(\|u\|_{\mathbf{A}_{p,w}^{\gamma+1}(T)} + \|f\|_{\mathbf{A}_{p,w}^\gamma(T)} + \|u\|_{L_p((0,T), w dt; L_\infty(\mathbb{R}^d))} \right). \quad (3.33)$$

when $p \in (1, \infty)$. Similarly, when $p = \infty$, by using

$$\|u\|_{\mathbf{A}_{p,w}^{\gamma+2}(T)} = \sup_{t \in (0,T)} \|u(t, \cdot)\|_{\Lambda^{\gamma+2}(\mathbb{R}^d)} \leq N(R_0) \sup_{t \in (0,T)} \sup_{y \in \mathbb{R}^d} \|v^y(t, \cdot)\|_{\Lambda^{\gamma+2}(\mathbb{R}^d)},$$

one can still obtain (3.33) with $p = \infty$ (and $w \equiv 1$). Using (3.27) and taking sufficiently small $\varepsilon \in (0, 1)$ such that $N(R_0)\varepsilon < 1$ yields the result for $\gamma \in (0, 1]$.

The general case $\gamma \in (n-1, n]$ follows by induction using Proposition 2.1-(ii). Assume that (3.26) holds when $\tilde{\gamma} \in (0, n-1]$ with $n \geq 2$. Proposition 2.1-(ii) yields $\pi_k := D_k u \in \mathbf{H}_{p,w}^{\gamma+1}(T)$ and π_k satisfies

$$\partial_t \pi_k = \sum_{i,j=1}^d a_{ij} D_{ij} \pi_k + D_k f + \sum_{i,j=1}^d D_k a_{ij} D_{ij} u.$$

By induction hypothesis with $\gamma - 1 \in (0, n-1]$, Proposition 2.1-(ii) and Lemma 2.5,

$$\begin{aligned} \|\pi_k\|_{\Lambda_{p,w}^{\gamma+1}(T)} &\leq N \|\pi_k\|_{L_p((0,T),w \, dt; L_\infty(\mathbb{R}^d))} + N \|D_k f\|_{\Lambda_{p,w}^{\gamma-1}(T)} + N \sum_{i,j=1}^d \|D_k a_{ij} D_{ij} u\|_{\Lambda_{p,w}^{\gamma-1}(T)} \\ &\leq N \left(\|u\|_{\Lambda_{p,w}^{\gamma+1}(T)} + N \|f\|_{\Lambda_{p,w}^{\gamma}(T)} \right). \end{aligned}$$

Since (3.26) is true when $\gamma - 1 \in (0, n-1]$,

$$\|u\|_{\Lambda_{p,w}^{\gamma+1}(T)} \leq N \left(\|f\|_{\Lambda_{p,w}^{\gamma-1}(T)} + \|u\|_{L_p((0,T),w \, dt; L_\infty(\mathbb{R}^d))} \right).$$

Thus, we arrive at (3.26) with $\gamma \in (0, \infty)$.

Step 2. We now show

$$\|u\|_{L_p((0,T),w \, dt; L_\infty(\mathbb{R}^d))} \leq N \|f\|_{\Lambda_{p,w}^{\gamma}(T)},$$

where $N = N(d, \nu, K, p, K_0, T)$. Let $s_l := \frac{lT}{m}$ for $l = 0, \dots, m$, and take cut-off functions $\eta_l \in C^\infty(\mathbb{R})$ ($l \geq 1$) such that $\eta_l = 1$ for $t > s_l$, $\eta_l = 0$ for $t \leq s_{l-1}$, and $|\eta'_l| \leq 2m/T$. For $l = 0$, set $s_{-1} := 0$ and $\eta_0 \equiv 1$. Then $q^l(t, x) := u(t, x)\eta_l(t) \in \Lambda_{p,w}^{\gamma+2}(s_{l-1}, s_{l+1})$ satisfies

$$\partial_t q^l = \sum_{i,j=1}^d a_{ij} D_{ij} q^l + \sum_{i=1}^d b_i D_i q^l + cq^l + f\eta_l + u\partial_t \eta_l, \quad t \in (s_{l-1}, s_{l+1}),$$

with zero initial condition $q^l(s_{l-1}, \cdot) = 0$. By (3.26) from Step 1, for $l \geq 1$,

$$\begin{aligned} &\|\partial_t q^l\|_{\Lambda_{p,w}^{\gamma}(s_{l-1}, s_{l+1})} + \|q^l\|_{\Lambda_{p,w}^{\gamma+2}(s_{l-1}, s_{l+1})} \\ &\leq N \|f\eta_l\|_{\Lambda_{p,w}^{\gamma}(s_{l-1}, s_{l+1})} + N \|u\partial_t \eta_l\|_{\Lambda_{p,w}^{\gamma}(s_{l-1}, s_{l+1})} + N \|q^l\|_{L_p((s_{l-1}, s_{l+1}), w \, dt; L_\infty(\mathbb{R}^d))} \\ &\leq N \|f\|_{\Lambda_{p,w}^{\gamma}(T)} + N \frac{m}{T} \|u\|_{\Lambda_{p,w}^{\gamma}(s_l)} + N \|u\|_{L_p((s_{l-1}, s_l), w \, dt; L_\infty(\mathbb{R}^d))} \\ &\quad + N \|u\|_{L_p((s_l, s_{l+1}), w \, dt; L_\infty(\mathbb{R}^d))} \end{aligned}$$

For $l = 0$, since $\partial_t \eta_0 = 0$,

$$\|u\|_{\Lambda_{p,w}^{\gamma+2}(s_1)} \leq N \|f\|_{\Lambda_{p,w}^{\gamma}(T)} + N \|u\|_{L_p((0, s_1), w \, dt; L_\infty(\mathbb{R}^d))}.$$

Here, we remark that, although the analysis is carried out on smaller time intervals, the constants N can be regarded as depending on T . Now apply Lemma 3.3 and Step 1 to get, for $l \geq 1$,

$$\begin{aligned} \|u\|_{L_p((s_l, s_{l+1}), w \, dt; L_\infty(\mathbb{R}^d))} &\leq \|q^l\|_{L_p((s_{l-1}, s_{l+1}), w \, dt; L_\infty(\mathbb{R}^d))} \\ &\leq N \frac{T}{m} \|\partial_t q^l\|_{L_p((s_{l-1}, s_{l+1}), w \, dt; L_\infty(\mathbb{R}^d))} \\ &\leq N \frac{T}{m} \|f\|_{\Lambda_{p,w}^{\gamma}(T)} + N \frac{T}{m} \|u\|_{L_p((s_{l-1}, s_l), w \, dt; L_\infty(\mathbb{R}^d))} \\ &\quad + N \frac{T}{m} \|u\|_{L_p((s_l, s_{l+1}), w \, dt; L_\infty(\mathbb{R}^d))} + N \|u\|_{\Lambda_{p,w}^{\gamma}(s_l)}, \end{aligned}$$

and for $l = 0$,

$$\|u\|_{L_p((0,s_1),w \, dt; L_\infty(\mathbb{R}^d))} \leq N \frac{T}{m} \left(\|f\|_{\Lambda_{p,w}^\gamma(T)} + \|u\|_{L_p((0,s_1),w \, dt; L_\infty(\mathbb{R}^d))} \right).$$

Choosing m sufficiently large so that $N T/m < 1$, for $l \geq 1$,

$$\begin{aligned} \|u\|_{L_p((s_l, s_{l+1}), w \, dt; L_\infty(\mathbb{R}^d))} &\leq N \left(\|f\|_{\Lambda_{p,w}^\gamma(T)} + \|u\|_{\Lambda_{p,w}^\gamma(s_l)} \right) \\ &\leq N \left(\|f\|_{\Lambda_{p,w}^\gamma(T)} + \|u\|_{L_p((0,s_l),w \, dt; L_\infty(\mathbb{R}^d))} \right), \end{aligned}$$

and for $l = 0$,

$$\|u\|_{L_p((0,s_1),w \, dt; L_\infty(\mathbb{R}^d))} \leq N \|f\|_{\Lambda_{p,w}^\gamma(T)}.$$

Hence, by using induction, the desired result is obtained. Combining Steps 1 and 2, the lemma is proved. \square

Proof of Theorem 3.1. The uniqueness and the estimate follow directly from Lemma 3.4.

For the existence, we apply the method of continuity. For $\lambda \in [0, 1]$, define

$$\mathcal{L}^\lambda := \sum_{i,j=1}^d (\lambda a_{ij} + (1-\lambda)\delta_{ij}) D_{ij} + \lambda \sum_{i=1}^d b_i D_i + \lambda c.$$

Let S denote the set of all $\lambda \in [0, 1]$ such that for every $f \in \Lambda_{p,w}^\gamma(T)$, there exists a unique $u \in \Lambda_{p,w}^{\gamma+2}(T)$ solving

$$\partial_t u = \mathcal{L}^\lambda u + f \quad \text{with zero initial condition.} \quad (3.34)$$

We claim that $S = [0, 1]$.

Step 1. S is nonempty. By previously known results (see, e.g., [7, Theorem 1.6]), the equation corresponding to $\lambda = 0$ is solvable; hence $0 \in S$.

Step 2. Uniform openness of S . Assumption 1.5(γ) holds for \mathcal{L}^λ for all $\lambda \in [0, 1]$ with the same ellipticity constant ν and the same bound K on the coefficients. Thus Lemma 3.4 yields the a priori estimate

$$\|u\|_{\mathbf{H}_{p,w}^{\gamma+2}(T)} \leq N \|f\|_{\Lambda_{p,w}^\gamma(T)} \quad (3.35)$$

for every solution u of (3.34), with a constant $N = N(d, \gamma, K_0, K, \nu, p, T)$ that is independent of λ .

Fix $\lambda_0 \in S$, and let

$$\mathcal{T}_{\lambda_0} : f \mapsto u$$

denote the solution operator associated with (3.34) at $\lambda = \lambda_0$. Then (3.35) implies

$$\|\mathcal{T}_{\lambda_0} f\|_{\mathbf{H}_{p,w}^{\gamma+2}(T)} \leq N \|f\|_{\Lambda_{p,w}^\gamma(T)}, \quad \forall f \in \Lambda_{p,w}^\gamma(T), \quad (3.36)$$

so that \mathcal{T}_{λ_0} is bounded with operator norm at most N . Now fix $f \in \Lambda_{p,w}^\gamma(T)$ and $\lambda \in [0, 1]$. We rewrite (3.34) as a fixed point problem around λ_0 :

$$\partial_t u = \mathcal{L}^{\lambda_0} u + f + (\mathcal{L}^\lambda - \mathcal{L}^{\lambda_0})u.$$

Equivalently, $u = \mathcal{T}_{\lambda_0}(f + (\mathcal{L}^\lambda - \mathcal{L}^{\lambda_0})u)$. Define the map

$$\Phi_{\lambda_0, \lambda}(u) := \mathcal{T}_{\lambda_0}(f + (\mathcal{L}^\lambda - \mathcal{L}^{\lambda_0})u), \quad u \in \mathbf{H}_{p,w}^{\gamma+2}(T).$$

Then fixed points of $\Phi_{\lambda_0, \lambda}$ are precisely the solutions of (3.34) for the parameter λ . For $u_1, u_2 \in \mathbf{H}_{p,w}^{\gamma+2}(T)$ we have, by the linearity of \mathcal{T}_{λ_0} and (3.36),

$$\|\Phi_{\lambda_0, \lambda}(u_1) - \Phi_{\lambda_0, \lambda}(u_2)\|_{\mathbf{H}_{p,w}^{\gamma+2}(T)} \leq N \|(\mathcal{L}^\lambda - \mathcal{L}^{\lambda_0})(u_1 - u_2)\|_{\Lambda_{p,w}^\gamma(T)}. \quad (3.37)$$

Since the coefficients of \mathcal{L}^λ depend linearly on λ and are uniformly bounded in $\lambda \in [0, 1]$, there exists a constant $C_0 = C_0(d, \gamma, K, K_0, \nu, p, T) > 0$ such that

$$\|(\mathcal{L}^\lambda - \mathcal{L}^{\lambda_0})v\|_{\Lambda_{p,w}^\gamma(T)} \leq C_0 |\lambda - \lambda_0| \|v\|_{\mathbf{H}_{p,w}^{\gamma+2}(T)}, \quad \forall v \in \mathbf{H}_{p,w}^{\gamma+2}(T), \lambda, \lambda_0 \in [0, 1]. \quad (3.38)$$

Combining (3.37) and (3.38) gives

$$\|\Phi_{\lambda_0, \lambda}(u_1) - \Phi_{\lambda_0, \lambda}(u_2)\|_{\mathbf{H}_{p,w}^{\gamma+2}(T)} \leq NC_0|\lambda - \lambda_0| \|u_1 - u_2\|_{\mathbf{H}_{p,w}^{\gamma+2}(T)}.$$

Choose $\varepsilon_0 > 0$ such that $NC_0\varepsilon_0 < 1$, where $\varepsilon_0 = \varepsilon_0(d, \gamma, K_0, K, \nu, p, T)$ is independent of λ_0 . Then, for all $\lambda \in [0, 1]$ with $|\lambda - \lambda_0| < \varepsilon_0$, the map $\Phi_{\lambda_0, \lambda}$ is a contraction on $\mathbf{H}_{p,w}^{\gamma+2}(T)$. By the Banach fixed-point theorem, $\Phi_{\lambda_0, \lambda}$ admits a unique fixed point $u \in \mathbf{H}_{p,w}^{\gamma+2}(T)$, which solves (3.34). Hence,

$$(\lambda_0 - \varepsilon_0, \lambda_0 + \varepsilon_0) \cap [0, 1] \subset S.$$

Thus every point of S has a neighborhood of radius ε_0 contained in S , with ε_0 independent of the point.

Step 3. Covering argument. Since $0 \in S$ and $\varepsilon_0 > 0$ is fixed, we have $[0, \varepsilon_0] \cap [0, 1] \subset S$. Iterating this argument, we obtain

$$[0, k\varepsilon_0] \cap [0, 1] \subset S \quad \text{for each } k \in \mathbb{N}.$$

Choosing $k_0 \in \mathbb{N}$ with $k_0\varepsilon_0 \geq 1$ yields $[0, 1] \subset S$, hence $S = [0, 1]$.

This proves the existence of a unique solution $u \in \mathbf{H}_{p,w}^{\gamma+2}(T)$ to (3.34) for $\lambda = 1$. Together with the a priori estimate and uniqueness, this completes the proof. \square

3.2. Proof of Theorem 1.6 with non-zero initial data. We now turn to the case of non-zero initial data. The following trace theorem plays a crucial role in our analysis. It shows that $\mathbf{H}_{p,w}^{\gamma+2}(T)$ admits a well-defined and continuous trace at $t = 0$, and moreover that every element of the space $\Lambda_p^{\gamma+2,w}(\mathbb{R}^d)$ arises as such a trace. In particular, this identifies $\Lambda_p^{\gamma+2,w}(\mathbb{R}^d)$ as the *optimal* initial data space for the regularity class $\mathbf{H}_{p,w}^{\gamma+2}(T)$, and provides the exact mechanism by which the non-zero initial value problem can be reduced to the zero initial data case.

Theorem 3.5 (Trace theorem). *Let $\eta \in (0, \infty)$, $p \in (1, \infty]$, $T > 0$, and $w \in A_p$ with $[w]_{A_p} \leq K_0$.*

(i) The space $\mathbf{H}_{p,w}^{\eta+2}(T)$ is continuously embedded into $C([0, T]; \Lambda_p^{\eta+2,w}(\mathbb{R}^d))$; that is,

$$\sup_{t \in [0, T]} \|u(t, \cdot)\|_{\Lambda_p^{\eta+2,w}(\mathbb{R}^d)} \leq N \|u\|_{\mathbf{H}_{p,w}^{\eta+2}(T)}.$$

(ii) For every $u_0 \in \Lambda_p^{\eta+2,w}(\mathbb{R}^d)$, there exists $v \in \mathbf{H}_{p,w}^{\eta+2}(T)$ satisfying $v(0, \cdot) = u_0$, and

$$\|v\|_{\mathbf{H}_{p,w}^{\eta+2}(T)} \leq N \|u_0\|_{\Lambda_p^{\eta+2,w}(\mathbb{R}^d)}. \quad (3.39)$$

Here N depends on K_0, p , and T .

Proof. Since the results for the case $p = \infty$ follow directly from the definition of $\mathbf{H}_{p,w}^{\eta+2}(T)$, we focus on the case $p \in (1, \infty)$. By [7, Lemma 3.1] with $\phi(\lambda) = \lambda^\eta$ and $\gamma = 2$, we have

$$(\Lambda^{\eta+2}(\mathbb{R}^d), \Lambda^\eta(\mathbb{R}^d))_{W^{1/p}, p} = \Lambda_p^{\eta+2,w}(\mathbb{R}^d), \quad (3.40)$$

where $W(t) := \int_0^t w(s) \, ds$, and $(X, Y)_{W^{1/p}, p}$ denotes the generalized real interpolation space (see [9]). With the characterization of the generalized real interpolation space (3.40), (i) and (ii) follow directly from [7, Theorem 1.8] and [9, Theorem 1.5], respectively. This completes the proof of the theorem. \square

Proof of Theorem 1.6. The uniqueness follows directly from Theorem 3.1.

For the existence, we first obtain $v \in \mathbf{H}_{p,w}^{\gamma+2}(T)$ satisfying $v(0, \cdot) = u_0$ and (3.39) by applying Theorem 3.5-(ii). Then $\mathcal{L}v - \partial_t v \in \Lambda_p^\gamma(T)$, where

$$\mathcal{L}u := \sum_{i,j=1}^d a_{ij} D_{ij} u + \sum_{i=1}^d b_i D_i u + cu.$$

Hence, by Theorem 3.1, there exists a unique solution $\bar{u} \in \mathbf{H}_{p,w}^{\gamma+2}(T)$ to

$$\partial_t \bar{u} = \mathcal{L}\bar{u} + (f + \mathcal{L}v - \partial_t v), \quad \text{with zero initial condition.}$$

Let $u := \bar{u} + v$. Then u is a solution to

$$\partial_t u = \mathcal{L}u + f, \quad u(0, \cdot) = u_0.$$

The desired estimates follow directly from Theorems 3.1 and 3.5. This completes the proof of the theorem. \square

REFERENCES

- [1] H. Bahouri, J.-Y. Chemin, and R. Danchin. *Fourier analysis and nonlinear partial differential equations*, volume 343 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer, Heidelberg, 2011.
- [2] M. Ben-Artzi. Global solutions of two-dimensional Navier-Stokes and Euler equations. *Arch. Rational Mech. Anal.*, 128(4):329–358, 1994.
- [3] J. Bergh and J. Löfström. *Interpolation spaces. An introduction*, volume No. 223 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, Berlin-New York, 1976.
- [4] S. Boccia and N. V. Krylov. On the fundamental matrix solution for higher-order parabolic systems. In *Proceedings of the St. Petersburg Mathematical Society. Vol. XV. Advances in mathematical analysis of partial differential equations*, volume 232 of *Amer. Math. Soc. Transl. Ser. 2*, pages 11–33. Amer. Math. Soc., Providence, RI, 2014.
- [5] A. Brandt. Interior Schauder estimates for parabolic differential- (or difference-) equations via the maximum principle. *Israel J. Math.*, 7:254–262, 1969.
- [6] H. Brezis. Remarks on the preceding paper by M. Ben-Artzi: “Global solutions of two-dimensional Navier-Stokes and Euler equations” [Arch. Rational Mech. Anal. **128** (1994), no. 4, 329–358]. *Arch. Rational Mech. Anal.*, 128(4):359–360, 1994.
- [7] J.-H. Choi. A regularity theory for evolution equations with time-measurable pseudo-differential operators in weighted mixed-norm Sobolev-Lipschitz spaces. *Potential Anal.*, 63(2):557–577, 2025.
- [8] J.-H. Choi, J. B. Lee, J. Seo, and K. Woo. Characterizations of weighted besov and triebel-lizorkin spaces with variable smoothness. *arXiv preprint arXiv:2309.01359 (To appear in Communications on Pure and Applied Analysis, 10.3934/cpaa.2026007)*, 2023.
- [9] J.-H. Choi, J. B. Lee, J. Seo, and K. Woo. On the trace theorem to volterra-type equations with local or non-local derivatives. *arXiv preprint arXiv:2309.00370 (To appear in Journal of Evolution Equations)*, 2023.
- [10] H. Dong and D. Kim. On L_p -estimates for elliptic and parabolic equations with A_p weights. *Trans. Amer. Math. Soc.*, 370(7):5081–5130, 2018.
- [11] H. Dong and S. Kim. Partial Schauder estimates for second-order elliptic and parabolic equations. *Calc. Var. Partial Differential Equations*, 40(3-4):481–500, 2011.
- [12] H. Dong and S. Kim. Partial Schauder estimates for second-order elliptic and parabolic equations: a revisit. *Int. Math. Res. Not. IMRN*, 2019(7):2085–2136, 2019.
- [13] H. Dong and H. Zhang. On Schauder estimates for a class of nonlocal fully nonlinear parabolic equations. *Calc. Var. Partial Differential Equations*, 58(2):Paper No. 40, 42, 2019.
- [14] X. Fernández-Real and X. Ros-Oton. *Regularity theory for elliptic PDE*, volume 28 of *Zurich Lectures in Advanced Mathematics*. EMS Press, Berlin, 2022.
- [15] I. Kim. An L_p -Lipschitz theory for parabolic equations with time measurable pseudo-differential operators. *Commun. Pure Appl. Anal.*, 17(6):2751–2771, 2018.
- [16] N. V. Krylov. *Lectures on elliptic and parabolic equations in Hölder spaces*, volume 12. American Mathematical Soc., 1996.
- [17] N. V. Krylov. The Calderón-Zygmund theorem and parabolic equations in $L_p(\mathbb{R}, C^{2+\alpha})$ -spaces. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 1(4):799–820, 2002.
- [18] N. V. Krylov. Parabolic equations in L_p -spaces with mixed norms. *Algebra i Analiz*, 14(4):91–106, 2002.
- [19] F. Kühn. Schauder estimates for Poisson equations associated with non-local Feller generators. *J. Theoret. Probab.*, 34(3):1506–1578, 2021.
- [20] L. Lorenzi. Optimal Schauder estimates for parabolic problems with data measurable with respect to time. *SIAM J. Math. Anal.*, 32(3):588–615, 2000.
- [21] A. Palczewski. Partial Schauder estimates for unbounded solutions of the Kolmogorov equation. *J. Elliptic Parabol. Equ.*, 11(2):615–641, 2025.
- [22] P. R. Stinga and J. L. Torrea. Hölder, Sobolev, weak-type and BMO estimates in mixed-norm with weights for parabolic equations. *Sci. China Math.*, 64(1):129–154, 2021.
- [23] H. Triebel. *Theory of function spaces*, volume 78 of *Monographs in Mathematics*. Birkhäuser Verlag, Basel, 1983.
- [24] J. Wei, W. Wang, G. Lv, and J. Duan. A new maximal regularity for parabolic equations and an application. *arXiv preprint arXiv:2411.13266*, 2024.

SCHOOL OF MATHEMATICS, KOREA INSTITUTE FOR ADVANCED STUDY, 85 HOEGI-RO, DONGDAEMUN-GU, SEOUL,
02455, REPUBLIC OF KOREA

Email address: jhchoi@kias.re.kr

SCHOOL OF MATHEMATICS, KOREA INSTITUTE FOR ADVANCED STUDY, 85 HOEGI-RO, DONGDAEMUN-GU, SEOUL,
02455, REPUBLIC OF KOREA

Email address: junhryu@kias.re.kr