

On the MLC Conjecture and the Renormalization Theory in Complex Dynamics

Dzmitry Dudko*

Abstract. In this Note, we present recent developments in the Renormalization Theory of quadratic polynomials and discuss their applications, with an emphasis on the MLC conjecture, the problem of local connectivity of the Mandelbrot set, and on its geometric counterparts.

1 Introduction. The Mandelbrot set \mathcal{M} (Figure 1) is one of the most recognizable fractals in mathematics and beyond. It encodes the dynamical dependence of quadratic polynomials on the parameter. A central question in Complex Dynamics is the MLC Conjecture, also known as the *Rigidity Conjecture*:

MLC Conjecture 1.1. The Mandelbrot set \mathcal{M} is locally connected.

This conjecture has various geometric and probabilistic counterparts; all of them are subjects of the Renormalization Theory of \mathcal{M} , which analyzes *first return maps* to small neighborhoods of special points. Nature and the classification of these first return maps is in focus of this Note. Let us start by outlining its context.

In Section 1 below, we give an informal introduction to renormalization ideas. Although the MLC is a topological statement, any substantial progress typically requires geometric input. Conjecture 1.2 states a simple geometric property of hyperbolic components that easily implies the MLC. A foundation of any renormalization theory is *a priori* bounds, see Conjecture 1.3: as we zoom into the Julia set and endow it with the first return map, we expect precompactness. A similar precompactness is expected in the parameter plane, see §1.4. In §1.3, we classify (sequences of) first return maps as either *renorm-expanding* or *renorm-balanced*. The exposition in Section 1 is conjectural, but it has been justified in many cases.

Historical background is presented in Section 2, where we also sketch the evolution of key ideas and briefly comment on the higher-degree cases. The development of the renormalization theory can be broadly divided into several stages. First, as already mentioned, initial *a priori* bounds are required. These are used in justifying various rigidity properties, e.g., the MLC. Next, a detailed geometric analysis of the convergence of first-return maps follows; a desired goal is to establish the *hyperbolicity* of the associated operator. Finally, deep applications can be derived; for instance, the “regular vs stochastic” dichotomy describing the *nature of chaos*.

Section 3 is dedicated to the theory of renormalization operators acting on infinite-dimensional spaces of maps. We recognize three types of renormalization regimes: *near-Neutral*, if some hyperbolic components become unbounded in the respective (small) parameter neighborhoods relative to the scale, *Quadratic-Like*, in the presence of intermediate hyperbolic components relative to the scale, and *Puzzle* (it is intertwined with the renorm-expanding notion), if all hyperbolic components become negligible relative to the scale. We outline the quadratic-like (“ql”) renormalization theory in its pure, bounded-type form. Its hyperbolicity program is now complete by Theorem 3.3, thanks to Theorem 4.1. We summarize the status of the near-Neutral Case and mention how Theorem 4.2 provides an opening towards the full hyperbolicity of the Neutral Renormalization and beyond, see §3.8. We will also discuss the puzzle regime in §3.9 and put forward Problem 3.1 on whether all these three regimes can be neatly integrated.

In Section 4, on the Near-Degenerate Regime, we illustrate the sources of *a priori* bounds on dynamical scales representing definite positive-entropy “ $\geq \varepsilon > 0$ ” (primitive ql and puzzle cases that are “ ε -away” from the main Molecule) and zero-entropy (Molecule: neutral and bounded-type satellite ql) settings. We conclude by summarizing the current status of the MLC as two problems: completing the remaining unbounded satellite cases and developing interpolation techniques between definite positive and zero entropy settings. The former relies on

*Department of Mathematics, Stony Brook University, USA




Figure 1: The Mandelbrot set (right) and the period-doubling cascade (left, colored in blue). The period-doubling Feigenbaum parameter $\lambda_* \simeq c_*$ marks the transition from zero entropy to positive entropy; it is where the “chaos” emerges. This transition is universal; its complexification is shown in Figure 2.

further development of the theory behind *almost-invariant pseudo-Siegel disks*, Figure 14. For the latter, we put forward a strategy based on partially invariant *virtual Julia sets* which control only *partial postcritical sets*. If these programs are achieved, they would establish *a priori* bounds for *all* quadratic polynomials and open a clear path towards the MLC (rigidity) and its geometric and probabilistic counterparts discussed in this Note.

1.1 Renormalization: summary and motivation. The renormalization $\mathcal{R}_{\eta}p_c$ of a quadratic polynomial $p_c: z \mapsto z^2 + c$ can be defined as the rescaled first return map to a small η -neighborhood of the critical value, see (1.2). A single iterate of $\mathcal{R}_{\eta}p_c$ controls a long finite orbit of p_c . Hence, instead of iterating p_c , we wish to iterate renormalization

$$(1.1) \quad p_c \rightsquigarrow \mathcal{R}_{\eta_1}p_c \rightsquigarrow \mathcal{R}_{\eta_2}p_c \rightsquigarrow \dots \rightsquigarrow \mathcal{R}_{\eta_n}p_c \rightsquigarrow \dots \xrightarrow{\text{“lim”}}$$

and then deduce strong dynamical consequences for p_c . To realize such a vision, initial *a priori* bounds (precompactness conditions) are required, see Conjecture 1.3.

We have two cases. If the derivative $[\mathcal{R}_{\eta}p_c]'(z)$ typically grows, then we should expect an almost linear phase-parameter relationship between \mathcal{M} and J_c at c as illustrated in Figure 8. We call this regime *renorm-expanding*, see §1.3; it is controlled by *puzzles*. On the other hand, if the derivative $[\mathcal{R}_{\eta}p_c]'(z)$ is bounded, then there should be geometric limits for the renormalization, see (1.3). We refer to this regime as *renorm-balanced*.

1.2 Hyperbolic components. For a hyperbolic component $\mathcal{H} \subset \mathcal{M}$, we denote by $\mathcal{M}_{\mathcal{H}}$ the associated small copy of \mathcal{M} centered at \mathcal{H} . We recall that \mathcal{H} consists of parameters with an attracting periodic cycle and can be of two types, see Figure 3: satellite, when $\bar{\mathcal{H}}$ is an analytic disk, and primitive, when $\bar{\mathcal{H}}$ is an analytic image of the filled cardioid. On pictures, hyperbolic components always have a flawless shape.

Conjecture 1.2 (Bounded geometry of $\mathcal{H} \subset \mathcal{M}_{\mathcal{H}}$). Every closed hyperbolic component $\bar{\mathcal{H}}$ is a uniform qc disk or a uniform qc filled cardioid; its diameter $\text{diam}(\mathcal{H})$ is uniformly comparable to $\text{diam}(\mathcal{M}_{\mathcal{H}})$.

Conjecture 1.2 easily implies the MLC: if $\mathcal{M}_{\mathcal{H}_n}$ is a sequence of strictly nested small copies of \mathcal{M} , then the diameter of $\mathcal{M}_{\mathcal{H}_n}$ shrinks because of finiteness of the area:

$$\sum_n [\text{diam}(\mathcal{M}_{\mathcal{H}_n})]^2 \sim \sum_n [\text{diam}(\mathcal{H}_n)]^2 \sim \sum_n \text{area}(\mathcal{H}_n) < \infty, \quad \text{thus } \bigcap_{n \geq 1} \mathcal{M}_{\mathcal{H}_n} \text{ is a singleton.}$$

This implies the MLC by Yoccoz’s results, see (2.1). For primitive \mathcal{H} , Conjecture 1.2 is expected to follow from *a priori* bounds of ql type §3.2; cf. Conjecture 1.3 below. For satellite \mathcal{H} , the hyperbolicity of the Molecule renormalization is likely necessary, see Conjecture 3.6.

1.3 Dynamical limits. Let us consider the dynamical plane of p_c for a *non-parabolic* $c \in \partial\mathcal{M}$. (Renormalizations for parabolic $c \in \partial\mathcal{M}$ arise through limits $c_n \rightarrow c$, see §3.5.) For a small $\eta > 0$ and the




Figure 2: The universality of the Mandelbrot set at the period-doubling Feigenbaum parameter c_* , see also Figure 1. The renormalization theory justifying this universality has been successfully developed in the 1990s, see §2. The MLC at c_* has been recently established in Theorem 4.1.

associated η -neighborhood of the critical value c , the *renormalization* $\mathcal{R}_\eta p_c$ of p_c is the *rescaled first return map* of p_c from this η -neighborhood back to itself; we write:

$$(1.2) \quad [\mathcal{R}_\eta p_c](z) = \left[(p_c^q)^{\eta\text{-rescaled}} \right](z) = \frac{1}{\eta} \left(p_c^q(\eta z + c) - c \right), \quad p_c(z) = z^2 + c,$$

where $q \equiv q(\eta z + c)$ is the first return time of the point $\eta z + c$ back to the η -neighborhood of c . The η -rescaling identifies the η -neighborhood of c with the 1-neighborhood of 0.

Conjecture 1.3 (Uniform *a priori* bounds). With appropriate understanding, the family of renormalizations (1.2) is precompact uniformly over all $\eta > 0$ and $c \in \partial\mathcal{M}$.

In particular, as we zoom into the Julia set J_c of p_c at c , the shape of J_c (e.g., its veins) should stay within a precompact class of configurations. Let us detail more what we expect from Conjecture 1.3.

In (1.2), different points can have different return times; and some points may not return to the selected neighborhood. Consider an iterate p_c^q realizing a first return of some point in the η -neighborhood of c back to the neighborhood. Rescaling p_c^q , we obtain a *globalized branch* $\mathcal{R}_\eta^{(q)} p_c \equiv (p_c^q)^{\eta\text{-rescaled}}: \mathbb{C} \rightarrow \mathbb{C}$ of $\mathcal{R}_\eta p_c$.

Consider now a sequence $\eta_n \rightarrow 0$ and the associated renormalizations $\mathcal{R}_{\eta_n} p_c$. We refer to the η_n 's as *dynamical scales*. Up to passing to a subsequence, we anticipate only two possibilities:

(Exp) *renorm-expanding* case: for all choices of branches, the derivative $[\mathcal{R}_{\eta_n}^{(q_n)} p_c]'(z)$ tends to infinity for a typical (roughly, non-critical) $z \in \mathbb{C}$; or

(Ren) *renorm-balanced* case: for some choices of globalized branches, there is a well-defined geometric limit, transcendental or polynomial (locally uniformly on U_G):

$$(1.3) \quad \exists [G: U_G \rightarrow \mathbb{C}] = \lim_{n \rightarrow \infty} [\mathcal{R}_{\eta_n}^{(q_n)} p_c: \mathbb{C} \rightarrow \mathbb{C}], \quad \eta_n \rightarrow 0.$$

We anticipate that the renorm-balanced case (Ren) occurs when certain hyperbolic components near c do not shrink in the respective parameter scales (see §1.4), so that the periodic cycles associated with such *non-negligible* hyperbolic components prevent expansion; cf. Conjecture 1.2.

By considering all sequences of dynamical scales $\eta_n \rightarrow 0$, we say that $c \in \mathcal{M}$ is

- *pure renorm-expanding*, if it is renorm-expanding with respect to any $\eta_n \rightarrow 0$; and
- *pure renorm-balanced*, if it is renorm-balanced with respect to any $\eta_n \rightarrow 0$.




Figure 3: Primitive and satellite hyperbolic components. Around every hyperbolic component, there is an associated primitive or satellite little copy of the Mandelbrot set. It is expected, cf. Conjecture 1.2, that all hyperbolic components have uniform geometry. The difference between the satellite (zero entropy) and primitive (positive entropy) cases is profound and has many implications; e.g., see Figures 12 and 13.

There are four main renormalization theories in the pure renorm-balanced case: primitive/satellite quadratic-like, near-parabolic, and near-Siegel; the last two are subtypes of near-neutral. These theories have many similarities as well as certain fundamental differences. The general case is a combination of all the above regimes, where various dynamical scales of p_c can exhibit different renormalization flavors; see Section 3 and Question 3.1.

1.4 Parameter limits. Recall [10] that the parameter and dynamical Böttcher functions provide conformal identification of $\widehat{\mathbb{C}} \setminus \mathcal{M}$ and $\widehat{\mathbb{C}} \setminus J_c$. Therefore, dynamical scales η_n as in §1.3 have associated parameter scales δ_n defined up to appropriate choices. Let us write the parameter counterpart to (1.2):

$$(1.4) \quad \mathcal{R}_{(\delta, \eta)} \{p_{c+\tau}\}_{\tau \in \mathbb{C}} = \left(\{p_{c+\tau}\}_{\tau \in \mathbb{C}} \right)^{(\delta, \eta)\text{-rescaled}} = \frac{1}{\eta} \left(p_{c+\delta\tau}^q(\eta z + c + \delta\tau) - c - \delta\tau \right),$$

i.e., all nearby maps $p_{c+\delta\tau}$ are η -rescaled with respect to the same return times as p_c , while the parameter dependence is δ -rescaled. The notion of branches is defined accordingly.

Similar to Conjecture 1.3, we expect the precompactness of (1.4) to hold uniformly over all c and across all scales. In fact, we anticipate much stronger properties: convergence toward the limits should be exponentially fast, under an appropriate understanding, both in the parameter and dynamical planes, see §3. Such statements are commonly referred to as *universality* and are illustrated in various figures throughout this Note.

2 Historical Perspective. The problem of local connectivity of the Mandelbrot set \mathcal{M} , known as the MLC Conjecture, was put forward in the 1980s by Douady and Hubbard, who also developed the foundational theory of \mathcal{M} , [10, 11]. It was the time when the first computer pictures of \mathcal{M} were generated; these pictures, ever since, have been providing invaluable guidance in research.

Yoccoz' results from the early 1990s reduced the MLC Conjecture to the following rigidity statement [23]:

$$(2.1) \quad \text{if } \mathcal{M}_1 \supsetneq \mathcal{M}_2 \supsetneq \dots \supsetneq \mathcal{M}_n \supsetneq \dots \quad \text{are nested small copies of } \mathcal{M},$$

$$\text{then } \bigcap_{n=1}^{\infty} \mathcal{M}_n = \{c\} \quad \text{is a singleton.}$$

This linked the MLC to the renormalization theory.

Before the Mandelbrot set and its pictures, there has been the *Logistic Family* which encodes the *real vein* $\mathcal{M} \cap \mathbb{R}$ of the Mandelbrot set:

$$(2.2) \quad x \mapsto \lambda x(1-x) : [0, 1] \hookrightarrow, \quad \lambda \in [0, 4].$$

The discovery of *chaos* is due to Poincaré (sensitive dependence combined with recurrence in the 3-body problem, the late 19th century). In an attempt to understand this phenomenon, much of the research activity in the second half of the 20th century was focused on the study of the Logistic Family (e.g., Ulam, Neumann in the 1940s,




Figure 4: Universality of the Mandelbrot set at the golden-mean Siegel parameter c_\bullet . The dynamical universality of $z^2 + c_\bullet$ is depicted in Figure 5. Theorem 3.4 completed the program towards the hyperbolicity of the near-Siegel renormalization of bounded-type.

Lorenz, Myrberg, Sharkovsky in the 1960s, Misiurewicz, Guckenheimer in the 1970s). The Combinatorial Theory of (2.2) was designed by Milnor and Thurston in the late 1970s in terms of kneading sequences.

Fifty years ago, the *period-doubling universality* in (2.2) was discovered by Feigenbaum [21] (the parameter universality) and, independently, by Tresser and Collet [9] (the dynamical universality); see Figures 1 and 2. This was the beginning of the Renormalization Theory in Dynamics; many renormalization ideas have been imported from Quantum Field Theory and Statistical Mechanics. The first computer-assisted proof of the period-doubling universality was presented by Lanford [33] in the early 1980s. A computer-free proof, with numerous extensions to other cases, was only achieved in the late 1990s using *complex* methods.

In his address to the ICM in 1986 [47], Sullivan proposed a program to approach the period-doubling universality based upon Teichmüller theory by employing the Douady-Hubbard quadratic-like theory. Sullivan partially realized this program in [48] by constructing a renormalization fixed point and justifying (non-exponential) convergence towards the fixed point within its hybrid class. The exponential convergence (the dynamical universality) was justified by McMullen in [41]. The program was completed by Lyubich in [36] who established the parameter universality.

Meanwhile, a *complete* renormalization theory has been successfully developed for the Logistic Family [40, 35, 22], culminating in the “regular vs stochastic” dichotomy [38] by Lyubich; i.e., the Logistic Family became the first natural class of maps outside of the hyperbolic setting for which *chaos was well understood*. Later, the dichotomy was refined to “regular vs Collet-Eckmann” by Avila and Moreira in [4].

In its foundation, the success with the Logistic Family relied on *real a priori* bounds which are unavailable for genuine complex maps. Moreover, various *near-Neutral* phenomena are not witnessed by the Logistic Family. For instance, there are Julia sets that are *not* Locally Connected – the *non-JLC Category*. Even if the Julia set of a map is locally connected, the map can have many renormalization “near non-JLC” levels where ql-bounds fail.

Interest in the near-Neutral/Rotational Dynamics is rooted in the classical Celestial Mechanics. The first breakthroughs in the linearization of circle diffeomorphisms and local maps were made by Denjoy (1932) and Siegel (1942). In the 1950s and 60s, the Kolmogorov-Arnold-Moser (KAM) theory emerged, resulting in the reenvision of the near-rotation phenomenon in Mathematics and Physics. The Renormalization Ideas introduced in the mid-1970s led, in particular, to numerous conjectures in Low-Dimensional Dynamics, including the nature of the KAM small divisor problem. The theory of analytic circle diffeomorphisms and *Local Theory* for neutral holomorphic germs and quadratic polynomials received an essentially complete treatment in the second half of the last century in the work by Arnold (in the KAM framework), Cherry, Bruno, Herman, Yoccoz, and Perez-Marco.

For quadratic polynomials, the *Neutral Family*

$$(2.3) \quad z \mapsto e^{2\pi i \theta} z + z^2 : \mathbb{C} \curvearrowright, \quad \theta \in \mathbb{R}/\mathbb{Z},$$

encodes the boundary of the main hyperbolic component of \mathcal{M} . There are two renormalization theories that




Figure 5: The universality of the Golden Siegel map $z^2 + c_\bullet$, where the parameter c_\bullet is depicted in Figure 4. The source of *a priori* bounds for the respective renormalization theory is the Douady-Ghys qc surgery; it transfers *real a priori* bounds from Blaschke maps to complex Siegel polynomials of bounded-type.

analyze the *Global Theory* of the Neutral Family.

Renormalization Theory of (bounded-type) Siegel maps, also initiated by physicists, was mathematically designed by McMullen in [42] in the mid-1990s. He established the dynamical universality as shown in Figure 5. The parameter counterpart, Figure 4, was developed two decades later in [20, 17], using the framework of the pacman renormalization. Theorem 3.4 states the hyperbolicity of the associated renormalization; its proof employs Transcendental Dynamics §3.7.

The significance of the near-parabolic renormalization (see Figures 6 and 7) has been demonstrated by Shishikura in the 1990s in [43], where he showed that the Hausdorff dimension of $\partial\mathcal{M}$ is equal to 2. (The area of $\partial\mathcal{M}$ is expected to be 0, see §3.8.) The argument in [43] involved two iterates of the near-parabolic renormalization. Control of *infinite iterates* became available in the mid-2000s when Inou and Shishikura established the hyperbolicity of the near-parabolic renormalization. New *sectorial a priori* bounds, see Figure 10, were put forward and justified in the perturbative regime; these were the first bounds in the *non-JLC Category*. Many applications followed. For instance, the first examples of positive area Julia sets were constructed by Buff and Chéritat in [6], and then, of a different type, by Avila and Lyubich [2]. In the mid-2010s, exact models of the postcritical sets were produced for neutral maps (Siegel and Cremer) [44, 8] in the Inou-Shishikura class.

Many global properties of the Neutral Family (2.3) are closely linked to those of unicritical circle maps; cf. the Douady-Ghys qc-surgery in Figure 5. The Renormalization Theory of unicritical circle maps was fully developed (up to uniform hyperbolicity) in the works of de Faria, de Melo, and Yampolsky by the early 2000s.

In the mid-2000s, motivated by W. Thurston's work on 3-manifolds and postcritically finite rational maps [50, 12], Kahn introduced the Near-Degenerate regime to Renormalization theory [25]. Jointly with Lyubich, they developed a machinery [30, 27, 28] that established quadratic-like *a priori* bounds for ql-combinatorics of positive core-entropy $\geq \varepsilon > 0$. It became apparent that this new theory provides an adequate replacement for the framework of *real a priori* bounds in the genuine complex setting. However, at that time, the theory could not handle the zero-entropy setting (e.g., the non-JLC Category), and its development halted for more than a decade.

The Near-Degenerate regime in the zero-entropy setting was designed in [16], see Theorem 4.2, using the framework of *almost-invariant* pseudo-Siegel disks. Theorem 4.2 established *a priori* bounds for all maps in the Neutral Family (2.3); in particular, providing the first *non-perturbative a priori* bounds in the non-JLC Category. Various ideas behind [16] shaped follow-up research activities. In particular, Theorem 4.1, [18] has finally confirmed the MLC at the classical period-doubling Feigenbaum parameter shown in Figures 1 and 2.

Advances in the renormalization theory of the Mandelbrot set have often provided guidance for more general settings. In the 2000s, various deep results for the Logistic Family have been extended to more general classes of real analytic maps; see [3, 32, 31]. For higher-degree polynomials, the Fibonacci-type renormalization discovered in [39] does not degenerate; employing the real bounds, this renormalization was successfully understood,




Figure 6: Parabolic Enrichment. Top: the Elephant structure in the parameter plane. Bottom: dynamical planes $p_{1/4}, p_{a_n}, p_{b_n}$. As $n \rightarrow \infty$, the pair $(p_{a_n}, p_{a_n}^{q_{1,n}})$ converges to the commuting pair $(p_{1/4}, p_{a_\infty}^{[1]})$, see Figure 7, where $p_{a_\infty}^{[1]}$ is the associated Lavaurs transcendental map. Similarly, the neutral pairs $(p_{b_n}, p_{b_n}^{q_{1,n}})$ converge to $(p_{1/4}, p_{b_\infty}^{[1]})$. Lavaurs limits in the neutral case are efficiently encoded by continued fractions, see §3.4.

see [45, 46]. With the introduction of the Near-Degenerate Regime, the complex Fibonacci case also became amenable to control, leading to generalizations of Yoccoz-type results to higher-degree polynomials [29, 1]. Recently, in my joint work with Drach [13], Yoccoz-type realization and rigidity results were established for Transcendental Dynamics (for maps of controlled finite order and without asymptotic values) by developing a transcendental version of the near-degenerate analysis.

In the 2020s, the Near-Degenerate regime has also been introduced for rational maps. In my joint work with Luo [15], *a priori* bounds were established for hyperbolic components of disjoint type in the space of rational maps, leading to various realization results. This confirms McMullen's conjecture in the disjoint-type case from the 1990s, which asserts that Sierpiński hyperbolic components are bounded. Simultaneously, Kahn [26] presented an independent approach with a different scope to *a priori* bounds for establishing McMullen's conjecture (for all Sierpiński cases). Meanwhile, Lim [34] established bounds for rotating Herman curves of bounded-type rotation speed, leading to the development of the corresponding renormalization theory.

A popular version of the MLC story has been presented in a 2024 Quanta Magazine article [7].

3 Renormalization operators. Following the discovery of the period-doubling universality (Figure 1), the framework of analytic renormalization operators acting on infinite-dimensional spaces of maps was put forward. The hyperbolicity of such operators always yields important consequences. There are two kinds of hyperbolicity frameworks:

(QL) *Quadratic-Like* Renormalization acting on ql maps as illustrated in Figure 9, top; and

(Neut) *Near-Neutral* (Cylinder or Sectorial) Renormalization, see Figures 10 and 7.

In addition to the (QL) and (Neut) Cases, let us also recognize:

(Puz) *Puzzle* (or generalized ql) Renormalization encoding the first return map to a puzzle, Figure 9, bottom.

The (QL) bounded-type case is classical and now complete. The *bounded-type* condition refers to parameters c within nested intersection of small copies $\bigcap_n \mathcal{M}_n$ as in (2.1), so that the relative periods of \mathcal{M}_{n+1} rel \mathcal{M}_n




Figure 7: Top: as $a_n \rightarrow 1/4$, see Figure 6, the first return map $p_n^{q_{1,n}}$ converges to the transcendental Lavaurs map $p_{a_\infty}^{[1]}$ commuting with $p_{1/4}$. Bottom: the associated cylinder renormalization. See §3.5 for details.

are uniformly bounded. Equivalently, in the dynamical planes of p_c , for every dynamical scale η (as in §1.3), p_c has a quadratic-like structure $p_c^q: U' \rightarrow V'$ with a non-escaping critical orbit so that $|V'| \asymp \eta$. Similarly, the name ‘‘bounded-type’’ is used elsewhere in the renormalization theory, e.g., for neutral maps. In the late 1990s, the hyperbolicity problem in the bounded-type ql case was reduced to the justification of quadratic-like bounds. They have been recently supplied by Theorem 4.1, completing the program; see Theorem 3.3.

The hyperbolicity theory for neutral maps emerged in the mid-2000s, when Inou and Shishikura established the hyperbolicity for near-parabolic renormalization. We will state this result in §3.6. The hyperbolicity of near-Siegel renormalization was established a decade later; see §3.7. The neutral and satellite ql renormalization naturally form the *Molecule renormalization*, see Figure 11 and §3.8. The hyperbolicity of such a renormalization is likely required for Conjecture 1.2 and the area-zero conjecture on $\partial\mathcal{M}$. Conjecturally, there is no renorm-expanding regime on the Molecule: every parameter there has substantial hyperbolic components in all scales.

In short, the Puzzle Renormalization deals with the renorm-expanding (see §1.3) and ‘‘virtual primitive ql’’ (central cascades §3.9) regimes. In the renorm-expanding regime, there are no geometric limits as in (1.3), but there is a strong phase-parameter relation between the Mandelbrot and Julia sets as illustrated in Figure 8. One may hope that this case can also be neatly integrated into the hyperbolicity framework:

Question 3.1 (Uniform hyperbolicity rel $\partial\mathcal{M}$). Is there a renormalization operator of a hyperbolic nature, associated with all parameters and scales of $\partial\mathcal{M}$, with dimension 1 unstable directions, such that the operator combines all the above kinds of renormalization regimes: (QL), (Neut), (Puz)?

Here is a hoped-for *universality statement* from a desired operator in Question 3.1. Consider a continuous zooming-in at $c \in \partial\mathcal{M}$. If in a small δ -scale of c , all hyperbolic components of \mathcal{M} are small rel δ , then we expect a form of Tan-Lei type similarity between \mathcal{M} and J_c as depicted in Figure 8. If there is an unbounded (of size $\gg \delta$) hyperbolic component of \mathcal{M} hitting the δ -scale of c , cf. Figure 4, then we anticipate strong near-Neutral effects. And if all non-negligible hyperbolic components of \mathcal{M} are intermediate in the δ -scale of c , cf. Figure 2, then the quadratic-like features emerge. Finally, if the δ_1 scale of c_1 is combinatorially close to the δ_2 -scale of c_2 , then these scales should also be close geometrically. The last statement is a standard application of the hyperbolicity framework of the renormalization.




Figure 8: It is natural to ask if a form of Tan-Lei type similarity [49] (e.g., for the vein structure) between respective small neighborhoods of \mathcal{M} and J_c at c occurs if and only if (parameter) hyperbolic components of \mathcal{M} become negligible; this is related to the notion of the renorm-expanding in §1.3. For puzzles and parapuzzles, the *Almost-Linear* relation (similarity) was established for Yoccoz maps in [37], see §3.9.

3.1 Analytic and laminar structures on the space of ql maps. A *quadratic-like map* $f: U \xrightarrow{2:1} V$ is a degree 2 holomorphic branched covering between open disks $U \Subset V \Subset \mathbb{C}$, see Figure 9. The set of its non-escaping points $K_f := \{z \in U : f^n(z) \in U \ \forall n \geq 1\}$ is called the *filled Julia set* of $f: U \rightarrow V$. The boundary $J_f := \partial K_f$ is the *Julia set* of f . The Julia set is connected if and only if the unique critical point of f is non-escaping. In this case, f is hybrid equivalent to a unique polynomial $p_c: z \mapsto z^2 + c$ with $c = c(f) \in \mathcal{M}$: there is a qc conjugacy h between $f: U \rightarrow V$ and a quadratic-like restriction $p_c: A \rightarrow B$ such that h is conformal on the non-escaping sets.

Here and elsewhere, by a slight thinning of V and U , we can assume that \overline{U} is a closed topological disk. A *small Banach neighborhood* of $f: U \rightarrow V$ is the set of all analytic maps close to f in the sup-norm:

$$(3.1) \quad \mathcal{B}_\varepsilon(f | U) = \{g: U \rightarrow V : \|f - g\|_{\overline{U}} \leq \varepsilon\}.$$

Let $V' \Subset V$ be a slight thinning of V . If ε is sufficiently small, then every $g \in \mathcal{B}_\varepsilon(f | U)$ admits a quadratic-like structure $g: U_g \xrightarrow{2:1} V'$, where U_g is the unique g -lift of V' . This naturally introduces complex analytic structure for quadratic-like maps. There are two natural equivalences on the space of quadratic-like maps $f: U \rightarrow V$: linear conjugacy (i.e., rescaling) and thickening/thinning U and V . In [36], respecting these equivalences, the space of *all* quadratic-like maps \mathcal{QL} was supplied with complex-analytic structure induced by small Banach balls as in (3.1). In this space, hybrid-equivalent maps form a codimension 1 lamination with analytic leaves. The holonomy along hybrid leaves is qc and is asymptotically conformal. Such a holonomy combined with the hyperbolicity rigorously justifies various universalities as in Figure 2. For near-Neutral Renormalization, establishing an appropriate version of the following consequence of the laminar structure is one of the key technical challenges, see §3.7, §3.8.

Lemma 3.2 ([36], codim = 1 stability of a ql f). Let $f: U \rightarrow V$ be a ql map with a connected Julia set as above. Then, for a small $\varepsilon > 0$, there is an analytic function $\tau: \mathcal{B}_\varepsilon(f | U) \rightarrow \mathbb{C}$ such that $f, g \in \mathcal{B}_\varepsilon(f | U)$ are hybrid conjugate if and only if $\tau(f) = \tau(g)$.




Figure 9: Top: Quadratic-Like Renormalization $f_1 = f^p: U_1 \xrightarrow{2:1} V_1$ is obtained from $f: U \xrightarrow{2:1} V$ by iterating, restricting, and then linear rescaling, see §3.3. Bottom: Puzzle (or generalized ql) Renormalization is the first return map to a puzzle Y_1 , see §3.9.

3.2 QL Compactness. Given a ql map $f: U \xrightarrow{2:1} V$, its *modulus* is either

$$(3.2) \quad \text{mod}(f) := \text{mod}(V \setminus U) \quad \text{or} \quad \text{mod}^+(f) := \text{mod}(V \setminus K_f);$$

both quantities are compatible. If K_f is a Cantor set, then $\text{mod}^+(f)$ is the supremum of moduli over all annuli in the correct homotopy class separating ∂V from K_f . Given a $\delta > 0$, the space of ql maps with $\text{mod}^+(f) \geq \delta$ is compact: every sequence of ql maps f_n with $\text{mod}^+(f_n) \geq \delta$ has a convergent subsequence within \mathcal{QL} ; the limit f_∞ also satisfies $\text{mod}^+(f_\infty) \geq \delta$. More generally, instead of separating ∂V and K_f , we can measure the separation between ∂V and the postcritical set. (In this framework, sequential limits of ql maps are, in general, outside the \mathcal{QL} space.) This approach is adopted for near-neutral maps where the classical ql bounds fail, see §3.4.

3.3 QL Renormalization operator. As Figure 9 illustrates, a ql renormalization $f_1 = f^p: U_1 \xrightarrow{2:1} V_1$ of $f: U \xrightarrow{2:1} V$ is obtained by iterating f followed by a ql-restriction and a linear rescaling. Since all three operations are analytic, the renormalization admits a natural analytic extension to nearby maps; we refer to such an extension as a *renormalization operator*. Below, we first introduce a non-dynamical analytic operator. By design, it is always compact: the image of a ball is precompact. If there are appropriate geometric bounds, then the operator becomes dynamical and hyperbolic.

Let $U' \Subset U$ and $V' \subset V$ be slight thinning of U and V with $\delta := \text{mod}(U \setminus U')$. It is a fundamental fact (follows from Cauchy's integral formula) that the restriction operator

$$(3.3) \quad \mathcal{B}_\varepsilon(f \mid U) \rightarrow \mathcal{B}_\varepsilon(f \mid U'), \quad [g: U \rightarrow \mathbb{C}] \mapsto [g: U' \rightarrow \mathbb{C}]$$

is compact. Here ε in $\mathcal{B}_\varepsilon(f \mid U)$, see (3.1), is chosen sufficiently small depending on δ . Viewing $f_1 \equiv f^p: U_1 \xrightarrow{2:1} V_1$ as a subsystem of $f: U \xrightarrow{2:1} V$, we assume the following the *compactness condition*:

$$(3.4) \quad \bigcup_{i=0}^{p-1} f^i(U_1) \Subset U' \quad \text{where} \quad \delta := \text{mod}(U \setminus U') \quad \text{as above;}$$




Figure 10: Sector Renormalization §3.4 is the first return map to a sector S whose sides are glued dynamically.

i.e., the forward orbits of points in U_1 stay in U' until their first return to $V_1 \subset V'$. Then, since the restriction operator (3.3) is compact, we obtain a compact non-dynamical analytic operator

$$(3.5) \quad \mathcal{R}: \mathcal{B}_\varepsilon(f_1 | U) \rightarrow \mathcal{B}_{\varepsilon_2}(f^p | U_1), \quad [g|U] \mapsto [g|U'] \mapsto [g_1 = g^p|U_1].$$

Suppose that $f \equiv f_0$ is infinitely ql-renormalizable: there is an infinite sequence $f_n: U_n \xrightarrow{2:1} V_n$ with $f_{n+1} = f_n^{p_{n+1}}$ as before. We also assume that $\delta > 0$ is selected uniformly over f_n and that the periods $p_n \leq \bar{p}$ are uniformly bounded. If the sequence $f_n: U_n \rightarrow V_n$ has geometric bounds $\text{mod}^+(f_n) \geq \eta > 0$, see (3.2), then, by the compactness of ql maps, the ω -limit set \mathfrak{H}_f of $(f_n)_n$ is compact. Such a limit set \mathfrak{H} is called a *renormalization horseshoe*. If the geometric bounds have an additional ‘‘beau’’ property (it is often the case), then \mathfrak{H}_f is a hyperbolic set of \mathcal{R} . Similarly, a renormalization horseshoe $\mathfrak{H}_\mathcal{X}$ is defined for a set of maps $\mathcal{X} = \{f_\tau \equiv f_{0,\tau}\}$. It is natural to construct a horseshoe as big as possible. Applying Theorem 4.1 together with [36], we obtain:

Theorem 3.3 (Hyperbolicity, (QL) Case). For every bound \bar{p} on renormalization periods, there is a hyperbolic renormalization horseshoe $\mathfrak{H}_{\bar{p}}$ with $\dim = 1$ unstable manifolds for the ql renormalization operator as above. The horseshoe $\mathfrak{H}_{\bar{p}}$ attracts exponentially fast all infinitely renormalizable ql maps $f \equiv f_0$ with relative periods $p_n \leq \bar{p}$.

3.4 Sector Renormalization. The Sector Renormalization was originally designed to study Local Dynamics around an indifferent fixed point by Douady and Yoccoz, with various ideas going back to the theory of Ecalle-Voronin’s invariants. For high-type rotation numbers, Inou and Shishikura extended the sector renormalization to a semi-global compact operator with geometric control of the critical orbit. Here, we provide more details on the semi-global case.

The idea of sector renormalization \mathcal{R}_{sec} is illustrated in Figure 10. In short, it is the first return map of f_n to the appropriately chosen sector S so that the critical orbit returns to S . Let us start with notation:

- $\Theta = \{\theta = [0; a_1, a_2, \dots] \mid a_i \in \mathbb{N}_{\geq 1}\} \simeq [0, 1] \setminus \mathbb{Q} \simeq (\mathbb{R}/\mathbb{Z}) \setminus \mathbb{Q}$ is the set of irrational rotation numbers written as continued fractions;
- $\Theta_{\text{bnd}} = \{\theta = [0; a_1, a_2, \dots] \mid a_i \leq M_\theta\} \subset \Theta$ is the set of bounded (irrational) rotation numbers;
- $\overline{\Theta} = \{\theta = [0; a_1, a_2, \dots] \mid a_i \in \mathbb{N}_{\geq 1} \cup \{\infty\}\} \supset \Theta$ is the formal compactification of Θ ;
- $\Theta_{>N} = \{\theta = [0; a_1, a_2, \dots] \mid a_i \in \mathbb{N}_{>N}\}$;
- $f_\theta: z \mapsto e^{2\pi i \theta} z + z^2$, $\theta \in \Theta$ is an irrationally-neutral quadratic polynomial as in (2.3),
- $\mathfrak{p}_n/\mathfrak{q}_n \approx \theta$ is the best approximants of $\theta \in \Theta$ starting with $\mathfrak{q}_0 = 1$,
- $f_n \equiv f_{n,\theta} := \mathcal{R}_{\text{sec}}^n f_\theta$ is the n th sector renormalization;
- if $\theta \in \Theta_{>1}$, then $\theta_n := (-1)^n [0; a_{n+1}, a_{n+2}, \dots]$ is the rotation number of f_n and $f_\theta^{[n]} := f_\theta^{\mathfrak{q}_n}$ is the sector prerenormalization of $f_n = f_\theta^{[n]} / f_\theta^{[n-1]}$.

The definitions of θ_n and $f_\theta^{[n]}$ require a minor adjustment for the general case “ $\theta \in \Theta$ ”. We view $f_n = \mathcal{R}_{\text{sec}}^n f_\theta$ as

$$(3.6) \quad f_n: \mathbb{D} \dashrightarrow \mathbb{D} \quad f_n(0) = 0, \quad f_n(0) = e^{2\pi i \theta_n}, \quad \theta_n \in \Theta.$$

The *sector renormalization* f_{n+1} of f_n as shown in Figure 10 is the first return map to the smallest sector S bounded by curves γ and $f_n(\gamma)$ emerging from 0 so that the sides $\gamma, f_n(\gamma) \subset S$ are dynamically glued by f_n :

$$(3.7) \quad \gamma \ni x \sim f_n(x) \in f_n(\gamma) \quad \text{so that} \quad S/f_n \simeq \mathbb{D}.$$

We assume that γ lands at 0 at a well-defined angle. The “smallest” condition on S refers to the requirement that S contains the smallest angle between γ and $f_n(\gamma)$. The sector S should be chosen appropriately so that the first return map to S controls the critical orbit. Before taking the quotient (3.7), the first return map is a commuting pair of the form $F_n = (f_n^{[1]} \equiv f_n^{\mathbf{q}_1(\theta_n)}, f_n^{\mathbf{q}_1(\theta_n) \pm 1})$. The quotient map by f_n identifies both branches in the commuting pair F_n ; we can write $f_{n+1} = \mathcal{R}_{\text{sec}} f_n = f_n^{[1]}/f_n \equiv F_n/f_{n+1}$, and we refer to both F_n and $f_n^{[1]}$ as the *sector prerenormalizations* of f_{n+1} . Similarly, $f_\theta^{[n]}$ is the n th sector prerenormalization of f_n , and we write: $f_n = f_\theta^{[n]}/f_\theta^{[n-1]}$. As in the quadratic-like case, see (3.4), we require a *compactness condition*:

$$(3.8) \quad \text{mod}(\mathbb{D} \setminus \text{orb}^{\mathbf{q}_1}(S)) \geq \delta,$$

where $\text{orb}^{\mathbf{q}_1}(S)$ is formed by orbit trajectories of points in S projecting to $\text{dom}(f_{n+1})$. We refer to (3.8) as *sectorial bounds*; they imply the compactness of \mathcal{R}_{sec} . As an application of Theorem 4.2, the compact sectorial renormalization operator \mathcal{R}_{sec} is justified in [19] for all neutral quadratic polynomials $(f_\theta)_{\theta \in \Theta}$ with a compactness condition (3.8) for a uniform δ and with well-defined parabolic limits parametrized by $\overline{\Theta}$.

3.5 Parabolic Enrichment and the Cylinder Renormalization. We recall that the filled Julia set is discontinuous at parabolic parameters: as shown in Figure 6, the limit of filled Julia sets can be smaller than the filled Julia set of the limiting parameter. This discontinuity can be resolved by introducing limiting Lavaurs maps. Let us illustrate a general idea with examples.

Consider a sequence of parameters $a_n \rightarrow 1/4$ as shown in Figure 6. In the dynamical plane of $p_{a_n}: z \mapsto z^2 + a_n$, see Figure 7, the critical orbit returns after $\mathbf{q}_{1,n} \equiv \mathbf{q}_1(a_n)$ iterates. The orbit spends $\mathbf{q}_{1,n} - s$ iterations traveling from the red cylinder to the blue cylinder, and then s iterations traveling back. We have a commuting pair $(p_{a_n}, p_{a_n}^{\mathbf{q}_{1,n}})$. In the limit, the parabolic gap is closed, and $(p_{a_n}, p_{a_n}^{\mathbf{q}_{1,n}})$ converges to the pair $(p_{1/4} = p_{a_\infty}, p_{a_\infty}^{[1]})$, where $p_{a_\infty}^{[1]}$ is a transcendental Lavaurs map defined on the attracting basin of the parabolic point of $p_{1/4}$. The red and blue cylinders converge to the limiting attracting and repelling cylinders of $p_{1/4}$.

The *cylinder renormalization* \mathcal{R}_{cyl} is a convenient and canonical extension of a sector renormalization \mathcal{R}_{sec} from §3.4 such that the renormalization sector of \mathcal{R}_{sec} is *extended to a cylinder* of \mathcal{R}_{cyl} . While \mathcal{R}_{sec} is defined up to conformal conjugacy, \mathcal{R}_{cyl} is defined up to a linear conjugacy. For p_{a_n} , its *cylinder renormalization* $\mathcal{R}_{\text{cyl}}(p_{a_n}) \simeq p_{a_n}^{\mathbf{q}_{1,n}}/p_{a_n}$ is the first return map of p_{a_n} back to the red cylinder, where its boundaries are identified using p_{a_n} . The limiting cylinder renormalization $\mathcal{R}_{\text{cyl}}(p_{1/4}, p_{a_\infty}^{[1]})$, also known as *parabolic*, is the quotient of $p_{a_\infty}^{[1]}$ under $p_{1/4}$. We write

$$(3.9) \quad \mathcal{R}_{\text{cyl}}(p_{a_n}) \simeq p_{a_n}^{\mathbf{q}_{1,n}}/p_{a_n} \longrightarrow \mathcal{R}_{\text{cyl}}(p_{1/4}, p_{a_\infty}^{[1]}) \simeq p_{1/4}^{[1]}/p_{1/4},$$

and refer to $\mathcal{R}_{\text{cyl}}(p_{a_n})$ as *near-parabolic* for $n \gg 1$. In fact, $p_{a_\infty}^{[1]}$ and $\mathcal{R}_{\text{cyl}}(p_{a_\infty}^{[1]})$ are uniquely determined by a linear map, called a *transitional map*, between red and blue cylinders, see the bottom-left part of Figure 7. Conversely, given a transitional map, one can construct the associated Lavaurs map and then realize it as a limit of nearby maps. A fundamental question is the dynamical control of this *parabolic enrichment*. As it was discussed in §3.4, renormalization of neutral quadratic polynomials is parametrized by $\theta = [0; a_1, a_2, \dots] \in \overline{\Theta}$. Parabolic renormalization applies to the cases where $a_i = \infty$. For illustration, see parameters b_n in Figure 6.

3.6 Hyperbolicity of the near-parabolic renormalizations. In [24], the hyperbolicity of the cylinder renormalization has been established for high-type parameters $\theta \in \Theta_{\leq N}$, where $N \gg 1$. The hyperbolicity implies the sectorial bounds as they were introduced in §3.4. As we mentioned in Section 2, this result had many




Figure 11: Left: the Main Molecule \mathfrak{M} of the Mandelbrot set and the induced by the Branner-Douady surgery Molecule map, see §3.8. The Molecule map is modeled by $z \mapsto z(z + 1)^2$, right.

applications. The strategy in [24] is as follows. First, with computer assistance, the *a priori* bounds (a variant of (3.8)) are established for the parabolic renormalization

$$[0; \infty, \infty, \infty, \dots] \in \overline{\Theta}, \quad \text{see §3.4.}$$

Using Teichmüller contraction, the geometric bounds are promoted to the hyperbolicity of the associated parabolic renormalization. By the stability of hyperbolicity, it also holds for all $\theta \in \Theta_{>N}$ for some $N \gg 1$.

3.7 Hyperbolicity of near-Siegel Renormalization. Following notations of 3.4, consider maps $f_\theta: z \mapsto e^{2\pi i \theta} z + z^2$. For $\theta \in \Theta_{\text{bnd}}$, the Douady-Ghys surgery transfers *real bounds* from circle maps (the Blaschke family) to quadratic polynomials, see Figure 5, left. Consequently, f_θ has a qc Siegel disk if and only if $\theta \in \Theta_{\text{bnd}}$. Moreover, such a qc Siegel disk contains a critical point on the boundary. Relying on such input, McMullen in [42] developed dynamical renormalization theory for $\theta \in \Theta_{\text{bnd}}$, and justified the dynamical universality: the exponential convergence of renormalizations towards the limit, see Figure 5, right.

For two decades, it remained an open question whether dynamical universality could be promoted to hyperbolicity. The key obstacle was to establish a version of Lemma 3.2: proving that Siegel maps remain stable under small perturbations, provided the perturbation does not change the multiplier of the Siegel fixed point. Here, perturbations are taken within Banach balls (cf. (3.1)) consisting of maps defined in neighborhoods of their Siegel disks. This obstacle was resolved by employing transcendental techniques on unstable manifolds. *A priori* bounds imply the compactness of the operator, thus unstable manifolds exist but may have a large dimension. It was shown in [20] that maps on the unstable manifolds admit maximal analytic extensions as σ -proper maps “ $W \rightarrow \mathbb{C}$,” cf. (1.3), where W is open and dense, with a single critical orbit. Such global σ -proper maps have $\text{codim} = 1$ stability as in Lemma 3.2; this stability is then promoted to $\text{codim} = 1$ stability in the original class of maps, and subsequently to:

Theorem 3.4 ([20]). Pacman/Siegel Renormalization operator is hyperbolic with $\dim = 1$ unstable manifold for parameters in Θ_{bnd} of periodic type.

For the Golden-mean Siegel map shown in Figure 5, the hyperbolicity was established earlier by Gaidashev and Yampolsky with computer-assistance. We also remark that the global σ -proper structure of renormalization fixed points (both of the quadratic-like and Siegel types) was discovered by McMullen in the 1990s. The theory of external transcendental rays for unstable manifolds was designed in [17], leading to further applications.

3.8 Molecule Renormalization. As we have indicated in §3.4, Theorem 4.2 implies the existence of a compact sectorial renormalization operator \mathcal{R}_{sec} rel $\overline{\Theta}$. The proof of Theorem 3.4, relying on the analysis of the transcendental dynamics on the unstable manifolds, provides a strategy towards:

Conjecture 3.5 (Hyperbolicity of Neutral Renormalization). There is a hyperbolic cylinder renormalization operator \mathcal{R}_{cyl} with $\dim = 1$ unstable manifolds for all parameters in $\overline{\Theta}$.

We recall from [20, Appendix C] that the satellite and neutral renormalizations are naturally unified into the Molecule Renormalization; its slowed-down version is illustrated in Figure 11. A natural extension of Conjecture 3.5:




Figure 12: Illustration to the positive entropy argument, §4.1, the airplane combinatorics.

Conjecture 3.6 (Hyperbolicity of Molecule Renormalization). There is a hyperbolic renormalization with $\dim = 1$ unstable manifolds parametrized by the sped-up Molecule map as in Figure 11.

The completion of Problem 4.3 is required for Conjecture 3.6. As we have already mentioned, Conjecture 3.6 is likely required for the satellite cases of Conjecture 1.2 and the Area Problem of $\partial\mathcal{M}$.

3.9 Puzzle Renormalization. Puzzle techniques were first employed by Branner and Hubbard in [5] to analyze cubic polynomials with one escaping critical point. In that setting, puzzles do not cut the Julia set. A few years later, Yoccoz introduced puzzles for quadratic polynomials and used them to reduce the MLC to the infinitely ql renormalizable case, see (2.1). Yoccoz puzzles cut a Julia set into smaller pieces. Then the Grötzsch inequality is used to show that such pieces shrink to points.

In [35, 37], in the 1990s, the puzzle techniques were incorporated into the renormalization framework as shown in Figure 9. Given a puzzle Y around a critical point so that the forward p_c -orbit of ∂Y is disjoint from Y , we can consider the first return map g of p_c to Y . The domain of g consists of countably many puzzles. One of them, say Y_1 , contains a critical point and is mapped with degree 2 into Y . All other pieces are mapped with degree 1. The *renormalization* g_1 of g is the first return map of g into Y_1 . Let $X \Subset Y_1$ be the next-level puzzle containing the unique critical point. If the return time of X rel g is bigger than 1, i.e., X travels around Y under g , then the “asymmetric modulus” grows; roughly, $g_1: Y_1 \dashrightarrow Y_1$ has more expansion than $g: Y \dashrightarrow Y$. On the other hand, g_1 belongs to a *central cascade* if the return time of X is 1, i.e., $g|X = g_1|X$.

If a central cascade is long but still finite, then the associated dynamical scales will be dominated by a certain primitive small copy \mathcal{M}' of \mathcal{M} that does not contain c . We refer to this phenomenon as *virtual primitive ql*.

If $g: Y \dashrightarrow Y$ is sufficiently expanding (i.e., moduli are big, cf. the renorm-expanding case in §1.3), then there is an *Almost-Linear puzzle-parapuzzle* relation [37]; i.e., the geometry of (puzzles of) $g: Y \dashrightarrow Y$ can be efficiently transferred to the parameter plane, cf. Figure 8. This is an essential ingredient for the “regular vs stochastic” dichotomy. Purely renorm-expanding real parameters (§1.3) are precisely Lebesgue density points for stochastic parameters: most of their small perturbations have exponential growth of the expansion along the critical orbit (the Collet-Eckmann condition), i.e., they are stochastic.

4 Near-Degenerate Regime. As we mentioned in §2, the Near-Degenerate Regime provides an adequate framework towards establishing *a priori* bounds. Since it operates at a basic topological level, arguments developed for different dynamical scales are often compatible and can be merged provided the setting is appropriately prepared. The non-dynamical foundations of the theory have been developed in [30]. Below, we will omit most of the technical discussion and indicate only the sources of geometric control.

On the dynamical scales with definite *positive core entropy* $\geq \varepsilon > 0$, the theory is fully developed, see §4.1 and §4.2. As Figure 12 illustrates, the positivity of entropy prevents dynamically invariant horizontal degeneration. Thus, if there is a substantial degeneration, then it can be amplified at shallower levels.

Consider the *zero core entropy* scales. On satellite ql levels, degeneration along non-invariant rectangles is amplified at shallower levels, while degeneration along combinatorially invariant rectangles is amplified at deeper levels; see Figure 13 and §4.3. In the (near-)neutral case, invariant degeneration can actually exist within parabolic

fjords because of the non-JLC; *pseudo-Siegel* disks are constructed to consume most of the invariant degeneration, see the left loop of Figure 14. Its right loop illustrates a strategy to develop pseudo-Siegel disks in the remaining near-neutral satellite cases. If achieved, this would fully unify the cases of §4.1–§4.4 and confirm, in particular, the Satellite Case of the MLC.

Let us now discuss the *Interpolation Problem* between the positive and zero entropy settings. It concerns parameters $c \in \mathcal{M}$ that are geometrically close to the main Molecule \mathfrak{M} of \mathcal{M} but are outside of satellite little copies centered at \mathfrak{M} . In the dynamical plane of $p_c: z^2 + c$, there are many geometric scales of zero-entropy nature associated with \mathfrak{M} . A key issue is that, on these scales, the postcritical set of p_c can look as if it is dense in the Julia sets; i.e., the entropy on the postcritical set can blow up in the limit as c approaches \mathfrak{M} . (The postcritical set is only lower semicontinuous.) This effectively compels us to deal with *virtual Julia sets* containing the critical orbit up to several first returns. We briefly formulate this *Virtual Renormalization* setting in §4.5.

4.1 Primitive QL levels. Assume that f_1 is a primitive ql renormalization of f with a period p as in Figure 9. Instead of $\text{mod}^+(f) = \text{mod}(V \setminus K_f)$ as in (3.2), it is more convenient to consider the dual quantity, called the *width* or the *degeneration* of f :

$$(4.1) \quad \mathcal{W}(f) := \frac{1}{\text{mod}^+(f)} \quad \text{so that} \quad \text{mod}^+(f) \leq \varepsilon \quad \text{iff} \quad \mathcal{W}(f) \geq K \equiv \frac{1}{\varepsilon},$$

and similar for f_1 . The width $\mathcal{W}(f)$ is the extremal width of the family of curves connecting K_f to ∂V .

Let us analyze how the degeneration of f_1 is developed in the dynamical plane of f . Since f has p small (filled) Julia sets of f_1 , the total degeneration around these small Julia sets is $\mathcal{W}^{\text{tot}} \asymp p\mathcal{W}(f_1) - O_p(1)$, where an additive error $O_p(1)$, that depends on p , comes from a variant of the Thin-Thick Decomposition. We can decompose $\mathcal{W}^{\text{tot}} = \mathcal{W}^{\text{ver}} + \mathcal{W}^{\text{hor}}$, where

- \mathcal{W}^{hor} is the *horizontal* degeneration between any pair of two small Julia sets; and
- \mathcal{W}^{ver} is the *vertical* degeneration between ∂V and any small Julia set.

A key claim in [25] is that $\mathcal{W}^{\text{ver}} \succeq \mathcal{W}^{\text{tot}} - O_p(1)$; i.e., the vertical part has a definite proportion. Since small Julia sets of f_1 are within the bigger Julia set of f , the claim implies

$$(4.2) \quad \mathcal{W}(f) \geq \mathcal{W}^{\text{ver}} \succeq p\mathcal{W}(f_1) - O_p(1).$$

The idea behind $\mathcal{W}^{\text{ver}} \succeq \mathcal{W}^{\text{hor}}$ is illustrated in Figure 12. Assume by contradiction that $\mathcal{W}^{\text{hor}} \gg \mathcal{W}^{\text{ver}}$, i.e., \mathcal{W}^{ver} is negligible. It is a meta-principle that a degeneration should be invariant under dynamics. Therefore, the degeneration \mathcal{W}^{hor} is (eventually) aligned with the Hubbard tree of f and is represented by two *wide rectangles* \mathcal{G}_1 and \mathcal{G}_2 depicted on the figure as narrow (for convenience) red and blue rectangles in the pencil style. Since \mathcal{W}^{ver} is negligible,

- (most of) \mathcal{G}_2 overflows a lift \mathcal{G}_1'' of \mathcal{G}_1 ; and
- (most of) \mathcal{G}_1 overflows a lift \mathcal{G}_2' of \mathcal{G}_2 and then a lift of \mathcal{G}_1' of \mathcal{G}_1 .

This is impossible by the Grötzsch inequality and (4.2) follows.

By applying (4.2) to f and its n th renormalization f_n with a big period $p_n \gg 1$; we obtain

$$(4.3) \quad \mathcal{W}(f) \succeq p_n \mathcal{W}(f_n) - O_{p_n}(1) \gg \mathcal{W}(f_n) \quad \text{if} \quad \mathcal{W}(f_n) \gg_{p_n} 1.$$

This implies *a priori* bounds in the primitive bounded-type ql case: the degeneration $\mathcal{W}(f_m)$ can not become big because, otherwise, the degeneration of $\mathcal{W}(f_{m-n})$ will be even bigger.

4.2 Puzzle levels. We emphasize the following subtlety in (4.2): the period of p_n is assumed to be big but *still bounded* as the additive error $O_{p_n}(1)$ depending on p_n needs to be bounded.

Assume now that the period p of f_1 rel f is big. We assume here that f_1 is the first ql renormalization of f of the primitive type; this renormalization is encoded by a maximal primitive copy \mathcal{M}_1 of \mathcal{M} . The renormalization $f \mapsto f_1$ can be factorized using the puzzle renormalization, see Figure 9,

$$(4.4) \quad f \mapsto g_1 \mapsto g_2 \mapsto \cdots \mapsto g_k \mapsto f_1.$$




Figure 13: Illustration to §4.3: if \mathcal{G} is invariant rel δ , then $\mathcal{W}(\mathcal{G} \setminus \mathcal{G}') \geq C_\delta \mathcal{W}(\mathcal{G}) \gg \mathcal{W}(\mathcal{G})$.

If the copy \mathcal{M}_1 is ε -away from the main molecule (the *anti-Molecule* property, cf. Figure 11) and $p \gg_\varepsilon 1$, then the (modified) Principal Nest of (4.4) is combinatorially substantial and provides amplification of the degeneration. Roughly, the argument goes as follows. Using the notations of Figure 9, as X travels around Y , the [30, Covering Lemma] efficiently spreads the degeneration around. Then the [30, Quasi-Additivity Law] collects the degeneration along the orbit of X ; this eventually leads to:

$$(4.5) \quad \mathcal{W}(f) \gg \mathcal{W}(f_1) \quad \text{if} \quad \mathcal{W}(f) \gg_\varepsilon 1 \quad \text{and} \quad p \gg_\varepsilon 1.$$

Together, (4.3) and (4.5) imply the *a priori* bounds and the MLC under the anti-molecule condition [27, 28].

4.3 Satellite QL levels. Assume now that f_1 is a satellite ql renormalization of f of bounded-type. The key difference with the primitive case (§4.1) is the existence of combinatorially invariant rectangles \mathcal{G} between small Julia sets as illustrated in Figure 13. Below, we sketch how to continue the argument behind (4.3) in the presence of such a \mathcal{G} .

Let us assume that most of the \mathcal{W}^{tot} is within \mathcal{G} . For a small $\delta > 0$, the $(1 - \delta)$ part of \mathcal{G} overflows its lift \mathcal{G}' as shown in the figure. However, before entering the rectangle \mathcal{G}' , curves forming the above $(1 - \delta)$ -part of \mathcal{G} must travel through preperiodic Julia sets as depicted using purple color. Let us denote by $\mathcal{G} \setminus \mathcal{G}'$ the family of such purple subcurves. Since $\mathcal{G}, \mathcal{G}'$ have almost the same width, up to “ $(1 - \delta)$ ”, the Grötzsch inequality implies

$$(4.6) \quad \mathcal{W}(\mathcal{G} \setminus \mathcal{G}') \geq C_\delta \mathcal{W}(\mathcal{G}), \quad \text{where } C_\delta \rightarrow \infty \text{ as } \delta \rightarrow 0.$$

The [18, Wave Lemma] allows us to interpret the quantity $\mathcal{W}(\mathcal{G} \setminus \mathcal{G}')$ as the degeneration $\mathcal{W}(f_2)$ on a deeper level. Considering now f, f_n and f_{n+1} , Estimate (4.6) gives us the following refinement of (4.3): either

- (i) $\mathcal{W}(f) \succeq_\delta p_n \mathcal{W}(f_n) - O_{p_n}(1)$; or
- (ii) $\mathcal{W}(f_{n+1}) \geq C_\delta \mathcal{W}(f_n)$.

The Alternative (ii) is prevented by the Teichmüller contraction. Indeed, given a bound \bar{p} on the relative period p_{m+1}/p_m of f_{m+1} rel f_m , the degeneration of $\mathcal{W}(f_m)$ can grow at most exponentially fast: $\mathcal{W}(f_m) = O(\Delta_{\bar{p}}^m)$. Therefore, by selecting $\delta > 0$ to be sufficiently small so that $C_\delta \gg \Delta_{\bar{p}}$ is sufficiently big, Alternative (ii) is eventually prevented on some levels. We also select n sufficiently big so that p_n dominates the multiplicative constant representing “ \succeq_δ ”. Alternative (i) implies

$$\mathcal{W}(f) \succeq_\delta p_n \mathcal{W}(f_n) - O_{p_n}(1) \gg \mathcal{W}(f_n) \quad \text{if } \mathcal{W}(f_n) \gg_{p_n} 1.$$

Theorem 4.1 ([18], §4.1+§4.3). *A priori* bounds and the MLC hold for all ql-combinatorics of bounded-type.

In particular, the MLC holds at the period-doubling Feigenbaum parameter illustrated in Figure 2. The arguments of §4.2 can also be integrated with those in Theorem 4.1; the combination, see [14], yields the MLC along a real line and along any vein of the Mandelbrot set.




Figure 14: Left loop: the interaction between pseudo-Siegel disks and neutral sector antirenormalization, see §4.4. Right loop: illustration to the strategy to extend the theory of pseudo-Siegel disks to the satellite case.

4.4 Near-Neutral levels. Let us first detail the following statement:

Theorem 4.2 ([16]). Uniform pseudo-Siegel a priori bounds hold for all neutral quadratic polynomials.

Using the notations of §3.4, consider a neutral polynomial $f_\theta: z \mapsto e^{2\pi i \theta} z + z^2$ for an irrational $\theta \in \Theta$. Theorem 4.2 asserts that it has a sequence of nested pseudo-Siegel disks

$$(4.7) \quad \widehat{Z}^{-1} \supseteq \widehat{Z}^0 \supseteq \widehat{Z}^{[1]} \supseteq \dots \supseteq \widehat{Z}^{[m]} \supseteq \dots \supseteq H_{f_\theta} \ni 0, \quad \bigcap_{m \geq -1} \widehat{Z}^{[m]} = H \equiv H_{f_\theta},$$

where H is the *Mother Hedgehog* (the filled postcritical set) of f_θ such that every $\widehat{Z}^{[m]}$ is almost invariant up to $q_{[m+1]}$ -iterations. The disks \widehat{Z}^{-1} and \widehat{Z}^0 are uniformly qc over all f_θ with $\theta \in \Theta$.

Under the n th sector renormalization $f_\theta \mapsto f_n$, cf. Figure 10, the pseudo-Siegel disk $\widehat{Z}_{f_\theta}^{[n+m]}$ of f_θ becomes $\widehat{Z}_{f_n}^{[m]}$ of f_n ; i.e., every $\widehat{Z}^{[m]}$ becomes uniformly qc after applying the m th renormalization.

Figure 14, the left loop, illustrates the interaction of \widehat{Z}^0 , \widehat{Z}^{-1} with sector antirenormalization. If the rotation number of f_n is close to 0, then the α and β fixed points are close to each other and $\partial\widehat{Z}_{f_n}^0$ travels through the gap between $\alpha = 0$ and β . The pseudo-Siegel disk $\widehat{Z}_{f_n}^{-1}$ is obtained by adding a parabolic fjord to $\widehat{Z}_{f_n}^0$. The fjord is bounded by a dam located inside an invariant rectangle, depicted in a pencil style. We say that $\widehat{Z}_{f_n}^0$ is *regularized* to $\widehat{Z}_{f_n}^{-1}$. Under the antirenormalization, $\widehat{Z}_{f_n}^{-1}$ become $\widehat{Z}_{f_{n-1}}^0$, and the procedure can be repeated. If the rotation number of f_n is away from 0, then there is no regularization: $\widehat{Z}_{f_n}^0 = \widehat{Z}_{f_n}^{-1}$.

Informally, since antirenormalization has contracting properties, we expect that the “non-invariance error” of $\widehat{Z}^{[m]}$ is uniformly bounded by a geometric series. The [16] justifies such a property by employing the near-degenerate analyses in the original plane of f_θ . The sectorial bounds (cf. Figure 10) are then justified in [19] in the *a posteriori* setting.

The right loop in Figure 14 illustrates a strategy, developed jointly with Lyubich, for extending the theory of pseudo-Siegel disks to the near-neutral setting. On this loop, the antirenormalization is taken rel β . For many combinatorics, this has been already justified and integrated with §4.1-§4.3. If the strategy is fully realized, then this would imply, in particular, the satellite case of the MLC.

Problem 4.3. Establish pseudo-Siegel a priori bounds in the remaining unbounded satellite ql cases.

4.5 Virtual near-Molecule Renormalization. Consider now a general case, when f_1 is the first ql renormalization of f . The satellite case has been discussed in §4.3 and §4.4. Let us assume that f_1 is primitive ql. The renormalization $f \mapsto f_1$ is encoded by a maximal primitive copy $\mathcal{M}_1 \subset \mathcal{M}$. There is a canonical sequence of satellite copies of \mathcal{M}

$$\mathcal{M} \equiv \mathcal{M}^{(0)} \supsetneq \mathcal{M}^{(1)} \supsetneq \cdots \supsetneq \mathcal{M}^{(n)} \supsetneq \mathcal{M}^{(n+1)}, \quad \mathcal{M}_1 \subset \mathcal{M} \setminus \mathcal{M}^{(1)}, \quad n \geq 0$$

such that $\mathcal{M}^{(m)}$ is a maximal satellite subcopy of $\mathcal{M}^{(m-1)}$, \mathcal{M}_1 is in the $1/2$ -decoration of $\mathcal{M}^{(n+1)}$, and \mathcal{M}_1 is in a small (i.e., not the $1/2$) decoration of $\mathcal{M}^{(m)}$ for all $m \leq n$. The copy \mathcal{M}_1 is close to the main Molecule if and only if the period of $\mathcal{M}^{(n+1)}$ is big.

The puzzle levels of §4.2 provide a satisfactory control on scales between $\mathcal{M}^{(n)}$ and \mathcal{M}_1 (the latter copy is smaller) and fail on scales between \mathcal{M} and $\mathcal{M}^{(n)}$, which we refer as *virtual Molecule*. The virtual Molecule case naturally consists of two subcases: *virtual bounded-type satellite ql* (when $n \gg 1$ and the relative periods of $\mathcal{M}^{(s+1)}$ rel $\mathcal{M}^{(s)}$ are bounded) and *virtual near-Neutral* ($n = 0$).

Interpolation Problem 4.4. Develop a Virtual Molecule version of the Near-Degenerate Regime.

Problems 4.3, 4.4 and the arguments in §4.1, §4.2, §4.3 §4.4 should imply the full MLC.

Jointly with Kahn and Lyubich, we put forward a strategy to approach Problem 4.4 by considering *partially invariant virtual Julia sets* of $\mathcal{M}^{(s)}$ – they are connected hulls of the corresponding Cantor small Julia sets within J_f and contain the critical orbit only up to an appropriate number of first returns. *A posteriori*, bounds for virtual Julia sets can be deduced by assuming a uniform hyperbolicity of the renormalization associated with \mathcal{M} ; cf. Problem 3.1. The strategy towards Problem 4.4 is to develop such control *a priori*.

Acknowledgments. I would like to thank Misha Lyubich, Yusheng Luo, and Kostya Drach for many useful discussions regarding the context of this Note.

References

- [1] A. Avila, J. Kahn, M. Lyubich, and W. Shen. Combinatorial rigidity for unicritical polynomials. *Annals of Mathematics*, 170, 783–797, 2009.
- [2] A. Avila and M. Lyubich. Lebesgue measure of Feigenbaum Julia sets. *Annals of Mathematics*, 195, 1–88, 2022.
- [3] A. Avila, M. Lyubich, and W. de Melo. Regular or stochastic dynamics in real analytic families of unimodal maps. *Inventiones Mathematicae*, 154, 451–550, 2003.
- [4] A. Avila and C. G. Moreira. Statistical properties of unimodal maps: the quadratic family. *Annals of Mathematics*, 161, 831–881, 2005.
- [5] B. Branner and J. H. Hubbard. The iteration of cubic polynomials. Part II. *Acta Mathematica*, 169, 229–325, 1992.
- [6] X. Buff and A. Chéritat. Quadratic Julia sets of positive area. *Annals of Mathematics*, 176, 673–746, 2012.
- [7] J. Cepelawicz. The Quest to Decode the Mandelbrot Set, Math’s Famed Fractal. *Quanta Magazine*, 2024.
- [8] D. Cheraghi. Topology of irrationally indifferent attractors. Accepted for publication in *Annales Scientifiques de l’École Normale Supérieure*, 2025. To appear.
- [9] P. Coullet and C. Tresser. Itérations d’endomorphismes et groupe de renormalisation. *Journal de Physique, Colloque*, 39, 5–25, 1978.

- [10] A. Douady and J. H. Hubbard. Étude dynamique des polynômes complexes. Publications Mathématiques d'Orsay, 1984–1985. Orsay preprints 84-02 and 85-04.
- [11] A. Douady and J. H. Hubbard. On the dynamics of polynomial-like mappings. *Annales Scientifiques de l'École Normale Supérieure* (4), 18, 287–343, 1985.
- [12] A. Douady and J. H. Hubbard. A proof of Thurston's topological characterization of rational functions. *Acta Mathematica*, 171, 263–297, 1993.
- [13] K. Drach and D. Dudko. Near-Degenerate Regime in Transcendental Dynamics. Manuscript, 2025.
- [14] D. Dudko, J. Kahn, and M. Lyubich. Local connectivity of the Mandelbrot set along the real line. Manuscript, 2022.
- [15] D. Dudko and Y. Luo. Sierpiński carpet hyperbolic components of disjoint type are bounded. Preprint, arXiv:2509.25658, 2025.
- [16] D. Dudko and M. Lyubich. Uniform a priori bounds for neutral renormalization. Preprint, arXiv:2210.09280, 2022.
- [17] D. Dudko and M. Lyubich. Local connectivity of the Mandelbrot set at some satellite parameters of bounded type. *Geometric and Functional Analysis*, 33, 1–136, 2023.
- [18] D. Dudko and M. Lyubich. MLC at Feigenbaum points. Accepted for publication in *Publications Mathématiques de l'IHÉS*, 2025. To appear.
- [19] D. Dudko and M. Lyubich. Uniform a priori bounds for neutral renormalization. Variation I: Sector Renormalization. Manuscript, 2024.
- [20] D. Dudko, M. Lyubich, and N. Selinger. Pacman renormalization and self-similarity of the Mandelbrot set near Siegel parameters. *Journal of the American Mathematical Society*, 33, 653–733, 2020.
- [21] M. Feigenbaum. Quantitative universality for a class of nonlinear transformations. *Journal of Statistical Physics*, 19, 25–52, 1978.
- [22] J. Graczyk and G. G. Świątek. Generic hyperbolicity in the logistic family. *Annals of Mathematics*, 146, 1–52, 1997.
- [23] J. H. Hubbard. Local connectivity of Julia sets and bifurcation loci: three theorems of J.-C. Yoccoz. In *Topological Methods in Modern Mathematics*, pages 467–511. Publish or Perish, Houston, 1992.
- [24] H. Inou and M. Shishikura. The renormalization for parabolic fixed points and their perturbations. Manuscript, <https://www.math.kyoto-u.ac.jp/~mitsu/pararenorm/>, 2008.
- [25] J. Kahn. A priori bounds for some infinitely renormalizable quadratics: I. Bounded primitive combinatorics. Preprint, arXiv:math/0609045, 2006.
- [26] J. Kahn. Hyperbolic Components and Limits of Extremal Length. MSRI Talks, <https://vimeo.com/704224013>, <https://vimeo.com/700148806>, 2022.
- [27] J. Kahn and M. Lyubich. A priori bounds for some infinitely renormalizable quadratics: II. Decorations. *Annales Scientifiques de l'École Normale Supérieure*, 41, 57–84, 2008.
- [28] J. Kahn and M. Lyubich. A priori bounds for some infinitely renormalizable quadratics: III. Molecules. In: *Complex Dynamics: Families and Friends* (ed.: D. Schleicher), A K Peters, Ltd., 2009.
- [29] J. Kahn and M. Lyubich. Local connectivity of Julia sets for unicritical polynomials. *Annals of Mathematics*, 170, 413–426, 2009.
- [30] J. Kahn and M. Lyubich. The quasi-additivity law in conformal geometry. *Annals of Mathematics*, 169, 561–593, 2009.

- [31] O. Kozlovski, W. Shen, and S. van Strien. Density of hyperbolicity in dimension one. *Annals of Mathematics*, 166, 145–182, 2007.
- [32] O. Kozlovski, W. Shen, and S. van Strien. Rigidity for real polynomials. *Annals of Mathematics*, 165, 749–841, 2007.
- [33] O. E. Lanford. A computer-assisted proof of the Feigenbaum conjectures. In *Universality in Chaos*, 2nd ed., pages 245–252. Routledge, 2017.
- [34] W. R. Lim. A priori bounds and degeneration of Herman rings with bounded type rotation number. *Inventiones Mathematicae*, 2025.
- [35] M. Lyubich. Dynamics of quadratic polynomials. I, II. *Acta Mathematica*, 178, 185–297, 1997.
- [36] M. Lyubich. Feigenbaum–Coullet–Tresser universality and Milnor’s Hairiness Conjecture. *Annals of Mathematics*, 149, 319–420, 1999.
- [37] M. Lyubich. Dynamics of quadratic polynomials III: parapuzzle and SBR measures. *Astérisque*, 261, 173–200, 2000.
- [38] M. Lyubich. Almost every real quadratic map is either regular or stochastic. *Annals of Mathematics*, 156, 1–78, 2002.
- [39] M. Lyubich and J. Milnor. The Fibonacci unimodal map. *Journal of the American Mathematical Society*, 6, 425–457, 1993.
- [40] C. T. McMullen. *Complex Dynamics and Renormalization*. Annals of Mathematics Studies, Vol. 135. Princeton University Press, Princeton, NJ, 1994.
- [41] C. T. McMullen. *Renormalization and Three Manifolds Which Fiber over the Circle*. Princeton University Press, Princeton, NJ, 1996.
- [42] C. T. McMullen. Self-similarity of Siegel disks and Hausdorff dimension of Julia sets. *Acta Mathematica*, 180, 247–292, 1998.
- [43] M. Shishikura. The Hausdorff dimension of the boundary of the Mandelbrot set and Julia sets. *Annals of Mathematics*, 147, 225–267, 1998.
- [44] M. Shishikura and F. Yang. The high type quadratic Siegel disks are Jordan domains. *Journal of the European Mathematical Society (JEMS)*, 27, 4501–4562, 2025.
- [45] D. Smania. Puzzle geometry and rigidity: the Fibonacci cycle is hyperbolic. *Journal of the American Mathematical Society*, 20, 629–673, 2007.
- [46] D. Smania. Solenoidal attractors with bounded combinatorics are shy. *Annals of Mathematics*, 191, 1–79, 2020.
- [47] D. Sullivan. Quasiconformal homeomorphisms and dynamics, topology and geometry. In *Proc. Int. Congr. Math. (ICM), Berkeley*, Vol. II, pages 1216–1228, 1986.
- [48] D. Sullivan. Bounds, quadratic differentials and renormalization conjectures. In *Mathematics Into the Twenty-first Century*, AMS Centennial Publications, Vol. 2, 1992.
- [49] L. Tan. Similarity Between the Mandelbrot Set and Julia Sets. *Communications in Mathematical Physics*, 134, 587–617, 1990.
- [50] W. P. Thurston. Hyperbolic structures on 3-manifolds I: Deformation of acylindrical manifolds. *Annals of Mathematics*, 124, 203–246, 1986.