

Gibbs conditioning principle
 for log-concave independent random variables
 Eric Cator
 Radboud University, Nijmegen
 Pablo A. Ferrari
 Universidad de Buenos Aires and Conicet

Abstract Let ν_1, ν_2, \dots be a sequence of probabilities on the nonnegative integers, and $\mathbf{X} = (X_1, X_2, \dots)$ be a sequence of independent random variables X_i with law ν_i . For $\lambda > 0$ denote $Z_i^\lambda := \sum_x \lambda^x \nu_i(x)$ and $\lambda^{\max} := \sup\{\lambda > 0 : Z_i^\lambda < \infty \text{ for all } i\}$, and assume $\lambda^{\max} > 1$. For $\lambda < \lambda^{\max}$, define the tilted probability $\nu_i^\lambda(x) := \lambda^x \nu_i(x) / Z_i^\lambda$, and let \mathbf{X}^λ be a sequence of independent variables X_i^λ with law ν_i^λ , and denote $S_n^\lambda := X_1^\lambda + \dots + X_n^\lambda$, with $S_n = S_n^1$. Choose $\lambda^* \in (1, \lambda^{\max})$ and denote $R_n^* := E(S_n^{\lambda^*})$. The Gibbs Conditioning Principle (GCP) holds if $P(\mathbf{X} \in \cdot | S_n > R_n^*)$ converges weakly to the law of \mathbf{X}^{λ^*} , as $n \rightarrow \infty$. We prove the GCP for log-concave ν_i 's, meaning $\nu_i(x+1) \nu_i(x-1) \leq (\nu_i(x))^2$, subject to a technical condition that prevents condensation. The canonical measures are the distributions of the first n variables, conditioned on their sum being k . Efron's theorem states that for log-concave ν_i 's, the canonical measures are stochastically ordered with respect to k . This, in turn, leads to the ordering of the conditioned tilted measures $P(\mathbf{X}^\lambda \in \cdot | S_n^\lambda > R_n^*)$ in terms of λ . This ordering is a fundamental component of our proof.

Keywords and phrases empirical measures , Gibbs conditioning principle, large deviations, log-concave random variables, Efron theorem.

AMS 2000 subject classifications 60F10, 82B26, 60K35

1 Introduction

Consider a sequence X_i of independent not necessarily identically distributed random variables taking values on \mathbb{Z}_+ , the nonnegative integers; denote ν_i the law of X_i and assume that ν_i is log-concave for all i . Let $\lambda^{\max} := \sup\{\lambda > 0 : \sup_i E(\lambda^{X_i}) < \infty\}$. For $\lambda \in (0, \lambda^{\max})$ let X_i^λ be a sequence of independent variables with marginals $P(X_i^\lambda = x) = \frac{1}{Z_i^\lambda} \lambda^x \nu_i(x)$, with $Z_i^\lambda := E(\lambda^{X_i})$. Consider the sums $S_n = X_1 + \dots + X_n$ and $S_n^\lambda = X_1^\lambda + \dots + X_n^\lambda$. Choose $\lambda^* \in (0, \lambda^{\max})$ and denote $R_n^* := E(S_n^{\lambda^*})$. To avoid condensation, we need a condition about the derivatives of the function $M_n(t) := \sum_{i=1}^n \log(Z_i^{e^t})$ around $t^* = \log \lambda^*$. We show that if $\lambda^* > 1$ then, the variables $(X_i)_{i \geq 1}$ conditioned to $S_n > R_n^*$ converge weakly (finite

dimensional distributions) to $(X_i^{\lambda^*})_{i \geq 1}$, as n grows to infinity. Analogous results hold for $\lambda^* < 1$ and/or the condition $S_n = \lfloor R_n^* \rfloor$.

This problem is well studied for independent and identically distributed (iid) variables Y_i . Let $\hat{\nu}_n$ be the empirical measure of the first n variables, $\hat{\nu}_n \varphi := \frac{1}{n}(\varphi(Y_1) + \dots + \varphi(Y_n))$, for test functions φ . Denote by Π a convex set of measures. In the case of iid random variables ($\nu_i \equiv \nu$), the Sanov property says that $\hat{\nu}_n$ satisfies $\lim_n \frac{1}{n} \log P(\hat{\nu}_n \in \Pi) = -D(\Pi|\nu)$, where $D(\Pi|\nu) = \inf_{\nu' \in \Pi} D(\nu'|\nu)$ and $D(\nu'|\nu) := \sum_x \log(\nu'(x)/\nu(x))\nu'(x)$. Vasicek [29] showed that the distribution of Y_1 given $\hat{\nu}_n \in \Pi$ converges to ν^* as $n \rightarrow \infty$, where ν^* satisfies $D(\Pi|\nu) = D(\nu^*|\nu)$, for some specific choices of Π . Czisar [4] shows that the variables (Y_1, \dots, Y_ℓ) converge in distribution to independent random variables with marginal distribution $Y_i \sim \nu^*$ for a wider set of models. This property was later called Gibbs conditioning principle by Dembo and Zeitouni [7], see also Seppäläinen and Rassoul-Agha [22]. Léonard and Najim [16] proved the GCP using the Sanov theorem. Diaconis and Freedman [9] consider the exponential family, and Liggett [17] the case of stable laws.

The Markov case was discussed by Bolthausen and Schmock [2], Dobrushin and Hrynniv [10] and Meda and Ney [19], see also Dembo and Zeitouni [6]. Messer and Spohn [20] and Deuschel, Stroock and Zessin [8] prove similar results for classical statistical mechanic models with pair interactions.

When there is no exponential moment the extra mass may go to a single site, a phenomenon known as condensation. See for instance Evans and Waclaw [12], Armendáriz and Loulakis [1]. Godrèche [14] and Ferrari, Landim and Sisko [13].

The proof of Theorem 1 has two ingredients. The first is Proposition 2 which establishes the stochastic order in λ of the conditioned tilted measures $P(\mathbf{X}_n^\lambda \in \cdot | S_n^\lambda > R_n)$. In turn this is a consequence of Efron theorem [11], which says that under log-concavity the canonical measures $\mu_n(\cdot|k) := P(\mathbf{X}_n \in \cdot | S_n = k)$ are stochastically ordered in k , for each n . The second ingredient is that the conditioned measures $P(\mathbf{X}_\ell^\lambda \in \cdot | S_n^\lambda > R_n)$ converge as $n \rightarrow \infty$ to the unconditioned measure $P(\mathbf{X}_\ell^\lambda \in \cdot)$ when the conditioning set gets full probability, $P(S_n^\lambda > R_n) \rightarrow_n 1$. This limit holds under condition (1), as we show in Lemma 3. Finally, the theorem follows from a sandwich argument, the conditioned measure is bounded above and below by unconditioned tilted measures, which converge to $P(\mathbf{X}_\ell^{\lambda^*} \in \cdot)$, as $\lambda \rightarrow \lambda^*$.

We do not make use of the entropy approach.

The paper is organized as follows. Section 2 introduces the notation and states the Gibbs conditioning principle for our setup, which is Theorem 1. In Section 3 we present the order in λ of the conditioned tilted measures and the convergence of the conditioned tilted measures to the unconditioned ones; we then show that Theorem 1 is a consequence of these results. Section 4 provides a coupling proof of Efron's theorem, and establishes the order of the conditioned tilted measures.

2 Gibbs conditioning principle

In this section we introduce notation and state our main result, Theorem 1.

Let ν_1, ν_2, \dots be probability measures on $\mathbb{Z}_+ := \{0, 1, 2, \dots\}$. Denote

$$Z_i^\lambda := \sum_z \lambda^z \nu_i(z), \quad \lambda^{\max} := \sup\{\lambda > 0 : Z_i^\lambda < \infty, \text{ for all } i\}.$$

For $\lambda \in (0, \lambda^{\max})$ define the λ -tilted measures ν_i^λ on \mathbb{Z}_+ by

$$\nu_i^\lambda(x) := \frac{\lambda^x \nu_i(x)}{Z_i^\lambda}.$$

Denote by $X_1^\lambda, X_2^\lambda, \dots$, independent random variables such that $X_i^\lambda \stackrel{\text{law}}{=} \nu_i^\lambda$. Define

$$S_n^\lambda := X_1^\lambda + \dots + X_n^\lambda, \quad S_n := S_n^1, \quad \mathbf{X}_n^\lambda := (X_1^\lambda, \dots, X_n^\lambda), \quad \mathbf{X}_n := \mathbf{X}_n^1.$$

Fix $\lambda^* \neq 1$ with $0 < \lambda^* < \lambda^{\max}$ and define for each $n \geq 1$

$$R_n^* := E(S_n^{\lambda^*}).$$

We are interested in the limiting distribution of (X_1, X_2, \dots) when we condition on $\{S_n > R_n^*\}$ if $\lambda^* > 1$, or on $\{S_n < R_n^*\}$ if $\lambda^* < 1$.

Before we can give our main result, we need a technical condition which is closely related to ensuring that the variance of $S_n^{\lambda^*}$ grows to infinity: define the functions

$$M_n(t) = \sum_{i=1}^n \log(Z_i^{e^t}) = \sum_{i=1}^n \log(E(e^{tX_i})).$$

Also define $t^* = \log(\lambda^*)$. It is well known that M_n is a convex function, and it is not hard to see that since $\lambda^* < \lambda^{\max}$, we have that M_n is twice differentiable at t^* , but we need a stronger condition around t^* :

$$\forall \varepsilon > 0 : \quad \begin{aligned} M_n(t^* - \varepsilon) - M_n(t^*) + \varepsilon M'_n(t^*) &\rightarrow +\infty, \text{ and} \\ M_n(t^* + \varepsilon) - M_n(t^*) - \varepsilon M'_n(t^*) &\rightarrow +\infty. \end{aligned} \tag{1}$$

Since $M''(t^*)$ is equal to the variance of $S_n^{\lambda^*}$, the condition is often implied by this variance going to $+\infty$, but this might not be enough: maybe the convex functions M_n approach another convex function with a discontinuity in the derivative at t^* . This might not satisfy condition (1).

Definition (Log-concavity). *Let ν_i be a probability measure on \mathbb{Z}_+ . Denote $\max_i = \sup\{x \in \mathbb{Z}_+ : \nu_i(x) > 0\}$, which could be infinite. We say that ν_i is log-concave if $\nu_i(x) > 0$ for all $x \in [0, \max_i] \cap \mathbb{Z}$, and*

$$\frac{\nu_i(x-1)}{\nu_i(x)} \leq \frac{\nu_i(x)}{\nu_i(x+1)}, \quad 0 < x < \max_i.$$

Theorem 1 (Gibbs conditioning principle). *Let ν_1, ν_2, \dots be a family of log-concave probability measures on \mathbb{Z}_+ . Let X_1, X_2, \dots be independent random variables with $X_i \stackrel{\text{law}}{=} \nu_i$. Choose $\lambda^* \in (0, \lambda^{\max})$, let $R_n^* = E(S_n^{\lambda^*})$ and assume Condition (1). Then, the following limits hold.*

$$\text{If } 1 < \lambda^* < \lambda^{\max}, \text{ then } P(\mathbf{X} \in \cdot \mid S_n > R_n^*) \xrightarrow[n \rightarrow \infty]{w} P(\mathbf{X}^{\lambda^*} \in \cdot), \quad (2)$$

$$\text{If } 0 < \lambda^* < 1, \text{ then } P(\mathbf{X} \in \cdot \mid S_n < R_n^*) \xrightarrow[n \rightarrow \infty]{w} P(\mathbf{X}^{\lambda^*} \in \cdot), \quad (3)$$

$$\text{If } 0 < \lambda^* < \lambda^{\max}, \text{ then } P(\mathbf{X} \in \cdot \mid S_n = \lfloor R_n^* \rfloor) \xrightarrow[n \rightarrow \infty]{w} P(\mathbf{X}^{\lambda^*} \in \cdot), \quad (4)$$

where $\xrightarrow[n \rightarrow \infty]{w}$ means convergence of the finite dimensional distributions, or weak convergence.

The theorem is proven at the end of Section 2. It is helpful to think of a continuous example where each ν_i corresponds to a normal random variable with expectation a_i and variance σ_i^2 . This means that

$$R_n^* = \sum_{i=1}^n a_i + \lambda^* \sum_{i=1}^n \sigma_i^2.$$

In this case Theorem 1 is true if $\sum_{i \geq 1} \sigma_i^2 = +\infty$. However, if $\sum_{i \geq 1} \sigma_i^2 < +\infty$, then the marginal of the first random variable does not converge to the tilted random variable $X_1^{\lambda^*}$, since in that case conditioning changes the variance of all variables, whereas tilting a normal random variable leaves the variance constant. Also Condition (1) is not true if the total variance is finite.

Log-concave distributions are connected with several areas of mathematics and statistics, see the survey by Saumard and Wellner [23].

3 Conditioned tilted measures

Stochastic order Consider the coordinatewise order in \mathbb{Z}^n , $\mathbf{x}_n \leq \mathbf{x}'_n$ if $x_i \leq x'_i, \forall i$. A measure μ' stochastically dominates μ , denoted $\mu \prec \mu'$, if there exists $\tilde{\mu}$ on $\mathbb{Z}^n \times \mathbb{Z}^n$ with marginals μ and μ' such that a vector $(\mathbf{X}, \mathbf{X}')$ with distribution $\tilde{\mu}$ satisfies $\mathbf{X} \leq \mathbf{X}'$ $\tilde{\mu}$ -a.s..

The next result says that conditioned tilted log-concave measures on \mathbb{Z}_+ are stochastically ordered on the tilt parameter λ . Denote $\mathbf{X}_n^\lambda := (X_1^\lambda, \dots, X_n^\lambda)$ and $\mathbf{X}_n = \mathbf{X}_n^1$.

Proposition 2 (Stochastic order of conditioned measures). *Let ν_1, ν_2, \dots be a family of log-concave probability measures on \mathbb{Z}_+ . Let X_1, X_2, \dots be independent random variables with $X_i \stackrel{\text{law}}{=} \nu_i$. Then, for each $n \geq 1$, $R_n < R_n^{\max} := \sup_{\lambda < \lambda^{\max}} E(S_n^\lambda)$, and $0 < \lambda < \lambda' < \lambda^{\max}$, the following stochastic dominations hold,*

$$P(\mathbf{X}_n^\lambda \in \cdot \mid S_n^\lambda > R_n) \prec P(\mathbf{X}_n^{\lambda'} \in \cdot \mid S_n^{\lambda'} > R_n), \quad (5)$$

$$P(\mathbf{X}_n^\lambda \in \cdot | S_n^\lambda < R_n) \prec P(\mathbf{X}_n^{\lambda'} \in \cdot | S_n^{\lambda'} < R_n), \quad (6)$$

$$P(\mathbf{X}_n^\lambda \in \cdot | S_n^\lambda < R_n) \prec P(\mathbf{X}_n^{\lambda'} \in \cdot | S_n^{\lambda'} > R_n). \quad (7)$$

The proposition is proven in Section 4.1 using two results, the first is Efron's Theorem [11], which states that the canonical measures $\mu_n(\cdot|k)$ of product measures with log-concave marginals are stochastically ordered in k . The second result says that the conditioned sum distributions $P(S_n^\lambda \in \cdot | S_n^\lambda \in I)$ are stochastically ordered in λ , for any interval I , a one-dimensional statement.

Convergence of conditioned tilted measures We now show that under Condition (1) the conditioned tilted measure converges to the unconditioned one as the conditioning set gets full probability. The following lemma does not use log-concavity.

Lemma 3. *Assume Condition (1). Then*

$$\text{If } \lambda \in (\lambda^*, \lambda^{\max}), \text{ then } P(\mathbf{X}^\lambda \in \cdot | S_n^\lambda > R_n^*) \xrightarrow[n \rightarrow \infty]{\text{w}} P(\mathbf{X}^\lambda \in \cdot), \quad (8)$$

$$\text{If } \lambda \in (0, \lambda^*), \text{ then } P(\mathbf{X}^\lambda \in \cdot | S_n^\lambda < R_n^*) \xrightarrow[n \rightarrow \infty]{\text{w}} P(\mathbf{X}^\lambda \in \cdot). \quad (9)$$

Proof. Suppose $\lambda > \lambda^*$. We want to prove that $P(S_n^\lambda > R_n^*) \rightarrow 1$. Define $t = \log(\lambda)$ and $t^* = \log(\lambda^*)$. Define $\varepsilon = t - t^*$.

$$P(S_n^\lambda \leq R_n^*) = P(e^{-\varepsilon S_n^\lambda} \geq e^{-\varepsilon R_n^*}) \leq \frac{E(e^{-\varepsilon S_n^\lambda})}{e^{-\varepsilon R_n^*}} = \frac{E(e^{(t-\varepsilon)S_n})}{e^{-\varepsilon R_n^*} E(e^{tS_n})}.$$

Therefore,

$$\begin{aligned} \log(P(S_n^\lambda \leq R_n)) &\leq M_n(t^*) - M_n(t^* + \varepsilon) + \varepsilon R_n^* \\ &= M_n(t^*) - M_n(t^* + \varepsilon) + \varepsilon M'_n(t^*) \rightarrow -\infty. \end{aligned}$$

For the last step we use Condition (1) and the fact that $M'_n(t^*) = E(S_n^{\lambda^*}) = R_n^*$. This proves (8). The proof for (9) is completely similar. \square

Proof of Theorem 1. We prove that the conditioned measure in (2) is dominated from above and below by measures converging to the distribution of \mathbf{X}^{λ^*} . Since we want the convergence of the finite dimensional distributions, it suffices to work with the marginal distribution of the vector \mathbf{X}_ℓ for each $\ell \geq 1$. The upperbound for $\lambda^* < \lambda < \lambda^{\max}$ and $n \geq \ell$ is obtained by

$$P(\mathbf{X}_\ell \in \cdot | S_n > R_n^*) \prec P(\mathbf{X}_\ell^\lambda \in \cdot | S_n^\lambda > R_n^*) \xrightarrow[n \rightarrow \infty]{} P(\mathbf{X}_\ell^\lambda \in \cdot) \xrightarrow[\lambda \rightarrow \lambda^*]{} P(\mathbf{X}_\ell^{\lambda^*} \in \cdot), \quad (10)$$

where the domination comes from (5) as $\mathbf{X}_n = \mathbf{X}_n^1$, the \lim_n from (8), and the \lim_λ from the continuity of $P(\mathbf{X}_\ell^\lambda \in \cdot)$ in λ . For the lowerbound, take $1 < \lambda < \lambda^*$ to get

$$P(\mathbf{X}_\ell \in \cdot \mid S_n > R_n^*) \succ P(\mathbf{X}_\ell^\lambda \in \cdot \mid S_n^\lambda > R_n^*) \xrightarrow[n \rightarrow \infty]{} P(\mathbf{X}_\ell^\lambda \in \cdot) \xrightarrow[\lambda \rightarrow \lambda_*]{} P(\mathbf{X}_\ell^{\lambda^*} \in \cdot), \quad (11)$$

where we used (7) for the domination, (9) for the \lim_n , and continuity for the \lim_λ . The bounds (10) and (11) show (2).

Proceed in the same way for the upperbound of (3), let $\lambda^* < \lambda < 1$,

$$P(\mathbf{X}_\ell \in \cdot \mid S_n < R_n^*) \prec P(\mathbf{X}_\ell^\lambda \in \cdot \mid S_n > R_n^*) \xrightarrow[n \rightarrow \infty]{} P(\mathbf{X}_\ell^\lambda \in \cdot) \xrightarrow[\lambda \rightarrow \lambda_*]{} P(\mathbf{X}_\ell^{\lambda^*} \in \cdot). \quad (12)$$

For the lowerbound, let $0 < \lambda < \lambda^*$,

$$P(\mathbf{X}_\ell \in \cdot \mid S_n < R_n^*) \succ P(\mathbf{X}_\ell^\lambda \in \cdot \mid S_n^\lambda < R_n^*) \xrightarrow[n \rightarrow \infty]{} P(\mathbf{X}_\ell^\lambda \in \cdot) \xrightarrow[\lambda \rightarrow \lambda_*]{} P(\mathbf{X}_\ell^{\lambda^*} \in \cdot). \quad (13)$$

The bounds (12) and (13) show (3).

For the canonical measure, observe that

$$P(\mathbf{X}_\ell \in \cdot \mid S_n \leq R_n) \prec P(\mathbf{X}_\ell \in \cdot \mid S_n = \lfloor R_n \rfloor) \prec P(\mathbf{X}_\ell \in \cdot \mid S_n > R_n),$$

so that (11) and (12) imply (4). \square

4 Measure dominations

In this section we show Proposition 2.

4.1 Mixture of canonical measures

Denote by π_n^λ the distribution of the sum S_n^λ and $\pi_n^\lambda(k|I)$ the distribution of the sum conditioned to belong to an integer set I ,

$$\pi_n^\lambda(k) := P(S_n^\lambda = k), \quad \pi_n^\lambda(k|I) := \frac{\pi_n^\lambda(k)}{\pi_n^\lambda(I)}.$$

Define the canonical measure $\mu_n(\cdot|k)$ as the distribution of \mathbf{X}_n conditioned to $S_n = k$,

$$\mu_n(\cdot|k) := \frac{1}{\pi_n^\lambda(k)} P(\mathbf{X}_n = \cdot \mid S_n = k). \quad (14)$$

The conditioned tilted measures are mixture of canonical measures,

$$P(\mathbf{X}_n^\lambda = \cdot \mid S_n^\lambda \in I) = \sum_{k \in I} \pi_n^\lambda(k|I) \mu_n(\cdot|k), \quad I \subset \mathbb{Z}_+.$$

In statistical mechanics this is called the grand canonical measure. The proof of Proposition 2 is based on the next two results.

Lemma 4 (Order of one-dimensional tilted measures). *Let $0 < \lambda \leq \lambda' < \lambda^{\max}$ and consider an integer interval $I \subset [0, R_n^{\max}] \cap \mathbb{Z}$. Then,*

$$\pi_n^\lambda(\cdot | I) \prec \pi_n^{\lambda'}(\cdot | I) \quad (15)$$

This Lemma, proven in Section 4.2, does not require log concavity.

Theorem 5 (Efron [11]. Order of canonical measures). *Consider independent random variables X_i with log-concave marginals $X_i \stackrel{\text{law}}{=} \nu_i$. Then, the canonical measures are stochastically ordered,*

$$\mu_n(\cdot | k) \prec \mu_n(\cdot | k+1), \quad k \geq 0, \quad n \geq 1.$$

We provide a coupling proof in Section 4.3.

Proof of Proposition 2. Take $\lambda \in (\lambda^*, \lambda^{\max})$ and denote $I_n := (R_n^*, R_n^{\max}] \cap \mathbb{Z}$. Use Lemma 4 to construct a coupling (K, K') satisfying that K and K' have marginal distributions $\pi_n^\lambda(\cdot | I_n)$, $\pi_n^{\lambda'}(\cdot | I_n)$, respectively, and $K \leq K'$. Use Theorem 5 to construct a coupling $(Y^k)_{k \geq 0}$ such that the marginal Y^k has canonical distribution $\mu_n(\cdot | k)$, and $Y^k \leq Y^{k+1}$ for all $k \geq 1$. (In both cases we have dropped the dependence on n , to simplify notation).

Consider the vector (K, K') independent of the family $(Y^k)_{k \geq 0}$ and notice that the marginals of the coupling $(Y^K, Y^{K'})$ satisfy

$$Y^K \stackrel{\text{law}}{=} P(X_n^\lambda \in \cdot | S_n^\lambda > R_n^*), \quad Y^{K'} \stackrel{\text{law}}{=} P(X_n^{\lambda'} \in \cdot | S_n^{\lambda'} > R_n^*).$$

Furthermore, $Y^K \leq Y^{K'}$. This proves (5).

The same proof works for (6). To see (7) observe that both measures are mixtures of canonical distributions, and the measure conditioned to $S_n^\lambda < R_n$ concentrates on canonicals with $k < R_n$, while the measure on the right concentrates on canonicals with $k > R_n$, so that the order of the canonicals implies the order of those measures. \square

4.2 Order of one dimensional tilted measures

Here we prove Lemma 4, a general one-dimensional result. Let π be a probability on \mathbb{Z}_+ , denote $Z^\lambda := \sum_k \lambda^k \pi(k)$ and $\lambda^{\max} := \sup\{\lambda : Z^\lambda < \infty\}$. For $\lambda < \lambda^{\max}$ and $I \subset \mathbb{Z}_+$, define

$$\pi^\lambda(k) := \frac{1}{Z^\lambda} \lambda^k \pi(k), \quad \pi^\lambda(k|I) := \frac{1}{\pi^\lambda(I)} \pi^\lambda(k) \mathbb{1}\{k \in I\}.$$

We will show that when I is a finite or semi infinite interval,

$$\pi^\lambda(\cdot | I) \prec \pi^{\lambda'}(\cdot | I), \quad 0 < \lambda \leq \lambda' < \lambda^{\max}.$$

Take $\lambda < \lambda^{\max}$ and consider a birth and death process $K^\lambda(t)$ with rates

$$q^\lambda(k, k+1) := \lambda, \quad q^\lambda(k+1, k) = q(k+1, k) := \frac{\pi(k)}{\pi(k+1)}, \quad k \geq 0;$$

the rates vanish otherwise. Notice that the death rates do not depend on λ . The measure π^λ satisfies the detailed balance equations,

$$\pi^\lambda(k) q^\lambda(k, k+1) = \frac{\lambda^k}{Z^\lambda} \pi(k) \lambda = \frac{\lambda^{k+1}}{Z^\lambda} \pi(k+1) \frac{\pi(k)}{\pi(k+1)} = \pi^\lambda(k+1) q^\lambda(k+1, k),$$

so that it is reversible for the process $K^\lambda(t)$. Let $K^{\lambda, I}(t)$ be the process constrained on $I \subset \mathbb{Z}_+$, with rates

$$q^{\lambda, I}(k, \check{k}) := q^\lambda(k, \check{k}) \mathbb{1}\{\check{k} \in I\}, \quad k \in I.$$

The conditioned measure $\pi^\lambda(\cdot | I)$ is reversible for $q^{\lambda, I}$, indeed, for $k, \check{k} \in I$,

$$\pi^\lambda(k|I) q^{\lambda, I}(k, \check{k}) = \frac{\pi^\lambda(k)}{\pi^\lambda(I)} q^\lambda(k, \check{k}) = \frac{\pi^\lambda(\check{k})}{\pi^\lambda(I)} q^\lambda(\check{k}, k) = \pi^\lambda(\check{k}|I) q^{\lambda, I}(\check{k}, k).$$

We now couple two birth and death processes with different λ 's.

Lemma 6. *There exists a coupling*

$$(\tilde{K}(t))_{t \geq 0} := (K(t), K'(t))_{t \geq 0}$$

with marginals distributed as $(K^{\lambda, I}(t))_{t \geq 0}$ and $(K^{\lambda', I}(t))_{t \geq 0}$, respectively, such that

$$K(0) \leq K'(0) \quad \text{implies} \quad K(t) \leq K'(t). \quad (16)$$

Proof. The death rate is the same for both marginals, while the birth rates are $\lambda < \lambda'$. Coupling: independent marginal jumps when $k \neq k' \in I$. When $k = k'$ both marginals jump down together at rate $q(k, k-1) \mathbb{1}\{k-1 \in I\}$, they jump up together at rate $\lambda \mathbb{1}\{k+1 \in I\}$, and the second marginal jumps up alone at rate $(\lambda' - \lambda) \mathbb{1}\{k'+1 \in I\}$. The explicit rates, denoted \tilde{b} , are given for $k, k' \in I$ by

$$\begin{aligned} \tilde{b}((k, k'), (k+1, k'+1)) &= \lambda, & k = k', \quad k+1 \in I \\ \tilde{b}((k, k'), (k, k'+1)) &= \lambda' - \lambda, & k = k', \quad k'+1 \in I \\ \tilde{b}((k, k'), (k-1, k'-1)) &= q(k, k-1), & k = k', \quad k-1 \in I \\ \tilde{b}((k, k'), (k+1, k')) &= \lambda, & k \neq k', \quad k+1 \in I \\ \tilde{b}((k, k'), (k, k'+1)) &= \lambda', & k \neq k', \quad k'+1 \in I \\ \tilde{b}((k, k'), (k, k'-1)) &= q(k', k'-1), & k \neq k', \quad k'-1 \in I \\ \tilde{b}((k, k'), (k-1, k')) &= q(k, k-1), & k \neq k', \quad k-1 \in I, \end{aligned}$$

and the rates vanish otherwise. The reader can check that the marginal rates are $q^{\lambda, I}$, and $q^{\lambda', I}$, respectively, and that jumps violating order have rate zero. \square

Proof of Lemma 4. Denote $E_{0,0}$ the expectation associated to the coupling $\tilde{K}(t)$ with initial condition $\tilde{K}(0) = (0, 0)$. Denote \tilde{T} the return time to $(0, 0)$, that is, the hitting time of $(0, 0)$ after visiting another state, defined by

$$\tilde{T} := \inf\{t > 0 : \tilde{K}(t) = (0, 0) \text{ and } \tilde{K}(t-) \neq (0, 0)\}.$$

The measure $\tilde{\pi}$ defined by

$$\tilde{\pi}(k, k') = \frac{1}{E_{0,0}(\tilde{T})} E_{0,0}\left(\int_0^{\tilde{T}} \mathbb{1}\{\tilde{K}(t) = (k, k')\} dt\right) \quad (17)$$

is invariant for the coupling $\tilde{K}(t)$, see Proposition 2.59 of Liggett [18]. By (16), the integrand in (17) vanishes when $k > k'$. This implies that $\tilde{\pi}$ concentrates on $\{(k, k') : k \leq k'\}$, implying (15). \square

4.3 Order of canonical measures

In this section we show that the canonical measures are stochastically ordered under log-concavity, Theorem 5.

When the measures ν_i are absolutely continuous, the result was proven by Efron [11] for a set of functions called PF₂ (Polya Frequency functions), which coincides with the set of log-concave functions, as observed by Schoenberg [25]. See also Pitman [21] and Theorem 6.1 in Saumard and Wellner [24], for the continuous case. Efron mentions that his proof works for the discrete case, which was later proven dynamically, in the framework of queuing theory, see Shantikumar [27], Van der Wal [28], Daduna and Szekli [5] and their references.

Our proof uses an interacting particle system which conserves the number of particles and has the canonical measures $\mu_n(\cdot|k)$ as invariant measures. We construct a monotone coupling with ordered initial configurations having k and $k+1$ particles, respectively. The monotonicity is possible as log-concavity guarantees that the rate a particle attempt to leave from any site is given by a non decreasing function of the number of particles in the site. This coupling was tacitly used by Van der Wal, we explain here how to handle the case where the queues have limited capacity. The “polite customer” mentioned by Van der Wal corresponds to a second-class particle, a concept widely used in zero-range and exclusion processes.

Queuing system with finite waiting room For each i consider a maximal value $\max_i \in \mathbb{N} \cup \{\infty\}$ and a rate function $g_i : \mathbb{Z} \cap [0, \max_i] \rightarrow \mathbb{R}_+$, with $g_i(0) = 0$. Denote

$$\mathcal{X}_n := (\mathbb{Z} \cap [0, \max_1]) \times \cdots \times (\mathbb{Z} \cap [0, \max_n]).$$

Denote by $\mathbf{x} = (x_1, \dots, x_n)$ the elements of \mathcal{X}_n . Define a pure jump continuous time Markov process $\mathbf{X}(t)$ on \mathcal{X}_n with rates

$$r(\mathbf{x}, \mathbf{z}) := \begin{cases} g(x_i) & \text{if } \mathbf{z} = \theta^{ij}\mathbf{x}, \text{ and } x_j < \max_j, \\ 0 & \text{else.} \end{cases}, \quad \mathbf{x}, \mathbf{z} \in \mathcal{X}_n, \quad (18)$$

$$(\theta^{ij}\mathbf{x})_h := x_h - \mathbb{1}\{h = i\} + \mathbb{1}\{h = j\}, \quad h \in \{1, \dots, n\}.$$

When $\max_j = \infty$ for all j , the jump rate at i is $g_i(x_i)$, and we have just a non-homogeneous zero range process [15]. Otherwise, when $x_j = \max_j$ the rate of jumping to j is zero, and the resulting dynamics is a misanthrope process, see for instance [3] and [12]. Taking $\max_i \equiv k$ and $g_i(x) \equiv 1$, the misanthrope process coincides with the (homogeneous) k -exclusion process, see for instance [26].

The Basic coupling The basic coupling $\tilde{\mathbf{X}}(t) = (\mathbf{X}(t), \mathbf{X}'(t)) \in \mathcal{X}_n \times \mathcal{X}_n$ is defined as the pure jump Markov process with rates

$$\tilde{r}((\mathbf{x}, \mathbf{x}'), (\mathbf{z}, \mathbf{z}')) = \begin{cases} g(x_i) \wedge g(x'_i) & \text{if } (\mathbf{z}, \mathbf{z}') = (\theta^{ij}\mathbf{x}, \theta^{ij}\mathbf{x}'), x_j < \max_j, x'_j < \max_j, \\ g(x_i) \wedge g(x'_i) & \text{if } (\mathbf{z}, \mathbf{z}') = (\mathbf{x}, \theta^{ij}\mathbf{x}'), x_j = \max_j, x'_j < \max_j, \\ g(x_i) \wedge g(x'_i) & \text{if } (\mathbf{z}, \mathbf{z}') = (\theta^{ij}\mathbf{x}, \mathbf{x}'), x_j < \max_j, x'_j = \max_j, \\ (g(x'_i) - g(x_i))^+ & \text{if } (\mathbf{z}, \mathbf{z}') = (\mathbf{x}, \theta^{ij}\mathbf{x}'), x'_j < \max_j, \\ (g(x_i) - g(x'_i))^+ & \text{if } (\mathbf{z}, \mathbf{z}') = (\theta^{ij}\mathbf{x}, \mathbf{x}'), x_j < \max_j, \\ 0 & \text{else,} \end{cases}$$

The marginals $\mathbf{X}(t)$ and $\mathbf{X}'(t)$ of the basic coupling are Markov processes with rates (18). The next lemma shows that the coupling is monotone.

Lemma 7 (The basic coupling is monotone). *Assume $g_i(x) \leq g_i(x+1)$ for $x, x+1 \in [0, \max_i]$. Then, the basic coupling with rates \tilde{r} is monotone, that is,*

$$\mathbf{X}(0) \leq \mathbf{X}'(0) \text{ implies } \mathbf{X}(t) \leq \mathbf{X}'(t).$$

Proof. When $\mathbf{x} \leq \mathbf{x}'$ the rates \tilde{r} are zero in the second and fifth lines. In the other three cases the configurations after the jump are ordered $\mathbf{z} \leq \mathbf{z}'$. \square

Canonical measures and queuing system Let ν_i be a family of probability measures on \mathbb{Z}_+ , denote

$$\max_i := \sup\{z : \nu_i(z) > 0\} (\leq \infty),$$

assume

$$\nu_i(z) > 0, \quad \text{for } z \in \mathbb{Z} \cap [0, \max_i], \quad (19)$$

and denote

$$g_i(z) := \begin{cases} \frac{\nu_i(z-1)}{\nu_i(z)}, & z \in \mathbb{Z} \cap [0, \max_i] \\ 0, & \text{else.} \end{cases} \quad (20)$$

The assumption (19) is necessary to construct particle systems with rates (18) and invariant measure $\mu_n(\cdot|k)$. The condition is satisfied under log-concavity, but the next two lemmas do not require that.

Lemma 8. *Let ν_i satisfy (19). Then, for each $k \geq 0$ the canonical measure $\mu_n(\cdot|k)$ associated to ν_1, \dots, ν_n is reversible for the process with rates (18) and g_i given by (20).*

Proof. Let $\mathbf{x} \in \mathcal{X}_n$ with sum $x_1 + \dots + x_n = k$ and denote $\mathbf{z} := \theta^{ij}\mathbf{x}$. It suffices to show that for each k , $\mu_n(\cdot|k)$ satisfies the detailed balance equations,

$$\mu_n(\mathbf{x}|k) g_i(x_i) = \mu_n(\mathbf{z}|k) g_j(z_j). \quad (21)$$

Let $x_i = x + 1$ and $x_j = y$, so that $z_i = x$, $z_j = y + 1$. Recalling (14) and (20), (21) is equivalent to the identity

$$\frac{(\prod_{h \notin \{i,j\}} \nu_h(x)) \nu_i(x+1) \nu_j(y)}{\pi_n(k)} \frac{\nu_i(x)}{\nu_i(x+1)} = \frac{(\prod_{h \notin \{i,j\}} \nu_h(x)) \nu_i(x) \nu_j(y+1)}{\pi_n(k)} \frac{\nu_j(y)}{\nu_j(y+1)}. \quad \square$$

Proof of Theorem 5. Log-concavity of ν_i implies that the condition (19) holds. Then, by Lemma 8, the measure $\mu(\cdot|k)$ is reversible for the process with rates

$$g_i(y) = \frac{\nu_j(y-1)}{\nu_j(y)} \mathbb{1}\{y \leq \max_i\}.$$

By log-concavity, each g_i is non decreasing in the range $[0, \max_i]$, which in turn implies that there is a monotone coupling with rates \tilde{r} , by Lemma 7.

Let $k < k'$ and $\tilde{\mathbf{X}}(t) := (\mathbf{X}(t), \mathbf{X}'(t))$ be a coupling of zero range processes starting with k and k' particles, respectively. Let A' be the event $\{X'_1(0) = k'\}$ and $X'_j(0) = 0$ for all $j \neq 1\}$, “all particles of the \mathbf{X}' -th marginal are at site 1”, and similarly A for the \mathbf{X} marginal. Denote $\tilde{A} := A \cap A'$, the event “all particles of both marginals are at site 1”. Denote the return times to those events by

$$\begin{aligned} \tilde{T} &:= \inf\{t > 0 : \tilde{\mathbf{X}}(t) \in \tilde{A} \text{ and } \tilde{\mathbf{X}}(t-) \notin \tilde{A}\}, \\ T' &:= \inf\{t > 0 : \mathbf{X}'(t) \in A' \text{ and } \mathbf{X}'(t-) \notin A'\}, \end{aligned}$$

Since the coupling is monotone, we have $\tilde{T} = T'$ which, using the argument in Lemma 4, shows that the coupling $\tilde{\mathbf{X}}(t)$ has a unique invariant measure $\tilde{\mu}((\cdot, \cdot) | k, k')$ with marginals $\mu(\cdot|k)$ and $\mu(\cdot|k')$ concentrating on the set of ordered marginals, $\tilde{\mu}_n(\mathbf{x} \leq \mathbf{x}' | k, k') = 1$, that is, $\mu_n(\cdot|k) \prec \mu_n(\cdot|k')$. \square

Acknowledgments

We thanks support by the Erasmus grant and Neuromat-FAPESP CEPID USP. We thank Milton Jara and Davide Gabrielli for enlightening discussions, and Senya Shlosman and Erwin Bolthausen for mentioning references about the Markovian case.

References

- [1] Inés Armendáriz and Michail Loulakis. Conditional distribution of heavy tailed random variables on large deviations of their sum. *Stochastic Processes Appl.*, 121(5):1138–1147, 2011.
- [2] Erwin Bolthausen and Uwe Schmock. On the maximum entropy principle for uniformly ergodic Markov chains. *Stochastic Process. Appl.*, 33(1):1–27, 1989.
- [3] Christiane Cocozza-Thivent. Processus des misanthropes. *Z. Wahrscheinlichkeitstheorie Verw. Geb.*, 70:509–523, 1985.
- [4] Imre Csiszár. Sanov property, generalized I -projection and a conditional limit theorem. *Ann. Probab.*, 12(3):768–793, 1984.
- [5] Hans Daduna and Ryszard Szekli. A queueing theoretical proof of increasing property of Polya frequency functions. *Stat. Probab. Lett.*, 26(3):233–242, 1996.
- [6] A. Dembo and O. Zeitouni. Refinements of the Gibbs conditioning principle. *Probab. Theory Relat. Fields*, 104(1):1–14, 1996.
- [7] Amir Dembo and Ofer Zeitouni. *Large deviations techniques and applications*. Boston, MA: Jones and Bartlett Publishers, 1993.
- [8] Jean-Dominique Deuschel, Daniel W. Stroock, and Hans Zessin. Microcanonical distributions for lattice gases. *Commun. Math. Phys.*, 139(1):83–101, 1991.
- [9] P. Diaconis and D. A. Freedman. Conditional limit theorems for exponential families and finite versions of de Finetti’s theorem. *J. Theor. Probab.*, 1(4):381–410, 1988.
- [10] R. Dobrushin and O. Hryniv. Fluctuations of shapes of large areas under paths of random walks. *Probab. Theory Related Fields*, 105(4):423–458, 1996.
- [11] Bradley Efron. Increasing properties of Pólya frequency functions. *Ann. Math. Statist.*, 36:272–279, 1965.
- [12] Martin Evans and Bartek Waclaw. Condensation in stochastic mass transport models: Beyond the zero-range process. *Journal of Physics A Mathematical and Theoretical*, 47, 02 2014.

- [13] Pablo A. Ferrari, Claudio Landim, and Valentin V. Sisko. Condensation for a fixed number of independent random variables. *J. Stat. Phys.*, 128(5):1153–1158, 2007.
- [14] Claude Godrèche. Condensation for random variables conditioned by the value of their sum. *J. Stat. Mech. Theory Exp.*, 2019(6):34, 2019. Id/No 063207.
- [15] Claude Kipnis and Claudio Landim. *Scaling limits of interacting particle systems*, volume 320 of *Grundlehren Math. Wiss.* Berlin: Springer, 1999.
- [16] Christian Léonard and Jamal Najim. An extension of Sanov’s theorem: Application to the Gibbs conditioning principle. *Bernoulli*, 8(6):721–743, 2002.
- [17] Thomas M. Liggett. An invariance principle for conditioned sums of independent random variables. *J. Math. Mech.*, 18:559–570, 1968.
- [18] Thomas M. Liggett. *Continuous time Markov processes. An introduction.*, volume 113 of *Grad. Stud. Math.* Providence, RI: American Mathematical Society (AMS), 2010.
- [19] Ana Meda and Peter Ney. The Gibbs conditioning principle for Markov chains. In *Perplexing problems in probability. Festschrift in honor of Harry Kesten*, pages 385–398. Boston: Birkhäuser, 1999.
- [20] Joachim Messer and Herbert Spohn. Statistical mechanics of the isothermal Lane-Emden equation. *J. Statist. Phys.*, 29(3):561–578, 1982.
- [21] Jim Pitman. Probabilistic bounds on the coefficients of polynomials with only real zeros. *J. Combin. Theory Ser. A*, 77(2):279–303, 1997.
- [22] Firas Rassoul-Agha and Timo Seppäläinen. *A course on large deviations with an introduction to Gibbs measures*, volume 162 of *Grad. Stud. Math.* Providence, RI: American Mathematical Society (AMS), 2015.
- [23] Adrien Saumard and Jon A. Wellner. Log-concavity and strong log-concavity: a review. *Stat. Surv.*, 8:45–114, 2014.
- [24] Adrien Saumard and Jon A. Wellner. Efron’s monotonicity property for measures on \mathbb{R}^2 . *J. Multivariate Anal.*, 166:212–224, 2018.
- [25] I. J. Schoenberg. On the zeros of the generating functions of multiply positive sequences and functions. *Ann. of Math. (2)*, 62:447–471, 1955.
- [26] Timo Seppäläinen. Existence of hydrodynamics for the totally asymmetric simple K -exclusion process. *Ann. Probab.*, 27(1):361–415, 1999.
- [27] J. George Shanthikumar. On stochastic comparison of random vectors. *J. Appl. Probab.*, 24:123–136, 1987.

- [28] J. van der Wal. Monotonicity of the throughput of a closed exponential queueing network in the number of jobs. *OR Spektrum*, 11(2):97–100, 1989.
- [29] Oldrich Alfons Vasicek. A conditional law of large numbers. *Ann. Probab.*, 8:142–147, 1980.