

# LIMIT THEOREMS FOR FIXED POINT BIASED PATTERN AVOIDING INVOLUTIONS

JUNGEUN PARK AND DOUGLAS RIZZOLO

**ABSTRACT.** We study fixed point biased involutions that avoid a pattern. For every pattern of length three we obtain limit theorems for the asymptotic distribution of the (appropriately centered and scaled) number of fixed points of a random fixed point biased involution avoiding that pattern. When the pattern being avoided is either 321, 132, or 213, we find a phase transition depending on the strength of the bias. We also obtain a limit theorem for distribution of fixed points when the pattern is  $123 \cdots k(k+1)$  for any  $k$  and partial results when the pattern is  $(k+1)k \cdots 321$ .

## 1. Introduction

If  $\pi$  and  $\sigma$  are permutations of  $[n] = \{1, \dots, n\}$  and  $[m]$  respectively with  $n > m$ , then  $\pi$  is said to contain  $\sigma$  if there exist indices  $i_1 < \dots < i_m$  such that for all  $j < k$ ,  $\pi(i_j) < \pi(i_k)$  if and only if  $\sigma(j) < \sigma(k)$ . The permutation  $\pi$  is said to avoid  $\sigma$  if it does not contain  $\sigma$ . Given a permutation  $\sigma$ , we let  $\mathbf{Av}_n(\sigma)$  be the set of permutations of  $[n]$  that avoid  $\sigma$  and  $\mathbf{Iv}_n(\sigma)$  be the subset of  $\mathbf{Av}_n(\sigma)$  of involutions. If  $\pi$  is a permutation of  $[n]$ , we call  $|\pi| = n$  the length of the permutation.

Recently there has been considerable interest in the study of random pattern avoiding permutations [1, 2, 4, 6, 9, 11, 14], and the fixed points of pattern avoiding permutations [5, 7, 8, 12, 13, 15]. Pattern avoiding involutions specifically have drawn attention [3, 7, 8, 15, 16] and are interesting because they shed light on the relationship between pattern avoidance and the structure of the symmetric group. In [15], the asymptotic distribution of the number of fixed points for uniformly random pattern avoiding involutions was studied for patterns of length 3 and monotone patterns. In this paper we extend these results to fixed point biased involutions.

Fixed point biased measures on pattern avoiding permutations were recently introduced in [5]. For a permutation  $\pi$  let  $\text{fp}(\pi)$  be the number of fixed points in  $\pi$  and  $q \in (0, \infty)$ , [5] introduced the measures  $\tilde{\mathbb{P}}_n^{q,\sigma}$  on  $\mathbf{Av}_n(\sigma)$  defined by

$$\tilde{\mathbb{P}}_n^{q,\sigma}(\pi) \propto q^{\text{fp}(\pi)}$$

and studied the distribution of  $\text{fp}(\pi)$  under  $\tilde{\mathbb{P}}_n^{q,\sigma}$  as  $n \rightarrow \infty$  when  $|\sigma| = 3$ . They were able to obtain limiting distributions except in the case  $\sigma \in \{231, 312\}$ .

We study the analogous problem for fixed point biased involutions. We define measures  $\mathbb{P}_n^{\sigma,q}$  on  $\mathbf{Iv}_n(\sigma)$  by

$$\mathbb{P}_n^{\sigma,q}(\pi) \propto q^{\text{fp}(\pi)}$$

and study the distribution of  $\text{fp}(\pi)$  under  $\mathbb{P}_n^{\sigma,q}$  as  $n \rightarrow \infty$  when either  $|\sigma| = 3$  or  $\sigma$  is a monotone pattern. The asymptotic distribution of  $\text{fp}(\pi)$  under  $\mathbb{P}_n^{1,\sigma}$ , which is the uniformly random case, for these choices of  $\sigma$  was studied in [15]. Our results extend those results to the fixed point biased case. In contrast to [5], we are able to obtain results for all patterns of length 3. Similar to [5], when  $\sigma \in \{321, 132, 213\}$  we find that there is a phase transition in the limiting distribution from Negative Binomial, to Rayleigh, to normal depending on the strength of the bias, see Theorem 3 below, however, the location of the phase transition is different.

## 2. Main Results

The following four theorems are our main results.

**Theorem 1.** Fix  $k \geq 1$ ,  $q > 0$ ,  $\sigma = 123 \cdots k(k+1)$ . Suppose that  $\Pi_n$  is a random element of  $\mathbf{I}_n(\sigma)$  distributed according to  $\mathbb{P}_n^{q,\sigma}$ . Then

$$\text{fp}(\Pi_{2n}) \rightarrow_d X_{\text{even}}^q$$

and

$$\text{fp}(\Pi_{2n-1}) \rightarrow_d X_{\text{odd}}^q$$

where  $X_{\text{even}}^q$  has probability mass function

$$\mathbb{P}(X_{\text{even}}^q = i) = \begin{cases} \frac{q^i}{2^{k-2}((q+1)^k + (q-1)^k)} \binom{k}{i} & i \text{ is even} \\ 0 & i \text{ is odd.} \end{cases}$$

and

$$\mathbb{P}(X_{\text{odd}}^q = i) = \begin{cases} \frac{q^i}{2^{k-2}((q+1)^k + (q-1)^k)} \binom{k}{i} & i \text{ is odd} \\ 0 & i \text{ is even.} \end{cases}$$

**Theorem 2.** Fix  $k \geq 1$ ,  $0 < q \leq 1$ ,  $\sigma = (k+1)k \cdots 321$ . Let  $q(n) = q^{\sqrt{k/n}}$ . Suppose that  $\Pi_n$  is a random element of  $\mathbf{I}_n(\sigma)$  distributed according to  $\mathbb{P}_n^{q(n),\sigma}$ . Let  $M$  be a random matrix drawn for the  $k \times k$  Gaussian Orthogonal Ensemble conditioned to have trace 0 and let  $\Lambda_1 \geq \Lambda_2 \geq \cdots \geq \Lambda_k$ . Define

$$S_k = \sum_{j=1}^k (-1)^{j+1} \Lambda_j,$$

and let  $\mathbb{P}_k$  be the distribution of  $\Lambda^k$ . Let  $X_k$  be a random variable with distribution  $\mathbb{P}_k$  that is absolutely continuous with respect to  $\mathbb{P}_k$  with density

$$\mathbb{P}_k(ds) = \frac{q^s}{\mathbb{E}[q^{S_k}]} d\mathbb{P}_k(ds).$$

(a) If  $k$  is even then

$$\sqrt{\frac{k}{n}} \text{fp}(\Pi_n) \rightarrow_d X_k.$$

(b) If  $k$  is odd then

$$\sqrt{\frac{k}{n}} \left( \text{fp}(\Pi_n) - \frac{n}{k} \right) \rightarrow_d X_k.$$

Note that Theorem 2 is qualitatively different from Theorem 1 and the results of [5] because  $q(n) \rightarrow 1$  as  $n \rightarrow \infty$  as opposed to being fixed. It would be interesting to investigate what happens when  $q$  is fixed, at the moment we are only able to do this in the case when  $k = 2$ .

**Theorem 3.** Fix  $\sigma \in \{321, 132, 213\}$ . Suppose that  $\Pi_n$  is a random element of  $\mathbf{I}_n(\sigma)$  distributed according to  $\mathbb{P}_n^{q,\sigma}$ .

(a) If  $0 < q < 1$  then

$$\text{fp}(\Pi_{2n}) \rightarrow_d N_1$$

where  $N_1$  is distributed like a Negative Binomial random variable with parameters  $p = 1 - q$  and  $r = 2$  conditioned to be even and

$$\text{fp}(\Pi_{2n+1}) \rightarrow_d N_2$$

where  $N_2$  is distributed like a Negative Binomial random variable with parameters  $p = 1 - q$  and  $r = 2$  conditioned to be odd.

(b) If  $q = 1$  then

$$\sqrt{\frac{1}{n}} \text{fp}(\Pi_n) \xrightarrow{d} \frac{1}{\sqrt{2}} X_2,$$

where  $X_2$  is as defined in Theorem 2. In fact,  $X_2/\sqrt{2}$  has a Rayleigh(1) distribution.

(c) If  $q > 1$  then

$$\frac{\text{fp}(\Pi_n) - \frac{q^2-1}{q^2+1}n}{\sqrt{\frac{4q^2}{q^2+1}n}} \xrightarrow{d} Z$$

where  $Z$  has a standard normal distribution.

For  $\sigma \in \{321, 132, 213\}$ , a similar phase transition was observed in [5] for fixed point biased permutations (not involutions), but in that case the phase transition happened at  $q = 3$ . In contrast, in our case the uniform case  $q = 1$  is the critical case.

**Theorem 4.** Fix  $\sigma \in \{231, 312\}$ . Suppose that  $\Pi_n$  is a random element of  $\mathbf{Iv}_n(\sigma)$  distributed according to  $\mathbb{P}_n^{q,\sigma}$ . Then

$$\frac{\text{fp}(\Pi_n) - \frac{q}{\sqrt{q^2+8}}n}{\sqrt{\frac{8q}{(8+q^2)^{3/2}}n}} \xrightarrow{d} Z$$

where  $Z$  has a standard normal distribution.

### 3. Proofs

An important observation that we will use in our proofs is that  $\mathbb{P}_n^{q,\sigma}$  is absolutely continuous with respect to  $\mathbb{P}_n^{1,\sigma}$  density

$$(1) \quad \mathbb{P}_n^{q,\sigma}(d\pi) = \frac{q^{\text{fp}(\pi)}}{\mathbb{E}_n^{1,\sigma} [q^{\text{fp}(\Pi)}]} \mathbb{P}_n^{1,\sigma}(d\pi).$$

The first few results follow quickly from Equation (1), the results of [15], and the continuous mapping theorem. The proofs of Theorems 3 and 4 require more calculation, but are based on standard techniques from analytic combinatorics.

*Proof of Theorem 1.* This follows immediately from Equation (1) and [15, Theorem 2] since  $\text{fp}(\Pi_n) \leq k$  (in particular, it is bounded).  $\blacksquare$

*Proof of Theorem 2.* This follows immediately from Equation (1) and [15, Theorem 1] since  $q < 1$  (in particular,  $f(x) = q^x$  for  $x \geq 0$  it is bounded and continuous).  $\blacksquare$

*Proof of Theorem 3.* When  $q = 1$  this is a direct consequence of Theorem 2.

For  $q \neq 1$ , we will approach the problem using analytic combinatorics. From [15], the bivariate generating function of involutions avoiding  $\sigma$  marked by their number of fixed points is

$$G(z, q) = \sum_n \sum_{\pi \in \mathbf{Iv}_n(\sigma)} q^{\text{fp}(\pi)} z^n = \frac{2}{1 - 2qz + \sqrt{1 - 4z^2}}.$$

Recall that  $[z^n]G(z, q)$  denotes the coefficient of  $z^n$  when treating  $G(z, q)$  as a formal power series in  $z$ , that is,

$$G(z, q) = \sum_{n=0}^{\infty} ([z^n]G(z, q)) z^n.$$

To understand the asymptotics of this function, we must understand its set of singularities. Due to the square root function, there are singularities at  $z = \pm 1/2$ . Furthermore, we see that subject to  $q > 0$  and  $|z| \leq 1/2$ , the equation

$$1 - 2qz + \sqrt{1 - 4z^2} = 0,$$

only has a solution for  $z$  when  $q \geq 1$ , in which case the solution is

$$\zeta = \frac{q}{q^2 + 1}.$$

Note that when  $q = 1$ ,  $|\zeta| = 1/2$  while  $|\zeta| < 1/2$  when  $q > 1$ . Thus the dominant singularity of  $G$  is at  $z = \pm 1/2$  for  $q \leq 1$  and at  $z = \zeta$  for  $q > 1$ . The singularity structure is the most complicated when  $q = 1$ , but fortunately for us, we have already addressed this case and need only work in the cases when  $q \neq 1$ .

Direct calculation shows that the probability generating function of  $\text{fp}(\Pi_n)$  is

$$p_n(u) = \mathbb{E}_n^{q,\sigma}[u^{\text{fp}(\Pi_n)}] = \frac{[z^n]G(z, uq)}{[z^n]G(z, q)}.$$

We now consider the case when  $q < 1$ . For any  $0 < \eta < 1$ , we define  $H_\eta(z) = G(z, \eta)$ . Direct calculation shows that at  $z = 1/2$

$$H_\eta(z) - \frac{2}{1-\eta} \sim \frac{-2\sqrt{2}}{(1-\eta)^2} \sqrt{1-2x}$$

and at  $z = -1/2$

$$H_\eta(z) - \frac{2}{1+\eta} \sim \frac{-2\sqrt{2}}{(1+\eta)^2} \sqrt{1+2x}.$$

Thus follows from [10, Theorem VI.5] (and the remark thereafter) that

$$[z^n]H_\eta(z) \sim \frac{2\sqrt{2}}{(1-\eta)^2} 2^n \left( \frac{1}{2\sqrt{\pi n^3}} \right) + \frac{2\sqrt{2}}{(1+\eta)^2} 2^n (-1)^n \left( \frac{1}{2\sqrt{\pi n^3}} \right).$$

Consequently,

$$p_n(u) = \frac{[z^n]H_{uq}(z)}{[z^n]H_q(z)} \sim \frac{\frac{1}{(1-uq)^2} + (-1)^n \frac{1}{(1+uq)^2}}{\frac{1}{(1-q)^2} + (-1)^n \frac{1}{(1+q)^2}}$$

Consequently

$$p_{2n}(u) \rightarrow \frac{\frac{1}{(1-uq)^2} + \frac{1}{(1+uq)^2}}{\frac{1}{(1-q)^2} + \frac{1}{(1+q)^2}}.$$

The function on the right hand side is the probability generating function of a Negative Binomial distribution with parameters  $p = 1 - q$  and  $r = 2$  conditioned to be even. Similarly

$$p_{2n+1}(u) \rightarrow \frac{\frac{1}{(1-uq)^2} - \frac{1}{(1+uq)^2}}{\frac{1}{(1-q)^2} - \frac{1}{(1+q)^2}},$$

and the function on the right hand side is the probability generating function of a Negative Binomial distribution with parameters  $p = 1 - q$  and  $r = 2$  conditioned to be odd.

For  $q > 1$ , we check the hypotheses of [10, Theorem IX.9]. In order for our notation to match that of [10, Theorem IX.9], we define  $F(z, u) = G(z, uq)$ , so that

$$p_n(u) = \frac{[z^n]F(z, u)}{[z^n]F(z, 1)}.$$

Note that there is an  $r$  such that  $F(z, 1)$  is meromorphic on a ball of radius  $r$  centered at the origin whose only pole is a simple pole at  $\rho = q/(q^2 + 1)$ . By rationalizing the denominator, we can write

$$F(z, u) = \frac{1 - 2uqz - \sqrt{1 - z^2}}{2z(z(u^2q^2 + 1) - uq)} = \frac{\frac{1-2uqz-\sqrt{1-4z^2}}{2z}}{z(u^2q^2 + 1) - uq}.$$

Letting

$$B(z, u) = \frac{1 - 2uqz - \sqrt{1 - 4z^2}}{2z} \quad \text{and} \quad C(z, u) = z(u^2q^2 + 1) - uq,$$

we have that  $F(z, u) = B(z, u)/C(z, u)$ . Note that the singularity of  $B$  at the origin is removable, so in fact  $B$  defines a bivariate analytic function on  $|z| < 1/2$ .

Letting

$$\rho(u) = \frac{uq}{u^2q^2 + 1}$$

we note that  $C(\rho(u), u) = 0$ . Define  $f(u) = \rho(1)/\rho(u)$ . By direct calculation, we find that

$$f'(1) = \frac{q^2 - 1}{q^2 + 1} \quad \text{and} \quad f''(1) + f'(1) - (f'(1))^2 = \frac{4q^2}{q^2 + 1}.$$

From [10, Theorem IX.9] it follows that

$$\mathbb{E}_n^{q, \sigma}[\text{fp}(\Pi_n)] = \frac{q^2 - 1}{q^2 + 1}n + O(1) \quad \text{and} \quad \text{Var}(\text{fp}(\Pi_n)) = \frac{4q^2}{q^2 + 1}n + O(1),$$

and

$$\frac{\text{fp}(\Pi_n) - \frac{q^2-1}{q^2+1}n}{\sqrt{\frac{4q^2}{q^2+1}n}} \rightarrow Z$$

where  $Z$  has a standard normal distribution. ■

*Proof of Theorem 4.* From [15], the bivariate generating function of involutions avoiding  $\sigma$  marked by their number of fixed points is

$$G(z, q) = \sum_n \sum_{\pi \in \mathbf{Iv}_n(\sigma)} q^{\text{fp}(\pi)} z^n = \frac{1 - z^2}{1 - 2z^2 - qz}.$$

Direct calculation shows that the probability generating function of  $\text{fp}(\Pi_n)$  is

$$p_n(u) = \mathbb{E}_n^{q, \sigma}[u^{\text{fp}(\Pi_n)}] = \frac{[z^n]G(z, uq)}{[z^n]G(z, q)}.$$

To prove our result, we check the hypotheses of [10, Theorem IX.9]. In order for our notation to match that of [10, Theorem IX.9], we define  $F(z, u) = G(z, uq)$ , so that

$$p_n(u) = \frac{[z^n]F(z, u)}{[z^n]F(z, 1)}.$$

Letting  $B(z, u) = 1 - z^2$  and  $C(z, u) = 1 - 2z^2 - quz$ , we have that  $F(z, u) = B(z, u)/C(z, u)$  where  $B$  and  $C$  are analytic. Furthermore, letting

$$\rho(u) = \frac{-qu + \sqrt{q^2u^2 + 8}}{4} \quad \text{and} \quad r(u) = \frac{-qu - \sqrt{q^2u^2 + 8}}{4}$$

we see that for  $u$  close to 1,  $0 < \rho(u) < |r(u)|$  and  $C(z, u)$  factors as

$$C(z, u) = (z - \rho(u))(z - r(u)).$$

Consequently  $C(z, 1)$  has a simple root at  $\rho(1)$  and  $F(z, 1)$  is meromorphic on a ball of radius  $q/4$  centered at the origin whose only singularity in that ball is a simple pole at  $\rho(1)$ . Furthermore, Note that  $\rho(u)$  is a non-constant function that is analytic near  $u = 1$  and  $C(\rho(u), u) = 0$ . We may also directly verify that

$$\partial_z C(\rho(1), 1) \cdot \partial_u C(\rho(1), 1) \neq 0$$

and that, letting  $f(u) = \rho(1)/\rho(u)$ , we have that

$$f''(1) + f'(1) - (f'(1))^2 = \frac{8q}{(8+q^2)^{3/2}} \neq 0.$$

Consequently, [10, Theorem IX.9] implies that

$$\mathbb{E}[\text{fp}(\Pi_n)] = f'(1)n + O(1) = \frac{q}{\sqrt{q^2+8}}n + O(1) \quad \text{and} \quad \text{Var}[\text{fp}(\Pi_n)] = \frac{8q}{(8+q^2)^{3/2}}n + O(1),$$

and

$$\frac{\text{fp}(\Pi_n) - \frac{q}{\sqrt{q^2+8}}n}{\sqrt{\frac{8q}{(8+q^2)^{3/2}}n}} \rightarrow_d Z$$

where  $Z$  has a standard normal distribution. ■

## References

- [1] Michael Albert, Mathilde Bouvel, Valentin Féray, and Marc Noy. A logical limit law for 231-avoiding permutations. *Discrete Mathematics & Theoretical Computer Science*, 26(Special issues), 2024.
- [2] Frédérique Bassino, Mathilde Bouvel, Valentin Féray, Lucas Gerin, and Adeline Pierrot. The Brownian limit of separable permutations. *The Annals of Probability*, 46(4):2134–2189, 2018.
- [3] Miklós Bóna, Cheyne Homberger, Jay Pantone, and Vincent Vatter. Pattern-avoiding involutions: exact and asymptotic enumeration. *Australasian Journal of Combinatorics*, 64(1):88–119, 2016.
- [4] Jacopo Borga and Mickaël Maazoun. Scaling and local limits of Baxter permutations and bipolar orientations through coalescent-walk processes. *The Annals of Probability*, 50(4):1359–1417, 2022.
- [5] Aksheytha Chelikavada and Hugo Panzo. Limit theorems for fixed point biased permutations avoiding a pattern of length three. *arXiv:2311.04623*, 2023.
- [6] Harry Crane, Stephen DeSalvo, and Sergi Elizalde. The probability of avoiding consecutive patterns in the mallows distribution. *Random Structures & Algorithms*, 53(3):417–447, 2018.
- [7] Emeric Deutsch, Aaron Robertson, and Dan Saracino. Refined restricted involutions. *European J. Combin.*, 28(1):481–498, 2007.
- [8] Sergi Elizalde and Igor Pak. Bijections for refined restricted permutations. *Journal of Combinatorial Theory, Series A*, 105:207–219, 2004.
- [9] Valentin Féray and Kelvin Rivera-Lopez. The permutoon limit of random recursive separable permutations. *Confluentes Mathematici*, 15:45–82, 2023.
- [10] Philippe Flajolet and Robert Sedgewick. *Analytic combinatorics*. Cambridge University Press, Cambridge, 2009.
- [11] Christopher Hoffman, Douglas Rizzolo, and Erik Slivken. Pattern-avoiding permutations and Brownian excursion Part I: shapes and fluctuations. *Random Structures & Algorithms*, 50(3):394–419, 2017.
- [12] Christopher Hoffman, Douglas Rizzolo, and Erik Slivken. Pattern-avoiding permutations and Brownian excursion, part II: fixed points. *Probab. Theory Related Fields*, 169(1-2):377–424, 2017.
- [13] Christopher Hoffman, Douglas Rizzolo, and Erik Slivken. Fixed points of 321-avoiding permutations. *Proceedings of the American Mathematical Society*, 147(2):861–872, 2019.
- [14] Svante Janson. Patterns in random permutations avoiding the pattern 132. *Combinatorics, Probability and Computing*, pages 24–51, 2017.
- [15] Samuel Miner, Douglas Rizzolo, and Erik Slivken. Asymptotic distribution of fixed points of pattern-avoiding involutions. *Discrete Mathematics & Theoretical Computer Science*, Vol. 19 no. 2, Permutation Patterns 2016, Dec 2017.
- [16] Rodica Simion and Frank Schmidt. Restricted permutations. *Euro. J. Combin.*, 6:383–406, 1985.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF DELAWARE  
*Email address:* `jungeun@udel.edu`

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF DELAWARE  
*Email address:* `drizzolo@udel.edu`