
THE MASCHKE OCTIC CONTAINS 96 PAIRWISE DISJOINT LINES

by

CÉDRIC BONNAFÉ

Abstract. — Miyaoka proved that a smooth surface of degree d in $\mathbf{P}^3(\mathbb{C})$ contains at most $2d(d-2)$ pairwise disjoint lines. In this note, we verify that the Maschke octic contains 96 pairwise disjoint lines, thereby proving that Miyaoka's bound is optimal for $d=8$.

Let i be a square root of -1 in the field \mathbb{C} of complex numbers. For better readability, we use dots to represent zeroes in sparse matrices (as is standard for character tables of finite groups). Let

$$s_1 = \begin{pmatrix} \cdot & 1 & \cdot & \cdot \\ 1 & \cdot & \cdot & \cdot \\ \cdot & \cdot & 1 & \cdot \\ \cdot & \cdot & \cdot & 1 \end{pmatrix}, \quad s_2 = \begin{pmatrix} 1 & \cdot & \cdot & \cdot \\ \cdot & \cdot & 1 & \cdot \\ \cdot & 1 & \cdot & \cdot \\ \cdot & \cdot & \cdot & 1 \end{pmatrix}, \quad s_3 = \begin{pmatrix} \cdot & -i & \cdot & \cdot \\ i & \cdot & \cdot & \cdot \\ \cdot & \cdot & 1 & \cdot \\ \cdot & \cdot & \cdot & 1 \end{pmatrix},$$

$$s_4 = \frac{1}{2} \begin{pmatrix} 1 & -1 & -1 & -1 \\ -1 & 1 & -1 & -1 \\ -1 & -1 & 1 & -1 \\ -1 & -1 & -1 & 1 \end{pmatrix} \quad \text{and} \quad s_5 = \begin{pmatrix} -1 & \cdot & \cdot & \cdot \\ \cdot & 1 & \cdot & \cdot \\ \cdot & \cdot & 1 & \cdot \\ \cdot & \cdot & \cdot & 1 \end{pmatrix}$$

be elements of $\mathbf{GL}_4(\mathbb{C})$. Then s_1, s_2, s_3, s_4 and s_5 are reflections of order 2: they generate a subgroup of $\mathbf{GL}_4(\mathbb{C})$ which we denote by G_{31} . It is well-known that G_{31} is finite and is the group denoted by G_{31} in Shephard-Todd classification of finite complex reflection groups [ShTo]. For its natural action on $V = \mathbb{C}^4$, it is irreducible and primitive.

We denote by (x, y, z, t) the dual basis of the canonical basis of V : the algebra $\mathbb{C}[V]$ of polynomial functions on V is the polynomial algebra $\mathbb{C}[x, y, z, t]$. If m is a monomial in x, y, z and t , we denote by $\Sigma_4(m)$ the sum of all the monomials obtained from m by permutation of these four variables. For instance,

$$\Sigma_4(xy) = xy + xz + xt + yz + yt + zt \quad \text{and} \quad \Sigma_4(xyzt) = xyzt.$$

We set

$$f = \Sigma_4(x^8) + 14 \Sigma_4(x^4y^4) + 168 x^2y^2z^2t^2.$$

Then, f is, up to a scalar, the unique invariant polynomial of G_{31} of degree 8. We denote by \mathcal{M} the surface in $\mathbf{P}(V) = \mathbf{P}^3(\mathbb{C})$ defined as the zero locus of f . It is straightforward to verify with MAGMA [Mag] that \mathcal{M} is smooth, which implies that the polynomial f and the surface \mathcal{M} are irreducible. The surface \mathcal{M} is called the *Maschke octic*.

The author is partly supported by the ANR (Project No ANR-24-CE40-3389, GRAW).

It was proved by Boissière and Sarti that \mathcal{M} contains 352 lines [BoiSa, Theo 3.1]. It was checked by Sarti and the author [BonSa, §6.5] that these 352 lines are divided into two G_{31} -orbits Ω_{160} and Ω_{192} of respective cardinality 160 and 192. More precisely, let L_{160} and L_{192} denote the lines in $\mathbf{P}^3(\mathbb{C})$ defined by

$$L_{160} : \begin{cases} 2x + (i+1)(\sqrt{3}+1)y = 0, \\ 2z + (i+1)(\sqrt{3}+1)t = 0 \end{cases} \quad \text{and} \quad L_{192} : \begin{cases} (i+1)(\sqrt{5}+1)x - 2(y+z) = 0, \\ 2(y-z) + (i-1)(\sqrt{5}+1)t = 0. \end{cases}$$

Here, \sqrt{n} denotes a square root of n (for $n \in \{3, 5\}$). Then L_{160} and L_{192} are contained in \mathcal{M} and Ω_k is the orbit of L_k (for $k \in \{160, 192\}$). A computer calculation with MAGMA allows to check easily the main result of this note:

Theorem. *The orbit Ω_{192} contains a family of 96 pairwise disjoint lines.*

Proof. — Let us give a few details about how this computation is made and some precision on the statement (again, everything is checked with MAGMA). First, let

$$a = \frac{1-i}{2} \begin{pmatrix} 1 & \cdot & \cdot & 1 \\ \cdot & 1 & 1 & \cdot \\ \cdot & -1 & 1 & \cdot \\ 1 & \cdot & \cdot & -1 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} \cdot & \cdot & -1 & \cdot \\ -1 & \cdot & \cdot & \cdot \\ \cdot & 1 & \cdot & \cdot \\ \cdot & \cdot & \cdot & 1 \end{pmatrix}.$$

Then $a, b \in G_{31}$. Let $G = \langle a, b \rangle$ and let Ω denote the G -orbit of the line L_{192} . Then $|G| = 1152$ and $|\Omega| = 96$. One can then check that, if $L, L' \in \Omega$ are such that $L \neq L'$, then $L \cap L' = \emptyset$. This shows that one can choose for the family of 96 pairwise distinct lines in Ω_{192} the orbit of L_{192} under the action of G . \square

Remarks. For $d \geq 5$, it is known that there exists a smooth surface of degree d in $\mathbf{P}^3(\mathbb{C})$ containing $d(d-2)+2$ pairwise disjoint lines [Ram2]⁽¹⁾. For $d = 8$, this gives a smooth octic surface with 50 pairwise disjoint lines: the above theorem improves drastically this record.

On the other hand, Miyaoka [Miy, §2.2] proved that a smooth surface of degree $d \geq 3$ in $\mathbf{P}^3(\mathbb{C})$ contains at most $2d(d-2)$ pairwise disjoint lines. The above theorem shows that Miyaoka's bound is optimal for $d = 8$.

Questions. (1) Miyaoka's bound is optimal for $d \in \{2, 3, 4, 8\}$. Is it optimal for all d ?

(2) Is the Maschke octic the unique octic surface of $\mathbf{P}^3(\mathbb{C})$ (up to the action of $\mathbf{GL}_4(\mathbb{C})$) containing 96 pairwise disjoint lines?

References

- [Bon] C. BONNAFÉ, *Some singular curves and surfaces arising from invariants of complex reflection groups*, Exp. Math. **30** (2021), 429-440.
- [BoiSa] S. BOISSIÈRE & A. SARTI, *Counting lines on surfaces*, Ann. Sc. Norm. Super. Pisa Cl. Sci. **6** (2007), 39-52.
- [BonSa] C. BONNAFÉ & A. SARTI, *Complex reflection groups and K3 surfaces II. The groups G_{29} , G_{30} and G_{31}* , J. Korean Math. Soc. **60** (2023), 695-743.
- [Mag] W. BOSMA, J. CANNON & C. PLAYOUST, *The Magma algebra system. I. The user language*, J. Symbolic Comput. **24** (1997), 235-265.

⁽¹⁾For d odd, this can be improved to $d(d-2)+4$; see [Ram1] for $d = 5$ and [BoiSa, Theo. 5.1] for $d \geq 7$.

- [Miy] Y. MIYAOKA, *The maximal number of quotient singularities on surfaces with given numerical invariants*, Math. Ann. **268** (1984), 159-171.
- [Ram1] S. RAMS, *Three-divisible families of skew lines on a smooth projective quintic*, Trans. Amer. Math. Soc. **354** (2002), 2359-2367.
- [Ram2] S. RAMS, *Projective surfaces with many skew lines*, Proc. Amer. Math. Soc. **133** (2005), 11-13.
- [ShTo] G. C. SHEPHARD & J. A. TODD, *Finite unitary reflection groups*, Canad. J. Math. **6** (1954), 274-304.

January 5, 2026

CÉDRIC BONNAFÉ, IMAG, Université de Montpellier, CNRS, Montpellier, France
E-mail : cedric.bonnafe@umontpellier.fr