

SPLITTING OF OPERATIONS FOR HOM-DIASSOCIATIVE ALGEBRAS AND HOM-TRIASSOCIATIVE ALGEBRAS

ABDELKADER HAMDOUNI¹, IMED BASDOURI², MARIEM JENDOUBI³, AHMED ZAHARI ABDOU⁴

ABSTRACT. Hom-quadri dendriform algebras and Hom-six-dendriform algebras are introduced and studied which is a splitting of a Hom-diassociative and Hom-triassociative algebras, respectively. Moreover we explore the connections between these categories of Hom-algebras. Finally We elaborate a classification of Hom-quadri-dendriform algebra in low dimension.

1. Introduction

The concept of diassociative algebras was first introduced by Loday in [21] inspired by his research on periodicity phenomena in algebraic K-theory and his study of Leibniz algebras. In [17] Loday and Ronco demonstrated that the family of chain modules over standard simplices can be endowed with an operad structure. The algebras over this operad are known as triassociative algebras characterized by three binary operations satisfying eleven defining identities. In the same work they introduced tridendriform algebras and established that triassociative algebras are Koszul dual to tridendriform algebras. Inspired by Loday's work [18] and Loday and Ronco [17]. Wen Teng [22] introduced new algebraic structures quadri-dendriform algebras and six-dendriform algebras as natural splittings of diassociative and triassociative algebras respectively. Dendriform algebras also introduced by Loday in [17, 5, 9, 10] decompose the associative product into two binary operations such that their sum defines an associative operation. These algebras are intimately connected to diassociative algebras via Koszul duality where the Koszul dual of the operad associated to diassociative algebras is precisely the operad of dendriform algebras. Dendriform algebras appear in several key areas of mathematics including algebraic K-theory, combinatorics (notably in the study of trees), cohomology and deformation theory. In recent years considerable attention has been devoted to Hom-type algebras due in part to their potential to provide a unified framework for constructing various types of natural deformations which are of interest in both mathematics and mathematical physics. Hom-dendriform algebras were introduced in [14, 4, 19] as twisted versions of dendriform algebras and as natural splittings of Hom-associative algebras. The notions of Hom-associative, Hom-diassociative and Hom-triassociative algebras which generalize classical associative structures by twisting the associativity condition through a linear map were developed in [26, 27, 13, 28]. Motivated by Loday's Ronco and Wen Teng [22, 18, 17] We introduce new algebraic structure Hom-quadri-dendriform algebras and Hom-six-dendriform algebras. The purpose of this paper is to introduce the notion of Hom-quadri dendriform algebras as a splitting of di-associative algebras and the notion of Hom-six-dendriform algebras as a splitting of tri-associative algebras. use relative averaging operators on Hom- algebras as a framework to construct and study these new Hom-algebraic structures. The idea of an averaging operator was first implicitly introduced by Reynolds in 1895. Throughout the twentieth century these operators were mainly investigated within the algebra of functions defined on a given space. In his doctoral dissertation In [2] Cao examined averaging operators

Date: January 5, 2026.

2010 Mathematics Subject Classification. 17A30 and 17A36 and 17B56 and 16W25 and 16S80 and 17D25.

Key words and phrases. Hom-dendriform algebra; Hom-Diassociative algebra; Hom-quadri-dendriform algebra; Hom-triassociative algebra; Hom-six-dendriform algebra, classification.

from both algebraic and combinatorial viewpoints. More recent research has further explored averaging operators on any binary operad highlighting their connections to concepts such as bisuccessors duplicators and Rota-Baxter operators in [23, 24, 11, 12]. The paper is structured as follows: In Section 1 we recall the basic definitions and properties of Hom-dendriform and Hom-diassociative algebras. In Section 2 we introduce the notion of Hom-quadri-dendriform algebras and define relative averaging operators on Hom-dendriform algebras. We show that such an operator gives rise to a Hom-quadri-dendriform structure and conversely that any such structure can be embedded into a Hom-dendriform algebra equipped with a relative averaging operator. In Section 3 we define Hom-six-dendriform algebras as splittings of Hom-triassociative algebras. We also demonstrate that homomorphic relative averaging operators yield Hom-six-dendriform structures. Finally we present a classification of low-dimensional Hom-quadri-dendriform algebras illustrating structural properties and examples.

Throughout this paper K denotes a field of characteristic zero. All the vector spaces and (multi)linear maps are taken over K .

2. Hom-dendriform algebra and Hom-diassociative algebra

In this section we introduce the notions of Hom-dendriform algebras and Hom-diassociative algebras generalizing the classical dendriform and diassociative algebras to Hom-algebras setting.

2.1. Hom-dendriform algebras. Dendriform algebras were introduced by Loday in [18]. Dendriform algebras are algebras with two operations which dichotomize the notion of associative algebra. We generalize now this notion by twisting the identities by a linear map.

Definition 2.1. A Hom-dendriform algebra is a quadruple $(D, \prec, \succ, \alpha)$ where D is a $\mathbb{K}$ -vector space the operations $\prec, \succ: D \times D \rightarrow D$ are bilinear maps and $\alpha: D \rightarrow D$ is a linear map (called the structure map) such that for all $x, y, z \in D$ the following identities hold:

$$\alpha(x) \prec (y \prec z + y \succ z) = (x \prec y) \prec \alpha(z) \quad (1)$$

$$\alpha(x) \succ (y \prec z) = (x \succ y) \prec \alpha(z) \quad (2)$$

$$\alpha(x) \succ (y \succ z) = (x \prec y + x \succ y) \succ \alpha(z). \quad (3)$$

$(D, \prec, \succ, \alpha)$ is said to be a multiplicative Hom-algebra if $\forall x, y \in D$ we have

$$\alpha(x \prec y) = \alpha(x) \prec \alpha(y) \quad \text{and} \quad \alpha(x \succ y) = \alpha(x) \succ \alpha(y).$$

Definition 2.2. Given two Hom-dendriform algebras $(D, \prec, \succ, \alpha)$ and $(D', \prec', \succ', \alpha')$. A Hom-dendriform homomorphism is a linear map $T: D \rightarrow D'$ that satisfies the following compatibility conditions $T(x \prec y) = T(x) \prec' T(y)$, $T(x \succ y) = T(x) \succ' T(y)$ and $T \circ \alpha = \alpha' \circ T$ for all $x, y \in D$.

In particular if T is bijective then $(D, \prec, \succ, \alpha)$ and $(D', \prec', \succ', \alpha')$ are said to be isomorphic.

Example 2.3. Let $(D, \prec, \succ, \alpha)$ be a Hom-dendriform algebra with a basis $\{e_1, e_2, e_3\}$ whose non-zero product are given by

$$D(\eta) = \begin{cases} e_1 \prec e_2 = \eta e_3 & e_2 \succ e_1 = e_3 \\ e_2 \prec e_2 = -\frac{1}{2}e_3 & e_2 \succ e_2 = e_1 + \frac{1}{4}e_3 \\ e_3 \prec e_2 = e_3 & e_3 \succ e_2 = \eta e_1 + e_3 \end{cases} \quad \alpha = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & b \end{pmatrix} \quad b, \eta \in \mathbb{R}.$$

Definition 2.4. Let $(D, \prec, \succ, \alpha)$ be a Hom-dendriform algebra. A representation of D is given by a vector space M together with four bilinear maps $\prec_l, \succ_l: D \times M \rightarrow M$, $\prec_r, \succ_r: M \times D \rightarrow M$ representing the left and right actions. These maps must satisfy the following nine compatibility conditions: $\forall x, y \in D, m \in M$ we have

(I) Left compatibility conditions

$$\begin{aligned}
(x \prec y) \prec_l \beta(m) &= \alpha(x) \prec_l (y \prec_l m + y \succ_l m), \\
(x \succ y) \prec_l \beta(m) &= \alpha(x) \succ_l (y \prec_l m), \\
\alpha(x) \succ_l (y \succ_l m) &= (x \prec y + x \succ y) \succ_l \beta(m).
\end{aligned}$$

(II) Right compatibility conditions

$$\begin{aligned}
\beta(m) \prec_r (x \prec y + x \succ y) &= (m \prec_r x) \prec_r \alpha(y), \\
\beta(m) \succ_r (x \prec y) &= (m \succ_r x) \prec_r \alpha(y), \\
(m \prec_r x + m \succ_r x) \succ_r \alpha(y) &= \beta(m) \succ_r (x \succ y).
\end{aligned}$$

(III) Mixed compatibility conditions

$$\begin{aligned}
(x \prec_l m) \prec_r \alpha(y) &= \alpha(x) \prec_l (m \prec_r y + m \succ_r y), \\
(x \succ_l m) \prec_r \alpha(y) &= \alpha(x) \succ_l (m \prec_r y), \\
(x \prec_l m + x \succ_l m) \succ_r \alpha(y) &= \alpha(x) \succ_l (m \succ_r y).
\end{aligned}$$

Any Hom-dendriform algebra $(D, \prec, \succ, \alpha)$ is a representation of itself with $\prec = \prec_l = \prec_r$ and $\succ = \succ_l = \succ_r$. This representation is known as the adjoint representation.

2.2. Hom-diassociative algebras. Diassociative algebras were introduced by Jean-Louis Loday [18] as algebras equipped with two associative multiplications that satisfy specific compatibility conditions. We generalize now this notion by twisting the identities by a linear map.

Definition 2.5. A Hom-diassociative algebras is a 4-tuple $(D, \dashv, \vdash, \alpha)$ consisting of a linear space D bilinear maps $\dashv, \vdash: D \times D \longrightarrow D$ and $\alpha: D \longrightarrow D$ satisfying for all $x, y, z \in D$ the following conditions:

$$(x \dashv y) \dashv \alpha(z) = \alpha(x) \dashv (y \dashv z) \quad (4)$$

$$(x \dashv y) \dashv \alpha(z) = \alpha(x) \dashv (y \vdash z) \quad (5)$$

$$(x \vdash y) \dashv \alpha(z) = \alpha(x) \vdash (y \dashv z) \quad (6)$$

$$(x \dashv y) \vdash \alpha(z) = \alpha(x) \vdash (y \vdash z) \quad (7)$$

$$(x \vdash y) \vdash \alpha(z) = \alpha(x) \vdash (y \vdash z). \quad (8)$$

Definition 2.6. A Hom-associative algebra is a triple (D, μ, α) where D is a $\mathbb{K}$ -vector space $\mu: D \times D \rightarrow D$, $\mu(x, y) = x \cdot y$ is a bilinear map and $\alpha: D \rightarrow D$ is a linear map such that the following Hom-associativity identity holds for all $x, y, z \in D$ we have $\alpha(x) \cdot (y \cdot z) = (x \cdot y) \cdot \alpha(z)$.

(D, μ, α) is said to be a multiplicative Hom-algebra if $\forall x, y \in D$ we have $\alpha(x \cdot y) = \alpha(x) \cdot \alpha(y)$.

Hom-diassociative algebras arise naturally from averaging operators on Hom-associative algebras:

Let (A, μ, α) be an Hom-associative algebra and H be an averaging operator of it.

In other words $H: A \rightarrow A$ is a linear map satisfying $\mu(Ha, Hb) = H\mu(a, Hb) = H\mu(Ha, b)$, $\forall a, b \in A$.

We generalize to Hom-algebras setting. the result given by Aguiar in [25] for diassociative algebra.

If H is an averaging operator on (A, μ, α) then the two operations $\dashv$ and $\vdash$ defined on A

$$a \dashv b = \mu(a, Hb) \quad a \vdash b = \mu(Ha, b)$$

define a Hom-diassociative algebra structure on A .

Definition 2.7. Let $(D, \dashv, \vdash, \alpha)$ be a Hom-diassociative algebra. A linear operator $R: D \rightarrow D$ is called a Rota-Baxter operator of weight zero if

$$R \circ \alpha = \alpha \circ R \quad (9)$$

$$R(x) \dashv R(y) = R(R(x) \dashv y + x \dashv R(y)) \quad (10)$$

$$R(x) \vdash R(y) = R(R(x) \vdash y + x \vdash R(y)) \quad (11)$$

for all $x, y \in D$.

Proposition 2.8. *Let $(D, \dashv, \vdash, \alpha)$ be a Hom-diassociative algebra and $R : D \longrightarrow D$ be a Rota-Baxter operator of weight 0 on D i.e R is linear and commutes with both α and*

$$\begin{aligned} R(x) \dashv R(y) &= R(R(x) \dashv y + x \dashv R(y)) \\ R(x) \vdash R(y) &= R(R(x) \vdash y + x \vdash R(y)). \end{aligned}$$

Then $(D, \preceq, \succeq, \alpha)$ is also Hom-diassociative algebra with:

$x \preceq y = R(x) \dashv y + x \dashv R(y)$ and $x \succeq y = R(x) \vdash y + x \vdash R(y)$, $\forall x, y \in D$.

Moreover R is a Rota-Baxter operator on $(D, \preceq, \succeq, \alpha)$.

Proof. We prove axiom (6) the other being proved in a similar way. Thus for any $x, y, z \in D$

$$\begin{aligned} (x \preceq y) \preceq \alpha(z) - \alpha(x) \preceq (y \succeq z) &= R(x \dashv R(y) + R(x) \dashv y) \dashv \alpha(z) + R(R(x) \dashv y + x \dashv R(y) \dashv \alpha(z) \\ &\quad - \alpha(x) \dashv R(R(y) \vdash z + y \vdash R(z)) + R(\alpha(x)) \dashv (R(y) \vdash z + y \vdash R(z)) \\ &= (x \dashv R(y)) \dashv R(\alpha(z)) + (R(x) \dashv y) \dashv R(\alpha(z)) + (R(x) \dashv R(y)) \dashv \alpha(z) \\ &\quad - \alpha(x) \dashv (R(y) \vdash R(z)) - R(\alpha(x)) \dashv (y \vdash R(z)) - R(\alpha(x)) \dashv (R(y) \vdash z) \\ &= 0 \end{aligned}$$

The left hand side vanishes by axiom (6). The ends the proof. $\square$

Example 2.9. Let $(D, \dashv, \vdash, \alpha)$ be a Hom-diassociative algebra with a basis $\{e_1, e_2, e_3\}$ whose non-zero product are given by

$$D = \begin{cases} e_1 \dashv e_2 = -2e_3 & e_1 \vdash e_1 = \frac{1}{4}e_3 \\ e_2 \dashv e_1 = e_3 & e_1 \vdash e_2 = e_3 \\ e_2 \dashv e_2 = e_3 & e_2 \vdash e_2 = e_3 \end{cases} \quad \alpha = \begin{pmatrix} a & 1 & 0 \\ 0 & a & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad a \in \mathbb{R}.$$

We have the operator Rota-Baxter:

$$\begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & r_{32} & r_{33} \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & r_{32} & 0 \end{pmatrix} \quad \begin{pmatrix} 2r_{33} & 0 & 0 \\ 0 & 2r_{33} & 0 \\ 0 & r_{32} & r_{33} \end{pmatrix}, \quad r_{ij} \in \mathbb{R}.$$

3. Hom-quadri-dendriform algebra

In this section we introduce the notions of Hom-quadri-dendriform algebras generalizing the classical quadri-dendriform algebras to Hom-algebras setting. Next we introduce relative averaging operators on Hom-dendriform algebras. We show that a relative averaging operator induces a Hom-quadri-dendriform algebra. Finally we show that any Hom-quadri-dendriform algebra can be embedded into an averaging Hom-dendriform algebra.

Definition 3.1. A Hom-quadri-dendriform algebra is a tuple $(D, \prec_+, \prec_-, \succ_+, \succ_-, \alpha)$ where D is a vector space the operations $\prec_+, \prec_-, \succ_+, \succ_- : D \times D \rightarrow D$ are bilinear and $\alpha : D \rightarrow D$ is a linear map such that for all $x, y, z \in D$ the following identities hold

$$(x \prec_+ y) \prec_+ \alpha(z) = (x \prec_- y) \prec_+ \alpha(z) = \alpha(x) \prec_+ (y \prec_- z + y \succ_+ z) \quad (12)$$

$$(x \succ_+ y) \prec_- \alpha(z) = (x \succ_- y) \prec_- \alpha(z) = \alpha(x) \succ_+ (y \prec_- z) \quad (13)$$

$$\alpha(x) \succ_+ (y \succ_+ z) = (x \prec_+ y + x \succ_+ y) \succ_+ \alpha(z) = (x \prec_- y + x \succ_- y) \succ_+ \alpha(z) \quad (14)$$

$$\alpha(x) \succ_- (y \succ_- z) = (x \prec_- y + x \succ_- y) \succ_- \alpha(z) = (x \prec_+ y + x \succ_+ y) \succ_- \alpha(z) \quad (15)$$

$$(x \prec_{\vdash} y) \prec_{\dashv} \alpha(z) = \alpha(x) \prec_{\vdash} (y \prec_{\dashv} z + y \succ_{\dashv} z) \quad (16)$$

$$(x \succ_{\vdash} y) \prec_{\dashv} \alpha(z) = \alpha(x) \succ_{\vdash} (y \prec_{\dashv} z) \quad (17)$$

$$\alpha(x) \succ_{\vdash} (y \succ_{\dashv} z) = (x \prec_{\vdash} y + x \succ_{\vdash} y) \succ_{\dashv} \alpha(z) \quad (18)$$

$$(x \prec_{\dashv} y) \prec_{\dashv} \alpha(z) = \alpha(x) \prec_{\dashv} (y \prec_{\vdash} z + y \succ_{\vdash} z) = \alpha(x) \prec_{\dashv} (y \prec_{\dashv} z + y \succ_{\dashv} z) \quad (19)$$

$$(x \prec_{\dashv} y) \prec_{\dashv} \alpha(z) = \alpha(x) \prec_{\dashv} (y \prec_{\vdash} z + y \succ_{\dashv} z) = \alpha(x) \prec_{\dashv} (y \prec_{\dashv} z + y \succ_{\vdash} z) \quad (20)$$

$$(x \succ_{\dashv} y) \prec_{\dashv} \alpha(z) = \alpha(x) \succ_{\dashv} (y \prec_{\vdash} z) = \alpha(x) \succ_{\dashv} (y \prec_{\dashv} z) \quad (21)$$

$$\alpha(x) \succ_{\dashv} (y \succ_{\vdash} z) = \alpha(x) \succ_{\dashv} (y \succ_{\dashv} z) = (x \prec_{\dashv} y + x \succ_{\dashv} y) \succ_{\dashv} \alpha(z) \quad (22)$$

$$(23)$$

$(D, \prec_{\vdash}, \prec_{\dashv}, \succ_{\vdash}, \succ_{\dashv}, \alpha)$ is said to be a multiplicative Hom-algebra if $\forall x, y, \in D$ we have

$$\begin{aligned} \alpha(x \prec_{\vdash} y) &= \alpha(x) \prec'_{\vdash} \alpha(y), & \alpha(x \prec_{\dashv} y) &= \alpha(x) \prec'_{\dashv} \alpha(y), \\ \alpha(x \succ_{\vdash} y) &= \alpha(x) \succ'_{\vdash} \alpha(y), & \alpha(x \succ_{\dashv} y) &= \alpha(x) \succ'_{\dashv} \alpha(y). \end{aligned}$$

Definition 3.2. Given two Hom-quadri-dendriform algebras $(D, \prec_{\vdash}, \prec_{\dashv}, \succ_{\vdash}, \succ_{\dashv}, \alpha)$ and $(D', \prec'_{\vdash}, \prec'_{\dashv}, \succ'_{\vdash}, \succ'_{\dashv}, \alpha')$. A Hom-quadri-dendriform homomorphism is a linear map $T : D \rightarrow D'$ that satisfies the following compatibility conditions

$$\begin{aligned} T(x \prec_{\vdash} y) &= T(x) \prec'_{\vdash} T(y), & T(x \prec_{\dashv} y) &= T(x) \prec'_{\dashv} T(y), \\ T(x \succ_{\vdash} y) &= T(x) \succ'_{\vdash} T(y), & T(x \succ_{\dashv} y) &= T(x) \succ'_{\dashv} T(y), \\ T \circ \alpha &= \alpha' \circ T. \end{aligned}$$

for all $x, y \in D$. In particular if T is bijective then

$(D, \prec_{\vdash}, \prec_{\dashv}, \succ_{\vdash}, \succ_{\dashv}, \alpha)$ and $(D', \prec'_{\vdash}, \prec'_{\dashv}, \succ'_{\vdash}, \succ'_{\dashv}, \alpha')$ are said to be isomorphic.

Proposition 3.3. Let $(D, \prec_{\vdash}, \prec_{\dashv}, \succ_{\vdash}, \succ_{\dashv}, \alpha)$ be an Hom-quadri-dendriform algebra. Then $(D, \vdash, \dashv, \alpha)$ is a Hom-diassociative algebra where

$$\begin{aligned} x \vdash y &= x \prec_{\vdash} y + x \succ_{\vdash} y \\ x \dashv y &= x \prec_{\dashv} y + x \succ_{\dashv} y \end{aligned}$$

for all $x, y \in D$.

Proof. $\forall x, y, z \in D$ we have

$$\begin{aligned} (x \dashv y) \dashv \alpha(z) &= (x \prec_{\dashv} y + x \succ_{\dashv} y) \succ_{\dashv} \alpha(z) + (x \prec_{\dashv} y + x \succ_{\dashv} y) \prec_{\dashv} \alpha(z) \\ &= (x \prec_{\dashv} y + x \succ_{\dashv} y) \succ_{\dashv} \alpha(z) + (x \prec_{\dashv} y) \prec_{\dashv} \alpha(z) + (x \succ_{\dashv} y) \prec_{\dashv} \alpha(z) \\ &= (x \prec_{\dashv} y + x \succ_{\dashv} y) \succ_{\dashv} \alpha(z) + \alpha(x) \prec_{\dashv} (y \prec_{\dashv} z + y \succ_{\dashv} z) + (x \succ_{\dashv} y) \prec_{\dashv} \alpha(z) \\ &= \alpha(x) \succ_{\dashv} (y \prec_{\dashv} z) + \alpha(x) \succ_{\dashv} (y \succ_{\dashv} z) + \alpha(x) \prec_{\dashv} (y \prec_{\dashv} z + y \succ_{\dashv} z) \\ &= \alpha(x) \dashv (y \dashv z), \end{aligned}$$

$$\begin{aligned} (x \dashv y) \dashv \alpha(z) &= (x \prec_{\dashv} y + x \succ_{\dashv} y) \succ_{\dashv} \alpha(z) + (x \prec_{\dashv} y) \prec_{\dashv} \alpha(z) \\ &= (x \prec_{\dashv} y + x \succ_{\dashv} y) \succ_{\dashv} \alpha(z) + \alpha(x) \prec_{\dashv} (y \prec_{\vdash} z + y \succ_{\vdash} z) + (x \succ_{\dashv} y) \prec_{\dashv} \alpha(z) \\ &= \alpha(x) \prec_{\dashv} (y \prec_{\vdash} z + y \succ_{\vdash} z) + \alpha(x) \succ_{\dashv} (y \prec_{\vdash} z) + \alpha(x) \succ_{\dashv} (y \succ_{\vdash} z) \\ &= \alpha(x) \dashv (y \vdash z), \end{aligned}$$

$$\begin{aligned}
(x \vdash y) \dashv \alpha(z) &= (x \prec_{\vdash} y + x \succ_{\vdash} y) \prec_{\dashv} \alpha(z) + (x \prec_{\vdash} y + x \succ_{\vdash} y) \succ_{\dashv} \alpha(z) \\
&= \alpha(x) \succ_{\vdash} (y \succ_{\dashv} z) + (x \prec_{\vdash} y) \prec_{\dashv} \alpha(z) + (x \succ_{\vdash} y) \succ_{\dashv} \alpha(z) \\
&= \alpha(x) \succ_{\vdash} (y \succ_{\dashv} z) + \alpha(x) \succ_{\vdash} (y \prec_{\dashv} z) + \alpha(x) \prec_{\vdash} (y \succ_{\dashv} z + y \prec_{\dashv} z) \\
&= \alpha(x) \vdash (y \dashv z),
\end{aligned}$$

$$\begin{aligned}
(x \dashv y) \vdash \alpha(z) &= (x \prec_{\dashv} y + x \succ_{\dashv} y) \prec_{\vdash} \alpha(z) + (x \prec_{\dashv} y + x \succ_{\dashv} y) \succ_{\vdash} \alpha(z) \\
&= (x \prec_{\dashv} y + x \succ_{\dashv} y) \prec_{\vdash} \alpha(z) + \alpha(x) \succ_{\vdash} (y \succ_{\vdash} z) \\
&= \alpha(x) \succ_{\vdash} (y \succ_{\vdash} z) + \alpha(x) \prec_{\vdash} (y \prec_{\vdash} z + y \succ_{\vdash} z) + \alpha(x) \succ_{\vdash} (y \succ_{\vdash} z) \\
&= \alpha(x) \vdash (y \vdash z),
\end{aligned}$$

$$\begin{aligned}
(x \vdash y) \vdash \alpha(z) &= (x \prec_{\vdash} y + x \succ_{\vdash} y) \prec_{\vdash} \alpha(z) + (x \prec_{\vdash} y + x \succ_{\vdash} y) \succ_{\vdash} \alpha(z) \\
&= (x \prec_{\vdash} y) \prec_{\vdash} \alpha(z) + (x \succ_{\vdash} y) \prec_{\vdash} \alpha(z) + \alpha(x) \succ_{\vdash} (y \succ_{\vdash} z) \\
&= \alpha(x) \prec_{\vdash} (y \prec_{\vdash} z + y \succ_{\vdash} z) + \alpha(x) \succ_{\vdash} (y \prec_{\vdash} z) + \alpha(x) \succ_{\vdash} (y \succ_{\vdash} z) \\
&= \alpha(x) \vdash (y \vdash z).
\end{aligned}$$

The end of the proof. □

Definition 3.4. Let $(D, \prec_{\vdash}, \prec_{\dashv}, \succ_{\vdash}, \succ_{\dashv}, \alpha)$ be a Hom-quadri-dendriform algebra and D_0 subset of D . We say that D_0 is a Hom-quadri-dendriform subalgebra of D if D_0 is stable under α , $\prec_{\vdash}$, $\prec_{\dashv}$, $\succ_{\vdash}$ and $\succ_{\dashv}$, i.e. $\forall x, y, z \in D_0$, $\alpha(x) \in D_0$, $x \prec_{\vdash} y \in D_0$, $x \prec_{\dashv} y \in D_0$, $x \succ_{\vdash} y \in D_0$ and $x \succ_{\dashv} y \in D_0$.

Example 3.5. Let $(D, \prec_{\vdash}, \prec_{\dashv}, \succ_{\vdash}, \succ_{\dashv}, \alpha)$ be a Hom-quadri-dendriform algebra. The vector subspace I_D spanned by

$$\{x \prec_{\dashv} y - x \prec_{\vdash} y, \quad x \succ_{\dashv} y - x \succ_{\vdash} y \mid x, y \in D\}$$

is an ideal of D . It is clear that D is a Hom-dendriform algebra if and only if $I_D = \{0\}$. Therefore the quotient algebra $(D/I_D, \prec, \succ, \bar{\alpha})$ is a Hom-dendriform algebra where $\bar{\alpha}(\bar{x}) = \overline{\alpha(x)}$, $\bar{x} \prec \bar{y} = \overline{x \prec_{\vdash} y} = \overline{x \prec_{\dashv} y}$, $\bar{x} \succ \bar{y} = \overline{x \succ_{\vdash} y} = \overline{x \succ_{\dashv} y}$, for all $x, y \in D$ where $\bar{x}$ denotes the class of an element $x \in D$.

Proposition 3.6. Let $(A, \prec_{\vdash_A}, \prec_{\dashv_A}, \succ_{\vdash_A}, \succ_{\dashv_A}, \alpha_A)$ and $(B, \prec_{\vdash_B}, \prec_{\dashv_B}, \succ_{\vdash_B}, \succ_{\dashv_B}, \alpha_B)$ be two Hom-quadri-dendriform algebras. Then there exists a Hom-quadri-dendriform algebra structure on $A \oplus B$ with the bilinear maps: $\triangleleft, \trianglelefteq, \triangleright, \trianglerighteq : (A \oplus B)^{\otimes 2} \longrightarrow A \oplus B$ given by

$$\begin{aligned}
(a_1 + b_1) \triangleleft (a_2 + b_2) &:= a_1 \prec_{\vdash_A} a_2 + b_2 \prec_{\vdash_B} b_2 \\
(a_1 + b_1) \trianglelefteq (a_2 + b_2) &:= a_1 \prec_{\dashv_A} a_2 + b_2 \prec_{\dashv_B} b_2 \\
(a_1 + b_1) \triangleright (a_2 + b_2) &:= a_1 \succ_{\vdash_A} a_2 + b_2 \succ_{\vdash_B} b_2 \\
(a_1 + b_1) \trianglerighteq (a_2 + b_2) &:= a_1 \succ_{\dashv_A} a_2 + b_2 \succ_{\dashv_B} b_2
\end{aligned}$$

and the linear map $\alpha = \alpha_A + \alpha_B : A \oplus B \longrightarrow A \oplus B$ given by

$$(\alpha_A + \alpha_B)(a + b) := \alpha_A(a) + \alpha_B(b), \quad \forall (a, b) \in A \times B.$$

Moreover if $\xi : A \longrightarrow B$ is a linear map. Then

$$\xi : (A, \prec_{\vdash_A}, \prec_{\dashv_A}, \succ_{\vdash_A}, \succ_{\dashv_A}, \alpha_A) \longrightarrow (B, \prec_{\vdash_B}, \prec_{\dashv_B}, \succ_{\vdash_B}, \succ_{\dashv_B}, \alpha_B)$$

is a morphism if and only if its graph $\Gamma_\xi = \{(x, \Gamma(x), x \in A)\}$ is a Hom-quadri-dendriform algebra of $(A \oplus B, \prec_+, \prec_-, \succ_+, \succ_-, \alpha)$.

Proof. The proof of the first part comes from a direct computation so we omit it. Now let us suppose that $\xi : (A, \prec_{+A}, \prec_{-A}, \succ_{+A}, \succ_{-A}, \alpha_A) \longrightarrow (B, \prec_{+B}, \prec_{-B}, \succ_{+B}, \succ_{-B}, \alpha_B)$ is a morphism of Hom-quadri-dendriform algebras. Then

$$\begin{aligned} (x + \xi(x)) \prec_+ (y + \xi(y)) &:= (x \prec_{+A} y + \xi(x) \prec_{+B} \xi(y)) = x \prec_{+A} y + \xi(x \prec_{+A} y) \\ (x + \xi(x)) \prec_- (y + \xi(y)) &:= (x \prec_{-A} y + \xi(x) \prec_{-B} \xi(y)) = x \prec_{-A} y + \xi(x \prec_{-A} y) \\ (x + \xi(x)) \succ_+ (y + \xi(y)) &:= (x \succ_{+A} y + \xi(x) \succ_{+B} \xi(y)) = x \succ_{+A} y + \xi(x \succ_{+A} y) \\ (x + \xi(x)) \succ_- (y + \xi(y)) &:= (x \succ_{-A} y + \xi(x) \succ_{-B} \xi(y)) = x \succ_{-A} y + \xi(x \succ_{-A} y). \end{aligned}$$

Thus the graph Γ_ξ is closed under the operations $\prec_+, \prec_-, \succ_+$ and $\succ_-$.

Furthermore since $\xi \circ \alpha_A = \alpha_B \circ \xi$ we have

$$(\alpha_A \oplus \alpha_B)(x, \xi(x)) = (\alpha_A(x), \alpha_B \circ \xi(x)) = (\alpha_A(x), \xi \circ \alpha_A(x)).$$

Which imples that Γ_ξ is closed $\alpha_A \oplus \alpha_B$. Thus Γ_ξ is a Hom-subalgebra of $(A \oplus B, \prec_+, \prec_-, \succ_+, \succ_-, \alpha)$. Conversely if the graph $\Gamma_\xi \subset A \oplus B$ is a Hom-subalgebra of $(A \oplus B, \prec_+, \prec_-, \succ_+, \succ_-, \alpha)$ then we have

$$\begin{aligned} (x + \xi(x)) \prec_+ (y + \xi(y)) &:= (x \prec_{+A} y + \xi(x) \prec_{+B} \xi(y)) \in \Gamma_\xi \\ (x + \xi(x)) \prec_- (y + \xi(y)) &:= (x \prec_{-A} y + \xi(x) \prec_{-B} \xi(y)) \in \Gamma_\xi \\ (x + \xi(x)) \succ_+ (y + \xi(y)) &:= (x \succ_{+A} y + \xi(x) \succ_{+B} \xi(y)) \in \Gamma_\xi \\ (x + \xi(x)) \succ_- (y + \xi(y)) &:= (x \succ_{-A} y + \xi(x) \succ_{-B} \xi(y)) \in \Gamma_\xi. \end{aligned}$$

Furthermore $(\alpha_A \oplus \alpha_B)(\Gamma_\xi) \subset \Gamma_\xi$ implies

$$(\alpha_A \oplus \alpha_B)(x, \xi(x)) = (\alpha_A(x), \alpha_B \circ \xi(x)) \in \Gamma_\xi$$

which is equivalent to the condition $\alpha_B \circ \xi(x) = \xi \circ \alpha_A(x)$. i.e $\alpha_B \circ \xi = \xi \circ \alpha_A$. Therefore ξ is a morphism of Hom-quadri-dendriform algebras. $\square$

Proposition 3.7. Let $(V, \prec_l, \succ_l, \prec_r, \succ_r, \beta)$ be a representation of a Hom-dendriform algebra $(D, \prec, \succ, \alpha)$. Then $(D \oplus V, \prec_+, \prec_-, \succ_+, \succ_-, \alpha \otimes \beta)$ is a Hom-quadri-dendriform algebra where

$$\begin{aligned} (x, u) \prec_+ (y, v) &:= (x \prec y, x \prec_l v) \\ (x, u) \prec_- (y, v) &:= (x \prec y, u \prec_r y) \\ (x, u) \succ_+ (y, v) &:= (x \succ y, x \succ_l v) \\ (x, u) \succ_- (y, v) &:= (x \succ y, u \succ_r y). \end{aligned}$$

$\forall x, y \in D$ and $u, v \in V$. This Hom-quadri-dendriform algebra is called the hemi-semidirect product Hom-quadri-dendriform algebra.

Proof. For any $(x, u), (y, v) (z, w) \in D \oplus V$ and $* \in \{+, -\}$ we have

$$\begin{aligned} ((x, u) \prec_* (y, v)) \prec_+ (\alpha(z), \beta(w)) &= ((x \prec y) \prec \alpha(z), (x \prec y) \prec_\ell \beta(w)) \\ &= (\alpha(x) \prec (y \prec z + y \succ z), \alpha(x) \prec_\ell (v \prec_\ell w + v \succ_\ell w)) \\ &= (\alpha(x), \beta(u)) \prec_+ ((y, v) \prec_+ (z, w) + (y, v) \succ_+ (z, w)), \end{aligned}$$

$$\begin{aligned} ((x, u) \succ_* (y, v)) \prec_+ (\alpha(z), \beta(w)) &= ((x \succ y) \prec \alpha(z), (x \succ y) \prec_\ell \beta(w)) \\ &= (\alpha(x) \succ (y \prec z), \alpha(x) \succ_\ell (y \prec_\ell w)) \end{aligned}$$

$$= (\alpha(x), \beta(u)) \succ_{\vdash} ((y, v) \prec_{\vdash} (z, w)),$$

$$\begin{aligned} (\alpha(x), \beta(u)) \succ_{\vdash} ((y, v) \succ_{\vdash} (z, w)) &= (\alpha(x) \succ (y \succ z), \alpha(x) \succ_{\ell} (y \succ_{\ell} w)) \\ &= ((x \prec y + x \succ y) \succ \alpha(z), (x \prec y + x \succ y) \succ_{\ell} \beta(w)) \\ &= ((x, u) \prec_{*} (y, v) + (x, u) \succ_{*} (y, v)) \succ_{\vdash} (\alpha(z), \beta(w)), \end{aligned}$$

$$\begin{aligned} ((x, u) \prec_{\vdash} (y, v)) \prec_{\vdash} (\alpha(z), \beta(w)) &= ((x \prec y) \prec \alpha(z), (x \prec_{\ell} v) \prec_r \alpha(z)) \\ &= (\alpha(x) \prec (y \prec z + y \succ z), \alpha(x) \prec_{\ell} (v \prec_r z + v \succ_r z)) \\ &= (\alpha(x), \beta(u)) \prec_{\vdash} ((y, v) \prec_{\vdash} (z, w) + (y, v) \succ_{\vdash} (z, w)). \end{aligned}$$

Thus the Eq (16) hold. The same for Eqs (12) à (22) . This completes the proof. $\square$

Definition 3.8. Let $(D, \prec, \succ, \alpha)$ be a Hom-dendriform algebra and $(V, \prec_{\ell}, \succ_{\ell}, \prec_r, \succ_r, \beta)$ be a representation of it. A relative averaging operator on D with respect to the representation $(V, \prec_{\ell}, \succ_{\ell}, \prec_r, \succ_r, \beta)$ is a linear map $T : V \rightarrow D$ that satisfies

$$\begin{aligned} Tu \prec Tv &= T(Tu \prec_{\ell} v) = T(u \prec_r Tv), \\ Tu \succ Tv &= T(Tu \succ_{\ell} v) = T(u \succ_r Tv), \\ T \circ \beta &= \alpha \circ T \end{aligned}$$

for all $u, v \in V$. The relative averaging operator on a Hom-dendriform algebra $(D, \prec, \succ, \alpha)$ with respect to the adjoint representation $(D, \prec_{\ell}=\prec, \succ_{\ell}=\succ, \prec_r=\prec, \succ_r=\succ, \alpha)$ is called an averaging operator. In this case the Eqs can be written as

$$\begin{aligned} Tx \prec Ty &= T(Tx \prec y) = T(x \prec Ty) \\ Tx \succ Ty &= T(Tx \succ y) = T(x \succ Ty) \\ T \circ \alpha &= \alpha \circ T. \end{aligned}$$

Definition 3.9. An averaging Hom-dendriform algebra is a pair (D, T) consisting of a Hom-dendriform algebra $(D, \prec, \succ, \alpha)$ endowed with an averaging operator T .

In the following we give a characterization of a relative averaging operator in terms of the graph of the operator.

Theorem 3.10. A linear map $T : V \rightarrow D$ is a relative averaging operator on a Hom-dendriform algebra $(D, \prec, \succ, \alpha)$ with respect to the representation $(V, \prec_{\ell}, \succ_{\ell}, \prec_r, \succ_r, \beta)$ if and only if the graph $Gr(T) = \{(Tu, u) \mid u \in V\}$ is a Hom-subalgebra of the hemi-semidirect product Hom-quadri-dendriform algebra $(D \oplus V, \prec_{\vdash}, \prec_{\vdash}, \succ_{\vdash}, \succ_{\vdash}, \alpha \otimes \beta)$.

Proof. Let $T : V \rightarrow D$ be a linear map. For any $u, v \in V$ we have

$$\begin{aligned} (Tu, u) \prec_{\vdash} (Tv, v) &= (Tu \prec Tv, Tu \prec_l v), \quad (Tu, u) \prec_{\vdash} (Tv, v) = (Tu \prec Tv, u \prec_r Tv), \\ (Tu, u) \succ_{\vdash} (Tv, v) &= (Tu \succ Tv, Tu \succ_l v), \quad (Tu, u) \succ_{\vdash} (Tv, v) = (Tu \succ Tv, u \succ_r Tv). \end{aligned}$$

The above five elements are in $Gr(T)$ if and only if

$$\begin{aligned} Tu \prec Tv &= T(Tu \prec_l v), \quad Tu \prec Tv = T(u \prec_r Tv), \quad Tu \succ Tv = T(Tu \succ_l v), \\ Tu \succ Tv &= T(u \succ_r Tv), \quad \alpha \otimes \beta((Tu, u)) = (\alpha(Tu), \beta(u)) = (T(\beta(u)), \beta(u)) \in Gr(T). \end{aligned}$$

Therefore the graph $Gr(T)$ is a Hom-subalgebra of the hemi semidirect product Hom-quadri-dendriform algebra $(D \oplus V, \prec_{\vdash}, \prec_{\vdash}, \succ_{\vdash}, \succ_{\vdash}, \alpha \otimes \beta)$ if and only if T is a relative averaging operator. $\square$

Proposition 3.11. *Let $T : V \rightarrow D$ be a relative averaging operator on a Hom-dendriform algebra $(D, \prec, \succ, \alpha)$ with respect to the representation $(V, \prec_\ell, \succ_\ell, \prec_r, \succ_r, \beta)$. Then V inherits a Hom-quadri-dendriform algebra structure with the operations:*

$$\begin{aligned} u \prec_{\vdash}^T v &= T(u) \prec_\ell v, & u \prec_{\dashv}^T v &= u \prec_r T(v) \\ u \succ_{\vdash}^T v &= T(u) \succ_\ell v, & u \succ_{\dashv}^T v &= u \succ_r T(v) \end{aligned}$$

for all $u, v \in V$. Moreover T is a homomorphism from the Hom-quadri-dendriform algebra $(V, \prec_{\vdash}^T, \prec_{\dashv}^T, \succ_{\vdash}^T, \succ_{\dashv}^T, \beta)$ to the Hom-dendriform algebra $(D, \prec, \succ, \alpha)$.

Corollary 3.12. *Let $(D, \prec, \succ, \alpha)$ be a Hom-dendriform algebra and let $T : D \rightarrow D$ be an averaging operator. Then D is also equipped with a Hom-quadri-dendriform algebra structure defined by $x \prec_{\vdash}^T y = T(x) \prec y$, $x \prec_{\dashv}^T y = x \prec T(y)$, $x \succ_{\vdash}^T y = T(x) \succ y$, $x \succ_{\dashv}^T y = x \succ T(y)$, $\forall x, y \in D$.*

Theorem 3.13. *Every Hom-quadri-dendriform algebra can induce a relative averaging operator on a Hom-dendriform algebra with respect to a representation.*

Proof. Let $(D, \prec_{\vdash}, \prec_{\dashv}, \succ_{\vdash}, \succ_{\dashv}, \alpha)$ be an Hom-quadri-dendriform algebra. Then by Example 3.5 $D_{\text{Dend}} = \frac{D}{\mathbb{I}_D}$ carries a Hom-dendriform algebra structure with the operations

$$\begin{aligned} \overline{\alpha(\bar{x})} &= \overline{\alpha(x)}, \\ \bar{x} \prec \bar{y} &= \overline{x \prec_{\vdash} y} = \overline{x \prec_{\dashv} y}, \\ \bar{x} \succ \bar{y} &= \overline{x \succ_{\vdash} y} = \overline{x \succ_{\dashv} y}. \end{aligned}$$

for all $x, y \in D$ here $\bar{x}$ denotes the class of an element $x \in D$. We define four bilinear maps $\prec_l, \succ_l : D_{\text{Dend}} \times D \rightarrow D$ and $\prec_r, \succ_r : D \times D_{\text{Dend}} \rightarrow D$ by

$$x \prec_l y = x \prec_{\vdash} y, \quad x \succ_l y = x \succ_{\vdash} y, \quad y \prec_r x = y \prec_{\dashv} x, \quad y \succ_r x = y \succ_{\dashv} x.$$

for all $\bar{x} \in D_{\text{Dend}}$ and $y \in D$. It is easy to verify that the maps $\prec_l, \succ_l, \prec_r, \succ_r$ define a representation of the Hom-dendriform algebra D_{Dend} on the vector space D . Moreover the quotient map $T : D \rightarrow D_{\text{Dend}}$ $x \mapsto \bar{x}$ is a relative averaging operator as

$$\begin{aligned} Tx \prec Ty &= \bar{x} \prec \bar{y} = \begin{cases} = \overline{x \prec_{\vdash} y} = \overline{\bar{x} \prec_l y} = T(Tx \prec_l y) \\ = \overline{x \prec_{\dashv} y} = \overline{\bar{x} \prec_r y} = T(x \prec_r Ty). \end{cases} \\ Tx \succ Ty &= \bar{x} \succ \bar{y} = \begin{cases} = \overline{x \succ_{\vdash} y} = \overline{\bar{x} \succ_l y} = T(Tx \succ_l y) \\ = \overline{x \succ_{\dashv} y} = \overline{\bar{x} \succ_r y} = T(x \succ_r Ty) \end{cases} \end{aligned}$$

for all $x, y \in D$. □

4. Hom-Six-dendriform algebras

In [17] Loday and Ronco introduced the notion of dendriform trialgebra. Inspired by Loday and Ronco's work we present the notion of Hom-six-dendriform algebras generalizing the classical six-dendriform algebras to Hom-algebras setting.

Definition 4.1. Let $(D, \prec, \succ, \alpha)$ and $(D', \prec', \succ', \alpha')$ be two Hom-dendriform algebras. We say that D acts on D' if there exist four bilinear maps $\prec_l, \succ_l : D \otimes D' \rightarrow D'$ and $\prec_r, \succ_r : D' \otimes D \rightarrow D'$ such that $(D', \prec_l, \succ_l, \prec_r, \succ_r, \alpha')$ is a representation of D and for any $x, y, z \in D$ and $u, v, w \in D'$ we have

$$(x \prec_l v) \prec' \alpha'(w) = \alpha(x) \prec_l (v \prec' w + v \succ' w) \tag{24}$$

$$(x \succ_l v) \prec' \alpha'(w) = \alpha(x) \succ_l (v \prec' w) \tag{25}$$

$$\alpha(x) \succ_l (v \succ' w) = (x \prec_l v + x \succ_l v) \succ' \alpha'(w) \quad (26)$$

$$(u \prec_r y) \prec' \alpha'(w) = \alpha'(u) \prec' (y \prec_l w + y \succ_l w) \quad (27)$$

$$(u \succ_r y) \prec' \alpha'(w) = \alpha'(u) \succ' (y \prec_l w) \quad (28)$$

$$\alpha'(u) \succ' (y \succ_l w) = (u \prec_r y + u \succ_r y) \succ' \alpha'(w) \quad (29)$$

$$(u \prec' v) \prec_r \alpha(z) = \alpha'(u) \prec' (v \prec_r z + v \succ_r z) \quad (30)$$

$$(u \succ' v) \prec_r \alpha(z) = \alpha'(u) \succ' (v \prec_r z) \quad (31)$$

$$\alpha'(u) \succ' (v \succ_r z) = (u \prec' v + u \succ' v) \succ_r \alpha(z). \quad (32)$$

Proposition 4.2. *Let $(D, \prec, \succ, \alpha)$ and $(D', \prec', \succ', \alpha')$ be two Hom-dendriform algebras and let $\prec_l, \succ_l: D \otimes D' \rightarrow D'$ and $\prec_r, \succ_r: D' \otimes D \rightarrow D'$ be bilinear maps. Then the tuple $(D', \prec_l, \succ_l, \prec_r, \succ_r, \alpha')$ is an action of D if and only if $(D \oplus D', \alpha \otimes \alpha')$ carries a Hom-dendriform algebra structure with operations given by*

$$(x, u) \prec_{\boxtimes} (y, v) = (x \prec y, x \prec_l v + u \prec_r y + u \prec' v)$$

$$(x, u) \succ_{\boxtimes} (y, v) = (x \succ y, x \succ_l v + u \succ_r y + u \succ' v).$$

for all $(x, u), (y, v) \in (D \oplus D')$.

$(D \oplus D', \alpha \otimes \alpha')$ is called the semi-direct product of D with D' .

Proof. Let $(D, \prec, \succ, \alpha)$ and $(D', \prec', \succ', \alpha')$ be two Hom-dendriform algebras.

Then for all $(x, u), (y, v), (z, w) \in (D \oplus D')$ we have

$$\begin{aligned} & ((x, u) \prec_{\boxtimes} (y, v)) \prec_{\boxtimes} (\alpha(z), \alpha'(w)) - (\alpha(x), \alpha'(u)) \prec_{\boxtimes} ((y, v) \prec_{\boxtimes} (z, w) + (y, v) \succ_{\boxtimes} (z, w)) \\ &= ((x \prec y) \prec \alpha(z) - \alpha(x) \prec (y \prec z + y \succ z), (x \prec y) \prec_l \alpha'(w) + (x \prec_l v + u \prec_r y + u \prec' v) \prec_r \alpha(z) \\ &+ (x \prec_l v + u \prec_r y + u \prec' v) \prec' \alpha'(w) - \alpha(x) \prec_l (y \prec_l w + y \succ_l w + v \prec_r z + v \succ_r z + v \prec' w + v \succ' w) \\ &- \alpha'(u) \prec_r (y \prec z + y \succ z) - \alpha'(u) \prec' (y \prec_l w + y \succ_l w + v \prec_r z + v \succ_r z + v \prec' w + v \succ' w)) \\ &= (0, (x \prec y) \prec_{\ell} \alpha'(w) + (x \prec_{\ell} v + u \prec_r y + u \prec' v) \prec_r \alpha(z) + (x \prec_{\ell} v + u \prec_r y) \prec' \alpha'(w) \\ &- \alpha(x) \prec_{\ell} (y \prec_{\ell} w + y \succ_{\ell} w + v \prec_r z + v \succ_r z + v \prec' w + v \succ' w) - \alpha'(u) \prec_r (y \prec z + y \succ z) - \\ &\alpha'(u) \prec' (y \prec_{\ell} w + y \succ_{\ell} w + v \prec_r z + v \succ_r z)) \\ & ((x, u) \succ_{\boxtimes} (y, v)) \succ_{\boxtimes} (\alpha(z), \alpha'(w)) - (\alpha(x), \alpha'(u)) \succ_{\boxtimes} ((y, v) \succ_{\boxtimes} (z, w) + (y, v) \prec_{\boxtimes} (z, w)) \\ &= ((x \succ y) \prec \alpha(z) - \alpha(x) \succ (y \prec z), (x \succ y) \prec_l \alpha'(w) + (x \succ_l v + u \succ_r y + u \succ' v) \prec_r \alpha(z) \\ &+ (x \succ_l v + u \succ_r y + u \succ' v) \prec' \alpha'(w) - \alpha(x) \succ_l (y \prec_l w + v \prec_r z + v \prec' w) \\ &- \alpha'(u) \succ_r (y \prec z) - \alpha'(u) \succ' (y \prec_l w + v \prec_r z + v \succ_r z + v \prec' w)) \\ &= (0, (x \succ y) \prec_{\ell} \alpha'(w) + (x \succ_l v + u \succ_r y + u \succ' v) \prec_r \alpha(z) \\ &+ (x \succ_l v + u \succ_r y + u \succ' v) \prec' \alpha'(w) - \alpha(x) \succ_l (y \prec_l w + v \prec_r z + v \prec' w) \\ &- \alpha'(u) \succ_r (y \prec z) - \alpha'(u) \succ' (y \prec_l w + v \prec_r z)) \\ & (\alpha(x), \alpha'(u)) \succ_{\boxtimes} ((y, v) \succ_{\boxtimes} (z, w)) - ((x, u) \prec_{\boxtimes} (y, v) + (x, u) \succ_{\boxtimes} (y, v)) \succ_{\boxtimes} (\alpha(z), \alpha'(w)) \\ &= (\alpha(x) \succ (y \succ z) - (x \prec y + x \succ y) \succ \alpha(z), \alpha(x) \succ_l (y \succ_l w + v \succ_r z + v \succ' w) + \alpha'(u) \succ_r (y \succ z) \\ &+ \alpha'(u) \succ' (y \succ_l w + v \succ_r z + v \succ' w) - (x \prec y) \succ_l \alpha'(w) - (x \prec_l v + u \prec_r y + u \prec' v) \succ_r \alpha(z) \\ &- (x \prec_l v + u \prec_r y + u \prec' v) \succ' \alpha'(w) - (x \succ y) \succ_l \alpha'(w) - (x \succ_l v + u \succ_r y + u \succ' v) \succ_r \alpha(z) \\ &- (x \succ_l v + u \succ_r y + u \succ' v) \succ' \alpha'(w)) \end{aligned}$$

$$\begin{aligned}
&= (0, \alpha(x) \succ_l (y \succ_l w + v \succ_r z + v \succ' w) + \alpha'(u) \succ_r (y \succ z) + \alpha'(u) \succ' (y \succ_l w + v \succ_r z) \\
&\quad - (x \prec y) \succ_l \alpha'(w) - (x \prec_l v + u \prec_r y + u \prec' v) \succ_r \alpha(z) - (x \prec_l v + u \prec_r y) \succ' \alpha'(w) \\
&\quad - (x \succ y) \succ_l \alpha'(w) - (x \succ_l v + u \succ_r y + u \succ' v) \succ_r \alpha(z) - (x \succ_l v + u \succ_r y) \succ' \alpha'(w)).
\end{aligned}$$

Then $(D \oplus D', \succ_{\boxtimes}, \prec_{\boxtimes}, \alpha \otimes \alpha')$ is a Hom-dendriform algebra if and only if $(D', \prec_l, \succ_l, \prec_r, \succ_r, \alpha')$ is a representation of D and Eqs.(24)-(32) are satisfied. $\square$

Definition 4.3. A linear map $T : D' \rightarrow D$ is called a homomorphic relative averaging operator on the Hom-dendriform algebra $(D, \prec, \succ, \alpha)$ with respect to an action $(D', \prec', \succ', \prec_l, \succ_l, \prec_r, \succ_r, \alpha')$ if T qualifies both as a relative averaging operator and a Hom-dendriform homomorphism.

A homomorphic relative averaging operator on a Hom-dendriform algebra is also called a relative averaging operator of weight 1 on a Hom-dendriform algebra. In particular a homomorphic relative averaging operator on a Hom-dendriform algebra $(D, \prec, \succ, \alpha)$ with respect to the adjoint representation is called a homomorphic averaging operator.

Definition 4.4. A Hom-six-dendriform algebra is a tuple $(D, \prec_{\perp}, \succ_{\perp}, \prec_{\vdash}, \succ_{\vdash}, \prec_{\dashv}, \succ_{\dashv}, \alpha)$ consisting of a Hom-dendriform algebra $(D, \prec_{\perp}, \succ_{\perp}, \alpha)$ and a Hom-quadri-dendriform algebra $(D, \prec_{\vdash}, \succ_{\vdash}, \prec_{\dashv}, \succ_{\dashv}, \alpha)$ such that for all $x, y, z \in D$

$$(x \prec_{\vdash} y) \prec_{\perp} \alpha(z) = \alpha(x) \prec_{\vdash} (y \prec_{\perp} z + y \succ_{\perp} z) \quad (33)$$

$$(x \succ_{\vdash} y) \prec_{\perp} \alpha(z) = \alpha(x) \succ_{\vdash} (y \prec_{\perp} z) \quad (34)$$

$$\alpha(x) \succ_{\vdash} (y \succ_{\perp} z) = (x \prec_{\vdash} y + x \succ_{\vdash} y) \succ_{\perp} \alpha(z) \quad (35)$$

$$(x \prec_{\dashv} y) \prec_{\perp} \alpha(z) = \alpha(x) \prec_{\perp} (y \prec_{\vdash} z + y \succ_{\vdash} z) \quad (36)$$

$$(x \succ_{\dashv} y) \prec_{\perp} \alpha(z) = \alpha(x) \succ_{\perp} (y \prec_{\vdash} z) \quad (37)$$

$$\alpha(x) \succ_{\perp} (y \succ_{\vdash} z) = (x \prec_{\dashv} y + x \succ_{\dashv} y) \succ_{\perp} \alpha(z) \quad (38)$$

$$(x \prec_{\perp} y) \prec_{\dashv} \alpha(z) = \alpha(x) \prec_{\dashv} (y \prec_{\vdash} z + y \succ_{\vdash} z) \quad (39)$$

$$(x \succ_{\perp} y) \prec_{\dashv} \alpha(z) = \alpha(x) \succ_{\dashv} (y \prec_{\vdash} z) \quad (40)$$

$$\alpha(x) \succ_{\dashv} (y \succ_{\vdash} z) = (x \prec_{\perp} y + x \succ_{\perp} y) \succ_{\dashv} \alpha(z) \quad (41)$$

$$(x \prec_{\perp} y) \prec_{\vdash} \alpha(z) = (x \prec_{\vdash} y) \prec_{\vdash} \alpha(z) = (x \prec_{\dashv} y) \prec_{\vdash} \alpha(z) \quad (42)$$

$$(x \succ_{\perp} y) \prec_{\vdash} \alpha(z) = (x \succ_{\vdash} y) \prec_{\vdash} \alpha(z) = (x \succ_{\dashv} y) \prec_{\vdash} \alpha(z) \quad (43)$$

$$(x \prec_{\perp} y) \succ_{\vdash} \alpha(z) = (x \prec_{\vdash} y) \succ_{\vdash} \alpha(z) = (x \prec_{\dashv} y) \succ_{\vdash} \alpha(z) \quad (44)$$

$$(x \succ_{\perp} y) \succ_{\vdash} \alpha(z) = (x \succ_{\vdash} y) \succ_{\vdash} \alpha(z) = (x \succ_{\dashv} y) \succ_{\vdash} \alpha(z) \quad (45)$$

$$\alpha(x) \prec_{\dashv} (y \prec_{\perp} z) = \alpha(x) \prec_{\dashv} (y \prec_{\vdash} z) = \alpha(x) \prec_{\dashv} (y \prec_{\dashv} z) \quad (46)$$

$$\alpha(x) \succ_{\dashv} (y \prec_{\perp} z) = \alpha(x) \succ_{\dashv} (y \prec_{\vdash} z) = \alpha(x) \succ_{\dashv} (y \prec_{\dashv} z) \quad (47)$$

$$\alpha(x) \prec_{\dashv} (y \succ_{\perp} z) = \alpha(x) \prec_{\dashv} (y \succ_{\vdash} z) = \alpha(x) \prec_{\dashv} (y \succ_{\dashv} z) \quad (48)$$

$$\alpha(x) \succ_{\dashv} (y \succ_{\perp} z) = \alpha(x) \succ_{\dashv} (y \succ_{\vdash} z) = \alpha(x) \succ_{\dashv} (y \succ_{\dashv} z). \quad (49)$$

$(D, \prec_{\perp}, \succ_{\perp}, \prec_{\vdash}, \succ_{\vdash}, \prec_{\dashv}, \succ_{\dashv}, \alpha)$ is said to be a multiplicative Hom-algebra if:

$$\alpha(x \prec_{\perp} y) = \alpha(x) \prec_{\perp} \alpha(y), \quad \alpha(x \succ_{\perp} y) = \alpha(x) \succ_{\perp} \alpha(y),$$

$$\alpha(x \prec_{\vdash} y) = \alpha(x) \prec_{\vdash} \alpha(y), \quad \alpha(x \succ_{\vdash} y) = \alpha(x) \succ_{\vdash} \alpha(y),$$

$$\alpha(x \prec_{\dashv} y) = \alpha(x) \prec_{\dashv} \alpha(y), \quad \alpha(x \succ_{\dashv} y) = \alpha(x) \succ_{\dashv} \alpha(y).$$

If $\prec_{\perp} = \prec_{\vdash} = \prec_{\dashv}$ and $\succ_{\perp} = \succ_{\vdash} = \succ_{\dashv}$ then we simply obtain a Hom-dendriform algebra.

If $\prec_{\perp} = \prec_{\vdash} = \prec_{\dashv}$ and $\succ_{\perp} = \succ_{\vdash} = \succ_{\dashv}$ then we simply obtain a Hom-dendriform algebra.

Definition 4.5. Given two Hom-six-dendriform algebras $(D, \prec_\perp, \succ_\perp, \prec_\vdash, \succ_\vdash, \prec_\dashv, \succ_\dashv, \alpha)$ and $(D', \prec'_\perp, \succ'_\perp, \prec'_\vdash, \succ'_\vdash, \prec'_\dashv, \succ'_\dashv, \alpha')$ a Hom-six-dendriform homomorphism is a linear map $T : D \rightarrow D'$ that satisfies the following compatibility conditions:

$$\begin{aligned} T(x \prec_\perp y) &= T(x) \prec'_\perp T(y), & T(x \succ_\perp y) &= T(x) \succ'_\perp T(y), \\ T(x \prec_\vdash y) &= T(x) \prec'_\vdash T(y), & T(x \succ_\vdash y) &= T(x) \succ'_\vdash T(y), \\ T(x \prec_\dashv y) &= T(x) \prec'_\dashv T(y), & T(x \succ_\dashv y) &= T(x) \succ'_\dashv T(y). \end{aligned}$$

In particular if T is bijective then $(D, \prec_\perp, \succ_\perp, \prec_\vdash, \succ_\vdash, \prec_\dashv, \succ_\dashv, \alpha)$ and $(D', \prec'_\perp, \succ'_\perp, \prec'_\vdash, \succ'_\vdash, \prec'_\dashv, \succ'_\dashv, \alpha')$ are said to be isomorphic.

Definition 4.6. A Hom-triassociative algebra is a vector space D equipped with three binary operations $\vdash, \dashv, \perp : D \otimes D \rightarrow D$ a linear map $\alpha : D \rightarrow D$ such that $(D, \vdash, \dashv, \alpha)$ constitutes a Hom-diassociative algebra $(D, \perp, \alpha)$ forms an Hom-associative algebra and for all $x, y, z \in D$ the following equations hold:

$$(x \dashv y) \dashv \alpha(z) = \alpha(x) \dashv (y \perp z) \quad (50)$$

$$(x \vdash y) \perp \alpha(z) = \alpha(x) \vdash (y \perp z) \quad (51)$$

$$(x \perp y) \dashv \alpha(z) = \alpha(x) \perp (y \dashv z) \quad (52)$$

$$(x \perp y) \vdash \alpha(z) = \alpha(x) \vdash (y \vdash z) \quad (53)$$

$$(x \dashv y) \perp \alpha(z) = \alpha(x) \perp (y \vdash z). \quad (54)$$

$(D, \vdash, \dashv, \perp, \alpha)$ is said to be a multiplicative Hom-algebra if $\forall x, y \in D$:

$$\alpha(x \vdash y) = \alpha(x) \vdash \alpha(y), \quad \alpha(x \dashv y) = \alpha(x) \dashv \alpha(y) \text{ and } \alpha(x \perp y) = \alpha(x) \perp \alpha(y).$$

Proposition 4.7. Let $(D, \prec_\perp, \succ_\perp, \prec_\vdash, \succ_\vdash, \prec_\dashv, \succ_\dashv, \alpha)$ be a Hom-six-dendriform algebra. Then $(D, \perp, \vdash, \dashv, \alpha)$ is a Hom-triassociative algebra where the operations are defined by:

$$\begin{aligned} x \perp y &= x \prec_\perp y + x \succ_\perp y, \\ x \vdash y &= x \prec_\vdash y + x \succ_\vdash y, \\ x \dashv y &= x \prec_\dashv y + x \succ_\dashv y, \end{aligned}$$

for all $x, y \in D$.

Proof. It follows from Proposition 3.3 that $(D, \vdash, \dashv, \alpha)$ is a Hom-di-associative algebra. The fact that $(D, \prec_\perp, \succ_\perp, \alpha)$ is a Hom-dendriform algebra means that $(D, \perp, \alpha)$ is a Hom-associative algebra. Next for any $x, y, z \in D$ by Eqs.(12)-(16) and (17)-(22) we have

$$\begin{aligned} (x \dashv y) \dashv \alpha(z) &= (x \prec_\dashv y + x \succ_\dashv y) \prec_\dashv \alpha(z) + (x \prec_\dashv y + x \succ_\dashv y) \succ_\dashv \alpha(z) \\ &= \alpha(x) \prec_\dashv (y \prec_\vdash z + y \succ_\vdash z) + \alpha(x) \succ_\dashv (y \prec_\vdash z) + \alpha(x) \succ_\dashv (y \succ_\vdash z) \\ &= \alpha(x) \prec_\dashv (y \prec_\perp z + y \succ_\perp z) + \alpha(x) \succ_\dashv (y \prec_\perp z + y \succ_\perp z) \\ &= \alpha(x) \dashv (y \perp z) \end{aligned}$$

$$\begin{aligned} (x \perp y) \dashv \alpha(z) &= (x \prec_\perp y + x \succ_\perp y) \prec_\dashv \alpha(z) + (x \prec_\perp y + x \succ_\perp y) \succ_\dashv \alpha(z) \\ &= \alpha(x) \prec_\perp (y \prec_\dashv z + y \succ_\dashv z) + \alpha(x) \succ_\perp (y \prec_\dashv z + y \succ_\dashv z) \\ &= \alpha(x) \perp (y \dashv z) \end{aligned}$$

$$(x \dashv y) \perp \alpha(z) = (x \prec_\dashv y + x \succ_\dashv y) \prec_\perp \alpha(z) + (x \prec_\dashv y + x \succ_\dashv y) \succ_\perp \alpha(z)$$

$$\begin{aligned}
&= \alpha(x) \prec_{\perp} (y \prec_{\vdash} z + y \succ_{\vdash} z) + \alpha(x) \succ_{\perp} (y \prec_{\vdash} z + y \succ_{\vdash} z) \\
&= \alpha(x) \perp (y \vdash z)
\end{aligned}$$

$$\begin{aligned}
(x \vdash y) \vdash \alpha(z) &= (x \prec_{\vdash} y + x \succ_{\vdash} y) \prec_{\vdash} \alpha(z) + (x \prec_{\vdash} y + x \succ_{\vdash} y) \succ_{\vdash} \alpha(z) \\
&= \alpha(x) \prec_{\vdash} (y \prec_{\vdash} z + y \succ_{\vdash} z) + \alpha(x) \succ_{\vdash} (y \prec_{\vdash} z + y \succ_{\vdash} z) \\
&= \alpha(x) \vdash (y \vdash z)
\end{aligned}$$

$$\begin{aligned}
(x \perp y) \vdash \alpha(z) &= (x \prec_{\perp} y + x \succ_{\perp} y) \prec_{\vdash} \alpha(z) + (x \prec_{\perp} y + x \succ_{\perp} y) \succ_{\vdash} \alpha(z) \\
&= \alpha(x) \prec_{\vdash} (y \prec_{\vdash} z + y \succ_{\vdash} z) + \alpha(x) \succ_{\vdash} (y \prec_{\vdash} z + y \succ_{\vdash} z) \\
&= \alpha(x) \vdash (y \vdash z)
\end{aligned}$$

This completes the proof. $\square$

Theorem 4.8. *Let $T : D' \rightarrow D$ be a homomorphic relative averaging operator on the Hom-dendriform algebra $(D, \prec, \succ, \alpha)$ with respect to an action $(D', \prec', \succ', \prec_{\ell}, \succ_{\ell}, \prec_r, \succ_r, \alpha')$. Then $(D', \prec_{\perp}, \succ_{\perp}, \prec_{\vdash}^T, \succ_{\vdash}^T, \prec_{\vdash}^T, \succ_{\vdash}^T, \alpha')$ is a Hom-six-dendriform algebra, where*

$$\begin{aligned}
u \prec_{\perp} v &= u \prec' v, \quad u \succ_{\perp} v = u \succ' v, \\
u \prec_{\vdash}^T v &= Tu \prec_l v, \quad u \prec_{\vdash}^T v = u \prec_r Tv, \\
u \succ_{\vdash}^T v &= Tu \succ_l v, \quad u \succ_{\vdash}^T v = u \succ_r Tv
\end{aligned}$$

Proof. We have already $(D', \prec_{\perp}, \succ_{\perp}, \alpha')$ is a Hom-dendriform algebra. Moreover it following from Proposition 3.11 that $(D', \prec_{\vdash}^T, \succ_{\vdash}^T, \prec_{\vdash}^T, \succ_{\vdash}^T, \alpha')$ is a Hom-quadri-dendriform algebra. Now for any $u, v, w \in D'$ according to Eqs.(24)-(32) we have

$$\begin{aligned}
(u \prec_{\vdash}^T v) \prec_{\perp} \alpha'(w) &= (Tu \prec_{\ell} v) \prec' \alpha'(w) = \alpha(Tu) \prec_{\ell} (v \prec_{\perp} w + v \succ_{\perp} w) \\
&= T\alpha'(u) \prec_{\ell} (v \prec_{\perp} w + v \succ_{\perp} w) = \alpha'(u) \prec_{\vdash}^T (v \prec_{\perp} w + v \succ_{\perp} w) \\
(u \succ_{\vdash}^T v) \prec_{\perp} \alpha'(w) &= (Tu \succ_{\ell} v) \prec' \alpha'(w) = \alpha(Tu) \succ_{\ell} (v \prec' w) \\
&= T\alpha'(u) \succ_{\ell} (v \prec' w) = \alpha'(u) \succ_{\vdash}^T (v \prec' w) = \alpha'(u) \succ_{\vdash}^T (v \succ_{\perp} w) \\
\alpha'(u) \succ_{\vdash}^T (v \succ_{\perp} w) &= T\alpha'(u) \succ_{\ell} (v \prec' w) = (Tu \prec_{\ell} v + Tu \succ_{\ell} v) \succ' \alpha'(w) \\
&= (u \succ_{\vdash}^T v + u \prec_{\vdash}^T v) \succ_{\perp} \alpha'(w) \\
(u \prec_{\vdash}^T v) \prec_{\perp} \alpha'(w) &= (u \prec_r Tv) \prec' \alpha'(w) = \alpha'(u) \prec' (Tv \prec_{\ell} w + Tv \succ_{\ell} w) \\
&= \alpha'(u) \prec_{\perp} (v \prec_{\vdash}^T w + v \succ_{\vdash}^T w) \\
(u \succ_{\vdash}^T v) \prec_{\perp} \alpha'(w) &= (u \succ_r Tv) \prec' \alpha'(w) = \alpha'(u) \succ' (Tv \prec_{\ell} w) = \alpha'(u) \succ_{\perp} (v \prec_{\vdash}^T w) \\
\alpha'(u) \succ_{\perp} (v \succ_{\vdash}^T w) &= \alpha'(u) \succ' (Tv \succ_{\ell} w) = (u \prec_r Tv + u \succ_r Tv) \succ' \alpha'(w) \\
&= (u \prec_{\vdash}^T v + u \succ_{\vdash}^T v) \succ_{\perp} \alpha'(w) \\
(u \prec_{\perp} v) \prec_{\vdash}^T \alpha'(w) &= (u \prec' v) \prec_r T\alpha'(w) = (u \prec' v) \prec_r \alpha(Tw) \\
&= \alpha'(u) \prec' (v \prec_r Tw + v \succ_r Tw) = \alpha'(u) \prec_{\perp} (v \prec_{\vdash}^T w + v \succ_{\vdash}^T w) \\
(u \succ_{\perp} v) \prec_{\vdash}^T \alpha'(w) &= (u \succ' v) \prec_r T\alpha'(w) = (u \succ' v) \prec_r \alpha(Tw) \\
&= \alpha'(u) \succ' (v \prec_r Tw) = \alpha'(u) \succ_{\perp} (v \prec_{\vdash}^T w)
\end{aligned}$$

Similarly it can be proved that other equations 40 à 49 are also true. This completes the proof. $\square$

Corollary 4.9. *Let $(D, \prec, \succ, \alpha)$ be a Hom-dendriform algebra and $T : D \rightarrow D$ be a homomorphic averaging operator. Then $(D', \prec_\perp, \succ_\perp, \prec_\perp^T, \succ_\perp^T, \prec_\perp^T, \succ_\perp^T, \alpha)$ is a Hom-six-dendriform algebra where $\prec_\perp = \prec, \succ_\perp = \succ, x \prec_\perp^T y = Tx \prec y, x \succ_\perp^T y = x \prec Ty, x \succ_\perp^T y = Tx \succ y, x \prec_\perp^T y = x \succ Ty$ for all $x, y \in D$.*

Theorem 4.10. *Every Hom-six-dendriform algebra can induce a homomorphic relative averaging operator on a Hom-dendriform algebra with respect to a action.*

Proof. Let $(D, \prec_\perp, \succ_\perp, \prec_\perp, \succ_\perp, \prec_\perp, \succ_\perp, \alpha)$ be a Hom-six-dendriform algebra. The vector subspace I_D generated by

$$\{x \prec_\perp y - x \prec_\perp y, x \succ_\perp y - x \succ_\perp y \mid x, y \in D\}$$

is an ideal of D . The quotient algebra $(D/I_D, \prec, \succ, \bar{\alpha})$ is a Hom-dendriform algebra denoted by D_{Dend} where

$$\bar{\alpha}(\bar{x}) = \overline{\alpha(x)}, \quad \bar{x} \prec \bar{y} = \overline{x \prec_\perp y} = \overline{x \prec_\perp y}, \quad \bar{x} \succ \bar{y} = \overline{x \succ_\perp y} = \overline{x \succ_\perp y}$$

for all $\bar{x}, \bar{y} \in D/I_D$. On the other hand $(D, \prec_\perp, \succ_\perp, \alpha)$ is a Hom-dendriform algebra denoted by $D_\perp$. $D_\perp$ is a D_{Dend} -module defined by the actions:

$$\begin{aligned} \bar{x} \prec_l \bar{y} &= x \prec_\perp y, & \bar{x} \succ_l y &= x \succ_\perp y, \\ y \prec_r \bar{x} &= y \prec_\perp x, & y \succ_r \bar{x} &= y \succ_\perp x. \end{aligned}$$

for all $x, y \in D_\perp$. Moreover the quotient map $T : D_\perp \rightarrow D_{Dend} \quad x \mapsto \bar{x}$ is a homomorphic relative averaging operator. $\square$

5. Classification

In this section we give the classification of Hom-quadri-dendriform algebra in low dimension.

For a selection of cases we classify the Hom-quadri-dendriform algebras. The computations for our classifications were done using Mathematica. Throughout we work over the complex field.

Theorem 5.1. *The isomorphism class of 2-dimensional Hom-quadri-dendriform algebras is given by the following representatives.*

- $D_1 : \begin{cases} e_1 \prec_\perp e_1 = e_2 & e_1 \succ_\perp e_1 = e_2 \\ e_1 \prec_\perp e_1 = e_2 & e_1 \succ_\perp e_1 = \frac{1}{2}e_2 \end{cases} \quad \alpha = \begin{pmatrix} a & 1 \\ 0 & a \end{pmatrix}; a \in \mathbb{R}.$
- $D_2 : \begin{cases} e_1 \prec_\perp e_1 = e_2 & e_1 \succ_\perp e_1 = -e_2 \\ e_1 \prec_\perp e_1 = e_2 & e_1 \succ_\perp e_1 = e_2 \end{cases} \quad \alpha = \begin{pmatrix} a & 0 \\ 0 & 0 \end{pmatrix}; a \in \mathbb{R}.$
- $D_3 : \begin{cases} e_1 \prec_\perp e_1 = e_1 & e_1 \succ_\perp e_1 = e_1 \\ e_2 \prec_\perp e_1 = e_1 & e_1 \succ_\perp e_1 = e_1 \\ e_1 \prec_\perp e_1 = e_1 & e_1 \succ_\perp e_1 = e_1 \\ e_2 \prec_\perp e_2 = e_1 & e_2 \succ_\perp e_2 = e_1 \end{cases} \quad \alpha = \begin{pmatrix} 0 & 0 \\ 0 & b \end{pmatrix}; b \in \mathbb{R}.$
- $D_4(\gamma) : \begin{cases} e_1 \prec_\perp e_2 = e_1 & e_1 \succ_\perp e_2 = e_1 \\ e_2 \prec_\perp e_2 = e_1 & e_2 \succ_\perp e_2 = e_1 \\ e_2 \prec_\perp e_2 = \gamma e_2 \end{cases} \quad \alpha = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}; \gamma \in \mathbb{R}.$

$$\bullet D_5(\eta) : \begin{cases} e_1 \lhd \vdash e_1 = e_1 & e_1 \rhd \vdash e_1 = e_1 \\ e_2 \lhd \vdash e_1 = e_1 & e_2 \rhd \vdash e_2 = e_1 \\ e_1 \lhd \dashv e_1 = e_1 & e_1 \rhd \dashv e_1 = e_1 \\ e_2 \lhd \dashv e_2 = e_1 & e_2 \rhd \dashv e_2 = \eta e_1 \end{cases} \quad \alpha = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}; \eta \in \mathbb{R}.$$

Theorem 5.2. *The isomorphism class of 3-dimensional Hom-quadri-dendriform algebras is given by the following representatives.*

$$\bullet D_1 : \begin{cases} e_1 \lhd \vdash e_1 = e_2 & e_1 \rhd \vdash e_1 = e_2 \\ e_1 \lhd \vdash e_3 = e_2 & e_1 \rhd \vdash e_3 = e_2 \\ e_3 \lhd \vdash e_3 = e_2 & e_3 \rhd \vdash e_3 = e_2 \\ e_1 \lhd \dashv e_1 = e_2 & e_1 \rhd \dashv e_3 = e_2 \\ e_1 \lhd \dashv e_3 = e_2 & e_3 \rhd \dashv e_1 = e_2 \\ e_3 \lhd \dashv e_3 = e_2 & e_3 \rhd \dashv e_3 = e_2 \end{cases} \quad \alpha = \begin{pmatrix} a & 1 & 0 \\ 0 & a & 0 \\ 0 & 0 & a \end{pmatrix}; a \in \mathbb{R}.$$

$$\bullet D_2 : \begin{cases} e_1 \lhd \vdash e_1 = e_3 & e_1 \rhd \vdash e_1 = e_3 \\ e_1 \lhd \vdash e_2 = e_3 & e_1 \rhd \vdash e_2 = e_3 \\ e_2 \lhd \vdash e_1 = e_3 & e_2 \rhd \vdash e_1 = e_3 \\ e_2 \lhd \vdash e_2 = e_3 & e_2 \rhd \vdash e_2 = e_3 \\ e_1 \lhd \dashv e_1 = e_3 & e_1 \rhd \dashv e_1 = e_3 \\ e_1 \lhd \dashv e_2 = e_3 & e_1 \rhd \dashv e_2 = e_3 \\ e_2 \lhd \dashv e_1 = e_3 & e_2 \rhd \dashv e_1 = e_3 \\ e_2 \lhd \dashv e_2 = e_3 & e_2 \rhd \dashv e_2 = e_3 \end{cases} \quad \alpha = \begin{pmatrix} a & 1 & 0 \\ 0 & a & 0 \\ 0 & 0 & a \end{pmatrix}; a \in \mathbb{R}.$$

$$\bullet D_3 : \begin{cases} e_1 \lhd \vdash e_2 = e_1 + e_3 & e_1 \rhd \vdash e_2 = e_1 + e_3 \\ e_2 \lhd \vdash e_1 = e_1 + e_3 & e_2 \rhd \vdash e_1 = e_1 + e_3 \\ e_2 \lhd \vdash e_3 = e_1 + e_3 & e_2 \rhd \vdash e_2 = \sigma e_3 \\ e_3 \lhd \vdash e_3 = e_1 + e_3 & e_2 \rhd \vdash e_3 = e_1 + e_3 \\ e_1 \lhd \dashv e_2 = e_3 & e_3 \rhd \vdash e_2 = e_1 + e_3 \\ e_2 \lhd \dashv e_1 = e_3 & e_3 \rhd \vdash e_3 = e_3 \\ e_2 \lhd \dashv e_2 = e_3 & e_2 \rhd \dashv e_1 = e_3 \\ e_3 \lhd \dashv e_2 = e_3 & e_2 \rhd \dashv e_2 = e_3 \\ e_3 \lhd \dashv e_3 = e_3 & e_2 \rhd \dashv e_3 = e_3 \\ & e_3 \rhd \dashv e_2 = -\eta e_3 \end{cases} \quad \alpha = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}; \sigma, \eta \in \mathbb{R}.$$

$$\bullet D_4 : \begin{cases} e_1 \lhd \vdash e_3 = e_1 + e_2 & e_1 \rhd \vdash e_3 = e_1 + e_2 \\ e_2 \lhd \vdash e_2 = e_1 + e_2 & e_2 \rhd \vdash e_2 = e_1 + e_2 \\ e_2 \lhd \vdash e_3 = e_1 + e_2 & e_2 \rhd \vdash e_3 = e_1 + e_2 \\ e_3 \lhd \vdash e_1 = e_1 + e_2 & e_3 \rhd \vdash e_1 = e_1 + e_2 \\ e_3 \lhd \vdash e_3 = e_1 + e_2 & e_3 \rhd \vdash e_2 = e_1 + e_2 \\ e_1 \lhd \dashv e_3 = e_2 & e_3 \rhd \vdash e_3 = e_1 + e_2 \\ e_2 \lhd \dashv e_2 = e_2 & e_2 \rhd \dashv e_2 = e_2 \\ e_2 \lhd \dashv e_3 = e_2 & e_2 \rhd \dashv e_3 = e_2 \\ e_3 \lhd \dashv e_1 = e_2 & e_3 \rhd \dashv e_1 = e_2 \\ e_3 \lhd \dashv e_2 = e_2 & e_3 \rhd \dashv e_2 = e_2 \\ e_3 \lhd \dashv e_3 = e_2 & e_3 \rhd \dashv e_3 = e_2 \end{cases} \quad \alpha = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}; \sigma, \eta \in \mathbb{R}.$$

$$\bullet D_5 : \left\{ \begin{array}{ll} e_1 \neg \vdash e_3 = e_2 & e_1 \neg \vdash e_3 = e_2 \\ e_2 \neg \vdash e_2 = e_2 & e_2 \neg \vdash e_2 = e_2 \\ e_3 \neg \vdash e_1 = e_2 & e_3 \neg \vdash e_1 = e_2 \\ e_3 \neg \vdash e_3 = e_2 & e_3 \neg \vdash e_3 = e_2 \\ e_1 \neg \vdash e_3 = e_2 & e_1 \neg \vdash e_3 = e_2 \\ e_2 \neg \vdash e_2 = e_2 & e_2 \neg \vdash e_2 = e_2 \\ e_3 \neg \vdash e_1 = e_2 & e_3 \neg \vdash e_1 = e_2 \\ e_3 \neg \vdash e_3 = e_2 & e_3 \neg \vdash e_3 = e_2 \end{array} \right. \quad \alpha = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & b \end{pmatrix}; b \in \mathbb{R}.$$

$$\bullet D_6 : \left\{ \begin{array}{ll} e_1 \neg \vdash e_2 = e_3 & e_1 \neg \vdash e_2 = e_3 \\ e_2 \neg \vdash e_1 = e_3 & e_2 \neg \vdash e_1 = e_3 \\ e_2 \neg \vdash e_2 = e_3 & e_2 \neg \vdash e_2 = e_3 \\ e_3 \neg \vdash e_3 = e_3 & e_3 \neg \vdash e_3 = e_3 \\ e_1 \neg \vdash e_2 = e_3 & e_1 \neg \vdash e_2 = e_3 \\ e_2 \neg \vdash e_1 = e_3 & e_2 \neg \vdash e_1 = e_3 \\ e_2 \neg \vdash e_2 = e_3 & e_2 \neg \vdash e_2 = e_3 \\ e_3 \neg \vdash e_3 = e_3 & e_3 \neg \vdash e_3 = e_3 \end{array} \right. \quad \alpha = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}.$$

$$\bullet D_7 : \left\{ \begin{array}{ll} e_1 \neg \vdash e_1 = e_3 & e_1 \neg \vdash e_1 = e_3 \\ e_1 \neg \vdash e_2 = e_3 & e_2 \neg \vdash e_1 = e_3 \\ e_2 \neg \vdash e_1 = e_3 & e_2 \neg \vdash e_2 = e_3 \\ e_2 \neg \vdash e_2 = e_3 & e_3 \neg \vdash e_3 = e_3 \\ e_1 \neg \vdash e_1 = e_3 & e_1 \neg \vdash e_1 = e_3 \\ e_1 \neg \vdash e_2 = e_3 & e_1 \neg \vdash e_2 = e_3 \\ e_2 \neg \vdash e_1 = e_3 & e_2 \neg \vdash e_1 = e_3 \\ e_2 \neg \vdash e_2 = e_3 & e_2 \neg \vdash e_2 = e_3 \end{array} \right. \quad \alpha = \begin{pmatrix} a & 1 & 0 \\ 0 & a & 1 \\ 0 & 0 & a \end{pmatrix}; a \in \mathbb{R}.$$

$$\bullet D_8 : \left\{ \begin{array}{ll} e_1 \neg \vdash e_1 = e_2 & e_1 \neg \vdash e_1 = e_2 \\ e_1 \neg \vdash e_3 = e_2 & e_1 \neg \vdash e_3 = e_2 \\ e_3 \neg \vdash e_1 = e_2 & e_3 \neg \vdash e_1 = e_2 \\ e_3 \neg \vdash e_3 = e_2 & e_3 \neg \vdash e_3 = e_2 \\ e_1 \neg \vdash e_1 = e_2 & e_1 \neg \vdash e_1 = e_2 \\ e_1 \neg \vdash e_3 = e_2 & e_1 \neg \vdash e_3 = e_2 \\ e_3 \neg \vdash e_1 = e_2 & e_3 \neg \vdash e_1 = e_2 \\ e_3 \neg \vdash e_3 = e_2 & e_3 \neg \vdash e_3 = e_2 \end{array} \right. \quad \alpha = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}.$$

$$\bullet D_9 : \left\{ \begin{array}{ll} e_1 \neg \vdash e_1 = e_3 & e_1 \neg \vdash e_1 = e_3 \\ e_1 \neg \vdash e_2 = e_3 & e_1 \neg \vdash e_2 = e_3 \\ e_2 \neg \vdash e_1 = e_3 & e_2 \neg \vdash e_2 = e_3 \\ e_3 \neg \vdash e_3 = e_3 & e_3 \neg \vdash e_3 = e_3 \\ e_1 \neg \vdash e_1 = e_3 & e_1 \neg \vdash e_2 = e_3 \\ e_2 \neg \vdash e_1 = e_3 & e_2 \neg \vdash e_1 = e_3 \\ e_2 \neg \vdash e_2 = e_3 & e_2 \neg \vdash e_2 = e_3 \\ e_3 \neg \vdash e_3 = e_3 & e_3 \neg \vdash e_3 = e_3 \end{array} \right. \quad \alpha = \begin{pmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

$$\begin{aligned}
\bullet D_{10} : & \begin{cases} e_1 \curlywedge_{\top} e_3 = e_1 + e_2 & e_1 \curlywedge_{\top} e_3 = e_1 + e_2 \\ e_3 \curlywedge_{\top} e_1 = e_1 + \lambda e_2 & e_3 \curlywedge_{\top} e_1 = e_1 + e_2 \\ e_3 \curlywedge_{\top} e_3 = e_1 + \frac{1}{2}e_2 & e_3 \curlywedge_{\top} e_3 = e_1 + e_2 \\ e_1 \curlywedge_{\perp} e_3 = e_1 + e_2 & e_1 \curlywedge_{\perp} e_3 = e_1 + e_2 \\ e_3 \curlywedge_{\perp} e_1 = e_1 + e_2 & e_3 \curlywedge_{\perp} e_1 = e_1 + e_2 \\ e_3 \curlywedge_{\perp} e_3 = e_1 + e_2 & e_3 \curlywedge_{\perp} e_3 = e_1 + e_2 \end{cases} & \alpha = \begin{pmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & 0 \end{pmatrix}; a, b, \lambda \in \mathbb{R}. \\
\bullet D_{11} : & \begin{cases} e_1 \curlywedge_{\top} e_2 = e_1 + e_3 & e_2 \curlywedge_{\top} e_1 = e_1 + e_3 \\ e_2 \curlywedge_{\top} e_1 = e_1 + e_3 & e_2 \curlywedge_{\top} e_2 = e_1 + e_3 \\ e_2 \curlywedge_{\top} e_2 = e_1 + e_3 & e_3 \curlywedge_{\top} e_2 = e_1 + e_3 \\ e_3 \curlywedge_{\top} e_2 = e_1 + e_3 & e_2 \curlywedge_{\perp} e_1 = e_1 + e_3 \\ e_1 \curlywedge_{\top} e_2 = e_1 + e_3 & e_2 \curlywedge_{\perp} e_2 = \eta e_1 + e_3 \end{cases} & \alpha = \begin{pmatrix} a & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & c \end{pmatrix}; \eta, a, c \in \mathbb{R}. \\
\bullet D_{12} : & \begin{cases} e_1 \curlywedge_{\top} e_2 = e_1 & e_2 \curlywedge_{\top} e_1 = e_1 \\ e_2 \curlywedge_{\top} e_1 = e_1 & e_2 \curlywedge_{\top} e_2 = e_1 \\ e_2 \curlywedge_{\top} e_2 = e_1 & e_2 \curlywedge_{\top} e_3 = e_1 \\ e_2 \curlywedge_{\top} e_3 = e_1 & e_3 \curlywedge_{\top} e_3 = e_1 \\ e_3 \curlywedge_{\top} e_3 = e_1 & e_2 \curlywedge_{\perp} e_2 = e_1 \\ e_1 \curlywedge_{\top} e_2 = e_1 & e_2 \curlywedge_{\perp} e_3 = \frac{1}{2}\eta e_1 \end{cases} & \alpha = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}; \eta \in \mathbb{R}. \\
\bullet D_{13} : & \begin{cases} e_1 \curlywedge_{\top} e_1 = e_3 & e_1 \curlywedge_{\top} e_1 = e_3 \\ e_1 \curlywedge_{\top} e_2 = e_3 & e_2 \curlywedge_{\top} e_1 = e_3 \\ e_2 \curlywedge_{\top} e_1 = e_3 & e_2 \curlywedge_{\top} e_1 = e_3 \\ e_2 \curlywedge_{\top} e_2 = e_3 & e_2 \curlywedge_{\top} e_2 = e_3 \\ e_1 \curlywedge_{\top} e_1 = e_3 & e_1 \curlywedge_{\perp} e_1 = e_3 \\ e_1 \curlywedge_{\top} e_2 = e_3 & e_1 \curlywedge_{\perp} e_2 = e_3 \\ e_2 \curlywedge_{\top} e_1 = e_3 & e_2 \curlywedge_{\perp} e_1 = e_3 \\ e_2 \curlywedge_{\top} e_2 = e_3 & e_2 \curlywedge_{\perp} e_2 = e_3 \end{cases} & \alpha = \begin{pmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{pmatrix}; b, c \in \mathbb{R}.
\end{aligned}$$

Proposition 5.3. *The following matrix forms characterize averaging operators of 2-dimensional Hom-quadri-dendriform algebras:*

$$\begin{aligned}
D_1 : & \begin{pmatrix} \vartheta_{11} & 0 \\ \vartheta_{21} & \vartheta_{11} \end{pmatrix} \begin{pmatrix} 0 & 0 \\ \vartheta_{21} & \vartheta_{22} \end{pmatrix}; \quad D_2 : \begin{pmatrix} \vartheta_{11} & 0 \\ \vartheta_{21} & \vartheta_{11} \end{pmatrix} \begin{pmatrix} 0 & 0 \\ \vartheta_{21} & \vartheta_{22} \end{pmatrix} \\
D_3 : & \begin{pmatrix} \vartheta_{22} & 0 \\ 0 & \vartheta_{22} \end{pmatrix}; \quad D_4 : \begin{pmatrix} \vartheta_{11} & 0 \\ 0 & \vartheta_{11} \end{pmatrix} \begin{pmatrix} 0 & \vartheta_{12} \\ 0 & 0 \end{pmatrix}; \quad D_5 : \begin{pmatrix} \vartheta_{22} & 0 \\ 0 & \vartheta_{22} \end{pmatrix}.
\end{aligned}$$

Proposition 5.4. *The following matrix forms characterize averaging operators of 3-dimensional Hom-quadri-dendriform algebras:*

$$\begin{aligned}
D_1 : & \begin{pmatrix} \vartheta_{11} & 0 & 0 \\ 0 & \vartheta_{11} & 0 \\ 0 & 0 & \vartheta_{11} \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 \\ 0 & \vartheta_{22} & \vartheta_{23} \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \vartheta_{23} \end{pmatrix}; \quad D_2 : \begin{pmatrix} \vartheta_{22} & 0 & 0 \\ 0 & \vartheta_{22} & 0 \\ 0 & 0 & \vartheta_{22} \end{pmatrix}; \\
D_3 : & \begin{pmatrix} \vartheta_{33} & 0 & 0 \\ 0 & \vartheta_{33} & 0 \\ 0 & 0 & \vartheta_{33} \end{pmatrix} \begin{pmatrix} 0 & \vartheta_{12} & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}; \quad D_4 : \begin{pmatrix} \vartheta_{22} & 0 & 0 \\ 0 & \vartheta_{22} & 0 \\ 0 & 0 & \vartheta_{22} \end{pmatrix} \begin{pmatrix} 0 & 0 & \vartheta_{13} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}; \\
D_5 : & \begin{pmatrix} \vartheta_{33} & 0 & 0 \\ 0 & \vartheta_{33} & 0 \\ 0 & 0 & \vartheta_{33} \end{pmatrix} \begin{pmatrix} \vartheta_{11} & 0 & \vartheta_{13} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 \\ 0 & \vartheta_{22} & 0 \\ 0 & 0 & 0 \end{pmatrix};
\end{aligned}$$

$$\begin{aligned}
D_6 &: \begin{pmatrix} \vartheta_{22} & 0 & 0 \\ 0 & \vartheta_{22} & 0 \\ 0 & 0 & \vartheta_{22} \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 \\ \vartheta_{21} & \vartheta_{22} & 0 \\ i\vartheta_{21} & i\vartheta_{22} & 0 \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \vartheta_{33} \end{pmatrix}; \\
D_7 &: \begin{pmatrix} \vartheta_{33} & 0 & 0 \\ 0 & \vartheta_{33} & 0 \\ 0 & 0 & \vartheta_{33} \end{pmatrix} \begin{pmatrix} 0 & -\vartheta_{33} & 0 \\ 0 & \vartheta_{33} & 0 \\ 0 & \vartheta_{33} & 0 \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \vartheta_{33} \end{pmatrix}; \\
D_8 &: \begin{pmatrix} \vartheta_{11} & 0 & \vartheta_{13} \\ \vartheta_{21} & \vartheta_{22} & \vartheta_{23} \\ \vartheta_{22} - \vartheta_{11} & 0 & \vartheta_{22} - \vartheta_{13} \end{pmatrix} \begin{pmatrix} \vartheta_{11} & -\vartheta_{12} & \vartheta_{13} \\ \vartheta_{21} & \vartheta_{22} & \vartheta_{23} \\ -\vartheta_{11} & -\vartheta_{12} & -\vartheta_{13} \end{pmatrix}; D_9 : \begin{pmatrix} \vartheta_{33} & 0 & 0 \\ 0 & \vartheta_{33} & 0 \\ 0 & 0 & \vartheta_{33} \end{pmatrix}; \\
D_{10} &: \begin{pmatrix} \vartheta_{33} & 0 & 0 \\ 0 & \vartheta_{33} & \vartheta_{23} \\ 0 & 0 & \vartheta_{33} \end{pmatrix} \begin{pmatrix} 0 & 0 & \vartheta_{13} \\ 0 & 0 & \vartheta_{23} \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} \vartheta_{33} & 0 & 0 \\ 0 & 0 & \vartheta_{23} \\ 0 & 0 & \vartheta_{33} \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 \\ \vartheta_{21} & \vartheta_{22} & \vartheta_{23} \\ 0 & 0 & 0 \end{pmatrix}; \\
D_{11} &: \begin{pmatrix} \vartheta_{22} & 0 & 0 \\ 0 & \vartheta_{22} & 0 \\ 0 & 0 & \vartheta_{22} \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & \vartheta_{32} & 0 \end{pmatrix} \begin{pmatrix} 0 & -\vartheta_{22} & 0 \\ 0 & \vartheta_{22} & 0 \\ 0 & \vartheta_{22} & 0 \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ -\vartheta_{33} & \vartheta_{32} & \vartheta_{33} \end{pmatrix}; \\
D_{12} &: \begin{pmatrix} \vartheta_{11} & 0 & 0 \\ 0 & \vartheta_{11} & 0 \\ 0 & 0 & \vartheta_{11} \end{pmatrix} \begin{pmatrix} 0 & \vartheta_{12} & \vartheta_{13} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}; \\
D_{13} &: \begin{pmatrix} \vartheta_{11} & \vartheta_{12} & \vartheta_{13} \\ -\vartheta_{11} & -\vartheta_{12} & -\vartheta_{31} \\ \vartheta_{31} & \vartheta_{32} & \vartheta_{33} \end{pmatrix} \begin{pmatrix} \vartheta_{11} & \vartheta_{12} & 0 \\ \vartheta_{21} & \vartheta_{11} - \vartheta_{12} + \vartheta_{21} & 0 \\ \vartheta_{12} & \vartheta_{32} & \vartheta_{11} + \vartheta_{22} \end{pmatrix}.
\end{aligned}$$

Acknowledgement

The authors thank the anonymous referees for their valuable suggestions and comments.

Conflicts of Interest

The authors declare no conflicts of interest.

REFERENCES

1. A. Das. *Cohomology and deformations of dendriform algebras and Dend_∞-algebras*. Comm. Algebra 202250(4) : 1544-1567.
2. Weili Cao *AN ALGEBRAIC STUDY OF AVERAGING OPERATORS* August 30 2000.
3. A. Makhlouf A. *Hom-dendriform algebras and Rota-Baxter Hom-algebras Operads and uni-versal algebra* 147-171 Nankai Ser. Pure Appl. Math. Theoret. Phys. 9 World Sci. Publ. Hackensack NJ 2012.
4. A. Makhlouf and A. Zahari *Structure and Classification of Hom-Associative Algebras*. Acta et commentationes universitatis Tartuensis de mathematica Vol. 24 No. 1 (2020). pp. 79-102.
5. Mosbahi, B. , Basdouri, I. and Zahari, A. (2025). Classification, derivations and centroids of low-dimensional associative trialgebras. Galois Journal of Algebra, 1(1), 118-136.
6. Basri W. Rakhimov I. Rikhsiboev I. *Classification of 3-Dimensional Complex Diassociative Algebras* Malaysian Journal of Mathematical Sciences Vol. 4 No. 2 (2010).
7. Basri W. Rakhimov I. Rikhsiboev I. *Four-Dimensional Nilpotent Diassociative Algebras* Journal of Generalized Lie Theory and Applications Vol. 9 No. 1 (2015).
8. Blokh A *generalization of the concept of a Lie algebra* . In Doklady Akademii Nauk (Vol. 165 No. 3 pp. 471-473) Russian Academy of Sciences. A. Y. (1965).
9. I. Basdouri, S. Benabdelhafidh, A. Saghrouni, S. Silvestrov, *Cohomology and deformations of crossed homomorphisms on Lie conformal algebras*, Adv. Stud. Euro-Tbilisi Math. J. **18** (3), 91-107.
10. B. Imed, G. Salima, S. Mohamed Amin, *Maurer-Cartan characterization and cohomology of compatible LieDer and AssDer pairs*, Adv. Stud. Euro-Tbilisi Math. J. 18(3): 173-205.

11. B. Imed, B. Sami, N. Saber, S. Mohamed Amin, *Cohomologies of modified Rota-Baxter Lie algebras with derivations and applications*, Adv. Stud. Euro-Tbilisi Math. J. 18 (2), 189-210.
12. I. Basdouri, E. Peyghan, M. A. Sadraoui, R. Saha, *Formal deformations and extensions of "twisted" Lie algebras*, Adv. Stud. Euro-Tbilisi Math. J. 18 (3), 225-239.
13. B. Mosbahi I. Basdouri A. Zahari *Classification α -Inner Derivations and α -Centroids of Finite-Dimensional Complex Hom-Trialgebras*, Pure and Applied Mathematics Journal Vol. 12 No. 5 (2023), pp. 86-97.
14. Daniel Larsson and Sergei D. Silvestrov *Quasi-hom-Lie algebras central extensions and 2-cocycle-like identities* Centre for Mathematical Sciences Department of Mathematics Lund Institute of Technology Lund University Sweden 288 (2005) 321-344.
15. K. Ebrahimi-Fard *Loday-type algebras and the Rota-Baxter relation* Lett. Math. Phys. 61 (2002) 139-147.
16. K. Ebrahimi-Fard D. Manchon and F. Patras *New identities in dendriform algebras* J. Algebra 320 (2008) 708-727.
17. J.-L. Loday M. Ronco. *Trialgebras and families of polytopes Homotopy theory: relations with algebraic geometry group cohomology and algebraic K-theory*. 369-398 Contemp. Math. 346 Amer. Math. Soc. Providence RI 2004.
18. J.-L. Loday *Dialgebras Dialgebras and related operads*. 7-66 Lecture Notes in Math. 1763 Springer Berlin 2001.
19. Makhlouf A. and Sergei D. Silvestrov *Hom-algebras and Hom-coalgebras* Journal of Algebra and Its Applications Vol. 9 (4) 553-589 (2010).
20. Y. Y. Zhang X. Gao and D. Manchon *Free (tri)dendriform family algebras* J. Algebra 547 (2020) 456-493.
21. Makhlouf A. and Silvestrov S. D. *Hom-algebra structures* J. Gen. Lie Theory Appl. 2 (2) (2008) 51-64.
22. Wen Teng. *Splitting of operations for di-associative algebras and tri-associative algebras* arXiv:2502.21070v1 (2025).
23. C. Bai O. Bellier L. Guo X. Ni *Splitting of operations Manin products and Rota Baxter operators* Int. Math. Res. Not. 2013 2013(3): 485-524.
24. C. Bai O. Bellier L. Guo X. Ni *Replicators Manin white product of binary operads and average operators*. . Conference: 3rd International Congress in Algebras and Combinatorics 2020.
25. M. Aguiar *Pre-Poisson algebras*. Lett. Math. Phys. 2000 54(4): 263-277.
26. A. Zahari *Classification α^k -Derivations and Centroids of 4-dimensional Complex Hom-associative Dialgebras* arXiv : 2305.04041 (2023).
27. A. Zahari and Bakayoko I. *On BiHom-Associative dialgebras* Open Journal of Mathematical Sciences Vol. 7 No. 1 (2023).pp. 96-117.
28. A. Zahari and A. Sania, *Classification of 3-dimensional Bihom-Associative and Bihom-Bialgebras*, Open J. Math. Sci. **9** (2025), 182-201.

¹ UNIVERSITY OF CARTHAGE· HIGHER INSTITUTE OF SCIENCES AND TECHNOLOGY OF ENVIRONMENT OF BORJ CEDRIA (IS-TEUB), TUNISIA

² UNIVERSITÉ DE GAFSA, FACULTÉ DES SCIENCES DE GAFSA, TUNIS

³DÉPARTEMENT DE MATHÉMATIQUES, FACULTÉ DES SCIENCES, UNIVERSITÉ DE SFAX, SFAX, TUNISIA

⁴IRIMAS - DÉPARTEMENT DE MATHÉMATIQUES, FACULTÉ DES SCIENCES, UNIVERSITÉ DE HAUTE ALSACE, MULHOUSE, FRANCE

Email address: ¹abdelkader.hamdouni@isimg.rnu.tn

Email address: ²basdourimed@yahoo.fr

Email address: ³mariemj090@gmail.com

Email address: ⁴abdou-damdji.ahmed-zahari@uha.fr