

Kazhdan groups of dimension 16 with an uncountable range of second ℓ^2 -Betti numbers

Francesco Fournier-Facio and Roman Sauer

January 5, 2026

Abstract

We construct a family of simple, lacunary hyperbolic groups with property (T) that have rational cohomological dimension 16 and uncountably many values of second ℓ^2 -Betti numbers. Along the way, we present new constructions of measurably diverse finitely generated groups, and we prove that the second ℓ^2 -Betti number is far from being semi-continuous in the space of marked groups, even assuming good finiteness properties.

1 Introduction

In relation to the Atiyah Conjecture [Ati76], there have been many constructions of matrices over group rings of finitely generated groups whose kernels have irrational von Neumann dimension [Aus13, Gra14, Gra16]. Much less is known about ℓ^2 -Betti numbers of groups: the main open question is whether there exists a group of type $FP_{n+1}(\mathbb{Q})$ with irrational n -th ℓ^2 -Betti number [Gra14, Question 4], but even without the finiteness conditions, we only know of one construction for finitely generated groups. Using a construction of Gaboriau [Gab00, Proposition VI.16], one can build finitely generated groups with an uncountable range of first ℓ^2 -Betti numbers [ITD25, Proposition 5.1].

We present a new construction of groups taking exotic values of second ℓ^2 -Betti numbers, and with several additional desirable properties. Given a set π of primes, we say that a group G has *cohomological dimension n modulo π* , if G has cohomological dimension n over every ring in which each prime from π is invertible. Note that if $n < \infty$, this implies that G only contains π -torsion. The ℓ^2 -*Euler characteristic* of a group, when defined, is the alternating sum of its ℓ^2 -Betti numbers [ITD25, Proposition 5.1].

Theorem A. *Let $p \neq q$ be primes. There exists an uncountable family of groups $(G_\alpha)_{\alpha \in A}$ and an integer $\chi \geq 1$ with the following properties.*

1. G_α is simple, has property (T), and is lacunary hyperbolic.
2. G_α has cohomological dimension 16 modulo $\{p, q\}$.
3. The second ℓ^2 -Betti numbers $b_2^{(2)}(G_\alpha)$ are pairwise distinct. Further, $b_n^{(2)}(G_\alpha) < \infty$ for every $n \in \mathbb{N}$ and every $\alpha \in A$.
4. The ℓ^2 -Euler characteristic is $\chi^{(2)}(G_\alpha) = b_2^{(2)}(G_\alpha) + \chi$.

By renouncing simplicity, we can also construct groups where a single prime obstructs the finiteness of the cohomological dimension: see Theorem 6.1. Property (T) distinguishes these groups from the examples obtained from Gaboriau's construction, which arise as amalgamated products [ITD25, Proposition 5.1]. Lacunary hyperbolicity distinguishes them from the counterexamples to the general Atiyah Conjecture, since these all contain lamplighters [Gra16, Problem 3], while lacunary hyperbolic groups cannot (Proposition 4.11).

As noticed by Ioana–Tucker-Drob, it follows easily from Gaboriau's construction [Gab00, Proposition VI.16] that there exist uncountably many pairwise non-measure equivalent groups [ITD25, Proposition 5.1]. However, in measurable group theory, property (T) is especially desirable, and last year two constructions of measurably diverse families of property (T) groups have appeared. One by the first author and Sun [FFS25], where these groups are distinguished by the whole sequence of their ℓ^2 -Betti numbers - in the same vein as the paper by López Neumann [LN23], which constructs an infinite family of finitely presented simple groups. Then very recently by Ioana–Tucker-Drob [ITD25], where the groups have vanishing ℓ^2 -Betti numbers, and arise from the technology of wreath-like products [CIOS23]. In particular, these groups have infinite rational cohomological dimension.

Corollary B. *The groups from Theorem A are pairwise non-measure equivalent.*

Proof. Let G_α and G_β be groups from the family. If they are measure equivalent with index $c > 0$, then $b_n^{(2)}(G_\alpha) = cb_n^{(2)}(G_\beta)$ for all $n \in \mathbb{N}$ by Gaboriau's theorem [Gab02], and thus $\chi^{(2)}(G_\alpha) = c\chi^{(2)}(G_\beta)$. Hence

$$cb_2^{(2)}(G_\beta) + \chi = b_2^{(2)}(G_\alpha) + \chi = \chi^{(2)}(G_\alpha) = c\chi^{(2)}(G_\beta) = cb_2^{(2)}(G_\beta) + c\chi.$$

Since χ is a positive integer, this shows that $c = 1$, and thus $b_2^{(2)}(G_\alpha) = b_2^{(2)}(G_\beta)$, so $\alpha = \beta$. \square

A version of the construction leading to Theorem A produces torsion-free hyperbolic groups with prescribed integer second ℓ^2 -Betti number (Theorem 5.1). This has the following consequence.

Theorem C. *Let $(n_i)_{i \in \mathbb{N}}$ be a sequence of natural numbers. There exist property (T) torsion-free hyperbolic groups G, G_i of cohomological dimension 16, with quotient maps $G \rightarrow G_i$ such that $G_i \xrightarrow{i \rightarrow \infty} G$ in the space of marked groups, and*

$$b_2^{(2)}(G_i) = n_i \text{ for all } i \in \mathbb{N} \text{ but } b_2^{(2)}(G) = 0.$$

Therefore the second ℓ^2 -Betti number is far from being semi-continuous in the space of marked groups, even with the strong finiteness properties that are enjoyed by torsion-free hyperbolic groups. This is in contrast with the semi-continuity of the first ℓ^2 -Betti number [Pic06].

Our computations of ℓ^2 -Betti numbers exploit the work of Petrosyan–Sun on group-theoretic Dehn filling [PS24a]. More recently [PS24b], they applied this theory to compute ℓ^2 -Betti numbers, by working in convenient classes where the Lück approximation and Atiyah conjectures are known to hold: virtually locally indicable and cocompactly cubulated groups. Note that these are incompatible with property (T).

We instead take a different route, and start from a hyperbolic group enjoying the strong rigidity property (T_2) , introduced by Bader and the second author [BS23, BS25]. This group

is a torsion-free cocompact lattice in the isometry group of the octonionic hyperbolic plane (Theorem 3.10), which is the source of the number 16 in our results. Under this condition, building on an observation from [FF25], we will see that the excision theorem from [PS24a] is sufficient to give exact computations of the second ℓ^2 -Betti number of iterated group-theoretic Dehn fillings.

All of the constructions mentioned above, both in this paper and in others, are non-elementary to various extents. Because of this, we feel it is worth presenting possibly the most elementary construction of finitely generated groups achieving uncountably many values of ℓ^2 -Betti numbers.

Theorem D. *Let p be a prime. There exists a family of finitely generated groups $(G_x)_{x \in \mathbb{R}_{>0}}$ such that $b_1^{(2)}(G_x) = 0$, $b_2^{(2)}(G_x) = \infty$, and $b_3^{(2)}(G_x) = x$. Moreover, G_x is residually finite and has cohomological dimension 6 modulo p .*

Taking a free product with \mathbb{Z} gives an uncountable family of pairwise non-measure equivalent groups (Corollary 7.5). Of course, none of these groups have property (T).

Acknowledgements. FFF is supported by the Herchel Smith Postdoctoral Fellowship Fund. The authors would like to thank the Isaac Newton Institute for Mathematical Sciences, Cambridge, for support and hospitality during the programme Operators, Graphs, Groups, where work on this paper was undertaken. This work was supported by EPSRC grant EP/Z000580/1.

2 Homology and cohomology

We start by collecting some preliminary facts about homology and cohomology of groups. We fix a ring R for the rest of this section. All rings in this paper are unital.

First is an easy application of the transfer, whose cohomological version is given in [Bro94, Proposition III.10.4 on p. 85]. The homological version can be obtained similarly via the transfer, alternatively it is a direct consequence of the Lyndon–Hochschild–Serre spectral sequence [Bro94, VII.6 on p. 171].

Proposition 2.1. *Let $N < G$ a normal subgroup of finite index, and suppose that $[G : N]$ is invertible in R . Then for every $R[G]$ -module M and every $n \in \mathbb{N}$, the corestriction induces an isomorphism*

$$H_n(N; M)_{G/N} \cong H_n(G; M).$$

Next, we look at homology and cohomology of colimit groups, which will be especially relevant for the groups from Theorem A.

Proposition 2.2 ([Bie81, Proposition 4.8 on p. 61]). *Let $G_1 \rightarrow G_2 \rightarrow \dots$ be a directed sequence of groups with colimit G_∞ , and let M be an $R[G_\infty]$ -module. Then there is an isomorphism*

$$H_n(G_\infty; M) \cong \operatorname{colim} H_n(G_i; M).$$

This is the standard fact that homology commutes with filtered colimits. This is of course not true in cohomology, but it does work under some additional assumptions. We point out that the following is only used to compute cohomological dimensions of the groups

from Theorem A and Theorem D. If the reader is happy with the standard estimate of cohomological dimension as one more than the homological dimension [Bie81, Theorem 4.6 on p. 60], then they can use Proposition 2.2, and move to the next section.

Proposition 2.3. *Let $G_1 \rightarrow G_2 \rightarrow \dots$ be a directed sequence of groups with colimit G_∞ , and let M be an $R[G_\infty]$ -module. Suppose that the pullback $H^n(G_{i+1}; M) \rightarrow H^n(G_i; M)$ is surjective for all $i \in \mathbb{N}$. Then*

$$H^{n+1}(G_\infty; M) \cong \varprojlim H^{n+1}(G_i; M).$$

Proof. A tower of abelian groups

$$\dots \rightarrow D_2 \rightarrow D_1$$

is said to satisfy the *Mittag-Leffler condition* if for each k there exists $j \geq k$ such that for all $i \geq j$ the image of $D_i \rightarrow D_k$ equals the image of $D_j \rightarrow D_k$ [Wei94, Definition 3.5.6 on p. 82]. We will replace this with the stronger and easier to check condition that each map $D_{i+1} \rightarrow D_i$ is surjective.

Let X_1 be a model of the classifying space BG_1 ¹. Using mapping cones, we may inductively define X_{i+1} as a model of BG_{i+1} such that the map $X_i \rightarrow X_{i+1}$ is an inclusion of CW-complexes. The colimit X , in this case a union, of the X_i is a model of BG_∞ . The universal covering \tilde{X} , which is a free G_∞ -CW complex, is a model of EG_∞ . Let \bar{X}_i be the restriction of the universal covering to X_i . By covering theory we have $\bar{X}_i \cong G_\infty \times_{G_i} \tilde{X}_i$. The cellular cochain complexes

$$D_i = \hom_{R[G_\infty]}(C_*(\bar{X}_i), M) \cong \hom_{R[G_i]}(C_*(\tilde{X}_i), M)$$

compute the cohomology $H^*(G_i; M)$, and fit into a tower of cochain complexes. Since \tilde{X} is the colimit of the G_∞ -CW complexes \tilde{X}_i , the projective limit of the D_i is the cochain complex $\hom_{R[G_\infty]}(C_*(\tilde{X}_i), M)$, which computes the cohomology of G_∞ .

By our choice of X_i , the maps $D_{i+1} \rightarrow D_i$ are degree-wise surjective because $C_*(\bar{X}_i)$ is degree-wise a direct $R[G_\infty]$ -summand of $C_*(\tilde{X}_i)$, so we can apply [Wei94, Theorem 3.5.8 on p. 83]² and obtain a short exact sequence

$$0 \rightarrow \varprojlim^1 H^n(G_i; M) \rightarrow H^{n+1}(G_\infty; M) \rightarrow \varprojlim H^{n+1}(G_i; M) \rightarrow 0.$$

By the assumption on surjectivity, the \varprojlim^1 term vanishes [Wei94, Proposition 3.5.7 on p. 83], so we obtained the desired isomorphism. \square

Corollary 2.4. *Let $G_1 \rightarrow G_2 \rightarrow \dots$ be a directed sequence of groups with colimit G_∞ . Suppose that the following hold for all $i \in \mathbb{N}$.*

- For every $R[G_\infty]$ -module M the pullback $H^n(G_{i+1}; M) \rightarrow H^n(G_i; M)$ is surjective.
- $cd_R(G_i) = n$;
- $H^n(G_1; R) \neq 0$ and the pullback $H^{n-1}(G_{i+1}; R) \rightarrow H^{n-1}(G_i; R)$ is surjective.

Then $cd_R(G_\infty) = n$.

¹also denoted by $K(G_1, 1)$.

²the citation is about homology of chain complexes, hence the left-hand side term being one degree below instead of above

Proof. By the assumption on surjectivity we can apply Proposition 2.3. First, for every $R[G_\infty]$ -module M :

$$H^{n+1}(G_\infty; M) \cong \varprojlim H^{n+1}(G_i; M) \cong \varprojlim 0 \cong 0,$$

thus $cd_R(G_\infty) \leq n$. Second,

$$H^n(G_\infty; R) \cong \varprojlim H^n(G_i; R) \twoheadrightarrow H^n(G_1; R) \neq 0,$$

and so $cd_R(G_\infty) \geq n$. □

3 ℓ^2 -Betti numbers of tracial unitary representations

We refer the reader to [Lück02] for details and background, where everything is developed for group von Neumann algebras. Lück formulated his dimension theory for arbitrary finite von Neumann algebras in [Lück98].

A unitary G -representation π on a Hilbert space V_π is called *tracial* if the associated von Neumann algebra $\mathcal{N}(\pi) := \pi(G)''$, which is the double commutant of $\pi(G)$ in the algebra of bounded operators on V_π , has a cyclic vector $\xi_\pi \in V_\pi$ such that

$$\tau(T) := \langle T\xi_\pi, \xi_\pi \rangle_{V_\pi}$$

defines a finite trace on $\mathcal{N}(\pi)$. The embedding

$$\mathcal{N}(\pi) \hookrightarrow V_\pi : T \mapsto T\xi_\pi \tag{1}$$

is isometric with respect to the norm $\|T\|^2 = \tau(T^*T)$ on $\mathcal{N}(\pi)$ and has dense image. The embedding is naturally a map of left $\mathcal{N}(\pi)$ -modules. It extends to an isometric isomorphism of the completion $L^2(\mathcal{N}(\pi), \tau) \cong V_\pi$. Via this isomorphism, V_π has a natural right $\mathcal{N}(\pi)$ -module structure that commutes with the left $\mathcal{N}(\pi)$ -module structure. On the dense set of vectors $T\xi_\pi$, the von Neumann algebra $\mathcal{N}(\pi)$ acts from the right via $(T\xi_\pi) \cdot S = TS\xi_\pi$.

We regard the cyclic vector, and therefore τ , as part of the data of a tracial unitary representation. Via the Gelfand–Naimark–Segal (GNS) construction, tracial unitary G -representations are in one-to-one correspondence with conjugation-invariant positive functions on G [BdlHV08, Theorem C.4.10 on p. 354].

Example 3.1. The von Neumann algebra associated to the regular representation λ_G on $\ell^2(G)$ with cyclic vector 1_G is the *group von Neumann algebra* of G , which we denote by $\mathcal{N}(G)$.

If $G \rightarrow Q$ is a surjective group homomorphism, then composing it with λ_Q yields a tracial unitary G -representation on $\ell^2(Q)$.

Lück introduced a dimension function

$$\dim_{\mathcal{N}} : \{\mathcal{N}\text{-modules}\} \rightarrow \mathbb{R}_{\geq 0} \cup \{\infty\}$$

for arbitrary (algebraic) modules over a finite von Neumann algebra \mathcal{N} with finite normalised trace τ . The choice of τ is understood to be specified, and we usually drop τ from the notation. Lück's dimension function is additive for short exact sequences of \mathcal{N} -modules and normalised, i.e. $\dim_{\mathcal{N}}(\mathcal{N}) = 1$. Moreover:

Lemma 3.2 ([Lü98, Theorem 0.6 (b) and (c)]). *Let $(V_i)_{i \in \mathbb{N}}$ be a sequence of \mathcal{N} -modules. Then*

$$\dim_{\mathcal{N}} \left(\bigoplus_{i \in \mathbb{N}} V_i \right) = \sum_{i \in \mathbb{N}} \dim_{\mathcal{N}}(V_i).$$

Let X be a topological G -space. We do not require a cellular structure on X nor freeness of the action. Let π be a tracial unitary G -representation. The equivariant homology

$$H_n^G(X, \mathcal{N}(\pi)) = H_n(\mathcal{N}(\pi) \otimes_{\mathbb{Z}[G]} C_*(X))$$

is naturally a left $\mathcal{N}(\pi)$ -module. Here $C_*(X)$ denotes the singular chain complex of X . We have

$$H_n(G; \mathcal{N}(\pi)) = H_n^G(EG; \mathcal{N}(\pi)).$$

Definition 3.3. Let π be a tracial unitary G -representation. The ℓ^2 -Betti numbers of π are defined as

$$b_n^{(2)}(G; \pi) = \dim_{\mathcal{N}(\pi)} H_n(G; \mathcal{N}(\pi)).$$

When π is the regular representation, we simply write $b_n^{(2)}(G)$. When $G \rightarrow Q$ is a quotient, and π is the pullback of the regular representation of Q , we write $b_n^{(2)}(G \rightarrow Q)$.

Definition 3.4. Let π be a tracial unitary G -representation such that $\sum_{n \in \mathbb{N}} b_n^{(2)}(G; \pi) < \infty$. Then we define the ℓ^2 -Euler characteristic of π as

$$\chi^{(2)}(G; \pi) = \sum_{n \in \mathbb{N}} (-1)^n b_n^{(2)}(G; \pi).$$

When π is the regular representation we just write $\chi^{(2)}(G)$.

The next result generalises the Euler–Poincaré formula for the regular representation to arbitrary tracial unitary representations.

Proposition 3.5. *Let G be a group of type $FP(\mathbb{Q})$. Then the Euler characteristic of G and the ℓ^2 -Euler characteristic of every tracial unitary G -representation π coincide:*

$$\chi(G) = \chi^{(2)}(G; \pi).$$

Proof. Let P_* be a finite projective $\mathbb{Q}[G]$ -resolution of the trivial $\mathbb{Q}[G]$ -module \mathbb{Q} . Then $C_* = \mathcal{N}(\pi) \otimes_{\mathbb{Q}[G]} P_*$ is a finite projective $\mathcal{N}(\pi)$ -chain complex. The proof that

$$\sum_{n \geq 0} (-1)^n \dim_{\mathcal{N}(\pi)}(C_n) = \sum_{n \geq 0} (-1)^n \dim_{\mathcal{N}(\pi)}(H_n(C_*))$$

is the same as the Euler–Poincaré formula for the classical Euler characteristic and basically follows from the additivity of the dimension. See the standard proof in Hatcher’s book [Hat02, Theorem 2.44 on p. 146]. \square

The following result is an analogue of [PT11, Theorem 2.2], which treated the cohomological version, and only for the regular representation. According to the discussion after Eq. (1), the Hilbert space V_π carries a left $\mathcal{N}(\pi)$ -module structure and a right $\mathcal{N}(\pi)$ -module structure, in particular, also a right $\mathbb{C}[G]$ -module structure. So the homology on the right hand side is naturally a left $\mathcal{N}(\pi)$ -module.

Proposition 3.6. *Let π be a tracial unitary G -representation on the Hilbert space V_π . Then we have*

$$b_n^{(2)}(G; \pi) = \dim_{\mathcal{N}(\pi)} H_n(G; V_\pi)$$

for every $n \in \mathbb{N}$.

Proof. The embedding $\mathcal{N}(\pi) \hookrightarrow V_\pi$ of from Eq. (1) is rank dense:

One calls a $\mathcal{N}(\pi)$ -linear homomorphism $f: M \rightarrow N$ of $\mathcal{N}(\pi)$ -modules *rank dense* if for every $x \in N$ there is an increasing sequence of projections $p_i \in \mathcal{N}(\pi)$, $i \in \mathbb{N}$, converging to id such that $p_i x \in N$. By [KPV15, Lemma A.5] the embedding $\mathcal{N}(\pi) \hookrightarrow V_\pi$ is rank dense because it is dense.

The embedding $\mathcal{N}(\pi) \hookrightarrow V_\pi$ is also right $\mathcal{N}(\pi)$ -linear, in particular, an embedding of right $\mathbb{C}[G]$ -modules, and the right $\mathbb{C}[G]$ -action commutes with the left $\mathcal{N}(\pi)$ -action. Now we consider the quotient $W := V_\pi / \mathcal{N}(\pi)$; this is an undesirable object from a functional-analytic perspective, however it makes sense to look at W as an (algebraic) $\mathcal{N}(\pi)$ -module. By rank density for every $x \in W$ there is an increasing sequence of projections $p_i \in \mathcal{N}(\pi)$ converging to id such that $p_i x = 0$ for every $i \in \mathbb{N}$. By the local criterion [Sau05, Theorem 2.4] we obtain that $\dim_{\mathcal{N}(\pi)}(W) = 0$. Let P_* be a free $\mathbb{C}[G]$ -resolution of \mathbb{C} of finite type. We have a short exact sequence of $\mathcal{N}(\pi)$ -chain complexes

$$0 \rightarrow \mathcal{N}(\pi) \otimes_{\mathbb{C}[G]} P_* \rightarrow V_\pi \otimes_{\mathbb{C}[G]} P_* \rightarrow W \otimes_{\mathbb{C}[G]} P_* \rightarrow 0.$$

By Lemma 3.2, the right hand side is a chain complex of modules of $\dim_{\mathcal{N}(\pi)}$ -dimension zero. By additivity and the long exact sequence in homology, the statement follows. \square

Proposition 3.6 will be especially useful in conjunction with higher property (T) .

Definition 3.7 ([BS23]). We say that a group G has *property $[T_n]$* if for every $i \leq n$ and for every unitary representation V , it holds $H^i(G; V) = 0$. We say that G has *property (T_n)* if the vanishing holds under the additional assumption that $V^G = 0$.

Since we will mostly be working with homology, the following will be more useful.

Proposition 3.8 ([BS25]). *Let G be a group with property $[T_n]$ (respectively, (T_n)) and of type $FP_{n+1}(\mathbb{Q})$. Then for every $i \leq n$ and for every unitary representation V (respectively, every unitary representation V with $V^G = 0$), it holds $H_i(G; V) = 0$.*

Proof. Combine [BS25, Lemma 29] and [BS25, Lemma 31]. \square

Corollary 3.9. *Let G be a group with property $[T_n]$ (respectively, (T_n)) and of type $FP(\mathbb{Q})$, and let Q be a quotient (respectively, an infinite quotient) of G . Then for every $i \leq n$ it holds $b_i^{(2)}(G \rightarrow Q) = 0$. Hence*

$$\chi^{(2)}(G) = \sum_{i>n} (-1)^i b_i^{(2)}(G \rightarrow Q).$$

We recall that hyperbolic groups are of type $FP(\mathbb{Q})$ [BH99, Theorem III.Γ.3.21 on p 468].

Proof. Combine Propositions 3.6 and 3.8. The second statement now follows from Proposition 3.5. \square

Our starting point will be the following example.

Theorem 3.10 ([BS23]). *Let G be a cocompact lattice in the isometry group of the octonionic hyperbolic plane. Then G has properties (T_3) and $[T_2]$.*

In the proofs of Theorems A and C, we will actually only need that G has property (T_2) : see Remark 5.3.

Proof. The ambient Lie group has property (T_3) by [BS23, Theorem B and Appendix A]. We emphasise that this relies only on a small part of [BS23]: It follows from a result of Kumaresan [Kum80, Theorem 3] for non-trivial irreducible unitary representation of the ambient Lie group, and the ultrapower technique in [BS23]. Finally, property (T_3) passes to cocompact lattices by [BS23, Lemma 3.13]. Moreover, the second Betti number of G vanishes, because it is the fundamental group of a non-Hermitian locally symmetric space of rank one [Mat62], so G also has $[T_2]$. \square

We conclude with a useful fact about subgroups.

Proposition 3.11 ([Lüc02, Theorem 6.29 on p. 253]). *Let $G \rightarrow Q$ be a homomorphism, and let $Q \rightarrow L$ be an injective homomorphism. Then*

$$\dim_{\mathcal{N}(Q)} H_n(G; \mathcal{N}(Q)) = \dim_{\mathcal{N}(L)} H_n(G; \mathcal{N}(L)),$$

for all $n \in \mathbb{N}$.

4 Group-theoretic Dehn filling

In this section we introduce group-theoretic Dehn filling, and do some preliminary work towards our main construction.

4.1 Cohen–Lyndon triples

Let $P < G$ and let N be a normal subgroup of P . We denote $\bar{P} := P/N$ and $\bar{G} := G/\langle\langle N \rangle\rangle$. Such a quotient is called a *group-theoretic Dehn filling of G* .

Definition 4.1. We say that (G, P, N) is a *Cohen–Lyndon triple* if there exists a set of transversals T of $G/P \langle\langle N \rangle\rangle$ such that

$$\langle\langle N \rangle\rangle = *_T t N t^{-1}.$$

This notion goes back to the work of Cohen–Lyndon on one-relator groups [CL63]; in this form it is defined in [Sun20]. The most important consequence is the following algebraic excision theorem.

Theorem 4.2 ([PS24a, Theorem A (ii)], see also [PS24b, Corollary 4.5]). *Let (G, P, N) be a Cohen–Lyndon triple. Then for all $\mathbb{Z}[\bar{G}]$ -modules M and all $n \in \mathbb{N}$, the quotient maps $G \rightarrow \bar{G}$ and $P \rightarrow \bar{P}$ induce isomorphisms*

$$H_n(G, P; M) \cong H_n(\bar{G}, \bar{P}; M) \quad \text{and} \quad H^n(G, P; M) \cong H^n(\bar{G}, \bar{P}; M).$$

We will only be concerned with an especially well-behaved case.

Corollary 4.3. *Let (G, P, N) be a Cohen–Lyndon triple. Suppose that P is virtually cyclic, N is infinite cyclic, and hence \bar{P} is finite. Let R be a ring where $[P : N]$ is invertible. Then for all $R[\bar{G}]$ -modules M and all $n \geq 3$, the quotient map $G \rightarrow \bar{G}$ induces isomorphisms*

$$H_n(G; M) \cong H_n(\bar{G}; M) \quad \text{and} \quad H^n(G; M) \cong H^n(\bar{G}; M).$$

Moreover, there is an exact sequence

$$0 \rightarrow H_2(G; M) \rightarrow H_2(\bar{G}; M) \rightarrow M_{\bar{P}} \rightarrow H_1(G; M).$$

Proof. By the assumption on invertibility of $[P : N]$, we have $cd_R(\bar{P}) = 0$ and $cd_R(P) = 1$. Therefore $H_n(\bar{G}, \bar{P}; M) \cong H_n(\bar{G}; M)$ for all $n \geq 2$, and $H_n(G, P; M) \cong H_n(G; M)$ for all $n \geq 3$. Using this, for all $n \geq 3$, the long exact sequences in homology for the pairs (\bar{G}, \bar{P}) and (G, P) , together with Theorem 4.2, give isomorphisms

$$H_n(\bar{G}; M) \cong H_n(\bar{G}, \bar{P}; M) \cong H_n(G, P; M) \cong H_n(G; M),$$

and similarly for cohomology.

For the second statement, the long exact sequence in homology of the pair (G, P) includes:

$$H_2(P; M) \rightarrow H_2(G; M) \rightarrow H_2(G, P; M) \rightarrow H_1(P; M) \rightarrow H_1(G; M).$$

The first term vanishes because $cd_R(P) = 1$. Using $cd_R(\bar{P}) = 0$ and Theorem 4.2, the third term becomes $H_2(\bar{G}; M)$. Finally, the fourth term can be replaced by $H_1(N; M)_{\bar{P}}$ by Proposition 2.1. But $N \cong \mathbb{Z}$ acts trivially on M , so $H_1(N; M) = M$. \square

4.2 (Relatively) hyperbolic groups

When G is hyperbolic relative to P , group-theoretic Dehn fillings are especially well-behaved [Osi07]. The following was proved in [GMS19], and then in the more general context of groups with hyperbolically embedded subgroups [Sun20].

Theorem 4.4 ([GMS19, Sun20]). *Let G be hyperbolic relative to P . Then for every sufficiently deep normal subgroup $N < P$, the triple (G, P, N) is Cohen–Lyndon.*

Here a property is said to hold for *sufficiently deep* normal subgroups $N < P$ if there exists a finite subset $F \subset P \setminus \{1\}$ such that the property holds for all normal subgroups $N < P$ such that $N \cap F = \emptyset$. In this context, group-theoretic Dehn fillings retain many geometric properties.

Theorem 4.5 ([Osi07]). *Let G be hyperbolic relative to $\{P_\lambda\}_{\lambda \in \Lambda}$. Let $B \subset G$ be a finite subset. Then for every sufficiently deep normal subgroups $N_\lambda < P_\lambda$, it holds*

1. *The group $G/\langle\langle \cup_\lambda N_\lambda \rangle\rangle$ is hyperbolic relative to $\{P_\lambda/N_\lambda\}_{\lambda \in \Lambda}$;*
2. *The quotient $G \rightarrow G/\langle\langle \cup_\lambda N_\lambda \rangle\rangle$ induces an embedding $P_\lambda/N_\lambda \rightarrow G/\langle\langle \cup_\lambda N_\lambda \rangle\rangle$;*
3. *The quotient map $G \rightarrow G/\langle\langle \cup_\lambda N_\lambda \rangle\rangle$ is injective when restricted to B .*

We will mostly be concerned with a very particular example of a relatively hyperbolic pair. Let G be a hyperbolic group, and let $g \in G$ be an element of infinite order. Then g is contained in a unique maximal virtually cyclic subgroup denoted $E_G(g)$.

Theorem 4.6 ([Bow12, Theorem 7.11]). *Let G be a hyperbolic group, and let $g \in G$ be an element of infinite order. Then G is hyperbolic relative to $E_G(g)$.*

Putting these results together we obtain the following, which will serve as the inductive step in our constructions.

Proposition 4.7. *Let G be a non-elementary hyperbolic group with no non-trivial finite normal subgroups, and let $B \subset G$ be a finite subset. Let $g \in G$ be an element of infinite order and let $P := E_G(g)$. Then for every sufficiently deep finite-index normal subgroups $N < P$, denoting by $\bar{G} := G/\langle\langle N \rangle\rangle$ and $\bar{P} := P/N$:*

1. *The triple (G, P, N) is Cohen–Lyndon;*
2. *The group \bar{G} is non-elementary hyperbolic with no non-trivial finite normal subgroups;*
3. *The quotient map $G \rightarrow \bar{G}$ is injective when restricted to B , and induces an embedding $\bar{P} \hookrightarrow \bar{G}$.*

Proof. By Theorem 4.6, G is hyperbolic relative to $E_G(g)$. The Cohen–Lyndon property follows from Theorem 4.4; the hyperbolicity follows from Theorem 4.5, since a group that is hyperbolic relative to a finite subgroup is hyperbolic; the injectivity follows again from Theorem 4.5.

It remains to prove that \bar{G} is non-elementary and without finite normal subgroups. To this end, we find an additional element of infinite order $h \in G$ such that G is hyperbolic relative to $\{E_G(g), E_G(h)\}$ [Osi07, Lemmas 5.5 and 5.6], see also [Osi06a, Corollaries 1.7 and 4.5]. Consider the normal subgroups $N < E_G(g)$ and $\{1\} < E_G(h)$, so that the quotient $G \rightarrow \bar{G}$ fits into the framework of Theorem 4.5. In particular, \bar{G} is hyperbolic relative to the image of $E_G(h)$, which maps injectively and is almost malnormal in \bar{G} by [Osi06b, Theorem 1.4]. If \bar{G} had a non-trivial finite normal subgroup, then a sufficiently high power of h would centralise it, contradicting malnormality. Similarly, if \bar{G} were virtually cyclic, then a sufficiently high power of h would generate a normal subgroup of \bar{G} , contradicting malnormality again. \square

The situation when $E_G(g) = \langle g \rangle$ is especially convenient. Following [OT13], we call such elements *special*.

Proposition 4.8 ([OT13, Proposition 3.4]). *Let G be a hyperbolic group with no non-trivial finite normal subgroups, and let $x, h \in G$ with $h \neq 1$. Then the coset $x \langle\langle h \rangle\rangle$ contains a special element.*

In the cases above, to obtain desireable properties we needed to pass to a sufficiently deep normal subgroup of the elementary closure of an element. Let us highlight one situation in which this is not necessary.

Lemma 4.9. *Let G be a torsion-free non-elementary hyperbolic group, and let $B \subset G$ be a finite subset. Then there exists $1 \neq g \in G$ such that denoting by $\bar{G} := G/\langle\langle g \rangle\rangle$, the following hold:*

1. *The triple $(G, \langle g \rangle, \langle g \rangle)$ is Cohen–Lyndon;*
2. *The group \bar{G} is torsion-free non-elementary hyperbolic;*
3. *The quotient map $G \rightarrow \bar{G}$ is injective when restricted to B .*

Proof. This obtained in the course of the proof of [FFS25, Theorem 3.1], the second and third statement are consequences of the theorem, the first one is [FFS25, Lemma 3.5]. \square

4.3 Lacunary hyperbolic groups

A group is said to be *lacunary hyperbolic* if one of its asymptotic cones is an \mathbb{R} -tree. We will use the following useful characterisation.

Theorem 4.10 ([OOS09, Theorem 1.1]). *Let $G_1 \rightarrow G_2 \rightarrow \dots$ be a directed sequence of epimorphisms of hyperbolic groups, with colimit G_∞ . Let S_1 be a finite generating set for G_1 , and let S_i be its pushforward to G_i ; let δ_i be the hyperbolicity constant of $\text{Cay}(G_i, S_i)$. For each $i \in \mathbb{N}$, let ρ_i be the radius of the largest ball centred at the identity on which the quotient $G_i \rightarrow G_{i+1}$ is injective.*

If $\delta_i = o(\rho_i)$, then G_∞ is lacunary hyperbolic.

As we mentioned in the introduction, many earlier constructions of exotic values of von Neumann dimensions associated to groups involved lamplighters. Hence we record the following:

Proposition 4.11 ([OOS09]). *Let H be a non-trivial group, then $H \wr \mathbb{Z}$ cannot be a subgroup of a lacunary hyperbolic group.*

Proof. Up to passing to a subgroup, it suffices to show this for $H = \mathbb{Z}$ or $H = \mathbb{Z}/p$ for a prime p . When $H = \mathbb{Z}$, this follows from the more general fact that a lacunary hyperbolic group cannot contain a copy of \mathbb{Z}^2 by [OOS09, Corollary 3.21(a)]. For $H = \mathbb{Z}/p$, it follows from [OOS09, Corollary 3.21(c)] and its proof, see [OOS09, Remark 3.22]. \square

5 Failure of semi-continuity

In this section we prove Theorem C. We start with a more precise statement.

Theorem 5.1. *Let G be a torsion-free cocompact lattice in the isometry group of the octonionic hyperbolic plane, and let $B \subset G$ be a finite subset. Then for every $n \in \mathbb{N}$, there exists a torsion-free hyperbolic quotient $G \rightarrow G_n$, injective on B , of cohomological dimension 16, such that for every unitary G_n -representation V , the second homology is $H_2(G_n; V) \cong V^n$.*

Cocompact lattices in the isometry group of the octonionic hyperbolic plane exist [BHC62], and they can be chosen torsion-free up to passing to a finite index subgroup [Sel60].

Proof. The group G is torsion-free hyperbolic of cohomological dimension 16, and by Theorem 3.10 it has property $[T_2]$. We construct G_n by induction on n , starting with $G_0 = G$, which works by Proposition 3.8.

Suppose by induction that G_{n-1} has been constructed. Lemma 4.9 gives an element $g \in G_{n-1}$ and we define G_n to be the corresponding group-theoretic Dehn filling. Lemma 4.9 gives injectivity on B and the fact that G_n is torsion-free hyperbolic. The first part of Corollary 4.3 implies that the cohomological dimension of G_n is 16. Moreover, since G_{n-1} has property (T) , applying Proposition 3.8 and the second part of Corollary 4.3, we obtain, for every unitary G_n -representation V , a short exact sequence

$$0 \rightarrow H_2(G_{n-1}; V) \rightarrow H_2(G_n; V) \rightarrow V \rightarrow 0.$$

By the induction hypothesis the first term is V^{n-1} . Therefore $H_2(G_n; V) \cong V^n$, which proves the claim and hence the theorem. \square

Remark 5.2. The groups G_n of course fail to have property (T_2) , despite being quotients of the (T_2) group G , as remarked in [FF25]. Still, the second homology (and similarly the second cohomology) functor enjoys some strong form of rigidity: in a way, its output can be read directly from its input. It would be interesting to explore whether a (co)homological rigidity of this form characterises quotients of (T_2) groups.

Proof of Theorem C. We have $b_2^{(2)}(G) = 0$ by Corollary 3.9. For each $i \in \mathbb{N}$, we let B_i be the ball of radius i in G , for a fixed choice of finite generating set, and $n_i \in \mathbb{N}$ as in the statement of Theorem C. Then Theorem 5.1 builds a group G_i , which by Proposition 3.6 has

$$b_2^{(2)}(G_i) = \dim_{\mathcal{N}(G_i)} H_2(G_i; \ell^2(G_i)) = \dim_{\mathcal{N}(G_i)} \ell^2(G_i)^{n_i} = n_i.$$

Since $G \rightarrow G_i$ is injective on larger and larger balls, $G_i \xrightarrow{i \rightarrow \infty} G$ in the space of marked groups. The groups G_i have property (T) being quotients of G . The other properties of G_i are all contained in the statement of Theorem 5.1. \square

Remark 5.3. In Theorem 5.1, we used the full power of property $[T_2]$, but the proof shows that if V is assumed to have no non-zero invariant vectors, then property (T_2) suffices. In the proof of Theorem C, we applied Theorem 5.1 only with $V = \ell^2(G_i)$, which has no non-zero invariant vectors, so property (T_2) would have been sufficient. Similarly, the computation of second ℓ^2 -Betti numbers in the next section could be obtained only using property (T_2) .

6 Proof of the main theorem

In this section we prove Theorem A. Because the proof has many moving parts, we start with a simpler version, that only involves one prime, and does not give simplicity.

Theorem 6.1. *Let p be a prime. There exists an uncountable family of groups $(G_\alpha)_{\alpha \in A}$ and an integer $\chi \geq 1$ with the following properties.*

1. G_α has property (T) and is lacunary hyperbolic.
2. G_α has cohomological dimension 16 modulo p .
3. The second ℓ^2 -Betti numbers $b_2^{(2)}(G_\alpha)$ are pairwise distinct.
4. The ℓ^2 -Euler characteristic is $\chi^{(2)}(G_\alpha) = b_2^{(2)}(G_\alpha) + \chi$.

Proof. We choose as an index set $A := \{2, 3\}^{\mathbb{N}}$, and fix an element $\alpha \in A$. The group G_α will arise as the colimit of a direct sequence of epimorphisms of hyperbolic groups $G_1 \rightarrow G_2 \rightarrow \dots$, where G_1 is a torsion-free cocompact lattice in the isometry group of the octonionic hyperbolic plane, which has Euler characteristic $\chi \geq 1$ [Lüc02, Corollary 5.16(1) on p. 231]. We fix a finite generating set S_1 for G_1 and let S_i be the pushforward to G_i . We also specify elements $g_i \in G_i$, and will prove the following additional properties.

1. G_i is non-elementary hyperbolic with no non-trivial finite normal subgroups.
2. $\text{Cay}(G_i, S_i)$ is δ_i -hyperbolic, and the quotient $G_i \rightarrow G_{i+1}$ is injective on the ball of radius $i\delta_i$ centred at the identity.

3. Each quotient $G_i \rightarrow G_{i+1}$ induces isomorphisms in homology and cohomology with coefficients in $R[G_{i+1}]$ -modules and degree at least 3, where R is any ring where p is invertible.
4. The element $g_i \in G_i$ has infinite order, and for all $j > i$ its image in G_j has order p^{a_i} , where $a_i \in \mathbb{N}$ is a sequence with the property that $a_{i+1} - a_i - 1$ is a positive power of $\alpha(i) \in \{2, 3\}$.
5. For every unitary G_i -representation V , the second homology is

$$H_2(G_i; V) = \bigoplus_{j < i} V_{g_j}.$$

Let us first prove that the colimit G_α has the desired properties. The group G_1 has property (T) by Theorem 3.10, hence so does G_α , being its quotient. Item 2 implies that G_α is lacunary hyperbolic, by Theorem 4.10. Now G_1 is the fundamental group of a closed 16-dimensional aspherical manifold, so it has cohomological dimension 16, and $H^{16}(G_1; R) \cong R$ for every ring R . Therefore Item 3 and Corollary 2.4 imply that G_α has cohomological dimension 16 modulo p .

Next, we compute $b_2^{(2)}(G_\alpha)$. By Proposition 2.2, we have

$$H_2(G_\alpha; \ell^2(G_\alpha)) \cong \bigoplus_{i \geq 1} \ell^2(G_\alpha)_{g_i}.$$

Each g_i has order p^{a_i} in G_α , hence by Proposition 3.6 and Proposition 3.11:

$$\begin{aligned} \dim_{\mathcal{N}(G_\alpha)} \ell^2(G_\alpha)_{g_i} &= \dim_{\mathcal{N}(G_\alpha)} H_0(\langle g_i \rangle, \ell^2(G_\alpha)) = \dim_{\mathcal{N}(G_\alpha)} H_0(\langle g_i \rangle, \mathcal{N}(G_\alpha)) \\ &= \dim_{\mathcal{N}(\langle g_i \rangle)} H_0(\langle g_i \rangle, \mathcal{N}(\langle g_i \rangle)) = b_0^{(2)}(\langle g_i \rangle) = p^{-a_i}. \end{aligned}$$

By Proposition 3.6 and Lemma 3.2, we therefore have

$$b_2^{(2)}(G_\alpha) = \dim_{\mathcal{N}(G_\alpha)} H_2(G_\alpha; \ell^2(G_\alpha)) = \sum_{i \geq 1} \frac{1}{p^{a_i}}.$$

Reading this number in base p , we see a 1 at position a_i , then a string of 0s of length $(a_{i+1} - a_i - 1)$ before the next 1 at position a_{i+1} . The length of this string of 0s is either a power of 2 or a power of 3 depending on whether $\alpha(i) = 2$ or 3. This shows that we can read α from $b_2^{(2)}(G_\alpha)$.

Finally, we compute $\chi^{(2)}(G_\alpha)$. By Item 3 and Proposition 2.2, for all $n \geq 3$ we have

$$b_n^{(2)}(G_\alpha) = \dim_{\mathcal{N}(G_\alpha)} H_n(G_\alpha; \mathcal{N}(G_\alpha)) = \dim_{\mathcal{N}(G_\alpha)} H_n(G_1; \mathcal{N}(G_\alpha)) = b_n^{(2)}(G_1 \rightarrow G_\alpha).$$

Since G_α has property (T), $b_1^{(2)}(G_\alpha) = 0$ by [BV97, Corollary 6]. Hence

$$\begin{aligned} \chi^{(2)}(G_\alpha) &= \sum_{n \in \mathbb{N}} (-1)^n b_n^{(2)}(G_\alpha) = b_2^{(2)}(G_\alpha) + \sum_{n \geq 3} (-1)^n b_n^{(2)}(G_\alpha) \\ &= b_2^{(2)}(G_\alpha) + \sum_{n \geq 3} (-1)^n b_n^{(2)}(G_1 \rightarrow G_\alpha) = b_2^{(2)}(G_\alpha) + \chi, \end{aligned}$$

where the last equality follows from Corollary 3.9.

It remains to construct the directed sequence $G_1 \rightarrow G_2 \rightarrow \dots$ and the elements $g_i \in G_i$ so that the items above are satisfied. Starting with G_1 , note that Item 5 holds by Proposition 3.8, because G_1 has property $[T_2]$, and there is nothing to check for the other items, before g_1 has been chosen. So suppose by induction that G_1, \dots, G_i and the elements g_1, \dots, g_{i-1} have been constructed so that all items above are satisfied. Using Proposition 4.8, we choose a special element $g_i \in G_i$. For an integer $k \in \mathbb{N}$ to be specified later, we define $G_{i+1} := G_i / \langle\langle g_i^k \rangle\rangle$. This is the group-theoretic Dehn filling arising from the triple $(G_i, \langle g_i \rangle, \langle g_i^k \rangle)$. By Proposition 4.7, if k is large enough, G_{i+1} is non-elementary hyperbolic with no non-trivial finite normal subgroups, giving Item 1, and we can choose the set B to be such that Item 2 is satisfied, and the elements g_1, \dots, g_{i-1} have the same order in G_{i+1} , giving part of Item 4. Choosing $k = p^{a_i}$, for a sufficiently large a_i such that $a_i - a_{i-1} - 1$ is a positive power of $\alpha(i-1)$ gives the rest of Item 4, thanks to the last statement of Proposition 4.7.

By Proposition 4.7 again, the triple $(G_i, \langle g_i \rangle, \langle g_i^k \rangle)$ is Cohen–Lyndon. Because k is a power of p , for every ring R in which p is invertible, we can apply Corollary 4.3. The first part immediately gives Item 3. Moreover, the second part, together with the fact that G_i has property (T) , gives a short exact sequence

$$0 \rightarrow H_2(G_i; V) \rightarrow H_2(G_{i+1}; V) \rightarrow V_{g_i} \rightarrow 0.$$

The induction hypothesis gives

$$H_2(G_{i+1}; V) = \bigoplus_{j \leq i} V_{g_j},$$

proving Item 5. This concludes the construction, hence the proof. \square

We now prove Theorem A. The proof will be similar, but more technical, in that we need to leverage an additional prime, and the full power of Proposition 4.8, to make the group G_α simple. This improvement employs a trick from [OT13], where it was used to construct simple groups with large first ℓ^2 -Betti numbers. The following will be our starting point.

Lemma 6.2. *Let G be a torsion-free cocompact lattice in the isometry group of the octonionic hyperbolic plane. Let q be a prime. Then there exists a quotient $G \rightarrow G_1$ with the following properties.*

1. G_1 is non-elementary hyperbolic with no non-trivial finite normal subgroups.
2. The quotient $G \rightarrow G_1$ induces isomorphisms in homology and cohomology with coefficients in $R[G_1]$ -modules and degree at least 3, where R is any ring where q is invertible.
3. G_1 is generated by elements $\{x_1, \dots, x_m\}$, each of which has order a power of q .
4. There is a finite set \mathcal{E} of finite subgroups of G_1 , such that for every unitary G_1 -representation V , the second homology is

$$H_2(G_1; V) = \bigoplus_{E \in \mathcal{E}} V_E.$$

Proof. We start by choosing a generating set $\{x_1^0, \dots, x_m^0\}$ of G . For $i = 0, \dots, m$, we construct by induction a quotient $G \rightarrow G_{i/m}$ with the following properties.

1. $G_{i/m}$ is non-elementary hyperbolic with no non-trivial finite normal subgroups.
2. The quotient $G \rightarrow G_{i/m}$ induces isomorphisms in homology and cohomology with coefficients in $R[G_{i/m}]$ -modules and degree at least 3, where R is any ring where q is invertible. In particular $G_{i/m}$ has cohomological dimension 16 modulo q , and so it only contains q -torsion [Bie81, Proposition 4.11 on p. 63].
3. $G_{i/m}$ is generated by elements $\{x_1^i, \dots, x_m^i\}$, where x_1^i, \dots, x_i^i have order a power of q .
4. There is a finite set of finite subgroups \mathcal{E}_i of $G_{i/m}$, such that for every unitary $G_{i/m}$ -representation V , the second homology is

$$H_2(G_{i/m}; V) = \bigoplus_{E \in \mathcal{E}_i} V_E.$$

Then the final group $G_{m/m}$ will be the group G_1 in the statement of the lemma, with $\mathcal{E} = \mathcal{E}_m$ and $x_j = x_j^m$. The base case for $i = 0$ is G itself, with $\mathcal{E} = \emptyset$, where we do not need to check anything for the first three items, and Item 4 follows from Proposition 3.8 and Theorem 3.10.

Suppose by induction that $G_{i/m}$ has been constructed. If x_{i+1}^i is torsion, then it is q -torsion, so we set $G_{(i+1)/m} = G_{i/m}$, $x_j^{i+1} = x_j^i$ for all j , and $\mathcal{E}_{i+1} = \mathcal{E}_i$. Otherwise, consider the infinite virtually cyclic group $E_{G_{i/m}}(x_{i+1})$. This has a unique maximal finite normal subgroup of order a power of q , and the quotient is isomorphic to \mathbb{Z} (or possibly to D_∞ , in case $q = 2$). We replace x_{i+1}^i with a root x_{i+1}^{i+1} that generates the \mathbb{Z} quotient (or the index-2 subgroup of the D_∞ quotient), so that $[E_{G_{i/m}}(x_{i+1}^{i+1}) : \langle x_{i+1}^{i+1} \rangle]$ is a power of q . We now let $G_{(i+1)/m}$ be the group-theoretic Dehn filling for the triple $(G_{i/m}, E_{G_{i/m}}(x_{i+1}^{i+1}), \langle (x_{i+1}^{i+1})^k \rangle)$, where k is a sufficiently large power of q . By Proposition 4.7, we see that $G_{(i+1)/m}$ satisfies Items 1 and 3, choosing $x_j^{i+1} = x_j^i$ for all $j \neq i+1$. We also choose k to be large enough so that the orders of x_1^i, \dots, x_i^i and their images $x_1^{i+1}, \dots, x_i^{i+1}$ have the same order, and each finite subgroup from \mathcal{E}_i embeds into $G_{(i+1)/m}$. Moreover, by Proposition 4.7 again, this triple is Cohen–Lyndon, so Corollary 4.3 applies, and proves Item 2. Finally, let $E := E_{G_{i/m}}(x_{i+1}^{i+1})/\langle (x_{i+1}^{i+1})^k \rangle$. Corollary 4.3 and property (T) for $G_{i/m}$ give a short exact sequence

$$0 \rightarrow H_2(G_{i/m}; V) \rightarrow H_2(G_{(i+1)/m}; V) \rightarrow V_E \rightarrow 0,$$

so setting $\mathcal{E}_{i+1} = \mathcal{E}_i \cup \{E\}$, we obtain Item 4 and conclude. \square

Proof of Theorem A. We proceed as in the proof of Theorem 6.1, using the prime $p \neq q$, but starting with the group G_1 from Lemma 6.2. The result will be a group G_α with the same properties as in Theorem 6.1, with the exception that cohomological dimension now has to be taken modulo $\{p, q\}$, and $b_2^{(2)}(G_\alpha)$ is computed starting from

$$b_2^{(2)}(G_1 \rightarrow G_\alpha) = \dim_{\mathcal{N}(G_\alpha)} \left(\bigoplus_{E \in \mathcal{E}} \mathcal{N}(G_\alpha)_E \right) = \sum_{E \in \mathcal{E}} |E|^{-1}.$$

Since this starting value is independent of α , the same computations go through. For the finiteness of the ℓ^2 -Betti numbers of G_α we refer to Remark 6.3.

In each induction step in the proof of Theorem 6.1, we chose arbitrary special elements $g_i \in G_i$. We will now explain how to use the full power of Proposition 4.8 in the construction so that this time the colimit is simple. Enumerate the set $\mathcal{X} := \{x_1, \dots, x_m\} \times G_1$. When the group G_i has been built, let (x, h) denote the i -th element of \mathcal{X} . If h is trivial, then we pick g_i

to be an arbitrary special element, and proceed as before. Otherwise, we use Proposition 4.8 to pick $g_i \in x^{-1} \langle\langle h \rangle\rangle_{G_i}$. We claim that now $x \in \langle\langle h \rangle\rangle_{G_{i+1}} \leq G_{i+1}$. Indeed, consider the image of x in the quotient $G_{i+1}/\langle\langle h \rangle\rangle_{G_{i+1}}$. On the one hand, it has order a power of q , because x has order a power of q in G_1 . On the other hand, it equals the image of g_i , which has order a power of p in G_{i+1} , by construction of the quotient $G_i \rightarrow G_{i+1}$. Because $p \neq q$, we see that x must be trivial in $G_{i+1}/\langle\langle h \rangle\rangle_{G_{i+1}}$, which proves our claim.

As a result, for every $(x, h) \in \mathcal{X}$, either h is trivial in G_α , or x lies in the normal closure of h in G_α . Because x ranges over a generating set for G_1 and h ranges over G_1 , this shows that every non-trivial element of G_α normally generates, that is, G_α is simple. \square

Remark 6.3. Note that the constructions give some information about higher ℓ^2 -Betti numbers as well. Indeed, G_α is obtained from a torsion-free cocompact lattice in the isometry group of the octonionic hyperbolic plane, denoted G , by a sequence of group-theoretic Dehn fillings satisfying Corollary 4.3. Using this and Proposition 2.2 we have

$$b_n^{(2)}(G_\alpha) = b_n^{(2)}(G \rightarrow G_\alpha)$$

for all $n \geq 3$. Since G is of type F , this implies that all ℓ^2 -Betti numbers of G_α are finite. We know from Theorem 3.10 that G has property (T_3) , so Corollary 3.9 gives $b_3^{(2)}(G_\alpha) = 0$.

Moreover, G satisfies the Singer conjecture [Lück02, Corollary 5.16 on p. 231], that is $b_n^{(2)}(G) = 0$ for all $n \neq 8$ and $b_8^{(2)}(G) = \chi \geq 1$. This does not give information about $b_n^{(2)}(G \rightarrow G_\alpha)$, however, assuming Lück's approximation conjecture [Lück02, Chapter 13], and repeating the construction ensuring that $G \rightarrow G_\alpha$ is injective on a large ball centred at the identity, we would have that $b_n^{(2)}(G \rightarrow G_\alpha)$ is close to $b_n^{(2)}(G)$. It would then follow that $b_n^{(2)}(G_\alpha)$ is close to 0, for $n \geq 4, n \neq 8$, and close to χ for $n = 8$.

Remark 6.4. Taking $V = \mathbb{C}$ to be the trivial unitary representation, the proofs of Theorems 6.1 and A, and Proposition 2.2, show that $H_2(G_\alpha; \mathbb{C})$ is infinite-dimensional. It follows that the groups G_α do not have type $FP_2(\mathbb{Q})$.

7 An elementary construction

We conclude with the elementary construction giving Theorem D. We start with two basic product formulas.

Proposition 7.1 ([Lück02, Theorem 6.54(5) on p. 266]). *Let G_1, G_2 be two groups. Then*

$$b_n^{(2)}(G_1 \times G_2) = \sum_{i+j=n} b_i^{(2)}(G_1) b_j^{(2)}(G_2).$$

Proposition 7.2. *Let $(G_i)_{i \in \mathbb{N}}$ be infinite groups, and let G be their free product. Then $b_1^{(2)}(G) = \infty$ and*

$$b_n^{(2)}(G) = \sum_{i \in \mathbb{N}} b_n^{(2)}(G_i)$$

for all $n \geq 2$.

This is usually stated for finite free products only [Lück02, Theorem 1.35(5) on p. 38] so we include a proof.

Proof. Let P_i denote the free product $G_1 * \cdots * G_i$, so $P_{i+1} = P_i * G_{i+1}$. By Mayer–Vietoris, for all $n \geq 2$ we have an isomorphism

$$H_n(P_{i+1}; \mathcal{N}(G)) \cong H_n(P_i; \mathcal{N}(G)) \oplus H_n(G_{i+1}; \mathcal{N}(G)),$$

which shows by induction that

$$H_n(P_i; \mathcal{N}(G)) = \bigoplus_{j \leq i} H_n(G_j; \mathcal{N}(G)).$$

By Mayer–Vietoris again, we have an exact sequence

$$0 \rightarrow H_1(P_{i+1}; \mathcal{N}(G)) \rightarrow H_1(P_i; \mathcal{N}(G)) \oplus H_1(G_{i+1}; \mathcal{N}(G)) \rightarrow \mathcal{N}(G) \rightarrow \mathcal{N}(G)_{P_{i+1}}.$$

By Proposition 3.11, the last term has dimension

$$\dim_{\mathcal{N}(G)} H_0(P_{i+1}; \mathcal{N}(G)) = \dim_{\mathcal{N}(P_{i+1})} H_0(P_{i+1}; \mathcal{N}(P_{i+1})) = b_0^{(2)}(P_{i+1}) = 0.$$

So by induction

$$H_1(P_i; \mathcal{N}(G)) = \left(\bigoplus_{j \leq i} H_1(G_j; \mathcal{N}(G)) \right) \oplus V_i,$$

where V_i is a submodule of $\mathcal{N}(G)^{i-1}$ of codimension 0. We conclude from Propositions 2.2 and 3.11, together with the fact that $\dim_{\mathcal{N}(G)}$ respects directed unions [Lü98, Theorem 0.6(b)]. \square

We can now begin the construction.

Proposition 7.3. *Let p be a prime. There exists a family of groups $(C_x)_{x \in \mathbb{R}_{>0}}$ with the following properties.*

1. *Each C_x is countable and residually finite.*
2. *$b_1^{(2)}(C_x) = \infty, b_2^{(2)}(C_x) = x$.*
3. *Each C_x has cohomological dimension 2 modulo p . In particular, $b_n^{(2)}(C_x) = 0$ for all $n \geq 3$.*

Proof. Write x in base- p as

$$x = \sum_{i \in \mathbb{N}} \frac{r_i}{p^i},$$

where $r_0 \in \mathbb{N}$; $r_i \in \{0, 1, \dots, p-1\}$ for $i \geq 1$. For elements in $\mathbb{Z} \left[\frac{1}{p} \right]$ we pick the periodic representation, which allows to have infinite sums for every x . Now for each i such that $r_i \neq 0$ we let $A_i := \mathbb{Z}/p^i \mathbb{Z} \times F_{r_{i+1}}^2$, where F_r is the free group of rank r . Then

$$b_0^{(2)}(A_i) = b_1^{(2)}(A_i) = 0; b_2^{(2)}(A_i) = \frac{r_i}{p^i},$$

by Proposition 7.1. Moreover, A_i has cohomological dimension 2 modulo p .

Now let

$$C_x := \ast_{i \geq 0} A_i.$$

By Proposition 7.2, we have the desired values of ℓ^2 -Betti numbers. Moreover, C_x has cohomological dimension 2 modulo p , by Corollary 2.4. Finally, C_x is residually finite, being a free product of residually finite groups. \square

Proof of Theorem D. Let C_x be the groups from Proposition 7.3. We embed each C_x into a finitely generated residually finite group R_x following [WZ96]. The group R_x is an HNN extension of $F_2 \times C_x$ over a free subgroup, therefore it has cohomological dimension 3 modulo p . Moreover, C_x is separable, that is closed in the profinite topology [WZ96, Lemma 4].

Consider the group $R_x \times F_2^3$: it has cohomological dimension 6 modulo p , and contains C_x as a separable subgroup. Moreover, $b_n^{(2)}(R_x \times F_2^3) = 0$ for $n \leq 3$, by Proposition 7.1. Define

$$G_x := (R_x \times F_2^3) *_{C_x} (R_x \times F_2^3),$$

where the double is over the copy of C_x coming from the first coordinate of each factor. Because C_x is separable, G_x is residually finite, and moreover it has cohomological dimension 6 modulo p . By Mayer–Vietoris:

$$H_n((R_x \times F_2^3); \mathcal{N}(G_x))^2 \rightarrow H_n(G_x; \mathcal{N}(G_x)) \rightarrow H_{n-1}(C_x; \mathcal{N}(G_x)) \rightarrow H_{n-1}(R_x \times F_2^3); \mathcal{N}(G_x))^2.$$

Taking \dim_{G_x} and $n \leq 3$, the first and last term give $2b_n^{(2)}(R_x \times F_2^3) = 2b_{n-1}^{(2)}(R_x \times F_2^3) = 0$, by Proposition 3.11, so

$$b_n^{(2)}(G_x) = b_{n-1}^{(2)}(C_x).$$

The value of ℓ^2 -Betti numbers are as desired, by Proposition 7.3. \square

Remark 7.4. The construction is quite flexible, and can be made more elementary, or less elementary, depending on what properties one wants to impose, if we forgo residual finiteness.

1. We can choose R_x to be a finitely generated group that is an HNN extension of $F_2 * C_x$ over a free subgroup, therefore has cohomological dimension 2 modulo p [HNN49]. This is the first and easiest embedding result for countable groups into finitely generated groups. Then $R_x \times F_2^3$ has cohomological dimension 5 modulo p , hence so does G_x .
2. We can bring the dimension down by one more, up to possibly adding some primes in the torsion. Let L be a generalised triangle group that is hyperbolic and has property (T) [CCKW22]. Then there exists a finite set of primes π (the ones appearing in the orders of the finite vertex groups) such that L has cohomological dimension 2 modulo π . Using [FFS25, Theorem 4.1], we can embed C_x into a group R_x that is a quotient of L and has cohomological dimension 2 modulo $\pi \cup \{p\}$. In particular R_x has property (T) and so $b_1^{(2)}(R_x) = 0$ by [BV97, Corollary 6]. So in the construction above it suffices to take $R_x \times F_2^3$ to get the correct vanishing of ℓ^2 -Betti numbers, and thus the resulting G_x has cohomological dimension 4 modulo $\pi \cup \{p\}$.
3. If we forgo the finiteness of the dimension, then we can gain control on all other ℓ^2 -Betti numbers. We first choose R'_x to be any finitely generated group containing C_x , and then let $R_x := R'_x \wr \mathbb{Z}$, which has vanishing ℓ^2 -Betti numbers in all degrees [WWZZ25, Remark 7.6]. So in the construction above we do not need to take products with F_2 , and directly get $b_n^{(2)}(G_x) = b_{n-1}^{(2)}(C_x)$ for all $n \in \mathbb{N}$. Therefore $b_n^{(2)}(G_x) = 0$ for all $n \geq 4$.

Corollary 7.5. Let $(G_x)_{x \in \mathbb{R}_{>0}}$ be as in Theorem D. Then the groups $G_x * \mathbb{Z}$ are pairwise non-measure equivalent.

Proof. If $G_x * \mathbb{Z}$ and $G_y * \mathbb{Z}$ are measure equivalent with index $c > 0$, then $b_n^{(2)}(G_x * \mathbb{Z}) = cb_n^{(2)}(G_y * \mathbb{Z})$ for all $n \in \mathbb{N}$ by Gaboriau's theorem [Gab02]. So $1 = b_1^{(2)}(G_x * \mathbb{Z}) = cb_1^{(2)}(G_y * \mathbb{Z}) = c$, that is $c = 1$, and so $x = b_3^{(2)}(G_x * \mathbb{Z}) = cb_3^{(2)}(G_y * \mathbb{Z}) = cy = y$. \square

References

- [Ati76] M. F. Atiyah, *Elliptic operators, discrete groups and von Neumann algebras*, Colloque “Analyse et Topologie” en l’Honneur de Henri Cartan (Orsay, 1974), Astérisque, No. 32-33, Soc. Math. France, Paris, 1976, pp. 43–72. MR 420729
- [Aus13] T. Austin, *Rational group ring elements with kernels having irrational dimension*, Proc. Lond. Math. Soc. (3) **107** (2013), no. 6, 1424–1448. MR 3149852
- [BdlHV08] B. Bekka, P. de la Harpe, and A. Valette, *Kazhdan’s property (T)*, New Mathematical Monographs, vol. 11, Cambridge University Press, Cambridge, 2008. MR 2415834
- [BH99] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 319, Springer-Verlag, Berlin, 1999. MR 1744486
- [BHC62] A. Borel and Harish-Chandra, *Arithmetic subgroups of algebraic groups*, Ann. of Math. (2) **75** (1962), 485–535. MR 147566
- [Bie81] R. Bieri, *Homological dimension of discrete groups*, second ed., Queen Mary College Mathematics Notes, Queen Mary College, Department of Pure Mathematics, London, 1981. MR 715779
- [Bow12] B. H. Bowditch, *Relatively hyperbolic groups*, Internat. J. Algebra Comput. **22** (2012), no. 3, 1250016, 66. MR 2922380
- [Bro94] K. S. Brown, *Cohomology of groups*, Graduate Texts in Mathematics, vol. 87, Springer-Verlag, New York, 1994, Corrected reprint of the 1982 original. MR 1324339
- [BS23] U. Bader and R. Sauer, *Higher Kazhdan property and unitary cohomology of arithmetic groups*, arXiv preprint arXiv:2308.06517, 2023.
- [BS25] ———, *Higher property T and below-rank phenomena of lattices*, arXiv preprint arXiv:2511.20192, 2025.
- [BV97] M. E. B. Bekka and A. Valette, *Group cohomology, harmonic functions and the first L^2 -Betti number*, Potential Anal. **6** (1997), no. 4, 313–326. MR 1452785
- [CCKW22] P.-E. Caprace, M. Conder, M. Kaluba, and S. Witzel, *Hyperbolic generalized triangle groups, property (T) and finite simple quotients*, J. Lond. Math. Soc. (2) **106** (2022), no. 4, 3577–3637. MR 4524205
- [CIO23] I. Chifan, A. Ioana, D. Osin, and B. Sun, *Wreath-like products of groups and their von Neumann algebras I: W^* -superrigidity*, Ann. of Math. (2) **198** (2023), no. 3, 1261–1303. MR 4660139
- [CL63] D. E. Cohen and R. C. Lyndon, *Free bases for normal subgroups of free groups*, Trans. Amer. Math. Soc. **108** (1963), 526–537. MR 170930
- [FF25] F. Fournier-Facio, *Stability, approximable quotients, and higher property (T)*, arXiv preprint arXiv:2512.09180, 2025.
- [FFS25] F. Fournier-Facio and B. Sun, *Dimensions of finitely generated simple groups and their subgroups*, arXiv preprint arXiv:2503.01987, 2025.
- [Gab00] D. Gaboriau, *Coût des relations d’équivalence et des groupes*, Invent. Math. **139** (2000), no. 1, 41–98. MR 1728876
- [Gab02] ———, *Invariants l^2 de relations d’équivalence et de groupes*, Publ. Math. Inst. Hautes Études Sci. (2002), no. 95, 93–150. MR 1953191
- [GMS19] D. Groves, J. F. Manning, and A. Sisto, *Boundaries of Dehn fillings*, Geom. Topol. **23** (2019), no. 6, 2929–3002. MR 4039183
- [Gra14] Ł. Grabowski, *On Turing dynamical systems and the Atiyah problem*, Invent. Math. **198** (2014), no. 1, 27–69. MR 3260857
- [Gra16] ———, *Irrational l^2 invariants arising from the lamplighter group*, Groups Geom. Dyn. **10** (2016), no. 2, 795–817. MR 3513118

[Hat02] A. Hatcher, *Algebraic topology*, Cambridge University Press, Cambridge, 2002. MR 1867354

[HNN49] G. Higman, B. H. Neumann, and N. Neumann, *Embedding theorems for groups*, J. London Math. Soc. **24** (1949), 247–254. MR 32641

[ITD25] A. Ioana and R. Tucker-Drob, *A continuum of non-measure equivalent groups*, arXiv preprint arXiv:2512.04531, 2025.

[KPV15] D. Kyed, H. D. Petersen, and S. Vaes, *L^2 -Betti numbers of locally compact groups and their cross section equivalence relations*, Trans. Amer. Math. Soc. **367** (2015), no. 7, 4917–4956. MR 3335405

[Kum80] S. Kumaresan, *On the canonical k -types in the irreducible unitary g -modules with nonzero relative cohomology*, Invent. Math. **59** (1980), no. 1, 1–11. MR 575078

[LN23] A. López Neumann, *Finitely presented simple groups and measure equivalence*, Colloq. Math. **172** (2023), no. 2, 261–279. MR 4589106

[Lück98] W. Lück, *Dimension theory of arbitrary modules over finite von Neumann algebras and L^2 -Betti numbers. I. Foundations*, J. Reine Angew. Math. **495** (1998), 135–162. MR 1603853

[Lück02] ———, *L^2 -invariants: theory and applications to geometry and K -theory*, *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*, vol. 44, Springer-Verlag, Berlin, 2002. MR 1926649

[Mat62] Y. Matsushima, *On Betti numbers of compact, locally symmetric Riemannian manifolds*, Osaka Math. J. **14** (1962), 1–20. MR 141138

[OOS09] A. Yu. Olshanskii, D. Osin, and M. V. Sapir, *Lacunary hyperbolic groups*, Geom. Topol. **13** (2009), no. 4, 2051–2140, With an appendix by M. Kapovich and B. Kleiner. MR 2507115

[Osi06a] D. Osin, *Elementary subgroups of relatively hyperbolic groups and bounded generation*, Internat. J. Algebra Comput. **16** (2006), no. 1, 99–118. MR 2217644

[Osi06b] ———, *Relatively hyperbolic groups: intrinsic geometry, algebraic properties, and algorithmic problems*, Mem. Amer. Math. Soc. **179** (2006), no. 843, vi+100. MR 2182268

[Osi07] ———, *Peripheral fillings of relatively hyperbolic groups*, Invent. Math. **167** (2007), no. 2, 295–326. MR 2270456

[OT13] D. Osin and A. Thom, *Normal generation and ℓ^2 -Betti numbers of groups*, Math. Ann. **355** (2013), no. 4, 1331–1347. MR 3037017

[Pic06] M. Pichot, *Semi-continuity of the first ℓ^2 -Betti number on the space of finitely generated groups*, Comment. Math. Helv. **81** (2006), no. 3, 643–652. MR 2250857

[PS24a] N. Petrosyan and B. Sun, *Cohomology of group theoretic Dehn fillings II*, Adv. Math. **437** (2024), Paper No. 109412, 56. MR 4669335

[PS24b] ———, *L^2 -Betti numbers of Dehn fillings*, arXiv preprint arXiv:2412.16090, 2024.

[PT11] J. Peterson and A. Thom, *Group cocycles and the ring of affiliated operators*, Invent. Math. **185** (2011), no. 3, 561–592. MR 2827095

[Sau05] R. Sauer, *L^2 -Betti numbers of discrete measured groupoids*, Internat. J. Algebra Comput. **15** (2005), no. 5–6, 1169–1188. MR 2197826

[Sel60] A. Selberg, *On discontinuous groups in higher-dimensional symmetric spaces*, Contributions to function theory (Internat. Colloq. Function Theory, Bombay, 1960), Tata Inst. Fund. Res., Bombay, 1960, pp. 147–164. MR 130324

[Sun20] B. Sun, *Cohomology of group theoretic Dehn fillings I: Cohen-Lyndon type theorems*, J. Algebra **542** (2020), 277–307. MR 4019787

[Wei94] C. A. Weibel, *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics, vol. 38, Cambridge University Press, Cambridge, 1994. MR 1269324

[WWZZ25] F. Wu, X. Wu, M. Zhao, and Z. Zhou, *Embedding groups into boundedly acyclic groups*, J. Lond. Math. Soc. (2) **111** (2025), no. 5, Paper No. e70164, 39. MR 4899448

[WZ96] J. S. Wilson and P. A. Zalesskii, *An embedding theorem for certain residually finite groups*, Arch. Math. (Basel) **67** (1996), no. 3, 177–182. MR 1402516

DEPARTMENT OF PURE MATHEMATICS AND MATHEMATICAL STATISTICS, UNIVERSITY OF CAMBRIDGE, UK

E-mail address: ff373@cam.ac.uk

FACULTY OF MATHEMATICS, KARLSRUHE INSTITUTE OF TECHNOLOGY, GERMANY

E-mail address: roman.sauer@kit.edu