

**CONSTRUCTION OF SIMPLE QUOTIENTS OF BERNSTEIN-ZELEVINSKY
DERIVATIVES AND HIGHEST DERIVATIVE MULTISEGMENTS III:
PROPERTIES OF MINIMAL SEQUENCES**

KEI YUEN CHAN

ABSTRACT. Let F be a non-Archimedean local field. For an irreducible smooth representation π of $\mathrm{GL}_n(F)$ and a multisegment \mathfrak{m} , one associates a simple quotient $D_{\mathfrak{m}}(\pi)$ of a Bernstein-Zelevinsky derivative of π . In the preceding article, we showed that

$$\mathcal{S}(\pi, \tau) := \{\mathfrak{m} : D_{\mathfrak{m}}(\pi) \cong \tau\},$$

has a unique minimal element under the Zelevinsky ordering, where \mathfrak{m} runs for all multisegments. The main result of this article includes commutativity and subsequent property of the minimal sequence. At the end of this article, we conjecture some module structure arising from the minimality.

Part 1. Introduction and preliminaries

1. INTRODUCTION

1.1. The poset $\mathcal{S}(\pi, \tau)$ and the minimal sequence. Let F be a non-Archimedean local field. Let $G_n = \mathrm{GL}_n(F)$, the general linear group over F . Fix a cuspidal representation ρ throughout the whole article. All the representations we consider are smooth and over \mathbb{C} . Let $\mathrm{Irr}(G_n)$ be the set of irreducible representations of G_n . We shall usually not distinguish isomorphic irreducible representations. For a representation π_1 of G_{n_1} and a representation π_2 of G_{n_2} , denote the normalized parabolic induction by $\pi_1 \times \pi_2$.

The complex representation theory of $\mathrm{GL}_n(F)$ has fruitful study in the literature, back to [Ze80]. One central object is the so-called multisegments that parametrize irreducible representations of $\mathrm{GL}_n(F)$. Recently, [Ch25, Ch25b] found interesting combinatorics on multisegments arising from certain sequence of derivatives of essentially square-integrable representations in the content of Bernstein-Zelevinsky derivatives [Ch25, Ch25b]. The notion of highest derivative multisegments is a key combinatorial notion to encode information for derivatives and is quite computable (c.f. explicit algorithms in [LM16, CP25+]). These results are motivated and have applications to quotient branching laws [Ch22+b], and also attempt to generalize some aspects of some segment cases in [AL25, LM25] to multisegment cases.

This article continues to study those sequences of derivatives. A main goal is to establish and prove some general properties of a sequence of derivatives, and explain why those properties are natural from some structure of Jacquet modules. In order to define the notion of derivatives precisely, we need more notations, following Zelevinsky [Ze80].

Let $\nu : G_n \rightarrow \mathbb{C}^\times$ be the character $\nu(g) = |\det(g)|_F$, where $|\cdot|_F$ is the normalized absolute value for F . Let $\mathrm{Irr}_\rho(G_n)$ be the set of irreducible representations of G_n which are an irreducible constituent of $\nu^{a_1} \rho \times \dots \times \nu^{a_r} \rho$ for some integers a_1, \dots, a_r . Let $\mathrm{Irr}_\rho = \sqcup_n \mathrm{Irr}_\rho(G_n)$.

We now define some combinatorial objects to parametrize and study representations. For $a, b \in \mathbb{Z}$ with $b - a \in \mathbb{Z}_{\geq 0}$, we call $[a, b]_\rho$ to be a *segment* (associated to ρ). We also set $[a, a - 1]_\rho = \emptyset$ for $a \in \mathbb{Z}$. For a segment $\Delta = [a, b]_\rho$, we write $a(\Delta) = a$ and $b(\Delta) = b$. We also write $[a]_\rho := [a, a]_\rho$. Let Seg_ρ be the set of segments. A *multisegment* (associated to ρ) is a multiset of non-empty segments.

Let Mult_ρ be the set of multisegments. For each segment $\Delta \in \text{Seg}_\rho$, we associate to the unique essentially square-integrable representation, $\text{St}(\Delta)$, of G_n as one of the irreducible composition factors in $\nu^a \rho \times \dots \times \nu^b \rho$ (also see [Ch25, Section 2.6]). One may refer to [Ze80] a more general notion of multisegments, which we shall not use in this article.

For $\pi \in \text{Irr}_\rho(G_n)$ and a segment $\Delta \in \text{Seg}_\rho$, there is at most one irreducible module $\tau \in \text{Irr}_\rho(G_{n-i})$ such that

$$\pi \hookrightarrow \tau \times \text{St}(\Delta).$$

If such τ exists, we denote such τ by $D_\Delta(\pi)$ and call Δ to be *admissible* to π . Otherwise, we set $D_\Delta(\pi) = 0$. We shall refer D_Δ to be a *derivative*.

A sequence of segments $[a_1, b_1]_\rho, \dots, [a_k, b_k]_\rho$ (all $a_j, b_j \in \mathbb{Z}$) is said to be in an *ascending order* if for any $i \leq j$, either $[a_i, b_i]_\rho$ and $[a_j, b_j]_\rho$ are unlinked; or $a_i < a_j$. For a multisegment $\mathbf{n} \in \text{Mult}_\rho$, which we write the segments in \mathbf{n} in an ascending order $\Delta_1, \dots, \Delta_k$. Define

$$D_{\mathbf{n}}(\pi) := D_{\Delta_k} \circ \dots \circ D_{\Delta_1}(\pi).$$

The derivative is independent of the choice of an ascending order [Ch25]. In particular, one may choose an ordering such that $a_1 \leq \dots \leq a_k$. We say that \mathbf{n} is *admissible* to π if $D_{\mathbf{n}}(\pi) \neq 0$. We refer the reader to [LM16, Ch24, Ch25] for more theory on derivatives.

For $\pi \in \text{Irr}_\rho$, denote its i -th Bernstein-Zelevinsky derivatives by $\pi^{(i)}$. We shall refer the reader [Ze80, Ch21, Ch25] for the precise definition, and the main discussions and proofs will not involve the use of Bernstein-Zelevinsky derivatives. The main relation of derivatives and Bernstein-Zelevinsky derivatives is that $D_{\mathbf{n}}(\pi)$ is a simple quotient of $\pi^{(i)}$ [Ch25], where $i = l_{abs}(\mathbf{n})$ (see Section 3.1 for the notation l_{abs}). The goal of this series of articles [Ch25, Ch25b] is to study constructions of Bernstein-Zelevinsky derivatives from $D_{\mathbf{n}}(\pi)$. In particular, [Ch25b] studies the following poset: for a simple quotient τ of $\pi^{(i)}$,

$$\mathcal{S}(\pi, \tau) := \{\mathbf{n} \in \text{Mult}_\rho : D_{\mathbf{n}}(\pi) \cong \tau\}.$$

The ordering \leq_Z on $\mathcal{S}(\pi, \tau)$ is the Zelevinsky ordering (see Section 4.1). We recall two fundamental combinatorial structure on the set $\mathcal{S}(\pi, \tau)$:

Theorem 1.1. [Ch25b, Theorem 1.1] *We use the notations above. Let $\mathbf{n}_1, \mathbf{n}_2 \in \mathcal{S}(\pi, \tau)$ with $\mathbf{n}_1 \leq_Z \mathbf{n}_2$. If $\mathbf{n} \in \text{Mult}_\rho$ with $\mathbf{n}_1 \leq_Z \mathbf{n} \leq_Z \mathbf{n}_2$, then $\mathbf{n} \in \mathcal{S}(\pi, \tau)$.*

Theorem 1.2. [Ch25b, Theorem 1.2] *We use the notations above. Suppose $\mathcal{S}(\pi, \tau) \neq \emptyset$. Then $\mathcal{S}(\pi, \tau)$ has a unique \leq_Z -minimal element.*

For $\pi \in \text{Irr}_\rho$, a multisegment $\mathbf{n} \in \text{Mult}_\rho$ is said to be *minimal* to π if $D_{\mathbf{n}}(\pi) \neq 0$ and \mathbf{n} is \leq_Z -minimal in $\mathcal{S}(\pi, D_{\mathbf{n}}(\pi))$. We shall sometimes refer such \mathbf{n} to be the *minimal multisegment* or *minimal sequence* (of derivatives) of $\mathcal{S}(\pi, \tau)$.

1.2. Main results. The main goal of this article is to obtain some properties of the minimal sequence, which are useful in [Ch22+b]. We also provide some (partly conjectural) representation theoretic interpretations in Part 3. The main results are the following subsequence and commutativity phenomenons:

Theorem 1.3. (=Theorem 8.6) *Let $\pi \in \text{Irr}_\rho$. If $\mathbf{n} \in \text{Mult}_\rho$ is minimal to π , then any submultisegment \mathbf{n}' of \mathbf{n} is also minimal to π and in particular, $D_{\mathbf{n}'}(\pi) \neq 0$.*

Theorem 1.4. (=Theorem 9.4) *Let $\pi \in \text{Irr}_\rho$. If $\mathbf{n} \in \text{Mult}_\rho$ is minimal to π , then for any submultisegment \mathbf{n}' of \mathbf{n} , we have:*

- (1) $\mathbf{n} - \mathbf{n}'$ is minimal to $D_{\mathbf{n}'}(\pi)$; and
- (2) $D_{\mathbf{n}-\mathbf{n}'} \circ D_{\mathbf{n}'}(\pi) \cong D_{\mathbf{n}}(\pi)$.

By using Theorem 1.4 multiple times, we have:

Corollary 1.5. *Let $\pi \in \text{Irr}_\rho$. Let $\mathbf{n} \in \text{Mult}_\rho$ be minimal to π . Write the segments in \mathbf{n} as $\{\Delta_1, \dots, \Delta_r\}$ in any order. Then,*

$$D_{\Delta_r} \circ \dots \circ D_{\Delta_1}(\pi) \cong D_{\mathbf{n}}(\pi).$$

One important ingredient in studying the commutativity is a notion of η -invariants (see Definition 4.1), which also plays an important role in studying "left-right" commutativity in [Ch22+c]. It seems that such commutativity phenomenon in derivatives plays a crucial role in quotient branching laws [Ch22+b], and the above results are not quite expected before, not even that the minimal sequences seem to be known before. It is interesting to see whether there are some deeper reasons behind such commutativity. The embedding model in Conjecture 11.3 is an attempt to provide some explanations on that, and the interplay with the removal process is conjectured in Section 13.3.

In order to demonstrate the above non-straightforward results, it is unavoidable to work on some details. The main idea of the proofs for Theorems 1.3 and 1.4 is to reduce checking elementary intersection-union processes by Theorem 1.1. Then one uses some more basic commutativity for two segment case (e.g. Proposition 5.4) to reduce to three segment cases in Section 7. There are also interactions between commutativity and minimality in the proofs.

While our proofs are largely combinatorial in nature, a motivation comes from a simple example from Lemma 10.2 below. There are some further results on minimality such as a construction from the removal process, which will be explored in another sequel [CZ26+].

The commutativity and minimality also play important roles in the branching law [Ch22+b]. The uniqueness of minimality is closely related to the layer of Bernstein-Zelevinsky filtration determining a branching law [Ch22+b].

We remark that it has been known that the representation theory of reductive p -adic groups has been known to have connections to many objects of type A. It is a classical result of Borel-Casselman to transfer results to the module category of affine Hecke algebras (see e.g. [Ro86, CS19]), and then transfer to the module category of quantum affine algebras of type A (e.g [Ch86, CP96]). One may expect that the results can be translated into those setting in suitable manners, and may have some other interpretations. It is also an interesting question to see whether the results have a geometric counterpart in the geometry for ABV packets, see e.g. a recent study in [CR24] as well as in [AL25].

We also remark that Bernstein-Zelevinsky derivatives can be viewed as certain kinds of degenerate Whittaker models. While the sequences of derivatives study structure of Bernstein-Zelevinsky derivatives, the information is more sensitive in the representation category rather than simply in its Grothendieck group. This is in contrast to some study on some behaviours of wavefront sets in the content of degenerate Whittaker model, see, e.g. [Mu03], more recent study [CMBO24] and references therein, and hence our study should also provide some complementary information.

1.3. Discussions on applications. For $\pi \in \text{Irr}_\rho$ and $\Delta \in \text{Seg}_\rho$, instead of studying $D_\Delta(\pi)$, one studies on a so-called big derivative in [Ch24] involving some higher structures. It is shown in [Ch24] to be useful to study a reduced decomposition [AL25] for π in the following sense:

$$(1.1) \quad \text{St}(\mathbf{p}) \times D_{\mathbf{p}}(\pi) \rightarrow \pi \quad (\text{equivalently, } \pi \hookrightarrow D_{\mathbf{p}}(\pi) \times \text{St}(\mathbf{p})),$$

where $\mathbf{p} = \mathfrak{mr}(\pi, \Delta)$ for some segment Δ (see (10.7) for the definition of \mathfrak{mrpt}^a). In Appendix B, we give a generalization to multisegment cases. Such reduced decomposition is also useful to study the relation between Bernstein-Zelevinsky derivatives and layers of Bernstein-Zelevinsky filtrations in [Ch22+b].

Another application is to give an inductive construction of some simple quotients of Bernstein-Zelevinsky derivatives (which is a main goal in this series of articles). For example, for $\pi \in \text{Irr}_\rho$ and $\Delta \in \text{Seg}_\rho$, if a simple quotient of $\pi^{(i)}$ is Δ -reduced in the sense that $\mathfrak{mr}(\pi, \Delta) = \emptyset$ (see (10.7) for a detailed notion), then one may construct such simple quotient from a simple quotient of

$(D_{\mathfrak{p}}(\pi))^{(i-l)}$ via (1.1), where $l = l_{abs}(\mathfrak{p})$. The idea of this construction is closely related to the commutativity discussed above and see Proposition 15.2 for a precise statement.

1.4. Organization. In first few sections, we recall some main ingredients: highest derivative multisegments in Section 2, removal processes in Section 3, and non-overlapping and intermediate segment properties in Section 4.

Sections 5 to 9 study the commutativity and subsequent property for minimal sequences. The approach is largely combinatorial using the overlapping property. Section 5 studies the two segment case while Section 7 studies the three segment case. Section 6 shows some preliminary results for general cases. Sections 8 and 9 prove the general case for the subsequent and commutativity property respectively.

Sections 10 to 13 study some representation-theoretic aspects of the minimality. Section 10 explains a representation-theoretic proof of commutativity of two segment case. Section 11 conjectures a representation-theoretic interpretation for the minimality and proves for the two segment case. Sections 12 and 13 study how the interpretation gives some applications and connections to removal processes.

1.5. Acknowledgements. This article is benefited from author's visit to NCTS at Taiwan in December 2023 and December 2024, and the author would like to thank the center for hospitality. This project is partly supported by the Research Grants Council of the Hong Kong Special Administrative Region, China (Project No: 17305223) and NSFC grant for Excellent Young Scholar (Project No.: 12322120).

2. HIGHEST DERIVATIVE MULTISEGMENTS

The highest derivative multisegment is introduced in [Ch25, Section 8.1] as a main tool for the entire study. In this section, we first recall the definition and then prove a new realization theorem, which is of independent interests.

2.1. Highest derivative multisegments. For $c \in \mathbb{Z}$ and $\pi \in \text{Irr}_\rho$, define $\text{mrpt}^a(\pi, c)$ to be the maximal multisegment such that

- (1) for any $\Delta \in \text{mrpt}^a(\pi, c)$, $a(\Delta) = c$; and
- (2) $D_{\text{mrpt}^a(\pi, c)}(\pi) \neq 0$.

Here the maximality is to take the lexicographical ordering on the $b(\Delta)$ values for all segments in $\text{mrpt}^a(\pi, c)$. See [Ch25] for details and examples. Define the *highest derivative multisegment* of $\pi \in \text{Irr}_\rho$ to be

$$\mathfrak{h}\mathfrak{d}(\pi) := \sum_{c \in \mathbb{Z}} \text{mrpt}^a(\pi, c).$$

It is shown in [Ch25, Theorem 7.3] that $D_{\mathfrak{h}\mathfrak{d}(\pi)}(\pi)$ is the highest derivative of π in the sense of [Ze80]. An efficient algorithm in computing the highest derivative multisegment is given in [CP25+]. However, a key point is that effects of derivatives can be reflected from the removal process on the highest derivative multisegment, and so we can directly study removal processes.

2.2. Realization Theorem. For $d, m \in \mathbb{Z}_{>0}$ and a cuspidal representation ρ , define $u_\rho(d, m)$ to be the unique simple quotient of

$$\text{St} \left(\left[-\frac{d-m}{2}, \frac{d+m-2}{2} \right]_\rho \right) \times \text{St} \left(\left[-\frac{d-m}{2} - 1, \frac{d+m-2}{2} - 1 \right]_\rho \right) \times \dots \times \text{St} \left(\left[-\frac{d+m-2}{2}, \frac{d-m}{2} \right]_\rho \right).$$

In other words, $u_\rho(d, m)$ is the Langlands quotient of the above parabolically induced module, see [Ch25, Section 2.6] for some notations. One special property of $u_\rho(d, m)$ is that it is unitarizable, while this is not the main property used in this content.

An irreducible representation π of G_n is said to be an *essentially Speh representation* if $\pi \cong \nu^c \cdot u_\rho(d, m)$ for some $c \in \mathbb{Z}$ and some $d, m \in \mathbb{Z}_{>0}$. Denote such representation by $u_\rho(c, d, m)$. We now show the following realization theorem by an explicit construction. The key idea is to exploit some commutativity of the product of some essentially Speh representations. For results in this theme, one sees e.g. [Ba08], [Ta15], [LM16] and [Ch21].

To facilitate discussions, for $\pi \in \text{Irr}_\rho$, define $\text{csupp}(\pi)$ to be the multiset of cuspidal representations $\{\rho_1, \dots, \rho_r\}$ such that π is an irreducible composition factor in $\rho_1 \times \dots \times \rho_r$.

Lemma 2.1. *Let $u_\rho(c, d, m)$ be an essentially Speh representation. Let $\pi \in \text{Mult}_\rho$. Suppose, for any $\rho' \in \text{csupp}(\pi)$, $\rho' \in \text{csupp}(u_\rho(c, d, m))$. Then $\pi \times u_\rho(c, d, m)$ is irreducible, and*

$$\pi \times u_\rho(c, d, m) \cong u_\rho(c, d, m) \times \pi.$$

Proof. See e.g. [LM16, Section 6], [Ch21, Section 9] and [Ch24, Proposition 5.2]. \square

The highest derivative multisegments for essentially Speh representations are particularly simple:

Lemma 2.2. *Let $u_\rho(c, d, m)$ be an essentially Speh representation. Then*

$$\mathfrak{hd}(u_\rho(c, d, m)) = \left\{ \left[c - \frac{d-m}{2}, c + \frac{d+m-2}{2} \right]_\rho \right\}.$$

Proof. See [Ch25, Section 11.3]. \square

Theorem 2.3. *Let $\mathfrak{m} \in \text{Mult}_\rho$. Then there exists $\pi \in \text{Irr}_\rho$ such that*

$$\mathfrak{hd}(\pi) = \mathfrak{m}.$$

Proof. We label the segments in \mathfrak{m} as:

$$\Delta_1, \dots, \Delta_r$$

such that $b(\Delta_1) \leq b(\Delta_2) \leq \dots \leq b(\Delta_r)$.

We simply let $\pi_1 = \text{St}(\Delta_1)$. It is clear that $\mathfrak{hd}(\pi_1) = \{\Delta_1\}$. Now, for $i \geq 2$, we recursively define π_i to be an essentially Speh representation $u_\rho(c_i, d_i, m_i)$ such that for any $\sigma \in \text{csupp}(\pi_{i-1})$, $\sigma \in \text{csupp}(\pi_i)$. We just have to justify such π_i exists. To see this, we write $\Delta_i = [a_i, b_i]_\rho$ and we can first choose d_i large enough such that any representation in $\text{csupp}(\pi_{i-1})$ lies in $[b_i - d_i + 1, b_i]_\rho$. (Such d_i exists by using $b(\Delta_i) \geq b(\Delta_{i-1})$.) For such fixed d_i , now we solve c_i and m_i such that $\mathfrak{hd}(u_\rho(c_i, d_i, m_i)) = \Delta_i$.

Now let

$$\pi = \pi_1 \times \dots \times \pi_r.$$

The cuspidal conditions guarantee that π is irreducible by applying Lemma 2.1 multiple times. By [Ch25, Proposition 11.1], (also see the Arthur type representation in [Ch25, Proposition 11.2]), we have that

$$\mathfrak{hd}(\pi) = \mathfrak{hd}(\pi_1) + \dots + \mathfrak{hd}(\pi_r) = \mathfrak{m}.$$

\square

Part 2. Combinatorial aspects

3. REMOVAL PROCESSES

In this section, we recall some results of removal processes in [Ch25].

3.1. More notations on multisegments. Let $\deg(\rho)$ be the number such that ρ is in $\text{Irr}(G_{\deg(\rho)})$. For a segment $\Delta = [a, b]_\rho$, let $l_{abs}(\Delta) = (b - a + 1)\deg(\rho)$. For a multisegment \mathfrak{m} in Mult_ρ and an integer c , let $\mathfrak{m}[c]$ be the submultisegment of \mathfrak{m} containing all the segments Δ satisfying $a(\Delta) = c$; and let $\mathfrak{m}\langle c \rangle$ be the submultisegment of \mathfrak{m} containing all the segments Δ satisfying $b(\Delta) = c$.

For a multisegment $\mathfrak{m} = \{\Delta_1, \dots, \Delta_k\}$, we also set:

$$l_{abs}(\mathfrak{m}) = l_{abs}(\Delta_1) + \dots + l_{abs}(\Delta_k).$$

3.2. Removal process. We write $[a, b]_\rho \prec^L [a', b']_\rho$ if either $a < a'$; or $a = a'$ and $b < b'$. A segment $\Delta = [a, b]_\rho$ is said to be *admissible* to a multisegment \mathfrak{h} if there exists a segment of the form $[a, c]_\rho$ in \mathfrak{h} for some $c \geq b$. We now recall the *removal process*.

Definition 3.1. [Ch25, Section 8.2] Let $\mathfrak{h} \in \text{Mult}_\rho$ and let $\Delta = [a, b]_\rho$ be admissible to \mathfrak{h} . The removal process on \mathfrak{h} by Δ is an algorithm to carry out the following steps:

- (1) Pick a shortest segment $[a, c]_\rho$ in $\mathfrak{h}[a]$ satisfying $b \leq c$. Set $\Delta_1 = [a, c]_\rho$. Set $a_1 = a$ and $b_1 = c$.
- (2) One recursively finds the \prec^L -minimal segment $\Delta_i = [a_i, b_i]_\rho$ in \mathfrak{h} such that $a_{i-1} < a_i$ and $b \leq b_i < b_{i-1}$ ¹. The process stops if one can no longer find those segments.
- (3) Let $\Delta_1, \dots, \Delta_r$ be all those segments. For $1 \leq i < r$, define $\Delta_i^{tr} = [a_{i+1}, b_i]_\rho$ and $\Delta_r^{tr} = [b_r + 1, b]_\rho$ (possibly empty).
- (4) Define

$$\mathfrak{r}(\Delta, \mathfrak{h}) := \mathfrak{h} - \sum_{i=1}^r \Delta_i + \sum_{i=1}^r \Delta_i^{tr}.$$

We call $\Delta_1, \dots, \Delta_r$ to be the *removal sequence* for (Δ, \mathfrak{h}) . We also define $\Upsilon(\Delta, \mathfrak{h}) = \Delta_1$, the first segment of the removal sequence. If Δ is not admissible to \mathfrak{h} , we set $\mathfrak{r}(\Delta, \mathfrak{h}) = \infty$, called the infinity multisegment. We also set $\mathfrak{r}(\Delta, \infty) = \infty$.

3.3. Properties of removal process. For a segment $\Delta = [a, b]_\rho \neq \emptyset$, let ${}^-\Delta = [a + 1, b]_\rho$. Two non-empty segments $\Delta = [a, b]_\rho$ and $\Delta' = [a', b']_\rho$ in Seg_ρ are said to be *linked* if one of the following conditions holds:

- (1) $a < a' \leq b + 1 \leq b'$; or
- (2) $a' < a \leq b' + 1 \leq b$.

Otherwise, Δ and Δ' are called to be not linked or unlinked. For two linked segments Δ, Δ' , we write $\Delta < \Delta'$ if it is in the first condition above. Otherwise, we write $\Delta' < \Delta$. For example, $[2, 3]_\rho < [4, 5]_\rho$.

We recall the following properties for computations:

Lemma 3.2. [Ch25] Let $\mathfrak{h} \in \text{Mult}_\rho$ and let $\Delta, \Delta' \in \text{Seg}_\rho$ be admissible to \mathfrak{h} . Then

- (1) [Ch25, Lemma 8.7] Let $\mathfrak{h}^* = \mathfrak{h} - \Upsilon(\Delta, \mathfrak{h}) + {}^-\Upsilon(\Delta, \mathfrak{h})$. Then $\mathfrak{r}(\Delta, \mathfrak{h}) = \mathfrak{r}({}^-\Delta, \mathfrak{h}^*)$.
- (2) [Ch25, Lemma 8.8] Write $\Delta = [a, b]_\rho$. For any $a' < a$, $\mathfrak{r}(\Delta, \mathfrak{h})[a'] = \mathfrak{h}[a']$.
- (3) [Ch25, Lemma 8.9] If $\Delta \in \mathfrak{h}$, then $\mathfrak{r}(\Delta, \mathfrak{h}) = \mathfrak{h} - \Delta$.
- (4) [Ch25, Lemma 8.10] Suppose $a(\Delta) = a(\Delta')$. Then

$$\Upsilon(\Delta, \mathfrak{h}) + \Upsilon(\Delta', \mathfrak{r}(\Delta, \mathfrak{h})) = \Upsilon(\Delta', \mathfrak{h}) + \Upsilon(\Delta, \mathfrak{r}(\Delta', \mathfrak{h})).$$

- (5) [Ch25, Lemma 8.12] If Δ, Δ' are unlinked, then $\mathfrak{r}(\Delta', \mathfrak{r}(\Delta, \mathfrak{h})) = \mathfrak{r}(\Delta, \mathfrak{r}(\Delta', \mathfrak{h}))$.

For $\mathfrak{h} \in \text{Mult}_\rho$, a multisegment $\mathfrak{n} = \{\Delta_1, \dots, \Delta_r\} \in \text{Mult}_\rho$ written in an ascending order, define:

$$\mathfrak{r}(\mathfrak{n}, \mathfrak{h}) = \mathfrak{r}(\Delta_r, \dots, \mathfrak{r}(\Delta_1, \mathfrak{h}) \dots).$$

We say that \mathfrak{n} is *admissible* to \mathfrak{h} if $\mathfrak{r}(\mathfrak{n}, \mathfrak{h}) \neq \infty$.

Theorem 3.3. [Ch25, Theorem 10.2] Let $\pi \in \text{Irr}_\rho$. Let $\mathfrak{m}, \mathfrak{m}' \in \text{Mult}_\rho$ be admissible to π . Then $\mathfrak{m}, \mathfrak{m}'$ are admissible to $\mathfrak{hd}(\pi)$, and furthermore, $D_{\mathfrak{m}}(\pi) \cong D_{\mathfrak{m}'}(\pi)$ if and only if $\mathfrak{r}(\mathfrak{m}, \pi) = \mathfrak{r}(\mathfrak{m}', \pi)$.

¹There is a missing condition in [Ch25] and [Ch25b]: $b \leq b_i$. The author would like to thank Peng Zhou for pointing out that. Alternatively, one can use Lemma 3.2(1) to compute the removal process. There are no changes in results and proofs in [Ch25] and [Ch25b].

3.4. More relations to derivatives. For $\mathfrak{h} \in \text{Mult}_\rho$ and $\Delta = [a, b]_\rho \in \text{Seg}_\rho$, set

$$(3.2) \quad \varepsilon_\Delta(\mathfrak{h}) = |\{\tilde{\Delta} \in \mathfrak{h}[a] : \Delta \subset \tilde{\Delta}\}|.$$

Define $\varepsilon_\Delta(\pi) := \varepsilon_\Delta(\mathfrak{h}\mathfrak{d}(\pi))$, which is equivalent to a different formulation of the same notation in [Ch25, Section 4.2]. We also remark that when Δ is a singleton, this number coincides with the number defined in [Ja07, Definition 2.1.1]. In terms of derivatives, we have the following:

Theorem 3.4. [Ch25, Theorem 9.3] *Let $\pi \in \text{Irr}_\rho$. Let $\Delta = [a, b]_\rho \in \text{Seg}_\rho$ be admissible to π . Let $\Delta' = [a', b']_\rho$ be another segment in Seg_ρ . If either $a' > a$; or Δ' and Δ are unlinked, then*

$$\varepsilon_{\Delta'}(D_\Delta(\pi)) = \varepsilon_{\Delta'}(\mathfrak{r}(\Delta, \pi)).$$

Theorem 3.5. [Ch25, Theorem 9.3] *Let $\pi \in \text{Irr}_\rho$. Let $\Delta = [a, b]_\rho \in \text{Seg}_\rho$ be admissible to π . Let $\Delta' = [a', b']_\rho$ be another segment. If $a' < a$, then*

$$\varepsilon_{\Delta'}(D_\Delta(\pi)) \geq \varepsilon_{\Delta'}(\mathfrak{r}(\Delta, \pi)) = \varepsilon_{\Delta'}(\pi).$$

We remark that the equality in Theorem 3.5 follows from Lemma 3.2(2).

4. NON-OVERLAPPING PROPERTY FOR A SEQUENCE

In [Ch25b], we have shown some characterizations for the minimality of two segment case. The goal of this section is to generalize a so-called non-overlapping property to a multisegment case. For this, we first recall some ingredients in [Ch25b]: fine chains and local minimizability. Those ingredients are combinatorial in nature and so most statements will be formulated for Mult_ρ rather than Irr_ρ .

4.1. Zelevinsky ordering. For two linked segments $\Delta = [a, b]_\rho$ and $\Delta' = [a', b']_\rho$ with $\Delta < \Delta'$, we define:

$$\Delta \cup \Delta' = [a, b']_\rho, \quad \Delta \cap \Delta' = [a', b]_\rho.$$

A multisegment \mathfrak{n} is said to be obtained from another multisegment \mathfrak{m} by an *elementary intersection-union process* if there exists a pair of linked segments Δ, Δ' in \mathfrak{m} such that

$$\mathfrak{n} = \mathfrak{m} - \Delta - \Delta' + \Delta \cup \Delta' + \Delta \cap \Delta'.$$

Here the subtractions mean the (multi-)set theoretic subtractions and additions mean the (multi-)set theoretic unions. We shall also use such notions for subtractions and additions later. Note that $\Delta \cap \Delta'$ is possibly the empty set and in such case, we simply drop the term. For example, $\{[1, 3]_\rho, [2]_\rho, [4, 5]_\rho\}$ and $\{[1, 2]_\rho, [2, 5]_\rho\}$ are obtained from $\{[1, 2]_\rho, [2, 3]_\rho, [4, 5]_\rho\}$ by elementary intersection-union processes.

For two multisegments $\mathfrak{m}, \mathfrak{n} \in \text{Mult}_\rho$, we write $\mathfrak{n} \leq_Z \mathfrak{m}$ if \mathfrak{n} is obtained from \mathfrak{m} by a sequence of intersection-union processes, or $\mathfrak{m} = \mathfrak{n}$. It is well-known from [Ze80] that \leq_Z defines a partial ordering on Mult_ρ .

4.2. Non-overlapping property and intermediate segment property. We first define the η -invariant, which is crucial in studying the minimality of sequences of derivatives:

Definition 4.1. (1) Let $\mathfrak{h} \in \text{Mult}_\rho$. Let $\Delta = [a, b]_\rho \in \text{Seg}_\rho$ be admissible to \mathfrak{h} . Define

$$\eta_\Delta(\mathfrak{h}) := (\varepsilon_{[a, b]_\rho}(\mathfrak{h}), \varepsilon_{[a+1, b]_\rho}(\mathfrak{h}), \dots, \varepsilon_{[b, b]_\rho}(\mathfrak{h})).$$

(2) Let $\pi \in \text{Irr}_\rho$. Let $\Delta \in \text{Seg}_\rho$ be admissible to π . Define $\eta_\Delta(\pi) = \eta_\Delta(\mathfrak{h}\mathfrak{d}(\pi))$.

Definition 4.2. Let $\Delta, \Delta' \in \text{Seg}_\rho$ with $\Delta < \Delta'$. Let $\mathfrak{h} \in \text{Mult}_\rho$. Suppose Δ is admissible to \mathfrak{h} .

(1) We say that $(\Delta, \Delta', \mathfrak{h})$ satisfies the *non-overlapping property* if

$$\eta_{\Delta'}(D_\Delta(\pi)) = \eta_{\Delta'}(\pi).$$

(2) We say that $(\Delta, \Delta', \mathfrak{h})$ satisfies the *intermediate segment property* if there exists a segment $\tilde{\Delta} \in \mathfrak{h}$ such that

$$a(\Delta) \leq a(\tilde{\Delta}) < a(\Delta'), \text{ and } a(\tilde{\Delta}) \leq b(\Delta) \leq b(\tilde{\Delta}) < b(\Delta').$$

We remark that the original formulation of the non-overlapping property in [Ch25b] is phrased in terms of some properties among segments related to intersections between segments. We shall use the above equivalent combinatorial formulation (see [Ch25b, Proposition 9.5]), which will be more useful for our study later.

Proposition 4.3. [Ch25b, Proposition 9.5] *Let $\Delta, \Delta', \mathfrak{h}$ be as in Definition 4.2. We further assume that Δ' is admissible to \mathfrak{h} . Then the following conditions are equivalent:*

- (1) $(\Delta, \Delta', \mathfrak{h})$ satisfies the non-overlapping property.
- (2) $(\Delta, \Delta', \mathfrak{h})$ satisfies the intermediate segment property.
- (3) $\{\Delta, \Delta'\}$ is the \leq_Z -minimal element in $\mathcal{S}(\mathfrak{h}, \mathfrak{r}(\{\Delta, \Delta'\}, \mathfrak{h}))$, where $\mathcal{S}(\mathfrak{h}, \mathfrak{r}(\{\Delta, \Delta'\}, \mathfrak{h}))$ is defined as:

$$\mathcal{S}(\mathfrak{h}, \mathfrak{r}(\{\Delta, \Delta'\}, \mathfrak{h})) := \{\mathfrak{n} \in \text{Mult}_\rho : \mathfrak{r}(\mathfrak{n}, \mathfrak{h}) = \mathfrak{r}(\{\Delta, \Delta'\}, \mathfrak{h})\}.$$

4.3. Fine chains.

Definition 4.4. Let $\mathfrak{h} \in \text{Mult}_\rho$ and let $\mathfrak{n} \in \text{Mult}_\rho$. Let a be the smallest integer such that $\mathfrak{n}[a] \neq \emptyset$. Write $\mathfrak{n}[a] = \{\Delta_1, \dots, \Delta_k\}$.

- Define $\mathfrak{r}_1 = \mathfrak{h}$. For $i \geq 2$, define

$$\mathfrak{r}_i := \mathfrak{r}(\Delta_{i-1}, \dots, \mathfrak{r}(\Delta_1, \mathfrak{h}) \dots).$$

Define

$$\mathfrak{fs}(\mathfrak{n}, \mathfrak{h}) := \{\Upsilon(\Delta_1, \mathfrak{r}_1), \dots, \Upsilon(\Delta_k, \mathfrak{r}_k)\}.$$

- Define

$$\mathfrak{trr}(\mathfrak{n}, \mathfrak{h}) := \mathfrak{h} - \mathfrak{fs}(\mathfrak{n}, \mathfrak{h}) + {}^-(\mathfrak{fs}(\mathfrak{n}, \mathfrak{h}))$$

and

$$\mathfrak{trd}(\mathfrak{n}, \mathfrak{h}) := \mathfrak{n} - \mathfrak{n}[a] + {}^-(\mathfrak{n}[a]).$$

Definition 4.5. Let $\mathfrak{h} \in \text{Mult}_\rho$. Let $\mathfrak{n} \in \text{Mult}_\rho$ be admissible to \mathfrak{h} . Set $\mathfrak{h}_0 = \mathfrak{h}$ and $\mathfrak{n}_0 = \mathfrak{n}$. Define recursively

$$\mathfrak{h}_i = \mathfrak{trr}(\mathfrak{n}_{i-1}, \mathfrak{h}_{i-1}), \quad \mathfrak{n}_i = \mathfrak{trd}(\mathfrak{n}_{i-1}, \mathfrak{h}_{i-1}).$$

The *fine chain for the removal process* for $(\mathfrak{n}, \mathfrak{h})$ (or simply fine chain for $(\mathfrak{n}, \mathfrak{h})$) is the sequence

$$\mathfrak{fs}(\mathfrak{n}_0, \mathfrak{h}_0), \mathfrak{fs}(\mathfrak{n}_1, \mathfrak{h}_1), \dots$$

The sequences $\mathfrak{h}_0, \mathfrak{h}_1, \dots$ and $\mathfrak{n}_0, \mathfrak{n}_1, \dots$ will also be useful later.

Lemma 4.6. *We use the notations in Definition 4.5. Then, for all i ,*

$$\mathfrak{r}(\mathfrak{n}, \mathfrak{h}) = \mathfrak{r}(\mathfrak{n}_i, \mathfrak{h}_i).$$

Proof. This follows from repeated uses of Lemma 3.2(1), or a slightly stronger statement of [Ch25b, Lemma 3.4]. \square

4.4. Local minimizability.

Definition 4.7. Let $\mathfrak{h} \in \text{Mult}_\rho$ and let $\mathfrak{n} \in \text{Mult}_\rho$. Let a be the smallest integer such that $\mathfrak{n}[a] \neq \emptyset$. We say that $(\mathfrak{n}, \mathfrak{h})$ is *locally minimizable* if there exists a segment Δ in $\mathfrak{n}[a+1]$ such that

$$|\{\tilde{\Delta} \in \mathfrak{n}[a] : \Delta \subset \tilde{\Delta}\}| < |\{\tilde{\Delta} \in \mathfrak{fs}(\mathfrak{n}, \mathfrak{h}) : \Delta \subset \tilde{\Delta}\}|.$$

Note that each $\tilde{\Delta} \in \mathfrak{n}[a]$ satisfying $\Delta \subset \tilde{\Delta}$ guarantees the corresponding first segment $\overline{\Delta}$ in the removal sequence satisfying $\Delta \subset \overline{\Delta}$. Roughly speaking, when the difference of two cardinalities in Definition 4.7 is non-zero, one can find a "short" segment in $\mathfrak{n}[a]$ to do the intersection-union process which still does not change the choices of first segments in the removal processes.

Theorem 4.8. [Ch25b, Lemma 6.1 and Lemma 6.2] *Let $\mathfrak{h} \in \text{Mult}_\rho$. Let $\mathfrak{n} \in \text{Mult}_\rho$ be admissible to \mathfrak{h} . Let $\mathfrak{h}_0, \mathfrak{h}_1, \dots$ and $\mathfrak{n}_0, \mathfrak{n}_1, \dots$ be as constructed in Definition 4.5. Then \mathfrak{n} is minimal to \mathfrak{h} if and only if $(\mathfrak{n}_i, \mathfrak{h}_i)$ is not locally minimizable for all i .*

4.5. Non-overlapping property for a sequence. We now generalize Proposition 4.3 to a multisegment situation, which will be used in Section 7.3. We first prove a lemma:

Lemma 4.9. *Let $\Delta = [a, b]_\rho$ and $\Delta' = [c, d]_\rho$ be two segments. Suppose $\Delta' < \Delta$. Let $\mathfrak{h} \in \text{Mult}_\rho$ such that Δ' is admissible to \mathfrak{h} . If $\Delta \subset \Upsilon(\Delta', \mathfrak{h})$, then $\eta_\Delta(\mathfrak{r}(\Delta', \mathfrak{h})) \neq \eta_\Delta(\mathfrak{h})$.*

Proof. Let $\Delta_1, \dots, \Delta_r$ be the removal sequence for (Δ', \mathfrak{h}) . Let i^* , which exists from our assumption, be the largest integer such that $b(\Delta) \leq b(\Delta_{i^*})$. Write $\Delta_{i^*}^{tr} = [a_{i^*}, b_{i^*}]_\rho$. Then $\Delta_{i^*}^{tr} \neq \emptyset$ contributes extra one to $\varepsilon_{[a_{i^*}, b]_\rho}$ for $\mathfrak{r}(\Delta', \mathfrak{h})$. However, the segments $\Delta_1, \dots, \Delta_r$ as well as Δ_j^{tr} ($j \neq i^*$) do not contribute to $\varepsilon_{[a_{i^*}, b]_\rho}$. This implies the desired inequality. \square

Proposition 4.10. *Let $\mathfrak{h} \in \text{Mult}_\rho$. Let $\mathfrak{n} \in \text{Mult}_\rho$ be minimal to \mathfrak{h} . Let $\Delta = [a, b]_\rho$ be a segment such that $a(\Delta') < a$ and $b(\Delta') < b$ for any $\Delta' \in \mathfrak{n}$. Then $\mathfrak{n} + \Delta$ is still minimal to \mathfrak{h} if and only if*

$$\eta_\Delta(\mathfrak{r}(\mathfrak{n}, \mathfrak{h})) = \eta_\Delta(\mathfrak{h}).$$

Proof. We construct a sequence of multisegments $\mathfrak{n}_0, \mathfrak{n}_1, \mathfrak{n}_2, \dots$ and $\mathfrak{h}_0, \mathfrak{h}_1, \mathfrak{h}_2, \dots$ as in Definition 4.5. By Lemma 4.6, we have:

$$(*) \quad \mathfrak{r}(\mathfrak{n}_0, \mathfrak{h}_0) = \mathfrak{r}(\mathfrak{n}_1, \mathfrak{h}_1) = \dots$$

Let a_i be the smallest integer such that $\mathfrak{n}_i[a_i] \neq \emptyset$.

Let i^* be the index such that $a_{i^*} = c-1$. If such index does not exist, it implies that $b(\tilde{\Delta}) < c-1$ for all $\tilde{\Delta} \in \mathfrak{n}$. In such case, by a direct computation on removal process using Definition 3.1(3) and (4), one has $\mathfrak{r}(\mathfrak{n}, \mathfrak{h})[x] = \mathfrak{h}[x]$ for $x \geq c$. In particular, $\eta_\Delta(\mathfrak{r}(\mathfrak{n}, \mathfrak{h})) = \eta_\Delta(\mathfrak{h})$; and $\mathfrak{n} + \Delta$ is still minimal since Δ is unlinked to any segment in \mathfrak{n} . In other words, both conditions are automatically satisfied if such i^* does not exist.

We now assume such i^* exists. We first prove the only if direction. By the minimality condition and Theorem 4.8, $(\mathfrak{n}_{i^*} + \Delta, \mathfrak{h}_{i^*})$ is not locally minimizable. On the other hand, the hypothesis in this proposition guarantees that any segment $\tilde{\Delta}$ in \mathfrak{n}_{i^*} satisfies $\Delta \not\subset \tilde{\Delta}$ and so

$$|\{\tilde{\Delta} \in \mathfrak{n}_{i^*}[a] : \Delta \subset \tilde{\Delta}\}| = 0.$$

The local non-minimizability on $(\mathfrak{n}_{i^*} + \Delta, \mathfrak{h}_{i^*})$ implies that

$$|\{\tilde{\Delta} \in \mathfrak{fs}(\mathfrak{n}_{i^*} + \Delta, \mathfrak{h}_{i^*}) : \Delta \subset \tilde{\Delta}\}| = 0.$$

In other words, for any segment $[x, y]_\rho$ in $\mathfrak{fs}(\mathfrak{n}_{i^*} + \Delta, \mathfrak{h}_{i^*}) = \mathfrak{fs}(\mathfrak{n}_{i^*}, \mathfrak{h}_{i^*})$ (the equality follows from Definition 4.5), it must take the form $[x', y']_\rho$ for $y' \leq b-1$. Thus the segments involved or produced in the removal process (by the nesting property) cannot contribute to $\eta_\Delta(\mathfrak{r}(\mathfrak{n}_{i^*}, \mathfrak{h}_{i^*}))$ and so

$$(**) \quad \eta_\Delta(\mathfrak{r}(\mathfrak{n}_{i^*}, \mathfrak{h}_{i^*})) = \eta_\Delta(\mathfrak{h}_{i^*}).$$

Moreover, \mathfrak{h}_{i^*} is obtained by truncating left points for $\nu^x \rho$ for some $x \leq c - 2$. Thus, $\eta_\Delta(\mathfrak{h}_0) = \dots = \eta_\Delta(\mathfrak{h}_{i^*})$ by Definitions 4.4 and 4.5. Combining with $(*)$ and $(**)$, we have the desired equation, proving the only if direction.

We now prove the if direction. By the construction of \mathfrak{h}_i (Definition 4.5), we have:

$$\eta_\Delta(\mathfrak{h}) = \eta_\Delta(\mathfrak{h}_{i^*})$$

Now with $(*)$ and the hypothesis $\eta_\Delta(\mathfrak{h}) = \eta_\Delta(\mathfrak{r}(\mathfrak{n}, \mathfrak{h}))$, we again have that

$$\eta_\Delta(\mathfrak{r}(\mathfrak{n}_{i^*}, \mathfrak{h}_{i^*})) = \eta_\Delta(\mathfrak{h}_{i^*}).$$

This condition says that $\mathfrak{f}\mathfrak{s}(\mathfrak{n}_{i^*}, \mathfrak{h}_{i^*})$ cannot take the form $[c - 1, y]_\rho$ for some $y \geq b$, by Lemma 4.9 and the assumption that any segment in \mathfrak{n} (and so \mathfrak{n}_{i^*}) satisfies $b(\Delta) < b$. In other words, $(\mathfrak{n}_{i^*}, \mathfrak{h}_{i^*})$ is not locally minimizable. The local non-minimizability for other pairs $(\mathfrak{n}_j, \mathfrak{h}_j)$ for $j < i^*$ follows from the minimality of \mathfrak{n} . Then the minimality of $\mathfrak{n} + \Delta$ to \mathfrak{h} follows from Theorem 4.8. \square

5. TWO SEGMENT BASIC CASE (COMMUTATIVITY)

5.1. Lemma for unlinked segments. We recall the following first commutativity (see e.g. [Ch25, Lemma 4.4]), which will be used later:

Lemma 5.1. *Let $\Delta, \Delta' \in \text{Seg}_\rho$ be unlinked. For any $\pi \in \text{Irr}_\rho$,*

$$D_{\Delta'} \circ D_\Delta(\pi) \cong D_\Delta \circ D_{\Delta'}(\pi).$$

5.2. Intermediate segment property under a derivative.

Lemma 5.2. *Let $\pi \in \text{Irr}_\rho$. Let $\Delta, \Delta'' \in \text{Seg}_\rho$ be admissible to π . Let $\Delta' \in \text{Seg}_\rho$. Suppose $(\Delta, \Delta', \mathfrak{h}\mathfrak{d}(\pi))$ satisfies the non-overlapping property or intermediate segment property. If $\Delta'' \subset \Delta'$, then $(\Delta, \Delta', \mathfrak{h}\mathfrak{d}(D_{\Delta''}(\pi)))$ also satisfies the non-overlapping property and the intermediate segment property.*

Proof. Since the non-overlapping property and the intermediate segment property are equivalent by Proposition 4.3, it suffices to see that $(\Delta, \Delta', \mathfrak{h}\mathfrak{d}(D_{\Delta'}(\pi)))$ also satisfies the intermediate segment property.

Write $\Delta = [a, b]_\rho$, $\Delta' = [a', b']_\rho$ and $\Delta'' = [a'', b'']_\rho$. By Theorems 3.4 and 3.5, we have that for any segment $\tilde{\Delta} = [\tilde{a}, \tilde{b}]_\rho$,

$$(*) \quad \varepsilon_{\tilde{\Delta}}(\mathfrak{r}(\Delta'', \pi)) \leq \varepsilon_{\tilde{\Delta}}(D_{\Delta''}(\pi)).$$

From this, one can recover $\mathfrak{h}\mathfrak{d}(\pi)[c]$ and $\mathfrak{r}(\Delta, \pi)[c]$ for each integer c . The missing part of $(*)$ between $\mathfrak{r}(\Delta', \mathfrak{h}\mathfrak{d}(\pi))$ and $\mathfrak{h}\mathfrak{d}(D_{\Delta''}(\pi))$ is on some values $a'' - 1, \dots, b'' - 1 < b'$. Thus, one obtains $\mathfrak{h}\mathfrak{d}(D_{\Delta''}(\pi))$ by prolonging some segments in $\mathfrak{r}(\Delta'', \mathfrak{h}\mathfrak{d}(\pi))$ using (possibly some of) $a'' - 1, \dots, b'' - 1$.

Now, by the intermediate segment property for $(\Delta, \Delta', \mathfrak{h}\mathfrak{d}(\pi))$, there exists a segment of the form $[c, d]_\rho$ satisfying:

$$a \leq c < a', \quad b \leq d < b'.$$

Now, by the above process of obtaining $\mathfrak{h}\mathfrak{d}(D_{\Delta''}(\pi))$, the segment $[c, d]_\rho$ can be prolonged to the form $[c, e]_\rho$ in $\mathfrak{r}(\Delta'', \pi)$ for some $e \leq b' - 1$. Then the segment $[c, e]_\rho$ gives the desired requirement for the intermediate segment property for $(\Delta, \Delta', D_{\Delta''}(\pi))$. Thus, by Proposition 4.3, the triple satisfies the two properties. \square

For a segment $\Delta = [a, b]_\rho \in \text{Seg}_\rho$ and $\pi \in \text{Irr}_\rho$, define:

$$|\eta|_\Delta(\pi) = \varepsilon_{[a, b]_\rho}(\pi) + \varepsilon_{[a+1, b]_\rho}(\pi) + \dots + \varepsilon_{[b, b]_\rho}(\pi).$$

In view of (3.2), $|\eta|_\Delta(\pi)$ measures the number of segments $\overline{\Delta}$ in $\mathfrak{h}\mathfrak{d}(\pi)$ satisfying that $\nu^a \rho \leq a(\overline{\Delta}) \leq \nu^b \rho$ and $\nu^b \rho \leq b(\overline{\Delta})$. For example, suppose we have $\pi \in \text{Irr}_\rho$ such that

$$\mathfrak{h}\mathfrak{d}(\pi) = \{[1, 4]_\rho, [1, 3]_\rho, [1, 2]_\rho, [2, 5]_\rho, [2, 4]_\rho\}.$$

Then $|\eta|_{[1,3]_\rho}(\pi) = 4$, which is contributed from $[1,4]_\rho, [1,3]_\rho$ for $\varepsilon_{[1,3]_\rho}(\pi)$ and from $[2,5]_\rho, [2,4]_\rho$ for $\varepsilon_{[2,3]_\rho}(\pi)$.

Lemma 5.3. *Let $\pi \in \text{Irr}_\rho$. Let $\Delta, \Delta' \in \text{Seg}_\rho$. Suppose Δ is admissible to π . Suppose $(\Delta, \Delta', \mathfrak{hd}(\pi))$ satisfies the non-overlapping property or the intermediate segment property. Let $\tilde{\Delta} = \Delta \cup \Delta'$.*

$$|\eta|_{\tilde{\Delta}}(\pi) - |\eta|_{\Delta'}(\pi) = |\eta|_{\tilde{\Delta}}(D_\Delta(\pi)) - |\eta|_{\Delta'}(D_\Delta(\pi)).$$

Proof. Write the removal sequence for (Δ, π) to be:

$$\Delta_1, \dots, \Delta_r.$$

By the intermediate segment property, there exists a segment Δ_k such that $a(\Delta_k) < a(\Delta')$ and $b(\Delta_k) < b(\Delta')$. Let k^* be the smallest such integer. Then, with the nesting property, only $\Delta_1, \dots, \Delta_{k^*-1}$ (among $\Delta_1, \dots, \Delta_r$) contribute to $|\eta|_{\tilde{\Delta}}(\pi) - |\eta|_{\Delta'}(\pi)$.

Write $\Delta = [a, b]_\rho$ and $\Delta' = [a', b']_\rho$. By Theorem 3.4, for $a \leq c \leq a' - 1$

$$\varepsilon_{[c,b']_\rho}(D_\Delta(\pi)) = \varepsilon_{[c,b']_\rho}(\mathfrak{r}(\Delta, \pi)).$$

Then, only $\Delta_1^{tr}, \dots, \Delta_{k^*-1}^{tr}$ (among $\Delta_1^{tr}, \dots, \Delta_r^{tr}$) can contribute to $|\eta|_{\tilde{\Delta}}(\pi) - |\eta|_{\Delta'}(\pi)$ and so this implies the equality. \square

5.3. Commutativity and minimality for two segment case.

Proposition 5.4. *Let $\pi \in \text{Irr}_\rho$. Let $\Delta_1, \Delta_2 \in \text{Seg}_\rho$ be admissible to π . Suppose $\Delta_1 < \Delta_2$ and $(\Delta_1, \Delta_2, \mathfrak{hd}(\pi))$ satisfies the non-overlapping property, or equivalently*

$$D_{\Delta_2} \circ D_{\Delta_1}(\pi) \not\cong D_{\Delta_1 \cup \Delta_2} \circ D_{\Delta_1 \cap \Delta_2}(\pi).$$

Then

$$D_{\Delta_2} \circ D_{\Delta_1}(\pi) \cong D_{\Delta_1} \circ D_{\Delta_2}(\pi).$$

Proof. The equivalence of the two conditions follows from Proposition 4.3.

We shall use the notations in the proof. Indeed, in view of a criteria of commutativity in [Ch25, Proposition 6.2], it suffices to prove

$$D_{\Delta_1} \circ D_{\Delta_2}(\pi) \not\cong D_{\Delta_1 \cup \Delta_2} \circ D_{\Delta_1 \cap \Delta_2}(\pi).$$

Let $\tilde{\Delta} = \Delta_1 \cup \Delta_2$. To this end, it suffices to show that

$$(5.3) \quad (|\eta|_{\tilde{\Delta}} - |\eta|_{\Delta_2})(D_{\Delta_1} \circ D_{\Delta_2}(\pi)) = (|\eta|_{\tilde{\Delta}} - |\eta|_{\Delta_2})(\pi).$$

Note that, by the unlinked part of Theorem 3.4,

$$(|\eta|_{\tilde{\Delta}} - |\eta|_{\Delta_2})(D_{\Delta_2}(\pi)) = (|\eta|_{\tilde{\Delta}} - |\eta|_{\Delta_2})(\pi)$$

On the other hand, by Proposition 4.3(2), $(\Delta_1, \Delta_2, \pi)$ satisfies the intermediate segment property and so $(\Delta_1, \Delta_2, D_{\Delta_2}(\pi))$ also satisfies the intermediate segment property by Lemma 5.2. Now, by Lemma 5.3,

$$(|\eta|_{\tilde{\Delta}} - |\eta|_{\Delta_2})(D_{\Delta_1} \circ D_{\Delta_2}(\pi)) = (|\eta|_{\tilde{\Delta}} - |\eta|_{\Delta_2})(D_{\Delta_2}(\pi)).$$

Combining above two equations, we have (5.3) as desired. \square

We shall give a proof of Proposition 5.4 from a representation-theoretic perspective (see Lemma 10.2).

6. SOME PRELIMINARY RESULTS FOR SUBSEQUENT AND COMMUTATIVITY PROPERTIES

6.1. Cancellative property.

Proposition 6.1. (*Cancellative property*) Let $\pi \in \text{Irr}_\rho$. Let \mathbf{n} and \mathbf{n}' be multisegments with respective segments in the following respective ascending sequences:

$$\Delta'_1, \dots, \Delta'_p, \Delta_1, \dots, \Delta_r$$

and

$$\Delta''_1, \dots, \Delta''_q, \Delta_1, \dots, \Delta_r.$$

Then $\mathbf{r}(\mathbf{n}', \pi) = \mathbf{r}(\mathbf{n}'', \pi)$ if and only if

$$\mathbf{r}(\{\Delta'_1, \dots, \Delta'_p\}, \pi) = \mathbf{r}(\{\Delta''_1, \dots, \Delta''_q\}, \pi).$$

Proof. The if direction is straightforward. We now consider the only if direction. By Theorem 2.3, let $\pi \in \text{Irr}_\rho$ such that $\mathbf{h}\mathbf{d}(\pi) = \mathbf{h}$. By Theorem 3.3, we have that

$$D_{\Delta_r} \circ \dots \circ D_{\Delta_1} \circ D_{\Delta'_p} \circ \dots \circ D_{\Delta'_1}(\pi) \cong D_{\Delta_r} \circ \dots \circ D_{\Delta_1} \circ D_{\Delta'_q} \circ \dots \circ D_{\Delta'_1}(\pi).$$

For any irreducible $\tau \in \text{Irr}_\rho$ and any segment $\Delta \in \text{Seg}_\rho$, denote by $I_\Delta(\tau)$ the unique irreducible submodule of $\pi \times \text{St}(\Delta)$. Now, by uniqueness, $I_{\Delta_i} \circ D_{\Delta_i}(\tau) \cong \tau$ if $D_{\Delta_i}(\tau) \neq 0$ for any i and irreducible τ . Hence, we cancel the derivatives $D_{\Delta_r}, \dots, D_{\Delta_1}$ to obtain:

$$D_{\Delta'_p} \circ \dots \circ D_{\Delta'_1}(\pi) \cong D_{\Delta'_q} \circ \dots \circ D_{\Delta'_1}(\pi).$$

□

6.2. **First subsequent property.** We shall frequently use the following simple fact:

Lemma 6.2. [Ze80, Section 6.7] Let $\Delta_1, \dots, \Delta_r$ be a sequence of segments in an ascending order. Suppose Δ_k and Δ_{k+1} are linked for some k . Then

$$\Delta_1, \dots, \Delta_{k-1}, \Delta_k \cup \Delta_{k+1}, \Delta_k \cap \Delta_{k+1}, \Delta_{k+2}, \dots, \Delta_r$$

is also in an ascending order.

A particular case of the lemma is that if $\Delta', \Delta'', \Delta'''$ are in an ascending order, then one has:

- (1) $\Delta' \cap \Delta'', \Delta' \cup \Delta'', \Delta'''$ is also in an ascending order;
- (2) $\Delta', \Delta'' \cap \Delta''', \Delta'' \cup \Delta'''$ is also in an ascending order.

From these two simple cases, one can deduce some other variations needed in the following Proposition 6.3.

Proposition 6.3. Let $\pi \in \text{Irr}_\rho$. Let \mathbf{n} be minimal to π . We write the segments in \mathbf{n} in an ascending order

$$\Delta_1, \dots, \Delta_r.$$

Then, for any $s \leq r$,

- (1) $\{\Delta_1, \dots, \Delta_s\}$ is still minimal to π ; and
- (2) $\{\Delta_{s+1}, \dots, \Delta_r\}$ is minimal to $\mathbf{r}(\{\Delta_1, \dots, \Delta_s\}, \pi)$.

Proof. We only prove (1), and (2) can be proved similarly.

The admissibility follows from definitions (and Lemma 3.2(5)). Let

$$\mathbf{n}' = \{\Delta_1, \dots, \Delta_s\}.$$

We pick two linked segments Δ_i and Δ_j in \mathbf{n}' and we set

$$\mathbf{n}'' = \mathbf{n}' - \{\Delta_i, \Delta_j\} + \Delta_i \cup \Delta_j + \Delta_i \cap \Delta_j.$$

It suffices to show that $\mathbf{r}(\mathbf{n}'', \mathbf{h}) \neq \mathbf{r}(\mathbf{n}, \mathbf{h})$. To this end, we first write the segments in \mathbf{n}'' in an ascending order:

$$\Delta'_1, \dots, \Delta'_s.$$

(There are $s - 1$ segments if $\Delta_i \cap \Delta_j = \emptyset$, but the below arguments could be still applied.) It follows from Lemma 6.2 that $\Delta'_1, \dots, \Delta'_s, \Delta_{s+1}, \dots, \Delta_r$ are still in an ascending order.

Now we return to the proof. The minimality of \mathbf{n} implies that

$$\mathbf{r}(\{\Delta'_1, \dots, \Delta'_s, \Delta_{s+1}, \dots, \Delta_r\}, \pi) \neq \mathbf{r}(\{\Delta_1, \dots, \Delta_s, \Delta_{s+1}, \dots, \Delta_r\}, \pi).$$

By Proposition 6.1,

$$\mathbf{r}(\{\Delta'_1, \dots, \Delta'_s\}, \pi) \neq \mathbf{r}(\{\Delta_1, \dots, \Delta_s\}, \pi),$$

as desired. \square

7. THREE SEGMENT BASIC CASES

The main goal of this section is to prove the subsequent property and commutativity for three segment cases. To show the minimality, the main strategy is to use the convex property for $\mathcal{S}(\pi, \tau)$ of Theorem 1.1 and the non-overlapping property of Theorem 4.3.

7.1. Case: $\{\Delta_1, \Delta_3\}$ minimal to $D_{\Delta_2}(\pi)$.

Lemma 7.1. *Let $\Delta, \Delta' \in \text{Seg}_\rho$ with $\Delta \subset \Delta'$. Let $\pi \in \text{Irr}_\rho$ with $D_{\Delta'}(\pi) \neq 0$. Then, the following holds:*

- (1) *If $a(\Delta) > a(\Delta')$, then $\eta_\Delta(D_{\Delta'}(\pi)) = \eta_\Delta(\pi)$.*
- (2) *If $a(\Delta) = a(\Delta')$, then $\eta_\Delta(D_{\Delta'}(\pi))$ is obtained from $\eta_\Delta(\pi)$ by decreasing the coordinate $\varepsilon_\Delta(\pi)$ by 1.*

Proof. By Theorem 3.4, it suffices to compare $\eta_\Delta(\mathbf{r}(\Delta', \pi))$ and $\eta_\Delta(\pi)$. Let the removal sequence for (Δ', π) be

$$\Delta_1, \dots, \Delta_r.$$

For (1), we consider two cases.

- (i) Suppose there does not exist an integer i^* such that $a(\Delta_{i^*}) \geq a(\Delta)$ and $a(\Delta_{i^*}) \leq b(\Delta)$. In such case, all $\Delta_1, \dots, \Delta_r$ and $\Delta_1^{tr}, \dots, \Delta_r^{tr}$ do not contribute η_Δ . Thus we have such equality.
- (ii) Suppose there exists an integer i^* such that $a(\Delta_{i^*}) \geq a(\Delta)$. Let $i^* > 1$ be the smallest such integer. Let j^* be the largest integer such that $a(\Delta_{j^*}) \leq b(\Delta)$. We have that $\Delta_{i^*}, \dots, \Delta_{j^*}$ are all the segments in the removal sequence contributing to $\eta_\Delta(\pi)$ and $\Delta_{i^*-1}^{tr}, \dots, \Delta_{j^*-1}^{tr}$ are all the segments in the truncated one contributing to $\eta_\Delta(\mathbf{r}(\Delta', \pi))$. Note that, for $i^* \leq k \leq j^*$, Δ_k and Δ_{k-1}^{tr} contribute to the same coordinate $\varepsilon_{\tilde{\Delta}}$ for some segment $\tilde{\Delta}$. This shows the equality to two η_Δ .

For (2), it is similar, but i^* in above notation becomes 1. Again, for $2 \leq k \leq j^*$, Δ_k and Δ_{k-1}^{tr} contribute to the same ε . The term Δ_1 explains $\varepsilon_\Delta(\pi)$ is decreased by 1 to obtain $\varepsilon_\Delta(D_{\Delta'}(\pi))$. \square

Lemma 7.2. *Let $\mathbf{m} = \{\Delta_1, \Delta_2, \Delta_3\} \in \text{Mult}_\rho$ in an ascending order. Let $\pi \in \text{Irr}_\rho$ be such that \mathbf{m} is minimal to π . Then $\{\Delta_1, \Delta_3\}$ is also minimal to $D_{\Delta_2}(\pi)$.*

Proof. By Proposition 6.3, $\{\Delta_1, \Delta_2\}$ is minimal to π . By Proposition 5.4 (for the linked case between Δ_1 and Δ_2) and Lemma 5.1 (for the unlinked case between Δ_1 and Δ_2), we have that

$$D_{\Delta_3} \circ D_{\Delta_2} \circ D_{\Delta_1}(\pi) \cong D_{\Delta_3} \circ D_{\Delta_1} \circ D_{\Delta_2}(\pi).$$

The minimality is automatic if Δ_1 and Δ_3 are unlinked. So we shall assume that Δ_1 and Δ_3 are linked. There are three possibilities:

- Δ_2 is unlinked to Δ_1 (and so $\{\Delta_2, \Delta_1, \Delta_3\}$ is still in an ascending order). Then the minimality of \mathbf{m} and Proposition 6.3 imply this case.
- Δ_2 is unlinked to Δ_3 . We consider following possibilities:

(i) $\Delta_3 \subset \Delta_2$. Then $\{\Delta_1, \Delta_3\}$ is also minimal to π by Lemma 5.1 and Proposition 6.3. Hence, $\eta_{\Delta_3}(D_{\Delta_1}(\pi)) = \eta_{\Delta_3}(\pi)$. On the other hand, if $a(\Delta_3) > a(\Delta_2)$, by Lemma 7.1(1),

$$\eta_{\Delta_3}(D_{\Delta_2}(\pi)) = \eta_{\Delta_3}(\pi)$$

and

$$\eta_{\Delta_3}(D_{\Delta_2} \circ D_{\Delta_1}(\pi)) = \eta_{\Delta_3}(D_{\Delta_1}(\pi)).$$

Combining two equations, we have

$$\eta_{\Delta_3}(D_{\Delta_2} \circ D_{\Delta_1}(\pi)) = \eta_{\Delta_3}(D_{\Delta_2}(\pi))$$

and so, by Proposition 5.4 and Lemma 5.1 again,

$$\eta_{\Delta_3}(D_{\Delta_1} \circ D_{\Delta_2}(\pi)) = \eta_{\Delta_3}(D_{\Delta_2}(\pi)).$$

Thus, we have the minimality by Proposition 4.3.

When $a(\Delta_3) \cong a(\Delta_2)$, the argument is similar. The only difference, by Lemma 7.1(2), is that $\eta_{\Delta_3}(D_{\Delta_2}(\pi))$ (resp. $\eta_{\Delta_3}(D_{\Delta_2} \circ D_{\Delta_1}(\pi))$) is obtained from $\eta_{\Delta_3}(\pi)$ (resp. $\eta_{\Delta_3}(D_{\Delta_1}(\pi))$) by decreasing the $\varepsilon_{\Delta_3}(\pi)$ (resp. $\varepsilon_{\Delta_3}(D_{\Delta_1}(\pi))$) by 1.

- (ii) $\Delta_2 \subset \Delta_3$. In such case, one first has that $(\Delta_1, \Delta_3, \mathfrak{hd}(\pi))$ satisfies the non-overlapping property by the minimality of $\{\Delta_1, \Delta_3\}$ to π . Then, by Lemma 5.2 to show that $(\Delta_1, \Delta_3, \mathfrak{hd}(D_{\Delta_2}(\pi)))$ still satisfies the non-overlapping property. Thus, $\{\Delta_1, \Delta_3\}$ is minimal to $D_{\Delta_2}(\pi)$.
- (iii) $b(\Delta_2) < a(\Delta_3)$. Then the ascending order and linkedness between Δ_1 and Δ_3 also give that Δ_1 and Δ_2 are not linked. This goes back to the above bullet.
- (iv) $b(\Delta_3) < a(\Delta_2)$. Then the ascending order and linkedness between Δ_1 and Δ_3 also give that Δ_1 and Δ_2 are not linked. This goes back to the above bullet.
- $\Delta_1 < \Delta_2 < \Delta_3$. If $\{\Delta_1, \Delta_3\}$ is not minimal to $D_{\Delta_2}(\pi)$, then

$$D_{\Delta_1 \cup \Delta_3} \circ D_{\Delta_1 \cap \Delta_3} \circ D_{\Delta_2}(\pi) \cong D_{\Delta_3} \circ D_{\Delta_1} \circ D_{\Delta_2}(\pi) \cong D_{\Delta_3} \circ D_{\Delta_2} \circ D_{\Delta_1}(\pi).$$

This contradicts the minimality of \mathfrak{m} . □

7.2. Case: $\{\Delta_2, \Delta_3\}$ minimal to π .

Lemma 7.3. *Let $\pi \in \text{Irr}_\rho$. Let $\Delta_1, \Delta_2, \Delta_3$ be segments in an ascending order. If $\{\Delta_1, \Delta_2, \Delta_3\}$ is minimal to π , then $\{\Delta_2, \Delta_3\}$ is also minimal to π .*

Proof. When Δ_2 and Δ_3 are unlinked, there is nothing to prove. We assume that Δ_2 and Δ_3 are linked and so $\Delta_2 < \Delta_3$. We consider the following cases:

- $b(\Delta_3) \leq b(\Delta_1)$. Then Δ_1 is unlinked to both Δ_2 and Δ_3 . Then the minimality of $\{\Delta_2, \Delta_3\}$ to π follows from the minimality of $\{\Delta_1, \Delta_2, \Delta_3\}$ to π by Proposition 6.3.
- $b(\Delta_1) < b(\Delta_3)$ and $a(\Delta_1) < a(\Delta_3)$, and Δ_1 and Δ_3 are linked. Then $\{\Delta_1, \Delta_2\}$ with Δ_3 is in the situation of Proposition 4.10. Then $\eta_{\Delta_3}(D_{\{\Delta_1, \Delta_2\}}(\pi)) = \eta_{\Delta_3}(\pi)$. On the other hand, we have $\eta_{\Delta_3}(D_{\Delta_1} \circ D_{\Delta_2}(\pi)) = \eta_{\Delta_3}(D_{\Delta_2}(\pi))$ by Lemma 7.2 with Proposition 4.3(1) \Leftrightarrow (3). Furthermore, $D_{\Delta_1} \circ D_{\Delta_2}(\pi) \cong D_{\{\Delta_1, \Delta_2\}}(\pi)$, which again follows from Proposition 5.4 and Lemma 5.1 (see the top of the proof of Lemma 7.2 for a bit more details).

Thus, combining above, we have $\eta_{\Delta_3}(\pi) = \eta_{\Delta_3}(D_{\Delta_2}(\pi))$. This implies $\{\Delta_2, \Delta_3\}$ is minimal to π by Proposition 4.3(1) \Leftrightarrow (3).

- $b(\Delta_1) < b(\Delta_3)$ and $a(\Delta_1) < a(\Delta_3)$, and Δ_1 and Δ_3 are not linked. Indeed, the argument in above bullet still works except that one has to directly observe from the removal process that $\eta_{\Delta_3}(D_{\Delta_1} \circ D_{\Delta_2}(\pi)) = \eta_{\Delta_3}(D_{\Delta_2}(\pi))$.

- $b(\Delta_1) < b(\Delta_3)$ and $a(\Delta_3) \leq a(\Delta_1)$. In particular, Δ_1 is unlinked to Δ_3 . Since we are assuming $\Delta_1, \Delta_2, \Delta_3$ are in an ascending order, and we are assuming that Δ_2 and Δ_3 are linked, we also have that Δ_1 and Δ_2 are unlinked. Then the minimality of $\{\Delta_2, \Delta_3\}$ to π follows from the minimality of $\{\Delta_1, \Delta_2, \Delta_3\}$ to π and Proposition 6.3.

□

7.3. Case: $\{\Delta_1, \Delta_3\}$ minimal to π .

Lemma 7.4. *Let $\pi \in \text{Irr}_\rho$. Let $\Delta_1, \Delta_2, \Delta_3$ be segments in an ascending order. If $\{\Delta_1, \Delta_2, \Delta_3\}$ is minimal to π , then $\{\Delta_1, \Delta_3\}$ is minimal to π .*

Proof. We may assume Δ_1 and Δ_3 are linked. Otherwise, there is nothing to prove. We consider the following cases.

- $\Delta_1 < \Delta_2 < \Delta_3$. By Proposition 5.4, we have

$$D_{\{\Delta_1, \Delta_3, \Delta_2\}}(\pi) \cong D_{\Delta_2} \circ D_{\{\Delta_1, \Delta_3\}}(\pi).$$

If the minimality does not hold, then we have

$$D_{\{\Delta_1, \Delta_2, \Delta_3\}}(\pi) \cong D_{\{\Delta_2, \Delta_1 \cup \Delta_3, \Delta_1 \cap \Delta_3\}}(\pi)$$

since $\Delta_1 \cup \Delta_3, \Delta_1 \cap \Delta_3, \Delta_2$ are still in an ascending order. This contradicts to the minimality of $\{\Delta_1, \Delta_2, \Delta_3\}$ to π .

- Δ_2 and Δ_3 are not linked. Then we can switch the labellings of Δ_2 and Δ_3 , which gives the minimality of $\{\Delta_1, \Delta_3\}$ to π by Proposition 6.3.
- Δ_1 and Δ_2 are not linked. In this case, we can switch the labellings for Δ_1 and Δ_2 by using linkedness. Then the result follows from Lemma 7.3.

□

7.4. Case: $\{\Delta_1, \Delta_2\}$ minimal to $D_{\Delta_3}(\pi)$. We now need some inputs from representation theory to prove a combinatorics result. Let $N_i \subset G_n$ (depending on n) be the unipotent radical containing matrices of the form $\begin{pmatrix} I_{n-i} & * \\ & I_i \end{pmatrix}$. For a smooth representation π of G_n , we write π_{N_i} to be its Jacquet module.

Lemma 7.5. *Let $\Delta = [a, b]_\rho, \Delta' = [a', b']_\rho$ be two segments such that $\Delta < \Delta'$. Let $\omega = \text{St}(\{\Delta, \Delta'\})$. Let \mathfrak{m} be a multisegment whose segments $\Delta'' = [a'', b'']_\rho$ satisfy that $b'' = b$ and $a'' < a$. Then $\text{St}(\mathfrak{m}) \times \omega$ is irreducible and*

$$\text{St}(\mathfrak{m}) \times \omega \cong \omega \times \text{St}(\mathfrak{m}).$$

Proof. It is well-known that the second assertion implies the first one. We only have to prove the first one. Since $\text{St}(\mathfrak{m})$ can be written as $\times_{\Delta \in \mathfrak{m}} \text{St}(\Delta)$, it reduces to the case that \mathfrak{m} contains only one segment and so now we consider $\mathfrak{m} = \{\tilde{\Delta}\}$.

We analyse possible composition factors of

$$\text{St}(\tilde{\Delta}) \times \omega.$$

Since we know that a composition factor of $\text{St}(\tilde{\Delta}) \times \omega$ is also a composition factor of $\text{St}(\tilde{\Delta}) \times \text{St}(\Delta) \times \text{St}(\Delta')$, the possible composition factors are

$$\text{St}(\{\tilde{\Delta}, \Delta, \Delta'\}), \text{St}(\tilde{\Delta} \cup \Delta' + \tilde{\Delta} \cap \Delta' + \Delta), \text{St}(\Delta \cup \Delta' + \Delta \cap \Delta' + \tilde{\Delta}).$$

We denote the three representations π_1, π_2, π_3 respectively.

Thus it suffices to show that the last two composition factors cannot appear in $\text{St}(\tilde{\Delta}) \times \omega$. We first consider π_2 . Note that π_2 is generic. However, ω is not generic and so $\text{St}(\tilde{\Delta}) \times \omega$ cannot contain a generic composition factor and so π_2 cannot appear in $\text{St}(\tilde{\Delta}) \times \omega$.

We now consider π_3 . Let $l = l_{abs}(\Delta \cup \Delta')$. Then $(\pi_3)_{N_l}$ has the composition factor $\text{St}(\Delta) \boxtimes \text{St}(\Delta \cup \Delta')$. Now we consider composition factors in $(\text{St}(\Delta) \times \omega)_{N_l}$. If $(\text{St}(\Delta) \times \omega)_{N_l}$ contains the factor $\text{St}(\Delta) \boxtimes \text{St}(\Delta \cup \Delta')$, a simple composition factor is a simple composition factor in

$$\text{St}(\Delta) \times \omega_1 \boxtimes \omega_2,$$

where $\omega_1 \boxtimes \omega_2$ is a simple composition factor in ω_{N_l} . However, the possibilities of those composition factors are well-known and it is impossible for ω_2 to be the factor $\text{St}(\Delta \cup \Delta')$. \square

Lemma 7.6. *Let $\pi \in \text{Irr}_\rho$. Let $\Delta_1, \Delta_2, \Delta_3$ be segments satisfying $\Delta_1 < \Delta_2 < \Delta_3$. Suppose $\{\Delta_1, \Delta_2, \Delta_3\}$ is minimal to π . Then $D_{\Delta_3}(\pi)$. Let $\tilde{\Delta} = \Delta_1 \cup \Delta_2$. Then the followings hold:*

- (1) $\eta_{\tilde{\Delta}}(\pi) - \eta_{\Delta_2}(\pi) = \eta_{\tilde{\Delta}}(D_{\Delta_3}(\pi)) - \eta_{\Delta_2}(D_{\Delta_3}(\pi))$;
- (2) *If $(\Delta_1, \Delta_2, \pi)$ satisfies the intermediate segment property, then $(\Delta_1, \Delta_2, D_{\Delta_3}(\pi))$ also satisfies the intermediate segment property.*

Here the subtraction in (1) means the subtraction entry-wise.

Proof. We have shown in Lemma 7.3 that $\{\Delta_2, \Delta_3\}$ is minimal to π . Thus, we have

$$D_{\Delta_3} \circ D_{\Delta_2}(\pi) \not\cong D_{\Delta_2 \cup \Delta_3} \circ D_{\Delta_2 \cap \Delta_3}(\pi).$$

By a standard argument (see e.g. proof of [Ch25, Proposition 6.2]), we have that π is the unique simple submodule of

$$D_{\Delta_3} \circ D_{\Delta_2}(\pi) \times \text{St}(\{\Delta_2, \Delta_3\}).$$

Write $\Delta_2 = [a_2, b_2]_\rho$. Let

$$\mathfrak{m} = \sum_{c < a_2} \varepsilon_{[c, b_2]_\rho}(D_{\Delta_3}(\pi)) \cdot [c, b_2]_\rho,$$

and

$$\mathfrak{p} = \sum_{c < a_2} \varepsilon_{[c, b_2]_\rho}(\pi) \cdot [c, b_2]_\rho.$$

Thus, we have:

$$D_{\Delta_3} \circ D_{\Delta_2}(\pi) \hookrightarrow D_{\mathfrak{m}} \circ D_{\Delta_3} \circ D_{\Delta_2}(\pi) \times \text{St}(\mathfrak{m})$$

and so

$$\pi \hookrightarrow D_{\Delta_2} \circ D_{\Delta_2}(\pi) \times \text{St}(\{\Delta_2, \Delta_3\}) \hookrightarrow D_{\mathfrak{m}} \circ D_{\Delta_3} \circ D_{\Delta_2}(\pi) \times \text{St}(\mathfrak{m}) \times \text{St}(\{\Delta_2, \Delta_3\}).$$

By Lemma 7.5,

$$\pi \hookrightarrow D_{\mathfrak{m}} \circ D_{\Delta_3} \circ D_{\Delta_2}(\pi) \times \text{St}(\{\Delta_2, \Delta_3\}) \times \text{St}(\mathfrak{m}).$$

This implies that $D_{\mathfrak{m}}(\pi) \neq 0$ and so \mathfrak{m} is a submultisegment of \mathfrak{p} . On the other hand, using Lemma 3.2 and Theorem 3.5, we also have that \mathfrak{p} is a submultisegment of \mathfrak{m} . Hence, $\mathfrak{m} = \mathfrak{p}$. Translating to η -invariants, we obtain (1).

We now prove (2). Write $\Delta_1 = [a_1, b_1]_\rho$. Suppose $(\Delta_1, \Delta_2, \pi)$ satisfies the intermediate segment property. Then there exists a segment $[a, b]_\rho$ in $\mathfrak{hd}(\pi)$ satisfying that $a_1 \leq a < a_2$ and $b_1 \leq b < b_2$. Then,

$$\varepsilon_{[a, b]_\rho}(\pi) > \varepsilon_{[a, b_2]_\rho}(\pi)$$

By (1), we have that

$$\varepsilon_{[a, b_2]_\rho}(\pi) = \varepsilon_{[a, b_2]_\rho}(D_{\Delta_3}(\pi)).$$

By Theorem 3.5,

$$\varepsilon_{[a, b]_\rho}(D_{\Delta_3}(\pi)) \geq \varepsilon_{[a, b]_\rho}(\pi).$$

Combining the above equalities and inequalities, we have:

$$\varepsilon_{[a, b]_\rho}(D_{\Delta_3}(\pi)) > \varepsilon_{[a, b_2]_\rho}(D_{\Delta_3}(\pi)).$$

This implies that there exists a segment $[a, b']_\rho$ in $\mathfrak{hd}(D_{\Delta_3}(\pi))$ with $b' < b_2$. Thus $(\Delta_1, \Delta_2, D_{\Delta_3}(\pi))$ satisfies the intermediate segment property. \square

It is of course desirable to also have a more combinatorial proof for Lemma 7.6 while it seems to require some further developments on removal process to do so.

Lemma 7.7. *Let $\pi \in \text{Irr}_\rho$. Let $\Delta_1, \Delta_2, \Delta_3$ be segments in an ascending order. If $\{\Delta_1, \Delta_2, \Delta_3\}$ is minimal to π , then $\{\Delta_1, \Delta_2\}$ is also minimal to $D_{\Delta_3}(\pi)$.*

Proof. If Δ_1 and Δ_2 are unlinked, then there is nothing to prove. If Δ_2 and Δ_3 are unlinked, then we use Lemma 5.1 to transfer to Lemma 7.2.

The remaining case is that Δ_1 and Δ_2 are linked, and Δ_2 and Δ_3 are linked. In other words, $\Delta_1 < \Delta_2 < \Delta_3$. This case follows from Proposition 4.3(2) \Leftrightarrow (3) and Lemma 7.6. \square

7.5. Case: $\{\Delta_1, \Delta_2\}$ minimal to π .

Lemma 7.8. *Let $\pi \in \text{Irr}_\rho$. Let $\Delta_1, \Delta_2, \Delta_3$ be segments in an ascending order. If $\{\Delta_1, \Delta_2, \Delta_3\}$ is minimal to π , then $\{\Delta_1, \Delta_2\}$ is also minimal to π .*

The above lemma is a special case of Proposition 6.3(1).

7.6. Case: $\{\Delta_2, \Delta_3\}$ minimal to $D_{\Delta_1}(\pi)$.

Lemma 7.9. *Let $\pi \in \text{Irr}_\rho$. Let $\Delta_1, \Delta_2, \Delta_3$ be segments in an ascending order. If $\{\Delta_1, \Delta_2, \Delta_3\}$ is minimal to π , then $\{\Delta_2, \Delta_3\}$ is also minimal to $D_{\Delta_1}(\pi)$.*

The above lemma is again a special case of Proposition 6.3(2).

8. SUBSEQUENT PROPERTY OF MINIMAL SEQUENCES

8.1. Consecutive pairs.

Definition 8.1. Let $\mathbf{m} \in \text{Mult}_\rho$. Two segments Δ_1 and Δ_2 in \mathbf{m} are said to be *consecutive* in \mathbf{m} if

- $\Delta_1 < \Delta_2$ i.e. Δ_1 and Δ_2 are linked with $a(\Delta_1) < a(\Delta_2)$
- there is no other segment Δ' in \mathbf{m} such that

$$a(\Delta_1) \leq a(\Delta') \leq a(\Delta_2), \quad b(\Delta_1) \leq b(\Delta') \leq b(\Delta_2)$$

and Δ' is linked to either Δ_1 or Δ_2 .

(The last linkedness condition guarantees that $\Delta' \neq \Delta_1 \cap \Delta_2$ and $\Delta' \neq \Delta_1 \cup \Delta_2$.)

Example 8.2. • Let $\mathbf{h} = \{[0, 3], [1, 4], [2, 5]\}$. Then $[0, 3], [1, 4]$ form a pair of consecutive segments. Similarly, $[1, 4], [2, 5]$ also form a pair of consecutive segments, but $[0, 3], [2, 5]$ do not form a pair of consecutive segments.

- Let $\mathbf{h} = \{[0, 4], [1, 2], [2, 5]\}$. Then $[0, 4], [2, 5]$ form a pair of consecutive segments; and $[1, 2], [2, 5]$ also form a pair of consecutive segments.
- Let $\mathbf{h} = \{[0, 3], [1, 3], [2, 4], [2, 5]\}$. Then $[1, 3], [2, 4]$ form a pair of consecutive segments, while $[0, 3], [2, 4]$ do not form a pair of consecutive segments.

The terminology of consecutive segments is suggested by its property in the intersection-union process.

Lemma 8.3. *Let Δ_1, Δ_2 be linked segments with $\Delta_1 < \Delta_2$. Suppose there exists a segment Δ' satisfying the conditions in the second bullet of Definition 8.1. Then, if Δ' is linked to Δ_i ($i = 1, 2$), then*

$$\{\Delta_1 \cap \Delta_2, \Delta_1 \cup \Delta_2, \Delta'\} \leq_Z \{\Delta_i \cap \Delta', \Delta_i \cup \Delta', \Delta_j\} \leq_Z \{\Delta_1, \Delta_2, \Delta'\}.$$

Here j is the index other than i i.e. $j \in \{1, 2\} - \{i\}$.

The above lemma follows from a direct checking and we omit the details. A simple combinatorics give the following:

Lemma 8.4. (1) Let $\mathbf{m}, \mathbf{m}' \in \text{Mult}_\rho$ such that $\mathbf{m}' \leq_Z \mathbf{m}$ with $\mathbf{m}' \neq \mathbf{m}$. Then there exists a pair of consecutive segments Δ, Δ' in \mathbf{m} such that for the multisegment \mathbf{m}'' obtained from \mathbf{m} by the elementary intersection-union process involving Δ and Δ' , $\mathbf{m}' \leq_Z \mathbf{m}'' \leq_Z \mathbf{m}$.
(2) Let $\mathbf{m} \in \text{Mult}_\rho$. Let Δ, Δ' be a pair of consecutive segments in \mathbf{m} with $\Delta < \Delta'$. Let \mathbf{m}' be the submultisegment of $\mathbf{m} - \Delta - \Delta'$ that contains all the segments $\tilde{\Delta}$ with $a(\Delta') \leq a(\tilde{\Delta})$ or $b(\Delta') \leq b(\tilde{\Delta})$. Write the segments in $\mathbf{m} - \mathbf{m}' - \Delta - \Delta'$ in an ascending order: $\Delta_1, \dots, \Delta_r$ and write the segments in \mathbf{m}' in an ascending order: $\Delta'_1, \dots, \Delta'_s$. Then the sequence:

$$\Delta_1, \dots, \Delta_r, \Delta, \Delta', \Delta'_1, \dots, \Delta'_s$$

is ascending.

Proof. For (1), it suffices to show for \mathbf{m}' obtained by a pair of elementary intersection-union operation involving $\overline{\Delta}$ and $\overline{\Delta}'$. If the segments involved in the operation are consecutive, then the statement is immediate. Otherwise, there exists a segment $\tilde{\Delta}$ such that $\tilde{\Delta}$ is linked to either $\overline{\Delta}$ or $\overline{\Delta}'$, and produce a multisegment $\tilde{\mathbf{m}}$ such that $\mathbf{m}' \leq_Z \tilde{\mathbf{m}} \leq_Z \mathbf{m}''$. We repeat the process if such linked pair is still not consecutive. Note that if $\tilde{\Delta}$ is linked to $\overline{\Delta}$ (resp. $\overline{\Delta}'$), we must have $\tilde{\Delta} \cap \overline{\Delta}$ (resp. $\tilde{\Delta} \cap \overline{\Delta}'$) strictly longer than $\overline{\Delta} \cap \overline{\Delta}'$, and hence after repeating the process several times, we obtain desired consecutive segments.

For (2), it is a direct check from the definition of an ascending order. \square

8.2. Minimality under commutativity (second basic case). We first prove a commutativity result of minimal sequences, which is useful in proving Theorem 8.6.

Lemma 8.5. Let $\pi \in \text{Irr}$. Let $\Delta, \Delta_1, \Delta_2, \dots, \Delta_r$ be in an ascending order and minimal to π . Then $\{\Delta, \Delta_{k+1}, \dots, \Delta_r\}$ is also minimal to $D_{\Delta_k} \circ \dots \circ D_{\Delta_1}(\pi) \neq 0$, and

$$D_{\Delta_r} \circ \dots \circ D_{\Delta_{k+1}} \circ D_\Delta \circ D_{\Delta_k} \circ \dots \circ D_{\Delta_1}(\pi) \cong D_{\Delta_r} \circ \dots \circ D_{\Delta_1} \circ D_\Delta(\pi).$$

Proof. By induction, it suffices to show when $k = 1$. The case that Δ and Δ_1 are unlinked is easy by Lemma 5.1 and Proposition 6.3(1). Suppose $\{\Delta, \Delta_2, \dots, \Delta_r\}$ is not minimal to $D_{\Delta_1}(\pi)$ to arrive a contradiction. By Theorem 1.1 and Lemma 8.4(1), there exists a pair of consecutive segments $\Delta' < \Delta''$ such that the multisegment obtained from the intersection-union operation of those two segments Δ', Δ'' gives the same derivative on π .

Note that $\{\Delta_2, \dots, \Delta_r\}$ is minimal to $D_{\Delta_1} \circ D_\Delta(\pi) \cong D_\Delta \circ D_{\Delta_1}(\pi)$ (the last isomorphism by Proposition 5.4). Thus the remaining possible cases to be considered could be that one of Δ', Δ'' is Δ and so $\Delta' = \Delta$. Let

$$\mathbf{n} = \{\Delta_i : a(\Delta_i) \leq a(\Delta) \text{ or } b(\Delta_i) \leq b(\Delta)\}$$

Then any Δ_i is unlinked to Δ and Δ_1 by using the ascending property for the sequence $\Delta, \Delta_1, \dots, \Delta_r$ (and $\Delta < \Delta_1$). (In particular, Δ_1 is not in \mathbf{n} .)

Let $\mathbf{m} = \{\Delta_2, \dots, \Delta_r\}$. Now, since we chose Δ and Δ'' to be consecutive, we can arrange and relabel the segments in an ascending order:

$$\tilde{\Delta}_1, \dots, \tilde{\Delta}_l, \Delta, \Delta'', \tilde{\Delta}_{k+1}, \dots, \tilde{\Delta}_{r-1},$$

where all $\tilde{\Delta}_1, \dots, \tilde{\Delta}_l$ are all elements in \mathbf{n} and $\tilde{\Delta}_{k+1}, \dots, \tilde{\Delta}_{r-1}$ are all elements in $\mathbf{m} - \mathbf{n}$.

Case 1: $\mathbf{n} = \emptyset$. We still have that $\Delta, \Delta_1, \Delta'', \tilde{\Delta}_1, \dots, \tilde{\Delta}_{r-1}$ form an ascending order and is minimal to π . In particular, we have $\Delta, \Delta_1, \Delta''$ is minimal to π by the cancellative property. Thus

$$D_{\Delta''} \circ D_\Delta \circ D_{\Delta_1}(\pi) \not\cong D_{\Delta'' \cup \Delta} \circ D_{\Delta'' \cap \Delta} \circ D_{\Delta_1}(\pi)$$

by Lemma 7.2 and Proposition 5.4. Since

$$\Delta'' \cap \Delta, \Delta'' \cup \Delta, \tilde{\Delta}_1, \dots, \tilde{\Delta}_{r-1}$$

still form an ascending order (Lemma 6.2), applying $D_{\tilde{\Delta}_1}, \dots, D_{\tilde{\Delta}_{r-1}}$ gives different derivatives on π (Proposition 6.1) and so this gives a contradiction.

Case 2: $\mathbf{n} \neq \emptyset$. This implies that $\Delta + \mathbf{m} - \mathbf{n}$ is not minimal to $D_{\mathbf{n}} \circ D_{\Delta_1}(\pi)$. However, we have that, by using unlinkedness discussed in the second paragraph,

$$D_{\mathbf{n}} \circ D_{\Delta_1}(\pi) \cong D_{\Delta_1} \circ D_{\mathbf{n}}(\pi).$$

and $\Delta + \Delta_1 + \mathbf{m} - \mathbf{n}$ is minimal to $D_{\mathbf{n}}(\pi)$ by Proposition 6.3. However, from Case 1, we have that $\Delta + \mathbf{m} - \mathbf{n}$ is minimal to

$$D_{\Delta_1} \circ D_{\mathbf{n}}(\pi) (\cong D_{\mathbf{n}} \circ D_{\Delta_1}(\pi)).$$

This gives a contradiction. \square

8.3. Minimality of a subsequent sequence.

Theorem 8.6. *Let $\pi \in \text{Irr}_\rho$. Let $\mathbf{n} \in \text{Mult}_\rho$ be minimal to π . Then any submultisegment of \mathbf{n} is also minimal to π .*

Proof. By an induction, it suffices to show the minimality for $\mathbf{n}' = \mathbf{n} - \Delta$ for any segment Δ in \mathbf{n} . By Theorem 1.1 and Lemma 8.4, it reduces to check that for any multisegment \mathbf{m}' obtained from \mathbf{n}' by an elementary intersection-union operations involving two consecutive segments,

$$D_{\mathbf{m}'}(\pi) \not\cong D_{\mathbf{n}'}(\pi).$$

Denote such two consecutive segments by $\tilde{\Delta} < \tilde{\Delta}'$.

Let \mathbf{m} be obtained by the intersection-union process from \mathbf{n} involving $\tilde{\Delta}$ and $\tilde{\Delta}'$. Then $\mathbf{m}' = \mathbf{m} - \Delta$. We consider following possibilities:

(1) Case 1: Suppose $\tilde{\Delta}$ and $\tilde{\Delta}'$ still form a pair of consecutive segments in \mathbf{n} . Then we write the segments in \mathbf{n} as in Lemma 8.4 (with obvious notation replacement):

$$\Delta_1, \dots, \Delta_r, \tilde{\Delta}, \tilde{\Delta}', \Delta'_1, \dots, \Delta'_s.$$

(a) Case 1(a): Δ appears in one of $\Delta'_1, \dots, \Delta'_s$. If $D_{\mathbf{n} - \Delta}(\pi) = D_{\mathbf{m} - \Delta}(\pi)$, then the cancellative property (Proposition 6.1) and Lemma 8.4 (also see Proposition 6.3) imply that

$$D_{\{\Delta_1, \dots, \Delta_r, \tilde{\Delta}, \tilde{\Delta}'\}}(\pi) \not\cong D_{\{\Delta_1, \dots, \Delta_r, \tilde{\Delta} \cup \tilde{\Delta}', \tilde{\Delta} \cap \tilde{\Delta}'\}}(\pi).$$

However, this implies $D_{\mathbf{n}}(\pi) \not\cong D_{\mathbf{m}}(\pi)$ by applying $D_{\Delta'_1}, \dots, D_{\Delta'_s}$ with D_Δ omitted.

(b) Case 1(b): Δ appears in one of $\Delta_1, \dots, \Delta_r$. Let

$$\mathbf{p} = \{\Delta_1, \dots, \Delta_r, \tilde{\Delta}, \tilde{\Delta}'\} - \Delta$$

and let \mathbf{q} be obtained from \mathbf{p} by an elementary intersection-union process on $\tilde{\Delta}$ and $\tilde{\Delta}'$. By the cancellative property (Proposition 6.1), it suffices to show that $D_{\mathbf{p}}(\pi) \not\cong D_{\mathbf{q}}(\pi)$. Let $\tau = D_{\mathbf{p} - \tilde{\Delta} - \tilde{\Delta}' - \Delta}(\pi)$. By repeatedly using Lemma 8.5, we have:

$$D_{\mathbf{p} + \Delta}(\pi) \cong D_{\tilde{\Delta}'} \circ D_{\tilde{\Delta}} \circ D_\Delta(\tau)$$

and $\{\Delta, \tilde{\Delta}, \tilde{\Delta}'\}$ is minimal to τ . Now by Lemma 7.3, we have that

$$D_{\mathbf{p}}(\pi) = D_{\tilde{\Delta}'} \circ D_{\tilde{\Delta}}(\tau) \not\cong D_{\tilde{\Delta} \cup \tilde{\Delta}'} \circ D_{\tilde{\Delta} \cap \tilde{\Delta}'}(\tau) = D_{\mathbf{q}}(\pi)$$

as desired.

(2) Case 2: $\tilde{\Delta}$ and $\tilde{\Delta}'$ do not form a consecutive pair. We then first write the segments in \mathbf{n}' as in Lemma 8.4:

$$\Delta_1, \dots, \Delta_r, \tilde{\Delta}, \tilde{\Delta}', \Delta'_1, \dots, \Delta'_s.$$

Since the pair is not consecutive, the segment Δ must take the form as in the second bullet of Definition 8.1. Then one can still check that

$$\Delta_1, \dots, \Delta_r, \tilde{\Delta}, \Delta, \tilde{\Delta}', \Delta'_1, \dots, \Delta'_s$$

is an ascending sequence. Now let $\tau = D_{\{\Delta_1, \dots, \Delta_r\}}(\pi)$. By the cancellative property (Proposition 6.1), it suffices to show that

$$D_{\{\tilde{\Delta}, \tilde{\Delta}'\}}(\tau) \not\cong D_{\{\tilde{\Delta} \cap \tilde{\Delta}', \tilde{\Delta} \cup \tilde{\Delta}'\}}(\tau).$$

This follows from the basic case of Lemma 7.4. \square

9. COMMUTATIVITY AND MINIMALITY

In this section, we study the commutativity for a minimal sequence.

9.1. Commutativity and minimality (another basic case).

Lemma 9.1. *Let $\pi \in \text{Irr}_\rho$. Let $\mathbf{m} \in \text{Mult}_\rho$ be minimal to π . Let $\Delta \in \mathbf{m}$. Then $D_{\mathbf{m} - \Delta} \circ D_\Delta(\pi) \cong D_{\mathbf{m}}(\pi)$.*

Proof. We write the segments in $\mathbf{m} - \Delta$ in an ascending order: $\Delta_1, \dots, \Delta_r$. By Proposition 6.3(1), we can reduce to the case that $\Delta_1, \dots, \Delta_r, \Delta$ still form an ascending order. By Proposition 6.3(2) and the basic case (Proposition 5.4),

$$D_{\mathbf{m}}(\pi) \cong D_\Delta \circ D_{\mathbf{m} - \Delta}(\pi) \cong D_{\Delta_r} \circ D_\Delta \circ D_{\mathbf{m} - \Delta - \Delta_r}(\pi).$$

By Theorem 8.6, $\mathbf{m} - \Delta_r$ is still minimal to π . We now inductively obtain the statement. \square

We first study a special case of commutativity and minimality, and we shall prove a full version in Theorem 9.3.

Lemma 9.2. *Let $\pi \in \text{Irr}_\rho$. Let $\mathbf{m} \in \text{Mult}_\rho$ be minimal to π . Let c (resp. d) be the largest integer such that $\mathbf{m}[c] \neq 0$ (resp. $\mathbf{m}\langle d \rangle \neq 0$). Let $\Delta \in \mathbf{m}[c]$ or $\in \mathbf{m}\langle d \rangle$. Then $\mathbf{m} - \Delta$ is minimal to $D_\Delta(\pi)$.*

Proof. The condition in the lemma guarantees that Δ can be arranged in the last one for an ascending order for \mathbf{m} .

Suppose $\mathbf{m} - \Delta$ is not minimal to $D_\Delta(\pi)$. Let $\mathbf{m}' = \mathbf{m} - \Delta$. By Theorem 1.1 and Lemma 8.4, it suffices to show that for any pair $\tilde{\Delta} < \tilde{\Delta}'$ of consecutive segments,

$$\mathbf{n}' := \mathbf{m}' - \{\tilde{\Delta}, \tilde{\Delta}'\} + \tilde{\Delta} \cup \tilde{\Delta}' + \tilde{\Delta} \cap \tilde{\Delta}'$$

does not give the same derivative on $D_\Delta(\pi)$ i.e.

$$(*) \quad D_{\mathbf{n}'} \circ D_\Delta(\pi) \not\cong D_{\mathbf{m}'} \circ D_\Delta(\pi).$$

Now we arrange and relabel the segments in \mathbf{m}' as in Lemma 8.4:

$$\Delta_1, \dots, \Delta_k, \tilde{\Delta}, \tilde{\Delta}', \Delta_{k+3}, \dots, \Delta_r,$$

which is in an ascending order. By Proposition 6.3, in order to show $(*)$, it suffices to show that $\{\Delta_1, \dots, \Delta_k, \tilde{\Delta}, \tilde{\Delta}'\}$ is still minimal to $D_\Delta(\pi)$. Let

$$\tau = D_{\Delta_k} \circ \dots \circ D_{\Delta_1}(\pi), \text{ and } \tau' = D_{\Delta_k} \circ \dots \circ D_{\Delta_1} \circ D_\Delta(\pi).$$

By Lemma 6.2 and Proposition 6.1, it suffices to prove

$$(9.4) \quad D_{\tilde{\Delta}'} \circ D_{\tilde{\Delta}}(\tau') \not\cong D_{\tilde{\Delta}' \cup \tilde{\Delta}} \circ D_{\tilde{\Delta}' \cap \tilde{\Delta}}(\tau').$$

Let $\mathfrak{p} = \{\Delta_1, \dots, \Delta_k\}$. To this end, by the subsequent property (Theorem 8.6), we have that $\mathfrak{p} + \tilde{\Delta}' + \tilde{\Delta} + \Delta$ is minimal to π . Thus, we also have that $\{\Delta, \tilde{\Delta}', \tilde{\Delta}\}$ is minimal to $\tau = D_{\mathfrak{p}}(\pi)$ by Proposition 6.3. Now, Lemma 7.7 implies:

$$D_{\tilde{\Delta}'} \circ D_{\tilde{\Delta}} \circ D_{\Delta}(\tau) \not\cong D_{\tilde{\Delta}' \cup \tilde{\Delta}} \circ D_{\tilde{\Delta}' \cap \tilde{\Delta}} \circ D_{\Delta}(\tau).$$

On the other hand, by the subsequent property (Theorem 8.6), it gives that $\{\Delta_1, \dots, \Delta_k, \Delta\}$ is minimal to π . Then combining with Lemma 9.1, we have $D_{\Delta}(\tau) \cong \tau'$. Combining, we have the desired non-isomorphism (9.4). \square

9.2. Commutativity and minimality (one segment case).

Theorem 9.3. *Let $\pi \in \text{Irr}_{\rho}$. Let $\mathfrak{n} \in \text{Mult}_{\rho}$ be minimal to π . Let $\Delta \in \mathfrak{n}$. Then $\mathfrak{n} - \Delta$ is minimal to $D_{\Delta}(\pi)$ and*

$$D_{\mathfrak{n} - \Delta} \circ D_{\Delta}(\pi) \cong D_{\mathfrak{n}}(\pi).$$

Proof. We first prove the second assertion. We write the segments in \mathfrak{n} in an ascending order:

$$\Delta_1, \dots, \Delta_k, \Delta, \Delta_{k+1}, \dots, \Delta_r$$

with $b(\Delta_1) \leq \dots \leq b(\Delta) \leq \dots \leq b(\Delta_r)$. Then, by Proposition 6.3, $\{\Delta_1, \dots, \Delta_k, \Delta\}$ is still minimal to \mathfrak{n} . Then, the second assertion follows from Lemma 9.2.

We now prove the first assertion. By Theorem 1.1 and Lemma 8.4, it suffices to consider for a consecutive pair of segments. Let $\tilde{\Delta}, \tilde{\Delta}'$ be a pair of consecutive segments in $\mathfrak{n} - \Delta$. Let

$$\mathfrak{n}' = \mathfrak{n} - \Delta - \{\tilde{\Delta}, \tilde{\Delta}'\} + \tilde{\Delta} \cup \tilde{\Delta}' + \tilde{\Delta} \cap \tilde{\Delta}'.$$

We are going to show

$$D_{\mathfrak{n} - \Delta} \circ D_{\Delta}(\pi) \not\cong D_{\mathfrak{n}'} \circ D_{\Delta}(\pi).$$

We consider the following two cases:

(1) $\tilde{\Delta}$ and $\tilde{\Delta}'$ still form a pair of consecutive segments in \mathfrak{n} . We arrange and relabel the segments as:

$$\Delta_1, \dots, \Delta_p, \tilde{\Delta}, \tilde{\Delta}', \Delta_{p+1}, \dots, \Delta_r$$

with $\Delta_{p+1}, \dots, \Delta_r$ to be all the segments with $a(\Delta_t) \geq a(\tilde{\Delta}')$ or $b(\Delta_t) \geq b(\tilde{\Delta}')$. Similar to Lemma 8.4, one has that

$$\Delta_1, \dots, \Delta_p, \tilde{\Delta} \cup \tilde{\Delta}', \tilde{\Delta} \cap \tilde{\Delta}', \Delta_{p+1}, \dots, \Delta_r$$

form an ascending sequence.

(i) Suppose Δ appears in one of $\Delta_1, \dots, \Delta_p$, say Δ_i . Let $\tau = D_{\Delta_p} \circ \dots \circ D_{\Delta_{i+1}} \circ D_{\Delta_{i-1}} \circ \dots \circ D_{\Delta_1} \circ D_{\Delta}(\pi)$. In such case, the proved second assertion gives that

$$\tau \cong D_{\Delta_p} \circ \dots \circ D_{\Delta_1}(\pi).$$

The isomorphism with the minimality of $\{\Delta_1, \dots, \Delta_p, \tilde{\Delta}, \tilde{\Delta}'\}$ to π and the discussion on the ascending sequence above gives that

$$D_{\tilde{\Delta}'} \circ D_{\tilde{\Delta}}(\tau) \not\cong D_{\tilde{\Delta}' \cup \tilde{\Delta}} \circ D_{\tilde{\Delta}' \cap \tilde{\Delta}}(\tau).$$

Applying $D_{\Delta_{p+1}}, \dots, D_{\Delta_r}$, by Proposition 6.1, we have that

$$D_{\mathfrak{n} - \Delta} \circ D_{\Delta}(\pi) \not\cong D_{\mathfrak{n}'} \circ D_{\Delta}(\pi).$$

(ii) Suppose Δ appears in one of $\Delta_{p+1}, \dots, \Delta_r$, say Δ_j . By rearranging and relabeling the segments in $\Delta_{p+1}, \dots, \Delta_r$ if necessary, we assume $b(\Delta_{p+1}) \leq \dots \leq b(\Delta_r)$ if $b(\tilde{\Delta}') \leq b(\Delta_j)$ and assume $a(\Delta_{p+1}) \leq \dots \leq a(\Delta_r)$ if $a(\tilde{\Delta}') \leq a(\Delta_j)$. Let

$$\mathfrak{n}_j = \{\Delta_1, \dots, \Delta_p, \tilde{\Delta}, \tilde{\Delta}', \Delta_{p+1}, \dots, \Delta_j = \Delta\},$$

$$\mathfrak{n}'_j = \left\{ \Delta_1, \dots, \Delta_p, \tilde{\Delta} \cup \tilde{\Delta}', \tilde{\Delta} \cap \tilde{\Delta}', \Delta_{p+1}, \dots, \Delta_j = \Delta \right\}.$$

Then, by Lemma 9.2,

$$D_{\mathfrak{n}'_j - \Delta} \circ D_{\Delta}(\pi) \not\cong D_{\mathfrak{n}_j - \Delta} \circ D_{\Delta}(\pi)$$

Thus, applying the derivatives $D_{\Delta_{j+1}}, \dots, D_{\Delta_r}$, we have $D_{\mathfrak{n}'} \circ D_{\Delta}(\pi) \not\cong D_{\mathfrak{n} - \Delta} \circ D_{\Delta}(\pi)$ as desired.

(2) $\tilde{\Delta}$ and $\tilde{\Delta}'$ do not form a consecutive pair in \mathfrak{n} . One uses similar argument in Theorem 8.6 to reduce to three segment case. Then one reduces to a basic case (see similar discussions in the proof of Theorem 8.6), that is Lemma 7.2. In detail, we arrange the segments in \mathfrak{n} in an ascending order:

$$\Delta_1, \dots, \Delta_p, \tilde{\Delta}, \Delta, \tilde{\Delta}', \Delta_{p+1}, \dots, \Delta_r.$$

Then, we still have

$$\Delta_1, \dots, \Delta_p, \tilde{\Delta}, \Delta, \tilde{\Delta}'$$

is still minimal to π by Theorem 8.6. Then we also have $\{\tilde{\Delta}, \Delta, \tilde{\Delta}'\}$ is minimal to

$$(9.5) \quad \tau' := D_{\Delta_p} \circ \dots \circ D_{\Delta_1}(\pi).$$

By Lemma 7.2, we then have that

$$D_{\tilde{\Delta}'} \circ D_{\tilde{\Delta}} \circ D_{\Delta}(\tau') \not\cong D_{\tilde{\Delta}' \cup \tilde{\Delta}} \circ D_{\tilde{\Delta}' \cap \tilde{\Delta}} \circ D_{\Delta}(\tau').$$

But now, by Theorem 8.6, $\{\Delta_1, \dots, \Delta_p, \Delta\}$ is minimal to π , and so by the proved second assertion, we also have

$$(9.6) \quad \tau' \cong D_{\Delta_p} \circ \dots \circ D_{\Delta_1} \circ D_{\Delta}(\pi).$$

Now, one applies $D_{\Delta_{p+1}}, \dots, D_{\Delta_r}$ on (9.5), and then uses (9.6) and Proposition 6.1 to see that $D_{\mathfrak{n} - \Delta} \circ D_{\Delta}(\pi) \not\cong D_{\mathfrak{n}'} \circ D_{\Delta}(\pi)$ as desired. \square

9.3. General form of commutativity and minimality.

Theorem 9.4. *Let $\pi \in \text{Irr}_{\rho}$. Let $\mathfrak{n} \in \text{Mult}_{\rho}$ be minimal to π . Let \mathfrak{n}' be a submultisegment of \mathfrak{n} . Then $D_{\mathfrak{n} - \mathfrak{n}'} \circ D_{\mathfrak{n}'}(\pi) \cong D_{\mathfrak{n}}(\pi)$ and $\mathfrak{n} - \mathfrak{n}'$ is minimal to $D_{\mathfrak{n}'}(\pi)$.*

Proof. This follows by repeatedly using Theorem 9.3. \square

Part 3. Representation-theoretic aspects

10. η -INVARIANT AND COMMUTATIVITY

We shall first discuss the representation-theoretic interpretation for η -invariants. Then we explain how to relate with the commutativity. The representation-theoretic approach provides different techniques, which are of independent interests.

10.1. Representation-theoretic counterpart of η_{Δ} . Let $\Delta = [a, b]_{\rho}$. For $\pi \in \text{Irr}_{\rho}$, let

$$(10.7) \quad \mathfrak{m}_{\mathfrak{r}}(\pi, \Delta) = \sum_{a \leq a' \leq b} \varepsilon_{[a', b]_{\rho}}(\pi) \cdot [a', b]_{\rho},$$

which means that $[a', b]_{\rho}$ appears with multiplicity $\varepsilon_{[a', b]_{\rho}}(\pi)$ in $\mathfrak{m}_{\mathfrak{r}}(\pi, \Delta)$. This is the multisegment analogue of the η -invariant.

The following is the key property:

Lemma 10.1. [Ch24, Proposition 11.1] *Let $l = l_{\text{abs}}(\mathfrak{m}_{\mathfrak{r}}(\pi, \Delta))$. Let $\mathfrak{m} = \mathfrak{m}_{\mathfrak{r}}(\pi, \Delta)$. Then $D_{\mathfrak{m}}(\pi) \boxtimes \text{St}(\mathfrak{m})$ is a direct summand in π_{N_l} .*

10.2. Commutativity. Here we explain how to view the commutativity of Proposition 5.4 from the perspective of Lemma 10.1. We first show the following new lemma, using Lemma 10.1:

Lemma 10.2. *Let $\pi \in \text{Irr}_\rho(G_n)$. Let $\Delta_1, \Delta_2 \in \text{Seg}_\rho$ be admissible to π . Suppose $\Delta_1 < \Delta_2$. Suppose $(\Delta_1, \Delta_2, \pi)$ satisfies the non-overlapping property. Let $\mathfrak{m} = \mathfrak{m}_\mathfrak{X}(\pi, \Delta_2)$. Then $D_{\Delta_1} \circ D_{\mathfrak{m}}(\pi) \cong D_{\mathfrak{m}} \circ D_{\Delta_1}(\pi)$.*

Proof. Let $l = l_{abs}(\Delta_1)$ and let $N = N_l$. We have that:

$$\pi \hookrightarrow D_{\mathfrak{m}}(\pi) \times \text{St}(\mathfrak{m}).$$

Then

$$D_{\Delta_1}(\pi) \boxtimes \text{St}(\Delta_1) \hookrightarrow \pi_N \hookrightarrow (D_{\mathfrak{m}}(\pi) \times \text{St}(\mathfrak{m}))_N.$$

An analysis on the layers in the geometric lemma, we have that

$$(*) \quad D_{\Delta_1}(\pi) \boxtimes \text{St}(\Delta_1) \hookrightarrow D_{\mathfrak{m}}(\pi)_{N_l} \dot{\times}^1 \text{St}(\mathfrak{m}),$$

where $\dot{\times}^1$ means that the induction to a $G_{n-l} \times G_l$ -representation. By Proposition 4.10 and Lemma 10.1, $D_{\mathfrak{m}} \circ D_{\Delta_1}(\pi) \boxtimes \text{St}(\mathfrak{m})$ is a direct summand in $D_{\Delta_1}(\pi)_{N_{l'}}$, where $l' = l_{abs}(\mathfrak{m})$. Furthermore, no other composition factors in $D_{\Delta_1}(\pi)_{N_{l'}}$ take the form $\tau \boxtimes \text{St}(\mathfrak{m})$. Now, via Frobenius reciprocity on the map in $(*)$, we obtain a non-zero map from $D_{\Delta_1}(\pi)_{N_{l'}} \boxtimes \text{St}(\Delta_1)$ to $(D_{\mathfrak{m}}(\pi)_{N_l} \boxtimes \text{St}(\mathfrak{m}))^\phi$, where ϕ is a twisting sending a $G_{n-l-l'} \times G_l \times G_{l'}$ -representation to a $G_{n-l-l'} \times G_{l'} \times G_l$ -representation. Then

$$D_{\mathfrak{m}} \circ D_{\Delta_1}(\pi) \boxtimes \text{St}(\Delta_1) \hookrightarrow D_{\mathfrak{m}}(\pi)_{N_l}.$$

Thus, we have $D_{\mathfrak{m}} \circ D_{\Delta_1}(\pi) \cong D_{\Delta_1} \circ D_{\mathfrak{m}}(\pi)$. \square

Lemma 10.2 can also be deduced from Proposition 5.4 and Lemma 5.2. On the other hand, one can also give another proof for Proposition 5.4 by using Lemmas 5.2 and 10.2.

11. CONJECTURAL INTERPRETATION FOR MINIMAL SEQUENCES

11.1. Minimality for two segments.

Proposition 11.1. *Let $\pi \in \text{Irr}_\rho$. Let Δ_1, Δ_2 be a pair of linked segments with $\Delta_1 < \Delta_2$. Suppose*

$$D_{\Delta_1 \cup \Delta_2} \circ D_{\Delta_1 \cap \Delta_2}(\pi) \not\cong D_{\Delta_2} \circ D_{\Delta_1}(\pi).$$

Then the unique non-zero map

$$D_{\Delta_2} \circ D_{\Delta_1}(\pi) \boxtimes (\text{St}(\Delta_1) \times \text{St}(\Delta_2)) \rightarrow \pi_N,$$

where $N = N_{l_{abs}(\Delta_1) + l_{abs}(\Delta_2)}$, is injective.

Proof. Suppose the map is not injective. Since $D_{\Delta_2} \circ D_{\Delta_1}(\pi) \boxtimes (\text{St}(\Delta_1) \times \text{St}(\Delta_2))$ is indecomposable and has length 2, the image of the map can only be isomorphic to $D_{\Delta_2} \circ D_{\Delta_1}(\pi) \boxtimes (\text{St}(\Delta_1 \cup \Delta_2) \times \text{St}(\Delta_1 \cap \Delta_2))$. This implies that

$$D_{\Delta_2} \circ D_{\Delta_1}(\pi) \boxtimes (\text{St}(\Delta_1 \cup \Delta_2) \times \text{St}(\Delta_1 \cap \Delta_2))$$

is a submodule of π_N . Then, applying Frobenius reciprocity, we have that π is the unique submodule of $D_{\Delta_2} \circ D_{\Delta_1}(\pi) \times \text{St}(\Delta_1 \cup \Delta_2 + \Delta_1 \cap \Delta_2)$ (see [LM16, Ch24]).

Recall that $\text{St}(\Delta_1 \cup \Delta_2 + \Delta_1 \cap \Delta_2) \cong \text{St}(\Delta_1 \cup \Delta_2) \times \text{St}(\Delta_1 \cap \Delta_2)$. Let τ be the unique submodule of $D_{\Delta_2} \circ D_{\Delta_1}(\pi) \times \text{St}(\Delta_1 \cup \Delta_2)$ so that

$$(*) \quad D_{\Delta_1 \cup \Delta_2}(\tau) \cong D_{\Delta_2} \circ D_{\Delta_1}(\pi).$$

Then the uniqueness of submodule above also forces that

$$\pi \hookrightarrow \tau \times \text{St}(\Delta_1 \cap \Delta_2)$$

and so $D_{\Delta_1 \cap \Delta_2}(\pi) \cong \tau$. Combinig with $(*)$, we have:

$$D_{\Delta_1 \cup \Delta_2} \circ D_{\Delta_1 \cap \Delta_2}(\pi) \cong D_{\Delta_2} \circ D_{\Delta_1}(\pi),$$

giving a contradiction. \square

In the Appendix (Section 14), we shall prove a converse of Proposition 11.1.

11.2. A representation-theoretic interpretation of minimal sequences.

Definition 11.2. For a multisegment $\mathfrak{h} = \{\Delta_1, \dots, \Delta_r\} \in \text{Mult}_\rho$ labelled in an ascending order, define

$$\tilde{\lambda}(\mathfrak{h}) := \text{St}(\Delta_1) \times \dots \times \text{St}(\Delta_r).$$

We shall call it a *co-standard representation*. Sometimes $\lambda(\mathfrak{h})$ is used for standard modules and so we prefer to use $\tilde{\lambda}$ here.

We conjecture that Proposition 11.1 can be generalized to general minimal multisegments.

Conjecture 11.3. Let $\pi \in \text{Irr}_\rho$. Let $\mathfrak{n} \in \text{Mult}_\rho$ be minimal to π . Then the unique non-zero map

$$D_{\mathfrak{n}}(\pi) \boxtimes \tilde{\lambda}(\mathfrak{n}) \rightarrow \pi_{N_l},$$

where $l = l_{\text{abs}}(\Delta_1) + \dots + l_{\text{abs}}(\Delta_r)$, is injective.

We remark that the uniqueness of the non-zero map in Conjecture 11.3 follows from the uniqueness of simple quotients of Bernstein-Zelevinsky derivatives, shown in [Ch25, Proposition 3.15].

12. SOME APPLICATIONS ON THE EMBEDDING MODEL

In this section, we shall discuss applications of the embedding model arising from the minimality in Proposition 11.1.

Lemma 12.1. Let $\Delta', \Delta'', \Delta''' \in \text{Seg}_\rho$. Suppose $\Delta' < \Delta''$. Let $\tau = D_{\Delta''} \circ D_{\Delta'}(\pi)$ and let $l' = l_{\text{abs}}(\Delta'), l'' = l_{\text{abs}}(\Delta''), l''' = l_{\text{abs}}(\Delta''')$. Let $\kappa = D_{\Delta'''}(\pi)$. Suppose the followings hold:

- $\dim \text{Hom}_{G_{n-l'-l''-l'''} \times G_{l'+l''}}(\tau \boxtimes (\text{St}(\Delta') \times \text{St}(\Delta'')), (\kappa \times \text{St}(\Delta'''))_{N_{l'+l'''}}) \leq 1$;
- The non-zero map in the first bullet factors through the natural embedding:

$$\kappa_{N_{l'+l'''}} \dot{\times}^1 \text{St}(\Delta''') \hookrightarrow (\kappa \times \text{St}(\Delta'''))_{N_{l'+l'''}}$$

from the bottom layer in the geometric lemma. Here $\dot{\times}^1$ again denotes a parabolic induction from a $G_{n-l'-l''-l'''} \times G_{l'''} \times G_{l'+l''}$ -representation to a $G_{n-l'-l''} \times G_{l'+l''}$ -representation.

- $D_{\Delta'''} \circ D_{\Delta''} \circ D_{\Delta'}(\pi) \cong D_{\Delta''} \circ D_{\Delta'''} \circ D_{\Delta'}(\pi)$.

Then, if $\{\Delta', \Delta''\}$ is minimal to π , then $\{\Delta', \Delta''\}$ is minimal to $D_{\Delta'''}(\pi)$.

Proof. Suppose $\{\Delta', \Delta''\}$ is not minimal to $D_{\Delta'''}(\pi)$. Let $i = l' + l''$. Let $\lambda_1 = \text{St}(\Delta' \cup \Delta'') \times \text{St}(\Delta' \cap \Delta'')$. Let $i = l_{\text{abs}}(\Delta') + l_{\text{abs}}(\Delta'')$. Then, by Proposition 11.1,

$$D_{\Delta''} \circ D_{\Delta'} \circ D_{\Delta'''}(\pi) \boxtimes \lambda_1 \hookrightarrow D_{\Delta'''}(\pi)_{N_i}.$$

On the other hand, we have the following embedding:

$$\pi \hookrightarrow D_{\Delta'''}(\pi) \times \text{St}(\Delta''').$$

Let $\omega = D_{\Delta''} \circ D_{\Delta'} \circ D_{\Delta'''}(\pi) \times \text{St}(\Delta''')$. Via the bottom layer in the geometric lemma, we have an embedding:

$$\omega \boxtimes \lambda_1 \hookrightarrow (D_{\Delta'''}(\pi) \times \text{St}(\Delta'''))_{N_i}.$$

By the third bullet of the hypothesis, $D_{\Delta''} \circ D_{\Delta'}(\pi)$ is a submodule of ω . Combining above, we have that:

$$D_{\Delta''} \circ D_{\Delta'}(\pi) \boxtimes \lambda_1 \hookrightarrow (D_{\Delta'''}(\pi) \times \text{St}(\Delta'''))_{N_i}.$$

Let $\lambda_2 = \text{St}(\Delta') \times \text{St}(\Delta'')$. Then, by the minimality of $\{\Delta', \Delta''\}$ to π and Proposition 11.1, we have

$$D_{\Delta''} \circ D_{\Delta'}(\pi) \boxtimes \lambda_2 \hookrightarrow \pi_{N_i}.$$

This induces another embedding:

$$D_{\Delta''} \circ D_{\Delta'}(\pi) \boxtimes \lambda_2 \hookrightarrow (D_{\Delta'''}(\pi) \times \text{St}(\Delta'''))_{N_i}.$$

Since λ_1 and λ_2 have no isomorphic submodules, the above two embeddings give:

$$D_{\Delta''} \circ D_{\Delta'}(\pi) \boxtimes (\lambda_1 \oplus \lambda_2) \hookrightarrow (D_{\Delta'''}(\pi) \times \text{St}(\Delta'''))_{N_i}.$$

However, this induces two non-zero maps from $D_{\Delta''} \circ D_{\Delta'}(\pi) \boxtimes \lambda_1$ to $(D_{\Delta'''}(\pi) \times \text{St}(\Delta'''))_{N_i}$, which are not scalar multiple of each other. This contradicts to the first bullet. \square

Lemma 12.1 provides another strategy for checking minimality in some three segment cases in Section 7 e.g. the second bullet case in the proof of Lemma 7.2 and the linked case in the proof of Lemma 7.7. Checking the second bullet usually involves analysis on the layers arising from the geometric lemma while checking the third bullet usually uses some known commutativity from minimality (Proposition 5.4) in some other cases. Checking the first bullet requires some inputs of multiplicity-one theorems from [AGRS10] and [Ch23].

13. EMBEDDING MODEL, MINIMALITY AND REMOVAL PROCESS

13.1. Combintorial preliminaries. Let S_n be the symmetric group permuting the integers $\{1, \dots, n\}$.

Definition 13.1. Let $w \in S_n$. For $1 \leq k \leq n$ and $1 \leq l \leq n$, define

$$w[k, l] := |\{a : 1 \leq a \leq k, w(a) \geq l\}|.$$

We shall write \leq_B to be the Bruhat ordering on S_n i.e. $w' \leq_B w$ if and only if w' is a subword of a reduced expression of w . We write $w' <_B w$ if $w' \leq_B w$ and $w' \neq w$.

Proposition 13.2. [BB05, Theorem 2.1.5] *Let $w, w' \in S_n$. Then the following statements are equivalent:*

- (1) $w' \leq_B w$;
- (2) $w'[k, l] \leq w[k, l]$ for any k, l .

We shall now discuss a special situation. Let $S^{n-i,i}$ be the set of minimal representatives in the cosets in $S_n/(S_{n-i} \times S_i)$. It is well-known that if $w \in S^{n-i,i}$, then

- $w(k) < w(l)$ if $k, l \in \{1, \dots, n-i\}$ and $k < l$;
- $w(k) < w(l)$ if $k, l \in \{n-i+1, \dots, n\}$ and $k < l$.

It is straightforward to obtain the following proposition from Proposition 13.2:

Proposition 13.3. *Let $w, w' \in S^{n-i,i}$. The following statements are equivalent:*

- (1) $w' \leq_B w$;
- (2) $w'(k) \leq w(k)$ for all $k = 1, \dots, n-i$;
- (3) $w'(k) \geq w(k)$ for all $k = n-i+1, \dots, n$.

13.2. Embedding model and removal process. We now explain how the removal process plays a role in the embedding model. We first provide a basic case in Proposition 13.4 and then conjecture more general case in Section 13.6.

Proposition 13.4. *Let $\mathfrak{h} \in \text{Mult}_\rho$. Let $l = l_{\text{abs}}(\Delta)$. Then there exists an embedding*

$$\tilde{\lambda}(\mathfrak{r}(\Delta, \mathfrak{h})) \boxtimes \text{St}(\Delta) \hookrightarrow \tilde{\lambda}(\mathfrak{h})_{N_l}.$$

Proof. Write the segments in \mathfrak{m} as $\Delta_i = [a_i, b_i]_\rho$ ($i = 1, \dots, r$). We arrange the segments in \mathfrak{m} satisfying:

- $b_1 \leq b_2 \leq \dots \leq b_r$;
- if $b_i = b_{i+1}$, then $a_{i+1} \geq a_i$.

We now apply the geometric lemma on $\tilde{\lambda}(\mathfrak{m})_{N_l}$. We first write the segments as:

$$\Delta_1 = [a_1, b_1]_\rho, \dots, \Delta_r = [a_r, b_r]_\rho.$$

and $\Delta = [a, b]_\rho$. Set

$$\Delta_k^{+,l_k} = [a_k + l_k, b_k]_\rho, \quad \Delta_k^{-,l_k} = [a_k, a_k + l_k - 1]_\rho$$

Those are possibly empty sets.

Then the layers arising from the geometric lemma takes the form:

$$(*) \quad (\text{St}(\Delta_1^{+,l_1}) \times \dots \times \text{St}(\Delta_r^{+,l_r})) \boxtimes (\text{St}(\Delta_1^{-,l_1}) \times \dots \times \text{St}(\Delta_r^{-,l_r})),$$

where l_1, \dots, l_r run for all integers such that $l_1 + \dots + l_r = l/\deg(\rho)$. Let $d = \deg(\rho)$. Let $t_k = l_{\text{abs}}(\Delta_k^{+,l_k})$. We now describe the underlying element in $(S_{t_1} \times \dots \times S_{t_r}) \setminus S_n / (S_{n-i} \times S_i)$ corresponding to the geometric lemma. The assignment takes the form:

$$\begin{aligned} n - (j-1) &\mapsto t_1 + \dots + t_{r-1} + t_r - (j-1) \quad \text{for } j = 1, \dots, l_r d, \\ (n - l_r d) - (j-1) &\mapsto t_1 + \dots + t_{r-1} - (j-1) \quad \text{for } j = 1, \dots, l_{r-1} d, \\ &\vdots \\ (n - l_2 d - \dots - l_r d) - (j-1) &\mapsto t_1 - (j-1) \quad \text{for } j = 1, \dots, l_1 d. \end{aligned}$$

The assignment for $x < n - l_1 d - l_2 d - \dots - l_r d$ is then uniquely determined by using the properties of elements in $S_n / (S_{n-i} \times S_i)$ stated before Proposition 13.2.

We now consider a specific layer from the geometric lemma. The segments are chosen in the order as the removal sequence for (Δ, \mathfrak{h}) :

$$(13.8) \quad \Delta_{p_1}, \dots, \Delta_{p_e}.$$

For those indexes, if there is more than one segment for Δ_{p_i} , we always choose the one of smaller index p_i . By using the nesting property and our arrangement of segments in \mathfrak{m} , we have that

$$p_1 > p_2 > \dots > p_e.$$

If $k \neq p_i$ for some $i = 1, \dots, e$, then we set $\tilde{l}_k = 0$. If $k = p_i$ for some $i = 1, \dots, e-1$, then we set $\tilde{l}_k = a_{p_{i-1}} - a_{p_i}$. If $k = p_e$, then we set $\tilde{l}_k = b - a_{p_e}$.

We now study layers of the geometric lemma of the form $(*)$ such that, as sets,

$$\Delta_1^{+,l_1} \cup \dots \cup \Delta_r^{+,l_r} = \Delta.$$

For such layer, we say it is in *standard order* if

$$a(\Delta_1^{+,l_x}) \geq a(\Delta_r^{+,l_y})$$

for any $x < y$ with $l_x, l_y \neq 0$.

We now analyse some behaviors of two following cases.

- (1) Case 1: $\Delta_1^{+,l_1}, \dots, \Delta_r^{+,l_r}$ are in standard order. Let $q_1 < \dots < q_k$ be all the indexes such that $l_{q_x} \neq 0$. We first prove the following claim:

Claim 1: The sequence $\Delta_{q_1}, \dots, \Delta_{q_k}$ satisfies the nesting property i.e.

$$\Delta_{q_1} \supset \dots \supset \Delta_{q_k}.$$

Proof of Claim 1: Suppose the sequence does not satisfy the nesting property. Since the nesting property is not transitive, we have that a pair $\Delta_{q_x}, \Delta_{q_{x+1}}$ does not satisfy the nesting property. Due to the arrangement of the segments in \mathfrak{m} , we must have that $b_{q_x} < b_{q_{x+1}}$ and so the violation of the nesting property implies that

$$a_{q_x} < a_{q_{x+1}}.$$

However, this then contradicts that the sequence is in standard order.

As a consequence of Claim 1, we also have that $l_{q_i} = a_{q_{i-1}} - a_{q_i}$.

Let w^* be the fixed element in $(S_{t_1} \times \dots \times S_{t_r}) \setminus S_n / (S_{n-i} \times S_i)$ associated to (13.8) and let w be the element in $(S_{t_1} \times \dots \times S_{t_r}) \setminus S_n / (S_{n-i} \times S_i)$ associated to any layer in standard order in the sense defined above with $w \neq w^*$.

Claim 2: $w \not\leq_B w^$.*

Proof of Claim 2: The strategy is to apply Proposition 13.3. Suppose not to derive a contradiction. Let i^* be the largest integer such that $p_{i^*} = q_{i^*}$. Set $i^* = 0$ if such integer does not exist i.e. $p_1 \neq q_1$.

If $i^* = r$, then q_{i^*+1} is also not defined. Otherwise, by the nesting property shown in Claim 1, we obtain a contradiction to choices of segments in the removal sequence for (Δ, \mathfrak{h}) . This implies that $w^* = w$, giving a contradiction.

Thus $i^* < r$. Now we compare $\Delta_{p_{i^*+1}}$ and $\Delta_{q_{i^*+1}}$. By using Definition 3.1(2) and Claim 1, we must have that

$$a_{p_{i^*+1}} \geq a_{q_{i^*+1}},$$

or q_{i^*+1} is not defined.

We further divide into two subcases.

- $a_{p_{i^*+1}} = a_{q_{i^*+1}}$. But now, we must have that $b_{q_{i^*+1}} \geq b_{p^*+1}$ by the nesting property in Claim 1 again. Then, from our arrangements and choices, $q_{i^*+1} > p_{i^*+1}$. As noted from above that $l_{q_1} = \tilde{l}_{p_1}, \dots, l_{q_{i^*}} = \tilde{l}_{p_{i^*}}$, one checks that

$$w(n - l_1 d - \dots - l_{i^*} d - 1) > w^*(n - l_1 d - \dots - l_{i^*} d - 1)$$

and so $w \not\leq_B w^*$ by Proposition 13.3.

- $a_{p_{i^*+1}} > a_{q_{i^*+1}}$ or q_{i^*+1} is not defined. In such case, $l_{q_{i^*}} > \tilde{l}_{p_{i^*}}$. Hence we also have that:

$$w(n - l_1 d - \dots - l_{i^*} d - 1) > w^*(n - l_1 d - \dots - l_{i^*} d - 1)$$

and so again $w \not\leq_B w^*$ by Proposition 13.3.

(2) Case 2: $\Delta_1^{+,l_1}, \dots, \Delta_r^{+,l_r}$ are not in standard order.

Let $\omega_{l_1, \dots, l_r} = \text{St}(\Delta_1^{-,l_1}) \times \dots \times \text{St}(\Delta_r^{-,l_r})$ and similarly let $\omega_{\tilde{l}_1, \dots, \tilde{l}_r}$.

Claim 3: Let $\tilde{l} = \tilde{l}_1 + \dots + \tilde{l}_r$. For all j ,

$$\text{Ext}_{G_{n-\tilde{l}} \times G_{\tilde{l}}}^j(\omega_{\tilde{l}_1, \dots, \tilde{l}_r} \boxtimes \text{St}(\Delta), \omega_{l_1, \dots, l_r} \boxtimes \text{St}(\Delta_1^{+,l_1}) \times \dots \times \text{St}(\Delta_r^{+,l_r})) = 0$$

Proof of Claim 3: We apply Frobenius reciprocity on the second factor. Then the Jacquet module of $\text{St}(\Delta)$ takes the form

$$\text{St}(\Delta'_1) \boxtimes \dots \boxtimes \text{St}(\Delta'_r)$$

with $\Delta'_1, \dots, \Delta'_r$ in standard order. Since $\Delta_1^{+,l_1}, \dots, \Delta_r^{+,l_r}$ are not in standard order, an argument on comparing cuspidal support gives

$$\text{Ext}_{G_{\tilde{l}}}^j(\text{St}(\Delta), \text{St}(\Delta'_1) \boxtimes \dots \boxtimes \text{St}(\Delta'_r)) = 0$$

for all j . Then Künneth formula then gives the claim.

We now go back to the proof. The element $w^* \in (S_{t_1} \times \dots \times S_{t_r}) \setminus S_n / (S_{n-i} \times S_i)$ is defined as above. For each $w \in (S_{t_1} \times \dots \times S_{t_r}) \setminus S_n / (S_{n-i} \times S_i)$, let $\kappa(w)$ be the associated layer taking the form (*). Note that $\omega_{\tilde{l}_1, \dots, \tilde{l}_r} \cong \tilde{\lambda}(\mathfrak{r}(\Delta, \mathfrak{h}))$.

Let $\kappa_{\leq w^*}$ (resp. $\kappa_{< w^*}$) be the submodule of $\tilde{\lambda}(\mathfrak{h})_{N_l}$ that consists of only the layers associated to $w' \in (S_{t_1} \times \dots \times S_{t_r}) \setminus S_n / (S_{n-i} \times S_i)$ satisfying $w' \leq_B w^*$ (resp. $w' <_B w^*$). We have the short exact sequence:

$$0 \rightarrow \kappa_{< w^*} \rightarrow \kappa_{\leq w^*} \rightarrow \kappa(w^*) \rightarrow 0.$$

On the other hand, there is an embedding

$$\omega_{\tilde{l}_1, \dots, \tilde{l}_r} \boxtimes \text{St}(\Delta) \hookrightarrow \kappa(w^*)$$

Hence, we have a submodule κ' of $\kappa_{\leq w^*}$ admitting a short exact sequence:

$$0 \rightarrow \kappa_{< w^*} \rightarrow \kappa' \rightarrow \omega_{\tilde{l}_1, \dots, \tilde{l}_r} \boxtimes \text{St}(\Delta) \rightarrow 0.$$

However, for $w' < w^*$, we can conclude that

$$\text{Ext}_{G_{n-i} \times G_l}^j((\mathfrak{r}(\Delta, \mathfrak{h})) \boxtimes \text{St}(\Delta), \kappa(w')) = 0$$

for all j . This follows from a standard cuspidal support argument if the G_l -part of $\kappa(w')$ does not have the same cuspidal support as $\text{St}(\Delta)$, and follows from Claim 2 and Claim 3 otherwise. In other words, we have:

$$\kappa' \cong \kappa_{< w^*} \oplus \tilde{\lambda}(\mathfrak{r}(\Delta, \mathfrak{h})) \boxtimes \text{St}(\Delta).$$

This then gives the following desired embedding

$$\tilde{\lambda}(\mathfrak{r}(\Delta, \mathfrak{h})) \boxtimes \text{St}(\Delta) \hookrightarrow \kappa' \hookrightarrow \kappa \hookrightarrow \tilde{\lambda}(\mathfrak{h})_{N_l}.$$

□

13.3. Conjectures. For $n_1 + \dots + n_s = n$, define P_{n_1, \dots, n_s} to be the parabolic subgroup of G_n generated by the matrices $\text{diag}(g_1, \dots, g_s)$ (each $g_i \in G_{n_i}$) and upper triangular matrices. Let N_{n_1, \dots, n_s} be the unipotent radical of P_{n_1, \dots, n_s} .

We end with some conjectures for the embedding model, which are possibly used to interpret some results in this article from representation-theoretic viewpoint:

Conjecture 13.5. *Let $\mathfrak{h} \in \text{Mult}_\rho$. The embedding in Proposition 13.4 is unique.*

We remark that Conjecture 13.5 is not a mere consequence of the multiplicity one theorem for standard representations in [Ch23].

Conjecture 13.6. *Let $\mathfrak{h} \in \text{Mult}_\rho$. Let $\mathfrak{n} \in \text{Mult}_\rho$ be minimal to \mathfrak{h} . Let $l = l_{\text{abs}}(\mathfrak{n})$. Then there exists a unique embedding:*

$$\tilde{\lambda}(\mathfrak{r}(\mathfrak{n}, \mathfrak{h})) \boxtimes \tilde{\lambda}(\mathfrak{n}) \hookrightarrow \tilde{\lambda}(\mathfrak{h})_{N_l}.$$

Proposition 13.4 is a special case of Conjecture 13.6.

Conjecture 13.7. *Let $\pi \in \text{Irr}_\rho$. Let $\mathfrak{n} \in \text{Mult}_\rho$ be minimal to π . Let $\mathfrak{h} = \mathfrak{h}\mathfrak{d}(\pi)$. Let $l_1 = l_{\text{abs}}(\mathfrak{n})$ and let $l_2 = l_{\text{abs}}(\mathfrak{h}\mathfrak{d}(\pi)) - l_{\text{abs}}(\mathfrak{n})$. Suppose Conjectures 11.3 and 13.6 hold. We have the following diagram of maps:*

$$D_{\mathfrak{h}}(\pi) \boxtimes \tilde{\lambda}(\mathfrak{r}(\mathfrak{n}, \pi)) \boxtimes \tilde{\lambda}(\mathfrak{n}) \xrightarrow{\iota_1} D_{\mathfrak{h}}(\pi) \boxtimes \tilde{\lambda}(\mathfrak{h})_{N_{l_2, l_1}} \xrightarrow{\iota_2} \pi_{N_{n-l_1-l_2, l_2, l_1}},$$

$\uparrow \iota_3$

$$D_{\mathfrak{n}}(\pi)_{N_{n-l_1-l_2, l_2}} \boxtimes \tilde{\lambda}(\mathfrak{n})$$

where

- ι_1 is the map indeed from the one in Conjecture 13.6;
- ι_2 is the map induced from the unique embedding $D_{\mathfrak{h}}(\pi) \boxtimes \tilde{\lambda}(\mathfrak{h})$ in Conjecture 11.3
- ι_3 is the map induced from the unique embedding $D_{\mathfrak{n}}(\pi) \boxtimes \tilde{\lambda}(\mathfrak{n}) \hookrightarrow \pi_{n-l_1, l_1}$ in Conjecture 11.3.

Then $\iota_2 \circ \iota_1$ factors through ι_3 .

Part 4. Appendices

14. APPENDIX A: NON-ISOMORPHIC DERIVATIVES

We prove a converse of Proposition 11.1 in this appendix. For a ladder representation or a generic representation σ of G_k , let $I_\sigma(\pi)$ be the unique submodule of $\pi \times \sigma$ (see [LM16], also see [Ch24, Ch25]). See [At24] for a generalization of ladder representations to some other classical groups.

14.1. Non-isomorphic integrals. For two segments Δ_1, Δ_2 in Seg_ρ ,

- (1) Suppose $\Delta_1 < \Delta_2$. Define $\text{St}(\{\Delta_1, \Delta_2\})$ to be the unique irreducible quotient of $\text{St}(\Delta_2) \times \text{St}(\Delta_1)$.
- (2) Suppose Δ_1 and Δ_2 are not linked. Define $\text{St}(\{\Delta_1, \Delta_2\})$ to be the irreducible module $\text{St}(\Delta_1) \times \text{St}(\Delta_2)$.

Proposition 14.1. *Let $\pi \in \text{Irr}_\rho$. Let Δ_1, Δ_2 be two linked segments. Let $\Delta'_1 = \Delta_1 \cup \Delta_2$ and let $\Delta'_2 = \Delta_1 \cap \Delta_2$ (possibly the empty set). Let $\sigma = \text{St}(\{\Delta_1, \Delta_2\})$ and let $\sigma' = \text{St}(\Delta'_1 + \Delta'_2)$. Then $I_\sigma(\pi) \not\cong I_{\sigma'}(\pi)$.*

Proof. We shall use the invariant $\text{mr}(., \Delta_1)$ to distinguish the two representations. Write $\Delta_1 = [a_1, b_1]_\rho$ and $\Delta_2 = [a_2, b_2]_\rho$. Switching labels if necessary, we may and shall assume $a_1 < a_2$. We shall prove by an induction on the sum $l_{abs}(\Delta_1) + l_{abs}(\Delta_2)$. When the sum is 2, the argument is similar to the cases below and we omit the details.

Case 1: $a_1 \neq a_2 - 1$. Let $k = \varepsilon_{[a_1]_\rho}(\sigma)$. We have that $\varepsilon_{[a_1]_\rho}(I_\sigma(\pi)), \varepsilon_{[a_2]_\rho}(I_{\sigma'}(\pi)) = \varepsilon_{[a_1]_\rho}(\pi) + 1$. Let $\kappa = \text{St}(\{-\Delta_1, \Delta_2\})$ and let $\kappa' = \text{St}(\{-\Delta'_1, \Delta'_2\})$. Furthermore,

$$D_{[a_1]_\rho}^{k+1}(I_\sigma(\pi)) = I_\kappa(D_{[a_1]_\rho}^k(\pi)), \quad D_{[a_1]_\rho}^{k+1}(I_{\sigma'}(\pi)) = I_{\kappa'}(D_{[a_1]_\rho}^k(\pi)).$$

Then, by induction, we have that $I_\kappa(D_{[a_1]_\rho}^k(\pi)) \not\cong I_{\kappa'}(D_{[a_1]_\rho}^k(\pi))$ as desired.

Case 2: $a_1 = a_2 - 1$. In this case, let $\mathfrak{m} = \text{mr}(\pi, \Delta_1)$. Then

$$\pi \hookrightarrow D_{\mathfrak{m}}(\pi) \times \text{St}(\mathfrak{m})$$

Now let $\tilde{\sigma} = D_{\Delta'_2}(\sigma)$. We have that

$$\begin{aligned} I_\sigma(\pi) &\hookrightarrow \pi \times \sigma \\ &\hookrightarrow D_{\mathfrak{m}}(\pi) \times \text{St}(\mathfrak{m}) \times \sigma \\ &\cong D_{\mathfrak{m}}(\pi) \times \sigma \times \text{St}(\mathfrak{m}) \\ &\hookrightarrow D_{\mathfrak{m}}(\pi) \times \tilde{\sigma} \times \text{St}(\Delta'_2 + \mathfrak{m}), \end{aligned}$$

where the isomorphism in the third line follows from $\text{St}(\Delta') \times \sigma \cong \sigma \times \text{St}(\Delta')$ for any Δ_1 -saturated segment Δ' (see e.g. [MW86, Lemme II 10.1]).

Since $\text{mr}(D_{\mathfrak{m}}(\pi), \Delta_1) = \emptyset$ and $\text{mr}(\tilde{\sigma}, \Delta_1) = \emptyset$, we have that

$$\text{mr}(I_\sigma(\pi), \Delta_1) = \mathfrak{m} + \Delta'_2$$

The last equality follows from an application on the geometric lemma (see details from the proof of [Ch24, Proposition 11.1]).

Similarly, let $\tilde{\sigma}' = D_{\Delta_1} \circ D_{\Delta_2'}(\sigma')$. We also have that:

$$\begin{aligned} I_{\sigma'}(\pi) &\hookrightarrow \pi \times \sigma' \\ &\hookrightarrow D_{\mathfrak{m}}(\pi) \times \text{St}(\mathfrak{m}) \times \sigma' \\ &\cong D_{\mathfrak{m}}(\pi) \times \sigma' \times \text{St}(\mathfrak{m}) \\ &\hookrightarrow D_{\mathfrak{m}}(\pi) \times \tilde{\sigma}' \times \text{St}(\Delta_1) \times \text{St}(\Delta_2') \times \text{St}(\mathfrak{m}) \\ &\hookrightarrow D_{\mathfrak{m}}(\pi) \times \tilde{\sigma}' \times \text{St}(\mathfrak{m} + \Delta_1 + \Delta_2') \end{aligned}$$

Again, $\mathfrak{m}\mathfrak{x}(D_{\mathfrak{m}}(\pi), \Delta_1) = \emptyset$ and $\mathfrak{m}\mathfrak{x}(\tilde{\sigma}', \Delta_1) = \emptyset$. Thus,

$$\mathfrak{m}\mathfrak{x}(I_{\sigma'}(\pi), \Delta_1) = \mathfrak{m} + \Delta_1 + \Delta_2'.$$

Thus, comparing the invariant $\mathfrak{m}\mathfrak{x}(\cdot, \Delta_1)$, we have $I_{\sigma}(\pi) \not\cong I_{\sigma'}(\pi)$. \square

It is an interesting to investigate if an analogue of Proposition 14.1 can be obtained if one replaces essentially square-integrable representations by other interesting representations such as Speh representations and ladder representations. The composition factors for parabolically induced from Speh representations and ladder representations are studied in [Ta15] and [Gu21], and so it is possible to develop a parallel theory from those via above approach.

14.2. Consequences. We similarly define those notions for derivatives for ladder representations (also see e.g. [Ch22+c]). If there exists $\omega \in \text{Irr}(G_{n-k})$ such that $\omega \boxtimes \sigma \hookrightarrow \pi_{N_k}$ for σ defined in Proposition 14.1, then denote such ω by $D_{\sigma}(\pi)$. Otherwise, set $D_{\sigma}(\pi) = 0$.

Corollary 14.2. *We use the notations in Proposition 14.1. Then $D_{\sigma}(\pi) \not\cong D_{\sigma'}(\pi)$ if both terms are non-zero.*

Proof. Let $\pi' = D_{\sigma}(\pi)$. Then $I_{\sigma}(\pi') \not\cong I_{\sigma'}(\pi')$ and so $\pi \not\cong I_{\sigma'} \circ D_{\sigma}(\pi)$. Applying $D_{\sigma'}$ on both sides, we obtain the corollary. \square

Corollary 14.3. *Let $\Delta_1, \Delta_2 \in \text{Seg}_{\rho}$ such that $\Delta_1 < \Delta_2$. Let $\pi \in \text{Irr}_{\rho}$. Suppose $D_{\Delta_2} \circ D_{\Delta_1}(\pi) \neq 0$. If the non-zero map*

$$D_{\Delta_2} \circ D_{\Delta_1}(\pi) \boxtimes (\text{St}(\Delta_1) \times \text{St}(\Delta_2)) \rightarrow \pi_{N_{l_a(\Delta_1)+l_a(\Delta_2)}}$$

is injective, then $\{\Delta_1, \Delta_2\}$ is minimal to π .

Proof. If the map is injective, then $D_{\Delta_2} \circ D_{\Delta_1}(\pi) \boxtimes \text{St}(\{\Delta_1 + \Delta_2\})$ is a submodule of $\pi_{N_{l_a(\Delta_1)+l_a(\Delta_2)}}$. This implies that

$$D_{\Delta_2} \circ D_{\Delta_1}(\pi) \cong D_{\text{St}(\Delta_1 + \Delta_2)}(\pi).$$

Then the corollary follows from Corollary 14.2. \square

15. APPENDIX B: APPLICATIONS

15.1. Minimality under Δ -reduced condition.

Corollary 15.1. *Let $\pi \in \text{Irr}_{\rho}$. Let $\mathfrak{n} \in \text{Mult}_{\rho}$ be minimal to π . Let Δ be a segment. Suppose $b(\Delta') < b(\Delta)$ for any segment $\Delta' \in \mathfrak{n}$ with $b(\Delta') \neq b(\Delta)$. If $\eta_{\Delta}(D_{\mathfrak{n}}(\pi)) = 0$, then $\mathfrak{m}\mathfrak{x}(\pi, \Delta) \subseteq \mathfrak{n}$.*

Proof. Let $\Delta = [a, b]_{\rho}$. Let $\mathfrak{p} = \mathfrak{m}\mathfrak{x}(\pi, \Delta)$. Let \mathfrak{p}' be all the segments Δ' in \mathfrak{n} such that $b(\Delta') = b$ and $\Delta' \subseteq \Delta$.

We first prove the following claim:

Claim: $|\mathfrak{p}'| \geq |\mathfrak{p}|$.

Proof of Claim: We also let \mathfrak{p}'' be all the segments Δ' in \mathfrak{n} such that $b(\Delta') = b$ and $\Delta \subsetneq \Delta'$. Note that

$$\eta_{\Delta}(D_{\mathfrak{p}''} \circ D_{\mathfrak{n}-\mathfrak{p}''}(\pi)) = \eta_{\Delta}(D_{\mathfrak{n}-\mathfrak{p}''}(\pi)).$$

Hence, we also have $\eta_\Delta(D_{\mathbf{n}-\mathbf{p}''}(\pi)) = 0$. But then, by counting the multiplicity at $b(\Delta)$, we must have that $|\mathbf{p}'| \geq |\mathbf{p}|$. This proves the claim.

If $\mathbf{p} \neq \mathbf{p}'$, then the Claim implies that there exists $a \leq c \leq b$ such that the number of segments $[c, b]_\rho$ in \mathbf{p}' is strictly greater than $\varepsilon_{[c, b]_\rho}(\pi)$. Let c^* be such smallest integer. Now let $\tilde{\mathbf{n}}$ be the submultisegment of \mathbf{n} whose segments Δ' satisfy $a(\Delta') < a$. By the subsequent property, $\tilde{\mathbf{n}} + [c^*, b]_\rho$ and $\tilde{\mathbf{n}} + \mathbf{p}'$ are still minimal to π . Now, by Proposition 4.10 and the minimality of $\tilde{\mathbf{n}} + [c^*, b]_\rho$, $\varepsilon_{[c^*, b]_\rho}(\mathbf{r}(\tilde{\mathbf{n}}, \pi)) = \varepsilon_{[c^*, b]_\rho}(\pi)$. But, by the admissibility of $\tilde{\mathbf{n}} + \mathbf{p}'$,

$$\varepsilon_{[c^*, b]_\rho}(\mathbf{r}(\tilde{\mathbf{n}}, \pi)) = \text{number of segments } [c^*, b]_\rho \text{ in } \mathbf{p}' > \varepsilon_{[c^*, b]_\rho}(\pi).$$

This gives a contradiction. Thus we must have $\mathbf{p}' = \mathbf{p}$. \square

15.2. Generalized reduced decomposition. The notion of reduced decomposition is introduced in [AL25, Section 7] for a segment Δ .

We now describe a generalization of reduced decomposition for multisegments. Let $\pi \in \text{Irr}_\rho(G_n)$. Let \mathbf{n} be minimal to π . Let b be the largest $b(\Delta)$ among all segments Δ in \mathbf{n} . We then choose the longest segment $\Delta_1 \in \mathbf{n}$ such that $b(\Delta_1) = b$. Let $\mathbf{p}_1 = \mathbf{m}\mathbf{r}(D_{\mathbf{n}}(\pi), \Delta)$. We now set $\mathbf{m}'_1 = \mathbf{n} + \mathbf{p}_1$. In general, \mathbf{n}_1 is not minimal to π and so one find the minimal element, denoted by \mathbf{m}_1 , in $\mathcal{S}(\pi, D_{\mathbf{m}_1}(\pi))$. Then, by Corollary 15.1, $\mathbf{m}_1 = \mathbf{q}_1 + \mathbf{n}_2$ for some multisegment \mathbf{n}_2 and $\mathbf{q}_1 = \mathbf{m}\mathbf{r}(\pi, \Delta)$. Thus, from commutativity, we now have that $D_{\mathbf{m}_1}(\pi) \cong D_{\mathbf{n}_2} \circ D_{\mathbf{q}_1}(\pi)$. Thus, one may consider that $\pi \hookrightarrow D_{\mathbf{q}_1}(\pi) \times \text{St}(\mathbf{q}_1)$ is the step for the reduction. One can repeat the same process for \mathbf{n}_2 and then repeat to obtain a sequence of triples $(\mathbf{p}_1, \mathbf{q}_1, \mathbf{n}_1), \dots, (\mathbf{p}_r, \mathbf{q}_r, \mathbf{n}_r)$ until the process terminates. Then we obtain a kind of reduced decompositon for π with respect to \mathbf{n} as follows:

$$\pi \hookrightarrow (D_{\mathbf{q}_r} \circ \dots \circ D_{\mathbf{q}_1})(\pi) \times \text{St}(\mathbf{q}_r) \times \dots \times \text{St}(\mathbf{q}_1).$$

Let $l = l_{abs}(\mathbf{n})$. One may expect there is a natural map:

$$D_{\mathbf{n}}(\pi) \boxtimes \text{St}(\mathbf{n}) \rightarrow (D_{\mathbf{q}_r} \circ \dots \circ D_{\mathbf{q}_1})(\pi) \dot{\times}^1 (\text{St}(\mathbf{q}_r) \times \dots \times \text{St}(\mathbf{q}_1))_{N_l},$$

where $\dot{\times}^1$ is a parabolic induction from a $G_{n_1} \times G_{n_2} \times G_l$ -representation to a $G_{n_1+n_2} \times G_l$ representation. Here $n_1 = n - l_{abs}(\mathbf{q}_1) - \dots - l_{abs}(\mathbf{q}_r)$ and $n_2 = l_{abs}(\mathbf{p}_1) + \dots + l_{abs}(\mathbf{p}_r)$.

15.3. An inductive construction of simple quotients of Bernstein-Zelevinsky derivatives. Let $\pi \in \text{Irr}_\rho$. Let $\Delta \in \text{Seg}_\rho$ and let $\mathbf{n} \in \text{Mult}_\rho$. Let $\mathbf{p} = \mathbf{m}\mathbf{r}(\pi, \Delta)$. Note that $D_{\mathbf{n}} \circ D_{\mathbf{p}}(\pi)$ is a simple quotient of the $l_{abs}(\mathbf{n} + \mathbf{p})$ -th Bernstein-Zelevinsky derivative of $\text{St}(\mathbf{p}) \times D_{\mathbf{p}}(\pi)$ if $\eta_\Delta(D_{\mathbf{n}} \circ D_{\mathbf{p}}(\pi)) = 0$. Then one may ask if $D_{\mathbf{n}} \circ D_{\mathbf{p}}(\pi)$ is also a simple quotient of the $l_{abs}(\mathbf{n} + \mathbf{p})$ -th Bernstein-Zelevinsky derivative of π . In a special case, we have the following criteria using commutativity:

Proposition 15.2. *Let $\pi \in \text{Irr}_\rho$. Let $\Delta \in \text{Seg}_\rho$ and let $\mathbf{n} \in \text{Mult}_\rho$. Let $\mathbf{p} = \mathbf{m}\mathbf{r}(\pi, \Delta)$.*

- (1) *Suppose \mathbf{n} is minimal to $D_{\mathbf{p}}(\pi)$ and $\eta_\Delta(D_{\mathbf{n}} \circ D_{\mathbf{p}}(\pi)) = 0$. Suppose further that for any segment $\Delta' \in \mathbf{n}$, $b(\Delta') < b(\Delta)$. If $D_{\mathbf{n}} \circ D_{\mathbf{p}}(\pi) \cong D_{\mathbf{m}}(\pi)$ for some $\mathbf{m} \in \text{Mult}_\rho$, then $\mathbf{n} + \mathbf{p}$ is minimal to π .*
- (2) *Suppose \mathbf{m} is minimal to π and $\eta_\Delta(D_{\mathbf{m}}(\pi)) = 0$. Suppose further that for any segment $\Delta' \in \mathbf{m}$, $b(\Delta') \leq b(\Delta)$. Then $D_{\mathbf{m}}(\pi) \cong D_{\mathbf{n}} \circ D_{\mathbf{p}}(\pi)$ and $\mathbf{m} = \mathbf{n} + \mathbf{p}$ for some multisegment \mathbf{n} minimal to $D_{\mathbf{p}}(\pi)$.*

Proof. We first consider (1). Suppose $D_{\mathbf{n}} \circ D_{\mathbf{p}}(\pi) \cong D_{\mathbf{m}}(\pi)$ for some $\mathbf{m} \in \text{Mult}_\rho$. Without loss of generality, we may assume that \mathbf{m} is minimal to π . Then, by Corollary 15.1, $\mathbf{m} = \mathbf{p} + \mathbf{n}'$. By the subsequent property and commutativity property (Theorem 9.4), \mathbf{n}' is also minimal to $D_{\mathbf{p}}(\pi)$ and $D_{\mathbf{n}'} \circ D_{\mathbf{p}}(\pi) \cong D_{\mathbf{m}}(\pi)$. By Theorem 1.2, we then have that $\mathbf{n} = \mathbf{n}'$ and this implies (1).

We now consider (2). By Corollary 15.1, $\mathbf{p} \subset \mathbf{m}$ and so $\mathbf{m} = \mathbf{p} + \mathbf{n}$ for some multisegment \mathbf{n} . By the subsequent property and commutativity property (Theorem 9.4), we also have that \mathbf{n} is minimal to $D_{\mathbf{p}}(\pi)$. \square

REFERENCES

- [AL25] A. Aizenbud and E. Lapid, A binary operation on irreducible components of Lusztig's nilpotent varieties I: definition and properties, *Pure and Applied Mathematics Quarterly* (2025), Volume 21 (2025) Number 1, pp. 5-41.
- [AGRS10] A. Aizenbud, D. Gourevitch, S. Rallis and G. Schiffmann, Multiplicity one theorems, *Ann. of Math.* (2) **172** (2010), no. 2, 1407-1434.
- [At24] H. Atobe, An Analogue of Ladder Representations for Classical Groups, *International Mathematics Research Notices*, Volume 2024, Issue 9, May 2024, Pages 7891-7913.
- [Ba08] A. Badulescu, Global Jacquet-Langlands correspondence, multiplicity one and classification of automorphic representations . *Invent. math.* 172, 383-438 (2008). <https://doi.org/10.1007/s00222-007-0104-8>.
- [BZ77] I. N. Bernstein and A. V. Zelevinsky, Induced representations of reductive p-adic groups, I, *Ann. Sci. Ecole Norm. Sup.* **10** (1977), 441-472.
- [BB05] A. Bjorner, F. Brenti, *Combinatorics of Coxeter Groups*, Graduate Texts in Mathematics (2005).
- [Ch21] K.Y. Chan, Homological branching law for $(\mathrm{GL}_{n+1}(F), \mathrm{GL}_n(F))$: projectivity and indecomposability, *Invent. Math.* (2021). <https://doi.org/10.1007/s00222-021-01033-5>.
- [Ch23] K.Y. Chan, Ext-multiplicity theorem for standard representations of $(\mathrm{GL}_{n+1}, \mathrm{GL}_n)$, *Math. Z.* 303, 45 (2023). <https://doi.org/10.1007/s00209-022-03198-y>.
- [Ch24] K.Y. Chan, On the product functor on inner forms of general linear group over a non-Archimedean local field, *Transformation Groups* (2024).
- [Ch22+b] K.Y. Chan, Quotient branching law for p -adic $(\mathrm{GL}_{n+1}, \mathrm{GL}_n)$ I: generalized Gan-Gross-Prasad relevant pairs, *arXiv:2212.05919* (2022).
- [Ch22+c] K.Y. Chan, Duality for the generalized Gan-Gross-Prasad relevant pairs for p -adic GL , *arXiv:2210.17249* (2022).
- [Ch25] K.Y. Chan, Construction of simple quotients of Bernstein-Zelevinsky derivatives and highest derivative multisegments I: reduction to combinatorics, *Transactions of the AMS (Series B)* (2025).
- [Ch25b] K.Y. Chan, Construction of simple quotients of Bernstein-Zelevinsky derivatives and highest derivative multisegments II: minimal sequences, *Manuscripta Mathematica* (2025).
- [CP25+] K. Y. Chan and B. Pattanayak, Algorithms for parabolic inductions and Jacquet modules in GL_n , *arXiv:2503.00886*, 2025.
- [CS19] K.Y. Chan and G. Savin, Bernstein-Zelevinsky derivatives: a Hecke algebra approach, *International Mathematics Research Notices*, Volume 2019, Issue 3, February 2019, Pages 731-760, <https://doi.org/10.1093/imrn/rnx138>.
- [CZ26+] K.Y. Chan and P. Zhou, Construction of simple quotients of Bernstein-Zelevinsky derivatives and highest derivative multisegments IV: removal processes, in preparation.
- [CMBO24] D. Ciubotaru, L. Mason-Brown and E. Okada, The wavefront sets of unipotent supercuspidal representations, *Algebra and Number Theory*, Vol. 18 (2024), No. 10, 1863-1889.
- [CP96] V. Chari and A. Pressley, Quantum affine algebras and affine Hecke algebras, *Pacific J. Math.* 174 (1996), no. 2, 295-326.
- [Ch86] I.V. Cherednik, A new interpretation of Gelfand-Tzetlin bases, *Duke Math. J.* (1987), Vol.54 (2), p.563-577.
- [CR24] C. Cunningham, M. Ray, Proof of Vogan's conjecture on Arthur packets: irreducible parameters of p -adic general linear groups. *manuscripta math.* 173, 1073-1097 (2024). <https://doi.org/10.1007/s00229-023-01490-7>.
- [Gu21] M. Gurevich, Quantum invariants for decomposition problems in type A rings of representations, *Journal of Combinatorial Theory, Series A*, **180** (2021), <https://doi.org/10.1016/j.jcta.2021.105431>.
- [Ja07] C. Jantzen, Jacquet modules of p -adic general linear groups, *Represent. Theory* **11** (2007), 45-83 <https://doi.org/10/1090/S1088-4165-0700316-0>.
- [LM16] E. Lapid, A. Minguez, On parabolic induction on inner forms of the general linear group over a non-Archimedean local field, *Sel. Math. New Ser.* (2016) 22, 2347-2400.
- [LM25] E. Lapid, Minguez, A binary operation on irreducible components of Lusztig's nilpotent varieties II: applications and conjectures for representations of GL_n over a non-archimedean local field, *Pure and Applied Mathematics Quarterly* (2025).
- [MW86] C. Moeglin, J.-L. Waldspurger, Sur l'involution de Zelevinski. *J. Reine Angew. Math.* 372, 136-177 (1986), doi.org/10.1515/crll.1986.372.136.
- [Mu03] F. Murnaghan, Local character expansions of admissible representations of p -adic general linear groups. In: *Journal für die reine und angewandte Mathematik* 2003.554 (2003), pp. 139-155. doi:10.1515/crll.2003.002.
- [Ro86] J.D. Rogawski, Representations of $GL(n)$ over a p -adic field with an Iwahori-fixed vector, *Israel J. Math.* **54** (1986), 242-256. <https://doi.org/10.1007/BF02764944>.
- [Ta15] M. Tadić: On the reducibility points beyond the ends of complementary series of p -adic general linear groups, *J. Lie Theory* (2015) 25(1), 147-183.

[Ze80] A. Zelevinsky, Induced representations of reductive p -adic groups II, Ann. Sci. Ecole Norm. Sup. **13** (1980), 154-210.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF HONG KONG
Email address: `kychan1@hku.hk`