

# SYMBOLIC INTEGRATION OF DIFFERENTIAL FORMS: FROM ABEL TO ZEILBERGER

SHAOSHI CHEN, DAVID A. COX, AND YISEN WANG

**ABSTRACT.** This paper focuses on symbolic integration of differential forms, with a particular emphasis on historical and modern developments, from Abel's addition theorems for Abelian integrals to Zeilberger's creative telescoping for parameterized integrals. It explores closed rational  $p$ -forms and provides algorithmic approaches for their integration, extending classical results like Hermite reduction and Liouville's theorem. The integration of closed differential forms with parameters is further examined through telescopers, offering a unified framework for handling both algebraic and transcendental cases.

## 1. INTRODUCTION

Differential forms play a central role in mathematics in both theoretical and computational aspects. They provide an elegant language to unify different kinds of integrals in multivariable calculus and are used in differential geometry [50], algebraic topology [11], and algebraic geometry [33] whenever integrals are involved. Symbolic integration of univariate functions has been well-developed starting with the celebrated work by Risch [48] and summarized comprehensively in Bronstein's book [12] and the foundational references collected in [47]. A long-standing challenging project in symbolic computation is developing symbolic algorithms for computing integrals of multivariate functions. In the multivariate setting, differential forms provide a natural language for studying integration problems. Symbolic computation of differential forms has been studied in computational algebraic geometry [13, 14] with applications to computing periods in [39]. Packages in computer algebra systems such as Maple and Mathematica rely on a variety of heuristic techniques to integrate differential forms. The goal of this paper is to enrich the algorithmic methods available for symbolic integration of differential forms, motivated by Abel's classical addition theorem on integrals of algebraic functions.

In 1826, Abel submitted a remarkable memoir [1] to the Paris Academy, who did not publish it until 1841, long after his death. His main focus is on what we now call *Abelian integrals*, which are integrals of the form

$$\int f(x, y) dx,$$

where  $x$  and  $y$  are related by a polynomial equation  $\chi(x, y) = 0$ . Abel considers an auxiliary equation  $\theta(x, y) = 0$  that depends on parameters  $a, a', a'', \dots$ , so that the

---

2010 *Mathematics Subject Classification.* 16S32, 35G35, 68W30.

*Key words and phrases.* Abel's theorem, differential forms, integration, telescopers.

S. Chen and Y. Wang were partially supported by National Key R&D Programs of China (No. 2023YFA1009401), the NSFC grants (No. 12271511), the CAS Funds of the Youth Innovation Promotion Association (No. Y2022001), and the Strategic Priority Research Program of the Chinese Academy of Sciences (No. XDB0510201).

solutions of the simultaneous equations

$$\chi(x, y) = \theta(x, y) = 0$$

are algebraic functions of the parameters. Abel focuses on the nonconstant solutions and writes them as  $(x_i, y_i)$ ,  $i = 1, \dots, \mu$ . In Section 2 of [1], Abel states the following theorem:

I say that if we denote by  $f(x, y)$  some rational function of  $x$  and  $y$ , and if we set

$$dv = f(x_1, y_1) dx_1 + f(x_2, y_2) dx_2 + \dots + f(x_\mu, y_\mu) dx_\mu,$$

the differential  $dv$  will be a rational function of the quantities  $a, a', a''$ , etc.

This is one of several results in [1] that go under the name “Abel’s Theorem” (see [37]). Abel’s first proof of the rationality of  $dv$  is very terse. A more detailed argument is given in his last paper [2], and modern explanations can be found in [27], [31, Ch. 9], and [35]. In Section 4 of [1], Abel notes that this method is “in general very long, and for functions that are a little complicated, almost impractical.” He then gives a direct proof of the rationality of  $dv$  by expressing it as an explicit rational differential of the parameters. His astonishing formula is described in [27].

Later in Section 2 of [1], Abel notes that:

If now  $dv$  is a rational function of the quantities  $a, a', a'' \dots$  its integral or the quantity  $v$  will be an algebraic and logarithmic function of  $a, a', a'' \dots$ <sup>1</sup>

For us, this is where things get interesting. In going from  $dv$  to  $v$ , Abel is clearly referring to the method of partial fractions, which is where the logarithms come from. But this method applies to rational 1-forms in one variable, while here,  $dv$  is a rational 1-form in the parameters  $a, a', a'', \dots$ . What’s missing is the recognition that  $dv$  is *closed* (which is easy to prove) and that one needs a result like Theorem 3.1 to guarantee that  $v$  is a rational and logarithmic function of the parameters. Abel avoids this difficulty since his explicit formula for  $dv$  leads immediately to an explicit formula for  $v$ .

In contrast, we show in Theorem 3.1 that this is the case for *any* closed rational 1-form. A natural question is whether Theorem 3.1 still holds for closed rational  $p$ -forms. This is answered affirmatively by Theorem 4.3 in Section 4.

Liouville’s classical theorem is crucial for determining whether an elementary function has an elementary integral. The first step towards symbolic integration of differential forms with elementary-function coefficients would be establishing Liouville’s theorem in this setting. In this direction, the first work was done by Caviness and Rothstein [16] who proved a version of Liouville’s theorem for integration in finite terms of line integrals. Using the techniques introduced in Section 3, we provide a more direct proof in the language of differential 1-forms. It is still challenging to extend this theorem to the case of  $p$ -forms.

---

<sup>1</sup>While Abel says “algebraic and logarithmic” in his result, Theorem 3.1 shows that “algebraic” can be replaced with “rational” (this is also proved in [27]). Abel’s preference for “algebraic” is related to a change made in Section 6 of [1]. If  $\alpha$  is the number of parameters  $a, a', a'', \dots$  and  $(x_i, y_i)$ ,  $i = 1, \dots, \mu$ , are the solutions considered above, then Abel claims that  $a, a', a'', \dots$  are algebraic functions of the first  $\alpha$  of  $x_1, \dots, x_\mu$ . This makes  $v$  an algebraic and logarithmic function of  $x_1, \dots, x_\alpha$ .

Symbolic computation of parameterized integrals is an active topic [46] with rich applications to Gauss–Manin connections and Picard–Fuchs equations in algebraic geometry [41, 39] and Feynmann diagrams in mathematical physics [42, 4]. A powerful tool for evaluating parameterized integrals is Zeilberger’s method of creative telescoping [3, 56]. The formulation of creative telescoping can be generalized from functions to differential forms. Given a differential form  $\omega$  in  $x_1, \dots, x_m$  with a parameter  $t$ , a linear differential operator  $L$  with respect to  $t$  is called a *telescopers* for  $\omega$  if  $L(\omega)$  is exact. The existence problem of telescopers was first studied by Manin in [41] for the case of  $m$ -forms with algebraic functions as coefficients and recently was investigated in [19, 20] for the case of differential forms with D-finite functions as coefficients. We will present an algorithm that uses Hermite reduction to compute minimal telescopers of rational differential 1-forms with one parameter.

The outline of the paper is as follows. After establishing some notation and preliminary lemmas in Section 2, we generalize the 1-forms  $dv$  studied by Abel to the case of closed rational 1-forms and develop the corresponding Hermite reduction in Section 3. The case of closed rational  $p$ -forms is considered in Section 4. In Section 5, we recall Picard’s problem about the exactness of differential 3-forms and Griffiths–Dwork reduction for solving the smooth case together with an example and a conjecture related to Fermat hypersurfaces. In Section 6, we generalize Liouville’s theorem to differential 1-forms. We conclude the paper by studying Zeilberger’s creative telescoping for closed rational 1-forms in Section 7.

## 2. PRELIMINARIES

In this section, we first recall some basic facts about differential fields and their extensions [12]. We also discuss differential forms.

**Definition 2.1.** Let  $R$  be a ring (resp. field). A *derivation* on  $R$  is a map  $D : R \rightarrow R$  such that for any  $a, b \in R$ , we have  $D(a + b) = D(a) + D(b)$  and  $D(ab) = aD(b) + bD(a)$ . The pair  $(R, D)$  is called a *differential ring* (resp. *differential field*). The set  $\text{Const}_D(R) := \{r \in R \mid D(r) = 0\}$  is a subring (resp. subfield) of  $R$ . A pair  $(R^*, D^*)$  is called a *differential extension* of  $(R, D)$  if  $R \subseteq R^*$  and  $D^*|_R = D$ .

The following fact is used frequently in symbolic integration [12].

**Lemma 2.2.** For a field  $F$  of characteristic zero, let  $F(y)$  be the field of rational functions in  $y$  over  $F$ . Let  $D_y$  denote the derivative  $d/dy$  on  $F(y)$  satisfying  $D_y(y) = 1$  and  $D_y(c) = 0$  for all  $c \in F$ . Given pairwise coprime polynomials  $u_1, \dots, u_n \in F[y] \setminus F$ , constants  $c_1, \dots, c_n \in F$ , and a rational function  $v \in F(y)$ , suppose that

$$\sum_{i=1}^n c_i \frac{D_y(u_i)}{u_i} + D_y(v) = 0.$$

Then  $c_1 = \dots = c_n = 0$  and  $v \in F$ .

*Proof.* Fix  $i$  and let  $\beta$  be a root of  $u_i$  of multiplicity  $\ell$  in some differential extension of  $F$ . By hypothesis,  $u_j(\beta) \neq 0$  for  $j \neq i$ . A standard calculation shows that

$$\text{res}_{y=\beta} \left( \frac{D_y(u_i)}{u_i} \right) = \ell \neq 0.$$

The residue of the derivative of a rational function always vanishes, so taking the residue at  $y = \beta$  of each side of the equation in the lemma implies  $c_i \ell = 0$ , and  $c_1 = \dots = c_n = 0$  follows. Then  $D_y(v) = 0$ , which implies  $v \in F$ .  $\square$

For a field  $k$  of characteristic zero, let  $K := k(x_1, \dots, x_m)$  be the field of rational functions in  $x_1, \dots, x_m$  over  $k$ . Let  $\partial_i$  denote the usual partial derivative  $\partial/\partial x_i$  that satisfies  $\partial_i \partial_j(f) = \partial_j \partial_i(f)$  for all  $f \in K$  and  $\partial_i(c) = 0$  for all  $c \in k(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_m)$ . Let  $\mathcal{U}$  be the universal differential extension of  $K$  in which any system of algebraic differential equations has a solution in  $\mathcal{U}$  if it has a solution in some extension of  $K$  (see [38, Chapter III, Section 7] for the existence of such universal fields). Note that the  $\partial_i$ 's also commute on  $\mathcal{U}$ .

We now recall some basic notions about exterior algebras over  $\mathcal{U}$  from Lang's book [40, Chapter XIX]. Given a vector space  $\mathcal{M}$  over  $\mathcal{U}$  of dimension  $m$  with basis  $\{\mathbf{a}_1, \dots, \mathbf{a}_m\}$ , let

$$\bigwedge \mathcal{M} = \bigoplus_{p=0}^m \bigwedge^p \mathcal{M}$$

be the exterior algebra of  $\mathcal{M}$ , where  $\bigwedge^p \mathcal{M}$  denotes the vector space over  $\mathcal{U}$  with basis  $\{\mathbf{a}_{i_1} \wedge \dots \wedge \mathbf{a}_{i_p} \mid 1 \leq i_1 < \dots < i_p \leq m\}$ . We call an element in  $\bigwedge^p \mathcal{M}$  a  $p$ -form, and an element  $f \in \mathcal{U}$  is called a 0-form. Let  $d: \mathcal{U} \rightarrow \mathcal{M}$  be the map defined by

$$df = \partial_1(f) \mathbf{a}_1 + \dots + \partial_m(f) \mathbf{a}_m.$$

Hence  $dx_i = \mathbf{a}_i$  by taking  $f = x_i$ . For the remainder of this paper, we will use  $\{dx_1, \dots, dx_m\}$  instead of  $\{\mathbf{a}_1, \dots, \mathbf{a}_m\}$ . We call  $\mathcal{M}$  the *space of differential 1-forms*.

We recall some properties of operations on differential forms from [54, 15]. The exterior product  $\omega \wedge \eta$  is bilinear, anticommutative, and associative as follows:

$$\begin{aligned} f_I dx_I \wedge g_J dx_J &= f_I g_J dx_I \wedge dx_J \\ (f_I dx_I + f'_{I'} dx_{I'}) \wedge g_J dx_J &= f_I g_J dx_I dx_J + f'_{I'} g_J dx_{I'} dx_J \\ f_I dx_I \wedge g_J dx_J &= (-1)^{|I| \cdot |J|} g_J dx_J \wedge f_I dx_I \\ (f_I dx_I \wedge g_J dx_J) \wedge h_R dx_R &= f_I dx_I \wedge (g_J dx_J \wedge h_R dx_R). \end{aligned}$$

Here,  $dx_I$  stands for  $dx_{i_1} \wedge \dots \wedge dx_{i_p}$ , where  $I = \{i_1, \dots, i_p\}$ , often written  $dx_I = dx_{i_1} \cdots dx_{i_p}$ . If  $dx_I$  is any string with some  $dx_i$  appearing more than once, then  $dx_I = 0$ . If  $i_1, \dots, i_k$  are distinct and  $\sigma: \{i_1, \dots, i_k\} \rightarrow \{i_1, \dots, i_k\}$  is a permutation, then

$$dx_{\sigma(i_1)} \cdots dx_{\sigma(i_k)} = \text{sign}(\sigma) dx_{i_1} \cdots dx_{i_k},$$

where  $\text{sign}(\sigma) = \pm 1$  is the sign of the permutation. The map  $d$  can be extended to a derivation on  $\bigwedge \mathcal{M}$  defined recursively by

$$d(\omega_1 \wedge \omega_2) = d\omega_1 \wedge \omega_2 + (-1)^p \omega_1 \wedge d\omega_2,$$

for any  $\omega_1 \in \bigwedge^p \mathcal{M}$  and  $\omega_2 \in \bigwedge^q \mathcal{M}$ . The difference between differential forms in  $\mathcal{M}$  and Kähler differentials is well explained in [32].

**Definition 2.3.** We say that  $\omega \in \bigwedge \mathcal{M}$  is *closed* if  $d\omega = 0$  and  $\omega \in \bigwedge \mathcal{M}$  is *exact* if there exists  $\eta \in \bigwedge \mathcal{M}$  such that  $d\eta = \omega$ .

Since  $d(d\eta) = 0$ , it follows that every exact differential form is closed.

We now define operators  $d_s$  and  $d^s$  that will be used in our recursive arguments. For any  $f \in E$  and  $s \in \{1, 2, \dots, m\}$ , define

$$d^s(f dx_I) = \partial_s(f) dx_s \wedge dx_I = \partial_s(f) dx_s dx_I.$$

The definition of  $d^s$  extends linearly to all of  $\bigwedge \mathcal{M}$ . Then define

$$d_s = d^1 + \dots + d^s,$$

and note that

$$d = d^1 + \cdots + d^m = d_{m-1} + d^m.$$

Given  $\omega \in \bigwedge^p M$ , we can always decompose  $\omega$  into

$$\omega = \omega_{m-1} + \omega^m,$$

where all summands  $f_{i_1, \dots, i_p} dx_{i_1} \cdots dx_{i_p}$  of  $\omega_{m-1}$  do not involve  $dx_m$  and  $\omega^m = \mu dx_m$  with  $\mu \in \bigwedge^{p-1} M$  and free of  $dx_m$ . The following lemma is an easy consequence of the above definitions.

**Lemma 2.4.** *If  $\omega = \omega_{m-1} + \omega^m \in \bigwedge^p M$  as above, then we have*

- (i)  $d^m(\omega^m) = 0$ .
- (ii)  $d(\omega dx_m) = d_{m-1}(\omega) dx_m$ .
- (iii)  $d\omega = 0$  if and only if  $d_{m-1}(\omega_{m-1}) = 0$  and  $d^m(\omega_{m-1}) + d_{m-1}(\omega^m) = 0$ .
- (iv) If  $\omega$  is free of  $dx_m$ , then  $d\omega = 0$  if and only if  $d_{m-1}(\omega) = 0$  and  $d^m(\omega) = 0$ .
- (v) If  $\omega$  is free of  $x_m$  and  $dx_m$ , then  $\omega = d\mu$  for some  $(p-1)$ -form  $\mu$  if and only if  $\omega = d_{m-1}(\tilde{\mu})$  for some  $(p-1)$ -form  $\tilde{\mu}$  also free of  $x_m$  and  $dx_m$ .

### 3. INTEGRATION OF CLOSED RATIONAL 1-FORMS

We first study the integration of closed 1-forms with coefficients in  $K = F(x_m)$ , where  $F = k(x_1, \dots, x_{m-1})$  and  $k$  is algebraically closed. Assume that  $\omega = f_1 dx_1 + \cdots + f_m dx_m$ ,  $f_1, \dots, f_m \in K$ , is a closed 1-form. Let  $\bar{F}$  denote the algebraic closure of  $F$ . By the classical integration formula for rational functions in the variable  $x_m$  (i.e., partial fractions with respect to  $x_m$ ), we have

$$(3.1) \quad f_m = \partial_m(g_m) \quad \text{with} \quad g_m = a_m + \sum_j c_{m,j} \log b_{m,j},$$

where  $a_m \in K$ ,  $c_{m,j} \in \bar{F}$ , and  $b_{m,j} \in F(c_{m,j})[x_m]$  are pairwise coprime polynomials of positive degree. Since  $\omega$  is closed, we have  $\partial_i(f_m) = \partial_m(f_i)$  for  $i = 1, \dots, m-1$ . Thus

$$(3.2) \quad \begin{aligned} \partial_m(f_i) &= \partial_i(f_m) = \partial_i(\partial_m(g_m)) = \partial_m(\partial_i(g_m)) \\ &= \partial_m \left( \partial_i(a_m) + \sum_j \left( c_{m,j} \frac{\partial_i(b_{m,j})}{b_{m,j}} + \partial_i(c_{m,j}) \log b_{m,j} \right) \right) \\ &= \partial_m \left( \partial_i(a_m) + \sum_j c_{m,j} \frac{\partial_i(b_{m,j})}{b_{m,j}} \right) + \sum_j \partial_i(c_{m,j}) \frac{\partial_m(b_{m,j})}{b_{m,j}}, \end{aligned}$$

where the last line follows because  $c_{m,j} \in \bar{F}$  implies that  $\partial_m(\partial_i(c_{m,j})) = 0$ .

For fixed  $j$ , Lemma 2.2 implies that  $\partial_i(c_{m,j}) = 0$  for all  $i = 1, \dots, m-1$ . This means that each  $c_{m,j}$  is a constant in  $k$  since  $k$  is algebraically closed. Let  $\tilde{\omega} = \omega - dg_m$ . Then  $\tilde{\omega} = \tilde{f}_1 dx_1 + \cdots + \tilde{f}_{m-1} dx_{m-1}$ , where for each  $i = 1, \dots, m-1$ ,

$$\tilde{f}_i = f_i - \partial_i(g_m) = f_i - \partial_i(a_m) - \sum_j c_{m,j} \frac{\partial_i(b_{m,j})}{b_{m,j}} \in K.$$

Since  $\omega$  is closed, so is  $\tilde{\omega}$ , which implies that

$$0 = d\tilde{\omega} = \sum_{i=1}^{m-1} d\tilde{f}_i \wedge dx_i = \sum_{i=1}^{m-1} \sum_{j=1}^{m-1} \partial_j(\tilde{f}_i) dx_j dx_i + \sum_{i=1}^{m-1} \partial_m(\tilde{f}_i) dx_m dx_i.$$

Thus  $\partial_m(\tilde{f}_i) = 0$  for  $i = 1, \dots, m-1$ . So  $\tilde{\omega}$  is a differential form that is free of  $dx_m$  and has coefficients in  $F = k(x_1, \dots, x_{m-1})$ . Repeating this process recursively, we obtain the following result.

**Theorem 3.1.** *Let  $\omega = f_1 dx_1 + \dots + f_m dx_m$  be a closed 1-form with  $f_i \in K$ . Then  $\omega = dg$  for some  $g$  of the form*

$$g = a + \sum_j c_j \log b_j,$$

where  $a \in K$ ,  $c_j \in k$ , and  $b_j \in F[x_m]$  are pairwise coprime polynomials of positive degree.

**Remark 3.2.** If the field  $k$  is not algebraically closed, then the above constants  $c_j$  are in  $k$  and  $b_j \in F(c_j)[x_m]$ . Note also that a version of Theorem 3.1 appears as Theorem 8 in [24]. In [35, (1.5)], Griffiths states a version of Theorem 3.1 and provides a proof in the case of two variables.

Ostrogradsky [43] and Hermite [36] introduced a reduction method, now called *Hermite reduction*, that decomposes  $f \in k(x)$  as

$$f = \frac{dg}{dx} + \frac{a}{b},$$

where  $g \in k(x)$ ,  $a, b \in k[x]$  are such that  $\gcd(a, b) = 1$ ,  $\deg(a) < \deg(b)$  and  $b$  is squarefree. Moreover,  $f$  is integrable in  $k(x)$  if and only if  $a = 0$ . Hermite reduction separates the rational part of the integral of a rational function from the logarithmic part that is transcendental over  $k(x)$ . This reduction has been extended to algebraic functions [51, 22], hyperexponential functions [6], and D-finite functions [8, 52, 18].

Based on the proof of Theorem 3.1, we now extend Hermite reduction from rational functions to closed rational 1-forms. We begin with a statement of the classical Hermite reduction algorithm.

**Algorithm HermiteRat:** Given a field  $k$  and  $A, D \in k[x]$  with  $D$  nonzero and coprime with  $A$ , return  $g, h \in k(x)$  such that

$$\frac{A}{D} = \frac{dg}{dx} + h,$$

where  $h$  has a squarefree denominator of degree greater than its numerator. Readers can refer to [36] and Section 2.2 in [12] for more details.

We summarize the generalized Hermite reduction algorithm as follows.

**Algorithm HermiteOneForm:** Given a closed 1-form  $\omega = f_1 dx_1 + \dots + f_m dx_m$  with  $f_i \in K = k(x_1, \dots, x_m)$ , return  $g \in K$  and a closed rational 1-form  $\tilde{\omega}$  such that

$$\omega = dg + \tilde{\omega},$$

where  $\tilde{\omega} = \tilde{f}_1 dx_1 + \dots + \tilde{f}_m dx_m$  and each  $\tilde{f}_i$  has a squarefree denominator as a multivariate polynomial in  $k[x_1, \dots, x_m]$ .

When  $m = 1$ , just apply **HermiteRat** to  $f_1$  and multiply the result by  $dx_1$ . When  $m > 1$ , first apply **HermiteRat** to  $f_m$  with respect to  $x_m$  to obtain

$$f_m = \partial_m(a_m) + \bar{f}_m, \quad \text{where } a_m \in K.$$

Note that the above  $a_m$  is the same  $a_m$  that appears in (3.1). This implies that

$$\bar{f}_m = \sum_j c_{m,j} \frac{\partial_m(b_{m,j})}{b_{m,j}},$$

where  $c_{m,j} \in k$  and  $b_{m,j} \in F[x_m]$  are as in (3.1). Then define

$$\tilde{r}_m := \sum_j c_{m,j} \frac{\partial_1(b_{m,j})}{b_{m,j}} dx_1 + \cdots + \sum_j c_{m,j} \frac{\partial_m(b_{m,j})}{b_{m,j}} dx_m.$$

Here  $\sum_j c_{m,j} \partial_i(b_{m,j})/b_{m,j}$  has a squarefree denominator in  $k[x_1, \dots, x_m]$  by normalization, and it is easy to verify that  $\tilde{r}_m$  is a closed 1-form. Thus  $\omega - da_m - \tilde{r}_m$  is a closed 1-form and free of  $x_m$  and  $dx_m$ . By induction,

$$\omega - da_m - \tilde{r}_m = dg_{m-1} + \tilde{\omega}_{m-1},$$

where  $g_{m-1} \in K$  and  $\tilde{\omega}_{m-1} = \tilde{f}_{m-1,1} dx_1 + \cdots + \tilde{f}_{m-1,m-1} dx_{m-1}$ , and each  $\tilde{f}_{m-1,i}$  has a squarefree denominator. Setting  $g := a_m + g_{m-1}$  and  $\tilde{\omega} := \tilde{r}_m + \tilde{\omega}_{m-1}$  gives the desired reduction  $\omega = dg + \tilde{\omega}$ .

**Remark 3.3.** Let  $\omega = dg + \tilde{\omega} \in \mathcal{M}$  be given from Algorithm `HermiteOneForm`. Then  $\omega$  is exact if and only if  $\tilde{\omega} = 0$ .

**Example 3.4.** Let

$$\omega = \frac{txyz - 1}{x^2yz} dx + \frac{txyz - 1}{xy^2z} dy + \frac{t^2xyz + xyz - 1}{xyz^2} dz.$$

If we let  $g := \frac{1}{xyz}$  and

$$\tilde{\omega} := \frac{t}{x} dx + \frac{t}{y} dy + \frac{t^2 + 1}{z} dz,$$

then  $\omega = dg + \tilde{\omega}$  is the result of Hermite reduction on  $\omega$ .

Example 7.3 below will revisit this example from the point of view of creative telescoping.

#### 4. INTEGRATION OF CLOSED RATIONAL $p$ -FORMS

In this section, we will extend Theorem 3.1 to the case of rational closed  $p$ -forms in  $m$  variables. The strategy is to reduce the problem from differential forms in  $m$  variables to forms in  $m - 1$  or fewer variables and then proceed recursively. The result for  $p$ -forms differs from the case of 1-forms because the coefficients of the logarithmic terms can be more complicated. We continue to use the notation  $K = F(x_m)$  with  $F = k(x_1, \dots, x_{m-1})$ , where  $k$  is now any field of characteristic 0.

We first give an example to show that the factors in front of the logarithms may fail to be constants when considering rational closed  $p$ -forms with  $p > 1$ .

**Example 4.1.** Let  $K = \mathbb{C}(x, y)$  and  $\omega = x^{-1}y^{-1} dx dy$ . Suppose that we can write  $\omega = d(A dy - B dx)$  with

$$A = a_0 + \sum_{i>0} \lambda_i \log a_i \quad \text{and} \quad B = b_0 + \sum_{i>0} \mu_i \log b_i,$$

where  $a_i, b_i \in \mathbb{C}(x, y)$  and  $\lambda_i, \mu_i$  are constants in  $\mathbb{C}$ . Then  $x^{-1}y^{-1} = \partial_x(A) + \partial_y(B)$ . If we view  $x^{-1}y^{-1}$ ,  $\partial_x(A)$  and  $\partial_y(B)$  as rational functions in  $y$  with coefficients in  $\overline{\mathbb{C}(x)}$ , then by arguments similar to the proof of Lemma 2.2, taking the residue at  $y = 0$  of both sides of  $x^{-1}y^{-1} = \partial_x(A) + \partial_y(B)$  leads to

$$\frac{1}{x} = \partial_x(\text{res}_{y=0}(a_0)) + c \quad \text{for some constant } c \in \mathbb{C},$$

which is impossible by Lemma 2.2. However, if we no longer require that the factors in front of the logarithms be constants, then we can write

$$\omega = d\left(\frac{1}{y} \log(x) dy\right) = d\left(-\frac{1}{x} \log(y) dx\right).$$

Before beginning our study of general  $p$ -forms, we need some preliminary work. Any rational function  $f \in \overline{F}(x_m)$ ,  $F = k(x_1, \dots, x_{m-1})$  can be written as

$$(4.1) \quad f = c_0 + c_1 x_m + \dots + c_\ell x_m^\ell + \frac{p}{q},$$

where  $c_i \in \overline{F}$ ,  $\ell \geq 0$  and  $p, q \in \overline{F}[x_m]$  with  $\deg_{x_m}(p) < \deg_{x_m}(q)$ . While there are many choices for  $p$  and  $q$ , the polynomial part  $c_0 + c_1 x_m + \dots + c_\ell x_m^\ell$  is unique, as can be seen from the series expansion of  $f$  in descending powers of  $x_m$ . We call  $c_0$  the *constant part* of  $f$ .

**Lemma 4.2.** *If  $f \in \overline{F}(x_m)$  and  $\partial_m(f) \in K = F(x_m)$ , then  $f = g + c_0$  where  $g \in K$  and  $c_0 \in \overline{F}$  is the constant part of  $f$ . Thus*

$$f \in K \iff c_0 \in F.$$

*Proof.* Let  $E \subseteq \overline{F}(x_m)$  be the minimal Galois extension of  $K$  that contains  $f$  and set  $d := [E : K] < \infty$ . If  $G$  is the Galois group of  $E$  over  $K$ , then the trace map  $\text{Tr}_{E/K} : E \rightarrow K$  is defined by  $\text{Tr}_{E/K}(a) = \sum_{\sigma \in G} \sigma(a)$  for  $a \in E$ . According to Theorem 3.2.4 in [12], each  $\sigma \in G$  commutes with  $\partial_i$  for all  $i \in \{1, \dots, m\}$ .

If  $\partial_m(f) = h \in K$ , then taking the trace yields

$$\partial_m(\text{Tr}_{E/K}(f)) = \text{Tr}_{E/K}(\partial_m(f)) = \text{Tr}_{E/K}(h) = d \cdot h = d \cdot \partial_m(f),$$

which implies that  $\partial_m(f - 1/d \cdot \text{Tr}_{E/K}(f)) = 0$ . Since  $\text{Const}_{\partial_m}(E) \subseteq \overline{F}$ ,  $f$  can be written as

$$f = \frac{1}{d} \text{Tr}_{E/K}(f) + c, \text{ for some } c \in \overline{F}.$$

Note that  $\tilde{g} := 1/d \cdot \text{Tr}_{E/K}(f)$  is in  $K$ . Thus  $f = \tilde{g} + c$ .

To bring  $c_0$  into the picture, note that writing  $f$  as in (4.1) gives  $\tilde{g} = f - c = c_0 - c + c_1 x_m + \dots + c_\ell x_m^\ell + \frac{p}{q}$ , which shows that  $c_0 - c$  is the constant part of  $\tilde{g}$ . Then  $\tilde{g} \in K$  implies  $c_0 - c \in F$ , so that  $g := \tilde{g} + c - c_0 \in K$  satisfies

$$g + c_0 = \tilde{g} + c = f.$$

The final assertion of the lemma follows immediately, and the proof is complete.  $\square$

Now let  $\omega$  be a closed  $p$ -form with coefficients in  $K = k(x_1, \dots, x_m) = F(x_m)$ . To explore its integration, write  $\omega = \omega_{m-1} + \omega^m$  and  $d = d_{m-1} + d^m$  as in Section 2. Following the proof of the Poincaré Lemma in [54], we write

$$\omega^m = dx_m(A_1 dx_{I_1} + \dots + A_n dx_{I_n}),$$

where  $A_i \in K$  and the differentials  $dx_{I_i}$  are free of  $dx_m$ . By the classical integration formula in  $x_m$  that we have already used several times, we have

$$A_i = \partial_m(B_i) \quad \text{with } B_i = b_{i,0} + \sum_{j>0} c_{i,j} \log b_{i,j},$$

where  $b_{i,0} \in K$ ,  $c_{i,j} \in \overline{F}$ , and  $b_{i,j} \in F(c_{i,j})[x_m]$  are monic and pairwise coprime polynomials of positive degree when  $j > 0$ . Then the  $p$ -form  $\Psi^m := B_1 dx_{I_1} + \dots + B_n dx_{I_n}$  satisfies  $\omega^m = d^m(\Psi^m)$ , and we can define

$$\tilde{\omega} := \omega - d\Psi^m = (\omega_{m-1} + \omega^m) - (d_{m-1} + d^m)(\Psi^m) = \omega_{m-1} - d_{m-1}(\Psi^m).$$

Since  $d\omega = 0$ , we have  $d\tilde{\omega} = 0$ . Note that  $\tilde{\omega}$  is free of  $dx_m$ . By Lemma 2.4, we have  $d_{m-1}(\tilde{\omega}) = 0$  and  $d^m(\tilde{\omega}) = 0$ , which implies that  $\tilde{\omega}$  is free of  $x_m$ , i.e., the coefficients of  $\tilde{\omega}$  are constants with respect to  $x_m$ .

As defined,  $\tilde{\omega}$  appears to involve logarithms and elements of  $\overline{F}$  since these appear in  $\Psi^m$ . We claim that the coefficients of  $\tilde{\omega}$  actually lie in  $k(x_1, \dots, x_{m-1})$ . This is proved as follows. For  $\ell = 1, \dots, m-1$  and  $i = 1, \dots, n$ , we have

$$\begin{aligned}\partial_\ell(B_i) &= \partial_\ell\left(b_{i,0} + \sum_{j>0} c_{i,j} \log b_{i,j}\right) \\ &= \partial_\ell(b_{i,0}) + \sum_{j>0} c_{i,j} \frac{\partial_\ell(b_{i,j})}{b_{i,j}} + \sum_{j>0} \partial_\ell(c_{i,j}) \log b_{i,j}.\end{aligned}$$

Thus

$$\begin{aligned}\tilde{\omega} &= \omega_{m-1} - d_{m-1}(\Psi^m) = \omega_{m-1} - d_{m-1}\left(\sum_{i=1}^n B_i dx_{I_i}\right) \\ &= \omega_{m-1} - \sum_{\ell=1}^{m-1} \sum_{i=1}^n \partial_\ell(B_i) dx_\ell dx_{I_i} \\ &= \omega_{m-1} - \sum_{\ell=1}^{m-1} \sum_{i=1}^n \left(\partial_\ell(b_{i,0}) + \sum_{j>0} c_{i,j} \frac{\partial_\ell(b_{i,j})}{b_{i,j}} + \sum_{j>0} \partial_\ell(c_{i,j}) \log b_{i,j}\right) dx_\ell dx_{I_i} \\ &= \tilde{\omega}_{m-1} - \sum_{\ell=1}^{m-1} \sum_{i=1}^n \left(\sum_{j>0} c_{i,j} \frac{\partial_\ell(b_{i,j})}{b_{i,j}} + \sum_{j>0} \partial_\ell(c_{i,j}) \log b_{i,j}\right) dx_\ell dx_{I_i},\end{aligned}$$

where on the last line,  $\tilde{\omega}_{m-1} := \omega_{m-1} - \sum_{\ell=1}^{m-1} \sum_{i=1}^n \partial_\ell(b_{i,0}) dx_\ell dx_{I_i}$  has coefficients in  $K$ . It follows that we can write  $\tilde{\omega}$  as

$$\tilde{\omega} = f_{\tilde{I}_1} dx_{\tilde{I}_1} + f_{\tilde{I}_2} dx_{\tilde{I}_2} + \dots,$$

where each  $dx_{\tilde{I}_i}$  is free of  $dx_m$  and  $\partial_m(f_{\tilde{I}_i}) = 0$  since  $d^m(\tilde{\omega}) = 0$ . Observe that each  $f_{\tilde{I}_i}$  is an element in  $K$  minus the sum

$$\sum_{\substack{i,\ell \\ dx_\ell \wedge dx_{I_i} = \varepsilon_{i,\ell} dx_{\tilde{I}_i}}} \varepsilon_{i,\ell} \left( \sum_{j>0} c_{i,j} \frac{\partial_\ell(b_{i,j})}{b_{i,j}} + \sum_{j>0} \partial_\ell(c_{i,j}) \log b_{i,j} \right),$$

where  $\varepsilon_{i,\ell} = \pm 1$ . By Lemma 2.2, the logarithmic part of  $f_{\tilde{I}_j}$  vanishes. Then  $\partial_{x_m}(f_{\tilde{I}_i}) = 0$  implies that

$$f := \sum_{\substack{i,\ell \\ dx_\ell \wedge dx_{I_i} = \varepsilon_{i,\ell} dx_{\tilde{I}_i}}} \varepsilon_{i,\ell} \sum_{j>0} c_{i,j} \frac{\partial_\ell(b_{i,j})}{b_{i,j}}$$

satisfies the hypothesis of Lemma 4.2. We claim that this sum lies in  $K$ . Note that  $\deg_{x_m}(\partial_{x_\ell}(b_{i,j})) < \deg_{x_m}(b_{i,j})$  since  $b_{i,j}$  is monic of positive degree in  $x_m$  and  $\ell < m$ . This makes it easy to see that the constant part of  $f$  is 0  $\in F$ . By Lemma 4.2, it follows that  $f$  and hence  $f_{\tilde{I}_j}$  lie in  $K$ . Thus  $\tilde{\omega} = \omega - d(\Psi^m)$  is free of  $x_m$  and  $dx_m$  and has coefficients in  $k(x_1, \dots, x_{m-1})$ . Furthermore, since  $d = d_m$  in the  $m$ -variable case, we see that  $\omega = \tilde{\omega} + d_m(\Psi^m)$ .

If we repeat the above process on the closed rational form  $\tilde{\omega}$  in  $x_1, \dots, x_{m-1}$ , we obtain  $\tilde{\tilde{\omega}}$  and  $\Psi^{m-1}$ , where  $\tilde{\tilde{\omega}}$  is a closed rational form in  $x_1, \dots, x_{m-2}$  and

$\Psi^{m-1}$  is similar to  $\Psi^m$  except that  $x_m$  no longer appears. The result is that  $\tilde{\omega} = \tilde{\tilde{\omega}} + d_{m-1}(\Psi^{m-1})$ , where  $d_{m-1}$  is defined in Section 2. Thus

$$\omega = \tilde{\tilde{\omega}} + d_{m-1}(\Psi^{m-1}) + d_m(\Psi^m)$$

Repeating this process recursively, we must stop when only  $p$  variables remain since there are no  $p$ -forms in fewer than  $p$  variables. Hence we have proved the following result.

**Theorem 4.3.** *Let  $\omega$  be a closed  $p$ -form with coefficients in  $K = k(x_1, \dots, x_m)$ . Then  $\omega$  can be written in the form*

$$\omega = d_p(\Psi^p) + \dots + d_m(\Psi^m) = d(\Psi^p + \dots + \Psi^m),$$

where for each  $i = p, \dots, m$ , the coefficients  $f_{i,k}$  of  $\Psi^i$  are of the form

$$f_{i,k} = b_{i,k,0} + \sum_{j>0} c_{i,k,j} \log b_{i,k,j}$$

with

$$\begin{aligned} b_{i,k,0} &\in k(x_1, \dots, x_i) \\ c_{i,k,j} &\in \overline{k(x_1, \dots, x_{i-1})} \\ b_{i,k,j} &\in k(x_1, \dots, x_{i-1})(c_{i,k,j})[x_i], \quad j > 0. \end{aligned}$$

**Remark 4.4.** The above theorem says that Theorem 3.1 can be extended to general  $p$ -forms provided that the coefficients of the logarithms are allowed to be more general (specifically, algebraic over  $k(x_1, \dots, x_{i-1})$  for some  $i \in \{p, \dots, m\}$ ).

**Example 4.5.** Let us apply Theorem 4.3 to the 2-form  $\omega = x^{-1}y^{-1}dxdy$  from Example 4.1. In the notation of the theorem with  $x, y$  in place of  $x_1, x_2$ , we have

$$\omega^2 = \omega = x^{-1}y^{-1}dxdy = dy(A_1 dx) \text{ with } A_1 = -x^{-1}y^{-1}.$$

Furthermore,

$$A_1 = \partial_y(B_1) \text{ with } B_1 = -x^{-1} \log(y),$$

and the above procedure gives

$$\omega = d_2(\Psi^2) = d(B_1 dx) = d\left(-\frac{1}{x} \log(y) dx\right),$$

exactly as in Example 4.1. Switching  $x$  and  $y$  gives the other representation of  $\omega$  in the example.

**Example 4.6.** Consider the closed 2-form

$$\omega = \left(\frac{1}{z^2 - x} + \frac{1}{xy}\right) dxdy + \frac{1}{z^2 - x} dydz + \frac{y - 2yz}{(z^2 - x)^2} dxdz.$$

In the notation of Theorem 4.3 with  $x, y, z$  in place of  $x_1, x_2, x_3$ , we have  $\omega = (1/(z^2 - x) + 1/(xy)) dxdy + dz(A_1 dx + A_2 dy)$  with

$$A_1 := -\frac{y - 2yz}{(z^2 - x)^2} \text{ and } A_2 := -\frac{1}{z^2 - x}.$$

Furthermore,  $A_1 = \partial_z(B_1)$  with

$$B_1 := \frac{\sqrt{xy}}{4x^2} \log \frac{z - \sqrt{x}}{z + \sqrt{x}} + \frac{2yz - 4xy}{4x(z^2 - x)},$$

and  $A_2 = \partial_z(B_2)$  with

$$B_2 := -\frac{\sqrt{x}}{2x} \log \frac{z - \sqrt{x}}{z + \sqrt{x}}.$$

Hence,  $\Psi^3 := B_1 dx + B_2 dy$  and

$$\tilde{\omega} := \omega - d(\Psi^3) = \left( \frac{1}{z^2 - x} + \frac{1}{xy} + \partial_y(B_1) - \partial_x(B_2) \right) dx dy = \frac{1}{xy} dx dy.$$

Example 4.5 shows that  $\tilde{\omega} = d(\Psi^2)$  with  $\Psi^2 := -\frac{1}{x} \log(y) dx$ , so  $\omega = \tilde{\omega} + d(\Psi^3)$  is given by

$$\begin{aligned} \omega &= d(\Psi^2) + d(\Psi^3) = d(\Psi^2 + \Psi^3) \\ &= d\left( \left( -\frac{1}{x} \log(y) + \frac{\sqrt{xy}}{4x^2} \log \frac{z - \sqrt{x}}{z + \sqrt{x}} + \frac{2yz - 4xy}{4x(z^2 - x)} \right) dx - \frac{\sqrt{x}}{2x} \log \frac{z - \sqrt{x}}{z + \sqrt{x}} dy \right). \end{aligned}$$

Notice that the coefficients of the logs in  $\Psi^3$  involve  $x$  and  $y$ , while the coefficient of the log in  $\Psi^2$  involves only  $x$ , exactly as described in Theorem 4.3.

## 5. PICARD'S PROBLEM AND GRIFFITHS-DWORK REDUCTION

Let  $k$  be a field of characteristic zero and  $k(x, y, z)$  be the field of rational functions in  $x, y$  and  $z$  over  $k$ . Let  $\partial_x, \partial_y, \partial_z$  denote the usual derivations on  $k(x, y, z)$  with respect to  $x, y, z$ , respectively. In his books [45, Vol. II, pp. 475–479] and [44, pp. 209–215], Picard posed the following problem.

**Picard's Problem.** *Given  $f \in k(x, y, z)$ , decide whether there exist  $u, v, w \in k(x, y, z)$  such that*

$$(5.1) \quad f = \partial_x(u) + \partial_y(v) + \partial_z(w).$$

In this section, we will study an explicit example and reformulate Picard's problem in terms of differential forms. In the case of  $m$  variables, we will then explain how the Griffiths-Dwork order of pole reduction relates to Picard's problem. Note that the discrete analogue of Picard's problem has been recently solved in [17].

Set  $K = k(x, y)$  and let  $\overline{K}$  be the algebraic closure of  $K$ . Any rational function  $f = P/Q \in K(z)$  can be decomposed into

$$f = p + \sum_{i=1}^m \sum_{j=1}^{n_i} \frac{\alpha_{i,j}}{(z - \beta_i)^j},$$

where  $p \in K[z]$ , the  $\beta_i$ 's are the roots in  $\overline{K}$  of the denominator  $Q$ , and  $\alpha_{i,j} \in \overline{K}$ . We call the element  $\alpha_{i,1}$  the *z-residue* of  $f$  at  $\beta_i$ .

A rational function  $f \in k(x, y, z)$  is said to be *rationally integrable* with respect to  $x, y, z$  if  $f = \partial_x(u) + \partial_y(v) + \partial_z(w)$  for some  $u, v, w \in k(x, y, z)$ . An algebraic function  $\alpha \in \overline{k(x, y)}$  is said to be *algebraically integrable* with respect to  $x, y$  if  $\alpha = \partial_x(\beta) + \partial_y(\gamma)$  for some  $\beta, \gamma \in \overline{k(x, y)}$ . We recall a lemma from [23] on the relation between those two notions of being integrable.

**Lemma 5.1.** *Let  $f \in k(x, y, z)$ . Then  $f$  is rationally integrable with respect to  $x, y, z$  if and only if all z-residues of  $f$  are algebraically integrable with respect to  $x, y$ .*

We now show that some rational functions are not rationally integrable.

**Example 5.2.** For a fixed  $n \in \mathbb{N}$ , consider the rational function

$$f = \frac{1}{x^n + y^n + z^n}.$$

We will prove that  $f$  is rationally integrable with respect to  $x, y, z$  if and only if  $n \neq 3$ . It is straightforward to check that when  $n \neq 3$  we have

$$f = \partial_x \left( \frac{(3-n)^{-1}x}{x^n + y^n + z^n} \right) + \partial_y \left( \frac{(3-n)^{-1}y}{x^n + y^n + z^n} \right) + \partial_z \left( \frac{(3-n)^{-1}z}{x^n + y^n + z^n} \right).$$

So it remains to show that  $f$  is not rationally integrable with respect to  $x, y, z$  when  $n = 3$ . A root (in  $\bar{k}(x, y)$ ) of the denominator of  $f = 1/(x^3 + y^3 + z^3)$  is  $\lambda(x^3 + y^3)^{1/3}$  with  $\lambda^3 = -1$ . Then the residue of  $f$  at this root is

$$\frac{1}{3\lambda^2(x^3 + y^3)^{2/3}}.$$

By Lemma 5.1, it suffices to show that the algebraic function

$$\alpha(x, y) = \frac{1}{(x^3 + y^3)^{2/3}}$$

is not algebraically integrable. Consider the birational transformation

$$\bar{x} = x \quad \text{and} \quad \bar{y} = y/x.$$

Then one computes that

$$(5.2) \quad \alpha(x, y) \, dx dy = \bar{\alpha}(\bar{x}, \bar{y}) \, d\bar{x} d\bar{y}, \quad \text{where } \bar{\alpha}(\bar{x}, \bar{y}) = \frac{1}{\bar{x}(1 + \bar{y}^3)^{2/3}}.$$

Note that  $\bar{\alpha}$  is a rational function in  $\bar{x}$ . By a similar result to Lemma 5.1,  $\bar{\alpha}$  is algebraically integrable with respect to  $\bar{x}, \bar{y}$  if and only if the residue of  $\bar{\alpha}$  at  $\bar{x} = 0$  is algebraically integrable with respect to  $\bar{y}$ . The residue of  $f$  at  $\bar{x} = 0$  is

$$\beta(\bar{y}) = \frac{1}{(1 + \bar{y}^3)^{2/3}}.$$

This is a simple radical and one can show that it is not algebraically integrable with respect to  $\bar{y}$ . In fact, if  $\beta(\bar{y})$  is algebraically integrable, then

$$\beta(\bar{y}) = D_{\bar{y}}(\gamma(\bar{y})),$$

where  $\gamma(\bar{y})$  is algebraic over  $k(\bar{y})$  and  $D_{\bar{y}} = d/d\bar{y}$ . By [12, Lemma 5.1.1], it follows that  $D_{\bar{y}}(\gamma(\bar{y}))$  is the derivative of an element of  $k(\bar{y}, \beta(\bar{y}))$ . Using  $D_{\bar{y}}(\beta) = \frac{-2\bar{y}^2}{1 + \bar{y}^3}\beta$  and the basis  $\{1, \beta(\bar{y}), \beta(\bar{y})^2\}$  of  $k(\bar{y}, \beta(\bar{y}))$  over  $k(\bar{y})$ , one can show that  $\gamma(\bar{y}) = r\beta(\bar{y}) + c$ , where  $r \in k(\bar{y})$  and  $c \in k$ . Thus

$$\beta(\bar{y}) = D_{\bar{y}} \left( \frac{r}{(1 + \bar{y}^3)^{2/3}} \right),$$

where  $r \in k(\bar{y})$ . It follows that

$$(5.3) \quad D_{\bar{y}}(r) - \frac{2\bar{y}^2}{1 + \bar{y}^3} \cdot r = 1.$$

A straightforward application of the Risch Algorithm from [12, Ch. 6] to (5.3) shows that the rational solution  $r$  is in fact a polynomial of degree at most 2 (see Def. 6.11, Thm. 6.1.2, and Cor. 6.3.1 of [12]). An easy computation now leads to a contradiction, so that (5.3) has no rational solution. As we have seen, this implies that  $\bar{\alpha}$  is not algebraically integrable.

We claim that the same is true for  $\alpha$ . Here is a classical argument using Picard and Simart [45] (see Remark 5.3 below for a modern proof). Suppose that  $\alpha = \partial_x(\beta) + \partial_y(\gamma)$  for  $\beta, \gamma \in \overline{k(x, y)}$ . In terms of integrals, this means that

$$\iint \alpha \, dx \, dy = \iint \partial_x(\beta) + \partial_y(\gamma) \, dx \, dy.$$

By [45, Vol. II, pp. 160–161], integrals such as the one on the right are preserved under birational transformations when  $\beta$  and  $\gamma$  are rational functions. The proof adapts easily to the case of algebraic functions, so that for the birational transformation given above, we have

$$\iint \partial_x(\beta) + \partial_y(\gamma) \, dx \, dy = \iint \partial_{\bar{x}}(\bar{\beta}) + \partial_{\bar{y}}(\bar{\gamma}) \, d\bar{x} \, d\bar{y}$$

for algebraic functions  $\bar{\beta}$  and  $\bar{\gamma}$ . Combining this with (5.2), we conclude that  $\bar{\alpha}$  is algebraically integrable, which contradicts what we have proved above.

It follows that  $\alpha$  is not algebraically integrable. As noted above, Lemma 5.1 then implies that  $f = \frac{1}{x^3 + y^3 + z^3}$  is not rationally integrable.

The 2-form  $(\partial_x(\beta) + \partial_y(\gamma)) \, dx \, dy$  appearing in Example 5.2 suggests that the 3-form  $(\partial_x(u) + \partial_y(v) + \partial_z(w)) \, dx \, dy \, dz$  may be relevant to Picard's problem. In fact,

$$d(u \, dy \, dz - v \, dx \, dz + w \, dx \, dy) = (\partial_x(u) + \partial_y(v) + \partial_z(w)) \, dx \, dy \, dz,$$

which implies that Picard's problem (5.1) is equivalent to saying that the 3-form  $f(x, y, z) \, dx \, dy \, dz$  is exact, i.e.,

$$f \, dx \, dy \, dz = d(u \, dy \, dz - v \, dx \, dz + w \, dx \, dy), \quad \text{where } u, v, w \in k(x, y, z).$$

**Remark 5.3.** Similar formulas also apply to algebraic functions in two variables, so that  $\alpha(x, y)$  from Example 5.2 is algebraically integrable if and only if the 2-form  $\alpha(x, y) \, dx \, dy$  is the exterior derivative of an algebraic 1-form. Since the exterior derivative is compatible with pull-backs, we get an immediate proof that in (5.2),  $\alpha(x, y)$  is not algebraically integrable if and only if the same is true for  $\bar{\alpha}(\bar{x}, \bar{y})$ .

We can also relate Example 5.2 to Theorem 4.3 as follows.

**Example 5.4.** By the previous example,  $f = 1/(x^3 + y^3 + z^3)$  is not rationally integrable with respect to  $x, y, z$ , so that  $f \, dx \, dy \, dz$  is not the exterior derivative of a rational 2-form. However, Theorem 4.3 shows how to write  $f \, dx \, dy \, dz$  as the exterior derivative of a rational and logarithmic 2-form as follows.

Let  $\alpha = -(x^3 + y^3)^{1/3}$ . Then the three roots in  $\overline{\mathbb{Q}(x, y)}$  of the equation  $z^3 + x^3 + y^3 = 0$  are  $z = \alpha, \alpha\omega, \alpha\omega^2$ , where  $\omega$  is a primitive cube root of unity. Then

$$f = \frac{1/(3\alpha^2)}{z - \alpha} + \frac{\omega/(3\alpha^2)}{z - \alpha\omega} + \frac{\omega^2/(3\alpha^2)}{z - \alpha\omega^2},$$

which implies that  $f = \partial_z(g)$  with

$$g = \frac{1}{3\alpha^2} \log(z - \alpha) + \frac{\omega}{3\alpha^2} \log(z - \alpha\omega) + \frac{\omega^2}{3\alpha^2} \log(z - \alpha\omega^2).$$

It follows easily that  $f \, dx \, dy \, dz = d(g \, dx \, dy)$ .

**Conjecture 5.5.** For a fixed  $n \in \mathbb{N}$ , the rational function

$$f = \frac{1}{x_1^n + \cdots + x_m^n}$$

is rationally integrable with respect to  $x_1, \dots, x_m$  with  $m \geq 4$  if and only if  $n \neq m$ .

The general case of Picard's problem asks when  $f \in K = k(x_1, \dots, x_m)$  can be written in the form

$$f = \partial_{x_1}(u_1) + \dots + \partial_{x_m}(u_m), \quad \text{where } u_1, \dots, u_m \in K.$$

Since

$$d\left(\sum_{i=0}^m (-1)^{i-1} u_i dx_1 \cdots \widehat{dx_i} \cdots dx_m\right) = \left(\sum_{i=1}^m \partial_i(u_i)\right) dx_1 \cdots dx_m,$$

Picard's problem is thus equivalent to deciding when the  $m$ -form

$$\omega = f dx_1 \cdots dx_m, \quad \text{where } f \in K,$$

is rationally integrable, i.e., is the exterior derivative of a rational  $(m-1)$ -form.

Under a regularity assumption, the Griffiths-Dwork method [29, §3], [30, §8], [34, §4] solves Picard's problem for a general  $m$ -variable case. Here we sketch a solution based on the modern presentation [10]. One difference is that we make explicit use of differential forms, which are suppressed in [10].

The regularity assumption is that the polar locus of  $\omega$  is a smooth hypersurface  $V \subseteq \mathbb{P}^m$ . Let  $\ell$  be the order of the pole of  $\omega$  along  $V$ , and write the defining equation of  $V$  as  $Q = 0$ , where  $Q$  is irreducible in the homogeneous coordinates  $\xi_0, \dots, \xi_m$  of  $\mathbb{P}^m$ . Here  $x_i = \xi_i/\xi_0$  for  $i = 1, \dots, m$ . By [34, Cor. 2.11], this implies that in homogeneous coordinates,  $\omega$  can be written as

$$(5.4) \quad \omega = \frac{P\Omega}{Q^\ell},$$

where  $Q$  and  $\ell$  are as above,  $P$  is homogeneous with  $\ell \deg(Q) = \deg(P) + m + 1$ , and  $\Omega$  is the  $m$ -form

$$\Omega := \sum_{i=0}^m (-1)^i \xi_i (d\xi_0 \cdots \widehat{d\xi_i} \cdots d\xi_m).$$

In [34], Griffiths studies  $\omega$  using *order of pole reduction*. A key tool is the *Jacobian ideal*

$$J(Q) := \langle \partial_0(Q), \dots, \partial_m(Q) \rangle \subseteq k[\xi_0, \dots, \xi_m],$$

where  $\partial_i = \partial_{\xi_i}$  for  $i = 0, \dots, m$ . Our regularity assumption implies that the quotient

$$A(Q) := k[\xi_0, \dots, \xi_m]/J(Q)$$

is a finite-dimensional graded algebra over  $k$ .

Now suppose that we have  $\omega = \frac{P\Omega}{Q^\ell}$  as in (5.4) with  $\ell \deg(Q) = \deg(P) + m + 1$ . If  $\ell > 1$ , consider the  $(m-1)$ -form

$$\varphi_1 = \frac{1}{\ell-1} \sum_{0 \leq i < j \leq m} (\xi_i A_j - \xi_j A_i) d\xi_0 \cdots \widehat{d\xi_i} \cdots \widehat{d\xi_j} \cdots d\xi_m$$

for homogeneous polynomials  $A_0, \dots, A_m$  with  $\deg(A_i) + m = (\ell-1) \deg(Q)$ . Griffiths [34, §2 and §4] shows that  $\varphi_1$  represents an  $(m-1)$  form on  $\mathbb{P}^m$  with exterior derivative

$$d\varphi_1 = \frac{(\sum_{i=0}^m A_i \partial_i(Q))\Omega}{Q^\ell} - \frac{\frac{1}{\ell-1} (\sum_{i=0}^m \partial_i(A_i))\Omega}{Q^{\ell-1}}.$$

Note that  $\sum_{i=0}^m A_i \partial_i(Q) \in J(Q)$ . The rough idea is that subtracting  $d\varphi_1$  from  $\omega = \frac{P\Omega}{Q^\ell}$  for carefully chosen  $A_i$  can reduce the order of the pole in part of  $\omega$ . To make this precise, let  $G$  be a Gröbner basis of  $J(Q)$  and divide  $P$  by  $G$  to obtain

$$P = \sum_{i=0}^m A_i \partial_i(Q) + r_1,$$

where  $r_1$  is the remainder (which can be identified with an element of  $A(Q)$ ). Combining this with the above formula for  $d\varphi_1$  gives

$$\omega = \frac{P\Omega}{Q^\ell} = \frac{(\sum_{i=0}^m A_i \partial_i(Q))\Omega}{Q^\ell} + \frac{r_1\Omega}{Q^\ell} = d\varphi_1 + \underbrace{\frac{(\frac{1}{\ell-1} \sum_{i=0}^m \partial_i(A_i))\Omega}{Q^{\ell-1}}}_{\omega_1} + \frac{r_1\Omega}{Q^\ell}.$$

Applying this recursively to  $\omega_1 = \frac{P_1\Omega}{Q^{\ell-1}}$  gives  $\omega_1 = d\varphi_2 + \omega_2 + \frac{r_2\Omega}{Q^{\ell-2}}$  with  $\omega_2 = \frac{P_2\Omega}{Q^{\ell-2}}$ . We can continue until we get  $\omega_{\ell-1} = \frac{P_{\ell-1}\Omega}{Q}$  with  $\deg(Q) = \deg(P_{\ell-1}) + m + 1$ . Then  $\deg(P_{\ell-1}) < \deg(Q) - 1 = \deg(\partial_i(Q))$ . It follows that  $P_{\ell-1}$  is already a remainder  $r_\ell$  modulo  $J(Q)$ , so that  $\omega_{\ell-1} = \frac{r_\ell\Omega}{Q}$  and  $d\varphi_\ell = \omega_\ell = 0$ . Thus we obtain

$$\omega = d\varphi_1 + \cdots + d\varphi_{\ell-1} + [\omega], \quad \text{where } [\omega] = \frac{r_1\Omega}{Q^\ell} + \frac{r_2\Omega}{Q^{\ell-1}} + \cdots + \frac{r_\ell\Omega}{Q}.$$

We call  $[\omega]$  the *Griffiths-Dwork reduction* of  $\omega$ .

This reduction algorithm is Algorithm 1 from [10] recast in the language of differential forms. The results of [34, §4] lead to Theorem 1 of [10], which we restate as follows.

**Proposition 5.6.** *Assume that  $\omega = \frac{P\Omega}{Q^\ell}$  satisfies the regularity condition that  $Q = 0$  defines a smooth hypersurface in  $\mathbb{P}^m$ . Then  $\omega$  is rationally integrable if and only if its Griffiths-Dwork reduction  $[\omega]$  vanishes.*

Since the summands in  $[\omega] = \frac{r_1\Omega}{Q^\ell} + \frac{r_2\Omega}{Q^{\ell-1}} + \cdots + \frac{r_\ell\Omega}{Q}$  are easily seen to be linearly independent, it follows that  $\omega$  is not rationally integrable as soon as one of the remainders  $r_1, r_2, \dots$  is nonzero. So if one is only interested in knowing whether  $\omega$  is rationally integrable, it is often not necessary to compute the full Griffiths-Dwork reduction.

**Example 5.7.** Consider the 3-form

$$\omega = \frac{dxdydz}{x^3 + y^3 + z^3}$$

from Example 5.4. Setting  $(x, y, z) = (\xi_1/\xi_0, \xi_2/\xi_0, \xi_3/\xi_0)$ , one computes that  $dxdydz = \frac{\Omega}{\xi_0^4}$ , so that

$$\omega = \frac{\Omega}{\xi_0(\xi_1^3 + \xi_2^3 + \xi_3^3)}.$$

The polar locus of this form is not smooth and irreducible, so the regularity assumption is violated. Hence we cannot use Proposition 5.6 to decide if  $\omega$  is exact.

In [39], Lairez uses results of Dimca [28] to solve Picard's problem without assuming regularity. In this setup, we instead assume that  $Q$  is squarefree. Then any  $m$ -form with polar locus  $V$  defined by  $Q = 0$  can be written

$$\omega = \frac{P\Omega}{Q^\ell}$$

for some  $\ell$ . Lairez [39, Sec. 4.2] defines a series of reductions  $[\omega]_r$  for  $r \geq 1$ , where  $[\omega]_1$  closely related to the Griffiths-Dwork reduction defined above (see [39, Rem. 17]). The challenges involved in computing  $[\omega]_r$  are discussed in [39, Sec. 4.2 and 4.3].

Using these reductions, we can solve Picard's problem as follows.

**Proposition 5.8.**

- (i)  $\omega$  is rationally integrable if and only if  $[\omega]_r = 0$  for some  $r \geq 1$ .
- (ii) There is an integer  $r_Q$  depending only on  $Q$  such that  $\omega$  is rationally integrable if and only if  $[\omega]_{r_Q} = 0$ .

*Proof.* Part (i) follows from [39, Cor. 16] after unwinding a lot of notation, and part (ii) then follows from [39, Cor. 24].  $\square$

In order to use Proposition 5.8, we need to find an effective bound for  $r_Q$ . One known bound comes from the proof of Theorem B of [28] and is computed using an embedded resolution of singularities of the polar locus  $V$  of  $\omega$ . A conjecture made in [28] would imply that  $r_Q \leq m + 1$  (see [39, Conj. 27]), but this is still an open question.

The upshot is that although the methods of [39] apply to examples like those in Example 5.7, a substantial amount of work would be involved. The Griffiths-Dwork reduction provides a partial resolution to the question of the rational integrability of  $m$ -forms. It also offers a novel approach to handling multiple integrals through creative telescoping, see [10]. Nevertheless, the problem of determining the rational integrability of general  $p$ -forms remains open.

## 6. LIOUVILLE'S THEOREM FOR DIFFERENTIAL 1-FORMS

We will extend Liouville's classical theorem to integrals of differential 1-forms in  $\mathcal{M}$ . Let  $K \subseteq \mathcal{U}$  be the universal differential extension from Section 2.

**Definition 6.1.** Let  $E \subseteq \mathcal{U}$  be a differential extension of  $K$  and  $t \in E$ .

- (1)  $t$  is a *logarithm* over  $K$  if there exists nonzero  $b \in K$  such that  $\partial_i t = \partial_i b/b$  for each  $i \in \{1, \dots, m\}$ .
- (2)  $t$  is an *exponential* over  $K$  if there exists  $b \in K$  such that  $\partial_i t/t = \partial_i b$  for each  $i \in \{1, \dots, m\}$ .
- (3)  $E$  is an *elementary extension* of  $K$  if there exist  $t_1, \dots, t_n \in E$  such that  $E = K(t_1, \dots, t_n)$  and for each  $j \in \{1, \dots, n\}$ ,  $t_j$  is either algebraic or a logarithm or an exponential over  $K(t_1, \dots, t_{j-1})$ .

The following theorem is Risch's strong version of the classical Liouville's theorem (see [49, 48] and Section 5.5 in [12] for its proof).

**Theorem 6.2** (Liouville's theorem). *Let  $(K, D)$  be a differential field and  $f \in K$ . If there exists an elementary extension  $E$  of  $K$  and  $g \in E$  such that  $D(g) = f$ , then*

there are  $v \in K, u_1, \dots, u_n \in K^*$  and  $c_1, \dots, c_n \in \overline{\text{Const}_D(K)}$  such that

$$f = D(v) + \sum_{i=1}^n c_i \frac{D(u_i)}{u_i}.$$

The following theorem was first proved by Caviness and Rothstein in [16]. We now present a more direct proof inspired by the recursive argument used in the proof of Theorem 3.1.

**Theorem 6.3.** *Let  $F \subseteq \mathcal{U}$  be a differential extension  $K$  and  $C := \text{Const}_{\partial_1, \dots, \partial_m}(F)$ . Let  $\omega = f_1 dx_1 + \dots + f_m dx_m \in \mathcal{M}$  be a closed 1-form with  $f_i \in F$ . If there exist an elementary extension  $E$  of  $F$  and  $g \in E$  such that  $\omega = dg$ , then there are  $a \in F$ ,  $c_1, \dots, c_n \in C$ , and  $b_1, \dots, b_n \in F(c_1, \dots, c_n)^*$  such that*

$$g = a + \sum_j c_j \log b_j.$$

*Proof.* Denote  $C_m := \text{Const}_{\partial_m}(F)$ . Since  $f_m$  has an elementary integral over  $F$  with respect to  $\partial_m$ , Theorem 6.2 implies that there exist  $a_m \in F$ ,  $c_{m,1}, \dots, c_{m,n} \in \overline{C_m}$ ,  $b_{m,1}, \dots, b_{m,n} \in F(c_{m,1}, \dots, c_{m,n})$  such that

$$f_m = \partial_m(g_m) \quad \text{with} \quad g_m = a_m + \sum_{j=1}^n c_{m,j} \log b_{m,j}$$

where  $\partial_m(b_{m,j}) \neq 0$  for each  $j$ . Note that  $F' := \{\partial_m(f) \mid f \in F\}$  is a linear subspace of  $F$  over  $C_m$ . By Corollary 3.3.1 from [12], we have  $\text{Const}_{\partial_m}(\overline{F}) = \overline{C_m}$ , so  $\overline{F}' := \{\partial_m(f) \mid f \in \overline{F}\}$  is a linear subspace of  $\overline{F}$  over  $\overline{C_m}$ . For every  $f \in \overline{F}$ , let  $\widehat{f}$  be the image of  $f$  in the quotient space  $\overline{F}/\overline{F}'$ . We can always select maximal many elements of the  $\widehat{\frac{\partial_m(b_{m,j})}{b_{m,j}}}$ 's so that they are linearly independent over  $\overline{C_m}$  and then rearrange the logarithmic part of  $g_m$ . After this process, we can always assume the  $\widehat{\frac{\partial_m(b_{m,j})}{b_{m,j}}}$ 's appearing in  $g_m$  are linearly independent over  $\overline{C_m}$ , though  $a_m$  may now be in  $\overline{F}$ . Since  $\omega$  is closed, we have  $\partial_m(f_i) = \partial_i(f_m)$  for all  $1 \leq i \leq m$ . Then (3.1) and (3.2) imply that

$$\partial_m(f_i) = \partial_m(\partial_i(a_m)) + \sum_{j=1}^n \partial_i(c_{m,j}) \frac{\partial_m(b_{m,j})}{b_{m,j}} + \partial_m \left( \sum_{j=1}^n c_{m,j} \frac{\partial_i(b_{m,j})}{b_{m,j}} \right).$$

Taking the image of both sides in the quotient space  $\overline{F}/\overline{F}'$  gives

$$\sum_{j=1}^n \partial_i(c_{m,j}) \widehat{\frac{\partial_m(b_{m,j})}{b_{m,j}}} = 0,$$

where  $\partial_i(c_{m,j}) \in \overline{C_m}$  because  $c_{m,j} \in \overline{C_m}$ . Since the  $\widehat{\frac{\partial_m(b_{m,j})}{b_{m,j}}}$ 's are linearly independent over  $\overline{C_m}$ ,  $\partial_i(c_{m,j}) = 0$  for each  $j$ , which implies that  $c_{m,j} \in \overline{C_i}$ . Since this holds for all  $i \in \{1, \dots, m-1\}$ , we see that  $c_{m,j} \in \overline{C}$ .

Let  $\tilde{\omega} := \omega - dg_m$ . Then  $\tilde{\omega} = \tilde{f}_1 dx_1 + \dots + \tilde{f}_{m-1} dx_{m-1}$ , where for each  $i$ ,

$$\tilde{f}_i := f_i - \partial_i(g_m) = f_i - \partial_i(a_m) - \sum_{j=1}^n c_{m,j} \frac{\partial_i(b_{m,j})}{b_{m,j}} \in \overline{F}.$$

This shows that the coefficient field is extended only by algebraically. And the constant field is also extended algebraically, i.e.,  $\text{Const}_{\partial_1, \dots, \partial_m}(\overline{F}) = \overline{C} \cap \overline{F} = \overline{C}$ . Since  $\omega$  is closed, so is  $\tilde{\omega}$ , which implies that  $\partial_m(\tilde{f}_i) = 0$  for  $i = 1, \dots, m-1$ . So  $\tilde{\omega}$  is a differential form that is free of  $dx_m$  and has coefficients in  $C_m(c_{m,1}, \dots, c_{m,n})$ .

Repeating this process recursively shows that there are  $a \in \overline{F}$ ,  $c_j \in \overline{C}$  and  $b_j \in \overline{C}F^*$  such that

$$f_i = \partial_i(a) + \sum_j c_j \frac{\partial_i(b_j)}{b_j}$$

for all  $i \in \{1, \dots, m\}$ . This almost has the required form. The problem is that  $a$  need not lie in  $F$  and the  $b_j$ 's need not lie in  $F(\{c_j\})$ . We have more work to do.

Let  $A$  be the minimal Galois extension of  $F$  generated by  $a$ , the  $c_j$ 's, and the  $b_j$ 's with  $d := [A : F] < \infty$ . As in the proof of Lemma 4.2, we have the trace map  $\text{Tr}_{A/F}(a) = \sum_{\sigma \in G} \sigma(a)$  for  $a \in A$ , where  $G$  be the Galois group of  $A$  over  $F$ . Using the trace map, we obtain

$$d \cdot f_i = \text{Tr}_{A/F}(f_i) = \partial_i(\text{Tr}_{A/F}(a)) + \sum_j \sum_{\sigma \in G} \sigma(c_j) \frac{\partial_i(\sigma(b_j))}{\sigma(b_j)}.$$

Thus

$$f_i = \partial_i \left( \frac{\text{Tr}_{A/F}(a)}{d} \right) + \sum_j \sum_{\sigma \in G} \frac{\sigma(c_j)}{d} \frac{\partial_i(\sigma(b_j))}{\sigma(b_j)},$$

where

$$\frac{\text{Tr}_{A/F}(a)}{d} \in F, \quad \frac{\sigma(c_j)}{d} \in \overline{C} \quad \text{and} \quad \sigma(b_j) \in \overline{C}F.$$

This is closer to what we want, but the  $\sigma(b_j)$ 's need not lie in  $F(\{\sigma(c_j)\}_{j,\sigma})$ . So there is still more work to do.

Now let  $L$  be the field extension of  $F$  generated by  $\sigma(c_j)$ 's. Note that  $F \subseteq L \subseteq A$ . For the extension  $L \subseteq A$  of degree  $\ell := [A : L]$ , let  $\text{Tr}_{A/L} : A \rightarrow L$  and  $N_{A/L} : A \rightarrow L$  be the trace and norm maps. Recall that for any  $b \in A^*$ , we have the well-known identity

$$\text{Tr}_{A/L} \left( \frac{\partial_i(b)}{b} \right) = \frac{\partial_i(N_{A/L}(b))}{N_{A/L}(b)}.$$

Then applying  $\text{Tr}_{A/L}$  to the above expression for  $f_i$  yields

$$\begin{aligned} \ell \cdot f_i &= \text{Tr}_{A/L}(f_i) = \ell \cdot \partial_i \left( \frac{\text{Tr}_{A/F}(a)}{d} \right) + \sum_j \sum_{\sigma \in G} \frac{\sigma(c_j)}{d} \text{Tr}_{A/L} \left( \frac{\partial_i(\sigma(b_i))}{\sigma(b_i)} \right) \\ &= \ell \cdot \partial_i \left( \frac{\text{Tr}_{A/F}(a)}{d} \right) + \sum_j \sum_{\sigma \in G} \frac{\sigma(c_j)}{d} \frac{\partial_i(N_{A/L}(\sigma(b_i)))}{N_{A/L}(\sigma(b_i))}, \end{aligned}$$

where the second line uses the above identity. Hence,

$$f_i = \partial_i \left( \frac{\text{Tr}_{A/F}(a)}{d} \right) + \sum_j \sum_{\sigma \in G} \frac{\sigma(c_j)}{d \cdot \ell} \frac{\partial_i(N_{A/L}(\sigma(b_i)))}{N_{A/L}(\sigma(b_i))},$$

where  $\text{Tr}_{A/F}(a)/d \in F$ ,  $\frac{\sigma(c_j)}{d \cdot \ell} \in \overline{C}$ , and  $N_{A/L}(\sigma(b_i)) \in L = F(\{\sigma(c_j)\}_{j,\sigma})$ . Since this holds for  $i \in \{1, \dots, m\}$ , it follows that  $\omega = f_1 dx_1 + \dots + f_m dx_m = dg$ , where

$$g = \frac{\text{Tr}_{A/F}(a)}{d} + \sum_j \sum_{\sigma \in G} \frac{\sigma(c_j)}{d \cdot \ell} \log N_{A/L}(\sigma(b_i))$$

has the required form. This completes the proof.  $\square$

The following example indicates the difficulty of extending Theorem 6.3 to  $p$ -forms for  $p > 1$ .

**Example 6.4.** Let  $\omega = (\exp(x + y^2) + \exp(y + x^2)) dx dy$ . It is easy to check that  $\omega = d(\eta)$  with  $\eta := \exp(x + y^2) dy - \exp(y + x^2) dx$ .

The recursive strategy used to prove Theorem 3.1 for 1-forms adapts nicely to prove Theorem 4.3 for  $p$ -forms. However, this strategy fails when we try to adapt the proof of Theorem 6.3 to the above 2-form, which we write as  $\omega = f dx dy$  for  $f := \exp(x + y^2) + \exp(y + x^2)$ . To see why, we follow the strategy of Example 4.5. The first step is to write  $\omega = dy(-f dx)$  and then find  $g$  such that  $-f = \partial_y(g)$ . But no elementary function has this property because  $f$  contains  $\exp(x + y^2)$ . Also, switching  $x$  and  $y$  does not help, since here  $\omega = dx(f dy)$ , but there is no elementary function  $g$  satisfying  $f = \partial_x(g)$  because  $f$  contains  $\exp(y + x^2)$ .

## 7. ZEILBERGER'S CREATIVE TELESCOPING FOR DIFFERENTIAL FORMS WITH ONE PARAMETER

The method of *creative telescoping* first appeared in van der Poorten's report [53] on Apéry's proof of the irrationality of  $\zeta(3)$ . Specifically, the method was used to show that the sum

$$b_n := \sum_{\ell} b_{n,\ell} := \sum_{\ell} \binom{n}{\ell}^2 \binom{n+\ell}{\ell}^2$$

satisfies the recurrence

$$n^3 b_n - (34n^3 - 51n^2 + 27n - 5)b_{n-1} + (n-1)^3 b_{n-2} = 0, \text{ for all } n \geq 2.$$

The essential idea is to "cleverly construct" [53] the product

$$B_{n,\ell} = 4(2n+1)(\ell(2\ell+1) - (2n+1)^2) \cdot b_{n,\ell}$$

and observe that if  $\Delta_{\ell}$  is the usual difference operator, then the identity

$$\begin{aligned} n^3 b_{n,\ell} + (n-1)^3 b_{n-2,\ell} - (34n^3 - 51n^2 + 27n - 5)b_{n-1,\ell} &= B_{n,\ell} - B_{n,\ell-1} \\ &= \Delta_{\ell}(B_{n,\ell}). \end{aligned}$$

follows from a straightforward algebraic computation since  $b_{n,\ell} = \binom{n}{\ell}^2 \binom{n+\ell}{\ell}^2$ . Note that both  $b_{n,\ell}$  and  $B_{n,\ell}$  vanish automatically if  $\ell < 0$  or  $\ell > n$ . If we fix  $n \geq 2$  and sum over all integers  $\ell$ , the right-hand side turns to zero, which proves the recurrence.

In general, for a given sequence  $b_{n,\ell}$ , the process of creative telescoping finds a nontrivial linear recurrence  $L \in k(n)\langle S_n \rangle$  and another operator  $Q \in k(n, \ell)\langle S_n, S_{\ell} \rangle$  such that

$$L(n, S_n)(b_{n,\ell}) = \Delta_{\ell}(Q(b_{n,\ell})).$$

Here,  $n$  is the operator of multiplication by  $n$  and  $S_n$  is the shift operator that maps  $n$  to  $n+1$ . The notation  $k(n)\langle S_n \rangle$  reflects the fact that  $n$  and  $S_n$  do not commute since  $S_n n = (n+1)S_n$ . In the above identity, the operator  $L$  is called a *telescopers* for  $b_{n,\ell}$  and  $Q(b_{n,\ell})$  is the *certificate* of  $L$  for  $b_{n,\ell}$ . Zeilberger's fast algorithm [55, 56] for creative telescoping was developed to solve this problem when restricted to so-called *hypergeometric terms*  $b_{n,\ell}$ , meaning that both  $S_n(b_{n,\ell})/b_{n,\ell}$  and  $S_{\ell}(b_{n,\ell})/b_{n,\ell}$  are rational functions of  $n$  and  $\ell$ .

In the differential analogue of creative telescoping [3], sums and difference operators are replaced with integrals and differential operators, respectively. Thus, given  $F(x, y)$  in place of  $b_{n,\ell}$ , we seek a linear differential equation satisfied by the integral

$$f(x) = \int_{-\infty}^{+\infty} F(x, y) dy,$$

where  $F(x, y)$  is *hyperexponential* with respect to  $x, y$ , i.e.,  $\partial_x(F(x, y))/F(x, y)$  and  $\partial_y(F(x, y))/F(x, y)$  are both rational functions of  $x$  and  $y$ . In this case, there always exists a nonzero telescopers  $L \in k(x) \langle \partial_x \rangle$  for  $F(x, y)$  with respect to  $y$  with certificate  $G(x, y)$  such that

$$L(F(x, y)) = \partial_y(G(x, y)).$$

(As above, the notation  $k(x) \langle \partial_x \rangle$  reflects the fact that  $x$  and  $\partial_x$  do not commute since  $\partial_x x = x\partial_x + 1$ .) Then integrating with respect to  $y$  leads to

$$L(f(x)) = 0.$$

In recent years, reduction-based creative telescoping method [5, 7, 10, 21, 22, 9, 26] is a new trend because of its efficiency in computing telescopers without certificates. The existence of telescopers for differential forms with D-finite function coefficients is studied in [20].

In this section, we assume that all integrals involve one parameter  $t$  and the derivation with respect to  $t$  is denoted by  $\partial_t$ . We will prove that telescopers exist for closed algebraic 1-forms and use `HermiteOneForm` from Section 3 to give a reduction-based creative telescoping algorithm to compute the telescopers for closed rational 1-forms.

**Definition 7.1.** Given a 1-form  $\omega = f_1 dx_1 + \cdots + f_m dx_m$  with  $f_i$  algebraic over  $k(t, x_1, \dots, x_m)$ , a linear differential operator  $L$  in  $k(t) \langle \partial_t \rangle$  is called a *telescopers* for  $\omega$  if  $L(\omega) := L(f_1) dx_1 + \cdots + L(f_m) dx_m = dg$  for some algebraic function  $g$  over  $k(t, x_1, \dots, x_m)$ . The function  $g$  is called a *certificate* of  $L$ . A nonzero telescopers for  $\omega$  of minimal degree in  $\partial_t$  is called a *minimal telescopers*.

Observe that  $L(\omega) = dg$  is equivalent to  $L(f_i) = \partial_i(g)$  for each  $i \in \{1, \dots, m\}$ . In the terminology of [19],  $L$  is called a *parallel telescopers*.

**Theorem 7.2.** Let  $\omega = f_1 dx_1 + \cdots + f_m dx_m$  be a 1-form with coefficients algebraic over  $k(t, x_1, \dots, x_m)$ . If  $\omega$  is closed, then it always has a nonzero telescopers.

*Proof.* We proceed by induction on  $m$ , the number of variables  $x_1, \dots, x_m$ . For  $m = 1$ , one can construct a nonzero telescopers for  $f_1$  with respect to  $x_1$  by the method in [23, 22]. Suppose now that  $m > 1$  and that the theorem holds for the case when the number of variables is less than  $m$ . For  $f_m \in \overline{k(t, x_1, \dots, x_m)}$ , there exists a telescopers  $L_m \in k(t, x_1, \dots, x_{m-1}) \langle \partial_t \rangle$  with respect to  $x_m$  and its certificate  $g_m$  such that

$$L_m(f_m) = \partial_m(g_m).$$

We claim that  $L_m$  can be chosen to be free of  $x_1, \dots, x_{m-1}$ . *W.L.O.G.*, assume that all coefficients of  $L_m$  lie in the polynomial ring  $k[t, x_1, \dots, x_{m-1}]$ , otherwise we can multiply both sides of the above equation by the least common multiple of the denominators. Take  $L_m$  with  $\deg_{x_1}(L_m)$  minimal, denoted by  $n$ , and write

$$L_m = x_1^n L_{m,n} + \cdots + x_1 L_{m,1} + L_{m,0},$$

where  $L_{m,j} \in k[t, x_2, \dots, x_{m-1}] \langle \partial_t \rangle$  for each  $j \in \{1, \dots, n\}$ . To make the following computation clearer, we use “ $\circ$ ” to represent the composition of operators. Observe that

$$\partial_1 \circ x_1^j = x_1^j \circ \partial_1 + jx_1^{j-1}.$$

Thus,

$$\partial_1 \circ L_m = x_1^n \circ \partial_1 \circ L_{m,n} + \dots + x_1 \circ \partial_1 \circ L_{m,1} + \partial_1 \circ L_{m,0} + \sum_{j=1}^n jx_1^{j-1} \circ L_{m,j}.$$

Since  $\omega$  is closed, we have  $\partial_1(f_m) = \partial_m(f_1)$ . For each  $j \in \{1, \dots, n\}$ ,

$$x_1^j \circ \partial_1 \circ L_{m,j}(f_m) = x_1^j \circ L_{m,j} \circ \partial_m(f_1) = \partial_m(x_1^j L_{m,j}(f_1)).$$

Now define  $\tilde{L}_m := \partial_1 \circ L_{m,0} + \sum_{j=1}^n jx_1^{j-1} \circ L_{m,j}$ . Putting this all together and using  $\partial_1(L_m(f_m)) = \partial_1(\partial_m(g_m)) = \partial_m(\partial_1(g_m))$ , we obtain

$$\tilde{L}_m(f_m) = \partial_m \left( \partial_1(g_m) - \sum_{j=1}^n x_1^j L_{m,j}(f_1) \right),$$

which implies that  $\tilde{L}_m$  is a telescopers for  $f_m$  with respect to  $x_m$  with the degree in  $x_1$  at most  $n-1$ , leading to a contradiction. So  $L_m$  is free of  $x_1$ . The same argument holds for  $x_2, \dots, x_{m-1}$ . Hence our claim is true, i.e.,  $L_m \in k(t) \langle \partial_t \rangle$ .

Since  $\partial_i \circ L_m = L_m \circ \partial_i$  for each  $i \in \{1, \dots, m\}$ , we see that  $L_m(\omega)$  is closed with coefficients algebraic over  $k(t, x_1, \dots, x_m)$ . Applying  $L_m$  to  $\omega$  leads to

$$L_m(\omega) = L_m(f_1) dx_1 + \dots + L_m(f_{m-1}) dx_{m-1} + \partial_m(g_m).$$

By subtracting  $dg_m$ , we see that  $L_m(\omega) - dg_m$  is a closed algebraic differential form free of  $x_m$ . By induction, there exists a nonzero telescopers  $\tilde{L}$  with certificate  $\tilde{g}$  such that

$$\tilde{L}(L_m(\omega) - dg_m) = d\tilde{g},$$

which implies that

$$\tilde{L} \circ L_m(\omega) = d(\tilde{L}(g_m) + \tilde{g}).$$

Thus  $L := \tilde{L} \circ L_m$  is a nonzero telescopers for  $\omega$  with certificate  $g = \tilde{L}(g_m) + \tilde{g}$ .  $\square$

Next we present a reduction-based algorithm to construct a minimal telescopers for any closed rational 1-form  $\omega = f_1 dx_1 + \dots + f_m dx_m$  with  $f_i \in K(t)$ . By Section 3, write

$$\omega = dg + \tilde{\omega},$$

where  $g \in K(t)$  and  $\tilde{\omega} = \tilde{f}_1 dx_1 + \dots + \tilde{f}_m dx_m$  and each  $\tilde{f}_i$  is a rational function of  $x_i$  with squarefree denominator. According to [25] and the analysis above, we can construct a minimal telescopers  $L_m \in k(t) \langle \partial_t \rangle$  for  $\tilde{f}_m$  with respect to  $x_m$  such that

$$L_m(\tilde{f}_m) = \partial_m(g_m),$$

for some  $g_m \in K(t)$ . Using these ingredients, we give a recursive algorithm of computing the minimal telescopers for a closed rational 1-form.

**Algorithm CTOneForm:** Given a closed rational 1-form  $\omega = f_1 dx_1 + \dots + f_m dx_m$ , where  $f_i \in K$ , compute a minimal telescopers  $L \in K(t) \langle \partial_t \rangle$  and a certificate  $g$  of  $L$ .

- (1) Decompose  $\omega$  as  $\omega = d\tilde{g} + \tilde{\omega}$  by `HermiteOneForm` from Section 3. Set

$$\tilde{\omega} := \tilde{f}_1 dx_1 + \dots + \tilde{f}_m dx_m.$$

- (2) Compute a minimal telescopers  $L_m \in k(t) \langle \partial_t \rangle$  for  $\tilde{f}_m$  with respect to  $x_m$  and its certificate  $g_m$  by algorithms in [3, 5].
- (3) If  $m = 1$ , then return  $(L_m, L_m(\tilde{g}) + g_m)$ , otherwise, set  $\bar{\omega} := L_m(\tilde{\omega}) - dg_m$  and write

$$\bar{\omega} = \bar{f}_1 dx_1 + \cdots + \bar{f}_{m-1} dx_{m-1}, \text{ where } \bar{f}_i := L_m(\tilde{f}_i) - \partial_i(g_m) \text{ for } i = 1, \dots, m-1.$$

- (4) Decompose  $\bar{\omega}$  as  $\bar{\omega} = d_{m-1}\tilde{g} + \tilde{\omega}$  by `HermiteOneForm` from Section 3.
- (5) Run `CTOneForm` for  $\tilde{\omega}$  to get  $(\bar{L}, \bar{g})$ , where  $\bar{L}$  is of minimal order and  $\bar{g}$  is its certificate.
- (6) Return  $L := \bar{L} \circ L_m$ , and  $g := \bar{g} + \bar{L}(\tilde{g}) + \bar{L}(g_m) + \bar{L}(L_m(\tilde{g}))$ .

The minimality of  $L$  follows from Proposition 13 of [19].

**Example 7.3.** Let  $\omega$  be as the same 1-form as in Example 3.4. Then Hermite reduction gives  $\tilde{g} = 1/(xyz)$  and

$$\tilde{\omega} := \frac{t}{x} dx + \frac{t}{y} dy + \frac{t^2 + 1}{z} dz.$$

Then we compute the minimal telescopers for  $(t^2 + 1)/z$  with respect to  $z$ :

$$L_3 := (t^2 + 1)\partial_t - 2t,$$

and its certificate is  $g_3 = 0$ . Update  $\omega$  as

$$\bar{\omega} := L_3(\tilde{\omega}) - dg_3 = \frac{1 - t^2}{x} dx + \frac{1 - t^2}{y} dy.$$

Note that each coefficient of  $\bar{\omega}$  is already squarefree, so Hermite reduction for  $\bar{\omega}$  is not needed. The minimal telescopers for  $(1 - t^2)/y$  with respect to  $y$  is

$$\bar{L} := (t^2 - 1)\partial_t - 2t,$$

and its certificate is  $\bar{g} = 0$ . By computation, we obtain

$$\bar{L}(\bar{\omega}) = 0.$$

Thus

$$\bar{L}L_3(\omega) = \bar{L}L_3(d\tilde{g} + \tilde{\omega}) = d\bar{L}L_3(\tilde{g}) + \bar{L}(L_3(\tilde{\omega})) = d\bar{L}L_3(\tilde{g}) + \bar{L}(\bar{\omega}) = d\bar{L}L_3(\tilde{g}).$$

It follows that the minimal telescopers for  $\omega$  is

$$L := \bar{L}L_3 = (t^4 - 1)\partial_t^2 - (2t^3 + 2t)\partial_t + (2t^2 + 2),$$

and its certificate is  $g = \bar{L}L_3(\tilde{g}) = (2t^2 + 2)/xyz$ .

#### ACKNOWLEDGEMENTS

The authors are very grateful to Michael Singer for making this joint work possible and also to Pierre Lairez for many discussions about Picard's problem.

## REFERENCES

- [1] N. H. Abel. Démonstration d'une propriété générale d'une certaine classe de fonctions transcendentes. *J. Reine Angew. Math.*, 4:200–201, 1829.
- [2] N. H. Abel. Mémoire sur une propriété générale d'une classe très étendue de fonctions transcendantes. In L. Sylow and S. Lie, editors, *Œuvres Complètes de Niels Henrik Abel. Tome I*, pages 145–211. Grøndahl & Son, Christiania, 1881.
- [3] G. Almkvist and D. Zeilberger. The method of differentiating under the integral sign. *J. Symbolic Comput.*, 10(6):571–591, 1990.
- [4] J. Blümlein and C. Schneider. *Anti-differentiation and the calculation of Feynman amplitudes*. Texts and Monographs in Symbolic Computation. Springer, Cham, [2021] ©2021. Edited by Johannes Blümlein and Carsten Schneider.
- [5] A. Bostan, S. Chen, F. Chyzak, and Z. Li. Complexity of creative telescoping for bivariate rational functions. In *ISSAC 2010—Proceedings of the 2010 International Symposium on Symbolic and Algebraic Computation*, pages 203–210. ACM, New York, 2010.
- [6] A. Bostan, S. Chen, F. Chyzak, Z. Li, and G. Xin. Hermite reduction and creative telescoping for hyperexponential functions. In *ISSAC 2013—Proceedings of the 38th International Symposium on Symbolic and Algebraic Computation*, pages 77–84. ACM, New York, 2013.
- [7] A. Bostan, S. Chen, F. Chyzak, Z. Li, and G. Xin. Hermite reduction and creative telescoping for hyperexponential functions. In *ISSAC 2013—Proceedings of the 2013 International Symposium on Symbolic and Algebraic Computation*, pages 77–84. ACM, New York, 2013.
- [8] A. Bostan, F. Chyzak, P. Lairez, and B. Salvy. Generalized Hermite reduction, creative telescoping and definite integration of D-finite functions. In *ISSAC'18—Proceedings of the 2018 ACM International Symposium on Symbolic and Algebraic Computation*, pages 95–102. ACM, New York, 2018.
- [9] A. Bostan, F. Chyzak, P. Lairez, and B. Salvy. Generalized Hermite reduction, creative telescoping and definite integration of D-finite functions. In *ISSAC 2018—Proceedings of the 2018 ACM International Symposium on Symbolic and Algebraic Computation*, pages 95–102. ACM, New York, 2018.
- [10] A. Bostan, P. Lairez, and B. Salvy. Creative telescoping for rational functions using the Griffiths-Dwork method. In *ISSAC 2013—Proceedings of the 38th International Symposium on Symbolic and Algebraic Computation*, pages 93–100. ACM, New York, 2013.
- [11] R. Bott and L. W. Tu. *Differential Forms in Algebraic Topology*, volume 82 of *Graduate Texts in Mathematics*. Springer-Verlag, New York-Berlin, 1982.
- [12] M. Bronstein. *Symbolic Integration. I*, volume 1 of *Algorithms and Computation in Mathematics*. Springer-Verlag, Berlin, second edition, 2005.
- [13] P. Bürgisser and P. Scheiblechner. Differential forms in computational algebraic geometry. In *ISSAC 2007*, pages 61–68. ACM, New York, 2007.
- [14] P. Bürgisser and P. Scheiblechner. On the complexity of counting components of algebraic varieties. *J. Symbolic Comput.*, 44(9):1114–1136, 2009.
- [15] H. Cartan. *Differential Forms*. Houghton Mifflin Co., Boston, MA, 1970.
- [16] B. F. Caviness and M. Rothstein. A Liouville theorem on integration in finite terms for line integrals. *Comm. Algebra*, 3(9):781–795, 1975.
- [17] S. Chen, L. Du, and H. Fang. Symbolic summation of multivariate rational functions. *Foundations of Computational Mathematics*, Apr. 2025.
- [18] S. Chen, L. Du, and M. Kauers. Hermite reduction for D-finite functions via integral bases. In *ISSAC 2023—Proceedings of the 2023 International Symposium on Symbolic and Algebraic Computation*, pages 155–163. ACM, New York, [2023].
- [19] S. Chen, R. Feng, Z. Li, and M. F. Singer. Parallel telescoping and parameterized Picard-Vessiot theory. In *ISSAC 2014—Proceedings of the 2014 International Symposium on Symbolic and Algebraic Computation*, pages 99–106. ACM, New York, 2014.
- [20] S. Chen, R. Feng, Z. Li, M. F. Singer, and S. M. Watt. Telescopers for differential forms with one parameter. *Selecta Math. (N.S.)*, 30(3):Paper No. 36, 23, 2024.
- [21] S. Chen, H. Huang, M. Kauers, and Z. Li. A modified Abramov-Petkovšek reduction and creative telescoping for hypergeometric terms. In *ISSAC'15—Proceedings of the 2015 ACM International Symposium on Symbolic and Algebraic Computation*, pages 117–124. ACM, New York, 2015.

- [22] S. Chen, M. Kauers, and C. Koutschan. Reduction-based creative telescoping for algebraic functions. In *ISSAC 2016—Proceedings of the 2016 ACM International Symposium on Symbolic and Algebraic Computation*, pages 175–182. ACM, New York, 2016.
- [23] S. Chen, M. Kauers, and M. F. Singer. Telescopers for rational and algebraic functions via residues. In *ISSAC 2012—Proceedings of the International Symposium on Symbolic and Algebraic Computation*, pages 130–137. ACM, New York, 2012.
- [24] S. Chen and C. Koutschan. Proof of the Wilf-Zeilberger conjecture for mixed hypergeometric terms. *J. Symbolic Comput.*, 93:133–147, 2019.
- [25] S. Chen and M. F. Singer. Residues and telescopers for bivariate rational functions. *Adv. in Appl. Math.*, 49(2):111–133, 2012.
- [26] S. Chen, M. van Hoeij, M. Kauers, and C. Koutschan. Reduction-based creative telescoping for fuchsian D-finite functions. *J. Symbolic Comput.*, 85:108–127, 2018.
- [27] D. A. Cox. Notes on three formulas of Abel. *Bull. Amer. Math. Soc. (N.S.)*, 62(3):481–536, 2025.
- [28] A. Dimca. On the De Rham cohomology of a hypersurface complement. *American Journal of Mathematics*, 113(4):763–771, 1991.
- [29] B. Dwork. On the zeta function of a hypersurface. *Inst. Hautes Études Sci. Publ. Math.*, (12):5–68, 1962.
- [30] B. Dwork. On the zeta function of a hypersurface. II. *Ann. of Math. (2)*, 80:227–299, 1964.
- [31] H. M. Edwards. *Essays in Constructive Mathematics*. Springer, Cham, second edition, 2022. With contribution by David A. Cox.
- [32] G. Fu, M. Halás, U. Kotta, and Z. Li. Some remarks on Kähler differentials and ordinary differentials in nonlinear control theory. *Systems Control Lett.*, 60(9):699–703, 2011.
- [33] P. Griffiths and J. Harris. *Principles of algebraic geometry*. Pure and Applied Mathematics. Wiley-Interscience [John Wiley & Sons], New York, 1978.
- [34] P. A. Griffiths. On the periods of certain rational integrals. I, II. *Ann. of Math. (2)*, 90:460–495; 90 (1969), 496–541, 1969.
- [35] P. A. Griffiths. Variations on a theorem of Abel. *Invent. Math.*, 35:321–390, 1976.
- [36] C. Hermite. Sur l'intégration des fractions rationnelles. *Ann. Sci. École Norm. Sup. (2)*, 1:215–218, 1872.
- [37] S. L. Kleiman. What is Abel's theorem anyway? In O. A. Laudal and R. Piene, editors, *The Legacy of Niels Henrik Abel*, pages 395–440. Springer, Berlin, 2004.
- [38] E. R. Kolchin. *Differential algebra and algebraic groups*. Academic Press, New York-London, 1973. Pure and Applied Mathematics, Vol. 54.
- [39] P. Lairez. Computing periods of rational integrals. *Math. Comp.*, 85(300):1719–1752, 2016.
- [40] S. Lang. *Algebra*, volume 211 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, third edition, 2002.
- [41] J. I. Manin. Algebraic curves over fields with differentiation. *Izv. Akad. Nauk SSSR Ser. Mat.*, 22:737–756, 1958.
- [42] S. Müller-Stach, S. Weinzierl, and R. Zayadeh. Picard-Fuchs equations for Feynman integrals. *Comm. Math. Phys.*, 326(1):237–249, 2014.
- [43] M. V. Ostrogradskii. De l'intégration des fractions rationnelles. *Bull. de la classe physico-mathématique de l'Acad. Impériale des Sciences de Saint-Pétersbourg*, 4:145–167, 286–300, 1845.
- [44] E. Picard. *Quelques Applications Analytiques de la Théorie des Courbes et des Surfaces Algébriques*. Paris, Gauthier-Villars, 1931.
- [45] E. Picard and G. Simart. *Théorie des Fonctions Algébriques de Deux Variables Indépendantes. Tome I, II*. Chelsea Publishing Co., Bronx, NY, 1971. Réimpression corrigée (en un volume) de l'édition en deux volumes de 1897 et 1906.
- [46] C. G. Raab. Symbolic computation of parameter integrals. In *Proceedings of the 2016 ACM International Symposium on Symbolic and Algebraic Computation*, ISSAC '16, page 13–15, New York, NY, USA, 2016. Association for Computing Machinery.
- [47] C. G. Raab and M. F. Singer, editors. *Integration in finite terms: fundamental sources*. Texts and Monographs in Symbolic Computation. Springer, Cham, [2022] ©2022.
- [48] R. H. Risch. The problem of integration in finite terms. *Trans. Amer. Math. Soc.*, 139:167–189, 1969.
- [49] M. Rosenlicht. Liouville's theorem on functions with elementary integrals. *Pacific J. Math.*, 24:153–161, 1968.

- [50] M. Spivak. Calculus on Manifolds. A Modern Approach to Classical Theorems of Advanced Calculus. W. A. Benjamin, Inc., New York-Amsterdam, 1965.
- [51] B. M. Trager. Integration of algebraic functions. In Integration in finite terms: fundamental sources, Texts Monogr. Symbol. Comput., pages 235–296. Springer, Cham, [2022] ©2022.
- [52] J. van der Hoeven. Constructing reductions for creative telescoping: the general differentially finite case. Appl. Algebra Engrg. Comm. Comput., 32(5):575–602, 2021.
- [53] A. van der Poorten. A proof that Euler missed ... Apéry’s proof of the irrationality of  $\zeta(3)$ . Math. Intelligencer, 1(4):195–203, 1978/79.
- [54] S. H. Weintraub. Differential Forms. Elsevier/Academic Press, Amsterdam, second edition, 2014.
- [55] D. Zeilberger. A fast algorithm for proving terminating hypergeometric identities. Discrete Math., 80(2):207–211, 1990.
- [56] D. Zeilberger. The method of creative telescoping. J. Symbolic Comput., 11(3):195–204, 1991.

KLMM, ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES, and SCHOOL OF MATHEMATICS, UNIVERSITY OF CHINESE ACADEMY OF SCIENCES, BEIJING, 100190, CHINA

*Email address:* schen@amss.ac.cn

DEPARTMENT OF MATHEMATICS, AMHERST COLLEGE, AMHERST, MA 01002, USA

*Email address:* dacox@amherst.edu

KLMM, ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES, and SCHOOL OF MATHEMATICS, UNIVERSITY OF CHINESE ACADEMY OF SCIENCES, BEIJING, 100190, CHINA

*Email address:* wangyisen@amss.ac.cn