

Avoiding Intersections of Dragon Curves

REIMUND ALBERS, ZONGYI GUO, HUAIYI GUO¹

Abstract: This article proves that there are no self-intersections in the dragon curve when the unfolding angle is greater than 98.195° . This is shown by constructing a hull for the dragon curve that is mapped onto itself by the generating mappings for the dragon curve. The treatment is purely geometric. The proof is supplemented by a conjecture that reduces the boundary for the unfolding angle to 96.241°

1. Introduction

The paperfolding polygon, when unfolded by exactly 90° , exhibits no intersections, meaning that no segment of the broken line doubles back on itself¹. The question if intersections are avoided or not when the unfolding angle exceeds 90° first came to attention² in 1990, stemming from M. Mendès France's intuitive assumption, that no intersections occur with unfolding angles greater than 90° . Computer experiments conducted around 1995 by H.-O. Peitgen and S. Michaelis provided indications of intersections occurring for unfolding angles between 90° and 95° . In his 2011 book¹, D. Knuth noted that he "*noticed in 1969 that 95° -angle folds would lead to paths that cross themselves*". In 2006, Albers presented a proof³ demonstrating the existence of crossing segments in iteration 10 for angles ranging between 90° and 95.126° . It is additionally demonstrated³ that certain vertices of the paper-folding curve coincide when the unfolding angle is approximately 93.06° . In 2014, S. Tabachnikov raises in "Dragon curves revisited"⁴ the question: "..., what is the value of the critical angle for which the curve starts to cross itself, and where does this self-crossing occur?" In 2018, J.-P. Allouche, M. Mendès France, and G. Skordev provide a comprehensive overview⁵ of the problem and its solutions to date. In 2022, Y. Kamiya constructs a sequence of angles⁶ with the limit of 93.7912° , where two vertices of the paper-folding curve precisely coincide. Finally, in January 2024, S. Akiyama, Y. Kamiya, and F. Wen establish that no intersections occur⁷ when the unfolding angle exceeds 99.3438° . Their work and the present article address the same problem and use the same big picture. While their approach employs complex calculations focused on the limit Dragon curve, this article presents a purely geometric analysis of the finite paperfolding polygons.


The current findings are summarized in Figure 1.

¹Universität Bremen. MZH Bibliothekstr. 5, 28359 Bremen, Germany


Gymnasium Horn, Vorkampsweg 97, 28359 Bremen, Germany

E-mail addresses: ralbers@uni-bremen.de, zy.guo@dxfund.org, hy.guo@dxfund.org

Keywords: Highway Dragon, Dragon Curve



In this article the gap of uncertainty is narrowed as follows.



2. Basics

2. 1. Defining the paperfolding process as a sequence of symbols

The paperfolding polygons are generated by iteratively folding a strip of paper in the same direction and analyzing the resulting creases. This process generates a sequence of up and downs or hills and valleys. Referring to the geometric interpretation (see Definition 3) a down (valley) is denoted as L and an up (hill) as R. If the strip is orientated so that the first crease forms a down (= L), the sequence of symbols starts with LLRLLRLLLRLRR

Definition 1

Let $\sigma: \mathbb{N} \rightarrow \{L, R\}$ be the function that assigns the symbol at position n in the sequence.

This function allows us to define the paperfolding sequence.

Definition 2 (Inflation Law)

The paperfolding sequence consists of symbols L and R determined by:

$$\sigma(n) = \begin{cases} L & \text{if } n \equiv 1 \pmod 4 \\ R & \text{if } n \equiv 3 \pmod 4 \\ \sigma\left(\frac{n}{2}\right) & \text{if } n \equiv 0 \pmod 2 \end{cases} \quad n \in \mathbb{N}$$

There is an alternative method to construct the sequence.

Lemma 1 (Reflection Law)

The paperfolding sequence can be defined by

$\sigma(2^k) = L, k \in \mathbb{N}_0$ and $\sigma(2^k - d) = \overline{\sigma(2^k + d)}, d \in \{1, 2, \dots, 2^k - 1\}$,
where \bar{S} denotes the opposite symbol to the symbol $S \in \{L, R\}$.

Proof

See R. Albers, Papierfalten ³, chapter 3

2. 2. Defining the paperfolding process as a sequence of polygons

For a more geometric perspective, consider the folded paper strip as a polygon where creases form vertices and paper segments form edges.

Definition 3 (sequence of paperfolding polygons)

The polygon is dependent on an angle α , called the unfolding angle.

Q_n is a polygon of $2^n + 1$ vertices

$P_{n,0}$ to $P_{n,2^n}$ and their connecting segments

$P_{n,k}P_{n,k+1}$, $k \in \{0, 1, 2, \dots, 2^n - 1\}$,

meeting at the angle α . (Figure 3)

Q_0 is the straight line from $P_{0,0}$ to

$P_{0,1}$.

Q_{n+1} is constructed from Q_n by

- Renumbering the vertices $P_{n,k}$ to

$$P_{n+1,2k}$$

- Replacing each segment $\overline{P_{n,k}P_{n,k+1}}$ by the sides of an isosceles triangle, meeting at

$$P_{n+1,2k+1} \text{ under the angle } \alpha.$$

- The isosceles triangle is added

i. to the left side, forming a right turn, if $\overline{P_{n,k}P_{n,k+1}}$ is for odd k to even $k+1$.

ii. to the right side, forming a left turn, if $\overline{P_{n,k}P_{n,k+1}}$ is for even k to odd $k+1$.




Figure 3: Geometric inflation


Conclusion

- Each polygon begins at $P_{0,0}$ and ends at $P_{0,1}$.
- This geometric construction of Q_n aligns with the inflation law (Definition 2).

Definition 4

$$\begin{aligned} a) \quad \beta &= |\angle P_{n,k+1}P_{n,k}P_{n+1,2k+1}| \\ &= \frac{180^\circ - \alpha}{2} = 90^\circ - \frac{\alpha}{2} \end{aligned}$$

$$b) \quad q = \frac{|P_{n,k}P_{n+1,2k+1}|}{|P_{n,k}P_{n,k+1}|} = \frac{1}{2\cos\beta}$$



With the parameters from Definition 4, two essential mappings can now be defined.

Figure 4: Definition of β and q

Definition 5

Mapping π_0 : The rotation around $P_{0,0}$ by $-\beta$ combined with the dilation at center $P_{0,0}$ and scale factor q .

Mapping π_1 : The rotation around $P_{0,0}$ by $\beta - 180^\circ$ combined with the dilation at center $P_{0,0}$ and scale factor q , then the translation by $\overrightarrow{P_{0,0}P_{0,1}}$.

Remark

Since the angles $-\beta$ and $\beta - 180^\circ$ are both negative, the rotations are clockwise.

Lemma 2

The sequence of polygons Q_n is generated recursively by

$$Q_0 = \overline{P_{0,0}P_{0,1}} \text{ and } Q_{n+1} = \pi_0(Q_n) \cup \pi_1(Q_n).$$

Proof

For the special case $\beta = 45^\circ$ see R. Albers, Papierfalten ³, chapter 8.

Remark

Lemma 2 corresponds to the reflection law (Lemma 1)

To facilitate further analysis, the segments get an orientation. The direction flows from vertices $P_{n,k}$ with an even index k to those with an odd index $k+1$ or $k-1$. As illustrated in Figure 5, alternate segments are oriented retrograde to their predecessors.




Figure 5: Orientated segments (k is even)

Lemma 3 (orientated segments)

When following the established segment orientation, all triangles in the construction of Q_{n+1} from Q_n are consistently inserted on the right-hand side of the directed segments.

Proof

This follows directly from Definition 3 c), see Figure 5

Preview: Avoiding intersections of the paperfolding polygons Q_n

This article shows (in Proposition 3 and Theorem 1), that for unfolding angles $98.195^\circ \leq \alpha \leq 108^\circ$ the paperfolding polygons Q_n avoid intersections.

The proof strategy relies on constructing a set A_{PF} that contains all polygons Q_n as subsets. If the images under the mappings π_0 and π_1 of this set are clearly separated,

i.e. $\pi_0(A_{PF}) \cap \pi_1(A_{PF}) = P_{1,1}$,
then no intersections occur in the paperfolding polygons.




Figure 6: The idea of the proof

The complete formal proof follows after Lemma 10.

3. Points on a straight line


Lemma 4a

Let k be an odd integer. If there is a left turn at $P_{n,k}$, then the three points $P_{n,k-1}, P_{n,k} = P_{n+4,16k}$ and $P_{n+4,16k+1}$ are collinear. (Figure 7)

Proof

As k is odd, the segment $\overline{P_{n,k}P_{n,k+1}}$ is orientated toward $P_{n,k}$. The inflation process, generating Q_{n+1} from Q_n , adds $\overline{P_{n+1,2k}P_{n+1,2k+1}}$ so that it is rotated from $\overline{P_{n,k}P_{n,k+1}}$ by β . Subsequent inflation steps generate segments rotated by $-\beta$. So, the angle (absolute value) between $\overline{P_{n,k}P_{n,k+1}}$ and $\overline{P_{n+4,16k}P_{n+4,16k+1}}$ is 2β .

As $\alpha + 2\beta = 180^\circ$, the points $P_{n,k-1}, P_{n,k} = P_{n+4,16k}$ and $P_{n+4,16k+1}$ are on a straight line. \square



Conclusion

The point $P_{n+4,16k+1}$ initiates another left turn (see

Figure 7: New segments after a left turn

Definition 2). By iterating this property, it yields $P_{n+8,256k+17}$ on the same straight line. This can be continued indefinitely.

Lemma 4b

Let k be an odd integer. If there is a right turn at $P_{n,k}$, then the three points $P_{n,k+1}, P_{n,k} = P_{n+4,16k}$ and $P_{n+4,16k-1}$ are collinear. (Figure 8)

Proof

As k is odd, the segment $\overline{P_{n,k}P_{n,k-1}}$ is orientated toward $P_{n,k}$. The inflation process, generating Q_{n+1} from Q_n , adds $\overline{P_{n+1,2k}P_{n+1,2k-1}}$ so that it is rotated from $\overline{P_{n,k}P_{n,k-1}}$ by β . Subsequent inflation steps generate segments rotated by $-\beta$. So, the angle (absolute value) between $\overline{P_{n,k}P_{n,k-1}}$ and $\overline{P_{n+4,16k}P_{n+4,16k-1}}$ is 2β .

As $\alpha + 2\beta = 180^\circ$, the points $P_{n,k+1}, P_{n,k} = P_{n+4,16k}$ and $P_{n+4,16k-1}$ are on a straight line. \square




Figure 8: New segments before a right turn

Conclusion

The point $P_{n+4,16k-1}$ initiates another right turn (see Definition 2). By iterating this property, it yields $P_{n+8,256k-17}$ on the same straight line. This can be continued indefinitely.

In the following part of the article logarithmic spirals are intensively used. So, it is useful to denote some

Basic facts for logarithmic spirals


In a polar coordinate system, a logarithmic spiral with the origin C as center is described by

$$r(\varphi) = ab^\varphi, a, b > 0$$

For every point P of the spiral the angle τ between the radius CP and the tangent to the spiral at P is constant and given by

$$\tan(\tau) = \frac{1}{\ln b}$$

This yields for two spirals with center C resp. C' and the same base b : If P and P' are two points of the spirals and collinear with C and C' , then the tangents t and t' to the spirals at P and P' are parallel. If $P = P'$, then the tangent is a common tangent to both spirals.



Let base $b > 1$ and φ_P be the polar angle of P measured from center C . Let t be the tangent to the spiral at P . Then all points of the spiral with a polar angle $\varphi \leq \varphi_P + \tau$ are with respect to t in the half-plane containing center C .

The dilation of a spiral at its center C by the factor s and the rotation around C by the angle ρ transforms the describing formula from

$$r(\varphi) = a b^\varphi, \varphi_{min} \leq \varphi \leq \varphi_{max} \text{ to } \tilde{r}(\varphi) = s a b^{\varphi-\rho}, \varphi_{min} + \rho \leq \varphi \leq \varphi_{max} + \rho$$

4. Constructing a borderline for the polygons Q_n

Construction 1: The maximal distance to $P_{0,0}$

According to Lemma 4a) the three points $P_{n,k-1}, P_{n,k} = P_{n+4,16k}$ and $P_{n+4,16k+1}$ are collinear. After four inflation steps, this pattern repeats with additional points on the same line. If the length of $\overline{P_{n,k-1}P_{n,k}}$ is L , the total length of consecutive segments on this line follows a geometric series: $L(1 + q^4 + q^8 + q^{12} + \dots) = L \frac{1}{1-q^4}$ (for q see Definition 4).

Applying this to $\overline{P_{n,k-1}P_{n,k}} = \overline{P_{1,0}P_{1,1}}$ there exists a straight line from $P_{0,0}$ to $P_{1,1}$ to $P_{5,17}, P_{9,273}, P_{13,4369}, \dots$ (see Figure 9). Given $|P_{0,0}P_{0,1}| = 1 \Rightarrow |P_{1,0}P_{1,1}| = q$, the distance from $P_{0,0}$ through $P_{1,1}$ has in the limit the length of $q \frac{1}{1-q^4}$.

Then applying it to $\overline{P_{n,k-1}P_{n,k}} = \overline{P_{2,0}P_{2,1}}$ and compared to $\overline{P_{1,0}P_{1,1}}$, this segment is rotated by $-\beta$ (clockwise) and dilated by q . Therefore, the limit length becomes $q^2 \frac{1}{1-q^4}$. Iterating this process generates a spiral, that serves as an outer boundary for the paperfolding polygons

around $P_{0,0}$. In a polar system with origin in $P_{0,0}$ and $\overrightarrow{P_{0,0}P_{0,1}}$ defines the direction of $\varphi = 0$ this spiral is given by

$$a_0(\varphi) = \frac{1}{1-q^4} q^{-\frac{\varphi}{\beta}}, \varphi \leq -\beta$$




Figure 9: The polygon Q_{12} with the outer and inner spiral at $P_{0,0}$

Construction 2: The minimal distance to $P_{0,0}$

According to Lemma 4b) the three points $P_{n,k+1}$, $P_{n,k} = P_{n+4,16k}$ and $P_{n+4,16k-1}$ are collinear. After four inflation steps, this pattern repeats with additional points on the same line. If the length of $\overline{P_{n,k+1}P_{n,k}}$ is L , the total length of consecutive segments on this line follows a geometric series: $L(1 + q^4 + q^8 + q^{12} + \dots) = L \frac{1}{1-q^4}$.

Applying this to $\overline{P_{n,k+1}P_{n,k}} = \overline{P_{2,4}P_{2,3}}$ (remember $P_{2,4} = P_{0,1}$) there exists a straight line from $P_{0,1}$ to $P_{2,3}$ to $P_{6,47}$, $P_{10,751}$, ... (see Figure 9). Given $|P_{0,0}P_{0,1}| = 1 \Rightarrow |P_{2,3}P_{0,1}| = q^2$ and the length is in the limit $\frac{q^2}{1-q^4}$. As $P_{0,0}$ lies on the same line, the distance from this limit to $P_{0,0}$ is $\left(1 - \frac{q^2}{1-q^4}\right)$.

Applying this to the next right turn at $P_{n,k} = P_{3,3}$. It yields $P_{n,k+1} = P_{3,4} = P_{1,1}$. Relative to $\overline{P_{2,4}P_{2,3}}$ the straight line is rotated by $-\beta$ (clockwise) and everything is shortened by q , resulting in a minimal distance to $P_{0,0}$ of $q \left(1 - \frac{q^2}{1-q^4}\right)$. Iterating this process generates a spiral that serves as the minimum distance of the paperfolding polygons to $P_{0,0}$.

In the same coordinate system of construction 1 it is given by

$$b_0(\varphi) = \left(1 - \frac{q^2}{1-q^4}\right) q^{-\frac{\varphi}{\beta}}, \varphi \leq 0$$

Construction 3: The maximal and minimal distance to $P_{0,1}$

According to Lemma 2 every polygon Q_n is generated by the mappings π_0 and π_1 .

Since $P_{0,1} = \pi_1(P_{0,0})$, the spirals a_0 and b_0 are transformed by π_1 .

π_1 : dilation by q , rotation by $\beta - 180^\circ = -\alpha - \beta$

$$a_0(\varphi) = \frac{1}{1-q^4} q^{-\frac{\varphi}{\beta}}, \varphi \leq -\beta \xrightarrow{\pi_1} c_0(\varphi) = \frac{1}{1-q^4} q^{-\frac{\varphi+\alpha}{\beta}}, \varphi \leq -180^\circ$$

For the further argumentation the domain for c_0 is extended to $\varphi \leq -180^\circ + \beta = -\alpha - \beta$.

$$b_0(\varphi) = \left(1 - \frac{q^2}{1-q^4}\right) q^{-\frac{\varphi}{\beta}}, \varphi \leq 0 \xrightarrow{\pi_1} d_0(\varphi) = \left(1 - \frac{q^2}{1-q^4}\right) q^{-\frac{\varphi+\alpha}{\beta}}, \varphi \leq -\alpha - \beta$$

The origin of the underlying coordinate system for c_0 and d_0 is translated to $P_{0,1}$.

Construction 4: Closing the gap between a_0 and d_0

Since $P_{1,1} = \pi_1(P_{0,1})$, the spiral c_0 is transformed by π_1 .

π_1 : dilation by q , rotation by $\beta - 180^\circ = -\alpha - \beta$

$$c_0(\varphi) = \frac{1}{1-q^4} q^{-\frac{\varphi+\alpha}{\beta}}, \varphi \leq -180^\circ \xrightarrow{\pi_1} e_0(\varphi) = \frac{1}{1-q^4} q^{-\frac{\varphi+2\alpha}{\beta}}, \varphi \leq -360^\circ + \beta$$

Applying the parameter-transformation $\varphi \rightarrow \varphi - 360^\circ = \varphi - (2\alpha + 4\beta)$ yields

$$e_0(\varphi) = \frac{1}{1-q^4} q^{-\frac{\varphi-4\beta}{\beta}}, \varphi \leq \beta$$

The origin of the underlying coordinate system is translated to $P_{1,1}$.

For a complete encompassing of the paperfolding polygons, only the part of the spiral is required until it reaches a_0 at the polar angle $-\beta$. Therefore, the interval for φ is restricted to $-\beta \leq \varphi \leq \beta$.




Figure 10: The five parts of the borderline

5. Defining a set, that is a hull for the paperfolding polygons




Figure 11: The four sets of points

Definition 6

a) F is the point on $\overline{P_{0,0}P_{0,1}}$ with $|P_{0,0}F| = b_0(0^\circ) = 1 - \frac{q^2}{1-q^4}$.

D is the point on $\overline{P_{0,0}P_{0,1}}$ with $|P_{0,0}D| = q$.

H is the point on c_0 with polar angle $-\alpha - \beta$.

b) For $90^\circ \leq \alpha \leq 108^\circ$ (corresponding to $45^\circ \geq \beta \geq 36^\circ, \frac{1}{2}\sqrt{2} \geq q \geq \frac{\sqrt{5}-1}{2}$)

the set of points, that is delimited by the encompassing spirals constructed above, consist of four parts:

$(d_{i,k}; \varphi)$ are the polar coordinates of a point, $d_{i,k}$ is the distance to the point $P_{i,k}$

$$S_1 = \{(d_{0,0}; \varphi) \in \mathbb{R}^2 \mid b_0(\varphi) \leq d_{0,0} \leq a_0(\varphi), \varphi \leq -\beta\}$$

$$S_2 = \{(d_{0,0}; \varphi) \in \mathbb{R}^2 \mid b_0(\varphi) \leq d_{0,0} \leq q, -\beta \leq \varphi \leq 0^\circ\}$$

$$S_3 = \{(d_{0,1}; \varphi) \in \mathbb{R}^2 \mid d_0(\varphi) \leq d_{0,1} \leq c_0(\varphi), \varphi \leq -\alpha - \beta\}$$

$$S_4 = \{(d_{1,1}; \varphi) \in \mathbb{R}^2 \mid 0 \leq d_{1,1} \leq e_0(\varphi), -\beta \leq \varphi \leq \beta\}$$

Then the total set of points is

$$A_{PF} = S_1 \cup S_2 \cup S_3 \cup S_4$$

This set constitutes a hull for all paperfolding polygons Q_n of a fixed unfolding angle α . Due to its characteristic shape, we will refer to it as the “duck-shaped hull” or, concisely, the “duck”.

Lemma 5

Let D and H be the points defined in Definition 6 (see Figure 11). For $\frac{1}{2}\sqrt{2} \geq q \geq \frac{\sqrt{5}-1}{2}$

- a) $D \in S_3$.
- b) $H \in S_1$

Proof

a) $D \in S_3$ holds, if $d_0(-180^\circ) \leq 1 - q \leq c_0(-180^\circ)$

$$\Leftrightarrow \left(1 - \frac{q^2}{1-q^4}\right)q^2 \leq 1 - q \leq \frac{1}{1-q^4}q^2 \text{ which holds for } \frac{1}{2}\sqrt{2} \geq q \geq \frac{\sqrt{5}-1}{2}.$$

b) $H \in S_1$ holds, if $a_0(\lambda) \leq |P_{0,0}H| \leq b_0(\lambda)$, where λ is the polar angle for $\overline{P_{0,0}H}$.

H is defined by $|P_{0,1}H| = c_0(-\alpha - \beta)$. Applying the law of cosine in the triangle $P_{0,0}HP_{0,1}$

yields $|P_{0,0}H| = q \frac{\sqrt{1 - q^2 + q^6}}{1-q^4}$ and $\tan|\lambda| = \frac{\sqrt{4q^2-1}}{1-2q^4}$. A numerical analysis shows that

$$a_0\left(-\arctan\left(\frac{\sqrt{4q^2-1}}{1-2q^4}\right)\right) \leq q \frac{\sqrt{1 - q^2 + q^6}}{1-q^4} \leq b_0\left(-\arctan\left(\frac{\sqrt{4q^2-1}}{1-2q^4}\right)\right) \text{ holds for } \frac{1}{2}\sqrt{2} \geq q \geq \frac{\sqrt{5}-1}{2}.$$

Definition 7

Generalization of A_{PF}

Let π be the composition of the maps π_0 and π_1 such that

$$\overline{P_{n,k}P_{n,k+1}} = \pi(\overline{P_{0,0}P_{0,1}}), \quad k \in \{0, 1, 2, \dots, 2^n - 1\}.$$

Then $A_{PF}(\overline{P_{n,k}P_{n,k+1}})$ is defined by $A_{PF}(\overline{P_{n,k}P_{n,k+1}}) = \pi(A_{PF})$.

So, $A_{PF}(\overline{P_{n,k}P_{n,k+1}})$ is a contracted copy of A_{PF} orientated at the segment $\overline{P_{n,k}P_{n,k+1}}$. The initial set A_{PF} can be written as $A_{PF}(\overline{P_{0,0}P_{0,1}})$.

For further argumentation additional subsets of A_{PF} are defined.

Definition 8 (see Figure 12)

a) Points:

A is the intersection of the straight line $P_{0,0}P_{1,1}$ with $a_0, |P_{0,0}A| = a_0(-\beta)$.

E is the intersection of the straight line $P_{0,0}P_{1,1}$ with $a_0, |P_{0,0}E| = a_0(-180^\circ - \beta)$.

B is the intersection of the straight line $P_{0,0}P_{1,1}$ with $b_0, |P_{0,0}B| = b_0(-\beta)$.

C is the intersection of the straight line $P_{0,1}P_{1,1}$ with $d_0, |P_{0,1}C| = d_0(-\alpha - \beta)$.

b) Lines:

As $|P_{1,1}B| = |P_{1,1}C| = \frac{q^3}{1-q^4}$ there exists a circular arc \widehat{BC} with midpoint $P_{1,1}$.

t is the tangent to b_0 at B.

e_0' extends e_0 to cover polar angles from $-\beta$ to $-180^\circ - \beta$ (center $P_{1,1}$).

The intersection of arc \widehat{BC} with the tangent t is G.

c) Sets:

$$T_1 = \{(d_{1,1}; \varphi) \in \mathbb{R}^2 \mid 0 \leq d_{1,1} \leq e_0'(\varphi), -\beta \geq \varphi \geq -180^\circ - \beta\}$$

T_2 is the set of points bounded by arc \widehat{CG} (center $P_{1,1}$, radius $\frac{q^3}{1-q^4}$), segment \overline{GB} , segment $\overline{BP_{1,1}}$ and segment $\overline{P_{1,1}C}$. Therefore, every point of T_2 has a polar angle φ to $P_{1,1}$ which satisfies $-180^\circ - \beta \geq \varphi \geq -180^\circ - \beta - \alpha$.




Figure 12: Subsets of A_{PF}

Lemma 6

For $90^\circ \leq \alpha \leq 108^\circ$, $\frac{1}{2}\sqrt{2} \geq q \geq \frac{\sqrt{5}-1}{2}$ (see Figure 13)

- a) $\text{arc } \widehat{BC} \cap \overline{P_{0,0}P_{0,1}} = \emptyset$
- b) Point G is an inner point on arc \widehat{BC} .

Proof

- a) The radius of arc \widehat{BC} is $\frac{q^3}{1-q^4}$, the distance of $P_{1,1}$ to $\overline{P_{0,0}P_{0,1}}$ is $q \sin \beta = q \sqrt{1 - \frac{1}{4q^2}}$.

The condition $\frac{q^3}{1-q^4} < q \sqrt{1 - \frac{1}{4q^2}}$ can be simplified to

$$1 - 4q^2 - 2q^4 + 12q^6 + q^8 - 4q^{10} < 0, \text{ which holds for } 0.724 \geq q \geq 0.524.$$

- b) τ is the angle between the tangent to the spiral and the ray $\overrightarrow{P_{0,0}B}$.

$$\text{Therefore, } \tan \tau = -\frac{\beta}{\ln q} = \frac{\beta}{\ln(2 \cos \beta)}$$

G lies between C and B is equivalent to $\tau > \beta$.

This can be transformed to

$$\frac{\beta}{\ln(2 \cos \beta)} > \tan \beta.$$

This holds for all $0^\circ < \beta < 45^\circ$




Figure 13: Angle τ and point G

Lemma 7

- a) $T_1 \subset S_1 \subset A_{PF}$
- b) $T_2 \subset (S_2 \cup S_3) \subset A_{PF}$

Proof (see Figure 12)

a) Let T_1^* be the area bounded by the spiral arc \widehat{AE} and the line segment \overline{EA} . The dilation with center A and scaling factor q^4 maps T_1^* to T_1 because:

$$|AP_{0,0}| = a_0(-\beta) = \frac{1}{1-q^4} q^{-\frac{-\beta}{\beta}} = \frac{1}{1-q^4} q \quad |AP_{1,1}| = e_0(-\beta) = \frac{1}{1-q^4} q^{-\frac{-\beta-4\beta}{\beta}} = \frac{1}{1-q^4} q^5$$

So, the center $P_{0,0}$ is mapped to the center $P_{1,1}$.

$$|P_{0,0}P| = a_0(\varphi) = \frac{1}{1-q^4} q^{-\frac{\varphi}{\beta}} \quad |P_{1,1}P'| = e_0'(\varphi) = \frac{1}{1-q^4} q^{-\frac{\varphi-4\beta}{\beta}} = \frac{1}{1-q^4} q^{-\frac{\varphi}{\beta}} q^4$$

So, every point P on a_0 is mapped to a point P' on e_0' .

T_1^* is convex and $A \in T_1^*$, so $T_1 \subset T_1^*$.

Now it has to be shown for all points of T_1 that the distance to $P_{0,0}$ is larger than the distance of an according point on b_0 to $P_{0,0}$.

Let P be a point of T_1 . Applying the triangle inequality:

$$|P_{0,0}P| \geq |P_{0,0}P_{1,1}| - |P_{1,1}P| \geq q - |P_{1,1}A| = q - \frac{1}{1-q^4} q^5$$

$$\max(b_0(\varphi): -\beta \geq \varphi \geq -180^\circ - \beta) = b_0(-\beta) = \left(1 - \frac{q^2}{1-q^4}\right) q = q - \frac{1}{1-q^4} q^3$$

Therefore, $|P_{0,0}P| \geq \max(b_0(\varphi): -\beta \geq \varphi \geq -180^\circ - \beta)$, which implies $T_1 \subset S_1$. \square

b) Let $P \in T_2$

Case 1: $|P_{0,0}P| \leq q$

The definition of S_2 directly yields $P \in S_2$.

Case 2: $|P_{0,0}P| > q$

The definition of T_2 yields $|P_{1,1}P| \leq \frac{q^3}{1-q^4}$.

Now it has to be shown for P that the distance to $P_{0,1}$ is equal or larger than the distance of an according point on d_0 to $P_{0,1}$. Applying the triangle inequality:

$$|PP_{0,1}| \geq |P_{1,1}P_{0,1}| - |P_{1,1}P| \geq q - \frac{q^3}{1-q^4}$$

$$\max(d_0(\varphi): -180^\circ + \beta \geq \varphi \geq -180^\circ) = d_0(-180^\circ + \beta) = q - \frac{q^3}{1-q^4}$$

Therefore, $|PP_{0,1}| \geq \max(d_0(\varphi): -180^\circ + \beta \geq \varphi \geq -180^\circ)$ which implies $P \in S_3$. \square

Proposition 1

a) $\pi_0(A_{PF}) \subset A_{PF}$

b) $\pi_1(A_{PF}) \subset A_{PF}$

Proof a)

Since π_0 preserves the ratio, it suffices to examine the mapping of the encompassing spirals.

π_0 : A dilation by q and a rotation by $-\beta$, both with center $P_{0,0}$

points: $P_{0,0} \rightarrow P_{0,0}$ $P_{0,1} \rightarrow P_{1,1}$ $P_{1,1} \rightarrow P_{2,1}$

spirals:

$$a_0(\varphi) = \frac{1}{1-q^4} q^{-\frac{\varphi}{\beta}}, \varphi \leq -\beta, \text{ center } P_{0,0}$$

$$\rightarrow a_1(\varphi) = \frac{1}{1-q^4} q^{-\frac{\varphi}{\beta}}, \varphi \leq -2\beta, \text{ center } P_{0,0}$$

$$b_0(\varphi) = \left(1 - \frac{q^2}{1-q^4}\right) q^{-\frac{\varphi}{\beta}}, \varphi \leq 0, \text{ center } P_{0,0}$$

$$\rightarrow b_1(\varphi) = \left(1 - \frac{q^2}{1-q^4}\right) q^{-\frac{\varphi}{\beta}}, \varphi \leq -\beta, \text{ center } P_{0,0}$$

$$\begin{aligned}
c_0(\varphi) &= \frac{1}{1-q^4} q^{-\frac{\varphi+\alpha}{\beta}}, \varphi \leq -180^\circ, \text{center } P_{0,1} \\
&\rightarrow c_1(\varphi) = \frac{1}{1-q^4} q^{-\frac{\varphi+\alpha}{\beta}}, \varphi \leq -180^\circ - \beta, \text{center } P_{1,1} \\
d_0(\varphi) &= \left(1 - \frac{q^2}{1-q^4}\right) q^{-\frac{\varphi+\alpha}{\beta}}, \varphi \leq -\alpha - \beta, \text{center } P_{0,1} \\
&\rightarrow d_1(\varphi) = \left(1 - \frac{q^2}{1-q^4}\right) q^{-\frac{\varphi+\alpha}{\beta}}, \varphi \leq -180^\circ, \text{center } P_{1,1} \\
e_0(\varphi) &= \frac{1}{1-q^4} q^{-\frac{\varphi-4\beta}{\beta}}, -\beta \leq \varphi \leq \beta, \text{center } P_{1,1} \\
&\rightarrow e_1(\varphi) = \frac{1}{1-q^4} q^{-\frac{\varphi-4\beta}{\beta}}, -2\beta \leq \varphi \leq 0^\circ, \text{center } P_{2,1}
\end{aligned}$$




Figure 14: $\pi_0(A_{PF})$

The dilation by q and the rotation by $-\beta$ cancel each other out. The spirals are just translated to different centers and the interval for φ is modified.

a_0 and b_0 are mapped to themselves.

c_1 is tangent to b_0 at B with common tangent line t . $|P_{0,0}B| = b_0(-\beta) = \left(1 - \frac{q^2}{1-q^4}\right) q$

$|BP_{1,1}| = c_1(-180^\circ - \beta) = \frac{1}{1-q^4} q^{-\frac{-180^\circ - \beta + \alpha}{\beta}} = \frac{1}{1-q^4} q^3$, yielding $|P_{0,0}B| + |BP_{1,1}| = q$.

For $-180^\circ - \beta \geq \varphi \geq -180^\circ - \beta - \alpha$ c_1 is a subset of T_2 , as

$c_1(\varphi) \leq |BP_{1,1}| = \frac{1}{1-q^4} q^3$ and c_1 is in the same half-plane as $P_{1,1}$ with respect to t .

For $-180^\circ - \beta - \alpha = -360^\circ + \beta \geq \varphi \geq -360^\circ - \beta$ c_1 is a subset of S_4 as

$c_1(\varphi) = \frac{1}{1-q^4} q^{-\frac{\varphi+\alpha}{\beta}} \leq e_0(\varphi + 360^\circ) = \frac{1}{1-q^4} q^{-\frac{\varphi+360^\circ-4\beta}{\beta}} = \frac{1}{1-q^4} q^{-\frac{\varphi+2\alpha}{\beta}}$

For $-360^\circ - \beta \geq \varphi \geq -540^\circ - \beta$ c_1 is a subset of T_1 as

$c_1(\varphi) = \frac{1}{1-q^4} q^{-\frac{\varphi+\alpha}{\beta}} \leq e_0'(\varphi + 360^\circ) = \frac{1}{1-q^4} q^{-\frac{\varphi+2\alpha}{\beta}}$

For $-540^\circ - \beta \geq \varphi$ the argumentation repeats with contracted radii.

For $-180^\circ \geq \varphi \geq -180^\circ - \beta$ d_1 is a subset of S_1 as:

d_1 lies in the same half-plane as $P_{0,0}$ with respect to $P_{2,1}P_{1,1}$, so only the distance to b_0 needs to be tested. Let P be a point on d_1 . Applying the triangle inequality:

$$|PP_{0,0}| \geq |P_{1,1}P_{0,0}| - |P_{1,1}P| \geq q - d_1(-180^\circ) = q - \left(1 - \frac{q^2}{1-q^4}\right) q^{\frac{180^\circ - \alpha}{\beta}} = q - \left(1 - \frac{q^2}{1-q^4}\right) q^2$$

The polar angle of line $PP_{0,0}$ is between $-\beta$ and -2β .

In this domain $b_0(\varphi) \leq \left(1 - \frac{q^2}{1-q^4}\right) q$. Numerical analysis implies that

$$b_0(\varphi) \leq \left(1 - \frac{q^2}{1-q^4}\right) q \leq q - \left(1 - \frac{q^2}{1-q^4}\right) q^2 \leq |PP_{0,0}| \text{ holds for } q > 0.57, \text{ corresponding to } \alpha \leq 122^\circ.$$

For $-180^\circ - \beta \geq \varphi$ d_1 is a subset of $T_2 \cup S_4 \cup T_1$ as

$$d_1(\varphi) = \left(1 - \frac{q^2}{1-q^4}\right) q^{-\frac{\varphi+\alpha}{\beta}} \leq c_1(\varphi) = \frac{1}{1-q^4} q^{-\frac{\varphi+\alpha}{\beta}}$$

e_1 :

Let K and A be the intersection points of $P_{0,0}P_{2,1}$ and $P_{0,0}P_{1,1}$ with a_0 . The straight-line KA intersects e_1 in A' . The dilation with center K and factor q^4 maps the spiral arc \widehat{KA} (part of a_0) to \widehat{KA}' (part of e_1):

$$|KP_{0,0}| = a_0(-2\beta) = \frac{1}{1-q^4} q^2 \xrightarrow{\text{dilation}} |KP_{0,0}| q^4 = \frac{1}{1-q^4} q^6 = e_1(-2\beta) = |KP_{2,1}|$$

So, $P_{0,0}$, the center of a_0 is mapped to $P_{2,1}$, the center of e_1 .

$$a_0(\varphi) = \frac{1}{1-q^4} q^{-\frac{\varphi}{\beta}} \xrightarrow{\text{dilation}} \frac{1}{1-q^4} q^{-\frac{\varphi}{\beta}} q^4 = \frac{1}{1-q^4} q^{-\frac{\varphi-4\beta}{\beta}} = e_1(\varphi)$$

So, every point on the spiral arc \widehat{KA} is mapped to a point on \widehat{KA}' .

Let T_4 be the area, bounded by the arc \widehat{KA} and the line segment \overline{KA} . T_4 is convex and is a subset of S_1 . Since $K \in T_4$ the image arc \widehat{KA}' is a subset of T_4 .

$$\text{So, } \widehat{KA}' = \left\{ (d_{2,1}; \varphi) \in \mathbb{R}^2 \mid d_{2,1} = \frac{1}{1-q^4} q^{-\frac{\varphi+4\beta}{\beta}}, -2\beta \geq \varphi \geq -\beta \right\}$$

Let P be a point on the remaining spiral arc \widehat{HA}' . Then

- i. The polar angle φ of P : $-\beta \leq \varphi \leq 0^\circ$.
- ii. The distance to $P_{2,1}$: $|P_{2,1}P| \leq |P_{2,1}H| = e_1(0^\circ) = \frac{1}{1-q^4} q^4 < q^2 = |P_{2,1}P_{1,1}|$, which holds for $q \leq \frac{1}{2}\sqrt{2}$.
- iii. P lies with respect to the tangent to e_1 at A in the halfplane that contains $P_{2,1}$.

The three properties yield that the spiral arc \widehat{HA}' is subset of the parallelogram $P_{1,1}P_{2,1}A'A \subset S_1$ (for $\overline{P_{2,1}P_{1,1}} \subset S_1$ see Lemma 9).

Therefore, $\pi_0(A_{PF}) \subset A_{PF}$. \square

Proof b)

Since π_1 preserves the ratio, it suffices to examine the mapping of the encompassing spirals.

π_1 : A dilation by q and a rotation by $\beta - 180^\circ$, both with center $P_{0,0}$, and a translation by

$$\overrightarrow{P_{0,0}P_{0,1}}.$$

points: $P_{0,0} \rightarrow P_{0,1}$ $P_{0,1} \rightarrow P_{1,1}$ $P_{1,1} \rightarrow P_{2,3}$

spirals:

$$a_0(\varphi) = \frac{1}{1-q^4} q^{-\frac{\varphi}{\beta}}, \varphi \leq -\beta, \text{ center } P_{0,0}$$

$$\rightarrow a_2(\varphi) = \frac{1}{1-q^4} q^{-\frac{\varphi+\alpha}{\beta}}, \varphi \leq -180^\circ, \text{ center } P_{0,1}$$

$$b_0(\varphi) = \left(1 - \frac{q^2}{1-q^4}\right) q^{-\frac{\varphi}{\beta}}, \varphi \leq 0, \text{ center } P_{0,0}$$

$$\rightarrow b_2(\varphi) = \left(1 - \frac{q^2}{1-q^4}\right) q^{-\frac{\varphi+\alpha}{\beta}}, \varphi \leq -\alpha - \beta, \text{ center } P_{0,1}$$

$$c_0(\varphi) = \frac{1}{1-q^4} q^{-\frac{\varphi+\alpha}{\beta}}, \varphi \leq -180^\circ, \text{ center } P_{0,1}$$

$$\rightarrow c_2(\varphi) = \frac{1}{1-q^4} q^{-\frac{\varphi+2\alpha}{\beta}}, \varphi \leq \beta - 360^\circ, \text{ center } P_{1,1}$$

$$d_0(\varphi) = \left(1 - \frac{q^2}{1-q^4}\right) q^{-\frac{\varphi+\alpha}{\beta}}, \varphi \leq -\alpha - \beta, \text{ center } P_{0,1}$$

$$\rightarrow d_2(\varphi) = \left(1 - \frac{q^2}{1-q^4}\right) q^{-\frac{\varphi+2\alpha}{\beta}}, \varphi \leq -\alpha - 180^\circ, \text{ center } P_{1,1}$$

$$e_0(\varphi) = \frac{1}{1-q^4} q^{-\frac{\varphi-4\beta}{\beta}}, -\beta \leq \varphi \leq \beta, \text{ center } P_{1,1}$$

$$\rightarrow e_2(\varphi) = \frac{1}{1-q^4} q^{-\frac{\varphi+\alpha-4\beta}{\beta}}, -180^\circ \leq \varphi \leq -\alpha, \text{ center } P_{2,3}$$




Figure 15: $\pi_1(A_{PF})$

a_2 is mapped to c_0 , b_2 is mapped to d_0 (see construction 3).

For $\beta \geq \varphi \geq -\beta$, c_2 is mapped to e_0 (see construction 4) with $c_2(\varphi) = e_0(\varphi + 360^\circ)$.

For $-\beta \geq \varphi \geq -180^\circ - \beta$, c_2 is a subset of T_1

For $-180^\circ - \beta \geq \varphi$, c_2 is a subset of $T_2 \cup S_4 \cup T_1$ (analogous to c_1 in Part a)).

For $-180^\circ - \alpha \geq \varphi \geq -360^\circ + \beta$, d_2 is a subset of T_2 as

$$d_2(\varphi) \leq d_2(-180^\circ - \alpha) = \left(1 - \frac{q^2}{1-q^4}\right) q^2 \leq \frac{q^3}{1-q^4}.$$

Numerical analysis implies that this holds for $q > 0.57$ corresponding to $\alpha \leq 122^\circ$.

For $-360^\circ + \beta \geq \varphi$ d_2 is a subset of $T_2 \cup S_4 \cup T_1$, as $d_2(\varphi) \leq c_2(\varphi)$.

e_2 is tangent to b_0 at F , as $|P_{0,0}F| = b_0(0^\circ) = 1 - \frac{q^2}{1-q^4}$

$$|FP_{2,3}| = e_2(-180^\circ) = q \frac{1}{1-q^4} q^5, |P_{2,3}P_{0,1}| = q^2 \text{ and } |P_{0,0}F| + |FP_{2,3}| + |P_{2,3}P_{0,1}| = 1.$$

All points of e_2 lie within the quadrilateral $P_{1,1}P_{2,3}FN$ which is for $90^\circ \leq \alpha \leq 108^\circ$ a subset of S_2 .

Therefore, $\pi_1(A_{PF}) \subset A_{PF}$. \square

Both maps together yield

$$\pi_0(A_{PF}) \cup \pi_1(A_{PF}) \subset A_{PF}$$

Corollary 1 (for generalized A_{PF})




Figure 16: The mapping of generalized A_{PF} by π_0 and π_1

$$A_{PF}(\overline{P_{n,2k\cdot 2^m}P_{n,(2k+1)\cdot 2^m}}) \subset A_{PF}(\overline{P_{n,2k\cdot 2^m}P_{n,(2k+2)\cdot 2^m}})$$

$$A_{PF}(\overline{P_{n,(2k+2)\cdot 2^m}P_{n,(2k+1)\cdot 2^m}}) \subset A_{PF}(\overline{P_{n,2k\cdot 2^m}P_{n,(2k+2)\cdot 2^m}})$$

$$k, m, n \in \mathbb{N}_0, n \geq 1, 0 \leq m \leq n-1, 0 \leq k \leq 2^{n-m-1} - 1$$

6. The paperfolding polygons Q_n are subsets of A_{PF}

Or simply said: All dragons are inside the duck.

Proposition 2

$$Q_n \setminus (\overline{P_{n,0}P_{n,1}} \cup \overline{P_{n,2^n-1}P_{n,2^n}}) \subset A_{PF}, n \in \mathbb{N}_0$$

The proof consists of two parts. First, all vertices of a paperfolding polygon are elements of A_{PF} (Lemma 8) and second, all segments excluding the first $\overline{P_{n,0}P_{n,1}}$ and the last $\overline{P_{n,2^n-1}P_{n,2^n}}$ are subsets of A_{PF} (Lemma 10).

Lemma 8

For all paperfolding polygons $Q_n, n \in \mathbb{N}_0$ the vertices $P_{n,k}, k \in \{0,1,2, \dots, 2^n\}$ are elements of A_{PF} .

Proof (by mathematical induction)

Base step: $P_{0,0}, P_{0,1}$ and $P_{1,1}$ are elements of A_{PF} .

Induction step: Let $P_{n,k}$ be an element of A_{PF} .

Then $\pi_0(P_{n,k}) \in \pi_0(A_{PF}) \subset A_{PF}$ and $\pi_1(P_{n,k}) \in \pi_1(A_{PF}) \subset A_{PF}$.

All vertices of the paperfolding polygons $Q_n, n \geq 2$ are obtained through iterative applications of π_0 and π_1 to $P_{1,1}$. \square

Regarding the segments: Those connected to $P_{1,1}$ are special, since they are pictures under π_0 or π_1 of segments, that end in $P_{0,1}$ and these segments are not subsets of A_{PF} .

Lemma 9

For $90^\circ \leq \alpha \leq 108^\circ$ all segments $\overline{P_{n,2^{n-1}-1}P_{n,2^n-1}}$ and $\overline{P_{n,2^{n-1}}P_{n,2^{n-1}+1}}, n \geq 2$ are subsets of A_{PF} . These are all segments which start or end at $P_{1,1} = P_{n,2^{n-1}}$.




Figure 17: Some segments connected to $P_{1,1}$

Proof for $\overline{P_{n,2^{n-1}-1}P_{n,2^n-1}}, n \geq 2$

With respect to $P_{1,1}$ the points $P_{n,2^{n-1}-1}$ have polar angle of $\varphi_n = -180^\circ - (n-2)\beta$ and distance $r_n = q^n$.

$n = 2$ Proof for $\overline{P_{2,1}P_{2,2}}$

Let P be a point on $\overline{P_{2,1}P_{2,2}}$. By the law of sines in triangle $PP_{2,2}P_{0,0}$ (Figure 18)

$$|P_{0,0}P| = s(\varphi) = \frac{q \sin(\beta)}{\sin(180^\circ - |\varphi|)} = \frac{q \sin(\beta)}{-\sin(\varphi)}$$

Segment $\overline{P_{2,1}P_{2,2}}$ is a subset of A_{PF} ,

if $b_0(\varphi) < s(\varphi) < a_0(\varphi)$ for $-2\beta \leq \varphi \leq -\beta$

$$s(-\beta) = q < a_0(-\beta) = \frac{1}{1-q^4}q$$

$$s(-2\beta) = q^2 < a_0(-2\beta) = \frac{1}{1-q^4}q^2$$

Since $a_0(\varphi)$ forms a convex subset of A_{PF} (with respect to $P_{0,0}$), $s(\varphi) < a_0(\varphi)$ for $-2\beta \leq \varphi \leq -\beta$.

For $-2\beta \leq \varphi \leq -\beta$ there is $\max\{b_0(\varphi)\} = b_0(-\beta) = \left(1 - \frac{q^2}{1-q^4}\right)q$

and $\min\{s(\varphi)\} = s(-2\beta) = q^2$.

The condition $\max\{b_0(\varphi)\} < \min\{s(\varphi)\} \Leftrightarrow \left(1 - \frac{q^2}{1-q^4}\right)q < q^2 \Leftrightarrow 1 - q - q^2 - q^4 + q^5 < 0$

Numerical analysis implies that the inequality holds for $0.6 < q < 1$.

This is given for $90^\circ \leq \alpha \leq 108^\circ$.




Figure 18: The segment $\overline{P_{2,1}P_{2,2}}$ (note: $P_{2,2} = P_{1,1}$)

$n = 3$ Proof for $\overline{P_{3,3}P_{3,4}}$

$P_{0,0}, P_{3,3}$ and $P_{3,4}$ are collinear (see Figure 17) and $b_0(-\beta) = \left(1 - \frac{q^2}{1-q^4}\right)q < q - q^3 < q$

$$4 \leq n \leq 5 \quad -180^\circ - \beta \geq \varphi_n \geq -180^\circ - \beta - \alpha \text{ and } r_n \leq \frac{q^3}{1-q^4}.$$

Therefore $\overline{P_{n,2^{n-1}-1}P_{n,2^{n-1}}} \subset T_2$

$$6 \leq n \leq \frac{360^\circ}{\beta} + 1 \quad -360^\circ + \beta \geq \varphi_n \geq -540^\circ - \beta \text{ and } r_n \leq e_0(\varphi_n + 360^\circ) = \frac{q^4}{1-q^4}q^{-\frac{\varphi_n}{\beta}}.$$

Therefore $\overline{P_{n,2^{n-1}-1}P_{n,2^{n-1}}} \subset S_4 \cup T_1$

$\frac{360^\circ}{\beta} + 1 < n \quad$ The argument repeats with smaller radii r_n . \square

Proof for $\overline{P_{n,2^{n-1}}P_{n,2^{n-1}+1}}, n \geq 2$

With respect to $P_{1,1}$ the points $P_{n,2^{n-1}+1}$ have polar angle of $\varphi_n = -360^\circ - (n-4)\beta$ and distance $r_n = q^n$.

$n = 2$ Proof (for $\overline{P_{2,2}P_{2,3}}$)

$90^\circ \leq \alpha \leq 108^\circ$ yields $1 - q^2 \leq q$, so for all points P on segment $\overline{P_{2,2}P_{2,3}}$ the inequality $|P_{0,0}P| \leq q$ holds.

Let F be the foot of the perpendicular from $P_{0,0}$ to the line $P_{2,2}P_{2,3}$. For $90^\circ < \alpha < 120^\circ$ F is an interior point on $\overline{P_{2,2}P_{2,3}}$ and $|P_{0,0}F|$ is the minimum distance from $P_{0,0}$ to $\overline{P_{2,2}P_{2,3}}$.

$$|P_{0,0}F| = q \cdot \sin(\alpha - \beta) = \frac{\sqrt{4q^2 - 1}}{2q^2} (1 - q^2)$$




Figure 19: The segment $\overline{P_{2,2}P_{2,3}}$ (note: $P_{2,2} = P_{1,1}$)

$$\max\{b_0(\varphi) : -\beta \leq \varphi \leq 0^\circ\} = b_0(0^\circ) = \left(1 - \frac{q^2}{1-q^4}\right).$$

The condition $|P_{0,0}F| > \max\{b_0(\varphi)\}$ is equivalent to $\frac{\sqrt{4q^2-1}}{2q^2}(1-q^2) > \left(1 - \frac{q^2}{1-q^4}\right)$ and can be transformed to $\sqrt{4q^2-1}(1-q^2-q^4+q^6) - 2(q^2-q^4-q^6) > 0$.

Numerical analysis implies that this holds for $q > 0.599$ corresponding to $\alpha \leq 113^\circ$.

Therefore, $\overline{P_{2,2}P_{2,3}} \subset S_2$ for $\frac{1}{2}\sqrt{2} \geq q \geq \frac{\sqrt{5}-1}{2}$.

$$3 \leq n \leq 5 + \frac{180^\circ}{\beta} \quad -360^\circ + \beta \geq \varphi_n \geq -540^\circ - \beta \text{ and } r_n \leq e_0(\varphi_n + 360^\circ) = \frac{q^4}{1-q^4} q^{-\frac{\varphi_n}{\beta}}.$$

Therefore, $\overline{P_{n,2^{n-1}}P_{n,2^{n-1}+1}} \subset S_4 \cup T_1$

$$5 + \frac{180^\circ}{\beta} \leq n \leq 5 + \frac{180^\circ + \alpha}{\beta} \quad -540^\circ - \beta \geq \varphi_n \geq -540^\circ - \beta - \alpha \text{ and } r_n \leq \frac{q^3}{1-q^4}.$$

Therefore, $\overline{P_{n,2^{n-1}}P_{n,2^{n-1}+1}} \subset T_2$

$$5 + \frac{180^\circ + \alpha}{\beta} < n \quad \text{The argument repeats with smaller radii } r_n. \quad \square$$

Lemma 10

For all paperfolding polygons $Q_n, n \in \mathbb{N}, n \geq 2$ the segments $\overline{P_{n,k}P_{n,k+1}}, k \in \{1, 2, \dots, 2^n - 2\}$ are subsets of A_{PF} .

Remark

The segments $\overline{P_{n,0}P_{n,1}}$ (first segment) and $\overline{P_{n,2^{n-1}}P_{n,2^n}}$ (last segment) are excluded.

Proof (by mathematical induction)

Base step: $n = 2$ According to Lemma 9 $\overline{P_{2,1}P_{2,2}}$ and $\overline{P_{2,2}P_{2,3}}$ are subsets of A_{PF} .

Induction step: Let $\overline{P_{n,k}P_{n,k+1}}, k \in \{1, 2, \dots, 2^n - 2\}$ be subset of A_{PF} .

Then $\pi_0(\overline{P_{n,k}P_{n,k+1}})$ is subset of $\pi_0(A_{PF})$. $\pi_0(A_{PF}) \subset A_{PF}$ (Proposition 1 a)),

therefore $\pi_0(\overline{P_{n,k}P_{n,k+1}}) = \overline{P_{n+1,k}P_{n+1,k+1}}, k \in \{1, 2, \dots, 2^n - 2\}$ is subset of A_{PF} .

$\pi_1(\overline{P_{n,k}P_{n,k+1}})$ is subset of $\pi_1(A_{PF})$. $\pi_1(A_{PF}) \subset A_{PF}$ (Proposition 1 b)),

therefore $\pi_1(\overline{P_{n,k}P_{n,k+1}}) = \overline{P_{n+1,2^{n+1}-(k+1)}P_{n+1,2^{n+1}-k}}, k \in \{1, 2, \dots, 2^n - 2\}$ is subset of A_{PF} .

For $k = 2^n - 1$, the segments $\overline{P_{n+1,2^{n-1}}P_{n+1,2^n}}$ and $\overline{P_{n+1,2^n}P_{n+1,2^{n+1}}}$ are subsets of A_{PF} , as proven in Lemma 9.

Lemma 8 and Lemma 10 yield the proof of Proposition 2.

Now everything is ready for the main sentences.

7. Avoiding intersections of the paperfolding polygons Q_n

Proposition 3

For unfolding angles $98.195^\circ \leq \alpha \leq 108^\circ$ $\pi_0(A_{PF}) \cap \pi_1(A_{PF}) = \{P_{1,1}\}$.

Proof

The main condition is $c_1(\varphi) < d_2(\varphi)$

$$\begin{aligned} &\Leftrightarrow \frac{1}{1-q^4} q^{-\frac{\varphi+\alpha}{\beta}} < \left(1 - \frac{q^2}{1-q^4}\right) q^{-\frac{\varphi+2\alpha}{\beta}} \\ &\Leftrightarrow q^{-\frac{\varphi+\alpha}{\beta}} < (1-q^2-q^4) q^{-\frac{\varphi+2\alpha}{\beta}} \\ &\Leftrightarrow 0 < 1-q^2-q^4-q^{\frac{\alpha}{\beta}} \end{aligned}$$

The right-hand side is in $[0,1]$ monotonously decreasing with a zero between $q = 0.6615289$ and $q = 0.6615339$. Therefore, $c_1(\varphi) < d_2(\varphi)$ holds for $q \leq 0.6615289$, which corresponds to $\alpha \geq 98.195^\circ$.

To prove that $\pi_0(A_{PF}) \cap \pi_1(A_{PF}) = \{P_{1,1}\}$ for $98.195^\circ \leq \alpha \leq 108^\circ$ in detail, the plane is partitioned into three pairs of half-planes:

- by line $P_{0,0}P_{1,1}$ into H_0 containing $P_{2,1}$ and H_1 containing $P_{2,3}$
- by line $P_{2,3}P_{1,1}$ into H_2 containing $P_{0,1}$ and H_3 containing $P_{0,0}$
- by line $P_{0,0}P_{0,1}$ into H_4 containing $P_{1,1}$ and H_5 not containing $P_{1,1}$




Figure 20: $\alpha = 99^\circ$ $\pi_0(A_{PF})$ and $\pi_1(A_{PF})$ are clearly separated

The proof is organized in the following table:

\cap	$\pi_0(S_1)$	$\pi_0(S_2)$	$\pi_0(S_3)$	$\pi_0(S_4)$
$\pi_1(S_1)$	1.	\emptyset	2.	\emptyset
$\pi_1(S_2)$	3.	\emptyset	4.	\emptyset
$\pi_1(S_3)$	5.	6.	7.	8.
$\pi_1(S_4)$	9.	\emptyset	10.	\emptyset

$\pi_0(S_2)$ and $\pi_0(S_4)$ are subsets of H_0 . $\pi_1(S_1)$, $\pi_1(S_2)$ and $\pi_1(S_4)$ are subsets of H_1 , because the distance between the point $P_{0,1}$ and the line $P_{0,0}P_{1,1}$ is greater than that between the point $P_{0,1}$

and any point of these three sets. This yields the \emptyset in the table.

1. $\pi_0(S_1) \cap \pi_1(S_1)$

Since $\pi_1(S_1) \subset H_1$, only $\pi_0(S_1) \cap H_1$ needs to be considered in the calculation.

$$P \in \pi_0(S_1) \cap H_1 \Rightarrow |P_{0,0}P| \leq a_1(-180^\circ - \beta) = \frac{1}{1-q^4} q^{-\frac{-180^\circ - \beta}{\beta}}$$

$$\text{Applying the triangle inequality: } |P_{0,1}P| \geq |P_{0,0}P_{0,1}| - |P_{0,0}P| \geq 1 - \frac{1}{1-q^4} q^{-\frac{-180^\circ - \beta}{\beta}}.$$

$$Q \in \pi_1(S_1) \Rightarrow |P_{0,1}Q| \leq a_2(-180^\circ) = \frac{1}{1-q^4} q^{-\frac{-180^\circ + \alpha}{\beta}}$$

$$\frac{1}{1-q^4} q^{-\frac{-180^\circ + \alpha}{\beta}} < 1 - \frac{1}{1-q^4} q^{-\frac{-180^\circ - \beta}{\beta}} \text{ holds for } q \leq \frac{1}{2}\sqrt{2}.$$

For any $P \in \pi_0(S_1) \cap H_1$ and $Q \in \pi_1(S_1)$, the inequality $|P_{0,1}Q| < |P_{0,1}P|$ holds.

This implies $(\pi_0(S_1) \cap H_1) \cap \pi_1(S_1) = \emptyset$ and therefore $\pi_0(S_1) \cap \pi_1(S_1) = \emptyset$.

2. $\pi_0(S_3) \cap \pi_1(S_1)$

Since $\pi_1(S_1) \subset H_1$ only $\pi_0(S_3) \cap H_1$ needs to be considered in the calculation.

$\pi_1(S_1)$ is split by line $P_{0,0}P_{0,1}$ into $\pi_1(S_1) \cap H_5$ and $\pi_1(S_1) \cap H_4$.

$$P \in \pi_0(S_3) \cap H_1 \Rightarrow |P_{1,1}P| \leq c_1(-180^\circ - \beta) = \frac{1}{1-q^4} q^3 \text{ Lemma 6 a) implies}$$

that $\pi_0(S_3) \cap H_1 \subset H_4$. So $(\pi_0(S_3) \cap H_1) \cap (\pi_1(S_1) \cap H_5) = \emptyset$.

$$\text{Applying the triangle inequality: } |P_{0,1}P| \geq |P_{1,1}P_{0,1}| - |P_{1,1}P| \geq q - \frac{1}{1-q^4} q^3$$

$$Q \in \pi_1(S_1) \cap H_4 \Rightarrow |P_{0,1}Q| \leq a_2(-360^\circ) = \frac{1}{1-q^4} q^{-\frac{-360^\circ + \alpha}{\beta}}$$

$$\frac{1}{1-q^4} q^{-\frac{-360^\circ + \alpha}{\beta}} < q - \frac{1}{1-q^4} q^3 \text{ holds for } q \leq \frac{1}{2}\sqrt{2}.$$

For any $P \in \pi_0(S_3) \cap H_1$ and $Q \in \pi_1(S_1) \cap H_4$, the inequality $|P_{0,1}Q| < |P_{0,1}P|$ holds.

This implies $(\pi_0(S_3) \cap H_1) \cap (\pi_1(S_1) \cap H_4) = \emptyset$ and therefore $\pi_0(S_3) \cap \pi_1(S_1) = \emptyset$.

3. $\pi_0(S_1) \cap \pi_1(S_2)$

$\pi_0(S_1) \subset H_3$ and $\pi_1(S_2) \subset H_2$ and as H_2 and H_3 are disjoint halfplanes with respect to line $P_{2,3}P_{1,1}$ this implies $\pi_0(S_1) \cap \pi_1(S_2) = \emptyset$

4. $\pi_0(S_3) \cap \pi_1(S_2)$

Since $\pi_1(S_2) \subset H_2$, only $\pi_0(S_3) \cap H_2$ needs to be considered in the calculation.

$$P \in \pi_0(S_3) \cap H_2 \Rightarrow |P_{1,1}P| \leq c_1(-180^\circ - \alpha) = \frac{1}{1-q^4} q^{-\frac{-180^\circ}{\beta}}$$

$$\text{Applying the triangle inequality: } |P_{0,1}P| \geq |P_{1,1}P_{0,1}| - |P_{1,1}P| \geq q - \frac{1}{1-q^4} q^{\frac{\alpha+2\beta}{\beta}}$$

$$Q \in \pi_1(S_2) \Rightarrow |P_{0,1}Q| \leq q^2 \text{ and } q^2 < q - \frac{1}{1-q^4} q^{\frac{\alpha+2\beta}{\beta}} \text{ holds for } q < 0.67.$$

For any $P \in \pi_0(S_3) \cap H_2$ and $Q \in \pi_1(S_2)$, the inequality $|P_{0,1}Q| < |P_{0,1}P|$ holds.

This implies $(\pi_0(S_3) \cap H_2) \cap \pi_1(S_2) = \emptyset$ and therefore $\pi_0(S_3) \cap \pi_1(S_2) = \emptyset$.

5. $\pi_0(S_1) \cap \pi_1(S_3)$

The line $P_{0,0}P_{1,1}$ divides both sets. Thus, there are two cases.

Case 1: $(\pi_0(S_1) \cap H_0) \cap (\pi_1(S_3) \cap H_0)$

The line $P_{0,0}P_{2,1}$ partitions the plane into two half-planes. $\pi_1(S_3) \cap H_0$ is entirely in one half-plane, say H' . Hence only $\pi_0(S_1) \cap H_0 \cap H'$ needs to be considered in the calculation.

Let P be a point of $\pi_0(S_1) \cap H_0 \cap H'$ with a polar angel φ to the center $P_{0,0}$,

$$-\beta - k \cdot 360^\circ \geq \varphi \geq -2\beta - k \cdot 360^\circ, k = 1, 2, 3, \dots.$$

$$\text{Then } |P_{0,0}P| \leq a_1(-\beta - 360^\circ) = \frac{1}{1-q^4} q^{-\frac{-5\beta-2\alpha}{\beta}}.$$

Applying the triangle inequality: $|P_{1,1}P| \geq |P_{0,0}P_{1,1}| - |P_{0,0}P| \leq q - \frac{1}{1-q^4}q^{-\frac{-5\beta-2\alpha}{\beta}}$

$$Q \in \pi_1(S_3) \cap H_0 \Rightarrow |P_{1,1}Q| \leq c_2(-\beta - 360^\circ) = \frac{1}{1-q^4}q^{-\frac{-\beta-360^\circ+2\alpha}{\beta}}$$

$$\frac{1}{1-q^4}q^{-\frac{-\beta-360^\circ+2\alpha}{\beta}} < q - \frac{1}{1-q^4}q^{-\frac{-5\beta-2\alpha}{\beta}} \text{ holds for } q \leq \frac{1}{2}\sqrt{2}.$$

For any $P \in \pi_0(S_1) \cap H_0 \cap H'$ and $Q \in \pi_1(S_3) \cap H_0$, the inequality $|P_{1,1}Q| < |P_{1,1}P|$ holds. This implies $(\pi_0(S_1) \cap H_0 \cap H') \cap (\pi_1(S_3) \cap H_0) = \emptyset$ and therefore $(\pi_0(S_1) \cap H_0) \cap (\pi_1(S_3) \cap H_0) = \emptyset$.

Case 2: $(\pi_0(S_1) \cap H_1) \cap (\pi_1(S_3) \cap H_1)$

Since $\pi_0(S_1) \cap H_1 \subset H_3$ only $\pi_1(S_3) \cap H_1 \cap H_3$ needs to be considered in the calculation.

$$P \in \pi_0(S_1) \cap H_1 \Rightarrow |P_{0,0}P| \leq a_1(-180^\circ - \beta) = \frac{1}{1-q^4} q^{-\frac{-180^\circ - \beta}{\beta}}.$$

Applying the triangle inequality: $|P_{1,1}P| \geq |P_{0,0}P_{1,1}| - |P_{0,0}P| \geq q - \frac{1}{1-q^4} q^{-\frac{-180^\circ - \beta}{\beta}}$.

$$Q \in \pi_1(S_3) \cap H_1 \cap H_3 \Rightarrow |P_{1,1}Q| \leq c_2(-\beta - 540^\circ) = \frac{1}{1-q^4}q^{-\frac{-\beta - 540^\circ + 2\alpha}{\beta}}$$

$$\frac{1}{1-q^4}q^{-\frac{-\beta-540^\circ+2\alpha}{\beta}} < q - \frac{1}{1-q^4}q^{-\frac{-180^\circ-\beta}{\beta}} \text{ holds for } q \leq \frac{1}{2}\sqrt{2}.$$

For any $P \in \pi_0(S_1) \cap H_1$ and $Q \in \pi_1(S_3) \cap H_1 \cap H_3$, the inequality $|P_{1,1}Q| < |P_{1,1}P|$ holds. This implies $(\pi_0(S_1) \cap H_1) \cap (\pi_1(S_3) \cap H_1 \cap H_3) = \emptyset$ and therefore $(\pi_0(S_1) \cap H_1) \cap (\pi_1(S_3) \cap H_1) = \emptyset$.

$$6. \quad \pi_0(S_2) \cap \pi_1(S_3)$$

Since $\pi_0(S_2) \subset H_0$, only $\pi_1(S_3) \cap H_0$ needs to be considered in the calculation.

$$P \in \pi_0(S_2) \Rightarrow |P_{0,0}P| \leq q^2$$

Applying the triangle inequality: $|P_{1,1}P| \geq |P_{0,0}P_{1,1}| - |P_{0,0}P| \geq q - q^2$

$$Q \in \pi_1(S_3) \cap H_0 \Rightarrow |P_{1,1}Q| \leq c_2(-\beta - 360^\circ) = \frac{1}{\frac{1}{1-q^4}} q^{-\frac{-\beta - 360^\circ + 2\alpha}{\beta}}$$

$$\frac{1}{1-q^4}q^{-\frac{-\beta-360^\circ+2\alpha}{\beta}} < q - q^2 \text{ holds for } q < 0.69.$$

For any $P \in \pi_0(S_2)$ and $Q \in \pi_1(S_3) \cap H_0$, the inequality $|P_{1,1}Q| < |P_{1,1}P|$ holds.

This implies $\pi_0(S_2) \cap (\pi_1(S_3) \cap H_0) = \emptyset$ and therefore $\pi_0(S_2) \cap \pi_1(S_3) = \emptyset$.

$$7. \quad \pi_0(S_3) \cap \pi_1(S_3)$$

For polar angles $-180^\circ \geq \varphi > -180^\circ - \alpha$ only $\pi_0(S_3)$ is defined.

For $-180^\circ - \alpha \geq \varphi$ a ray from $P_{1,1}$ at polar angle φ intersects $\pi_0(S_3)$ and $\pi_1(S_3)$ infinitely many times. The situation is illustrated in Figure 21.




Figure 21: The infinitely intersection of $\pi_0(S_0)$ and $\pi_1(S_0)$ (case 7)

Points between $c_2(\varphi - k \cdot 360^\circ)$ and $d_2(\varphi - k \cdot 360^\circ)$, $k \in \mathbb{N}_0$ belong to $\pi_1(S_3)$, points between $c_1(\varphi - k \cdot 360^\circ)$ and $d_1(\varphi - k \cdot 360^\circ)$, $k \in \mathbb{N}_0$ belong to $\pi_2(S_3)$.

These intervals are pairwise disjoint if $c_2(\varphi) > d_2(\varphi) > c_1(\varphi) > d_1(\varphi) > c_2(\varphi - 360^\circ)$ for all $-180^\circ - \alpha \geq \varphi$.

$$\frac{c_2(\varphi)}{d_2(\varphi)} = \frac{1}{1-q^4-q^2} > 1 \text{ holds for } q \leq \frac{1}{2}\sqrt{2}$$

$d_2(\varphi) > c_1(\varphi)$ was calculated in the beginning,

yielding the condition $q \leq 0.6615289$, which corresponds to $\alpha \geq 98.195^\circ$.

$$\frac{c_1(\varphi)}{d_1(\varphi)} = \frac{1}{1-q^4-q^2} > 1 \text{ holds for } q \leq \frac{1}{2}\sqrt{2}$$

$$d_1(\varphi) = \left(1 - \frac{q^2}{1-q^4}\right) q^{-\frac{\varphi+\alpha}{\beta}}, c_2(\varphi - 360^\circ) = \frac{1}{1-q^4} q^{-\frac{\varphi-4\beta}{\beta}}$$

$d_1(\varphi) > c_2(\varphi - 360^\circ)$ simplifies to $1 - q^2 - q^4 - q^{\frac{4+\alpha}{\beta}} > 0$, which holds for $q \leq \frac{1}{2}\sqrt{2}$.

Therefore, $\pi_0(S_3) \cap \pi_1(S_3) = \{P_{1,1}\}$.

8. $\pi_0(S_4) \cap \pi_1(S_3)$

Since $\pi_0(S_4) \subset H_0$ only $\pi_1(S_3) \cap H_0$ needs to be considered in the calculation.

$$P \in \pi_0(S_4) \Rightarrow |P_{2,1}P| \leq e_1(0^\circ) = \frac{1}{1-q^4} q^{-\frac{-4\beta}{\beta}}.$$

Applying the triangle inequality: $|P_{1,1}P| \geq |P_{2,1}P_{1,1}| - |P_{2,1}P| \geq q^2 - \frac{1}{1-q^4} q^{-\frac{-4\beta}{\beta}}$.

$$Q \in \pi_1(S_3) \cap H_0 \Rightarrow |P_{1,1}Q| \leq c_2(-\beta - 360^\circ) = \frac{1}{1-q^4} q^{-\frac{-\beta-360^\circ+2\alpha}{\beta}}$$

$$\frac{1}{1-q^4} q^{-\frac{-\beta-360^\circ+2\alpha}{\beta}} < q^2 - \frac{1}{1-q^4} q^{-\frac{-4\beta}{\beta}} \text{ holds for } q < 0.68.$$

For any $P \in \pi_0(S_4)$ and $Q \in \pi_1(S_3) \cap H_0$, the inequality $|P_{1,1}Q| < |P_{1,1}P|$ holds.

This implies $\pi_0(S_4) \cap (\pi_1(S_3) \cap H_0) = \emptyset$ and therefore $\pi_0(S_4) \cap \pi_1(S_3) = \emptyset$.

9. $\pi_0(S_1) \cap \pi_1(S_4)$

Since $\pi_1(S_4) \subset H_1$ only $\pi_0(S_1) \cap H_1$ needs to be considered in the calculation.

$$P \in \pi_0(S_1) \cap H_1 \Rightarrow |P_{0,0}P| \leq a_1(-180^\circ - \beta) = \frac{1}{1-q^4} q^{-\frac{-180^\circ - \beta}{\beta}}.$$

Applying the triangle inequality: $|P_{2,3}P| \geq |P_{0,0}P_{2,3}| - |P_{0,0}P| \geq 1 - q^2 - \frac{1}{1-q^4} q^{-\frac{-180^\circ - \beta}{\beta}}$.

$$Q \in \pi_1(S_4) \Rightarrow |P_{2,3}Q| \leq e_2(-\alpha) = \frac{1}{1-q^4} q^{-\frac{-4\beta}{\beta}}$$

$$\frac{1}{1-q^4} q^{-\frac{-4\beta}{\beta}} < 1 - q^2 - \frac{1}{1-q^4} q^{-\frac{-180^\circ - \beta}{\beta}} \text{ holds for } q < 0.69.$$

For any $P \in \pi_0(S_1) \cap H_1$ and $Q \in \pi_1(S_4)$, the inequality $|P_{2,3}Q| < |P_{2,3}P|$ holds.

This implies $(\pi_0(S_1) \cap H_1) \cap \pi_1(S_4) = \emptyset$ and therefore $\pi_0(S_1) \cap \pi_1(S_4) = \emptyset$.

10. $\pi_0(S_3) \cap \pi_1(S_4)$

$c_1(-180^\circ - \alpha)$ defines the point R_1 on the spiral c_1 and $e_2(-\alpha)$ defines the point R_2 on the spiral e_2 .

Let t_1 be the tangent line to c_1 at R_1 .

Let t_2 be the tangent line to e_2 at R_2 .

Since $P_{1,1}, R_1, R_2$ and $P_{2,3}$ are collinear, t_1 and t_2 are parallel.

$c_1(-180^\circ - \alpha) < d_2(-180^\circ - \alpha) = |P_{2,3}P_{1,1}| - e_2(-\alpha)$ implies that t_1 and t_2 are distinct lines.

c_1 and t_2 lie in different half-planes determined by t_1 .

e_2 and t_1 lie in different half-planes determined by t_2 .

Therefore, $\pi_0(S_3) \cap \pi_1(S_4) = \emptyset$.

□




Figure 22: $\pi_0(S_3)$ and $\pi_1(S_4)$ (case 10)

Corollary 2 (for generalized A_{PF})

For unfolding angles $98.195^\circ \leq \alpha \leq 108^\circ$

$$A_{PF}(\overline{P_{n,2k \cdot 2^m} P_{n,(2k+1) \cdot 2^m}}) \cap A_{PF}(\overline{P_{n,(2k+2) \cdot 2^m} P_{n,(2k+1) \cdot 2^m}}) = \{P_{n,(2k+1) \cdot 2^m}\}.$$

Theorem 1

For unfolding angles $98.195^\circ \leq \alpha \leq 108^\circ$ the paperfolding polygons Q_n avoid intersections.

Proof

The paperfolding polygon Q_n is constructed by $Q_n = \pi_0(Q_{n-1}) \cup \pi_1(Q_{n-1})$.

$\pi_0(A_{PF}) \cap \pi_1(A_{PF}) = \{P_{1,1}\}$ implies that all segments of $\pi_0(Q_{n-1})$, which are subsets of $\pi_0(A_{PF})$ do not intersect any segment of $\pi_1(Q_{n-1})$, which is a subset of $\pi_1(A_{PF})$.

This argument excludes the segment $\overline{P_{n,0}P_{n,1}}$ at the beginning and $\overline{P_{n,2^{n-1}-1}P_{n,2^{n-1}}}$ at the end of $\pi_0(Q_{n-1})$, respectively $\overline{P_{n,2^{n-1}}P_{n,2^{n-1}+1}}$ at the beginning and $\overline{P_{n,2^n-1}P_{n,2^n}}$ at the end of $\pi_1(Q_{n-1})$.

Discussing these four segments:

1. $\overline{P_{n,0}P_{n,1}}$ does not intersect with any segment of $\pi_1(Q_{n-1})$, as for $n \geq 3$ the circle with center $P_{0,0}$ and radius q^n does not intersect $\pi_1(A_{PF})$.

2. An analogous argumentation holds for the segment $\overline{P_{n,2^{n-1}}P_{n,2^n}}$ at the end of $\pi_1(Q_{n-1})$.




Figure 23: The situation at $P_{n,2^{n-1}}$

3. Assume toward a contradiction that segment $\overline{P_{n,2^{n-1}-1}P_{n,2^{n-1}}}$ intersects in point S a segment $\overline{P_{n,k}P_{n,k+1}}$, which is a segment of $\pi_1(Q_{n-1})$. Therefore, $2^{n-1} \leq k \leq 2^n - 1$.

Case 1: $S \notin \pi_1(A_{PF})$

Then S can only be the intersection point with the first segment $\overline{P_{n,2^{n-1}}P_{n,2^{n-1}+1}}$ of $\pi_1(Q_{n-1})$ or with the last segment $\overline{P_{n,2^{n-1}}P_{n,2^n}}$.

First segment $\overline{P_{n,2^{n-1}}P_{n,2^{n-1}+1}}$: If $S = \overline{P_{n,2^{n-1}}P_{n,2^{n-1}+1}} \cap \overline{P_{n,2^{n-1}-1}P_{n,2^{n-1}}}$ then $S = P_{n,2^{n-1}}$. This is not an intersection in the discussed sense.

Last segment $\overline{P_{n,2^{n-1}}P_{n,2^n}}$: Let Q be a point on $\overline{P_{n,2^{n-1}}P_{n,2^n}}$. Then $|QP_{n,2^n}| \leq q^n$.

$S \in \overline{P_{n,2^{n-1}-1}P_{n,2^{n-1}}}$ yields $|SP_{n,2^{n-1}}| \leq q^n$.

Applying the triangle inequality: $|SP_{n,2^n}| \geq |P_{n,2^{n-1}}P_{n,2^n}| - |SP_{n,2^{n-1}}| = q - q^n$

$q - q^n > q^n$ holds for $q \leq \frac{1}{2}\sqrt{2}$ and $n \geq 4$.

For $S \in \overline{P_{n,2^{n-1}-1}P_{n,2^{n-1}}}$ and any $Q \in \overline{P_{n,2^{n-1}}P_{n,2^n}}$, the inequality

$|QP_{n,2^n}| < |SP_{n,2^n}|$ holds. This implies $S \notin \overline{P_{n,2^{n-1}-1}P_{n,2^n}}$ and therefore S is not the intersection point of $\overline{P_{n,2^{n-1}}P_{n,2^n}}$ and $\overline{P_{n,2^{n-1}-1}P_{n,2^{n-1}}}$.

Case 2: $S \in \pi_1(A_{PF})$

$S \in \overline{P_{n,2^{n-1}-1}P_{n,2^{n-1}}}$ and $\overline{P_{n,2^{n-1}-1}P_{n,2^{n-1}}}$ has a polar angle of $-180^\circ - (n-2)\beta$, with respect to the point $P_{n,2^{n-1}}$. Therefore

$$|SP_{n,2^{n-1}}| \leq c_2(-540^\circ - (n-2)\beta) = \frac{1}{1-q^4} q^{-\frac{-540^\circ - (n-2)\beta + 2\alpha}{\beta}} = \frac{1}{1-q^4} q^n q^4 q^{\frac{\alpha}{\beta}}$$

Hence $\frac{1}{1-q^4} q^n q^4 q^{\frac{\alpha}{\beta}} < q^n$ the point S is an element of $A_{PF}(\overline{P_{n,2^{n-1}}P_{n,2^{n-1}+1}})$ which

implies that S belongs to all $A_{PF}(\overline{P_{n,2^{n-1}}P_{n,2^{n-1}+2^m}})$, $0 \leq m \leq n-2$ (Corollary 1 of

Proposition 1).

Now the main idea is: Because $S \in A_{PF}(\overline{P_{n,2^{n-1}}P_{n,2^{n-1}+2^m}})$, $0 \leq m \leq n-2$ and $A_{PF}(\overline{P_{n,2^{n-1}}P_{n,2^{n-1}+2^m}}) \cap A_{PF}(\overline{P_{n,2^{n-1}+2^{m+1}}P_{n,2^{n-1}+2^m}}) = \{P_{n,2^{n-1}+2^m}\}$ (Corollary 2) and $S \neq P_{n,2^{n-1}+2^m}$ (not an intersection in the discussed sense),

it follows that S cannot be an element of $A_{PF}(\overline{P_{n,2^{n-1}+2^{m+1}}P_{n,2^{n-1}+2^m}})$.

Therefore, S cannot be the intersection point between the segment $\overline{P_{n,2^{n-1}-1}P_{n,2^{n-1}}}$ and any segment $\overline{P_{n,2^{n-1}+k}P_{n,2^{n-1}+k+1}}$, $2^m+1 \leq k \leq 2^{m+1}-2$, i.e. any segment that is a subset of $A_{PF}(\overline{P_{n,2^{n-1}+2^{m+1}}P_{n,2^{n-1}+2^m}})$. The special segments

$\overline{P_{n,2^{n-1}+2^m}P_{n,2^{n-1}+2^{m+1}}}$ (first) and $\overline{P_{n,2^{n-1}+2^{m+1}-1}P_{n,2^{n-1}+2^{m+1}}}$ (last) have to be treated separately.

The proof proceeds by case analysis on the parameter m :

$m = 0$:

Since $S \in A_{PF}(\overline{P_{n,2^{n-1}}P_{n,2^{n-1}+1}})$ it follows $S \notin A_{PF}(\overline{P_{n,2^{n-1}+1}P_{n,2^{n-1}+2}})$

The special case of $A_{PF}(\overline{P_{n,2^{n-1}+1}P_{n,2^{n-1}+2}})$ concerns the segment $\overline{P_{n,2^{n-1}+1}P_{n,2^{n-1}+2}}$.

All points of this line segment have polar angles φ with respect to $P_{n,2^{n-1}}$ satisfying $-180^\circ - (n-2)\beta - \alpha + \beta \geq \varphi \geq -180^\circ - (n-2)\beta - \alpha$. S lies on segment

$\overline{P_{n,2^{n-1}-1}P_{n,2^{n-1}}}$ which has a polar angle of $\varphi = -180^\circ - (n-2)\beta$ with respect to $P_{n,2^{n-1}}$. Since these angle ranges are disjoint, S cannot be the intersection point between both segments.

$m = 1$:

Since $S \in A_{PF}(\overline{P_{n,2^{n-1}}P_{n,2^{n-1}+2}})$ it follows $S \notin A_{PF}(\overline{P_{n,2^{n-1}+2}P_{n,2^{n-1}+4}})$

Only the special segments have to be considered.

First segment $\overline{P_{n,2^{n-1}+2}P_{n,2^{n-1}+3}}$: All points of this line segment have polar angles φ with respect to $P_{n,2^{n-1}}$ satisfying

$-180^\circ - (n-2)\beta - \alpha + 2\beta \geq \varphi \geq -180^\circ - (n-2)\beta - \alpha + \beta$. S lies on segment $\overline{P_{n,2^{n-1}-1}P_{n,2^{n-1}}}$ which has a polar angle of $\varphi = -180^\circ - (n-2)\beta$ with respect to $P_{n,2^{n-1}}$. Since these angle ranges are disjoint, S cannot be the intersection point between both segments.

Last segment $\overline{P_{n,2^{n-1}+3}P_{n,2^{n-1}+4}}$: All points of this line segment have a polar angle $\varphi = -180^\circ - (n-2)\beta - \alpha + 2\beta$ with respect to $P_{n,2^{n-1}}$. S lies on segment

$\overline{P_{n,2^{n-1}-1}P_{n,2^{n-1}}}$ which has a polar angle of $\varphi = -180^\circ - (n-2)\beta$ with respect to $P_{n,2^{n-1}}$. Since these angles are distinct, S cannot be the intersection point between both segments.

General case : $2 \leq m \leq n-2$

Since $S \in A_{PF}(\overline{P_{n,2^{n-1}}P_{n,2^{n-1}+2^m}})$ it follows that $S \notin A_{PF}(\overline{P_{n,2^{n-1}+2^m}P_{n,2^{n-1}+2^{m+1}}})$.

Thus, S is not the intersection point between $\overline{P_{n,2^{n-1}-1}P_{n,2^{n-1}}}$ and any segment $\overline{P_{n,k}P_{n,k+1}}$, $2^{n-1}+2^m+1 \leq k \leq 2^{n-1}+2^{m+1}-2$.

First segment $\overline{P_{n,2^{n-1}+2^m}P_{n,2^{n-1}+2^{m+1}}}$:

Let Q be a point on $\overline{P_{n,2^{n-1}+2^m}P_{n,2^{n-1}+2^{m+1}}}$. Then $|QP_{n,2^{n-1}+2^m}| \leq q^n$.

Applying the triangle inequality to S yields

$$|SP_{n,2^{n-1}+2^m}| \geq |P_{n,2^{n-1}}P_{n,2^{n-1}+2^m}| - |SP_{n,2^{n-1}}| = q^{n-m} - \frac{1}{1-q^4}q^nq^4q^{\frac{\alpha}{\beta}}$$

$$q^{n-m} - \frac{1}{1-q^4}q^nq^4q^{\frac{\alpha}{\beta}} > q^n \text{ holds for } q \leq \frac{1}{2}\sqrt{2} \text{ and } m \geq 2.$$

For $S \in \overline{P_{n,2^{n-1}-1}P_{n,2^n}}$ and any $Q \in \overline{P_{n,2^{n-1}+2^m}P_{n,2^{n-1}+2^{m+1}}}$, the inequality

$|QP_{n,2^{n-1}+2^m}| < |SP_{n,2^{n-1}+2^m}|$ holds. This implies $S \notin \overline{P_{n,2^{n-1}+2^m}P_{n,2^{n-1}+2^{m+1}}}$ and therefore S is not the intersection point of $\overline{P_{n,2^{n-1}-1}P_{n,2^n}}$ and $\overline{P_{n,2^{n-1}+2^m}P_{n,2^{n-1}+2^{m+1}}}$.

Last segment $\overline{P_{n,2^{n-1}+2^{m+1}-1}P_{n,2^{n-1}+2^{m+1}}}$:

Let Q be a point on $\overline{P_{n,2^{n-1}+2^{m+1}-1}P_{n,2^{n-1}+2^{m+1}}}$. Then $|QP_{n,2^{n-1}+2^{m+1}}| \leq q^n$.

For S the triangle inequality yields

$$|SP_{n,2^{n-1}+2^{m+1}}| \geq |P_{n,2^{n-1}}P_{n,2^{n-1}+2^{m+1}}| - |SP_{n,2^{n-1}}| = q^{n-(m+1)} - \frac{1}{1-q^4}q^nq^4q^{\frac{\alpha}{\beta}}$$

$$q^{n-(m+1)} - \frac{1}{1-q^4}q^nq^4q^{\frac{\alpha}{\beta}} > q^n \text{ holds for } q \leq \frac{1}{2}\sqrt{2} \text{ and } m \geq 2.$$

For $S \in \overline{P_{n,2^{n-1}-1}P_{n,2^n}}$ and any $Q \in \overline{P_{n,2^{n-1}+2^{m+1}-1}P_{n,2^{n-1}+2^{m+1}}}$, the inequality

$|QP_{n,2^{n-1}+2^{m+1}}| < |SP_{n,2^{n-1}+2^{m+1}}|$ holds. This implies $S \notin \overline{P_{n,2^{n-1}+2^{m+1}-1}P_{n,2^{n-1}+2^{m+1}}}$ and therefore S is not the intersection point of $\overline{P_{n,2^{n-1}-1}P_{n,2^n}}$ and

$\overline{P_{n,2^{n-1}+2^{m+1}-1}P_{n,2^{n-1}+2^{m+1}}}$.

Therefore, segment $\overline{P_{n,2^{n-1}}P_{n,2^{n-1}-1}}$ does not intersect any segment of $\pi_1(Q_{n-1})$.

4. An analogous argumentation like 3. holds for $\overline{P_{n,2^{n-1}}P_{n,2^{n-1}+1}}$ and its intersection with segments of $\pi_0(Q_{n-1})$.

Applying the duck to the gap between $P_{4,7}$ and $P_{4,11}$

Each segment of a paperfolding polygon develops by the iterative process into a smaller copy of a paperfolding polygon. Consequently, the local duck-shaped hull of any segment serves as a boundary for all following segments between its start and endpoint.

This principle is applied to the loop formed by the points $P_{4,7}, P_{4,8}, \dots, P_{4,11}$ and the gap between $P_{4,7}$ and $P_{4,11}$. If the iterative process of generating the paperfolding polygons avoids intersection in this region, the assumption is, that intersections are avoided for all paperfolding polygons².




Figure 24: $A_{PF}(\overrightarrow{P_{4,6}P_{4,7}})$ and $A_{PF}(\overrightarrow{P_{4,10}P_{4,11}})$

Lemma 11

If the unfolding angle $\alpha \geq 96.241^\circ$ then $A_{PF}(\overrightarrow{P_{4,6}P_{4,7}})$ and $A_{PF}(\overrightarrow{P_{4,10}P_{4,11}})$ will not intersect.

Proof

The length of all segments in Q_4 is q^4 . For points S_1 and S_2 see Figure 24.

$$\begin{aligned} |P_{4,7}P_{4,11}| &= 2(|P_{4,8}S_1| - |P_{4,8}S_2|) \\ &= 2q^4(\cos\beta - \cos(\alpha - \beta)) = 2q^4(\cos\beta - \cos(180^\circ - 3\beta)) = 2q^4(\cos\beta + \cos 3\beta) \end{aligned}$$

² D. Knuth writes ¹: „The most troublesome self-crossing points appear to lie near $7 \cdot 2^n$ and $11 \cdot 2^n$ “




Figure 25: Enlargement of Fig. 24

For points S_3 and S_4 see Figure 25.

An intersection of both ducks is avoided, if $|P_{4,7}S_3| + |P_{4,11}S_4| < |P_{4,7}P_{4,11}|$

Segment $\overline{P_{4,6}P_{4,7}}$ is, compared to $\overline{P_{0,0}P_{0,1}}$, dilated by q^4 and not rotated.

So, the outer spiral at $P_{4,7}$ is given by $c_3(\varphi) = q^4 \frac{1}{1-q^4} q^{-\frac{\varphi+\alpha}{\beta}}$, $\varphi \leq -180^\circ$, center $P_{4,7}$

Therefore, $|P_{4,7}S_3| = c_3(-360^\circ + \beta) = c_3(-2\alpha - 4\beta + \beta) = q^4 \frac{1}{1-q^4} q^{-\frac{-\alpha-3\beta}{\beta}}$.

Segment $\overline{P_{4,10}P_{4,11}}$ is, compared to $\overline{P_{0,0}P_{0,1}}$, dilated by q^4 and rotated by 4β .

So, the outer spiral around $P_{4,11}$ is given by

$$c_4(\varphi) = q^4 \frac{1}{1-q^4} q^{-\frac{\varphi+\alpha-4\beta}{\beta}}, \varphi \leq -180^\circ + 4\beta, \text{ center } P_{4,11}$$

Therefore, $|P_{4,11}S_4| = c_4(-180^\circ + \beta) = c_4(-\alpha - 2\beta + \beta) = q^4 \frac{1}{1-q^4} q^5$

This yields

$$\begin{aligned} & |P_{4,7}S_3| + |P_{4,11}S_4| < |P_{4,7}P_{4,11}| \\ \Leftrightarrow & \frac{1}{1-q^4} q^{-\frac{-\alpha-3\beta}{\beta}} + \frac{1}{1-q^4} q^5 < 2(\cos\beta + \cos 3\beta) \\ \Leftrightarrow & \frac{q^3}{1-q^4} \left(q^{\frac{\alpha}{\beta}} + q^2 \right) < 2(\cos\beta + \cos 3\beta) \end{aligned}$$

The inequality is solved numerically (3 digits):

alpha(deg)	beta(deg)	q	left side	right side	<
96.235	41.883	0.6715777	0.3238188	0.3234373	FALSE
96.236	41.882	0.6715724	0.3238001	0.3234915	FALSE
96.237	41.882	0.6715672	0.3237814	0.3235457	FALSE
96.238	41.881	0.6715619	0.3237627	0.3235999	FALSE
96.239	41.881	0.6715567	0.3237439	0.3236541	FALSE
96.240	41.880	0.6715514	0.3237252	0.3237083	FALSE
96.241	41.880	0.6715462	0.3237065	0.3237625	TRUE
96.242	41.879	0.6715409	0.3236878	0.3238167	TRUE
96.243	41.879	0.6715357	0.3236691	0.3238709	TRUE
96.244	41.878	0.6715304	0.3236504	0.3239251	TRUE
96.245	41.878	0.6715252	0.3236317	0.3239793	TRUE

Conjecture

If intersections of $A_{PF}(\overrightarrow{P_{4,6}P_{4,7}})$ and $A_{PF}(\overrightarrow{P_{4,10}P_{4,11}})$ are avoided in the gap between $P_{4,7}$ and $P_{4,11}$ then intersections are avoided for all paperfolding polygons. By Lemma 11, this is satisfied when the unfolding angle $\alpha \geq 96,241^\circ$.

Summary

For the unfolding angle $\alpha \leq 90^\circ$ of the paperfolding polygons we know:




Figure 26: Diagram of the result of this article

$90^\circ < \alpha \leq 95.126^\circ$	Intersections (from 10 th iteration) ³
$95.126 < \alpha < 96.241$	Interval of uncertainty
$96.241 \leq \alpha < 98.195$	Conjecture, that there are no intersections
$98.195 \leq \alpha \leq 108^\circ$	No intersections, proved in theorem 1

In addition

$99.3438 < \alpha \leq 180^\circ$	No intersections ⁷
-----------------------------------	-------------------------------

References

1. D. E. Knuth, Selected Papers on Fun and Games, pp. 571–614. CSLI Lecture Notes 192, CSLI Publications, Stanford, California (2011)
2. M. Mendès France, The inhomogeneous Ising chain and paperfolding, in Number Theory and Physics (Les Houches, 1989), Springer Proc. Phys. 47, Springer (1990), pp. 195–202.
3. R. Albers, Papierfalten, Doctoral thesis, Bremen University (2006). Available at <https://nbn-resolving.de/urn:nbn:de:gbv:46-diss000105747>
4. S. Tabachnikov, Dragon curves revisited, Math. Intelligencer 36 (2014), 13–17
5. J.-P. Allouche, M. Mendès France, G. Skordev, Non-intersectivity of paperfolding dragon curves and curves generated by automatic sequences, Integers 18A (2018)
6. Y. Kamiya, On dragon curves which have two corners just meeting, Theoret. Comput. Sci., 938 (2022), 65–80
7. S. Akiyama, Y. Kamiya, F. Wen, Non-intersective dragon curves (2024). Available at <https://arxiv.org/html/2401.07002v1>
8. F. Wen, Convex Hulls of Dragon Curves (2022). Available at <https://arxiv.org/abs/2208.12434v1>