

**MOHAMED BOUDIAF UNIVERSITY OF M'SILA**  
**FACULTY OF MATHEMATICS AND COMPUTER SCIENCES**

**DEPARTMENT OF MATHEMATICS**

# **Study of Composition Operators in Certain Functional Spaces**

**A Thesis Submitted in Partial Fulfillment  
of the Requirements for the Degree of  
MAGISTER  
in Functional Analysis**

**By**  
**Mahdi Tahar BRAHIMI**

**Supervised by**  
**Professor Madani MOUSSAI**

**Academic Year: 2007/2008**

## **Abstract**

In this thesis we study three problems , the first is the superposition of the operators and their proprieties , such as boundedness, continuity, regularity and the inequalities of the norms of the composition of functions in some functional spaces , the second is to generalize some results of the composition of more than two functions , and the third is to give a generalization of Peetre's theorem.

# Table of Contents

<b>Notations</b>	ii
<b>Introduction</b>	v
<b>1 Some preliminary results</b>	<b>1</b>
1.1 Littlewood-Paley series	1
1.1.1 Partition of unity	1
1.1.2 Decomposition of a function $f \in \mathcal{S}'(\mathbb{R}^n)$	5
1.1.3 Decomposition of $\Delta_k(f \cdot g)$	6
1.2 Finite difference operators $\Delta_h^m$	6
1.3 Norms in Besov spaces	8
1.3.1 By Littlewood-Paley theory	9
1.3.2 By finite differences	11
1.4 Functions of bounded p-variation	15
1.4.1 General notions	15
1.4.2 Functions of bounded p-variation as distributions	17
<b>2 Composition of Operators in <math>BV_p^1(I)</math></b>	<b>20</b>
2.1 Reminder	20
2.2 Properties of the spaces $BV_p^1(I)$	21
2.3 Solution of the S.O.P $BV_p^1(I)$	22
2.4 Statement of results	29
<b>3 Composition of Operators in Homogeneous Besov Spaces <math>\dot{B}_p^{s,q}(\mathbb{R}^n)</math></b>	<b>34</b>
3.1 Composition Problem in Homogeneous Besov Spaces $\dot{B}_p^{s,q}(\mathbb{R}^n)$	36
3.1.1 Properties of Homogeneous Besov Spaces	36
3.1.2 Realizations of Homogeneous Besov Spaces	39
3.1.3 Statement of Results	41
3.2 Peetre's Theorem	43
<b>4 Composition of Operators in <math>BV_p^\alpha(I)</math></b>	<b>48</b>
4.1 Basic Notions	48
4.1.1 The Spaces $\mathcal{V}_p^\alpha(I)$	48
4.1.2 The Spaces $BV_p^\alpha(I)$	51
4.2 The Spaces $W^{1,p}(\Omega)$ , $\Omega$ an open subset of $\mathbb{R}^n$	53
4.2.1 Properties of the Spaces $W^{1,p}(\Omega)$	53
4.2.2 Composition $W^{1,p}(\Omega)$	56
4.3 Peetre's theorem $BV_p^\alpha(I)$	58
4.4 Examples	61

# Notations

- For all  $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n) \in \mathbb{N}^n$ ,  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ , we have:

$$D^\alpha f = \frac{\partial^{|\alpha|} f}{\partial x_1^{\alpha_1} \cdots \partial x_n^{\alpha_n}}, \quad \text{where } |\alpha| = \alpha_1 + \alpha_2 + \dots + \alpha_n,$$

- $C = C(\mathbb{R}^n)$ : Denotes the space of continuous functions.

$$C(\mathbb{R}^n) = \{u : \mathbb{R}^n \rightarrow \mathbb{R}; \forall x_0 \in \mathbb{R}^n, \forall \varepsilon > 0, \exists \delta > 0, \forall x \in \mathbb{R}^n, \|x - x_0\| < \delta \Rightarrow |u(x) - u(x_0)| < \varepsilon\}$$

- $C^r(\mathbb{R}^n) = \{f \in C(\mathbb{R}^n); D^\alpha f \in C(\mathbb{R}^n), \text{ for all } |\alpha| \leq r\}$
- $\mathcal{E}(\mathbb{R}^n) = C^\infty(\mathbb{R}^n) = \{f \in C(\mathbb{R}^n); D^\alpha f \in C(\mathbb{R}^n), \text{ for all } \alpha \in \mathbb{N}^n\}$
- $\text{supp}(f) = \overline{\{x \in \mathbb{R}^n; f(x) \neq 0\}}$ : Support of the function  $f$
- $\mathcal{D}(\mathbb{R}^n) = C_0^\infty(\mathbb{R}^n) = \{f \in C^\infty(\mathbb{R}^n); \text{supp}(f) \text{ is compact}\}$
- $\mathcal{S}(\mathbb{R}^n)$ : The Schwartz space and  $\mathcal{S}'(\mathbb{R}^n)$  the space of tempered distributions

$$\mathcal{S}(\mathbb{R}^n) = \left\{ \varphi \in C^\infty(\mathbb{R}^n); \quad \|\varphi\|_{\mathcal{S}(\mathbb{R}^n)} \sim p_{k,m}(\varphi) = \sup_{x \in \mathbb{R}^n} \left( \sup_{\alpha \in \mathbb{N}^n, |\alpha| \leq m} \left( \sup_{\beta \in \mathbb{N}^n, |\beta| \leq k} |x^\beta \partial^\alpha \varphi(x)| \right) \right) < \infty \right\}$$

$$\mathcal{S}'(\mathbb{R}^n) = \{T : \mathcal{S} \rightarrow \mathbb{R}, \exists k \in \mathbb{N}, \exists m \in \mathbb{N}, \exists C_{k,m} > 0, \forall \varphi \in \mathcal{S}(\mathbb{R}^n); |\langle T, \varphi \rangle| < C_{k,m} p_{k,m}(\varphi)\}$$

- We define the norm of the Lebesgue space  $L^p(\mathbb{R}^n)$  by

$$\begin{aligned}\|f\|_{L^p(\mathbb{R}^n)} &= \left( \int_{\mathbb{R}^n} |f(x)|^p dx \right)^{\frac{1}{p}}, \quad \text{if } p \geq 1 \\ \|f\|_{L^\infty(\mathbb{R}^n)} &= \inf\{c > 0; |f(x)| \leq c, \text{ (a.e.)}\} = \text{ess sup}_{x \in \mathbb{R}^n} |f(x)|\end{aligned}$$

- For all  $f, g \in \mathcal{S}(\mathbb{R}^n)$ , the convolution  $f * g$  satisfies  $f * g \in \mathcal{S}(\mathbb{R}^n)$ , such that

$$\forall \xi \in \mathbb{R}^n; f * g(\xi) = \int_{\mathbb{R}^n} \tau_t f(\xi) g(t) dt = \int_{\mathbb{R}^n} f(u) \tau_u g(\xi) du, \quad \text{where } \tau_u g(\xi) = g(\xi - u)$$

- If  $f \in \mathcal{S}(\mathbb{R}^n)$ , we define the Fourier transform and its inverse by

$$\begin{aligned}\mathcal{F}f(\xi) &= \hat{f}(\xi) \sim \int_{\mathbb{R}^n} e^{-i\langle x, \xi \rangle} f(x) dx, \quad \text{for all } \xi \in \mathbb{R}^n \\ \mathcal{F}^{-1}f(\xi) &= \check{f}(\xi) \sim (2\pi)^{-n} \int_{\mathbb{R}^n} e^{i\langle x, \xi \rangle} f(x) dx, \quad \text{for all } \xi \in \mathbb{R}^n \\ (\mathcal{F}T)(f) &= \hat{T}(f) = T(\hat{f}), \quad (\mathcal{F}^{-1}T)(f) = \check{T}(f) = T(\check{f}), \quad \text{for all } T \in \mathcal{S}'(\mathbb{R}^n)\end{aligned}$$

- $[s]$ : Denotes the integer part of  $s \in \mathbb{R}$ .

- $p'$ : Denotes the conjugate exponent of  $p \geq 1$ , defined by  $\frac{1}{p} + \frac{1}{p'} = 1$ .

- $\mathcal{P}_m(\mathbb{R}^n)$ : The set of polynomials of degree at most  $m$ ,  $m \in \mathbb{N}$ , such that

$$\mathcal{P}_m(\mathbb{R}^n) = \left\{ P \in \mathcal{S}'(\mathbb{R}^n); P(x) = \sum_{|\alpha| \leq m} a_\alpha x^\alpha, x \in \mathbb{R}^n \right\}, \quad \mathcal{P}_{-1}(\mathbb{R}^n) = \{0\}$$

- $\mathcal{P}(\mathbb{R}^n) = \mathcal{P}_\infty(\mathbb{R}^n)$ : The subspace of  $\mathcal{S}'(\mathbb{R}^n)$  of polynomials or multinomials.

- $[f]$ : The equivalence class of a distribution  $f \in \mathcal{S}'(\mathbb{R}^n)$  modulo  $\mathcal{P}_\infty(\mathbb{R}^n)$ .

- We agree to take  $x^\alpha = x_1^{\alpha_1} x_2^{\alpha_2} \cdots x_n^{\alpha_n}$ ,  $\alpha! = \alpha_1! \alpha_2! \cdots \alpha_n!$ ,  $t^\alpha x = (t^{\alpha_1} x_1, t^{\alpha_2} x_2, \dots, t^{\alpha_n} x_n)$ ,

$(t \geq 0)$ , and  $\langle x, y \rangle = x_1 y_1 + x_2 y_2 + \cdots + x_n y_n$ .

- We introduce the space  $\mathcal{S}_0(\mathbb{R}^n)$  defined by functions in  $\mathcal{S}(\mathbb{R}^n)$  with zero moments

$$\begin{aligned}\mathcal{S}_0(\mathbb{R}^n) &= \left\{ f \in \mathcal{S}(\mathbb{R}^n); \partial^\alpha \mathcal{F}f(0) = \int_{\mathbb{R}^n} x^\alpha f(x) dx = 0, \text{ for all } \alpha \in \mathbb{N}^n \right\} \\ \widehat{\mathcal{S}}_0(\mathbb{R}^n) &= \left\{ \hat{f}; f \in \mathcal{S}_0(\mathbb{R}^n) \right\} = \left\{ f \in \mathcal{S}(\mathbb{R}^n); \partial^\alpha f(0) = 0, \text{ for all } \alpha \in \mathbb{N}^n \right\}\end{aligned}$$

$$\mathcal{S}'_0(\mathbb{R}^n) = \mathcal{S}'(\mathbb{R}^n)/\mathcal{S}_0^\perp(\mathbb{R}^n) = \mathcal{S}'(\mathbb{R}^n)/\mathcal{P}(\mathbb{R}^n)$$

- For each  $s \in \mathbb{R}^+ \setminus \mathbb{N}$ , we define the Holder space by

$$\begin{aligned}\mathcal{C}^s(\mathbb{R}^n) &= \left\{ f \in L^\infty; \|f\|_{\mathcal{C}^s(\mathbb{R}^n)} = \|f\|_\infty + \sup_{x,y \in \mathbb{R}^n, x \neq y} \frac{|f(x) - f(y)|}{|x - y|^s} < \infty \right\}, \quad \text{if } 0 < s < 1 \\ \mathcal{C}^s(\mathbb{R}^n) &= \left\{ f \in \mathcal{C}^{[s]}(\mathbb{R}^n); \|f\|_{\mathcal{C}^s(\mathbb{R}^n)} = \sum_{|\beta| \leq [s]} \|D^\beta f\|_{\mathcal{C}^\alpha(\mathbb{R}^n)} < \infty \right\}, \quad \text{if } s = [s] + \alpha, 0 < \alpha < 1\end{aligned}$$

For each  $s \in \mathbb{N}$ , we define the Zygmund space by

$$\begin{aligned}\mathcal{C}^1(\mathbb{R}^n) &= \left\{ f \in C^0(\mathbb{R}^n); \|f\|_{\mathcal{C}^1(\mathbb{R}^n)} = \sup_{h \neq 0, x \in \mathbb{R}^n} \frac{|u(x+h) + u(x-h) - 2u(x)|}{|h|} < \infty \right\} \\ \mathcal{C}^s(\mathbb{R}^n) &= \left\{ f \in \mathcal{C}^1(\mathbb{R}^n); D^\alpha f \in \mathcal{C}^1, \text{ for all } |\alpha| \leq s-1 \right\}, \quad \text{if } s \neq 1\end{aligned}$$

- $A \lesssim B$ : Means that for two parametric expressions  $A$  and  $B$ , there exists an independent constant  $c > 0$ , such that  $A \leq cB$ .
  - $A \sim B$ : If  $A \lesssim B$  and  $B \lesssim A$ , then there exist  $c_1, c_2 > 0$ , such that  $c_1 B \leq A \leq c_2 B$ .

# Introduction

The main results of this thesis are related to three problems.

1. The first problem has been studied by several authors, G. Bourdaud, [1], [2], [3], W. Sickel [16], D. Kateb [11], S. Igari [10], and consists in solving the superposition operator problem by finding a necessary and sufficient condition for a composition operator  $T_G : E \rightarrow E$ , such that  $T_G(f) = G \circ f$ , where  $G$  is a real-valued function, to satisfy  $T_G(E) \subset E$ . Several researchers have tried to find non-trivial operators (associated with non-affine functions) verifying the condition  $T_G(E) \subset E$ , for a given functional space  $E$ .
2. The second problem consists in giving a general formula for the derivative of the composition of several functions, using a basic inequality introduced in [2] to generalize some results, studying the properties of composition operators (bounded in particular), such as differentiability, and verifying norm inequalities for the composition of more than two functions.
3. The third problem consists in giving an extension to the famous theorem of Peetre, based on the works introduced in [2] and [14].

Our work is organized into four chapters:

- In the first chapter, we present known functional spaces, such as Besov spaces and their properties. We define the spaces  $BV_p(\mathbb{R})$ ,  $BV_p^1(\mathbb{R})$  from the spaces of functions of bounded p-variation for  $p \geq 1$  and their norms.
- In the second chapter, we study the composition operator problem in the spaces  $BV_p^1(\mathbb{R})$ ,  $1 \leq p < \infty$ , based on the works introduced in [2], [3], as well as some multiplication and continuity properties, and we will extend some results concerning a basic inequality for the composition of several functions. Thus, we obtain a basic algorithm giving the derivative of the composition of several functions.
- In the third chapter, we study functional calculus in homogeneous Besov spaces  $\dot{B}_p^{s,q}(\mathbb{R}^n)$ , presenting the necessary and sufficient conditions for composition operators.
- In the fourth chapter, we present a new functional space  $BV_p^\alpha(\mathbb{R})$  defined thanks to the space  $\mathcal{V}_p^\alpha(\mathbb{R})$ , introduced in [14], and we generalize Peetre's Theorem (3.2.4) to the spaces  $BV_p^\alpha(\mathbb{R})$ ,  $p \in ]1, +\infty[$ ,  $0 \leq \alpha < 1$ , as well as application examples to affirm the stated results.

# Chapter 1

## Some preliminary results

In this chapter, we will recall essential notions, namely the Littlewood-Paley series as well as finite differences, which will allow us to construct equivalent norms for Besov spaces; finally, we will give an overview of functions of bounded  $p$ -variation, as well as their properties that will be very useful later.

### 1.1 Littlewood-Paley series

In this paragraph, we give a definition of the partition of unity in  $C_0^\infty(\mathbb{R}^n)$ , followed by an example, and for more details see [18].

#### 1.1.1 Partition of unity

Let the sequence of reals  $A = \{A_j\}_{j \in \mathbb{N}} \subset \mathbb{R}^+$ , such that there exist  $0 < \lambda_0 \leq \lambda_1$  with

$$\lambda_0 A_j \leq A_{j+1} \leq \lambda_1 A_j, \quad j \in \mathbb{N}.$$

Then there exists  $\ell_0 \in \mathbb{N}$  such that

$$2A_j \leq A_k, \quad \text{for all } j, k \text{ and } j + \ell_0 \leq k.$$

We define  $\Omega_A = \{\Omega_{j,A}\}_{j \in \mathbb{N}}$ , a covering of  $\mathbb{R}^n$ , associated with  $A$ , such that

$$\Omega_{j,A} = \begin{cases} \{\xi \in \mathbb{R}^n; |\xi| \leq A_{j+\ell_0}\}, & \text{if } j = 0, 1, \dots, \ell_0 - 1 \\ \{\xi \in \mathbb{R}^n; A_{j-\ell_0} \leq |\xi| \leq A_{j+\ell_0}\}, & \text{if } j \geq \ell_0 \end{cases}$$

### Definition 1.1.1.

We say that the sequence of functions  $\varphi_A = \{\varphi_{j,A}\}_{j \in \mathbb{N}} \subset C_0^\infty(\mathbb{R}^n)$  is a partition of unity with respect to  $\Omega_A$  if the following conditions are satisfied:

- (i)  $\varphi_{j,A}(\xi) \geq 0, \quad \xi \in \mathbb{R}^n, \quad j \in \mathbb{N}$
- (ii)  $\text{supp}(\varphi_{j,A}) \subset \Omega_{j,A}, \quad j \in \mathbb{N}$
- (iii) For all  $\alpha \in \mathbb{N}^n$ , there exists a constant  $c_\alpha > 0$ , such that

$$|D^\alpha \varphi_{j,A}(\xi)| \leq c_\alpha (1 + |\xi|^2)^{-|\alpha|/2}, \quad \xi \in \mathbb{R}^n, \quad j \in \mathbb{N}$$

- (iv) There exists a constant  $c_\varphi > 0$ , such that

$$0 < \sum_{j=0}^{\infty} \varphi_{j,A}(\xi) = c_\varphi < \infty, \quad \xi \in \mathbb{R}^n.$$

For  $A_j = 2^j$ ,  $\ell_0 = 1$ ,  $c_\varphi = 1$ , we have a dyadic partition of unity.

The partition  $\varphi_A = \{\varphi_{j,A}\}_{j \in \mathbb{N}}$  is called inhomogeneous and can be replaced by the partition  $\{\varphi_{j,A}\}_{j \in \mathbb{Z}}$ , called homogeneous.

### Example 1.1.2.

Let  $K > 1$ , consider the covering  $\{C_p\}_{-1}^{+\infty}$ , defined by

$$\begin{cases} C_p &= \{\xi \in \mathbb{R}^n; K^{-1}2^p \leq |\xi| \leq K2^{p+1}\} \\ C_{-1} &= \overline{B}(0, K) = \{\xi \in \mathbb{R}^n; |\xi| \leq K\} \end{cases}$$

$\{C_p\}_{-1}^{+\infty}$  is a uniformly finite covering for  $\mathbb{R}^n$ , i.e., the set  $\{q \in \mathbb{N}; C_q \cap C_p \neq \emptyset\}$  is finite.

We can construct  $\{\phi_0, \psi_\nu\}$ ,  $\nu \in \mathbb{N}$ , an inhomogeneous Littlewood-Paley decomposition of unity such that

$$\psi_\nu(\xi) = \phi_\nu(\xi) - \phi_{\nu-1}(\xi), \quad \text{and} \quad \phi_\nu(\xi) = \phi_0(2^{-\nu}\xi),$$

where  $\phi_0 \in C_0^\infty(\mathbb{R}^n)$  is real, and  $\phi_0 \equiv 1$  on the closed ball  $\overline{B}(0, 1)$ ,  $\text{supp}(\phi_0) \subset \overline{B}(0, K)$ , such that

$$\left( \phi_0 + \sum_{\nu \in \mathbb{N}} \psi_\nu \right) u = u, \quad u \in \mathcal{S}'(\mathbb{R}^n).$$

Indeed, such a decomposition exists because we have the following lemma.

**Lemma 1.1.3.**

There exist  $\varphi, \psi \in C_0^\infty(\mathbb{R}^n)$ , with  $\text{supp}(\psi) \subset C_{-1}$ ,  $\text{supp}(\varphi) \subset C_0$ , such that

$$\begin{aligned}\varphi(\xi) + \sum_{p=0}^{\infty} \varphi(2^{-p}\xi) &= 1, \\ \psi(\xi) + \sum_{p=0}^N \varphi(2^{-p}\xi) &= \psi(2^{-(N+1)}\xi).\end{aligned}$$

**Proof.**

Let  $\theta \in C_0^\infty(\mathbb{R}^n)$ , with  $0 \leq \theta \leq 1$ ,  $\text{supp}(\theta) \subset C_0$ ,  $\theta(\xi) = 1$  for  $1 \leq |\xi| \leq 2$ .

Define  $s(\xi) = \sum_{p=-\infty}^{\infty} \theta(2^{-p}\xi)$ ,  $\xi \in \mathbb{R}^n \setminus \{0\}$ .

Since  $\{C_p\}_{-1}^{+\infty}$  is a uniformly finite covering for  $\mathbb{R}^n$ , then  $s \in C^\infty(\mathbb{R}^n \setminus \{0\})$ .

Define  $\varphi \in C_0^\infty(\mathbb{R}^n)$  by

$$\varphi(\xi) = \frac{\theta(\xi)}{s(\xi)},$$

such that  $s(2^{-p}\xi) = s(\xi)$ .

For  $|\xi| \geq K$ ,  $p \leq -1$ , we have  $2^{-p}|\xi| = 2^{|p|}|\xi| \geq 2^{|p|}K \geq 2K$ , and  $2^{-p}\xi \notin C_0$ , hence  $\theta(2^{-p}\xi) = 0$ .

Therefore, if  $|\xi| \geq K$ ,  $\xi \in \mathbb{R}^n \setminus C_{-1}$ , then

$$\sum_{p=0}^{\infty} \varphi(2^{-p}\xi) = \sum_{p=-\infty}^{\infty} \varphi(2^{-p}\xi) = \sum_{p=-\infty}^{\infty} \frac{\theta(2^{-p}\xi)}{s(2^{-p}\xi)} = \frac{\sum_{p=-\infty}^{\infty} \theta(2^{-p}\xi)}{\sum_{p=-\infty}^{\infty} s(2^{-p}\xi)} = 1.$$

Take  $\psi(\xi) = 1 - \sum_{p=0}^{\infty} \varphi(2^{-p}\xi)$ , then  $\psi \in C_0^\infty(\mathbb{R}^n)$ , with  $\text{supp}(\psi) \subset C_{-1}$ .

For all  $j \in \mathbb{Z}$ , let  $\psi_j(\xi) = \varphi(2^{-j}|\xi|)$  and  $\psi_0(\xi) = 1 - \sum_{k=1}^{\infty} \psi_k(\xi)$ .

Then we have the following properties:

$$\begin{aligned}\psi_j \text{ is even such that } \text{supp}(\psi_j) &\subset \overline{C}_j, \quad \text{for all } j \in \mathbb{N} \\ \sum_{j=0}^{\infty} \psi_j(\xi) &= 1, \quad \text{for all } \xi \in \mathbb{R}^n \\ C_m \cap \text{supp}(\psi_j) &= \emptyset, \quad \text{if } |m - j| > 1 \\ \psi_{j-1}(\xi) + \psi_j(\xi) + \psi_{j+1}(\xi) &= 1, \quad \text{for all } \xi \in \text{supp}(\psi_j), \quad j \in \mathbb{N}.\end{aligned}$$

To construct a homogeneous partition on  $\mathbb{Z}$ , we introduce the sequence  $\Phi_j \in \mathcal{S}(\mathbb{R}^n)$ , such that:

$$\begin{aligned}\hat{\Phi}_j(\xi) &= \frac{\psi_j(\xi)}{\sum_{k \in \mathbb{Z}} \psi_k(\xi)} \\ \Phi_j(\xi) &= 2^{jn} \Phi_0(2^j \xi), \quad \text{for all } \xi \in \mathbb{R}^n \\ \int_{\mathbb{R}^n} x^k \Phi_j(x) dx &= \hat{\Phi}_j^{(k)}(0) = 0, \quad (k \in \mathbb{N}^n).\end{aligned}$$

We have

$$\sum_{j \in \mathbb{Z}} \hat{\Phi}_j(\xi) = \sum_{j \in \mathbb{Z}} \left( \frac{\psi_j(\xi)}{\sum_{k \in \mathbb{Z}} \psi_k(\xi)} \right) = \frac{\sum_{j \in \mathbb{Z}} \psi_j(\xi)}{\sum_{k \in \mathbb{Z}} \psi_k(\xi)} = 1, \quad (\xi \in \mathbb{R}^n \setminus \{0\}).$$

Since  $1 - \sum_{j=1}^{\infty} \hat{\Phi}_j$  is infinitely differentiable and compactly supported,

we can choose a function  $\Psi \in \mathcal{S}(\mathbb{R}^n)$ , such that  $\hat{\Psi} = 1 - \sum_{j=1}^{\infty} \hat{\Phi}_j$ , where  $\hat{\Psi} \neq 0$  on  $|\xi| \leq 1$ , and thus:

$$\hat{\Psi} + \sum_{j=1}^{\infty} \hat{\Phi}_j = 1.$$

Set  $\psi(\xi) = \hat{\Psi}(\xi)$  and  $\varphi(2^{-j}\xi) = \hat{\Phi}_j(\xi)$ , then we have

$$\psi(\xi) + \sum_{j=1}^{\infty} \varphi(2^{-j}\xi) = 1, \quad (\xi \in \mathbb{R}^n). \quad (1.1)$$

We have  $\psi(2^{-N}\xi) + \sum_{p=0}^{\infty} \varphi(2^{-p-N}\xi) = \psi(\xi) + \sum_{p=0}^{N-1} \varphi(2^{-p}\xi) + \sum_{p=N}^{\infty} \varphi(2^{-p}\xi) = 1$ , hence

$$\begin{aligned}\sum_{p=0}^{\infty} \varphi(2^{-p-N}\xi) &= \sum_{p=N}^{\infty} \varphi(2^{-p}\xi), \\ \psi(2^{-N}\xi) &= \psi(\xi) + \sum_{p=0}^{N-1} \varphi(2^{-p}\xi).\end{aligned}$$

By induction, if  $\varphi(2^{-N}\xi) = \psi(2^{-(N+1)}\xi) - \psi(2^{-N}\xi)$ , and since  $\psi(2^{-(N+1)}\xi) = \psi(\xi) + \sum_{p=0}^N \varphi(2^{-p}\xi)$ , then we have

$$\begin{aligned}\psi(2^{-(N+2)}\xi) &= \psi(\xi) + \sum_{p=0}^{N+1} \varphi(2^{-p}\xi) \\ &= \psi(2^{-(N+1)}\xi) + \varphi(2^{-(N+1)}\xi),\end{aligned}$$

hence  $\varphi(2^{-(N+1)}\xi) = \psi(2^{-(N+2)}\xi) - \psi(2^{-(N+1)}\xi)$ . □

### 1.1.2 Decomposition of a function $f \in \mathcal{S}'(\mathbb{R}^n)$

Since  $\Phi_j$  is real and even,  $\hat{\Phi}_j$  is real and even and its support is  $C_j$ , hence  $\Phi_j \in \mathcal{S}_0(\mathbb{R}^n)$  and thus  $\Phi_j * f$  is defined for all  $f \in \mathcal{S}'_0(\mathbb{R}^n)$ .

Multiplying both sides of equality (1.1.1) by  $u \in \mathcal{S}(\mathbb{R}^n)$  and applying the inverse Fourier transform, we obtain

$$u = \Psi * u + \sum_{j \in \mathbb{N}} \Phi_j * u, \quad \text{for all } u \in \mathcal{S}(\mathbb{R}^n),$$

and we deduce its dual relation

$$\begin{aligned} f &= \Psi * f + \sum_{j \in \mathbb{N}} \Phi_j * f, \quad \text{for all } f \in \mathcal{S}'(\mathbb{R}^n), \\ u &= \sum_{j \in \mathbb{Z}} \Phi_j * u, \quad \text{for all } u \in \mathcal{S}_0(\mathbb{R}^n), \\ f &= \sum_{j \in \mathbb{Z}} \Phi_j * f, \quad \text{for all } f \in \mathcal{S}'_0(\mathbb{R}^n). \end{aligned}$$

For all  $j, k \in \mathbb{Z}$ , we define the continuous operators  $\Delta_k, Q_j$  by

$$\Delta_k : \mathcal{S}'(\mathbb{R}^n) \rightarrow \mathcal{S}'(\mathbb{R}^n) \quad \text{and} \quad Q_j : \mathcal{S}'(\mathbb{R}^n) \rightarrow \mathcal{S}'(\mathbb{R}^n),$$

such that

$$\begin{aligned} Q_j f &= (\psi(2^{-j} \cdot))^\vee * f = 2^{jn} \psi(2^j \cdot) * f, \quad \text{for } j = 0, 1, 2, \dots \\ \Delta_k f &= (\varphi(2^{-k} \cdot))^\vee * f = 2^{kn} \varphi(2^k \cdot) * f, \quad \text{for } k = 1, 2, \dots \end{aligned}$$

We have  $\widehat{\Delta_k f}(\xi) = \hat{\Phi}_k(\xi) \hat{f}(\xi)$ , hence  $\Delta_k f(\xi) = \Phi_k(\xi) * f(\xi)$ , and we set  $Q_0 f(\xi) = F(\xi)$ .

For all  $f \in \mathcal{S}'(\mathbb{R}^n)$  and  $k \in \mathbb{N}$ , (convergence in  $\mathcal{S}'(\mathbb{R}^n)$ ), we have

$$\begin{aligned} Q_k f &= \sum_{j \in \mathbb{Z}, j \leq k} \Delta_j f, \\ f &= \sum_{j \in \mathbb{Z}} \Delta_j f = F + \sum_{j=1}^{\infty} \Delta_j f, \quad \text{for all } f \in \mathcal{S}'_0(\mathbb{R}^n), \end{aligned}$$

such that  $\widehat{\Delta_j f} \neq 0$  on  $A_j$  and  $\hat{F}(\xi) \neq 0$  for  $|\xi| \leq 1$ .

If  $\xi = 0$ , then the expression  $\sum_{j \in \mathbb{Z}} \hat{\Phi}_j(0) = 0$  implies (1.1.1), because if  $\phi^{(k)} = 0$  for all  $k \in \mathbb{N}^n$ , then (1.1.1) is verified for all  $\phi \in \widehat{\mathcal{S}}_0(\mathbb{R}^n)$ .

### 1.1.3 Decomposition of $\Delta_k(f \cdot g)$

We compute

$$\begin{aligned}
\Delta_k(f \cdot g) &= \Delta_k \left( \sum_{j \in \mathbb{N}} \sum_{\ell \in \mathbb{N}} \Delta_j f \cdot \Delta_\ell g \right) \\
&= \sum_{j \in \mathbb{N}} \sum_{\ell \in \mathbb{N}} \Delta_k(\Delta_j f \cdot \Delta_\ell g) \\
&= \left( \prod_{k,1} + \prod_{k,2} + \prod_{k,3} \right) (f, g),
\end{aligned}$$

where

$$\begin{aligned}
\prod_{k,1}(f, g) &= \Delta_{k(1)}(f \cdot g) = \Delta_k(\tilde{\Delta}_k f \cdot Q_{k+1} g) \\
\prod_{k,2}(f, g) &= \Delta_{k(2)}(f \cdot g) = (Q_{k+1} f \cdot \tilde{\Delta}_k g) = \prod_{k,2}(g, f) \\
\prod_{k,3}(f, g) &= \Delta_{k(3)}(f \cdot g) = \sum_{j=k}^{\infty} \Delta_k(\Delta_j f \cdot \bar{\Delta}_j g),
\end{aligned}$$

with  $\tilde{\Delta}_k = \sum_{j=k-2}^{k+4} \Delta_j$ , and  $\bar{\Delta}_k = \sum_{j=k-1}^{k+1} \Delta_j$  (support calculations).

## 1.2 Finite difference operators $\Delta_h^m$

### Definition 1.2.1.

Let  $x, h \in \mathbb{R}^n$ ,  $m \in \mathbb{N}$ , and  $f$  an arbitrary function, introduce the finite difference operator  $\Delta_h$  such that

$$\begin{aligned}
\Delta_h f &= \tau_{-h} f - f, \quad \text{where } \tau_h f(x) = f(x - h), \\
\Delta_h^1 f(x) &= \Delta_h f(x) = f(x + h) - f(x).
\end{aligned}$$

The operators  $\Delta_h^m f$  are defined by the recurrence relation

$$\Delta_h^m f(x) = \Delta_h(\Delta_h^{m-1} f(x)), \quad m \geq 2. \tag{1.2}$$

We deduce that

$$\Delta_h^2 f(x) = \Delta_h(\Delta_h^1 f(x)) = f(x + 2h) - 2f(x + h) + f(x).$$

**Lemma 1.2.2.**

Let  $x, h \in \mathbb{R}^n$ ,  $m \in \mathbb{N}$ , and  $f$  an arbitrary function defined on a subset of  $\mathbb{R}^n$ , then the general term of  $\Delta_h^m f(x)$  can be written as

$$\Delta_h^m f(x) = \sum_{\ell=0}^m \binom{m}{\ell} (-1)^{\ell \pm m} \tau_{-\ell h} f(x) = \sum_{\ell=0}^m \binom{m}{\ell} (-1)^\ell \tau_{(\ell-m)h} f(x).$$

**Proof.**

(i) Since we have  $\binom{m}{\ell-1} + \binom{m}{\ell} = \binom{m+1}{\ell}$ , and  $(-1)^{m-\ell} = (-1)^{m+\ell} = -(-1)^{m+\ell-1} = -(-1)^{m+\ell+1}$ , we deduce by induction that if  $\Delta_h^m f(x) = \sum_{\ell=0}^m \binom{m}{\ell} (-1)^{m+\ell} \tau_{-\ell h} f(x)$ , then according to (1.1.2),

$$\begin{aligned} \Delta_h^{m+1} f(x) &= \Delta_h(\Delta_h^m f(x)) = \Delta_h^m f(x+h) - \Delta_h^m f(x) \\ &= \sum_{k=0}^m \binom{m}{k} (-1)^{m+k} f(x+(k+1)h) - \sum_{k=0}^m \binom{m}{k} (-1)^{m+k} f(x+kh) \\ &= \sum_{\ell=1}^{m+1} \binom{m}{\ell-1} (-1)^{m+\ell-1} f(x+\ell h) - \sum_{k=0}^m \binom{m}{k} (-1)^{m+k} f(x+kh) \\ &= f(x+(m+1)h) - (-1)^m f(x) + \sum_{\ell=1}^m \left( \binom{m}{\ell-1} (-1)^{m+\ell-1} - \binom{m}{\ell} (-1)^{m+\ell} \right) f(x+\ell h) \\ &= f(x+(m+1)h) - (-1)^m f(x) + \sum_{\ell=1}^m \binom{m+1}{\ell} (-1)^{m+\ell+1} f(x+\ell h) \\ &= \sum_{\ell=0}^{m+1} \binom{m+1}{\ell} (-1)^{m+1+\ell} f(x+\ell h). \end{aligned}$$

(ii) From (i) and since  $\binom{m}{\ell} = \binom{m}{m-\ell}$ , then we have

$$\Delta_h^m f(x) = \sum_{\ell=0}^m \binom{m}{m-\ell} (-1)^{m+\ell} f(x+\ell h).$$

Since  $(-1)^{m-\ell} = (-1)^{m+\ell}$ , and for  $0 \leq \ell \leq m$ , we have  $0 \leq m-\ell \leq m$ , so for  $m-\ell = k$ , we have

$$\Delta_h^m f(x) = \sum_{\ell=0}^m \binom{m}{m-\ell} (-1)^{m-\ell} f(x+\ell h) = \sum_{k=0}^m \binom{m}{k} (-1)^k f(x+(m-k)h).$$

Thus, we agree to set

$$\Delta_h^m f(x) = \sum_{k=0}^m \binom{m}{k} (-1)^k f(x + (m-k)h) = \sum_{\ell=0}^m \binom{m}{\ell} (-1)^{m+\ell} \tau_{-\ell h} f(x).$$

□

**Definition 1.2.3.**

Let  $h \in \mathbb{R}^n$ ,  $m \in \mathbb{N}$ ,  $t > 0$ , and  $f \in L^p(\mathbb{R}^n)$ , we define

$$\omega_p^m(t, f) = \sup_{|h| \leq t, h \in \mathbb{R}^n} \|\Delta_h^m f\|_{L^p(\mathbb{R}^n)}, \quad (1.3)$$

as the  $m$ -th order modulus of continuity of  $f$  in  $L^p(\mathbb{R}^n)$ .

This modulus of continuity is used to find equivalent norms.

**Remark 1.2.4.a)** The continuity of a function  $f$  at  $x$  is defined by

$$\lim_{h \rightarrow 0} |\Delta_h^1 f(x)| = \lim_{h \rightarrow 0} |f(x + h) - f(x)| \rightarrow 0.$$

**b)** The differentiability of a function  $f$  at  $x$  is described by

$$\lim_{h \rightarrow 0} \frac{|\Delta_h^1 f(x)|}{|h|} \leq c < \infty, \quad (c > 0).$$

**c)** The moduli of continuity defined in (1.1.3) converge in the  $L^p$  norm, are monotone with respect to the sup operator, and are  $m$ -differentiable (regularity of order  $m$ ).

**d)**  $\omega_p^r(t, f) = 0$  if and only if  $f$  is a polynomial of degree  $\leq r - 1$ .

**e)** For all  $s > 0$ , we have an equivalent norm for the Holder-Zygmund space  $\mathcal{C}^s(\mathbb{R}^n)$

$$\|f\|_{\mathcal{C}^s(\mathbb{R}^n)} \sim \sup_{x \in \mathbb{R}^n} |f(x)| + \sup_{x \in \mathbb{R}^n, 0 < |h| \leq 1} |h|^{-s} |\Delta_h^k f(x)|, \quad \text{for all } k \in \mathbb{N}, k > s.$$

### 1.3 Norms in Besov spaces

In this paragraph, we give the definitions of norms for Besov spaces, using the **Littlewood-Paley** theory and **finite differences**.

### 1.3.1 By Littlewood-Paley theory

#### Definition 1.3.1.

Let  $s \in \mathbb{R}$ ,  $p, q \in [1, +\infty]$ , then we say that a function  $f$  belongs to  $\dot{B}_p^{s,q}(\mathbb{R}^n)$ , the homogeneous Besov space, if and only if  $f \in \mathcal{S}'(\mathbb{R}^n)/\mathcal{P}(\mathbb{R}^n)$  and  $f = \sum_{j \in \mathbb{Z}} \Delta_j f$  such that

$$\|f\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} = \begin{cases} \left( \sum_{j \in \mathbb{Z}} (2^{sj} \|\Phi_j * f\|_p)^q \right)^{1/q} < \infty, & \text{for } 1 \leq q < \infty \\ \sup_{j \in \mathbb{Z}} (2^{sj} \|\Phi_j * f\|_p) < \infty, & \text{for } q = \infty \end{cases}$$

#### Proposition 1.3.2 ([18]).

For all  $f \in \dot{B}_p^{s,q}(\mathbb{R}^n)$  and all  $\lambda > 0$ , there exist two constants  $0 < c_1 \leq c_2$ , such that

$$c_1 \|f\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} \leq \lambda^{n/p-s} \|f(\lambda \cdot)\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} \leq c_2 \|f\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)}. \quad (1.4)$$

#### Proof.

- Let  $g(x) = f(\lambda x)$ ,  $\gamma = s - n/p$ ,  $\lambda = 2^N$ ,  $u = 2^N x$ , then  $du = 2^{nN} dx$ , and we have

$$\begin{aligned} \Phi_j * g(x) &= \int_{\mathbb{R}^n} \Phi_j(t) g(x-t) dt = \int_{\mathbb{R}^n} 2^j \Phi_0(2^j t) f(2^N(x-t)) dt \\ &= \int_{\mathbb{R}^n} 2^{j-N} \Phi_0(2^{j-N} u) f(2^N x - u) du = \int_{\mathbb{R}^n} \Phi_{j-N}(u) f(2^N x - u) du \\ &= \Phi_{j-N} * f(2^N x). \end{aligned}$$

Then

$$\begin{aligned} \|g\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} &= \left( \sum_{j \in \mathbb{Z}} \left( 2^{js} \left( \int_{\mathbb{R}^n} |\Phi_{j-N} * f(2^N x)|^p dx \right)^{1/p} \right)^q \right)^{1/q} \\ &= \left( \sum_{j \in \mathbb{Z}} \left( 2^{js} 2^{-nN/p} \left( \int_{\mathbb{R}^n} |\Phi_{j-N} * f(u)|^p du \right)^{1/p} \right)^q \right)^{1/q} \\ &= (2^N)^\gamma \left( \sum_{k=j-N \in \mathbb{Z}} \left( 2^{ks} \|\Phi_k * f\|_p \right)^q \right)^{1/q} = \lambda^\gamma \|f\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)}. \end{aligned}$$

- Consider  $\lambda > 0$  such that  $2^{N-1} < \lambda \leq 2^N$ , ( $N \in \mathbb{Z}$ ). To prove (1.2.1), we prove one of its

inequalities because if  $\|f(\lambda \cdot)\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} \leq c \|f\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)}$ , then

$$\|f\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} = \|f(\lambda^{-1}(\lambda \cdot))\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} \leq c' \|f(\lambda \cdot)\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)}.$$

We have  $f = \sum_{j \in \mathbb{Z}} \Delta_j f$ , so  $\Delta_k f = \Phi_k * f = \sum_{j \in \mathbb{Z}} \Phi_k * \Delta_j f$ , but  $\text{supp}(\Delta_k f) \subset [2^{k-1}, 2^{k+1}]$ , hence

$$\Delta_k f = \sum_{j \in \mathbb{Z}} \Phi_k * \Delta_j f = \Phi_k * \Delta_{k-1} f + \Phi_k * \Delta_{k+1} f + \Phi_k * \Delta_k f.$$

By induction, if  $\Delta_k f$ ,  $\Delta_{k-1} f$ ,  $\Phi_0$  are decreasing, then  $\Phi_k$ ,  $\Phi_k * \Delta_{k-1} f$ ,  $\Phi_k * \Delta_k f$  are decreasing, and thus  $\Phi_k * \Delta_{k+1} f$  is decreasing as well as  $\Delta_{k+1} f$ , whence

$$\begin{aligned} \Delta_k f(\lambda(x-t)) &= \Phi_k * \Delta_{k-1} f + \Phi_k * \Delta_{k+1} f + \Phi_k * \Delta_k f(\lambda(x-t)) \\ &\leq \Phi_k * \Delta_{k-1} f + \Phi_k * \Delta_{k+1} f + \Phi_k * \Delta_k f(2^{N-1}(x-t)) \\ &= \Delta_k f(2^{N-1}(x-t)), \end{aligned}$$

and  $f(\lambda(x-t)) = \sum_{j \in \mathbb{Z}} \Delta_j f(\lambda(x-t)) \leq \sum_{j \in \mathbb{Z}} \Delta_j f(2^{N-1}(x-t)) = f(2^{N-1}(x-t))$ .

We deduce that

$$\begin{aligned} f(2^N(x-t)) &\leq f(\lambda(x-t)) \leq f(2^{N-1}(x-t)), \\ \Phi_j * f(2^{N-1} \cdot) &\leq \Phi_j * f(\lambda \cdot) \leq \Phi_j * f(2^N \cdot), \\ |\Phi_j * f(\lambda \cdot)| &\leq \max(|\Phi_j * f(2^N \cdot)|, |\Phi_j * f(2^{N-1} \cdot)|), \\ \|\Phi_j * f(\lambda \cdot)\|_p &\leq \max(\|\Phi_j * f(2^N \cdot)\|_p, \|\Phi_j * f(2^{N-1} \cdot)\|_p), \end{aligned}$$

hence

$$\begin{aligned} \|g\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} &\leq \max\left(\|f(2^N \cdot)\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)}, \|f(2^{N-1} \cdot)\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)}\right) \\ &\leq 2^\gamma (2^{N-1})^\gamma \|f\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} \leq C \lambda^\gamma \|f\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)}, \quad C = 2^\gamma. \end{aligned}$$

Thus, we can write  $\|\cdot\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} \sim \lambda^{n/p-s} \|(\lambda \cdot)\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)}$ .

If  $\lambda = 2^m$ , then  $\|f(2^m \cdot)\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} = (2^m)^{s-n/p} \|f\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)}$ .

If  $q = 2$ , then  $\|f(\lambda \cdot)\|_{\dot{B}_p^{s,2}(\mathbb{R}^n)} = \|f(\lambda \cdot)\|_{\dot{H}^{s,p}(\mathbb{R}^n)} = \lambda^{s-n/p} \|f\|_{\dot{H}^{s,p}(\mathbb{R}^n)} = \lambda^{s-n/p} \|f\|_{\dot{B}_p^{s,2}(\mathbb{R}^n)}$ .

We also have  $\|f(\lambda \cdot)\|_{L^p(\mathbb{R}^n)} = \lambda^{-n/p} \|f\|_{L^p(\mathbb{R}^n)}$ .

□

### Definition 1.3.3.

Let  $s \in \mathbb{R}$ ,  $p, q \in [1, +\infty]$ , then we say that a function  $f$  belongs to  $B_p^{s,q}(\mathbb{R}^n)$ , the non-homogeneous Besov space, if and only if  $f \in \mathcal{S}'(\mathbb{R}^n)$  and  $f = Q_0 f + \sum_{j \geq 1} \Delta_j f$ , such that

$$\|f\|_{B_p^{s,q}(\mathbb{R}^n)} = \begin{cases} \|\Psi * f\|_p + \left( \sum_{j \in \mathbb{N}} (2^{sj} \|\Phi_j * f\|_p)^q \right)^{1/q} < \infty, & \text{for } 1 \leq q < \infty \\ \|\Psi * f\|_p + \sup_{j \in \mathbb{N}} (2^{sj} \|\Phi_j * f\|_p) < \infty, & \text{for } q = \infty \end{cases}$$

### 1.3.2 By finite differences

**Lemma 1.3.4** ([18]).

For all  $0 < s < 1$ , we have

$$\sum_{j \in \mathbb{Z}} 2^{js} |\Phi_j(x)| \leq cx^{-(1+s)}, \quad \text{for all } x > 0, \quad (c > 0).$$

**Proof.**

- If  $x = 2^{-N}$ , then  $\Phi_0 \in \mathcal{S}$  implies  $|\Phi_0(x)| \leq cx^{-2}$ , whence

$$\sum_{j=N+1}^{\infty} 2^{js} |\Phi_j(x)| = \sum_{j=N+1}^{\infty} 2^{j(s+1)} |\Phi_0(2^j x)| \leq \sum_{n=N+1}^{\infty} 2^{n(\alpha-1)} x^{-2} = \frac{1}{1-2^{1-\alpha}} \cdot 2^{N(\alpha-1)} x^{-2} = cx^{-(1+s)}.$$

Since  $\Phi_0$  is bounded, then

$$\sum_{j=-\infty}^N 2^{js} |\Phi_j(x)| \leq c \sum_{j=-\infty}^N 2^{j(1+s)} = c2^{N(1+s)} = cx^{-(1+s)}.$$

- If  $2^{-N} < x \leq 2^{-N+1}$ ,  $N \in \mathbb{Z}$ , and by the decrease of  $\Phi_j$ , we have

$$\begin{aligned} \sum_{j \in \mathbb{Z}} 2^{js} |\Phi_j(x)| &\leq \sum_{j \in \mathbb{Z}} 2^{js} |\Phi_j(2^{-N})| \\ &\leq c2^{N(1+s)} = (2^{1+s}c)2^{(N-1)(1+s)} \leq c^{-(1+s)}. \end{aligned}$$

□

**Lemma 1.3.5** ([18]).

Let  $1 \leq p \leq \infty$ ,  $k \in \mathbb{N}$ ,  $f \in \mathcal{S}(\mathbb{R}^n)$ , then

- (i)  $\|\tau_{-h}f - f\|_p \leq |h| \|f'\|_p$
- (ii)  $\|\Delta_h^k f\|_p \leq |h|^k \|f^{(k)}\|_p$ ,  $(k \geq 2)$

**Proof.**

(i) Using the inequality  $|f(x) - f(y)| \leq |x - y|^{1-1/p} \|f'\|_p$ , and since  $p \geq 1$  gives  $0 \leq 1 - 1/p \leq 1$ , taking  $h = x - y$ , we obtain

$$\|\tau_{-h}f - f\|_p \leq |h|^{1-1/p} \|f'\|_p \leq |h| \|f'\|_p.$$

(ii) We obtain by induction

$$\begin{aligned} \|\Delta_h^{k+1} f\|_p &= \|\Delta_h(\Delta_h^k f)\|_p \leq |h| \|(\Delta_h^k f)'\|_p = |h| \|\Delta_h^k f'\|_p \\ &\leq |h| |h|^k \|f'^{(k)}\|_p \leq |h|^{k+1} \|f^{(k+1)}\|_p. \end{aligned}$$

□

**Proposition 1.3.6** ([18]).

Let  $h \in \mathbb{R}^n$ ,  $0 < s < 1$ ,  $1 \leq p \leq \infty$ ,  $1 \leq q < \infty$ , then  $f \in B_p^{s,q}(\mathbb{R}^n)$  if and only if  $f \in L^p(\mathbb{R}^n)$ , such that

$$\int_{\mathbb{R}^n} (|h|^{-s} \|\tau_{-h}f - f\|_p)^q \frac{dh}{|h|^n} < \infty, \quad (1.5)$$

and  $f \in \dot{B}_p^{s,q}(\mathbb{R}^n)$  if and only if  $f \in \mathcal{S}'(\mathbb{R}^n)/\mathcal{P}(\mathbb{R}^n)$ , such that (1.2.2) is verified.

**Proof.**

First, take the simplest case  $q = n = 1$ , because the case  $q = \infty$ , as well as the general case of  $\mathbb{R}^n$ , can easily be deduced using the interpolation theorem of two Besov spaces which gives another Besov space, as well as the Marcinkiewicz theorem.

(1) Let  $f = F + \sum_{j \in \mathbb{N}} \Delta_j f \in B_p^{s,1}(\mathbb{R})$ , then  $f \in L^p$ , such that  $\sum_{j \in \mathbb{N}} \|\Delta_j f\|_p < \infty$ , and thus by Lemma

1.3.5, we have

$$\begin{aligned} \int_0^{2^{-j}} |h|^{-s} \|\Delta_h^1(\Delta_j f)\|_p \frac{dh}{|h|} &\leq \int_0^{2^{-j}} |h|^{1-s} \|(\Delta_j f)'\|_p \frac{dh}{|h|} \\ &\leq c 2^j \|(\Delta_j f)'\|_p \int_0^{2^{-j}} |h|^{-s} dh = c 2^{js} \|(\Delta_j f)'\|_p, \end{aligned}$$

and

$$\int_{2^{-j}}^{\infty} |h|^{-s} \|\Delta_h^1(\Delta_j f)\|_p \frac{dh}{|h|} \leq 2 \|(\Delta_j f)'\|_p \int_{2^{-j}}^{\infty} |h|^{-s} \frac{dh}{|h|} \leq c 2^{js} \|(\Delta_j f)'\|_p,$$

hence

$$\int_0^{\infty} |h|^{-s} \|\Delta_h^1(f)\|_p \frac{dh}{|h|} = \int_0^{\infty} |h|^{-s} \|\Delta_h^1 \left( \sum_{j \in \mathbb{N}} \Delta_j f \right)\|_p \frac{dh}{|h|} \leq c \sum_{j \in \mathbb{N}} 2^{js} \|(\Delta_j f)'\|_p < \infty.$$

(2) Let  $f \in L^p$ , and  $\int_0^{\infty} |h|^{-s} \|\Delta_h^1(f)\|_p \frac{dh}{|h|} < \infty$ , then  $f * \Psi \in L^p$ .

We have  $\Phi_j * f(x) = \int_{\mathbb{R}} \Phi_j(y) (f(x-y) - f(x)) dy$ , hence by Minkowski's inequality

$$\begin{aligned} \|\Phi_j * f\|_p &= \left\| \int_{-\infty}^0 \Phi_j(y) (f(x-y) - f(x)) dy + \int_0^{\infty} \Phi_j(y) (f(x-y) - f(x)) dy \right\|_p \\ &\leq 2 \int_0^{\infty} |\Phi_j(y)| \|\tau_{-y} f - f\|_p dy. \end{aligned}$$

Using Lemma 1.3.4, we obtain  $f \in B_p^{s,1}(\mathbb{R})$ , because

$$\begin{aligned} \sum_{j \in \mathbb{N}} 2^{js} \|\Phi_j * f\|_p &\leq 2 \int_0^{\infty} \sum_{j \in \mathbb{N}} 2^{js} |\Phi_j(y)| \|\tau_{-y} f - f\|_p dy \\ &\leq c \int_0^{\infty} |y|^{-(1+s)} \|\tau_{-y} f - f\|_p dy = c \int_0^{\infty} |h|^{-s} \|\Delta_h^1(f)\|_p \frac{dh}{|h|} < \infty. \end{aligned}$$

Note that (1.2.2) is equivalent to  $\int_{|h| \leq 1} (|h|^{-s} \|\tau_{-h} f - f\|_p)^q \frac{dh}{|h|^n}$ , because

$$\int_{\mathbb{R}^n} (|h|^{-s} \|\tau_{-h} f - f\|_p)^q \frac{dh}{|h|^n} = \int_{|h| < 1} (...) dh + \int_{|h| \geq 1} (...) dh,$$

and the integral  $\int_{|h| \geq 1} (|h|^{-s} \|\tau_{-h} f - f\|_p)^q \frac{dh}{|h|^n}$  is transformed by a change of variable into

$$\int_{|h| \leq 1} (|h|^{-s} \|\tau_{-h} f - f\|_p)^q \frac{dh}{|h|^n}.$$

□

**Theorem 1.3.7** ([16]).

Let  $h \in \mathbb{R}^n$ ,  $s > 0$ ,  $1 \leq q \leq \infty$ ,  $1 \leq p < \infty$ ,  $M \in \mathbb{N}$ ,  $M \leq s < M + 1$ , then  $f \in \dot{B}_p^{s,q}(\mathbb{R}^n)$  if and only if  $f \in \mathcal{S}'(\mathbb{R}^n)/\mathcal{P}(\mathbb{R}^n)$ , and such that

$$\|f\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} \sim \left( \int_{\mathbb{R}^n} |h|^{-sq} \|\Delta_h^{M+1} f\|_p^q \frac{dh}{|h|^n} \right)^{1/q} < \infty.$$

The term  $\int_{\mathbb{R}^n} \dots dh$  can be replaced by  $\int_{|h|<\varepsilon} \dots dh$ , for any  $\varepsilon > 0$ , and in general we take for  $\varepsilon = 1$ , the term  $\int_{|h|<1} \dots dh$  in the sense of equivalent semi-norms.

**Proof.**

As in the proof of Proposition 1.3.6, suppose  $f \in B_p^{s,1}(\mathbb{R})$ ,  $0 < s < 1$ , thus  $f \in L^p$  and

$$\int_0^\infty |h|^{-s} \|\Delta_h^1 f\|_p \frac{dh}{|h|} < \infty.$$

Since  $\Psi * f \in L^p$ , then for all  $j \in \mathbb{N}$ , we have  $\|\Phi_j * f\|_p \leq c \|\Delta_h^1 f\|_p$ .

By Lemma 1.3.5, and for  $2^{-(j+1)} \leq h \leq 2^{-j}$ , we obtain

$$\int_{2^{-(j+1)}}^{2^{-j}} |h|^{-s} \|\Delta_h^1 f\|_p \frac{dh}{|h|} \geq c \|\Phi_j * f\|_p \int_{2^{-(j+1)}}^{2^{-j}} |h|^{-s} \frac{dh}{|h|} = c 2^{js} \|\Phi_j * f\|_p,$$

hence

$$\sum_{j=0}^{\infty} 2^{js} \|\Phi_j * f\|_p \leq c \sum_{j=0}^{\infty} \int_{2^{-(j+1)}}^{2^{-j}} |h|^{-s} \|\Delta_h^1 f\|_p \frac{dh}{|h|} \leq c \int_0^1 |h|^{-s} \|\Delta_h^1 f\|_p \frac{dh}{|h|} < \infty.$$

– If  $s \geq 1$ , then  $M \leq s < M + 1$ , whence  $0 \leq s - M < 1$ , and thus

$$\|f\|_{\dot{B}_p^{s,q}(\mathbb{R})} \sim \int_0^\infty |h|^{-(s-M)} \|\Delta_h^1 f\|_p \frac{dh}{|h|} \sim \int_0^\infty |h|^{-s} \|\Delta_h^{M+1} f\|_p \frac{dh}{|h|}.$$

□

**Theorem 1.3.8** ([16]).

Let  $h \in \mathbb{R}^n$ ,  $0 < q \leq \infty$ ,  $0 < p < \infty$ ,  $M \in \mathbb{N}$ , and denote  $\sigma_p$  by

$$\sigma_p = \max \left( 0, \frac{n}{p} - n \right).$$

(i) Suppose  $\sigma_p < s < M$ , then

$$\|f\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} \sim \left( \int_{\mathbb{R}^n} |h|^{-sq} \|\Delta_h^M f\|_p^q \frac{dh}{|h|^n} \right)^{1/q}.$$

(ii) If  $0 < q \leq \infty$ ,  $0 < p < \infty$ ,  $s > \sigma_p$ ,  $M \leq s < M + 1$ , then

$$\|f\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} \sim \left( \int_{\mathbb{R}^n} |h|^{-sq} \|\Delta_h^{M+1} f\|_p^q \frac{dh}{|h|^n} \right)^{1/q}.$$

**Proof.**

For the proof, see for example [17], paragraph 3.5.3. □

## 1.4 Functions of bounded p-variation

Throughout this paragraph,  $I$  denotes an interval of  $\mathbb{R}$ .

### 1.4.1 General notions

In this subsection, we give the definitions of functions of bounded p-variation  $\mathcal{V}_p(I)$  and the spaces  $\mathcal{BV}_p(I)$ ,  $BV_p(I)$ , of their equivalence classes with the equivalence relation of equality almost everywhere, as well as the space of their primitives  $BV_p^1(I)$ , and their properties that we will use later.

#### Definition 1.4.1.

Let  $p \in [1, +\infty[$ , then a function  $f : I \rightarrow \mathbb{R}$  is said to be of bounded p-variation, or briefly p-v b, if for all finite strict real sequences  $t_0 < t_1 < \dots < t_N$  of  $I$ , there exists  $c > 0$ , such that

$$\begin{aligned} \sum_{k=1}^N |f(t_k) - f(t_{k-1})|^p &\leq c^p, \quad \text{or equivalently} \\ \sup_{\{t_k\} \subset I} \left[ \sum_{k=1}^N |f(t_k) - f(t_{k-1})|^p \right] &< \infty. \end{aligned}$$

We denote the set of these functions by  $\mathcal{V}_p(I)$ , ( $\mathcal{V}_p$  if  $I = \mathbb{R}$ ), and the minimum of such constants  $c$  with respect to  $f$  by  $\nu_p(f, I)$ , ( $\nu_p(f)$  if  $I = \mathbb{R}$ )

$$\nu_p(f, I) = \inf_c \left\{ c > 0 : \sum_{k=1}^N |f(t_k) - f(t_{k-1})|^p \leq c^p, \{t_i\}_{1 \leq i \leq n} \subset I \right\}.$$

Definition 1.4.1 is equivalent to the fact that for any family of disjoint intervals  $I_k = [a_k, b_k] \subset I$ , we have

$$\left( \sum_{I_k} |f(a_k) - f(b_k)|^p \right)^{1/p} \leq c < \infty, \quad (c > 0).$$

The space  $\mathcal{V}_1(I)$ , or simply  $\mathcal{V}(I)$ , is called the space of functions of bounded variation on  $I$ , and  $\mathcal{V}_\infty(I)$  is a Banach space for the norm

$$\|f\|_{\mathcal{V}_\infty(I)} = \nu_\infty(f, I) = \sup_{x \in I} |f(x)|.$$

**Definition 1.4.2.**

A function  $f : A \rightarrow \mathbb{R}^n$ ,  $A \subset \mathbb{R}^n$ , is said to be  $\gamma$ -Lipschitz,  $\gamma \geq 0$ , of order  $\alpha$ ,  $0 < \alpha \leq 1$ , if and only if

$$|f(x) - f(y)| \leq \gamma|x - y|^\alpha, \quad \text{for all } x, y \in A.$$

The set of these functions is denoted by  $\text{Lip}_\alpha(A)$ , ( $\text{Lip}_\alpha$  if  $A = \mathbb{R}^n$ ). We also say that a function is Lipschitz if it is  $\gamma$ -Lipschitz for some  $\gamma \geq 0$ , and we equip  $\text{Lip}_\alpha(A)$  with the following norm

$$\|f\|_{\text{Lip}_\alpha(A)} = \sup_{x, y \in A, x \neq y} \frac{|f(x) - f(y)|}{|x - y|^\alpha}.$$

A function  $f : A \rightarrow \mathbb{R}^n$  is said to be locally  $\gamma$ -Lipschitz if at every point in  $A$  there exists a neighborhood where  $f$  is  $\gamma$ -Lipschitz. Note that

$$B_\infty^{s, \infty}(\mathbb{R}) = \mathcal{C}^s(\mathbb{R}) = \text{Lip}_s(\mathbb{R}), \quad \text{if } 0 < s < 1.$$

**Proposition 1.4.3** ([2]).

For all  $x, y \in I$  and  $p \in [1, +\infty[$ , each element of  $\mathcal{V}_p(I)$  is a bounded function; moreover,  $\mathcal{V}_p(I)$  becomes a Banach space if equipped with the norm

$$\|f\|_{\mathcal{V}_p(I)} = \sup_{x \in I} |f(x)| + \nu_p(f, I). \quad (1.6)$$

**Proof.**

By a sequence with only two terms, we obtain  $|f(t_k) - f(t_{k-1})|^p \leq (\nu_p(f, I))^p$ , for all  $t_k, t_{k-1} \in I$ .

If we take  $t_{k-1} = 0$  and  $t_k = x$ , then  $|f(x) - f(0)| \leq \nu_p(f, I)$ , hence

$$\begin{aligned} |f(x)| &= |f(x) - f(0) + f(0)| \\ &\leq |f(x) - f(0)| + |f(0)| \\ &\leq \nu_p(f, I) + |f(0)| = C < \infty. \end{aligned}$$

Thus, each function in  $\mathcal{V}_p(I)$  is bounded, and the norm (1.6) satisfies all conditions making  $\mathcal{V}_p(I)$  a Banach space.  $\square$

### 1.4.2 Functions of bounded p-variation as distributions

**Definition 1.4.4** ([2]).

Let  $p \in [1, +\infty]$ , we denote by  $\mathcal{BV}_p(I)$  the set of functions  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that there exists a function  $g \in \mathcal{V}_p(I)$  that coincides with  $f$  almost everywhere,

$$\begin{aligned} \mathcal{BV}_p(I) &= \{f : I \rightarrow \mathbb{R}; \exists g \in \mathcal{V}_p(I), \text{ such that } f = g \text{ (a.e.)}\}, \\ \varepsilon_p(f, I) &= \inf \{\nu_p(g, I); g \in \mathcal{V}_p(I), \text{ such that } g = f \text{ (a.e.)}\}. \end{aligned}$$

– We denote by  $BV_p(I)$  the quotient set with respect to the equivalence relation "equality in  $\mathcal{BV}_p(I)$  almost everywhere", such that

$$\begin{aligned} \dot{f} &= \{g \in \mathcal{BV}_p(I); g = f \text{ (a.e.)}\}, \\ BV_p(I) &= \left\{ \dot{f}; f \in \mathcal{BV}_p(I) \right\} = \mathcal{BV}_p(I)/a.e. \end{aligned}$$

– If  $h \in BV_p(I)$ , we denote by  $\varepsilon_p(h, I)$  the number  $\varepsilon_p(f, I)$ , for any representative  $f$  of  $h$ .

**Definition 1.4.5.**

Let  $f : I \rightarrow \mathbb{R}$  be a function having discontinuities only of the first kind, then we say that  $f$  is normalized if

$$\begin{aligned} f(x) &= \frac{1}{2}(f(x^+) + f(x^-)), \quad \text{for all } x \in \mathring{I}, \\ f(x) &= \lim_{y \rightarrow x, y \in \mathring{I}} f(y), \quad \text{for all } y \in I \cap \partial I, \end{aligned}$$

where  $\partial I = \bar{I} \setminus \mathring{I}$  is the boundary of  $I$ ,  $\bar{I}$  is the closure, and  $\mathring{I}$  is the interior, such that

$$f(x^+) = \lim_{h>0, h \rightarrow 0} f(x+h), \quad \text{and} \quad f(x^-) = \lim_{h>0, h \rightarrow 0} f(x-h).$$

**Proposition 1.4.6** ([2]).

If  $f$  is a function in  $\mathcal{V}_p(I)$ , then the function  $\tilde{f}$  defined by

$$\begin{aligned}\tilde{f}(x) &= \frac{1}{2}(f(x^+) + f(x^-)), \quad \text{for all } x \in \mathring{I}, \\ \tilde{f}(x) &= \lim_{y \rightarrow x, y \in \bar{I}} f(y), \quad \text{for all } x \in I \cap \partial I,\end{aligned}$$

is normalized, and belongs to  $\mathcal{V}_p(I)$ , and satisfies the following inequalities

$$\nu_p(\tilde{f}, I) \leq \nu_p(f, I), \quad \text{and} \quad \sup_I |\tilde{f}| \leq \sup_I |f|.$$

**Proof.**

For the proof, see [2], [3]. □

**Proposition 1.4.7** ([2]).

Let  $p \in [1, +\infty]$ , and  $f \in BV_p(\mathbb{R})$ , then  $f$  has a unique normalized representative  $\tilde{f} \in \mathcal{V}_p$ , such that  $\varepsilon_p(f) = \nu_p(\tilde{f})$ .

We thus consider the space  $BV_p(\mathbb{R})$  as a Banach space of distributions, equipped with the following norm

$$\begin{aligned}\|f\|_{BV_p(\mathbb{R})} &= \varepsilon_p(f) + \|f\|_\infty = \nu_p(\tilde{f}) + \sup_{x \in \mathbb{R}} |\tilde{f}(x)|, \quad \text{if } p < \infty, \\ \|f\|_{BV_\infty(\mathbb{R})} &= \|f\|_\infty = \sup_{x \in \mathbb{R}} |\tilde{f}(x)|.\end{aligned}$$

**Definition 1.4.8.**

Let  $p \in [1, +\infty]$ , then a function  $f : I \rightarrow \mathbb{R}$  belongs to  $BV_p^1(I)$  if there exist  $\alpha, x_0 \in \mathbb{R}$  and  $g \in BV_p(I)$ , such that for all  $x \in I$ , we have

$$f(x) = \alpha + \int_{x_0}^x g(t) dt. \tag{1.7}$$

If (1.2.4) is verified, then  $f$  is a Lipschitz continuous function, and we equip  $BV_p^1(I)$  with the norm  $\|f\|_{BV_p^1(I)} = |f(x_0)| + \|f'\|_{BV_p(I)}$ , for which  $BV_p^1(I)$  becomes a Banach space, and for each point  $x_0 \in I$ ,

we obtain an equivalent norm.

**Proposition 1.4.9** ([3]).

For any interval  $I$  of  $\mathbb{R}$ , the space  $\mathcal{V}_p(I)$ ,  $p \geq 1$ , is a Banach algebra for pointwise multiplication of functions, such that

$$\|fg\|_{\mathcal{V}_p(I)} \leq \|f\|_{\mathcal{V}_p(I)}\|g\|_{\mathcal{V}_p(I)}, \quad \text{for all } f, g \in \mathcal{V}_p(I).$$

**Proof.**

Let  $x_0 < x_1 < \dots < x_n$  be a finite sequence in  $I$  and  $f, g \in \mathcal{V}_p(I)$ . Since  $p \geq 1$ , and using Minkowski's inequality, we obtain

$$\begin{aligned} \left( \sum_{j=1}^N |fg(x_j) - fg(x_{j-1})|^p \right)^{1/p} &\leq \left( \sum_{j=1}^N |f(x_j)(g(x_j) - g(x_{j-1}))|^p \right)^{1/p} \\ &+ \left( \sum_{j=1}^N |g(x_{j-1})(f(x_j) - f(x_{j-1}))|^p \right)^{1/p} \\ &\leq \sup_I |f| \left( \sum_{j=1}^N |g(x_j) - g(x_{j-1})|^p \right)^{1/p} + \sup_I |g| \left( \sum_{j=1}^N |f(x_j) - f(x_{j-1})|^p \right)^{1/p} \\ &\leq \sup_I |f| \nu_p(g, I) + \sup_I |g| \nu_p(f, I), \end{aligned}$$

hence

$$\nu_p(fg, I) = \sup_I \left( \sum_{j=1}^N |fg(x_j) - fg(x_{j-1})|^p \right)^{1/p} \leq \sup_I |f| \nu_p(g, I) + \sup_I |g| \nu_p(f, I),$$

and thus

$$\begin{aligned} \|fg\|_{\mathcal{V}_p(I)} &= \nu_p(fg, I) + \sup_I |f| \sup_I |g| \\ &\leq \sup_I |f| \nu_p(g, I) + \sup_I |g| \nu_p(f, I) + \sup_I |f| \sup_I |g| \\ &\leq \sup_I |f| \nu_p(g, I) + \sup_I |g| \nu_p(f, I) + \sup_I |f| + \sup_I |g| + \nu_p(g, I) \nu_p(f, I) \\ &= \left( \sup_I |f| + \nu_p(f, I) \right) \left( \sup_I |g| + \nu_p(g, I) \right) = \|f\|_{\mathcal{V}_p(I)}\|g\|_{\mathcal{V}_p(I)}. \end{aligned}$$

Therefore,  $\|fg\|_{\mathcal{V}_p(I)} \leq \|f\|_{\mathcal{V}_p(I)}\|g\|_{\mathcal{V}_p(I)}$ , for all  $f, g \in \mathcal{V}_p(I)$ .  $\square$

# Chapter 2

## Composition of Operators in $BV_p^1(I)$

In this chapter, we give a reminder of some basic notions concerning composition operators, then we present the properties of the spaces  $BV_p^1(I)$ , where  $I$  is an interval of  $\mathbb{R}$ , then we present some fundamental results of functional calculus on these spaces, then we present two fundamental Theorems 2.3.2 and 2.3.6, due to the works of [2].

Finally, we give our contribution, namely Theorem 2.4.2 which is a generalization of a fundamental inequality (Theorem 2.3.2), based on Lemma 2.4.1 which is an algorithm giving the derivative of the composition of  $n$  functions.

### 2.1 Reminder

#### Definition 2.1.1.

Let  $E$  be a functional space and let  $\phi$  be a real-valued function, we define the composition operator  $T_\phi$  associated with  $\phi$  by

$$T_\phi(f) = \phi \circ f, \quad \text{for all } f \in E.$$

In general,  $T_\phi$  is nonlinear, and the **Problem of Superposition Operators (P.S.O)** for  $E$  consists in finding the set  $S(E)$  of real-valued functions  $\phi$  such that  $T_\phi(E) \subseteq E$

$$S(E) = \{\phi : \mathbb{R} \rightarrow \mathbb{R}; T_\phi(E) \subseteq E\}.$$

If  $T_\phi(E) \subseteq E$ , then we say that the composition operator

$$T_\phi : E \rightarrow E$$

operates on the functional space  $E$ .

**Remark 2.1.2** ([2]).

We agree to say that a superposition operator  $T_f : E \rightarrow E$  satisfies the norm composition inequality property for a normed space  $E$  if it verifies

$$\|T_f(g)\|_E \leq c_f(1 + \|g\|_E), \quad (c_f > 0), \quad \text{for all } g \in E.$$

Note that if a composition operator satisfies the norm inequality property in the normed space  $E$ , this implies that it operates on  $E$ .

## 2.2 Properties of the spaces $BV_p^1(I)$

In all that follows,  $I$  denotes an interval of  $\mathbb{R}$ .

All results in this paragraph are due to [2] and [3].

**Proposition 2.2.1** ([3]).

If  $f \in BV_p^1(I)$ , then  $f$  can be extended uniquely to a function in  $BV_p^1(\bar{I})$  with the same norm as  $f$  in  $BV_p^1(I)$ .

**Proof.**

For the proof, see [3]. □

By the above proposition, and exploiting affine transformations, the study of the spaces  $BV_p^1(I)$  can be reduced to the following three cases:

$$I = \mathbb{R}, \quad I = [0, +\infty[, \quad I = [0, 1].$$

**Proposition 2.2.2.**

The homogeneity property

$$\|f(\lambda \cdot)'\|_{BV_p} = \lambda \|f'\|_{BV_p}, \quad \text{for all } \lambda > 0$$

is verified for each function  $f$  in  $BV_p^1(\mathbb{R})$  or  $BV_p^1([0, +\infty[)$ .

Thus, the spaces  $BV_p^1(\mathbb{R})$  and  $BV_p^1([0, +\infty[)$  can be seen as analogous to the homogeneous Sobolev

space

$$\dot{W}^{1,p}(\mathbb{R}) = \{f : f'^p(\mathbb{R})\},$$

equipped with the semi-norm  $\|f'\|_p$ , while the usual Sobolev space  $W^{1,p}(\mathbb{R})$  is non-homogeneous:

$$W^{1,p}(\mathbb{R}) = \dot{W}^{1,p}(\mathbb{R}) \cap L^p(\mathbb{R}).$$

**Proposition 2.2.3** ([3]).

- (i)  $BV_p^1(I) \cap L^p(I) = BV_p^1(I)$  if and only if  $I$  is bounded.
- (ii)  $BV_p^1(I) \cap L^p(I)$  injects continuously into  $BV_p(I)$ .

**Proof.**

For the proof, see [3].

□

**Theorem 2.2.4** ([3]).

$BV_p^1(I) \cap L^p(I)$  is a Banach algebra.

**Proof.**

For the proof, see [2] and [3].

□

**Proposition 2.2.5** ([3]).

Let  $I$  be a compact interval of  $\mathbb{R}$ , then we have the following multiplication properties

If  $f, g \in BV_p^1(\mathbb{R})$ , and  $\text{supp}(g) \subseteq I$ , then  $fg \in BV_p^1(\mathbb{R})$ ,

moreover, there exists  $c > 0$ , such that for all  $f, g \in BV_p^1(\mathbb{R})$ ,  $\text{supp}(g) \subseteq I$ , we have

$$\|fg\|_{BV_p^1(\mathbb{R})} \leq c\|f\|_{BV_p^1(\mathbb{R})}\|g\|_{BV_p^1(\mathbb{R})}.$$

**Proof.**

For the proof, see [2] and [3].

□

## 2.3 Solution of the S.O.P $BV_p^1(I)$

Theorems 2.3.2 and 2.3.6 are due to [2].

**Lemma 2.3.1** ([8]).

Let  $a, b, c \in \mathbb{R}$ ,  $a < b < c$ , and let  $h$  be a measurable function defined on  $[a, c]$  with real values such that

$$\begin{aligned}\int_a^b h(x) dx &\geq 0, \\ \int_b^c h(x) dx &< 0,\end{aligned}$$

then there exist  $u, v \in ]a, c[$ , such that  $h(u)h(v) \leq 0$ .

**Proof.**

Applying Bonnet's theorem (second mean value theorem for integrals) to the function  $h$  on the interval  $[a, b]$ , then there exists an average value  $\langle h \rangle_{[a,b]} = u \in ]a, b[$ , such that

$$\int_a^b h(x) dx = h(u)(b - a).$$

Since  $\int_a^b h(x) dx \geq 0$ , and  $b - a \geq 0$ , then  $h(u) \geq 0$ . Similarly, for the interval  $[b, c]$ , there exists an average value  $\langle h \rangle_{[b,c]} = v \in ]b, c[$ , such that

$$\int_b^c h(x) dx = h(v)(c - b).$$

Since  $\int_b^c h(x) dx \leq 0$ , and  $c - b \geq 0$ , then  $h(v) \leq 0$ , whence the existence of  $u, v$  in  $]a, c[$ , such that  $h(u)h(v) \leq 0$ .  $\square$

**Theorem 2.3.2** ([2], (Basic inequality)).

If  $p \in [1, +\infty[$ ,  $t_0 \in I$ ,  $\alpha \in \mathbb{R}$ ,  $h \in \mathcal{V}_p(I)$ , such that

$$\begin{aligned}g(t) &= \alpha + \int_{t_0}^t h(x) dx, \quad f \in \mathcal{V}_p(g(I)), \text{ then} \\ \nu_p((f \circ g) \cdot h, I) &\leq \nu_p(f, g(I)) \cdot (\sup_I |h| + 2^{1/p} \cdot \nu_p(h, I)) + \nu_p(h, I) \cdot \sup_{g(I)} |f|, \\ \|(f \circ g) \cdot h\|_{\mathcal{V}_p(I)} &\leq 2^{1/p} \|f\|_{\mathcal{V}_p(g(I))} \|h\|_{\mathcal{V}_p(I)}.\end{aligned}$$

**Proof.**

1) We have

$$\begin{aligned} \sum_{j=1}^N |(f \circ g) \cdot h(t_j) - (f \circ g) \cdot h(t_{j-1})| &= \sum_{k=0}^{N-1} |(f \circ g) \cdot h(t_{k+1}) - (f \circ g) \cdot h(t_k)| \\ &= \sum_{k=0}^{N-1} |[f \circ g(t_{k+1}) - f \circ g(t_k)] \cdot h(t_k) + f \circ g(t_{k+1}) \cdot (h(t_{k+1}) - h(t_k))|, \end{aligned}$$

hence by Minkowski's inequality

$$\left( \sum_{j=1}^N |(f \circ g) \cdot h(t_j) - (f \circ g) \cdot h(t_{j-1})|^p \right)^{1/p} \leq A_1^{1/p} + B_1^{1/p},$$

such that

$$\begin{aligned} A_1 &= \sum_{k=0}^{N-1} |[f \circ g(t_{k+1}) - f \circ g(t_k)] \cdot h(t_k)|^p, \\ B_1 &= \sum_{k=0}^{N-1} |f \circ g(t_{k+1}) \cdot [h(t_{k+1}) - h(t_k)]|^p. \end{aligned}$$

Since  $|f \circ g(t_{k+1})| \leq \sup_{g(I)} |f|$ , for all  $k \in \mathbb{N}$ , then

$$B_1^{1/p} \leq (\sup_{g(I)} |f|) \cdot \left( \sum_{k=0}^{N-1} |h(t_{k+1}) - h(t_k)|^p \right)^{1/p} \leq (\sup_{g(I)} |f|) \cdot \nu_p(h, I).$$

There exists a sequence  $0 = n_0 < n_1 < \dots < n_J = N$ , such that for all indices  $j = 1, \dots, J$ , we have

- (i) The restriction  $s_j$  of  $(g(t_k))_{0 \leq k \leq N}$  to  $\{k\}_{n_{j-1} \leq k \leq n_j, k \in \mathbb{N}}$  is monotone.
- (ii) The restriction of  $(g(t_k))_{0 \leq k \leq N}$  to  $\{k\}_{n_{j-1} \leq k \leq n_j + 1}$  is not monotone.

Thus,  $[0, N] = [n_0, n_1 - 1] \cup [n_1, n_2 - 1] \cup \dots \cup [n_{J-1}, n_J - 1]$ .

Set  $A_1 = A_2 + B_2$ , such that

$$\begin{aligned} A_2 &= \sum_{j=1}^{J-1} \sum_{k=n_{j-1}}^{n_j-1} |f(g(t_{k+1})) - f(g(t_k))|^p \cdot |h(t_k)|^p, \\ B_2 &= \sum_{k=n_{J-1}}^{n_J-1} |f(g(t_{k+1})) - f(g(t_k))|^p \cdot |h(t_k)|^p. \end{aligned}$$

- If  $J \neq 1$ , then  $B_2 \leq \nu_p^p(f, g(I)) \cdot \sup_I |h|^p$ , and  $A_2 \leq \nu_p^p(f, g(I)) \cdot \sum_{j=1}^{J-1} |h(t_{k_j})|^p$ .

There always exists some  $a_j \in ]t_{k_j}, t_{n_j+1}[$ , such that  $h(a_j)h(t_{k_j}) \leq 0$ . If  $h(t_{k_j}) = 0$  or if  $h$  changes sign on the interval  $]t_{k_j}, t_{n_j+1}[$ , the result is direct; otherwise, we can consider a function  $\hat{h}$  defined from functions of the form  $\pm h$  such that

$$\int_{t_{k_j}}^b \hat{h}(x) dx \geq 0 \quad \text{and} \quad \int_b^{t_{n_j+1}} \hat{h}(x) dx < 0, \quad \text{where } b \in ]t_{k_j}, t_{n_j+1}[,$$

then according to Lemma 2.3.1, there exist  $a'_j, a''_j \in ]t_{k_j}, t_{n_j+1}[$ , such that  $\hat{h}(a'_j)\hat{h}(a''_j) < 0$ , hence

$$\forall j = 1, \dots, J-1, \exists a_j \in ]t_{k_j}, t_{n_j+1}[ : \hat{h}(a_j)\hat{h}(t_{k_j}) \leq 0,$$

such that

$$\sum_{j=1}^{J-1} |h(t_{k_j})|^p = \sum_{j=1}^{J-1} |\hat{h}(t_{k_j})|^p \quad \text{and} \quad \nu_p^p(h, I) = \nu_p^p(\hat{h}, I).$$

Let  $M = \max\{m \in \mathbb{N} : 2m + 1 \leq J\}$ , then

$$\begin{aligned} \sum_{j=1}^{J-1} |h(t_{k_j})|^p &= \begin{cases} \sum_{j=1}^{2M} |h(t_{k_j})|^p, & \text{if } J-1 \text{ is even,} \\ \sum_{j=1}^{2M} |h(t_{k_j})|^p + |h(t_{k_{J-1}})|^p, & \text{if } J-1 \text{ is odd,} \end{cases} \\ \text{hence } \sum_{j=1}^{2M} |h(t_{k_j})|^p &\leq \sum_{l=1}^M |h(t_{k_{2l}}) - h(a_{2l})|^p + \sum_{l=1}^M |h(t_{k_{2l-1}}) - h(a_{2l-1})|^p, \\ \text{and } |h(t_{k_{J-1}})|^p &\leq |h(t_{k_{J-1}}) - h(a_{J-1})|^p. \end{aligned}$$

By the inequalities  $t_{n_{j-1}} \leq t_{k_j} < t_{n_j}$ , and  $t_{k_j} < a_j < t_{n_j+1}$ , for all  $j = 1, \dots, J-1$ , we deduce that  $a_j < t_{n_j+1} \leq t_{n_{j+1}} \leq t_{k_{j+2}}$ . - If  $J \geq 4$  and  $j = 1, \dots, J-3$ , then the intervals  $[t_{k_{2l}}, a_{2l}]$  are pairwise disjoint for  $l = 1, \dots, M$ , and the intervals  $[t_{k_{2l-1}}, a_{2l-1}]$  are pairwise disjoint if  $J$  is odd, for  $l = 1, \dots, M+1$ .

- If  $J$  is even, then

$$\begin{aligned} \sum_{j=1}^{J-1} |\hat{h}(t_{k_j})|^p &\leq \sum_{l=1}^M |\hat{h}(t_{k_{2l}}) - \hat{h}(t_{k_{2l+2}})|^p + \sum_{l=1}^M |\hat{h}(t_{k_{2l-1}}) - \hat{h}(t_{k_{2l+1}})|^p \\ &\leq 2\nu_p^p(\hat{h}, I), \end{aligned}$$

hence  $\sum_{j=1}^{J-1} |h(t_{k_j})|^p \leq 2\nu_p^p(h, I)$ .

Since  $(s_J)$  is monotone and  $p \geq 1$ , and given the above inequalities, we have

$$\begin{aligned} \nu_p((f \circ g) \cdot h, I) &\leq \nu_p(f, g(I)) \cdot \left( 2\nu_p^p(h, I) + \sup_I |h|^p \right)^{1/p} + \nu_p(h, I) \cdot \sup_{g(I)} |f|, \\ \text{and } \left( 2\nu_p^p(h, I) + \sup_I |h|^p \right)^{1/p} &\leq 2^{1/p} \nu_p(h, I) + \sup_I |h|, \end{aligned}$$

hence:

$$\nu_p((f \circ g) \cdot h, I) \leq \nu_p(f, g(I)) \cdot \left( 2^{1/p} \nu_p(h, I) + \sup_I |h| \right) + \nu_p(h, I) \cdot \sup_{g(I)} |f|.$$

2) Inequality (2.1.2) follows directly from inequality (2.1.1).  $\square$

**Theorem 2.3.3** ( [13]).

Let  $f_1(x)$  and  $f_2(x)$  be two functions defined, continuous and differentiable on a compact set of  $[a, b]$ , such that  $f'_1(x) = f'_2(x)$ , (a.e.), then  $f_1(x) - f_2(x)$  is constant.

**Proof.**

This theorem is due to De la Valle de-Poussin, and for the proof, see [13].  $\square$

**Theorem 2.3.4** ( [9] ).

Let  $\Omega \subset \mathbb{R}^n$  be an open set, and let  $f : \Omega \rightarrow \mathbb{R}^n$  be a Lipschitz function, then  $f$  is differentiable almost everywhere on  $\Omega$ ; moreover, the following assertions are verified:

(i)  $f$  is differentiable almost everywhere on the set  $L(f)$  such that

$$\begin{aligned} L(f) &= \{x \in \Omega : \text{Lip } f(x) < \infty\}, \\ \text{where } \text{Lip } f(x) &= \limsup_{\substack{y \rightarrow x \\ y \in \Omega}} \frac{|f(x) - f(y)|}{|x - y|}. \end{aligned}$$

(ii) Every function  $f \in W^{1,p}(\Omega)$ ,  $p \in [1, +\infty]$ , is differentiable (a.e.).

**Proof.**

This theorem of Rademacher is a generalization of Lebesgue's theorem to the case

$$n = m = 1, \quad \Omega = ]a, b[$$

for functions of bounded variation, and for the proof, see [9]; assertion (i) is due to a theorem of Stepanov, and (ii) to a theorem of Calderon.  $\square$

**Proposition 2.3.5** ([9]).

Let  $f : [a, b] \rightarrow \mathbb{R}$  be a Lipschitz function, then

$$f(b) - f(a) = \int_a^b f'(t) dt,$$

moreover,  $f : [a, b] \rightarrow \mathbb{R}$  is absolutely continuous if and only if there exists a function  $g \in L^1[a, b]$  such that  $f(x) = f(a) + \int_a^x g(t) dt$ ,  $x \in [a, b]$ , and in this case the derivative  $\frac{df}{dx}$  exists for almost every  $x \in [a, b]$  such that  $\frac{df}{dx} = g \in L^1[a, b]$ .

**Theorem 2.3.6** ([2]).

Let  $1 \leq p < \infty$ , then the following assertions are verified:

(i) If  $f, g \in BV_p^1(\mathbb{R})$ , then  $f \circ g \in BV_p^1(\mathbb{R})$ , and

$$\|f \circ g\|_{BV_p^1(\mathbb{R})} \leq \|f\|_{BV_p^1(\mathbb{R})} \left(1 + 2^{1/p} \|g\|_{BV_p^1(\mathbb{R})}\right).$$

(ii) Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a Borel measurable function, then:

The operator  $T_f$  operates on  $BV_p^1(\mathbb{R})$  if and only if  $f \in BV_p^1(\mathbb{R})$ .

This Theorem 2.3.6 is fundamental because it solves the composition operator problem such that

$$S(BV_p^1(\mathbb{R})) = BV_p^1(\mathbb{R}) = \{\phi : \mathbb{R} \rightarrow \mathbb{R}; T_\phi(BV_p^1(\mathbb{R})) \subseteq BV_p^1(\mathbb{R})\}.$$

**Proof.**

(i) Since  $f, g \in BV_p^1(\mathbb{R})$ , then  $f, g$  are continuous and Lipschitz, and by Rademacher's theorem, they are differentiable almost everywhere, and we thus have  $(f \circ g)'(x) = (f' \circ g) \cdot g'(x)$ . By De la Vallée Poussin's Theorem 2.3.3, the quantity  $(f \circ g)(x) - \int_0^x f'(g(t))g'(t) dt$  is constant, and applying Proposition 2.3.5, we obtain

$$(f \circ g)(x) = (f \circ g)(0) + \int_0^x f'(g(t))g'(t) dt.$$

Since  $f', g' \in \mathcal{V}_p(\mathbb{R})$ , then we can apply Theorem 2.3.2, and we conclude that  $(f' \circ g)g' \in \mathcal{V}_p(\mathbb{R})$ , such that

$$\|(f' \circ g)g'\|_{\mathcal{V}_p(\mathbb{R})} \leq 2^{1/p} \|f'\|_{\mathcal{V}_p(\mathbb{R})} \|g'\|_{\mathcal{V}_p(\mathbb{R})} < \infty.$$

We have

$$\begin{aligned} \|f \circ g\|_{BV_p^1(\mathbb{R})} &\approx |f \circ g(x_0)| + |(\widetilde{f \circ g})'|_{\mathcal{V}_p(\mathbb{R})} \\ &\lesssim |f \circ g(x_0)| + 2^{1/p} \|(\tilde{f})'\|_{\mathcal{V}_p(\mathbb{R})} \|(\tilde{g})'\|_{\mathcal{V}_p(\mathbb{R})}. \end{aligned}$$

Set  $g(x_0) = y_0$ , then we have

$$\begin{aligned} \|f \circ g\|_{BV_p^1(\mathbb{R})} &\lesssim |\tilde{f}(y_0)| + 2^{1/p} \|\tilde{f}'\|_{\mathcal{V}_p(\mathbb{R})} \|\tilde{g}'\|_{\mathcal{V}_p(\mathbb{R})} \\ &\lesssim |\tilde{f}(y_0)| + \|\tilde{f}'\|_{\mathcal{V}_p(\mathbb{R})} + 2^{1/p} \|\tilde{f}'\|_{\mathcal{V}_p(\mathbb{R})} \cdot (\|\tilde{g}'\|_{\mathcal{V}_p(\mathbb{R})} + |g(y_0)|) \\ &+ 2^{1/p} (|g(y_0)| + \|\tilde{g}'\|_{\mathcal{V}_p(\mathbb{R})}) |f(y_0)| \\ &\lesssim \left( |f(y_0)| + \|\tilde{f}'\|_{\mathcal{V}_p(\mathbb{R})} \right) \cdot \left[ 1 + 2^{1/p} (|g(y_0)| + \|\tilde{g}'\|_{\mathcal{V}_p(\mathbb{R})}) \right] \\ &\approx \|f\|_{BV_p^1(\mathbb{R})} \cdot \left( 1 + 2^{1/p} \|g\|_{BV_p^1(\mathbb{R})} \right), \end{aligned}$$

$$\text{hence } \|f \circ g\|_{BV_p^1(\mathbb{R})} \lesssim \|f\|_{BV_p^1(\mathbb{R})} \left( 1 + 2^{1/p} \|g\|_{BV_p^1(\mathbb{R})} \right).$$

(ii) – If  $f \in BV_p^1(\mathbb{R})$ , then according to (i),

$$\begin{aligned} \|f \circ g\|_{BV_p^1(\mathbb{R})} &\lesssim \|f\|_{BV_p^1(\mathbb{R})} \left( 1 + 2^{1/p} \|g\|_{BV_p^1(\mathbb{R})} \right) < \infty, \quad \text{for all } g \in BV_p^1(\mathbb{R}), \\ \text{hence } f \circ g &\in BV_p^1(\mathbb{R}), \quad \text{for all } g \in BV_p^1(\mathbb{R}), \end{aligned}$$

and thus  $f$  operates on  $BV_p^1(\mathbb{R})$ .

– If  $f$  operates on  $BV_p^1(\mathbb{R})$ , then  $T_f(BV_p^1(\mathbb{R})) \subseteq BV_p^1(\mathbb{R})$ .

We have

$$\text{id}_{\mathbb{R}}(x) = x = x_0 + \int_{x_0}^x dt = x_0 + \int_{x_0}^x 1 dt,$$

such: that

$$1 \in BV_p(\mathbb{R}).$$

Moreover:

$$\begin{aligned}
\|\text{id}_{\mathbb{R}}\|_{BV_p^1(\mathbb{R})} &\approx |f(x_0)| + \|(\text{id}_{\mathbb{R}})'\|_{BV_p(\mathbb{R})} \\
&\approx |x_0| + \|1\|_{BV_p(\mathbb{R})} \\
&\approx |x_0| + \nu_p(1) + \sup_{x \in \mathbb{R}} |1| \\
&= |x_0| + 0 + 1 < \infty,
\end{aligned}$$

hence  $\text{id}_{\mathbb{R}} \in BV_p^1(\mathbb{R})$ , and thus  $f = f \circ \text{id}_{\mathbb{R}} = T_f(\text{id}_{\mathbb{R}}) \in BV_p^1(\mathbb{R})$ .

□

## 2.4 Statement of results

In this paragraph, we present Theorem 2.4.2, which represents our contribution, namely a generalization of the basic inequality introduced in Theorem 2.3.2 ([2]) by Lemma 2.4.1, giving the derivative of  $n$  functions, followed by Example 2.4.3 to affirm the obtained results, and we agree to take

$$\begin{aligned}
\mathop{\circ}\limits_{i=n}^m g_i &= \begin{cases} g_n \circ g_{n+1} \circ \cdots \circ g_k \circ \cdots \circ g_m, & \text{if } m \geq n \\ \text{id}, & \text{if } m < n \end{cases} \\
\text{and} \\
\prod_{i=n}^m A_i &= \begin{cases} A_n \times A_{n+1} \times \cdots \times A_m, & \text{if } m \geq n \\ 1, & \text{if } m < n \end{cases}
\end{aligned}$$

### Lemma 2.4.1.

(Basic algorithm) Let  $(I_k)_{2 \leq k \leq n}$  be a sequence of intervals of  $\mathbb{R}$ , and let  $(g_k)_{1 \leq k \leq n}$  be a sequence of differentiable functions such that  $g_k : I_k \rightarrow I_{k-1}$ , then for all  $n \geq 2$ ,

$$\left[ \mathop{\circ}\limits_{i=1}^n g_i \right]' = g'_n \times \prod_{i=1}^{n-1} \left[ g'_i \circ \left( \mathop{\circ}\limits_{j=i+1}^n g_j \right) \right] = \prod_{i=1}^n \left[ g'_i \circ \left( \mathop{\circ}\limits_{j=i+1}^n g_j \right) \right].$$

### Proof.

For  $n = 3$ , we have

$$\begin{aligned}(g_1 \circ g_2 \circ g_3)' &= (g_1 \circ (g_2 \circ g_3))' = (g_2 \circ g_3)' \times (g_1' \circ (g_2 \circ g_3)) \\ &= g_3' \times (g_2' \circ g_3) \times (g_1' \circ (g_2 \circ g_3)) = g_3' \times (g_1' \circ g_2 \circ g_3) \times (g_2' \circ g_3).\end{aligned}$$

By induction, for  $n$  terms,

$$(g_1 \circ g_2 \circ \cdots \circ g_n)' = g_n' \times (g_1' \circ g_2 \circ \cdots \circ g_n) \times \cdots \times (g_k' \circ g_{k+1} \circ \cdots \circ g_n) \times \cdots \times g_{n-1}' \circ g_n.$$

We deduce that for  $n+1$  terms, taking as the  $n$ -th term  $(g_n \circ g_{n+1})$ ,

$$\begin{aligned}(g_1 \circ \cdots \circ (g_n \circ g_{n+1}))' &= (g_n \circ g_{n+1})' \times (g_1' \circ \cdots \circ (g_n \circ g_{n+1})) \times \cdots \times g_{n-1}' \circ (g_n \circ g_{n+1}) \\ &= g_{n+1}' \times (g_n' \circ g_{n+1}) \times (g_1' \circ \cdots \circ g_n \circ g_{n+1}) \times \cdots \times g_{n-1}' \circ (g_n \circ g_{n+1}) \\ &= g_{n+1}' \times \left[ g_n' \circ g_{n+1} \times \prod_{i=1}^{n-1} g_i' \circ \left( \underset{j=i+1}{\overset{j=n+1}{\circ}} g_j \right) \right] \\ &= g_{n+1}' \times \left[ \prod_{i=1}^n g_i' \circ \left( \underset{j=i+1}{\overset{j=n+1}{\circ}} g_j \right) \right].\end{aligned}$$

□

### Theorem 2.4.2.

Let  $p \in [1, +\infty[$ ,  $n \geq 2$ ,  $h \in \mathcal{V}_p(I)$ , such that for all  $t \in I$ ,  $t_0 \in I$ ,  $\alpha \in \mathbb{R}$ ,

$$\underset{i=1}{\overset{i=n}{\circ}} g_i(t) = \alpha + \int_{t_0}^t h(x) \, dx \quad \text{and} \quad f \in \mathcal{V}_p \left( \underset{i=1}{\overset{i=n}{\circ}} g_i(I) \right), \text{ then}$$

$$(i) \quad \left\| \left[ \underset{i=1}{\overset{i=n}{\circ}} g_i \right]' \right\|_{\mathcal{V}_p(I)} = \left\| g_n' \times \prod_{i=1}^{n-1} \left[ g_i' \circ \left( \underset{j=i+1}{\overset{j=n}{\circ}} g_j \right) \right] \right\|_{\mathcal{V}_p(I)} \leq 2^{\frac{n-1}{p}} \|g_n'\|_{\mathcal{V}_p(I)} \times \prod_{k=1}^{n-1} \|g_k'\|_{\mathcal{V}_p \left( \underset{\ell=k+1}{\overset{\ell=n}{\circ}} g_\ell(I) \right)}.$$

$$(ii) \quad \left\| f \circ \left( \underset{i=1}{\overset{i=n}{\circ}} g_i \right) \times h \right\|_{\mathcal{V}_p(I)} \leq 2^{\frac{n}{p}} \|f\|_{\mathcal{V}_p \left( \underset{i=1}{\overset{i=n}{\circ}} g_i(I) \right)} \times \|g_n'\|_{\mathcal{V}_p(I)} \times \prod_{k=1}^{n-1} \|g_k'\|_{\mathcal{V}_p \left( \underset{\ell=k+1}{\overset{\ell=n}{\circ}} g_\ell(I) \right)}.$$

**Proof.**

(i) According to Lemma 2.4.1,

$$\left\| \left[ \begin{smallmatrix} i=n \\ i=1 \\ \circ \\ i=n \end{smallmatrix} g_i \right]' \right\|_{\mathcal{V}_p(I)} = \left\| \left[ \prod_{i=1}^{n-1} g_i' \circ \left( \begin{smallmatrix} j=n \\ j=i+1 \\ \circ \\ j=n \end{smallmatrix} g_j \right) \right] \times g_n' \right\|_{\mathcal{V}_p(I)},$$

hence

$$\begin{aligned} \left\| \left[ \begin{smallmatrix} i=n \\ i=1 \\ \circ \\ i=n \end{smallmatrix} g_i \right]' \right\|_{\mathcal{V}_p(I)} &= \left\| \left[ \left( \prod_{i=1}^{n-1} g_i' \circ \left( \begin{smallmatrix} j=n-1 \\ j=i+1 \\ \circ \\ j=n-1 \end{smallmatrix} g_j \right) \right) \circ g_n \right] \times g_n' \right\|_{\mathcal{V}_p(I)} \\ &\leq 2^{1/p} \left\| \prod_{i=1}^{n-1} g_i' \circ \left( \begin{smallmatrix} j=n-1 \\ j=i+1 \\ \circ \\ j=n-1 \end{smallmatrix} g_j \right) \right\|_{\mathcal{V}_p(g_n(I))} \times \|g_n'\|_{\mathcal{V}_p(I)}, \quad \text{see (2.1.2).} \end{aligned}$$

And

$$\begin{aligned} \left\| \prod_{i=1}^{n-1} g_i' \circ \left( \begin{smallmatrix} j=n-1 \\ j=i+1 \\ \circ \\ j=n-1 \end{smallmatrix} g_j \right) \right\|_{\mathcal{V}_p(g_n(I))} &= \left\| \left[ \prod_{i=1}^{n-2} g_i' \circ \left( \begin{smallmatrix} j=n-1 \\ j=i+1 \\ \circ \\ j=n-1 \end{smallmatrix} g_j \right) \right] \times g_{n-1}' \circ \left( \begin{smallmatrix} j=n-1 \\ j=n \\ \circ \\ j=n-1 \end{smallmatrix} g_j \right) \right\|_{\mathcal{V}_p(g_n(I))} \\ &= \left\| \left[ \left( \prod_{i=1}^{n-2} g_i' \circ \left( \begin{smallmatrix} j=n-2 \\ j=i+1 \\ \circ \\ j=n-2 \end{smallmatrix} g_j \right) \right) \circ g_{n-1} \right] \times g_{n-1}' \right\|_{\mathcal{V}_p(g_n(I))}, \quad \text{since } \begin{smallmatrix} j=n-1 \\ j=n \\ \circ \\ j=n-1 \end{smallmatrix} g_j = \text{id} \\ &\leq 2^{1/p} \left\| \prod_{i=1}^{n-2} g_i' \circ \left( \begin{smallmatrix} j=n-2 \\ j=i+1 \\ \circ \\ j=n-2 \end{smallmatrix} g_j \right) \right\|_{\mathcal{V}_p(g_{n-1}(g_n(I)))} \times \|g_{n-1}'\|_{\mathcal{V}_p(g_n(I))}. \end{aligned}$$

Step by step, we obtain

$$\begin{aligned} \left\| \left[ \begin{smallmatrix} i=n \\ i=1 \\ \circ \\ i=n \end{smallmatrix} g_i \right]' \right\|_{\mathcal{V}_p(I)} &= \left\| g_n' \times \prod_{i=1}^{n-1} \left[ g_i' \circ \left( \begin{smallmatrix} j=n \\ j=i+1 \\ \circ \\ j=n \end{smallmatrix} g_j \right) \right] \right\|_{\mathcal{V}_p(I)} \\ &\leq 2^{\frac{n-1}{p}} \|g_n'\|_{\mathcal{V}_p(I)} \times \prod_{k=1}^{n-1} \|g_k'\|_{\mathcal{V}_p\left(\begin{smallmatrix} n \\ \circ \\ \ell=k+1 \\ g_\ell(I) \end{smallmatrix}\right)}. \end{aligned}$$

On the other hand, by induction, we have

$$\begin{aligned} \left\| \left[ \begin{smallmatrix} i=n+1 \\ i=1 \\ \circ \\ i=n+1 \end{smallmatrix} g_i \right]' \right\|_{\mathcal{V}_p(I)} &= \left\| \left[ \left( \begin{smallmatrix} i=n \\ i=1 \\ \circ \\ i=n \end{smallmatrix} g_i \right) \circ g_{n+1} \right]' \right\|_{\mathcal{V}_p(I)} = \left\| \left[ \left( \begin{smallmatrix} i=n \\ i=1 \\ \circ \\ i=n \end{smallmatrix} g_i \right)' \circ g_{n+1} \right] \times g_{n+1}' \right\|_{\mathcal{V}_p(I)} \\ &\leq 2^{1/p} \left\| \left( \begin{smallmatrix} i=n \\ i=1 \\ \circ \\ i=n \end{smallmatrix} g_i \right)' \right\|_{\mathcal{V}_p(g_{n+1}(I))} \times \|g_{n+1}'\|_{\mathcal{V}_p(I)}, \quad \text{according to (2.1) of Theorem 2.3.2} \\ &\leq 2^{\frac{n-1}{p}} 2^{1/p} \|g_n'\|_{\mathcal{V}_p(g_{n+1}(I))} \times \prod_{k=1}^{n-1} \|g_k'\|_{\mathcal{V}_p\left(\begin{smallmatrix} n \\ \circ \\ \ell=k+1 \\ g_\ell(g_{n+1}(I)) \end{smallmatrix}\right)} \times \|g_{n+1}'\|_{\mathcal{V}_p(I)} \end{aligned}$$

$$\begin{aligned}
&= 2^{\frac{n}{p}} \|g'_{n+1}\|_{\mathcal{V}_p(I)} \times \left[ \prod_{k=1}^{n-1} \|g'_k\|_{\mathcal{V}_p\left(\ell=\frac{n}{k+1} g_{\ell}(g_{n+1}(I))\right)} \times \|g'_n\|_{\mathcal{V}_p(g_{n+1}(I))} \right] \\
&= 2^{\frac{n}{p}} \|g'_{n+1}\|_{\mathcal{V}_p(I)} \times \prod_{k=1}^n \|g'_k\|_{\mathcal{V}_p\left(\ell=\frac{n+1}{k+1} g_{\ell}(I)\right)}.
\end{aligned}$$

(ii) Since  $h(t) = \left( \bigcirc_{i=1}^{i=n} g_i(t) \right)'$ , and according to (2.1.2), we have

$$\left\| \left[ f \circ \left( \bigcirc_{i=1}^{i=n} g_i \right) \right] \times h \right\|_{\mathcal{V}_p(I)} \leq 2^{1/p} \|f\|_{\mathcal{V}_p\left(\bigcirc_{i=1}^{i=n} g_i(I)\right)} \times \left\| \left( \bigcirc_{i=1}^{i=n} g_i \right)' \right\|_{\mathcal{V}_p(I)},$$

$$\begin{aligned}
&\leq \left[ 2^{1/p} \|f\|_{\mathcal{V}_p\left(\bigcirc_{i=1}^{i=n} g_i(I)\right)} \right] \times \left[ 2^{\frac{n-1}{p}} \|g'_n\|_{\mathcal{V}_p(I)} \times \prod_{k=1}^{n-1} \|g'_k\|_{\mathcal{V}_p\left(\ell=\frac{n}{k+1} g_{\ell}(I)\right)} \right] \\
&= 2^{\frac{n}{p}} \|f\|_{\mathcal{V}_p\left(\bigcirc_{i=1}^{i=n} g_i(I)\right)} \times \|g'_n\|_{\mathcal{V}_p(I)} \times \prod_{k=1}^{n-1} \|g'_k\|_{\mathcal{V}_p\left(\ell=\frac{n}{k+1} g_{\ell}(I)\right)}.
\end{aligned}$$

□

### Example 2.4.3.

- For  $n = 2$ , let  $g_1 \circ g_2 = h$ , then we have

$$\text{(i)} \quad \|(g_1 \circ g_2)'\|_{\mathcal{V}_p(I)} = \|g'_2 \times (g'_1 \circ g_2)\|_{\mathcal{V}_p(I)} = \|(g'_1 \circ g_2) \times g'_2\|_{\mathcal{V}_p(I)} \leq 2^{1/p} \|g'_1\|_{\mathcal{V}_p(g_2(I))} \|g'_2\|_{\mathcal{V}_p(I)}.$$

$$\text{(ii)} \quad \|f \circ (g_1 \circ g_2) \times h\|_{\mathcal{V}_p(I)} \leq 2^{1/p} \|f\|_{\mathcal{V}_p(g_1 \circ g_2(I))} \|h\|_{\mathcal{V}_p(I)} = 2^{1/p} \|f\|_{\mathcal{V}_p(g_1 \circ g_2(I))} \|(g_1 \circ g_2)'\|_{\mathcal{V}_p(I)}.$$

$$\begin{aligned}
&\leq \left[ 2^{1/p} \|f\|_{\mathcal{V}_p(g_1 \circ g_2(I))} \right] \cdot \left[ 2^{1/p} \|g'_1\|_{\mathcal{V}_p(g_2(I))} \|g'_2\|_{\mathcal{V}_p(I)} \right] \\
&\leq 2^{2/p} \|f\|_{\mathcal{V}_p(g_1 \circ g_2(I))} \|g'_1\|_{\mathcal{V}_p(g_2(I))} \|g'_2\|_{\mathcal{V}_p(I)},
\end{aligned}$$

and we have

$$2^{2/p} \|f\|_{\mathcal{V}_p\left(\bigcirc_{i=1}^{i=2} g_i(I)\right)} \|g'_2\|_{\mathcal{V}_p(I)} \prod_{k=1}^1 \|g'_k\|_{\mathcal{V}_p\left(\ell=\frac{2}{k+1} g_{\ell}(I)\right)} = 2^{2/p} \|f\|_{\mathcal{V}_p(g_1 \circ g_2(I))} \|g'_2\|_{\mathcal{V}_p(I)} \|g'_1\|_{\mathcal{V}_p(g_2(I))}$$

- For  $n = 3$ , let  $g_1 \circ g_2 \circ g_3 = h$ , then we have

(i)

$$\begin{aligned}
\|(g_1 \circ g_2 \circ g_3)'\|_{\mathcal{V}_p(I)} &= \|((g_1 \circ g_2) \circ g_3)'\|_{\mathcal{V}_p(I)} = \|g'_3 \times (g_1 \circ g_2)'\circ g_3\|_{\mathcal{V}_p(I)} \\
&= \|[g_1 \circ g_2]'\circ g_3] \times g'_3\|_{\mathcal{V}_p(I)} \\
&\leq 2^{1/p} \|(g_1 \circ g_2)'\|_{\mathcal{V}_p(g_3(I))} \|g'_3\|_{\mathcal{V}_p(I)} \\
&\leq 2^{1/p} \left[ 2^{1/p} \|g'_1\|_{\mathcal{V}_p(g_2(g_3(I)))} \|g'_2\|_{\mathcal{V}_p(g_3(I))} \right] \|g'_3\|_{\mathcal{V}_p(I)} \\
&= 2^{2/p} \|g'_1\|_{\mathcal{V}_p(g_2(g_3(I)))} \|g'_2\|_{\mathcal{V}_p(g_3(I))} \|g'_3\|_{\mathcal{V}_p(I)},
\end{aligned}$$

$$\text{and we have } 2^{\frac{3-1}{p}} \|g'_3\|_{\mathcal{V}_p(I)} \prod_{k=1}^{3-1} \|g'_k\|_{\mathcal{V}_p\left(\substack{3 \\ \circ \\ \ell=k+1} g_\ell(I)\right)} = 2^{2/p} \|g'_3\|_{\mathcal{V}_p(I)} \prod_{k=1}^2 \|g'_k\|_{\mathcal{V}_p\left(\substack{3 \\ \circ \\ \ell=k+1} g_\ell(I)\right)}$$

$$= 2^{2/p} \|g'_3\|_{\mathcal{V}_p(I)} \|g'_1\|_{\mathcal{V}_p\left(\substack{3 \\ \circ \\ \ell=2} g_\ell(I)\right)} \|g'_2\|_{\mathcal{V}_p\left(\substack{3 \\ \circ \\ \ell=3} g_\ell(I)\right)} = 2^{2/p} \|g'_3\|_{\mathcal{V}_p(I)} \|g'_1\|_{\mathcal{V}_p(g_2 \circ g_3(I))} \|g'_2\|_{\mathcal{V}_p(g_3(I))}.$$

$$\text{(ii)} \quad \|f \circ (g_1 \circ g_2 \circ g_3) \times h\|_{\mathcal{V}_p(I)} \leq 2^{1/p} \|f\|_{\mathcal{V}_p(g_1 \circ g_2 \circ g_3(I))} \|h\|_{\mathcal{V}_p(I)}$$

$$\begin{aligned}
&= 2^{1/p} \|f\|_{\mathcal{V}_p(g_1 \circ g_2 \circ g_3(I))} \|(g_1 \circ g_2 \circ g_3)'\|_{\mathcal{V}_p(I)} \\
&\leq \left[ 2^{1/p} \|f\|_{\mathcal{V}_p(g_1 \circ g_2 \circ g_3(I))} \right] \left[ 2^{2/p} \|g'_3\|_{\mathcal{V}_p(I)} \|g'_1\|_{\mathcal{V}_p(g_2 \circ g_3(I))} \|g'_2\|_{\mathcal{V}_p(g_3(I))} \right] \\
&\leq 2^{3/p} \|f\|_{\mathcal{V}_p(g_1 \circ g_2 \circ g_3(I))} \|g'_3\|_{\mathcal{V}_p(I)} \|g'_1\|_{\mathcal{V}_p(g_2 \circ g_3(I))} \|g'_2\|_{\mathcal{V}_p(g_3(I))}.
\end{aligned}$$

On the other hand, since  $\prod_{k=1}^2 \|g'_k\|_{\mathcal{V}_p\left(\substack{3 \\ \circ \\ \ell=k+1} g_\ell(I)\right)} = \|g'_1\|_{\mathcal{V}_p\left(\substack{3 \\ \circ \\ \ell=1+1} g_\ell(I)\right)} \|g'_2\|_{\mathcal{V}_p\left(\substack{3 \\ \circ \\ \ell=2+1} g_\ell(I)\right)}$ ,  
then

$$2^{3/p} \|f\|_{\mathcal{V}_p\left(\substack{i=3 \\ \circ \\ i=1} g_i(I)\right)} \|g'_3\|_{\mathcal{V}_p(I)} \prod_{k=1}^2 \|g'_k\|_{\mathcal{V}_p\left(\substack{3 \\ \circ \\ \ell=k+1} g_\ell(I)\right)} = 2^{3/p} \|f\|_{\mathcal{V}_p(g_1 \circ g_2 \circ g_3(I))} \|g'_3\|_{\mathcal{V}_p(I)} \|g'_1\|_{\mathcal{V}_p(g_2 \circ g_3(I))} \|g'_2\|_{\mathcal{V}_p(g_3(I))}.$$

## Chapter 3

# Composition of Operators in Homogeneous Besov Spaces $\dot{B}_p^{s,q}(\mathbb{R}^n)$

In this chapter, we first recall some basic notions, then define the norms of homogeneous Besov spaces and their realizations, as well as functional calculus by presenting two fundamental theorems (3.1.9) and (3.1.10). Finally, we give an extension of Peetre's Theorem (3.2.4) along with its proof by introducing certain functional spaces based on the works of [2], [3].

### Definition 3.0.1.

Let  $(X, \|\cdot\|_X)$ ,  $(Y, \|\cdot\|_Y)$  be two normed vector spaces. We denote by  $X \hookrightarrow Y$  the injection of  $X$  into  $Y$ , which is defined by the following conditions:

- (i)  $X$  is a vector subspace of  $Y$ .
- (ii) The identity map  $J : X \rightarrow Y$ , such that  $J(x) = x$  for all  $x \in X$ , is continuous.

Being linear, the continuity of the identity operator is equivalent to the existence of a constant  $C > 0$  such that

$$\|f\|_Y \leq C \|f\|_X , \quad \text{for all } f \in X, \quad (3.1)$$

- If (3.0.1) holds, then we say that  $X$  embeds into  $Y$ . We say that the space  $X$  is embedded in  $Y$  if there exists a continuous operator  $U : X \rightarrow Y$  such that

$$\|U(f)\|_Y \leq C \|f\|_X , \quad \text{for all } f \in X.$$

**Definition 3.0.2** ([14]).

Let  $A_0$  and  $A_1$  be two compatible quasi-Banach spaces embedded in a Hausdorff locally convex space. Set  $A = (A_0, A_1)$ , such that

$$\sum(A) = A_0 + A_1, \|a\|_{A_0+A_1} = \inf_{a=a_0+a_1} (\|a_0\|_{A_0} + \|a_1\|_{A_1})$$

and

$$\Delta(A) = A_0 \cap A_1, \|a\|_{A_0 \cap A_1} = \max_{a \in A_0 \cap A_1} (\|a\|_{A_0}, \|a\|_{A_1}),$$

and we define for each  $t > 0$ , the Peetre functionals

$$K(t, a) = \inf_{a=a_0+a_1} (\|a_0\|_{A_0} + t \|a_1\|_{A_1}), \text{ for } a \in \sum(A)$$

and

$$J(t, a) = \max (\|a\|_{A_0}, t \|a\|_{A_1}), \text{ for } a \in \Delta(A).$$

The function  $t \mapsto K(t, a)$  is positive, increasing, and concave. For each  $0 < \theta < 1$ ,  $0 < p \leq \infty$ , we define the interpolation space  $(A)_{\theta, p}$  of the spaces  $A_0$  and  $A_1$  by

$$(A)_{\theta, p} = (A_0, A_1)_{\theta, p} = \left\{ a \in \sum(A) : \|a\|_{(A)_{\theta, p}} < \infty \right\},$$

and we define its quasi-norm by

$$\|a\|_{(A)_{\theta, p}} = \left( \int_0^\infty (t^{-\theta} K(t, a))^p \frac{dt}{t} \right)^{\frac{1}{p}}, \text{ if } 1 \leq p < +\infty$$

and

$$\|a\|_{(A)_{\theta, \infty}} = \sup_{0 < t < \infty} t^{-\theta} K(t, a).$$

The quasi-norm of  $(A)_{\theta, p}$  can be replaced by the norm

$$\|a\|_{A_0+A_1} + \left( \int_0^\infty (t^{-\theta} K(t, a))^p \frac{dt}{t} \right)^{\frac{1}{p}}.$$

**Theorem 3.0.3** ([14]).

Let  $(A_0, A_1)$  and  $(B_0, B_1)$  be two compatible couples of Banach spaces, where the bounded linear operator

$T$  embeds  $A_i$  into  $B_i$  ( $i = 0, 1$ ), such that

$$\|Tf\|_{B_0} \leq M_0 \|f\|_{A_0}, \text{ and } \|Tf\|_{B_1} \leq M_1 \|f\|_{A_1},$$

then for all  $0 < \theta < 1$ ,  $0 < p \leq +\infty$ , the operator  $T$  embeds the interpolation space  $(A_0, A_1)_{\theta, p}$  into the interpolation space  $(B_0, B_1)_{\theta, p}$ , such that

$$\|Tf\|_{(B_0, B_1)_{\theta, p}} \leq M_\theta \|f\|_{(A_0, A_1)_{\theta, p}}, \text{ and } M_\theta \leq M_0^{1-\theta} M_1^\theta.$$

**Proof.**

From this main Theorem 3.0.3, all classical theorems such as Riesz-Thorin or Marcinkiewicz interpolation theorems, and classical inequalities such as Young's and Bernstein's inequalities follow. For the proof, see [12], as well as Peetre's monograph [14], Chap 1.  $\square$

### 3.1 Composition Problem in Homogeneous Besov Spaces $\dot{B}_p^{s, q}(\mathbb{R}^n)$

In this section, we study functional calculus in homogeneous Besov spaces and introduce the spaces  $\dot{B}_p^{s, q}(\mathbb{R}^n)$  defined via the set of distributions tending to 0 at infinity  $\widetilde{C}_0(\mathbb{R}^n)$ , which greatly facilitates functional calculus by allowing the truncation of polynomials.

#### 3.1.1 Properties of Homogeneous Besov Spaces

Let  $\psi$  be an infinitely differentiable, even, and positive function, with support a compact subset of  $\mathbb{R}^n \setminus \{0\}$ , and such that

$$\sum_{j \in \mathbb{Z}} \psi(2^j \xi) = 1, \text{ for all } \xi \neq 0. \quad (3.2)$$

The operator  $Q_j : \mathcal{S}'(\mathbb{R}^n) \rightarrow \mathcal{S}'(\mathbb{R}^n)$  is defined by the identity

$$\widehat{Q_j f}(\xi) = \psi(2^{-j} \xi) \widehat{f}(\xi), \quad (j \in \mathbb{Z}).$$

**Definition 3.1.1.**

Let  $s \in \mathbb{R}$ ,  $p, q \in [1, +\infty]$ . Then the homogeneous Besov space  $\dot{B}_p^{s, q}(\mathbb{R}^n)$  is the set of equivalence classes

of distributions  $f \in \mathcal{S}'(\mathbb{R}^n) / \mathcal{P}_\infty(\mathbb{R}^n)$  such that

$$\|f\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} = \left( \sum_{j \in \mathbb{Z}} \left( 2^{js} \|Q_j f\|_p \right)^q \right)^{1/q} < +\infty. \quad (3.3)$$

The norm (3.0.3) makes  $\dot{B}_p^{s,q}(\mathbb{R}^n)$  a homogeneous Banach space, satisfying homogeneity properties such that for all  $\lambda > 0$ , and for all  $a \in \mathbb{R}^n$ , we have:

$$\|\tau_a f\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} = \|f\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)}, \quad (3.4)$$

$$c_1 \|f\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} \leq \lambda^{(n/p)-s} \|f(\lambda(\cdot))\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} \leq c_2 \|f\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)}. \quad (3.5)$$

We can replace the norm  $\|-\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)}$  by an equivalent norm  $\|-\|'$ , which satisfies (3.1.1) and improves (3.1.2), by replacing the discrete partition (3.0.2) with a continuous partition such that

$$\lambda^{(n/p)-s} \|f(\lambda(\cdot))\|' = \|f\|', \quad \text{for all } \lambda > 0.$$

### Definition 3.1.2.

Let  $p, q \in [1, +\infty]$ . If  $s \in ]0, 1]$ , then we denote by  $\dot{B}_p^{s,q}(\mathbb{R}^n)$  the set of tempered distributions  $f$  in  $L_{loc}^p(\mathbb{R}^n)$  such that  $\|f\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} < +\infty$ , with

$$\begin{aligned} \|f\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} &\sim \left( \int_{\mathbb{R}^n} \frac{\omega^q(h)}{|h|^{sq}} \frac{dh}{|h|^n} \right)^{\frac{1}{q}}, \quad \text{if } s \neq 1 \\ &\sim \left( \int_{\mathbb{R}^n} \left( \frac{1}{|h|^s} \left( \int_{\mathbb{R}^n} |f(x+h) - f(x)|^p dx \right)^{1/p} \right)^q \frac{dh}{|h|^n} \right)^{1/q}, \\ \text{and } \|f\|_{\dot{B}_p^{1,q}(\mathbb{R}^n)} &\sim \left( \int_{\mathbb{R}^n} \left( \frac{1}{|h|} \left( \int_{\mathbb{R}^n} |f(x+h) + f(x-h) - 2f(x)|^p dx \right)^{\frac{1}{p}} \right)^q \frac{dh}{|h|^n} \right)^{\frac{1}{q}}, \end{aligned}$$

with the usual modifications for  $p = +\infty$  or  $q = +\infty$ ,

where  $\omega(h) = \|f(x+h) - f(x)\|_p$  denotes the modulus of continuity.

If  $s \in ]1, 2]$ , then we denote by  $\dot{B}_p^{s,q}(\mathbb{R}^n)$  the set of tempered distributions  $f$  in  $L_{loc}^p(\mathbb{R}^n)$  such that  $\partial_j f \in$

$\dot{B}_p^{s-1,q}(\mathbb{R}^n)$  for all  $j = 1, \dots, n$ , and we define its semi-norm by

$$\|f\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} \sim \sum_{j=1}^n \|\partial_j f\|_{\dot{B}_p^{s-1,q}(\mathbb{R}^n)}.$$

We distinguish between the homogeneous Besov space  $\dot{B}_p^{s,q}(\mathbb{R}^n)$  included in  $\mathcal{S}'(\mathbb{R}^n)/\mathcal{P}_\infty(\mathbb{R}^n)$  from Definition 3.1.1 and the space  $\dot{B}_p^{s,q}(\mathbb{R}^n)$  from Definition 3.1.2 included in  $L_{loc}^p(\mathbb{R}^n)$ .

**Lemma 3.1.3** ([2]).

Let  $f \in L_{loc}^p(\mathbb{R}^n)$ ,  $a \in \mathbb{R}^n$ ,  $\lambda > 0$ . Then the space  $\dot{B}_p^{s,q}(\mathbb{R}^n)$  satisfies the following homogeneity properties

$$\|f(\cdot - a)\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} = \lambda^{(n/p)-s} \|f(\lambda(\cdot))\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} = \|f\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)},$$

moreover,  $\|f\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} = 0$  if and only if  $f \in \mathcal{P}_{[s]}(\mathbb{R}^n)$ .

**Proof.**

– For  $s \in ]0, 1]$ , take  $x - a = y$ , so  $dx = dy$ ,

$$\text{hence } \|f(\cdot - a)\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} = \|f\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)}.$$

– For  $s \in ]1, 2]$ , we have

$$\|f(\cdot - a)\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} = \sum_{j=1}^n \|\partial_j f(\cdot - a)\|_{\dot{B}_p^{s-1,q}(\mathbb{R}^n)} = \sum_{j=1}^n \|\partial_j f\|_{\dot{B}_p^{s-1,q}(\mathbb{R}^n)} = \|f\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)},$$

and similarly by induction, for all  $s > 0$ .

– Set  $\lambda \cdot x = y$  and  $\lambda \cdot h = t$ , then

$$dy = \lambda^n \cdot dx, \quad dt = \lambda^n \cdot dh, \quad |t| = |\lambda|^n \cdot |h|,$$

applying Definition 3.1.2 we obtain

$$\begin{aligned} \|f(\lambda(\cdot))\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} &= \left( \int_{\mathbb{R}^n} \left( \frac{1}{|\lambda|^{-s} |t|^s} \left( \int_{\mathbb{R}^n} \lambda^{-n} |f(y+t) - f(y)|^p dy \right)^{1/p} \right)^q \frac{\lambda^{-n} dt}{|\lambda|^{-n} |t|^n} \right)^{1/q} \\ &= \lambda^{-(n/p)+s} \|f\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)}, \end{aligned}$$

- If  $\|f\|_{\dot{B}_p^{1,q}(\mathbb{R}^n)} = 0$ , then

$$f(x+h) + f(x-h) - 2f(x) = 0,$$

hence  $\omega_p^2(h, f) = 0$ , and so  $f$  is a polynomial of first degree.

- If  $\|f\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} = 0$ , then

$$\|\partial_j f\|_{\dot{B}_p^{s-1,q}(\mathbb{R}^n)} = 0, \quad (j = 1, \dots, n),$$

hence by induction  $\partial_j f$  is a polynomial of degree  $[s-1] = [s] - 1$ , and so  $f$  is a polynomial of degree  $[s]$ .

□

**Proposition 3.1.4** ([2]).

Let  $s \in \mathbb{R}$ ,  $p, q \in [1, +\infty]$ . Then the quotient space  $\dot{B}_p^{s,q}(\mathbb{R}^n) / \mathcal{P}_{[s]}(\mathbb{R}^n)$  can be identified with the homogeneous Besov space, which we can denote by  $\dot{\dot{B}}_p^{s,q}(\mathbb{R}^n)$ , and can be taken as the set of equivalence classes of tempered distributions modulo polynomials  $f \in \mathcal{S}'(\mathbb{R}^n) / \mathcal{P}_\infty(\mathbb{R}^n)$  such that  $\partial_j f \in \dot{B}_p^{s-1,q}(\mathbb{R}^n)$  for all  $j = 1, \dots, n$ . Moreover, the following expression is an equivalent norm for  $\dot{\dot{B}}_p^{s,q}(\mathbb{R}^n)$

$$\|f\|_{\dot{\dot{B}}_p^{s,q}(\mathbb{R}^n)} \approx \sum_{j=1}^n \|\partial_j f\|_{\dot{B}_p^{s-1,q}(\mathbb{R}^n)}.$$

**Proof.**

For the proof, see [4].

□

### 3.1.2 Realizations of Homogeneous Besov Spaces

**Definition 3.1.5.**

Let  $\sigma : \dot{B}_p^{s,q}(\mathbb{R}^n) \rightarrow \mathcal{S}'(\mathbb{R}^n) / \mathcal{P}_m(\mathbb{R}^n)$  be a continuous linear map such that  $\sigma(f) = [f]$  is the equivalence class of  $f$  modulo  $\mathcal{P}_m(\mathbb{R}^n)$ . Then for all  $f \in \dot{B}_p^{s,q}(\mathbb{R}^n)$ , we say that  $\sigma$  is a **realization** modulo  $\mathcal{P}_m(\mathbb{R}^n)$  of  $\dot{B}_p^{s,q}(\mathbb{R}^n)$ . It is a linear isomorphism from  $\dot{B}_p^{s,q}(\mathbb{R}^n)$  onto its image, such that the space  $\sigma(\dot{B}_p^{s,q}(\mathbb{R}^n))$  endowed with the norm

$$\|\sigma(f)\| = \|f\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)}$$

becomes a Banach space.

If  $f \in \mathcal{S}'(\mathbb{R}^n)/\mathcal{P}_\infty(\mathbb{R}^n)$  and if the series  $\sum_{j \in \mathbb{Z}} Q_j f$  converges in  $\mathcal{S}'(\mathbb{R}^n)/\mathcal{P}_m(\mathbb{R}^n)$ , then we have

$$\sigma_m(f) = \sum_{j \in \mathbb{Z}} Q_j f \in \mathcal{S}'(\mathbb{R}^n)/\mathcal{P}_m(\mathbb{R}^n).$$

**Definition 3.1.6.**

We say that a tempered distribution  $f \in \mathcal{S}'(\mathbb{R}^n)$  **tends to 0 at infinity** if we have

$$\lim_{\lambda \rightarrow 0} f\left(\frac{\cdot}{\lambda}\right) = 0, \text{ in } \mathcal{S}'(\mathbb{R}^n).$$

The set of such distributions is denoted by  $\tilde{C}_0(\mathbb{R}^n)$ .

If  $C(\mathbb{R})$  denotes the set of real-valued continuous functions and  $C_b(\mathbb{R})$  the Banach space of bounded functions in  $C(\mathbb{R})$  endowed with the sup norm, and  $C_0(\mathbb{R})$  the Banach subspace of functions in  $C_b(\mathbb{R})$  with limit zero at infinity, then the following distributions tend to 0 at infinity.

- Functions belonging to  $C_0(\mathbb{R}^n)$  or to  $L^p(\mathbb{R}^n)$ ,  $1 \leq p < +\infty$ .
- Bounded Borel measures.
- Derivatives of bounded continuous functions.
- Derivatives of distributions belonging to  $\tilde{C}_0(\mathbb{R}^n)$ .

**Proposition 3.1.7** ([4]).

Let  $1 \leq p < \infty$ , such that

$$(0 < s < 1 + \frac{1}{p} \quad \text{and} \quad 1 \leq q \leq +\infty) \quad \text{or} \quad (s = 1 + \frac{1}{p} \quad \text{and} \quad q = 1).$$

If we denote by  $\dot{\mathcal{B}}_p^{s,q}(\mathbb{R}^n)$  the set of tempered distributions  $f$  such that

$$[f] \in \dot{\mathcal{B}}_p^{s,q}(\mathbb{R}^n) \text{ and } \partial_j f \in \tilde{C}_0, \text{ for } j = 1, \dots, n,$$

then every element of  $\dot{\mathcal{B}}_p^{s,q}(\mathbb{R}^n)$  admits a representative in  $\dot{\mathcal{B}}_p^{s,q}(\mathbb{R}^n)$  unique up to addition of a constant, and the space  $\dot{\mathcal{B}}_p^{s,q}(\mathbb{R}^n)$  thus defined can be endowed with the semi-norm  $\|-\|_{\dot{\mathcal{B}}_p^{s,q}(\mathbb{R}^n)}$ .

### 3.1.3 Statement of Results

In this subsection, we compare the different spaces  $B_p^{s,q}$ ,  $\dot{B}_p^{s,q} \approx \dot{B}_p^{s,q}$ ,  $\dot{\mathcal{B}}_p^{s,q}$ , concerning functional calculus to solve the composition problem by specifying the conditions related to each of them and presenting two comparative theorems 3.1.9 and 3.1.10.

**Definition 3.1.8** ([4]).

Let  $p \in [1, +\infty[$  and  $J$  an interval of  $\mathbb{R}$ . Then we denote by  $\mathcal{U}_p(J)$  the set of measurable functions  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that

$$\sup_{|h| \leq t} |f(x+h) - f(x)| \text{ is measurable on } J, \text{ for all } t > 0,$$

endowed with the norm

$$\|f\|_{\mathcal{U}_p(J)}^p = \sup_{t > 0} t^{-1} \int_J \sup_{|h| \leq t} |f(x+h) - f(x)|^p \, dx < +\infty.$$

We say that a continuous function  $f$  belongs to  $U_p^1(\mathbb{R})$  if there exists a bounded Borel function  $h \in \mathcal{U}_p(\mathbb{R})$  such that

$$f(x) - f(0) = \int_0^x h(t) \, dt, \quad \text{for all } x \in \mathbb{R},$$

and for every bounded Borel measurable function  $h$  of class  $\mathcal{U}_p(\mathbb{R})$ , we endow  $U_p^1(\mathbb{R})$  with the semi-norm

$$\|f\|_{U_p^1} = \inf \left\{ \sup_{\mathbb{R}} |h| + \|h\|_{\mathcal{U}_p}, h = f' \text{ (p.p.)} \right\} \sim |f(0)| + \|f'\|_{\infty} + \|f'\|_{U_p(\mathbb{R})}.$$

And just as we defined  $BV_p(\mathbb{R})$  from  $\mathcal{V}_p(\mathbb{R})$  (cf. Definition 1.4.4), we also define  $U_p(\mathbb{R})$  from  $\mathcal{U}_p(\mathbb{R})$ , taking functions that are almost everywhere equal to at least one element of  $\mathcal{U}_p(\mathbb{R})$ , and we endow  $U_p(\mathbb{R})$  with the semi-norm

$$\|f\|_{U_p(\mathbb{R})} = \inf \left\{ \|g\|_{\mathcal{U}_p(\mathbb{R})}; g \in \mathcal{U}_p(\mathbb{R}), g = f \text{ (p.p.)} \right\}.$$

**Theorem 3.1.9** ([2]).

Let

$$p \in ]1, +\infty[, s \in ]0, 1 + \frac{1}{p}[, q \in [1, +\infty[, f \in U_p^1(\mathbb{R}).$$

(i) If  $f(0) = 0$ , then  $T_f \left( B_p^{1+(1/p),1}(\mathbb{R}^n) \right) \subseteq B_p^{1+(1/p),\infty}(\mathbb{R}^n)$ ,

and  $\|f \circ g\|_{B_p^{1+(1/p),\infty}(\mathbb{R}^n)} \leq c \|f\|_{U_p^1(\mathbb{R})} \|g\|_{B_p^{1+(1/p),1}(\mathbb{R}^n)}$ , for all  $g \in B_p^{1+(1/p),1}(\mathbb{R}^n)$ .

(ii) If  $f(0) = 0$ , then  $T_f(B_p^{s,q}(\mathbb{R}^n)) \subseteq B_p^{s,q}(\mathbb{R}^n)$ ,

and  $\|f \circ g\|_{B_p^{s,q}(\mathbb{R}^n)} \leq c \|f\|_{U_p^1(\mathbb{R})} \|g\|_{B_p^{s,q}(\mathbb{R}^n)}$ , for all  $g \in B_p^{s,q}(\mathbb{R}^n)$ .

(iii) If  $g \in \dot{B}_p^{1+\frac{1}{p},1}(\mathbb{R})$ , then  $f \circ g \in \dot{B}_p^{1+\frac{1}{p},\infty}(\mathbb{R})$ ,

and  $\|f \circ g\|_{\dot{B}_p^{1+\frac{1}{p},\infty}(\mathbb{R})} \leq c_p \|f\|_{U_p^1(\mathbb{R})} (\|g\|_{\dot{B}_p^{1+\frac{1}{p},1}(\mathbb{R})} + |L_{g'}|)$ , ( $c_p > 0$ ).

(iv) If  $g \in BV_p^1(\mathbb{R})$ , then  $f \circ g \in \dot{B}_p^{1+\frac{1}{p},\infty}(\mathbb{R})$ ,

and  $\|f \circ g\|_{\dot{B}_p^{1+\frac{1}{p},\infty}(\mathbb{R})} \leq c_p \|f\|_{U_p^1(\mathbb{R})} \|g'\|_{BV_p(\mathbb{R})}$ , ( $c_p > 0$ ).

### Proof.

For the proof, see [2]. Theorem 3.1.9 can be generalized to spaces  $B_p^{s,q}(\mathbb{R}^n)$ , but we do not know a generalization to the n-dimensional case for spaces  $BV_p^1$ .  $\square$

### Theorem 3.1.10 ([4]).

Let

$$1 \leq p < \infty, q \in [1, +\infty], 0 < s < 1 + \frac{1}{p} \text{ and } f \in U_p^1(\mathbb{R}).$$

(i) If  $g \in \dot{B}_p^{1+\frac{1}{p},1}(\mathbb{R}^n)$ , then we have

$$[f \circ g] \in \dot{B}_p^{1+\frac{1}{p},\infty}(\mathbb{R}^n), \partial_j(f \circ g) \in \widetilde{C}_0(\mathbb{R}^n), (j = 1, \dots, n),$$

$$\text{and } \|f \circ g\|_{\dot{B}_p^{1+\frac{1}{p},\infty}(\mathbb{R}^n)} \leq c \|f\|_{U_p^1} \|g\|_{\dot{B}_p^{1+\frac{1}{p},1}(\mathbb{R}^n)}, (c > 0).$$

(ii) If  $g \in \dot{B}_p^{s,q}(\mathbb{R}^n)$ , then we have

$$f \circ g \in \dot{B}_p^{s,q}(\mathbb{R}^n), \text{ and } \|f \circ g\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)} \leq c \|f\|_{U_p^1} \|g\|_{\dot{B}_p^{s,q}(\mathbb{R}^n)}, (c > 0).$$

### Proof.

For the proof, see [4], page 13. □

## 3.2 Peetre's Theorem

**Lemma 3.2.1** ([8]).

Let  $\phi : [a, b] \rightarrow \mathbb{R}$  be an absolutely continuous function.

Then:

$$\nu_1(\phi, [a, b]) = \int_a^b |\phi'(x)| dx, \text{ where } \nu_1(f, I) = \sup_{\{t_i\} \subset I} \sum_{k=1}^N |f(t_k) - f(t_{k-1})|.$$

**Theorem 3.2.2** ([14]).

Let  $1 \leq p_0, p_1 \leq \infty$ ,  $0 < q_0, q_1 \leq \infty$ ,  $0 < \theta < 1$ .

(i) If  $s = (1 - \theta)s_0 + \theta s_1$ ,  $\frac{1}{p} = \frac{1 - \theta}{p_0} + \frac{\theta}{p_1}$ ,  $\frac{1}{q} = \frac{1 - \theta}{q_0} + \frac{\theta}{q_1}$ , then

$$B_p^{s, \min(q, p)}(\mathbb{R}^n) \hookrightarrow (B_{p_0}^{s_0, q_0}(\mathbb{R}^n), B_{p_1}^{s_1, q_1}(\mathbb{R}^n))_{\theta, p} \hookrightarrow B_p^{s, \max(q, p)}(\mathbb{R}^n).$$

(ii) If  $s = (1 - \theta)s_0 + \theta s_1$ ,  $0 < \theta < 1$ ,  $r \in \mathbb{R}$ , then

$$(B_p^{s_0, q_0}(\mathbb{R}^n), B_p^{s_1, q_1}(\mathbb{R}^n))_{\theta, r} = B_p^{s, r}(\mathbb{R}^n).$$

(iii) If  $\frac{1}{q} = \frac{1 - \theta}{q_0} + \frac{\theta}{q_1}$ , then

$$(B_p^{s_0, q_0}(\mathbb{R}^n), B_p^{s_1, q_1}(\mathbb{R}^n))_{\theta, q} = B_p^{s, q}(\mathbb{R}^n).$$

### Proof.

For the proof, see Peetre's monograph, [14] Chap 5, P107. Theorem 3.2.2 remains true for homogeneous Besov spaces. □

**Proposition 3.2.3** ([2]).

(i) Let  $p \in ]1, +\infty[$ . If  $g$  is an element of  $\dot{B}_p^{\frac{1}{p}, 1}(\mathbb{R})$ ,

then  $g$  is a continuous function and the limit  $L_g = \lim_{x \rightarrow \infty} g(x)$  exists in  $\mathbb{R}$ .

In particular, each element of  $\dot{B}_p^{\frac{1}{p}, 1}(\mathbb{R})$  is congruent modulo  $\mathcal{P}_0$  to exactly one element of  $C_0(\mathbb{R})$ .

(ii) Let  $p \in [1, +\infty[$ . Then the space  $\dot{B}_p^{\frac{1}{p},1}(\mathbb{R}) \cap C_0(\mathbb{R})$ , endowed with the restriction of the semi-norm  $\|\cdot\|_{\dot{B}_p^{\frac{1}{p},1}(\mathbb{R})}$ , is a Banach space isometric to  $\dot{\tilde{B}}_p^{1/p,1}(\mathbb{R})$ .

**Theorem 3.2.4** ([2]).

Let  $p \in ]1, +\infty[$ . Then the following continuous inclusions hold:

$$\begin{aligned} \dot{\tilde{B}}_p^{1/p,1}(\mathbb{R}) &\hookrightarrow (L^\infty(\mathbb{R}), BV_1(\mathbb{R}))_{1/p,p} = (BV_\infty(\mathbb{R}), BV_1(\mathbb{R}))_{1/p,p} \\ &\hookrightarrow BV_p(\mathbb{R}) \hookrightarrow U_p(\mathbb{R}) \hookrightarrow \dot{\tilde{B}}_p^{1/p,\infty}(\mathbb{R}). \end{aligned}$$

**Proof.**

(1) Applying Theorem 3.2.2, (i) with

$$\theta = s = 1/p, p_0 = \infty, s_0 = 0, s_1 = q_0 = q_1 = p_1 = 1, \quad \text{then}$$

$$\dot{\tilde{B}}_p^{1/p,1}(\mathbb{R}) \hookrightarrow (\dot{\tilde{B}}_\infty^{0,1}(\mathbb{R}), \dot{\tilde{B}}_1^{1,1}(\mathbb{R}))_{1/p,p} \hookrightarrow \dot{\tilde{B}}_p^{1/p,p}(\mathbb{R}),$$

and according to Proposition 3.2.3, (ii) we have

$$\dot{B}_p^{1/p,1}(\mathbb{R}) \cap C_0(\mathbb{R}) \approx \dot{\tilde{B}}_p^{1/p,1}(\mathbb{R}) \quad \text{and} \quad \dot{B}_\infty^{0,1}(\mathbb{R}) \cap C_b(\mathbb{R}) \approx \dot{\tilde{B}}_\infty^{0,1}(\mathbb{R}),$$

hence  $\dot{B}_p^{1/p,1}(\mathbb{R}) \cap C_0(\mathbb{R}) \hookrightarrow \left( \dot{B}_\infty^{0,1}(\mathbb{R}) \cap C_b(\mathbb{R}), \dot{\tilde{B}}_1^{1,1}(\mathbb{R}) \cap C_0(\mathbb{R}) \right)_{1/p,p}.$

– Let us prove the injections

$$\dot{\tilde{B}}_1^{1,1}(\mathbb{R}) \hookrightarrow \dot{W}^{1,1}(\mathbb{R}) \hookrightarrow BV_1(\mathbb{R}).$$

– If  $f \in \dot{\tilde{B}}_1^{1,1}(\mathbb{R})$ , then using Bernstein's inequality we obtain

$$\|f'\|_1 \leq cR \|f\|_1 = cR \left\| \sum_{j \in \mathbb{Z}} Q_j f \right\|_1 \leq cR \sum_{j \in \mathbb{Z}} 2^j \|Q_j f\|_1 = C \|f\|_{\dot{\tilde{B}}_1^{1,1}},$$

where  $supp(\hat{f}) \subset \{\xi \in \mathbb{R}^n : |\xi| \leq R\},$

hence  $\|f\|_{\dot{W}^{1,1}(\mathbb{R})} \leq C \|f\|_{\dot{B}_1^{1,1}(\mathbb{R})}$ , and so  $\dot{B}_1^{1,1}(\mathbb{R}) \hookrightarrow \dot{W}^{1,1}(\mathbb{R})$ . – If  $f \in \dot{W}^{1,1}(\mathbb{R})$ , then  $f$  is absolutely continuous, and by Lemma 3.2.1 we have

$$\|f\|_{BV_1(\mathbb{R})} = \nu_1(\tilde{f}, (\mathbb{R})) + \sup_{x \in \mathbb{R}} |\tilde{f}(x)| \leq 2\nu_1(\tilde{f}, \mathbb{R}) \leq 2 \int_{\mathbb{R}} |\tilde{f}'(x)| dx \leq 2 \|f\|_{\dot{W}^{1,1}(\mathbb{R})},$$

hence  $\dot{W}^{1,1}(\mathbb{R}) \hookrightarrow BV_1(\mathbb{R})$ , and therefore  $\dot{B}_1^{1,1}(\mathbb{R}) \hookrightarrow BV_1(\mathbb{R})$ . – We also prove that  $\dot{B}_{\infty}^{0,1}(\mathbb{R}) \hookrightarrow L^{\infty}(\mathbb{R})$ , because we have

$$\|f\|_{\infty} = \left\| \sum_{j \in \mathbb{Z}} Q_j f \right\|_{\infty} \leq \sum_{j \in \mathbb{Z}} \|Q_j f\|_{\infty} = \|f\|_{\dot{B}_{\infty}^{0,1}(\mathbb{R})},$$

– applying the interpolation Theorem 3.0.3 we obtain

$$(\dot{B}_{\infty}^{0,1}(\mathbb{R}), \dot{B}_1^{1,1}(\mathbb{R}))_{1/p,p} \hookrightarrow (L^{\infty}(\mathbb{R}), BV_1(\mathbb{R}))_{1/p,p},$$

$$\text{hence } \dot{B}_p^{1/p,1}(\mathbb{R}) \hookrightarrow (L^{\infty}(\mathbb{R}), BV_1(\mathbb{R}))_{1/p,p}.$$

(2) If  $E$  is the closure of  $BV_1(\mathbb{R})$  with respect to  $L^{\infty}(\mathbb{R})$ , then

$$(L^{\infty}(\mathbb{R}), BV_1(\mathbb{R}))_{1/p,p} = (E, BV_1(\mathbb{R}))_{1/p,p},$$

and since  $E \hookrightarrow BV_{\infty}(\mathbb{R})$ , then

$$(E, BV_1(\mathbb{R}))_{1/p,p} \hookrightarrow (BV_{\infty}(\mathbb{R}), BV_1(\mathbb{R}))_{1/p,p},$$

$$\text{hence } (L^{\infty}(\mathbb{R}), BV_1(\mathbb{R}))_{1/p,p} \hookrightarrow (BV_{\infty}(\mathbb{R}), BV_1(\mathbb{R}))_{1/p,p}.$$

(3) We define the norms of the space  $l_N^p$  of finite sequences  $\{t_i\}_{1 \leq i \leq N} \subset \mathbb{R}$  by

$$\|t_i\|_{l_N^p} = \left( \sum_{i=1}^{i=N} |t_i|^p \right)^{\frac{1}{p}} \text{ and } \|t_i\|_{l_N^{\infty}} = \sup_{1 \leq i \leq N} |t_i|.$$

Consider  $t_0 < t_1 < \dots < t_N$ , and let the continuous function  $U$  defined by

$$U : BV_\infty(\mathbb{R}) \longrightarrow l_N^\infty(\mathbb{R})$$

$$f \mapsto U(f) = \left( \tilde{f}(t_k) - \tilde{f}(t_{k-1}) \right)_{1 \leq k \leq N},$$

we then have

$$\begin{aligned} \|U(f)\|_{l_N^\infty} &= \sup_{1 \leq i \leq N} \left| \tilde{f}(t_i) - \tilde{f}(t_{i-1}) \right| \leq \sup_{1 \leq i \leq N} \left( \left| \tilde{f}(t_i) \right| + \left| \tilde{f}(t_{i-1}) \right| \right) \\ &\leq 2 \sup \left| \tilde{f} \right| = 2 \|f\|_{BV_\infty(\mathbb{R})}, \end{aligned}$$

$$\text{and } \|U(f)\|_{l_N^1} = \sum_{k=1}^{k=N} \left| \tilde{f}(t_k) - \tilde{f}(t_{k-1}) \right| \leq \nu_1(\tilde{f}) \leq \|f\|_{BV_1(\mathbb{R})}.$$

We have  $(L^\infty(X), L^1(X))_{\frac{1}{p}, p} = L^p(X)$ , and taking as a particular case the spaces  $l_N^p$ , we obtain by the interpolation Theorem 3.0.3

$$\|U(f)\|_{l_N^p} \leq c_p \|f\|_{(BV_\infty(\mathbb{R}), BV_1(\mathbb{R}))_{\frac{1}{p}, p}},$$

and since  $\|f\|_{BV_p(\mathbb{R})} = \inf_a \left\{ a > 0 : \|U(\tilde{f})\|_{l_N^p} \leq a \right\}$ , then

$$\|f\|_{BV_p(\mathbb{R})} \leq c_p \|f\|_{(BV_\infty(\mathbb{R}), BV_1(\mathbb{R}))_{\frac{1}{p}, p}},$$

which gives  $(BV_\infty(\mathbb{R}), BV_1(\mathbb{R}))_{\frac{1}{p}, p} \hookrightarrow BV_p(\mathbb{R})$ .

(4)

$$\begin{aligned} \text{We have } \int_{\mathbb{R}} \sup_{|h| \leq t} |f(x+h) - f(x)|^p dx &= \sum_{m \in \mathbb{Z}} \int_{mt}^{(m+1)t} \sup_{|h| \leq t} |f(x+h) - f(x)|^p dx \\ &= \int_0^t \sum_{m \in \mathbb{Z}} \sup_{|h| \leq t} |f(y+mt+h) - f(y+mt)|^p dy. \end{aligned}$$

For each  $m \in \mathbb{Z}$ ,  $y \in \mathbb{R}$ , there exists  $h_m(y) \in [-t, t]$ , such that

$$|f(y + mt + h_m(y)) - f(y + mt)|^p \geq \sup_{|h| \leq t} |f(y + mt + h) - f(y + mt)|^p - \frac{\varepsilon}{4} 2^{-|m|}.$$

Since the family of intervals  $\{[y + kt, y + kt + h(y)]\}$  is disjoint, then

$$\begin{aligned} \int_0^t \sum_{m \in \mathbb{Z}} \sup_{|h| \leq t} |f(y + mt + h) - f(y + mt)|^p dy &\leq \int_0^t \sum_{m \in \mathbb{Z}} \left\{ |f(y + mt + h_m(y)) - f(y + mt)|^p + \frac{\varepsilon}{4} 2^{-|m|} \right\} dy \\ &\leq \int_0^t (2\nu_p^p(f) + \varepsilon) dy = t((2\nu_p^p(f) + \varepsilon)). \end{aligned}$$

Since  $\varepsilon > 0$  is arbitrary, we obtain the inequality  $\|f\|_{\mathcal{U}_p(\mathbb{R})} \leq 2^{1/p} \|f\|_{\mathcal{V}_p(\mathbb{R})}$ , which gives

$$\mathcal{V}_p(\mathbb{R}) \hookrightarrow \mathcal{U}_p(\mathbb{R}),$$

and we deduce that

$$BV_p(\mathbb{R}) \hookrightarrow U_p(\mathbb{R}).$$

(5) We have for all  $g = f$ , (a.e.)

$$\begin{aligned} \|f\|_{\dot{B}_p^{1/p,\infty}(\mathbb{R})} &= \sup_{|h| \geq 1} \left[ h^{-\frac{1}{p}} \|\tau_{-h} f - f\|_{L^p} \frac{1}{|h|} \right], \\ &\leq \left[ \sup_{t > 0} t^{-1} \int_{\mathbb{R}} \left( \sup_{|h| \leq t} |g(x + h) - g(x)|^p \right) dx \right]^{\frac{1}{p}} = \|g\|_{\mathcal{U}_p(\mathbb{R})}, \end{aligned}$$

$$\text{hence } \|f\|_{\dot{B}_p^{1/p,\infty}(\mathbb{R})} \leq \inf \left\{ \|g\|_{\mathcal{U}_p(\mathbb{R})} ; g \in \mathcal{U}_p(\mathbb{R}), g = f \text{ (p.p.)} \right\} = \|f\|_{U_p(\mathbb{R})},$$

$$\text{therefore } \|f\|_{\dot{B}_p^{1/p,\infty}(\mathbb{R})} \leq \|f\|_{U_p(\mathbb{R})},$$

$$\text{which gives } U_p(\mathbb{R}) \hookrightarrow \dot{B}_p^{1/p,\infty}(\mathbb{R}).$$

□

# Chapter 4

## Composition of Operators in $BV_p^\alpha(I)$

In this chapter, we give the definitions of the spaces  $\mathcal{V}_p^\alpha(I)$ ,  $BV_p^\alpha(I)$ , of functions defined on intervals  $I$  of  $\mathbb{R}$  with bounded  $p$ -variation of order  $\alpha$ , and the space of their equivalence classes with respect to almost everywhere equality. Then we study the particular cases  $\alpha = 1 - \frac{1}{p}$ , ( $p \geq 1$ ) which give the spaces  $W^{1,p}(I)$ . We also give a generalization of Peetre's Theorem 3.2.4 to the spaces  $BV_p^\alpha(I)$ ,  $0 \leq \alpha < 1$ ,  $1 < p < \infty$ . Finally, we present some examples of composition operators defined in certain functional spaces.

### 4.1 Basic Notions

In this section, we present some basic notions on the spaces  $\mathcal{V}_p^\alpha(I)$  of functions with bounded  $p$ -variation of order  $\alpha$  where  $p \geq 1$ ,  $\alpha \geq 0$ , as well as the space of their primitives  $BV_p^\alpha(I)$ , in order to generalize some results from previous chapters.

#### 4.1.1 The Spaces $\mathcal{V}_p^\alpha(I)$

##### Definition 4.1.1.

Let  $p \in [1, +\infty]$ . We define the norm of the space  $l_N^p$  of finite real sequences  $\{t_i\}_{1 \leq i \leq N}$  by

$$\left\{ \begin{array}{ll} \|t_i\|_{l_N^p} = \left( \sum_{i=1}^{i=N} |t_i|^p \right)^{1/p}, & \text{if } p \neq \infty \\ \|t_i\|_{l_N^\infty} = \sup_{1 \leq i \leq N} |t_i|, & \text{if } p = \infty \end{array} \right.$$

##### Definition 4.1.2.

Let  $p \in [1, +\infty]$ ,  $\alpha \geq 0$ . Then a function  $f : I \rightarrow \mathbb{R}$  is said to be of **bounded  $p$ -variation of order  $\alpha$** ,

or briefly of  $p$ - $v$ .  $b$  of order  $\alpha$ , if for all finite, strictly increasing real sequences  $t_0 < t_1 < \dots < t_N$  in  $I$ , there exists  $c > 0$  such that

$$\left\| \frac{f(t_k) - f(t_{k-1})}{(t_k - t_{k-1})^\alpha} \right\|_{l_N^p} \leq c < \infty \quad (4.1)$$

or equivalently, if the number  $\sup_{\{t_k\}_{1 \leq k \leq n, t_{k-1} < t_k} \subset I} \left[ \sum_{k=1}^N \left| \frac{f(t_k) - f(t_{k-1})}{(t_k - t_{k-1})^\alpha} \right|^p \right]$  is finite.

The set of these functions is denoted by  $\mathcal{V}_p^\alpha(I)$ , ( $\mathcal{V}_p^\alpha$  if  $I = \mathbb{R}$ ), and the minimum of such constants  $c$  is denoted by  $\nu_p^\alpha(f, I)$ , ( $\nu_p^\alpha$  if  $I = \mathbb{R}$ ):

$$\nu_p^\alpha(f, I) = \inf_c \left\{ c > 0 : \sum_{k=1}^N \left| \frac{f(t_k) - f(t_{k-1})}{(t_k - t_{k-1})^\alpha} \right|^p \leq c^p, \text{ for all } \{t_k\}_{1 \leq k \leq n, t_{k-1} < t_k} \subset I \right\}.$$

### Remark 4.1.3.

- We agree to take  $\mathcal{V}_p^0(I) = \mathcal{V}_p(I)$ .
- Condition (4.1.1) is equivalent to the fact that for every family of disjoint intervals  $[a_k, b_k]$ ,  $a_k \neq b_k$  of  $I$ , we have  $\sum_{k=1}^N \left| \frac{f(a_k) - f(b_k)}{(a_k - b_k)^\alpha} \right|^p \leq c < \infty$ , ( $c > 0$ ).
- $\mathcal{V}_1^\alpha(I)$  or simply  $\mathcal{V}^\alpha(I)$  is called the space of functions of bounded variation of order  $\alpha$  on  $I$ , and  $\mathcal{V}_\infty^\alpha(I)$  is a Banach space for the norm

$$\|f\|_{\mathcal{V}_\infty^\alpha(I)} = \nu_\infty^\alpha(f, I) = \sup_{x \in I \setminus \{0\}} \left| \frac{f(x)}{x^\alpha} \right|,$$

and we have  $\mathcal{V}_\infty^\alpha(I) = \text{Lip}_\alpha(I)$ , for  $0 \leq \alpha < 1$ .

### Proposition 4.1.4.

Let  $x, y \in I$ ,  $p \in [1, +\infty[$ , and  $0 \leq \alpha < 1$ . Then each element of  $\mathcal{V}_p^\alpha(I)$  is a Lipschitz function of order  $\alpha$ , and  $\mathcal{V}_p^\alpha(I)$  becomes a Banach space when endowed with the following norm:

$$\|f\|_{\mathcal{V}_p^\alpha(I)} = \sup_{x \in I} |f(x)| + \nu_p^\alpha(f, I). \quad (4.2)$$

### Proof.

If  $f \in \mathcal{V}_p^\alpha(I)$ , then  $f$  is  $\nu_p^\alpha$ -Lipschitz of order  $\alpha$  because

$$\left| \frac{f(x) - f(y)}{(x - y)^\alpha} \right| \leq \nu_p^\alpha(f, I) \quad \text{for all } x, y \in I.$$

and the norm (4.1.2) satisfies all conditions making  $\mathcal{V}_p^\alpha(I)$  a Banach space.  $\square$

**Proposition 4.1.5.**

The space  $\mathcal{V}_p^\alpha(I)$  is a Banach algebra for pointwise multiplication of functions,

$$\text{such that } \|f \cdot g\|_{\mathcal{V}_p^\alpha(I)} \leq \|f\|_{\mathcal{V}_p^\alpha(I)} \|g\|_{\mathcal{V}_p^\alpha(I)} \quad \text{for all } f, g \in \mathcal{V}_p^\alpha(I).$$

**Proof.**

Set  $x_0 < x_1 < \dots < x_n$  a finite sequence in  $I$  and  $f, g \in \mathcal{V}_p(I)$ . Since  $p \geq 1$  and by using Minkowski's inequality we obtain:

$$\begin{aligned} \left( \sum_{j=1}^N \left| \frac{fg(x_j) - fg(x_{j-1})}{(x_j - x_{j-1})^\alpha} \right|^p \right)^{\frac{1}{p}} &\leq \left( \sum_{j=1}^N \left| f(x_j) \left( \frac{g(x_j) - g(x_{j-1})}{(x_j - x_{j-1})^\alpha} \right) \right|^p \right)^{1/p} + \left( \sum_{j=1}^N \left| g(x_{j-1}) \left( \frac{f(x_j) - f(x_{j-1})}{(x_j - x_{j-1})^\alpha} \right) \right|^p \right)^{1/p} \\ &\leq \sup_I |f| \cdot \left( \sum_{j=1}^N \left| \frac{g(x_j) - g(x_{j-1})}{(x_j - x_{j-1})^\alpha} \right|^p \right)^{\frac{1}{p}} + \sup_I |g| \cdot \left( \sum_{j=1}^N \left| \frac{f(x_j) - f(x_{j-1})}{(x_j - x_{j-1})^\alpha} \right|^p \right)^{\frac{1}{p}} \\ &\leq \sup_I |f| \cdot \nu_p^\alpha(g, I) + \sup_I |g| \cdot \nu_p^\alpha(f, I), \end{aligned}$$

hence

$$\nu_p^\alpha(f \cdot g, I) = \sup_I \left( \sum_{j=1}^N \left| \frac{fg(x_j) - fg(x_{j-1})}{(x_j - x_{j-1})^\alpha} \right|^p \right)^{\frac{1}{p}} \leq \sup_I |f| \cdot \nu_p^\alpha(g, I) + \sup_I |g| \cdot \nu_p^\alpha(f, I).$$

We deduce that:

$$\begin{aligned}
\|f \cdot g\|_{\mathcal{V}_p^\alpha(I)} &= \nu_p^\alpha(f \cdot g, I) + \sup_I |f| \cdot \sup_I |g| \leq \sup_I |f| \cdot \nu_p^\alpha(g, I) + \sup_I |g| \cdot \nu_p^\alpha(f, I) + \sup_I |f| \cdot \sup_I |g| \\
&\leq \sup_I |f| \cdot \nu_p^\alpha(g, I) + \sup_I |g| \cdot \nu_p^\alpha(f, I) + \sup_I |f| + \sup_I |g| + \underbrace{[\nu_p^\alpha(g, I) \cdot \nu_p^\alpha(f, I)]}_{\geq 0} \\
&= \left( \sup_I |f| + \nu_p^\alpha(f, I) \right) \cdot \left( \sup_I |g| + \nu_p^\alpha(g, I) \right) = \|f\|_{\mathcal{V}_p^\alpha(I)} \cdot \|g\|_{\mathcal{V}_p^\alpha(I)},
\end{aligned}$$

□

#### 4.1.2 The Spaces $BV_p^\alpha(I)$

In this subsection, we generalize the space of classes  $BV_p(I)$  to the space of classes  $BV_p^\alpha(I)$  such that  $0 \leq \alpha < 1$ . Then we define the space of distributions  $BV_p^\alpha$  for  $\alpha \geq 1$  as their primitives.

##### Definition 4.1.6.

Let  $p \in [1, +\infty]$ ,  $0 \leq \alpha < 1$ .

Then we denote by  $\mathcal{BV}_p^\alpha(I)$  the set of functions  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that there exists a function  $g \in \mathcal{V}_p^\alpha(I)$  which coincides with  $f$  almost everywhere.

$$\mathcal{BV}_p^\alpha(I) = \{f : I \rightarrow \mathbb{R} ; \exists g \in \mathcal{V}_p^\alpha, \text{ such that } f = g \text{ (p.p.)}\},$$

and we set

$$\varepsilon_p^\alpha(f) = \inf \{ \nu_p^\alpha(g) ; g \in \mathcal{V}_p^\alpha, \text{ such that } g = f \text{ (p.p.)} \}.$$

We denote by  $BV_p^\alpha(I)$  the quotient set with respect to the equivalence relation "equality in  $\mathcal{BV}_p^\alpha(I)$  almost everywhere", such that

$$\dot{f} = \{g \in \mathcal{BV}_p^\alpha(I) ; g = f \text{ (p.p.)}\},$$

$$\text{and } BV_p^\alpha(I) = \{ \dot{f} ; f \in \mathcal{BV}_p^\alpha(I) \} = \mathcal{BV}_p^\alpha(I) / \text{e.p.p.}$$

If  $h \in BV_p^\alpha(I)$ , we denote by  $\varepsilon_p^\alpha(h)$  the number  $\varepsilon_p^\alpha(f)$  for any representative  $f$  of  $h$ .

We agree to take  $BV_p^0(I) = BV_p(I)$ .

**Proposition 4.1.7.**

Let  $p \in [1, +\infty]$ ,  $0 \leq \alpha < 1$ . If  $f \in BV_p^\alpha(\mathbb{R})$ , then  $f$  has a unique normal representative  $\tilde{f} \in \mathcal{V}_p^\alpha$ , and we have

$$\varepsilon_p^\alpha(f) = \nu_p^\alpha(\tilde{f}).$$

We will therefore consider  $BV_p^\alpha(\mathbb{R})$  as a Banach space of distributions endowed with the following norm

$$\begin{aligned} \|f\|_{BV_p^\alpha(\mathbb{R})} &= \varepsilon_p^\alpha(f) + \left\| \frac{f(x)}{x^\alpha} \right\|_\infty \\ &= \nu_p^\alpha(\tilde{f}) + \sup_{x \in \mathbb{R} \setminus \{0\}} \left| \frac{\tilde{f}(x)}{x^\alpha} \right|. \end{aligned}$$

Note that for  $p = \infty$ , we obtain

$$\|f\|_{BV_\infty^\alpha(I)} \sim \sup_{x \in I \setminus \{0\}} \left| \frac{\tilde{f}(x)}{x^\alpha} \right|.$$

**Definition 4.1.8.**

Let  $p \in [1, +\infty]$ ,  $\alpha \geq 1$ . Then we say that a function  $f : I \rightarrow \mathbb{R}$  belongs to  $BV_p^\alpha(I)$  if there exist  $c, x_0 \in \mathbb{R}$  and  $g \in BV_p^{\alpha-1}(I)$  such that

$$f(x) = c + \int_{x_0}^x g(t) dt \quad \text{for all } x \in I. \quad (4.3)$$

If (4.1.3) holds, then  $f$  is a Lipschitz continuous function, and we endow  $BV_p^\alpha(I)$  with the norm

$$\|f\|_{BV_p^\alpha(I)} = |f(x_0)| + \|f'\|_{BV_p^{\alpha-1}(I)},$$

for which  $BV_p^\alpha(I)$  becomes a Banach space. For each point  $x_0$  in  $I$ , we obtain an equivalent norm.

## 4.2 The Spaces $W^{1,p}(\Omega)$ , $\Omega$ an open subset of $\mathbb{R}^n$

In this section, we study the space  $W^{1,p}(\Omega)$ , where  $\Omega$  is an open subset of  $\mathbb{R}^n$ , which proves to be very interesting due to its regularity with respect to composition operators. See the works of H. Brezis [6].

### 4.2.1 Properties of the Spaces $W^{1,p}(\Omega)$

**Definition 4.2.1** ([15]).

\* Let  $\Omega$  be an open subset of  $\mathbb{R}^n$ , and  $1 \leq p \leq \infty$ . Then we define the space  $W^{1,p}(\Omega)$  by

$$W^{1,p}(\Omega) = \left\{ u \in L^p(\Omega), \frac{\partial u}{\partial x_i} \text{ (in the sense of distributions)} \in L^p(\Omega), i = 1, \dots, n \right\}.$$

\* We set  $H^1(\Omega) = W^{1,2}(\Omega)$ , which is endowed with the scalar product

$$\langle u, v \rangle_{H^1(\Omega)} = \langle u, v \rangle_{L^2(\Omega)} + \sum_{i=1}^n \left\langle \frac{\partial u}{\partial x_i}, \frac{\partial v}{\partial x_i} \right\rangle_{L^2(\Omega)},$$

and the associated norm is

$$\|u\|_{H^1(\Omega)} = \left( \|u\|_{L^2(\Omega)}^2 + \sum_{i=1}^n \left\| \frac{\partial u}{\partial x_i} \right\|_{L^2(\Omega)}^2 \right)^{\frac{1}{2}}.$$

\* The space  $W^{1,p}(\Omega)$  is endowed with the norm

$$\|u\|_{W^{1,p}(\Omega)} = \|u\|_{L^p(\Omega)} + \sum_{i=1}^n \left\| \frac{\partial u}{\partial x_i} \right\|_{L^p(\Omega)}.$$

\* If  $1 < p < +\infty$ , then

$$\|u\|_{W^{1,p}(\Omega)} \sim \left[ \|u\|_{L^p(\Omega)}^p + \sum_{i=1}^n \left\| \frac{\partial u}{\partial x_i} \right\|_{L^p(\Omega)}^p \right]^{1/p}.$$

\* If we denote  $g_i = \frac{\partial u}{\partial x_i}$ , then  $u \in W^{1,p}(\Omega)$  if and only if

$$\forall \varphi \in \mathcal{D}(\Omega), \forall i = 1, \dots, n ; \quad \int_{\Omega} u \frac{\partial \varphi}{\partial x_i} = - \int_{\Omega} g_i \varphi, .$$

**Definition 4.2.2** ([15]).

\* Let  $\Omega$  be an open subset of  $\mathbb{R}^n$ ,  $1 \leq p \leq \infty$ ,  $m \geq 2$ , then we define the Sobolev space  $W^{m,p}(\mathbb{R}^n)$ , by

$$W^{m,p}(\Omega) = \left\{ u \in W^{m-1,p}(\Omega) ; \quad \frac{\partial u}{\partial x_i} \in W^{m-1,p}(\Omega), \text{ for all } i = 1, \dots, N \right\}.$$

\* We set  $H^m(\Omega) = W^{m,2}(\Omega)$  which becomes a Hilbert space, if equipped with the inner product

$$\langle u, v \rangle_{H^m(\Omega)} = \sum_{0 \leq |\alpha| \leq m} \langle D^\alpha u, D^\alpha v \rangle_{L^2(\Omega)}.$$

\* The space  $W^{m,p}(\Omega)$  is equipped with the norm

$$\|u\|_{W^{m,p}} = \sum_{0 \leq |\alpha| \leq m} \|D^\alpha u\|_{L^p}.$$

\* If we denote  $g_\alpha = D^\alpha u = \frac{D^{|\alpha|}}{\partial x_1^{\alpha_1} \cdots \partial x_n^{\alpha_n}}$ , then for all  $u \in L^p(\Omega)$

$u \in W^{m,p}(\Omega)$ , if and only if  $\forall \alpha \in \mathbb{N}^n$ ,  $|\alpha| \leq m$ ,  $\exists g_\alpha \in L^p(\Omega)$ , such that

$$\int_{\Omega} u D^\alpha \varphi = (-1)^{|\alpha|} \int_{\Omega} g_\alpha \varphi, \quad \text{for all } \varphi \in \mathcal{D}(\Omega).$$

\* If  $W_0^{1,p}(\Omega)$  denotes the closure of  $\mathcal{D}(\mathbb{R}^n)$  in  $W^{1,p}(\Omega)$ , then we have the following inequality called Poincare's inequality

$$\forall u \in W_0^{1,p}(\Omega); \quad \int_{\Omega} |u|^p \leq c(\Omega)^p \int_{\Omega} \|\nabla u\|^p, \quad c(\Omega) > 0, \quad p \geq 1.$$

$$\text{Where } \nabla u = \left( \frac{\partial u}{\partial x_1}, \dots, \frac{\partial u}{\partial x_k}, \dots, \frac{\partial u}{\partial x_n} \right), \quad \frac{\partial u}{\partial x_k} \in L^p(\Omega), \quad k = 1, \dots, n.$$

**Proposition 4.2.3** ([8]).

Let  $I = [a, b]$  such that  $a, b \in \mathbb{R}$ , and consider a function  $f : I \rightarrow \mathbb{R}$ , and a partition  $\mathcal{P} \subset I$  such that

$$\mathcal{P} = \{x_0, x_1, x_2, \dots, x_n\}, \text{ where } a = x_0 < x_1 < x_2 < \dots < x_n = b,$$

if we define the sums  $S(f, \mathcal{P})$ ,  $s(f, \mathcal{P})$ ,  $V(f, \mathcal{P})$  by

$$\begin{aligned}
S(f, \mathcal{P}) &= \sum_{k=1}^n \sup \{f(x) ; x_{k-1} \leq x \leq x_k\} \cdot (x_k - x_{k-1}) \\
s(f, \mathcal{P}) &= \sum_{k=1}^n \inf \{f(x) ; x_{k-1} \leq x \leq x_k\} \cdot (x_k - x_{k-1}) \\
V(f, \mathcal{P}) &= \sum_{k=1}^n |f(x_k) - f(x_{k-1})| \text{ and } \nu(f, [a, b]) = \sup \{V(f, \mathcal{P}) ; \mathcal{P} \subset [a, b]\}, \\
\text{then } \int_a^b f(x) \, dx &= \inf \{S(f, \mathcal{P}) ; \mathcal{P} \subset [a, b]\} = \sup \{s(f, \mathcal{P}) ; \mathcal{P} \subset [a, b]\},
\end{aligned}$$

and if  $f$  is absolutely continuous then  $\nu(f, [a, b]) = \int_a^b |f'(x)| \, dx$

**Proposition 4.2.4** ([6]).

Let  $1 < p < \infty$ ,  $s \in \mathbb{N}$ ,  $0 < \sigma \leq s$ ,  $\Omega \subset \mathbb{R}^n$ , then we can find equivalent norms for the space  $W^{s,p}(\Omega)$  defined by

$$\|f\|_{W^{s,p}(\Omega)} = \|f\|_{L^p(\Omega)} + \sup_{h \in \mathbb{R}^n, h \neq 0} \frac{\|\Delta_h^s f\|_{L^p(\Omega)}}{|h|^\sigma}.$$

**Lemma 4.2.5** ([8]).

Let  $1 < p < \infty$ , then for every interval  $I$  of  $\mathbb{R}$ , we have

$$W^{1,p}(I) = \mathcal{V}_p^{1-\frac{1}{p}}(I).$$

**Proof.**

– If  $f \in \mathcal{V}_p^{1-\frac{1}{p}}([a, b])$ , then for any partition  $\mathcal{P} = \{a \leq x_{k-1} < x_k \leq \dots, b\}$ ,

$$\text{we have } \left( \sum_{k \geq 0} |x_k - x_{k-1}| \cdot \left( \frac{|f(x_k) - f(x_{k-1})|}{|x_k - x_{k-1}|} \right)^p \right)^{1/p} \leq c < \infty,$$

fix the difference  $|x_k - x_{k-1}| = h$ , and consider the function  $g_h$ , defined by  $g_h(x) = (|f(x+h) - f(x)|)^p$ , then according to Proposition 4.2.3 we have

$$\int_a^b g_h(x) \, dx = \inf_{\mathcal{P} \subset [a, b]} \left\{ \sum_{k=1}^n \sup_x \{g_h(x) ; x_{k-1} \leq x \leq x_k\} \cdot (x_k - x_{k-1}) \right\}, \text{ hence}$$

$$\int_a^b (|f(x+h) - f(x)|)^p \leq \sum_{k \geq 0} |x_k - x_{k-1}| (|f(x_k) - f(x_{k-1})|)^p,$$

dividing both sides by  $|x_k - x_{k-1}|^p$ , we obtain

$$\left[ \frac{\int_a^b (|f(x+h) - f(x)|)^p dx}{|h|^p} \right]^{\frac{1}{p}} \leq \left[ \sum_{k \geq 0} |x_k - x_{k-1}| \left( \frac{|f(x_k) - f(x_{k-1})|}{|x_k - x_{k-1}|} \right)^p \right]^{\frac{1}{p}} \leq \|f\|_{\mathcal{V}_p^{1-\frac{1}{p}}},$$

and thus  $\mathcal{V}_p^{1-\frac{1}{p}}(I) \subset W^{1,p}(I)$ .

- Consider a sequence of disjoint intervals  $[a_k, b_k]$  of  $I$ , we can always choose a subsequence of intervals such that  $[a'_k, b'_k] \subset [a_k, b_k]$ ,  $|b'_k - a'_k| = |h| \neq 0$ .

If  $f \in W^{1,p}(I)$ , then according to Proposition 4.2.4

$$\begin{aligned} \frac{\left[ \sum_{k \geq 0} (|f(b_k) - f(a_k)|)^p \right]^{1/p}}{|b_k - a_k|^{1-\frac{1}{p}}} &= \frac{\left( \sum_{k \geq 0} (|f(a_k + h) - f(a_k)|)^p \right)^{1/p}}{|h|^{1-\frac{1}{p}}} \\ &\leq \frac{\left( \int_a^b (|f(x+h) - f(x)|)^p dx \right)^{1/p}}{|h|^{1-\frac{1}{p}}} = \frac{\|\Delta_h^1 f\|_{L^p}}{|h|^\sigma} \leq c < \infty, \end{aligned}$$

with  $0 < \sigma = 1 - \frac{1}{p} < 1$  because  $p > 1$ , hence  $W^{1,p}(I) \subset \mathcal{V}_p^{1-\frac{1}{p}}(I)$ .

□

## 4.2.2 Composition $W^{1,p}(\Omega)$

**Corollary 4.2.6** ([15]).

Let  $G \in C^1(\mathbb{R})$ ,  $G(0) = 0$ ,  $1 \leq p < \infty$ ,  $u \in W^{1,p}(I)$ , then

$$G \circ u \in W^{1,p}(I) \quad \text{and} \quad (G \circ u)' = (G' \circ u) u'.$$

**Proof.**

By the mean value theorem we have

$$\frac{|G(s) - G(0)|}{|s - 0|} \leq 1_{\text{supp } G}(s) \cdot \|G'\|_\infty.$$

If  $G(s) = 0$ , the result is immediate; otherwise  $1_{\text{supp } G}(s) = 1$ ,

and since  $G \in C^1(\mathbb{R})$  then  $\|G'\|_\infty \leq c < \infty$ , and taking  $M = \|u\|_{L^\infty}$ , so for every  $M > 0$ , we have

$$\forall s \in [-M, +M], \quad |G(s)| \leq c|s| \leq c\|u\|_{L^\infty},$$

but  $\|u\|_{L^\infty} \leq c'\|u\|_{W^{1,p}}$ ,  $c' > 0$ , hence

$$|G \circ u| \leq c\|u\|_{L^\infty} \leq c''\|u\|_{W^{1,p}} < \infty,$$

and since  $u \in L^p(I)$ ,  $u'^p(I)$ , then

$$G \circ u \in L^p(I), \quad (G \circ u') \in u'^p(I),$$

and since  $1 \leq p < \infty$ , then there exists a sequence  $u_n$  in  $\mathcal{D}(\mathbb{R})$  such that  $u_n \rightarrow u$  in  $W^{1,p}(I)$  and in  $L^\infty(I)$ , and thus

$$G \circ u_n \rightarrow G \circ u \in L^\infty(I)$$

$$\text{and } (G' \circ u_n)u'_n \rightarrow (G' \circ u)u'^p(I),$$

$$\text{but we have } \forall \varphi \in \mathcal{D}(I), \quad \int_I (G \circ u_n)\varphi' = - \int_I (G' \circ u_n)u'_n \varphi,$$

hence the result, because it suffices to take  $\varphi(t) = e^{-t} \in \mathcal{D}(I)$ .  $\square$

**Lemma 4.2.7** ([15]).

Let  $\Omega$  be an open subset of  $\mathbb{R}^n$ ,  $1 \leq p < \infty$ ,  $G \in C^1(\mathbb{R})$ , such that

$$G(0) = 0, \quad \text{and} \quad |G'(s)| \leq M, \quad (M > 0) \text{ for all } s \in \mathbb{R},$$

we have if  $u \in W^{1,p}(\Omega)$ , then  $G \circ u \in W^{1,p}(\Omega)$  and  $\frac{\partial}{\partial x_i}(G \circ u) = (G' \circ u) \frac{\partial u}{\partial x_i}$ .

**Proof.**

The same steps as in the proof of Corollary 4.2.6 give us Lemma 4.2.7.  $\square$

**Proposition 4.2.8** ([6]).

Let  $\Omega$  be an open subset of  $\mathbb{R}^n$ ,  $s > 1$ ,  $1 < p < \infty$ ,  $sp = n$ ,  $k = [s] + 1$ , such that

$$G \in C^k(\mathbb{R}), \quad G(0) = 0 \quad \text{and} \quad D^j G \in L^\infty(\mathbb{R}), \quad \text{for all } j \leq k,$$

we have, if  $u \in W^{s,p}(\Omega)$  then  $G \circ u \in W^{s,p}(\Omega)$ .

This Proposition 4.2.8 is a generalization of Lemma 4.2.7,

but it is not verified for  $p = 1$ , see [6].

**Theorem 4.2.9** ([5]).

Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a Borel measurable function such that  $f(0) = 0$ .

- (i) If  $1 \leq p \leq \infty$ , then  $T_f$  operates on  $W^{1,p}(\mathbb{R})$  if and only if  $f$  is a continuous and locally Lipschitz function.
- (ii) If  $1 \leq p \leq \infty$  and  $m \geq 2$ , then

$$T_f \text{ operates on } W^{m,p}(\mathbb{R}) \text{ if and only if } f \in W_{loc}^{m,p}(\mathbb{R}).$$

It follows that if the composition operator  $T_f$  operates on a Sobolev space then it must be bounded. This Theorem 4.2.9 is due to Marcus and Mizel, [5].

### 4.3 Peetre's theorem $BV_p^\alpha(I)$

**Theorem 4.3.1.**

Let  $p \in ]1, +\infty[$ ,  $0 \leq \alpha < 1$ , then we have the following continuous injections which are verified

$$\begin{aligned} \dot{B}_p^{1/p,1}(\mathbb{R}) &\hookrightarrow (L^\infty(\mathbb{R}), BV_1^\alpha(\mathbb{R}))_{1/p,p} = (BV_\infty^\alpha(\mathbb{R}), BV_1^\alpha(\mathbb{R}))_{1/p,p} \\ &\hookrightarrow BV_p^\alpha(\mathbb{R}) \hookrightarrow U_p(\mathbb{R}) \hookrightarrow \dot{B}_p^{1/p,\infty}(\mathbb{R}). \end{aligned}$$

**Proof.**

1) As we saw in the proof of Theorem 3.2.4 we have

$$\dot{B}_p^{1/p,1}(\mathbb{R}) \cap C_0(\mathbb{R}) \hookrightarrow \left( \dot{B}_\infty^{0,1}(\mathbb{R}) \cap C_b(\mathbb{R}), \dot{B}_1^{1,1}(\mathbb{R}) \cap C_0(\mathbb{R}) \right)_{1/p,p}.$$

– Let us prove the injections  $\dot{B}_1^{1,1} \hookrightarrow \dot{W}^{1,1} \hookrightarrow BV_1^\alpha$ .

\* The injection  $\dot{B}_1^{1,1} \hookrightarrow \dot{W}^{1,1}$  is verified (Theorem 3.2.4).

\* Since  $\nu_1^\alpha(f, \mathbb{R}) = \sup_{\{t_0 < t_1 < \dots < t_k < \dots < t_n\} \subset \mathbb{R}, n \in \mathbb{N}} \left[ \sum_{k=1}^n \frac{|f(t_k) - f(t_{k-1})|}{|t_k - t_{k-1}|^\alpha} \right]$ , then

$$\begin{aligned} \|f\|_{BV_1^\alpha(\mathbb{R})} &= \nu_1^\alpha(\tilde{f}) + \sup_{x \in \mathbb{R} \setminus \{0\}} \left| \frac{\tilde{f}'(x)}{x^\alpha} \right| \leq 2\nu_1^\alpha(\tilde{f}), \text{ because } 0 \leq \alpha < 1 \\ &\leq 2 \int_{\mathbb{R} \setminus \{0\}} \left| \frac{\tilde{f}'(x)}{x^\alpha} \right| dx \sim 2 \|f\|_{\dot{W}^{1,1}(\mathbb{R})}, \end{aligned}$$

hence  $\dot{W}^{1,1} \hookrightarrow BV_1^\alpha$ , and thus  $\dot{B}_1^{1,1} \hookrightarrow BV_1^\alpha$ .

\* We also prove that  $\dot{B}_\infty^{0,1} \hookrightarrow L^\infty$  (Theorem 3.2.4).

– Then applying the interpolation Theorem 3.0.3 we obtain

$$\left( \dot{B}_\infty^{0,1}, \dot{B}_1^{1,1} \right)_{1/p,p} = \dot{B}_p^{1/p,1} \hookrightarrow (L^\infty, BV_1^\alpha)_{1/p,p}.$$

(2) If  $E_\alpha$  is the closure of  $BV_1^\alpha(\mathbb{R})$  with respect to  $L^\infty(\mathbb{R})$ , then

$$(L^\infty(\mathbb{R}), BV_1^\alpha(\mathbb{R}))_{1/p,p} = (E_\alpha, BV_1^\alpha(\mathbb{R}))_{1/p,p}$$

and since  $E_\alpha \hookrightarrow BV_\infty^\alpha(\mathbb{R})$ , then

$$(E_\alpha, BV_1^\alpha(\mathbb{R}))_{1/p,p} \hookrightarrow (BV_\infty^\alpha(\mathbb{R}), BV_1^\alpha(\mathbb{R}))_{1/p,p},$$

hence  $(L^\infty(\mathbb{R}), BV_1^\alpha(\mathbb{R}))_{1/p,p} \hookrightarrow (BV_\infty^\alpha(\mathbb{R}), BV_1^\alpha(\mathbb{R}))_{1/p,p}$ .

(3) We define the norms of the space  $l_N^p$  of sequences  $\{t_i\}_{1 \leq i \leq N}$  by

$$\|t_i\|_{l_N^p} = \left( \sum_{i=1}^N |t_i|^p \right)^{1/p} \text{ and } \|t_i\|_{l_N^\infty} = \sup_{1 \leq i \leq N} |t_i|.$$

Fix a finite strictly increasing real sequence  $\{t_i\}_{0 \leq i \leq N}$  such that  $t_0 < t_1 < \dots < t_N$ , and associate for every  $0 \leq \alpha < 1$ , a function  $U_\alpha$ , defined by

$$U_\alpha : BV_\infty^\alpha(\mathbb{R}) \longrightarrow l_N^\infty$$

$$f \mapsto \left( \frac{\tilde{f}(t_i) - \tilde{f}(t_{i-1})}{|t_i - t_{i-1}|^\alpha} \right)_{1 \leq i \leq N},$$

$$\begin{aligned} \text{thus } \|U_\alpha(f)\|_{l_N^\infty} &= \sup_{t_k \in \{t_i\}} \left| \frac{\tilde{f}(t_k) - \tilde{f}(t_{k-1})}{|t_k - t_{k-1}|^\alpha} \right| \leq \sup_{t_k \in \{t_i\}} \left( \left| \frac{\tilde{f}(t_k)}{|t_k - t_{k-1}|^\alpha} \right| + \left| \frac{\tilde{f}(t_{k-1})}{|t_k - t_{k-1}|^\alpha} \right| \right) \\ &\leq 2 \sup_{t \in \mathbb{R} \setminus \{0\}} \left| \frac{\tilde{f}(t)}{t^\alpha} \right| = 2 \|f\|_{BV_\infty^\alpha(\mathbb{R})}, \end{aligned}$$

$$\text{hence } \|U_\alpha(f)\|_{l_N^\infty} \leq 2 \|f\|_{BV_\infty^\alpha(\mathbb{R})},$$

$$\text{and thus } BV_\infty^\alpha(\mathbb{R}) \hookrightarrow l_N^\infty.$$

On the other hand we have

$$\|U_\alpha(f)\|_{l_N^1} = \sum_{k=1}^N \left| \frac{\tilde{f}(t_k) - \tilde{f}(t_{k-1})}{|t_k - t_{k-1}|^\alpha} \right| \leq \nu_1^\alpha(\tilde{f}) + \sup |\tilde{f}| = \|f\|_{BV_1^\alpha(\mathbb{R})},$$

$$\text{hence } \|U_\alpha(f)\|_{l_N^1} \leq \|f\|_{BV_1^\alpha(\mathbb{R})},$$

$$\text{and thus } BV_1^\alpha(\mathbb{R}) \hookrightarrow l_N^1.$$

We have  $(L^\infty(X), L^1(X))_{\frac{1}{p}, p} = L^p(X)$ , and by the interpolation Theorem 3.0.3 we obtain

$$\|U_\alpha(f)\|_{l_N^p} \leq c_{p,\alpha} \|f\|_{(BV_\infty^\alpha(\mathbb{R}), BV_1^\alpha(\mathbb{R}))_{\frac{1}{p}, p}}, \quad c_{p,\alpha} > 0.$$

Since  $\|f\|_{BV_p^\alpha(\mathbb{R})} = \inf_a \left\{ a > 0 : \|U_\alpha(f)\|_{l_N^p} \leq a \right\}$ , then

$$\|f\|_{BV_p^\alpha(\mathbb{R})} \leq c_{p,\alpha} \|f\|_{(BV_\infty^\alpha(\mathbb{R}), BV_1^\alpha(\mathbb{R}))_{\frac{1}{p}, p}}, \quad c_{p,\alpha} > 0,$$

which gives  $(BV_\infty^\alpha(\mathbb{R}), BV_1^\alpha(\mathbb{R}))_{\frac{1}{p}, p} \hookrightarrow BV_p^\alpha(\mathbb{R})$ .

(4) According to the proof of Theorem 3.2.4, and since  $0 \leq \alpha < 1$ , then we have

$$\|f\|_{\mathcal{U}_p(\mathbb{R})} \leq 2^{1/p} \|f\|_{\mathcal{V}_p(\mathbb{R})} \leq 2^{1/p} \|f\|_{\mathcal{V}_p^\alpha(\mathbb{R})},$$

$$\text{hence } \mathcal{V}_p^\alpha(\mathbb{R}) \hookrightarrow \mathcal{V}_p(\mathbb{R}) \hookrightarrow \mathcal{U}_p(\mathbb{R}),$$

and we deduce that  $BV_p^\alpha(\mathbb{R}) \hookrightarrow BV_p(\mathbb{R}) \hookrightarrow U_p(\mathbb{R})$ ,

$$\text{which gives } BV_p^\alpha \hookrightarrow U_p.$$

(5) The continuous inclusion  $U_p \hookrightarrow \dot{B}_p^{1/p, \infty}$  was proved in the proof of Theorem 3.2.4.

□

## 4.4 Examples

The following examples are taken from the works of: [2, 3, 11, 16, 5, 7, 10, 1, 4].

We denote by  $\mathcal{E}_p$  the closure of  $BV_p(\mathbb{R}) \cap C^\infty(\mathbb{R})$  in  $BV_p(\mathbb{R})$  and by  $\mathcal{E}_p^1$  the closure of  $BV_p^1(\mathbb{R}) \cap C^\infty(\mathbb{R})$  in  $BV_p^1(\mathbb{R})$ .

#### 4.4.1 Example 1 [11, 5, 16]

- [11] Let  $1 < p < \infty$ ,  $1 < s < 1 + 1/p$ ,  $1 \leq q \leq \infty$ , and let  $F$  be a real variable function, Lipschitz,  $F(0) = 0$ ,  $F \in \dot{B}_p^{1+(1/p),\infty}(\mathbb{R})$ , then we have

$$F(B_p^{s,q}(\mathbb{R}^n)) \subset B_p^{s,q}(\mathbb{R}^n)$$

and

$$\exists C(n, s, p, q) > 0 : \|F(f)\|_{B_p^{s,q}(\mathbb{R}^n)} \leq C \max\left(\|F'\|_\infty, \|F\|_{\dot{B}_p^{1+(1/p),\infty}}\right) \cdot \|f\|_{B_p^{s,q}(\mathbb{R}^n)}$$

- [11] Let the nonlinear operator  $F_\mu : f \mapsto |f|^\mu$ ,  $\mu > 0$ ,

– If  $\mu > 1$ ,  $1 \leq p \leq \infty$ ,  $1 \leq q \leq \infty$ ,  $0 < s < \mu + \frac{1}{p}$ , then there exists a constant

$$C(s, p, q, n, \mu) > 0,$$

such that:

- \*  $\forall f \in B_p^{s,q}(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$ ,  $\|F_\mu(f)\|_{B_p^{s,q}(\mathbb{R}^n)} \leq C \|f\|_{B_p^{s,q}(\mathbb{R}^n)} \|f\|_\infty^{\mu-1}$
- \*  $\forall g \in B_p^{\mu+(1/p),\infty}(\mathbb{R})$ ,  $\| |g|^\mu \|_{B_p^{\mu+(1/p),\infty}(\mathbb{R})} \leq C \|g\|_{B_p^{\mu+(1/p),\infty}(\mathbb{R})}^\mu$

– If  $\mu > 1$ ,  $1 < p < \infty$ ,  $m < n/p$ , then we have an equivalence between

- \*  $\{|f|^\mu : f \in W^{m,t}\} \subset W^{m,p}$ , and
- \*  $m < \mu + \frac{1}{p}$ ,  $\frac{n\mu}{m(\mu-1) + \frac{n}{p}} \leq t \leq p\mu$ ,

the result remains true if we replace  $F_\mu : f \mapsto |f|^\mu$ , by

$$\tilde{F}_\mu : f \mapsto f |f|^{\mu-1}, \quad \text{or by } \bar{F}_\mu : f \mapsto (\max(f, 0))^\mu.$$

- If  $m \in \mathbb{N}$ ,  $1 < p < \infty$ ,  $m < n/p$ ,  $k \in \mathbb{N}$ ,  $k \geq 2$ , then we have an equivalence between

$$\{f^k : f \in W^{m,t}\} \subset W^{m,p} \quad \text{and} \quad \frac{nk}{m(k-1) + \frac{n}{p}} \leq t \leq pk.$$

The operator  $f \mapsto f |f|^\mu$  plays an essential role in the study of the Cauchy initial value problem for nonlinear Schrödinger partial differential equations.

#### 4.4.2 Example 2 [2, 3]

- Let  $p \geq 1$ , and consider the absolute value operator  $T_{F_1} = T_{|\cdot|}$  such that

$$T_{F_1}(g) = F_1(g) = |g|,$$

then  $|\cdot|$  belongs to  $BV_p^1(\mathbb{R}) \setminus \mathcal{E}_{p,loc}^1$ , and it verifies

- $T_{|\cdot|}$  operates on  $BV_p^1(I)$ , such that

$$\| |g| \|_{BV_p^1(I)} \leq c \| g \|_{BV_p^1(I)}, \quad (c > 0)$$

- $T_{|\cdot|}$  sends  $B_p^{1+1/p,1}(\mathbb{R}^n)$  to  $B_p^{1+1/p,\infty}(\mathbb{R}^n)$  such that

$$\| |g| \|_{B_p^{1+1/p,\infty}(\mathbb{R}^n)} \leq c \| g \|_{B_p^{1+1/p,1}(\mathbb{R}^n)}, \quad (c > 0)$$

- The operator  $T_{|\cdot|}$  is not continuous from  $B_p^{1+1/p,\infty}(\mathbb{R}^n)$  to  $B_p^{1+1/p,\infty}(\mathbb{R}^n)$ .

- If  $0 < s < 1 + (1/p)$ ,  $1 \leq q \leq \infty$ , it operates on  $B_p^{s,q}(\mathbb{R}^n)$ , such that

$$\| |g| \|_{B_p^{s,q}(\mathbb{R}^n)} \leq c \| g \|_{B_p^{s,q}(\mathbb{R}^n)}, \quad (c > 0)$$

- If  $0 < s < \infty$ ,  $1 \leq q < \infty$ , it operates continuously on  $B_p^{s,q}(\mathbb{R}^n)$ .

- Consider the family of functions

$$u_\alpha(x) = |x + \alpha| - |\alpha|, \quad x, \alpha \in \mathbb{R},$$

then

$$\| u_\alpha(g) \|_{BV_p^1(\mathbb{R})} = \| |g + \alpha| - |\alpha| \|_{BV_p^1(\mathbb{R})} \leq c_p \| g \|_{BV_p^1(\mathbb{R})}.$$

#### 4.4.3 Example 3 [3]

- Let  $p \geq 1$ , and the operator  $T_\psi : g \mapsto \psi \circ g$ , such that

- $\rho \in \mathcal{D}(\mathbb{R})$ ,  $\text{supp } \rho \subseteq [-1/2, 1/2]$ ,  $\rho(t) = 1$  in  $[-1/e, 1/e]$ ,
- $\psi : \mathbb{R} \rightarrow \mathbb{R}$ , and  $\psi(t) = |t| \frac{\rho(t)}{\log|t|}$ , if  $t \neq 0$ ,  $\psi(0) = 0$ ,

then the following assertions are verified

a) The function  $\psi$  belongs to  $\mathcal{E}_p^1$ .

b)  $T_\psi$  operates on  $BV_p^1(I)$  such that  $\|\psi \circ g\|_{BV_p^1(I)} \leq c \|g\|_{BV_p^1(I)}$ ,  $(c > 0)$ .

c)  $T_\psi$  sends  $B_p^{1+1/p,1}(\mathbb{R}^n)$  to  $B_p^{1+1/p,\infty}(\mathbb{R}^n)$  such that

$$\|\psi \circ g\|_{B_p^{1+1/p,\infty}(\mathbb{R}^n)} \leq c \|g\|_{B_p^{1+1/p,1}(\mathbb{R}^n)}, \quad (c > 0).$$

d)  $T_\psi$  is continuous from  $B_p^{1+1/p,1}(\mathbb{R}^n)$  to  $B_p^{1+1/p,\infty}(\mathbb{R}^n)$ .

e) If  $0 < s < 1 + (1/p)$ ,  $1 \leq q \leq \infty$ , then  $T_\psi$  operates on  $B_p^{s,q}(\mathbb{R}^n)$ , such that

$$\|\psi \circ g\|_{B_p^{s,q}(\mathbb{R}^n)} \leq c \|g\|_{B_p^{s,q}(\mathbb{R}^n)}, \quad (c > 0).$$

f) If  $q \in [1, \infty[$ , then the operator in (e) is continuous. The above properties do not change if  $\log|t|$  is replaced by iterated logarithms like  $\log|\log|t||$  or  $\log|\log|\log|t|||$ .

#### 4.4.4 Example 4 [3]

- Since the spaces  $BV_p^1(\mathbb{R})$  decrease with respect to  $p$ , we look for functions that belong to  $BV_{p_0}^1(\mathbb{R})$  such that  $p > p_0$ , and considering thus the family of functions  $\psi_{\alpha,\beta} : \mathbb{R} \rightarrow \mathbb{R}$ , defined by

$$\psi_{\alpha,\beta}(t) = |t|^{\alpha+1} \rho(t) \sin(|t|^{-\beta}) \text{ if } t \neq 0, \text{ and } \psi_{\alpha,\beta}(0) = 0, \quad 0 < \beta < \alpha,$$

Then we obtain an equivalence between the following assertions (a), (b), (c) such that

a)  $\frac{1}{p} < \frac{\alpha}{\beta} - 1$

b) The function  $\psi_{\alpha,\beta}$  belongs to  $BV_p^1(\mathbb{R})$

c) The operator  $T_{\psi_{\alpha,\beta}}$  sends  $BV_p^1(\mathbb{R})$  to  $BV_p^1(\mathbb{R})$ ,

and if  $\frac{1}{p} < \frac{\alpha}{\beta} - 1$ , then the operator  $T_{\psi_{\alpha,\beta}}$  satisfies properties (c)-(f) of the third example if we replace  $\psi$  by  $\psi_{\alpha,\beta}$ .

#### 4.4.5 Example 5 [7, 10]

- For the problem of composition of operators we have

i) A classical theorem due to B.E.J. Dahlberg (cf [7]) for the Sobolev space  $W^{m,p}(\mathbb{R}^n)$  stating that

If  $1 + \frac{1}{p} < m < \frac{n}{p}$ ,  $p \neq 0$ , or  $1 < p < \infty$ ,  $2 \leq m < \frac{n}{p}$ , then

$$T_G(W^{m,p}(\mathbb{R}^n)) = \{G(f); f \in W^{m,p}(\mathbb{R}^n)\} \subset W^{m,p}(\mathbb{R}^n),$$

implies  $G(t) = c \cdot t$ ,  $c \in \mathbb{R}$ .

ii) A classical result due to S. Igari ([10]) for the Besov space  $B_p^{s,q}(\mathbb{R}^n)$ , which states that

If  $1 \leq p < \infty$ ,  $1 \leq q \leq \infty$ ,  $0 < s < 1/p$  then

$$T_G(B_p^{s,q}) \subset B_p^{s,q}, \quad \text{if and only if } G \text{ is Lipschitz and } G(0) = 0.$$

#### 4.4.6 Example 6 [2, 1]

- Operators related to continuous Lipschitz functions satisfy the norm inequality property on  $W^{1,p}(\mathbb{R}^n)$  but not on  $W^{2,p}(\mathbb{R}^n)$ , this restriction is due to the following result

**Proposition 4.4.1** ([6]).

Let  $1 < p \leq +\infty$ ,  $s > 1 + (1/p)$  and  $N$  a norm on  $\mathcal{D}(\mathbb{R}^n)$ .

If  $E$  is a normed space such that  $\mathcal{D}(\mathbb{R}^n) \subset E \subset W_{loc}^{1,1}(\mathbb{R}^n)$

$$\sup_{h \neq 0} |h|^{1-s} \left( \int_{\mathbb{R}^n} \left| \frac{\partial g}{\partial x_i}(x+h) - \frac{\partial g}{\partial x_i}(x) \right|^p dx \right)^{1/p} \leq A \|g\|_E, \quad (A > 0)$$

for all  $g \in E$ , and  $i = 1, \dots, n$ , and if there exists a continuously differentiable function  $f : \mathbb{R} \rightarrow \mathbb{R}$ , and a constant  $B > 0$ , such that  $T_f$  injects  $\mathcal{D}(\mathbb{R}^n)$  into  $E$ , and such that the following inequality is verified, then  $f$  must be an affine function.

$$\|f \circ g\|_E \leq B(N(g) + 1) \quad \text{for all } g \in \mathcal{D}(\mathbb{R}^n).$$

- Superposition operators related to affine functions trivially satisfy the norm inequality properties with respect to function composition for every normed functional space.

- For the Sobolev space  $W^{m,p}(\mathbb{R}^n)$ ,  $m \in \mathbb{N}$ ,  $1 \leq p \leq \infty$ , nontrivial superposition operators that satisfy the norm inequality property exist if and only if

$$(m = 0) \quad \text{or} \quad (m = 1) \quad \text{or} \quad (m = 2 \text{ and } p = 1).$$

- For every  $\mu > 1$ ,  $m < \mu + \frac{1}{p}$ ,  $f \in W^{m,p}(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$  we have

$$\|F_\mu(f)\|_{W^{m,p}(\mathbb{R}^n)} \leq c \|f\|_{W^{m,p}(\mathbb{R}^n)} \|f\|_\infty^{\mu-1}, \quad (c > 0).$$

The norm inequality property for the operator  $F_\mu = |\cdot|^\mu$  is not entirely verified on  $W^{m,p}(\mathbb{R}^n)$  but partially on  $W^{m,p}(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$ .

- [1] - Let  $p \geq 1$ ,  $m > \max(n/p, 1)$ ,  $m \in \mathbb{N}$ , then

the composition operator  $T_f$  operates on  $W^{m,p}(\mathbb{R}^n)$ , if and only if  $f \in W_{loc}^{m,p}(\mathbb{R}^n)$ .

#### 4.4.7 Example 7 [4]

- We obtain a solution for the **P.S.O.**, with an additional norm inequality property in the Sobolev spaces  $W^{s,p}(\mathbb{R}^n)$  by Theorem 4.4.2, by introducing the space of functions of bounded variation  $BV(\mathbb{R}^n)$  equipped with the following seminorm:

$$\|f\|_{BV(\mathbb{R}^n)} \sim \nu(f) = \sum_{j=1}^n \|\partial_j f\|_M \sim \sup_{h \in \mathbb{R}^n \setminus \{0\}} \frac{1}{|h|} \int_{\mathbb{R}^n} |f(x+h) - f(x)| \, dx < +\infty,$$

where  $\|g\|_M$  denotes the total variation of the measure  $g$ , and the space  $BH(\mathbb{R})$  of distributions whose derivative belongs to  $BV(\mathbb{R})$ , equipped with the seminorm

$$\|f\|_{BH} = \nu(f') + \|f'\|_\infty.$$

#### Theorem 4.4.2 ([4]).

Let  $1 \leq p \leq \infty$ ,  $0 < s < 1 + (1/p)$ , then every function  $f \in BH(\mathbb{R})$ ,  $f(0) = 0$ , operates on  $W^{s,p}(\mathbb{R}^n)$ , moreover there exists  $c(s, p, n) > 0$ , such that:

$$\text{For all } g \in W^{s,p}(\mathbb{R}^n), \quad \|f \circ g\|_{W^{s,p}(\mathbb{R}^n)} \leq c(s, p, n) \|f\|_{BH} \|g\|_{W^{s,p}(\mathbb{R}^n)}.$$

#### 4.4.8 Example 8 [1]

- Every composition operator such that  $T_f(g) = f \circ g$  and  $f : \mathbb{R} \rightarrow \mathbb{R}$ , defined on

$$B_{\infty}^{s,\infty}(\mathbb{R}^n) = \mathcal{C}^s(\mathbb{R}^n), s > 0$$

the Holder-Zygmund space, verifies

$$T_f(\mathcal{C}^s(\mathbb{R}^n)) \subset \mathcal{C}^s(\mathbb{R}^n)$$

if and only if we have the following conditions

- $f$  is continuous and locally Lipschitz, for  $0 < s < 1$ .
- $f$  belongs locally to  $\mathcal{C}^s(\mathbb{R}^n)$ , for  $s > 1$ .
- $f$  is continuous and locally Lipschitz and satisfies the condition

$$f(x+t) + f(x-t) - 2f(x) = O\left(\frac{t}{|\log t|}\right), \quad \text{with } t \rightarrow 0^+,$$

uniformly on each compact subset of  $\mathbb{R}$ , for  $s = 1$ .

## References

- [1] G. Bourdaud and M. Lanza de Cristoforis, *Functional calculus in Hölder-Zygmund spaces*, Trans. Amer. Math. Soc. **354** (2002), 4109–4129.
- [2] G. Bourdaud, M. Lanza de Cristoforis, and W. Sickel, *Superposition operators and functions of bounded  $p$ -variation*, Rev. Mat. Iberoamer (Feb., 2004), (to appear). Preprint 362. Institut de Mathématiques de Jussieu. Unité Mixte de Recherche 7586. Université Paris VI et Paris VII / CNRS.
- [3] G. Bourdaud, Massimo Lanza de Cristoforis, and W. Sickel, *Superposition operators and functions of bounded  $p$ -variation II*, Nonlinear Analysis Series A **Volume. 62** (2005), 483–518.
- [4] G. Bourdaud and Y. Meyer, *Le calcul fonctionnel sous-linéaire dans les espaces de Besov homogènes*, (1st November 2004).
- [5] G. Bourdaud, M. Moussai, and W. Sickel, *Towards Sharp Superposition Theorems in Besov and Lizorkin-Triebel Spaces*, (March 29, 2006).
- [6] H. Brezis and P. Mironescu, *Composition in fractional Sobolev spaces*, Discrete and Continuous Dynamical Systems **Number 2. Volume 7** (April 2001), 241–246, Website: <http://math.smsu.edu/journal>.
- [7] B. E. J. Dahlberg, *A note on Sobolev spaces*, Proc. Symp. Pure Math. **N. 35 Vol. 1** (1979), 183–185, MR 81h:46030.
- [8] A. Giroux, *Notes de cours. Mesure et intégration*, Département de Mathématiques et Statistique. Université de Montréal, Mai 2004.
- [9] J. Heinonen, *Lectures on Lipschitz Analysis*, Lectures at the 14th Jyväskylä Summer School in August 2004. Supported by NSF grant DMS 0353549 and DMS 0244421, April 28, 2005.
- [10] S. Igari, *Sur les fonctions qui opèrent sur l'espace  $\widehat{A^2}$* , Ann. Inst. Fourier. Grenoble **15-21** (1965), 525–533.

- [11] D. Kateb, *On the boundedness of the mapping  $f \mapsto |f|^\mu$ ,  $\mu > 1$ , on Besov Spaces*, Math.Nachr **248-249** (2003), 110–128.
- [12] A. Kufner, *Some difference inequalities with weights and interpolation*, L. E. Persson **Volume 1. Number 3** (1998), 437–444, Mathematical inequalities Applications.
- [13] B. Maurey and J. P. Tacchi, *Ludwing Scheffer et les extensions du théorème des accroissements finis*, Version of 27/10/01.
- [14] J. Peetre, *New Thoughts on Besov Spaces*, Mathematics Series. No.1, Mathematics Department. Duke University. Durham. N.C, 1976. MR [57:1108](#).
- [15] C. Portenier, *Cours d'analyse fonctionnelle*, Version of November 29, 2004.
- [16] W. Sickel, *Necessary conditions on composition operators acting on Sobolev spaces of fractional order*, Forum Math. 9 (1997), 267–302.
- [17] H. Triebel, *Theory of Function Spaces III*, Birkhäuser Verlag, P. O. Box : 133. CH-4010 Basel. Switzerland. e-ISBN : 3-7643-7643-7582-5, 2006.
- [18] D. C. Ullrich, *Besov spaces. A primer*, Department of Mathematics Oklahoma State University. Stillwater OK 74078.