

A Novel Multiple Imputation Approach For Parameter Estimation in Observation-Driven Time Series Models With Missing Data

Guilherme Pumi^{a,*}, Taiane Schaedler Prass^a and Douglas Krauthein Verдум^a

Abstract

Handling missing data in time series is a complex problem due to the presence of temporal dependence. General-purpose imputation methods, while widely used, often distort key statistical properties of the data, such as variance and dependence structure, leading to biased estimation and misleading inference. These issues become more pronounced in models that explicitly rely on capturing serial dependence, as standard imputation techniques fail to preserve the underlying dynamics. This paper proposes a novel multiple imputation method specifically designed for parameter estimation in observation-driven models (ODM). The approach takes advantage of the iterative nature of the systematic component in ODM to propagate the dependence structure through missing data, minimizing its impact on estimation. Unlike traditional imputation techniques, the proposed method accommodates continuous, discrete, and mixed-type data while preserving key distributional and dependence properties. We evaluate its performance through Monte Carlo simulations in the context of GARMA models, considering time series with up to 70% missing data. An application to the proportion of stocked energy stored in South Brazil further demonstrates its practical utility.

Keywords: time series analysis, regression models, partial maximum likelihood, non-gaussian time series.

MSC: 62M10, 62F12, 62E20, 62G20, 60G15.

1 Introduction

The literature on missing data is extensive, with numerous books covering a wide range of methods, primarily in the context of independent data (e.g., Little and Rubin, 2020; van Buren, 2012; Carpenter and Kenward, 2012). As noted in van Buren (2012), relatively little is known about imputation for time series. Unlike independent data, handling missing values in time series is significantly more challenging due to temporal dependence. There are two primary approaches to dealing with missing data in time series when inference and forecasting are of interest. The first, imputation, is the simplest and most widely used. It involves replacing missing values with plausible estimates and proceeding with analysis as if no data were missing. The second approach relies on estimators specifically designed to handle missing data without requiring a fully completed time series. When available, these estimators typically yield better results, but they are difficult to derive and often highly model-dependent.

This work focuses on imputation in the context where missing data mechanism is completely random. Various methods exist, with many implemented in software packages. For

*Corresponding author. This Version: January 7, 2026

^aMathematics and Statistics Institute and Graduate Program in Statistics - Universidade Federal do Rio Grande do Sul.

E-mails: guilherme.pumi@ufrgs.br (G. Pumi), taiane.prass@ufrgs.br (T.S. Prass), dkverdum@gmail.com (D.K. Verдум)

ORCIDs: 0000-0002-6256-3170 (Pumi); 0000-0003-3136-909X (Prass); 0009-0009-4136-6731 (Verдум).

instance, the R package `imputeTS` (Moritz and Bartz-Beielstein, 2017) provides 18 general-purpose imputation methods, including mean, median, and mode imputation, last observation carried forward, next observation carried backward, spline interpolation, and random imputation. These methods are broadly applicable, but they may fail in specific contexts. For example, in time series constrained to the unit interval $(0, 1)$, spline interpolation often generate imputed values outside the boundaries, making it unsuitable for GARMA models for double-bounded data.

General-purpose imputation methods have the advantage of being quick and simple, which adds to its appeal, but they are not without drawbacks. First, they tend to underestimate variance, introducing biases in parameters that depend on it, such as correlations. Mean or median substitution, for instance, replaces distinct missing values with a single fixed number, artificially reducing variance and introducing a point mass in an otherwise continuous distribution. Second, imputed values alter the dependence structure of the time series in complex and often unpredictable ways, which is problematic since many statistical models rely on accurately capturing this structure. Third, standard errors and p -values are distorted, as standard inferential procedures assume that all observed values are accurate, failing to account for the additional uncertainty introduced by imputation. As the proportion of missing values increases, these issues worsen, requiring more specialized techniques.

A promising strategy to mitigate these issues is multiple imputation, in which missing values are replaced by plausible random draws from a carefully chosen distribution. This process is repeated $m > 1$ times, generating m complete time series. Inference is performed on each time series separately, and results are combined using simple pooling rules to obtain a final estimate with an associated standard error. The advantage of multiple imputation is that it accounts for the additional uncertainty introduced by the imputation process, but it presents two major problems: how to choose the distribution from which the imputed values are to be drawn from, and how to do it smartly, aiming at minimizing the aforementioned shortcomings of imputation methods?

A simple but naive multiple imputation approach is to draw missing values uniformly from the observed range of the time series. However, this fails to address the problems outlined earlier. A more refined approach was proposed by Pumi et al. (2025), in which missing values are drawn from a truncated normal distribution centered at the previous observed value, with variance calibrated to match the sample variance of the observed data. This method aims to preserve the local dependence structure while maintaining variance consistency. However, the results in Pumi et al. (2025) showed only marginal improvements over standard imputation methods, which may be attributed to the specific application context - namely, long-range dependent processes, where the method's local nature may not be sufficient to enhance estimation.

In this work, we introduce a novel multiple imputation method designed specifically for parameter estimation in observation-driven models. Our approach takes advantage of the systematic component's iterative nature in these models to propagate the conditional dependence structure through missing values, reducing imputation-induced distortions in dependence. The method is applicable to continuous, discrete, or mixed random components and can be used with any estimator in observation-driven models.

Informally, given an initial parameter estimate, missing values are imputed by reconstructing the systematic component of the model and sampling from the assumed conditional distribution. This ensures that the dependence structure of the imputed series closely matches that of the observed data. After each imputation step, model parameters are reestimated using the newly completed dataset. This process is iterated several times, with estimates from

multiple imputed datasets combined to produce a final parameter estimate. The method is compatible with any estimator of any quantity of interest in observation-driven models, making it widely applicable.

We evaluate the proposed method through a series of Monte Carlo simulation studies in the context of GARMA models, considering time series with up to 70% missing data. Finally, we illustrate its practical application by analyzing the proportion of stocked energy stored in South Brazil.

2 Preliminaries

In this work we consider observation-driven models as discussed in Cox (1981). More specifically, let $\{Y_t\}_{t \in \mathbb{Z}}$ be a time series of interest and let $\{\mathbf{X}_t\}_{t \in \mathbb{Z}}$ denote a set of r -dimensional exogenous time dependent (possibly random) covariates. Let \mathcal{F}_t denote the information (σ -field) available to the observer at time t , that is, $\mathcal{F}_t := \sigma\{\mathbf{X}_{t+1}, Y_t, \mathbf{X}_t, Y_{t-1}, \dots\}$, where, by convention, \mathbf{X}_t denotes the observed values at time t for deterministic covariates, and at time $t-1$ for stochastic ones. The model's random component is specified assuming that Y_t given \mathcal{F}_{t-1} is distributed according to a distribution depending on an identifiable parameter vector $\boldsymbol{\nu} \in \mathbb{R}^v$ and a quantity of interest μ_t satisfying

$$g(\mu_t) = A(\mathbf{X}'_t, Y_{t-1}, \mathbf{X}'_{t-1}, Y_{t-2}, \dots; \boldsymbol{\lambda}), \quad (1)$$

for some twice continuously differentiable with respect to $\boldsymbol{\lambda}$, and integrable real function A , where $\boldsymbol{\lambda}$ is an identifiable finite dimensional parameter vector and g is a twice continuously differentiable link function. Observe that, specification (1) implies that $g(\mu_t)$ is \mathcal{F}_{t-1} -measurable. Let $\boldsymbol{\gamma} = (\boldsymbol{\nu}', \boldsymbol{\lambda}')'$ be the parameter of interest and denote by $f(\cdot; \boldsymbol{\gamma} | \mathcal{F}_t)$ be the conditional density, mass or density-mass hybrid (mixed type) related to Y_t given \mathcal{F}_t . Equation (1) determines the model's systematic component. The observation-driven model we are interested in this paper is given by the specification of the random component and the systematic component (1).

One of the main advantages of observation-driven models is that its dynamic is sequentially determined given the model's past knowledge. This allows for sequential conditional inference regarding model parameters without requiring full knowledge of the joint distribution, leading to the partial likelihood approach as discussed in Cox (1975) and Wong (1986) (see also Fokianos and Kedem, 2004).

The full generality presented in (1) is sufficient for developing the method proposed and studied in this paper, albeit of limited practical use without a complete specification of the function A and its interaction with parameter $\boldsymbol{\lambda}$. Also, the distributional parameter $\boldsymbol{\nu}$ in most cases is uni-dimensional. Model (1) encompasses many interesting particular cases of interest:

- By taking

$$A(\mathbf{X}'_t, Y_{t-1}, \mathbf{X}'_{t-1}, Y_{t-2}, \dots; \boldsymbol{\lambda}) = \mathbf{X}'_t \boldsymbol{\beta} + \sum_{i=1}^p \phi_i \mathcal{A}(Y_{t-i}, \mathbf{X}_{t-i}, \boldsymbol{\beta}) + \sum_{i=1}^q \theta_i \mathcal{M}(Y_{t-i}, \mu_{t-i}),$$

for \mathcal{A} and \mathcal{M} representing the autoregressive and moving average terms with associated parameters $\boldsymbol{\phi} = (\phi_1, \dots, \phi_p)'$ and $\boldsymbol{\theta} = (\theta_1, \dots, \theta_q)'$, respectively, and $\boldsymbol{\beta} = (\beta_1, \dots, \beta_r)'$ is the parameter associated to the covariates. This specification for $g(\mu_t)$ resembles a set of ARMA equations with exogenous covariates, often referred to as ARMAX, but it will only satisfy a set of ARMA recurrence equations if \mathcal{M} is carefully chosen. This

general form, briefly mentioned in Benjamin et al. (2003), is still too general for practical purposes.

- A specialization of the previous specification is the base of all the models known in the literature as the Generalized ARMA models (Benjamin et al., 2003), abbreviated GARMA when the random component is a distribution member of the exponential family in canonical form, and called GARMA-like models otherwise, has the form

$$g(\mu_t) = \mathbf{X}'_t \boldsymbol{\beta} + \sum_{j=1}^p \phi_j (g(Y_{t-j}) - \mathbf{X}'_{t-j} \boldsymbol{\beta}) + \sum_{i=1}^q \theta_i r_{t-i}, \quad (2)$$

with r_{t-j} taking one of two commonly used forms. In the predictive scale, $r_t = g(Y_t) - g(\mu_t)$ (Rocha and Cribari-Neto, 2009; Bayer et al., 2017; Pumi et al., 2019, 2024, to cite just a few). With this choice (1) can be shown to satisfy an ARMA set of difference equations and the AR and MA parameters have their usual interpretation. However, $\mathbb{E}(r_t) \neq 0$ and nothing is known about the distribution of r_t ; Considering the response scale instead, namely $r_t = Y_t - \mu_t$, the pair $\{(r_t, \mathcal{F}_t)\}_{t \in \mathbb{Z}}$ is a martingale difference, provided $\mathbb{E}(Y_t) < \infty$. This fact is often used for goodness-of-fit purposes (see Bayer et al., 2023; Benaduce and Pumi, 2023; Prass et al., 2025).

- The models studied in the literature of Poisson Autoregression (PAR) (see Fokianos et al., 2009, and references therein), are often observation-driven and satisfy (1). In PAR models, the distribution of Y_t given \mathcal{F}_{t-1} is assumed to be Poisson with parameter μ_t . Davis et al. (2003) considers a log-linear PAR model by specifying

$$A(\mathbf{X}'_t, Y_{t-1}, \mathbf{X}'_{t-1}, Y_{t-2}, \dots; \beta_1, \beta_2) = \beta_0 + \beta_1 \left(\frac{Y_{t-1} - \mu_{t-1}}{\mu_{t-1}^\alpha} \right),$$

for β_0 and β_1 regression parameter and $\alpha \in (0, 1]$. The log-linear autoregressive models introduced in Fokianos and Tjøtheim (2011) considers

$$A(\mathbf{X}'_t, Y_{t-1}, \mathbf{X}'_{t-1}, Y_{t-2}, \dots; a, b, d) = d + a \log(\mu_{t-1}) + b \log(Y_{t-1} - 1),$$

where a , b and d are parameters. See Fokianos and Tjøtheim (2011) and Fokianos et al. (2009) for more details and related works.

- PARX models introduced by Agosto et al. (2016) consider a term depending of exogenous covariates (hence the X in the name) with specification

$$A(\mathbf{X}'_{t-1}, Y_{t-1}, \mathbf{X}'_{t-2}, Y_{t-2}, \dots; \boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}) = \omega + \sum_{i=1}^p \alpha_i Y_{t-i} + \sum_{j=1}^q \beta_j \mu_{t-j} + f(\mathbf{X}_{t-1}, \boldsymbol{\gamma}),$$

where $f(\cdot; \boldsymbol{\gamma}) : \mathbb{R}^r \rightarrow [0, \infty)$ is a user chosen function depending on a parameter $\boldsymbol{\gamma} \in \mathbb{R}^g$. The additive specification $f(\mathbf{X}_{t-1}, \boldsymbol{\gamma}) = \sum_{i=1}^g \gamma_i f_i(\mathbf{X}_{t-i})$ for real functions f_i is often used in practice.

- The Beta Autoregressive Chaotic (β ARC) models, introduced by Pumi et al. (2021), considers a systematic component that resembles (2) but substituting the MA term by a chaotic process, yielding the specification

$$g_1(\mu_t) = \mathbf{X}'_t \boldsymbol{\beta} + \sum_{j=1}^p \phi_j (g_1(Y_{t-j}) - \mathbf{X}'_{t-j} \boldsymbol{\beta}) + g_2(\mathbf{T}_\theta^{t-1}(U_0)),$$

where g_1 and g_2 are twice differentiable link functions, $T_\theta : [0, 1] \rightarrow [0, 1]$ is a dynamical system, depending on a parameter θ and $T^t = T \circ T^{t-1}$ denotes the t -fold composition. The dynamical system T_θ is assumed to posses an absolutely continuous invariant measure from which U_0 (a random variable) is distributed accordingly. β ARC models present several attractive and interesting properties, allowing the modeling of many dependence structures that are not possible to achieve with other GARMA structures. See Pumi et al. (2021), for more details.

- The Rayleigh Generalized Autoregressive Score (Ray-GAS) model of Peña Ramírez et al. (2024) considers a random component following the Rayleigh distribution parameterized in terms of its mean μ_t and systematic component following a GAS structure of the form

$$\log(\mu_t) = \mathbf{X}'_t \boldsymbol{\beta} + \frac{1}{8} \sum_{i=1}^p \phi_i \left(\frac{\pi y_{t-i}}{\mu_{t-i}} - 4\mu_{t-i} \right) + \sum_{j=1}^q \theta_j \log(\mu_{t-j}).$$

In this case, parameters ϕ_i and θ_j do not have an ARMA-like interpretation.

3 Multiple imputation approach for parameter estimation.

In this section we present the proposed multiple imputation approach for parameter estimation in the context of ODMs, considering that missing data occurs completely at random. Let Y_1, \dots, Y_n be a sample drawn from a ODM for which the random component is distributed according to $f(\cdot | \boldsymbol{\gamma}; \mathcal{F}_{t-1})$, whereas the systematic component follows (1), for some quantity of interest μ_t . The identifiability of $\boldsymbol{\lambda}$ implies that the dependence of $f(\cdot | \boldsymbol{\gamma}; \mathcal{F}_{t-1})$ on $\boldsymbol{\lambda}$ is uniquely tied to the specification of μ_t and conversely. Hence, writing $f(\cdot | \boldsymbol{\gamma}; \mathcal{F}_{t-1})$ is equivalent to $f(\cdot | \mu_t, \boldsymbol{\nu}; \mathcal{F}_{t-1})$. This notation will prove useful in what follows. Suppose that, given a complete sample from $\{Y_t\}_{t \in \mathbb{Z}}$, a method for estimating $\boldsymbol{\gamma}$ is available. At this point, no further assumption is required.

Let \mathcal{I} denote the subset of indexes for which Y_t is observed (not-missing). Naturally, its complement, \mathcal{I}^c , is the set of indexes for which Y_t is missing. The idea of the proposed multiple imputation method for parameter estimation in ODMs is as follows.

Given an initial estimate $\hat{\boldsymbol{\gamma}}_0 = (\hat{\boldsymbol{\nu}}'_0, \hat{\boldsymbol{\lambda}}'_0)'$, missing values are imputed sequentially using a random draw from $f(\cdot | \hat{\mu}_t, \hat{\boldsymbol{\nu}}_0; \mathcal{F}_{t-1})$, where $\hat{\mu}_t$ is reconstructed using the previous information through

$$\hat{\mu}_t = g^{-1}(A(\mathbf{X}_t, Y_{t-1}^*, \mathbf{X}_{t-1}, Y_{t-2}^*, \dots; \hat{\boldsymbol{\lambda}}_0)). \quad (3)$$

with Y_t^* denoting Y_t , if $t \in \mathcal{I}$, or its imputed value, otherwise. This imputation process generates a sample y_1^1, \dots, y_n^1 , referred to as the imputed sample, from which we estimate $\hat{\boldsymbol{\gamma}}_0^1$.

This procedure is repeated $K - 1$ more times, resulting in a set of estimated parameters $\hat{\boldsymbol{\gamma}}_0^1, \dots, \hat{\boldsymbol{\gamma}}_0^K$. We define $\hat{\boldsymbol{\gamma}}_1 := K^{-1} \sum_{k=1}^K \hat{\boldsymbol{\gamma}}_0^k$, as the new reference value for $\boldsymbol{\gamma}$. Then, the procedure discussed so far is iterated using $\hat{\boldsymbol{\gamma}}_1$ as reference, yielding a new estimate $\hat{\boldsymbol{\gamma}}_2$ in the end, and so on. These steps are repeated until convergence is achieved according to a predetermined criteria, or until a predetermined maximum number of iterations, say H , is reached. In any cases, the last estimate obtained, say $\hat{\boldsymbol{\gamma}}$, is taken as the final estimated value for $\boldsymbol{\gamma}$. This process is outlined below in the form of an algorithm, which systematically illustrates these steps.

Algorithm 1: Proposed multiple imputation approach for parameter estimation.

1. Given the observed time series $\{y_t\}_{t \in \mathcal{I}}$, compute the initial estimate $\hat{\gamma}_0$.
2. Set $k = 1$. Starting from $\hat{\gamma}_0$, sequentially impute each missing value y_t for $t \in \mathcal{I}^c$ by sampling from $f(\cdot | \hat{\mu}_t, \hat{\nu}_0)$, where $\hat{\mu}_t$ is recursively computed using (3). Denote the imputed series by y_1^1, \dots, y_n^1 .
3. Using the imputed series y_1^1, \dots, y_n^1 , compute the updated estimate $\hat{\gamma}_0^1$.
4. Repeat steps 2 and 3 for $k \in \{2, \dots, K\}$, obtaining a set of estimates $\hat{\gamma}_0^2, \dots, \hat{\gamma}_0^K$.
5. Define $\hat{\gamma}_1$ as the component-wise sample mean of the estimates $\hat{\gamma}_0^1, \dots, \hat{\gamma}_0^K$.
6. Iterate steps 2 to 5, updating the estimate at each iteration, until convergence is achieved according to some criteria or until a predefined maximum number of iterations H is reached. In any case, the last estimate, say $\hat{\gamma}$ is set as the final estimate.

If we let $\hat{\gamma}_h^1, \dots, \hat{\gamma}_h^K$ be the estimates from which $\hat{\gamma}$ was obtained, we can estimate the uncertainty in the estimation of γ associated with the data imputation by calculating its standard deviation, for instance.

The proposed methodology depends on some prespecified quantities, namely, the starting value $\hat{\gamma}_0$; K , the number of multiple imputations to be performed; a stopping criteria and H , the maximum number of iterations allowed. In what follows we discuss some strategies to determine these values.

3.1 Starting value

The first step in applying the algorithm is to provide the initial estimate of γ , $\hat{\gamma}_0$. There are several ways to obtain this estimate. One simple solution is to use the estimate of γ obtained upon applying the estimation procedure to the longest sequence of contiguous observed values in the sample.

More formally, suppose that \mathcal{I} can be partitioned into U sub-intervals of consecutive indices, say $\mathcal{I}_1, \dots, \mathcal{I}_U$, with $\mathcal{I}_u = \{t_{u,1}, \dots, t_{u,v_u}\}$ denoting the u -th interval of size v_u . Define

$$u^* := \underset{1 \leq u \leq U}{\operatorname{argmax}} \{v_u\}.$$

When u^* is unique, $\mathcal{T} := \mathcal{I}_{u^*}$ is the set of indices corresponding to the longest sequence of non-missing data in the time series. We propose estimating $\hat{\gamma}_0$ using the subset $\{Y_t\}_{t \in \mathcal{T}}$. In cases where two or more sequences have the same maximum length, one can either arbitrarily select one of them or average the estimates obtained from each maximum-length subset to define $\hat{\gamma}_0$.

In some cases, obtaining $\hat{\gamma}_0$ from the longest sequence of contiguous observed values may result in a poor initial value for the algorithm, particularly when the sequence is too short. In such scenarios, an alternative approach is necessary. One viable solution is to consider the longest sequence obtainable from the original time series by imputing at most L consecutive points, where L is small (e.g., 1 or 2), using any automatic imputation method. This approach is valid as long as the number of imputed values is small relative to the total length of the time series.

3.2 Number of multiple imputations

The next step in the implementation of the proposed method is to determine a value for the number of multiple imputations (K) to be used in Step 4. After m iterations of Steps 2 to 5, the estimates $\hat{\gamma}_{m-1}^1, \dots, \hat{\gamma}_{m-1}^K$ are obtained and averaged to define $\hat{\gamma}_m$. Therefore, K should be sufficiently large to ensure that the average of the K estimates converges to a stable value, which will then be used as the updated parameter value for the next iteration of Steps 2 to 5.

Note that the sequence $\hat{\gamma}_{m-1}^1, \dots, \hat{\gamma}_{m-1}^K$ is not independent, as these estimates are computed from time series that differ only in the set \mathcal{I}^c . Results regarding the convergence of the sample mean to a non-random value are well established. For instance, the sample mean computed from a stationary short-range dependent process converges almost surely to a non-random value at a rate of \sqrt{n} , whereas for long-range dependent stationary sequences, convergence still holds but at a slower rate depending on the strength of dependence (Brockwell and Davis, 1991; Palma, 2007).

In the context of GARMA models presented in the simulation, the value of K was determined through a pilot simulation study. Even when the proportion of missing values was as high as 70%, we found that $K = 25$ provided a good balance between computational efficiency, while ensuring the convergence of the sample mean to a stable value. Additionally, our simulation revealed evidence of short-range dependence in the sequence $\hat{\gamma}_{m-1}^1, \dots, \hat{\gamma}_{m-1}^K$. While a much larger value of K could also be used, especially in practical applications where there is no need for thousands of replications, $K = 25$ proved to be sufficient for our purposes.

3.3 Stopping Criteria

We proposed two convergence criteria, as described in the following subsections. Both criteria consider the reduction in variability of the estimated parameter as the algorithm iterates. Low variability in the estimated parameter is naturally associated with the convergence of the iterative process. The question we aim to answer is which criterion is more efficient in detecting process convergence and what the difference is in terms of the estimated value once both criteria have been satisfied.

3.3.1 Coefficient of Variation

The coefficient of variation-based stopping criterion (CVSC) relies on the coefficient of variation of the Euclidean distance between the estimate vectors. At each iteration of the algorithm, after computing the component-wise mean of the estimate vectors $\hat{\gamma}_{h-1}^1, \dots, \hat{\gamma}_{h-1}^K$, the Euclidean distance is calculated between the estimated vector of the current iteration and that of the immediately preceding iteration. The coefficient of variation of these distances is computed at each iteration, the stopping criterion is considered met once the relative distances fall below a pre-specified tolerance, in which case the estimate vector of that iteration is taken as the final estimate associated with that multiple imputation. A pseudo-algorithm for CVSC is as follows:

Algorithm 2: Coefficient of Variation-based stopping criteria - CVSC

1. Obtain the initial estimate vector $\hat{\gamma}_0$;
2. Compute $\hat{\gamma}_1$ following steps 2 to 5 of Algorithm 1;
3. Calculate $d_1 = d(\hat{\gamma}_0, \hat{\gamma}_1)$, where d denotes the Euclidean distance;
4. Compute $\hat{\gamma}_2$ by iterating steps 2 to 5 of Algorithm 1;
5. Calculate $d_2 = d(\hat{\gamma}_1, \hat{\gamma}_2)$ and C_1 , the coefficient of variation of the set $\{d_1, d_2\}$, that is,

$$m_1 = \frac{1}{2} \sum_{i=1}^2 d_i, \quad S_1^2 = \sum_{i=1}^2 (d_i - m_1)^2, \quad C_1 = \frac{S_1}{m_1};$$

6. Iterate the algorithm until $|C_h/C_{h-1} - 1| < \tau$, for some tolerance $\tau \in (0, 1)$.

The tolerance value τ in step 6 represents the relative rate of decline in the variability of the Euclidean distances between successive parameter estimates and controls how close consecutive coefficient of variations must be in terms of proportion in order to declare convergence of the algorithm.

3.3.2 Variance reduction stopping criteria

The second criterion we consider is based on the reduction of variance in consecutive estimates (VRSC). The idea is that as the algorithm iterates, we track the behavior of the variance vector associated with the estimated parameters at each iteration. Specifically, at each step, we compute the Euclidean distance between the current variance vector and that of the previous iteration. The algorithm is considered to have met the stopping criterion when the absolute difference between consecutive distances falls below a predefined tolerance. The VRSC stopping criterion can be systematized as follows:

Algorithm 3: Variance reduction stopping criteria - VRSC

1. Obtain the initial estimate vector $\hat{\gamma}_0$;
2. Obtain $\hat{\gamma}_1$ by following steps 2 to 5 of Algorithm 1;
3. Calculate the vector \mathbf{S}_1^2 containing the component-wise sample variance of the set $\{\hat{\gamma}_0, \hat{\gamma}_1\}$;
4. Obtain $\hat{\gamma}_2$ by iterating steps 2 to 5 of Algorithm 1;
5. Calculate \mathbf{S}_2^2 , the vector containing the component-wise sample variances of the set $\{\hat{\gamma}_0, \hat{\gamma}_1, \hat{\gamma}_2\}$. Compute $d_1 = d(\mathbf{S}_1^2, \mathbf{S}_2^2)$, i.e., the Euclidean distance between the vectors \mathbf{S}_1^2 and \mathbf{S}_2^2 ;
6. Iterate the algorithm until $|d_h - d_{h-1}| < \tau$, for some tolerance $\tau > 0$.

The tolerance τ in the VRSC has a different meaning to the CVSC as it considers the difference between consecutive euclidean distances calculated from consecutive variance vectors. As such, the CVSC tends to be more conservative regarding convergence, often requiring more iterations than VRSC for the same threshold. A comparison between these two criteria

in the context of GARMA model is presented in Section 5.

3.4 Handling Non-Convergence and the Maximum Number of Iterations

Algorithm 1 may not always converge within the predefined maximum number of iterations. In such cases, simple diagnostic measures can be employed to assess and address the issue. A practical approach is to restart the algorithm using the last estimated value as the initial state – this is equivalent to increasing H from the start but is computationally more efficient.

Another factor that may affect convergence is the choice of the initial state, particularly in time series with a high percentage of missing values, where obtaining a reliable initial parameter estimate can be challenging. In these cases, restarting the algorithm using the last obtained parameter may improve convergence. Additionally, the stopping criteria can be reviewed, either by adjusting the threshold or modifying the criterion itself.

For likelihood-based estimation procedures, the algorithm may fail to converge if it becomes trapped in a region of the likelihood surface with little variation, preventing updates from triggering the stopping criteria. This can be diagnosed by plotting the sequences obtained in step 4. If necessary, increasing the value of K or modifying the initial state might help.

In our simulations, we found that setting $H = 30$ iterations was generally sufficient to achieve high convergence rates across all scenarios with the applied thresholds. As expected, the proportion of non-convergent cases increased with the percentage of missing data. However, even at 70% missing values, the highest observed non-convergence rate was 2.3% for the CVSC and 3.7% for the VRSC, which we consider acceptably low. For further details, please refer to Section 5.5.

4 Monte Carlo Simulation Study

In this section we present a simulation study in the context of GARMA models. To the best of our knowledge, this is the first study on GARMA models with missing data in the literature. We will consider 3 GARMA-like models with systematic components given by (2) with errors in the predictor scale, meaning $r_t := g(y_t) - g(\mu_t)$, with g being the logit function. In the following we present the models from the oldest to newest. The β ARMA (Beta ARMA) model was introduced in Rocha and Cribari-Neto (2009). For the β ARMA, the Beta distribution is parametrized in terms of its conditional mean, denoted by $\mu_t \in (0, 1)$, and a shape parameter $\nu > 0$, resulting in the density

$$f(y; \mu_t, \nu | \mathcal{F}_{t-1}) = \frac{\Gamma(\nu)}{\Gamma(\nu\mu_t)\Gamma(\nu(1-\mu_t))} y^{\nu\mu_t-1} (1-y)^{\nu(1-\mu_t)-1} I(0 < y < 1).$$

The KARMA (Kumaraswamy ARMA) model was introduced in Bayer et al. (2017). For the KARMA, the Kumaraswamy distribution is parametrized in terms of its conditional median, represented by $\mu_t \in (0, 1)$, and a shape parameter $\nu > 0$. The conditional density is given by

$$f(y; \mu_t, \nu | \mathcal{F}_{t-1}) = \frac{\varphi \log(0.5)}{\log(1 - \mu_t^\nu)} y^{\nu-1} (1-y)^\frac{\log(0.5)}{\log(1 - \mu_t^\nu)} I(0 < y < 1).$$

The UWARMA (Unit-Weibull ARMA) model was introduced in Pumi et al. (2024). For the UWARMA, the Unit-Weibull distribution is parameterized in terms of its ρ th quantile,

represented by $\mu_t \in (0, 1)$, and a shape parameter $\nu > 0$, for any fixed $\rho \in (0, 1)$. The conditional density on a UWARMA model is given by

$$f_\rho(y; \mu_t, \nu | \mathcal{F}_{t-1}) = \frac{\nu}{y} \left[\frac{\log(\rho)}{\log(\mu_t)} \right] \left[\frac{\log(y)}{\log(\mu_t)} \right]^{\nu-1} \rho^{\left[\frac{\log(y_t)}{\log(\mu_t)} \right]^\nu} I(0 < y < 1).$$

The β ARMA, KARMA and UWARMA were chosen because these models are featured in the package **BTSR** (Prass and Pumi, 2025) for R (R Core Team, 2023), which makes them freely available for anyone to use. Although β ARC models are also available in **BTSR**, they were not included in the simulation as they present systematic component whose behavior is radically different from (2), making comparison impossible with the other models.

It is worth mentioning that inference in KARMA models considered the more restrictive approach of conditional maximum likelihood (CMLE). Originally in Rocha and Cribari-Neto (2009), the authors also considered CMLE, but the β ARMA models are a particular case of the β ARFIMA models of Pumi et al. (2019), where the authors considered inference via partial maximum likelihood (PMLE), adopted here. Computationally speaking, the difference between using PMLE or CMLE is immaterial.

4.1 Data Generating Process

Using the **BTSR** package (Prass and Pumi, 2025, version 1.0.0), time series of size $n = 500$ were generated from models β ARMA, KARMA, and UWARMA models, with order $p = q = 1$, considering a burn-in of size 100. Three scenarios were established for the simulations, each corresponding to a parameter vector $\gamma = (\alpha, \phi, \theta, \nu)'$ with $\gamma_1 = (0.5, -0.4, -0.6, 20)'$, $\gamma_2 = (0.5, -0.5, -0.3, 30)'$, and $\gamma_3 = (0.5, -0.4, -0.2, 20)'$. For simplicity, the same parameter ν was used to denote the fixed parameter of the three models. Missing data were introduced at proportions of $r \in \{0.1, 0.4, 0.7\}$, assuming that they occur completely at random. For simplicity, we assume that the first and last values of the time series were never missing.

These time series with missing data were then subjected to the multiple imputation process described in Algorithm 1, resulting in imputed series, on which parameter estimation using PMLE was performed. The data were generated using the stopping criteria described in Section 3.3. In both cases, the tolerance was fixed at $\tau = 0.01$ and the maximum number of iterations was set to $H = 30$ and K was set to 25 following the results of a pilot simulation (not shown). Within the **BTSR** package, optimization was conducted using Nelder-Mead. The simulation was carried out using the R language (R Core Team, 2023), version 4.4.1. For each combination of model, scenario, and proportion of missing data, the process was replicated $R = 1,000$ times.

5 Simulation Results

The simulation results are presented in Figures 1 to 8, displayed sequentially in this section. Each figure consists of twelve blocks. Each block represents a combination of parameter and proportion of missing data, described respectively in the columns and rows. Each block contains three box plots of the estimated parameters, two of which correspond to the stopping criteria (CVSC and VRSC, discussed in Section 3.3), and one referring to the estimates based on the complete series. It is worth noting that the latter remain the same across the rows and serve only as a reference.

5.1 Scenario 1

Simulation results are presented in Figures 1 for the β ARMA and KARMA (displayed together due to similarity) and 2 for the UWARMA. For the three models, as the proportion of missing values increase, we observe an increase in bias and variance across all cases, except for parameter ϕ in the context of β ARMA and $r \in \{0.1, 0.4\}$, where the bias was not significantly affected by the missing values. Notably, the results for θ approached 0 as r increased to 0.7.

As for the stopping criteria, for $r \in \{0.1, 0.4\}$ both present similar behavior in terms of bias and variance for all models. For $r = 0.7$, the results are mixed. Both stopping criteria presented quite similar behavior for the UWARMA, the CVSC presented slightly better results for the β ARMA whereas VRSC presented smaller variance and slightly smaller bias in the case of KARMA models.




Figure 1: Simulation results for the BARMA and KARMA models under scenario 1.




Figure 2: Simulation results for the UWARMA model under scenario 1.

5.2 Scenario 2

Figure 3 illustrates the simulation results for the β ARMA model. The introduction of missing data in the proportion of $r = 0.10$ did not cause a significant difference between the median of the estimates of α and ϕ and the variance was not significantly affected either. Overall θ was overestimated whereas ν was underestimated. As r increases over 0.1, the bias and variance for all parameter increase as well. For $r \in \{0.1, 0.4\}$ both stopping criteria presented similar results, while the CVSC performed slightly better for $r = 0.7$.


Figure 3: Simulation results for the β ARMA model under scenario 2.

Figure 4 presents the results for the KARMA model. Unlike the β ARMA model, a significant difference was observed between the median of the estimates and all true parameters, except for ν . For the complete series, the estimates of θ were concentrated around zero, with median exactly 0. It is important to note that the BTSR initializes θ exactly at 0. The fact that the median of the estimates for θ is exactly equal to zero indicates that the optimization algorithm did not move from the initial point for most of the estimated series, suggesting the existing of a flat likelihood surface in the vicinity of $\theta = 0$. This is a behavior frequently observed in practice, especially for low values of ν . The introduction of missing data at a proportion of $r = 0.10$ caused the median of the estimates for θ to move away from 0 toward the true value, suggesting that the imputation had a sufficient effect on the likelihood surface to allow the optimization algorithm to move away from the initial value of 0 and into the correct direction.

The bias on the estimates of α and ϕ had an inverted sign compared to those of the complete series, with the magnitude increasing as the proportion of missing data increased. For θ , the difference between the median estimates and the true parameter decreased for $r = 0.1$, unlike the case for $r \in \{0.4, 0.7\}$, where the estimates remained centered around zero. ν is always underestimated with bias increasing with r . As for the stopping criteria, both presented similar results for $r \in \{0.1, 0.4\}$ and the VRSC performed slightly better for $r = 0.7$.




Figure 4: Simulation results for the KARMA model under scenario 2

Figure 5 presents the simulation results for the UWARMA model. The introduction of missing data, for all values of r , led to an increase in the difference between the median estimates and the true parameter, except for θ . Parameter α and ν tended to be underestimated whereas ϕ tended to be overestimated. For $r = 0.1$, the estimates for α and ϕ were nearly identical to the ones obtained from the complete series. When $r = 0.1$, the estimated values for θ obtained from the imputed time series were significantly better than the ones obtained from the complete time series. However, for $r \in \{0.4, 0.7\}$, the estimates of θ for the imputed series were moving toward zero as r increased. As for the stopping criteria, both present a comparable performance with no clear winner.




Figure 5: Simulation results for the UWARMA model under scenario 2

5.3 Scenario 3

Figure 6 presents the simulation results for the β ARMA model, which show a behavior similar to other cases, with bias and variances increasing as r increases. For $r = 0.1$ the estimated values from the imputed and the complete time series are nearly identical. The estimates obtained from both stopping criteria are nearly identical, with no clear winner.

Figure 7 presents the results for the KARMA model. Similarly to scenario 2, the estimated values for θ obtained from the complete time series are concentrated at 0. For $r = 0.1$, the estimated values move away from 0, but ultimately tend to fluctuate around zero as r increases. The estimates for α and ϕ obtained from the complete series exhibit considerable bias, but when missing data are inserted at rate $r = 0.1$, the estimates obtained from the imputed time series are more accurate than those obtained from the complete series, albeit with greater variability.

For $r = 0.4$, the estimates of α and ϕ showed bias of similar magnitude to that of the complete series but with an inverted signal. For the parameter ν , a clear pattern was observed: as the proportion of missing data increased, the bias of the estimates grew negatively. Considering the stopping criterias, the estimates obtained when $r \in \{0.1, 0.4\}$ are nearly identical, whereas for $r = 0.7$ the VRSC was superior.



Figure 6: Simulation results for the β ARMA model under scenario 3.

Figure 7: Simulation results for the KARMA model under scenario 3.




Figure 8: Simulation results for the UWARMA model under scenario 3.

The results for the UWARMA model, are shown in Figure 8. Estimates of α and ϕ obtained from the complete time series and from the imputed time series when $r = 0.1$ are nearly identical. For θ , the estimates obtained from the imputed time series presented smaller bias, but as r increased, the situation was reversed. For r increased away from 0.1, bias and variance of the estimates obtained from the imputed time series increased. Once more, ν was underestimated with bias increasing with r . The estimated values obtained with both criteria are nearly identical.

5.4 Discussion

The bias and variance of the estimates tended to increase as the proportion of missing data moves away from 10%, which is expected due to the dominance of possibly inaccurate information due to imputation and the loss of information from the observed values. This effect is especially noticeable for $r = 0.7$, whereas for $r = 0.4$ often the effect was less noticeable for parameter α and ϕ . Also noteworthy is the tendency of the PMLE to underestimate ν , for all models and scenarios, with rare exceptions. For the UWARMA, both stopping criteria yielded nearly identical results, but overall both criteria performed similarly, especially when $r \in \{0.1, 0.4\}$. For $r = 0.7$, a marginal advantage for VRSC was observed.

An important facet of imputation in the context of time series analysis under massive amounts of missing data was also observed here. For all analyzed models, it was observed that as the percentage of missing data increases, the estimates of ϕ and θ tend to approach zero. This behavior is, to some extent, expected, since, although the proposed method takes into account the dependence structure reconstructed from the estimates obtained at each step to impute the series, the presence of a large volume of missing data – as in the case of $r = 0.7$ – ends up limiting the effectiveness of the imputation. In this context, imputation based on approximate parameter values is unable to adequately restore the correlation structure in μ_t , resulting in an imputed series close to a uncorrelated sample. Consequently, the estimates of ϕ and θ , which are responsible for this structure, tend to zero. This effect was, overall more evident for the parameter θ . This is also a consequence of the loss of information that

occurs under massive percentage of missing values, which ultimately affect every imputation method. Nevertheless, the results obtained can still be considered satisfactory, given the extreme nature of the problem when $r = 0.7$.

5.5 Iterations to Convergence

Table 1 presents the average number of iterations until convergence for the replications in which the maximum number of iterations, $H = 30$, was not reached. The table includes results for all combinations of model type, stopping criterion, percentage of missing data, and scenario. The percentage of replications that reached the maximum number of iterations is provided in parentheses. Notably, the average number of iterations until convergence is systematically lower for the VRSC stopping criterion, with the CVSC required over 40% more iterations to attain convergence in most cases.

As for non-converge, this percentage is quite low overall peaking at 2.3% for the CVSC and 3.7% for the VRSC. For $r = 0.1$ the CVSC present uniformly higher percentage of non-convergence compared to VRSC, peaking at 0.8%. The situation is reversed when considering scenario 2, for $r \in \{0.4, 0.7\}$. For other cases, no clear pattern was found, with CVSC and VRSC alternating their position. Interestingly, the average number of iterations for the CVSC criterion increases with the proportion of missing data, r . In contrats, the average number of iterations to convergence for for the VRSC criterion shows homogeneity across all values of r .

Table 1: Average number of iterations for convergence and the percentage of replications that reached the maximum number of iterations (in parenthesis).

| r | Scenario | β ARMA | | KARMA | | UWARMA | |
|-----|----------|--------------|------------|-------------|------------|-------------|------------|
| | | CVSC | VRSC | CVSC | VRSC | CVSC | VRSC |
| 0.1 | 1 | 11.58 (0.3) | 7.61 (0.0) | 11.24 (0.3) | 6.37 (0.0) | 11.22 (0.8) | 6.70 (0.0) |
| | 2 | 11.55 (0.6) | 9.99 (0.2) | 11.41 (1.2) | 8.33 (0.0) | 11.22 (0.4) | 8.76 (0.0) |
| | 3 | 11.21 (0.5) | 7.31 (0.1) | 11.30 (0.2) | 5.58 (0.0) | 11.24 (0.5) | 6.06 (0.0) |
| 0.4 | 1 | 12.71 (0.8) | 6.62 (0.3) | 12.23 (0.3) | 6.06 (0.1) | 12.63 (0.6) | 6.71 (0.2) |
| | 2 | 12.77 (0.6) | 9.57 (1.6) | 12.50 (0.5) | 8.25 (0.7) | 12.64 (0.6) | 8.43 (1.2) |
| | 3 | 12.60 (0.6) | 7.95 (0.8) | 12.14 (0.4) | 6.00 (0.0) | 12.63 (0.6) | 6.70 (0.2) |
| 0.7 | 1 | 13.82 (1.4) | 5.70 (1.7) | 13.08 (1.2) | 5.56 (0.9) | 14.19 (1.4) | 6.27 (0.7) |
| | 2 | 13.75 (2.3) | 9.11 (3.7) | 13.71 (1.7) | 7.60 (3.1) | 14.61 (2.0) | 7.95 (2.7) |
| | 3 | 13.73 (1.9) | 7.99 (2.7) | 13.25 (0.8) | 5.98 (1.2) | 14.16 (1.3) | 6.30 (0.6) |

5.6 Joint Asymptotic Behavior

To heuristically assess whether the asymptotic normality of the PMLE computed using the proposed methodology may hold, we examine pairwise scatter plots and the marginal behavior of the estimates (histograms and boxplots). Figures 9, 10, and 11 present the results for the β ARMA, KARMA, and UWARMA models, respectively, corresponding to scenario 1 of the simulation ($\alpha = 0.5$, $\phi = -0.4$, $\theta = -0.6$, and $\nu = 20$) and using the VRSC stopping criterion.




Figure 9: Pairwise joint and marginal behavior of the estimated values for the β ARMA model. Solid lines in the scatter plot represent the true values.




Figure 10: Pairwise joint and marginal behavior of the estimated values for the KARMA model. Solid lines in the scatter plot represent the true values.




Figure 11: Pairwise joint and marginal behavior of the estimated values for the UWARMA model. Solid lines in the scatter plot represent the true values.

Focusing first on the marginal distributions, the histograms for parameter ν closely resemble a normal distribution but are centered at increasingly underestimated values as r increases across all models. A similar pattern is observed for parameters α and ϕ , with both bias (in absolute value) and variance increasing as r grows. Parameter θ exhibits the worst behavior, with estimates frequently concentrating around zero.

The scatter plots for parameter pairs (α, ϕ) , (α, θ) , and (ϕ, θ) indicate a strong correlation among these estimates across all models, except for (α, ϕ) in the β ARMA case. For $r \in \{0.1, 0.4\}$, the scatter plots generally display a well-defined bivariate normal pattern. Scatter plots involving parameter ν exhibit a recognizable bivariate normal pattern, with weaker correlations between ν and the other parameters.

6 Empirical Analysis

In order to showcase the usefulness of the proposed methodology, we conducted an empirical analysis on a real dataset. We consider the same data used in Scher et al. (2020), referring to the proportion of hydroelectric energy stored in southern Brazil from January 2001 to April 2017, with a sample size $n = 196$. Missing data were introduced at random, following a uniform distribution on the time stamps $2, \dots, 195$ (for simplicity the first and last observation were assumed never missing) at proportions of $r \in \{0.2, 0.4, 0.7\}$. The time series plot of the data is presented in Figure 13(a), while Figure 13(b) present the same time series after randomly removing 70% of the data. Next, the multiple imputation procedure was applied to these artificially created time series, along with the estimation of the parameters for models β ARMA(1, 1) and KARMA(1, 1). We used $K = 25$, $H = 30$ and $\tau = 0.01$, as in the simulation. This procedure was repeated 1,000 times.

The results of the empirical analysis are presented in Figure 12, which consists of eight blocks arranged in four columns corresponding to the parameters and two rows corresponding to the models. Each block contains three box plots, one for each proportion of missing data, along with a vertical line indicating the parameter estimate obtained from the complete series.




Figure 12: Estimation results for the β ARMA and KARMA models, for proportion of missing data $r \in \{0.2, 0.4, 0.7\}$.


For the β ARMA model, α presented higher estimated values when compared to the estimate obtained from the complete time series, with magnitude increasing with r . For parameters ϕ and ν the situation was reversed. The estimates of θ oscillate between the estimated

value in the complete time series.


The KARMA model yielded estimates closer to the ones obtained from the complete time series for α and ϕ when $r = 0.2$. However, introducing missing data at proportions $r \in \{0.4, 0.7\}$ pulled the estimates for these parameters away from the estimates obtained from the complete time series. Regarding θ , the estimates were concentrated around zero for $r \in \{0.2, 0.4\}$, but for $r = 0.7$, its median was closer to the estimate obtained from the complete series.

The estimates of ν in the KARMA model exhibited a pattern similar to that observed in the β ARMA model. A significant increase in the variability of the estimates for all parameters when $r = 0.7$ was common to both models. For $r \in \{0.2, 0.4\}$, the increase in variance was less noticeable, especially for the KARMA.

We emphasize that, in some cases, obtaining $\hat{\gamma}_0$ from the longest sequence of contiguous observed values can negatively impact the initial value results, which plays an important role in the proposed methodology. This occurs when the value of n is small and when the proportion of missing data r is large, resulting in the longest sequence of observed data being very short, complicating estimation by the PMLE. Figures 13(a) and 13(b) illustrate this point. In this example, the longest sequences of contiguous values has only 4 points, a situation in which obtaining a good initial estimate is challenging.



(a) Original time series ($n = 196$).



(b) Time series with 70% of missing data.

Figure 13: Visual comparison between (a) the complete time series and (b) the same time series after randomly removing 70% of the data.

To overcome this problem, we can employ the solution presented in Section 3.1. The idea is to consider the longest sequence of the original time series imputed with at most L consecutive points, using any imputation method. To assess the impact of implementing this strategy, we conduct the same experiment as before, but considering this strategy for $L \in \{0, 1, 2\}$, imputing values using the linear interpolation method implemented in package `imputeTS`. Observe that $L = 0$ is equivalent to using the longest sequence without any modification presented in Figure 12.

The results are presented in Figures 14, for the β ARMA, and 15 for the KARMA. Each figure contains twelve blocks, which are arranged in four columns representing the parameters and three rows representing the values of r . Each block contains three boxplots, each associated with one of the considered values of L . The vertical dashed blue lines represent the parameter estimates obtained from the complete series for the respective model, presented for reference.




Figure 14: Experiment results considering the β ARMA model.




Figure 15: Experiment results considering the KARMA model.

For simplicity, we begin by analyzing the results for the KARMA model, presented in Figure 15. For almost all parameters and all values of r , increasing the value of L substantially improved the final estimated value, in the sense that it was closer to the values obtained from the complete time series. The only exception is θ when $r = 0.7$, in which case increasing L yielded worse results.

For the β ARMA model, the behavior is more complex. For the parameters α and ϕ , when $r \in \{0.2, 0.4\}$, using $L = 0$ or $L = 1$ yielded similar results, whereas $L = 2$ resulted in slightly worse estimates. However, for $r = 0.7$, increasing L improved the estimates for both parameters.

For θ , when $r \in \{0.2, 0.4\}$, increasing L improved the estimates. For $r = 0.7$, $L = 1$ yielded substantially better results than $L = 0$, whereas for $L = 2$, the results were worse than the previous ones.

For ν , $L = 0$ consistently produced the worst results. When $r = 0.1$, $L = 1$ yielded the best results, whereas for $r \in \{0.4, 0.7\}$, $L = 2$ provided the best estimates. Table 2 presents the average length of the longest contiguous sequence for each value of L used and $r \in \{0.2, 0.4, 0.7\}$. The average length is naturally a function of r . For $r = 0.2$, the average length is 36, which is relatively low, though not extremely so. For $r = 0.7$, however, this value drops to less than 12. In the most extreme case, $r = 0.7$, the average length increases by approximately 50% for each unitary increase in L , with this percentage increase being higher for lower values of L .

Table 2: Average length of the longest contiguous sequence.

| $L \backslash r$ | 0.2 | 0.4 | 0.7 |
|------------------|-------|------|------|
| 0 | 36.0 | 22.3 | 11.9 |
| 1 | 84.9 | 42.4 | 17.6 |
| 2 | 149.0 | 73.8 | 25.8 |

References

Agosto, A., Cavaliere, G., Kristensen, D., and Rahbek, A. (2016). Modeling corporate defaults: Poisson autoregressions with exogenous covariates (PARX). *Journal of Empirical Finance*, 38:640–663. Recent developments in financial econometrics and empirical finance.

Bayer, F. M., Bayer, D. M., and Pumi, G. (2017). Kumaraswamy autoregressive moving average models for double bounded environmental data. *Journal of Hydrology*, 555:385–396.

Bayer, F. M., Pumi, G., Pereira, T. L., and Souza, T. C. (2023). Inflated beta autoregressive moving average models. *Computational And Applied Mathematics*, 42:183.

Benaduce, H. S. C. and Pumi, G. (2023). SYMARFIMA: a dynamical model for conditionally symmetric time series with long range dependence mean structure. *Journal of Statistical Planning and Inference*, 225:71–88.

Benjamin, M. A., Rigby, R. A., and Stasinopoulos, D. M. (2003). Generalized autoregressive moving average models. *Journal of the American Statistical Association*, 98(461):214–223.

Brockwell, P. J. and Davis, R. A. (1991). *Time Series: Theory and Methods*. Springer-Verlag, 2 edition.

Carpenter, J. R. and Kenward, M. G. (2012). *Multiple Imputation and Its Application*. Statistics in Practice. John Wiley & Sons.

Cox, D. R. (1975). Partial likelihood. *Biometrika*, 62:69–76.

Cox, D. R. (1981). Statistical analysis of time series: some recent developments. *Scandinavian Journal of Statistics*, 8:93–115.

Davis, R. A., Dunsmuir, W. T. M., and Streett, S. B. (2003). Observation-driven models for Poisson counts. *Biometrika*, 90(4):777–790.

Fokianos, K. and Kedem, B. (2004). Partial likelihood inference for time series following generalized linear models. *Journal of Time Series Analysis*, 25(2):173–197.

Fokianos, K., Rahbek, A., and Tøstheim, D. (2009). Poisson autoregression. *Journal of the American Statistical Association*, 104(488):1430–1439.

Fokianos, K. and Tjøtheim, D. (2011). Log-linear Poisson autoregression. *Journal of Multivariate Analysis*, 102(3):563–578.

Little, R. and Rubin, D. (2020). *Statistical Analysis with Missing Data*. Wiley Series in Probability and Statistics. John Wiley & Sons, 3rd edition.

Moritz, S. and Bartz-Beielstein, T. (2017). imputeTS: Time series missing value imputation in R. *The R Journal*, 9(1):207–218.

Palma, W. (2007). *Long-Memory Time Series: Theory and Methods*. Wiley Series in Probability and Statistics. Wiley.

Peña Ramírez, M., Rojas Guerra, R., and Bayer, F. M. (2024). The Rayleigh generalized autoregressive score model for SAR data interpretation. *IEEE Geoscience and Remote Sensing Letters*, 21:1–5.

Prass, T. S. and Pumi, G. (2025). *BTSR: Bounded Time Series Regression*. R package version 1.0.0.

Prass, T. S., Pumi, G., Taufembach, C. G., and Carlos, J. H. (2025). Positive time series regression models: theoretical and computational aspects. *Computational Statistics*, 40:1185–1215.

Pumi, G., Prass, T. S., and Souza, R. R. a. (2021). A dynamic model for double-bounded time series with chaotic-driven conditional averages. *Scandinavian Journal of Statistics*, 48(1):68–86.

Pumi, G., Prass, T. S., and Taufembach, C. G. (2024). Unit-Weibull autoregressive moving average models. *TEST*, 33:204–209.

Pumi, G., Ulloa, G. C., and Prass, T. S. (2025). Estimation of long-range dependent models with missing data: to impute or not to impute? arXiv:2303.04754.

Pumi, G., Valk, M., Bisognin, C., Bayer, F. M., and Prass, T. S. (2019). Beta autoregressive fractionally integrated moving average models. *Journal of Statistical Planning and Inference*, 200:196 – 212.

R Core Team (2023). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.

Rocha, A. V. and Cribari-Neto, F. (2009). Beta autoregressive moving average models. *TEST*, 18(3):529–545.

Scher, V. T., Cribari-Neto, F., Pumi, G., and Bayer, F. M. (2020). Goodness-of-fit tests for arma hydrological time series modeling. *Environmetrics*, 31.

van Buren, S. (2012). *Flexible Imputation of Missing Data*. Interdisciplinary Statistics. CRC Press.

Wong, W. H. (1986). Theory of partial likelihood. *The Annals of Statistics*, 14:88–123.