

# THE HOMOCLINIC STRUCTURE OF GENERIC SECTIONAL-HYPERBOLIC CLASSES.

ELIAS REGO AND KENDRY J. VIVAS

**ABSTRACT.** The notion of sectional-hyperbolicity is a weakened form of hyperbolicity introduced for vector fields in order to understand the dynamic behavior of certain higher-dimensional systems such as the multidimensional Lorenz attractor. In this paper we explore the questions proposed in [*Math. Z.*, **298** (2021), 469-488] and we provide an affirmative answer by proving that  $C^1$ -generic non-trivial sectional-hyperbolic chain-recurrent class is robustly a homoclinic class. We then apply our findings to the dynamics of star flows providing a positive answer to Conjecture 1.2 in [*Tran. Amer. Math. Soc.* **376** (2023), 6845-6871]. We remark that our arguments do not assume Lyapunov stability.

## 1. INTRODUCTION

Differentiable dynamical systems have become a fruitful and important research field in recent decades. The birth of this discipline is attributed to the seminal work of S. Smale [[Sm67](#)], where the notion of hyperbolic set was introduced, presenting Smale's Horseshoe as a representative example. Since then, a very strong program towards the understanding of the global dynamics of dynamical systems has been initiated, yielding a rich theory where several important advances were achieved, including a better understanding of stability phenomena and of chaotic dynamics from both topological and statistical point of view. Later, motivated by the chaotic dynamics of the strange attractor in Lorenz's polynomial system

$$\begin{cases} \dot{x} = \sigma(y - x) \\ \dot{y} = \rho x - y - xz \\ \dot{z} = xy - \beta z, \end{cases}$$

where  $\sigma \approx 10$ ,  $\beta \approx 8/3$  and  $\rho \approx 28$ , [[ABS77](#)] and [[Gu76](#)] introduced, independently, a geometric model now known as the geometric Lorenz attractor (GLA). One of its main features is the presence of a unique singularity that is accumulated by regular orbits, which causes the system to fail to be hyperbolic. Nevertheless, the GLA exhibits rich dynamical behavior reminiscent of hyperbolic sets, including a dense set of periodic orbits and robust transitivity. Moreover, it was shown in [[Ba04](#)] that the GLA is a homoclinic class. Motivated by this example, the notion of singular-hyperbolicity was introduced in [[MPP99](#)], where it is proven that singular-hyperbolic sets properly extend the classical notion of hyperbolic sets, incorporating the GLA as a prototypical case.

---

2020 *Mathematics Subject Classification.* Primary: 37C70, 37C10, 37C20, 37D45.

*Key words and phrases.* Flows, Robustly Transitive, Homoclinic Class, Sectional Hyperbolic, Star Flows.

In order to gain a better understanding of the dynamics of higher-dimensional sets such as the multidimensional Lorenz attractor [BPV97], C. Morales and R.J. Metzger introduced the concept of sectional-hyperbolic set in [MM08]. Both notions, sectional-hyperbolicity and singular-hyperbolicity, agree for three-dimensional vector fields, but in the higher-dimensional setting sectional-hyperbolicity is stronger than singular-hyperbolicity (see [MM08] and [Sa19] for more details).

Now, in topological dynamics, the chain-recurrent set constitutes an important subject of study because it encompasses all the interesting dynamics of the system. Moreover, by considering an equivalence relation, this set is decomposed into pieces called chain-recurrence classes. The fundamental theorem of dynamical systems, due to Conley [Con78], asserts the existence of a Lyapunov function which is constant along these classes. In this way, this result provides a general procedure for describing the global dynamics of a system. These sets are the central focus of this work and hence we now recall this concept.

Hereafter, we denote by  $M$  a  $n$ -dimensional compact Riemannian manifold, endowed with a Riemannian metric  $\|\cdot\|$ . We will always assume  $n \geq 4$ . Denote by  $d$  the metric on  $M$  induced by its Riemannian metric. Throughout this text,  $\mathcal{X}^1(M)$  denotes the set of  $C^1$ -vector fields on  $M$  endowed with the  $C^1$ -topology. It is well known that any  $X \in \mathcal{X}^1(M)$  induces a  $C^1$ -flow that will be denoted by the one-parameter family of maps  $\{X_t\}_{t \in \mathbb{R}}$ . The *orbit* of a point  $x \in M$  is the set

$$\mathcal{O}(x) = \{X_t(x) : t \in \mathbb{R}\}.$$

For  $a, b \in \mathbb{R}$ , the *orbit segment from  $a$  to  $b$*  of a point  $x$  is defined by  $X_{[a,b]}(x) = \{X_t(x) : t \in [a, b]\}$ . A point  $x \in M$  is said to be a *singularity* of  $X$  if  $X(x) = 0$ . We will denote the set of singularities of  $X$  by  $Sing(X)$ . A point  $x \in M \setminus Sing(X)$  is a *periodic point* of  $X$  if there is  $t > 0$  such that  $X_t(x) = x$ . Denote by  $Per(X)$  the set of periodic points of  $X$ . The set of *critical elements* of  $X$  is given by  $Crit(X) = Sing(X) \cup Per(X)$ . An orbit that does not belong to  $Crit(X)$  is called a *regular orbit*. As usual, we say that a subset  $\Lambda$  of  $M$  is *invariant* if  $X_t(\Lambda) = \Lambda$  for any  $t \in \mathbb{R}$ . We say that a compact invariant set  $\Lambda$  is *Lyapunov stable* if for every neighborhood  $U$  of  $\Lambda$  there exists a neighborhood  $V \subset U$  of  $\Lambda$  such that  $X_t(V) \subset U$ , for any  $t > 0$ . We say that a compact and invariant set  $\Lambda$  is *transitive* if it contains a point whose orbit is dense in  $\Lambda$ .

Let us now define the chain-recurrent sets. There are several equivalent ways of defining the chain-recurrent set (see [AN07, Theorem 2.7.18]), here we choose the form that is most suitable to our purposes. For  $\varepsilon > 0$  and  $T > 0$ , we say that a finite sequence  $(x_i, t_i)_{i=0}^n$  is an  $(\varepsilon, T)$ -chain if  $t_0 + \dots + t_n \geq T$ ,  $t_i \geq 1$ , and  $d(X_{t_i}(x_i), x_{i+1}) < \varepsilon$  for any  $i = 0, \dots, n-1$ . Besides, we say that  $y$  is *chain attainable from  $x$* , and we denote it by  $x \sim y$ , if for any  $\varepsilon, T > 0$  there exists an  $(\varepsilon, T)$ -chain from  $x$  to  $y$ , i.e.  $x_0 = x$  and  $x_n = y$ . When  $x \sim x$ , we say that  $x$  is a *chain-recurrent point*. The chain-recurrent set of  $X$  is defined as

$$CR(X) = \{x \in M : x \sim x\}.$$

It is well known that  $CR(X)$  is a compact and invariant set. Moreover,  $\sim$  is an equivalence relation on  $CR(X)$ . So, each equivalent class under this relation is called *chain-recurrent class*. It is easy to see that each chain-recurrent class of  $X$  is also a compact and invariant

set. When  $x \in CR(X)$ , we denote by  $C(x)$  its chain-recurrence class. We say that a chain-recurrent class is *non-trivial* if it is not reduced to either a periodic orbit or a singularity. We say that a chain-recurrent class is an *aperiodic class* if it does not contain periodic orbits.

Several generic properties of these chain-recurrence classes were obtained. For instance, the well-known Kupka-Smale Theorem states that critical orbits of generic vector fields are hyperbolic. Later, in [BC04], it was shown that for  $C^1$ -generic systems, chain-recurrence classes with periodic orbits coincide with the homoclinic class of some of such periodic orbits. Moreover, when some form of hyperbolicity is present, we obtain more interesting dynamical properties. Indeed, according to [PY21], for  $C^1$ -generic vector fields, a non-trivial Lyapunov stable chain-recurrence class that is sectional-hyperbolic is necessarily a transitive attractor, and hence, a homoclinic class. In [GYZ22], the same conclusion was obtained for chain-recurrence classes, not necessarily singular-hyperbolic, associated to  $C^1$ -generic vector fields away from homoclinic tangencies. In [CY21], S. Crovisier and D. Yang proved that transitivity is present in a robust way for sectional-hyperbolic Lyapunov stable chain-recurrence classes associated to vector fields  $X$  in a certain  $C^1$ -generic set. In light of this result, the authors posed the following question:

**Question:** Let  $M$  be a compact Riemannian manifold with dimension at least 4. Does there exist an open and dense set  $\mathcal{U} \subset \mathcal{X}^1(M)$  such that for any  $X \in \mathcal{U}$ , any non-trivial sectional-hyperbolic chain-recurrent class is robustly a homoclinic class?

In this work, we explore this question and some interesting implications. One important aspect of all of the aforementioned results is that they assert the non-existence of aperiodic classes under good generic conditions. Indeed, the existence of aperiodic classes represents a major obstacle in several important conjectures. See for instance the weak Palis conjecture which asserts that any  $C^1$ -system is approximated by either a Morse-Smale system or a system containing a hyperbolic horseshoe. This conjecture was verified for diffeomorphisms in [Cro10], for nonsingular flows in [XZ08] and for singular three-dimensional flows in [GY18].

On the other hand, in [CY21] and [GYZ22] the existence of periodic orbits for generic chain-recurrence classes was obtained under the assumption that the class is Lyapunov stable. Also, in [PY23] a dichotomy for chain-recurrence classes of generic star flows was obtained: either they have periodic orbits and positive entropy or they are sectional-hyperbolic aperiodic classes, and hence non-Lyapunov stable, with zero entropy. This evidences that obtaining periodic orbits for generic chain-recurrence classes is challenging, without Lyapunov stability. In this work, we shall see that in many cases, sectional-hyperbolicity is enough to rule out aperiodic classes.

We now state our main results. Let us begin by recalling the concept of sectional-hyperbolic set. We say that a compact invariant set  $\Lambda$  has a *dominated splitting* if there is a continuous invariant splitting  $T_\Lambda M = E \oplus F$  (with respect to the tangent flow  $\Phi_t = DX_t$ ,

$t \in \mathbb{R}$ ) and constants  $K, \lambda > 0$  satisfying the relation

$$\frac{\|\Phi_t(x)|_{E_x}\|}{m(\Phi_t(x)|_{F_x})} \leq K e^{-\lambda t}, \quad \forall x \in \Lambda, \forall t > 0,$$

where  $m(A)$  denotes the co-norm of a linear transformation  $A$ . In this case, we say that  $E$  is *dominated* by  $F$ . When the subbundle  $E$  is uniformly contracting, i.e.,  $\|\Phi_t(x)|_{E_x}\| \leq K e^{-\lambda t}$  for every  $t > 0$  and  $x \in \Lambda$ , we say that  $\Lambda$  is *partially hyperbolic*. In [PY21] it was defined the notion of *sectional-hyperbolic* set as follows:

**Definition 1.1.** A compact invariant set  $\Lambda \subset M$  is *sectional-hyperbolic* if it is partially hyperbolic and its central bundle  $F$  is *sectional-expanding*, i.e, there are  $K, \lambda > 0$  such that for every two-dimensional subspace  $L_x$  of  $F_x$  one has

$$|\det \Phi_t(x)|_{L_x}| \geq K e^{\lambda t}, \quad \forall x \in \Lambda, \forall t > 0.$$

**Remark 1.2.** The notion of sectional-hyperbolicity was first introduced in [MM08] to describe the dynamical behavior of higher-dimensional systems, such as the multidimensional Lorenz attractor (see [BPV97]). That definition differs from Definition 1.1 in that it assumes the hyperbolicity of singularities, a condition we do not require here. However, as we shall show in Lemma 2.2, every singularity contained in a sectional-hyperbolic chain-recurrent class (under Definition 1.1) is necessarily hyperbolic. Thus, this distinction poses no real restriction when dealing with chain-recurrence classes.

Next, let us clarify the notion of a *robustly transitive sectional-hyperbolic chain-recurrent class*. It is immediate to verify that  $\text{Crit}(X) \subset CR(X)$ , so the chain-recurrent class of any critical element is well defined. Moreover, by the previous remark and the hyperbolic lemma (see Lemma 2.1), every critical element of a sectional-hyperbolic chain-recurrent class is hyperbolic. If  $\gamma_X$  is a critical element of  $X$  and  $C(\gamma_X)$  is sectional-hyperbolic, then for every  $Y \in \mathcal{X}^1(M)$  sufficiently  $C^1$ -close to  $X$ , there exists a continuation  $\gamma_Y$  of  $\gamma_X$ . The corresponding chain-recurrent class  $C(\gamma_Y)$  is called the *continuation* of  $C(\gamma_X)$ . We say that the chain-recurrent class  $C(\gamma_X)$  is *robustly a homoclinic class* if there exists an open  $C^1$ -neighborhood  $\mathcal{U}$  of  $X$  such that for every  $Y \in \mathcal{U}$ , the continuation  $C(\gamma_Y)$  is a homoclinic class. The first main result of this paper is the following:

**Theorem A.** *There exists a  $C^1$ -generic set  $\mathcal{R} \subset \mathcal{X}^1(M)$  such that if  $\Lambda$  is a non-trivial sectional-hyperbolic chain-recurrent class, then it is robustly a homoclinic class.*

Notice that in the previous result we dropped out any stability assumption on  $\Lambda$ . In particular, Theorem A is a direct improvement of [PY21, Corollary E]. Also, an implicit fact behind the statement of Theorem A is that any non-trivial sectional-hyperbolic chain-recurrent set within  $\mathcal{R}$  must contain periodic orbits; in other words, sectional-hyperbolicity is enough to rule out the existence of aperiodic classes.

Another application of Theorem A concerns the setting of star flows. Recall that a vector field  $X \in \mathcal{X}^1(M)$  is *star* if there exists a neighborhood  $\mathcal{U}$  such that all critical elements of every  $Y \in \mathcal{U}$  are hyperbolic. Star systems play a central role in stability theory. For diffeomorphisms, it was proved in [Ma78, Li81] that  $\Omega$ -stability implies the star condition,

and later in [Ha92] that the star condition is equivalent to Axiom A with the no-cycle condition. For regular vector fields, the same equivalence was obtained in [GW06].

Beyond stability, singular star flows are known to exhibit rich dynamics, which has motivated substantial research. In [ZGW08], it was conjectured that  $C^1$ -generic star flows and sectional-hyperbolicity should be intrinsically related; namely, they were expected to have finitely many chain-recurrent classes, each sectional-hyperbolic. A partial confirmation was obtained in [SGW14] under the assumption of Lyapunov stability. However, [BdL21, dL17] disproved half of the conjecture by showing that  $C^1$ -generic star flows need not be sectional-hyperbolic, thus opening a new line of research with the introduction of multi-singular hyperbolic sets. Nevertheless, the other half of the conjecture remains open:

**Conjecture 1.3** (Conjecture 1.2 in [PYY23]).  *$C^1$ -generic singular star flows do not admit singular aperiodic classes. Consequently, they have only finitely many chain-recurrent classes, all of which are homoclinic classes of periodic orbits.*

A partial answer was given in [PYY23], proving that generic star flows admit finitely many Lyapunov chain-recurrent classes and establishing a dichotomy for the non-Lyapunov stable case, based on entropy. As a consequence of our techniques, we can fully confirm the conjecture:

**Theorem B.** *Let  $M$  be an  $n$ -dimensional manifold with  $n \geq 4$ . There exists a  $C^1$ -generic set  $\mathcal{R} \subset \mathcal{X}_*^1(M)$  such that for every  $X \in \mathcal{R}$  and every non-trivial chain-recurrent class  $\Lambda$  of  $X$ , the set  $\Lambda$  has positive topological entropy, contains a periodic orbit, and is isolated. Consequently,  $C^1$ -generic singular star flows have only finitely many chain-recurrent classes, all of which are either trivial or homoclinic classes of hyperbolic periodic orbits.*

The remainder of this text is organized as follows: In Section 2, we present a collection of preliminary results concerning sectional-hyperbolic chain-recurrence classes. In Section 3, we prove that any chain-recurrent class associated with  $X \in \mathcal{R}$  is robustly periodic, which constitutes a key step toward proving Theorem A. Finally, Section 4 is devoted to proving the main results of this article.

## 2. PRELIMINARY RESULTS ON SECTIONAL-HYPERBOLIC CHAIN-CLASSES

In this section, we collect some tools that will be used in our analysis. Let us begin by recalling that an important fact about chain-recurrence classes is that they are chain transitive: If  $C$  is a chain-recurrent class, then for any pair of points  $x, y \in C$  the point  $y$  is chain attainable from  $x$  through points in  $C$ , i.e., for every  $\varepsilon, T > 0$ , there is an  $(\varepsilon, T)$ -chain  $(x_i, t_i)_{i=0}^n$  such that  $x_0 = x, x_n = y$  and  $x_i \in C$  for any  $i = 0, \dots, n$  (see for instance [FS76, Proposition 2.1]).

Next, we recall the concept of hyperbolic set. A compact invariant set  $\Lambda$  is said to be a *hyperbolic set* if there is a  $\Phi_t$ -invariant splitting  $T_\Lambda M = E^s \oplus \langle X \rangle \oplus E^u$ , such that  $E^s$  is of contracting type,  $E^u$  is of expanding type and  $\langle X \rangle$  is the direction generated by  $X$ . A hyperbolic set is of *saddle type* if both  $E^s$  and  $E^u$  are non-trivial. A critical element of  $X$  is hyperbolic if its orbit is a hyperbolic set. Notice that when  $x$  is a singularity, the

subspace  $\langle X \rangle$  is trivial. It is well known that sectional-hyperbolic sets satisfy the so-called hyperbolic lemma, whose proof can be found in [BM11]:

**Lemma 2.1** (Hyperbolic Lemma). *Every compact invariant set without singularities contained in a sectional-hyperbolic set is hyperbolic of saddle type.*

The above lemma shows that this notion generalizes the concept of uniform hyperbolicity introduced by Smale in [Sm67]. Furthermore, it tells us that this notion is good enough to obtain interesting dynamical results.

In what follows, we prove some basic lemmas that will be instrumental in the proof of our main theorems. First, we prove the assertion made in Remark 1.2.

**Lemma 2.2.** *Let  $\Lambda$  be a non trivial sectional-hyperbolic chain-recurrent class for a  $C^1$  vector field  $X$  containing singularities. Then, every  $\sigma \in \Lambda$  is hyperbolic. Moreover, if  $T_\sigma M = E_\sigma^s \oplus E_\sigma^u$  is the hyperbolic splitting of  $\sigma$  and  $T_\sigma M = E_\sigma \oplus F_\sigma$  is the hyperbolic splitting of  $\sigma$  given by sectional-hyperbolicity, we have*

$$\dim E_\sigma + 1 = \dim E_\sigma^s.$$

*Proof.* Assume that there is a complex eigenvalue  $\lambda$  of  $DX(\sigma)$  with zero real part, and let  $L_\lambda = \text{span}\{v_1, v_2\}$  be the subspace spanned by the eigenvectors associated to  $\lambda$ . Then,  $L_\lambda$  is invariant for  $\Phi_t(\sigma)$  and there are  $a, b > 0$  such that  $a < \|\Phi_t(\sigma)v_i\| < b$ ,  $i = 1, 2$ , for every  $t > 0$ , so that  $L_\lambda \cap E_\sigma = \{0\}$ . Therefore, since  $E_\sigma$  is dominated by  $F_\sigma$ , we conclude that  $L_\lambda \subset F_\sigma$ , which contradicts the sectional expanding property of  $F$ . This shows that real part of any eigenvalue of  $DX(\sigma)$  must be non-zero.

Now let consider the hyperbolic splitting  $T_\sigma M = E_\sigma^s \oplus E_\sigma^u$  of  $\sigma$ . Note that each one of this subspaces is non trivial, otherwise we have that  $\sigma$  is either a source or a sink, so that  $\Lambda$  would be trivial which is impossible. Besides, since vectors in  $E(\sigma)$  contract uniformly, we see by the domination property that  $E \subseteq E_\sigma^s$  and  $E_\sigma^u \subseteq F$ .

We claim that  $E \neq E_\sigma^s$ . Indeed, since  $\Lambda$  is invariant and the splitting of  $T_\Lambda M$  is dominated, it follows that the flow direction  $X(x)$  belongs to  $F_x$  for any regular point  $x \in \Lambda$ . Indeed, without loss of generality, we can assume that the Riemannian metric left the subbundles  $E$  and  $F$  orthogonal. In this way, for  $x \in \Lambda$  we write  $X(x) = v_E + v_F$ , with  $v_E \in E_x$  and  $v_F \in F_x$ . Assume that  $X(x) \notin F_x$  for some  $x \in \Lambda$ , then  $v_E \neq 0$ . Since  $E$  and  $E^c$  are  $\Phi_t$ -invariant and orthogonal we have that for every  $t \in \mathbb{R}$  that

$$\begin{aligned} K &\geq \|X(X_t(x))\|^2 \\ &= \|\Phi_t(x)X(x)\|^2 \\ &= \|\Phi_t(x)v_E\|^2 + \|\Phi_t v_F\|^2 \\ &\geq \|\Phi_t(x)v_E\|^2 \geq \frac{1}{K} e^{\lambda t} \|v_E\| \rightarrow \infty, \quad t \rightarrow -\infty. \end{aligned}$$

which is a contradiction.

On the other hand, since  $\Lambda$  is partially-hyperbolic it is well known that any point  $x \in \Lambda$  has associated its strong stable manifold  $\mathcal{F}_{ss}(x)$  which is invariant and it is tangent to  $E$  at  $x$  (see [HPPS70] for instance). Thus, if  $x \in \Lambda \cap (\mathcal{F}_{ss}(\sigma) \setminus \{\sigma\})$ , then  $X(x) \in T_x \mathcal{F}_{ss}(\sigma) = E_x^s$ ,

which is absurd. Therefore, if  $E = E_\sigma^s$  we have that  $W^s(\sigma) = \mathcal{F}_{ss}(\sigma)$ , so that

$$\Lambda \cap (W^s(\sigma) \setminus \{\sigma\}) = \Lambda \cap (\mathcal{F}_{ss}(\sigma) \setminus \{\sigma\}) = \emptyset,$$

which is a contradiction because  $\Lambda$  is a non trivial chain-recurrent class. So, by domination property we see that  $E_\sigma^s \cap F_\sigma \neq \{0\}$ . Hence, since  $F$  is area expanding, it follows that  $\dim(E_\sigma^s \cap F_\sigma) = 1$ , which proves the result.  $\square$

**Remark 2.3.** A singularity  $\sigma$  satisfying the conditions of above lemma is called *Lorenz-like singularity*. Moreover, the conditions  $\dim E_\sigma^s \geq 2$  and  $\dim(E_\sigma^s \cap F_\sigma) = 1$  provide an alternative definition of Lorenz-like singularity: A singularity  $\sigma$  for  $X$  is Lorenz-like if there are at least two eigenvalues with negative real part, one of them is real, say  $\lambda_-$ , and the real part of remaining eigenvalues are outside the interval  $[\lambda_-, -\lambda_-]$ .

Notice that by Lemma 2.2 and Remark 2.3, we can consider a useful tool for proving Theorem A, coming from [PY23]. More precisely, let  $E^{cs} \oplus E^u = E_\sigma^s \oplus F_\sigma$ , where  $E^{cs} = E_\sigma \oplus G$  and  $F_\sigma = G \oplus E_\sigma^u$ , with  $G \subset F$  and  $\dim G = 1$ , be the hyperbolic splitting of  $T_\sigma M$ . Up to a change of coordinates, we think  $\sigma$  as the origin in  $\mathbb{R}^n$ , and the subspaces  $E^*$ ,  $* = cs, u$ , are perpendicular. As in the aforementioned reference, for a small neighborhood  $U_\sigma$  of  $\sigma$  and  $x \in U_\sigma$ , we denote by  $x^{cs}$  and  $x^u$  its distances to  $E^u$  and  $E^{cs}$  respectively. In this way, for  $\alpha, \beta > 0$  define the  *$\alpha$ -cone on the manifold*  $D_\alpha^{cs}(\sigma)$  as

$$D_\alpha^{cs}(\sigma) = \{x \in U : x^u < \alpha x^{cs}\}.$$

Now, for a non singular point  $x \in M$ , the *orthogonal complement*  $N_x$  of  $X(x)$  is defined by

$$N_x = \{v \in T_x M : \langle v, X(x) \rangle = 0\}.$$

In this way, the *linear Poincaré flow*  $\psi_t$  on the normal bundle  $N_{M \setminus Sing(X)}$  is given by

$$\psi_t(v) = \pi_{N_{X_t(x)}}(\Phi_t(x)v), \quad \forall x \in M \setminus Sing(X), v \in N_x,$$

where  $\pi_{N_z}(w)$  is the orthogonal projection of  $w$  on  $N_z$ . The *scaled linear Poincaré flow*, denoted by  $\psi_t^*$ , is defined as the normalization of  $\psi_t$  by the flow speed, i.e.,

$$\psi_t^*(v) = \frac{\|X(x)\|}{\|X(X_t(x))\|} \psi_t(v).$$

In [BM11] it is showed that a partially hyperbolic splitting  $T_\Lambda M = E \oplus F$  of a compact invariant set  $\Lambda$  induces a partially hyperbolic splitting  $T_x N_x = \mathcal{E}_x \oplus \mathcal{F}_x$  of the normal subspace  $N_x$  for any  $x \in \Lambda \setminus Sing(X)$ , where  $\mathcal{E}_x = \mathcal{F}_x = \pi_x(E_x)$  for every  $x \in \Lambda$ . Indeed, since  $F_x$  contains the flow direction, it follows that  $\pi(v) \in F_x$  for any  $v \neq X(x)$  in  $F_x$ , hence  $\mathcal{F}_x \subset F_x$  because  $\pi$  is linear. So, since the splitting  $E \oplus F$  is dominated for  $\Phi_t$ , we deduce the dominance of the splitting  $\mathcal{E} \oplus \mathcal{F}$ . The uniform contracting property for  $\pi(E)$  follows because  $\|\pi(v)\| \leq \|v\|$  for every  $v \in E$ . It is important to observe that the above property remains valid for the rescaled linear Poincaré flow  $\psi_t^*$ .

Recall the notion of  $(\eta, T, E)$ -contracting points.

**Definition 2.4.** Let  $X \in X^1(M)$ ,  $\Lambda$  a compact invariant set of  $X$ , and  $\mathcal{E} \subset N_{\Lambda \setminus Sing(X)}$  an invariant bundle of the rescaled linear Poincaré flow  $\psi_t^*$ . For  $0 < \eta < 1$  and  $T_0 > 0$ , we say that a point  $x \in \Lambda \setminus Sing(X)$  is  $(\eta, T_0)$ -contracting for the bundle  $\mathcal{E}$  if there is a partition  $0 = t_0 < t_1 < t_2 < \dots < t_k < \dots$ , where  $t_{i+1} - t_i \in [T_0, 2T_0]$  for every  $i = 0, 1, \dots$  such that

$$\prod_{i=0}^{k-1} \|\psi_{t_{i+1}-t_i}^*|_{\mathcal{E}(X_{t_i}(x))}\| \leq \eta^k, \quad \forall k \geq 1.$$

Similarly,  $x \in \Lambda \setminus Sing(X)$  is said to be  $(\eta, T_0)$ -expanding for the bundle  $\mathcal{F}$  if it is  $(\eta, T_0)$ -contracting for  $-\mathcal{F}$ .

We are now ready to state the result mentioned above.

**Lemma 2.5** (Lemma 3.11 in [PY23]). *Let  $\sigma$  be a Lorenz-like singularity. Then for  $r > 0$  small enough, for every  $\lambda \in (0, 1)$  and  $T_0 > 0$  there exists  $\alpha_2 > 0$  such that for all  $\alpha < \alpha_2$ , if  $y \in B_r(\sigma) \cap \bar{D}_\alpha^{cs}(\sigma)$  is a  $(\lambda, T_0)$ -expanding time on its orbit, then  $W_{loc}^u(y) \pitchfork W_{loc}^{cs}(\sigma) \neq \emptyset$ .*

Next, we recall a remarkable theorem due to Liao regarding the existence of stable (unstable) manifolds for contracting (expanding) points.

**Theorem 2.6.** *Let  $X \in X^1(M)$ ,  $\Lambda$  a compact invariant set of  $X$ . Given  $\eta \in (0, 1)$ ,  $T_0 > 0$ , there is  $\delta = \delta(\eta, T_0) > 0$  such that if  $x$  is  $(\eta, T)$ -contracting for the bundle  $E \subset N$ , then there is a  $C^1$  map  $\kappa : E_x(\delta \|X(x)\|) \rightarrow N_x$  such that*

- $d_{C^1}(\kappa, id) < \varepsilon$ .
- $\kappa(0) = 0$ .
- $W_{\delta \|X(x)\|}^s(x) \subset W^s(\mathcal{O}(x))$ , where  $W_{\delta \|X(x)\|}^s(x) = \exp_x(\text{Image}(\kappa))$ .

A similar result holds for  $(\eta, T_0)$ -expanding points for the bundle  $F \subset N$ .

Finally, we end this subsection by showing that sectional-hyperbolicity of a compact invariant set can be extended to the maximal invariant set of a neighborhood of it.

**Lemma 2.7.** *Let  $\Lambda$  be a sectional-hyperbolic set for a  $C^1$  vector field  $X$ . Then, there is a neighborhood  $U$  of  $\Lambda$  such that the maximal invariant set  $\tilde{\Lambda}$  in  $U$  is also sectional-hyperbolic for  $X$ .*

*Proof.* The proof of this result is based in the proof of Propositions 3.1 and 3.2 in [AM17] and Proposition 2.10 in [AM19]. First, there is a neighborhood  $U_0$  of  $\Lambda$  and a continuous (not necessarily invariant) extension  $T_{U_0}M = \tilde{E} \oplus \tilde{F}$  of the splitting  $T_\Lambda M = E \oplus F$ . Moreover, if  $\Lambda'$  is the maximal invariant set for the flow of  $X$  contained in  $U_0$ , there are constants  $a, T > 0$  such that for every  $x \in \Lambda'$  the cone fields

$$\mathcal{C}_x^s(a) = \{v = v^{\tilde{E}} + v^{\tilde{F}} \in \tilde{E} \oplus \tilde{F} : \|v^{\tilde{F}}\| \leq a \|v^{\tilde{E}}\|\}$$

and

$$\mathcal{C}_x^c(a) = \{v = v^{\tilde{E}} + v^{\tilde{F}} \in \tilde{E} \oplus \tilde{F} : \|v^{\tilde{E}}\| \leq a \|v^{\tilde{F}}\|\}$$

satisfy the following conditions:

(1) Invariance:

$$\Phi_{-t}(\mathcal{C}_{\phi_t(x)}^s(a)) \subset \mathcal{C}_x^s(a) \text{ and } \Phi_t(\mathcal{C}_x^c(a)) \subset \mathcal{C}_{\phi_t(x)}^c(a), \quad \forall t \geq T.$$

(2) Domination: There are constants  $c > 0$  and  $\lambda_0 \in (0, 1)$  such that

$$\frac{\|\Phi_t(x)u\|}{\|\Phi_t(x)v\|} \leq c\lambda_0^t \frac{\|u\|}{\|v\|}, \quad \forall t > 0,$$

for every nonzero vectors  $v \in \mathcal{C}_x^c(a)$  and  $u \in \Phi_{-t}(\mathcal{C}_{\phi_t(x)}^s(a))$ .

Now, for  $x \in \tilde{\Lambda}$  define the bundles

$$E^s = \bigcap_{t \geq 0} \Phi_{-t}(\mathcal{C}_{X_t(x)}^s(a)) \text{ and } E^c = \bigcap_{t \geq 0} \Phi_t(\mathcal{C}_{X_{-t}(x)}^c(a)).$$

By Proposition 3.2 in [AM17] one has that  $E^s$  is a  $\Phi_t$ -invariant and uniformly contracting continuous bundle satisfying  $E_x^s \subset \mathcal{C}_x^s(a)$  for every  $x \in \Lambda'$  and  $E_x^s = E_x$  for every  $x \in \Lambda$ . Besides, since  $\Phi_t(\mathcal{C}_{X_{-t}(x)}^c(a)) \subset \mathcal{C}_x^c(a)$  for every  $t \geq T$  we have that  $E_x^c \subset \mathcal{C}_x^c(a)$ . So, by construction of the bundles and domination property, we have that  $T_{\Lambda}M = E^s \oplus E^c$  is a dominated splitting for  $\Lambda'$ .

**Claim:**  $E^c$  is a  $\Phi_t$ -invariant sectional expanding continuous bundle such that  $E_x^c = F_x$  for every  $x \in \Lambda$ .

On one hand, let's see that  $E_x^c$  is an element of the space  $\mathcal{G}_x$  of  $d_F$ -dimensional subspaces of  $T_x M$ . First, since  $\Lambda'$  is compact, the space  $\mathcal{G}_x$  is compact, so that the sequence  $\{\Phi_t(\tilde{F}_{X_{-t}(x)})\}_{t>0} \subset \mathcal{G}_x$  has a subsequence  $\Phi_{t_n}(\tilde{F}_{X_{-t_n}(x)})$ ,  $t_n \geq T$ , whose limit  $G_x$  belongs to  $\mathcal{G}_x$ . Second, since  $\Phi_t(\mathcal{C}_{X_{-t}(x)}^c(a))$  is a nested family of subspaces of  $T_x M$ , by definition of  $E^c$  we have that  $E^c = \lim_{t \rightarrow \infty} \Phi_t(\mathcal{C}_{X_{-t}(x)}^c(a))$ . Then, as  $\tilde{F}_x \subset \mathcal{C}_x^c(a)$ , it follows that  $G_x \subset E_x^c$ . In fact,  $E_x^c = G_x$  for every  $x \in \Lambda'$ . Indeed, assume that  $E_x^c \neq G_x$  for some  $x \in \Lambda'$ , so that  $E_x^c$  is a non-trivial cone. Let  $u_n \in \tilde{F}_{X_{-t_n}(x)}$ ,  $n \geq 1$ , such that  $\|\Phi_{t_n}(X_{-t_n}(x))u_n\| = 1$ . So, there is a nonzero vector  $v \in E_x^s$  such that  $w_n = \Phi_{t_n}(X_{-t_n}(x))u_n + v \in E_x^c$  for every  $n \geq 1$ , so that  $\Phi_{-t_n}(x)w_n = u_n + \Phi_{-t_n}(x)v \in \mathcal{C}_{X_{-t_n}(x)}^c(a)$  by definition of  $E_x^c$ . Hence,

$$\|\Phi_{-t_n}(x)\tilde{E}v\| \leq a(1 - a^2)^{-1}\|u_n\|.$$

Then, by domination and the definition of  $u_n$ ,

$$\begin{aligned} c^{-1}(\lambda_0)^{-t_n}\|u_n\|\|v\| &\leq \|\Phi_{-t_n}(x)v\| \leq (1 + a)\|\Phi_{-t_n}(x)\tilde{E}v\| \\ &\leq a(1 + a)(1 - a^2)^{-1}\|u_n\|, \end{aligned}$$

which implies that

$$\|v\| \leq c(\lambda_0)^{t_n}a(1 + a)(1 - a^2)^{-1} \rightarrow 0, \quad n \rightarrow \infty.$$

This contradicts the choice of  $v$ .

On the other hand, since  $F$  is invariant on  $\Lambda$  it follows from definition of  $E^c$  that  $E_x^c = F_x$  for every  $x \in \Lambda$ . By continuity of  $\Phi_t$  and the definition of  $E^c$  we have that  $E^c$  is continuous and invariant.

Finally, we show the sectional expanding property of  $E^c$ . Let  $T_0 > 0$  such that  $\lambda' = \ln((1/2)Ke^{T_0}) > 0$ , where  $K$  is given by sectional-hyperbolicity of the bundle  $F$ . By continuity of  $E^c$  and  $\Phi_t$  for  $x \in \Lambda$  there is a neighborhood  $V_x \subset G(2, F)$ , where  $G(2, F)$  is the Grassmannian bundle of two-planes contained in  $E^c$ , of  $(x, L_x)$  such that for any  $(y, L_y) \in V_x$ ,

$$|\det \Phi_t(x)|_{L_y}| \geq \frac{1}{2} |\det \Phi_t(x)|_{L_x}| \geq Ke^{\lambda t}, \quad \forall t \in [0, T_0],$$

where  $K = C/2 > 0$ . So, by compactness of  $\Lambda$  and  $G(2, F)$ , and by shrinking  $U_0$  if it is necessary, we have that if  $y \in \Lambda'$  and  $L_y \subset E_y^c$ , there are  $x \in \Lambda$  and a two-dimensional subspace  $L_x \subset F_x$  such that

$$(1) \quad |\det \Phi_t(y)|_{L_y}| \geq Ke^{\lambda t}, \quad \forall t \in [0, T_0].$$

So, if  $t > 0$ , we write  $t = mT_0 + r$ , where  $0 \leq r < T_0$ , and, by applying (1) repeatedly, we obtain that

$$|\det \Phi_t(y)|_{L_y}| \geq K^{1+\frac{t}{T_0}} e^{\lambda t} = Ke^{\lambda' t}.$$

Since  $L_y$  was chosen arbitrarily, we get the desired result.  $\square$

### 3. ROBUSTLY PERIODIC CHAIN-RECURRENT CLASSES

In this section, we shall derive one of the main results of this paper. Namely, we prove that  $C^1$ -generically every sectional-hyperbolic non-trivial chain-recurrent class robustly periodic. The statement is as follows:

**Theorem 3.1.** *There exists a  $C^1$ -generic set  $\mathcal{R} \subset \mathcal{X}^1(M)$  such that if  $\Lambda$  is a non-trivial sectional-hyperbolic chain-recurrent class then it is robustly periodic, that is, there is an open neighborhood  $\mathcal{U}$  of  $X$  such that for every  $Y \in \mathcal{U}$ , one has  $\text{Per}(Y|_{\Lambda_Y}) \neq \emptyset$ , where  $\Lambda_Y$  denotes the continuation of the chain recurrent class  $\Lambda$  for  $X$ .*

In order to prove Theorem 3.1, we shall show first that  $C^1$ -generically every sectional-hyperbolic non-trivial chain-recurrent class actually contains a periodic orbit, and therefore it is a homoclinic class. As a direct consequence, it has positive topological entropy.

Recall that Hausdorff distance between two compact subsets  $A, B$  of a metric space  $(M, d)$  is defined as

$$d_H(A, B) = \max \left\{ \sup_{x \in A} d(x, B), \sup_{y \in B} d(y, A) \right\}, \quad d(a, C) = \inf_{c \in C} d(a, c).$$

Denote by  $\mathcal{K}(M)$  the set of compact sets of  $M$ . In this case,  $(\mathcal{K}(M), d_H)$  is a metric space.

Now, recall that for a hyperbolic periodic point  $p$  associated to a  $C^1$  vector field  $X$ , with splitting  $T_{\mathcal{O}(p)}M = E^s \oplus \langle X \rangle \oplus E^u$ , the strong stable and strong unstable manifolds of  $p$  are defined, respectively, by

$$W^{ss}(p) = \{y \in M; \lim_{t \rightarrow \infty} d(X_t(p), X_t(y)) = 0\},$$

and

$$W^{uu}(p) = \{y \in M; \lim_{t \rightarrow -\infty} d(X_t(p), X_t(y)) = 0\}.$$

Then, the stable and unstable manifolds of  $p$ , respectively, are given by

$$W^s(p) = \bigcup_{t \in \mathbb{R}} W^{ss}(X_t(p)) \text{ and } W^u(p) = \bigcup_{t \in \mathbb{R}} W^{uu}(X_t(p)).$$

These manifolds are tangent to  $E^s$ ,  $E^u$ ,  $E^s \oplus \langle X \rangle$ , and  $E^u \oplus \langle X \rangle$ , respectively, at each one of the points of  $\mathcal{O}(p)$ . Thus, the *homoclinic class* of  $p$ , denoted by  $H(p)$ , is defined as:

$$H(p) := \overline{W^s(p) \pitchfork W^u(p)}.$$

We say that the homoclinic class  $H(p)$  is *non trivial* if it is not reduced to  $\mathcal{O}(p)$ .

Next, consider the following generic set that is obtained by combining some items from [CY21, Proposition 2.1] and [GYZ22, Lemma 3.12].

**Lemma 3.2.** *There is a residual subset  $\mathcal{R} \subset \mathcal{X}^1(M)$  such that for every  $X \in \mathcal{R}$  it holds:*

- (1) *Any non-trivial chain-recurrent class  $C$  of  $X$  containing a hyperbolic periodic orbit  $\gamma$  is a homoclinic class.*
- (2) *For every non trivial compact chain-transitive set  $C$  of  $X$ , there is a sequence of periodic orbits  $\gamma_n$  of  $X$  such that  $\gamma_n$  converges to  $C$  in the Hausdorff topology.*
- (3) *There is a  $C^1$ -open neighborhood  $\mathcal{U}_X$ , if  $C(\sigma)$  is a non-trivial chain-recurrent class of  $X$  such, then the map  $Y \in \mathcal{U}_X \rightarrow C(\sigma_Y)$  is continuous.*

The generic property given in the second item of Lemma 3.2 was not under consideration in the proof of Theorem A in [CY21]. It will be the key to eliminating the assumption of Lyapunov stability in our proofs.

We now present the following lemma, which will be used to prove Theorem 3.1.

**Lemma 3.3.** *Let  $\mathcal{R}$  be the residual set of Lemma 3.2. Let  $X \in \mathcal{R}$ ,  $\sigma_X \in \text{Sing}(X)$  and  $C(\sigma)$  be a non-trivial sectional-hyperbolic chain-recurrent class. There is a neighborhood  $\mathcal{U}_0 \subset \mathcal{U}_X$  of  $X$  such that for every  $Y \in \mathcal{U}_0$ , the continuation  $C(\sigma_Y)$  is sectional-hyperbolic.*

*Proof.* Let  $X \in \mathcal{R}$  and  $\Lambda$  a non-trivial sectional-hyperbolic class of  $X$ . Recall that sectional-hyperbolicity is a robust property, i.e., there are neighborhoods  $U$  of  $\Lambda$  and  $\mathcal{U}'_X$  of  $X$  such that for every  $Y \in \mathcal{U}'_X$  the set  $\cap_{t \in \mathbb{R}} Y_t(U)$  is sectional-hyperbolic. Let  $\mathcal{U}_X$  be given by item 3 of Lemma 3.2. Denote  $\mathcal{U}_0 = \mathcal{U}_X \cap \mathcal{U}'_X$ . Since the map  $Y \in \mathcal{U}_X \rightarrow C(\sigma)$  is continuous, after possibly shrinking  $\mathcal{U}_X$ , we can assume  $C(\sigma_Y) \subset U$ . Therefore,  $C(\sigma_Y)$  is contained in the maximal invariant set of  $U$  under  $Y$ . In particular,  $C(\sigma_Y)$  is sectional-hyperbolic.  $\square$

**Remark 3.4.** By the above lemma, we see that Lemma 2.2 holds for every  $Y \in \mathcal{U}_0$ .

Next, we present the aforementioned result.

**Lemma 3.5.** *If  $X \in \mathcal{R}$  and  $\Lambda \subset M$  is a non-trivial sectional-hyperbolic chain-recurrent class, then it is a homoclinic class.*

*Proof.* To begin with, let  $U$  be the neighborhood of  $\Lambda$  given by Lemma 2.7. By Lemma 3.2 there is a sequence of periodic orbits of  $\gamma_n \subset M$  converging to  $\Lambda$  in the Hausdorff metric. So, we can assume  $\gamma_n \subset U$  for every  $n \geq 1$ . Denote

$$\Lambda' = \Lambda \cup \bigcup_{n \geq 1} \gamma_n.$$

**Claim 1.**  $\Lambda'$  is a compact and invariant subset of  $U$ .

*Proof.* First, observe that  $\Lambda' \subset U$  and  $\Lambda'$  is invariant by definition. Next, let  $\{x_k\}_{k \geq 1} \subset \Lambda'$  be a sequence such that  $x_k \rightarrow x \in M$ , when  $k \rightarrow +\infty$ . If  $x_k \in \Lambda$  for sufficiently large  $k$ , then  $x \in \Lambda$  since  $\Lambda$  is compact. Similarly, if  $x_k \subset \gamma_n$  for sufficiently large  $k$  and some  $n \geq 1$  fixed, then  $x \in \gamma_n \subset \Lambda'$ . It remains to verify the case where there are sequences  $k_l, n_l \rightarrow \infty$ ,  $n_l \neq n_{l'}$ , for  $l \neq l'$  and  $x_{k_l} \subset \gamma_{n_l}$ . In this case, since  $d^H(\gamma_n, \Lambda) \rightarrow 0$ , we have  $d(x_{k_l}, \Lambda) \rightarrow 0$ . Hence,  $x \in \Lambda \subset \Lambda'$  and the claim holds.  $\square$

Notice that by Lemma 2.7 the set  $\Lambda'$  is sectional hyperbolic. Now, we consider the following cases:

**Case 1:**  $\Lambda$  does not contain singularities

In this case,  $\Lambda$  is hyperbolic by Lemma 2.1. We could now be tempted to conclude the existence of a periodic orbit contained in  $\Lambda$  via the Shadowing lemma. Nevertheless, we cannot follow this approach, because we do not know whether  $\Lambda$  is isolated or not, and this may prevent the periodic orbits given by the shadowing lemma from being contained in  $\Lambda$ . Instead, we shall use Lemma 3.2 to derive our conclusion.

First, let  $U$  be the neighborhood given by Lemma 2.7 and let  $\Lambda'$  be defined as above. Since  $\Lambda' \subset U$ , Claim 1 and the hyperbolic lemma imply that  $\Lambda'$  is a compact invariant hyperbolic set. Let  $T\Lambda' = E'^s \oplus \langle X \rangle \oplus E'^u$  be the hyperbolic splitting of  $\Lambda'$ . It is well known that dominated splittings vary continuously and hence  $\dim(E'^s)$  and  $\dim(E'^u)$  are constant along  $\Lambda'$ , because  $\Lambda'$  is a chain transitive set. Moreover, up to possibly reducing  $U$ , we can assume  $\dim(E'^s)$  and  $\dim(E'^u)$  are constant along  $\Lambda'$  because  $d^H(\gamma_n, \Lambda) \rightarrow 0$ . Also, the continuity of the splitting and the compactness of  $\Lambda'$  implies that the angles between the fibers  $E'^s$ ,  $\langle X \rangle$  and  $E'^u$  are uniformly limited away from zero.

Since  $\Lambda'$  is hyperbolic,  $W_{loc}^s(x)$  and  $W_{loc}^u(x)$  are immersed manifolds, respectively tangent to  $E'^s \oplus \langle X \rangle$  and  $\langle X \rangle \oplus E'^u$ , for any  $x \in \Lambda'$ . Moreover, those manifolds have a uniform size. For each  $n \geq 1$ , let  $p_n$  be a point of  $\gamma_n$ . By possibly taking a subsequence, we can assume that  $p_n \rightarrow x \in \Lambda$ . By the uniform size of the local stable and unstable manifolds, the constant dimension of the invariant subspaces and the angle uniformly bounded away from zero, we get

$$W^u(p_n) \pitchfork W^s(x) \neq \emptyset \text{ and } W^s(p_n) \pitchfork W^u(x) \neq \emptyset,$$

for  $n$  sufficiently large. Consequently,  $\Lambda$  contains a periodic orbit.

**Case 2:**  $\Lambda$  contains singularities

Now we assume that  $\Lambda = C(\sigma_X)$ , for some singularity  $\sigma \in \Lambda$ . We shall proceed similarly to Case 1, but here the proof is more delicate, since  $C(\sigma_X)$  is not hyperbolic. Again, consider the neighborhood  $U$  from Lemma 2.7 and  $\Lambda'$ . Thus,  $\Lambda'$  is sectional hyperbolic. Let  $T\Lambda' = E \oplus F$  be the sectional-hyperbolic splitting of  $\Lambda'$ . Similarly to Case 1, the continuity of the splitting implies that  $\dim(E)$  and  $\dim(F)$  are constant along  $\Lambda'$  and that the angles between  $E$  and  $F$  are uniformly bounded away from zero. Since  $E$  is uniformly

contracted, by the classical stable manifold theory, for every  $x \in \Lambda'$  there is a well defined local stable manifold  $W_{loc}^{ss}(x)$  with uniform size along  $\Lambda'$ .

In contrast with Case 1, since  $\Lambda'$  is not hyperbolic, there may not be well defined for local unstable manifolds for every point in  $\Lambda'$ . Moreover, even though these local unstable manifolds may exist for some points (e.g. along non-singular compact and invariant subsets) they do not need to have uniform size. To overcome this issue, we firstly recall that the periodic orbits  $\gamma_n$  are hyperbolic by the hyperbolic lemma. So, fix  $\gamma_n$  and let

$$T\gamma_n = E'^s \oplus \langle X \rangle \oplus E'^u$$

be the hyperbolic splitting of  $\gamma_n$ . Since  $E$  is uniformly contracted, we have  $E(x) \subset E'^s(x)$ , for every  $x \in \gamma_n$ . Now, suppose there is  $x \in \gamma_n$  so that  $E'^s(x) \not\subset E(x)$ . In this case, there is a vector  $v \in E'^s(x)$  such that  $v = v^E + v^F$ ,  $v^E \in E(x)$ ,  $v^F \in F(x)$  and  $v^F \neq 0$ . In particular  $v^F \subset E(x)$ . Recall that  $\langle X(x) \rangle \subset F$  for any  $x \in \gamma_n$ . Since  $\gamma_n \cap Sing(X) = \emptyset$ , there are  $A_2 > A_1 > 0$  such that  $A_1 \leq X(x) \leq A_2$ , for any  $x \in \gamma_n$ . But this implies that the flow direction along  $\gamma_n$  can neither be uniformly contracted or expanded by  $DX_t$ . Consequently,  $v^F \not\subset \langle X(x) \rangle$ . On the other hand, by considering the subspace  $L \subset F$  generated by  $v^F$  and  $\langle X(x) \rangle$ , the sectional-expansion of  $L$  implies,  $v^F \notin E'^s(x)$ . This contradiction implies  $E(x) = E'^s(x)$ , for every  $x \in \gamma_n$ . Consequently,

$$\langle X(x) \rangle \oplus E'^u(x) = F(x),$$

for any  $x \in \gamma_n$ .

Let  $r > 0$  be such that  $B_r(\sigma_X) \cap Sing(X) = \{\sigma_X\}$ . Since any singularity contained in  $C(\sigma_X)$  is hyperbolic by Lemma 2.2, there are only finitely many of them. By shrinking  $U$  if it is necessary, we have that  $Sing(X|_U) = Sing(X|_{C(\sigma_X)})$  and hence, there is a neighborhood  $V$  of  $Sing(X|_{C(\sigma_X)})$  contained in  $U$  satisfying

$$W_{loc}^s(\sigma) \not\subset V \text{ and } W_{loc}^u(\sigma) \not\subset V.$$

Moreover, since  $Sing(X|_V)$  is finite, we can assume there is a family of pairwise disjoint open sets  $V_1, \dots, V_j$  such that:

- $V = V_1 \cup \dots \cup V_j$ .
- Each  $V_i$  is a neighborhood of a unique singularity in  $X|_U$ .
- $V_i \subset B_r(\sigma_X)$ , for any  $i = 1, \dots, j$ .
- There is  $0 < L_0 < 1$  such that  $\|X(z)\| < L_0$ , for every  $z \in V$ .

Recall we are assuming  $C(\sigma_X)$  is non-trivial. Since  $\gamma_n$  converges to  $C(\sigma_X)$  in the Hausdorff topology, then we can assume that each  $\gamma_n$  contains a point  $p_n$  so that  $p_n \rightarrow \sigma_X$ . So, by a classical Grobman-Hartman argument, there is a sequence of times  $t_n^+ > 0$  so that, by denoting  $p_n^s = \phi_{t_n^+}(p_n)$ , we can assume, by possibly taking subsequences, that

$$p_n^s \rightarrow p^u,$$

where  $p^u \in W^u(\sigma_X)$ . By possibly reducing  $V$ , we can assume  $p^u \in B_r(\sigma_X) \setminus V$ . On the other hand, since  $d^H(C(\sigma_X), \gamma_n) \rightarrow 0$ , there is a sequence of regular points  $x_n^u \in C(\sigma_X)$  such that  $d(x_n^u, p_n^s) \rightarrow 0$ . Then,  $x_n^u \rightarrow p^u$ . Hence,  $p^u \in C(\sigma_X) \setminus Sing(X|_{C(\sigma_X)})$ , because  $C(\sigma_X)$  is closed.

Let

$$K_0 = \inf\{|X(x)| : x \in U \setminus V\},$$

and notice that  $K_0 > 0$ . Let  $K_1 > 0$  be such that  $K_0 < |X(x)| < K_1$ , for every  $x \in U \setminus V$ . Let  $K, \lambda > 0$  the constants given by sectional hyperbolicity of  $\Lambda'$ , and let  $T_0 > 0$  in a such way that

$$(2) \quad K_2 e^{\lambda t} \geq e^{\lambda' t} \text{ and } K_3^2 e^{\lambda' t} \geq e^{\lambda'' t}, \quad \forall t \geq T_0,$$

where

$$K_2 = \min \left\{ K, \frac{K}{K_1} \right\} > 0, \quad K_3 = \min \left\{ \frac{K_0}{L_0}, \frac{K_0}{K_1} \right\} > 0 \text{ and } 0 < \lambda'' < \lambda' < \lambda.$$

Since  $p_n \rightarrow \sigma$ , one has  $\tau_n \rightarrow \infty$ , where  $\tau_n$  is the period of  $\gamma_n$ . In particular, there is a natural number  $k_0 > 0$  so that such that  $\tau_n > T_0$ , for every  $n \geq N_1$ .

Now, let  $\eta = (e^{\lambda'' T_0})^{-1} \in (0, 1)$  and let  $0 = t_0 < t_1 < \dots < t_m < \dots$  such that  $t_{i+1} - t_i = T_0$  for every  $i = 0, \dots, k-1$ . On one hand, notice that for every unitary vector  $v_i \in \mathcal{F}_{X_{t_i}(p_n^s)}$  one has

$$\begin{aligned} K e^{\lambda(t_{i+1} - t_i)} &\leq |\det \Phi_{t_{i+1} - t_i}(X_{t_i}(p_n^s))| \\ &= \frac{\|\Phi_{t_{i+1} - t_i}(X_{t_i}(p_n^s))\| \cdot \|X(X_{t_{i+1}}(p_n^s))\| \cdot \sin(\theta_t)}{\|X(X_{t_i}(p_n^s))\|} \\ &= \frac{\|X(X_{t_{i+1}}(p_n^s))\|}{\|X(X_{t_i}(p_n^s))\|} \cdot \|\psi_{t_{i+1} - t_i}(X_{t_i}(p_n^s))v_i\|, \end{aligned}$$

where  $\theta_t = \angle(\Phi_{t_{i+1} - t_i}(X_{t_i}(p_n^s)), X(X_{t_{i+1}}(p_n^s)))$ , so that by the choice of  $K_2$  in (2),

$$\begin{aligned} \|\psi_{t_{i+1} - t_i}^*(X_{t_i}(p_n^s))|_{\mathcal{F}_{X_{t_i}(p_n^s)}}\| &\geq \|\psi_{t_{i+1} - t_i}^*(X_{t_i}(p_n^s))v_i\| \\ &= \left[ \frac{\|X(X_{t_i}(p_n^s))\|}{\|X(X_{t_{i+1}}(p_n^s))\|} \right]^2 |\det \Phi_{t_{i+1} - t_i}(X_{t_i}(p_n^s))| \\ &\geq \left[ \frac{\|X(X_{t_i}(p_n^s))\|}{\|X(X_{t_{i+1}}(p_n^s))\|} \right]^2 e^{\lambda'(t_{i+1} - t_i)}, \quad \forall i = 0, \dots, k-1. \end{aligned}$$

Thus, since  $X_{t_k}(p_n^s)$  belongs either  $V$  or  $U \setminus V$ , we have by the choice of  $K_3$  in (2) that

$$\begin{aligned} \prod_{i=0}^{k-1} \|\psi_{t_{i+1} - t_i}^*(X_{t_i}(p_n^s))|_{\mathcal{F}_{X_{t_i}(p_n^s)}}\| &\geq \prod_{i=0}^{k-1} \left[ \frac{\|X(X_{t_i}(p_n^s))\|}{\|X(X_{t_{i+1}}(p_n^s))\|} \right]^2 e^{\lambda'(t_{i+1} - t_i)} \\ &\geq \left[ \frac{\|X(p_n^s)\|}{\|X(X_{t_k}(p_n^s))\|} \right]^2 e^{\lambda' k T_0} \\ &\geq (K_3)^2 e^{\lambda' k T_0} \\ &\geq (e^{\lambda'' T_0})^k = (\eta^{-1})^k. \end{aligned}$$

So, by Theorem 2.6, the local unstable manifold  $W_{\delta_1}^u(p_n^s)$  contains a segment of size  $\delta_1 \|X(p_n^s)\| \geq \delta_1 K_0$ , with  $\delta_1 = \delta_1(\eta, T_0)$ , for any  $n \geq N_1$ . On the other hand, since  $E$  is uniformly contracting on  $\Lambda'$  there are local stable manifolds  $W_{\varepsilon_0}^s(p_n^s)$  of uniform size and

tangent to  $E(p_n) \oplus \langle X(p_n) \rangle = E'^s(p_n) \oplus \langle X(p_n) \rangle$ , we say  $\varepsilon_0 > 0$ , on  $p_n^s$  for every  $n \geq 1$ . Hence, if we choose  $\varepsilon = \min\{\varepsilon_0, \delta_1\}$ , we have by the convergence of  $p_n^s$  to  $p^u$  that there is a natural number  $N > N_1$  such that

$$W_\varepsilon^u(p_n^s) \pitchfork W_\varepsilon^s(p_m^s) \text{ and } W_\varepsilon^s(p_n^s) \pitchfork W_\varepsilon^u(p_m^s), \quad n, m \geq N.$$

We have proved that  $p_n \in C(p_N)$  for every  $n \geq N$  and hence  $p_n^s \in C(p_N)$  for every  $n \geq N$ . To finish the proof we need to show that  $\sigma_X \in C(p_N)$ . For this sake, recall that  $p_n^s \rightarrow p^u \in C(\sigma_X) \cap W^u(\sigma_X)$ . Since  $W^u(\sigma_X)$  is invariant and uniformly contracted in the past, we conclude  $\sigma_X \sim p_N$ . Analogously, since  $p_n^u \rightarrow p^s \in C(\sigma_X) \cap W^s(\sigma_X)$  we conclude  $p_N \sim \sigma_X$ . This implies  $C(\sigma) \supset C(p_N)$  proving that  $p_N \in C(\sigma)$ . Then, the result is obtained by item (2) of Lemma 3.2.  $\square$

According to Lemma 3.5, it remains to prove that the continuation  $\Lambda_Y$  of  $\Lambda$  contains a periodic orbit. When  $\Lambda$  contains singularities, the argument is divided into two steps:

- (1) To find a transverse intersection between the stable manifold of  $\sigma_Y \in \Lambda_Y$  and the unstable manifold of some periodic orbit  $\gamma_Y$  for  $Y$ .
- (2) To find a transverse intersection between the stable manifold of a periodic orbit  $\gamma_Y$  for  $Y$  (whose dimension is  $\dim E^s$ ) and some disk  $D^c$  of dimension  $\dim F$ , tangent to the central cone.

The next result gives us step 1 in our argument.

**Lemma 3.6.** *Let  $X \in \mathcal{R}$  and let  $C(\sigma_X) \subset M$  be a non-trivial sectional-hyperbolic chain-recurrent class for  $X$ . Then, there is a neighborhood  $\mathcal{U}_1 \subset \mathcal{U}_0$  of  $X$  such that for every  $Y \in \mathcal{U}_1$ ,*

$$W^s(\sigma_Y) \pitchfork W^u(\gamma_Y) \neq \emptyset,$$

for some periodic orbit  $\gamma_Y \in \text{Per}(Y)$ .

*Proof.* By Lemma 3.5, there are a neighborhood  $V$  of  $\text{Sing}(C(\sigma_X))$  a point  $p^s \in C(\sigma_X) \cap W^s(\sigma_X)$  and a sequence of periodic orbits  $\gamma_n \in C(\sigma_X)$ ,  $n \geq 1$ , such that  $p_n^u \in \gamma_n$  converges to  $p^s$  when  $n$  goes to infinity.

Let  $\xi_k$ ,  $k \geq 1$ , be a decreasing sequence of positive numbers such that  $\xi_k \rightarrow 0$ . For every  $k > 0$ , choose  $N_k > 0$  such that  $d(p_n^u, p^s) < \xi_k$  for every  $n \geq N_k$ . Fix  $\alpha > 0$  as in Lemma 2.5, with respect to  $r$ ,  $C(\sigma)$  and  $T_0$ . Following the argument given in the proof of Case 2 of Lemma 3.5, there are  $N \in \mathbb{N}$  and  $\delta_1 > 0$  such that the local unstable manifold  $W_{\delta_1}^u(p_n^u)$  contains a segment of size  $\delta_1 \|X(p_n^s)\| > \delta_1 K_0$ , with  $\delta_1 = (\eta, T_0)$ , for any  $n \geq N$ . Since  $d(p_n^u, p^s) \leq \xi_k$ , for every  $n \geq N_k$  and  $\xi_k \rightarrow 0$ , we conclude there is a natural number  $k_1 \geq k_0$  such that  $p_n^u \in B_r(\sigma) \cap \overline{D_\alpha^{cs}(\sigma)}$ , for every  $n \geq N_{k_1}$ . So, Lemma 2.5 gives

$$(3) \quad W^u(p_n^u) \pitchfork W_{loc}^s(\sigma_X) \neq \emptyset,$$

for every  $n \geq N_{k_1}$ .

Now, since  $X \in \mathcal{R}$ , by item (3) of Lemma 3.2 and Lemma 2.7 we can find a neighborhood  $U$  of  $C(\sigma_X)$  and a neighborhood  $\mathcal{U}_X$  of  $X$  such that the map  $Y \in \mathcal{U}_X \rightarrow C(\gamma_Y)$  is continuous and the maximal invariant set  $\Lambda_Y$  for  $Y \in \mathcal{U}_X$  in  $U$  is sectional-hyperbolic for  $Y$  and it

contains  $C(\sigma_Y)$ . In particular, there are  $\delta_0 \approx \delta$  and a neighborhood  $\mathcal{U}_1 \subset \mathcal{U}_X$  of  $X$  such that  $W^s(\sigma_Y) \pitchfork W_{\delta_0}^u(\gamma_Y) \neq \emptyset$  for any  $Y \in \mathcal{U}_1$ . This proves the result.  $\square$

**Remark 3.7.** From the above lemma we derive the following observations:

- In the proof of Case 2, we connect a periodic orbit  $\gamma$  of  $X$  with the singularity  $\sigma \in C(\sigma)$ . Since the periodic orbits accumulate on  $C(\sigma)$  in the Hausdorff topology, according to item 2 of Lemma 3.2, we deduce that (3) holds for  $\gamma_n = \gamma$  and any singularity  $\rho \in C(\sigma)$ .
- It should be noted that not every non-trivial chain transitive sectional-hyperbolic is encompassed by the set  $\mathcal{R}$ , as illustrated by the Venice mask. Indeed, the Venice mask is a non transitive sectional-hyperbolic chain-transitive class having a dense set of periodic orbits, so that it cannot be a homoclinic class. The reason is that since the Venice mask is the union non disjoint of two homoclinic classes  $H_0$  and  $H_1$  whose intersection is the closure of the unstable manifold of a singularity, any periodic of its periodic orbits belongs either  $H_0$  or  $H_1$ , implying that it is not accumulated by the periodic orbits in the Hausdorff topology.

Now, recall that for a non singular point  $x \in M$ , the *orthogonal complement*  $N_x$  of  $X(x)$  is defined by

$$N_x = \{v \in T_x M : \langle v, X(x) \rangle = 0\}.$$

Let  $\Lambda = C(\sigma_X)$  and  $\Lambda_Y = C(\sigma_Y)$ . Fix a neighborhood  $U$  for  $\Lambda$  and  $\Lambda_Y$ , given by the continuity of  $Y \in \mathcal{U} \rightarrow C(\gamma_Y)$ . Since  $\Lambda$  is partially hyperbolic, it is possible to extend the SH splitting  $T_\Lambda M$  to  $U$  (by shrinking  $U$  if it is possible), that we will keep written as  $T_U M = E^s \oplus F$ . It should be noted that since  $\Lambda$  is sectional-hyperbolic one has  $1 \leq \dim E^s \leq n-2$  and  $1 \leq \dim F \leq n-1$ . Thus, the subspace  $N_x$  is of codimension one and has a coordinate system induced by the SH splitting for every  $x \in U$  in the following way:

$$\mathcal{E}_x^s = \pi(E_x^s) \text{ and } \mathcal{F}_x = \pi(F) \text{ or } F_x \cap N_x, \quad x \in U,$$

where  $\pi : TM \rightarrow N$  denotes the orthogonal projection. From partial hyperbolicity we have that the angle between these subspaces is uniformly bounded away from zero. In this way, denote by  $N(\eta)$  the standard cube  $(-\eta, \eta)^{n-1} \subset N_x$ , according to this coordinate system.

For any point  $x \in U$ , there is a number  $\eta(x) > 0$  such that the exponential map  $Exp_x : N_x \rightarrow M$  is an isometry. Note that for every  $\delta > 0$ , there is  $\eta_\delta > 0$  such that  $\eta(x) > \eta_\delta$  for any  $x \in U \setminus B(Sing(\Lambda), \delta)$ . Let  $\eta^* : U \rightarrow \mathbb{R}^+$  such that  $\eta^*(x) = \min\{\eta(x), \eta_\delta\}$  for every  $x \in U$ . Denote by  $\mathcal{N}_x = Exp_x(N_x(\eta^*(x))) \subset M$ , a codimension 1 cross section on  $x$  of size  $\eta^*(x)$ .

For a regular point  $x$ , the coordinates  $\mathcal{E}_x^s$  and  $\mathcal{F}_x$  allows us to define boxes  $B \subset N_x$  with respect to these axis as follows: Denote by  $\hat{\pi}_s : N_x \rightarrow \mathcal{E}_x^s$  and  $\hat{\pi}_c : N_x \rightarrow \mathcal{F}_x$  the projections along the  $\mathcal{E}_x^s$ -axis and the  $\mathcal{F}_x$ -axis respectively. The stable (central) volume of  $B$  is given by

$$|B|_{s(c)} := vol_{s(c)}(\pi_{s(c)}(B)),$$

where  $vol_{s(c)}(A_{s(c)})$  denotes the volume of a subset  $A \subset \mathbb{R}^{\dim \hat{E}_x^s(\hat{F}_x)}$ . We shall use a vector  $v = (\varepsilon_1^s, \varepsilon_2^s, \dots, \varepsilon_{n_s}^s, \varepsilon_1^c, \varepsilon_2^c, \dots, \varepsilon_{n_c}^c)$ , where  $1 \leq n_s \leq n-2$  and  $1 < n_c \leq n-1$  satisfy  $n_s + n_c + 1 = n$ , to describe the box  $B[v]$  with stable volume  $\varepsilon_s$  and central volume  $\varepsilon_c$ .

Let  $x \in U$  be a regular point does not belong to the stable manifold of the singularities of  $X$ , and  $t > 0$ . By the Implicit Function Theorem that there is an open connected subset  $D \subset \mathcal{N}_x$  containing  $x$  and a continuous function  $\tau : D \rightarrow N$  such that  $\tau(x) = t$  and  $X_{\tau(y)}(y) \in \mathcal{N}_{X_t(x)}$ , for any  $y \in D$ . The map  $\tau$  induces a diffeomorphism

$$H_x^t : \text{dom}(G_x^t) \subset \mathcal{N}_x \rightarrow \mathcal{N}_{X_t(x)},$$

where  $\text{dom}(H_x^t)$  is the maximal connected component of  $\mathcal{N}_x$ , containing  $x$  in its interior, where  $\tau$  is defined. the map  $H_x^t$  is called *the holonomy map between  $\mathcal{N}_x$  and  $\mathcal{N}_{X_t(x)}$  of time  $t > 0$* .

Before proceeding to step 2 of our argument, we present the next result about the non-trivial dynamical behavior of the chain-recurrence classes  $C(\sigma_Y)$  sufficiently  $C^1$ -close to  $X \in \mathcal{R}$ .

**Lemma 3.8.** *Let  $X \in \mathcal{R}$  and let  $C(\sigma_X) \subset M$  be a non-trivial sectional-hyperbolic chain-recurrent class for  $X$ . There is a neighborhood  $\mathcal{U}_2 \subset \mathcal{U}_1$  of  $X$  such that for every  $Y \in \mathcal{U}_2$  one has  $C(\sigma_Y) \not\subset W^s(\text{Sing}(Y))$ .*

*Proof.* By lemma 3.5 we have that  $C(\sigma_X)$  is a homoclinic class. Take two different periodic orbits  $\gamma_0, \gamma_1$  in  $C(\sigma_X)$ , and let  $d_\gamma = d_H(\gamma_0, \gamma_1) > 0$ . By continuity of  $Y \in \mathcal{U}_X \rightarrow C(\gamma_Y)$  there is a neighborhood  $\mathcal{U}_2 \subset \mathcal{U}_1$  of  $X$  such that

$$(4) \quad d_H(C(\sigma_X), C(\sigma_Y)) < \frac{d_\gamma}{2}, \quad \forall Y \in \mathcal{U}_2.$$

Hence, if  $C(\sigma_Y) \subset W^s(\text{Sing}(Y))$ , then either it is a homoclinic loop or it is contained into finitely many saddle connections  $\gamma_{\alpha, \rho}$ , associated to singularities  $\alpha, \rho \in C(\sigma_Y)$ . In any case, by (4) one has  $d_\gamma = d_H(\gamma_0, \gamma_1) < \frac{d_\gamma}{2}$ , which is impossible.  $\square$

Next we present, in a precise way, the step 2 in our argument.

**Lemma 3.9.** *Let  $X \in \mathcal{R}$  and let  $C(\sigma_X) \subset M$  be a non-trivial sectional-hyperbolic chain-recurrent class for  $X$ . There is a neighborhood  $\mathcal{U} \subset \mathcal{U}_2$  of  $X$  such that, for every  $Y \in \mathcal{U}$ , the periodic orbit  $\gamma_Y$  given by Lemma 3.6 is chain-attainable from  $\sigma_Y$ .*

*Proof.* To begin with, we recall that by Lemma 2.2 and Lemma 3.5,  $C(\sigma_X)$  is a homoclinic class whose singularities are Lorenz-like. Let us denote

$$\text{Sing}(X|_{C(\sigma_X)}) = \{\sigma_X = \sigma^0, \sigma^1, \dots, \sigma^k\}.$$

Notice that for any vector field  $Y$  sufficiently  $C^1$ -close to  $X$ , the continuations  $\text{Sing}(X|_{C(\sigma_Y)})$  consist entirely of Lorenz-like singularities.

Fix  $\delta > 0$  so that

$$\|X(z)\| \leq \|X(w)\|, \quad \forall z \in B(\text{Sing}(X|_{C(\sigma_x)}), \delta), w \notin B(\text{Sing}(X|_{C(\sigma_x)}), \delta).$$

Also, from the proof of Lemma 3.5, there is a sequence of periodic orbits  $\gamma_n \subset C(\sigma_X)$  such that  $d_H(\gamma_n, C(\sigma_X)) \rightarrow 0$ . In this case, let us assume

$$d_H(\gamma_n, C(\sigma_X)) < \delta/4,$$

for every  $n \geq 0$ . In particular, for each  $i = x, 1, 2, 3, \dots, k$ , we can obtain sequences of points  $(p_{n,i}^s)_{n \geq 1}$  and  $(p_{n,i}^u)_{n \geq 1}$  such that for every  $i = 0, 1, 2, \dots, k$ :

- $p_{n,i}^s, p_{n,i}^u$  are contained in  $\gamma_n$ , for every  $n \geq 1$  .
- $p_{n,i}^u \rightarrow p_{u,i} \in C(\sigma_X) \cap W^u(\sigma^i)$ ,
- $p_{n,i}^s \rightarrow p_{s,i} \in C(\sigma_X) \cap W^s(\sigma^i)$ ,
- $d(p_{s,i}, \sigma_i), d(p_{u,i}, \sigma_i) = \delta$ ,

Next, for such a value of  $\delta > 0$ , take a family of cross-sections

$$\mathcal{N}_i^{in} = \mathcal{N}_{p_{s,i}}^{in} \subset B(\sigma_i, \delta) \text{ and } \mathcal{N}_i^{out} = \mathcal{N}_{p_{u,i}}^{out} \subset B(\sigma_i, \delta)$$

at  $p_{s,i}$  and  $p_{u,i}$  respectively. Note that the size  $\eta$  of both  $\mathcal{N}_i^{in}$  and  $\mathcal{N}_i^{out}$  is bigger or equal than  $\eta_\delta$ . Moreover, for any  $i = 0, \dots, k$ , the holonomy map

$$H_i : Dom(H_i) \subset \mathcal{N}_i^{in} \setminus W_{loc}^s(\sigma^i) \rightarrow \mathcal{N}_i^{out}$$

is well defined. Since  $p_{n,i}^s \rightarrow p_{s,i}$  and  $p_{n,i}^u \rightarrow p_{u,i}$ , we can assume, without loss of generality, that  $p_{n,i}^s \in \mathcal{N}_i^{in}$  and  $p_{n,i}^u \in \mathcal{N}_i^{out}$ , for every  $n \geq 0$ . Take  $N \in \mathbb{N}$  satisfying

$$d_s^i = d(Q_{X,i}^s, p_{N,i}^s) < \eta/2 \text{ and } d_u^i = d(Q_{X,i}^u, p_{N,i}^u) < \eta/2,$$

where

$$Q_{X,i}^s = \mathcal{N}_i^{in} \cap W^s(\sigma^i) \text{ and } Q_{X,i}^u = \mathcal{N}_i^{out} \cap W^u(\sigma^i).$$

By the continuity of  $Y \in \mathcal{U}_X \rightarrow C(\gamma_Y)$ , there is a  $C^1$ -neighborhood  $\mathcal{U} \subset \mathcal{U}_2$  of  $X$  such that

(a) For every  $Y \in \mathcal{U}$ , the singularities of  $Y$  are Lorenz-like and

$$\|Y(z)\| \leq \|Y(w)\|, \quad \forall z \in B(Sing(Y|_{C(\sigma_Y)}), \delta), w \notin B(Sing(Y|_{C(\sigma_Y)}), \delta).$$

(b) For every  $Y \in \mathcal{U}$  and  $i = 0, 1, \dots, k$ , the submanifolds  $\mathcal{N}_i^{in}$  and  $\mathcal{N}_i^{out}$  are, respectively, cross-sections for  $Y$  at

$$p_{s,i}^Y \in C(\sigma_Y^i) \cap W^s(\sigma_Y^i) \text{ and } p_{u,i}^Y \in C(\sigma_Y^i) \cap W^u(\sigma_Y^i),$$

where  $p_{s,i}^Y, p_{u,i}^Y \in \gamma_Y$  and  $\gamma_Y$  is the periodic orbit given by Lemma 3.6.

(c) For every  $Y \in \mathcal{U}$  and  $i = 0, 1, \dots, k$ , the holonomy map  $H_i^Y$ , and the cross-sections  $\mathcal{N}_i^{in}$  and  $\mathcal{N}_i^{out}$ , with respect to the flow of  $Y$ , satisfy:

$$(\mathcal{N}_i^{in} \setminus W_{loc}^s(\sigma_Y^i)) \subset dom(H_i^Y) \text{ and } H_i^Y(Dom(H_i^Y)) \subset \mathcal{N}_i^{out}.$$

(d) For every  $Y \in \mathcal{U}$  and  $i = 0, 1, \dots, k$ , there exist  $y_{0,i}^s, y_{0,i}^u \in C(\sigma_Y)$  such that  $y_{0,i}^s \in \mathcal{N}_i^{in}$ ,  $y_{0,i}^u \in \mathcal{N}_i^{out}$  and

$$d(Q_{Y,i}^{s,u}, y_{0,i}^{s,u}), d(Q_{Y,i}^{s,u}, \gamma_Y) \approx d_{s,u}^i,$$

where  $Q_{Y,i}^{s,u}$  is the continuation of  $Q_{X,i}^{s,u}$ . This shows, in particular, that

$$d_{Y,i}^{s,u} = \sup\{d(y, Q_{Y,i}^{s,u}) : y \in C(\sigma_Y)\} \geq d(Q_{Y,i}^{s,u}, y_{0,i}^{s,u}) \approx d_{s,u}^i.$$

Also, define

$$d_Y^{s,u} = \min \left\{ d_{Y,i}^{s,u} : i = 0, \dots, k \right\}.$$

Now, we are ready to proceed with the proof. Let us split it into two cases.

**Case 1:** There is  $x \in C(\sigma_Y)$  such that  $\omega(x) \cap \text{Sing}(Y) = \emptyset$ .

Since  $\omega(x)$  has no singularities, it follows by Lemma 2.1 that it is hyperbolic of saddle type. In particular, has also local stable and unstable manifolds of size  $\varepsilon > 0$ . So, by shrinking  $\mathcal{U}$  if it is necessary, we obtain that

$$W^u(z) \pitchfork W^s(\gamma_Y) \neq \emptyset$$

for some  $z \in \omega(x) \subset C(\sigma_Y)$ . Thus, since  $z$  is chain attainable from  $\sigma_Y$ , one has that  $\gamma_Y$  is chain attainable from  $\sigma_Y$ .

**Case 2:**  $\omega(x) \cap \text{Sing}(Y) \neq \emptyset$  for any  $x \in C(\sigma_Y)$ .

In this case, we denote

$$\mathcal{N}^{in} = \bigcup_{i=0}^k \mathcal{N}_i^{in} \text{ and } \mathcal{N}^{out} = \bigcup_{i=0}^k \mathcal{N}_i^{out}.$$

The compactness of  $C(\sigma_Y) \setminus B(\text{Sing}(Y|_{C(\sigma_Y)}, \delta)$  implies the existence of a positive number  $T_\delta$  such that if

$$z \in C(\sigma_Y) \setminus B(\text{Sing}(Y|_{C(\sigma_Y)}, \delta),$$

there is  $0 < t_z \leq T_\delta$  such that  $Y_{t_z}(z) \in \mathcal{N}^{in}$  and  $Y_{(0,t_z)}(z) \cap \mathcal{N}^{in} = \emptyset$ .

By Lemma 3.8, there is a point  $x \in C(\sigma_Y) \setminus W^s(\text{Sing}(Y))$ . Moreover, since the positive orbit of  $x$  accumulates  $\text{Sing}(Y|_{C(\sigma_Y)})$ , there is a sequence of times  $t_n \rightarrow \infty$  and a singularity  $\sigma^i \in \text{Sing}(Y|_{C(Y)})$  such that  $X_{t_n}(x) \rightarrow \sigma^i$ . In particular, the positive orbit of  $x$  crosses  $\mathcal{N}^{in}$  infinitely many times. We may assume  $x \in \mathcal{N}_i^{in}$  for some  $i \in \{0, 1, \dots, k\}$ . Let us consider a box  $B \subset \mathcal{N}_i^{in}$  containing  $x$  with central volume  $0 < b < \frac{d_Y^s}{2}$ .

**Claim:** For any  $T > 0$ , there is  $T_b > T$ , depending on  $b$ , such that  $Y_{[0,T_b]}(B) \pitchfork W^s(\gamma_Y) \neq \emptyset$ .

Indeed, let consider a unitary vector  $\hat{v} \in \mathcal{F}_x$ , and take the plane  $L_x$  spanned by  $\hat{v}$  and  $Y(x)$ . Note that  $\hat{v}$  is either contained in  $F_x$  or has the form  $\hat{v} = \pi(v)$  for some  $v \in F_x$ . So, by denoting

$$K = \inf \left\{ \frac{\|Y(z)\|}{\|Y(w)\|} : z, w \notin B(\sigma_Y, \delta) \right\},$$

the sectional-hyperbolicity implies

$$\begin{aligned}
 \|\pi_{N_{Y_t(x)}}(\Phi_t(\hat{v}))\| &= \frac{\|Y(x)\|}{\|Y(Y_t(x))\|} \cdot \det(\Phi_t^Y(x)|_{L_x}) \\
 &\geq \frac{\|Y(x)\|}{\|Y(Y_t(x))\|} \cdot e^{\lambda t} \\
 (5) \quad &\geq Ke^{\lambda t},
 \end{aligned}$$

whenever  $t > 0$ ,  $Y_{[0,t)}(x) \in U$  and  $Y_t(x) \notin B(\sigma, \delta)$ .

Recall by the construction of the cross sections  $\mathcal{N}^{in}$  and  $\mathcal{N}^{out}$  that if  $z \in \mathcal{N}_i^{in}$ , then  $Y_{[0,\tau(z))}(z) \in B(\sigma_Y, \delta) \subset U$  and  $H_i^Y(z) = Y_{\tau(z)}(z) \in \mathcal{N}_i^{out}$ . In this case, define

$$\tau_i = \inf\{\tau(z) : z \in \mathcal{N}_i^{in}\}.$$

Notice that we can choose  $\delta > 0$  in a such way that

$$Ke^{\lambda t} \geq e^{\lambda_u t}, \quad \forall i = 0, 1, \dots, k, \forall t > \tau_i,$$

where  $0 < \lambda_u < \lambda$ . Let  $\mathcal{N}_{X_t(x)}$ ,  $t > \tau(x)$ , be a cross-section through  $X_t(x)$  of radius at least  $\eta_\delta$ . Since  $C(\sigma_Y)$  is not necessarily Lyapunov stable, the set  $H_x^t(B)$  may not be entirely contained in  $B(C(\sigma_Y), 2\delta)$  for some  $t > \tau_i$ . Nevertheless, there is a connected component  $B_1$  of  $H_x^t(B)$  whose central volume is bigger than  $b$  and  $\overline{B} \subset B(C(\sigma_Y), 2\delta)$ . In this case, we set

$$r_1 = \min\{r > 0 : Y_r(z) \in \partial B(C(\sigma_Y), 2\delta) \text{ for some } z \in (H_x^r)^{-1}(B_1)\}.$$

Then, if  $t = \tau(x) + r_1$ , by shrinking  $\mathcal{N}^{in}$  if it is necessary we have by (5) that

$$\|\pi_{N_{X_{\tau(x)+r}(x)}}(\Phi_{\tau(x)+r_1}(\hat{v}))\| \geq e^{\lambda_u(\tau+r_1)}, \quad \forall \hat{v} \in \hat{F}_x.$$

Hence, the map

$$H_x^{r_1} : Dom(H_x^{r_1}) \subset \mathcal{N}_i^{out} \rightarrow \mathcal{N}_{X_t(x)}$$

satisfies

$$H_i^Y(B) \subset Dom(H_x^{r_1}) \text{ and } |H_x^t(B)|_c = |H_x^{r_1} \circ H_i(B)|_c \geq e^{\lambda'(\tau_i+r)}b.$$

In particular, one has

$$|H_x^t(B)|_c \geq e^{\tau_i+r_1}b > 2b.$$

In this way, since  $T_\delta < \infty$ , we proceed in an inductively way to find a maximal  $\ell \in \mathbb{N}$ , positive numbers  $r_1, r_2, \dots, r_\ell$  such that  $\tau + r_1 + \dots + r_\ell \leq t_x^1$ , where  $t_x^1 > 0$  is the first return time of the orbit of  $x$  to  $\mathcal{N}^{in}$ , and a connected component  $B_\ell$  of  $H_x^{\tau(x)+r_1+\dots+r_\ell}(B)$  whose size is bigger than  $2b$ .

By the choice of  $\ell$ , there is  $s \geq 0$  such that  $\tau + r_1 + \dots + r_\ell + s = t_x^1$ . Besides, if  $\hat{w} = \pi_{N_{Y_{t_x^1-s}(x)}}(\Phi_{t_x^1-s}(\hat{v}))$ , we have by (5) and (1) that

$$\|\pi_{N_{Y_{t_x^1}(x)}}(\Phi_s(\hat{w}))\| \geq \frac{\|Y(x)\|}{\|Y(Y_t(x))\|} \cdot e^{\lambda s} \geq e^{\lambda s} > 1.$$

Hence,  $|H_x^{t_x^1}(B)|_c > 2b$ . In particular, since  $Y_{t_x^1}(x) \in \mathcal{N}^{in}$ , it follows that there is a connected component  $C_1 \subset \mathcal{N}^{in}$  of  $H_x^{t_x^1}(B)$  containing  $x$  whose central volume is bigger than  $2b$ .

By repeating the previous process in an inductively way, there is  $N \in \mathbb{N}$ , and a connected component  $C_N \subset \mathcal{N}^{in}$  of  $H_x^{t_x^N}(B)$  whose central volume is bigger than  $Nb > \frac{3}{2}d_Y^s$ . Furthermore, we can choose  $N$  and  $C_N$  in a such way that  $t_x^N > T$  and  $C_N \subset \mathcal{N}_i^{in}$ . So, we have that  $Y_{[0,s]}(C_N)$ ,  $s > 0$ , is a submanifold of dimension  $\dim F$  which intersects  $W^s(\sigma_j)$  transversely for some  $j \in \{0, 1, \dots, k\}$ . Moreover, since the central volume of  $C_N$  is bigger than  $d_Y^s$ , one has by (b) that  $W^s(\gamma_j) \pitchfork Y_{[0,s_0]}(C_N) \neq \emptyset$  for some  $s_0 > 0$ . So, the claim follows by taking  $T_b = t_x^N + s_0$ .

To finish the proof, recall that the positive orbit of  $x$  crosses  $\mathcal{N}^{in}$  infinitely many times and, therefore, there is a sequence of times  $t_n \rightarrow \infty$  and a singularity  $\sigma^i \in \text{Sing}(Y|_{C(Y)})$  such that  $X_{t_n}(x) \rightarrow x_0$  and  $x_0 \in W_{loc}^s(\sigma^i)$ . Without loss of generality, we can assume  $x_0, X_{t_n}(x) \in \mathcal{N}_i^{out}$ , for every  $n > 0$ . Fix  $\varepsilon, T > 0$  and fix  $N > 0$  such that  $d(X_{T_n}(x), x_0) \leq \varepsilon$ , for every  $n \geq N$ . Fix  $s_\varepsilon > T$  satisfying  $d(X_{-s_\varepsilon}(x_0), \sigma^i) \leq \varepsilon$ . Let  $x_n \in \mathcal{N}^{in}$  be such that  $x_n = X_{t_n+s}(x)$  and  $s > T$ . We we consider a box  $B$  around  $x_n$  with diameter  $\varepsilon$  and volume  $b$ , by the claim and the invariance of stable sets, there is  $z \in B \cap W^s(\gamma_Y)$ . Therefore, there exist  $s_z > T$  and  $x_z \in \gamma_Y$  such that  $d(X_{s_z}(z), x_z) \leq \varepsilon$ . Finally, the set

$$\{\sigma^i, X_{-s_\varepsilon}(x_0), X_{t_n}(x), z, x_z\}$$

is an  $\varepsilon$ - $T$ -chain from  $\sigma^i$  to  $\gamma_Y$ . Since  $\sigma_Y$  and  $\sigma^i$  are chain related, the proof is complete.  $\square$

Next, we are ready to prove Theorem 3.1.

*Proof of Theorem 3.1.* Let  $\mathcal{R}$  be given by Lemma 3.2. First, by Lemma 3.5 we have that  $C(\sigma_X)$  is a homoclinic class, which implies by Lemma 3.8 that the continuation  $C(\sigma_Y)$  does not consist of a homoclinic loop around  $\sigma_Y$  for every  $Y \in \mathcal{U}$ . Second, by Lemma 3.6 and Lemma 3.9, there is a periodic orbit  $\gamma_Y$  such that  $\sigma_Y \sim \gamma_Y$  and  $\gamma_Y \sim \sigma_Y$ , showing that  $\gamma_Y \in C(\sigma_Y)$  by definition of chain class.  $\square$

#### 4. PROOF OF THEOREMS A AND B

This section is devoted to prove our main results. Before proving Theorem A, we briefly outline the main idea behind the argument. The strategy combines the density of periodic orbits, in non-trivial sectional-hyperbolic chain-recurrence classes, and the denseness of its stable and unstable manifolds. In this way, for proving Theorem A we need to prove the following lemma:

**Lemma 4.1.** *Let  $\mathcal{R}$  be the generic set given by Lemma 3.2,  $X \in \mathcal{R}$  and  $C(\sigma)$  be a non-trivial chain-recurrent class of  $X$  containing a periodic orbit  $\gamma$ . There is an open neighborhood  $\mathcal{U}_X$  of  $X$  such that if  $Y \in \mathcal{U}_X$ , then for every  $x \in C(\sigma_Y)$  and every neighborhood  $V_x$  of  $x$ , we have*

$$W^s(\gamma_Y) \cap V_x \neq \emptyset \text{ and } W^u(\gamma_Y) \cap V_x \neq \emptyset,$$

where  $\gamma_Y \subset C(\sigma_Y)$  is a periodic orbit.

To improving our exposition, we shall split the proof of Lemma 4.1 into four separate lemmas. Let us begin with the first of them which is stated as follows:

**Lemma 4.2** (Theorem 3.1 in [CY21]). *There is a residual set  $\mathcal{R} \subset \mathcal{X}^1(M)$  such that if  $X \in \mathcal{R}$  and  $C(\sigma)$  is a non-trivial sectional-hyperbolic chain-recurrent class of  $X$  and  $\gamma \subset C(\sigma)$ , then there are neighborhood  $\mathcal{U}$  of  $X$  and a neighborhood  $U$  of  $C(\sigma)$  such that:*

*If  $Y \subset \mathcal{U}$  and  $x \in C(\sigma_Y)$ , then either  $x$  is on the unstable manifold of a singularity or  $W_{loc,Y}^{ss}(x) \pitchfork W_Y^u(\gamma_Y) \neq \emptyset$ .*

**Remark 4.3.** The attentive reader will notice that Lemma 4.2 is stated under slightly different assumptions than in [CY21], where it is assumed that  $C(\sigma)$  is a Lyapunov-stable class. However, a careful analysis of the proof in [CY21] reveals that this additional assumption is not actually used. Therefore, the same argument applies verbatim to the version we present here. For this reason, we omit the proof of Lemma 4.2 to avoid unnecessary repetition.

The next lemma gives us one of the inequalities in Lemma 4.1:

**Lemma 4.4.** *Under the assumption of Theorem 3.1,  $W^u(\gamma_Y)$  is dense on  $C(\sigma_Y)$ .*

*Proof.* Let  $Y \in \mathcal{U}$ ,  $x \in C(\sigma_Y)$  and let  $V$  be a neighborhood of  $x$ . We claim that  $W^u(\gamma_Y) \cap V \neq \emptyset$ . Indeed, we have the following cases:

**Case 1:** Assume  $x \in W^u(\rho)$ , for some  $\rho \in C(\sigma_Y)$ . In this case, for  $\varepsilon > 0$  there is  $t > 0$  such that  $d(\sigma, Y_{-t}(x)) < \varepsilon$ . Take a neighborhood  $W \subset M$  of  $Y_{-t}(x)$  such that  $Y_t(W) \subset V$ . By Remark 3.7 one has  $W^u(\gamma_Y) \pitchfork W^s(\rho)$ . So, if  $\varepsilon$  is small enough, we have by the inclination lemma that there is a point  $y_t \in W^u(\gamma_Y) \cap W$ , so that  $Y_t(y_t) \in V \cap W^u(\gamma_Y)$ .

**Case 2:** Assume  $x \notin W^u(\sigma'_Y)$  and let  $\varepsilon > 0$  such that  $B(x, \varepsilon) \subset V$ . By Lemma 4.2, for every  $t > 0$ , one has there is  $y_{-t} \in W_{loc}^{ss}(Y_{-t}(x)) \pitchfork W^u(\gamma_Y)$ . Since  $W_{loc}^{ss}(Y_t(x))$  is uniformly contracted by sectional-hyperbolicity, with rate  $\lambda_Y > 0$ , we take  $t > 0$  in a such way that  $d(Y_t(y_t), x) \leq Ce^{-\lambda_Y t} < \varepsilon$ . So,  $Y_t(W_{loc}^{ss}(Y_{-t}(x))) \subset V$ . This shows that  $Y_t(y_t) \in V \cap W^u(\gamma_Y)$ , and therefore  $W^u(\gamma_Y) \cap V \neq \emptyset$ .  $\square$

Now, since  $M$  is a compact manifold, there is a well-defined injectivity radius  $e_0 > 0$  for the exponential map  $\exp_x$ , for every  $x \in M$ . For every regular point  $x \in C(\sigma_Y)$  and  $\delta > 0$ , we say that  $D_\delta(x) \subset M$  is a *center-unstable section disc* (or simply *cu-section disc*) of  $Y$ , if it is given by  $\exp_x(B_\delta^{c-1}(0))$ , where  $B_\delta^{c-1}(0) \subset T_x M$  is the  $c-1$ -dimensional ball centered at  $x$  and radius  $\delta$ , and  $T_y D_\delta(x) \oplus \langle Y(y) \rangle \subset \mathcal{C}_y^c(a)$  for every  $y \in D_\delta(x)$ .

Let  $e_0 > 0$  be the injectivity radius of  $\exp$ . Fix  $0 < \varepsilon < e_0$  so that  $B_\varepsilon(C(\sigma_Y)) \subset U$ , where  $Y \in \mathcal{U}$  and  $U \subset M$  are given by Theorem 3.1. Given  $T > 0$ , we can shrinking  $U$  in a such way that  $Y_{[0,T]}(x) \in U$ . Hence, in a similar way that of [CY21, Section 4.2], there are finitely many, and well defined, cross sections  $D_Y \subset M$ , which are uniformly away from the singularities of  $C(\sigma_Y)$  for every  $Y \in \mathcal{U}$ . The next lemma shows that these cu-section discs expands, in a certain way, along the orbit of points of  $C(\sigma_Y)$ .

**Lemma 4.5.** *Suppose we are under the assumptions of Theorem 3.1. Let  $Y \in \mathcal{U}$  and fix  $\varepsilon > 0$  so that  $B_\varepsilon(C(\sigma_Y)) \subset U$ . Then, for every regular point  $x \in C(\sigma_Y)$ ,  $\delta > 0$ , and a*

cu-section disc  $D_\delta(x)$ , with  $\text{diam}(D_\delta(x)) \leq \delta$ , containing  $x$ , there is  $T' = T'(x, \delta) > 0$  such that  $Y_t(D_x)$  contains a cu-disc section  $D_\varepsilon(Y_t(x))$ , for every  $t \geq T'$ .

*Proof.* Let  $e_0 > 0$  be the injectivity radius of  $\exp$ . Fix  $0 < \varepsilon < e_0$  so that  $B_\varepsilon(C(\sigma_Y)) \subset U$ , and let consider  $0 < \delta < \varepsilon$ . Then, for every regular point  $x \in C(\sigma_Y)$  there is  $t'_x > 0$  such that  $Y_{t'_x}(x)$  belongs to a cross-section  $S_x$ . In particular, there are  $\alpha_1, \alpha_2, \beta > 0$  such that  $\alpha_1 \leq |Y(Y_{t'_x}(x))| \leq \alpha_2$ , and the angle between  $Y(Y_{t'_x}(x))$  and  $S_x$  is bigger than  $\beta$ . Therefore, we have  $DY_t(\mathcal{C}^c(y)) \subset \mathcal{C}^c(Y_t(y))$ , for every  $y \in D_\delta(x)$  and every  $t > 0$  satisfying  $Y_t(y) \subset U$ . Hence,  $D_{r_t}(Y_t(x)) = Y_t(D_\delta(x)) \cap \exp_{Y_t(x)} B_\varepsilon(0)$ , where  $t > 0$  and  $0 < r_t \leq \varepsilon$ , is a cu-section disc of  $Y$ . In other words, the cu-section discs are invariant for the flow.

Now, without loss of generality we assume that  $t'_x = 0$ ,  $D = D_\delta(x) \subset S_x$  for some cross section  $S_x$  for  $Y$ . Note that, by definition,  $D$  is a connected set. Then, if there is  $y \in D$  and  $t > 0$  such that  $Y_t(y) \notin B_\varepsilon(C(\sigma_Y))$ , there is a connected component  $D' \subset D$  such that  $\text{diam } Y_t(D') \geq \varepsilon$ , so the result is obtained in this case.

Assume that  $Y_t(D) \subset B_\varepsilon(C(\sigma_Y))$  for every  $t > 0$ . Let  $y \in D$ ,  $v \in T_y D$  be a unit vector, and denote by  $L_y$  the parallelogram formed by  $\langle Y(y) \rangle$  and  $v$ . Take  $t_0 > 0$  such that  $C'' \sin(\beta) e^{\lambda t_0} \geq \frac{\varepsilon}{\delta}$ , where  $C'' = C' \frac{\alpha_1}{\alpha_2} \sin(\beta) > 0$  and  $C' = CK$ , where  $C, K > 0$  are uniform constants given by sectional-hyperbolicity and the choice of the cross sections, and  $Y_{t_0}(x)$  belongs to some cross section  $S'$  for  $Y$ . Therefore, we have that

$$\begin{aligned} \|\Phi_{t_0}(v)\| &= K \cdot \frac{\text{Area}(\Phi_{t_0}^Y(L_y))}{\|Y(Y_{t_0}(x))\|} \\ &= K \cdot \frac{\det(\Phi_{t_0}^Y(x)) \cdot \text{Area}(L_y)}{\|Y(Y_{t_0}(y))\|} \\ &\geq KC e^{\lambda t_0} \cdot \frac{\|Y(y)\| \cdot \sin(\theta)}{\|Y(Y_{t_0}(y))\|} \\ &\geq C' \frac{\alpha_1}{\alpha_2} \sin(\beta) e^{\lambda t_0} \\ &= C'' e^{\lambda t_0} \geq \frac{\varepsilon}{\delta}, \end{aligned}$$

where  $\theta$  is the angle between  $Y(y)$  and  $v$ . So,  $\text{diam } Y_t(D) \geq \frac{\varepsilon}{\delta} \cdot \text{diam } D = \varepsilon$ , for every  $t \geq t_0$ . This concludes the proof.  $\square$

**Lemma 4.6.** *Under the assumption of Theorem 3.1,  $W^s(\gamma_Y)$  is dense on  $C(\sigma_Y)$ .*

*Proof.* Let  $Y \in \mathcal{U}$ ,  $x \in C(\sigma_Y)$  and let  $V$  be a neighborhood of  $x$ . We will prove that  $W^s(\gamma_Y) \cap V \neq \emptyset$ . First, we assume that  $x$  is regular. In this case, fix  $0 < \varepsilon < e_0$  such that  $B_\varepsilon(C(\sigma_Y)) \subset U$ , let consider a cu-section disc  $D = D_\delta(x)$ ,  $0 < \delta < \varepsilon$ , such that  $D \subset V$ , and let  $T' = T'(x, \delta) > 0$  be given by Lemma 4.5. According to the choice of the neighborhood  $U$  and the invariance of the cu-discs, there is  $\delta_0 > 0$  such that the disc  $D_{\delta_0} = \bigcup_{|t| < \delta_0} Y_t(D)$ , of dimension  $\dim F$ , is tangent to  $\mathcal{C}^c$  and it is contained in  $V$ . Moreover, by following the proof of Lemma 4.5, one has that the inner radius of  $Y_t(D_{\delta_0})$  increases with  $t > 0$ .

Second, recall that the angle between  $E$  and  $F$  is uniformly bounded away from zero. So, up to possibly reducing  $\varepsilon$ , there is  $\eta > 0$  satisfying the following property: If  $y, z \in C(\sigma_Y)$

satisfy  $d(y, z) \leq \eta$  and  $D(y), D(z)$  are discs tangent to the center unstable cone  $\mathcal{C}^c$  with inner radius  $\varepsilon$  and  $0 < r_z < \varepsilon/2$  respectively, then

$$(6) \quad W_{loc}^s(q) \pitchfork D(y) \neq \emptyset, \quad \forall q \in D(z).$$

Now, let  $t > 0$  such that the inner radius of  $Y_t(D_{\delta_0})$  contains a *cu*-section disc at  $x_t = Y_t(x)$  with inner radius  $\varepsilon$ . Since  $C(\sigma_Y)$  is a chain-recurrent class, it is also chain-transitive. Consequently, for every  $\eta > 0$  there is a  $\eta$ -chain  $(x_i, t_i)_{i=0}^k$ , with  $t_i \geq 1$ , such that  $x_0 = x_t, x_k = p \in \gamma_Y, d(Y_{t_i}(x_i), x_{i+1}) \leq \eta$ , and  $x_i \in C(\sigma_Y)$  for every  $i = 0, \dots, k$ . Then, if we take a *cu*-section disc  $D_\varepsilon(Y_{t_{k-1}}(x_{k-1}))$ , by the choice of  $\eta$  and (6) we obtain a point  $z_k$  in  $W_{loc}^s(\gamma_Y) \pitchfork D_\varepsilon(Y_{t_{k-1}}(x_{k-1})) \neq \emptyset$ . Moreover, since the inner radius of  $Y_{-t_i}(D_\varepsilon(Y_{t_i}(x_i)))$  is less than  $\varepsilon/2$  for every  $i = 1, \dots, k-1$ , we obtain in a recursive way, points  $z_i$  in  $W_{loc}^s(\gamma_Y) \pitchfork D_\varepsilon(Y_{t_i}(x_i))$ . The Figure 3 illustrates how these points are obtained.




FIGURE 1. Proof of Lemma 4.6.

In this way, since the inner radius of  $Y_{t+t_0}(D_{\delta_0})$  is at least  $\varepsilon$ , we have by definition of  $\eta$  a point  $z_0 \in W_{loc}^s(\gamma_Y) \pitchfork Y_{t+t_0}(D_{\delta_0})$ . So, the point  $Y_{-(t+t_0)}(z_0) \in W^s(\gamma_Y) \cap D_{\delta_0} \subset V$ .

Finally, if  $\rho$  is a singularity of  $C(\sigma_Y)$  and  $V \subset M$  is an open set containing  $\rho$ , there is a regular point  $x \in C(\sigma_Y)$  and an open neighborhood  $V_x$  of  $x$  satisfying  $V_x \subset V$ . So, by the above argument, there is a point  $z \in W^s(\gamma_Y) \cap V_x \subset V$ . This concludes the proof.  $\square$

Next, we are ready to prove Theorem A.

*Proof of Theorem A.* Consider the residual set  $\mathcal{R}$  given by Lemma 3.2. Let  $X \in \mathcal{R}$  and let  $C(\gamma)$  be a non-trivial sectional-hyperbolic chain-recurrent of  $X$ .

By Lemma 3.5 we have that  $C(\gamma)$  is a homoclinic class. Consequently, the stable manifold of  $\gamma$  is dense in  $C(\sigma)$  and meets transversally every submanifold of dimension  $\dim(F)$  that is tangent to  $\mathcal{C}^c$ , has inner radius  $\varepsilon$ , and intersects a sufficiently small neighborhood  $U' \subset U$  of  $C(\sigma)$ . Then, following the arguments in the proof of [CY21, Theorem 4.1], we can see that this property holds for vector fields  $Y$  in a small  $C^1$ -neighborhood  $\mathcal{U}_X$  of  $X$ . By Theorem 3.1 we have that  $C(\sigma_Y)$  contains periodic orbits. Moreover, for every  $\gamma' \subset C(\sigma_Y)$  and a neighborhood  $U_{\gamma'} \subset U'$  of  $\gamma'$ , one has that  $W_\varepsilon^u(\gamma')$  is a submanifold tangent to  $\mathcal{C}^c$  with inner radius  $\varepsilon$ . Thus, by Lemma 4.1 we have  $W^s(\gamma_Y) \cap W_\varepsilon^u(\gamma') \neq \emptyset$ . In particular, this shows that the set of transverse intersections  $W^s(\gamma_Y) \pitchfork W^u(\gamma')$  is dense in  $W^u(\gamma')$ .

Therefore, if  $x \in C(\sigma_Y)$  and  $V_x \subset U$  is a neighborhood of  $x$ , by Lemma 4.1 there is a point  $y \in W^u(\gamma')$  for some periodic orbit  $\gamma' \subset C(\sigma_Y)$ , which in turns is accumulated by elements of  $W^s(\gamma_Y) \pitchfork W^u(\gamma')$ . In other words,  $C(\gamma)$  is robustly a homoclinic class. This proves the result.  $\square$

Once we have proved Theorem A, let us provide the reader with the proof of Theorem B.

*Proof of Theorem B.* To begin with, notice that the star property is a  $C^1$ -open property, so that  $\mathcal{X}_*^1(M)$  is a  $C^1$ -open subset of  $\mathcal{X}^1(M)$ . Consequently,  $\mathcal{R}' = \mathcal{R} \cap \mathcal{X}_*^1(M)$  is a residual subset of  $\mathcal{X}^1(M)$ , where  $\mathcal{R}$  is the residual set given by Theorem A.

On the other hand, by [PY23, Theorem A], there is a residual subset  $\mathcal{S}' \subset \mathcal{X}_*^1(M)$  such that for any  $Y \in \mathcal{S}'$  and  $\Lambda$  is a non-trivial chain-recurrent class, one of the following holds:

- (1)  $h_{top}(X|_\Lambda) > 0$ ,  $\Lambda$  contains a periodic orbit and is isolated; or
- (2)  $h_{top}(\Lambda) = 0$  and  $\Lambda$  is sectional-hyperbolic aperiodic class.

By setting  $\mathcal{S} = \mathcal{R}' \cap \mathcal{S}'$ , any non-trivial sectional-hyperbolic chain-recurrent class for  $Y \in \mathcal{S}$  contains a periodic orbit by Theorem A. Hence, option 2 cannot occur, and the proof is complete.  $\square$

**Acknowledgments.** The authors would like to thank Professor Alexander Arbieto for pointing a gap in the proof of a crucial lemma in an earlier version of this work and for valuable suggestions that helped us to figure out a solution for this gap. K. J. Vivas was supported by ANID Proyecto FONDECYT Iniciación 11250633, Chile.



This research is part of a project that has received funding from the European Union's European Research Council Marie Skłodowska-Curie Project No. 101151716 – TMSHADS – HORIZON–MSCA–2023–PF–01.

## REFERENCES

- [AN07] J. M. Aroldi and G. S. Nelson. *Recurrence and topology*, Graduate Studies in Mathematics, 85, Amer. Math. Soc., Providence, RI, 2007;
- [ABS77] Afraimovich V. S., Bykov V. V. and Shilnikov L. P. On the appearance and structure of the Lorenz attractor, *Dokl. Acad. Sci. USSR*. 234: 336–339, 1977.
- [AM17] V. Araujo and I. Melbourne. Existence and smoothness of the stable foliation for sectional-hyperbolic attractors. *Bull. London Math. Soc.* 49: 351–367, 2017.
- [AM19] V. Araujo and I. Melbourne. Mixing properties and statistical limit theorems for singular hyperbolic flows without a smooth stable foliation. *Adv. in Math.* 349: 212–245, 2019.
- [Ba04] S. Bautista. The geometric Lorenz attractor is a homoclinic class. *Bol. Mat. (N.S.)* 11: 69–78, 2004.
- [BM11] S. Bautista and C. Morales. Lectures on sectional-Anosov flows. *Preprint IMPA Série D.* 84, 2011.

- [BdL21] C. Bonatti; A. da Luz. Star flows and multisingular hyperbolicity, *Journal of the European Mathematical Society* 23, no. 8, pp. 2649–2705 2021
- [BPV97] C. Bonatti; A. Pumaríño and M. Viana. Lorenz attractors with arbitrary expanding dimension, *C. R. Acad. Sci. Paris Sér. I Math.* 325: 883–888, 1997.
- [BC04] C. Bonatti and S. Crovisier. Récurrence et généricité. *Invent. Math.*, 158: 33–104, 2004.
- [Con78] C. Conley Isolated invariant sets and Morse index. *CBMS Regional Conference Series in Mathematics*, 38, AMS (1978)
- [Cro06] S. Crovisier. Periodic orbits and chain-transitive sets of  $C^1$ -diffeomorphisms. *Publ. Math. Inst. Hautes Études Sci.*, 104: 87–141, 2006.
- [Cro10] S. Crovisier. Birth of homoclinic intersections: a model for the central dynamics of partially hyperbolic systems. *Ann. of Math.*, 172: 1641–1677, 2010.
- [CY21] S. Crovisier and D. Yang. Robust transitivity of singular hyperbolic attractors. *Mathematische Zeitschrift*, 298: 469–488, 2021.
- [FS76] J.E. Franke and J. F. Selgrade, Abstract  $\omega$ -limit sets, chain-recurrent sets, and basic sets for flows. *Proc. Amer. Math. Soc.*, 60 (1976), 309–316 (1977)
- [GW06] Gan S. and Wen L. Nonsingular star flows satisfy Axiom A and the no-cycle condition, *Invent. Math.* 164, 279–315, 2006.
- [GY18] Gan S. and Yang D. Morse-Smale systems and horseshoes for three dimensional singular flows *Annales Scienti ques de l'École Normale Supérieure* 51, 39–112 (2018).
- [GYZ22] Gan S., Yang J. and Zheng R. Lyapunov stable chain-recurrence classes for singular flows. Pre-Print arXiv:2202.09742.
- [Gu76] Guckenheimer J. A strange, strange attractor. The Hopf bifurcation and its application, *Applied Mathematical Sciences* 19, 1976.
- [Ha92] Hayashi S. Diffeomorphisms in  $\mathcal{F}^1(M)$  satisfy Axiom A *Ergodic Theory & Dynam. Systems* 12, 233–253, 1992.
- [HPPS70] M. Hirsch, J. Palis, C. Pugh, M. Shub. Neighborhoods of hyperbolic sets. *Invent. Math.* 9: 121–134, 1970.
- [Li81] Liao S. Obstruction sets. II. (in Chinese) *Beijing Daxue Xuebao* 2, 1-36, 1981.
- [dL17] da Luz A. Hyperbolic sets that are not contained in a locally maximal one. *Discrete Contin. Dyn. Syst.* 37, no. 9, 4923–4941, 2017.
- [Ma78] Mañé R. Contributions to the stability conjecture. *Topology* 17, no. 4, 383–396, 1978.
- [MM08] R. Metzger and C. Morales. Sectional-Hyperbolic systems. *Ergodic Theory Dynam. Systems*. 28:1587–1597, 2008.
- [MPP99] C. Morales, M. J. Pacifico and E. Pujals. Singular hyperbolic systems. *Proc. Amer. Math. Soc.* 127: 3393–3401, 1999.
- [New72] S. Newhouse. Hyperbolic Limit Sets. *Trans. Amer. Math. Soc.* 167: 125–150, 1972.

- [PYY21] M. J. Pacifico, F. Yang and J. Yang. Entropy theory for sectional hyperbolic flows. *Annales de L'Institut Henri Poincaré C, Analyse non linéaire*. 38: 1001–1030, 2021.
- [PYY23] M. J. Pacifico, F. Yang and J. Yang. An entropy dichotomy for singular star flows. *Trans. Amer. Math. Soc.* 376: 6845–6871, 2023.
- [Sa19] L. Salgado. Singular hyperbolicity and sectional Lyapunov exponents of various orders. *Proc. Amer. Math. Soc.* 147: no. 2, 735–749, 2019.
- [SGW14] Y. Shi, S. Gan and L. Wen. On the singular hyperbolicity of star flows. *Journal of Modern Dynamics* 8, 191–219, 2014.
- [Sm67] S. Smale. Differentiable dynamical systems. *Bull. Amer. Math. Soc.* 73: 747–817, 1967.
- [XZ08] Xiao Q. and Zheng Z.  $C^1$  weak Palis conjecture for nonsingular flows. *Discrete & Continuous Dynamical Systems* 38, 1809–1832 (2008).
- [ZGW08] Zhu S., Gan S. and Wen. L. Indices of singularities of robustly transitive sets. *Discrete & Continuous Dynamical Systems* 21, 945–957 (2008).

DEPARTMENT OF APPLIED MATHEMATICS, AGH UNIVERSITY OF SCIENCE AND TECHNOLOGY, KRAKOW, POLAND.

*Email address:* `rego@agh.edu.pl`

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD CATÓLICA DEL NORTE, ANTOFAGASTA, CHILE.  
*Email address:* `kendry.vivas01@ucn.cl`