

THE EXISTENCE OF VALUATIVE INTERPOLATION AT A SINGULAR POINT

SHIJIE BAO, QI'AN GUAN, ZHITONG MI, AND ZHENG YUAN

ABSTRACT. The present paper studies the existence of valuative interpolation on the local ring of an irreducible analytic subvariety at singular points. We firstly develop the concepts and methods of Zhou weights and Tian functions near singular points of irreducible analytic subvarieties. By applying these tools, we establish the necessary and sufficient conditions for the existence of valuative interpolations on the rings of germs of holomorphic functions and weakly holomorphic functions at a singular point.

As applications, we characterize the existence of valuative interpolations on the quotient ring of the ring of convergent power series in real variables. We also present separated necessary and sufficient conditions for the existence of valuative interpolations on the quotient ring of polynomial rings with complex coefficients and real coefficients. Furthermore, we show that the conditions become both necessary and sufficient under certain conditions on the zero set of the given polynomials.

CONTENTS

1. Introduction	2
1.1. Zhou weights and Zhou valuations near singular points	2
1.2. Interpolation problem	5
2. Preparation	7
2.1. Basic knowledge	7
2.2. Concavity	10
2.3. Tian functions and Zhou numbers	11
2.4. Relation between relative types and valuations	14
2.5. Jumping number in the singular case	25
2.6. Convergence results for valuations and for relative types	31
3. Proof of Remark 1.3	32
4. Proof of Theorem 1.6	33
5. Proof of Theorem 1.8	35
6. Proofs of Theorem 1.9, Theorem 1.10 and their Corollaries	36
Appendix A. The universal denominators for algebraic varieties	39
References	42

Date: January 6, 2026.

2020 Mathematics Subject Classification. 13A18, 14B05, 32U05.

Key words and phrases. Singular point, valuation, relative type, interpolation.

1. INTRODUCTION

Valuation theory is deeply connected to the theory of singularities in several complex variables and complex algebraic geometry (see e.g. [11, 12, 13, 8, 21, 2]). Let X be an analytic subvariety and x a point of X . Denote the ring of germs of holomorphic functions at x in X by $\mathcal{O}_{X,x}$, and $\mathcal{O}_{X,x}^* := \mathcal{O}_{X,x} \setminus \{0\}$. Recall that a *valuation* on $\mathcal{O}_{X,x}$ is a nonconstant map $\nu: \mathcal{O}_{X,x}^* \rightarrow \mathbb{R}_{\geq 0}$ satisfying the following:

- (1) $\nu(fg) = \nu(f) + \nu(g)$;
- (2) $\nu(f+g) \geq \min\{\nu(f), \nu(g)\}$;
- (3) $\nu(c) = 0$, where $c \neq 0$ is a constant function.

Similarly, we can define the valuations on the polynomial ring $\mathbb{C}[z_1, \dots, z_n]/I$, where I is a proper ideal of $\mathbb{C}[z_1, \dots, z_n]$.

A natural problem is how to characterize the existence of valuative interpolation:

Question 1.1. *Given any positive integer m , a finite set of elements $\{f_j\}_{1 \leq j \leq m}$ in $\mathcal{O}_{X,x}$ (or $\mathbb{C}[z_1, \dots, z_n]/I$), and any finite nonnegative real numbers $\{a_j\}_{1 \leq j \leq m}$, can one find necessary and sufficient conditions for the existence of the valuation ν on $\mathcal{O}_{X,x}$ (or on $\mathbb{C}[z_1, \dots, z_n]/I$) such that $\nu(f_j) = a_j$ for all $j \in \{1, \dots, m\}$?*

When x is a regular point, we [3] established the necessary and sufficient conditions for the existence of valuative interpolations on $\mathcal{O}_{X,x}$. The proof in [3] relies heavily on the properties of Zhou weights and Tian functions, where the concepts and properties of Zhou weights and Tian functions in the smooth case were established by the authors in [2]. As applications, characterizations of the existence of valuative interpolation in the polynomial ring case and the real case were also established in [3].

In the present paper, we consider the valuative interpolation problem when x is a singular point of an irreducible analytic subvariety X . To do this, we generalize the concepts and properties of Zhou weights, Zhou valuations and Tian functions from the smooth case (presented in [2]) to the singular case. The properties of jumping numbers and relative types in the singular case are also studied. Based on these preparations, we present necessary and sufficient conditions for the existence of valuative interpolations on rings of germs of holomorphic functions and weakly holomorphic functions at a singular point x of an irreducible analytic subvariety X . As an application, we also give the necessary and sufficient conditions separately for the existence of valuative interpolations on the quotient ring of polynomial rings with complex coefficients and we show that the conditions become both necessary and sufficient when the intersection of the zero sets of the given polynomials is the origin o contained in X . As a further application, we derive necessary and sufficient conditions for the valuative interpolation problem in the real case.

1.1. Zhou weights and Zhou valuations near singular points. Let X be an analytic subset of $\Omega \in \mathbb{C}^n$ with pure dimension d . Let

$$dV_X := \frac{1}{2^d d!} \bigwedge^d \left(\sum_{1 \leq j \leq n} \sqrt{-1} dz_j \wedge d\bar{z}_j \right)$$

be the volume form on X induced by the standard volume form on \mathbb{C}^n .

We call a Lebesgue function φ on an analytic set X a *plurisubharmonic* function, if φ is plurisubharmonic on X_{reg} and is bounded above near any $z \in X_{\text{sing}}$. In this

article, when we consider the values of plurisubharmonic functions on analytic sets, we ignore their values on singular points.

Let (X, z_0) be an irreducible germ of an analytic set. Let $f_0 = (f_{0,1}, \dots, f_{0,m})$ be a vector, where $f_{0,1}, \dots, f_{0,m}$ are holomorphic functions near z_0 . Denote by $|f_0|^2 = |f_{0,1}|^2 + \dots + |f_{0,m}|^2$. Let φ_0 be a plurisubharmonic function near z_0 , such that $|f_0|^2 e^{-2\varphi_0}$ is integrable near z_0 .

Definition 1.2. *We call a plurisubharmonic function $\Phi_{z_0, \max}^{f_0, \varphi_0}$ ($\Phi_{z_0, \max}$ for short) near z_0 on X a local Zhou weight related to $|f_0|^2 e^{-2\varphi_0}$ near z_0 , if the following three statements hold*

- (1) $|f_0|^2 e^{-2\varphi_0} |z|^{2N_0} e^{-2\Phi_{z_0, \max}}$ is integrable near z_0 for large enough $N_0 \gg 0$, where $|z|^2 := \sum_{1 \leq j \leq n} |z_j|^2$;
- (2) $|f_0|^2 e^{-2\varphi_0} e^{-2\Phi_{z_0, \max}}$ is not integrable near z_0 ;
- (3) for any plurisubharmonic function $\varphi' \geq \Phi_{z_0, \max} + O(1)$ near z_0 such that $|f_0|^2 e^{-2\varphi_0} e^{-2\varphi'}$ is not integrable near z_0 , $\varphi' = \Phi_{z_0, \max} + O(1)$ holds.

Let φ be a plurisubharmonic function near z_0 . The existence of local Zhou weights follows from the strong openness property of multiplier ideal sheaves [17].

Remark 1.3. *Assume that $|f_0|^2 e^{-2\varphi_0} |z|^{2N_0} e^{-2\varphi}$ is integrable near o for large enough $N_0 \gg 0$, and $|f_0|^2 e^{-2\varphi_0-2\varphi}$ is not integrable near z_0 .*

Then there exists a local Zhou weight $\Phi_{z_0, \max}$ related to $|f_0|^2 e^{-2\varphi_0}$ near z_0 such that $\Phi_{z_0, \max} \geq \varphi$.

Moreover, for any local Zhou weight $\Phi_{z_0, \max}$, $\Phi_{z_0, \max} \geq N \log |z| + O(1)$ near z_0 holds for some $N \gg 0$.

We recall the definition of *weakly holomorphic functions* on a complex space.

Definition 1.4 (see [10, Definition 7.1]). *Let X be a complex space. A weakly holomorphic function f on X is a holomorphic function on X_{reg} such that any point of X_{sing} has a neighborhood V for which f is bounded on $X_{\text{reg}} \cap V$. We denote by $\mathcal{O}_{X,x}^w$ the ring of germs of weakly holomorphic functions on neighborhoods of x and \mathcal{O}_X^w the associated sheaf.*

When X is normal at some point x , we know that $\mathcal{O}_{X,x}^w = \mathcal{O}_{X,x}$. In general, we only have $\mathcal{O}_{X,x} \subset \mathcal{O}_{X,x}^w$.

When $f_{0,i}$ ($i = 1, \dots, m$) are weakly holomorphic functions near z_0 , we can also define the local Zhou weight related to $|f_0|^2 e^{-2\varphi_0}$.

Remark 1.5. *Let $f_{0,i}$ ($i = 1, \dots, m$) be weakly holomorphic functions near z_0 and φ_0 be a plurisubharmonic function near z_0 , such that $|f_0|^2 e^{-2\varphi_0}$ is integrable near z_0 . It follows from Theorem 2.15 that there exists a holomorphic function δ near o such that $g_{0,i} := \delta f_{0,i}$ ($i = 1, \dots, m$) is holomorphic near z_0 . Then we define that a plurisubharmonic function Φ near o is a local Zhou weight related to $|f_0|^2 e^{-2\varphi_0}$ if and only if Φ is a local Zhou weight related to $|g_0|^2 e^{-2\log |\delta| - 2\varphi_0}$.*

Thus, by Remark 1.3, the local Zhou weights exists for weakly holomorphic functions.

When z_0 is a smooth point of X , to study the singularity of the plurisubharmonic functions near z_0 , Rashkovskii [27] introduced the concept of **relative type**

$$\sigma(\psi, \varphi) := \sup\{c \geq 0: \psi \leq c\varphi + O(1) \text{ near } o\},$$

where ψ is a plurisubharmonic function near z_0 and φ is a maximal weights with an isolated singularity at z_0 . The notation of relative type generalizes the classical Lelong number [25] and Kiselman number [23].

When z_0 is a singular point of X , we can generalize the definition of relative type to the singular case. Let ψ, φ be any plurisubharmonic functions defined near z_0 . Define the **relative type** by

$$\begin{aligned} \sigma(\psi, \varphi) := \sup \{c \geq 0: \psi \leq c\varphi + O(1) \text{ holds on } U \setminus X_{\text{sing}} \\ \text{for some open neighborhood } U \text{ of } z_0 \text{ in } X\}. \end{aligned}$$

When the $\varphi = \Phi_{z_0, \max}$ is a local Zhou weight, we call the relative type $\sigma(\cdot, \Phi_{z_0, \max})$ the **Zhou number**.

Note that for any $b < \sigma(\psi, \Phi_{z_0, \max})$, $|f_0|^2 e^{-2\varphi_0} e^{-2\max\{\Phi_{z_0, \max}, \frac{1}{b}\psi\}}$ is not integrable near z_0 . Then it follows from the *strong openness property* of multiplier ideal sheaves (Theorem 2.7) that $|f_0|^2 e^{-2\varphi_0} e^{-2\max\{\Phi_{z_0, \max}, \frac{1}{\sigma(\psi, \Phi_{z_0, \max})}\psi\}}$ is not integrable near z_0 . Note that

$$\max \left\{ \Phi_{z_0, \max}, \frac{1}{\sigma(\psi, \Phi_{z_0, \max})} \psi \right\} \geq \Phi_{z_0, \max}.$$

Then

$$\max \left\{ \Phi_{z_0, \max}, \frac{1}{\sigma(\psi, \Phi_{z_0, \max})} \psi \right\} = \Phi_{z_0, \max} + O(1),$$

which implies that

$$\Phi_{z_0, \max} \geq \frac{1}{\sigma(\psi, \Phi_{z_0, \max})} \psi + O(1),$$

i.e.,

$$\psi \leq \sigma(\psi, \Phi_{z_0, \max}) \Phi_{z_0, \max} + O(1).$$

For the Lelong numbers and the Kiselman numbers, there are expressions in integral form (see [10]). Now, we show that Zhou numbers also have expressions in integral form.

Theorem 1.6. *Let $\Phi_{z_0, \max}$ be a local Zhou weight related to $|f_0|^2 e^{-2\varphi_0}$ near z_0 . Then for any plurisubharmonic function ψ near z_0 satisfying $\psi \leq c \log |z| + O(1)$ near z_0 for some $c > 0$, we have*

$$\sigma(\psi, \Phi_{z_0, \max}) = \lim_{t \rightarrow +\infty} \frac{\int_{\{\Phi_{z_0, \max} < -t\}} |f_0|^2 e^{-2\varphi_0} (-\psi)}{t \int_{\{\Phi_{z_0, \max} < -t\}} |f_0|^2 e^{-2\varphi_0}}.$$

Denote by

$$\nu(f, \Phi_{z_0, \max}) := \sigma(\log |f|, \Phi_{z_0, \max})$$

for any $(f, z_0) \in \mathcal{O}_{X, z_0}$. By Theorem 1.6 and the definition of $\nu(\cdot, \Phi_{z_0, \max})$, we see that $\nu(\cdot, \Phi_{z_0, \max})$ is a *valuation* of \mathcal{O}_{X, z_0} for any local Zhou weight $\Phi_{z_0, \max}$, and we call it **Zhou valuation**.

Corollary 1.7. *For any local Zhou weight $\Phi_{z_0, \max}$ near o , $\nu(\cdot, \Phi_{z_0, \max}) : \mathcal{O}_{X, z_0} \rightarrow \mathbb{R}_{\geq 0}$ satisfies the following:*

- (1) $\nu(fg, \Phi_{z_0, \max}) = \nu(f, \Phi_{z_0, \max}) + \nu(g, \Phi_{z_0, \max})$;
- (2) $\nu(f + g, \Phi_{z_0, \max}) \geq \min \{\nu(f, \Phi_{z_0, \max}), \nu(g, \Phi_{z_0, \max})\}$;
- (3) $\nu(f, \Phi_{z_0, \max}) = 0$ if and only if $f(z_0) \neq 0$.

Let G be a holomorphic function near z_0 . We recall the definition of jumping number (see [21, 22])

$$c_{z_0}^G(\Phi_{z_0, \max}) := \sup \{c : |G|^2 e^{-2c\Phi_{z_0, \max}} \text{ is integrable near } z_0\}.$$

When $G = 1$, the jumping number was also called complex singularity exponent and denote $c_{z_0}(\Phi_{z_0, \max}) := c_{z_0}^1(\Phi_{z_0, \max})$ (see [29, 9]).

Theorem 1.8. *Let $\Phi_{z_0, \max}$ be a local Zhou weight near z_0 . For any holomorphic function G near z_0 , we have the following relation between $c_{z_0}^G(\Phi_{z_0, \max})$ and $\sigma(G, \Phi_{z_0, \max})$,*

$$\begin{aligned} \nu(G, \Phi_{z_0, \max}) + c_{z_0}(\Phi_{z_0, \max}) &\leq c_{z_0}^G(\Phi_{z_0, \max}) \\ &\leq \nu(G, \Phi_{z_0, \max}) - \sigma(\log |f_0|, \Phi_{z_0, \max}) + 1 + \sigma(\varphi_0, \Phi_{z_0, \max}). \end{aligned}$$

Especially, if $|f_0|^2 e^{-2\varphi_0} \equiv 1$, we have

$$\nu(G, \Phi_{z_0, \max}) + 1 = c_{z_0}^G(\Phi_{z_0, \max}).$$

1.2. Interpolation problem. Let X be an analytic variety with pure dimension d contained in \mathbb{C}^n . Assume that the origin $o \in X_{\text{sing}}$ and (X, o) is an irreducible germ of an analytic set. The following theorem presents a criterion for the valuative interpolation problem on $\mathcal{O}_{X, o}$.

Theorem 1.9. *Let $f_j \in \mathcal{O}_{X, o}^*$ for $0 \leq j \leq m$. Let $a_0 = 0$ and $\{a_j\}_{1 \leq j \leq m}$ be positive numbers. The following two statements are equivalent:*

- (1) *There exists a valuation ν on $\mathcal{O}_{X, o}$ such that $\nu(f_j) = a_j$ for all $0 \leq j \leq m$;*
- (2) *$\sigma(\log |F|, \varphi) = \sum_{1 \leq j \leq m} a_j$, where*

$$F := \prod_{0 \leq j \leq m} f_j, \quad \varphi := \log \left(\sum_{1 \leq j \leq m} |f_j|^{\frac{1}{a_j}} \right).$$

Note that $\nu(f_1 f_2) = 0$ if and only if $\nu(f_1) = 0$ and $\nu(f_2) = 0$. Thus, in the above theorem, it suffices to consider the case only one function f_0 satisfies $\nu(f_0) = 0$. As $\nu(1) = 0$ holds for any valuation ν , in the above theorem, we can actually ignore f_0 when we take $f_0 \equiv 1$.

For valuations on the germs of weakly holomorphic functions $\mathcal{O}_{X, o}^w$, we also have the following interpolation result.

Theorem 1.10. *Let $\{f_j\}_{0 \leq j \leq m}$ be weakly holomorphic functions near o . Let $a_0 = 0$ and $\{a_j\}_{1 \leq j \leq m}$ be positive numbers. The following two statements are equivalent:*

- (1) *There exists a valuation ν on $\mathcal{O}_{X, o}^w$ such that $\nu(f_j) = a_j$ for all $0 \leq j \leq m$;*
- (2) *$\sigma(\log |F|, \varphi) = \sum_{1 \leq j \leq m} a_j$, where*

$$F := \prod_{0 \leq j \leq m} f_j, \quad \varphi := \log \left(\sum_{1 \leq j \leq m} |f_j|^{\frac{1}{a_j}} \right).$$

Let us consider the valuative interpolation problem on the polynomial ring $\mathbb{C}[z_1, \dots, z_n]/I$, where I is a prime ideal in $\mathbb{C}[z_1, \dots, z_n]$. Let $X := V(I)$ be the affine variety defined by I and $o \in X$. Denote the germ of the set X at o by (X, o) and we assume that (X, o) is irreducible as a germ of analytic set.

Remark 1.11. *It follows from the main theorem in [32] that, when the affine variety X is normal at o , (X, o) is irreducible as a germ of analytic set. So there exists many affine varieties X such that (X, o) is irreducible as a germ of analytic set.*

Let $\{f_j\}_{0 \leq j \leq m}$ be polynomials in $\mathbb{C}[z_1, \dots, z_n]/I$, $a_0 = 0$ and $\{a_j\}_{1 \leq j \leq m}$ be m positive numbers. Denote $F := \prod_{0 \leq j \leq m} f_j$ and $\varphi := \log(\sum_{1 \leq j \leq m} |f_j|^{\frac{1}{a_j}})$.

Corollary 1.12. *If $\sigma(\log |F|, \varphi) = \sum_{1 \leq j \leq m} a_j$, then there exists a valuation ν on $\mathbb{C}[z_1, \dots, z_n]/I$ such that $\nu(f_j) = a_j$ for every $1 \leq j \leq m$.*

Conversely, if there exists a valuation ν on $\mathbb{C}[z_1, \dots, z_n]/I$ satisfying that $\nu(f_j) = a_j$ for every $1 \leq j \leq m$ and $\nu(z_l) > 0$ for every $1 \leq l \leq n$, then we have $\sigma(\log |F|, \varphi) = \sum_{1 \leq j \leq m} a_j$.

When $\cap_{1 \leq j \leq m} \{f_j = 0\} = \{o\}$, Theorem 1.12 gives a necessary and sufficient condition for the existence of valuative interpolations on $\mathbb{C}[z_1, \dots, z_n]$.

Corollary 1.13. *If $\cap_{1 \leq j \leq m} \{f_j = 0\} = \{o\}$, then there exists a valuation ν on $\mathbb{C}[z_1, \dots, z_n]/I$ such that $\nu(f_j) = a_j$ for every j if and only if $\sigma(\log |F|, \varphi) = \sum_{1 \leq j \leq m} a_j$.*

The following example (see [3]) shows that the condition “ $\cap_{1 \leq j \leq m} \{f_j = 0\} = \{o\}$ ” can not be removed.

Example 1.14. *Let $I = (z_1^3 - z_2^2)$. Denote $X := \mathbb{C}[z_1, z_2]/I$. Then o and $(1, 1)$ are points in X where o is a singular point and $(1, 1)$ is a smooth point. Note that $z_1^3 - z_2^2$ is irreducible in $\mathbb{C}[z_1, z_2]$, $\mathcal{O}_{\mathbb{C}^2, o}$ and $\mathcal{O}_{\mathbb{C}^2, (1,1)}$. We know that I is a prime ideal in $\mathbb{C}[z_1, z_2]$ and (X, o) , $(X, (1,1))$ is irreducible as a germ of analytic set.*

Let $h_1 = [z_1]$, $h_2 = [z_2]$, $h_3 = [z_1 z_2]$, $g_1 = [z_1 - 1]$ and $g_2 = [z_2 - 1]$ on X . Take $\{f_1, f_2, f_3, f_4, f_5, f_6\} = \{h_1 g_1, h_1 g_2, h_2 g_1, h_2 g_2, h_3 g_1, h_3 g_2\}$ and $a_1 = \dots = a_6 = 1$. It is clear that

$$\cap_{1 \leq l \leq 6} \{f_l = 0\} = \{o, (1, 1)\}.$$

Let $t \in \mathbb{C} \setminus \{0\}$ and $z_1 = t^2$ and $z_2 = t^3$. Then t is a coordinate on X_{reg} . Note that the point $(1, 1) \in X$ corresponds to $t = 1$. As $\log F = \log(\prod_{1 \leq l \leq 6} f_l) = 3 \log(|z_1 - 1| |z_2 - 1|) + O(1)$ and $\varphi = \log(\sum_{1 \leq l \leq 6} |f_l|^{\frac{1}{a_l}}) = \log(|z_1 - 1| + |z_2 - 1|) + O(1)$ near $(1, 1)$, then we know that $\log |F| = 6 \log |t - 1| + O(1)$ and $\varphi = \log |t - 1| + O(1)$ near $t = 1$. Then we have

$$\sigma_{(1,1)}(\log |F|, \varphi) = 6 = \sum_{1 \leq l \leq 6} a_l$$

holds on X near $(1, 1)$.

By Theorem 1.12, there exists a valuation ν on $\mathbb{C}[z_1, z_2]/I$ such that $\nu(f_l) = a_l$ for every $1 \leq l \leq 6$. Since $F = 4 \log |z_1 z_2| + O(1)$ near o and $\varphi = \log(|z_1| + |z_2| + |z_1 z_2|) + O(1)$ near o in \mathbb{C}^2 , we know that

$$\sup\{c: \log |F| \leq c\varphi + O(1) \text{ near } o \text{ on } \mathbb{C}^2\} > 6 = \sum_{1 \leq l \leq 6} a_l.$$

Note that the relative type may increase after restriction. We have

$$\sigma_o(\log |F|, \varphi) > 6 = \sum_{1 \leq l \leq 6} a_l.$$

Thus, the condition “ $\cap_{1 \leq j \leq m} \{f_j = 0\} = \{o\}$ ” in Theorem 1.13 can not be removed.

Denote the set of all germs of real analytic functions near the origin $o' \in \mathbb{R}^n$ by $C_{o'}^{\text{an}}$. There exists an injective ring homomorphism $P: C_{o'}^{\text{an}} \rightarrow \mathcal{O}_o$ that satisfies

$$P\left(\sum_{\alpha \in \mathbb{Z}_{\geq 0}^n} a_\alpha x^\alpha\right) = \sum_{\alpha \in \mathbb{Z}_{\geq 0}^n} a_\alpha z^\alpha,$$

where (x_1, \dots, x_n) and (z_1, \dots, z_n) are the standard coordinates on \mathbb{R}^n and \mathbb{C}^n respectively, and $\sum_{\alpha \in \mathbb{Z}_{\geq 0}^n} a_\alpha x^\alpha$ is the power series expansion of an arbitrarily given real analytic function near o' . It is clear that, for any $(h, o) \in \mathcal{O}_o$, there exists a unique pair of real analytic functions (h_1, h_2) near o' such that

$$h = P(h_1) + iP(h_2) \quad \text{near } o.$$

Assume that I is an ideal in $C_{o'}^{\text{an}}$. Denote by $(P(I))$ the ideal generated by $P(I)$ in \mathcal{O}_o . Assume that $(P(I))$ is prime ideal in \mathcal{O}_o . Note that $(P(I))$ is a prime ideal in \mathcal{O}_o implies I is a prime ideal in $C_{o'}^{\text{an}}$ (see Remark 2.24). Then we have an induced ring homomorphism $\tilde{P}: C_{o'}^{\text{an}}/I \rightarrow \mathcal{O}_o/(P(I))$ defined by $\tilde{P}([f]) = [P(f)]$. It is easy to check that \tilde{P} is well defined. Let X be the zero variety defined by $(P(I))$.

Corollary 1.15. *Let $f_j \in C_{o'}^{\text{an}}/I$ ($0 \leq j \leq m$) defined near o' . Given $a_0 = 0$ and m positive numbers $\{a_j\}_{1 \leq j \leq m}$, the following two statements are equivalent:*

- (1) *There exists a valuation ν on $C_{o'}^{\text{an}}$ such that $\nu(f_j) = a_j$ for all $0 \leq j \leq m$;*
- (2) *$\sigma(\log |F|, \varphi) = \sum_{1 \leq j \leq m} a_j$, where*

$$F := \prod_{0 \leq j \leq m} \tilde{P}(f_j), \quad \varphi := \log \left(\sum_{1 \leq j \leq m} |\tilde{P}(f_j)|^{\frac{1}{a_j}} \right).$$

Remark 1.16. *Theorem 1.9 and Corollary 1.15 show that for any valuation ν on $C_{o'}^{\text{an}}/I$ and any finite collection of real analytic functions $\{f_j\}_{0 \leq j \leq m}$ near o' with $\nu(f_j) > 0$ for all j , there exists a valuation $\tilde{\nu}$ on $\mathcal{O}_o/(\tilde{P}(I))$ such that $\tilde{\nu}(\tilde{P}(f_j)) = \nu(f_j)$ for all j .*

Let I be an ideal in $\mathbb{R}[x_1, \dots, x_n]$. Denote the restriction of P on $\mathbb{R}[x_1, \dots, x_n]$ also by P and \tilde{P} on $\mathbb{R}[x_1, \dots, x_n]/I$ also by \tilde{P} . Assume that $(P(I))$ is a prime ideal in $\mathbb{C}[x_1, \dots, x_n]$. Note that $(P(I))$ is a prime ideal in $\mathbb{C}[z_1, \dots, z_n]$ implies I is a prime ideal in $\mathbb{R}[z_1, \dots, z_n]$ (see Remark 2.28). Denote X be the affine variety defined by $(P(I))$. Assume that $o \in X$ and (X, o) is irreducible as a germ of analytic set. Let $\{f_j\}_{0 \leq j \leq m} \subset \mathbb{R}[x_1, \dots, x_n]/I$, $a_0 = 0$ and $\{a_j\}_{1 \leq j \leq m}$ be m positive numbers. Set $F := \prod_{0 \leq j \leq m} \tilde{P}(f_j)$ and $\varphi := \log(\sum_{1 \leq j \leq m} |\tilde{P}(f_j)|^{\frac{1}{a_j}})$.

Corollary 1.17. *If $\sigma(\log |F|, \varphi) = \sum_{1 \leq j \leq m} a_j$, then there exists a valuation ν on $\mathbb{R}[x_1, \dots, x_n]/I$ such that $\nu(f_j) = a_j$ for all $0 \leq j \leq m$.*

Conversely, if there exists a valuation ν on $\mathbb{R}[x_1, \dots, x_n]/I$ satisfying that $\nu(f_j) = a_j$ for all $0 \leq j \leq m$ and $\nu(x_l) > 0$ for every $1 \leq l \leq n$, then we have $\sigma(\log |F|, \varphi) = \sum_{1 \leq j \leq m} a_j$.

2. PREPARATION

2.1. Basic knowledge. In this subsection, we recall some basic results which will be used when we discuss Zhou weights near singular points.

Let X be an analytic subset of $\Omega \in \mathbb{C}^n$ with pure dimension d . Let

$$dV_X := \frac{1}{2^d d!} \bigwedge^d \left(\sum_{1 \leq j \leq n} \sqrt{-1} dz_j \wedge d\bar{z}_j \right)$$

be the volume form on X induced by the standard volume form on \mathbb{C}^n .

Let (u_α) be a family of plurisubharmonic functions on an open subset of X , and we assume that the family (u_α) is locally uniformly bounded from above. We recall the following Choquet's Lemma.

Lemma 2.1 (see [10]). *Every family (u_α) has a countable subfamily $(v_j) = (u_{\alpha(j)})$, such that its upper-envelope $v = \sup_j v_j$ satisfies $v \leq u \leq u^* = v^*$, where $u = \sup_\alpha u_\alpha$, $u^*(z) := \lim_{\varepsilon \rightarrow 0} \sup_{U_{z,\varepsilon}} u$ and $v^*(z) := \lim_{\varepsilon \rightarrow 0} \sup_{U_{z,\varepsilon}} v$ are the regularizations of u and v , where $U_{z,\varepsilon} := X \cap \mathbb{B}^n(z, \varepsilon)$.*

Moreover, the upper regularization u^* is plurisubharmonic and equals almost everywhere to u .

Remark 2.2. Note that the original version of Lemma 2.1 is stated for smooth X . When X has singularities, since we ignore the values of plurisubharmonic functions on singularities and (u_α) is locally uniformly bounded from above, we apply upper semicontinuous regularization on X_{reg} and the same argument as the smooth case shows the Lemma 2.1 still holds in the singular case.

Let $z \in X$ be a smooth point. Let φ and φ_0 be plurisubharmonic functions on X defined near z and f be a holomorphic function defined on X near z . The following result considers the growth of the volume of sub-level sets of plurisubharmonic functions.

Lemma 2.3 (see [18]). *Assume that $|f_0|^2 e^{-2(\varphi+\varphi_0)}$ is not integrable near z , and $|f_0|^2 e^{-2\varphi_0}$ is integrable near z . Then for any small enough neighborhood U of z , there exists $C > 0$ such that*

$$e^{2t} \int_{\{\varphi < -t\} \cap U} |f_0|^2 e^{-2\varphi_0} > C$$

for any $t \geq 0$.

We recall the following desingularization theorem due to Hironaka.

Theorem 2.4 ([20], see also [5]). *Let Y be a complex manifold, and M be an analytic sub-variety in Y . Then there is a local finite sequence of blow-ups $\mu_j: Y_{j+1} \rightarrow Y_j$ ($Y_1 := Y, j = 1, 2, \dots$) with smooth centers S_j such that:*

- (1) *Each component of S_j lies either in $(M_j)_{\text{sing}}$ or in $M_j \cap E_j$, where $M_1 := M$, M_{j+1} denotes the strict transform of M_j by μ_j , $(M_j)_{\text{sing}}$ denotes the singular set of M_j , and E_{j+1} denotes the exceptional divisor $\mu_j^{-1}(S_j \cup E_j)$;*
- (2) *Let M' and E' denote the final strict transform of M and the exceptional divisor respectively. Then:
 - (a) *The underlying point-set $|M'|$ is smooth;*
 - (b) *$|M'|$ and E' simultaneously have only normal crossings;**

The (b) in the above theorem means that, locally, there is a coordinate system in which E' is a union of coordinate hyperplanes and $|M'|$ is a coordinate subspace.

We recall that every locally bounded plurisubharmonic function can be extended across closed pluripolar sets.

Lemma 2.5 (see [10, Theorem 5.24]). *Let φ be a plurisubharmonic function on $\Omega \setminus E$, where Ω is a domain in \mathbb{C}^n and E is a closed pluripolar set in Ω . If φ is locally bounded above near E , then φ extends uniquely to a plurisubharmonic function on Ω .*

Recall that X is an analytic subset of $\Omega \in \mathbb{C}^n$ with pure dimension d . Let $z_0 \in X$ be a singular point. Let φ and φ_0 be plurisubharmonic functions on X defined near z_0 and f be a holomorphic function defined on X near z_0 . Recall that

$$dV_X := \frac{1}{2^d d!} \bigwedge^d \left(\sum_{1 \leq j \leq n} \sqrt{-1} dz_j \wedge d\bar{z}_j \right)$$

be the volume form on X induced by the standard volume form on \mathbb{C}^n .

Using Lemma 2.3, Lemma 2.5 and Theorem 2.4, we can generalize Lemma 2.3 to the singular case.

Lemma 2.6. *Assume that $|f_0|^2 e^{-2(\varphi+\varphi_0)}$ is not integrable near z_0 , and $|f_0|^2 e^{-2\varphi_0}$ is integrable near z_0 . Then for any small enough neighborhood $U \subset X$ of z_0 , there exists $C > 0$ such that*

$$e^{2t} \int_{\{\varphi < -t\} \cap U} |f_0|^2 e^{-2\varphi_0} > C$$

for any $t \geq 0$.

Proof. We use Theorem 2.4 on Ω to resolve the singularities of X , and denote the corresponding proper modification by $\mu: \tilde{\Omega} \rightarrow \Omega$. Denote the strict transform of X by \tilde{X} and denote $Z_0 := \tilde{X} \cap \mu^{-1}(\{z_0\})$.

As $|f_0|^2 e^{-2(\varphi+\varphi_0)}$ is not integrable near z_0 , there exists $\tilde{z}_0 \in Z_0$ such that

$$\int_U \mu^*(|f_0|^2 e^{-2(\varphi+\varphi_0)} dV_X) = +\infty$$

for any neighborhood $U \subset \tilde{X}$ of \tilde{z}_0 . Note that $\varphi \circ \mu$ and $\varphi_0 \circ \mu$ are plurisubharmonic functions on \tilde{X} (where we can use Lemma 2.5 to extend $\varphi \circ \mu$ and $\varphi_0 \circ \mu$ from $X \setminus \mu^{-1}(X_{\text{sing}})$ to X), and there exist holomorphic functions $\tilde{f}_1, \dots, \tilde{f}_s$ near \tilde{z}_0 such that $\mu^*(|f_0|^2 dV_X) = \sum_{1 \leq j \leq s} |\tilde{f}_j|^2 dV_{\tilde{X}}$, where $dV_{\tilde{X}}$ is a volume form on \tilde{X} . By Lemma 2.3, for any small neighborhood $U' \subset \tilde{X}$ of \tilde{z}_0 , there exists $C > 0$ such that

$$e^{2t} \int_{\{\mu^{-1}(\varphi) < -t\} \cap U'} \mu^*(|f_0|^2 e^{-2\varphi_0} dV_X) > C$$

for any $t \geq 0$, which proves Lemma 2.6. \square

The following result is a corollary of Guan–Zhou’s lower semicontinuity property of plurisubharmonic functions with a multiplier ideal in [18].

Theorem 2.7 ([18]). *Let f be a holomorphic function near z_0 , and $\{\varphi_j\}$ be a sequence of plurisubharmonic functions on a neighborhood $V \subset X$ of z_0 . Assume that $\{\varphi_j\}$ is increasingly convergent to a plurisubharmonic function φ almost everywhere with respect to Lebesgue measure. If $|f_0|^2 e^{-2\varphi_j}$ is not integrable near z_0 for any j , then $|f_0|^2 e^{-2\varphi}$ is not integrable near z_0 .*

Let φ and φ_0 be plurisubharmonic functions near $z_0 \in X$ such that $|f_0|^2 e^{-2\varphi_0}$ is integrable near z_0 . Assume that $\varphi \geq N \log |z| + O(1)$ near z_0 for large enough $N \gg 0$.

Lemma 2.8. *Assume that $(f_0, z_0) \notin \mathcal{I}(\varphi + \varphi_0)_{z_0}$, and $(f_0, z_0) \in \mathcal{I}((1-\varepsilon)\varphi + \varphi_0)_{z_0}$ for any $\varepsilon \in (0, 1)$. Then for any neighborhood $U \subset X$ of z_0 ,*

$$\lim_{t \rightarrow +\infty} \frac{-\log \int_{\{\varphi < -t\} \cap U} |f_0|^2 e^{-2\varphi_0}}{2t} = 1.$$

Proof. As $(f_0, z_0) \notin \mathcal{I}(\varphi + \varphi_0)_{z_0}$, we have

$$\liminf_{t \rightarrow +\infty} e^{2t} \int_{\{\varphi < -t\} \cap U} |f_0|^2 e^{-2\varphi_0} > 0$$

(see Lemma 2.6), which implies that

$$\limsup_{t \rightarrow +\infty} \frac{-\log \int_{\{\varphi < -t\} \cap U} |f_0|^2 e^{-2\varphi_0}}{2t} \leq 1.$$

As $\varphi \geq N \log |z| + O(1)$ for large enough $N \gg 0$ and $(f_0, o) \in \mathcal{I}((1-\varepsilon)\varphi + \varphi_0)_o$ for any $\varepsilon > 0$, we have

$$\begin{aligned} & \limsup_{t \rightarrow +\infty} e^{2(1-\varepsilon)t} \int_{\{\varphi < -t\} \cap U} |f_0|^2 e^{-2\varphi_0} \\ & \leq \limsup_{t \rightarrow +\infty} \int_{\{\varphi < -t\} \cap U} |f_0|^2 e^{-2\varphi_0 - 2(1-\varepsilon)\varphi} \\ & = 0, \end{aligned}$$

which implies that

$$\liminf_{t \rightarrow +\infty} \frac{-\log \int_{\{\varphi < -t\} \cap U} |f_0|^2 e^{-2\varphi_0}}{2t} \geq 1 - \varepsilon.$$

Then we have

$$\lim_{t \rightarrow +\infty} \frac{-\log \int_{\{\varphi < -t\} \cap U} |f_0|^2 e^{-2\varphi_0}}{2t} = 1.$$

□

2.2. Concavity. Let (X, z_0) be an irreducible germ of an analytic set. Let u and v be Lebesgue measurable functions on X with upper-bounds near z_0 . Let g be a nonnegative Lebesgue measurable function on X .

The following Lemma 2.9 and 2.10 come from real analysis. When z_0 is a singular point of X , the proof of Lemma 2.9 and 2.10 are the same as the original proof in the smooth case and hence we omit the proofs.

Lemma 2.9 (see [9]). *Assume that $g^2 e^{2(l_1 v - (1+l_2)u)}$ is integrable near z_0 , where $l_1, l_2 > 0$. Then $g^2 e^{-2u} - g^2 e^{-2 \max\{u, \frac{l_1}{l_2}v\}}$ is integrable on a small enough neighborhood of z_0 .*

Denote by

$$A_{u,v}(t) := \sup \{c: g^2 e^{2(tv - cu)} \text{ is integrable near } z_0\}.$$

Note that $A_{u,v}(t)$ is increasing with respect to t (maybe $+\infty$ or $-\infty$). Assume that $A_{u,v}(t) \in (0, +\infty]$ on $(t_0 - \delta, t_0 + \delta)$. It follows from Hölder's inequality that $A_{u,v}(t)$ is concave on $(t_0 - \delta, t_0 + \delta)$. We recall the following basic property of $A_{u,v}(t)$.

Lemma 2.10 (see [2, Lemma 2.14]). *Assume that $A_{u,v}(t)$ is strictly increasing on $(t_0 - \delta, t_0 + \delta)$. Then for any $b > 0$,*

$$A_{\max\{u, \frac{1}{b}v\}, v}(t_0) = A_{u,v}(t_0)$$

holds if and only if $b \in \left(0, \lim_{\Delta t \rightarrow 0-0} \frac{A_{u,v}(t_0 + \Delta t) - A_{u,v}(t_0)}{\Delta t}\right]$.

2.3. Tian functions and Zhou numbers. Let φ, ψ, φ_0 be plurisubharmonic functions near z_0 , and let $f_0 = (f_{0,1}, \dots, f_{0,m})$ be a vector, where $f_{0,1}, \dots, f_{0,m}$ are holomorphic functions near z_0 .

Denote by

$$c_{z_0}(\varphi, t\psi) := \sup \{c: |f_0|^2 e^{-2\varphi_0} e^{2t\psi} e^{-2c\varphi} \text{ is integrable near } z_0\},$$

which is a generalization of the jumping number (see [21, 22]). Define the *Tian function*

$$\text{Tn}(t) := c_{z_0}(\varphi, t\psi)$$

for any $t \in \mathbb{R}$. The Hölder inequality shows that $\text{Tn}(t)$ is concave with respect to $t \in (-\infty, +\infty)$.

In the following lemma, assume that the following three statements hold

- (1) $|f_0|^2 e^{-2\varphi_0}$ is integrable near z_0 ;
- (2) There exists integer $N_0 \gg 0$ such that $|f_0|^2 e^{-2\varphi_0} |z|^{2N_0} e^{-2\text{Tn}(0)\varphi}$ is integrable near z_0 ;
- (3) There exists $s_0 > 0$, such that $\psi \leq s_0 \log |z| + O(1)$ near $z_0 \in X$.

In this subsection, we discuss the derivatives of Tian functions $\text{Tn}(t)$.

We give the strictly increasing property of Tian functions in the following lemma.

Lemma 2.11. *$\text{Tn}(t)$ is strictly increasing near 0.*

Proof. Theorem 2.7 implies that there exists $\varepsilon_0 > 0$ such that

$$|f_0|^2 e^{-2\varphi_0} |z|^{2N_0} e^{-2(1+\varepsilon_0)\text{Tn}(0)\varphi}$$

is integrable near z_0 . Lemma 2.9 shows that

$$|f_0|^2 e^{-2\varphi_0} e^{-2\left(1+\frac{\varepsilon_0}{2}\right)\text{Tn}(0)\varphi} - |f_0|^2 e^{-2\varphi_0} e^{-2 \max \left\{ \left(1+\frac{\varepsilon_0}{2}\right)\text{Tn}(0)\varphi, \frac{2+\varepsilon_0}{\varepsilon_0} \log |z|^{N_0} \right\}}$$

is integrable near z_0 . Note that for any $t > 0$,

$$|f_0|^2 e^{-2\varphi_0} \left(e^{2t\psi} e^{-2\left(1+\frac{\varepsilon_0}{2}\right)\text{Tn}(0)\varphi} - e^{2t\psi} e^{-2 \max \left\{ \left(1+\frac{\varepsilon_0}{2}\right)\text{Tn}(0)\varphi, \frac{2+\varepsilon_0}{\varepsilon_0} \log |z|^{N_0} \right\}} \right)$$

is integrable near z_0 and

$$\begin{aligned} & |f_0|^2 e^{-2\varphi_0} e^{2t\psi} e^{-2 \max \left\{ \left(1+\frac{\varepsilon_0}{2}\right)\text{Tn}(0)\varphi, \frac{2+\varepsilon_0}{\varepsilon_0} \log |z|^{N_0} \right\}} \\ & \leq C |f_0|^2 e^{-2\varphi_0} |z|^{2s_0 t} e^{-2 \max \left\{ \left(1+\frac{\varepsilon_0}{2}\right)\text{Tn}(0)\varphi, \frac{2+\varepsilon_0}{\varepsilon_0} \log |z|^{N_0} \right\}} \\ & \leq C |f_0|^2 e^{-2\varphi_0} |z|^{2s_0 t} e^{-2 \frac{(2+\varepsilon_0)N_0}{\varepsilon_0} \log |z|} \end{aligned}$$

near z_0 . Then it is clear that for any $t > \frac{2+\varepsilon_0}{\varepsilon_0} \frac{N_0}{s_0}$,

$$|f_0|^2 e^{-2\varphi_0} e^{2t\psi} e^{-2\left(1+\frac{\varepsilon_0}{2}\right)\text{Tn}(0)\varphi}$$

is integrable near z_0 , which shows $\text{Tn}(t) > \left(1+\frac{\varepsilon_0}{2}\right)\text{Tn}(0)$ for any $t > \frac{2+\varepsilon_0}{\varepsilon_0} \frac{N_0}{s_0}$. Then the concavity of $\text{Tn}(t)$ implies that $\text{Tn}(t)$ is strictly increasing near 0. \square

The following property of Tian functions $Tn(t)$ will be used in the proof of Theorem 1.6.

Proposition 2.12. *Assume that there exists $N \gg 0$ such that $\varphi \geq N \log |z|$ near o . The following inequality holds*

$$\begin{aligned} & \frac{1}{Tn(0)} \lim_{t \rightarrow 0+0} \frac{Tn(0) - Tn(t)}{-t} \\ & \leq \liminf_{t_1 \rightarrow +\infty} \frac{1}{2t_1} \frac{\int_{\{Tn(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0} (-2\psi)}{\int_{\{Tn(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0}} \\ & \leq \limsup_{t_1 \rightarrow +\infty} \frac{1}{2t_1} \frac{\int_{\{Tn(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0} (-2\psi)}{\int_{\{Tn(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0}} \\ & \leq \frac{1}{Tn(0)} \lim_{t \rightarrow 0+0} \frac{Tn(0) - Tn(-t)}{t}. \end{aligned}$$

Proof. We prove Proposition 2.12 in two steps.

Step 1. Theorem 2.7 shows that for a small enough constant $t > 0$ (t is dependent on $|f_0|^2 e^{-2\varphi_0}$ and ψ), $|f_0|^2 e^{-2\varphi_0} e^{-2t\psi}$ is integrable near z_0 . Since $e^{-2t\psi} \geq (-2t\psi)$, we have

$$|f_0|^2 e^{-2\varphi_0} (-2\psi) = \frac{1}{t} |f_0|^2 e^{-2\varphi_0} (-2t\psi) \leq \frac{1}{t} |f_0|^2 e^{-2\varphi_0} e^{-2t\psi}$$

is integrable near z_0 . Note that there exists $N \gg 0$ such that $\varphi \geq N \log |z|$ near z_0 . There exists a neighborhood U of z_0 such that for any small enough $t > 0$ and $\varepsilon > 0$ (ε depends on t),

$$\limsup_{t_1 \rightarrow +\infty} \int_{\{Tn(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0} e^{-2(t\psi + (1-\varepsilon)Tn(-t)\varphi)} = 0,$$

which implies that

$$\limsup_{t_1 \rightarrow +\infty} e^{2t_1} \int_{\{Tn(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0} e^{-2(t\psi + (1-\varepsilon)Tn(-t)\varphi + t_1)} = 0.$$

Then for large enough $t_1 > 0$,

$$\int_{\{Tn(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0} e^{-2(t\psi + (1-\varepsilon)Tn(-t)\varphi + t_1)} < e^{-2t_1},$$

i.e.,

$$\log \left(\int_{\{Tn(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0} e^{-2(t\psi + (1-\varepsilon)Tn(-t)\varphi + t_1)} \right) < -2t_1.$$

Combining with Lemma 2.8, we obtain

$$\begin{aligned} & \limsup_{t_1 \rightarrow +\infty} \frac{1}{2t_1} \log \frac{\int_{\{Tn(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0} e^{-2(t\psi + (1-\varepsilon)Tn(-t)\varphi + t_1)}}{\int_{\{Tn(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0}} \\ (2.1) \quad & = \limsup_{t_1 \rightarrow +\infty} \frac{1}{2t_1} \log \int_{\{Tn(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0} e^{-2(t\psi + (1-\varepsilon)Tn(-t)\varphi + t_1)} \\ & - \lim_{t_1 \rightarrow +\infty} \frac{1}{2t_1} \log \int_{\{Tn(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0} \\ & \leq 1 - 1 = 0. \end{aligned}$$

It follows from Jensen's inequality and the concavity of the logarithm function that

$$\begin{aligned}
& \log \frac{\int_{\{\mathrm{Tn}(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0} e^{-2(t\psi + (1-\varepsilon)\mathrm{Tn}(-t)\varphi + t_1)}}{\int_{\{\mathrm{Tn}(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0}} \\
& \geq \frac{\int_{\{\mathrm{Tn}(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0} \log(e^{-2(t\psi + (1-\varepsilon)\mathrm{Tn}(-t)\varphi + t_1)})}{\int_{\{\mathrm{Tn}(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0}} \\
& = \frac{\int_{\{\mathrm{Tn}(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0} (-2(t\psi + (1-\varepsilon)\mathrm{Tn}(-t)\varphi + t_1))}{\int_{\{\mathrm{Tn}(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0}} \\
& \geq \frac{\int_{\{\mathrm{Tn}(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0} (-2(t\psi + (1-\varepsilon)\mathrm{Tn}(-t)(-t_1)\frac{1}{\mathrm{Tn}(0)} + t_1))}{\int_{\{\mathrm{Tn}(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0}}.
\end{aligned}$$

Combining with inequality (2.1), we obtain that

$$\limsup_{t_1 \rightarrow +\infty} \frac{1}{2t_1} \frac{\int_{\{\mathrm{Tn}(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0} (-2(t\psi + (1-\varepsilon)\mathrm{Tn}(-t)(-t_1)\frac{1}{\mathrm{Tn}(0)} + t_1))}{\int_{\{\mathrm{Tn}(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0}} \leq 0.$$

Letting $\varepsilon \rightarrow 0+0$ and $t \rightarrow 0+0$, we obtain

$$(2.2) \quad \limsup_{t_1 \rightarrow +\infty} \frac{1}{2t_1} \frac{\int_{\{\mathrm{Tn}(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0} (-2\psi)}{\int_{\{\mathrm{Tn}(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0}} \leq \frac{1}{\mathrm{Tn}(0)} \lim_{t \rightarrow 0+0} \frac{\mathrm{Tn}(0) - \mathrm{Tn}(-t)}{t}.$$

Step 2. By a similar discussion in Step 1 (only replacing t by $-t$), we obtain that

$$\limsup_{t_1 \rightarrow +\infty} \frac{1}{2t_1} \frac{\int_{\{\mathrm{Tn}(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0} (-2(-t\psi + (1-\varepsilon)\mathrm{Tn}(t)(-t_1)\frac{1}{\mathrm{Tn}(0)} + t_1))}{\int_{\{\mathrm{Tn}(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0}} \leq 0.$$

Letting $\varepsilon \rightarrow 0+0$ and $t \rightarrow 0+0$, we obtain

$$(2.3) \quad \frac{1}{\mathrm{Tn}(0)} \lim_{t \rightarrow 0+0} \frac{\mathrm{Tn}(0) - \mathrm{Tn}(t)}{-t} \leq \liminf_{t_1 \rightarrow +\infty} \frac{1}{2t_1} \frac{\int_{\{\mathrm{Tn}(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0} (-2\psi)}{\int_{\{\mathrm{Tn}(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0}}.$$

Combining inequality (2.3) and inequality (2.2), we get Proposition 2.12. \square

Let $\Phi_{z_0, \max}$ be a local Zhou weight related to $|f_0|^2 e^{-2\varphi_0}$ near z_0 . Taking $\varphi = \Phi_{z_0, \max}$, we have $\mathrm{Tn}(0) = 1$.

Proposition 2.13. *For any plurisubharmonic function ψ satisfying that $\psi \leq s_0 \log |z| + O(1)$ near $z_0 \in X$ for some $s_0 > 0$, the Tian function is differentiable at $t = 0$, and*

$$\mathrm{Tn}(t) = \mathrm{Tn}(0) + \sigma(\psi, \Phi_{z_0, \max})t$$

for any $t \geq 0$. Especially,

$$\sigma(\psi, \Phi_{z_0, \max}) = \lim_{t \rightarrow 0} \frac{\mathrm{Tn}(t) - \mathrm{Tn}(0)}{t}.$$

Proof. As $\psi \leq s_0 \log |z| + O(1)$ near z_0 and $\Phi_{z_0, \max} \geq N \log |z| + O(1)$ near z_0 for some N , we have $\sigma(\psi, \Phi_{z_0, \max}) > 0$. Lemma 2.10 and the strong openness property (Theorem 2.7) show that $|f_0|^2 e^{-2\varphi_0} e^{-2 \max\{\Phi_{z_0, \max}, \frac{1}{b}\psi\}}$ is not integrable near z_0 if and only if $b \leq \lim_{t \rightarrow 0+0} \frac{\mathrm{Tn}(0) - \mathrm{Tn}(-t)}{t}$.

If $b < \sigma(\psi, \Phi_{z_0, \max})$, it follows from $\psi \leq b\Phi_{z_0, \max} + O(1)$ that $\max\{\frac{1}{b}\psi, \Phi_{z_0, \max}\} = \Phi_{z_0, \max} + O(1)$. Then $|f_0|^2 e^{-2\varphi_0} e^{-2\max\{\Phi_{z_0, \max}, \frac{1}{b}\psi\}}$ is not integrable near z_0 . If $b > \sigma(\psi, \Phi_{z_0, \max})$, by definition of $\sigma(\psi, \Phi_{z_0, \max})$, we have $\max\{\frac{1}{b}\psi, \Phi_{z_0, \max}\} \neq \Phi_{z_0, \max} + O(1)$ and $\max\{\frac{1}{b}\psi, \Phi_{z_0, \max}\} \geq \Phi_{z_0, \max} + O(1)$ near z_0 . Following from the definition of local Zhou weights, we know that $|f_0|^2 e^{-2\varphi_0} e^{-2\max\{\Phi_{z_0, \max}, \frac{1}{b}\psi\}}$ is integrable near z_0 . Then we have

$$(2.4) \quad \sigma(\psi, \Phi_{z_0, \max}) = \lim_{t \rightarrow 0+0} \frac{\text{Tn}(0) - \text{Tn}(-t)}{t}.$$

It follows from that $\psi \leq b\Phi_{z_0, \max} + O(1)$ near z_0 for any $b < \sigma(\psi, \Phi_{z_0, \max})$ that $\text{Tn}(t) \geq \text{Tn}(0) + \sigma(\psi, \Phi_{z_0, \max})t$ for any $t \geq 0$. By and equality (2.4) and the concavity of $\text{Tn}(t)$, we obtain that $\text{Tn}(t) = \text{Tn}(0) + \sigma(\psi, \Phi_{z_0, \max})t$ for any $t \geq 0$ and $\lim_{t \rightarrow 0} \frac{\text{Tn}(t) - \text{Tn}(0)}{t} = \sigma(\psi, \Phi_{z_0, \max})$.

Thus, Proposition 2.13 holds. \square

2.4. Relation between relative types and valuations. In this subsection, we consider the relation between relative types and valuations for holomorphic and plurisubharmonic functions.

We recall Skoda's division theorem.

Theorem 2.14 (see [9]). *Let Ω be a weakly pseudoconvex Kähler manifold of dimension n and φ be a plurisubharmonic function on Ω . Set $m = \min\{n, r - 1\}$. Then for every holomorphic function f on Ω such that*

$$I := \int_{\Omega} |f|^2 |g|^{-2(m+1+\epsilon)} e^{-\varphi} < +\infty,$$

there exist holomorphic functions (h_1, \dots, h_r) on Ω such that $f = \sum_{1 \leq j \leq r} g_j h_j$ and

$$\int_{\Omega} |h|^2 |g|^{-2(m+\epsilon)} e^{-\varphi} \leq \left(1 + \frac{m}{\epsilon}\right) I.$$

Recall that X is an analytic variety with pure dimension d contained in \mathbb{C}^n . Let $z = (z_1, \dots, z_n)$ be the coordinate on \mathbb{C}^n . Assume that the origin $o \in X_{\text{sing}}$ and (X, o) is an irreducible germ of an analytic set.

Denote $\mathcal{O}_{X,x}^w$ be the ring of germs of weakly holomorphic functions defined on X near x and $\mathfrak{M}_{X,x}$ be the ring of germs of meromorphic functions defined on X near x .

Theorem 2.15 (see [10]). *For every point $x \in X$, there is a neighborhood V of x and $h \in \mathcal{O}_X(V)$ such that $h^{-1}(0)$ is nowhere dense in V and $h_y \mathcal{O}_{X,y}^w \subseteq \mathcal{O}_{X,y}$ for all $y \in V$; such a function h is called a universal denominator on V . In particular $\mathcal{O}_{X,x}^w$ is contained in the ring $\mathfrak{M}_{X,x}$.*

Using Theorem 2.15, we have following lemma.

Lemma 2.16. *Let ν be a valuation on $\mathcal{O}_{X,x}$ and ν naturally extended to be a valuation on the quotient field $\mathfrak{M}_{X,x}$. For any $s \in \mathcal{O}_{X,x}^w \subset \mathfrak{M}_{X,x}$, the value $\nu(s)$ has a universal lower bound $-\nu(h)$, where h is the universal denominator of $\mathcal{O}_{X,x}^w$ near x .*

Proof. Let $s \in \mathcal{O}_{X,x}^w$ be the germ of any weakly holomorphic function defined near x . It follows from Theorem 2.15 that we can find an $h_x \in \mathcal{O}_{X,x}$ such that $h_x s_x \in \mathcal{O}_{X,x}$. Note that $\nu|_{\mathcal{O}_{X,x}} \geq 0$. Then $\nu(s) = \nu(hs) - \nu(h) \geq -\nu(h)$.

Lemma 2.16 has been proved. \square

Lemma 2.17. *For any valuation ν on $\mathcal{O}_{X,o}$ where $o \in X_{\text{sing}}$ and holomorphic functions f_0, f_1, \dots, f_r belonging to $\mathcal{O}_{X,o}$ with $\nu(f_j) > 0$ for $1 \leq j \leq r$, we have $\sigma(\log |f_0|, \varphi) \leq \nu(f_0)$, where $\varphi := \log \left(\sum_{1 \leq j \leq r} |f_j|^{\frac{1}{\nu(f_j)}} \right)$.*

Proof. We prove this lemma by contradiction: if not, there exists $\delta_1 > 0$ such that $\log |f_0| \leq (\nu(f_0) + \delta_1)\varphi + O(1)$ near o . Then we can find some $\delta_2 > 0$ and a set of positive integers $\{m_0, \dots, m_r\}$ such that

$$\log |f_0| \leq \frac{\nu(f_0) + \delta_2}{m_0} \tilde{\varphi} + O(1) \quad \text{near } o,$$

where $\tilde{\varphi} := \log \left(\sum_{1 \leq j \leq r} |f_j|^{m_j} \right)$ and $\nu(f_j^{m_j}) \geq m_0$ for $1 \leq j \leq r$. For any positive integer l , we have

$$\log |f_0^l| \leq \frac{\nu(f_0^l) + l\delta_2}{m_0} \tilde{\varphi} + O(1)$$

holds on $(U \cap X) \setminus X_{\text{sing}}$, where $U := \{x \in \mathbb{C}^n : \log |z| < \delta\}$.

We use Theorem 2.4 on Ω to resolve the singularities of X , and denote the corresponding proper modification by $\mu: \tilde{\Omega} \rightarrow \Omega$. Denote the strict transform of X by \tilde{X} and denote $Z_0 := \tilde{\Omega} \cap \mu^{-1}(\{o\})$. Then $\mu^{-1}(U) \cap \tilde{X}$ contains $Z_0 \cap \tilde{X}$ and

$$(2.5) \quad \log |f_0^l \circ \mu| \leq \frac{\nu(f_0^l) + l\delta_2}{m_0} \tilde{\varphi} \circ \mu + O(1)$$

holds on $\mu^{-1}(U) \cap \tilde{X}$. Note that $Z_0 \cap \tilde{X}$ is compact in \tilde{X} , taking δ small enough, we may assume that $\mu^{-1}(U) \cap \tilde{X}$ is relatively compact in \tilde{X} . Inequality (2.5) and the relative compactness of $\mu^{-1}(U) \cap \tilde{X}$ imply that there exists a small $\epsilon > 0$ (independent of l) such that

$$(2.6) \quad \int_{\mu^{-1}(U) \cap \tilde{X}} |f_0^l \circ \mu|^2 e^{-2\frac{\nu(f_0^l) + l\delta_2 + \epsilon}{m_0} \tilde{\varphi} \circ \mu} < +\infty.$$

Let $|F| := \sum_{1 \leq j \leq r} |f_j|^{m_j} \circ \mu$, $\epsilon_0 := \frac{\epsilon}{m_0}$ and $m := \min\{n, r-1\}$. By inequality (2.6), we have

$$\begin{aligned} & \int_{\mu^{-1}(U) \cap \tilde{X}} |f_0^l \circ \mu|^2 |F|^{-2(m+1)-2\epsilon_0} |F|^{-2\left(\frac{\nu(f_0^l) + l\delta_2}{m_0} - (n+1)\right)} \\ & \leq C \int_{\mu^{-1}(U) \cap \tilde{X}} |f_0^l \circ \mu|^2 |F|^{-2(n+1)-2\epsilon_0} |F|^{-2\left(\frac{\nu(f_0^l) + l\delta_2}{m_0} - (n+1)\right)} \\ & = C \int_{\mu^{-1}(U) \cap \tilde{X}} |f_0^l \circ \mu|^2 e^{-2\frac{\nu(f_0^l) + l\delta_2 + \epsilon}{m_0} \tilde{\varphi} \circ \mu} < +\infty. \end{aligned}$$

Note that $(\log |z \circ \mu|)|_{\tilde{X}}$ is a smooth plurisubharmonic function on \tilde{X} and $\mu^{-1}(U) \cap \tilde{X} = \{x \in \tilde{X} : (\log |z \circ \mu|)|_{\tilde{X}} < \delta\}$. We know that $\mu^{-1}(U) \cap \tilde{X}$ is a weakly pseudoconvex Kähler manifold.

By Theorem 2.14, there exist holomorphic functions (h_1, \dots, h_r) on $\mu^{-1}(U) \cap \tilde{X}$ such that $f_0^l \circ \mu = \sum_{1 \leq j \leq r} (f_j \circ \mu)^{m_j} h_j$ holds on $\mu^{-1}(U) \cap \tilde{X}$ and

$$(2.7) \quad \begin{aligned} & \int_{\mu^{-1}(U) \cap \tilde{X}} |h|^2 |F|^{-2(m+1)-2\epsilon_0} |F|^{-2\left(\frac{\nu(f_0^l) + l\delta_2}{m_0} - 1 - (n+1)\right)} \\ & = \int_{\mu^{-1}(U) \cap \tilde{X}} |h|^2 |F|^{-2m-2\epsilon_0} |F|^{-2\left(\frac{\nu(f_0^l) + l\delta_2}{m_0} - (n+1)\right)} < +\infty. \end{aligned}$$

It follows from the estimate (2.7) that, using Theorem 2.14 for each h_j ($1 \leq j \leq r$), there exist holomorphic functions $(h_{j,1}, \dots, h_{j,r})$ on $\mu^{-1}(U) \cap \tilde{X}$ such that $h_j = \sum_{1 \leq k \leq r} h_{j,k} (f_k \circ \mu)^{m_k}$ holds on $\mu^{-1}(U) \cap \tilde{X}$ and

$$\int_{\mu^{-1}(U) \cap \tilde{X}} \sum_{1 \leq k \leq r} |h_{j,k}|^2 |F|^{-2m-2\epsilon_0} |F|^{-2\left(\frac{\nu(f_0^l)+l\delta_2}{m_0}-1-(n+1)\right)} < +\infty.$$

Since $f_0^l \circ \mu = \sum_{1 \leq j \leq r} (f_j \circ \mu)^{m_j} h_j$ and $h_j = \sum_{1 \leq k \leq r} h_{j,k} (f_k \circ \mu)^{m_k}$ hold on $\mu^{-1}(U) \cap \tilde{X}$, we have

$$f_0^l \circ \mu = \sum_{1 \leq j, k \leq r} (f_j \circ \mu)^{m_j} (f_k \circ \mu)^{m_k} h_{j,k}$$

holds on $\mu^{-1}(U) \cap \tilde{X}$.

Denote $k := \lfloor \frac{\nu(f_0^l)+l\delta_2}{m_0} \rfloor - n - 1$. Using Theorem 2.14 k times, we have

$$(2.8) \quad f_0^l \circ \mu = \sum_{1 \leq j_1, j_2, \dots, j_k \leq r} (f_{j_1} \circ \mu)^{m_{j_1}} (f_{j_2} \circ \mu)^{m_{j_2}} \cdots (f_{j_k} \circ \mu)^{m_{j_k}} h_J$$

hold on $\mu^{-1}(U) \cap \tilde{X}$, where $J = (j_1, j_2, \dots, j_k)$ is a multi-index and h_J is an L^2 integrable holomorphic function (depending on l) on $\mu^{-1}(U) \cap \tilde{X}$ for any J . As h_J is L^2 integrable on $\mu^{-1}(U) \cap \tilde{X}$, we know that $|h_J|$ is bounded near $Z_0 \cap \tilde{X}$. Then $\mu_*(h_J)$ is a weakly holomorphic function defined on $U \cap X$. It follows from the decomposition (2.8) that we have

$$f_0^l = \sum_{1 \leq j_1, j_2, \dots, j_k \leq r} (f_{j_1})^{m_{j_1}} (f_{j_2})^{m_{j_2}} \cdots (f_{j_k})^{m_{j_k}} \mu_*(h_J)$$

holds on $U \cap X$. Hence, for the germ level, we know

$$(f_0^l)_o = \sum_{1 \leq j_1, j_2, \dots, j_k \leq r} (f_{j_1}^{m_{j_1}})_o (f_{j_2}^{m_{j_2}})_o \cdots (f_{j_m}^{m_{j_m}})_o (\mu_*(h_J))_o$$

holds in $\mathfrak{M}_{X,o}$, where $\mathfrak{M}_{X,o}$ is the ring of germs of meromorphic functions near o .

Lemma 2.16 tells that $\nu(h_J) \geq -C$ for some constant $C > 0$ (independent of f_0 and l) for any J . Note that $\nu(f_j^{m_j}) \geq m_0$ for $1 \leq j \leq r$ and $k := \lfloor \frac{\nu(f_0^l)+l\delta_2}{m_0} \rfloor - n - 1$. We have

$$\begin{aligned} \nu(f_0^l) &\geq \min_{1 \leq j_1, j_2, \dots, j_k \leq r} \{ \nu(f_{j_1}^{m_{j_1}} f_{j_2}^{m_{j_2}} \cdots f_{j_m}^{m_{j_m}} \mu_*(h_J)) \} \\ &\geq m_0 k - C \\ &\geq m_0 \left(\frac{\nu(f_0^l) + l\delta_2}{m_0} - n - 2 \right) - C \\ &= \nu(f_0^l) + l\delta_2 - m_0(n+2) - C, \end{aligned}$$

which implies $l \leq \frac{m_0(n+2)}{\delta_2} + C$. This is a contradiction since we can choose l arbitrarily large. Then we must have $\sigma(\log |f_0|, \varphi) \leq \nu(f_0)$. \square

When we consider valuations on the ring of germs of weakly holomorphic functions, we have the following similar result as Lemma 2.17.

Lemma 2.18. *For any valuation ν on $\mathcal{O}_{X,o}^w$ where $o \in X_{\text{sing}}$ and holomorphic functions f_0, f_1, \dots, f_r belonging to $\mathcal{O}_{X,o}$ with $\nu(f_j) > 0$ for $1 \leq j \leq r$, we have $\sigma(\log |f_0|, \varphi) \leq \nu(f_0)$, where $\varphi := \log \left(\sum_{1 \leq j \leq r} |f_j|^{\frac{1}{\nu(f_j)}} \right)$.*

Proof. The proof of Lemma 2.18 is almost the same as the proof of Lemma 2.17. We just give the sketch, and when there are some differences, we will discuss in detail.

We prove Lemma 2.18 by contradiction. Assume that the conclusion does not hold. Then there exists $\delta_1 > 0$ such that $\log |f_0| \leq (\nu(f_0) + \delta_1)\varphi + O(1)$ near o . Then we can find some $\delta_2 > 0$ and a set of positive integers $\{m_0, \dots, m_r\}$ such that

$$\log |f_0| \leq \frac{\nu(f_0) + \delta_2}{m_0} \tilde{\varphi} + O(1) \quad \text{near } o,$$

where $\tilde{\varphi} := \log (\sum_{1 \leq j \leq r} |f_j^{m_j}|)$ and $\nu(f_j^{m_j}) \geq m_0$ for $1 \leq j \leq r$. For any positive integer l , we have

$$\log |f_0^l| \leq \frac{\nu(f_0^l) + l\delta_2}{m_0} \tilde{\varphi} + O(1)$$

holds on $(U \cap X) \setminus X_{\text{sing}}$, where $U := \{x \in \mathbb{C}^n : \log |z| < \delta\}$. Using the same argument and notations as the proof of Lemma 2.17, when l is big enough and denote $k := \lfloor \frac{\nu(f_0^l) + l\delta_2}{m_0} \rfloor - n - 1$, we have following decomposition

$$(2.9) \quad f_0^l \circ \mu = \sum_{1 \leq j_1, j_2, \dots, j_k \leq r} (f_{j_1} \circ \mu)^{m_{j_1}} (f_{j_2} \circ \mu)^{m_{j_2}} \cdots (f_{j_k} \circ \mu)^{m_{j_k}} h_J$$

holds on $\mu^{-1}(U) \cap \tilde{X}$, where $J = (j_1, j_2, \dots, j_k)$ is a multi-index and h_J is an L^2 integrable holomorphic functions (depending on l) on $\mu^{-1}(U) \cap \tilde{X}$ for any J . As h_J is L^2 integrable on $\mu^{-1}(U) \cap \tilde{X}$, we know that $|h_J|$ is bounded near $Z_0 \cap \tilde{X}$. Then $\mu_*(h_J)$ is a weakly holomorphic function defined on $U \cap X$. It follows from the decomposition (2.9) that we have

$$f_0^l = \sum_{1 \leq j_1, j_2, \dots, j_k \leq r} (f_{j_1})^{m_{j_1}} (f_{j_2})^{m_{j_2}} \cdots (f_{j_k})^{m_{j_k}} \mu_*(h_J)$$

holds on $U \cap X$. Hence, for the germ level, we know

$$(f_0^l)_o = \sum_{1 \leq j_1, j_2, \dots, j_k \leq r} (f_{j_1}^{m_{j_1}})_o (f_{j_2}^{m_{j_2}})_o \cdots (f_{j_k}^{m_{j_k}})_o (\mu_*(h_J))_o$$

holds in $\mathcal{O}_{X,o}^w$.

Note that $\nu(f_j^{m_j}) \geq m_0$ for $1 \leq j \leq r$ and $k := \lfloor \frac{\nu(f_0^l) + l\delta_2}{m_0} \rfloor - n - 1$. We have

$$\begin{aligned} \nu(f_0^l) &\geq \min_{1 \leq j_1, j_2, \dots, j_k \leq r} \{ \nu(f_{j_1}^{m_{j_1}} f_{j_2}^{m_{j_2}} \cdots f_{j_k}^{m_{j_k}} \mu_*(h_J)) \} \\ &\geq m_0 k \\ &\geq m_0 \left(\frac{\nu(f_0^l) + l\delta_2}{m_0} - n - 2 \right) \\ &= \nu(f_0^l) + l\delta_2 - m_0(n + 2), \end{aligned}$$

which implies $l \leq \frac{m_0(n+2)}{\delta_2}$. This is a contradiction since we can choose l arbitrarily large. Then we must have $\sigma(\log |f_0|, \varphi) \leq \nu(f_0)$. \square

Now we turn to the valuations on $\mathbb{C}[z_1, \dots, z_n]/I$ a quotient ring of the polynomial ring, where I is a prime ideal in $\mathbb{C}[z_1, \dots, z_n]$.

Let $X := V(I)$ be the affine variety defined by I and $o \in X$, where o is the origin in \mathbb{C}^n . Denote the germ of the set X at x by (X, x) . We firstly recall the following basic result

Remark 2.19. $\mathbb{C}[z_1, \dots, z_n]/I$ is a subring of $\mathcal{O}_{X,o} := \mathcal{O}_{\mathbb{C}^n,o}/(I \cdot \mathcal{O}_{\mathbb{C}^n,o})$.

Proof. Note that $\mathbb{C}[z_1, \dots, z_n]$ can be naturally embedded into $\mathcal{O}_{\mathbb{C}^n,o}$. Then we have an induced morphism

$$\begin{aligned}\Phi: \mathbb{C}[z_1, \dots, z_n]/I &\longrightarrow \mathcal{O}_{\mathbb{C}^n,o}/(I \cdot \mathcal{O}_{\mathbb{C}^n,o}) \\ f + I &\longmapsto f + (I \cdot \mathcal{O}_{\mathbb{C}^n,o}).\end{aligned}$$

Now we show that Φ is injective. Assume that $\Phi(f_1 + I) = 0 + (I \cdot \mathcal{O}_{\mathbb{C}^n,o})$, where $f_1 \in \mathbb{C}[z_1, \dots, z_n]$. We would like to show that $f_1 \in I$ and hence $f_1 + I = 0$ in $\mathbb{C}[z_1, \dots, z_n]/I$. Let $\{g_j\}_{j=1}^m$ be the generators of I in $\mathbb{C}[z_1, \dots, z_n]$. Then we know that $\{g_j\}_{j=1}^m$ also generate $(I \cdot \mathcal{O}_{\mathbb{C}^n,o})$ in $\mathcal{O}_{\mathbb{C}^n,o}$. By definition and $\Phi(f_1 + I) = 0 + (I \cdot \mathcal{O}_{\mathbb{C}^n,o})$, there exists $h_j \in \mathcal{O}_{\mathbb{C}^n,o}$, where $j = 1, \dots, m$, such that

$$f_1 = \sum_{j=1}^m h_j g_j$$

in $\mathcal{O}_{\mathbb{C}^n,o}$. Hence there exists an open neighborhood U of o such that h_k is defined on U for each $k = 1, 2, \dots, m$. Recall that $X = V(I)$ and $\{g_j\}_{j=1}^m$ are generators of I . We know, for any $x \in X$, $g_k(x) = 0$ and hence $f_1(x) = 0$ for any $x \in X \cap U$. It follows from I is a prime ideal and Hilbert's Nullstellensatz theorem that X is irreducible, i.e., X cannot be written as the union of two analytic varieties $X_1, X_2 \subset \mathbb{C}^n$ with $X_1, X_2 \neq V$. Hence X_{reg} is connected (see chapter 0 of [14]). As $f_1(x) = 0$ for any $x \in X \cap U$ and X_{reg} is connected, we know that $f_1|_X \equiv 0$. Hence $f_1 \in \mathbb{I}(X)$, where $\mathbb{I}(X) := \{f \in \mathbb{C}[z_1, \dots, z_n] : f(x) = 0 \text{ for all } x \in X\}$. It follows from Hilbert's Nullstellensatz theorem, $X = V(I)$ and I is a prime ideal that $\mathbb{I}(X) = \sqrt{I} = I$. Thus $f \in I$. \square

Let $\mathcal{O}_{X,x}^w$ be the ring of germs of weakly holomorphic functions defined near x . When X is algebraic and (X, o) is irreducible as a germ of analytic set, we show that the universal denominator for $\mathcal{O}_{X,x}^w$ can be chosen be a polynomial for any $x \in V$, where V is a neighborhood of o .

Theorem 2.20 (=Theorem A.12 in the appendix). *Assume that X is algebraic and (X, o) is irreducible as a germ of analytic set. Let V be a neighborhood of o in X . There exists a polynomial $\delta \in \mathbb{C}[z_1, \dots, z_d]$, where $d = \dim X$, such that $\delta_x \mathcal{O}_{X,x}^w \subset \mathcal{O}_{X,x}$ for any $x \in V$. Such δ is called a universal denominator for weakly holomorphic functions on V .*

Proof. See the proof of Theorem A.12 in the appendix. \square

Lemma 2.21. *Let ν be a valuation on $\mathbb{C}[z_1, \dots, z_n]/I$ such that $\nu(z_i) > 0$ for any $i = 1, \dots, n$. For any elements $f_0, f_1, \dots, f_r \in \mathbb{C}[z_1, \dots, z_n]/I$ satisfying $f_j(o) = 0$ for $0 \leq j \leq r$, we have $\sigma(\log|f_0|, \varphi) \leq \nu(f_0)$, where $\varphi := \log(\sum_{1 \leq j \leq r} |f_j|^{\frac{1}{\nu(f_j)}})$ is plurisubharmonic function defined on X .*

Proof. We also prove Lemma 2.21 by contradiction. Assume that the conclusion does not hold. Then there exists $\delta_1 > 0$ such that $\log|f_0| \leq (\nu(f_0) + \delta_1)\varphi + O(1)$ near o . Then we can find some $\delta_2 > 0$ and a set of positive integers $\{m_0, \dots, m_r\}$ such that

$$\log|f_0| \leq \frac{\nu(f_0) + \delta_2}{m_0} \tilde{\varphi} + O(1) \quad \text{near } o,$$

where $\tilde{\varphi} := \log(\sum_{1 \leq j \leq r} |f_j^{m_j}|)$ and $\nu(f_j^{m_j}) \geq m_0$ for $1 \leq j \leq r$. For any positive integer l , we have

$$\log |f_0^l| \leq \frac{\nu(f_0^l) + l\delta_2}{m_0} \tilde{\varphi} + O(1)$$

holds on $(U \cap X) \setminus X_{\text{sing}}$, where $U := \{x \in \mathbb{C}^n : \log|z| < \delta\}$. Using the same argument and notations as the proof of Lemma 2.17, when l is big enough and denote $k := \lfloor \frac{\nu(f_0^l) + l\delta_2}{m_0} \rfloor - n - 1$, we have following decomposition

$$(2.10) \quad f_0^l \circ \mu = \sum_{1 \leq j_1, j_2, \dots, j_k \leq r} (f_{j_1} \circ \mu)^{m_{j_1}} (f_{j_2} \circ \mu)^{m_{j_2}} \cdots (f_{j_k} \circ \mu)^{m_{j_k}} h_J$$

hold on some neighborhood U of $p \in \mu^{-1}(o) \Subset \tilde{X}$, where $J = (j_1, j_2, \dots, j_k)$ is a multi-index and h_J are L^2 integrable holomorphic functions (depending on l) on U for any J . As h_J are L^2 integrable on $\mu^{-1}(U) \cap \tilde{X}$, we know that $|h_J|$ is bounded near $Z_0 \cap \tilde{X}$. Then $\mu_*(h_J)$ is a weakly holomorphic function defined on $U \cap X$. It follows from decomposition (2.10) that we have

$$(2.11) \quad f_0^l = \sum_{1 \leq j_1, j_2, \dots, j_k \leq r} (f_{j_1})^{m_{j_1}} (f_{j_2})^{m_{j_2}} \cdots (f_{j_k})^{m_{j_k}} \mu_*(h_J)$$

holds on $U \cap X$, where $\mu_*(h_J)$ is a weakly holomorphic function defined on $U \cap X$.

It follows from Theorem 2.15 and Theorem 2.20 that, for any J , there exists a holomorphic function g_J on $U \cap X$ such that $\mu_*(h_J) = \frac{g_J}{\delta}$, where δ is a polynomial. Thus, we have

$$(2.12) \quad \delta f_0^l = \sum_{1 \leq j_1, j_2, \dots, j_k \leq r} (f_{j_1})^{m_{j_1}} (f_{j_2})^{m_{j_2}} \cdots (f_{j_k})^{m_{j_k}} g_J$$

holds on $U \cap X$.

Let $\tilde{m} = (z_1, \dots, z_n)$ be the ideal generated by z_1, \dots, z_n in $\mathbb{C}[z_1, \dots, z_n]/I$ and $\hat{m} = (z_1, \dots, z_n)$ be the ideal generated by z_1, \dots, z_n in $\mathcal{O}_{X,o} = \mathcal{O}_{\mathbb{C}^n,o}/I$. For any integer $N > 0$, let $g_{J,N}$ be the finite terms of Taylor expansion of g_J such that $[g_{J,N}] = [g_J]$ in $\mathcal{O}_{X,o}/\hat{m}^N$. Note that $g_{J,N}$ can be viewed as an element in $\mathbb{C}[z_1, \dots, z_n]/I$. Denote

$$R_N := \delta f_0^l - \sum_{1 \leq j_1, j_2, \dots, j_k \leq r} (f_{j_1})^{m_{j_1}} (f_{j_2})^{m_{j_2}} \cdots (f_{j_k})^{m_{j_k}} g_{J,N}.$$

Then R_N is a polynomial in $\mathbb{C}[z_1, \dots, z_n]/I$. It follows from equality (2.12) and $[g_{J,N}] = [g_J]$ in $\mathcal{O}_{X,o}/\hat{m}^N$ that $R_N \in \hat{m}^N$. As R_N is a polynomial, we know R_N actually belong to \tilde{m}^N . And we also know that

$$\delta f_0^l = \sum_{1 \leq j_1, j_2, \dots, j_k \leq r} (f_{j_1})^{m_{j_1}} (f_{j_2})^{m_{j_2}} \cdots (f_{j_k})^{m_{j_k}} g_{J,N} + R_N$$

holds in $\mathbb{C}[z_1, \dots, z_n]/I$. Note that $\nu(f_j^{m_j}) \geq m_0$ for $1 \leq j \leq r$, $v(\tilde{m}) > 0$ and $k := \lfloor \frac{\nu(f_0^l) + l\delta_2}{m_0} \rfloor - n - 1$. When N is big enough, we have

$$\begin{aligned} \nu(\delta) + \nu(f_0^l) &\geq \left\{ \min_{1 \leq j_1, j_2, \dots, j_k \leq r} \nu(f_{j_1}^{m_{j_1}} f_{j_2}^{m_{j_2}} \cdots f_{j_k}^{m_{j_k}} g_{J,N}), R_N \right\} \\ &\geq \left\{ \min_{1 \leq j_1, j_2, \dots, j_k \leq r} \nu(f_{j_1}^{m_{j_1}} f_{j_2}^{m_{j_2}} \cdots f_{j_k}^{m_{j_k}}), R_N \right\} \\ &\geq \min_{1 \leq j_1, j_2, \dots, j_k \leq r} \nu(f_{j_1}^{m_{j_1}} f_{j_2}^{m_{j_2}} \cdots f_{j_k}^{m_{j_k}}) \\ &\geq m_0 k \\ &\geq m_0 \left(\frac{\nu(f_0^l) + l\delta_2}{m_0} - n - 2 \right) \\ &= \nu(f_0^l) + l\delta_2 - m_0(n + 2), \end{aligned}$$

which implies $l \leq \frac{m_0(n+2)+\nu(\delta)}{\delta_2}$. This is a contradiction since we can choose l arbitrarily large. Then we must have $\sigma(\log |f_0|, \varphi) \leq \nu(f_0)$. \square

To prove Corollary 1.13, we also recall the following Hilbert's Nullstellensatz theorem for finitely generated \mathbb{C} -algebras. For example, the following Lemma can be referred to Theorem 5.5 in [26].

Lemma 2.22 (see [26]). *Let A be a finitely generated \mathbb{C} -algebra and $J \subset A$ be a proper ideal. Then one has $\sqrt{J} = \bigcap_{J \subset m} m$, where the intersection is over all maximal ideals m containing J .*

Let $A = \mathbb{C}[z_1, \dots, z_n]/I$, where I is an ideal in $\mathbb{C}[z_1, \dots, z_n]$. Denote $X = V(I)$ be the zero set of I . Let $J \subset A$ be a proper ideal. Define $V_X(J) := \{x \in X \mid P(x) = 0, \text{ for any } P \in J\}$. For any subset $Y \subset X$, denote $I(Y) := \{P \in A \mid P(x) = 0 \text{ for any } x \in Y\}$.

Using Lemma 2.22, we immediately have

Lemma 2.23. *Let $A = \mathbb{C}[z_1, \dots, z_n]/I$, where I is an ideal in $\mathbb{C}[z_1, \dots, z_n]$ and $J \subset A$ be a proper ideal. Then one has $I(V_X(J)) = \sqrt{J}$.*

For the convenience of the readers, we recall the proof of Lemma 2.23.

Proof of Lemma 2.23. It is obvious that $\sqrt{J} \subset I(V_X(J))$. If m is an maximal ideal in A , then $m = \sqrt{m} \subset I(V(m)) = m$, where the last “=” holds since m is maximal.

Note that $J \subset \sqrt{J} = \bigcap_{J \subset m} m$. For any maximal ideal m containing J , we have $V(J) \supset V(m)$. Hence $I(V_X(J)) \subset I(V(m)) = m$. By the arbitrariness of m and Lemma 2.22, we have $I(V_X(J)) = \bigcap_{J \subset m} m = \sqrt{J}$. This completes the proof. \square

Denote the set of all germs of real analytic functions near the origin $o' \in \mathbb{R}^n$ by $C_{o'}^{\text{an}}$. There exists an injective ring homomorphism $P: C_{o'}^{\text{an}} \rightarrow \mathcal{O}_{o'}$, which satisfies

$$P \left(\sum_{\alpha \in \mathbb{Z}_{\geq 0}^n} a_{\alpha} x^{\alpha} \right) = \sum_{\alpha \in \mathbb{Z}_{\geq 0}^n} a_{\alpha} z^{\alpha},$$

where (x_1, \dots, x_n) and (z_1, \dots, z_n) are the standard coordinates in \mathbb{R}^n and \mathbb{C}^n respectively, and $\sum_{\alpha \in \mathbb{Z}_{\geq 0}^n} a_{\alpha} x^{\alpha}$ is the power series expansion of arbitrary real analytic function near o' . Assume that I is an ideal in $C_{o'}^{\text{an}}$. Denote $(P(I))$ be the ideal generated by $P(I)$ in $\mathcal{O}_{o'}$. Note that P is obvious injective. Hence we have

Remark 2.24. *If $(P(I))$ is prime ideal in \mathcal{O}_o , then I is prime in $C_{o'}^{\text{an}}$.*

Proof. Let $ab \in I$. Then $P(ab) = P(a)P(b) \in (P(I))$. As $(P(I))$ is prime ideal, we may assume that $P(a) \in (P(I))$. As P is injective, we know that $a \in I$. \square

Denote the elements in $C_{o'}^{\text{an}}/I$ and $\mathcal{O}_o/(P(I))$ by $[f]$ and \tilde{g} for any $f \in C_{o'}^{\text{an}}$ and $g \in \mathcal{O}_o$ respectively. Then we have an induced ring homomorphism $\tilde{P} : C_{o'}^{\text{an}}/I \rightarrow \mathcal{O}_o/(P(I))$ defined by $\tilde{P}([f]) = \widetilde{P(f)}$. It is easy to check that \tilde{P} is well defined. Denote X be the zero variety defined by $(P(I))$.

We have following properties of the homomorphism \tilde{P} .

Remark 2.25. *\tilde{P} is injective.*

Proof. Assume $[F_1], [F_2] \in C_{o'}^{\text{an}}/I$ such that $\tilde{P}([F_1]) = \tilde{P}([F_2])$ in $\mathcal{O}_o/(P(I))$. Then there exist $g_k \in \mathcal{O}_o$ ($k = 1, \dots, m$) such that $P(F_1) - P(F_2) = \sum_{k=1}^m g_k P(f_k)$. For any $g \in \mathcal{O}_o$, g can be written as a convergent power series $g = \sum_{\alpha} a_{\alpha} z_1^{\alpha_1} z_2^{\alpha_2} \cdots z_n^{\alpha_n}$, where α is a multi-index, $a_{\alpha} \in \mathbb{C}$ and (z_1, \dots, z_n) is the coordinate near o . We define

$$g_{1,\text{Re}} := \sum_{\alpha} \text{Re}(a_{\alpha}) z_1^{\alpha_1} z_2^{\alpha_2} \cdots z_n^{\alpha_n}$$

and

$$g_{1,\text{Im}} := \sum_{\alpha} \text{Im}(a_{\alpha}) z_1^{\alpha_1} z_2^{\alpha_2} \cdots z_n^{\alpha_n}.$$

Hence, We have

$$P(F_1) - P(F_2) = \sum_{k=1}^m (P(g_{k,\text{Re}}) + iP(g_{k,\text{Im}})) P(f_k),$$

which implies that

$$P(F_1) - P(F_2) = \sum_{k=1}^m P(g_{k,\text{Re}}) P(f_k).$$

As P is injective, one has $F_1 - F_2 \in I$ and hence $[F_1] = [F_2]$ in $C_{o'}^{\text{an}}/I$. This means \tilde{P} is injective. \square

Remark 2.26. *For any $\tilde{g} \in \mathcal{O}_o/(P(I))$, there exists a unique pair of $[g_{\text{Re}}], [g_{\text{Im}}] \in C_{o'}^{\text{an}}/I$ such that*

$$\tilde{g} = \tilde{P}([g_{\text{Re}}]) + i\tilde{P}([g_{\text{Im}}])$$

Proof. Let g_1, g_2 belong to \mathcal{O}_o such that $g_1 - g_2 \in (P(I))$ and $\tilde{g} = \tilde{g}_1 = \tilde{g}_2$ in $\mathcal{O}_o/(P(I))$. Let $\{f_k\}_{k=1}^m$ be the generators of I in $C_{o'}^{\text{an}}$. Then $(P(I))$ was generated by $P(f_k)$. We know that there exists $h_k \in \mathcal{O}_o$ ($k = 1, \dots, m$) such that $g_1 - g_2 = \sum_{k=1}^m h_k P(f_k)$. Then we have

$$(2.13) \quad g_{1,\text{Re}} + ig_{1,\text{Im}} - g_{2,\text{Re}} - ig_{2,\text{Im}} = \sum_{k=1}^m (h_{k,\text{Re}} + ih_{k,\text{Im}}) P(f_k).$$

Note that $P(f_k)$ has real coefficients as a power series of z_1, \dots, z_n . Equality (2.13) tells that

$$(2.14) \quad \begin{aligned} g_{1,\text{Re}} - g_{2,\text{Re}} &= \sum_{k=1}^m h_{k,\text{Re}} P(f_k) \\ g_{1,\text{Im}} - g_{2,\text{Im}} &= \sum_{k=1}^m h_{k,\text{Im}} P(f_k) \end{aligned}$$

Thus we have $g_{1,\text{Re}} - g_{2,\text{Re}} \in (P(I))$ and $g_{1,\text{Im}} - g_{2,\text{Im}} \in (P(I))$ hold. It follows from P is injective and equalities (2.14) that $[P^{-1}(g_{1,\text{Re}})] = [P^{-1}(g_{2,\text{Re}})] \in C_{o'}^{\text{an}}/I$ and $[P^{-1}(g_{1,\text{Im}})] = [P^{-1}(g_{2,\text{Im}})] \in C_{o'}^{\text{an}}/I$. Denote $[g_{\text{Re}}] = [P^{-1}(g_{1,\text{Re}})]$ and $[g_{\text{Im}}] = [P^{-1}(g_{1,\text{Im}})]$. Hence we know there exist pair $[g_{\text{Re}}], [g_{\text{Im}}] \in C_{o'}^{\text{an}}/I$ such that $\tilde{g} = \tilde{P}(g_{\text{Re}}) + i\tilde{P}(g_{\text{Im}})$.

The uniqueness of the pair $[g_{\text{Re}}], [g_{\text{Im}}]$ follows from the fact \tilde{P} is injective, see Remark 2.25. \square

Denote X be the zero variety defined by $(P(I))$. Assume that $(P(I))$ is a prime ideal in \mathcal{O}_o .

Lemma 2.27. *Let ν be a valuation on $C_{o'}^{\text{an}}/I$. For any $[f_0], [f_1], \dots, [f_r]$ in $C_{o'}^{\text{an}}/I$ satisfying $\nu([f_j]) > 0$ for $1 \leq j \leq r$, we have $\sigma(\log |\tilde{f}_0|, \varphi) \leq \nu([f_0])$ holds near $o \in X$, where $\varphi := \log \left(\sum_{1 \leq j \leq r} |\tilde{f}_j|^{\frac{1}{\nu([f_j])}} \right)$ and $\tilde{f}_j := \tilde{P}([f_j])$ for $0 \leq j \leq r$.*

Proof. We prove Lemma 2.27 by contradiction. Suppose to the contrary that the conclusion does not hold. There exists $\delta_1 > 0$ such that $\log |\tilde{f}_0| \leq (\nu([f_0]) + \delta_1)\varphi + O(1)$ near o . Then we can find some $\delta_2 > 0$ and a set of positive integers $\{m_0, \dots, m_r\}$ such that

$$\log |\tilde{f}_0| \leq \frac{\nu([f_0]) + \delta_2}{m_0} \tilde{\varphi} + O(1) \quad \text{near } o,$$

where $\tilde{\varphi} := \log \left(\sum_{1 \leq j \leq r} |\tilde{f}_j|^{m_j} \right)$ and $\nu([f_j]^{m_j}) \geq m_0$ for $1 \leq j \leq r$. For any positive integer l , we have

$$\log |\tilde{f}_0^l| \leq \frac{\nu(\tilde{f}_0^l) + l\delta_2}{m_0} \tilde{\varphi} + O(1)$$

holds on $(U \cap X) \setminus X_{\text{sing}}$, where $U := \{x \in \mathbb{C}^n : \log |z| < \delta\}$.

Following the argument and notation of Lemma 2.17, we see that for sufficiently large l , by defining $k := \lfloor \frac{\nu(f_0^l) + l\delta_2}{m_0} \rfloor - n - 1$, we obtain that the following decomposition

$$(2.15) \quad \tilde{f}_0^l = \sum_{1 \leq j_1, j_2, \dots, j_k \leq r} (\tilde{f}_{j_1})^{m_{j_1}} (\tilde{f}_{j_2})^{m_{j_2}} \cdots (\tilde{f}_{j_k})^{m_{j_k}} \frac{g_J}{\delta}$$

holds on U , where U is an open neighborhood of o in X , and g_J (depending on l) and δ belong to $\mathcal{O}_{X,o} = \mathcal{O}_o/(P(I))$. Thus we have

$$(2.16) \quad \delta \tilde{f}_0^l = \sum_{1 \leq j_1, j_2, \dots, j_k \leq r} (\tilde{f}_{j_1})^{m_{j_1}} (\tilde{f}_{j_2})^{m_{j_2}} \cdots (\tilde{f}_{j_k})^{m_{j_k}} g_J.$$

Combining with Remark 2.26, we have

$$\begin{aligned} & (\tilde{P}([\delta_{\text{Re}}]) + i\tilde{P}([\delta_{\text{Im}}]))\tilde{P}([f_0]^l) \\ &= \sum_{1 \leq j_1, j_2, \dots, j_k \leq r} \tilde{P}([f_{j_1}]^{m_{j_1}})\tilde{P}([f_{j_2}]^{m_{j_2}}) \cdots \tilde{P}([f_{j_k}]^{m_{j_k}})(\tilde{P}([g_{J, \text{Re}}]) + i\tilde{P}([g_{J, \text{Im}}])). \end{aligned}$$

Hence we have

$$[\delta_{\text{Re}}][f_0]^l = \sum_{1 \leq j_1, j_2, \dots, j_k \leq r} [f_{j_1}]^{m_{j_1}} [f_{j_2}]^{m_{j_2}} \cdots [f_{j_k}]^{m_{j_k}} [g_{J, \text{Re}}]$$

holds in $C_{o'}^{\text{an}}/I$. As $\nu([f_j]^{m_j}) \geq m_0$ for $1 \leq j \leq r$, we get

$$\begin{aligned} \nu([f_0]^l) &\geq m_0 \left(\left\lfloor \frac{\nu(f_0^l) + l\delta_2}{m_0} \right\rfloor - n - 1 \right) - \nu([\delta_{\text{Re}}]) \\ &\geq m_0 \left(\frac{\nu(f_0^l) + l\delta_2}{m_0} - n - 2 \right) - \nu([\delta_{\text{Re}}]) \\ &= \nu(f_0^l) + l\delta_2 - m_0(n + 2) - \nu([\delta_{\text{Re}}]), \end{aligned}$$

which implies $l \leq \frac{m_0(n+2) + \nu([\delta_{\text{Re}}])}{\delta_2}$, contradiction. Then we have $\sigma(\log |\tilde{f}_0|, \varphi) \leq \nu([f_0])$. \square

Let I be an ideal in $\mathbb{R}[x_1, \dots, x_n]$. Denote the restriction of P on $\mathbb{R}[x_1, \dots, x_n]$ also by P and \tilde{P} on $\mathbb{R}[x_1, \dots, x_n]/I$ also by \tilde{P} . Let $(P(I))$ be the ideal in $\mathbb{C}[x_1, \dots, x_n]$ generated by $P(I)$.

Note that P is obvious injective. Similarly as Remark 2.24, we have

Remark 2.28. *If $(P(I))$ is prime ideal in $\mathbb{C}[x_1, \dots, x_n]$, then I is prime in $\mathbb{R}[x_1, \dots, x_n]$.*

Proof. Since the proof of Remark 2.28 is the same as the proof of Remark 2.24, we omit the proof. \square

We also have following remark, which is an analogy of Remark 2.25 and Remark 2.26 in polynomial case.

Remark 2.29. *The map $\tilde{P} : \mathbb{R}[x_1, \dots, x_n]/I \rightarrow \mathbb{C}[x_1, \dots, x_n]/(P(I))$ is injective. For any $\tilde{g} \in \mathbb{C}[x_1, \dots, x_n]/(P(I))$, there exists a unique pair of $[g_{\text{Re}}], [g_{\text{Im}}] \in \mathbb{R}[x_1, \dots, x_n]/I$ such that*

$$\tilde{g} = \tilde{P}([g_{\text{Re}}]) + i\tilde{P}([g_{\text{Im}}])$$

Proof. The proof of the injectiveness is almost the same as the proof of Remark 2.25. The proof of existence and uniqueness of the decomposition is almost the same as the proof of Remark 2.26. So we omit the proof of Remark 2.29. \square

Assume that $(P(I))$ is a prime ideal in $\mathbb{C}[x_1, \dots, x_n]$ and (X, o) is irreducible as a germ of analytic set, where $o \in X$ and X is the affine variety defined by $(P(I))$.

Lemma 2.30. *Let ν be a valuation on $\mathbb{R}[x_1, \dots, x_n]/I$ satisfying $\nu([x_j]) > 0$ for any $1 \leq j \leq n$. For any $[f_0], [f_1], \dots, [f_r]$ in $\mathbb{R}[x_1, \dots, x_n]_{o'}/I$ satisfying $\nu([f_j]) > 0$ for $1 \leq j \leq r$, we have $\sigma(\log |\tilde{f}_0|, \varphi) \leq \nu([f_0])$ holds near $o \in X$, where $\varphi := \log \left(\sum_{1 \leq j \leq r} |\tilde{f}_j|^{\frac{1}{\nu([f_j])}} \right)$ and $\tilde{f}_j := \tilde{P}([f_j])$ for $0 \leq j \leq r$.*

Proof. We prove this lemma by contradiction. Suppose to the contrary that the conclusion does not hold. There exists $\delta_1 > 0$ such that $\log |\tilde{f}_0| \leq (\nu([f_0]) + \delta_1)\varphi + O(1)$ near o . Then we can find some $\delta_2 > 0$ and a set of positive integers $\{m_0, \dots, m_r\}$ such that

$$\log |\tilde{f}_0| \leq \frac{\nu([f_0]) + \delta_2}{m_0} \tilde{\varphi} + O(1) \quad \text{near } o,$$

where $\tilde{\varphi} := \log \left(\sum_{1 \leq j \leq r} |\tilde{f}_j^{m_j}| \right)$ and $\nu([f_j]^{m_j}) \geq m_0$ for $1 \leq j \leq r$. For any positive integer l , we have

$$\log |\tilde{f}_0^l| \leq \frac{\nu(\tilde{f}_0^l) + l\delta_2}{m_0} \tilde{\varphi} + O(1)$$

holds on $(U \cap X) \setminus X_{\text{sing}}$, where $U := \{x \in \mathbb{C}^n : \log |z| < \delta\}$.

Following the argument and notation of Lemma 2.17, we see that for sufficiently large l , by defining $k := \lfloor \frac{\nu(f_0^l) + l\delta_2}{m_0} \rfloor - n - 1$, we obtain that the following decomposition

$$(2.17) \quad \delta \tilde{f}_0^l = \sum_{1 \leq j_1, j_2, \dots, j_k \leq r} (\tilde{f}_{j_1})^{m_{j_1}} (\tilde{f}_{j_2})^{m_{j_2}} \dots (\tilde{f}_{j_k})^{m_{j_k}} g_J.$$

holds on U , where U is an open neighborhood of o in X , δ is a polynomial with complex coefficients and g_J (depending on l) belongs to $\mathcal{O}_{X,o} = \mathcal{O}_o / (P(I))$.

Combining equality (2.17) and Remark 2.29, we have

$$\begin{aligned} & (\tilde{P}([\delta_{\text{Re}}]) + i\tilde{P}([\delta_{\text{Im}}])) \tilde{P}([f_0]^l) \\ &= \sum_{1 \leq j_1, j_2, \dots, j_k \leq r} \tilde{P}([f_{j_1}]^{m_{j_1}}) \tilde{P}([f_{j_2}]^{m_{j_2}}) \dots \tilde{P}([f_{j_k}]^{m_{j_k}}) (\tilde{P}([g_{J,\text{Re}}]) + i\tilde{P}([g_{J,\text{Im}}])) \end{aligned}$$

Hence we have

$$(2.18) \quad [\delta_{\text{Re}}][f_0]^l = \sum_{1 \leq j_1, j_2, \dots, j_k \leq r} [f_{j_1}]^{m_{j_1}} [f_{j_2}]^{m_{j_2}} \dots [f_{j_k}]^{m_{j_k}} [g_{J,\text{Re}}]$$

holds in $C_{o'}^{\text{an}} / I$.

Let $m_P = (x_1, \dots, x_n)$ be the ideal generated by x_1, \dots, x_n in $\mathbb{R}[x_1, \dots, x_n] / I$ and $\hat{m} = (x_1, \dots, x_n)$ be the ideal generated by x_1, \dots, x_n in $C_{o'}^{\text{an}} / I$. For any integer $N > 0$, let $g_{J,N}$ be the finite terms of the Taylor expansion of $g_{J,\text{Re}}$ such that $[g_{J,N}] = [g_{J,\text{Re}}]$ in $C_{o'}^{\text{an}} / \hat{m}^N$. Note that $g_{J,N}$ can be viewed as an element in $\mathbb{R}[x_1, \dots, x_n] / I$. Denote

$$R_N := [\delta_{\text{Re}}][f_0]^l - \sum_{1 \leq j_1, j_2, \dots, j_k \leq r} [f_{j_1}]^{m_{j_1}} [f_{j_2}]^{m_{j_2}} \dots [f_{j_k}]^{m_{j_k}} [g_{J,N}].$$

Then R_N is a polynomial in $\mathbb{R}[x_1, \dots, x_n] / I$. It follows from equality (2.18) and $[g_{J,N}] = [g_J]$ in $C_{o'}^{\text{an}} / \hat{m}^N$ that $R_N \in \hat{m}^N$. As R_N is a polynomial, we know R_N actually belongs to $(m_P)^N$. And we also know that

$$[\delta_{\text{Re}}][f_0]^l = \sum_{1 \leq j_1, j_2, \dots, j_k \leq r} [f_{j_1}]^{m_{j_1}} [f_{j_2}]^{m_{j_2}} \dots [f_{j_k}]^{m_{j_k}} [g_{J,N}] + R_N$$

holds in $\mathbb{R}[x_1, \dots, x_n]/I$. Note that $\nu([f_j]^{m_j}) \geq m_0$ for $1 \leq j \leq r$, $\nu(m_P) > 0$ and $k := \lfloor \frac{\nu(f_0^l) + l\delta_2}{m_0} \rfloor - n - 1$. When N is large enough, we have

$$\begin{aligned} \nu([\delta_{\text{Re}}]) + \nu([f_0]^l) &\geq \left\{ \min_{1 \leq j_1, j_2, \dots, j_k \leq r} \nu([f_{j_1}]^{m_{j_1}} [f_{j_2}]^{m_{j_2}} \cdots [f_{j_k}]^{m_{j_k}} [g_{J,N}]), R_N \right\} \\ &\geq \left\{ \min_{1 \leq j_1, j_2, \dots, j_k \leq r} \nu([f_{j_1}]^{m_{j_1}} [f_{j_2}]^{m_{j_2}} \cdots [f_{j_k}]^{m_{j_k}}), R_N \right\} \\ &\geq \min_{1 \leq j_1, j_2, \dots, j_k \leq r} \nu([f_{j_1}]^{m_{j_1}} [f_{j_2}]^{m_{j_2}} \cdots [f_{j_k}]^{m_{j_k}}) \\ &\geq m_0 k \\ &\geq m_0 \left(\frac{\nu(f_0^l) + l\delta_2}{m_0} - n - 2 \right) \\ &= \nu(f_0^l) + l\delta_2 - m_0(n+2), \end{aligned}$$

which implies $l \leq \frac{m_0(n+2) + \nu([\delta_{\text{Re}}])}{\delta_2}$. This is a contradiction since we can choose l arbitrarily large. Then we must have $\sigma(\log |f_0|, \varphi) \leq \nu(f_0)$. \square

We also note that we have the following basic result.

Remark 2.31. $\mathbb{R}[z_1, \dots, z_n]/I$ is a subring of $C_{o'}^{an}/(I \cdot C_{o'}^{an})$.

Proof. The proof of Remark 2.31 is almost the same as the proof of Remark 2.19. So we omit the proof. \square

2.5. Jumping number in the singular case. In this section, we discuss the properties of the jumping number $c_o^f(\varphi)$, where $o \in X$ can be a singular point of an irreducible analytic set X .

Definition 2.32. Let $X \subset \mathbb{C}^n$ be an analytic set and $K \subset X$ a compact subset. For any plurisubharmonic function φ defined on K and any weakly holomorphic function f defined on K , denote the jumping number $c_K^f(\varphi)$ by

$$c_K^f(\varphi) := \sup \left\{ c > 0 : \int_{U \cap X_{\text{reg}}} |f|^2 e^{-2c\varphi} dV_X < +\infty, \right. \\ \left. \text{for some open neighborhood } U \text{ of } K \text{ in } X \right\}.$$

When $K = \{o\}$ is a singular point of X , $c_o^f(\varphi)$ generalizes the ordinary jumping number to the singular case. We resolve the singularities of (X, o) and consider the jumping number $c_o^f(\varphi)$ after a sequence of blow-ups.

Remark 2.33. Using Theorem 2.4, we can resolve the singularities of X , and denote the corresponding proper modification by $\mu: \tilde{\Omega} \rightarrow \Omega$, where $\Omega \subset \mathbb{C}^n$ open such that $o \in X \cap \Omega$. Denote the strict transform of X by \tilde{X} and denote $Z_0 := \tilde{X} \cap \mu^{-1}(\{o\})$. Note that $f \circ \mu$ and $\varphi \circ \mu$ are well defined holomorphic function and plurisubharmonic function correspondingly on $\pi^{-1}(U) \cap \tilde{X}$ for any open neighborhood U of o in X respectively. Since μ is proper, Z_0 is compact in \tilde{X} . Note that μ is biholomorphic outside an analytic subset, we know that $c_o^f(\varphi) = c_{Z_0}^{\det \mu \cdot f \circ \mu}(\varphi \circ \mu)$, where $\det \mu$ is the determinant of the Jacobian of the modification μ (which is a holomorphic function on $\tilde{\Omega}$) and $c_{Z_0}^{\det \mu \cdot f \circ \mu}(\varphi \circ \mu)$ is the usual jumping number on a compact set in the regular case.

Recall that X is an analytic subset of $\Omega \in \mathbb{C}^n$ with pure dimension d and denote $|z|^2 := \sum_{1 \leq j \leq n} |z_j|^2$. Assume that (X, o) is singular, we have the following approximation result for jumping numbers.

Lemma 2.34. *Let φ be a plurisubharmonic function near o , where $c > 0$ is a constant and $f_j \in \mathcal{O}_{X,o}^w$ for $1 \leq j \leq m$. Then*

$$\lim_{N \rightarrow +\infty} c_o^f(\max\{\varphi, N \log |z|\}) = c_o^f(\varphi)$$

for any holomorphic function $f \in \mathcal{O}_{X,o}^*$.

Proof. By definition, we have $c_o^f(\max\{\varphi, N \log |z|\}) \geq c_o^f(\varphi)$.

Using Theorem 2.4, we can resolve the singularity of X and get a proper holomorphic map $\mu: \tilde{\Omega} \rightarrow \Omega$ where $\Omega \subset \mathbb{C}^n$ is an open subset such that $o \in X \cap \Omega$. Remark 2.33 tells that $c_o^f(\varphi) = c_{Z_0}^{\det \mu \cdot f \circ \mu}(\varphi \circ \mu)$ and $c_o^f(\max\{\varphi, N \log |z|\}) = c_{Z_0}^{\det \mu \cdot f \circ \mu}(\max\{\varphi \circ \mu, N \log |z \circ \mu|\})$. Note that $(N \log |z \circ \mu|)|_{\tilde{X}}$ is a plurisubharmonic function with $\{(N \log |z \circ \mu|)|_{\tilde{X}} = -\infty\} = Z_0$.

Assume that $c_{Z_0}^{\det \mu \cdot f \circ \mu}(\varphi \circ \mu) < 1$. Note that $c_{Z_0}^{\det \mu \cdot f \circ \mu}(\varphi \circ \mu) = \inf_{z \in Z_0} c_z^{\det \mu \cdot f \circ \mu}(\varphi \circ \mu)$. We know there exists a point $p \in Z_0$, such that $c_p^{\det \mu \cdot f \circ \mu}(\varphi \circ \mu) < 1$. As (\tilde{X}, p) is regular, we can find a local coordinate neighborhood (W_p, w) centered at p such that there exists an $A > 0$ such that $(\log |z \circ \mu|)|_{\tilde{X}} \leq A \log |w| + O(1)$ on W_p . Then $c_p^{\det \mu \cdot f \circ \mu}(\max\{\varphi \circ \mu, N \log |z \circ \mu|\}) \leq c_p^{\det \mu \cdot f \circ \mu}(\max\{\varphi \circ \mu, NA \log |w|\})$.

The proof in the smooth case (see [2]) shows that, $c_p^{\det \mu \cdot f \circ \mu}(\varphi \circ \mu) < 1$ implies there exists $N \gg 0$ such that

$$c_p^{\det \mu \cdot f \circ \mu}(\max\{\varphi \circ \mu, NA \log |w|\}) \leq 1.$$

Hence $c_p^{\det \mu \cdot f \circ \mu}(\max\{\varphi \circ \mu, N \log |z \circ \mu|\}) \leq 1$, which implies

$$c_{Z_0}^{\det \mu \cdot f \circ \mu}(\max\{\varphi \circ \mu, N \log |z \circ \mu|\}) \leq 1.$$

Thus we know $\lim_{N \rightarrow +\infty} c_{Z_0}^{\det \mu \cdot f \circ \mu}(\max\{\varphi \circ \mu, N \log |z \circ \mu|\}) \leq c_{Z_0}^{\det \mu \cdot f \circ \mu}(\varphi \circ \mu)$, which implies $\lim_{N \rightarrow +\infty} c_o^f(\max\{\varphi, N \log |z|\}) \leq c_o^f(\varphi)$. \square

We also need the following definition of relative types on compact subsets of complex subvarieties.

Definition 2.35. *For plurisubharmonic functions ψ, φ defined on a neighborhood of compact subset $K \subset X$, define the relative type on K by*

$$\begin{aligned} \sigma_K(\psi, \varphi) := \sup\{c \geq 0: \psi \leq c\varphi + O(1) \text{ holds on } U \setminus X_{\text{sing}} \\ \text{for some open neighborhood } U \text{ of } K \text{ in } X\}. \end{aligned}$$

We may omit K if there is no confusion.

Remark 2.36. *When $K = \{z_0\}$ is a single point, we have the relative type $\sigma_{z_0}(\psi, \varphi)$ at a point z_0 . Moreover, we have $\sigma_K(\psi, \varphi) = \inf_{p \in K} \sigma_p(\psi, \varphi)$.*

Proof. By definition, it is easy to verify that $\sigma_K(\psi, \varphi) \leq \inf_{p \in K} \sigma_p(\psi, \varphi)$.

Denote $c_1 = \inf_{p \in K} \sigma_p(\psi, \varphi)$. Let $\delta > 0$ be any small real number. Then for any $p \in K$, there exists an open neighborhood U_p of p in X such that $\psi \leq (c_1 - \delta)\varphi + O(1)$ on $U_p \setminus X_{\text{sing}}$. By the compactness of K , we know that there exist finite points $\{p_i\}_{i=1}^k$ such that $\cup_{i=1}^k U_{p_i}$ covers K . Thus $\psi \leq (c_1 - \delta)\varphi + O(1)$ on $\cup_{i=1}^k U_{p_i} \setminus X_{\text{sing}}$. By the arbitrariness of δ , we have $\sigma_K(\psi, \varphi) \geq \inf_{p \in K} \sigma_p(\psi, \varphi)$.

Thus, we know $\sigma_K(\psi, \varphi) = \inf_{p \in K} \sigma_p(\psi, \varphi)$. \square

Let f_j, h_k be holomorphic functions defined on a open set $U \subset X$ for any $1 \leq j \leq m$ and any $1 \leq k \leq q$. Let $\varphi_N := c \log \left(\sum_{1 \leq j \leq m} |f_j|^{\frac{1}{a_j}} + \sum_{1 \leq k \leq q} |h_k|^N \right)$, where $c > 0$, N is a positive integer and $\{a_j\}_{1 \leq j \leq m}$ are positive numbers. Let $K \subset U \subset X$ be a compact subset of X . We have the following result for relative types $\sigma(\cdot, \varphi_N)$.

Lemma 2.37. *Let f and g be holomorphic functions defined on K . Let $z_0 \in K$ be any point such that $z_0 \in \{\varphi_N = -\infty\}$. Then there exists a constant $C > 0$ (relies on g , $\{f_j\}$ and $\{h_k\}$ and is independent of N , f and z_0) such that*

$$\sigma_{z_0}(\log |fg|, \varphi_N) \leq \sigma_{z_0}(\log |f|, \varphi_N) + C.$$

Proof. Using Theorem 2.4 and blow-ups, we can resolve the singularities of $(\cup_{i=1}^m \{f_j = 0\}) \cup (\cup_{k=1}^q \{h_k = 0\})$, and denote the corresponding proper modification by $\mu: \tilde{X} \rightarrow X$. Denote $Z_0 := \mu^{-1}(z_0)$. Note that $\sigma_{z_0}(\log |f|, \varphi_N) = \sigma_{Z_0}(\log |f \circ \mu|, \varphi_N \circ \mu)$ for any $f \in \mathcal{O}_{X, o}$. Remark 2.36 tells that $\sigma_{Z_0}(\log |f \circ \mu|, \varphi_N \circ \mu) = \inf_{p \in Z_0} \sigma_p(\log |f \circ \mu|, \varphi_N \circ \mu)$.

Denote $\{f_j \circ \mu = 0\} \cap \mu^{-1}(K) = \sum_{\alpha_j} a_{j, \alpha_j} D_{j, \alpha_j}$ and $\{h_k \circ \mu = 0\} \cap \mu^{-1}(K) = \sum_{\beta_k} b_{k, \beta_k} D_{k, \beta_k}$ for any $1 \leq j \leq m$, $1 \leq k \leq q$ where $a_{j, \alpha_j}, b_{k, \beta_k}$ are positive integers. Note that the divisor D_{j, α_j} and D_{k, β_k} can be the same prime divisors. By the construction of μ , we know that all D_{j, α_j} and D_{k, β_k} are simple normal crossing divisors on \tilde{X} . Denote $D = (\cup_{i=1}^m \{f_j \circ \mu = 0\}) \cup (\cup_{k=1}^q \{h_k \circ \mu = 0\})$ be the support of all divisors appearing in $\{f_j \circ \mu = 0\}$ and $\{h_k \circ \mu = 0\}$.

Let $p \in Z_0$. Let U be a small neighborhood of p such that $U \cap D = \cup_{1 \leq l \leq n_p} \{w_l = 0\} = \cup_{1 \leq l \leq n_p} D_l$, where $w = (w_1, \dots, w_n)$ is the coordinate on U centered at p and $n_p \leq n$ is a positive integer. It follows from the construction of μ that one has

$$\begin{aligned} \varphi_N \circ \mu &= c \log \left(\sum_{1 \leq j \leq m} |f_j \circ \mu|^{\frac{1}{a_j}} + \sum_{1 \leq k \leq q} |h_k \circ \mu|^N \right) \\ &= c \log \left(\sum_{1 \leq j \leq m} |w_{j, \alpha_j}|^{\frac{a_{j, \alpha_j}}{a_j}} + \sum_{1 \leq k \leq q} |w_{k, \beta_k}|^{Nb_{k, \beta_k}} \right) \\ &= c \log \left(\prod_{1 \leq l \leq n_p} |w_l|^{\tau_{l, N}} \right) + O(1), \end{aligned}$$

where $\tau_{l, N} = \min_{(j, \alpha_j), (k, \beta_k)} \left\{ \frac{a_{j, \alpha_j}}{a_j}, Nb_{k, \beta_k} \right\}$ and the minimum is taken among all indices (j, α_j) and (k, β_k) satisfying $D_l = D_{j, \alpha_j}$ and $D_l = D_{k, \beta_k}$ respectively. For any $F \in \mathcal{O}_{\tilde{X}, p}$, we have

$$\sigma_p(\log |F|, \varphi_N) = \min_{1 \leq l \leq n_p} \left\{ \frac{\text{ord}_{D_l}(F)}{c\tau_{l, N}} \right\}.$$

Hence

$$\begin{aligned}
\sigma_p(\log |fg \circ \mu|, \varphi_N) &= \min_{1 \leq l \leq n_p} \left\{ \frac{\text{ord}_{D_l}(fg \circ \mu)}{c\tau_{l,N}} \right\} \\
&= \min_{1 \leq l \leq n_p} \left\{ \frac{\text{ord}_{D_l}(f \circ \mu) + \text{ord}_{D_l}(g \circ \mu)}{c\tau_{l,N}} \right\} \\
(2.19) \quad &\leq \min_{1 \leq l \leq n_p} \left\{ \frac{\text{ord}_{D_l}(f \circ \mu)}{c\tau_{l,N}} \right\} + \max_{1 \leq l \leq n_p} \left\{ \frac{\text{ord}_{D_l}(g \circ \mu)}{c\tau_{l,N}} \right\} \\
&= \sigma_p(\log |f \circ \mu|, \varphi_N) + \max_{1 \leq l \leq n_p} \left\{ \frac{\text{ord}_{D_l}(g \circ \mu)}{c\tau_{l,N}} \right\}
\end{aligned}$$

Note that $\tau_{l,N}$ is increasing with respect to N and thus $\max_{1 \leq l \leq n_p} \left\{ \frac{\text{ord}_{D_l}(g \circ \mu)}{c\tau_{l,N}} \right\}$ is decreasing with respect to N . Combining with the estimate (2.19), we have

$$\begin{aligned}
\sigma_p(\log |fg \circ \mu|, \varphi_N) &= \sigma_p(\log |f \circ \mu|, \varphi_N) + \max_{1 \leq l \leq n_p} \left\{ \frac{\text{ord}_{D_l}(g \circ \mu)}{c\tau_{l,N}} \right\} \\
(2.20) \quad &\leq \sigma_p(\log |f \circ \mu|, \varphi_N) + \max_{1 \leq l \leq n_p} \left\{ \frac{\text{ord}_{D_l}(g \circ \mu)}{c\tau_{l,1}} \right\}
\end{aligned}$$

Set $C := \max_{D_\gamma \subset D} \left\{ \frac{\text{ord}_{D_\gamma}(g \circ \mu)}{c\tau_{\gamma,1}} \right\}$, where D_γ is any prime divisor contained in D . Note that D contains finite prime divisors (depending on $\{f_j\}$ and $\{h_k\}$) and hence C is a finite number only depending on g , $\{f_j\}$ and $\{h_k\}$ and independent of N , f , p and $z_0 \in K$. It follows from inequality (2.20) and the definition of C that we have

$$\inf_{p \in Z_0} \sigma_p(\log |(f \circ \mu) \cdot (g \circ \mu)|, \varphi_N \circ \mu) \leq \inf_{p \in Z_0} \sigma_p(\log |f \circ \mu|, \varphi_N \circ \mu) + C.$$

Hence we know there exists a constant $C > 0$ (independent of N and K) such that

$$\sigma_{z_0}(\log |fg|, \varphi_N) \leq \sigma_{z_0}(\log |f|, \varphi_N) + C$$

holds for any positive N and $z_0 \in K$. \square

When o is a smooth point of X , we have the following relation between the jumping number and the relative type of Hölder continuous plurisubharmonic functions.

Lemma 2.38 (see [8]). *Let φ be a plurisubharmonic function near $o \in \mathbb{C}^n$ such that e^φ is α -Hölder for some $\alpha > 0$. Then for any $(f, o) \in \mathcal{O}_o^*$, we have*

$$c_o^f(\varphi) \leq \sigma(\log |f|, \varphi) + \frac{n}{\alpha}.$$

Recall that X is an analytic subset of $\Omega \in \mathbb{C}^n$ with pure dimension d and denote $|z|^2 := \sum_{1 \leq j \leq n} |z_j|^2$. Assume that (X, o) is singular, we have the following relation between the jumping number and the relative type when φ has analytic singularity.

Lemma 2.39. *Let $\varphi_0 := c \log \left(\sum_{1 \leq j \leq m} |f_j|^{\frac{1}{a_j}} \right)$ and $\varphi_N := c \log \left(\sum_{1 \leq j \leq m} |f_j|^{\frac{1}{a_j}} + |z|^{2N} \right)$ near o , where $c > 0$ is a constant, $\{a_j\}_{1 \leq j \leq m}$ are positive numbers and $f_j \in \mathcal{O}_{X,o}^w$ for $1 \leq j \leq m$. Then there exists a constant C (independent of N) such that for any $(f, o) \in \mathcal{O}_{X,o}^w \setminus \{0\}$ and $N \geq 0$, we have*

$$c_o^f(\varphi_N) \leq \sigma_o(\log |f|, \varphi_N) + C.$$

Proof. Using Theorem 2.4, we can resolve the singularities of X , and denote the corresponding proper modification by $\mu: \tilde{\Omega} \rightarrow \Omega$, where $\Omega \subset \mathbb{C}^n$ such that $o \in X \cap \Omega$. Denote the strict transform of X by \tilde{X} and denote $Z_0 := \tilde{X} \cap \mu^{-1}(\{o\})$. Note that $f \circ \mu$ and $\varphi_N \circ \mu$ are well defined holomorphic function and plurisubharmonic function correspondingly on $\pi^{-1}(U) \cap \tilde{X}$ for any open neighborhood U of o in X respectively. Since μ is proper, Z_0 is compact in \tilde{X} . Note that μ is holomorphic and biholomorphic outside an analytic subset. We know that $c_o^f(\varphi_N) = c_{Z_0}^{\det \mu \cdot f \circ \mu}(\varphi \circ \mu)$ and $\sigma_o(\log |f|, \varphi_N) = \sigma_{Z_0}(\log |f \circ \mu|, \varphi \circ \mu)$.

By adjusting c , we may assume all $\frac{1}{a_j} \geq 2$ and then e^{φ_N} is 1-Hölder continuous. For any $p \in Z_0$, Lemma 2.38 tells that

$$c_p^{\det \mu \cdot f \circ \mu}(\varphi_N \circ \mu) \leq \sigma_p(\log |\det \mu \cdot f \circ \mu|, \varphi_N \circ \mu) + C_1,$$

where C_1 is constant only depending on c and independent of p and N . Lemma 2.37 tells that, for any $p \in Z_0$, there exists a constant $C_2 > 0$ only depending on $\{f_j \circ \mu\}$, $\{z \circ \mu\}$ and μ and independent of p and N such that, for any $N > 0$,

$$\sigma_p(\log |\det \mu \cdot f \circ \mu|, \varphi_N \circ \mu) \leq \sigma_p(\log |f \circ \mu|, \varphi_N \circ \mu) + C_2.$$

Thus, for any $p \in Z_0$ and $N > 0$, we have

$$c_p^{\det \mu \cdot f \circ \mu}(\varphi_N \circ \mu) \leq \sigma_p(\log |f \circ \mu|, \varphi_N \circ \mu) + C,$$

where $C := C_1 + C_2 > 0$ is a constant independent of p and N .

Recall $c_{Z_0}^{\det \mu \cdot f \circ \mu}(\varphi_N \circ \mu) = \inf_{p \in Z_0} c_p^{\det \mu \cdot f \circ \mu}(\varphi_N \circ \mu)$ and $\sigma_{Z_0}(\log |f \circ \mu|, \varphi_N \circ \mu) = \inf_{p \in Z_0} \sigma_p(\log |f \circ \mu|, \varphi_N \circ \mu)$. Note that $c_o^f(\varphi_N) = c_{Z_0}^{\det \mu \cdot f \circ \mu}(\varphi \circ \mu)$ and $\sigma_o(\log |f|, \varphi_N) = \sigma_{Z_0}(\log |f \circ \mu|, \varphi \circ \mu)$. We know there exists a constant C (independent of N) such that for any $(f, o) \in \mathcal{O}_{X,o}^w \setminus \{0\}$ and $N \geq 0$,

$$\begin{aligned} c_o^f(\varphi_N) &= c_{Z_0}^{\det \mu \cdot f \circ \mu}(\varphi_N \circ \mu) \\ &\leq \sigma_{Z_0}(\log |f \circ \mu|, \varphi_N \circ \mu) + C \\ &= \sigma_o(\log |f|, \varphi_N) + C \end{aligned}$$

The proof is complete. \square

Remark 2.40. Let φ be a plurisubharmonic function near o satisfying that

$$\sup_{(f,o) \in \mathcal{O}_{X,o}^w \setminus \{0\}} (c_o^f(\varphi) - \sigma(\log |f|, \varphi)) < +\infty.$$

Then for any $(f, o) \in \mathcal{O}_{X,o}^w \setminus \{0\}$ with $c_o^f(\varphi) < +\infty$, considering the Tian function $\text{Tn}(t) := \{c \in \mathbb{R} : |f|^{2t} e^{-2c\varphi} \text{ is integrable near } o\}$, we have

$$\lim_{t \rightarrow +\infty} \frac{\text{Tn}(t)}{t} = \text{Tn}'_-(t) = \text{Tn}'_+(t) = \sigma(\log |f|, \varphi),$$

where $\text{Tn}'_-(t)$ and $\text{Tn}'_+(t)$ are the left and right derivatives of function $\text{Tn}(t)$, respectively.

Proof. By the definitions of $\text{Tn}(t)$ and $\sigma(\log |f|, \varphi)$, we have $\text{Tn}'_+(t) \geq \sigma(\log |f|, \varphi)$. It follows from the concavity of $\text{Tn}(t)$ that $\frac{\text{Tn}(t)}{t}$, $\text{Tn}'_-(t)$ and $\text{Tn}'_+(t)$ are decreasing and

$$\lim_{t \rightarrow +\infty} \frac{\text{Tn}(t)}{t} = \lim_{t \rightarrow +\infty} \text{Tn}'_-(t) = \lim_{t \rightarrow +\infty} \text{Tn}'_+(t).$$

As $\sup_{(f,o) \in \mathcal{O}_o^*} (c_o^f(\varphi) - \sigma(\log |f|, \varphi)) < +\infty$, there exists $C > 0$ such that $Tn(m) = c_o^m(\varphi) \leq m\sigma(\log |f|, \varphi) + C$ for any $m \in \mathbb{Z}_{>0}$, which implies $\lim_{m \rightarrow +\infty} \frac{Tn(m)}{m} \leq \sigma(\log |f|, \varphi)$. As $\frac{Tn(t)}{t} \geq \sigma(\log |f|, \varphi)$ for any $t \geq 0$ and $\lim_{t \rightarrow +\infty} \frac{Tn(t)}{t}$ exists, we have $\lim_{t \rightarrow +\infty} \frac{Tn(t)}{t} = \sigma(\log |f|, \varphi)$. \square

We recall the following basic property of holomorphic functions.

Lemma 2.41. *Let X be a connected complex manifold. Let $f \not\equiv 0$ be a holomorphic function defined on an open neighborhood of a compact set $K \subset X$. Then $\sup_{p \in K} \text{ord}_p f < +\infty$.*

Proof. If not, then $\sup_{p \in K} \text{ord}_p f = +\infty$. There exists a sequence of points $\{p_m\}_{m=1}^{+\infty}$ contained in K such that $\lim_{m \rightarrow +\infty} \text{ord}_{p_m} f = +\infty$. As K is compact, there exists a point $p \in K$ such that some subsequence of $\{p_m\}$ (also denoted by $\{p_m\}$) converges to p .

Let $(U; z_1, z_2, \dots, z_n)$ be a small local coordinated neighborhood near p . As $\lim_{m \rightarrow +\infty} \text{ord}_{p_m} f = +\infty$, for any multiple index $\alpha = (\alpha_1, \dots, \alpha_n)$, we have $\frac{\partial^\alpha f}{\partial z^\alpha}(p_m) = 0$ when m is large enough. By the continuity of $\frac{\partial^\alpha f}{\partial z^\alpha}$ and that p_m converges to p , we know $\frac{\partial^\alpha f}{\partial z^\alpha}(p) = 0$ for any multiple index $\alpha = (\alpha_1, \dots, \alpha_n)$. Thus, by the rigidity of holomorphic function, we know that f is identical zero which is a contradiction. \square

We recall the following result due to Skoda.

Lemma 2.42 ([28]; see also [9]). *For any plurisubharmonic function φ near o , if the Lelong number $\nu_o(\varphi) < 1$, then $e^{-2\varphi}$ is locally integrable near o .*

Using Lemma 2.42, we present the following lemma.

Lemma 2.43. *Let (X, o) be an irreducible germ of an analytic set. Let $C > 0$, and φ a plurisubharmonic function defined near (X, o) such that $c_o(\varphi) < C$. Then for any holomorphic function $f \in \mathcal{O}_{X,o}^w$, there exists a constant $C_f \geq 0$ (depending on f but independent of φ) such that*

$$\sigma_o(\log |f|, \varphi) \leq CC_f.$$

Proof. Using Theorem 2.4, we can resolve the singularities of X , and denote the corresponding proper modification by $\mu: \tilde{\Omega} \rightarrow \Omega$, where $\Omega \subset \mathbb{C}^{n+l}$ such that $o \in X \cap \Omega$. Denote the strict transform of X by \tilde{X} and denote $Z_0 := \tilde{\Omega} \cap \mu^{-1}(\{o\})$. Note that $c_o(\varphi) = c_{Z_0 \cap \tilde{X}}^{\det \mu}(\varphi \circ \mu)$. Hence $c_o(\varphi) < C$ implies that

$$c_{Z_0 \cap \tilde{X}}^{\det \mu}(\varphi \circ \mu) < C.$$

Recall that $c_{Z_0 \cap \tilde{X}}^{\det \mu}(\varphi \circ \mu) = \inf_{p \in Z_0 \cap \tilde{X}} c_p^{\det \mu}(\varphi \circ \mu)$. There exists a point $p \in Z_0 \cap \tilde{X}$ such that $c_p^{\det \mu}(\varphi \circ \mu) < C$. As $c_p^{\det \mu}(\varphi \circ \mu) \geq c_p(\varphi \circ \mu)$, we know that $c_p(\varphi \circ \mu) < C$. It follows from Lemma 2.42 that the Lelong number of $\varphi \circ \mu$ at p

$$\nu_p(\varphi \circ \mu) > \frac{1}{C},$$

which implies $\sigma_p(\log |f \circ \mu|, \varphi \circ \mu) \leq C \text{ord}_p(f)$.

Recall that $\sigma_o(\log|f|, \varphi) = \sigma_{Z_0 \cap \tilde{X}}(\log|f \circ \mu|, \varphi \circ \mu)$ and $\sigma_{Z_0 \cap \tilde{X}}(\log|f \circ \mu|, \varphi \circ \mu) = \inf_{p \in Z_0 \cap \tilde{X}} \sigma(\log|f \circ \mu|, \varphi \circ \mu)$. Hence we know

$$\sigma_o(\log|f|, \varphi) \leq \sigma_p(\log|f \circ \mu|, \varphi \circ \mu) \leq C \operatorname{ord}_p(f \circ \mu) \leq C \sup_{p \in Z_0 \cap \tilde{X}} \operatorname{ord}_p(f \circ \mu).$$

Note that $Z_0 \cap \tilde{X}$ is compact in \tilde{X} , Lemma 2.41 tells that $C_f := \sup_{p \in Z_0 \cap \tilde{X}} \operatorname{ord}_p(f \circ \mu) < +\infty$. Hence we know $\sigma_o(\log|f|, \varphi) \leq C C_f$. \square

2.6. Convergence results for valuations and for relative types.

Let $\{\nu_j\}_{j \in \mathbb{Z}_{>0}}$ be a sequence of valuations on \mathcal{O}_o . We need the following convergence result for valuations, which will be used to prove the main theorem.

Proposition 2.44. *Assume that $\sup_j \nu_j(g) < +\infty$ for any holomorphic function g near o . There exists a subsequence of $\{\nu_j\}$ denoted by $\{\nu_{j_l}\}$ such that $\{\nu_{j_l}\}$ converges to a valuation ν_0 on \mathcal{O}_o .*

Proof. Denote $I_k^j := \{(f, o) \in \mathcal{O}_o : \nu_j(f) \geq k\}$ for every $j, k \in \mathbb{Z}_{>0}$.

Step 1. Find a subsequence $\{I_k^{j_l}\}$ of I_k^j such that $\cup_{l \geq 1} (\cap_{l \geq l_1} I_k^{j_l})$ is maximal for any k . It means that for any subsequence $\{I_k^{j'_l}\}$ of $\{I_k^{j_l}\}$, $\cup_{l \geq 1} (\cap_{l \geq l_1} I_k^{j'_l}) = \cup_{l \geq 1} (\cap_{l \geq l_1} I_k^{j_l})$.

Note that

$$\cup_{j_1 \geq 1} (\cap_{j \geq j_1} I_k^j) = \{(f, o) \in \mathcal{O}_o : \exists j_1 \text{ s.t. } \nu_j(f) \geq k \text{ holds for any } j \geq j_1\}.$$

As \mathcal{O}_o is a Noetherian ring, there exists a subsequence $\{I_1^{j^{(1)}(l)}\}$ of I_1^j such that $\cup_{l \geq 1} (\cap_{l \geq l_1} I_1^{j^{(1)}(l)})$ is maximal. Similarly, there exists a subsequence $\{I_1^{j^{(2)}(l)}\}$ of $I_1^{j^{(1)}(l)}$ such that $\cup_{l \geq 1} (\cap_{l \geq l_1} I_2^{j^{(2)}(l)})$ is maximal. By induction, for any $r \in \mathbb{Z}_{\geq 1}$, there exists a sequence $\{I_1^{j^{(r)}(l)}\}$ such that $\cup_{l \geq 1} (\cap_{l \geq l_1} I_r^{j^{(r)}(l)})$ is maximal and $\{I_1^{j^{(r+1)}(l)}\}$ is a subsequence of $\{I_1^{j^{(r)}(l)}\}$. Using the diagonal method, we obtain a subsequence $\{I_k^{j_l}\}$ of I_k^j such that $\cup_{l \geq 1} (\cap_{l \geq l_1} I_k^{j_l})$ is maximal for any k .

Step 2. We prove that $\lim_{l \rightarrow +\infty} \nu_{j_l}(f)$ exists for any $(f, o) \in \mathcal{O}_o$.

We prove it by contradiction: if not, there exist two subsequences of valuations $\{\nu_{j'_l}\}$ and $\{\nu_{j''_l}\}$ of $\{\nu_{j_l}\}$ such that

$$\lim_{l \rightarrow +\infty} \nu_{j'_l}(f) = a_1 < a_2 = \lim_{l \rightarrow +\infty} \nu_{j''_l}(f).$$

Note that $\nu_j(f^m) = m \nu_j(f)$. Then, without loss of generality, assume that there exists a positive integer $k_0 \in (a_1, a_2)$. Then we have

$$(f, o) \in \cup_{l \geq 1} (\cap_{l \geq l_1} I_{k_0}^{j'_l})$$

and

$$(f, o) \notin \cup_{l \geq 1} (\cap_{l \geq l_1} I_{k_0}^{j_l}),$$

which contradicts to the maximality of $\cup_{l \geq 1} (\cap_{l \geq l_1} I_{k_0}^{j_l})$. Thus, $\lim_{l \rightarrow +\infty} \nu_{j_l}(f)$ exists for any $(f, o) \in \mathcal{O}_o$.

Denote $\nu_0(f) := \lim_{l \rightarrow +\infty} \nu_{j_l}(f)$ for any $(f, o) \in \mathcal{O}_o$. It follows from every ν_j is a valuation on \mathcal{O}_o that ν_0 is a valuation on \mathcal{O}_o . Proposition 2.44 has been proved. \square

Let us consider valuations on the ring of germs of weakly holomorphic functions. We recall the following basic property of the ring.

Lemma 2.45 (see [31]). *The ring of germs of weakly holomorphic functions \mathcal{O}_o^w is Noetherian.*

As \mathcal{O}_o^w is Noetherian, using the same proof as Proposition 2.44, we also have the following convergence result.

Proposition 2.46. *Let $\{\nu_j\}_{j \in \mathbb{Z}_{>0}}$ be a sequence of valuations on \mathcal{O}_o^w . Assume that $\sup_j \nu_j(g) < +\infty$ for any weakly holomorphic function g near o . Then there exists a subsequence $\{\nu_{j_l}\}$ of $\{\nu_j\}$ which converges to a valuation ν_0 on \mathcal{O}_o^w .*

3. PROOF OF REMARK 1.3

Proof of Remark 1.3. Theorem 2.7 shows that there exists $\varepsilon > 0$, such that

$$|f_0|^2 e^{-2\varphi_0} |z|^{2N_0} e^{-2(1+\varepsilon)\varphi}$$

is integrable near z_0 . We have

$$\begin{aligned} & \int_U |f_0|^2 e^{-2\varphi_0} e^{-2\varphi} - |f_0|^2 e^{-2\varphi_0} e^{-2 \max\{\varphi, \frac{N_0}{\varepsilon} \log |z|\}} \\ & \leq \int_{U \cap \left\{ \frac{N_0}{\varepsilon} \log |z| \geq \varphi \right\}} |f_0|^2 e^{-2\varphi_0} e^{-2\varphi} \\ & \leq \int_{U \cap \left\{ \frac{N_0}{\varepsilon} \log |z| \geq \varphi \right\}} |f_0|^2 e^{-2\varphi_0} |z|^{2N_0} e^{-2(1+\varepsilon)\varphi} \\ & < +\infty, \end{aligned}$$

where $U \subset X$ is a neighborhood of z_0 . Then it suffices to consider that there exists N large enough such that

$$N \log |z| \leq \varphi$$

near z_0 . By a similar discussion, for any local Zhou weight $\Phi_{z_0, \max}$, $\Phi_{z_0, \max} \geq N_1 \log |z| + O(1)$ near o for some $N_1 \gg 0$.

Let $V \subset X$ be a small neighborhood of z_0 such that $e^{-\varphi}$ is integrable near any $z \in V \setminus \{z_0\}$ and $\varphi < 0$ on V . Let $(u_\alpha)_\alpha$ be the negative plurisubharmonic functions on V such that $u_\alpha \geq \varphi + O(1)$ near z_0 and $|f_0|^2 e^{-2\varphi_0} e^{-2u_\alpha}$ is not integrable near z_0 .

Zorn's Lemma shows that there exists Γ which is a maximal set such that for any $\alpha, \alpha' \in \Gamma$, $u_\alpha \leq u_{\alpha'} + O(1)$ or $u_{\alpha'} \leq u_\alpha + O(1)$ holds near o , where (u_α) are negative plurisubharmonic functions on V .

Let $u(z) := \sup_{\alpha \in \Gamma} u_\alpha(z)$ on V , and let $u^*(z) = \lim_{\varepsilon \rightarrow 0} \sup_{U_{z, \varepsilon}} u$, where $U_{z, \varepsilon} := X \cap \mathbb{B}^{n+l}(z, \varepsilon)$. Lemma 2.1 shows that there exists subsequence (v_j) of (u_α) such that $(\max_j v_j)^* = u^*$. Moreover one can choose v_j ($\sup_{j' \leq j} v_j$) increasing with respect to j such that $|f_0|^2 e^{-2\varphi_0} e^{-2v_j}$ is not integrable near z_0 .

Lemma 2.1 shows that (v_j) is convergent to v^* with respect to j almost everywhere with respect to Lebesgue measure, and v^* is a plurisubharmonic function on V . Theorem 2.7 shows that $|f_0|^2 e^{-2\varphi_0} e^{-2v^*}$ is not integrable near z_0 .

In the following part, we prove that v^* is a local Zhou weight related to $|f_0|^2 e^{-2\varphi_0}$ near z_0 by contradiction. If not, then there exists a plurisubharmonic function \tilde{v}

near z_0 such that $\tilde{v} \geq v^*$, $|f_0|^2 e^{-2\varphi_0} e^{-2\tilde{v}}$ is not integrable near z_0 , and

$$\limsup_{z \rightarrow z_0} (\tilde{v}(z) - v^*(z)) = +\infty.$$

As $\varphi < 0$ on V , by the definition of v^* , we have $v^* \geq \varphi$, which shows $v^* \geq N \log |z|$. Then for $U_{z_0, \varepsilon} := \mathbb{B}(z_0, \varepsilon) \cap X$ (ε is small enough), there exists $M \ll 0$ such that $\tilde{v} + M < N \log |z| \leq v^*$ near the boundary of $U_{z_0, \varepsilon}$, which implies that $\max\{\tilde{v} + M, v^*\} = v^*$ near the boundary of $U_{z_0, \varepsilon}$. Let

$$\tilde{\varphi} := \begin{cases} \max\{\tilde{v} + M, v^*\} & \text{on } U_{z_0, \varepsilon}, \\ v^* & \text{on } V \setminus U_{z_0, \varepsilon}. \end{cases}$$

$\tilde{v} \geq v^*$ implies that $\tilde{\varphi} = \tilde{v} + O(1)$ near z_0 . Then $\tilde{\varphi}$ is a plurisubharmonic function on V such that

$$\limsup_{z \rightarrow z_0} (\tilde{\varphi}(z) - u^*(z)) = \limsup_{z \rightarrow z_0} (\tilde{\varphi}(z) - v^*(z)) \geq \limsup_{z \rightarrow z_0} (\tilde{v}(z) + M - v^*(z)) = +\infty,$$

and $|f_0|^2 e^{-2\varphi_0} e^{-2\tilde{\varphi}}$ is not integrable near z_0 , which contradicts the definition of u^* .

This proves Remark 1.3. \square

4. PROOF OF THEOREM 1.6

Proof of Theorem 1.6. Remark 1.3 shows that $\Phi_{z_0, \max} \geq N \log |z| + O(1)$ near z_0 for some $N > 0$. Following from Propositions 2.12 and 2.13, we know that for any plurisubharmonic function ψ satisfying that $\psi \leq s_0 \log |z| + O(1)$ near $z_0 \in X$ for some $s_0 > 0$, the limit $\lim_{t_1 \rightarrow +\infty} \frac{1}{2t_1} \frac{\int_{\{\mathrm{Tn}(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0} (-2\psi)}{\int_{\{\mathrm{Tn}(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0}}$ exists and

$$\lim_{t_1 \rightarrow +\infty} \frac{1}{2t_1} \frac{\int_{\{\mathrm{Tn}(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0} (-2\psi)}{\int_{\{\mathrm{Tn}(0)\varphi < -t_1\} \cap U} |f_0|^2 e^{-2\varphi_0}} = \sigma(\psi, \Phi_{z_0, \max}).$$

Thus, Theorem 1.6 holds. \square

Proof of Corollary 1.7. If $f(z_0) = 0$, it is clear that $\log |f| \leq C_1 \log |z| + O(1)$ near z_0 for some $C_1 > 0$. Thus, $\nu(f, \Phi_{z_0, \max}) > 0$ since $\Phi_{z_0, \max} \geq C_2 \log |z| + O(1)$ near z_0 .

If $f(z_0) \neq 0$, then there exists a neighborhood U such that $\inf_U |f| > 0$. As $|f_0|^2 e^{-2\varphi_0 - 2\Phi_{z_0, \max}}$ is not integrable near z_0 and $|f_0|^2 e^{-2\varphi_0}$ is integrable near z_0 , we know that $\liminf_{X_{\mathrm{reg}} \ni z \rightarrow z_0} \Phi_{z_0, \max} = -\infty$. Then $\nu(f, \Phi_{z_0, \max}) = 0$. The statement (3) holds.

Note that $\log |fg| = \log |f| + \log |g|$. It follows from Theorem 1.6 and statement (3) that $\nu(fg, \Phi_{z_0, \max}) = \nu(f, \Phi_{z_0, \max}) + \nu(g, \Phi_{z_0, \max})$. The statement (1) holds.

As $\log |f + g| \leq \log(|f| + |g|) = \max\{\log |f|, \log |g|\} + O(1)$, we have

$$\nu(f + g, \Phi_{z_0, \max}) \geq \sigma(\max\{\log |f|, \log |g|\}, \Phi_{z_0, \max}).$$

For any positive number $c < \min\{\nu(f, \Phi_{z_0, \max}), \nu(g, \Phi_{z_0, \max})\}$, by definition, we have $\log |f| \leq c\Phi_{z_0, \max} + O(1)$ and $\log |g| \leq c\Phi_{z_0, \max} + O(1)$ near z_0 , which shows $\max\{\log |f|, \log |g|\} \log c\Phi_{z_0, \max} + O(1)$ near z_0 . Hence, $\nu(f + g, \Phi_{z_0, \max}) \geq \sigma(\max\{\log |f|, \log |g|\}, \Phi_{z_0, \max}) \geq \min\{\nu(f, \Phi_{z_0, \max}), \nu(g, \Phi_{z_0, \max})\}$. The statement (2) holds.

Thus, Corollary 1.7 holds. \square

Proposition 4.1. *Let $f = (f_1, \dots, f_{m'})$ be a vector, where $f_1, \dots, f_{m'}$ are holomorphic functions near z_0 . Denote by $|f| := (\sum |f_i|^2)^{1/2}$. Let $\Phi_{z_0, \max}$ be a local Zhou weight related to $|f_0|^2 e^{-2\varphi_0}$ near z_0 . Then the following two statements hold:*

- (1) *for any $\alpha > 0$, $(1 + \alpha \sigma(\log |f|, \Phi_{z_0, \max})) \Phi_{z_0, \max}$ is a local Zhou weight related to $|f|^{2\alpha} |f_0|^2 e^{-2\varphi_0}$ near z_0 ;*
- (2) *$(1 + \sigma(\varphi_0, \Phi_{z_0, \max})) \Phi_{z_0, \max}$ is a local Zhou weight related to $|f_0|^2$.*

Proof. Firstly, we prove the statement (1) in Proposition 4.1.

Denote by

$$\text{Tn}(t) := \sup \{c : |f_0|^2 |f|^{2t} e^{-2\varphi_0 - 2c\Phi_{z_0, \max}} \text{ is integrable near } o\}.$$

By Proposition 2.13, we have $\text{Tn}(\alpha) = 1 + \sigma(\log |f|, \Phi_{z_0, \max})\alpha$, which implies that

$$|f_0|^2 |f|^{2\alpha} e^{-2\varphi_0 - 2(1 + \alpha \sigma(\log |f|, \Phi_{z_0, \max})) \Phi_{z_0, \max}}$$

is not integrable near z_0 by Theorem 2.7. As $\log |f| \leq \sigma(\log |f|, \Phi_{z_0, \max})\Phi_{z_0, \max} + O(1)$ and there exists $N \gg 0$ such that $|f_0|^2 |z|^{2N} e^{-2\varphi_0 - 2\Phi_{z_0, \max}}$ is integrable near z_0 , we know that

$$|f_0|^2 |f|^{2\alpha} |z|^{2N} e^{-2\varphi_0 - 2(1 + \alpha \sigma(\log |f|, \Phi_{z_0, \max})) \Phi_{z_0, \max}}$$

is integrable near z_0 .

Let $\tilde{\varphi}$ be a plurisubharmonic function near z_0 satisfying that

$$\tilde{\varphi} \geq (1 + \alpha \sigma(\log |f|, \Phi_{z_0, \max})) \Phi_{z_0, \max} + O(1)$$

and $|f|^{2\alpha} |f_0|^2 e^{-2\varphi_0 - 2\tilde{\varphi}}$ is not integrable near z_0 . Note that

$$\log |f| \leq \sigma(\log |f|, \Phi_{z_0, \max})\Phi_{z_0, \max} + O(1),$$

then

$$\tilde{\varphi} \geq \frac{1 + \alpha \sigma(\log |f|, \Phi_{z_0, \max})}{\sigma(\log |f|, \Phi_{z_0, \max})} \log |f| + O(1)$$

and

$$|f|^{2\alpha} |f_0|^2 e^{-2\varphi_0 - 2\tilde{\varphi}} \leq C |f_0|^2 e^{-2\varphi_0} e^{-\frac{2}{1 + \alpha \sigma(\log |f|, \Phi_{z_0, \max})} \tilde{\varphi}}.$$

As $|f|^{2\alpha} |f_0|^2 e^{-2\varphi_0 - 2\tilde{\varphi}}$ is not integrable near z_0 , we know that

$$|f_0|^2 e^{-2\varphi_0 - \frac{2}{1 + \alpha \sigma(\log |f|, \Phi_{z_0, \max})} \tilde{\varphi}}$$

is not integrable near z_0 . Note that $\Phi_{z_0, \max}$ is a local Zhou weight related to $|f_0|^2 e^{-2\varphi_0}$. Then we obtain

$$\tilde{\varphi} = (1 + \alpha \sigma(\log |f|, \Phi_{z_0, \max})) \Phi_{z_0, \max} + O(1),$$

which shows that $(1 + \alpha \sigma(\log |f|, \Phi_{z_0, \max})) \Phi_{z_0, \max}$ is a local Zhou weight related to $|f|^{2\alpha} |f_0|^2 e^{-2\varphi_0}$.

Next, we give the proof of statement (2), which is similar to the proof of statement (1).

Denote by

$$\text{Tn}(t) := \sup \{c : |f_0|^2 e^{-2\varphi_0} e^{2t\varphi_0} e^{-2c\Phi_{z_0, \max}} \text{ is integrable near } o\}.$$

By Proposition 2.13, we have $\text{Tn}(1) = 1 + \sigma(\varphi_0, \Phi_{z_0, \max})$, which implies that

$$|f_0|^2 e^{-2(1 + \sigma(\varphi_0, \Phi_{z_0, \max})) \Phi_{z_0, \max}}$$

is not integrable near z_0 . As $\varphi_0 \leq \sigma(\varphi_0, \Phi_{z_0, \max})\Phi_{z_0, \max} + O(1)$ and there exists $N \gg 0$ such that $|f_0|^2|z|^{2N}e^{-2\varphi_0-2\Phi_{z_0, \max}}$ is integrable near z_0 , we know that $|f_0|^2|z|^{2N}e^{-2(1+\sigma(\varphi_0, \Phi_{z_0, \max}))\Phi_{z_0, \max}}$ is integrable near z_0 .

Let $\tilde{\varphi}$ be a plurisubharmonic function near z_0 satisfying that

$$\tilde{\varphi} \geq (1 + \sigma(\varphi_0, \Phi_{z_0, \max}))\Phi_{z_0, \max} + O(1)$$

and $|f_0|^2e^{-2\tilde{\varphi}}$ is not integrable near z_0 . It follows from $\varphi_0 \leq \sigma(\varphi_0, \Phi_{z_0, \max})\Phi_{z_0, \max} + O(1)$ that

$$\begin{aligned} |f_0|^2e^{-2\tilde{\varphi}} &= e^{2\varphi_0}|f_0|^2e^{-2\varphi_0}e^{-2\tilde{\varphi}} \\ &\leq Ce^{2\sigma(\varphi_0, \Phi_{z_0, \max})\Phi_{z_0, \max}}|f_0|^2e^{-2\varphi_0}e^{-2\tilde{\varphi}} \\ &\leq C_1|f_0|^2e^{-2\varphi_0}e^{2\frac{\sigma(\varphi_0, \Phi_{z_0, \max})}{1+\sigma(\varphi_0, \Phi_{z_0, \max})}\tilde{\varphi}}e^{-2\tilde{\varphi}} \\ &= C_1|f_0|^2e^{-2\varphi_0}e^{-\frac{2}{1+\sigma(\varphi_0, \Phi_{z_0, \max})}\tilde{\varphi}}. \end{aligned}$$

As $|f_0|^2e^{-2\tilde{\varphi}}$ is not integrable near z_0 , we know that $|f_0|^2e^{-2\varphi_0-\frac{2}{1+\sigma(\varphi_0, \Phi_{z_0, \max})}\tilde{\varphi}}$ is not integrable near z_0 . Note that $\Phi_{z_0, \max}$ is a local Zhou weight related to $|f_0|^2e^{-2\varphi_0}$. Then we obtain

$$\tilde{\varphi} = (1 + \sigma(\varphi_0, \Phi_{z_0, \max}))\Phi_{z_0, \max} + O(1),$$

which shows that $(1 + \sigma(\varphi_0, \Phi_{z_0, \max}))\Phi_{z_0, \max}$ is a local Zhou weight related to $|f_0|^2$. \square

5. PROOF OF THEOREM 1.8

It is clear that

$$\sigma\left(\log|G|, (1 + \sigma(\varphi_0, \Phi_{z_0, \max}))\Phi_{z_0, \max}\right) = \frac{\nu(G, \Phi_{z_0, \max})}{1 + \sigma(\varphi_0, \Phi_{z_0, \max})}$$

and

$$c_{z_0}^G\left((1 + \sigma(\varphi_0, \Phi_{z_0, \max}))\Phi_{z_0, \max}\right) = \frac{c_{z_0}^G(\Phi_{z_0, \max})}{1 + \sigma(\varphi_0, \Phi_{z_0, \max})}.$$

Proposition 4.1 shows that it suffices to consider the case $\varphi_0 \equiv 0$. Let us consider a mixed Tian function

$$\text{Tn}(s, t) := \sup \{c : |G|^{2s}|f_0|^{2t}e^{-2c\Phi_{z_0, \max}} \text{ is integrable near } z_0\}.$$

By definitions of $\text{Tn}(s, t)$ and Zhou numbers, we have

$$\text{Tn}(1, 0) = c_{z_0}^G(\Phi_{z_0, \max}) \geq c_{z_0}(\Phi_{z_0, \max}) + \nu(G, \Phi_{z_0, \max})$$

and $\text{Tn}(1, 1) \geq \text{Tn}(1, 0) + \sigma(\log|f_0|, \Phi_{z_0, \max})$. Since $\Phi_{z_0, \max}$ is a local Zhou weight related to $|f_0|^2$ near z_0 , we have $\text{Tn}(0, 1) = 1$ and $\text{Tn}(1, 1) = \text{Tn}(0, 1) + \nu(G, \Phi_{z_0, \max})$ (by Proposition 4.1). Then we have

$$\begin{aligned} c_{z_0}^G(\Phi_{z_0, \max}) &= \text{Tn}(1, 0) \leq \text{Tn}(1, 1) - \sigma(\log|f_0|, \Phi_{z_0, \max}) \\ &= \text{Tn}(0, 1) + \nu(G, \Phi_{z_0, \max}) - \sigma(\log|f_0|, \Phi_{z_0, \max}) \\ &= \nu(G, \Phi_{z_0, \max}) - \sigma(\log|f_0|, \Phi_{z_0, \max}) + 1. \end{aligned}$$

Thus, Theorem 1.8 holds.

6. PROOFS OF THEOREM 1.9, THEOREM 1.10 AND THEIR COROLLARIES

We firstly proof Theorem 1.9.

Proof. We prove this theorem in two steps.

Step 1. (1) \Rightarrow (2).

Note that $F := \prod_{0 \leq j \leq m} f_j$. If there exists a valuation ν such that $\nu(f_j) = a_j$ for any j , then by Lemma 2.17 and $a_0 = 0$,

$$\sigma(\log |F|, \varphi) \leq \nu(F) = \sum_{0 \leq j \leq m} \nu(f_j) = \sum_{1 \leq j \leq m} a_j.$$

On the other hand, by the definition of relative type, clearly we also have

$$\sigma(\log |F|, \varphi) \geq \sum_{0 \leq j \leq m} \sigma(\log |f_j|, \varphi) \geq \sum_{1 \leq j \leq m} a_j.$$

Thus, $\sigma(\log |F|, \varphi) = \sum_{1 \leq j \leq m} a_j$ holds.

Step 2. (2) \Rightarrow (1).

Recall that X is an analytic subset of $\Omega \in \mathbb{C}^n$ with pure dimension n and denote $|z|^2 := \sum_{1 \leq j \leq n} |z_j|^2$

As $\sigma(\log |F|, \varphi) < +\infty$, we have $f_j(o) = 0$ for any $1 \leq j \leq m$. Denote $\varphi_N := \max\{\varphi, N \log |z|\}$ for any $N > 0$. Then we have $\sigma(\log |f_j|, \varphi_N) \geq \sigma(\log |f_j|, \varphi) \geq a_j$ for all $0 \leq j \leq m$. It follows from Lemma 2.39 that there exists $C > 0$ such that $c_o^f(\varphi) \leq \sigma(\log |f|, \varphi) + C$ for any $(f, o) \in \mathcal{O}_{X,o}$. Thus, for any $k > 0$, there exists N_k such that

$$\begin{aligned} \sigma(\log |F|, \varphi) &\geq \frac{c_o^{F^k}(\varphi) - C}{k} \\ &\geq \frac{c_o^{F^k}(\varphi_{N_k}) - 1 - C}{k} \\ &\geq \frac{k\sigma(\log |F|, \varphi_{N_k}) - 1 - C}{k} \\ &\geq \lim_{N \rightarrow +\infty} \sigma(\log |F|, \varphi_N) - \frac{1 + C}{k}, \end{aligned}$$

where the second “ \geq ” follows from Lemma 2.34. Then we have

$$\lim_{N \rightarrow +\infty} \sigma(\log |F|, \varphi_N) = \sigma(\log |F|, \varphi).$$

Consider the Tian function

$$\text{Tn}(t) := \sup \{c \geq 0 : |F|^{2t} e^{-2c\varphi_N} \text{ is integrable near } o\}.$$

It follows from Lemma 2.39 and Remark 2.40 that

$$(6.1) \quad \lim_{t \rightarrow +\infty} \text{Tn}'_-(t) = \lim_{t \rightarrow +\infty} \text{Tn}'_+(t) = \sigma(\log |F|, \varphi_N).$$

Then for every positive integer s , $|F|^{2s} e^{-2\text{Tn}(s)\varphi_N}$ is not integrable near o (because of the strong openness property of multiplier ideal sheaves [17]), then by Theorem 1.3, there exists a local Zhou weight $\Phi_{N,s}$ near o related to $|F|^{2s}$ such that

$$\Phi_{N,s} \geq \text{Tn}(s)\varphi_N.$$

Consider the Tian function

$$\widetilde{\text{Tn}}(t) := \sup \left\{ c \geq 0 : |F|^{2t} e^{-2c\frac{\Phi_{N,s}}{\text{Tn}(s)}} \text{ is integrable near } o \right\}.$$

Thus, we have $\widetilde{\text{Tn}}(s) = \text{Tn}(s)$ and $\widetilde{\text{Tn}}(t) \geq \text{Tn}(t)$ for any $t \geq 0$. It follows from Proposition 2.13 that $\widetilde{\text{Tn}}'(s) = \sigma\left(\log|F|, \frac{\Phi_{N,s}}{\text{Tn}(s)}\right)$. Then we have

$$(6.2) \quad \text{Tn}'_+(s) \leq \sigma\left(\log|F|, \frac{\Phi_{N,s}}{\text{Tn}(s)}\right) \leq \text{Tn}'_-(s).$$

Denote the corresponding Zhou valuation of $\Phi_{N,s}$ by $\nu_{N,s}$. Then we have

$$\text{Tn}(s)\nu_{N,s}(f_j) = \text{Tn}(s)\sigma(\log|f_j|, \Phi_{N,s}) \geq \sigma(\log|f_j|, \varphi_N) \geq a_j$$

for all $1 \leq j \leq m$ and

$$\text{Tn}(s)\nu_{N,s}(F) = \text{Tn}(s)\sigma(\log|F|, \Phi_{N,s}) = \sigma\left(\log|F|, \frac{\Phi_{N,s}}{\text{Tn}(s)}\right).$$

Note that

$$\varphi_N = \log\left(\sum_{1 \leq j \leq m} |f_j|^{\frac{1}{a_j}} + \sum_{1 \leq k \leq n} |z_k|^N\right) + O(1)$$

near o . By Lemma 2.39, there exists a constant C (independent of N) such that $\sup_{(f,o) \in \mathcal{O}_o^*} (c_o^f(\varphi_N) - \sigma(\log|f|, \varphi_N)) < C$. As $\widetilde{\text{Tn}}(t)$ is a concave function, we have

$$(6.3) \quad c_o\left(\frac{\Phi_{N,s}}{\text{Tn}(s)}\right) = \widetilde{\text{Tn}}(0) \leq \text{Tn}(s) - s\text{Tn}'_+(s).$$

It follows from the concavity of $\text{Tn}(t)$ and equality (6.1) that $\text{Tn}(s) - s\text{Tn}'_+(s) \leq c_o^{F^s}(\varphi_N) - \sigma(\log|F^s|, \varphi_N) < C$. Then the inequality (6.3) implies $c_o\left(\frac{\Phi_{N,s}}{\text{Tn}(s)}\right) < C$ for any N and s . It follows from Lemma 2.43 that for any (f,o) ,

$$(6.4) \quad \text{Tn}(s)\nu_{N,s}(f) = \sigma\left(\log|f|, \frac{\Phi_{N,s}}{\text{Tn}(s)}\right)$$

is uniformly bounded with respect to N and s .

By equality (6.1), inequalities (6.2), (6.4), and Proposition 2.44, there exists a subsequence of the valuations $\{\text{Tn}(s)\nu_{N,s}\}_s$ (also denoted by $\{\text{Tn}(s)\nu_{N,s}\}_s$), which converges to a valuation ν_N . Then we have

$$\nu_N(f_j) \geq a_j, \quad j = 0, \dots, m,$$

and

$$\nu_N(F) = \sigma(\log|F|, \varphi_N).$$

By the same reason, there exists a subsequence of the valuations $\{\nu_N\}_N$ (also denoted by $\{\nu_N\}_N$), which converges to a valuation ν . Then we have $\nu(f_j) \geq a_j$ for any $0 \leq j \leq m$ and

$$\sum_{0 \leq j \leq m} \nu(f_j) = \nu(F) = \lim_{N \rightarrow +\infty} \sigma(\log|F|, \varphi_N) = \sigma(\log|F|, \varphi) = \sum_{1 \leq j \leq m} a_j.$$

Combining the equality above and $a_0 = 0$, we have

$$\nu(f_j) = a_j, \quad j = 0, \dots, m.$$

□

We now prove Theorem 1.10.

Proof. The idea for the proof of Theorem 1.10 is the same as the proof of Theorem 1.9. We just give the different Lemmas or Propositions used in the two proof and omit the step by step proof of Theorem 1.10.

We prove this theorem in two steps.

Step 1. (1) \Rightarrow (2).

Replace Lemma 2.17 by Lemma 2.18. Using the same argument as (1) \Rightarrow (2) in the proof of Theorem 1.9, we know (1) \Rightarrow (2) in Theorem 1.10 holds.

Step 2. (2) \Rightarrow (1).

Replace Theorem 1.3 by Theorem 1.3 and Theorem 1.5, Proposition 2.44 by Proposition 2.46. Using the same argument as (2) \Rightarrow (1) in the proof of Theorem 1.9, we know (2) \Rightarrow (1) of Theorem 1.10 holds. \square

Now we prove Corollary 1.12.

Proof of Corollary 1.12. If $\sigma(\log |F|, \varphi) = \sum_{1 \leq j \leq m} a_j$, Theorem 1.9 shows that there exists a valuation ν on $\mathcal{O}_{X,o} = \mathcal{O}_{\mathbb{C}^n,o}/(I \cdot \mathcal{O}_{\mathbb{C}^n,o})$ such that $\nu(f_j) = a_j$ for any $0 \leq j \leq m$. Remark 2.19 tells $\mathbb{C}[z_1, \dots, z_n]/I$ is a subring of $\mathcal{O}_{X,o}$. Thus ν is also a valuation on $\mathbb{C}[z_1, \dots, z_n]/I$.

Conversely, if there exists a valuation ν on $\mathbb{C}[z_1, \dots, z_n]/I$ satisfying that $\nu(f_j) = a_j$ for any $0 \leq j \leq m$ and $\nu(z_l) > 0$ for any $1 \leq l \leq n$, by Lemma 2.21 and the definition of $\sigma(\log |F|, \varphi)$, we have $\sigma(\log |F|, \varphi) = \sum_{1 \leq j \leq m} a_j$. \square

Proof of Corollary 1.13. The sufficient part follows from Corollary 1.12, then it suffices to prove the necessary part. If there exists a valuation ν on $\mathbb{C}[z_1, \dots, z_n]/I$ satisfying that $\nu(f_0) = 0$ and $\nu(f_j) = a_j > 0$ for any $1 \leq j \leq m$. Let J be the ideal in $\mathbb{C}[z_1, \dots, z_n]/I$ generated by $\{f_j\}_{1 \leq j \leq m}$. As $\cap_{1 \leq j \leq m} \{f_j = 0\} = \{o\}$, it follows from Hilbert's Nullstellensatz (see Lemma 2.23) that $z_k^N \in J$ for large N and any $1 \leq k \leq n$, which implies $\nu(z_k) > 0$ for any k . Thus, we have $\sigma(\log |F|, \varphi) = \sum_{1 \leq j \leq m} a_j$ by Corollary 1.12. \square

Now we prove Corollary 1.15.

Proof of Corollary 1.15. If $\sigma(\log |F|, \varphi) = \sum_{1 \leq j \leq m} a_j$, Theorem 1.9 shows that there exists a valuation $\tilde{\nu}$ on $\mathcal{O}_{X,o} = \mathcal{O}_{\mathbb{C}^n,o}/(\tilde{P}(I))$ such that $\tilde{\nu}(\tilde{P}(f_j)) = a_j$ for any $0 \leq j \leq m$. Remark 2.25 tells $\tilde{P} : C_{o'}^{\text{an}}/I \rightarrow \mathcal{O}_{\mathbb{C}^n,o}/(\tilde{P}(I))$ is an injective ring homomorphism. Then there exists a valuation ν on $C_{o'}^{\text{an}}/I$ such that $\nu(f_j) = a_j$ for any j .

Conversely, if there exists a valuation ν on $C_{o'}^{\text{an}}/I$ such that $\nu(f_j) = a_j$ for any j , by Lemma 2.27 and the definition of $\sigma(\log |F|, \varphi)$, we have $\sigma(\log |F|, \varphi) = \sum_{1 \leq j \leq m} a_j$. \square

Now we prove Corollary 1.17.

Proof of Corollary 1.17. If $\sigma(\log |F|, \varphi) = \sum_{1 \leq j \leq m} a_j$, Corollary 1.12 shows that there exists a valuation $\tilde{\nu}$ on $\mathbb{C}[z_1, \dots, z_n]/(P(I))$ such that $\tilde{\nu}(\tilde{P}(f_j)) = a_j$ for any $0 \leq j \leq m$. Remark 2.29 tells $\tilde{P} : \mathbb{R}[x_1, \dots, x_n]/I \rightarrow \mathbb{C}[x_1, \dots, x_n]/(P(I))$ is an injective ring homomorphism. Then there exists a valuation ν on $\mathbb{R}[x_1, \dots, x_n]/I$ such that $\nu(f_j) = a_j$ for any j .

Conversely, if there exists a valuation ν on $\mathbb{R}[x_1, \dots, x_n]/I$ satisfying that $\nu(f_j) = a_j$ for any $1 \leq j \leq m$ and $\nu(x_l) > 0$ for any $1 \leq l \leq n$, by Lemma 2.30 and the definition of $\sigma(\log |F|, \varphi)$, we have $\sigma(\log |F|, \varphi) = \sum_{1 \leq j \leq m} a_j$. \square

Acknowledgements. The first-named author completed part of this work during a visit to the School of Mathematical Sciences at Peking University and would like to thank the School for its hospitality and support. The second author was supported by National Key R&D Program of China 2021YFA1003100 and NSFC-12425101. The third author was supported by NSFC-12401099 and the Talent Fund of Beijing Jiaotong University 2024-004. The fourth author was supported by NSFC-12501106.

APPENDIX A. THE UNIVERSAL DENOMINATORS FOR ALGEBRAIC VARIETIES

It is known (see [10]; see also Theorem A.11) that for any complex analytic subvariety X and $x \in X$, there exists locally a holomorphic function $\delta_{X,x}$ such that $\delta_{X,x}\mathcal{O}_{X,x}^w \subset \mathcal{O}_{X,x}$, where $\mathcal{O}_{X,x}^w$ is the ring of germs of weakly holomorphic functions. Such $\delta_{X,x}$ is called universal denominator of weakly holomorphic functions.

Now we turn to the algebraic case. Let I be a prime ideal in $\mathbb{C}[z_1, \dots, z_n]$. Let $X := V(I)$ be the affine variety defined by I and $o \in X$ where o is the origin in \mathbb{C}^n . Denote the germ of the set X at o by (X, o) and assume that (X, o) is irreducible as a germ of analytic set. In this appendix, we show that when (X, o) is irreducible as a germ of analytic set, one can choose the universal denominator to be a polynomial, see Theorem A.12.

We firstly recall some basic results.

Lemma A.1 (Noether's normalization lemma, see [1]). *Let k be a field and let $A \neq 0$ be a finitely generated k -algebra which is generated by x_1, \dots, x_n . Then there exist elements $y_1, \dots, y_r \in A$ which are algebraically independent over k and such that A is integral over $k[y_1, \dots, y_r]$. Moreover, one can choose y_1, \dots, y_r to be linear combinations of x_1, \dots, x_n .*

Proof. For the convenience of the readers, we recall the proof of Lemma A.1 which can be referred to Section 5 of [1].

Since $k = \mathbb{C}$ in our case, we may assume that k is infinite. As x_1, x_2, \dots, x_n generate A as a k -algebra. We can renumber the $\{x_i\}_{i=1}^n$ such that $\{x_1, \dots, x_r\}$ is a maximal algebraically independent system of $\{x_i\}_{i=1}^n$ over k . If $r = n$, there is nothing to prove. We may assume $r < n$. Now we prove Lemma A.1 by induction on n . Assume that Lemma A.1 holds when the finitely generated k -algebra has $n-1$ generators.

As $r < n$, then x_n is algebraic over $k[x_1, \dots, x_{n-1}]$, i.e., there exists a polynomial $f \in k[T_1, \dots, T_n]$ such that $f(x_1, x_2, \dots, x_n) = 0$ in A . Decompose f as a sum of homogeneous polynomials and let F be the homogeneous part of the highest degree in the decomposition of f . Denote $\deg F = m$. As k is infinite, we can find $(\lambda_1, \lambda_2, \dots, \lambda_{n-1}, 1) \in k^n$ such that $F(\lambda_1, \lambda_2, \dots, \lambda_{n-1}, 1) \neq 0$ in k . Let $w_i = x_i - \lambda_i x_n$ for $i = 1, \dots, n-1$. Then denote

$$\tilde{f}(T) = f(w_1 + \lambda_1 T, w_2 + \lambda_2 T, \dots, w_{n-1} + \lambda_{n-1} T, T) \in k[w_1, \dots, w_{n-1}][T].$$

We know that $\deg \tilde{f} = m$ and $\tilde{f}(x_n) = 0$ in A . The coefficient of T^m in \tilde{f} is just $F(\lambda_1, \lambda_2, \dots, \lambda_{n-1}, 1) \neq 0$. Then $\frac{1}{F(\lambda_1, \lambda_2, \dots, \lambda_{n-1}, 1)}\tilde{f}$ is a unitary polynomial that annihilates x_n . Hence x_n is integral over $A' = k[w_1, \dots, w_{n-1}]$ which implies A is integral over $A' = k[w_1, \dots, w_{n-1}]$. Note that w_1, \dots, w_{n-1} can be written as linear combinations of x_1, \dots, x_n . Now applying the induction hypothesis to

$A' = k[w_1, \dots, w_{n-1}]$, we know that there exist $y_1, \dots, y_r \in A'$ which are algebraically independent over k and such that A' is integral over $k[y_1, \dots, y_r]$. Moreover, y_1, \dots, y_r can be chosen to be linear combinations of w_1, \dots, w_{n-1} . By the transitivity of integral dependence (see [1, Corollary 5.4]), we get that A is integral over $k[y_1, \dots, y_r]$ and y_1, \dots, y_r can be chosen to be linear combinations of x_1, \dots, x_n . \square

Lemma A.2 (Gauss lemma). *Let R be a unique factorization domain and F its field of fractions. Any $f \neq 0$ in $R[x]$ is irreducible in $R[x]$ if and only if it is both irreducible in $F[x]$ and primitive in $R[x]$.*

Using Noether's normalization lemma A.1, after linear change of coordinate, we may assume that there exists an integer $d > 0$ such that $A := \mathbb{C}[z_1, \dots, z_n]/I$ is a finite integral extension of $\mathbb{C}[z_1, \dots, z_d]$.

Recall that $X = V(I)$ and $o \in X$. Denote by \tilde{f} the class of any $f \in \mathbb{C}[z_1, \dots, z_n]$ in $A = \mathbb{C}[z_1, \dots, z_n]/I$. For any $d+1 \leq k \leq n$, there exists a unitary polynomial $Q_k(z', T) \in \mathbb{C}[z_1, \dots, z_d][T]$ such that $Q_k(z', \tilde{z}_k) = 0$ in A which means $Q_k(z', z_k) \in I$ as an element of $\mathbb{C}[z_1, \dots, z_d][z_k] \subset \mathbb{C}[z_1, \dots, z_n]$. In a conclusion, we have

Proposition A.3. *There exist an integer d , a coordinates (z_1, \dots, z_n) with the following properties: $\mathcal{I}_d := I \cap \mathbb{C}[z_1, \dots, z_d] = \{0\}$ and for every integer $k = d+1, \dots, n$ there is a Weierstrass polynomial $Q_k \in \mathcal{I}_k := I \cap \mathbb{C}[z_1, \dots, z_d][z_k]$ of the form*

$$Q_k(z', z_k) = z_k^{s_k} + \sum_{1 \leq j \leq s_k} a_{k,j}(z') z_k^{s_k-j}, \quad a_{j,k}(z') \in \mathbb{C}[z_1, \dots, z_d].$$

Let $\pi: \mathbb{C}^n \rightarrow \mathbb{C}^d$ be the mapping which sends (z_1, \dots, z_n) to (z_1, \dots, z_d) .

Lemma A.4. *Let $\pi_X: X \rightarrow \mathbb{C}^d$ be the restriction of π to X . Then π_X is proper and surjective.*

Proof. Note that π_X is continuous. To prove that π_X is proper, it suffices to prove that, for any $K \subset \mathbb{C}^d$ a bounded subset, $\pi_X^{-1}(K)$ is bounded in \mathbb{C}^n . Note that $X \subset \cap_{k=d+1}^n \{Q_k = 0\}$ and $Q_k \in \mathbb{C}[z_1, \dots, z_d][z_k]$. If $z' \in \mathbb{C}^d$ is bounded, we know that the roots of each Q_k are bounded. Hence $\pi_X^{-1}(K)$ is bounded in \mathbb{C}^n .

To prove that π_X is surjective, without loss of generality, it suffices to prove that $o' \in \mathbb{C}^d$ belongs to $\pi_X(X)$. Denote by $m' = (z_1, \dots, z_d)$ the maximal ideal in $\mathbb{C}_d := \mathbb{C}[z_1, \dots, z_d]$. It follows from A is a finite integral extension of \mathbb{C}_d and [1, Corollary 5.2] that \mathbb{C}_d is a subring of A and A is a finitely-generated \mathbb{C}_d -module.

We claim that $A/m'A \neq 0$, where $m'A$ is the ideal generated by m' in A . If $A/m'A = 0$, then $m'A = A$. It follows from A is a finitely-generated \mathbb{C}_d -module and Nakayama's Lemma (cf. [1, Corollary 2.5]) that there exists an $r \in \mathbb{C}_d$ such that $r \equiv 1 \pmod{m'}$ in \mathbb{C}_d and $rA = 0$. View r as a function on \mathbb{C}^d , then $r \equiv 1 \pmod{m'}$ in \mathbb{C}_d implies that $r(o') = 1$ and hence $r \neq 0$ in \mathbb{C}_d . $rA = 0$ implies $r \cdot 1 = 0$ in A and thus $r \in I$. Then $r \in I \cap \mathbb{C}_d = \{0\}$. This contradicts to $r \neq 0$ in \mathbb{C}_d . Hence we must have $A/m'A \neq 0$.

Note that $A/m'A \cong \mathbb{C}[z_1, \dots, z_n]/(I + m'\mathbb{C}[z_1, \dots, z_n])$. It follows from $A/m'A \neq 0$ that $\mathbb{C}[z_1, \dots, z_n]/(I + m'\mathbb{C}[z_1, \dots, z_n]) \neq 0$ and hence $I + m'\mathbb{C}[z_1, \dots, z_n] \neq (1)$. Thus $V(I) \cap V(m'\mathbb{C}[z_1, \dots, z_n]) = V(I + m'\mathbb{C}[z_1, \dots, z_n])$ is not empty. This implies that $o' \in \pi_X(X)$. \square

Denote \mathcal{O}_n and \mathcal{O}_d be the germ of holomorphic functions at the origin of \mathbb{C}^n and \mathbb{C}^d respectively. Using Lemma A.4, we have

Lemma A.5. $(I \cdot \mathcal{O}_n) \cap \mathcal{O}_d = 0$.

Proof. We prove Lemma A.5 by contradiction. Assume there exists a nonzero holomorphic function $f \in (I \cdot \mathcal{O}_n) \cap \mathcal{O}_d$. It follows from $f \in (I \cdot \mathcal{O}_n) \cap \mathcal{O}_d$ that $\pi_X(X) \subset \{f = 0\}$ is a subset in \mathbb{C}^d . However Lemma A.4 tells that π_X is surjective, which is a contradiction. \square

Denote by \tilde{f} the class of any $f \in \mathbb{C}[z_1, \dots, z_n]$ in $A = \mathbb{C}[z_1, \dots, z_n]/I$. Denote by \mathfrak{R}_A and \mathfrak{R}_d the quotient fields of A and $\mathbb{C}[z_1, \dots, z_d]$ respectively. Then $\mathfrak{R}_A = \mathfrak{R}_d[\tilde{z}_{d+1}, \dots, \tilde{z}_n]$ is a finite algebraic extension of \mathfrak{R}_d . Primitive element theorem tells that there exist $c_i \in \mathbb{C}$ ($i = d+1, \dots, n$) such that $\tilde{u} = \sum_{i=d+1}^n c_i \tilde{z}_i$ and $\mathfrak{R}_A = \mathfrak{R}_d[\tilde{u}]$. Hence \tilde{u} belongs to A .

Lemma A.6. *Let R be a unique factorization domain and F its field of fractions. Assume that $R \subset A$ and A is a ring. For any $f \in A$ integral over $R[T]$, denote the minimal polynomial of \tilde{f} over R by $W_f(z', T)$. Then $W_f(z', T)$ is also the minimal polynomial of \tilde{f} over F .*

Proof. As R is UFD and $f \in A$ is integral over $R[T]$, we know that $R[T]$ is UFD and there exists a unitary irreducible polynomial $W_f(T) \in R[T]$ such that $W_f(f) = 0$. Denote by $M_f(T) \in F[T]$ the minimal polynomial of F over F . It follows from Gauss Lemma A.2 that $W_f(z', T)$ is also irreducible in $F[T]$. Then by the definition of minimal polynomial and that $W_f(T)$ is irreducible in $F[T]$, we have $M_f(T) \mid W_f(T)$ and hence $M_f(T) = W_f(T) \in R[T]$. \square

Remark A.7. *In this appendix, we will use Lemma A.6 in the following cases*

- (1) $R = \mathbb{C}[z_1, \dots, z_d]$, $A = \mathbb{C}[z_1, \dots, z_n]/I$ and $F = \mathfrak{R}_d$;
- (2) $R = \mathcal{O}_d$, $A = \mathcal{O}_n/(I \cdot \mathcal{O}_n)$ and $F = \mathfrak{M}_d$, where \mathfrak{M}_d is the quotient fields of $R = \mathcal{O}_d$.

Denote the minimal polynomial of \tilde{u} over \mathfrak{R}_d by $W_u(z', T)$. It follows from Lemma A.6 and Remark A.7 that the discriminant $\delta(z_1, \dots, z_d)$ of $W_u(z', T)$ belongs to $\mathbb{C}[z_1, \dots, z_d]$. Hence we conclude that

Lemma A.8. *There exists a element $\tilde{u} \in A$ such that $\mathfrak{R}_A = \mathfrak{R}_d[\tilde{u}]$. Denote the minimal polynomial of \tilde{u} over \mathfrak{R}_d by $W_u(z', T)$. Then $W_u(z', T) \in \mathbb{C}[z_1, \dots, z_d][T]$ and the discriminant $\delta(z_1, \dots, z_d)$ of $W_u(z', T)$ belongs to $\mathbb{C}[z_1, \dots, z_d]$.*

Let $q := [\mathfrak{R}_A : \mathfrak{R}_d]$ be the degree of the filed extension. As $\mathfrak{R}_A = \mathfrak{R}_d[\tilde{u}]$, we know that $W_u(z', T)$ is of degree q .

Denote by \mathfrak{M}_A the quotient fields of $A = \mathcal{O}_n/(I \cdot \mathcal{O}_n)$ and by \mathfrak{M}_d the the quotient fields of $R = \mathcal{O}_d$. It follows from Proposition A.3, Lemma A.5, Proposition 4.13 (and its proof) in [10, p. 93] and the arguments in [10, pp. 93–94] that we know

- (1) \mathcal{O}_n/I is a finite integral extension of \mathcal{O}_d ;
- (2) There exists a linear form $u_1 = \sum_{d+1 \leq k \leq n} c'_k z_k$ where $c'_k \in \mathbb{C}$ such that $\mathfrak{M}_A = \mathfrak{M}_d[u_1]$.

Recall that \mathfrak{R}_A and \mathfrak{R}_d are the rational function fields of X and \mathbb{C}^d respectively, which can be naturally viewed as a subset of \mathfrak{M}_A and \mathfrak{M}_d respectively. Note that $\mathfrak{R}_A = \mathfrak{R}_d[\tilde{u}]$.

Remark A.9. *We have $\mathfrak{M}_A = \mathfrak{M}_d[\tilde{u}]$.*

Proof. Since each \tilde{z}_k belongs to \mathfrak{R}_A and $\mathfrak{R}_A = \mathfrak{R}_d[\tilde{u}]$, we know that, for $d+1 \leq k \leq n$, \tilde{z}_k can be represented by the linear combination of \tilde{u}^j ($0 \leq j \leq q-1$) with coefficients in \mathfrak{R}_d . As $u_1 = \sum_{d+1 \leq k \leq n} c'_k z_k$ where $c'_k \in \mathbb{C}$, we know that u_1 can be also represented by the linear combination of \tilde{u}^j ($0 \leq j \leq q-1$) with coefficients in $\mathfrak{R}_d \subset \mathfrak{M}_d$. Thus we have $\mathfrak{M}_A = \mathfrak{M}_d[\tilde{u}]$. \square

Denote by $H_u(z', T) \in \mathcal{O}_d[T]$ the minimal polynomial of $\tilde{u} \in \mathcal{O}_n/I$ over \mathcal{O}_d . It follows from Lemma A.6 and Remark A.7 that we know $H_u(z', T)$ is also the minimal polynomial of \tilde{u} over \mathfrak{M}_d . Recall that $W_u(z', T) \in \mathbb{C}[z_1, \dots, z_d][T] \subset \mathcal{O}_d[T]$ is the minimal polynomial of \tilde{u} over $\mathbb{C}[z_1, \dots, z_d]$ and $H_u(z', T) \in \mathcal{O}_d[T]$ is the minimal polynomial of \tilde{u} over \mathcal{O}_d . Denote the discriminants of $W_u(z', T)$ and $H_u(z', T)$ by $\delta_{\mathfrak{R}}(z') \in \mathbb{C}[z_1, \dots, z_d] \subset \mathcal{O}_d$ and $\delta_{\mathfrak{M}}(z') \in \mathcal{O}_d$ respectively. We have the following relation between $\delta_{\mathfrak{R}}(z')$ and $\delta_{\mathfrak{M}}(z')$.

Lemma A.10. $\delta_{\mathfrak{M}}(z') \mid \delta_{\mathfrak{R}}(z')$ in \mathcal{O}_d .

Proof. As $H_u(z', T) \in \mathcal{O}_d[T]$ is the minimal polynomial of $\tilde{u} \in \mathcal{O}_n/I$ over \mathcal{O}_d and $W_u(z', T) \in \mathcal{O}_d[T]$ annihilates \tilde{u} , we have $H_u(z', T) \mid W_u(z', T)$ in $\mathcal{O}_d[T]$, i.e., there exists a polynomial $P_u(z', T) \in \mathcal{O}_d[T]$ such that $W_u(z', T) = H_u(z', T)P_u(z', T)$. Denote by $R_{H,P}$ the resultant of the polynomials H and P , and denote by $\delta_P(z')$ the discriminant of $P_u(z', T)$.

Recall that the resultant of two polynomials is the determinant of the Sylvester matrix of the two polynomials. Then it follows from $H_u(z', T), P_u(z', T) \in \mathcal{O}_d[T]$ that $R_{H,P}$ and $\delta_P(z')$ belong to \mathcal{O}_d . By the definition of discriminant, we know that $\delta_{\mathfrak{R}}(z') = \delta_{\mathfrak{M}}(z')\delta_P(z')R_{H,P}^2$. Thus, $\delta_{\mathfrak{M}}(z') \mid \delta_{\mathfrak{R}}(z')$ in \mathcal{O}_d . \square

Now we recall the following existence of “universal denominators” of weakly holomorphic functions on an irreducible complex variety X .

Theorem A.11 (see [10]). *For every point $x \in X$, there is a neighborhood V of x in X , such that $\delta_{\mathfrak{M},y}\mathcal{O}_{X,y}^w \subset \mathcal{O}_{X,y}$ for all $y \in V$, where $\delta_{\mathfrak{M}}(z')$ is the discriminant of the minimal polynomial $H_u(z', T)$ of \tilde{u} over \mathcal{O}_d and $\delta_{\mathfrak{M}}(z')$ is called a universal denominator on V .*

Using Lemma A.10 and Theorem A.11, we can show that, the universal denominator for \mathcal{O}_X^w can be chosen to be a polynomial.

Theorem A.12. *Assume that X is irreducible and algebraic. Let V be a neighborhood of o in X and $y \in V$. Then for any $f \in \mathcal{O}_{X,y}^w$, one has $\delta_{\mathfrak{R},y}(z')f \in \mathcal{O}_{X,y}$, where $\delta_{\mathfrak{R}}(z') \in \mathbb{C}[z_1, \dots, z_d]$ is the discriminant of the minimal polynomial $W_u(z', T)$ of \tilde{u} over $\mathbb{C}[z_1, \dots, z_d]$.*

Proof. It follows from Theorem A.11 that, for any $f \in \mathcal{O}_{X,y}^w$, one has $\delta_{\mathfrak{M},y}f \subset \mathcal{O}_{X,y}$. Lemma A.10 tells $\delta_{\mathfrak{R}}(z') = \delta_{\mathfrak{M}}(z')\delta_P(z')R_{H,P}$, where $\delta_P(z'), R_{H,P} \in \mathcal{O}_d$. Thus we know $\delta_{\mathfrak{R},y}(z')f = \delta_P(z')R_{H,P}\delta_{\mathfrak{M}}(z')f \in \mathcal{O}_{X,y}$.

Theorem A.12 has been proved. \square

REFERENCES

- [1] M.F. Atiyah and I.G. Macdonald, Introduction to Commutative Algebra, Addison-Wesley Publishing Co., Reading (1969).
- [2] S.J. Bao, Q.A. Guan, Z.T. Mi and Z. Yuan, Tame maximal weights, relative types and valuations, Adv. Math. 477 (2025), Paper No. 110364.

- [3] S.J. Bao, Q.A. Guan, Z.T. Mi and Z. Yuan, The existence of valuative interpolation, preprint.
- [4] E. Bedford and B.A. Taylor, The Dirichlet problem for a complex Monge-Ampère operator, *Invent. Math.* 37 (1976) 1–44.
- [5] E. Bierstone and P.D. Milman, A simple constructive proof of canonical resolution of singularities, pp. 11–30 in *Effective methods in algebraic geometry* (Castiglioncello, 1990), edited by T. Mora and C. Traverso, *Progr. Math.* 94, Birkhäuser, Boston, 1991.
- [6] Z. Blocki, Estimates for the complex Monge-Ampère operator, *Bull. Polish Acad. Sci. Math.* 41 (1993), no. 2, 151–157 (1994).
- [7] Z. Blocki, The complex Monge-Ampère operator in pluripotential theory, electronically accessible at <http://gamma.im.uj.edu.pl/~blocki/publ/ln/index.html>.
- [8] S. Boucksom, C. Favre and M. Jonsson, Valuations and plurisubharmonic singularities, *Publ. Res. Inst. Math. Sci.* 44 (2008), no. 2, 449–494.
- [9] J.-P. Demailly, *Analytic Methods in Algebraic Geometry*, Higher Education Press, Beijing, 2010.
- [10] J.-P. Demailly, Complex analytic and differential geometry, electronically accessible at <https://www-fourier.ujf-grenoble.fr/~demailly/manuscripts/agbook.pdf>.
- [11] C. Favre and M. Jonsson, The valuative tree, *Lecture Notes in Mathematics*, 1853, Springer, Berlin, 2004.
- [12] C. Favre and M. Jonsson, Valuations and multiplier ideals, *J. Amer. Math. Soc.* 18 (2005), no. 3, 655–684.
- [13] C. Favre and M. Jonsson, Valuative analysis of planar plurisubharmonic functions, *Invent. Math.* 162 (2005), no. 2, 271–311.
- [14] P. Griffiths and J. Harris, *Principles of Algebraic Geometry*, John Wiley, Ltd, 2011.
- [15] Q.A. Guan, A sharp effectiveness result of Demailly’s strong openness conjecture, *Adv. Math.* 348 (2019), 51–80.
- [16] Q.A. Guan, Decreasing equisingular approximations with analytic singularities, *J. Geom. Anal.* 30 (2020), no. 1, 484–492.
- [17] Q.A. Guan and X.Y. Zhou, A proof of Demailly’s strong openness conjecture, *Ann. of Math.* (2) 182 (2015), no. 2, 605–616. See also arXiv:1311.3781.
- [18] Q.A. Guan and X.Y. Zhou, Effectiveness of Demailly’s strong openness conjecture and related problems, *Invent. Math.* 202 (2015), no. 2, 635–676.
- [19] P.H. Hiep, The weighted log canonical threshold, *C. R. Math. Acad. Sci. Paris* 352 (2014), no. 4, 283–288.
- [20] H. Hironaka, Resolution of singularities of an algebraic variety over a field of characteristic zero. I, II, *Ann. Math.* 79 (1964), 109–203; ibid. 205–326.
- [21] M. Jonsson and M. Mustaţă, Valuations and asymptotic invariants for sequences of ideals, *Ann. Inst. Fourier (Grenoble)* 62 (2012), no. 6, 2145–2209.
- [22] M. Jonsson and M. Mustaţă, An algebraic approach to the openness conjecture of Demailly and Kollar, *J. Inst. Math. Jussieu* 13 (2014), no. 1, 119–144.
- [23] C.O. Kiselman, Plurisubharmonic functions and potential theory in several complex variables, in: *Development of Mathematics*, Birkhäuser, Basel, 1950–2000, pp. 655–714.
- [24] L. Lempert, Modules of square integrable holomorphic germs. Analysis meets geometry, 311–333, *Trends Math.*, Birkhäuser/Springer, Cham, 2017.
- [25] P. Lelong, Intégration sur un ensemble analytique complexe (in French) *Bull. Soc. Math. Fr.* 85 (1957) 239–262.
- [26] H. Matsumura, *Commutative Ring Theory*. Reid M, trans. Cambridge University Press; 1987.
- [27] A. Rashkovskii, Relative types and extremal problems for plurisubharmonic functions, *Int. Math. Res. Not.* 2006, Art. ID 76283, 26 pp.
- [28] H. Skoda, Sous-ensembles analytiques d’ordre fini ou infini dans \mathbb{C}^n , (French) *Bull. Soc. Math. France* 100 (1972), 353–408.
- [29] G. Tian, On Kähler-Einstein metrics on certain Kähler manifolds with $C_1(M) > 0$, *Invent. Math.* 89 (1987), no. 2, 225–246.
- [30] G. Tian, On Calabi’s conjecture for complex surfaces with positive first Chern class, *Invent. Math.* 101 (1990), no. 1, 101–172.
- [31] H. Whitney, *Complex Analytic Varieties*, Addison–Wesley Publ. Co., 1972.
- [32] O. Zariski, Analytical irreducibility of normal varieties, *Ann. of Math.* (2) 49 (1948), 352–361.

SHIJIE BAO: ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, CHINA.

Email address: bsjie@amss.ac.cn

QI'AN GUAN: SCHOOL OF MATHEMATICAL SCIENCES, PEKING UNIVERSITY, BEIJING, 100871, CHINA.

Email address: guanqian@math.pku.edu.cn

ZHITONG MI: SCHOOL OF MATHEMATICS AND STATISTICS, BEIJING JIAOTONG UNIVERSITY, BEIJING, 100044, CHINA.

Email address: zhitongmi@amss.ac.cn

ZHENG YUAN: STATE KEY LABORATORY OF MATHEMATICAL SCIENCES, ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, CHINA.

Email address: yuanzheng@amss.ac.cn